



US009899610B2

(12) **United States Patent**
Lee et al.

(10) **Patent No.:** **US 9,899,610 B2**
(45) **Date of Patent:** **Feb. 20, 2018**

(54) **COMPOUND AND ORGANIC
LIGHT-EMITTING DEVICE**

- (71) Applicant: **SAMSUNG DISPLAY CO., LTD.**,
Yongin-si, Gyeonggi-do (KR)
- (72) Inventors: **Jungsub Lee**, Yongin-si (KR); **Seulong Kim**, Yongin-si (KR); **Younsun Kim**, Yongin-si (KR); **Dongwoo Shin**, Yongin-si (KR); **Naoyuki Ito**, Yongin-si (KR); **Jino Lim**, Yongin-si (KR)
- (73) Assignee: **Samsung Display Co., Ltd.**, Yongin-si (KR)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 81 days.

(21) Appl. No.: **15/174,918**

(22) Filed: **Jun. 6, 2016**

(65) **Prior Publication Data**

US 2017/0155056 A1 Jun. 1, 2017

(30) **Foreign Application Priority Data**

Nov. 26, 2015 (KR) 10-2015-0166413

(51) **Int. Cl.**

H01L 51/00 (2006.01)
C07D 401/04 (2006.01)
C07F 9/572 (2006.01)
C07D 403/10 (2006.01)
C07D 403/04 (2006.01)
H01L 27/32 (2006.01)
H01L 51/50 (2006.01)
H01L 51/56 (2006.01)

(52) **U.S. Cl.**

CPC **H01L 51/0072** (2013.01); **C07D 401/04** (2013.01); **C07D 403/04** (2013.01); **C07D 403/10** (2013.01); **C07F 9/5728** (2013.01); **H01L 27/3248** (2013.01); **H01L 51/0058** (2013.01); **H01L 51/0067** (2013.01); **H01L 51/506** (2013.01); **H01L 51/5072** (2013.01); **H01L 51/56** (2013.01)

(58) **Field of Classification Search**

CPC C07D 401/04; H01L 51/0058; H01L 51/0067
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2012/0292603 A1 * 11/2012 Kwak H01L 51/0059
257/40
2012/0305903 A1 * 12/2012 Kai C07D 487/04
257/40
2013/0126856 A1 5/2013 Yokoyama et al.
2013/0256634 A1 * 10/2013 Cho H01L 51/0067
257/40
2017/0110669 A1 * 4/2017 Im H01L 51/0071

FOREIGN PATENT DOCUMENTS

KR 10-2012-0101620 A 9/2012
KR 10-2013-0093106 A 8/2013
KR 10-2013-0129543 A 11/2013
KR 10-2014-0128891 A 11/2014
KR 10-2014-0128892 A 11/2014

* cited by examiner


Primary Examiner — Melissa A Rioja

(74) *Attorney, Agent, or Firm* — Lewis Roca Rothgerber Christie LLP

(57)

ABSTRACT

An organic light-emitting device including: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes a compound represented by Formula 1:



Formula 1

When the compound of Formula 1 is used as an electron transport material, an organic light-emitting device including the compound may have high efficiency, low driving voltage, high brightness, and long lifespan.

20 Claims, 2 Drawing Sheets

FIG. 1




FIG. 2

20

190
150
110
210

FIG. 3

30

220
190
150
110

FIG. 4

40

220
190
150
110
210

COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE

CROSS-REFERENCE TO RELATED APPLICATION

This application claims priority to and the benefit of Korean Patent Application No. 10-2015-0166413, filed on Nov. 26, 2015, in the Korean Intellectual Property Office, the entire content of which is incorporated herein by reference.

BACKGROUND

Field

One or more aspects of example embodiments of the present disclosure are related to a compound and an organic light-emitting device including the same.

Organic light-emitting devices are self-emission devices that have wide viewing angles, high contrast ratios, short response times, and/or excellent brightness, driving voltage, and/or response speed characteristics, and may produce full-color images.


An organic light-emitting device may include a first electrode on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode sequentially positioned on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers (such as holes and electrons) may recombine in the emission layer to produce excitons. These excitons may transition (e.g., radiatively decay) from an excited state to a ground state to thereby generate light.

SUMMARY

One or more aspects of example embodiments of the present disclosure are directed toward a compound suitable for use as an electron injection and/or transport material, due to its excellent electron injection and/or transport capability and material stability, and an organic light-emitting device including the compound that accordingly has high efficiency, low driving voltage, high brightness, and long lifespan.

Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments.

One or more example embodiments of the present disclosure provide a compound represented by Formula 1:




In Formula 1, R₂ and R₃ may each independently be selected from hydrogen, deuterium, a halogen atom, a nitro group, a cyano

group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,


Ar₁ to Ar₃ may each independently be selected from hydrogen, deuterium, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, a substituted or unsubstituted C₅-C₆₀ carbocyclic group, and a substituted or unsubstituted C₁-C₆₀ heterocyclic group,

a to c may each independently be an integer selected from 0 to 2, and

R₁ may be one selected from Formulae 1-1 and 1-2:



Formula 1-1



Formula 1-2

In Formulae 1-1 and 1-2,

X₁ to X₅ may each independently be selected from nitrogen (N) and CR₁₁,

at least one selected from X₁ to X₅ may be N;

R₁₁, Y₁, and Y₂ may each independently be selected from a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

L₁ and L₂ may each independently be selected from a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

* may indicate a binding site, and

at least one substituent selected from the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₂-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀

arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted C_5 - C_{60} carbocyclic group, the substituted C_1 - C_{60} heterocyclic group, the substituted C_6 - C_{60} arylene group, the substituted C_1 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and the substituted divalent non-aromatic condensed heteropolycyclic group may be selected from the group consisting of:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q_{11})(Q_{12}), —Si(Q_{13})(Q_{14})(Q_{15}), and —B(Q_{16})(Q_{17});

a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q_{21})(Q_{22}), —Si(Q_{23})(Q_{24})(Q_{25}), and —B(Q_{26})(Q_{27}),

wherein Q_{11} to Q_{17} and Q_{21} to Q_{27} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60}

alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

One or more aspects of embodiments of the present disclosure provide an organic light-emitting device including: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes the compound.

One or more aspects of embodiments of the present disclosure provide a display apparatus including the organic light-emitting device, wherein the first electrode of the organic light-emitting device is electrically connected to a source electrode or drain electrode of a thin film transistor.

BRIEF DESCRIPTION OF THE DRAWINGS

These and/or other aspects will become apparent and more readily appreciated from the following description of the example embodiments, taken in conjunction with the accompanying drawings in which:

FIG. 1 is a schematic diagram of a structure of an organic light-emitting device according to an embodiment of the present disclosure;

FIG. 2 is a schematic diagram of a structure of an organic light-emitting device according to another embodiment of the present disclosure;

FIG. 3 is a schematic diagram of a structure of an organic light-emitting device according to another embodiment of the present disclosure; and

FIG. 4 is a schematic diagram of a structure of an organic light-emitting device according to another embodiment of the present disclosure.


DETAILED DESCRIPTION

Reference will now be made in more detail to embodiments, examples of which are illustrated in the accompanying drawings, wherein like reference numerals refer to like elements throughout and duplicative descriptions thereof may not be provided. In this regard, the present embodiments may have different forms and should not be construed as being limited to the descriptions set forth herein. Accordingly, the embodiments are merely described below, by referring to the drawings, to explain aspects of the present description. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of”, “one of”, and “selected from”, when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

In the drawings, the thicknesses of layers, films, panels, regions, etc., may be exaggerated for clarity. It will be understood that when an element such as a layer, film, region, or substrate is referred to as being “on” another element, it can be directly on the other element or intervening element(s) may also be present. In contrast, when an element is referred to as being “directly on” another element, no intervening elements are present.

One or more aspects of embodiments of the present disclosure provide a compound represented by Formula 1:

5




In Formula 1,

R₂ and R₃ may each independently be selected from hydrogen, deuterium, a halogen atom, a nitro group, a cyano group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,


Ar₁ to Ar₃ may each independently be selected from hydrogen, deuterium, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, a substituted or unsubstituted C₅-C₆₀ carbocyclic group, and a substituted or unsubstituted C₁-C₆₀ heterocyclic group,

a to c may each independently be an integer selected from 0 to 2, and

R₁ may be one selected from Formulae 1-1 and 1-2:



Formula 1-1



Formula 1-2

In Formulae 1-1 and 1-2,

X₁ to X₅ may each independently be selected from nitrogen (N) and CR₁₁,

at least one selected from X₁ to X₅ may be N,

R₁₁, Y₁, and Y₂ may each independently be selected from a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or

6

unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

- 5 L₁ and L₂ may each independently be selected from a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

- 10 * may indicate a binding site, and

at least one substituent selected from the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₂-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted C₅-C₆₀ carbocyclic group, the substituted C₁-C₆₀ heterocyclic group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and the substituted divalent non-aromatic condensed heteropolycyclic group may be selected from the group consisting of:

- 30 deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

- a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

- a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and


- a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a

7


hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇),

wherein Q₁₁ to Q₁₇ and Q₂₁ to Q₂₇ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.


As used herein, in Formula 1, the dashed line in the phenylene moiety indicates that a double bond may be included on either side of the dashed portion. For example, a moiety shown as




may include one selected from



and



In one or more embodiments, the compound represented by Formula 1 may be represented by Formula 2:




Formula 2

8


In one or more embodiments, the compound represented by Formula 1 may be represented by Formula 3:

Formula 3



In one or more embodiments, the compound represented by Formula 1 may be represented by Formula 4:

Formula 4




In Formulae 2 to 4, R₁ to R₃, Ar₁ to Ar₃, and a to c may each independently be the same as described herein in connection with Formula 1.

In one or more embodiments, in Formula 1-1, one selected from X₁ and X₃ may be N, X₁ and X₅ may each independently be N, X₁ and X₃ may each independently be N, or X₁, X₃, and X₅ may each independently be N.

In one or more embodiments, the group represented by Formula 1-2 may be represented by Formula 1-3:

Formula 1-3





In Formula 1-3, L₂ may be the same as described herein in connection with Formula 1-2.

In one or more embodiments, R₂ and R₃ may each independently be a substituted or unsubstituted C₁-C₂₀ alkyl group. For example, R₂ and R₃ may each independently be a methyl group.

9

In one or more embodiments, R_{11} may be represented by one selected from Formulae 2a to 2b:



In Formulae 2a to 2b,



Z_1 may be selected from hydrogen, deuterium, a halogen group (e.g., atom), a cyano group, a nitro group, a hydroxyl group, a carboxyl group, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_6 - C_{20} aryl group, a substituted or unsubstituted C_1 - C_{20} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

p may be an integer selected from 1 to 7,

when p is 2 or greater, a plurality of Z_1 (s) may be identical to or different from each other, and


* may indicate a binding site.

In one or more embodiments, L_1 and L_2 may each independently be represented by one selected from Formulae 3a to 3b:



In Formulae 3a to 3b, * may indicate a binding site.

In one or more embodiments, the compound represented by Formula 1 may be one selected from Compounds ET1 to ET42, but embodiments of the present disclosure are not limited thereto:



ET-1

55

60

65

10


-continued

ET-2

2a 5

2b 10

15




ET-3

20

25

30




35

3a

ET-4

3b 40

45




ET-5


55

60






65



11
-continued




12
-continued



13

-continued

ET-12




5

10

15

20

ET-13




25

30

35

ET-14




40

45

50

ET-15



55


60

65


14

-continued


ET-16




ET-17



ET-18




ET-19



15

-continued


ET-20




16

-continued


ET-24




ET-25




ET-26




ET-27




ET-21



ET-22




ET-23



17

-continued

ET-28




5

10

15

20

ET-29




25

30

35

40

ET-30



50

55


60

65


18

-continued


ET-31




ET-32



ET-33




ET-34



19

-continued

ET-35




5

10

15

ET-36




20

25

30

ET-37




35

40

45

ET-38



50


55

60


20

-continued


ET-39




ET-40



ET-41



ET-42



Description of FIG. 1

FIG. 1 is a schematic cross-sectional view of a structure of an organic light-emitting device 10 according to an embodiment of the present disclosure. The organic light-emitting device 10 includes a first electrode 110, an organic layer 150, and a second electrode 190.

21

Hereinafter, the structure of the organic light-emitting device **10** according to an embodiment of the present disclosure and a method of manufacturing the organic light-emitting device **10** will be described in connection with FIG. 1.

First Electrode **110**

In FIG. 1, a substrate may be under the first electrode **110** and/or above the second electrode **190**. The substrate may be a glass substrate or a plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and/or water-resistance.

The first electrode **110** may be formed by depositing or sputtering a material for forming the first electrode **110** on the substrate. When the first electrode **110** is an anode, the material for forming the first electrode may be selected from materials with a high work function to facilitate hole injection.

The first electrode **110** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. The first electrode **110** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode **110** is a transmissive electrode, the material for forming a first electrode **110** may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), zinc oxide (ZnO), and combinations thereof, but embodiments of the present disclosure are not limited thereto. In one or more embodiments, when the first electrode **110** is a semi-transmissive electrode or a reflective electrode, the material for forming the first electrode **110** may be selected from magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and combinations thereof, but embodiments of the present disclosure are not limited thereto.

The first electrode **110** may have a single-layer structure, or a multi-layer structure including two or more layers. For example, the first electrode **110** may have a three-layered structure of ITO/Ag/ITO, but embodiments of the structure of the first electrode **110** are not limited thereto.

Organic Layer **150**

The organic layer **150** is on the first electrode **110**. The organic layer **150** may include an emission layer.

The organic layer **150** may further include a hole transport region between the first electrode **110** and the emission layer, and an electron transport region between the emission layer and the second electrode **190**.

Hole Transport Region in Organic Layer **150**

The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.


The hole transport region may include at least one layer selected from a hole injection layer (HIL), a hole transport layer (HTL), an emission auxiliary layer, and an electron blocking layer (EBL).

For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, a multi-layered structure having a structure of hole injection layer/hole transport layer, a


22

structure of hole injection layer/hole transport layer/emission auxiliary layer, a structure of hole injection layer/emission auxiliary layer, a structure of hole transport layer/emission auxiliary layer, or a structure of hole injection layer/hole transport layer/electron blocking layer, wherein layers of each structure are sequentially stacked on the first electrode **110** in each stated order, but embodiments of the structure of the hole transport region are not limited thereto.

The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (also referred to as NPD), β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl) triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:




m-MTDATA



TDATA

23


-continued




2-TNATA

24


-continued




Spiro-TPD





Spiro-NPB




NPB




methylated NPB

 β -NPB

TAPC




TPD




HMTPD

Formula 201



Formula 202



a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene

each independently be selected from a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a

27

dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

wherein Q₃₁ to Q₃₃ may each independently be the same as described above.

In one or more embodiments, in Formula 201, at least one selected from R₂₀₁ to R₂₀₃ may be selected from the group consisting of:

a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, in Formula 202, i) R₂₀₁ and R₂₀₂ may be connected (e.g., coupled) via a single bond, and/or ii) R₂₀₃ and R₂₀₄ may be connected (e.g., coupled) via a single bond.

28


In one or more embodiments, in Formula 202, at least one selected from R₂₀₁ to R₂₀₄ may be selected from the group consisting of:

a carbazolyl group; and

a carbazolyl group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, but embodiments of the present disclosure are not limited thereto.


The compound represented by Formula 201 may be represented by Formula 201A:

Formula 201A




For example, the compound of Formula 201 may be represented by Formula 201A(1):

Formula 201A(1)



For example, the compound of Formula 201 may be represented by Formula 201A-1:


Formula 201A-1



The compound of Formula 202 may be represented by Formula 202A:


29

Formula 202A



In one or more embodiments, the compound of Formula 202 may be represented by Formula 202A-1:

Formula 202A-1



In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,

L_{201} to L_{203} , $xa1$ to $xa3$, $xa5$, and R_{202} to R_{204} may each independently be the same as described above,


30

R_{211} and R_{212} may each independently be the same as described above in connection with R_{203} , and


R_{213} to R_{217} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-F$, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

The hole transport region may include at least one selected from Compounds HT1 to HT30, but embodiments of the present disclosure are not limited thereto:

HT1



HT2




31


32

-continued


HT3



HT4



HT5




33


34

-continued

HT6



HT7




35


36

-continued

HT8



HT9




37


38

-continued


HT10




HT11



HT12



HT13




39


40

-continued

HT14





HT15



HT16

HT17





41

42


-continued

HT18

HT19



HT20




43


44

-continued


HT21



HT22



HT23




45

46



-continued

HT24




HT25

HT26



HT27




47


48

-continued


HT28



HT29



HT30




49


50

-continued


HT31



HT32



HT33




51


52

-continued


HT34





HT35



HT36



53


-continued
HT37

The thickness of the hole transport region may be about 100 Å to about 10,000 Å, and in some embodiments, about 100 Å to about 1,000 Å. When the hole transport region includes at least one selected from a hole injection layer and a hole transport layer, the thickness of the hole injection layer may be about 100 Å to about 9,000 Å, and in some embodiments, about 100 Å to about 1,000 Å. The thickness of the hole transport layer may be about 50 Å to about 2,000 Å, and in some embodiments, about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are each within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer (e.g., adjusting the optical resonance distance to match the wavelength of light emitted from the emission layer). The electron blocking layer may block or reduce the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include those materials as described above.

54

HT38



HT39

p-Dopant




The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

The charge-generation material may be, for example, a p-dopant.

The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but embodiments of the present disclosure are not limited thereto.

For example, the p-dopant may include at least one selected from a quinone derivative (such as a tetracyanoquinodimethane (TCNQ) and/or 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ)); a metal oxide (such as a tungsten oxide and/or a molybdenum oxide); 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and a compound represented by Formula 221, but embodiments of the present disclosure are not limited thereto:

55



In Formula 221,

R_{221} to R_{223} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

wherein at least one selected from R_{221} to R_{223} may include at least one substituent selected from a cyano group, —F, —Cl, —Br, —I, a C_1 - C_{20} alkyl group substituted with —F, a C_1 - C_{20} alkyl group substituted with —Cl, a C_1 - C_{20} alkyl group substituted with —Br, and a C_1 - C_{20} alkyl group substituted with —I.

Emission Layer in Organic Layer 150

When the organic light-emitting device 10 is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, and/or a blue emission layer, according to a sub pixel. In various embodiments, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers may contact each other or may be separated from each other. In one or more embodiments, the emission layer may include two or more materials selected from a red-light emission material, a green-light emission material, and a blue-light emission material, in which the two or more materials may be mixed with each other in a single layer to thereby emit white light.

The emission layer may include a host and a dopant. The dopant may be at least one selected from a phosphorescent dopant and a fluorescent dopant.


56

The amount of the dopant in the emission layer may be about 0.01 to about 15 parts by weight based on 100 parts by weight of the host, but embodiments of the present disclosure are not limited thereto.

5 The thickness of the emission layer may be about 100 Å to about 1,000 Å, and in some embodiments, about 200 Å to about 600 Å. When the thickness of the emission layer is within these ranges, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

Host in Emission Layer

In one or more embodiments, the host may include a compound represented by Formula 301:



In Formula 301,

Ar_{301} may be selected from a substituted or unsubstituted C_5 - C_{60} carbocyclic group and a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

20 $xb11$ may be selected from 1, 2, and 3,

L_{301} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

$xb1$ may be an integer selected from 0 to 5,

R_{301} may be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_{301})(Q_{302})(Q_{303}), —N(Q_{301})(Q_{302}), —B(Q_{301})(Q_{302}), —C(=O)(Q_{301}), —S(=O)₂(Q_{301}), and —P(=O)(Q_{301})(Q_{302}), and

$xb21$ may be an integer selected from 1 to 5,

wherein Q_{301} to Q_{303} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, in Formula 301, Ar_{301} may be selected from the group consisting of:


60 a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenylene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothio-
 65 phene group; and

a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenylene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),


wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

In Formula 301, when xb11 is two or more, two or more Ar₃₀₁(s) may be connected (e.g., coupled) via a single bond.

In one or more embodiments, the compound represented by Formula 301 may be represented by one selected from Formula 301-1 and 301-2:



Formula 301-1



Formula 301-2

In Formulae 301-1 to 301-2,

A₃₀₁ to A₃₀₄ may each independently be selected from a benzene, a naphthalene, a phenanthrene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a pyridine, a pyrimidine, an indene, a fluorene, a spiro-bifluorene, a benzofluorene, a dibenzofluorene, an indole, a carbazole, a benzocarbazole, a dibenzocarbazole, a furan, a benzofuran, a dibenzofuran, a naphthofuran, a benzonaphthofuran, a dinaphthofuran, a thiophene, a benzothiophene, a dibenzothiophene, a naphthothiophene, a benzonaphthothiophene, and a dinaphthothiophene,

X₃₀₁ may be selected from O, S, and N-[(L₃₀₄)_{xb4}-R₃₀₄],

R₃₁₁ to R₃₁₄ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

xb22 and xb23 may each independently be selected from 0, 1, and 2,

L₃₀₁, xb1, R₃₀₁, and Q₃₁ to Q₃₃ may each independently be the same as described above,

L₃₀₂ to L₃₀₄ may each independently be the same as described above in connection with L₃₀₁,

xb2 to xb4 may each independently be the same as described above in connection with xb1, and

R₃₀₂ to R₃₀₄ may each independently be the same as described above in connection with R₃₀₁.

For example, in Formulae 301, 301-1, and 301-2, L₃₀₁ to L₃₀₄ may each independently be selected from the group consisting of:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a

dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazininylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a

triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indoyl group, an isoindoyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosiloyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolynyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),

wherein Q_{31} to Q_{33} may each independently be the same as described above.

In one or more embodiments, in Formulae 301, 301-1, and 301-2, R_{301} to R_{304} may each independently be selected from the group consisting of:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indoyl group, an isoindoyl group, a

benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosiloyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolynyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indoyl group, an isoindoyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosiloyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolynyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indoyl group, an isoindoyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosiloyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolynyl group, an isoquinolynyl group, a benzo-


61

quinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinoxalinyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,


wherein Q_{31} to Q_{33} may each independently be the same as described above.

In one or more embodiments, the host may include an alkaline earth metal complex. For example, the host may be selected from a beryllium (Be) complex (for example, Compound H55), a Mg complex, and a Zn complex.


The host may include at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55, but embodiments of the present disclosure are not limited thereto:




H1



H2



H3




H4

62


-continued

H5




5

H6




15

H7




30

H8




35

H9



45

H10








60






65

63

-continued

**64**


-continued



65

-continued

H21




5

10

15


H22



25

30

H23




35

40

45

50

H24



55


60

65


66

-continued




H25



H26



67
-continued



68
-continued

H27

H30

5

10

15

H31

20

25

H28

30

H32

35

40

45

H33

50






H29

55

H34


60

65



69

-continued




H35

5

10

15

H36




20

25

30

35

H37




40

45

50

55


H38




65

70


-continued



H39







H40



H41

71

-continued



72

-continued

H42

H46

5

10

15

H43

20

25

30

35

H44

40

45






H45

50

55






60

65



73


-continued




74

Phosphorescent Dopant in Organic Layer 150

The phosphorescent dopant may include an organometallic complex represented by Formula 401:



Formula 401



Formula 402

In Formulae 401 and 402,

H53 25 M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

L_{401} may be a ligand represented by Formula 402,

$xc1$ may be selected from 1, 2, and 3,

30 when $xc1$ is two or more, two or more L_{401} (s) may be identical to or different from each other,

L_{402} may be an organic ligand,

$xc2$ may be an integer selected from 0 to 4,

35 when $xc2$ is two or more, two or more L_{402} (s) may be identical to or different from each other,

H54 X_{401} to X_{404} may each independently be selected from nitrogen (N) and carbon (C),

40 X_{401} and X_{403} may be connected (e.g., coupled) via a single bond or a double bond, and X_{402} and X_{404} may be connected (e.g., coupled) via a single bond or a double bond,

A_{401} and A_{402} may each independently be selected from a C_5 - C_{60} carbocyclic group and a C_1 - C_{60} heterocyclic group,

X_{405} may be selected from a single bond, $*-O-*$, $*-S-*$, $*-C(=O)-*$, $*-N(Q_{411})-*$, $*-C(Q_{411})$

45 $(Q_{412})-*$, $*-C(Q_{411})=C(Q_{412})-*$, $*-C(Q_{411})=*$, and $*=C(Q_{411})=*$, wherein Q_{411} and Q_{412} may each independently be selected from hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

50 X_{406} may be selected from a single bond, O, and S,

H55 R_{401} and R_{402} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_1 - C_{20} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})$

75

(Q₄₀₂), —B(Q₄₀₁)(Q₄₀₂), —C(=O)(Q₄₀₁), —S(=O)₂(Q₄₀₁), and —P(=O)(Q₄₀₁)(Q₄₀₂), wherein Q₄₀₁ to Q₄₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a C₆-C₂₀ aryl group, and a C₁-C₂₀ heteroaryl group,

xc11 and xc12 may each independently be an integer selected from 0 to 10, and

in Formula 402, * and *' may each independently indicate a binding site to M in Formula 401.

In one or more embodiments, in Formula 402, A₄₀₁ and A₄₀₂ may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothio-phenylene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

In one or more embodiments, in Formula 402, i) X₄₀₁ may be nitrogen and X₄₀₂ may be carbon, or ii) both X₄₀₁ and X₄₀₂ may be nitrogen.

In one or more embodiments, in Formula 402, R₄₀₁ and R₄₀₂ may each independently be selected from the group consisting of:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, and C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantyl group, a norborn-
anyl group, and a norbornenyl group;

a cyclopentyl group, a cyclohexyl group, an adamantyl group, a norbornenyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

a cyclopentyl group, a cyclohexyl group, an adamantyl group, a norbornenyl group, a norbornenyl group a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantyl group, a norbornenyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl

76

group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

—Si(Q₄₀₁)(Q₄₀₂)(Q₄₀₃), —N(Q₄₀₁)(Q₄₀₂), —B(Q₄₀₁)(Q₄₀₂), —C(=O)(Q₄₀₁), —S(=O)₂(Q₄₀₁), and —P(=O)(Q₄₀₁)(Q₄₀₂),


wherein Q₄₀₁ to Q₄₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group.

In one or more embodiments, in Formula 401, when xc1 is two or more, two or more L₄₀₁ (s) may be optionally connected (e.g., coupled) through a linking group (for example, X₄₀₇), or two A₄₀₂(s) may be optionally connected (e.g., coupled) through a linking group (for example, X₄₀₈) (see Compounds PD1 to PD4 and PD7). X₄₀₇ and X₄₀₈ may each independently be selected from a single bond, *—O—*, *—S—*, *—C(=O)—*, *—N(Q₄₁₃)—*, *—C(Q₄₁₃)(Q₄₁₄)—*, and *—C(Q₄₁₃)=C(Q₄₁₄)—* (wherein Q₄₁₃ and Q₄₁₄ may each independently be selected from hydrogen, deuterium, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group), but embodiments of the present disclosure are not limited thereto.


In Formula 401, L₄₀₂ may be a monovalent, divalent, or trivalent organic ligand. For example, L₄₀₂ may be selected from a halogen atom, a diketone (for example, an acetylacetonate), a carboxylic acid (for example, a picolinate), —C(=O), an isonitrile, —CN, and a phosphorus ligand (for example, a phosphine and/or a phosphite), but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, the phosphorescent dopant may be selected from, for example, Compounds PD1 to PD25, but embodiments of the present disclosure are not limited thereto:

PD1









PD2



77

-continued



78

-continued

PD3

5

10

PD4

15

20

25

PD5

30

35

PD6

40

45

PD7






50

55

PD8

60

65



PD9

PD10






PD11

PD12

PD13

79

-continued




80

-continued

PD14

5


10



PD15 15

20

25



PD16

30


35

40

PD17

45

50




PD18

55

60

65



PD19


PD20

PD21

PD22


81

-continued



Fluorescent Dopant in Emission Layer

The fluorescent dopant may include a compound represented by Formula 501:



Formula 501

In Formula 501,

Ar₅₀₁ may be selected from a substituted or unsubstituted C₅-C₆₀ carbocyclic group and a substituted or unsubstituted C₁-C₆₀ heterocyclic group,

L₅₀₁ to L₅₀₃ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent

82

non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

xd1 to xd3 may each independently be an integer selected from 0 to 3,

R₅₀₁ and R₅₀₂ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and xd4 may be an integer selected from 1 to 6.

In one or more embodiments, in Formula 501, Ar₅₀₁ may be selected from the group consisting of:

a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group; and

a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, in Formula 501, L₅₀₁ to L₅₀₃ may each independently be selected from the group consisting of:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, and a pyridinylenylene group.

83

nylene group, a dibenzothiophenylene group, a benzocarbazolyne group, a dibenzocarbazolyne group, a dibenzosilolylene group, and a pyridinylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, and a pyridinyl group.

In one or more embodiments, in Formula 501, R_{501} and R_{502} may each independently be selected from the group consisting of:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, and a pyridinyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, and a pyridinyl group, and —Si(Q_{31})(Q_{32})(Q_{33}),


wherein Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

84


In one or more embodiments, in Formula 501, $xd4$ may be two, but embodiments of the present disclosure are not limited thereto.

For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:


FD1



FD2




FD3



85

-continued



FD4

5

10


15

20

25

86

-continued



FD7


30

35

40

45

FD8



FD5



50

55

60


65

FD9



87

-continued

PD11

15

20

PD12

30

35

PD13

45

50

PD14





55

60

65

88


-continued



89

-continued


FD19



90


-continued

FD21




FD22


FD20



In one or more embodiments, the fluorescent dopant may be selected from the compounds illustrated below, but embodiments of the present disclosure are not limited thereto:




DPVBi




DPAVB

91




-continued


92




DCM



DCJTb



Coumarin 6



C545T

Electron Transport Region in Organic Layer 150

The electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer (ETL), and an electron injection layer, but embodiments of the present disclosure are not limited thereto.

For example, the electron transport region may have a structure of electron transport layer/electron injection layer, a structure of hole blocking layer/electron transport layer/electron injection layer, a structure of electron control layer/electron transport layer/electron injection layer, or a structure of buffer layer/electron transport layer/electron injection layer, wherein layers of each structure are sequentially stacked on the emission layer in each stated order. However, embodiments of the structure of the electron transport layer are not limited thereto.

The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, and/or an electron transport layer in the electron transport region) may include a metal-free compound containing at least one π electron-depleted nitrogen-containing ring.

As used herein, " π electron-depleted nitrogen-containing ring" indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

For example, the " π electron-depleted nitrogen-containing ring" may be i) a 5-membered to 7-membered hetero monocyclic group having at least one $*-N=*$ moiety, ii) a heteropolycyclic group in which two or more 5-membered to 7-membered hetero monocyclic groups each having at least one $*-N=*$ moiety are condensed with each other, or iii) a heteropolycyclic group in which at least one 5-membered to 7-membered hetero monocyclic group having at least one $*-N=*$ moiety is condensed with at least one C_5 - C_{60} carbocyclic group.

Non-limiting examples of the π electron-depleted nitrogen-containing ring may include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, a thiadiazole, an imidazopyridine, an imidazopyrimidine, and an azacarbazole, but embodiments of the present disclosure are not limited thereto.

The electron transport region may include the compound represented by Formula 1 according to an embodiment of

the present disclosure. In one or more embodiments, the electron transport layer may include the compound represented by Formula 1 according to an embodiment of the present disclosure.

The electron transport region and/or the electron transport layer may further include, in addition to the compound of Formula 1, other known compounds. For example, the electron transport region may include a compound represented by Formula 601:



In Formula 601,

Ar_{601} may be selected from a substituted or unsubstituted $\text{C}_5\text{-C}_{60}$ carbocyclic group and a substituted and unsubstituted $\text{C}_1\text{-C}_{60}$ heterocyclic group,

$\text{xe}11$ may be selected from 1, 2, and 3,

L_{601} may be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkylene group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenylene group, a substituted or unsubstituted $\text{C}_6\text{-arylene}$ group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

$\text{xe}1$ may be an integer selected from 0 to 5,

R_{601} may be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryloxy group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylthio group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_{601})(\text{Q}_{602})(\text{Q}_{603})$, $-\text{C}(=\text{O})(\text{Q}_{601})$, $-\text{S}(=\text{O})_2(\text{Q}_{601})$, and $-\text{P}(=\text{O})(\text{Q}_{601})(\text{Q}_{602})$, wherein Q_{601} to Q_{603} may each independently be selected from a $\text{C}_1\text{-C}_{10}$ alkyl group, a $\text{C}_1\text{-C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, and

$\text{xe}21$ may be an integer selected from 1 to 5.

In one or more embodiments, at least one selected from the $\text{xe}11$ $\text{Ar}_{601}(\text{s})$ and the $\text{xe}21$ $\text{R}_{601}(\text{s})$ may include the π electron-depleted nitrogen-containing ring as described above.

In one or more embodiments, in Formula 601, ring Ar_{601} may be selected from the group consisting of:

a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a

phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{20}$ alkyl group, a $\text{C}_1\text{-C}_{20}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,


wherein Q_{31} to Q_{33} may each independently be selected from a $\text{C}_1\text{-C}_{10}$ alkyl group, a $\text{C}_1\text{-C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In Formula 601, when $\text{xe}11$ is two or more, two or more $\text{Ar}_{601}(\text{s})$ may be connected (e.g., coupled) via a single bond.

In one or more embodiments, in Formula 601, Ar_{601} may be an anthracene group.

In one or more embodiments, the compound represented by Formula 601 may be represented by Formula 601-1:

Formula 601-1



In Formula 601-1,

X_{614} may be selected from N and C(R_{614}), X_{615} may be selected from N and C(R_{615}), and X_{616} may be selected from N and C(R_{616}), wherein at least one selected from X_{614} to X_{616} may be N,

L_{611} to L_{613} may each independently be the same as described above in connection with L_{601} ,

$\text{xe}611$ to $\text{xe}613$ may each independently be the same as described above in connection with $\text{xe}1$,

R_{611} to R_{613} may each independently be the same as described above in connection with R_{601} , and

R₆₁₄ to R₆₁₆ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, in Formulae 601 and 601-1, L₆₀₁ and L₆₁₁ to L₆₁₃ may each independently be selected from the group consisting of:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a

spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indoyl group, an isoindoyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyrimidinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group, but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, in Formulae 601 and 601-1, xe1 and xe611 to xe613 may each independently be selected from 0, 1, and 2.

In one or more embodiments, in Formulae 601 and 601-1, R₆₀₁ and R₆₁₁ to R₆₁₃ may each independently be selected from the group consisting of:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indoyl group, an isoindoyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyrimidinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indoyl group, an isoindoyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a

97


pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

—S(=O)₂(Q₆₀₁) and —P(=O)(Q₆₀₁)(Q₆₀₂),

wherein Q₆₀₁ and Q₆₀₂ may each independently be the same as described above.

The electron transport region may include at least one selected from Compounds ET1 to ET36, but embodiments of the present disclosure are not limited thereto:

ET1



50

55


60

65


98

-continued


ET2



ET3




ET4




99

-continued

**100**

-continued

ET8

ET6

25

30

35

40

ET7

45


50

55

60


65

ET9



101

-continued



ET10

5


10

15

20

25

ET11




30

35

40

45

ET12



50


55

60

65


102

-continued




ET13

ET14






ET15



103

-continued

**104**

-continued

ET16

5

10

15

20

ET17

25

30

35

40

ET18 45

50

55




60


65

ET19

ET20

ET21



105
-continued

ET22

5


10

15

20


106
-continued

ET25



25

ET26




ET23

30

35


40

45



ET24


50



55

60

65




ET27

107

-continued

ET28




5

10

15

20

ET29



25


30

35

40

45

ET30



50

55


60

65


108

-continued


ET31



ET32






ET33



109


-continued




In one or more embodiments, the electron transport region may include at least one compound selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq₃, Balq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ:

110


ET34

Alq₃


ET35



BALq



TAZ



ET36

NTAZ

50

The thicknesses of the buffer layer, the hole blocking layer, and/or the electron control layer may be about 20 Å to about 1,000 Å, and in some embodiments, about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are each within these ranges, the electron blocking layer may have excellent electron blocking characteristics and/or electron control characteristics without a substantial increase in driving voltage.


The thickness of the electron transport layer may be about 100 Å to about 1,000 Å, and in some embodiments, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within these ranges, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

111


The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.

The metal-containing material may include at least one selected from an alkali metal complex and an alkaline earth metal complex. The alkali metal complex may include a metal ion selected from a lithium (Li) ion, a sodium (Na) ion, a potassium (K) ion, a rubidium (Rb) ion, and a cesium (Cs) ion, and the alkaline earth metal complex may include a metal ion selected from a beryllium (Be) ion, a magnesium (Mg) ion, a calcium (Ca) ion, a strontium (Sr) ion, and a barium (Ba) ion. One or more ligands coordinated with the metal ion of the alkali metal complex and/or the alkaline earth metal complex may be selected from a hydroxyquinoline, a hydroxyisoquinoline, a hydroxybenzoquinoline, a hydroxyacridine, a hydroxyphenanthridine, a hydroxyphenyl oxazole, a hydroxyphenyl thiazole, a hydroxydiphenyl oxadiazole, a hydroxydiphenyl thiadiazole, a hydroxyphenyl pyridine, a hydroxyphenyl benzimidazole, a hydroxyphenyl benzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

For example, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) and/or Compound ET-D2:



ET-D1



ET-D2

The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode **190**. The electron injection layer may directly contact the second electrode **190**.

The electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

The electron injection layer may include a reducing dopant.

The reducing dopant may include at least one selected from an alkali metal, an alkaline earth metal, a rare earth based metal, an alkali metal compound, an alkaline earth metal compound, a rare earth based metal compound, an alkali metal complex, an alkaline earth metal complex, and a rare earth based metal complex.

112

The alkali metal may be selected from Na, K, Rb, and Cs. In one or more embodiments, the alkali metal may be selected from K, Rb, and Cs. In one or more embodiments, the alkali metal may be selected from Rb and Cs, but embodiments of the present disclosure are not limited thereto.

The alkaline earth metal may be selected from Ca, Sr, and Ba.

The rare earth metal may be selected from scandium (Sc), yttrium (Y), cerium (Ce), ytterbium (Yb), gadolinium (Gd), and terbium (Tb).

The alkali metal compound, the alkaline earth metal compound, and the rare earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, and/or iodines) of the alkali metal, the alkaline earth metal, and the rare earth metal.

The alkali metal compound may be selected from alkali metal oxides (such as Li_2O , Cs_2O , and/or K_2O) and alkali metal halides (such as LiF , NaF , CsF , KF , LiI , NaI , CsI , and/or KI). In one or more embodiments, the alkali metal compound may be selected from LiF , Li_2O , NaF , LiI , NaI , CsI , and KI , but embodiments of the present disclosure are not limited thereto.

The alkaline earth metal compound may be selected from alkaline earth metal compounds (such as BaO , SrO , CaO , $\text{Ba}_x\text{Sr}_{1-x}\text{O}$ ($0 < x < 1$), and/or $\text{Ba}_x\text{Ca}_{1-x}\text{O}$ ($0 < x < 1$)). In one or more embodiments, the alkaline earth metal compound may be selected from BaO , SrO , and CaO , but embodiments of the present disclosure are not limited thereto.

The rare earth metal compound may be selected from YbF_3 , ScF_3 , ScO_3 , Y_2O_3 , Ce_2O_3 , GdF_3 , and TbF_3 . In one or more embodiments, the rare earth metal compound may be selected from YbF_3 , ScF_3 , TbF_3 , YbI_3 , ScI_3 , and TbI_3 , but embodiments of the present disclosure are not limited thereto.

The alkali metal complex, the alkaline earth metal complex, and the rare earth metal complex may each include a metal ion selected from an alkali metal, an alkaline earth metal, and a rare earth metal as described above, respectively. Each ligand coordinated with a metal ion in the alkali metal complex, the alkaline earth metal complex, and/or the rare earth metal complex may independently be selected from a hydroxyquinoline, a hydroxyisoquinoline, a hydroxybenzoquinoline, a hydroxyacridine, a hydroxyphenanthridine, a hydroxyphenyl oxazole, a hydroxyphenyl thiazole, a hydroxydiphenyl oxadiazole, a hydroxydiphenyl thiadiazole, a hydroxyphenyl pyridine, a hydroxyphenyl benzimidazole, a hydroxyphenyl benzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

The electron injection layer may include only the reducing dopant as described above. In one or more embodiments, the electron injection layer may further include, in addition to the reducing dopant, an organic material. When the electron injection layer includes the reducing dopant and an organic material, the reducing dopant may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

The thickness of the electron injection layer may be about 1 Å to about 100 Å, and in some embodiments, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within these ranges, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

Second Electrode 190

The second electrode 190 may be on the organic layer 150. The second electrode 190 may be a cathode, which is an electron injection electrode, and in this regard, a material for forming the second electrode 190 may be selected from a metal, an alloy, an electrically conductive compound, and a mixture thereof, each having a relatively low work function.

The second electrode 190 may include at least one selected from Li, Mg, Al, Al—Li, Ca, Mg—In, Mg—Ag, ITO, and IZO, but embodiments of the present disclosure are not limited thereto. The second electrode 190 may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

The second electrode 190 may have a single-layered structure, or a multi-layered structure including two or more layers.

According to an embodiment of the present disclosure, the organic light-emitting device may be suitably included in one or more types or kinds of display apparatuses, for example, a passive matrix organic light-emitting display apparatus and an active matrix organic light-emitting display apparatus. For example, when the organic light-emitting device is included in an active matrix organic light-emitting display apparatus, a first electrode on a substrate may be a pixel electrode, and the first electrode may be electrically connected to a source electrode or a drain electrode of a thin film transistor 200. In some embodiments, the organic light-emitting device may be included in a display apparatus that may display images on both sides.

Description of FIGS. 2 to 4

The organic light-emitting device 20 of FIG. 2 may include a first capping layer 210, a first electrode 110, an organic layer 150, and a second electrode 190, which are sequentially stacked in this stated order. The organic light-emitting device 30 of FIG. 3 may include a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220, which are sequentially stacked in this stated order. The organic light-emitting device 40 of FIG. 4 includes a first capping layer 210, a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220, which are sequentially stacked in this stated order.

In FIGS. 2 to 4, the first electrode 110, the organic layer 150, and the second electrode 190 may each be the same as described herein in connection with FIG. 1.

In the organic layer 150 of each of the organic light-emitting devices 20 and 40, light generated in an emission layer may pass through the first electrode 110 (which is a semi-transmissive electrode or a transmissive electrode) and the first capping layer 210 toward the outside, and in the organic layer 150 of each of the organic light-emitting devices 30 and 40, light generated in an emission layer may pass through the second electrode 190 (which is a semi-transmissive electrode or a transmissive electrode) and the second capping layer 220 toward the outside.

The first capping layer 210 and the second capping layer 220 may increase the external luminescent efficiency of the organic light emitting device according to the principle of constructive interference. 4


The first capping layer 210 and the second capping layer 220 may each independently be an organic capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.

At least one selected from the first capping layer 210 and the second capping layer 220 may include at least one material selected from carbocyclic compounds, heterocyclic compounds, amine-based compounds, porphyrin derivatives, phthalocyanine derivatives, naphthalocyanine derivatives, alkali metal complexes, and alkaline earth metal complexes. The carbocyclic compound, the heterocyclic compound, and the amine-based compound may be optionally substituted with a substituent containing at least one element selected from O, N, sulfur (S), selenium (Se), silicon (Si), fluorine (F), chlorine (Cl), bromine (Br), and iodine (I).

In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may include an amine-based compound.


In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may include the compound represented by Formula 201 or the compound represented by Formula 202.

In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may include a compound selected from Compounds HT28 to HT33 and Compounds CP1 to CP5, but embodiments of the present disclosure are not limited thereto:



115

-continued



Hereinbefore, the organic light-emitting device according to an embodiment of the present disclosure has been described in connection with FIGS. 1 to 4. However, embodiments of the present disclosure are not limited thereto.

The layers constituting the hole transport region, the emission layer, and the layers constituting the electron transport region may be formed in a specific region using one or more methods selected from vacuum deposition, spin coating, casting, a Langmuir-Blodgett (LB) method, ink-jet printing, laser-printing, and laser-induced thermal imaging (LITI).

When the layers constituting the hole transport region, the emission layer, and the layers constituting the electron transport region are each formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of about 100 to about 500° C., at a vacuum degree of about 10⁻⁸ to about 10⁻³ torr, and at a deposition rate of about 0.01 to about 100 Å/sec, depending on the compound to be included in each layer, and the intended structure of each layer to be formed.

When the layers constituting the hole transport region, the emission layer, and the layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to 200° C., depending on the compound to be included in each layer, and the intended structure of each layer to be formed.

General Definition of Substituents

Hereinafter, definitions of representative substituents of compounds used herein will be presented. The number of carbon atoms used to restrict a substituent is not limited, and does not limit the properties of the substituent. Unless defined otherwise, the definition of the substituent is consistent with a general definition thereof.

116

The term “C₁-C₆₀ alkyl group” as used herein refers to a linear or branched aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and non-limiting examples thereof may include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “C₁-C₆₀ alkenylene group” as used herein refers to a divalent group having substantially the same structure as the C₁-C₆₀ alkyl group.

The term “C₂-C₆₀ alkenyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon double bond in the body (e.g., middle) or at the terminus of the C₂-C₆₀ alkyl group, and non-limiting examples thereof may include an ethenyl group, a propenyl group, and a butenyl group. The term “C₂-C₆₀ alkenylene group” as used herein refers to a divalent group having substantially the same structure as the C₂-C₆₀ alkenyl group.

The term “C₂-C₆₀ alkynyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the body (e.g., middle) or at the terminus of the C₂-C₆₀ alkyl group, and non-limiting examples thereof may include an ethynyl group and a propynyl group. The term “C₂-C₆₀ alkynylene group” as used herein refers to a divalent group having substantially the same structure as the C₂-C₆₀ alkynyl group.

The term “C₁-C₆₀ alkoxy group” as used herein refers to a monovalent group represented by —O-A₁₀₁ (wherein A₁₀₁ is a C₁-C₆₀ alkyl group), and non-limiting examples thereof may include a methoxy group, an ethoxy group, and an isopropoxy group.

The term “C₃-C₁₀ cycloalkyl group” as used herein refers to a monovalent hydrocarbon monocyclic group having 3 to 10 carbon atoms, and non-limiting examples thereof may include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group” as used herein refers to a divalent group having substantially the same structure as the C₃-C₁₀ cycloalkyl group.

The term “C₁-C₁₀ heterocycloalkyl group” as used herein refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom in addition 1 to 10 carbon atoms. Non-limiting examples thereof may include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuranyl group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group” as used herein refers to a divalent group having substantially the same structure as the C₁-C₁₀ heterocycloalkyl group.

The term “C₃-C₁₀ cycloalkenyl group” as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one double bond in the ring thereof, and does not have aromaticity (e.g., is non-aromatic). Non-limiting examples thereof may include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group” as used herein refers to a divalent group having substantially the same structure as the C₃-C₁₀ cycloalkenyl group.

The term “C₁-C₁₀ heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one double bond in its ring. Non-limiting examples of the C₁-C₁₀ heterocycloalkenyl group may include a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-hydrofuranyl group, and a 2,3-hydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group” as used herein refers to a divalent group having substantially the same structure as the C₁-C₁₀ heterocycloalkenyl group.

The term “C₆-C₆₀ aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system including 6 to 60 carbon atoms, and the term “C₆-C₆₀ arylene group” as used herein refers to a divalent group having a carbocyclic aromatic system including 6 to 60 carbon atoms. Non-limiting examples of the C₆-C₆₀ aryl group may include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be fused (e.g., condensed).

The term “C₁-C₆₀ heteroaryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom in addition to 1 to 60 carbon atoms. The term “C₁-C₆₀ heteroarylene group” as used herein refers to a divalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Non-limiting examples of the C₁-C₆₀ heteroaryl group may include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be fused (e.g., condensed).

The term “C₆-C₆₀ aryloxy group” as used herein indicates —O-A₁₀₂ (wherein A₁₀₂ is a C₆-C₆₀ aryl group), and the term “C₆-C₆₀ arylthio group” as used herein indicates —S-A₁₀₃ (wherein A₁₀₃ is a C₆-C₆₀ aryl group).

The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group that has two or more rings condensed with each other, only carbon atoms (for example, 8 to 60 carbon atoms) as ring-forming atoms, and non-aromaticity in the entire molecular structure (e.g., the molecule as a whole is non-aromatic). A non-limiting example of the monovalent non-aromatic condensed polycyclic group may include a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group” as used herein refers to a divalent group having substantially the same structure as the monovalent non-aromatic condensed polycyclic group.

The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group that has two or more rings condensed, has at least one heteroatom selected from N, O, Si, P, and S in addition to carbon atoms (for example, 1 to 60 carbon atoms) as ring-forming atoms, and has non-aromaticity in the entire molecular structure. A non-limiting example of the monovalent non-aromatic condensed heteropolycyclic group may include a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein refers to a divalent group having substantially the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

The term “C₅-C₆₀ carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which the ring-forming atoms are all carbons. The C₅-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₅-C₆₀ carbocyclic group may be a ring (such as benzene), a monovalent group (such as a phenyl group), or a divalent group (such as a phenylene group). In one or more embodiments, depending on the number of substituents connected (e.g., coupled) to the C₅-C₆₀ carbocyclic group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

The term “C₁-C₆₀ heterocyclic group” as used herein refers to a group having substantially the same structure as the C₅-C₆₀ carbocyclic group, except that at least one heteroatom selected from N, O, Si, P, and S is used in addition to the carbon atoms as a ring-forming atom. The number of carbon atoms may be 1 to 60.

At least one substituent of the substituted C₅-C₆₀ carbocyclic group, the substituted C₁-C₆₀ heterocyclic group, the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from the group consisting of:

deuterium(-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

119

ovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_{21})(\text{Q}_{22})(\text{Q}_{23})$, $-\text{N}(\text{Q}_{21})(\text{Q}_{22})$, $-\text{B}(\text{Q}_{21})(\text{Q}_{22})$, $-\text{C}(=\text{O})(\text{Q}_{21})$, $-\text{S}(=\text{O})_2(\text{Q}_{21})$, and $-\text{P}(=\text{O})(\text{Q}_{21})(\text{Q}_{22})$; and

$-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,

wherein Q_{11} to Q_{13} , Q_{21} to Q_{23} , and Q_{31} to Q_{33} may each independently be selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group and a terphenyl group.

The term “Ph” as used herein refers to a phenyl group, the term “Me” as used herein refers to a methyl group, the term “Et” as used herein refers to an ethyl group, the term “ter-Bu” or “Bu” as used herein refers to a tert-butyl group, and the term “OMe” as used herein refers to a methoxy group.


The term “biphenyl group” as used herein refers to “a phenyl group substituted with a phenyl group”. In other words, a “biphenyl group” refers to “a substituted phenyl group having a C₆-C₆₀ aryl group as a substituent”.

The term “terphenyl group” as used herein refers to “a phenyl group substituted with a biphenyl group”. In other words, a “terphenyl group” refers to “a substituted phenyl group having a C₆-C₆₀ aryl group substituted with a C₆-C₆₀ aryl group as a substituent”.


Symbols * and * as used herein, unless defined otherwise, refer to binding sites to neighboring atoms in a corresponding formula.

Hereinafter, a compound represented by Formula 1 according to an embodiment of the present disclosure and an organic light-emitting device according to an embodiment of the present disclosure will be described in more detail with reference to Synthesis Examples and Examples. The wording “B was used instead of A” used in describing Synthesis Examples refers to that an identical number of molar equivalents of A was used in place of B.

SYNTHESIS EXAMPLES



120



Synthesis Example 1: Synthesis of Intermediate-2 (In-2)

Synthesis of Intermediate-1 (In-1)

$\text{Pd}(\text{PPh}_3)_4$ (0.02 equivalent (eq)), Na_2CO_3 (1.2 eq), and a mixture of toluene, distilled water, and ethanol (5:3:2 (v/v) based on 0.1 M reagent (1 eq)) were added to a flask containing 2-bromo-12,12-dimethyl-4,12-dihydroindeno[2,1-a]phenylene (1 eq) and (2-nitrophenyl)boronic acid (1.2 eq), and the mixed solution was stirred under reflux for 12 hours. The reaction solution was cooled to room temperature, and extraction was performed using methylene chloride (MC), followed by washing using distilled water. The resulting solution was dried with MgSO_4 , filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound In-1 (yield: 87%).

High Resolution Mass Spectrometry (HRMS) for $\text{C}_{28}\text{H}_{21}\text{NO}_2$ $[\text{M}]^+$: calc: 403.48, found: 402.

Synthesis of In-2

5 g of In-1 was dissolved in 15 g of triethylphosphite, and the mixed solution was stirred under reflux in a nitrogen atmosphere for 12 hours. After the reaction was completed, the reaction solution was subjected to a vacuum distillation process to remove unreacted triethylphosphite. The resulting solution was purified by column chromatography using a mixture of hexane and MC (4:1 (v/v)), thereby obtaining the desired compound, In-2 (yield: 51.4%).

HRMS for $\text{C}_{28}\text{H}_{21}\text{N}$ $[\text{M}]^+$: calc: 371.48, found: 370.

Synthesis Example 2: Synthesis of Compound ET-1

$\text{Pd}(\text{dba})_3$ (0.03 eq), $(t\text{-Bu})_3\text{P}$ (0.06 eq), and toluene (based on 0.1 M reagent (1 eq)) were added to a flask containing

121

Int-2 (1 eq) and 2-bromo-4,6-diphenyl-1,3,5-triazine (1.2 eq). The mixed solution was stirred under reflux for 5 hours. The reaction solution was cooled to room temperature, and extraction was performed using MC, followed by washing using distilled water. The resulting solution was dried with MgSO_4 , filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound ET-1 (yield: 76.1%).

HRMS for $\text{C}_{43}\text{H}_{30}\text{N}_4$ $[\text{M}]^+$: calc: 602.74, found: 601.

Elemental Analysis for $\text{C}_{43}\text{H}_{30}\text{N}_4$ calc: C, 85.69; H, 5.02; N, 9.30

Synthesis Example 3: Synthesis of Compound ET-2

$\text{Pd}(\text{dba})_3$ (0.03 eq), $(\text{t-Bu})_3\text{P}$ (0.06 eq), and toluene (based on 0.1 M reagent (1 eq)) were added to a flask containing Int-2 (1 eq) and 2-bromo-4,6-diphenylpyrimidine (1.2 eq), and the mixed solution was stirred under reflux for 5 hours. The reaction solution was cooled to room temperature, and extraction was performed using MC, followed by washing using distilled water. The resulting solution was dried with MgSO_4 , filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound ET-2 (yield: 77.8%).

HRMS for $\text{C}_{44}\text{H}_{31}\text{N}_3$ $[\text{M}]^+$: calc: 601.75, found: 600.

Elemental Analysis for $\text{C}_{44}\text{H}_{31}\text{N}_3$ calc: C, 87.82; H, 5.19; N, 6.98

Synthesis Example 4: Synthesis of Compound ET-3

$\text{Pd}(\text{dba})_3$ (0.03 eq), $(\text{t-Bu})_3\text{P}$ (0.06 eq), and toluene (based on 0.1 M reagent (1 eq)) were added to a flask containing Int-2 (1 eq) and 4-bromo-2,6-diphenylpyrimidine (1.2 eq), and the mixed solution was stirred under reflux for 5 hours. The reaction solution was cooled to room temperature, followed by extraction using methylene chloride (MC) and washing using distilled water. The resultant was dried with MgSO_4 , filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound ET-3 (yield: 71.6%).

HRMS for $\text{C}_{44}\text{H}_{31}\text{N}_3$ $[\text{M}]^+$: calc: 601.75, found: 600.

Elemental Analysis for $\text{C}_{44}\text{H}_{31}\text{N}_3$ calc: C, 87.82; H, 5.19; N, 6.98

Synthesis Example 5: Synthesis of Compound ET-5

$\text{Pd}(\text{dba})_3$ (0.03 eq), $(\text{t-Bu})_3\text{P}$ (0.06 eq), and toluene (based on 0.1 M reagent (1 eq)) were added to a flask containing Int-2 (1 eq) and 4-bromo-2,6-diphenylpyridine (1.2 eq), and the mixed solution was stirred under reflux for 5 hours. The reaction solution was cooled to room temperature, and extraction was performed using MC, followed by washing using distilled water. The resulting solution was dried with MgSO_4 , filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound ET-5 (yield: 74.1%).

HRMS for $\text{C}_{45}\text{H}_{32}\text{N}_2$ $[\text{M}]^+$: calc: 600.77, found: 599.

Elemental Analysis for $\text{C}_{45}\text{H}_{32}\text{N}_2$ calc: C, 89.97; H, 5.37; N, 4.66

Synthesis Example 6: Synthesis of Compound ET-7

$\text{Pd}(\text{dba})_3$ (0.03 eq), $(\text{t-Bu})_3\text{P}$ (0.06 eq), and toluene (based on 0.1 M reagent (1 eq)) were added to a flask containing Int-2 (1 eq) and 2-bromo-4,6-di(naphthalen-2-yl)pyridine (1.2 eq), and the mixed solution was stirred under reflux for 5 hours. The reaction solution was cooled to room tempera-

122

ture, and extraction was performed using MC, followed by washing using distilled water. The resulting solution was dried with MgSO_4 , filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound ET-7 (yield: 71.9%).

HRMS for $\text{C}_{53}\text{H}_{36}\text{N}_2$ $[\text{M}]^+$: calc: 700.89, found: 699.

Elemental Analysis for $\text{C}_{53}\text{H}_{36}\text{N}_2$ calc: C, 90.83; H, 5.18; N, 4.00

Synthesis Example 7: Synthesis of Compound ET-13

$\text{Pd}(\text{dba})_3$ (0.03 eq), $(\text{t-Bu})_3\text{P}$ (0.06 eq), and toluene (0.1 M reagent (1 eq)) were added to a flask containing Int-2 (1 eq) and 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (1.2 eq), and the mixed solution was stirred under reflux for 5 hours. The reaction solution was cooled to room temperature, and extraction was performed using MC, followed by washing using distilled water. The resulting solution was dried with MgSO_4 , filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound ET-13 (yield: 69.4%).

HRMS for $\text{C}_{49}\text{H}_{34}\text{N}_4$ $[\text{M}]^+$: calc: 678.84, found: 677.

Elemental Analysis for $\text{C}_{49}\text{H}_{34}\text{N}_4$ calc: C, 86.70; H, 5.05; N, 8.25

Synthesis Example 8: Synthesis of Compound ET-14

$\text{Pd}(\text{dba})_3$ (0.03 eq), $(\text{t-Bu})_3\text{P}$ (0.06 eq), and toluene (0.1 M reagent (1 eq)) were added to a flask containing Int-2 (1 eq) and 2-(4-bromophenyl)-4,6-diphenylpyrimidine (1.2 eq), and the mixed solution was stirred under reflux for 5 hours. The reaction solution was cooled to room temperature, and extraction was performed using MC, followed by washing using distilled water. The resulting solution was dried with MgSO_4 , filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound ET-14 (yield: 66.7%).

HRMS for $\text{C}_{50}\text{H}_{35}\text{N}_3$ $[\text{M}]^+$: calc: 677.85, found: 676.

Elemental Analysis for $\text{C}_{50}\text{H}_{35}\text{N}_3$ calc: C, 88.60; H, 5.20; N, 6.20

Synthesis Example 9: Synthesis of Compound ET-16

$\text{Pd}(\text{dba})_3$ (0.03 eq), $(\text{t-Bu})_3\text{P}$ (0.06 eq), and toluene (0.1 M reagent (1 eq)) were added to a flask containing Int-2 (1 eq) and 2-(3-bromophenyl)-4,6-diphenyl-1,3,5-triazine (1.2 eq), and the mixed solution was stirred under reflux for 5 hours. The reaction solution was cooled to room temperature, and extraction was performed using MC, followed by washing using distilled water. The resulting solution was dried with MgSO_4 , filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound ET-16 (yield: 69.4%).

HRMS for $\text{C}_{49}\text{H}_{34}\text{N}_4$ $[\text{M}]^+$: calc: 678.84, found: 677.

Elemental Analysis for $\text{C}_{49}\text{H}_{34}\text{N}_4$ calc: C, 86.70; H, 5.05; N, 8.25

123

Synthesis Example 10: Synthesis of Compound ET-19

Pd(dba)₃ (0.03 eq), (t-Bu)₃P (0.06 eq), and toluene (0.1 M reagent (1 eq)) were added to a flask containing Int-2 (1 eq) and (4-bromophenyl)diphenylphosphine oxide (1.2 eq), and the mixed solution was stirred under reflux for 5 hours. The reaction solution was cooled to room temperature, and extraction was performed using MC, followed by washing using distilled water. The resulting solution was dried with MgSO₄, filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound ET-19 (yield: 69.4%).

HRMS for C₄₆H₃₄NOP [M]⁺: calc: 647.76, found: 646.

Elemental Analysis for C₄₉H₃₄N₄ calc: C, 85.30; H, 5.29; N, 2.16; O, 2.47; P, 4.78

Synthesis Example 11: Synthesis of Compound ET-20

Pd(dba)₃ (0.03 eq), (t-Bu)₃P (0.06 eq), and toluene (0.1 M reagent (1 eq)) were added to a flask containing Int-2 (1 eq) and (3-bromophenyl)diphenylphosphine oxide (1.2 eq), and the mixed solution was stirred under reflux for 5 hours. The reaction solution was cooled to room temperature, and extraction was performed using MC, followed by washing using distilled water. The resulting solution was dried with MgSO₄, filtered, and distilled under reduced pressure. The residue obtained therefrom was purified by column chromatography, thereby obtaining Compound ET-20 (yield: 66.4%).

HRMS for C₄₆H₃₄NOP [M]⁺: calc: 647.76, found: 646.

Elemental Analysis for C₄₉H₃₄N₄ calc: C, 85.30; H, 5.29; N, 2.16; O, 2.47; P, 4.78

Example 1



An anode was prepared by cutting an indium tin oxide (ITO) glass substrate (manufactured by Corning) with ITO at a thickness of 15 Ω/cm² (500 Å) to a size of 50 mm×50 mm×0.5 mm, ultrasonically cleaning the ITO glass substrate using isopropyl alcohol and pure water for 10 minutes each, exposing to UV irradiation for 10 minutes, and cleaning with ozone. Then, the ITO glass substrate was loaded into a vacuum deposition apparatus.

2-TNATA was vacuum deposited on the ITO glass substrate to form a hole injection layer having a thickness of 600 Å, and Compound NPB (which is a hole-transporting compound), was vacuum deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å.

Tris[2-phenylpyridinato]iridium(III) (Ir(ppy)₃) (which is a green phosphorescent dopant) and CBP were co-deposited on the hole transport layer at a weight ratio of 15:85 to form an emission layer having a thickness of 300 Å.

Subsequently, Compound E-1 was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å, and Al was deposited on the electron transport layer to form an Al electrode (i.e., a negative electrode) having a thickness of 1,200 Å, thereby completing the manufacture of an organic light-emitting device.

124



Example 2

An organic light-emitting device was manufactured in substantially the same manner as in Example 1, except that Compound E-2 was used instead of Compound E-1 in forming the electron transport layer.

Example 3

An organic light-emitting device was manufactured in substantially the same manner as in Example 1, except that Compound E-3 was used instead of Compound E-1 in forming the electron transport layer.

Example 4

An organic light-emitting device was manufactured in substantially the same manner as in Example 1, except that

125

Compound E-5 was used instead of Compound E-1 in forming the electron transport layer.

Example 5

An organic light-emitting device was manufactured in substantially the same manner as in Example 1, except that Compound E-7 was used instead of Compound E-1 in forming the electron transport layer.

Example 6

An organic light-emitting device was manufactured in substantially the same manner as in Example 1, except that Compound E-13 was used instead of Compound E-1 in forming the electron transport layer.

Example 7

An organic light-emitting device was manufactured in substantially the same manner as in Example 1, except that Compound E-14 was used instead of Compound E-1 in forming the electron transport layer.

Example 8

An organic light-emitting device was manufactured in substantially the same manner as in Example 1, except that Compound E-16 was used instead of Compound E-1 in forming the electron transport layer.

Example 9


An organic light-emitting device was manufactured in substantially the same manner as in Example 1, except that Compound E-19 was used instead of Compound E-1 in forming the electron transport layer.

Example 10

An organic light-emitting device was manufactured in substantially the same manner as in Example 1, except that Compound E-20 was used instead of Compound E-1 in forming the electron transport layer.

Comparative Example 1

An organic light-emitting device was manufactured in substantially the same manner as in Example 1, except that Compound Alq₃ was used instead of Compound E-1 in forming the electron transport layer.



The efficiency and lifespan (T₉₅) of each of the organic light-emitting devices of Examples 1 to 10 and Comparative Example were measured, and the results thereof are shown in Table 1.

126

TABLE 1

	Material for forming an electron transport layer	Efficiency (cd/A)	Lifespan (T ₉₅) (@ 3700 nit)
Example 1	ET-1	51	923
Example 2	ET-2	48.8	911
Example 3	ET-3	47.7	897
Example 4	ET-5	47.9	921
Example 5	ET-7	49.2	943
Example 6	ET-13	48.0	937
Example 7	ET-14	47.9	956
Example 8	ET-16	42	942
Example 9	ET-19	46.4	965
Example 10	ET-20	42.7	970
Comparative Example	Alq ₃	38.4	789

Compounds having the structure of Formula 1 according to an embodiment of the present disclosure were used as electron-transporting materials, and consequently, all the organic light-emitting devices including these compounds showed significantly improved characteristics, for example, increased lifespan characteristics, compared to the organic light-emitting device of Comparative Example 1. Accordingly, it was confirmed that the compounds having the structure of Formula 1 according to the present disclosure were excellent electron-transporting materials with significant effects.

An organic light-emitting device according to an embodiment of the present disclosure may have high efficiency, low driving voltage, high brightness, and long lifespan.

It should be understood that the embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as being available for other similar features or aspects in other embodiments.

The use of “may” when describing embodiments of the present disclosure refers to “one or more embodiments of the present disclosure”. In addition, as used herein, the terms “use”, “using”, and “used” may be considered synonymous with the terms “utilize”, “utilizing”, and “utilized”, respectively.

As used herein, the terms “substantially”, “about”, and similar terms are used as terms of approximation and not as terms of degree, and are intended to account for the inherent deviations in measured or calculated values that would be recognized by those of ordinary skill in the art.

Also, any numerical range recited herein is intended to include all subranges of the same numerical precision subsumed within the recited range. For example, a range of “1.0 to 10.0” is intended to include all subranges between (and including) the recited minimum value of 1.0 and the recited maximum value of 10.0, that is, having a minimum value equal to or greater than 1.0 and a maximum value equal to or less than 10.0, such as, for example, 2.4 to 7.6. Any maximum numerical limitation recited herein is intended to include all lower numerical limitations subsumed therein and any minimum numerical limitation recited in this specification is intended to include all higher numerical limitations subsumed therein. Accordingly, Applicant reserves the right to amend this specification, including the claims, to expressly recite any sub-range subsumed within the ranges expressly recited herein.


While one or more embodiments have been described with reference to the drawings, it will be understood by those of ordinary skill in the art that various changes in form

127

and details may be made therein without departing from the spirit and scope of the present disclosure as defined by the following claims and equivalents thereof.

What is claimed is:

1. A compound represented by Formula 1:




wherein, in Formula 1,

R₂ and R₃ are each independently selected from hydrogen, deuterium, a halogen atom, a nitro group, a cyano group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

Ar₁ to Ar₃ are each independently selected from hydrogen, deuterium, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, a substituted or unsubstituted C₅-C₆₀ carbocyclic group, and a substituted or unsubstituted C₁-C₆₀ heterocyclic group,

a to c are each independently an integer selected from 0 to 2,

R₁ is selected from Formulae 1-1 and 1-2:




Formula 1-1

128

-continued

Formula 1-2



wherein, in Formulae 1-1 and 1-2,

X₁ to X₅ are each independently selected from nitrogen (N) and CR₁₁,

at least one selected from X₁ to X₅ is N,

R₁₁, Y₁, and Y₂ are each independently selected from a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

L₁ and L₂ are each independently selected from a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

* indicates a binding site, and

at least one substituent selected from the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₂-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted C₅-C₆₀ carbocyclic group, the substituted C₁-C₆₀ heterocyclic group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and/or the substituted divalent non-aromatic condensed heteropolycyclic group, when present, is selected from the group consisting of:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group; a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl

129

group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-\text{N}(\text{Q}_{11})(\text{Q}_{12})$, $-\text{Si}(\text{Q}_{13})(\text{Q}_{14})(\text{Q}_{15})$, and $-\text{B}(\text{Q}_{16})(\text{Q}_{17})$,


a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a $\text{C}_1\text{-C}_{60}$ alkyl group, a $\text{C}_2\text{-C}_{60}$ alkenyl group, a $\text{C}_2\text{-C}_{60}$ alkynyl group, a $\text{C}_1\text{-C}_{60}$ alkoxy group, a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-\text{N}(\text{Q}_{21})(\text{Q}_{22})$, $-\text{Si}(\text{Q}_{23})(\text{Q}_{24})(\text{Q}_{25})$, and $-\text{B}(\text{Q}_{26})(\text{Q}_{27})$,

wherein Q_{11} to Q_{17} and Q_{21} to Q_{27} are each independently selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a $\text{C}_1\text{-C}_{60}$ alkyl group, a $\text{C}_2\text{-C}_{60}$ alkenyl group, a $\text{C}_2\text{-C}_{60}$ alkynyl group, a $\text{C}_1\text{-C}_{60}$ alkoxy group, a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

2. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 2:

Formula 2




wherein Ar_1 , Ar_2 , Ar_3 , R_1 , R_2 , R_3 , a , b , and c are the same as described with respect to Formula 1.

130

3. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 3:


Formula 3



wherein Ar_1 , Ar_2 , Ar_3 , R_1 , R_2 , R_3 , a , b , and c are the same as described with respect to Formula 1.

4. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 4:

Formula 4



wherein R_1 , R_2 , and R_3 are the same as described with respect to Formula 1.

5. The compound of claim 1, wherein R_1 corresponds to Formula 1-1 and X_1 or X_3 is N.


6. The compound of claim 1, wherein R_1 corresponds to Formula 1-1 and X_1 and X_5 are both N.

7. The compound of claim 1, wherein R_1 corresponds to Formula 1-1 and X_1 and X_3 are both N.

8. The compound of claim 1, wherein R_1 corresponds to Formula 1-1 and X_1 , X_3 , and X_5 are all N.

9. The compound of claim 1, wherein R_1 corresponds to a species of Formula 1-2 represented by Formula 1-3:



Formula 1-3



10. The compound of claim 1, wherein R_2 and R_3 are each independently selected from a substituted or unsubstituted $\text{C}_1\text{-C}_{20}$ alkyl group.

131

11. The compound of claim 1, wherein R_{11} corresponds to Formula 1-1, at least one of X_1 to X_5 corresponds to CR_{11} , and R_{11} is represented by one selected from Formulae 2a to 2b:



wherein, in Formulae 2a to 2b,



Z_1 is selected from hydrogen, deuterium, a halogen group, a cyano group, a nitro group, a hydroxyl group, a carboxyl group, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_6 - C_{20} aryl group, a substituted or unsubstituted C_1 - C_{20} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

p is an integer selected from 1 to 7,

when p is 2 or greater, a plurality of Z_1 (s) are identical to or different from each other, and

* indicates a binding site.

12. The compound of claim 1, wherein L_1 and L_2 are each independently represented by one selected from Formulae 3a to 3b:




wherein, in Formulae 3a to 3b, * indicates a binding site.

13. The compound of claim 1, wherein the compound of Formula 1 is one selected from Compounds ET1 to ET42:

132


-continued

ET-2




15

ET-3



30


ET-4



45

ET-5


ET-1




55

60






65



133
-continued




134
-continued



135

-continued

ET-12




5

10

15

20

ET-13




25

30

35

ET-14




40

45

50

ET-15



55


60

65


136

-continued


ET-16




ET-17



ET-18




ET-19



137

-continued


ET-20




138

-continued


ET-24




ET-25




ET-26




ET-27




ET-21



ET-22




ET-23



139

-continued

ET-28




5

10

15

20

ET-29




25

30

35

40

ET-30



55


60

65


140

-continued


ET-31




ET-32



ET-33




ET-34



141

-continued

ET-35




5

10

15

ET-36




20

25

30

ET-37




35

40

45

ET-38



50

55


60

65


142

-continued


ET-39




ET-40



ET-41



ET-42



14. An organic light-emitting device comprising:
 a first electrode;
 a second electrode facing the first electrode; and
 an organic layer between the first electrode and the second
 electrode, the organic layer comprising an emission
 layer,

wherein the organic layer comprises the compound of claim 1.

15. The organic light-emitting device of claim 14, wherein:

the first electrode is an anode,

5

the second electrode is a cathode, and

the organic layer comprises:

i) a hole transport region between the first electrode and the emission layer, the hole transport region comprising at least one selected from a hole transport layer, a hole injection layer, and an electron blocking layer; and

10

ii) an electron transport region between the emission layer and the second electrode, the electron transport region comprising an electron transport layer and at least one selected from a hole blocking layer and an electron injection layer.

15

16. The organic light-emitting device of claim 15, wherein the electron transport region comprises the compound represented by Formula 1.

17. The organic light-emitting device of claim 15, wherein the electron transport layer comprises the compound represented by Formula 1.

20

18. The organic light-emitting device of claim 15, wherein the hole transport region comprises a charge-generation material.

25

19. The organic light-emitting device of claim 18, wherein the charge-generation material is a p-dopant.

20. A display apparatus comprising the organic light-emitting device of claim 14, wherein the first electrode of the organic light-emitting device is electrically connected to a source electrode or a drain electrode of a thin film transistor.


30

* * * * *

专利名称(译)	复合和有机发光器件		
公开(公告)号	US9899610	公开(公告)日	2018-02-20
申请号	US15/174918	申请日	2016-06-06
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	LEE JUNGSUB KIM SEULONG KIM YOUNSUN SHIN DONGWOO ITO NAOYUKI LIM JINO		
发明人	LEE, JUNGSUB KIM, SEULONG KIM, YOUNSUN SHIN, DONGWOO ITO, NAOYUKI LIM, JINO		
IPC分类号	H01L51/00 C07D403/04 C07D403/10 H01L51/50 C07D401/04 H01L27/32 H01L51/56 C07F9/572		
CPC分类号	H01L51/0072 C07D401/04 C07D403/04 C07D403/10 C07F9/5728 H01L51/0058 H01L51/0067 H01L27/3248 H01L51/56 H01L51/506 H01L51/5072 H01L51/0085 H01L2251/308		
优先权	1020150166413 2015-11-26 KR		
其他公开文献	US20170155056A1		
外部链接	Espacenet USPTO		

摘要(译)

一种有机发光装置，包括：第一电极;面向第一电极的第二电极;第一电极和第二电极之间的有机层，有机层包括发光层，其中有机层包括由式1表示的化合物：当式1的化合物用作电子传输材料时，包含该化合物的有机发光装置可具有高效率，低驱动电压，高亮度和长寿命。



Formula 1