



US 20180053902A1

(19) **United States**

(12) **Patent Application Publication**

**KIM et al.**

(10) **Pub. No.: US 2018/0053902 A1**

(43) **Pub. Date: Feb. 22, 2018**

(54) **COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

(52) **U.S. Cl.**  
CPC ..... *H01L 51/0071* (2013.01); *C07D 325/00* (2013.01); *H01L 27/32* (2013.01); *C09K 11/06* (2013.01)


(71) Applicant: **Samsung Display Co., Ltd.**, Yongin-si (KR)

(72) Inventors: **Haejin KIM**, Yongin-si (KR); **Soobyung KO**, Yongin-si (KR); **Mieun JUN**, Yongin-si (KR); **Youngkook KIM**, Yongin-si (KR); **Seokhwan HWANG**, Yongin-si (KR)

(57) **ABSTRACT**

A compound represented by Formula 1 is provided:

<Formula 1>



(21) Appl. No.: **15/650,077**

(22) Filed: **Jul. 14, 2017**

(30) **Foreign Application Priority Data**

Aug. 19, 2016 (KR) ..... 10-2016-0105582

**Publication Classification**

(51) **Int. Cl.**  
*H01L 51/00* (2006.01)  
*C09K 11/06* (2006.01)  
*H01L 27/32* (2006.01)

wherein Formula 1 is the same as described in the detailed description of the present disclosure. The compound of Formula 1 may be included in a hole transport layer and/or an emission layer of an organic light-emitting device (OLED).

10




FIG. 1

10

190
150
110

**COMPOUND AND ORGANIC  
LIGHT-EMITTING DEVICE INCLUDING  
THE SAME**

CROSS-REFERENCE TO RELATED  
APPLICATION

[0001] This application claims the benefit of Korean Patent Application No. 10-2016-0105582 filed on Aug. 19, 2016 in the Korean Intellectual Property Office, the disclosure of which is incorporated herein in its entirety by reference.

BACKGROUND

1. Field

[0002] One or more embodiments relate to a compound and an organic light-emitting device including the same.

2. Description of the Related Art

[0003] Organic light-emitting devices are self-emission devices that produce full-color images, and also have wide viewing angles, high contrast ratios, short response times, as well as excellent characteristics in terms of brightness, driving voltage, and response speed.

[0004] An example of such organic light-emitting devices may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, recombine in the emission layer to produce excitons. These excitons transition from an excited state to a ground state, thereby generating light.


SUMMARY

[0005] An aspect of one or more embodiments provides a compound having a high glass transition temperature and excellent electrical stability, hole transport capability, and light emission efficiency, and introducing the compound to a hole transport layer and an emission layer of an organic light-emitting device.

[0006] Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments.

[0007] According to one or more embodiments, the compound is represented by Formula 1:

<Formula 1>



[0008] In Formula 1,

[0009]  $R_1$ ,  $R_2$ ,  $Ar_1$ , and  $Ar_2$  may each independently be selected from hydrogen, deuterium, halogen, an amino group, a nitro group, a nitrile group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkenyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkynyl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkoxy

group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

[0010]  $L_1$  and  $L_2$  may each independently be selected from a substituted or unsubstituted  $C_6$ - $C_{60}$  arylene group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0011]  $Y_1$  and  $Y_2$  may each independently be selected from N, B, and Si,

[0012] X may be selected from B, C, Si, Ge, Sn, S, SO, S(O)<sub>2</sub>, P, PO, N, and O,

[0013]  $A_1$  and  $A_2$  may each independently be a phenylene group or a naphthylene group,

[0014]  $a_1$ ,  $a_2$ ,  $b_1$ ,  $b_2$ ,  $c_1$ , and  $c_2$  may each independently be an integer from 0 to 3, and

[0015] at least one substituent of the substituted  $C_1$ - $C_{60}$  alkyl group, the substituted  $C_2$ - $C_{60}$  alkenyl group, the substituted  $C_2$ - $C_{60}$  alkynyl group, the substituted  $C_1$ - $C_{60}$  alkoxy group, the substituted  $C_3$ - $C_{10}$  cycloalkyl group, the substituted  $C_2$ - $C_{10}$  heterocycloalkyl group, the substituted  $C_3$ - $C_{10}$  cycloalkenyl group, the substituted  $C_2$ - $C_{10}$  heterocycloalkenyl group, the substituted  $C_6$ - $C_{60}$  aryl group, the substituted  $C_6$ - $C_{60}$  aryloxy group, the substituted  $C_6$ - $C_{60}$  arylthio group, the substituted  $C_1$ - $C_{60}$  heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted  $C_6$ - $C_{60}$  arylene group, the substituted  $C_1$ - $C_{60}$  heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and the substituted divalent non-aromatic condensed heteropolycyclic group may be selected from:

[0016] deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_1$ - $C_{60}$  alkyl group, a  $C_2$ - $C_{60}$  alkenyl group, a  $C_2$ - $C_{60}$  alkynyl group, and a  $C_1$ - $C_{60}$  alkoxy group;

[0017] a  $C_1$ - $C_{60}$  alkyl group, a  $C_2$ - $C_{60}$  alkenyl group, a  $C_2$ - $C_{60}$  alkynyl group, and a  $C_1$ - $C_{60}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazine group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_3$ - $C_{10}$  cycloalkyl group, a  $C_1$ - $C_{10}$  heterocycloalkyl group, a  $C_3$ - $C_{10}$  cycloalkenyl group, a  $C_2$ - $C_{10}$  heterocycloalkenyl group, a  $C_6$ - $C_{60}$  aryl group, a  $C_6$ - $C_{60}$  aryloxy group, a  $C_6$ - $C_{60}$  arylthio group, a  $C_2$ - $C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N( $Q_{11}$ )( $Q_{12}$ ), —Si( $Q_{13}$ )( $Q_{14}$ )( $Q_{15}$ ), and —B( $Q_{16}$ )( $Q_{17}$ );

[0018] a  $C_3$ - $C_{10}$  cycloalkyl group, a  $C_2$ - $C_{10}$  heterocycloalkyl group, a  $C_3$ - $C_{10}$  cycloalkenyl group, a  $C_2$ - $C_{10}$  heterocycloalkenyl group, a  $C_6$ - $C_{60}$  aryl group, a  $C_6$ - $C_{60}$

aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

[0019] a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q<sub>21</sub>)(Q<sub>22</sub>), —Si(Q<sub>23</sub>)(Q<sub>24</sub>)(Q<sub>25</sub>), and —B(Q<sub>26</sub>)(Q<sub>27</sub>),

[0020] wherein Q<sub>11</sub> to Q<sub>17</sub> and Q<sub>21</sub> to Q<sub>27</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0021] According to one or more embodiments, an organic light-emitting device includes: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes the compound described above.

[0022] According to one or more embodiments, a flat panel display apparatus includes the organic light-emitting device, wherein the first electrode of the organic light-emitting device is electrically connected to a source electrode or a drain electrode of a thin film transistor.


#### BRIEF DESCRIPTION OF THE DRAWING

[0023] These and/or other aspects will become apparent and more readily appreciated from the following description of the embodiments, taken in conjunction with FIG. 1 which is a schematic cross-sectional view of an organic light-emitting device according to an embodiment.

#### DETAILED DESCRIPTION

[0024] A compound according to an embodiment may be represented by Formula 1:

<Formula 1>



[0025] In Formula 1,

[0026] R<sub>1</sub>, R<sub>2</sub>, Ar<sub>1</sub>, and Ar<sub>2</sub> may each independently be selected from hydrogen, deuterium, halogen, an amino group, a nitro group, a nitrile group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

[0027] L<sub>1</sub> and L<sub>2</sub> may each independently be selected from a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylene group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0028] Y<sub>1</sub> and Y<sub>2</sub> may each independently be selected from N, B, and Si,

[0029] X may be selected from B, C, Si, Ge, Sn, S, SO, S(O)<sub>2</sub>, P, PO, N, and O,

[0030] A1 and A2 may each independently be a phenylene group or a naphthylene group,

[0031] a1, a2, b1, b2, c1, and c2 may each independently be an integer from 0 to 3, and

[0032] at least one substituent of the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted C<sub>6</sub>-C<sub>60</sub> arylene group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and the substituted divalent non-aromatic condensed heteropolycyclic group may be selected from:

[0033] deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

[0034] a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a

C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q<sub>11</sub>)(Q<sub>12</sub>), —Si(Q<sub>13</sub>)(Q<sub>14</sub>)(Q<sub>15</sub>), and —B(Q<sub>16</sub>)(Q<sub>17</sub>);

[0035] a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

[0036] a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q<sub>21</sub>)(Q<sub>22</sub>), —Si(Q<sub>23</sub>)(Q<sub>24</sub>)(Q<sub>25</sub>), and —B(Q<sub>26</sub>)(Q<sub>27</sub>),

[0037] wherein Q<sub>11</sub> to Q<sub>17</sub> and Q<sub>21</sub> to Q<sub>27</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0038] A blue emission compound having a diphenylanthracene structure in its center and an aryl group substituted at its end, which is a blue emission material, and an organic light-emitting device using the blue emission compound are disclosed. However, luminescent efficiency and brightness of the organic light-emitting device are insufficient.


[0039] An organic light-emitting device using a pyrene-based arylamine compound, various diamine compounds, or a chrysene-based arylamine compound is disclosed. However, blue color purity of the organic light-emitting device is low. Thus, it is difficult to realize a deep blue color. Consequently, there is a limitation in implementing a full-color display capable of producing natural colors.

[0040] The compound represented by Formula 1, according to one or more embodiments, is a compound having a novel structure with distinctly different characteristics in terms of the structure itself as well as a synthesis method. An aspect provides a deep blue compound having high efficiency and a long lifespan by using an arylamine compound including various condensed polycyclic groups.

[0041] Another aspect provides a hole injection layer material and a hole transport layer material using an arylamine compound including a condensed polycyclic group, which has a novel structure based on excellent thermal stability and a fast hole injection speed, and provides an organic light-emitting device using the same and having high efficiency and a long lifespan.

[0042] In one or more embodiments, X in Formula 1 may be selected from B, C, Si, Ge, Sn, S, SO, S(O)<sub>2</sub>, P, and PO.

[0043] In one or more embodiments, R<sub>1</sub> and R<sub>2</sub> in Formula 1 may each independently be represented by one of Formulae 2a and 2b:



[0044] Z<sub>1</sub> in Formulae 2a and 2b may each independently be selected from hydrogen, deuterium, a cyano group, a halogen group, a substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> alkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> alkoxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>20</sub> aryl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

[0045] p is an integer from 1 to 5, and \* indicates a binding site.


[0046] In one or more embodiments, L<sub>1</sub> and L<sub>2</sub> in Formula 1 may each independently be a group represented by Formula 3a:




[0047] \* in Formula 3a indicates a binding site.

[0048] In one or more embodiments, Y<sub>1</sub> and Y<sub>2</sub> in Formula 1 may be N.

[0049] In one or more embodiments, Ar<sub>1</sub> to Ar<sub>4</sub> in Formula 1 may each independently be represented by one of Formulae 4a to 4e:



-continued



4d


4e

[0050] H<sub>1</sub> in Formulae 4a to 4e may be O, S, CR<sub>11</sub>R<sub>12</sub>, or NR<sub>13</sub>,



[0051] R<sub>11</sub> to R<sub>13</sub> and Z<sub>1</sub> may each independently be selected from hydrogen, deuterium, a cyano group, a halogen group, a substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> alkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> alkoxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>20</sub> aryl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

[0052] p may be an integer from 1 to 9, and \* indicates a binding site.


[0053] In one or more embodiments, the compound represented by Formula 1 may be represented by Formula 2:




[0054] In one or more embodiments, the compound represented by Formula 1 may be represented by Formula 3, 4, 5, 6, 7, 8, 9, or 10:




-continued




<Formula 6>




<Formula 7>



<Formula 8>




<Formula 9>



-continued

<Formula 10>




[0055] Substituents and symbols in Formulae 2 to 10 are the same as described above.

[0056] In one or more embodiments, the compound represented by Formula 1 may be one of compounds illustrated below:



-continued

3





1

4




2

5




6

-continued




6

-continued




9

7




10


8



11




-continued




12


-continued




15




13



16




14




17

-continued




18

-continued




21


19




22




20



23



-continued




24


-continued




27




25



28




26




29


-continued




-continued




33




34



32




-continued




35

-continued




37


36



38




-continued




39


-continued



41




40




42

-continued




43


-continued



45




44




46


-continued




-continued



48







50







-continued

-continued







-continued

-continued







-continued

-continued





-continued




-continued




-continued



-continued




-continued




72


-continued




75




73



76




74




77

-continued




78


-continued




81




79



82




80




83

-continued




84


-continued



86




85






87


-continued






-continued



-continued







-continued




-continued

-continued




-continued




100


-continued




103




101



104




102




105

-continued




106


-continued



108







107




109

-continued

-continued




-continued




-continued

116


115

Cc1cc(C)c(N(c2ccc(C)cc2)c3c4ccccc4O3)c(C)c1N(c5ccc(C)cc5)c6ccc(C)cc6


117




-continued




-continued



119







121







-continued

-continued




-continued

-continued




-continued




130

-continued




131


133




132




134



135




-continued





-continued

138




139


137



-continued



-continued



[0057] The term “organic layer” used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of an organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

[0058] FIG. 1 is a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 includes a first electrode 110, an organic layer 150, and a second electrode 190.

[0059] Hereinafter, the structure of an organic light-emitting device according to an embodiment and a method of manufacturing an organic light-emitting device according to an embodiment will be described in connection with FIG. 1.

[0060] In FIG. 1, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be a glass substrate or a transparent plastic substrate, each having excellent mechanical strength,

thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

**[0061]** The first electrode **110** may be formed by depositing or sputtering a material for forming the first electrode **110** on the substrate. When the first electrode **110** is an anode, the material for forming the first electrode **110** may be selected from materials with a high work function to facilitate hole injection. The first electrode **110** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. The material for forming the first electrode **110** may be a transparent and highly conductive material, and examples of such a material are indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO<sub>2</sub>), and zinc oxide (ZnO). In one or more embodiments, to form the first electrode **110** as a semi-transmissible electrode or a reflective electrode, at least one selected from magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag) may be used as a material for forming a first electrode.

**[0062]** The first electrode **110** may have a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode **110** may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode **110** is not limited thereto.

**[0063]** The organic layer **150** is disposed on the first electrode **110**. The organic layer **150** may include an emission layer.

**[0064]** The organic layer **150** may further include a hole transport region between the first electrode and the emission layer, and an electron transport region between the emission layer and the second electrode.

**[0065]** In one or more embodiments, the hole transport region may include at least one selected from a hole transport layer (HTL), a hole injection layer (HIL), a buffer layer, and an electron blocking layer, and an electron transport region may include at least one selected from a hole blocking layer (HBL), an electron transport layer, and an electron injection layer (EIL). However, it may be understood that embodiments are not limited thereto.

**[0066]** The hole transport region may have a single-layered structure formed of a single material, a single-layered structure formed of a plurality of different materials, or a multi-layered structure having a plurality of layers formed of a plurality of different materials.

**[0067]** For example, the hole transport region may have a single-layered structure formed of a plurality of different materials, or a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/buffer layer structure, a hole injection layer/buffer layer structure, a hole transport layer/buffer layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure, wherein layers of each structure are sequentially stacked from the first electrode **110** in this stated order, but are not limited thereto.

**[0068]** When the hole transport region includes a hole injection layer, the hole injection layer may be formed on the first electrode **110** by using one or more suitable methods selected from vacuum deposition, spin coating, casting,


Langmuir-Blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

**[0069]** When a hole injection layer is formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of about 100° C. to about 500° C., at a vacuum degree of about 10<sup>-8</sup> torr to about 10<sup>-3</sup> torr, and at a deposition rate of about 0.01 Å/sec to about 100 Å/sec by taking into account the compound for the hole injection layer to be deposited, and the structure of the hole injection layer to be formed.

**[0070]** When a hole injection layer is formed by spin coating, the spin coating may be performed at a coating rate of about 2,000 rpm to about 5,000 rpm, and at a temperature of about 80° C. to about 200° C. by taking into account the compound for the hole injection layer to be deposited, and the structure of the hole injection layer to be formed.


**[0071]** When the hole transport region includes a hole transport layer, the hole transport layer may be formed on the first electrode **110** or the hole injection layer by using one or more suitable methods selected from vacuum deposition, spin coating, casting, a LB method, ink-jet printing, laser-printing, and laser-induced thermal imaging. When the hole transport layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the hole transport layer may be the same as the deposition and coating conditions for the hole injection layer.

**[0072]** The hole transport region may include m-MT-DATA, TDATA, 2-TNATA, NPB, β-NPB, TPD, Spiro-TPD, Spiro-NPB, alpha-NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201 below, a compound represented by Formula 202 below, or the compound of Formula 1 according to an embodiment of the present disclosure.




m-MT-DATA


-continued



TDATA




2-TNATA




NPB


-continued




$\beta$ -NPB




TPD



Spiro-TPD



Spiro-NPB




$\alpha$ -NPB




dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group, but they are not limited thereto.


[0087] The compound represented by Formula 201 may be represented by Formula 201A:



[0088] In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A-1 below, but embodiments of the present disclosure are not limited thereto:



[0089] For example, the compound represented by Formula 202 may be represented by Formula 202A below, but embodiments of the present disclosure are not limited thereto:



[0090]  $L_{201}$  to  $L_{203}$ ,  $x_{a1}$  to  $x_{a3}$ ,  $x_{a5}$ , and  $R_{202}$  to  $R_{204}$  in Formulae 201A, 201A-1, and 202A are already described in detail above, and  $R_{211}$  may be understood by referring to the description provided herein in connection with  $R_{203}$ , and  $R_{213}$  to  $R_{216}$  may each independently be selected from

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_1$ - $C_{60}$  alkyl group, a  $C_2$ - $C_{60}$  alkenyl group, a  $C_2$ - $C_{60}$  alkynyl group, a  $C_1$ - $C_{60}$  alkoxy group, a  $C_3$ - $C_{10}$  cycloalkyl group, a  $C_2$ - $C_{10}$  heterocycloalkyl group, a  $C_3$ - $C_{10}$  cycloalkenyl group, a  $C_2$ - $C_{10}$  heterocycloalkenyl group, a  $C_6$ - $C_{60}$  aryl group, a  $C_6$ - $C_{60}$  aryloxy group, a  $C_6$ - $C_{60}$  arylthio group, a  $C_1$ - $C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0091] For example, in Formulae 201A, 201A-1, and 202A,

[0092]  $L_{201}$  to  $L_{203}$  may each independently be selected from:

[0093] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylylene group, a carbazolylene group, and a triazinylene group; and

[0094] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylylene group, a carbazolylene group, and a triazinylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group;

[0095]  $x_{a1}$  to  $x_{a3}$  may each independently be 0 or 1;

[0096]  $R_{203}$ ,  $R_{211}$ , and  $R_{212}$  may each independently be selected from:

[0097] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group; and

[0098] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group,



an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group; and


[0109]  $x_{a5}$  is 1 or 2.

[0110]  $R_{213}$  and  $R_{214}$  in Formulae 201A and 201A-1 may be linked to each other to form a saturated or unsaturated ring.


[0111] The compound represented by Formula 201, and the compound represented by Formula 202 may each include compounds HT1 to HT20 illustrated below, but are not limited thereto.

-continued


HT3




HT1



HT4



HT2





-continued

-continued



HT5

HT7




HT6

HT8




-continued



HT9



-continued




HT11

HT12





HT10



HT13






-continued



-continued

HT18




[0112] A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes a hole injection layer and a hole transport layer, the thickness of the hole injection layer may be in a range of about 100 Å to about 10,000 Å, and for example, about 100 Å to about 1,000 Å, and the thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, and for example, about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.


[0113] The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-

generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0114] The charge-generation material may be, for example, a p-dopant. The p-dopant may be one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but embodiments are not limited thereto. For example, non-limiting examples of the p-dopant are a quinone derivative, such as tetracyanoquinonedimethane (TCNQ) or 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonedimethane (F4-TCNQ); a metal oxide, such as a tungsten oxide or a molybdenum oxide, and Compound HT-D1 illustrated below, but are not limited thereto.



&lt;Compound HT-D1&gt;



&lt;F4-TCNQ&gt;

[0115] The hole transport region may further include a buffer layer, in addition to an electron blocking layer, a hole injection layer, and a hole transport layer. Since the buffer layer may compensate for an optical resonance distance according to a wavelength of light emitted from the emission layer, light emission efficiency of a formed organic light-emitting device may be improved. For use as a material included in the buffer layer, materials that are to be included in the hole transport region may be used. The electron blocking layer prevents injection of electrons from the electron transport region.


[0116] An emission layer is formed on the first electrode 110 or the hole transport region by using one or more suitable methods selected from vacuum deposition, spin coating, casting, a LB method, ink-jet printing, laser-printing, and laser-induced thermal imaging. When an emission layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the emission layer may be the same as those for the hole injection layer.

[0117] When the organic light-emitting device 10 is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In one or more embodiments, the emission layer may have a stacked structure including a red emission layer, a green emission layer, and a blue emission layer, or may include a


red light-emitting material, a green light-emitting material, and a blue light-emitting material, which are mixed with each other in a single layer, to emit white light.

[0118] The emission layer may include a host and a dopant.


[0119] For example, the host may include at least one selected from TPBi, TBADN, ADN (also referred to as "DNA"), CBP, CDBP, and TCP:




TPBi




TBADN



ADN



CBP



[0120] In one or more embodiments, the host may include a compound represented by Formula 301 below.



[0121] In Formula 301,

[0122]  $\text{Ar}_{301}$  may be selected from:

[0123] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene; and

[0124] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $\text{C}_1$ - $\text{C}_{60}$  alkyl group, a  $\text{C}_2$ - $\text{C}_{60}$  alkenyl group, a  $\text{C}_2$ - $\text{C}_{60}$  alkynyl group, a  $\text{C}_1$ - $\text{C}_{60}$  alkoxy group, a  $\text{C}_3$ - $\text{C}_{10}$  cycloalkyl group, a  $\text{C}_2$ - $\text{C}_{10}$  heterocycloalkyl group, a  $\text{C}_3$ - $\text{C}_{10}$  cycloalkenyl group, a  $\text{C}_2$ - $\text{C}_{10}$  heterocycloalkenyl group, a  $\text{C}_6$ - $\text{C}_{60}$  aryl group, a  $\text{C}_6$ - $\text{C}_{60}$  aryloxy group, a  $\text{C}_6$ - $\text{C}_{60}$  arylthio group, a  $\text{C}_1$ - $\text{C}_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, monovalent non-aromatic condensed heteropolycyclic group, and —Si( $\text{Q}_{301}$ )( $\text{Q}_{302}$ )( $\text{Q}_{303}$ ) ( $\text{Q}_{301}$  to  $\text{Q}_{303}$  may each independently be selected from hydrogen, a  $\text{C}_1$ - $\text{C}_{60}$  alkyl group, a  $\text{C}_2$ - $\text{C}_{60}$  alkenyl group, a  $\text{C}_6$ - $\text{C}_{60}$  aryl group, and a  $\text{C}_1$ - $\text{C}_{60}$  heteroaryl group);

[0125]  $\text{L}_{301}$  may be selected from a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorene group, a dibenzofluorene

group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylenylene group, a carbazolylene group, and a triazinylene group; and [0126] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorene group, a dibenzofluorene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylenylene group, a carbazolylene group, and a triazinylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $\text{C}_1$ - $\text{C}_{20}$  alkyl group, a  $\text{C}_1$ - $\text{C}_{20}$  alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinylyl group, a carbazolyl group, and a triazinyl group;

[0127]  $\text{R}_{301}$  may be selected from:

[0128] a  $\text{C}_1$ - $\text{C}_{20}$  alkyl group and a  $\text{C}_1$ - $\text{C}_{20}$  alkoxy group;

[0129] a  $\text{C}_1$ - $\text{C}_{20}$  alkyl group and a  $\text{C}_1$ - $\text{C}_{20}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinylyl group, a carbazolyl group, and a triazinyl group;

[0130] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinylyl group, a carbazolyl group, and a triazinyl group; and

[0131] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinylyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a

carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and

[0132]  $xb1$  may be selected from 0, 1, 2, and 3; and

[0133]  $xb2$  may be selected from 1, 2, 3, and 4.

[0134] For example, in Formula 301,

[0135]  $L_{301}$  may be selected from a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, and a chrysenylene group; and

[0136] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, and a chrysenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group;

[0137]  $R_{301}$  may be selected from:

[0138] a  $C_1$ - $C_{20}$  alkyl group and a  $C_1$ - $C_{20}$  alkoxy group;


[0139] a  $C_1$ - $C_{20}$  alkyl group and a  $C_1$ - $C_{20}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group;

[0140] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group; and

[0141] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a

dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group, but embodiments of the present disclosure are not limited thereto.

[0142] For example, the host may include a compound represented by Formula 301A below:




[ $L_{301}$ ]<sub>xb1</sub>— $R_{301}$ ]<sub>xb2</sub>


<Formula 301A>

[0143] Substituents of Formula 301A may be understood by referring to the descriptions provided herein.


[0144] The compound represented by Formula 301 may include at least one of Compounds H1 to H42, but embodiments of the present disclosure are not limited thereto:



H1




H2




H3


-continued




H4




H5




H6



H7




H8




H9


-continued




H10




H11




H12



H13











H14








H15

-continued






-continued




-continued




-continued




-continued




-continued




H35




H38



H36




H39




-continued


H40




H43




H41




H44



H42




H45



[0145] In one or more embodiments, the host may include at least one of Compounds H43 to H49 below, but embodiments of the present disclosure are not limited thereto:

-continued




H46

[0146] The dopant may include a known fluorescent dopant, a known phosphorescent dopant, or the compound of Formula 1 of according to an embodiment of the present disclosure.

[0147] The phosphorescent dopant may include an organometallic complex represented by Formula 401 below:

&lt;Formula 401&gt;



H47



[0148] In Formula 401,

[0149] M may be selected from iridium (Ir), platinum (Pt), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), and thulium (Tm);


[0150] X<sub>401</sub> to X<sub>404</sub> may each independently be nitrogen or carbon;

[0151] A<sub>401</sub> and A<sub>402</sub> rings may each independently be selected from a substituted or unsubstituted benzene, a substituted or unsubstituted naphthalene, a substituted or unsubstituted fluorene, a substituted or unsubstituted spirofluorene, a substituted or unsubstituted indene, a substituted or unsubstituted pyrrole, a substituted or unsubstituted thiophene, a substituted or unsubstituted furan, a substituted or unsubstituted imidazole, a substituted or unsubstituted pyrazole, a substituted or unsubstituted thiazole, a substituted or unsubstituted isothiazole, a substituted or unsubstituted oxazole, a substituted or unsubstituted isoxazole, a substituted or unsubstituted pyridine, a substituted or unsubstituted pyrazine, a substituted or unsubstituted pyrimidine, a substituted or unsubstituted pyridazine, a substituted or unsubstituted quinoline, a substituted or unsubstituted isoquinoline, a substituted or unsubstituted benzoquinoline, a substituted or unsubstituted quinoxaline, a substituted or unsubstituted quinazoline, a substituted or unsubstituted carbazol, a substituted or unsubstituted benzimidazole, a substituted or unsubstituted benzofuran, a substituted or unsubstituted benzothiophene, a substituted or unsubstituted isobenzothiophene, a substituted or unsubstituted benzoxazole, a substituted or unsubstituted isobenzoxazole, a substituted or unsubstituted triazole, a substituted or unsubstituted oxadiazole, a substituted or unsubstituted triazine, a substituted or unsubstituted dibenzofuran, and a substituted or unsubstituted dibenzothiophene; and


[0152] at least one substituent of the substituted benzene, substituted naphthalene, substituted fluorene, substituted spirofluorene, substituted indene, substituted pyrrole, substituted thiophene, substituted furan, substituted imidazole, substituted pyrazole, substituted thiazole, substituted isothiazole, substituted oxazole, substituted isoxazole, substituted pyridine, substituted pyrazine, substituted pyrimidine, substituted pyridazine, substituted quinoline, substituted isoquinoline, substituted benzoquinoline, substituted quinoxaline,



H48



H49



substituted quinazoline, substituted carbazol, substituted benzimidazole, substituted benzofuran, substituted benzothiophene, substituted isobenzothiophene, substituted benzoxazole, substituted isobenzoxazole, substituted triazole, substituted oxadiazole, substituted triazine, substituted dibenzofuran, and substituted dibenzothiophene may be selected from:

**[0153]** deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

**[0154]** a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q<sub>401</sub>)(Q<sub>402</sub>), —Si(Q<sub>403</sub>)(Q<sub>404</sub>)(Q<sub>405</sub>), and —B(Q<sub>406</sub>)(Q<sub>407</sub>);

**[0155]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, and a non-aromatic condensed polycyclic group;

**[0156]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>1</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q<sub>411</sub>)(Q<sub>412</sub>), —Si(Q<sub>413</sub>)(Q<sub>414</sub>)(Q<sub>415</sub>) and —B(Q<sub>416</sub>)(Q<sub>417</sub>); and

**[0157]** —N(Q<sub>421</sub>)(Q<sub>422</sub>), —Si(Q<sub>423</sub>)(Q<sub>424</sub>)(Q<sub>425</sub>), and —B(Q<sub>426</sub>)(Q<sub>427</sub>);

**[0158]** L<sub>401</sub> is an organic ligand;

**[0159]** xc1 is 1, 2, or 3; and

**[0160]** xc2 is 0, 1, 2, or 3.


**[0161]** L<sub>401</sub> may be a monovalent, divalent, or trivalent organic ligand. For example, L<sub>401</sub> may be selected from a halogen ligand (for example, Cl or F), a diketone ligand (for example, acetylacetonate, 1,3-diphenyl-1,3-propanedionate,

2,2,6,6-tetramethyl-3,5-heptanedionate, or hexafluoroacetate), a carboxylic acid ligand (for example, picolinate, dimethyl-3-pyrazolecarboxylate, or benzoate), a carbon monoxide ligand, an isonitrile ligand, a cyano ligand, and a phosphorous ligand (for example, phosphine, and phosphite), but embodiments of the present disclosure are not limited thereto.

**[0162]** When A<sub>401</sub> in Formula 401 has two or more substituents, the substituents of A<sub>401</sub> may be linked to each other to form a saturated or unsaturated ring.


**[0163]** When A<sub>401</sub> in Formula 402 has two or more substituents, the substituents of A<sub>402</sub> may be linked to each other to form a saturated or unsaturated ring.

**[0164]** When xc1 in Formula 401 is two or more, a plurality of ligands




in Formula 401 may be identical or different. When xc1 in Formula 401 is two or more, A<sub>401</sub> and A<sub>402</sub> may be respectively directly connected to A<sub>401</sub> and A<sub>402</sub> of other neighboring ligands with or without a linker (for example, a C<sub>1</sub>-C<sub>5</sub> alkylene, or —N(R')— (wherein R' may be a C<sub>1</sub>-C<sub>10</sub> alkyl group or a C<sub>6</sub>-C<sub>20</sub> aryl group) or —C(=O)—) therebetween.

**[0165]** The phosphorescent dopant may include at least one of Compounds PD1 to PD74 below, but embodiments of the present disclosure are not limited thereto:








PD1




PD2

-continued




-continued

PD3




PD8

PD4




PD9

PD5




PD10

PD6




PD11

PD7




PD12


-continued




PD13




PD18




PD14




PD19




PD15




PD20



PD16




PD17




PD21


-continued




PD22




PD23




PD24



PD25




PD26




PD27


-continued




PD28




PD29



PD30








PD31




PD32

-continued




-continued

PD33




PD38

PD34




PD39

PD35




PD40

PD36








PD41

PD37




PD42

-continued




-continued

PD43




PD48

PD44




PD49

PD45




PD50

PD46








PD51

PD47



PD52


-continued



-continued


PD53

PD58




PD54

PD59




PD55

PD60




PD61

PD56




PD62


PD57




-continued




PD63




PD64



PD65




PD66




PD67


-continued




PD68




PD69



PD70




PD71



PD72


-continued




PD73

[0166] In one or more embodiments, the phosphorescent dopant may include PtOEP:

PtOEP




PD74




[0167] The fluorescent dopant may include at least one selected from DPAVBi, BDAVBi, TBPe, DCM, DCJTb, Coumarin 6, and C545T.


DPVBi




DPAVBi




TBPe




DCM




-continued  
DCJTB


Coumarin 6



C545T



[0168] In one or more embodiments, the fluorescent dopant may include a compound represented by Formula 501 below.



&lt;Formula 501&gt;

[0169] In Formula 501,

[0170]  $\text{Ar}_{501}$  may be selected from:

[0171] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene; and

[0172] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $\text{C}_1$ - $\text{C}_{60}$  alkyl group, a  $\text{C}_2$ - $\text{C}_{60}$  alkenyl group, a  $\text{C}_2$ - $\text{C}_{60}$  alkynyl group, a  $\text{C}_1$ - $\text{C}_{60}$  alkoxy group, a  $\text{C}_3$ - $\text{C}_{10}$  cycloalkyl group, a  $\text{C}_2$ - $\text{C}_{10}$  heterocycloalkyl group, a  $\text{C}_3$ - $\text{C}_{10}$  cycloalkenyl group, a  $\text{C}_2$ - $\text{C}_{10}$  heterocycloalkenyl group, a  $\text{C}_6$ - $\text{C}_{60}$  aryl group, a  $\text{C}_6$ - $\text{C}_{60}$  aryloxy group, a  $\text{C}_6$ - $\text{C}_{60}$  arylthio

group, a  $\text{C}_1$ - $\text{C}_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, monovalent non-aromatic condensed heteropolycyclic group, and —Si( $\text{Q}_{501}$ )( $\text{Q}_{502}$ )( $\text{Q}_{503}$ ) ( $\text{Q}_{501}$  to  $\text{Q}_{503}$  may each independently be selected from hydrogen, a  $\text{C}_1$ - $\text{C}_{60}$  alkyl group, a  $\text{C}_2$ - $\text{C}_{60}$  alkenyl group, a  $\text{C}_6$ - $\text{C}_{60}$  aryl group, and a  $\text{C}_1$ - $\text{C}_{60}$  heteroaryl group);

[0173] Descriptions of  $\text{L}_{501}$  to  $\text{L}_{503}$  are the same as the descriptions provided herein in connection with  $\text{L}_{203}$ ;

[0174]  $\text{R}_{501}$  and  $\text{R}_{502}$  may each independently be selected from:

[0175] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and


[0176] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof,

a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuran-yl group, and a dibenzothiophenyl group;


[0177] xd1 to xd3 may each independently be selected from 0, 1, 2, and 3; and

[0178] xb4 may be selected from 1, 2, 3, and 4.


[0179] The fluorescent dopant may include at least one of Compounds FD1 to FD8:




FD1



FD2






FD3




FD4

-continued

-continued



-continued



[0180] An amount of the dopant in the emission layer may be, in general, in a range of about 0.01 parts to about 15 parts by weight based on 100 parts by weight of the host, but embodiments of the present disclosure are not limited thereto.


[0181] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0182] Then, an electron transport region may be disposed on the emission layer.


[0183] The electron transport region may include at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer, but embodiments of the present disclosure are not limited thereto.

[0184] When the electron transport region includes a hole blocking layer, the hole blocking layer may be formed on the emission layer by using one or more suitable methods selected from vacuum deposition, spin coating, casting, a Langmuir-Blodgett (LB) deposition, ink-jet printing, laser-printing, and/or laser-induced thermal imaging. When the hole blocking layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the hole blocking layer may be determined by referring to the deposition and coating conditions for the hole injection layer.

[0185] The hole blocking layer may include, for example, at least one of BCP and Bphen, but embodiments of the present disclosure are not limited thereto.




-continued



Bphen

-continued




TAZ

[0186] A thickness of the hole blocking layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thickness of the hole blocking layer is within these ranges, the hole blocking layer may have excellent hole blocking characteristics without a substantial increase in driving voltage.

[0187] For example, the electron transport region may have an electron transport layer/electron injection layer structure or a hole blocking layer/electron transport layer/electron injection layer structure, wherein layers of each structure are sequentially stacked from the emission layer in the stated order, but the structure thereof is not limited thereto.

[0188] According to an embodiment, the organic layer 150 of the organic light-emitting device includes an electron transport region between the emission layer and the second electrode 190, and the electron transport region may include an electron transport layer. The electron transport layer may include a plurality of layers. For example, the electron transport layer may include a first electron transport layer and a second electron transport layer.

[0189] The electron transport layer may further include at least one selected from BCP, Bphen, Alq<sub>3</sub>, BAIq, TAZ, and NTAZ.



NTAZ

[0190] In one or more embodiments, the electron transport layer may include at least one compound selected from a compound represented by Formula 601 and a compound represented by Formula 602 illustrated below:



[0191] In Formula 601,

[0192] Ar<sub>601</sub> may be selected from:


[0193] a naphthalene, a heptalene, a fluorene, a spirofluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene;

[0194] a naphthalene, a heptalene, a fluorene, a spirofluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>3</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, monovalent non-aromatic condensed heteropolycyclic group, and —Si(Q<sub>301</sub>)(Q<sub>302</sub>)(Q<sub>303</sub>) (Q<sub>301</sub> to Q<sub>303</sub> may each independently be selected from hydrogen, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, and a C<sub>1</sub>-C<sub>60</sub> heteroaryl group);


[0195] a description of L<sub>601</sub> may be understood by referring to the description provided in connection with L<sub>203</sub>;

[0196] E<sub>601</sub> may be selected from:

[0197] a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a qui-

Alq<sub>3</sub>

BAIq



nolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranlyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thia-diazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group; and


[0198] a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranlyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thia-diazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranlyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl

group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thia-diazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0199] xe1 may be selected from 0, 1, 2, and 3; and

[0200] xe2 may be selected from 1, 2, 3, and 4.

<Formula 602>



[0201] In Formula 602,

[0202] X<sub>611</sub> may be N or C-(L<sub>611</sub>)<sub>xe611</sub>-R<sub>611</sub>, X<sub>612</sub> may be N or C-(L<sub>612</sub>)<sub>xe612</sub>-R<sub>612</sub>, X<sub>613</sub> may be N or C-(L<sub>613</sub>)<sub>xe613</sub>-R<sub>613</sub>, and at least one selected from X<sub>611</sub> to X<sub>613</sub> may be N;

[0203] L<sub>611</sub> to L<sub>616</sub> may be understood by referring to the description provided herein in connection with L<sub>203</sub>;

[0204] R<sub>611</sub> to R<sub>616</sub> may each independently be selected from:

[0205] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and


[0206] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and

[0207] xe611 to xe616 may each independently be selected from 0, 1, 2, and 3.


[0208] The compound represented by Formula 601 and the compound represented by Formula 602 may each be selected from Compounds ET1 to ET15 illustrated below.

-continued


ET1




ET4




ET2




ET5



ET3



ET6







-continued

-continued



ET14

ET11




ET15

ET12



ET13




**[0209]** A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

**[0210]** Also, the electron transport layer may further include, in addition to the materials described above, a metal-containing material.


**[0211]** The metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.

ET-D1



-continued

ET-D2



[0212] The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode **190**. The electron injection layer may be formed on the electron transport layer by using one or more suitable methods selected from vacuum deposition, spin coating, casting, a LB method, ink-jet printing, laser-printing, and laser-induced thermal imaging. When an electron injection layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the electron injection layer may be the same as those for the hole injection layer.

[0213] The electron injection layer may include at least one selected from LiF, NaCl, CsF, Li<sub>2</sub>O, BaO, and LiQ.

[0214] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

[0215] The second electrode **190** may be disposed on the organic layer **150** having such a structure. The second electrode **190** may be a cathode which is an electron injection electrode, and in this regard, a material for forming the second electrode **190** may be selected from metal, an alloy, an electrically conductive compound, and a mixture thereof, which have a relatively low work function. Examples of the second electrode **190** are lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag). In one or more embodiments, the material for forming the second electrode **190** may be ITO or IZO. The second electrode **190** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode.

[0216] Also, an organic layer according to an embodiment may be formed by depositing the compound according to an embodiment, or may be formed by using a wet method in which the compound according to an embodiment is prepared in the form of solution and then the solution of the compound is used for coating.

[0217] An organic light-emitting device according to an embodiment may be used in various flat panel display apparatuses, such as a passive matrix organic light-emitting display apparatus or an active matrix organic light-emitting display apparatus. For example, when the organic light-emitting device is included in an active matrix organic light-emitting display apparatus, a first electrode disposed on a substrate acts as a pixel and may be electrically connected to a source electrode or a drain electrode of a thin

film transistor. In addition, the organic light-emitting device may be included in a flat panel display apparatus that emits light in opposite directions.

[0218] Hereinbefore, the organic light-emitting device has been described with reference to FIG. 1, but embodiments of the present disclosure are not limited thereto.

[0219] Hereinafter, definitions of substituents of compounds used herein will be presented (the number of carbon atoms used to restrict a substituent is not limited, and does not limit properties of the substituent, and unless defined otherwise, the definition of the substituent is consistent with a general definition thereof).

[0220] The term “C<sub>1</sub>-C<sub>60</sub> alkyl group” as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “C<sub>1</sub>-C<sub>60</sub> alkylene group” as used herein refers to a divalent group having the same structure as the C<sub>1</sub>-C<sub>60</sub> alkyl group.

[0221] The term “C<sub>1</sub>-C<sub>60</sub> alkoxy group” as used herein refers to a monovalent group represented by —OA<sub>101</sub> (wherein A<sub>101</sub> is the C<sub>1</sub>-C<sub>60</sub> alkyl group), and examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

[0222] The term “C<sub>2</sub>-C<sub>60</sub> alkenyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon double bond in the middle or at the terminus of the C<sub>2</sub>-C<sub>60</sub> alkyl group, and examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C<sub>2</sub>-C<sub>60</sub> alkenylene group” as used herein refers to a divalent group having the same structure as the C<sub>2</sub>-C<sub>60</sub> alkenyl group.

[0223] The term “C<sub>2</sub>-C<sub>60</sub> alkynyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the middle or at the terminus of the C<sub>2</sub>-C<sub>60</sub> alkyl group, and examples thereof include an ethynyl group and a propynyl group. The term “C<sub>2</sub>-C<sub>60</sub> alkynylene group” as used herein refers to a divalent group having the same structure as the C<sub>2</sub>-C<sub>60</sub> alkynyl group.

[0224] The term “C<sub>3</sub>-C<sub>10</sub> cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C<sub>3</sub>-C<sub>10</sub> cycloalkylene group” as used herein refers to a divalent group having the same structure as the C<sub>3</sub>-C<sub>10</sub> cycloalkyl group.

[0225] The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group,” used herein, refers to a monovalent saturated monocyclic group having at least one heteroatom selected from N, O, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a tetrahydrofuranlyl group, and a tetrahydrothiophenyl group. The term “C<sub>2</sub>-C<sub>10</sub> heterocycloalkylene group,” used herein, refers to a divalent group having the same structure as the C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group.

[0226] The term “C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group,” used herein, refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and no aromaticity, and examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C<sub>3</sub>-C<sub>10</sub>

cycloalkenylene group,” used herein, refers to a divalent group having the same structure as the C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group.

**[0227]** The term “C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group,” used herein, refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, P, and S as a ring-forming atom, 2 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Examples of the C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group include a 2,3-dihydrofuran-2-yl group and a 2,3-dihydrothiophen-2-yl group. The term “C<sub>2</sub>-C<sub>10</sub> heterocycloalkenylene group,” used herein, refers to a divalent group having the same structure as the C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group.

**[0228]** The term “C<sub>6</sub>-C<sub>60</sub> aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and the term “C<sub>6</sub>-C<sub>60</sub> arylene group” as used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Examples of the C<sub>6</sub>-C<sub>60</sub> aryl group include a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C<sub>6</sub>-C<sub>60</sub> aryl group and the C<sub>6</sub>-C<sub>60</sub> arylene group each include two or more rings, the rings may be fused to each other.

**[0229]** The term “C<sub>1</sub>-C<sub>60</sub> heteroaryl group,” used herein, refers to a monovalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. A C<sub>1</sub>-C<sub>60</sub> heteroarylene group used herein refers to a divalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Examples of the C<sub>1</sub>-C<sub>60</sub> heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C<sub>1</sub>-C<sub>60</sub> heteroaryl group and the C<sub>1</sub>-C<sub>60</sub> heteroarylene group each include two or more rings, the rings may be fused to each other.

**[0230]** The term “C<sub>6</sub>-C<sub>60</sub> aryloxy group,” used herein, indicates —OA<sub>102</sub> (wherein A<sub>102</sub> is the C<sub>6</sub>-C<sub>60</sub> aryl group), and a C<sub>6</sub>-C<sub>60</sub> arylthio group indicates —SA<sub>103</sub> (wherein A<sub>103</sub> is the C<sub>6</sub>-C<sub>60</sub> aryl group).

**[0231]** The term “monovalent non-aromatic condensed polycyclic group,” used herein, refers to a monovalent group (for example, having 8 to 60 carbon atoms) having two or more rings condensed to each other, only carbon atoms as ring-forming atoms, and no aromaticity in its entire molecular structure. Examples of the monovalent non-aromatic condensed polycyclic group include a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group,” used herein, refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

**[0232]** The term “monovalent non-aromatic condensed heteropolycyclic group,” used herein, refers to a monovalent group (for example, having 2 to 60 carbon atoms) having two or more rings condensed to each other, has a heteroatom selected from N, O, P, and S, other than carbon atoms, as a ring-forming atom, and has no aromaticity in its entire molecular structure. Non-limiting examples of the monovalent non-aromatic condensed heteropolycyclic group include a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group,” used herein, refers to a

divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

**[0233]** In the present specification, at least one substituent of the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, the substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkylene group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, the substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenylene group, the substituted C<sub>6</sub>-C<sub>60</sub> arylene group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

**[0234]** deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

**[0235]** a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q<sub>11</sub>)(Q<sub>12</sub>), —Si(Q<sub>13</sub>)(Q<sub>14</sub>)(Q<sub>15</sub>), and —B(Q<sub>16</sub>)(Q<sub>17</sub>);

**[0236]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

**[0237]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>1</sub>-C<sub>60</sub> alkynyl group,

a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q<sub>21</sub>)(Q<sub>22</sub>), —Si(Q<sub>23</sub>)(Q<sub>24</sub>)(Q<sub>25</sub>) and —B(Q<sub>26</sub>)(Q<sub>27</sub>);

**[0238]** —N(Q<sub>31</sub>)(Q<sub>32</sub>), —Si(Q<sub>33</sub>)(Q<sub>34</sub>)(Q<sub>35</sub>), and —B(Q<sub>36</sub>)(Q<sub>37</sub>); and

**[0239]** Q<sub>11</sub> to Q<sub>17</sub>, Q<sub>21</sub> to Q<sub>27</sub> and Q<sub>31</sub> to Q<sub>37</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

**[0240]** For example, at least one substituent of the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, the substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkylene group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, the substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenylene group, the substituted C<sub>6</sub>-C<sub>60</sub> arylene group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

**[0241]** deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

**[0242]** a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl

group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —N(Q<sub>11</sub>)(Q<sub>12</sub>), —Si(Q<sub>13</sub>)(Q<sub>14</sub>)(Q<sub>15</sub>), and —B(Q<sub>16</sub>)(Q<sub>17</sub>);

**[0243]** a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

**[0244]** a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl

group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyly group, a pentacenyly group, a rubicenyly group, a coronenyly group, an ovalenyly group, a pyrrolyly group, a thiophenyly group, a furanyly group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranly group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranly group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyly group, a heptalenyl group, an indacenyly group, an acenaphthyl group, a fluorenyly group, a spiro-fluorenyly group, a benzofluorenyly group, a dibenzofluorenyly group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyly group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyly group, a pentacenyly group, a rubicenyly group, a coronenyly group, an ovalenyly group, a pyrrolyly group, a thiophenyly group, a furanyly group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranly group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranly group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, —N(Q<sub>21</sub>)(Q<sub>22</sub>), —Si(Q<sub>23</sub>)(Q<sub>24</sub>)(Q<sub>25</sub>), and —B(Q<sub>26</sub>)(Q<sub>27</sub>);

[0245] —N(Q<sub>31</sub>)(Q<sub>32</sub>), —Si(Q<sub>33</sub>)(Q<sub>34</sub>)(Q<sub>35</sub>) and —B(Q<sub>36</sub>)(Q<sub>37</sub>); and

[0246] Q<sub>11</sub> to Q<sub>17</sub>, Q<sub>21</sub> to Q<sub>27</sub>, and Q<sub>31</sub> to Q<sub>37</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro

group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyly group, a heptalenyl group, an indacenyly group, an acenaphthyl group, a fluorenyly group, a spiro-fluorenyly group, a benzofluorenyly group, a dibenzofluorenyly group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyly group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyly group, a pentacenyly group, a rubicenyly group, a coronenyly group, an ovalenyly group, a pyrrolyly group, a thiophenyly group, a furanyly group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranly group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranly group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.


[0247] The expression “Ph” used herein refers to a phenyl group, the expression “Me” used herein refers to a methyl group, the expression “Et” used herein refers to an ethyl group, and the expression “ter-Bu” or “Bu” used herein refers to a tert-butyl group.


[0248] Hereinafter, an organic light-emitting device according to an embodiment will be described in detail with reference to Synthesis Examples and Examples.

### Synthesis Example

#### Synthesis Example: Synthesis of Compound 8

[0249]





8

#### Synthesis of Intermediate A-1

**[0250]** As a starting material, 1.58 g (5.0 mmol) of 1-bromo-4-chloro-2-iodobenzene, 916 g (5.5 mmol) of 4-chloro-1-ethynyl-2-methoxybenzene, 70 mg (0.1 mmol) of  $\text{PdCl}_2(\text{PPh}_3)_2$ , 19 mg (0.05 mmol) of  $\text{CuI}$ , and 20 mL of  $\text{Et}_3\text{N}$  were stirred at room temperature for 24 hours in a

nitrogen atmosphere. When the reaction was completed, an organic layer was extracted three times by using diethyl ether. The extracted organic layer collected therefrom was dried by using magnesium sulfate and a solvent was evaporated. The residue obtained therefrom was purified by silica gel column chromatography to obtain 1.60 g (4.5 mmol, yield 90%) of Intermediate A-1. The obtained compound was identified by MS/FAB.  $\text{C}_{15}\text{H}_9\text{BrCl}_2\text{O}$  cal. 356.04, found 353.92.

#### Synthesis of Intermediate A-2

**[0251]** 1.60 g (4.5 mmol) of Intermediate A-1 and 4.56 g (18.0 mmol) of iodine were added to 40 mL of toluene and stirred under reflux for 24 hours. When the reaction was completed, 100 mL of saturated aqueous sodium thiosulfate was added thereto and stirred at room temperature for 30 minutes, and a separated organic layer was extracted by using toluene. The extracted organic layer was dried by using magnesium sulfate and a solvent was evaporated. The residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 1.89 g (4.1 mmol, yield 90%) of Intermediate A-2. The obtained compound was identified by MS/FAB.  $\text{C}_{14}\text{H}_6\text{BrCl}_2\text{IO}$  cal. 467.91, found 465.80.

#### Synthesis of Intermediate A-3

**[0252]** 1.89 g (4.1 mmol) of Intermediate A-2 and 1.88 g (16 mmol) of  $\text{N,N,N',N'}$ -tetramethylethylenediamine (TMEDA) were added to a flame-dried Schlenk flask, and 100 mL of diethyl ether was added thereto and stirred in a nitrogen atmosphere. The reaction solution was stirred at a temperature of  $-78^\circ\text{C}$ . for 30 minutes and  $n\text{-BuLi}$  (1.6M solution in hexane, 5.1 mL, 8.1 mmol) was added dropwise thereto. The reaction solution was stirred at the same temperature for 15 minutes, and 0.95 mL (5.3 mmol) of  $\text{PPhCl}_2$  was added thereto all at once. The resultant was slowly heated to room temperature and stirred for 15 minutes. After the reaction was completed, 300 mL of  $\text{CH}_2\text{Cl}_2$  was added and filtered by using neutral alumina and a solvent was evaporated. The residue obtained therefrom was separated and purified by neutral alumina chromatography in a nitrogen atmosphere to obtain 745 mg (2.0 mmol, yield 50%) of Intermediate A-3. The obtained compound was identified by  $^1\text{H}$  NMR.

#### Synthesis of Intermediate A-4




**[0253]** 745 mg (2.0 mmol) of Intermediate A-3 was added to a flame-dried Schlenk flask in a nitrogen atmosphere and dissolved in 100 mL of  $\text{CH}_2\text{Cl}_2$ . 3.2 mL of  $\text{H}_2\text{O}_2$  (a 35% aqueous solution) was added to the mixed solution all at once and stirred at room temperature for 30 minutes. After the reaction was completed, the mixture was stirred while 20 mL of a saturated aqueous  $\text{Na}_2\text{S}_2\text{O}_3$  was slowly added dropwise thereto. Then, an organic layer was separated from the resultant mixture, and a water layer was combined with the organic layer from which an oil layer was extracted by using  $\text{AcOEt}$ . Then, the organic layer was dried by using magnesium sulfate and a solvent was evaporated. The residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 423 mg (1.1 mmol, yield 55%) of Intermediate A-4. The obtained compound was identified by MS/FAB.  $\text{C}_{20}\text{H}_{11}\text{Cl}_2\text{O}_2\text{P}$  cal. 385.18, found 383.99.

## Synthesis of Compound 8

**[0254]** 423 mg (1.1 mmol) of Intermediate A-4, 410 mg (2.4 mmol) of diphenylamine, 50.5 mg (0.05 mmol) of tris(dibenzylideneacetone)dipalladium (0), 30.5 mg (0.15 mmol) of tri(*tert*-butyl)phosphine, and 317 mg (3.3 mmol) of sodium *tert*-butoxide were added to 10 mL of toluene and stirred at a temperature of 80° C. for 12 hours. The reaction solution was cooled to room temperature and an extraction process was performed thereon twice by using water and diethyl ether. An organic layer obtained therefrom was dried by using magnesium sulfate and a solvent was evaporated. The residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 515 mg (0.79 mmol, yield 72%) of Compound 8. The obtained compound was identified by MS/FAB.  $C_{44}H_{31}N_2O_2P$  cal. 650.72, found 650.21.

## Synthesis Example: Synthesis of Compound 10

**[0255]**



10

## Synthesis of Intermediate B-1

**[0256]** 564 mg (1.92 mmol, yield 47%) of Intermediate B-1 was synthesized in the same manner as in Synthesis of Intermediate A-3, except that 1.89 g (4.1 mmol) of Compound A-2 was used instead as a starting material and  $\text{SCl}_2$  was used instead of  $\text{PPhCl}_2$ . The obtained compound was identified by  $^1\text{H}$  NMR.

## Synthesis of Intermediate B-2


**[0257]** 343 mg (1.05 mmol, yield 55%) of Intermediate B-2 was synthesized in the same manner as in Synthesis of Intermediate A-4, except that 564 mg (1.92 mmol) of Compound B-1 was used as a starting material. The obtained compound was identified by MS/FAB.  $C_{14}H_6C_{12}O_3S$  cal. 325.16, found 323.94.

## Synthesis of Compound 10

**[0258]** 446 mg (0.75 mmol, yield 72%) of Compound 10 was synthesized in the same manner as in Synthesis of Compound 8, except that 343 mg (1.05 mmol) of Compound B-2 was used as a starting material. The obtained compound was identified by MS/FAB.  $C_{38}H_{26}N_2O_3S$  cal. 590.70, found 590.17.

## Synthesis Example: Synthesis of Compounds 12 and 13

**[0259]**



#### Synthesis of Intermediate C-1

**[0260]** 1.04 g (3.28 mmol, yield 80%) of Intermediate C-1 was synthesized in the same manner as in Synthesis of Intermediate A-3, except that 1.89 g (4.1 mmol) of Intermediate A-2 was used as a starting material and  $\text{Si}(\text{CH}_3)_2\text{Cl}_2$  was used instead of  $\text{PPhCl}_2$ . The obtained compound was identified by MS/FAB.  $\text{C}_{16}\text{H}_{12}\text{Cl}_2\text{OSi}$  cal. 319.26, found 318.00.

#### Synthesis of Intermediate C-2

**[0261]** 1.60 g (3.60 mmol, yield 88%) of Intermediate C-2 was synthesized in the same manner as in Synthesis of Intermediate A-3, except that 1.89 g (4.1 mmol) of Intermediate A-2 was used as a starting material and  $\text{SiPh}_2\text{Cl}_2$  was used instead of  $\text{PPhCl}_2$ . The obtained compound was identified by MS/FAB.  $\text{C}_{26}\text{H}_{16}\text{Cl}_2\text{OSi}$  cal. 443.40, found 442.03.

#### Synthesis of Compound 12

**[0262]** 1.43 g (2.46 mmol, yield 75%) of Compound 12 was synthesized in the same manner as in Synthesis of Compound 8, except that 1.04 g (3.28 mmol) of Intermediate

C-1 was used as a starting material. The obtained compound was identified by MS/FAB.  $\text{C}_{40}\text{H}_{32}\text{N}_2\text{OSi}$  cal. 584.79, found 584.23.

#### Synthesis of Compound 13

**[0263]** 2.14 g (3.02 mmol, yield 84%) of Compound 13 was synthesized in the same manner as in Synthesis of Compound 8, except that 1.60 g (3.60 mmol) of Intermediate C-2 was used as a starting material. The obtained compound was identified by MS/FAB.  $\text{C}_{50}\text{H}_{36}\text{N}_2\text{OSi}$  cal. 708.94, found 708.26.

#### Synthesis of Compound 85

**[0264]** 2.44 g (3.52 mmol, yield 75%) of Compound 85 was synthesized in the same manner as in Synthesis of Compound 8, except that 1.50 g (4.70 mmol) of Intermediate C-1 was used as a starting material and dimesitylamine was used instead of diphenylamine. The obtained compound was identified by MS/FAB.  $\text{C}_{52}\text{H}_{56}\text{N}_2\text{OSi}$  cal. 753.12, found 752.42.

#### Synthesis of Compound 87

**[0265]** 2.65 g (3.20 mmol, yield 68%) of Compound 87 was synthesized in the same manner as in Synthesis of

Compound 8, except that 1.50 g (4.70 mmol) was used as a starting material and N-phenyldibenzo[b,d]furan-4-amine was used instead of diphenylamine. The obtained compound was identified by MS/FAB.  $C_{52}H_{36}N_2O_3Si$  cal. 764.96, found 764.25.

#### Synthesis of Compound 97



**[0266]** 2.38 g (2.91 mmol, yield 62%) of Compound 97 was synthesized in the same manner as in Synthesis of Compound 8, except that 1.50 g (4.70 mmol) of Intermediate C-1 was used as a starting material and 9,9-dimethyl-N-phenyl-9H-fluoren-1-amine was used instead of diphenylamine. The obtained compound was identified by MS/FAB.  $C_{58}H_{48}N_2OSi$  cal. 817.12, found 816.35.

#### Synthesis of Compound 104

**[0267]** 2.58 g (2.82 mmol, yield 60%) of Compound 104 was synthesized in the same manner as in Synthesis of Compound 8, except that 1.50 g (4.70 mmol) of Intermediate C-1 was used as a starting material and N-([1,1'-biphenyl]-2-yl)dibenzo[b,d]furan-1-amine was used instead of diphenylamine. The obtained compound was identified by MS/FAB.  $C_{64}H_{44}N_2O_3Si$  cal. 917.15, found 916.31.

#### Synthesis Example: Synthesis of Compound 61

##### [0268]



61

#### Synthesis of Intermediate D-1

**[0269]** 1.22 g (3.44 mmol, yield 86%) of Intermediate D-1 was synthesized in the same manner as in Synthesis of Intermediate A-1, except that 4-chloro-2-ethynyl-1-methoxybenzene was used instead of 4-chloro-1-ethynyl-2-methoxybenzene. The obtained compound was identified by MS/FAB.  $C_{15}H_9BrCl_2O$  cal. 356.04, found 353.92.

#### Synthesis of Intermediate D-2

**[0270]** 1.32 g (2.82 mmol, yield 82%) of Intermediate D-2 was synthesized in the same manner as in Synthesis of Intermediate A-2, except that 1.22 g (3.44 mmol) of Intermediate D-1 was used as a starting material. The obtained compound was identified by MS/FAB.  $C_{14}H_6BrCl_2IO$  cal. 467.91, found 465.80.

#### Synthesis of Intermediate D-3

**[0271]** 428 mg (1.46 mmol, yield 52%) of Intermediate D-3 was synthesized in the same manner as in Synthesis of Intermediate B-1, except that 1.32 g (2.82 mmol) of Intermediate D-2 was used as a starting material. The obtained compound was identified by MS/FAB.  $C_{14}H_6Cl_2OS$  cal. 293.16, found 291.95.

#### Synthesis of Intermediate D-4


**[0272]** 260 mg (0.80 mmol, yield 55%) of Intermediate D-4 was synthesized in the same manner as in Synthesis of Intermediate B-2, except that 428 mg (1.46 mmol) of Intermediate D-3 was used as a starting material. The obtained compound was identified by MS/FAB.  $C_{14}H_6Cl_2O_3S$  cal. 325.16, found 323.94.

## Synthesis of Compound 61

[0273] 260 mg (0.80 mmol) of Intermediate D-4 and (4-(diphenylamino)phenyl)boronic acid (578 mg, 0.2 mmol), 46 mg (0.04 mol) of Pd(PPh<sub>3</sub>)<sub>4</sub>, and 331 mg (2.4 mmol) of K<sub>2</sub>CO<sub>3</sub> were added to 10 mL of a mixture of THF/H<sub>2</sub>O (volume ratio: 9/1) and stirred at a temperature of 80° C. for 12 hours. Then, the resultant mixture was cooled to room temperature and an extraction process was performed thereon three times by using water and diethyl ether. An organic layer obtained therefrom was dried by using magnesium sulfate and a solvent was evaporated. The residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 421 mg (0.57 mmol, yield 71%) of Compound Compound 61. The obtained compound was identified by MS/FAB. C<sub>50</sub>H<sub>34</sub>N<sub>2</sub>O<sub>3</sub>S cal. 742.89, found 742.23.

## Synthesis Example: Synthesis of Compound 64

[0274]



## Synthesis of Intermediate E-1

[0275] 943 mg (2.11 mmol, yield 78%) of Intermediate E-1 was synthesized in the same manner as in Synthesis of


Intermediate A-3, except that 468 mg (2.70 mmol) of Intermediate D-2 was used as a starting material and SiPh<sub>2</sub>Cl<sub>2</sub> was used instead of PPhCl<sub>2</sub>. The obtained compound was identified by MS/FAB. C<sub>26</sub>H<sub>16</sub>Cl<sub>2</sub>OS cal. 447.37, found 446.03.

## Synthesis of Compound 64

[0276] 1.40 g (1.62 mmol, yield 77%) of Compound 64 was synthesized in the same manner as in Synthesis of Compound 61, except that 943 mg (2.11 mmol) of Intermediate E-1 was used as a starting material. The obtained compound was identified by MS/FAB. C<sub>62</sub>H<sub>44</sub>N<sub>2</sub>O<sub>3</sub>Si cal. 861.13, found 860.32.

## Synthesis Example: Synthesis of Compound 130

[0277]



## Synthesis of Intermediate F-1

[0278] 610 mg (1.89 mmol, yield 68%) of Intermediate F-1 was synthesized in the same manner as in Synthesis of Intermediate A-3, except that 1.30 g (2.77 mmol) of Intermediate D-2 was used as a starting material and Si(CH<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>

was used instead of PPhCl<sub>2</sub>. The obtained compound was identified by MS/FAB. C<sub>16</sub>H<sub>12</sub>Cl<sub>2</sub>OS cal. 323.23, found 322.00.

#### Synthesis of Compound 130

**[0279]** 972 mg (1.66 mmol, yield 88%) of Compound 130 was synthesized in the same manner as in Synthesis of Compound 8, except that 610 mg (1.89 mmol) of Intermediate F-1 was used as a starting material. The obtained compound was identified by MS/FAB. C<sub>40</sub>H<sub>32</sub>N<sub>2</sub>OSi cal. 584.79, found 584.23.

#### Synthesis of Compound 141

**[0280]** 21.94 g (1.80 mmol, yield 45%) of Compound 141 was synthesized in the same manner as in Synthesis of Compound 8, except that 1.29 g (4.0 mmol) of Intermediate F-1 was used as a starting material and N-([1,1'-biphenyl]-2-yl)-5'-fluoro-[1,1':3',1''-terphenyl]-4'-amine was used instead of diphenylamine. The obtained compound was identified by MS/FAB. C<sub>76</sub>H<sub>54</sub>F<sub>2</sub>N<sub>2</sub>OSi cal. 1077.36, found 1076.40.

**[0281]** NMR and MS data of Compounds synthesized in Synthesis Examples are shown in Table 1:

#### Example 1




**[0282]** An anode was prepared by cutting an ITO glass substrate (manufactured by Corning), on which an ITO layer was deposited to a thickness of 15 Ωcm<sup>2</sup> (1,200 Å), to a size of 50 mm×50 mm×0.7 mm, sonicating the ITO glass substrate by using isopropyl alcohol and pure water each for 5 minutes, and then, exposing the ITO glass substrate to UV irradiation and ozone for 30 minutes to clean the ITO glass substrate. Then, the ITO glass substrate (anode) was loaded into a vacuum deposition apparatus.

**[0283]** Compound 2-TNATA, which is a known compound, was vacuum-deposited on the ITO glass substrate to form a hole injection layer having a thickness of 600 Å, and Compound 8, which is a hole transport compound according to one or more embodiments, was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å.


**[0284]** 9,10-di-naphthalene-2-yl-anthracene (ADN), which is a known blue fluorescent host, and N,N,N',N'-tetraphenyl-pyrene-1,6-diamine (TPD), which is a known blue fluorescent dopant, were co-deposited on the hole transport layer at a weight ratio of 98:2 to form an emission layer having a thickness of 300 Å.

TABLE 1

Compound	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400 MHz)	MS/FAB	
		calc.	found
8	δ = 7.82(dd, 1H), 7.8-7.7(m, 2H), 7.56-7.51(m, 1H), 7.49(dd, 1H), 7.41-7.35(m, 2H), 7.29(dd, 1H), 7.17(dd, 1H), 7.12-7.05(m, 8H), 6.8-6.74(m, 1H), 6.7-6.65(m, 5H), 6.33-6.23(m, 8H)	650.72	650.21
10	δ = 7.66(dd, 1H), 7.49(d, 1H), 7.43(dd, 1H), 7.18(dd, 1H), 7.18(d, 1H), 7.11-7.05(m, 8H), 6.88(dd, 1H), 6.7-6.62(m, 4H), 6.33-6.23(m, 8H)	590.70	590.17
12	δ = 7.73(d, 1H), 7.37(d, 1H), 7.26-7.21(m, 2H), 7.11-7.04(m, 8H), 6.83(d, 1H), 6.8-6.62(m, 5H), 6.33-6.23(m, 8H), 0.56(s, 6H)	584.79	584.23
13	δ = 7.65(d, 1H), 7.53-7.46(m, 5H), 7.36-7.26(m, 7H), 7.15(dd, 1H), 7.11-(m, 8H), 6.85(d, 1H), 6.69-6.64(m, 4H), 6.52(dd, 1H), 6.33-6.23(m, 8H)	708.94	708.26
61	δ = 8.49(d, 1H), 8.07(d, 1H), 8(d, 1H), 7.96(d, 1H), 7.7-7.62(m, 6H), 7.09-7.04(m, 8H), 6.9-6.84(m, 4H), 6.7-6.64(m, 4H), 6.21-6.15(m, 8H)	742.89	42.23
64	δ = 8.27(d, 1H), 7.88-7.82(m, 2H), 7.68(d, 1H), 7.66(d, 1H), 7.51-7.41(m, 9H), 7.36-7.26(m, 6H), 7.09-7.04(m, 8H), 6.9-6.82(m, 4H), 6.7-6.64(m, 4H), 6.21-6.15(m, 8H)	861.13	860.32
85	δ = 7.7(d, 1H), 7.32(d, 1H), 7.18(d, 1H), 7.03(d, 1H), 6.8-6.72(m, 9H), 6.65(d, 1H), 2.39(s, 12H), 2.26(s, 24H), 0.56(s, 6H)	753.12	752.42
87	δ = 7.85-7.82(m, 2H), 7.75-7.7(m, 3H), 7.56-7.54(m, 2H), 7.49-7.41(m, 4H), 7.33-7.23(m, 3H), 7.09-7.04(m, 4H), 7.02-6.94(m, 5H), 6.74-6.62(m, 3H), 6.4-6.3(m, 4H), 0.56(s, 6H)	764.96	764.25
97	δ = 7.76(d, 1H), 7.76(d, 1H), 7.69(d, 1H), 7.48(d, 1H), 7.47(d, 1H), 7.39(d, 1H), 7.38(d, 1H), 7.29-7.24(m, 4H), 7.14-7.02(m, 7H), 6.97(d, 1H), 6.96(d, 1H), 6.82-6.62(m, 5H), 6.56(dd, 1H), 6.26-6.18(m, 2H), 6.18-6.04(m, 2H), 1.71(s, 6H), 1.7(s, 6H), 0.56(s, 6H)	817.12	816.35
104	δ = 7.83-7.8(m, 2H), 7.73-7.7(m, 2H), 7.63(d, 1H), 7.6-7.53(m, 8H), 7.49-7.44(m, 4H), 7.34-7.27(m, 6H), 7.24(d, 1H), 7.21-7.16(m, 5H), 7.1(d, 1H), 7.05-6.99(m, 2H), 6.84-6.79(m, 3H), 6.62-6.53(m, 3H), 0.56(s, 6H)	917.15	916.31
130	δ = 7.45(d, 1H), 7.37(d, 1H), 7.17(d, 1H), 7.11-7.05(m, 9H), 6.88(dd, 1H), 6.7-6.62(m, 5H), 6.28-6.23(m, 8H), 0.53(s, 6H)	584.79	584.23
141	δ = 7.73-7.7(m, 5H), 7.66-7.61(m, 6H), 7.61-7.56(m, 12H), 7.56-7.51(m, 6H), 7.49-7.41(m, 4H), 7.3(d, 1H), 7.21-7.16(m, 5H), 7.12(d, 1H), 7.11(d, 1H), 6.98-7(m, 2H), 6.93-6.85(m, 3H), 6.47(dd, 1H), 6.19(dd, 1H), 0.53(s, 6H)	1077.36	1076.40



ADN



[0285] Then,  $\text{Alq}_3$  was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å. LiF, which is an alkali metal halide, was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited on the electron injection layer to form a LiF/Al electrode (cathode electrode) having a thickness of 3,000 Å, thereby completing the manufacture of an organic light-emitting device.

#### Example 2

[0286] An organic light-emitting device of Example 2 was manufactured in the same manner as in Example 1, except that Compound 10 was used instead of Compound 8 in forming a hole transport layer.

#### Example 3

[0287] An organic light-emitting device of Example 3 was manufactured as in Example 1, except that Compound 13 was used instead of Compound 8 in forming a hole transport layer.

#### Example 4

[0288] An organic light-emitting device of Example 4 was manufactured as in Example 1, except that Compound 61 was used instead of Compound 8 in forming a hole transport layer.

#### Example 5


[0289] An organic light-emitting device of Example 5 was manufactured as in Example 1, except that Compound 64 was used instead of Compound 8 in forming a hole transport layer.

#### Example 6

[0290] An organic light-emitting device of Example 6 was manufactured as in Example 1, except that Compound 130 was used instead of Compound 8 in forming a hole transport layer.

#### Comparative Example 1

[0291] An organic light-emitting device of Comparative Example 1 was manufactured as in Example 1, except that 4,4'-bis[N-(1-naphthyl)-N-phenylamino]biphenyl (NPB), which is a known compound, was used in forming a hole transport layer.



NPB

[0292] Results of Examples 1 to 6 and Comparative Example 1 are shown in Table 2.

TABLE 2

	Hole transport layer	Driving voltage (V)	Current density (mA/cm <sup>2</sup> )	Luminance (cd/m <sup>2</sup> )	Efficiency (cd/A)	Emission color	Half-life (hr @100 mA/cm <sup>2</sup> )
Example 1	Compound 8	5.43	50	3474	6.43	Blue	319
Example 2	Compound 10	5.91	50	3396	6.63	Blue	351
Example 3	Compound 13	5.01	50	3125	6.68	Blue	356
Example 4	Compound 61	5.35	50	3699	6.23	Blue	325
Example 5	Compound 64	4.99	50	3253	6.25	Blue	369
Example 6	Compound 130	6.00	50	3110	6.39	Blue	318
Comparative Example 1	NPB	6.99	50	2745	5.22	Blue	256

#### Example 7

[0293] An organic light-emitting device of Example 7 was manufactured in the same manner as in Example 1, except that NPB, which is a known compound, was used instead of Compound 8 in forming a hole transport layer, and Compound 8, which is a known dopant compound, was used instead of TPD in forming an emission layer.

#### Example 8

[0294] An organic light-emitting device of Example 8 was manufactured in the same manner as in Example 7, except that Compound 10 was used instead of Compound 8 in forming an emission layer.

#### Example 9

[0295] An organic light-emitting device of Example 9 was manufactured as in the same manner as in Example 7, except that Compound 12 was used instead of Compound 8 in forming an emission layer.

#### Example 10

[0296] An organic light-emitting device of Example 10 was manufactured in the same manner as in Example 7, except that Compound 13 was used instead of Compound 8 in forming an emission layer.

#### Example 11

[0297] An organic light-emitting device of Example 11 was manufactured in the same manner as in Example 7, except that Compound 85 was used instead of Compound 8 in forming an emission layer.

#### Example 12

[0298] An organic light-emitting device of Example 12 was manufactured in the same manner as in Example 7, except that Compound 87 was used instead of Compound 8 in forming an emission layer.

#### Example 13

[0299] An organic light-emitting device of Example 13 was manufactured in the same manner as in Example 7,

except that Compound 97 was used instead of Compound 8 in forming an emission layer.

#### Example 14

[0300] An organic light-emitting device of Example 14 was manufactured in the same manner as in Example 7, except that Compound 104 was used instead of Compound 8 in forming an emission layer.

#### Example 15


[0301] An organic light-emitting device of Example 15 was manufactured in the same manner as in Example 7, except that Compound 130 was used instead of Compound 8 in forming an emission layer.

#### Example 16

[0302] An organic light-emitting device of Example 16 was manufactured in the same manner as in Example 7, except that Compound 141 was used instead of Compound 8 in forming an emission layer.

#### Comparative Example 2

[0303] An organic light-emitting device of Comparative Example 2 was manufactured in the same manner as in Example 7, except that Compound A, which is a known compound, was used instead of Compound 8 in forming an emission layer.



A

[0304] Results of Comparative Example 2 and Examples 7 to 16 are shown in Table 3.

TABLE 3

	Emission layer	Driving voltage (V)	Current density (mA/cm <sup>2</sup> )	Brightness (cd/m <sup>2</sup> )	Efficiency (cd/A)	Emission color	Half-life (hr @100 mA/cm <sup>2</sup> )
Example 7	Compound 8	5.08	50	2962	7.26	Blue	299
Example 8	Compound 10	5.89	50	2969	7.26	Blue	341
Example 9	Compound 12	5.22	50	3337	7.41	Blue	358
Example 10	Compound 13	5.25	50	3255	7.27	Blue	348
Example 11	Compound 85	5.34	50	3456	7.31	Blue	341
Example 12	Compound 87	5.70	50	3463	6.98	Blue	300
Example 13	Compound 97	5.20	50	3497	7.62	Blue	355
Example 14	Compound 104	5.05	50	3555	7.31	Blue	371
Example 15	Compound 130	5.70	50	3463	6.98	Blue	299
Example 16	Compound 141	5.30	50	3497	7.62	Blue	325
Comparative Example 2	A	6.33	50	3560	6.15	Blue	335

[0305] When the structure of Formula 1 according to one or more embodiments is used as the materials of the hole transport layer and the emission layer, a driving voltage may be reduced and efficiency may be improved, as compared to the known compounds. In particular, the half-life of the device may be increased.

[0306] While the inventive concept has been particularly shown and described with reference to example embodiments thereof, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope as defined by the following claims.


[0307] An organic light-emitting device according to an embodiment has high efficiency, a low voltage, high luminance, and a long lifespan.

[0308] It should be understood that embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as available for other similar features or aspects in other embodiments.

[0309] While one or more embodiments have been described with reference to the FIGURES, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope as defined by the following claims.

What is claimed is:

1. A compound represented by Formula 1:



wherein, in Formula 1,

$R_1$ ,  $R_2$ ,  $Ar_1$ , and  $Ar_2$  are each independently selected from hydrogen, deuterium, halogen, an amino group, a nitro group, a nitrile group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkenyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkynyl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkoxy group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic

condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

$L_1$  and  $L_2$  are each independently selected from a substituted or unsubstituted  $C_6-C_{60}$  arylene group, a substituted or unsubstituted  $C_1-C_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

$Y_1$  and  $Y_2$  are each independently selected from N, B, and Si,

X is selected from B, C, Si, Ge, Sn, S, SO,  $S(O)_2$ , P, PO, N, and O,

A1 and A2 are each independently a phenylene group or a naphthylene group,

a1, a2, b1, b2, c1, and c2 are each independently an integer from 0 to 3, and

at least one substituent of the substituted  $C_1-C_{60}$  alkyl group, the substituted  $C_2-C_{60}$  alkenyl group, the substituted  $C_2-C_{60}$  alkynyl group, the substituted  $C_1-C_{60}$  alkoxy group, the substituted  $C_3-C_{10}$  cycloalkyl group, the substituted  $C_2-C_{10}$  heterocycloalkyl group, the substituted  $C_3-C_{10}$  cycloalkenyl group, the substituted  $C_2-C_{10}$  heterocycloalkenyl group, the substituted  $C_6-C_{60}$  aryl group, the substituted  $C_6-C_{60}$  aryloxy group, the substituted  $C_6-C_{60}$  arylthio group, the substituted  $C_1-C_{60}$  heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted  $C_6-C_{60}$  arylene group, the substituted  $C_1-C_{60}$  heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and the substituted divalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_1-C_{60}$  alkyl group, a  $C_2-C_{60}$  alkenyl group, a  $C_2-C_{60}$  alkynyl group, and a  $C_1-C_{60}$  alkoxy group;

a  $C_1-C_{60}$  alkyl group, a  $C_2-C_{60}$  alkenyl group, a  $C_2-C_{60}$  alkynyl group, and a  $C_1-C_{60}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_3-C_{10}$  cycloalkyl group, a  $C_1-C_{10}$  heterocycloalkyl group, a  $C_3-C_{10}$  cycloalkenyl group, a  $C_2-C_{10}$  heterocycloalkenyl group, a  $C_6-C_{60}$  aryl group, a  $C_6-C_{60}$  aryloxy group, a  $C_6-C_{60}$  arylthio group, a  $C_2-C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N( $Q_{11}$ )( $Q_{12}$ ), —Si( $Q_{13}$ )( $Q_{14}$ )( $Q_{15}$ ), and —B( $Q_{16}$ )( $Q_{17}$ );

a  $C_3-C_{10}$  cycloalkyl group, a  $C_2-C_{10}$  heterocycloalkyl group, a  $C_3-C_{10}$  cycloalkenyl group, a  $C_2-C_{10}$  heterocycloalkenyl group, a  $C_6-C_{60}$  aryl group, a  $C_6-C_{60}$  aryloxy group, a  $C_6-C_{60}$  arylthio group, a  $C_1-C_{60}$  heteroaryl group, a monovalent non-aromatic condensed

polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

a  $C_3-C_{10}$  cycloalkyl group, a  $C_2-C_{10}$  heterocycloalkyl group, a  $C_3-C_{10}$  cycloalkenyl group, a  $C_2-C_{10}$  heterocycloalkenyl group, a  $C_6-C_{60}$  aryl group, a  $C_6-C_{60}$  aryloxy group, a  $C_6-C_{60}$  arylthio group, a  $C_1-C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_1-C_{60}$  alkyl group, a  $C_2-C_{60}$  alkenyl group, a  $C_2-C_{60}$  alkynyl group, a  $C_1-C_{60}$  alkoxy group, a  $C_3-C_{10}$  cycloalkyl group, a  $C_2-C_{10}$  heterocycloalkyl group, a  $C_3-C_{10}$  cycloalkenyl group, a  $C_2-C_{10}$  heterocycloalkenyl group, a  $C_6-C_{60}$  aryl group, a  $C_6-C_{60}$  aryloxy group, a  $C_6-C_{60}$  arylthio group, a  $C_1-C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N( $Q_{21}$ )( $Q_{22}$ ), —Si( $Q_{23}$ )( $Q_{24}$ )( $Q_{25}$ ), and —B( $Q_{26}$ )( $Q_{27}$ ),

wherein  $Q_{11}$  to  $Q_{17}$  and  $Q_{21}$  to  $Q_{27}$  are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_1-C_{60}$  alkyl group, a  $C_2-C_{60}$  alkenyl group, a  $C_2-C_{60}$  alkynyl group, a  $C_1-C_{60}$  alkoxy group, a  $C_3-C_{10}$  cycloalkyl group, a  $C_2-C_{10}$  heterocycloalkyl group, a  $C_3-C_{10}$  cycloalkenyl group, a  $C_2-C_{10}$  heterocycloalkenyl group, a  $C_6-C_{60}$  aryl group, a  $C_1-C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

2. The compound of claim 1, wherein

X in Formula 1 is selected from B, C, Si, Ge, Sn, S, SO,  $S(O)_2$ , P, and PO.

3. The compound of claim 1, wherein

$R_1$  and  $R_2$  in Formula 1 are each independently represented by one of Formulae 2a and 2b:




wherein, in Formulae 2a and 2b,  $Z_1$  is selected from hydrogen, deuterium, a cyano group, a halogen group, a substituted or unsubstituted  $C_1-C_{20}$  alkyl group, a substituted or unsubstituted  $C_1-C_{20}$  alkoxy group, a substituted or unsubstituted  $C_6-C_{20}$  aryl group, a substituted or unsubstituted  $C_1-C_{20}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic

condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

p is an integer from 1 to 5, and

\* indicates a binding site.






4. The compound of claim 1, wherein each of  $L_1$  and  $L_2$  in Formula 1 is a group represented by Formula 3a:



wherein \* in Formula 3a indicates a binding site.

5. The compound of claim 1, wherein each of  $Y_1$  and  $Y_2$  in Formula 1 is N.

6. The compound of claim 1, wherein  $Ar_1$  to  $Ar_4$  in Formula 1 are each independently represented by one of Formulae 4a to 4e:



Wherein, in Formulae 4a to 4e,  $H_1$  is O, S,  $CR_{11}R_{12}$ , or  $NR_{13}$ ,


$R_{11}$  to  $R_{13}$  and  $Z_1$  are each independently selected from hydrogen, deuterium, a cyano group, a halogen group, a substituted or unsubstituted  $C_1$ - $C_{20}$  alkyl group, a substituted or unsubstituted  $C_1$ - $C_{20}$  alkoxy group, a substituted or unsubstituted  $C_6$ - $C_{20}$  aryl group, a substituted or unsubstituted  $C_1$ - $C_{20}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

p is an integer from 1 to 9, and

\* indicates a binding site.


7. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 2:

<Formula 2>




8. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 3:

<Formula 3>




9. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 4:

<Formula 4>




10. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 5:

<Formula 5>




11. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 6:

<Formula 6>




12. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 7:

<Formula 7>




13. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 8:

<Formula 8>




14. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 9:

<Formula 9>




15. The compound of claim 1, wherein the compound represented by Formula 1 is represented by Formula 10:


<Formula 10>



16. The compound of claim 1, wherein the compound represented by Formula 1 is any one of compounds below:




1




2

-continued




3


-continued




6




4



7




5




8

-continued




9


-continued




12




10



13




11




14

-continued




15


-continued




18




16



19




17




20

-continued




21


-continued




24




22



25




23




26

-continued




27


-continued




30




28



31




29




32


-continued




-continued




34



36




-continued




37

-continued




39

38




40







-continued




-continued




46



48




-continued




49

-continued




51

50







52







-continued

-continued







-continued

-continued







-continued

-continued




-continued

-continued




-continued




69


-continued




72




70



73






71






74

-continued



-continued



75

78


76

79


77

80


-continued



81




82




83

-continued




84






85


-continued



-continued




-continued




90


-continued



92




91




93

-continued




94


-continued



96





95






97


-continued



-continued




-continued




103

-continued





104

106




105




107


-continued




-continued



109




111




-continued


-continued




114



113







115




-continued

-continued




-continued




120

-continued




122

121







123




-continued

-continued




-continued




128


-continued



130




129




131

-continued




132


-continued




134



133



135





136

-continued

-continued



137

139




138

140




-continued

141




142



-continued

143



17. An organic light-emitting device comprising:  
 a first electrode;  
 a second electrode facing the first electrode; and  
 an organic layer between the first electrode and the second electrode, the organic layer comprising an emission layer,  
 wherein the organic layer comprises the compound of claim 1.
18. The organic light-emitting device of claim 17, wherein the first electrode is an anode, the second electrode is a cathode, and the organic layer comprises:  
 i) a hole transport region between the first electrode and the emission layer, the hole transport region comprising a hole transport layer; and at least one of a hole injection layer, and an electron blocking layer; and  
 ii) an electron transport region between the emission layer and the second electrode, the electron transport region comprising at least one of an electron transport layer, a hole blocking layer, and an electron injection layer.
19. The organic light-emitting device of claim 18, wherein the emission layer or the hole transport layer comprises the compound of claim 1.
20. A display apparatus comprising:  
 the organic light-emitting device of claim 17, wherein the first electrode of the organic light-emitting device is electrically connected to a source electrode or a drain electrode of a thin film transistor.

\* \* \* \* \*

专利名称(译)	包含相同的复合和有机发光装置		
公开(公告)号	<a href="#">US20180053902A1</a>	公开(公告)日	2018-02-22
申请号	US15/650077	申请日	2017-07-14
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	KIM HAEJIN KO SOOBYUNG JUN MIEUN KIM YOUNGKOOK HWANG SEOKHWAN		
发明人	KIM, HAEJIN KO, SOOBYUNG JUN, MIEUN KIM, YOUNGKOOK HWANG, SEOKHWAN		
IPC分类号	H01L51/00 C09K11/06 H01L27/32		
CPC分类号	H01L51/0071 C07D325/00 H01L27/32 C09K11/06		
优先权	1020160105582 2016-08-19 KR		
其他公开文献	US10439149		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

提供由式1表示的化合物：其中公式1相同如在本公开的详细描述中所描述的。式1的化合物可以包含在有机发光器件(OLED)的空穴传输层和/或发光层中。

10

190
150
110