




US 20170186952A1

(19) **United States**(12) **Patent Application Publication**
HAN et al.(10) **Pub. No.: US 2017/0186952 A1**(43) **Pub. Date: Jun. 29, 2017**(54) **COMPOUND AND ORGANIC
LIGHT-EMITTING DEVICE INCLUDING
THE SAME***C07D 209/88* (2006.01)*C07D 409/12* (2006.01)(52) **U.S. Cl.**CPC *H01L 51/006* (2013.01); *H01L 51/0072*
(2013.01); *H01L 51/005* (2013.01); *H01L*
51/0077 (2013.01); *C07D 209/88* (2013.01);
H01L 51/0061 (2013.01); *C07D 409/12*
(2013.01); *C07D 209/86* (2013.01); *C07D*
405/12 (2013.01); *C07D 409/14* (2013.01);
H01L 51/5012 (2013.01)(71) Applicant: **Samsung Display Co., Ltd**, Yongin-si
(KR)(72) Inventors: **Sanghyun HAN**, Yongin-si (KR);
Sooyon KIM, Yongin-si (KR);
Jongwoo KIM, Yongin-si (KR);
Youngkook KIM, Yongin-si (KR);
Seokhwan HWANG, Yongin-si (KR)(57) **ABSTRACT**A compound represented by Formula 1 and an organic
light-emitting device including the same are provided:(21) Appl. No.: **15/393,086**(22) Filed: **Dec. 28, 2016**(30) **Foreign Application Priority Data**

Dec. 28, 2015 (KR) 10-2015-0187638

Publication Classification(51) **Int. Cl.***H01L 51/00* (2006.01)*C07D 409/14* (2006.01)*C07D 209/86* (2006.01)*C07D 405/12* (2006.01)

<Formula 1>

40

220
190
150
110
210

FIG. 110

190
150
110

FIG. 2

20

190
150
110
210

FIG. 3

30

220
190
150
110

FIG. 440

220
190
150
110
210

COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application claims the benefit of Korean Patent Application No. 10-2015-0187638 filed on Dec. 28, 2015 in the Korean Intellectual Property Office, the disclosure of which is incorporated herein in its entirety by reference.

BACKGROUND

[0002] 1. Field

[0003] One or more embodiments relate to a compound and an organic light-emitting device including the same.

[0004] 2. Description of the Related Art

[0005] Organic light-emitting devices are self-emission devices that offer wide viewing angles, high contrast ratios, short response times, and excellent luminance, driving voltage, and response speed characteristics. Organic light-emitting devices produce full-color images.


[0006] An example of such organic light-emitting devices may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, recombine in the emission layer to produce excitons. These excitons transition from an excited state to a ground state, thereby generating light.

SUMMARY

[0007] One or more embodiments include a material for use in a hole transport region and an organic light-emitting device that has improved characteristics due to the inclusion of the material.

[0008] Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments.

[0009] According to one or more embodiments, a compound is represented by Formula 1:



[0010] wherein, in Formula 1,

[0011] R₁, Ar₁, and Ar₂ may each independently be selected from hydrogen, deuterium, a halogen, an amino

group, a nitro group, a nitrile group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

[0012] L may each independently be selected from a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0013] m and n may each independently be an integer selected from 0 to 4,

[0014] when m and n are each independently two or more, two or more R₁(s) and two or more L(s) may each independently be identical to or different from each other, and

[0015] at least one substituent selected from a substituent (s) of the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and the substituted divalent non-aromatic condensed heteropolycyclic group may be selected from one of a first group, a second group, a third group, and a fourth group:

[0016] the first group including deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0017] the second group including a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio

group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

[0018] the third group including a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

[0019] the fourth group including a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇),

[0020] wherein Q₁₁ to Q₁₇ and Q₂₁ to Q₂₇ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0021] According to one or more embodiments, an organic light-emitting device includes: a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode and includes an emission layer, wherein the organic layer includes the compound of Formula 1.

[0022] According to one or more embodiments, a panel display apparatus includes the organic light-emitting device, wherein the first electrode of the organic light-emitting device is electrically connected to a source electrode or a drain electrode of a thin film transistor.

BRIEF DESCRIPTION OF THE DRAWINGS

[0023] These and/or other aspects will become apparent and more readily appreciated from the following description of the embodiments, taken in conjunction with the accompanying drawings in which:

[0024] FIG. 1 is a schematic view of an organic light-emitting device according to an embodiment;


[0025] FIG. 2 is a schematic view of an organic light-emitting device according to an embodiment;

[0026] FIG. 3 is a schematic view of an organic light-emitting device according to an embodiment; and

[0027] FIG. 4 is a schematic view of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION

[0028] A compound according to an embodiment may be represented by Formula 1:



[0029] wherein, in Formula 1,

[0030] R₁, Ar₁, and Ar₂ may each independently be selected from hydrogen, deuterium, a halogen, an amino group, a nitro group, a nitrile group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0031] L may independently be selected from a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.

[0032] m and n may each independently be an integer selected from 0 to 4,

[0033] when m and n are each independently two or more, two or more R₁(s) and two or more L(s) may each independently be identical to or different from each other, and

[0034] at least one substituent selected from a substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the

substituted C_6-C_{60} arylthio group, the substituted C_1-C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted C_6-C_{60} arylene group, the substituted C_1-C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and the substituted divalent non-aromatic condensed heteropolycyclic group may be selected from one of a first group, a second group, a third group, and a fourth group:

[0035] the first group including deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, and a C_1-C_{60} alkoxy group;

[0036] the second group including a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, and a C_1-C_{60} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_2-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_{11})(Q_{12})$, $-Si(Q_{13})(Q_{14})(Q_{15})$, and $-B(Q_{16})(Q_{17})$;

[0037] the third group including a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_2-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

[0038] the fourth group including a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_2-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_1-C_{60} alkoxy group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_2-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_{21})(Q_{22})$, $-Si(Q_{23})(Q_{24})(Q_{25})$, and $-B(Q_{26})(Q_{27})$,

[0039] wherein Q_{11} to Q_{17} and Q_{21} to Q_{27} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro

group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_1-C_{60} alkoxy group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_2-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0040] As materials for use in a hole injection layer and a hole transport layer of an organic light-emitting device, various types of arylamine compounds have been reported and applied thereto.


[0041] A representative material for use in the hole injection layer and the hole transport layer is an NPB compound structure, and compounds including biphenyl are used in the hole injection layer and the hole transport layer. The existing materials show excellent efficiency and lifespan characteristics when applied to an organic light-emitting device, but continuous improvement is required because the existing materials have problems such as denaturalization due to excessive thermal stress during deposition and problems in various processes due to melting of materials.

[0042] In particular, it is necessary to develop a high heat-resisting material that is not denaturalized even in a high temperature environment. One or more embodiments of the present disclosure are directed to propose a material for use in a hole injection layer and a hole transport layer, which uses a compound having a novel structure, and to implement an organic light-emitting device having high efficiency and a long lifespan based on thermal stability of the proposed material.


[0043] Substituents of Formula 1 will now be described in detail.

[0044] In one or more embodiments, n in Formula 1 may be 0 or 1.


[0045] In one or more embodiments, Ar_1 and Ar_2 in Formula 1 may each independently be one selected from groups represented by Formulae 2a to 2d:



2a




2b



2c

-continued



[0046] wherein H_1 in Formula 2c may be O, S, NR_{21} , or $CR_{22}R_{23}$,


[0047] R_{21} to R_{23} and Z_1 may each independently be selected from hydrogen, deuterium, a halogen, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_6 - C_{20} aryl group, a substituted or unsubstituted C_1 - C_{20} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

[0048] p may be an integer selected from 1 to 7,


[0049] when p is two or more, two or more $Z_1(s)$ may be identical to or different from each other, and

[0050] * indicates a binding site.

[0051] In one or more embodiments, R_1 in Formula 1 may each independently be one selected from hydrogen, deuterium, a cyano group, a halogen, or groups represented by Formulae 3a and 3b:



3a



3b


[0052] In Formulae 3a and 3b, $Z_1(s)$ may each independently be selected from hydrogen, deuterium, a halogen, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_6 - C_{20} aryl group, a substituted or unsubstituted C_1 - C_{20} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

[0053] p may be an integer selected from 1 to 7,

[0054] when p is two or more, two or more $Z_1(s)$ may be identical to or different from each other, and

[0055] * indicates a binding site.

[0056] In one or more embodiments, L in Formula 1 may be represented by Formula 4a:




4a

[0057] wherein * in Formula 4a indicates a binding site.

[0058] In one or more embodiments, the composition represented by Formula 1 may be represented by Formula 2:


<Formula 2>



[0059] wherein n in Formula 2 may be 0 or 1.


[0060] In one or more embodiments, the compound represented by Formula 1 may be represented by Formula 3:

<Formula 3>




[0061] In one or more embodiments, the compound represented by Formula 1 may be represented by Formula 4:

<Formula 4>



[0062] In one or more embodiments, the compound represented by Formula 1 may be represented by Formula 5:







<Formula 5>



[0063] In one or more embodiments, the compound represented by Formula 1 may be one selected from compounds illustrated below:

-continued


4



6


-continued

7




-continued


10



8





11




12

9




-continued

13




-continued


16




14




15



17




18




-continued


19



20




21




-continued


22



23




24




-continued


25



26




27




-continued

28




29




-continued

30




-continued


33




31




32



34




35




-continued

36




-continued


39




40




41



37




38




-continued


42



43




44




-continued

45




46




-continued

47




-continued


50



48





51




52

49




-continued


53



54




55




-continued


56



57




58




-continued


59



60




61




-continued


62




63







64



-continued




-continued




-continued


70



71




72




-continued


73



74




75




-continued


76



77




78




-continued

79



80



[0064] [Description of FIG. 1]

[0065] FIG. 1 is a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 includes a first electrode 110, an organic layer 150, and a second electrode 190.

[0066] Hereinafter, the structure of the organic light-emitting device 10 according to an embodiment and a method of manufacturing the organic light-emitting device 10 will be described in connection with FIG. 1.

[First Electrode 110]

[0067] In FIG. 1, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be a glass substrate or a plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water-resistance.

[0068] The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode 110 on the substrate. When the first electrode 110 is an anode, the material for the first electrode 110 may be selected from materials with a high work function to facilitate hole injection.

[0069] The first electrode 110 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode 110 is a transmissible electrode, a material for forming the first electrode 110 may be selected from indium tin oxide (ITO), indium zinc oxide

(IZO), tin oxide (SnO_2), zinc oxide (ZnO), and any combinations thereof, but embodiments of the present disclosure are not limited thereto. In one or more embodiments, when the first electrode **110** is a semi-transmissible electrode or a reflectable electrode, a material for forming the first electrode **110** may be selected from magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and any combinations thereof, but embodiments of the present disclosure are not limited thereto.

[0070] The first electrode **110** may have a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode **110** may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode **110** is not limited thereto.

[Organic Layer **150**]

[0071] The organic layer **150** is disposed on the first electrode **110**. The organic layer **150** may include an emission layer.

[0072] The organic layer **150** may further include a hole transport region between the first electrode **110** and the emission layer, and an electron transport region between the emission layer and the second electrode **190**.

[Hole Transport Region in Organic Layer **150**]

[0073] The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0074] The hole transport region may include at least one layer selected from a hole injection layer, a hole transport layer, an emission auxiliary layer, and an electron blocking layer.



[0075] For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure, wherein for each structure, constituting layers are sequentially stacked from the first electrode **110** in this stated order, but the structure of the hole transport region is not limited thereto.

[0076] In one or more embodiments, the hole transport region may include the compound of Formula 1 according to an embodiment.


[0077] In one or more embodiments, the hole transport layer or the hole injection layer may include the compound of Formula 1 according to an embodiment.

[0078] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB

(NPB), β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTDP, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), polyaniline/poly(4-styrenesulfonate) (Pani/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:




-continued




2-TNATA


-continued




Spiro-TPD





Spiro-NPB




NPB




methylated NPB

 β -NPB


TAPC



TPD



HMTDP



[0079] In Formulae 201 and 202,

[0080] L₂₀₁ to L₂₀₄ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

[0081] L₂₀₅ may be selected from *—O—*, *—S—*, *—N(Q₂₀₁)—*, a substituted or unsubstituted C₁-C₂₀ alkylene group, a substituted or unsubstituted C₂-C₂₀ alkenylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.

[0082] xa1 to xa4 may each independently be an integer selected from 0 to 3.

[0083] xa5 may be an integer selected from 1 to 10, and

[0084] R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0085] For example, R₂₀₁ and R₂₀₂ in Formula 202 may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group, and R₂₀₃ and R₂₀₄ may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group.

[0086] In one or more embodiments, regarding Formulae 201 and 202,

[0087] L₂₀₁ to L₂₀₅ may each independently be selected from one of a fifth group and a sixth group:

[0088] the fifth group including a phenylene group, a pentalenylene group, an indenylene group, a naphthylene

group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolyene group, an indolyene group, an isoindolyene group, a benzofuranylene group, a benzothiofenylene group, a dibenzofuranylene group, a dibenzothiofenylene group, a benzocarbazolyene group, a dibenzocarbazolyene group, a dibenzosilolylene group, and a pyridinylene group; and

[10089] the sixth group including a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenylyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranlyl group, a benzothiophenyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q_{31})(Q_{32})(Q_{33}), and —N(Q_{31})(Q_{32}).

[0090] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0091] In one or more embodiments, xa1 to xa4 may each independently be 0, 1, or 2.

[0092] In one or more embodiments, xa5 may be 1, 2, 3, or 4.

[0093] In one or more embodiments, R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from one of a seventh group and an eighth group;

[0094] the seventh group including a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0095] the eighth group including a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl

group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂).

[0096] wherein Q₃₁ to Q₃₃ are the same as described above.

[0097] In one or more embodiments, at least one material selected from R₂₀₁ to R₂₀₃ in Formula 201 may each independently be selected from one of a ninth group and a tenth group:

[0098] the ninth group including a fluorenyl group, a spiro-bifluorenyl group, a carbazoyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0099] the tenth group including a fluorenyl group, a spiro-bifluorenyl group, a carbazoyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazoyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0100] but embodiments of the present disclosure are not limited thereto.

[0101] In one or more embodiments, in Formula 202, i) R₂₀₁ and R₂₀₂ may be linked via a single bond, and/or ii) R₂₀₃ and R₂₀₄ may be linked via a single bond.


[0102] In one or more embodiments, at least one of R₂₀₁, R₂₀₂, R₂₀₃, and R₂₀₄ in Formula 202 may be selected from an eleventh group and a twelfth group:

[0103] the eleventh group including a carbazoyl group; and


[0104] the twelfth group including a carbazoyl group, substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazoyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0105] but embodiments of the present disclosure are not limited thereto.


[0106] The compound represented by Formula 201 may be represented by Formula 201A:



[0107] In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A(1) below, but embodiments of the present disclosure are not limited thereto:




[0108] In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A-1 below, but embodiments of the present disclosure are not limited thereto:




[0109] In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A:

<Formula 202A>



[0110] In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A-1:

<Formula 202A-1>



[0111] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,


[0112] L_{201} to L_{203} , $xa1$ to $xa3$, $xa5$, and R_{202} to R_{204} are the same as described above,

[0113] R_{211} and R_{212} are the same as described in connection with R_{203} , and


[0114] R_{213} to R_{217} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-F$, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0115] The hole transport region may include at least one compound selected from Compounds HT1 to HT39, but embodiments of the present disclosure are not limited thereto:


HT1




HT2



HT3





HT4





-continued
HT5

HT6



HT7



HT8



-continued



HT9

HT10




HT11


HT12




-continued
HT13




HT14




HT15




HT16




HT17




HT18




-continued
HT19




HT20




HT21



HT22





HT23



-continued



HT24

HT25





HT26

HT27




HT28

HT29




-continued


HT30




HT31




HT32




HT33



HT34





HT35





-continued
HT36

HT37



HT38

HT39



[0116] A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 9,000 Å, for example, about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, for example about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0117] The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer, and the electron blocking layer may block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include the materials as described above.

[p-Dopant]

[0118] The hole transport region may further include, in addition to these materials, a charge-generation material for

the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0119] The charge-generation material may be, for example, a p-dopant.

[0120] In one or more embodiments, a lowest unoccupied molecular orbital (LUMO) level of the p-dopant may be -3.5 eV or less.

[0121] The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but embodiments of the present disclosure are not limited thereto.

[0122] For example, the p-dopant may include at least one selected from:


[0123] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) and 2,3,5,6-Tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);

[0124] a metal oxide, such as a tungsten oxide or a molybdenum oxide;


[0125] 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and

[0126] a compound represented by Formula 221 below:


[0127] but embodiments of the present disclosure are not limited thereto:



<HAT-CN>



<F4-TCNQ>



<Formula 221>

In Formula 221,

[0128] R₂₂₁ to R₂₂₃ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and at least one selected from R₂₂₁ to R₂₂₃ may have at least one substituent selected from a cyano group, —F, —Cl, —Br, —I, a C₁-C₂₀ alkyl group substituted with —F, a C₁-C₂₀ alkyl group substituted with —Cl, a C₁-C₂₀ alkyl group substituted with —Br, and a C₁-C₂₀ alkyl group substituted with —I.

[Emission Layer in Organic Layer 150]

[0129] When the organic light-emitting device 10 is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In one or more embodiments, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers contact each other or are separated from each other. In one or more embodiments,

the emission layer may include two or more materials selected from a red-light emission material, a green-light emission material, and a blue-light emission material, in which the two or more materials are mixed with each other in a single layer to emit white light.


[0130] The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant and a fluorescent dopant.

[0131] An amount of the dopant in the emission layer may be, in general, in a range of about 0.01 to about 15 parts by weight based on 100 parts by weight of the host, but embodiments of the present disclosure are not limited thereto.

[0132] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[Host in Emission Layer]

[0133] In one or more embodiments, the host may include a compound represented by Formula 301 below.



<Formula 301>

[0134] In Formula 301,

[0135] Ar₃₀₁ may be a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₁-C₆₀ heterocyclic group,

[0136] xb11 may be 1, 2, or 3,

[0137] L₃₀₁ may be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0138] xb1 may be an integer selected from 0 to 5,

[0139] R₃₀₁ may be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₃₀₁)(Q₃₀₂)(Q₃₀₃), —N(Q₃₀₁)(Q₃₀₂), —B(Q₃₀₁)(Q₃₀₂), —C(=O)(Q₃₀₁), —S(=O)₂(Q₃₀₁), and —P(=O)(Q₃₀₁)(Q₃₀₂),

[0140] xb21 may be an integer selected from 1 to 5,

[0141] Q₃₀₁ to Q₃₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl

group, a biphenyl group, a terphenyl group, and a naphthyl group, but are not limited thereto.

[0142] In one or more embodiments, Ar_{301} in Formula 301 may be selected from:



[0143] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group and a dibenzothiophene group; and

[0144] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})(Q_{32})$,

[0145] wherein Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but are limited thereto.

[0146] When $xb11$ in Formula 301 is two or more, two or more $Ar_{301}(s)$ may be linked via a single bond.

[0147] In one or more embodiments, the compound represented by Formula 301 may be represented by one of Formulae 301-1 and 301-2:



[0148] In Formulae 301-1 to 301-2,

[0149] A_{301} to A_{304} may each independently be selected from a benzene, a naphthalene, a phenanthrene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a pyridine, a pyrimidine, an indene, a fluorene, a spiro-bifluorene, a benzofluorene, a dibenzofluorene, an indole, a carbazole,

benzocarbazole, dibenzocarbazole, a furan, a benzofuran, a dibenzofuran, a naphthofuran, a benzonaphthofuran, a dinaphthofuran, a thiophene, a benzothiophene, a dibenzothiophene, a naphthothiophene, a benzonaphthothiophene, and a dinaphthothiophene,

[0150] X_{301} may be O, S, or $N-[(L_{304})_{xb4}-R_{304}]$,

[0151] R_{311} to R_{314} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})(Q_{32})$,

[0152] $xb22$ and $xb23$ may each independently be 0, 1, or 2,

[0153] L_{301} , $xb1$, R_{301} , and Q_{31} to Q_{33} are the same as described above,

[0154] L_{302} to L_{304} are each independently the same as described in connection with L_{301} ,

[0155] $xb2$ to $xb4$ are each independently the same as described in connection with $xb1$, and

[0156] R_{302} to R_{304} are each independently the same as described in connection with R_{301} .

[0157] For example, L_{301} to L_{304} in Formulae 301, 301-1, and 301-2 may each independently be selected from one of a thirteenth and a fourteenth group:

[0158] the thirteenth group including a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an

isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an

oxazolylenes group, an isoxazolylenes group, a thiadiazolylenes group, an oxadiazolylenes group, a pyrazinylenes group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylenes group, an isobenzothiazolylenes group, a benzoxazolylenes group, an isobenzoxazolylenes group, a triazolylenes group, a tetrazolylenes group, an imidazopyrimidinylene group, an imidazopyrimidinylene group, and an azacarbazolylenes group; and

[0159] the fourteenth group including a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylenes group, an indolylenes group, an isoindolylenes group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenes group, a dibenzocarbazolylenes group, a dibenzosilolylenes group, a pyridinylene group, an imidazolylenes group, a pyrazolylenes group, a thiazolylenes group, an isothiazolylenes group, an oxazolylenes group, an isoxazolylenes group, a thiadiazolylenes group, an oxadiazolylenes group, a pyrazinylenes group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylenes group, an isobenzothiazolylenes group, a benzoxazolylenes group, an isobenzoxazolylenes group, a triazolylenes group, a tetrazolylenes group, an imidazopyrimidinylene group, and an azacarbazolylenes group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolylenes group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolylenes group, a dibenzocarbazolylenes group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyrimidinyl group, an imidazopyrimidinyl group, and an azacarbazolylenes group, each substi-

group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyrimidinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}).

[0160] wherein Q_{31} to Q_{33} are the same as described above.

[0161] In one or more embodiments, R_{301} to R_{304} in Formulae 301, 301-1, and 301-2 may each independently be selected from one of a fifteenth group and a sixteenth group:

[0162] the fifteenth group including a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolylenes group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolylenes group, a dibenzocarbazolylenes group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyrimidinyl group, an imidazopyrimidinyl group, and an azacarbazolylenes group; and


[0163] the sixteenth group including a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolylenes group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolylenes group, a dibenzocarbazolylenes group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyrimidinyl group, an imidazopyrimidinyl group, and an azacarbazolylenes group, each substi-

tuted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),


[0164] wherein Q_{31} to Q_{33} are the same as described above.

[0165] In one or more embodiments, the host may include an alkaline-earth metal complex. For example, the host may be selected from a Be complex (for example, Compound H55), an Mg complex, and a Zn complex.

[0166] The host may include at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di-(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and one of Compounds H1 to H55, but embodiments of the present disclosure are not limited thereto:




H1




H2


-continued




H3




H4




H5



H6




H7




H8

-continued


H9




H10




H11



H12




H13




-continued


H14




H15




H16



H17




H18




-continued


H19




H20



H21




H22




-continued


H23



H24




H25




-continued


H26



H27




H28




-continued


H29




H30



H31




H32




-continued

H33




-continued


H37




H34




H38




H35




H36



H39




-continued




H40


-continued




H43




H41




H44



H42




H45




H46

-continued


H47




H48




H49



H50




H51




-continued


H52




H53



H54




H55




[Phosphorescent Dopant included in Emission Layer in Organic Layer **150**]

[0167] The phosphorescent dopant may include an organometallic complex represented by Formula 401 below:



<Formula 401>

<Formula 402>



[0168] In Formulae 401 and 402,

[0169] M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

[0170] L_{401} may be selected from ligands represented by Formula 402, and xc1 may be 1, 2, or 3, wherein, when xc1 is two or more, two or more L_{401} (s) may be identical to or different from each other,

[0171] L_{402} may be an organic ligand, and xc2 may be an integer selected from 0 to 4, wherein, when xc2 is two or more, two or more L_{402} (s) may be identical to or different from each other,

[0172] X_{401} to X_{404} may each independently be nitrogen or carbon,

[0173] X_{401} and X_{403} may be linked via a single bond or a double bond, and X_{402} and X_{404} may be linked via a single bond or a double bond,

[0174] A_{401} and A_{402} may each independently be a C_5 - C_{60} carbocyclic group or a C_1 - C_{60} heterocyclic group,

[0175] X_{405} may be a single bond, $*-O-*$, $*-S-*$, $*-C(=O)-*$, $*-N(Q_{411})-*$, $*-C(Q_{411})(Q_{412})-*$, $*-C(Q_{411})=C(Q_{412})-*$, $*-C(Q_{411})=*$, or $*=C(Q_{411})=*$, wherein Q_{411} and Q_{412} may be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

[0176] X_{406} may be a single bond, O, or S,

[0177] R_{401} and R_{402} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_1 - C_{20} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group $-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})(Q_{402})$, $-B(Q_{401})(Q_{402})$, $-C(=O)(Q_{401})$, $-S(=O)_2(Q_{401})$, and $-P(=O)(Q_{401})(Q_{402})$, wherein Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a C_6 - C_{20} aryl group, and a C_1 - C_{20} heteroaryl group,

[0178] xc11 and xc12 may each independently be an integer selected from 0 to 10, and

[0179] $*$ and $*'$ in Formula 402 may each be a binding site to M in Formula 401.

[0180] In one or more embodiments, A_{401} and A_{402} in Formula 402 may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiothiophene group, an isobenzothiothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiothiophene group.

[0181] In one or more embodiments, in Formula 402, i) X_{401} may be nitrogen, and X_{402} may be carbon, or ii) X_{401} and X_{402} may each be nitrogen at the same time.

[0182] In one or more embodiments, R_{401} and R_{402} in Formula 402 may each independently be selected from:

[0183] hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0184] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

[0185] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0186] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and


[0187] $-\text{Si}(\text{Q}_{401})(\text{Q}_{402})(\text{Q}_{403})$, $-\text{N}(\text{Q}_{401})(\text{Q}_{402})$, $-\text{B}(\text{Q}_{401})(\text{Q}_{402})$, $-\text{C}(=\text{O})(\text{Q}_{401})$, $-\text{S}(=\text{O})_2(\text{Q}_{401})$, and $-\text{P}(=\text{O})(\text{Q}_{401})(\text{Q}_{402})$,

[0188] wherein Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.


[0189] In one or more embodiments, when xc1 in Formula 401 is two or more, two $\text{A}_{401}(\text{s})$ in two or more $\text{L}_{401}(\text{s})$ may optionally be linked via X_{407} , which is a linking group, or two $\text{A}_{402}(\text{s})$ in two or more $\text{L}_{401}(\text{s})$ may optionally be linked via X_{408} , which is a linking group (see Compounds PD1 to PD4 and PD7). X_{407} and X_{408} may each independently be a single bond, $^*\text{—O—}^*$, $^*\text{—S—}^*$, $^*\text{—C(=O)—}^*$, $^*\text{—N}(\text{Q}_{413})\text{—}^*$, $^*\text{—C}(\text{Q}_{413})(\text{Q}_{414})\text{—}^*$, or $^*\text{—C}(\text{Q}_{413})=\text{C}(\text{Q}_{414})\text{—}^*$ (wherein Q_{413} and Q_{414} may each independently be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group), but embodiments of the present disclosure are not limited thereto.

[0190] L_{402} in Formula 401 may be a monovalent, divalent, or trivalent organic ligand. For example, L_{402} may be selected from a halogen, a diketone (for example, acetylacetonate), a carboxylic acid (for example, picolinate), $-\text{C}(=\text{O})$, an isonitrile, $-\text{CN}$, and phosphorus (for example, phosphine or phosphite), but embodiments of the present disclosure are not limited thereto.

[0191] In one or more embodiments, the phosphorescent dopant may be selected from, for example, Compounds PD1 to PD25, but embodiments of the present disclosure are not limited thereto:




PD1




PD2


-continued




PD3




PD4




PD5



PD6








PD7




PD8

-continued




-continued

PD9




PD14

PD10




PD15

PD11




PD16

PD12







PD17

PD13




PD18

-continued




-continued

PD19



PD23


PD20



PD24

PD25

PD21




[Fluorescent Dopant in Emission Layer]

[0192] The fluorescent dopant may include an arylamine compound or a styrylamine compound.

PD22

[0193] The fluorescent dopant may include a compound represented by Formula 501 below.

<Formula 501>



[0194] In Formula 501,

[0195] Ar_{501} may be a substituted or unsubstituted $\text{C}_5\text{-C}_{60}$ carbocyclic group or a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heterocyclic group,

[0196] L_{501} to L_{503} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0197] $xd1$ to $xd3$ may each independently be an integer of 0 to 3;

[0198] R_{501} and R_{502} may each independently be a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

[0199] $xd4$ may be an integer selected from 1 to 6.

[0200] In one or more embodiments, Ar_{501} in Formula 501 may be selected from:

[0201] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthacene group, a picene group, a perylene group, a pentaphene group, an indenanthracene group, and an indenophenanthrene group; and

[0202] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthacene group, a picene group, a perylene group, a pentaphene group, an indenanthracene group, and indenophenanthrene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0203] In one or more embodiments, L_{501} to L_{503} in Formula 501 may each independently be selected from:

[0204] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group; and

[0205] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0206] In one or more embodiments, R_{501} and R_{501} in Formula 502 may each independently be selected from:

[0207] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and


[0208] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$.


[0209] wherein Q_{31} to Q_{33} may be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0210] In one or more embodiments, xd4 in Formula 501 may be 2, but embodiments of the present disclosure are not limited thereto.


[0211] For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:




FD1



FD2




FD3



FD4


-continued

FD5




-continued


FD8




FD9




FD6




FD10




FD7



FD11




-continued




FD12


-continued




FD16




FD13




FD17




FD18



FD14




FD15




FD19

-continued

FD20




FD21




-continued

FD22





[0212] In one or more embodiments, the fluorescent dopant may be selected from the following compounds, but embodiments of the present disclosure are not limited thereto.

DPVBi



DPAVB





[Electron Transport Region in Organic Layer 150]

[0213] The electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0214] The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer, but embodiments of the present disclosure are not limited thereto.

[0215] For example, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, wherein for each structure, constituting layers are sequentially stacked from an emission layer. However, embodiments of the structure of the electron transport region are not limited thereto.

[0216] The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may

include a metal-free compound containing at least one π electron-depleted nitrogen-containing ring.

[0217] The “ π electron-depleted nitrogen-containing ring” indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

[0218] For example, the “ π electron-depleted nitrogen-containing ring” may be i) a 5-membered to 7-membered hetero monocyclic group having at least one $*-N=*$ moiety, ii) a heteropoly cyclic group in which two or more 5-membered to 7-membered hetero monocyclic groups each having at least one $*-N=*$ moiety are condensed with each other, or iii) a heteropoly cyclic group in which at least one of 5-membered to 7-membered hetero monocyclic groups, each having at least one $*-N=*$ moiety, is condensed with at least one C_5 - C_{60} carbocyclic group.

[0219] Examples of the π electron-depleted nitrogen-containing ring include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a

triazine, thiadiazol, an imidazopyridine, an imidazopyrimidine, and an azacarbazole, but are not limited thereto.

[0220] For example, the electron transport region may include a compound represented by Formula 601:



[0221] In Formula 601,

[0222] Ar_{601} may be a substituted or unsubstituted $\text{C}_5\text{-C}_{60}$ carbocyclic group or a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heterocyclic group,

[0223] xe11 may be 1, 2, or 3,

[0224] L_{601} is selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkylene group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenylene group, a substituted or unsubstituted heterocycloalkenylene group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0225] xe1 may be an integer selected from 0 to 5,

[0226] R_{601} may be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryloxy group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylthio group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_{601})(\text{Q}_{602})(\text{Q}_{603})$, $-\text{C}(=\text{O})(\text{Q}_{601})$, $-\text{S}(=\text{O})_2(\text{Q}_{601})$, and $-\text{P}(=\text{O})(\text{Q}_{601})(\text{Q}_{602})$,

[0227] Q_{601} to Q_{603} may each independently be a $\text{C}_1\text{-C}_{10}$ alkyl group, a $\text{C}_1\text{-C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

[0228] xe21 may be an integer selected from 1 to 5.

[0229] In one or more embodiments, at least one selected from $\text{Ar}_{601}(\text{s})$ in the number of xe11 and $\text{R}_{601}(\text{s})$ in the number of xe21 may include a π electron-depleted nitrogen-containing ring as described above.

[0230] In one or more embodiments, Ar_{601} in Formula 601 may be selected from:

[0231] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthalene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a

triazine group, a thiadiazol group, an imidazo pyridine group, an imidazo pyrimidine group, and an azacarbazole group; and


[0232] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthalene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, phenanthroline group, phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazol group, an imidazo pyridine group, an imidazo pyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{20}$ alkyl group, a $\text{C}_1\text{-C}_{20}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,

[0233] wherein Q_{31} to Q_{33} may each independently be selected from a $\text{C}_1\text{-C}_{10}$ alkyl group, a $\text{C}_1\text{-C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0234] When xe11 in Formula 601 is two or more, two or more $\text{Ar}_{601}(\text{s})$ may be linked via a single bond.

[0235] In one or more embodiments, Ar_{601} in Formula 601 may be an anthracene group.

[0236] In one or more embodiments, a compound represented by Formula 601 may be represented by Formula 601-1:



[0237] In Formula 601-1,

[0238] X_{614} may be N or C(R_{614}), X_{615} may be N or C(R_{615}), X_{616} may be N or C(R_{616}), and at least one selected from X_{614} to X_{616} may be N,

[0239] L_{611} to L_{613} may be each independently substantially the same as described in connection with L_{601} ,

[0240] xe611 to xe613 may be each independently substantially the same as described in connection with xe1 ,

[0241] R_{611} to R_{613} may be each independently substantially the same as described in connection with R_{601} ,

[0242] R_{614} to R_{616} may each independently be selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl

group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0243] In one or more embodiments, L₆₀₁ and L₆₁₁ to L₆₁₃ in Formulae 601 and 601-1 may each independently be selected from:

[0244] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylenylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylene group, and an azacarbazolylenylene group; and

[0245] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylenylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylene group, and an

azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indoyl group, an isoindoyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosiloyl group, a pyridinyl group, an imidazoyl group, a pyrazoyl group, a thiazoyl group, an isothiazoyl group, an oxazoyl group, an isoxazoyl group, a thiadiazoyl group, an oxadiazoyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazoyl group, an isobenzothiazoyl group, a benzoxazoyl group, an isobenzoxazoyl group, a triazoyl group, a tetrazoyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group;

[0246] but embodiments of the present disclosure are not limited thereto.

[0247] In one or more embodiments, xe1 and xe611 to xe613 in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

[0248] In one or more embodiments, R₆₀₁ and R₆₁₁ to R₆₁₃ in Formula 601 and 601-1 may each independently be selected from:

[0249] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indoyl group, an isoindoyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosiloyl group, a pyridinyl group, an imidazoyl group, a pyrazoyl group, a thiazoyl group, an isothiazoyl group, an oxazoyl group, an isoxazoyl group, a thiadiazoyl group, an oxadiazoyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazoyl group, an isobenzothiazoyl group, a benzoxazoyl group, an isobenzoxazoyl group, a triazoyl group, a tetrazoyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group;


[0250] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacetyl group, a pentacacetyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacetyl group, a pentacacetyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

[0251] —S(=O)₂(Q₆₀₁) and —P(=O)(Q₆₀₁)(Q₆₀₂),


[0252] wherein Q₆₀₁ and Q₆₀₂ are the same as described above.

[0253] The electron transport region may include at least one compound selected from Compounds ET1 to ET36, but embodiments of the present disclosure are not limited thereto:


ET1



ET2




ET3




-continued

ET4




-continued

ET7





ET5




ET8

ET6




-continued




ET9

-continued





ET11

ET12




ET10

ET13




-continued




ET14

-continued




ET17


ET15




ET18




ET16



ET19




-continued



ET20



-continued



ET23



ET24

ET21




ET25

ET22




-continued

ET26




-continued


ET29




ET27




ET30




ET28



ET31




-continued




ET32

-continued




ET35

ET33






ET36



[0254] In one or more embodiments, the electron transport region may include at least one compound selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq₃, BAlq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ.

ET34

Alq₃



[0255] A thickness of the buffer layer, the hole blocking layer, or the electron control layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.

[0256] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.


[0257] The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.

[0258] The metal-containing material may include at least one selected from alkali metal complex and alkaline earth-metal complex. The alkali metal complex may include a metal ion selected from an Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion, and the alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, an Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy


phenylan oxazole, a hydroxy phenylthiazole, a hydroxy diphenylan oxadiazole, a hydroxy diphenylthiadiazol, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

[0259] For example, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.

ET-D1



ET-D2



[0260] The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode 190. The electron injection layer may directly contact the second electrode 190.

[0261] The electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0262] The electron injection layer may include alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof.

[0263] The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one or more embodiments, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs, but embodiments of the present disclosure are not limited thereto.

[0264] The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

[0265] The rare-earth metal may be selected from Sc, Y, Ce, Tb, Yb, Gd, and Tb.

[0266] The alkali metal compound, the alkaline earth-metal compound, and the rare-earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal and rare-earth metal.

[0267] The alkali metal compound may be selected from alkali metal oxides, such as Li_2O , Cs_2O , or K_2O , and alkali metal halides, such as LiF, NaF, CsF, KF, LiI, NaI, CsI, or

KI. In one or more embodiments, the alkali metal compound may be selected from LiF, Li₂O, NaF, LiI, NaI, CsI, and KI, but embodiments of the present disclosure are not limited thereto.

[0268] The alkaline earth-metal compound may be selected from alkaline earth-metal compounds, such as BaO, SrO, CaO, Ba_xSr_{1-x}O (0<x<1), and Ba_xCa_{1-x}O (0<x<1). In one or more embodiments, the alkaline earth-metal compound may be selected from BaO, SrO, and CaO, but embodiments of the present disclosure are not limited thereto.

[0269] The rare-earth metal compound may be selected from YbF₃, ScF₃, ScO₃, Y₂O₃, Ce₂O₃, GdF₃, and TbF₃. In one or more embodiments, the rare-earth metal compound may be selected from YbF₃, ScF₃, TbF₃, YbI₃, ScI₃, and TbI₃, but embodiments of the present disclosure are not limited thereto.

[0270] The alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare-earth metal as described above, and a ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may each independently be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenylan oxazole, hydroxy phenylthiazole, hydroxy diphenylan oxadiazole, hydroxy diphenylthiadiazol, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

[0271] The electron injection layer may consist of alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof, as described above. In one or more embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

[0272] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

[Second Electrode 190]

[0273] The second electrode 190 may be disposed on the organic layer 150 having such a structure. The second electrode 190 may be a cathode which is an electron injection electrode, and in this regard, a material for forming the second electrode 190 may be selected from metal, an alloy, an electrically conductive compound, and a mixture thereof, which have a relatively low work function.

[0274] The second electrode 190 may include at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO, but embodiments of the present disclosure are not limited thereto. The second electrode 190 may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

[0275] The second electrode 190 may have a single-layered structure, or a multi-layered structure including two or more layers.

[Description of FIGS. 2 to 4]

[0276] An organic light-emitting device 20 of FIG. 2 includes a first capping layer 210, a first electrode 110, an organic layer 150, and a second electrode 190 which are sequentially stacked in this stated order, an organic light-emitting device 30 of FIG. 3 includes a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220 which are sequentially stacked in this stated order, and an organic light-emitting device 40 of FIG. 4 includes a first capping layer 210, a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220.

[0277] Regarding FIGS. 2 to 4, the first electrode 110, the organic layer 150, and the second electrode 190 may be understood by referring to the description presented in connection with FIG. 1.

[0278] In the organic layer 150 of each of the organic light-emitting devices 20 and 40, light generated in an emission layer may pass through the first electrode 110, which is a semi-transmissive electrode or a transmissive electrode, and the first capping layer 210 toward the outside, and in the organic layer 150 of each of the organic light-emitting devices 30 and 40, light generated in an emission layer may pass through the second electrode 190, which is a semi-transmissive electrode or a transmissive electrode, and the second capping layer 220 toward the outside.

[0279] The first capping layer 210 and the second capping layer 220 may increase external luminescent efficiency according to the principle of constructive interference.


[0280] The first capping layer 210 and the second capping layer 220 may each independently be an organic capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.

[0281] At least one selected from the first capping layer 210 and the second capping layer 220 may each independently include at least one material selected from carbocyclic compounds, heterocyclic compounds, amine-based compounds, porphyrine derivatives, phthalocyanine derivatives, naphthalocyanine derivatives, alkali metal complexes, and alkaline earth-based complexes. The carbocyclic compound, the heterocyclic compound, and the amine-based compound may be optionally substituted with a substituent containing at least one element selected from O, N, S, Se, Si, F, Cl, Br, and I. In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include an amine-based compound.

[0282] In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer


220 may each independently include the compound represented by Formula 201 or the compound represented by Formula 202.

[0283] In one or more embodiments, at least one selected from the first capping layer **210** and the second capping layer **220** may each independently include a compound selected from Compounds HT28 to HT33 and Compounds CP1 to CP5, but embodiments of the present disclosure are not limited thereto.



-continued

CP5



[0284] Hereinbefore, the organic light-emitting device according to an embodiment has been described in connection with FIGS. 1-4. However, embodiments of the present disclosure are not limited thereto.

[0285] Layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region may be formed in a certain region by using one or more suitable methods selected from vacuum deposition, spin coating, casting, langmuir-blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

[0286] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of about 100 to about 500° C., at a vacuum degree of about 10^{-8} to about 10^{-3} torr, and at a deposition rate of about 0.01 to about 100 Å/sec by taking into account a compound to be included in a layer to be formed, and the structure of a layer to be formed.

[0287] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to 200° C. by taking into account a compound to be included in a layer to be formed, and the structure of a layer to be formed.

[General Definition of Substituents]

[0288] Hereinafter, definitions of substituents of compounds used herein will be presented (the number of carbon atoms used to restrict a substituent is not limited, and does not limit properties of the substituent, and unless defined otherwise, the definition of the substituent is consistent with a general definition thereof).

[0289] The term “C₁-C₆₀ alkyl group” as used herein refers to a linear or branched saturated aliphatic saturated hydrocarbon monovalent group having 1 to 60 carbon atoms, and examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “C₁-C₆₀ alkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₆₀ alkyl group.

[0290] The term “C₂-C₆₀ alkenyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon double bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethenyl

group, a propenyl group, and a butenyl group. The term “C₂-C₆₀ alkenylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkenyl group.

[0291] The term “C₂-C₆₀ alkynyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethynyl group and a propynyl group. The term “C₂-C₆₀ alkynylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

[0292] The term “C₁-C₆₀ alkoxy group” as used herein refers to a monovalent group represented by —OA101 (wherein A101 is the C₁-C₆₀ alkyl group), and examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

[0293] The term “C₃-C₁₀ cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

[0294] The term “C₁-C₁₀ heterocycloalkyl group” used herein refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuranyl group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group,” used herein, refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

[0295] The term “C₃-C₁₀ cycloalkenyl group” used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and does not have aromaticity, and examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group,” used herein, refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

[0296] The term “C₁-C₁₀ heterocycloalkenyl group” used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Examples of the C₁-C₁₀ heterocycloalkenyl group are a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranyl group and a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group,” used herein, refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkenyl group.

[0297] The term “C₆-C₆₀ aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C₆-C₆₀ arylene group used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Examples of the C₆-C₆₀ aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be fused to each other.

[0298] The term “C₁-C₆₀ heteroaryl group” used herein refers to a monovalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C₁-C₆₀ heteroarylene group” used herein refers to a divalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Examples of the C₁-C₆₀ heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinoxalinyl group, and an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be fused to each other.

[0299] The term “C₆-C₆₀ aryloxy group,” used herein, indicates —OA102 (wherein A102 is the C₆-C₆₀ aryl group), and a C₆-C₆₀ arylthio group indicates —SA103 (wherein A103 is the C₆-C₆₀ aryl group).

[0300] The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed with each other, only carbon atoms as a ring-forming atom, and non-aromaticity in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group,” used herein, refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0301] The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group (for example, having 1 to 60 carbon atoms) that has two or more rings condensed to each other, has at least one heteroatom selected from N, O, Si, P, and S, other than carbon atoms, as a ring-forming atom, and has non-aromaticity in the entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group,” used herein, refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0302] The term “C₅-C₆₀ carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which a ring-forming atom is a carbon atom only. The C₅-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₅-C₆₀ carbocyclic group may be a ring, such as a benzene group, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In one or more embodiments, depending on the number of substituents connected to the C₅-C₆₀ carbocyclic group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

[0303] The term “C₁-C₆₀ heterocyclic group” used herein refers to a group having the same structure as the C₅-C₆₀ carbocyclic group, except that as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used in addition to carbon (the number of carbon atoms may be in a range of 1 to 60).

[0304] At least one substituent selected from a substituent (s) of the substituted C₅-C₆₀ carbocyclic group, the substituted C₁-C₆₀ heterocyclic group, the substituted C₃-C₁₀

cycloalkylene group, the substituted C_1 - C_{10} heterocycloalkylene group, the substituted C_3 - C_{10} cycloalkenylene group, the substituted C_1 - C_{10} heterocycloalkenylene group, the substituted C_6 - C_{60} arylene group, the substituted C_1 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_1 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0305] deuterium(-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0306] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_{11})(Q_{12})(Q_{13}), —N(Q_{11})(Q_{12}), —B(Q_{11})(Q_{12}), —C(=O)(Q_{11}), —S(=O)₂(Q_{11}), and —P(=O)(Q_{11})(Q_{12});

[0307] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0308] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_{21})(Q_{22})(Q_{23}), —N(Q_{21})(Q_{22}), —B(Q_{21})(Q_{22}), —C(=O)(Q_{21}), —S(=O)₂(Q_{21}), and —P(=O)(Q_{21})(Q_{22}); and

[0309] —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}).

[0310] wherein Q_{11} to Q_{13} , Q_{21} to Q_{23} and Q_{31} to Q_{33} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

[0311] The term “Ph” as used herein represents a phenyl group, the term “Me” as used herein represents a methyl group, the term “Et” as used herein represents an ethyl group, the term “ter-Bu” or “But” as used herein, represents a tert-butyl group, and the term “OMe” as used herein represents a methoxy group.

[0312] The term “biphenyl group” used herein refers to a “phenyl group substituted with a phenyl group”. The “biphenyl group” is a “substituted phenyl group” having a “ C_6 - C_{60} aryl group” as a substituent.

[0313] The term “terphenyl group” used herein refers to a “phenyl group substituted with a biphenyl group”. The term “terphenyl group” is a “substituted phenyl group” having a “ C_6 - C_{60} aryl group substituted with a C_6 - C_{60} aryl group” as a substituent.

[0314] * and * used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.


[0315] Hereinafter, a compound according to embodiments and an organic light-emitting device according to embodiments will be described in detail with reference to Synthesis Examples and Examples. The wording “B was used instead of A” used in describing Synthesis Examples indicates that an identical molar equivalent of B was used in place of A.


SYNTHESIS EXAMPLE

Synthesis Example

Synthesis of Intermediate A

[0316]





Synthesis of Intermediate A-1


[0317] 9 g (50 mmol) of 9-fluorenone was dissolved in 200 ml of anhydrous tetrahydrofuran (THF) under a nitrogen atmosphere at a temperature of 0° C., and 11.3 g of 4-nitrophenyl magnesium bromide was added thereto for 1 hour in a dropwise fashion. The obtained reaction solution was heated to room temperature and was stirred for 3 hours. After 3 hours, organic layers were extracted from the reaction solution three times by using 100 ml of water and 10 ml of diethyl ether. The obtained organic layers were dried by using magnesium sulfate and the solvent was evaporated. The obtained residue was separated and purified by silica gel chromatography, thereby completing the preparation of 13.635 g (45 mmol, yield=90%) of Intermediate A-1.

Synthesis of Intermediate A-2

[0318] 13.635 g (45 mmol) of Intermediate A-1 was dissolved in 200 ml of anhydrous THF under a nitrogen atmosphere at a temperature of 0° C., and 8.46 g (60 mmol) of methyl iodide was added thereto for 1 hour in a dropwise fashion. The obtained reaction solution was heated to room temperature and was stirred for 3 hours. After 3 hours, organic layers were extracted from the reaction solution three times by using 100 ml of water and 10 ml of diethyl ether. The obtained organic layers were dried by using magnesium sulfate and the solvent was evaporated. The obtained residue was separated and purified by silica gel chromatography, thereby completing the preparation of 12.04 g (45 mmol, yield=88%) of Intermediate A-2.

Synthesis of Intermediate A

[0319] 12.04 g (40 mmol) of Intermediate A-2, 11.8 g (100 mmol) of Tin, and 20 ml of HCl were dissolved in 200 ml of ethanol at a temperature of 80° C. and were stirred for 8 hours. Then, 8.46 g (60 mmol) of methyl iodide was added thereto for 1 hour in a dropwise fashion. The obtained reaction solution was heated to room temperature and was stirred for 3 hours. The reaction solution was cooled to room temperature and 10 g of sodium hydroxide was added thereto. Then, organic layers were extracted three times by using 100 ml of water and 10 ml of diethyl ether. The obtained organic layers were dried by using magnesium sulfate and the solvent was evaporated. The obtained residue was separated and purified by silica gel chromatography, thereby completing the preparation of 9.485 g (35 mmol, yield=87.5%) of Intermediate A.



Synthesis of Intermediate B-1


[0320] 13.635 g (45 mmol, yield=90%) of Intermediate B-1 was synthesized in the same manner as in Synthesis of Intermediate A-1, except that 3-nitrophenyl magnesium bromide was used instead of 4-nitrophenyl magnesium bromide.


Synthesis of Intermediate B-2

[0321] 12.04 g (45 mmol, yield=88%) of Intermediate B-2 was synthesized in the same manner as in Synthesis of Intermediate A-2, except that Intermediate B-1 was used instead of Intermediate A-1.

Synthesis of Intermediate B

[0322] 9.485 g (35 mmol, yield=87.5%) of Intermediate B was synthesized in the same manner as in Synthesis of Intermediate A, except that Intermediate B-2 was used instead of Intermediate A-2.





Synthesis of Intermediate C-1


[0323] 13.635 g (45 mmol, yield=90%) of Intermediate C-1 was synthesized in the same manner as in Synthesis of Intermediate A-1, except that 3-nitrophenyl magnesium bromide was used instead of 4-nitrophenyl magnesium bromide.

Synthesis of Intermediate B-2

[0324] 12.04 g (45 mmol, yield=88%) of Intermediate C-2 was synthesized in the same manner as in Synthesis of Intermediate A-2, except that Intermediate C-1 was used instead of Intermediate A-1.

Synthesis of Intermediate C

[0325] 9.485 g (35 mmol, yield=87.5%) of Intermediate C was synthesized in the same manner as in Synthesis of Intermediate A, except that Intermediate B-2 was used instead of Intermediate A-2.






Synthesis of Intermediate 1-1

[0326] 2.71 g (10 mmol) of Intermediate A, 1.57 g (10 mmol) of bromobenzene, 0.091 g (0.01 mmol) of $\text{Pd}_2(\text{dba})_3$ (tris(dibenzylideneacetone)dipalladium), 0.020 g (0.01 mmol) of $t\text{-Bu}_3\text{P}$, and 1.92 g (20 mmol) of $t\text{-BuONa}$ were dissolved in 50 ml of toluene under a nitrogen atmosphere and were stirred at a temperature of 90° C. for 4 hours. After 4 hours, organic layers were extracted from the reaction solution three times by using 100 ml of water and 10 ml of diethyl ether. The obtained organic layers were dried by using magnesium sulfate and the solvent was evaporated. The obtained residue was separated and purified by silica gel chromatography, thereby completing the preparation of 3.123 g (9 mmol, yield=90%) of Intermediate 1-1.

Synthesis of Compound 1

[0327] 4.704 g (8 mmol, yield=88.8%) of Compound 1 was synthesized in the same manner as in Synthesis of Intermediate 1-1, except that Intermediate 1-1 was used instead of Intermediate A and 3-bromo-9-phenylcarbazole was used instead of bromobenzene.

[0328] ^1H NMR (CDCl_3 , 300 MHz): 8.10 (d, 1H), 8.00-7.85 (m, 6H), 7.80-7.50 (m, 8H), 7.40-7.20 (m, 7H), 7.10-6.90 (m, 7H), 2.20 (s, 3H)






Synthesis of Intermediate 16-1

[0329] 4.077 g (9 mmol, yield=90%) of Intermediate 16-1 was synthesized in the same manner as in Synthesis of Intermediate 1-1, except that Intermediate B was used instead of Intermediate A and 3-bromo-dibenzothiophene was used instead of bromobenzene.

Synthesis of Compound 16

[0330] 5.5524 g (8 mmol, yield=88.8%) of Compound 16 was synthesized in the same manner as in Synthesis of Compound 1, except that Intermediate 16-1 was used instead of Intermediate 1-1.

[0331] ^1H NMR (CDCl_3 , 300 MHz): 8.30-8.10 (m, 3H), 8.00-7.80 (m, 4H), 7.75-7.45 (m, 14H), 7.40-7.10 (m, 10), 2.20 (s, 3H)




Synthesis of Intermediate 18-1

[0332] 3.573 g (9 mmol, yield=90%) of Intermediate 18-1 was synthesized in the same manner as in Synthesis of Intermediate 1-1, except that Intermediate 2 was used instead of Intermediate A and 2-naphthalene was used instead of bromobenzene.

Synthesis of Compound 18

[0333] 5.108 g (8 mmol, yield=88.8%) of Compound 18 was synthesized in the same manner as in Synthesis of Compound 1, except that Intermediate 18-1 was used instead of Intermediate 1-1.

[0334] ^1H NMR (CDCl_3 , 300 MHz): 8.30 (d, 1H), 8.00-7.85 (m, 4H), 7.80-7.20 (m, 18H), 7.10-7.00 (m, 4H), 6.80-6.7000 (m, 4H), 2.20 (s, 3H)



Synthesis of Intermediate 28-1


[0335] 3.807 g (9 mmol, yield=90%) of Intermediate 28-1 was synthesized in the same manner as in Synthesis of Intermediate 1-1, except that 4-bromobiphenyl was used instead of bromobenzene.

Synthesis of Compound 28

[0336] 5.920 g (8 mmol, yield=88.8%) of Compound 28 was synthesized in the same manner as in Synthesis of

Compound 1, except that Intermediate 28-1 was used instead of Intermediate 1-1 and 3-(4-bromophenyl)-9-phenylcarbazole was used instead of 3-bromo-9-phenylcarbazole.

[0337] ^1H NMR (CDCl_3 , 300 MHz): 8.30 (dd, 2H), 8.10-7.50 (m, 23H), 7.35-7.20 (m, 6H), 7.10-6.90 (m, 6H), 2.20 (s, 3H)




Synthesis of Intermediate 37-1

[0338] 3.969 g (9 mmol, yield=90%) of Intermediate 37-1 was synthesized in the same manner as in Synthesis of

Intermediate 16-1, except that 4-bromo-4'-fluoro-biphenyl was used instead of 3-bromo-dibenzothiophene. Synthesis of Compound 37

[0339] 6.0640 g (8 mmol, yield=88.8%) of Compound 37 was synthesized in the same manner as in Synthesis of Compound 28, except that Intermediate 37-1 was used instead of Intermediate 28-1.

[0340] ^1H NMR (CDCl_3 , 300 MHz): 8.30 (dd, 2H), 8.05-7.55 (m, 20H), 7.45-7.20 (m, 8H), 7.10 (d, 1H), 2.20 (s, 3H)




Synthesis of Intermediate 47-1

[0341] 3.933 g (9 mmol, yield=90%) of Intermediate 47-1 was synthesized in the same manner as in Synthesis of Intermediate 18-1, except that 4-bromo-dibenzofuran was used instead of 2-naphthalene.

Synthesis of Compound 47

[0342] 6.032 g (8 mmol, yield=88.8%) of Compound 47 was synthesized in the same manner as in Synthesis of Compound 28, except that Intermediate 47-1 was used instead of Intermediate 28-1.


[0343] ^1H NMR (CDCl_3 , 300 MHz): 8.30 (dd, 2H), 8.05-7.50 (m, 24H), 7.40 (dd, 4H), 7.20 (dd, 3H), 7.10 (d, 2H), 2.20 (s, 3H)




Synthesis of Compound 52

[0344] 5.456 g (8 mmol, yield=88.8%) of Compound 52 was synthesized in the same manner as in Synthesis of Compound 28, except that 3-bromo-9-(4-fluorophenyl)carbazole was used instead of 3-(4-bromophenyl)-9-phenylcarbazole.


[0345] ^1H NMR (CDCl_3 , 300 MHz): 8.25 (d, 1H), 8.05-7.90 (m, 4H), 7.80-7.40 (m, 14H), 7.35-7.20 (m, 6H), 7.10-6.90 (m, 7H), 2.20 (s, 3H)




16-1



18-1



64



66

Synthesis of Compound 66


[0348] 6.112 g (8 mmol, yield=88.8%) of Compound 66 was synthesized in the same manner as in Synthesis of Compound 18, except that 3-(3-bromophenyl)-9-(1-naphthyl)carbazole was used instead of 3-bromo-9-phenylcarbazole.

[0349] ^1H NMR (CDCl_3 , 300 MHz): 8.35-8.20 (m, 3H), 8.10-7.80 (m, 13H), 7.75-7.10 (m, 17H), 7.05 (dd, 3H), 6.90 (d, 1H), 2.20 (s, 3H)



Synthesis of Compound 64

[0346] 6.880 g (8 mmol, yield=88.8%) of Compound 64 was synthesized in the same manner as in Synthesis of Compound 16, except that 3-(4-bromophenyl)-9-(3-dibenzofuranyl)carbazole was used instead of 3-bromo-9-phenylcarbazole.

[0347] ^1H NMR (CDCl_3 , 300 MHz): 8.35-8.15 (m, 5H), 8.00-7.70 (m, 8H), 7.60-7.30 (m, 12H), 7.20-7.00 (m, 8H), 6.90-6.70 (m, 4H), 2.20 (s, 3H)



B



Synthesis of Intermediate 78-1

[0350] 3.933 g (9 mmol, yield=90%) of Intermediate 78-1 was synthesized in the same manner as in Synthesis of Intermediate 16-1, except that 3-bromo-dibenzofuran was used instead of 3-bromo-dibenzothiophene.

Synthesis of Compound 78

[0351] 6.176 g (8 mmol, yield=88.8%) of Compound 78 was synthesized in the same manner as in Synthesis of Compound 1, except that Intermediate 78-1 was used instead of Intermediate 1-1 and 3-(4-bromophenyl)-6-fluoro-9-phenylcarbazole was used instead of 3-bromo-9-phenylcarbazole.

[0352] ^1H NMR (CDCl_3 , 300 MHz): 8.25 (s, 1H), 8.10-7.50 (m, 20H), 7.45-7.10 (m, 9H), 6.90 (dd, 3H), 6.80 (d, 1H), 2.20 (s, 3H)

EXAMPLE


Example 1

[0353] An anode was prepared by cutting an ITO glass substrate (manufactured by Corning), on which an ITO layer was deposited to a thickness of $15\ \Omega/\text{cm}^2$ (1,200 Å), to a size of 50 mm×50 mm×0.7 mm, ultrasonically cleaning the ITO glass substrate (anode) by using isopropyl alcohol and pure water each for 5 minutes, and then, exposing the ITO glass substrate (anode) to irradiation of UV light for 30 minutes and ozone to clean the ITO glass substrate. Then, the ITO glass substrate (anode) was loaded into a vacuum deposition apparatus.

[0354] 2-TNATA was vacuum-deposited on the ITO glass substrate (anode) to form a hole injection layer having a thickness of 600 Å, and Compound 1 as a hole transport compound was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å.


[0355] 9,10-di-naphthalene-2-yl-anthracene (ADN) (blue fluorescent host) and N,N,N',N'-tetraphenyl-pyrene-1,6-diamine (TPD) (blue fluorescent dopant) were co-deposited on the hole transport layer at a weight ratio of 98:2 to form an emission layer having a thickness of 300 Å.

[0356] Then, Alq_3 was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å, LiF (alkali metal halide) was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited on the electron injection layer to form an LiF/Al electrode (cathode electrode) having a thickness of 3,000 Å, thereby completing the manufacture of an organic light-emitting device.




2-TNATA

-continued



ADN



TPD

Example 2

[0357] An organic light-emitting device of Example 2 was manufactured in the same manner as in Example 1, except that Compound 16 was used instead of Compound 1 in forming a hole transport layer.

Example 3

[0358] An organic light-emitting device of Example 3 was manufactured in the same manner as in Example 1, except that Compound 18 was used instead of Compound 1 in forming a hole transport layer.

Example 4

[0359] An organic light-emitting device of Example 4 was manufactured in the same manner as in Example 1, except that Compound 28 was used instead of Compound 1 in forming a hole transport layer.

Example 5

[0360] An organic light-emitting device of Example 5 was manufactured in the same manner as in Example 1, except

that Compound 37 was used instead of Compound 1 in forming a hole transport layer.

Example 6

[0361] An organic light-emitting device of Example 6 was manufactured in the same manner as in Example 1, except that Compound 47 was used instead of Compound 1 in forming a hole transport layer.

Example 7

[0362] An organic light-emitting device of Example 7 was manufactured in the same manner as in Example 1, except that Compound 52 was used instead of Compound 1 in forming a hole transport layer.

Example 8

[0363] An organic light-emitting device of Example 8 was manufactured in the same manner as in Example 1, except that Compound 64 was used instead of Compound 1 in forming a hole transport layer.

Example 9


[0364] An organic light-emitting device of Example 9 was manufactured in the same manner as in Example 1, except that Compound 66 was used instead of Compound 1 in forming a hole transport layer.

Example 10

[0365] An organic light-emitting device of Example 10 was manufactured in the same manner as in Example 1, except that Compound 78 was used instead of Compound 1 in forming a hole transport layer.

Comparative Example 1

[0366] An organic light-emitting device of Comparative Example 1 was manufactured in the same manner as in Example 1, except that well-known Compound NPB was used in forming a hole transport layer.





NPB

Comparative Example 2

[0367] An organic light-emitting device of Comparative Example 2 was manufactured in the same manner as in Example 1, except that Compound 314 was used in forming a hole transport layer.

314

(102)



Comparative Example 3

[0368] An organic light-emitting device of Comparative Example 3 was manufactured in the same manner as in Example 1, except that Compound 102 was used in forming a hole transport layer.

[0369] Results of Comparative Examples and Examples are shown in Table 1 below.

[0370] Referring to Table 1, when the compound according to one or more embodiments was used in the hole transport layer, a driving voltage was improved and efficiency was increased. In particular, the luminance and half lifespan of the organic light-emitting device were increased.

TABLE 1

	Hole transport layer	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Haft Lifespan (hr @ 100 mA/cm ²)
Example 1	Compound 1	5.29	50	3435	6.87	Blue	382 hr
Example 2	Compound 16	5.36	50	3135	6.27	Blue	358 hr
Example 3	Compound 18	5.44	50	3160	6.32	Blue	345 hr
Example 4	Compound 28	5.32	50	3285	6.57	Blue	354 hr
Example 5	Compound 37	5.83	50	3095	6.19	Blue	319 hr
Example 6	Compound 47	5.74	50	3115	6.23	Blue	337 hr
Example 7	Compound 52	6.17	50	3025	6.05	Blue	323 hr
Example 8	Compound 64	5.27	50	3095	6.19	Blue	349 hr
Example 9	Compound 66	5.38	50	3305	6.61	Blue	347 hr
Example 10	Compound 78	5.46	50	3215	6.43	Blue	352 hr
Comparative Example 1	NPB	7.07	50	2545	5.09	Blue	268 hr
Comparative Example 2	Compound 314	5.94	50	2715	5.43	Blue	290 hr
Comparative Example 3	Compound 102	6.02	50	2605	5.21	Blue	260 hr


[0371] Due to the inclusion of the compound of Formula 1 according to an embodiment, characteristics of an organic light-emitting device may be improved.

[0372] It should be understood that embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as available for other similar features or aspects in other embodiments.

[0373] While one or more embodiments have been described with reference to the figures, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope as defined by the following claims.

What is claimed is:

1. A compound represented by Formula 1:



wherein, in Formula 1,

R_1 , Ar_1 , and Ar_2 are each independently selected from hydrogen, deuterium, a halogen, an amino group, a nitro group, a nitrile group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

L is each independently selected from a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

m and n are each independently an integer selected from 0 to 4,

when m and n are each independently two or more, two or more $R_1(s)$ and two or more $L(s)$ are each independently identical to or different from each other; and at least one substituent selected from a substituent(s) of the substituted C_1 - C_{60} alkyl group, the substituted

C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_2 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted C_6 - C_{60} arylene group, the substituted C_1 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and the substituted divalent non-aromatic condensed heteropolycyclic group is selected from one of a first group, a second group, a third group, and a fourth group:

the first group including deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

the second group including a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_{11})(Q_{12})$, $-Si(Q_{13})(Q_{14})(Q_{15})$, and $-B(Q_{16})(Q_{17})$;

the third group including a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

the fourth group including a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt

thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇),


wherein Q₁₁ to Q₁₇ and Q₂₁ to Q₂₇ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

2. The compound of claim 1, wherein


n in Formula 1 is 0 or 1.

3. The compound of claim 1, wherein


Ar₁ and Ar₂ in Formula 1 are each independently one selected from groups represented by Formulae 2a to 2d:




2a



2b



2c



2d

wherein H₁ in Formula 2c is O, S, NR₂₁, or CR₂₂R₂₃,

R₂₁ to R₂₃ and Z₁ are each independently selected from hydrogen, deuterium, a halogen, a substituted or unsubstituted C₁-C₂₀ alkyl group, a substituted or unsubstituted C₆-C₂₀ aryl group, a substituted or unsubstituted C₁-C₂₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,


p is an integer selected from 1 to 7,

when p is two or more, two or more Z₁(s) may be identical to or different from each other, and


* indicates a binding site.

4. The compound of claim 1, wherein

R₁ in Formula 1 is each independently one selected from hydrogen, deuterium, a cyano group, a halogen, or groups represented by Formulae 3a and 3b:



3a



3b

wherein Z₁(s) in Formulae 3a and 3b are each independently hydrogen, deuterium, a halogen, a substituted or unsubstituted C₁-C₂₀ alkyl group, a substituted or unsubstituted C₆-C₂₀ aryl group, a substituted or unsubstituted C₁-C₂₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,


p is an integer selected from 1 to 7,

when p is two or more, two or more Z₁(s) are identical to or different from each other, and

* indicates a binding site.

5. The compound of claim 1, wherein

L in Formula 1 is represented by Formula 4a:




4a

wherein * in Formula 4a indicates a binding site.

6. The compound of claim 1, wherein


the compound represented by Formula 1 is represented by Formula 2:

<Formula 2>




wherein n in Formula 2 is 0 or 1.


7. The compound of claim 1, wherein
the compound represented by Formula 1 is represented by
Formula 3:






8. The compound of claim 1, wherein
the compound represented by Formula 1 is represented by
Formula 4:



9. The compound of claim 1, wherein
the compound represented by Formula 1 is represented by
Formula 5:




10. The compound of claim 1, wherein
the compound represented by Formula 1 is any one of the
compounds below:




-continued

4




-continued


7




5




8



6




9




-continued

10




-continued


13




11




14




12



15




-continued




16


-continued




19




17



20




18




21

-continued




22


-continued




25




23



26




24



27


-continued

28




-continued

30





31

29




32




-continued

33





-continued

36





37

34




38

35




-continued


39



40




41




-continued


42



43




44




-continued

45




-continued

48





49

46




50

47




-continued




51


-continued




54




52



55




53




56

-continued


57



58




59




-continued

60




61




-continued

62




-continued


65




63



64




66




-continued

67




-continued


69



68




70




-continued

71




-continued


74




72




75



73




76




-continued

77



-continued

80



11. An organic light-emitting device comprising:


a first electrode;

a second electrode facing the first electrode; and


an organic layer that is disposed between the first electrode and the second electrode and comprises an emission layer,

wherein the organic layer comprises a compound represented by Formula 1:

78




<Formula 1>



wherein, in Formula 1,

R_1 , Ar_1 , and Ar_2 are each independently selected from hydrogen, deuterium, a halogen, an amino group, a nitro group, a nitrile group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

79



L is each independently selected from a substituted or unsubstituted C_6-C_{60} arylene group, a substituted or unsubstituted C_1-C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

m and n are each independently an integer selected from 0 to 4,

when m and n are each independently two or more, two or more $R_1(s)$ and two or more L(s) are each independently identical to or different from each other; and

at least one substituent selected from a substituent(s) of the substituted C_1-C_{60} alkyl group, the substituted C_2-C_{60} alkenyl group, the substituted C_2-C_{60} alkynyl group, the substituted C_1-C_{60} alkoxy group, the substituted C_3-C_{10} cycloalkyl group, the substituted C_1-C_{10} heterocycloalkyl group, the substituted C_3-C_{10} cycloalkenyl group, the substituted C_2-C_{10} heterocycloalkenyl group, the substituted C_6-C_{60} aryl group, the substituted C_6-C_{60} aryloxy group, the substituted C_6-C_{60} arylthio group, the substituted C_1-C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted C_6-C_{60} arylene group, the substituted C_1-C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and the substituted divalent non-aromatic condensed heteropolycyclic group is selected from one of a first group, a second group, a third group, and a fourth group:

the first group including deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, and a C_1-C_{60} alkoxy group;

the second group including a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, and a C_1-C_{60} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_2-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q_{11})(Q_{12}), —Si(Q_{13})(Q_{14})(Q_{15}), and —B(Q_{16})(Q_{17});

the third group including a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_2-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-

aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

the fourth group including a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_2-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_1-C_{60} alkoxy group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_2-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q_{21})(Q_{22}), —Si(Q_{23})(Q_{24})(Q_{25}), and —B(Q_{26})(Q_{27}),

wherein Q_{11} to Q_{17} and Q_{21} to Q_{27} are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_1-C_{60} alkoxy group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_2-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

12. The organic light-emitting device of claim 11, wherein the emission layer comprises a blue fluorescent emission layer.

13. The organic light-emitting device of claim 11, wherein the first electrode is an anode, the second electrode is a cathode, and the organic layer comprises:


- i) a hole transport region that is disposed between the first electrode and the emission layer and comprises a hole transport layer; and at least one of a hole injection layer, and an electron blocking layer; and
- ii) an electron transport region that is disposed between the emission layer and the second electrode and comprises an electron transport layer, a hole blocking layer, and an electron injection layer.

14. The organic light-emitting device of claim 13, wherein the hole transport layer comprises the compound of Formula 1.

15. The organic light-emitting device of claim 13, wherein the hole transport region comprises a charge-generation material.

16. The organic light-emitting device of claim 15, wherein the charge-generation material comprises a p-dopant.


17. The organic light-emitting device of claim 15, wherein the charge-generation material is Compound HT-D1 or Compound F4-TCNQ:



18. The organic light-emitting device of claim 13, wherein the electron transport region comprises a metal-containing material.

19. The organic light-emitting device of claim 13, wherein the electron transport region comprises a Li complex.

20. A display apparatus comprising:
an organic light-emitting device that includes:
a first electrode;
a second electrode facing the first electrode; and
an organic layer that is disposed between the first electrode and the second electrode and comprises an emission layer,
wherein the organic layer comprises a compound represented by Formula 1:



wherein, in Formula 1,

R_1 , Ar_1 , and Ar_2 are each independently selected from hydrogen, deuterium, a halogen, an amino group, a nitro group, a nitrile group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted

C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

L is each independently selected from a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

m and n are each independently an integer selected from 0 to 4,

when m and n are each independently two or more, two or more R_1 (s) and two or more L (s) are each independently identical to or different from each other; and

at least one substituent selected from a substituent(s) of the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_2 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted C_6 - C_{60} arylene group, the substituted C_1 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, and the substituted divalent non-aromatic condensed heteropolycyclic group is selected from one of a first group, a second group, a third group, and a fourth group:

the first group including deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

the second group including a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60}

aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

the third group including a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

the fourth group including a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group,

a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇),

wherein Q₁₁ to Q₁₇ and Q₂₁ to Q₂₇ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

wherein the first electrode of the organic light-emitting device is electrically connected to a source electrode or a drain electrode of a thin film transistor.

* * * * *

专利名称(译)	包括其的化合物和有机发光装置		
公开(公告)号	US20170186952A1	公开(公告)日	2017-06-29
申请号	US15/393086	申请日	2016-12-28
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	HAN SANGHYUN KIM SOOYON KIM JONGWOO KIM YOUNGKOOK HWANG SEOKHWAN		
发明人	HAN, SANGHYUN KIM, SOOYON KIM, JONGWOO KIM, YOUNGKOOK HWANG, SEOKHWAN		
IPC分类号	H01L51/00 C07D409/14 C07D209/86 C07D405/12 C07D209/88 C07D409/12		
CPC分类号	H01L51/006 H01L51/0073 H01L51/005 H01L51/0077 C07D209/88 H01L51/0061 C07D409/12 C07D209/86 C07D405/12 C07D409/14 H01L51/5012 H01L51/5056 H01L51/5072 H01L51/0052 H01L51/0074 H01L51/0058 H01L51/0072		
优先权	1020150187638 2015-12-28 KR		
其他公开文献	US10622564		
外部链接	Espacenet USPTO		

摘要(译)

提供由式1表示的化合物和包含其的有机发光装置：

40

220
190
150
110
210