



US 20180240982A1

(19) **United States**(12) **Patent Application Publication**
KIM et al.(10) **Pub. No.: US 2018/0240982 A1**(43) **Pub. Date: Aug. 23, 2018**(54) **POLYCYCLIC COMPOUND AND ORGANIC
LIGHT-EMITTING DEVICE INCLUDING
THE SAME**(71) Applicant: **SAMSUNG DISPLAY CO., LTD.**,
Yongin-Si (KR)(72) Inventors: **SEHUN KIM**, Yongin-Si (KR);
Sooyoung Park, Seoul (KR); **Chihyun
Ryoo**, Seoul (KR); **Ilhun Cho**,
Yongin-Si (KR); **Junghoon Yang**, Seoul
(KR); **Seyoung Jung**, Seoul (KR);
Hyein Jeong, Seoul (KR)(73) Assignee: **SEOUL NATIONAL UNIVERSITY
R&DB FOUNDATION**, Seoul (KR)(21) Appl. No.: **15/700,344**(22) Filed: **Sep. 11, 2017**(30) **Foreign Application Priority Data**

Feb. 23, 2017 (KR) 10-2017-0024250

Publication Classification(51) **Int. Cl.**
H01L 51/00 (2006.01)
C07D 265/38 (2006.01)
C07D 213/16 (2006.01)
(52) **U.S. Cl.**
CPC **H01L 51/0071** (2013.01); **H01L 51/0059**
(2013.01); **C07D 213/16** (2013.01); **C07D**
265/38 (2013.01)(57) **ABSTRACT**

A polycyclic compound and an organic light-emitting device including the same are provided. The polycyclic compound is represented by Formula 1:

<Formula 1>



10

FIG. 1

10

190
150
110

FIG. 2





FIG. 3



**POLYCYCLIC COMPOUND AND ORGANIC
LIGHT-EMITTING DEVICE INCLUDING
THE SAME**

**CROSS-REFERENCE TO RELATED
APPLICATION**

[0001] This application claims priority under 35 U.S.C. § 119 to Korean Patent Application No. 10-2017-0024250, filed on Feb. 23, 2017, the Korean Intellectual Property Office, the disclosure of which is incorporated by reference herein in its entirety.

1. Technical Field

[0002] Exemplary embodiments of the present invention relate to a polycyclic compound, and more particularly to an organic light-emitting device including the same.

2. Discussion of Related Art



[0003] Organic light-emitting devices may be self-emission devices that have relatively wide viewing angles, relatively high contrast ratios, relatively short response times, relatively high brightness and relatively fast response speeds.

[0004] An example of such organic light-emitting devices may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, may recombine in the emission layer to produce excitons. These excitons may fall from an excited state to a ground state, thus generating light.

SUMMARY


[0005] One or more exemplary embodiments of the present invention include a polycyclic compound and an organic light-emitting device including the same.

[0006] According to one or more exemplary embodiments of the present invention, a polycyclic compound is represented by Formula 1:



-continued

<Formula 3>



[0007] In Formulae 1 to 4,

[0008] X_1 may be O or S,

[0009] Y_{11} , Y_{21} , and Y_{22} may each independently be selected from a single bond, $C(R_{31})(R_{32})$, $Si(R_{31})(R_{32})$, O, S, and $N(R_{31})$, and Y_{21} and Y_{22} are not a single bond at the same time,

[0010] A_{11} , A_{12} , A_{21} , and A_{22} may be each independently a C_3 - C_{60} group or a C_1 - C_{60} heterocyclic group,

[0011] L_1 , L_{11} , and L_{21} may each independently be a substituted or unsubstituted C_3 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0012] a_1 , a_{11} , and a_{21} may each independently be an integer from 0 to 3, and, when a_1 is two or more, two or more $L_1(s)$ may be identical to or different from each other, when a_{11} is two or more, two or more $L_{11}(s)$ may be identical to or different from each other, and when a_{21} is two or more, two or more $L_{21}(s)$ may be identical to or different from each other,

[0013] Ar_1 may be selected from $-F$, $-Cl$, $-Br$, $-I$, a cyano group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, $-N(Q_{101})(Q_{102})$, and $-S(Q_{101})$,

[0014] R_1 to R_{12} , R_{21} , R_{22} , R_{31} , and R_{32} may each independently be selected from a group represented by one of Formulae 2 to 4, hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group, $-Si(Q_1)(Q_2)(Q_3)$, $-N(Q_1)(Q_2)$, $-B(Q_1)(Q_2)$, $-S(=O)_2(Q_1)$, and $-P(=O)(Q_1)(Q_2)$,

[0015] provided that each of R_{11} to R_{12} , R_{21} , R_{22} , R_{31} , and R_{32} is not a group represented by one of Formulae 2 to 4,

[0016] at least one of R_1 to R_5 may be a group represented by Formula 2 or Formula 3,

[0017] at least one of R_6 to R_{10} may be a group represented by Formula 4,

[0018] b11, b12, b21, and b22 may each independently be an integer selected from 1 to 15, and, when b11 is two or more, two or more $R_{11}(s)$ may be identical to or different from each other, when b12 is two or more, two or more $R_{12}(s)$ may be identical to or different from each other, when b21 is two or more, two or more $R_{21}(s)$ may be identical to or different from each other, and when b22 is two or more, two or more $R_{22}(s)$ may be identical to or different from each other,

[0019] at least one substituent of the substituted C_3 - C_{60} carbocyclic group, the substituted C_1 - C_{60} heterocyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_1 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0020] deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0021] a C_1 - C_{60} group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{11})(Q_{12})(Q_{13})$, $-N(Q_{11})(Q_{12})$, $-B(Q_{11})(Q_{12})$, $-C(=O)(Q_{11})$, $-S(=O)_2(Q_{11})$, and $-P(=O)(Q_{11})(Q_{12})$;

[0022] C_3 - C_{10} cycloalkyl group, a C_1 - C_{20} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group;

[0023] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl

group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, $-Si(Q_{21})(Q_{22})(Q_{23})$, $-N(Q_{21})(Q_{22})$, $-B(Q_{21})(Q_{22})$, $-C(=O)(Q_{21})$, $-S(=O)_2(Q_{21})$, and $-P(=O)(Q_{21})(Q_{22})$; and

[0024] $-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})(Q_{32})$,

[0025] Q_{101} , Q_{102} , Q_1 to Q_3 , Q_{11} to Q_{13} , Q_{21} to Q_{23} , and Q_{31} to Q_{33} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

[0026] * indicates a binding site to a neighboring atom.

[0027] According to one or more exemplary embodiments of the present invention, an organic light-emitting device includes a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer and at least one of the polycyclic compound represented by Formula 1.

BRIEF DESCRIPTION OF THE DRAWINGS

[0028] The above and other features of the present invention will become more apparent by describing in detail exemplary embodiments thereof with reference to the accompanying drawings, in which:

[0029] FIG. 1 is a schematic view of an organic light-emitting device according to an exemplary embodiment of the present invention;

[0030] FIG. 2 is a graph showing light emission characteristics of organic light-emitting devices including TADF emitters, according to Example 1 and Comparative Examples 1 and 2; and

[0031] FIG. 3 is a graph showing external quantum efficiency (EQE) and emission color with respect to a driving current density in organic light-emitting devices including TADF emitters, according to Example 1 and Comparative Example 1 and 2.




DETAILED DESCRIPTION

[0032] Exemplary embodiments of the present invention will be described below in more detail with reference to the accompanying drawings. In this regard, the exemplary embodiments may have different forms and should not be construed as being limited to the exemplary embodiments of the present invention described herein.

[0033] It is to be understood that values described as substantially equal to each other may be equal to each other to within a measurement error, or if measurably unequal, are

close enough in value to be functionally equal to each other as would be understood by a person having ordinary skill in the art.

[0034] A polycyclic compound according to an exemplary embodiment of the present invention may be represented by Formula 1:



[0035] In Formula 1,

[0036] X_1 may be O or S.

[0037] For example, X_1 may be O.

[0038] In Formulae 2 and 3, Y_{11} , Y_{21} , and Y_{22} may each independently be selected from a single bond, $C(R_{31})(R_{32})$, $Si(R_{31})(R_{32})$, O, S, and $N(R_{31})$, and a case where both Y_{21} and Y_{22} are a single bond at the same time is excluded.

[0039] In an exemplary embodiment of the present invention, Y_{11} may be selected from a single bond, $C(R_{31})(R_{32})$, O, and S, and at least one of Y_{21} or Y_{22} may be $N(R_{31})$.

[0040] A_{11} , A_{12} , A_{21} , and A_{22} in Formulae 2 and 3 may each independently be a C_3 - C_{60} carbocyclic group or a C_1 - C_{60} heterocyclic group.

[0041] In an exemplary embodiment of the present invention, A_{11} , A_{12} , A_{21} , and A_{22} may be selected from:

[0042] a phenyl group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a spiro-fluorene-benzofluorene group, a benzofluorene group, a dibenzofluorene group, a phenanthrene group, an anthracene group, a triphenylene group, a pyrenylene group, a chrysenylene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, an indole group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an iso-benzothiazole group, a benzoxazole group, an isoben-

zoxazole group, an oxadiazole group, a triazine group, a dibenzofuran group, a benzonaphthofuran group, a dibenzothiophene group, a dibenzosilole group, a benzocarbazole group, an indolocarbazole group, a dibenzocarbazole group, an imidazopyridine group, and an imidazopyrimidine group; and

[0043] a phenyl group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a spiro-fluorene-benzofluorene group, a benzofluorene group, a dibenzofluorene group, a phenanthrene group, an anthracene group, a triphenylene group, a pyrenylene group, a chrysenylene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, an indole group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an iso-benzothiazole group, a benzoxazole group, an isoben-zoxazole group, an oxadiazole group, a triazine group, a dibenzofuran group, a benzonaphthofuran group, a dibenzothiophene group, a dibenzosilole group, a benzocarbazole group, an indolocarbazole group, a dibenzocarbazole group, an imidazopyridine group, and an imidazopyrimidine group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, an indolyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a benzonaphthofuranyl group, a benzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, an indolocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}).

[0044] For example, A_{11} , A_{12} , A_{21} , and A_{22} may be selected from: a phenyl group and a carbazole group; and a phenyl group substituted with at least one carbazolyl group.

[0045] L_1 , L_{11} , and L_{21} in Formulae 2 to 4 may each independently be a substituted or unsubstituted C_3 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group.

[0046] In an exemplary embodiment of the present invention, L_1 , L_{11} , and L_{21} may each independently be selected from:

[0047] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a spiro-fluorene-benzofluorenylene group, a benzo-

fluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picanylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylene group, a thiophenylenylene group, a furanylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylenylene group, an indolylenylene group, an indazolylenylene group, a purinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a carbazolylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, a benzofuranylenylene group, a benzothioophenylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylenylene group, a dibenzofuranylenylene group, a dibenzothioophenylenylene group, a dibenzosilolylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, and an imidazopyrimidinylenylene group; and

[0048] a phenylene group, a pentatenylenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylene group, a spiro-bifluorenylene group, a spiro-fluorene-benzofluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylene group, a thiophenylenylene group, a furanylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylenylene group, an indolylenylene group, an indazolylenylene group, a purinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a carbazolylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, a benzofuranylenylene group, a benzothioophenylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylenylene group, a dibenzofuranylenylene group, a dibenzothioophenylenylene group, a dibenzosilolylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and

[0049] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a biphenyl group, a terphenyl group, a naphthyl group, and a dibenzofuranyl group.

[0050] a1, a11, and a21 in Formulae 2 to 4 may each independently be an integer selected from 0 to 3, a1, a11, and a21 respectively indicate the number of L₁(s), the number of L₁₁(s), and the number of L₂₁(s), and, when a1 is two or more, two or more L₁(s) may be identical to or different from each other, when a11 is two or more, two or more L₁₁(s) may be identical to or different from each other, and when a21 is two or more, two or more L₂₁(s) may be identical to or different from each other. For example, a1 may be zero and a21 may be zero, but exemplary embodiments of the present invention are not limited thereto. For example, a1 may be zero, but exemplary embodiments of the present invention are not limited thereto.






[0051] In an exemplary embodiment of the present invention, the group represented by Formula 2 may be selected from groups represented by Formulae 2-1 to 2-7:

benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, and an imidazopyrimidinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothioophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothioophenyl group, a dibenzosilolylene group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and



[0049] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a biphenyl group, a terphenyl group, a naphthyl group, and a dibenzofuranyl group.

[0050] a1, a11, and a21 in Formulae 2 to 4 may each independently be an integer selected from 0 to 3, a1, a11, and a21 respectively indicate the number of L₁(s), the number of L₁₁(s), and the number of L₂₁(s), and, when a1 is two or more, two or more L₁(s) may be identical to or different from each other, when a11 is two or more, two or more L₁₁(s) may be identical to or different from each other, and when a21 is two or more, two or more L₂₁(s) may be identical to or different from each other. For example, a1 may be zero and a21 may be zero, but exemplary embodiments of the present invention are not limited thereto. For example, a1 may be zero, but exemplary embodiments of the present invention are not limited thereto.

[0051] In an exemplary embodiment of the present invention, the group represented by Formula 2 may be selected from groups represented by Formulae 2-1 to 2-7:






-continued



[0052] In Formulae 2-1 to 2-7:


[0053] Y_{11} , L_{11} , a_{11} , R_{11} , R_{12} , b_{11} , and b_{12} may each independently be the same as described above, and[0054] R_{13} may be the same as R_{11} and R_{12} , described above.

[0055] In an exemplary embodiment of the present invention, the group represented by Formula 3 may be selected from groups represented by Formulae 3-1 to 3-13:




-continued

3-4





-continued

3-10





3-11

3-5





3-12

3-6




3-13


3-7



3-8



3-9



[0056] In Formulae 3-1 to 3-13.

[0057] Y_{21} , Y_{22} , L_{21} , a_{21} , R_{21} , R_{22} , b_{21} , and b_{22} may each independently be the same as described above, and[0058] R_{23} may be the same as R_{21} and R_{22} , described above.

[0059] b_{11} , b_{12} , b_{21} , and b_{22} may each independently be an integer selected from 1 to 15. b_{11} , b_{12} , b_{21} , and b_{22} respectively indicate the number of $R_{11}(s)$, the number of $R_{12}(s)$, the number of $R_{21}(s)$, and the number of $R_{22}(s)$, and, when b_{11} is two or more, two or more $R_{11}(s)$ may be identical to or different from each other, when b_{12} is two or more, two or more $R_{12}(s)$ may be identical to or different from each other, when b_{21} is two or more, two or more $R_{21}(s)$ may be identical to or different from each other, and when b_{22} is two or more, two or more $R_{22}(s)$ may be identical to or different from each other. For example, b_{11} , b_{12} , b_{21} , and b_{22} may each independently be an integer selected from 1 to 6, but exemplary embodiments of the present invention are not limited thereto.

[0060] Ar₁ in Formula 4 may be selected from —F, —Cl, —Br, —I, a cyano group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, —N(Q₁₀₁)(Q₁₀₂), and —S(Q₁₀₁).

[0061] In an exemplary embodiment of the present invention, Ar₁ in Formula 4 may be selected from:

[0062] —F, —Cl, —Br, —I, a cyano group, a C₁-C₂₀ alkyl group, a C₂-C₂₀ alkenyl group, a C₂-C₂₀ alkynyl group, a C₁-C₂₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₆-C₂₀ aryl group, —N(Q₁₀₁)(Q₁₀₂), and —S(Q₁₀₁); and

[0063] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from —F, —Cl, —Br, —I, and a C₁-C₂₀ alkyl group.

[0064] For example, Ar₁ in Formula 4 may be selected from —F, a cyano group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₁-C₁₀ heterocycloalkyl group, N(Q₁₀₁)(Q₁₀₂), a C₁-C₂₀ alkyl group substituted with at least one —F, and a C₁-C₂₀ alkoxy group substituted with at least one C₁-C₂₀ alkyl group, but exemplary embodiments of the present invention are not limited thereto.

[0065] R₁ to R₁₂, R₂₁, R₂₂, R₃₁, and R₃₂ may each independently be selected from a group represented by one of Formulae 2 to 4, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂),

[0066] provided that each of R₁₁ to R₁₂, R₂₁, R₂₂, R₃₁, and R₃₂ is not a group represented by one of Formulae 2 to 4.

[0067] at least one of R₁ to R₅ to may be a group represented by Formula 2 or Formula 3, and

[0068] at least one of R₆ to R₁₀ may be a group represented by Formula 4.

[0069] In exemplary embodiment of the present invention, R₁ to R₁₀ in Formula 1 may each independently be selected from:

[0070] a group represented by one of Formulae 2 to 4, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C₁-C₂₀ alkyl group, or a C₁-C₂₀ alkoxy group;

[0071] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a ben-

zofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a benzonaphthofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0072] a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a benzonaphthofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazinyl group, a dibenzofuranyl group, a benzonaphthofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂); and

[0073] —Si(Q₁)(Q₂)(Q₃), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂), and

[0074] Q₁ to Q₃ and Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group,

a phenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a biphenyl group, a terphenyl group, a naphthyl group, and a carbazolyl group.

[0075] In exemplary embodiment of the present invention, R₃ may be a group represented by Formula 2 or Formula 3.









[0076] In an exemplary embodiment of the present invention, RR₁, R₂, R₄ to R₇, R₉, and R₁₀ may each be hydrogen.

[0077] For example, may be a group represented by Formula 2 or Formula 3, and R₈ may be a group represented by Formula 4, but exemplary embodiments of the present invention are not limited thereto. For example, R₃ may be a group represented by Formula 2, and R₈ may be a group represented by Formula 4, but exemplary embodiments of the present invention are not limited thereto.


[0078] A difference between a singlet energy level and a triplet energy level of the polycyclic compound may be 0.3 eV or less.

[0079] The polycyclic compound represented by Formula 1 may be selected from Compounds 1 to 9, but exemplary embodiments of the present invention are not limited thereto:

-continued



-continued



[0080] The polycyclic compound may have both a substituent exhibiting characteristics of an electron withdrawing group (EWG) and a substituent exhibiting characteristics of an electron donating group (EDG) within the compound, and a difference between singlet energy and triplet energy of the entire compound may be adjusted by introducing these substituents at appropriate positions. In this manner, thermally activated delayed fluorescence (TADF) may be exhibited.

[0081] As described above, the singlet energy and the triplet energy of the polycyclic compound satisfy the following equation:

$$\Delta E_{ST} = S_1 - T_1 \leq 0.3 \text{ eV.}$$

[0082] An existing TADF emitter may be manufactured through a combination of substituents having various characteristics of the EWG and the EDG. Even though the TADF emitter can exhibit high-efficiency TADF characteristics, there may be limitations in adjusting electronic structures of molecules. For example, since an emission wavelength region may be relatively wide, color purity may be relatively low.

[0083] The polycyclic compound may include a structure of Formulae 1 to 4. As an example, the polycyclic compound may have a structure in which left and right sides thereof are asymmetrical with respect to a carbonyl group. Due to a benzophenone structure, TADF characteristics similar to an existing TADF emitter may be maintained. Due to the lateral asymmetrical structure, color purity may be increased and the emission wavelength region may be adjusted.

[0084] The polycyclic compound may include, in addition to the lateral symmetrical structure, the structure of Formula 4 as a substituent. Thus, it is possible to finely adjust the emission region and adjust CIE color coordinates.

[0085] Since the structure of Formula 1 includes a substituent having characteristics of an EWG or an EDG at a parallel position with respect to the carbonyl group, the introduction of the substituent may exhibit an effect of changing only charge movement characteristics within the compound while not changing the basic light emission of the TADF emitter.

[0086] Synthesis methods of the polycyclic compound represented by Formula 1 may be recognizable by those of ordinary skill in the art by referring to Examples provided below.

[0087] At least one polycyclic compound represented by Formula 1 may be used between a pair of electrodes of an organic light-emitting device. For example, the polycyclic compound may be included in at least one of a hole transport region, an electron transport region, and an emission layer.

[0088] Thus, an organic light-emitting device may include: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode. The organic layer may include an emission layer and at least one of the polycyclic compound (e.g., the polycyclic compound represented by Formula 1).

[0089] The expression “(an organic layer) includes at least one of the polycyclic compound” used herein may include a case in which (an organic layer) includes one or more of the polycyclic compounds represented by Formula 1 or a case in which (an organic layer) includes two or more different polycyclic compounds.

[0090] In an exemplary embodiment of the present invention, the first electrode may be an anode and the second electrode may be a cathode. The organic layer may include a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode. The hole transport region may include a hole injection layer, a hole transport layer, an emission auxiliary layer, an electron blocking layer, or any combination thereof. The electron transport region may include a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, an electron injection layer, or any combination thereof.

[0091] In an exemplary embodiment of the present invention, the emission layer may include the polycyclic compound (e.g., the polycyclic compound represented by Formula 1).

[0092] For example, the polycyclic compound included in the emission layer may be a TADF emitter, and

[0093] the emission layer may emit delayed fluorescence.

[0094] As an example, the emission layer may include the polycyclic compound (e.g., the polycyclic compound represented by Formula 1).

[0095] As an example, the emission layer may include a host, and an amount of the polycyclic compound (e.g., the polycyclic compound represented by Formula 1) may be in a range of 0.1 parts by weight to about 50 parts by weight based on 100 parts by weight of the emission layer.

[0096] In an exemplary embodiment of the present invention, the hole transport region may include a p-dopant, of which a LUMO energy level is -3.5 eV or less.

[0097] The term “organic layer” used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of the organic light-emitting device. However, a material included in the “organic layer” is not limited to an organic material.

[0098] FIG. 1 is a schematic view of an organic light-emitting device according to an exemplary embodiment of the present invention. FIG. 1 is a schematic view of an organic light-emitting device 10 according to an exemplary embodiment of the present invention. The organic light-emitting device 10 may include a first electrode 110, an organic layer 150, and a second electrode 190.

[0099] A structure of the organic light-emitting device 10 according to an exemplary embodiment of the present invention and a method of manufacturing the organic light-emitting device 10 will be described in more detail below with reference to FIG. 1.

[0100] Referring to FIG. 1, a substrate may be disposed under the first electrode 110 or above the second electrode 190. The substrate may be a glass substrate or a plastic substrate, each having excellent mechanical strength, ther-

mal stability, transparency, surface smoothness, ease of handling, and water resistance.

[0101] The first electrode **110** may be formed by depositing or sputtering a material for forming the first electrode **110** on the substrate. When the first electrode **110** is an anode, the material for a first electrode may be selected from materials with a relatively high work function to facilitate hole injection.

[0102] The first electrode **110** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode **110** is a transmissive electrode, a material for forming the first electrode **110** may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO_2), zinc oxide (ZnO), and any combinations thereof, but exemplary embodiments of the present invention are not limited thereto. In an exemplary embodiment of the present invention, when the first electrode **110** is a semi-transmissive electrode or a reflectable electrode, a material for forming the first electrode **110** may be selected from magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and any combinations thereof, but exemplary embodiments of the present invention are not limited thereto.

[0103] The first electrode **110** may have a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode **110** may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode **110** is not limited thereto.

[0104] The organic layer **150** is disposed on the first electrode **110**. The organic layer **150** may include an emission layer.

[0105] The organic layer **150** may further include a hole transport region between the first electrode **110** and the emission layer, and an electron transport region between the emission layer and the second electrode **190**.



[0106] The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0107] The hole transport region may include at least one layer selected from a hole injection layer (HIL), a hole transport layer (HTL), an emission auxiliary layer, and an electron blocking layer (EBL).


[0108] As an example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure. For each structure of the hole transport region, the included layers may be sequen-

tially stacked from the first electrode **110** in this stated order, but the structure of the hole transport region is not limited thereto.








[0109] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPD), β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTDP: 4,4',4"-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:




-continued



-continued




<Formula 201>



-continued

<Formula 202>



[0110] In Formulae 201 and 202:

[0111] L_{201} to L_{204} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group; a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0112] L_{205} may be selected from $*-O-*$, $*-S-*$, $*-N(Q_{201})-*$, a substituted or unsubstituted C_1 - C_{20} alkylene group, a substituted or unsubstituted C_2 - C_{20} alkenylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0113] $xa1$ to $xa4$ may each independently be an integer selected from 0 to 3,

[0114] $xa5$ may be an integer selected from 1 to 10,

[0115] R_{201} to R_{204} and Q_{201} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0116] For example, in Formula 202, R_{201} and R_{202} may be chemically linked to each other via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group, and R_{203} and R_{204} may also be chemically linked to each other via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group.

[0117] In an exemplary embodiment of the present invention, in Formulae 201 and 202:

[0118] L_{201} to L_{205} may each independently be selected from:

[0119] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene

group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group;

[0120] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group; a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-F$, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranlyl group, a benzothiophenyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, $-Si(Q_{31})(Q_{32})(Q_{33})$, and $-N(Q_{31})(Q_{32})$; and

[0121] Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0122] In an exemplary embodiment of the present invention, $xa1$ to $xa4$ may each independently be 0, 1, or 2.

[0123] In an exemplary embodiment of the present invention, $xa5$ may be 1, 2, 3, or 4.

[0124] In an exemplary embodiment of the present invention, R_{201} to R_{204} and Q_{201} may each independently be selected from a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group;

[0125] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q_{31})(Q_{32})(Q_{33}), and —N(Q_{31})(Q_{32}); and

[0126] Q_{31} to Q_{33} may be the same as described above.

[0127] In an exemplary embodiment of the present invention, at least one selected from R_{201} to R_{203} in Formula 201 may each independently be selected from:

[0128] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0129] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, but exemplary embodiments of the present invention are not limited thereto.

[0130] In an exemplary embodiment of the present invention, in Formula 202, i) R_{201} and R_{202} may be linked via a single bond, and/or ii) R_{203} and R_{204} may be linked via a single bond.


[0131] In an exemplary embodiment of the present invention, at least one selected from R_{201} to R_{204} in Formula 202 may be selected from:

[0132] a carbazolyl group; and

[0133] a carbazolyl group, substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, but exemplary embodiments of the present invention are not limited thereto.


[0134] The compound represented by Formula 201 may be represented by Formula 201A.

<Formula 201A>




[0135] For example, the compound represented by Formula 201 may be represented by Formula 201A(1), but exemplary embodiments of the present invention are not limited thereto:

<Formula 201A(1)>




[0136] In an exemplary embodiment of the present invention, the compound represented by Formula 201 may be represented by Formula 201A-1, but exemplary embodiments of the present invention are not limited thereto:

<Formula 201A-1>




[0137] In an exemplary embodiment of the present invention, the compound represented by Formula 202 may be represented by Formula 202A:

<Formula 202A>



[0138] In an exemplary embodiment of the present invention, the compound represented by Formula 202 may be represented by Formula 202A-1:

<Formula 202A-1>



[0139] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1:


[0140] L₂₀₁ to L₂₀₃, xa1 to xa3, xa5, and R₂₀₂ to R₂₀₄ may be the same as described above,

[0141] R₂₁₁ and R₂₁₂ may be the same as described in connection with R₂₀₃, and


[0142] R₂₁₃ to R₂₁₇ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0143] The hole transport region may include at least one compound selected from Compounds HT1 to HT39, but exemplary embodiments of the present invention are not limited thereto:


HT1




HT2



HT3




HT4




-continued


HT5




HT6



HT7




HT8




-continued


HT9




HT10



HT11





HT12



-continued



HT13

HT14





HT15

HT16




HT17

HT18




-continued


HT19




HT20



HT21




HT22





-continued

HT23





HT24

HT25




HT26


HT27




-continued
HT28




HT30




HT32




HT34




HT29




HT31



HT33





HT35





-continued
HT36

HT37



HT38

HT39



[0144] A thickness of the hole transport region may be in a range of from about 100 Å to about 10,000 Å, for example, from about 100 Å to about 1,000 Å. When the hole transport region includes at least one selected from a hole injection layer or a hole transport layer the thickness of the hole injection layer may be in a range of from about 100 Å to about 9,000 Å, for example, from about 100 Å to about 1,000 Å, and the thickness of the hole transport layer may be in a range of from about 50 Å to about 2,000 Å, for example, from about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0145] The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer, and the electron blocking layer may block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include the materials as described above.

[0146] The hole transport region may further include, in addition to these materials, a charge-generation material for increasing conductive properties. The charge-generation material may be substantially homogeneously or non-homogeneously dispersed in the hole transport region.

[0147] The charge-generation material may be, for example, a p-dopant.

[0148] In an exemplary embodiment of the present invention, a lowest unoccupied molecular orbital (LUMO) energy level of the p-dopant may be -3.5 eV or less.

[0149] The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a compound including a cyano group, but exemplary embodiments of the present invention are not limited thereto.


[0150] For example, the p-dopant may include at least one selected from;

[0151] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) or 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);


[0152] a metal oxide, such as tungsten oxide or molybdenum oxide;

[0153] 1,4,5,8,9,12-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and


[0154] a compound represented by Formula 221, but exemplary embodiments of the present invention are not limited thereto:



<HAT-CN>



<F4-TCNQ>



<Formula 221>

[0155] In Formula 221:

[0156] R_{221} to R_{223} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and at least one selected from R_{221} to R_{223} may have at least one substituent selected from a cyano group, $-F$, $-Cl$, $-Br$, $-I$, a C_1 - C_{20} alkyl group substituted with $-F$, a C_1 - C_{20} alkyl group substituted with $-Cl$, a C_1 - C_{20} alkyl group substituted with $-Br$, and a C_1 - C_{20} alkyl group substituted with $-I$.

[0157] When the organic light-emitting device 10 is a full-color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In an exemplary embodiment of the present invention, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more

layers contact each other or are separated from each other. In an exemplary embodiment of the present invention, the emission layer may include two or more materials selected from a red light-emitting material, a green light-emitting material, and a blue light-emitting material, in which the two or more materials are mixed with each other in a single layer to emit white light.


[0158] The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant or a fluorescent dopant.

[0159] An amount of the dopant in the emission layer may be in a range of about 0.01 parts by weight to about 15 parts by weight based on 100 parts by weight of the host, but exemplary embodiments of the present invention are not limited thereto.

[0160] A thickness of the emission layer may be in a range of from about 100 Å to about 1,000 Å, for example, from about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0161] The host may include a compound represented by Formula 1.

[0162] In an exemplary embodiment of the present invention, the host may further include a compound represented by Formula 301.



[0163] In Formula 301:

[0164] Ar_{301} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0165] $xb11$ may be 1, 2, or 3,

[0166] L_{301} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0167] $xb1$ may be an integer selected from 0 to 5,

[0168] R_{301} may be selected from: deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(Q_{301})(Q_{302})(Q_{303})$, $-\text{N}(Q_{301})(Q_{302})$, $-\text{B}(Q_{301})(Q_{302})$, $-\text{C}(=\text{O})(Q_{301})$, $-\text{S}(=\text{O})_2(Q_{301})$, and $-\text{P}(=\text{O})(Q_{301})(Q_{302})$.

[0169] xb21 may be an integer selected from 1 to 5,

[0170] Q_{301} to Q_{303} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but exemplary embodiments of the present invention are not limited thereto.

[0171] In an exemplary embodiment of the present invention, Ar_{301} in Formula 301 may be selected from:

[0172] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group;

[0173] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}); and

[0174] Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but exemplary embodiments of the present invention are not limited thereto,

[0175] When xb11 in Formula 301 is two or more, two or more Ar_{301} (s) may be chemically linked to each other via a single bond.

[0176] In an exemplary embodiment of the present invention, the compound represented by Formula 301 may be represented by Formula 301-1 or 301-2:

[0177] In Formulae 301-1 to 301-2:

[0178] A_{301} to A_{304} may each independently be selected from benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, an indene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, an indole group, a carbazole group, a benzocarbazole group, a dibenzocarbazole group, a furan group, a benzofuran group, a dibenzofuran group, a naphthofuran group, a benzonaphthofuran group, a dinaphthofuran group, a thiophene group, a benzothiophene group, a dibenzothiophene group, a naphthothiophene group, a benzonaphthothiophene group group/ and a dinaphthothiophene group group,

[0179] X_{301} may be O, S, or N—[(L_{304})_{xb4}— R_{304}],

[0180] R_{311} to R_{314} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),

[0181] xb22 and xb23 may each independently be 0, 1, or 2,

[0182] L_{301} , xb1, R_{301} , Q_{31} to Q_{33} may be the same as described above,

[0183] L_{302} to L_{304} may each independently be the same as L_{301} , described above,


[0184] xb2 to xb4 may each independently be the same as xb1, described above, and

[0185] R_{302} to R_{304} may each independently be the same as R_{301} , described above.


[0186] For example, L_{301} to L_{304} in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0187] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a

<Formula 301-1>



<Formula 301-2>



chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylene group, a benzofuranylenylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group;

[0188] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenyline group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylene group, a benzofuranylenylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a benzofuranyl group, a dibenzothiophenyl group, a benzocarbazoly group, a dibenzocarbazoly group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoly group;

thiophenyl group, a benzocarbazoly group, a dibenzocarbazoly group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazoly group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}), and

[0189] Q_{31} to Q_{33} may be the same as described above.

[0190] In an exemplary embodiment of the present invention, R_{301} to R_{304} in Formulae 301, 301-1 and 301-2 may each independently be selected from:

[0191] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a benzofuranyl group, a dibenzothiophenyl group, a benzocarbazoly group, a dibenzocarbazoly group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoly group;


[0192] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a benzofuranyl group, a dibenzothiophenyl group, a benzocarbazoly group, a dibenzocarbazoly group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a

quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridiny group, an acridiny group, a phenanthroliny group, a phenaziny group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridiny group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridiny group, an acridiny group, a phenanthroliny group, a phenaziny group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridiny group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}); and

[0193] Q_{31} and Q_{33} may be the same as described above.


[0194] In an exemplary embodiment of the present invention, the host may include an alkaline earth metal complex. For example, the host may be selected from a Be complex (for example, Compound H55), a Mg complex, or a Zn complex.

[0195] The host may include at least one selected from 9,10-di(2-naphthyl)anthracene (ADM), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55, but exemplary embodiments of the present invention are not limited thereto:




H1


-continued




H2




H3




H4



H5




H6




H7

-continued


H8




H9




H10




H11



H12




H13




-continued


H14




H15




H16







H17






H18




-continued



-continued




-continued




H26


-continued




H29




H27




H30



H28




H31




H32

-continued


H33




H34



H35




H36




-continued


H37




H38




H39



H40




-continued




H41

-continued




H44


H42




H46




H43




H47



H48




-continued




H49

-continued




H54


H50



H55




H51




[0196] The phosphorescent dopant may include an organometallic complex represented by Formula 401 below:


H52




<Formula 401>



<Formula 402>



H53



[0197] In Formulae 401 and 402:

[0198] M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm):

[0199] L_{401} may be selected from ligands represented by Formula 402, $xc1$ may be 1, 2, or 3, and, when $xc1$ is two or more, two or more $L_{401}(s)$ may be identical to or different from each other,

[0200] L_{402} may be an organic ligand, $xc2$ may be an integer selected from 0 to 4, and, when $xc2$ is two or more, two or more $L_{402}(s)$ may be identical to or different from each other,

[0201] X_{401} to X_{404} may each independently be nitrogen or carbon,

[0202] X_{401} and X_{403} may be chemically linked to each other via a single bond or a double bond, and X_{402} and X_{404} may be chemically linked to each other via a single bond or a double bond,

[0203] A_{401} and A_{402} may each independently be selected from a C_5 - C_{60} carbocyclic group or a C_1 - C_{60} heterocyclic group,

[0204] X_{405} may be a single bond, $*-O-*$, $*-S-*$, $*-C(=O)-*$, $*-N(Q_{411})-*$, $*-C(Q_{411})(Q_{412})-*$, $*-C(Q_{411})=C(Q_{412})-*$, $*-C(Q_{411})=*$, or $*=C-*$, Q_{411} and Q_{412} may be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

[0205] X_{406} may be a single bond, O, or S,

[0206] R_{401} and R_{402} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_1 - C_{20} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})(Q_{402})$, $-B(Q_{401})(Q_{402})$, $-C(=O)(Q_{401})$, $-S(=O)_2(Q_{401})$, and $-P(=O)(Q_{401})(Q_{402})$;

[0207] Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a C_6 - C_{20} aryl group, and a C_1 - C_{20} heteroaryl group,

[0208] $xc11$ and $xc12$ may each independently be an integer selected from 0 to 10, and

[0209] $*$ and $*$ ' in Formula 402 each indicate a binding site to M in Formula 401.

[0210] In an exemplary embodiment of the present invention, A_{401} and A_{402} in Formula 402 may each independently be a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole(isoxazole) group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran (benzofuran) group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

[0211] In an exemplary embodiment of the present invention, in Formula 402, i) X_{401} may be nitrogen, and X_{402} may be carbon, or ii) X_{401} and X_{402} may each be nitrogen at the same time.

[0212] In an exemplary embodiment of the present invention, R_{401} and R_{402} in Formula 401 may each independently be selected from:

[0213] hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0214] a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

[0215] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0216] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0217] $-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})(Q_{402})$, $-B(Q_{401})(Q_{402})$, $-C(=O)(Q_{401})$, $-S(=O)_2(Q_{401})$ and $-P(=O)(Q_{401})(Q_{402})$; and










[0218] Q_{401} to Q_{403} may be each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group, but exemplary embodiments of the present invention are not limited thereto.

[0219] In an exemplary embodiment of the present invention, when $xc1$ in Formula 401 is two or more, two $A_{401}(s)$ in two or more $L_{401}(s)$ may be chemically linked to each other via X_{407} , which is a linking group, or two $A_{402}(s)$ in two or more $L_{401}(s)$ may be chemically linked to each other via X_{408} , which is a linking group (see Compounds PD1 to PD4 and PD7). X_{407} and X_{408} may each independently be a single bond, $*-O-*$, $*-S-*$, $*-C(=O)-*$, $*-N(Q_{413})-*$, $*-C(Q_{413})(Q_{414})-*$, or $*-C(Q_{413})=C(Q_{414})-*$ (e.g., in which Q_{413} and Q_{414} may each independently be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a


C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group), but exemplary embodiments of the present invention are not limited thereto.

[0220] L₄₀₂ in Formula 401 may be a monovalent, divalent, or trivalent organic ligand. For example, L₄₀₂ may be selected from halogen, diketone (for example, acetylacetonate), carboxylic acid (for example, picolinate), —C(=O), isonitrile, —CN, and a phosphorus-containing material (for example, phosphine, or phosphite), but exemplary embodiments of the present invention are not limited thereto.

[0221] In an exemplary embodiment of the present invention, the phosphorescent dopant may be selected from, for example, Compounds PD1 to PD25, but exemplary embodiments of the present invention are not limited thereto:




-continued




PD10


-continued




PD15




PD11




PD16




PD12




PD17




PD13



PD18







PD14




PD19

-continued



-continued


PD20



PD24

PD25

PD21




[0222] The fluorescent dopant may include a compound represented by Formula 1.

[0223] In an exemplary embodiment of the present invention, the fluorescent dopant may include an arylamine compound or a styrylamine compound.

[0224] The fluorescent, dopant may include a compound represented by Formula 501 below.

PD22

<Formula 501>



[0225] In Formula 501:

[0226] Ar_{501} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

PD23

[0227] L_{501} to L_{503} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0228] $xd1$ to $xd3$ may each independently be an integer selected from 0 to 3,

[0229] R_{501} and R_{502} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl

group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

[0230] $xd4'$ may be an integer selected from 1 to 6.

[0231] In an exemplary embodiment of the present invention, Ar_{501} in Formula 501 may be selected from:

[0232] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenanthracene group, and an indenophenanthrene group; and

[0233] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0234] In an exemplary embodiment of the present invention, L_{501} to L_{503} in Formula 501 may each independently be selected from:

[0235] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group; and

[0236] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro

group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoly group, a dibenzocarbazoly group, a dibenzosilolyl group, and a pyridinyl group.

[0237] In an exemplary embodiment of the present invention, R_{501} and R_{502} in Formula 501 may each independently be selected from:


[0238] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoly group, a dibenzocarbazoly group, a dibenzosilolyl group, and a pyridinyl group;

[0239] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoly group, a dibenzocarbazoly group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoly group, a dibenzocarbazoly group, a dibenzosilolyl group, and a pyridinyl group, and —Si(Q_{31})(Q_{32})(Q_{33}); and


[0240] Q_{31} to Q_{33} may be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0241] In an exemplary embodiment of the present invention, $xd4'$ in Formula 501 may be 2, but exemplary embodiments of the present invention are not limited thereto.

[0242] For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:




FD1




-continued


FD4




FD2



FD5




FD3



FD6


-continued

FD7




-continued


FD10




FD11




FD8




FD12




FD13




FD14




FD9




-continued




FD15



FD16




FD17




FD18

-continued


FD19




FD20




FD21




FD22





[0243] In an exemplary embodiment of the present invention, the fluorescent dopant may be selected from the following compounds, but exemplary embodiments of the present invention are not limited thereto.




DPVBi




DPAVB




DCM



DCJT



Coumarin 6



C545T

[0244] The electron transport region (e.g., the electron transport region in organic layer 150) may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers each including a plurality of different materials.

[0245] The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, or an electron injection layer, but exemplary embodiments of the present invention are not limited thereto.

[0246] For example, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, and for each structure, the layers may be sequentially stacked from an emission layer. However, exemplary embodiments of the present invention are not limited thereto.


[0247] The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include a metal-free compound including at least one π electron-depleted nitrogen-containing ring.

[0248] The “ π electron-depleted nitrogen-containing ring” indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

[0249] For example, the “ π electron-depleted nitrogen-containing ring” may be i) a 60-membered to 7-membered heteromonocyclic group having at least one moiety, ii) a heteropolycyclic group in which two or more 5-membered to 7-membered heteromonocyclic groups each having at least one $*-N=*$ moiety are condensed with each other, or iii) a heteropolycyclic group in which at least one of 5-membered to 7-membered heteromonocyclic groups, each having at least one $*-N=*$ moiety, is condensed with at least one C_5 - C_{60} carbocyclic group.

[0250] Examples of the π electron-depleted nitrogen-containing ring include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoxaline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, thiadiazol, an imidazopyridine, an imidazopyrimidine, and an azacarbazole, but exemplary embodiments of the present invention are not limited thereto.

[0251] For example, the electron transport region may include a compound represented by formula 601:



[0252] In Formula 601:

[0253] Ar_{601} may be selected from a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0254] $xe11$ may be 1, 2, or 3,

[0255] L_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or

unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0256] $xe1$ may be an integer selected from 0 to 5,

[0257] R_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{601})(Q_{602})(Q_{603})$, $-C(=O)(Q_{601})$, $-S(=O)_2(Q_{601})$, and $-P(=O)(Q_{601})(Q_{602})$,

[0258] Q_{601} to Q_{603} may each independently be a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

[0259] $xe21$ may be an integer selected from 1 to 5.

[0260] In an exemplary embodiment of the present invention, at least one of $Ar_{601}(s)$ in the number of $xe11$ and $R_{601}(s)$ in the number of $xe21$ may include the π electron-depleted nitrogen-containing ring.

[0261] In an exemplary embodiment of the present invention, ring Ar_{601} in Formula 601 may be selected from:

[0262] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoxaline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group;

[0263] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a

pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, phenanthroline group, phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a —Si(Q₃₁)(Q₃₂)(Q₃₃), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂); and


[0264] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0265] When xe11 In Formula 601 is two or more, two or more Ar₆₀₁(s) may be chemically linked to each other via a single bond.

[0266] In an exemplary embodiment of the present invention, Ar₆₀₁ in Formula 601 may be an anthracene group.

[0267] In an exemplary embodiment of the present invention, a compound represented by Formula 601 may be represented by Formula 601-1:

<Formula 601-1>



[0268] In Formula 601-1:

[0269] X₆₁₄ may be N or C(R₆₁₄), X₆₁₅ may be N or C(R₆₁₅), X₆₁₆ may be N or C(R₆₁₆), at least one selected from X₆₁₄ to X₆₁₆ may be N,

[0270] L₆₁₁ to L₆₁₃ may each independently be the same as L₆₀₁,

[0271] xe611 to xe613 may each independently be the same as xe1,

[0272] R₆₁₁ to R₆₁₃ may each independently be the same as R₆₀₁, and

[0273] R₆₁₄ to R₆₁₆ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0274] In an exemplary embodiment of the present invention, L₆₀₁ and L₆₁₁, L₆₁₂ and L₆₁₃ in Formulae 601 and 601-1 may each independently be selected from:

[0275] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a

chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

[0276] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzo-

thiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, but exemplary embodiments of the present invention are not limited thereto.

[0277] In an exemplary embodiment of the present invention, xe1 and xe611 to xe613 in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

[0278] In an exemplary embodiment of the present invention, R₆₀₁ and R₆₁₁, R₆₁₂ and R₆₁₃ in Formula 801 and 801-1 may each independently be selected from:

[0279] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;

[0280] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quino-


linyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylene group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;

[0281] —S(=O)₂(Q₆₀₁) and —P(=O)(Q₆₀₁)(Q₆₀₂); and


[0282] Q₆₀₁ and Q₆₀₂ may be the same as described above.

[0283] The electron transport region may include at least one compound selected from Compounds ET1 to ET36, but exemplary embodiments of the present invention are not limited thereto:

ET1




-continued




ET2


-continued




ET5




ET3



ET6



ET4





ET7

-continued

-continued



ET10

ET8




ET11

ET9




ET12




-continued

ET13




-continued


ET16




ET17




ET14






ET15



ET18




-continued




ET19

-continued




ET22

ET20



ET23


ET21



ET24


-continued

ET25





-continued

ET28




ET29


ET26




ET27



ET30




-continued




ET31


-continued




ET34




ET32



ET35




ET33



ET36

[0284] In an exemplary embodiment of the present invention, the electron transport region may include at least one selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), AlQ₃, BAQ, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), or NTAZ.



[0285] Thicknesses of the buffer layer, the hole blocking layer, and the electron control layer may each be in a range of about 20 Å to about 1,000 Å, for example, from about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.

[0286] A thickness of the electron transport layer may be in a range of from about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described


above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

[0287] The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.


[0288] The metal-containing material may include at least one selected from alkali metal complex or alkaline earth-metal complex. The alkali metal complex may include a metal ion selected from a Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion, and the alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, a Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenylan oxazole, a hydroxy phenylthiazole, a hydroxy diphenylan oxadiazole, a hydroxy diphenylthiadiazol, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bypyridine, a phenanthroline, and a cyclopentadiene, but exemplary embodiments of the present invention are not limited thereto.

[0289] For example, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.

ET-D1



ET-D2



[0290] The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode 190. The electron injection layer may be in direct contact with the second electrode 190.

[0291] The electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers, each including a plurality of different materials.

[0292] The electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare

earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combinations thereof.

[0293] The alkali metal may be selected from Li, Na, K, Rb, and Cs. In an exemplary embodiment of the present invention, the alkali metal may be Li, Na, or Cs. In an exemplary embodiment of the present invention, the alkali metal may be Li or Cs, but exemplary embodiments of the present invention are not limited thereto.

[0294] The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

[0295] The rare earth metal may be selected from Sc, Y, Ge, Tb, Yb, and Gd.

[0296] The alkali metal compound, the alkaline earth-metal compound, and the rare earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal, and the rare earth metal.

[0297] The alkali metal compound may be selected from alkali metal oxides, such as Li_2O , Cs_2O , or K_2O , and alkali metal halides, such as LiF, NaF, CsF, KF, LiI, NaI, CsI, or KI. In an exemplary embodiment of the present invention, the alkali metal compound may be selected from LiF, Li_2O , NaF, LiI, NaI, CsI, or KI, but exemplary embodiments of the present invention are not limited thereto.

[0298] The alkaline earth-metal compound may be selected from alkaline earth-metal oxides, such as BaO, SrO, CaO, $\text{Ba}_x\text{Sr}_{1-x}\text{O}$ ($0 < x < 1$), or $\text{Ba}_x\text{Ca}_{1-x}\text{O}$ ($0 < x < 1$). In an exemplary embodiment of the present invention, the alkaline earth-metal compound may be selected from BaO, SrO, and CaO, but exemplary embodiments of the present invention are not limited thereto.

[0299] The rare earth metal compound may be selected from YbF_3 , ScF_3 , ScO_3 , Y_2O_3 , Ce_2O_3 , GdF_3 , and TbF_3 . In an exemplary embodiment of the present invention, the rare earth metal compound may be selected from YbF_3 , ScF_3 , TbF_3 , YbI_3 , ScI_3 , and TbI_3 , but exemplary embodiments of the present invention are not limited thereto.

[0300] The alkali metal complex, the alkaline earth-metal complex, and the rare earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare earth metal as described above, and a ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, or the rare earth metal complex may be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenylan oxazole, hydroxy phenylthiazole, hydroxy diphenylan oxadiazole, hydroxy diphenylthiadiazol, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene, but exemplary embodiments of the present invention are not limited thereto.

[0301] The electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combinations thereof, as described above. In an exemplary embodiment of the present invention, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal com-

plex, an alkaline earth-metal complex, a rare earth metal complex, or any combinations thereof may be substantially homogeneously or non-homogeneously dispersed in a matrix including the organic material.

[0302] A thickness of the electron injection layer may be in a range of from about 1 Å to about 100 Å, for example, from about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

[0303] The second electrode 190 may be disposed on the organic layer 150. The second electrode 190 may be a cathode which is an electron injection electrode, and in this regard, a material included in the second electrode 190 may be selected from metal, an alloy, an electrically conductive compound, and a combination thereof, which may have a relatively low work function.

[0304] The second electrode 190 may include at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO, but exemplary embodiments of the present invention are not limited thereto. The second electrode 190 may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

[0305] The second electrode 190 may have a single-layered structure, or a multi-layered structure including two or more layers.

[0306] The organic light-emitting device according to an exemplary embodiment of the present invention is described above in more detail with reference to FIG. 1, and thus duplicative descriptions may be omitted.

[0307] Layers included in the hole transport region, an emission layer, and layers included in the electron transport region may be formed in a certain region by using one or more methods selected from, for example, vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

[0308] When layers included in the hole transport region, an emission layer, and layers included in the electron transport region are formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of from about 100° C. to about 500° C., at a vacuum degree of about 10^{-8} torr to about 10^{-3} torr, and at a deposition rate of from about 0.01 Å/sec to about 100 Å/sec by taking into account a material to be included in a layer to be formed, and the structure of a layer to be formed.

[0309] When layers included in the hole transport region, an emission layer, and layers included in the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of from about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to about 200° C. by taking into account a material to be included in a layer to be formed, and the structure of a layer to be formed.

[0310] The term “ $\text{C}_1\text{-C}_{60}$ alkyl group” used herein refers to a linear or branched aliphatic saturated hydrocarbon monovalent group having 1 to 60 carbon atoms, and examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, or a hexyl

group. The term “C₁-C₆₀ alkylene group” used herein refers to a divalent group having substantially the same structure as the C₁-C₆₀ alkyl group.

[0311] The term “C₂-C₆₀ alkenyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon double bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C₂-C₆₀ alkenylene group” as used herein refers to a divalent group having substantially the same structure as the C₂-C₆₀ alkenyl group.

[0312] The term “C₂-C₆₀ alkynyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethynyl group, or a propynyl group. The term “C₂-C₆₀ alkynylene group” as used herein refers to a divalent group having substantially the same structure as the C₂-C₆₀ alkynyl group.

[0313] The term “C₁-C₆₀ alkoxy group” as used herein refers to a monovalent group represented by —OA₁₀₁ (e.g., in which A₁₀₁ is the C₁-C₆₀ alkyl group), and examples thereof include a methoxy group, an ethoxy group, or an isopropoxy group.

[0314] The term “C₃-C₁₀ cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, or a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group” as used herein refers to a divalent group having the substantially same structure as the C₃-C₁₀ cycloalkyl group.

[0315] The term C₁-C₁₀ heterocycloalkyl group used herein refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, Si, P, and B as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuranlyl group, or a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group” as used herein refers to a divalent group having substantially the same structure as the C₁-C₁₀ heterocycloalkyl group.

[0316] The term C₃-C₁₀ cycloalkenyl group used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and no aromaticity, and examples thereof include a cyclopentenyl group, a cyclohexenyl group, or a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group” as used herein refers to a divalent group having substantially the same structure as the C₃-C₁₀ cycloalkenyl group.

[0317] The term “C₁-C₁₀ heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Non-limiting examples of the C₁-C₁₀ heterocycloalkenyl group include a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranlyl group, or a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group” as used herein refers to a divalent group having substantially the same structure as the C₁-C₁₀ heterocycloalkenyl group.

[0318] The term “C₆-C₆₀ aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C₁-C₆₀ arylene group used herein refers to a divalent group having a carbocyclic

aromatic system having 6 to 60 carbon atoms. Non-limiting examples of the C₆-C₆₀ aryl group include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, or a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be chemically bonded to each other.

[0319] The term “C₁-C₆₀ heteroaryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C₁-C₆₀ heteroarylene group” as used herein refers to a divalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Non-limiting examples of the C₁-C₆₀ heteroaryl group include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, or an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be chemically bonded with each other.

[0320] The term “C₆-C₆₀ aryloxy group” as used herein refers to —OA₁₀₂ (e.g., in which A₁₀₂ is the C₆-C₆₀ aryl group), and a C₆-C₆₀ arylthio group used herein indicates —SA₁₀₃ (e.g., in which A₁₀₃ is the C₆-C₆₀ aryl group).

[0321] The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) having two or more rings chemically bonded with each other, only carbon atoms as ring-forming atoms, and no aromaticity in its entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group,” used herein, refers to a divalent group having substantially the same structure as the monovalent non-aromatic condensed polycyclic group.

[0322] The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group (for example, having 1 to 60 carbon atoms) having two or more rings condensed to each other, at least one heteroatom selected from N, O, Si, P, and S, other than carbon atoms, as a ring-forming atom, and no aromaticity in its entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein refers to a divalent group having substantially the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0323] The term “C₅-C₆₀ carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which a ring-forming atom is a carbon atom only. The C₁-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₅-C₆₀ carbocyclic group may be a ring, such as benzene, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In an exemplary embodiment of the present invention, depending on the number of substituents connected to the C₅-C₆₀ carbocyclic group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

[0324] The term “C₁-C₆₀ heterocyclic group” as used herein refers to a group having substantially the same structure as the C₁-C₆₀ carbocyclic group, except that as a

ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used in addition to carbon (e.g., the number of carbon atoms may be in a range of 1 to 60).

[0325] At least one substituent of the substituted C_5 - C_{60} carbocyclic group, substituted C_1 - C_{60} heterocyclic group, substituted C_3 - C_{10} cycloalkylene group, substituted C_1 - C_{10} heterocycloalkylene group, substituted C_3 - C_{10} cycloalkenylene group, substituted C_1 - C_{10} heterocycloalkenylene group, substituted C_6 - C_{60} arylene group, substituted C_1 - C_{60} heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C_1 - C_{60} alkyl group, substituted C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group, substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10} cycloalkyl group, substituted C_1 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substituted C_1 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_1 - C_{60} heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, or substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0326] deuterium ($-D$), $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0327] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{11})(Q_{12})(Q_{13})$, $-N(Q_{11})(Q_{12})$, $-B(Q_{11})(Q_{12})$, $-C(=O)(Q_{11})$, $-S(=O)_2(Q_{11})$, and $-P(=O)(Q_{11})Q_{12}$;

[0328] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0329] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a

monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{21})(Q_{22})(Q_{23})$, $-N(Q_{21})(Q_{22})$, $-B(Q_{21})(Q_{22})$, $-C(=O)(Q_{21})$, $-S(=O)_2(Q_{21})$ and $-P(=O)(Q_{21})Q_{22}$;

[0330] $-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})Q_{32}$; and

[0331] Q_{11} to Q_{13} , Q_{21} to Q_{23} and Q_{31} to Q_{33} may be each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

[0332] The term “Ph” used herein refers to a phenyl group, the term “Me” used herein refers to a methyl group, the term “Et” used herein refers to an ethyl group, the term “ter-Bu” or “Bu” used herein refers to a tert-butyl group, and the term “OMe” used herein refers to a methoxy group.

[0333] The term “biphenyl group” as used herein refers to “a phenyl group substituted with a phenyl group.” As an example, the “biphenyl group” is a substituted phenyl group having a C_6 - C_{60} aryl group as a substituent.

[0334] The term “terphenyl group” as used herein refers to “a phenyl group substituted with a biphenyl group.” As an example, the “terphenyl group” is a phenyl group having, as a substituent, a C_6 - C_{60} aryl group substituted with a C_6 - C_{60} aryl group.

[0335] * and *' used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.

[0336] A compound according to an exemplary embodiment of the present invention and an organic light-emitting device according to an exemplary embodiment of the present invention will be described in more detail with reference to Synthesis Examples and Examples. The wording “B was used instead of A” used in describing Synthesis Examples refers to that an identical molar equivalent of B was used in place of A.


[0337] FIG. 2 is a graph showing light emission characteristics of organic light-emitting devices including TADF emitters, according to Example 1 and Comparative Examples 1 and 2. FIG. 3 is a graph showing external quantum efficiency (EQE) and emission color with respect to a driving current density in organic light-emitting devices including TADF emitters, according to Example 1 and Comparative Example 1 and 2.

EXAMPLE

Synthesis Examples 1 to 3

Synthesis of Compounds 1 to 3

[0338]




[0339] 10.1 mmol of phenoxazine and 20.2 mmol of potassium tert-butoxide may be added to a round-bottom flask, and the round bottom flask was filled with Ar gas. The resultant mixture may be stirred under reflux for about 30 minutes by using 20 ml of anhydrous dimethylformamide (DMF) as a solvent. A solution in which 7.49 mmol of 4,4'-difluorobenzophenone is dissolved in 10 ml of anhydrous DMF may be added to the round-bottom flask and stirred under reflux at a temperature of about 75° C. for about 24 hours. After the reaction is completed, the reaction solution may be dissolved in water, and an organic material may be extracted therefrom by using methylene chloride and purified by column chromatography to obtain Compound 3 of a yellow solid phase. Compounds 1 and 2 may be synthesized in substantially the same manner as in Compound 3, except that potassium methoxide and potassium ethoxide may each be used instead of potassium tert-butoxide.

Synthesis Example 4

Synthesis of Compound A

[0340]





[0341] 8.24 mmol of phenoxazine and 15.0 mmol of potassium tert-butoxide may be added to a round-bottom flask, and the round-bottom flask may be filled with Ar gas. The resultant mixture may be stirred under reflux for about 30 minutes by using 20 ml of anhydrous DMF as a solvent. A solution in which 7.49 mmol of 4-fluorobenzophenone is dissolved in 10 ml of anhydrous DMF may be added to the round-bottom flask and stirred under reflux at a temperature of about 110° C. for about 12 hours. After the reaction was completed, the reaction solution may be dissolved in water, and an organic material may be extracted therefrom by using methylene chloride and purified by column chromatography to obtain Compound A of a yellow solid phase.

Synthesis Example 5

Synthesis of Compound B

[0342]






[0343] 13.3 mmol of phenoxazine and 26.6 mmol of potassium tert-butoxide may be added to a round-bottom flask, and the round-bottom flask may be filled with Ar gas. The resultant mixture may be stirred under reflux for about 30 minutes by using 15 ml of anhydrous DMF as a solvent. A solution in which 13.3 mmol of 4,4'-difluorobenzophenone is dissolved in 5 ml of anhydrous DMF may be added to the round-bottom flask and stirred under reflux at a temperature of about 110° C. for about 12 hours. After the reaction is completed, the reaction solution may be dissolved in water, and an organic material may be extracted therefrom by using methylene chloride and purified by column chromatography to obtain Compound B of a yellow solid phase.

Example 1


[0344] As an ITO anode, an ITO glass substrate (e.g., manufactured by Samsung Corning Co., Ltd.), on which an ITO layer is deposited to a thickness of 15 Ω/cm^2 (e.g., about 1,200 Å), may be cut to a size of about 25 mm×25 mm×0.7 mm, sonicated with isopropyl alcohol and pure water each for 5 minutes, and then cleaned by exposure to ultraviolet rays and ozone for about 30 minutes. Then, the resultant ITO glass substrate may be positioned in a vacuum deposition apparatus.

[0345] α -NPD may be vacuum-deposited on the ITO anode formed on the ITO glass substrate to form a hole transport layer having a thickness of about 65 nm.


[0346] mCP and Compound 3 (5 wt %) may be co-deposited on the hole transport layer to form an emission layer having a thickness of about 30 nm. Then, 2,2',2''-(1,3,5-benzinetriyl)-bis(1-phenyl-1-Hbenzimidazole) (TPBi) may be deposited on the emission layer to form an electron transport layer having a thickness of about 30 nm. LiF may be deposited on the electron transport layer to form an electron injection layer having a thickness of about 1 nm, and Al may be deposited on the electron injection layer to form a cathode having a thickness of about 100 nm, thus forming an organic light-emitting device. The deposition apparatus may be Suicel plus 200 (manufactured by Sunic System Co., Ltd.).



< α -NPD>



< m CP>




<TPBi>

Comparative Example 1

[0347] An organic light-emitting device may be manufactured in substantially the same manner as in Example 1, except that Compound A may be used instead of Compound 3 in forming an emission layer.

<Compound A>




Comparative Example 2

[0348] An organic light-emitting device may be manufactured in substantially the same manner as in Example 1,

except that Compound B may be used instead of Compound 3 in forming an emission layer.

<Compound B>



[0349] A current-voltage meter (e.g., Keithley SMU 236) may supply power to the organic light-emitting devices manufactured according to Example 1 and Comparative Examples 1 to 2, and a luminance meter may be used to measure luminance, color coordinates, current efficiency, emission wavelength range, external quantum efficiency, and emission color. Exemplary results thereof are shown in Table 1 and FIGS. 2 and 3.

TABLE 1

	Driving voltage	Maximum current efficiency	Maximum external Quantum efficiency	Maximum emission wavelength	FWHM	Color coordinates	
	[V]	[cd/A]	[%]	[nm]	[nm]	CIE _x	CIE _y
Example 1	3.6	29.5	10.4	509	94	0.28	0.52
Comparative Example 1	3.6	31.5	10.6	516	103	0.31	0.53
Comparative Example 2	3.2	29.0	9.2	541	113	0.39	0.55

[0350] Referring to Table 1 and FIGS. 2 and 3, when a compound according to an exemplary embodiment of the present invention is used as a host of an emission layer, the driving voltage and efficiency similar to those of an existing TADF material are exhibited and a full width at half maximum (FWHM) is reduced to narrow an emission wavelength region thus increasing color purity, but in the case of Comparative Example 1 or 2 that uses a TADF material having an existing symmetrical structure or a compound in which only phenoxazine is substituted, an emission wavelength region is excessively wide, and thus, it is difficult to exhibit color purity of a desired level.

[0351] As an example, a main emission wavelength region of the organic light-emitting device of Example 1 may be shifted to a shorter wavelength, as compared with the organic light-emitting device of Comparative Example 2 using a TADF material having a symmetrical structure.

[0352] According to an exemplary embodiment of the present invention, an organic light-emitting device may have relatively high efficiency and relatively high color purity.

[0353] It should be understood that embodiments described herein should be considered in a descriptive and not for purposes of limitation. Descriptions of features or


aspects within each embodiment should typically be considered as available for other similar features or aspects in other embodiments.

[0354] While the present invention has been shown and described with reference to the exemplary embodiments thereof, it will be apparent to those of ordinary skill in the art that various changes in form and detail may be made thereto without departing from the spirit and scope of the present invention.


What is claimed is:

1. A polycyclic compound represented by Formula 1:


<Formula 1>




<Formula 2>



<Formula 3>



<Formula 4>



wherein, in Formulae 1 to 4,

X₁ is O or S,

Y₁₁, Y₂₁, and Y₂₂ are each independently selected from a single bond, C(R₃₁)(R₃₂), Si(R₃₁)(R₃₂), O, S, and

$N(R_{31})$, and wherein Y_{21} and Y_{22} are not a single bond at the same time,

A_{11} , A_{12} , A_{21} , and A_{22} are each independently a C_3 - C_{60} carbocyclic group or a C_1 - C_{60} heterocyclic group,

L_1 , L_{11} , and L_{21} are each independently a substituted or unsubstituted C_6 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

a_1 , a_{11} , and a_{21} are each independently an integer selected from 0 to 3, wherein, when a_1 is two or more, two or more $L_1(s)$ are identical to or different from each other, when a_{11} is two or more, two or more $L_{11}(s)$ are identical to or different from each other, and when a_{21} is two or more, two or more $L_{21}(s)$ are identical to or different from each other,

Ar_1 is selected from $-F$, $-Cl$, $-Br$, $-I$, a cyano group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, $-N(Q_{101})(Q_{102})$, and $-S(Q_{101})$,

R_1 to R_{12} , R_{21} , R_{22} , R_{31} , and R_{32} are each independently selected from a group represented by one of Formulae 2 to 4, hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group, $-Si(Q_1)(Q_2)(Q_3)$, $-N(Q_1)(Q_2)$, $-B(Q_1)(Q_2)$, $-(S=O)_2(Q_1)$, and $-P(=O)(Q_1)(Q_2)$,

provided that each of R_{11} to R_{12} , R_{21} , R_{22} , R_{31} , and R_{32} is not a group represented by one of Formulae 2 to 4,

at least one of R_1 to R_5 is a group represented by Formula 2 or Formula 3,

at least one of R_6 to R_{10} is a group represented by Formula 4,

b_{11} , b_{12} , b_{21} , and b_{22} are each independently an integer selected from 1 to 15, wherein, when b_{11} is two or more, two or more $R_{11}(s)$ are identical to or different from each other, when b_{12} is two or more, two or more $R_{12}(s)$ are identical to or different from each other, when b_{21} is two or more, two or more $R_{21}(s)$ are identical to or different from each other, and when b_{22} is two or more, two or more $R_{22}(s)$ are identical to or different from each other,

at least one substituent of the substituted C_3 - C_{60} carbocyclic group, the substituted C_1 - C_{60} heterocyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_1 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{11})(Q_{12})(Q_{13})$, $-N(Q_{11})(Q_{12})$, $-B(Q_{11})(Q_{12})$, $-C(=O)(Q_{11})$, $-S(=O)_2(Q_{11})$, and $-P(=O)(Q_{11})(Q_{12})$;

a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group;

a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a

terphenyl group, $-\text{Si}(\text{Q}_{21})(\text{Q}_{22})(\text{Q}_{23})$, $-\text{N}(\text{Q}_{21})(\text{Q}_{22})$, $-\text{B}(\text{Q}_{21})(\text{Q}_{22})$, $-\text{C}(=\text{O})(\text{Q}_{21})$, $-\text{S}(=\text{O})_2(\text{Q}_{21})$, and $-\text{P}(=\text{O})(\text{Q}_{21})(\text{Q}_{22})$; and

$-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$.

Q_{101} , Q_{102} , Q_1 to Q_3 , Q_{11} to Q_{13} , Q_{21} to Q_{23} , and Q_{31} to Q_{33} are each independently selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

* indicates a binding site to a neighboring atom.

2. The polycyclic compound of claim 1, wherein, in Formulae 2 and 3,

Y_{11} is selected from a single bond, $\text{C}(\text{R}_{31})(\text{R}_{32})$, O, and S, and at least one of Y_{21} or Y_{22} is $\text{N}(\text{R}_{31})$.

3. The polycyclic compound of claim 1, wherein, in Formulae 2 to 4,

L_1 , L_{11} , and L_{21} are each independently selected from:

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a spiro-fluorene-benzofluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylene group, a benzothioiophenylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a dibenzosilolylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thia-

diazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group; and

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a spiro-fluorene-benzofluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylene group, a benzothioiophenylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a dibenzosilolylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thia-

group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolylene group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$, and

Q_{31} to Q_{33} are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a biphenyl group, a terphenyl group, a naphthyl group, and a dibenzofuranyl group.

4. The polycyclic compound of claim 1, wherein

A_{11} , A_{12} , A_{21} , and A_{22} are each independently selected from:

a phenyl group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a spiro-fluorene-benzofluorene group, a benzofluorene group, a dibenzofluorene group, a phenanthrene group, an anthracene group, a triphenylene group, a pyrenylene group, a chrysenylene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, an indole group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, an oxadiazole group, a triazine group, a dibenzofuran group, a benzonaphthofuran group, a dibenzothiophene group, a dibenzosilole group, a benzocarbazole group, an indolocarbazole group, a dibenzocarbazole group, an imidazopyridine group, and an imidazopyrimidine group; and

a phenyl group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a spiro-fluorene-benzofluorene group, a benzofluorene group, a dibenzofluorene group, a phenanthrene group, an anthracene group, a triphenylene group, a pyrenylene group, a chrysenylene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, an indole group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, an oxadiazole group, a triazine group, a dibenzofuran group, a benzonaphthofuran group, a dibenzothiophene group, a dibenzosilole


group, a benzocarbazole group, an indolocarbazole group, a dibenzocarbazole group, an imidazopyridine group, and an imidazopyrimidine group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, an indolyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a benzonaphthofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, an indolocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$, and

Q_{31} to Q_{33} are each independently selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.


5. The polycyclic compound of claim 1, wherein

the group represented by Formula 2 is selected from groups represented by Formulae 2-1 to 2-7;


2-1



-continued












-continued









wherein, in Formulae 2-1 to 2-7,
 Y_{11} , L_{11} , a_{11} , R_{11} , r_{12} , b_{11} , and b_{12} are each independently the same as in claim 1, and
 R_{13} is the same as R_{11} and R_{12} ; and
 * indicates a binding site to a neighboring atom.



6. The polycyclic compound of claim 1; wherein
 the group represented by Formula 3 is selected from
 groups represented by Formulae 3-1 to 3-13:



-continued



-continued



wherein, in Formulae 3-1 to 3-13,

Y_{21} , Y_{22} , L_{21} , a_{21} , R_{21} , R_{22} , b_{21} , and b_{22} are each independently the same as in claim 1, and

R_{23} is the same as R_{21} and R_{22} , and

* indicates a binding site to a neighboring atom.

7. The polycyclic compound of claim 1, wherein b_{11} , b_{12} , b_{21} , and b_{22} are each independently an integer selected from 1 to 6.

8. The polycyclic compound of claim 1, wherein

Ar_1 in Formula 4 from:

—F, —Cl, —Br, —I, a cyano group, a C_1 - C_{20} alkyl group, a C_2 - C_{20} alkenyl group, a C_2 - C_{20} alkynyl group, a C_1 - C_{20} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_6 - C_{20} aryl group, —N(Q_{101})(Q_{102}), and —S(Q_{101}); and

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from —F, —Cl, —Br, —I, and a C_1 - C_{20} alkyl group, and

Q_{101} and Q_{102} are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

9. The polycyclic compound of claim 1, wherein

Ar_1 in Formula 4 is selected from —F, a cyano group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a C_1 - C_{10} heterocycloalkyl group, N(Q_{101})(Q_{102}), a C_1 - C_{20} alkyl group substituted with at least one —F, and a C_1 - C_{20} alkoxy group substituted with at least one C_1 - C_{20} alkyl group, and

Q_{101} , Q_{102} , Q_1 to Q_3 , Q_{11} to Q_{13} , Q_{21} to Q_{23} , and Q_{31} to Q_{33} are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group,

10. The polycyclic compound of claim 1, wherein

R_1 to R_{10} in Formula 1 are each independently selected from:

a group represented by one of Formulae 2 to 4, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a benzonaphthofuranyl group, a dibenzothiophenyl group, a benzosilolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a benzonaphthofuranyl group, a dibenzothiophenyl group, a benzosilolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from deute-

rium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a benzonaphthofuranyl group, a dibenzothiophenyl group, a benzosilolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —(Si(Q_{31})(Q_{32})(Q_{33}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}); and

—Si(Q_1)(Q_2)(Q_3), B(Q_1)(Q_2), —C(=O)(Q_1), —S(=O)₂(Q_1), and —P(=O)(Q_1)(Q_2), and

Q_1 to Q_3 and Q_{31} to Q_{33} are each independently, selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a biphenyl group, a terphenyl group, a naphthyl group, and a carbazolyl group.

11. The polycyclic compound of claim 1, wherein

R_3 is a group represented by Formula 2 or Formula 3.

12. The polycyclic compound of claim 1, wherein

R_3 is a group represented by Formula 2 or Formula 3, and


R_6 is a group represented by Formula 4.

13. The polycyclic compound of claim 1, wherein


a difference between a singlet energy level and a triplet energy level of the polycyclic compound is 0.3 eV or less.

14. The polycyclic compound of claim 1, wherein


the polycyclic compound is selected from Compounds 1 to 9:




-continued




2




3



4




5




6


-continued



7



8



9

15. An organic light-emitting device comprising:
a first electrode;

a second electrode facing the first electrode; and

an organic layer between the first electrode and the second electrode, the organic layer comprising an emission layer and at least one of the polycyclic compound of claim 1.

18. The organic light-emitting device of claim 15, wherein the first electrode is an anode, the second electrode is a cathode, the organic layer further comprises a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode,

the hole transport region comprises a hole injection layer, a hole transport layer, an emission auxiliary layer, an electron blocking layer, or any combination thereof, and

the electron transport region comprises a hole blocking layer, an electron transport layer, an electron injection layer, or any combination thereof.

17. The organic light emitting device of claim 16, wherein the emission layer comprises the polycyclic compound.

18. The organic light-emitting device of claim 17, wherein the polycyclic compound comprised in the emission layer is a thermally activated delayed fluorescence (TADF) emitter, and

the emission layer emits delayed fluorescence.

19. The organic light-emitting device of claim **17**, wherein the emission layer consists of the polycyclic compound;
or

the emission layer further comprises a host, and an amount of the polycyclic compound is in a range of about 0.1 parts by weight to about 50 parts based on 100 parts by weight of the emission layer.

20. The organic light-emitting device of claim **16**, wherein the hole transport region comprises a p-dopant, of which a lower unoccupied molecular orbital (LUMO) energy level is -3.5 eV

* * * * *

专利名称(译)	多环化合物和包括其的有机发光装置		
公开(公告)号	US20180240982A1	公开(公告)日	2018-08-23
申请号	US15/700344	申请日	2017-09-11
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	首尔国立大学R&DB基金会		
[标]发明人	KIM SEHUN PARK SOOYOUNG RYOO CHIHYUN CHO ILLHUN YANG JUNGHOON JUNG SEYOUNG JEONG HYEIN		
发明人	KIM, SEHUN PARK, SOOYOUNG RYOO, CHIHYUN CHO, ILLHUN YANG, JUNGHOON JUNG, SEYOUNG JEONG, HYEIN		
IPC分类号	H01L51/00 C07D265/38 C07D213/16		
CPC分类号	H01L51/0071 H01L51/0059 C07D213/16 C07D265/38 C09K11/06 C09K2211/1007 C09K2211/1033 H01L51/5012 H01L2251/30 H01L2251/50		
优先权	1020170024250 2017-02-23 KR		
其他公开文献	US10593888		
外部链接	Espacenet USPTO		

摘要(译)

提供多环化合物和包含其的有机发光装置。多环化合物由式 1 表示：

