



US010305046B2

(12) **United States Patent**
Sim et al.

(10) **Patent No.: US 10,305,046 B2**

(45) **Date of Patent: May 28, 2019**

(54) **HETEROCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

(71) Applicant: **SAMSUNG DISPLAY CO., LTD.**,
Yongin-si, Gyeonggi-do (KR)

(72) Inventors: **Munki Sim**, Yongin-si (KR);
Youngkook Kim, Yongin-si (KR);
Junha Park, Yongin-si (KR);
Hyoyoung Lee, Yongin-si (KR);
Eunjae Jeong, Yongin-si (KR);
Seokhwan Hwang, Yongin-si (KR)

(73) Assignee: **SAMSUNG DISPLAY CO., LTD.**,
Yongin-si, Gyeonggi-do (KR)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 50 days.

(21) Appl. No.: **15/496,120**

(22) Filed: **Apr. 25, 2017**

(65) **Prior Publication Data**

US 2017/0338419 A1 Nov. 23, 2017

(30) **Foreign Application Priority Data**

May 19, 2016 (KR) 10-2016-0061439

(51) **Int. Cl.**

H01L 51/00 (2006.01)

H01L 27/32 (2006.01)

C07D 491/153 (2006.01)

H01L 51/52 (2006.01)

C07F 9/6561 (2006.01)

H01L 51/50 (2006.01)

(52) **U.S. Cl.**

CPC **H01L 51/0071** (2013.01); **C07D 491/153**
(2013.01); **C07F 9/6561** (2013.01);

(Continued)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,023,261 A 6/1991 Kamijo et al.

5,645,948 A 7/1997 Shi et al.

(Continued)

FOREIGN PATENT DOCUMENTS

JP 10-017860 A 1/1998

JP 11-087067 A 3/1999

(Continued)

OTHER PUBLICATIONS

Tang, et al. "Organic Electroluminescent Diodes" Applied Physics Letters vol. 51 (1987) pp. 913-915.

(Continued)

Primary Examiner — Jay Yang


(74) Attorney, Agent, or Firm — Lee & Morse, P.C.

(57) **ABSTRACT**

A heterocyclic compound and an organic light-emitting device including the same, the compound being represented by Formula 1:

$A_1-(A_2)_{a21}$, <Formula 1>

wherein, in Formula 1, A_1 is a group represented by Formula 2, A_2 is a group represented by Formula 3, and $a21$ is an integer of 1 to 11, wherein, when $a21$ is 2 to 11, two or more of A_2 are identical to or different from each other:



<Formula 2>

* —(L)_n—(Ar)_m.

<Formula 3>

20 Claims, 4 Drawing Sheets

10

190

150

110

(52) **U.S. Cl.**
 CPC **H01L 27/3211** (2013.01); **H01L 51/0052**
 (2013.01); **H01L 51/0058** (2013.01); **H01L**
51/0067 (2013.01); **H01L 51/0072** (2013.01);
H01L 51/5218 (2013.01); **H01L 51/5056**
 (2013.01); **H01L 51/5072** (2013.01); **H01L**
51/5088 (2013.01); **H01L 51/5092** (2013.01);
H01L 51/5096 (2013.01)

KR 10-2004-0057862 A 7/2004
 KR 10-2013-0135178 A 12/2013

OTHER PUBLICATIONS

Adachi, et al. "Confinement of Charge Carriers and Molecular Excitons within 5nm Thick Emitter Layer in Organic Electroluminescent Devices with a Double Heterostructure" Applied Physics Letters vol. 57 (1990) pp. 531-533.
 Johansson, et al. "Solid-State Amplified Spontaneous Emission in Some Spiro-Type Molecules: A New Concept for the Design of Solid-State Lasing Moelcules" Advanced Materials vol. 10 (1998) pp. 1136-1141.
 Tao, et al. "Sharp Green Electroluminescence from 1H-pyrazolo[3,4-b]quinolone-based Light-Emitting Diodes" Applied Physics Letters vol. 77 (2000) pp. 1575-1577.
 Sakamoto, et al. "Synthesis, Characterization, and Electron-Transport Property of Perfluorinated Phenylene Dendrimers" Journal of American Chemical Society vol. 122 (2000) pp. 1832-1833.
 Yamaguchi, et al. "Diphenylamino-Substituted 2,5-Diarylsiloles for Single-Layer Organic Electroluminescent Devices" Chemistry Letters (2001) pp. 98-99.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2004/0053069 A1 3/2004 Sotoyama et al.
 2010/0004220 A1 1/2010 Jagtap et al.

FOREIGN PATENT DOCUMENTS

JP 2004-083507 A 3/2004

FIG. 1

10

190
150
110

FIG. 2

20

190
150
110
210

FIG. 3

30

220
190
150
110

FIG. 4

40

220
190
150
110
210

1

HETEROCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

Korean Patent Application No. 10-2016-0061439, filed on May 19, 2016, in the Korean Intellectual Property Office, and entitled: "Heterocyclic Compound and Organic Light-Emitting Device Including the Same," is incorporated by reference herein in its entirety.

BACKGROUND

1. Field

Embodiments relate to a heterocyclic compound and an organic light-emitting device including the same.

2. Description of the Related Art

Organic light-emitting devices are self-emission devices and have wide viewing angles, high contrast ratios, short response times, and excellent brightness, driving voltage, and response speed characteristics, compared to devices in the art, and produce full-color images.

An example of such organic light-emitting devices may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode in this stated order. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, may recombine in the emission layer to produce excitons. These excitons transition from an excited state to a ground state, thereby generating light.



SUMMARY

Embodiments are directed to a heterocyclic compound and an organic light-emitting device including the same.

The embodiments may be realized by providing a heterocyclic compound represented by Formula 1:



wherein, in Formula 1, A_1 is a group represented by Formula 2, A_2 is a group represented by Formula 3, and $a21$ is an integer of 1 to 11, wherein, when $a21$ is 2 to 11, two or more of A_2 are identical to or different from each other:



wherein F_1 in Formula 2 is a dibenzofuran moiety or a dibenzothiophene moiety, R_1 to R_5 and R_{12} in Formula 2 are each independently selected from a binding site to A_2 ,

2

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_1)(Q_2)(Q_3), and —S(=O)₂(Q_1), b_6 in Formula 2 is an integer of 0 to 6, wherein, when b_6 is 2 to 6, two or more of R_6 are identical to or different from each other, X_1 in Formula 2 is O or S, at least one of R_1 to R_5 and R_{12} is a binding site to A_2 , a number of R_1 to R_5 and R_{12} being the binding site to A_2 being the same as $a21$, L in Formula 3 is selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{100} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group, n in Formula 3 is an integer of 0 to 7, wherein, when n is 2 to 7, two or more of L are identical to or different from each other, Ar in Formula 3 is selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_4)(Q_5)(Q_6), —C(=O)(Q_4), —S(=O)₂(Q_4), and —P(=O)(Q_4)(Q_5), m in Formula 3 is an integer of 1 to 7, wherein, when m is 2 to 7, two or more of Ar are identical to or different from each other, and at least one substituent of the substituted C_3 - C_{10} cycloalkylene group, the substituted C_1 - C_{10} heterocycloalkylene group, the substituted C_3 - C_{10} cycloalkenylene group, the substituted C_1 - C_{10} heterocycloalkenylene group, the substituted C_6 - C_{60} arylene group, the substituted C_1 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_1 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent

non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, and a substituted or unsubstituted C₁-C₆₀ alkoxy group; a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂); a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), wherein Q₁ to Q₆, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group.

The embodiments may be realized by providing an organic light-emitting device including a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode

and comprises an emission layer, wherein the organic layer includes the heterocyclic compound according to an embodiment

BRIEF DESCRIPTION OF THE DRAWINGS

Features will be apparent to those of skill in the art by describing in detail exemplary embodiments with reference to the attached drawings in which:

FIG. 1 illustrates a schematic cross-sectional diagram of an organic light-emitting device according to an embodiment;

FIG. 2 illustrates a schematic cross-sectional diagram of an organic light-emitting device according to an embodiment;

FIG. 3 illustrates a schematic cross-sectional diagram of an organic light-emitting device according to an embodiment; and


FIG. 4 illustrates a schematic cross-sectional diagram of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION

Example embodiments will now be described more fully hereinafter with reference to the accompanying drawings; however, they may be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey exemplary implementations to those skilled in the art.


In the drawing figures, the dimensions of layers and regions may be exaggerated for clarity of illustration. It will also be understood that when a layer or element is referred to as being “on” another layer or element, it can be directly on the other layer or element, or intervening layers may also be present. In addition, it will also be understood that when a layer is referred to as being “between” two layers, it can be the only layer between the two layers, or one or more intervening layers may also be present. Like reference numerals refer to like elements throughout.

A heterocyclic compound according to an embodiment may be represented by Formula 1.




In Formula 1, A₁ may be a group represented by Formula 2; A₂ may be a group represented by Formula 3; and a21 may be an integer of 1 to 11. In an implementation, when a21 is 2 to 11, two or more of A₂ may be identical to or different from each other:

<Formula 2>



<Formula 3>



F₁, b6, R₁ to R₅, R₁₂, X₁, L, Ar, n, and m in Formulae 2 and 3 are described below.

In an implementation, F_1 in Formula 2 may be, e.g., a dibenzofuran moiety or a dibenzothiophene moiety. For example, F_1 may be a dibenzofuran.

R_1 to R_5 and R_{12} in Formula 2 may each independently be selected from or include, e.g., a binding site to A_2 , hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_1)(Q_2)(Q_3)$, and $-S(=O)_2(Q_1)$.

b_6 in Formula 2 may be an integer of 0 to 6. In an implementation, when b_6 is 2 to 6, two or more of R_6 may be identical to or different from each other.

X_1 in Formula 2 may be, e.g., O or S. In an implementation, X_1 may be O. In an implementation, X_1 may be S.

In an implementation, X_1 in Formula 2 may be O, and F_1 in Formula 2 may be a dibenzofuran moiety. In an implementation, X_1 in Formula 2 may be O, and F_1 in Formula 2 may be a dibenzothiophene moiety. In an implementation, X_1 in Formula 2 may be S, and F_1 in Formula 2 may be a dibenzofuran moiety. In an implementation, X_1 in Formula 2 may be S, and F_1 in Formula 2 may be a dibenzothiophene moiety.

In an implementation, at least one selected from R_1 to R_5 and R_{12} may be a binding site to A_2 Formula 1. For example, a number of R_1 to R_5 and R_{12} that are the binding site to A_2 may be the same number as the value of a_{21} . For example, when a_{21} is 2, two of R_1 to R_5 and R_{12} are binding sites to A_2 .

a_{21} in Formula 1 may be an integer of 1 to 6. In an implementation, a_{21} may be an integer of 1 to 3. In an implementation, a_{21} may be 1.

L in Formula 3 may be selected from or include, e.g., a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.

In an implementation, L in Formula 3 may be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylene group, an indolylene group, an isoindolylene group, a benzofura-








nylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, a pyridinylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a thiadiazolylene group, an oxadiazolylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylene group, an indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, a pyridinylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a thiadiazolylene group, an oxadiazolylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzo-

7

quinoliny group, a phthalaziny group, a naphthyridiny group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridiny group, an acridiny group, a phenanthroliny group, a phenaziny group, a benzoimidazoliny group, an isobenzothiazoliny group, a benzoxazoliny group, an isobenzoxazoliny group, a triazoliny group, a tetrazoliny group, an imidazopyridiny group, an imidazopyrimidiny group, an azacarbazoliny group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$.


In an implementation, L in Formula 3 may be a group represented by one of the following Formulae 5-1 to 5-34:




8

-continued


Formula 5-8




Formula 5-9




Formula 5-10




Formula 5-11




Formula 5-12




Formula 5-13




Formula 5-14



Formula 5-15



Formula 5-16



Formula 5-1

Formula 5-2

Formula 5-3

Formula 5-4

Formula 5-5

Formula 5-6

Formula 5-7

5

10

15

20

25

30

35

40

45











50

55

60

65


9
-continued



10
-continued


Formula 5-17

5




Formula 5-18

10



Formula 5-19

15




Formula 5-20

25


Formula 5-21

30



Formula 5-22

35




Formula 5-23

45

Formula 5-24

50




Formula 5-25

55

Formula 5-26

60



65

Formula 5-27

Formula 5-28

Formula 5-29


Formula 5-30

Formula 5-31

Formula 5-32

Formula 5-33

-continued



In Formulae 5-1 to 5-34,

Y_1 may be O, S, $C(Z_3)(Z_4)$, $N(Z_5)$, or $Si(Z_6)(Z_7)$,

Z_1 to Z_7 may each independently be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazolinyl group, a benzoquinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenaziny group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafuorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indenocarbazolyl group, an indolocarbazolyl group, and $-Si(Q_{31})(Q_{32})(Q_{33})$.

d_2 may be an integer selected from 0 to 2,

d_3 may be an integer selected from 0 to 3,

d_4 may be an integer selected from 0 to 4,

d_5 may be an integer selected from 0 to 5,

d_6 may be an integer selected from 0 to 6,

d_8 may be an integer selected from 0 to 8, and

* and *' each indicate a binding site to a neighboring atom.

Formula 5-34

In an implementation, L may be a group represented by one of Formula 5-1, 5-2, or 5-13, and Z_1 in Formula 5-1, 5-2, or 5-13 may be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group.

n in Formula 3 may be an integer selected from 0 to 7. In an implementation, n in Formula 3 may be 0, 1, or 2.

In an implementation, n may be 1. When n is two or more, two or more of L may be identical to or different from each other, and when n is zero, “-(L)_n” indicates a single bond.

Ar in Formula 3 may be selected from or include, e.g., a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_4)(Q_5)(Q_6)$, $-C(=O)(Q_4)$, $-S(=O)_2(Q_4)$, and $-P(=O)(Q_4)(Q_5)$.

In an implementation, Ar in Formula 3 may be selected from:


a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenaziny group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group,


a perylenyl group, a pentaphenyl group, a hexacenyly group, a pentacenyly group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyly group, a pentacenyly group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl

group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and —Si(Q₄)(Q₅)(Q₆), —C(=O)(Q₄), —S(=O)₂(Q₄), and —P(=O)(Q₄)(Q₅).


In an implementation, Ar in Formula 3 may be a group represented by one of the following Formulae 6-1 to 6-37:




Formula 6-1




Formula 6-2




Formula 6-3




Formula 6-4




Formula 6-5



Formula 6-6











Formula 6-7




Formula 6-8

15
-continued




16
-continued


Formula 6-9




Formula 6-10




Formula 6-11




Formula 6-12




Formula 6-13




Formula 6-14



Formula 6-15



Formula 6-16



Formula 6-17

Formula 6-18

Formula 6-19








Formula 6-20

Formula 6-21

Formula 6-22

17


-continued



18


-continued

Formula 6-23




5

Formula 6-24




15

Formula 6-25




20

Formula 6-26




30

Formula 6-27




40

Formula 6-28



50

Formula 6-29



60

65

Formula 6-30

Formula 6-31

Formula 6-32

Formula 6-33

Formula 6-34

Formula 6-35

Formula 6-36

Formula 6-37

45 In Formulae 6-1 to 6-37,

Y₃₁, Y₃₃ and Y₃₇ may each independently be O, S, C(Z₃₅)(Z₃₆), N(Z₃₇), or Si(Z₃₈)(Z₃₉).Y₃₂, Y₃₄, Y₃₆, and Y₃₈ may each independently be C(Z₃₅) or N,Z₃₁ to Z₃₉ may each independently be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cyclo-

heptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group,

19

a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazolinyl group, a benzoquinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indenocarbazolyl group, an indolocarbazolyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$.

e2 may be an integer selected from 0 to 2,

e3 may be an integer selected from 0 to 3,

e4 may be an integer selected from 0 to 4,



e5 may be an integer selected from 0 to 4,

e6 may be an integer selected from 0 to 6, and

* indicates a binding site to a neighboring atom.

In an implementation, Ar in Formula 3 may be a group represented by one of Formulae 6-1 to 6-6, 6-13 to 6-29, 6-31, 6-32, and 6-37.


In an implementation, Ar in Formula 3 may be a group represented by one of the following Formulae 7-1 to 7-25:



Formula 7-1


Formula 7-2

5




Formula 7-3

10




Formula 7-4

15




Formula 7-5

20




Formula 7-6

25




Formula 7-7

30




Formula 7-8

35




Formula 7-9

40




Formula 7-10

45



Formula 7-11

50



Formula 7-12








60

20

-continued

65


21
-continued



22
-continued

Formula 7-13

5




Formula 7-14

10

Formula 7-15

15




Formula 7-16

25

Formula 7-17


30



Formula 7-18

35

40




Formula 7-19

45

50

55



Formula 7-20

Formula 7-21

Formula 7-22


Formula 7-23

Formula 7-24

23

-continued

Formula 7-25




* in Formulae 7-1 to 7-25 indicates a binding site to a neighboring atom.

In an implementation, m in Formula 3 may be an integer selected from 1 to 7. In an implementation, m in Formula 3 may be 1, 2, or 3. When m is 2 to 7, two or more of Ar may be identical to or different from each other.


In an implementation, m in Formula 3 may be 1 or 2.

In an implementation, A₁ in Formula 1 may be a group represented by one of the following Formulae 2A to 2F:


Formula 2A



Formula 2B




Formula 2C




24

-continued


Formula 2D



Formula 2E



Formula 2F







In Formulae 2A to 2F, X₁ and X₂ may each independently be, e.g., O or S. R₁ to R₁₁ may each independently be selected from or include, e.g., a binding site to A₂ in Formula 1, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), and —S(=O)₂(Q₁).

In an implementation, in Formulae 2A to 2F, X₁ and X₂ may be O, and R₅ may be a binding site to A₂.

In an implementation, both X₁ and X₂ in Formulae 2A to 2F may be O. In an implementation, X₁ and X₂ in Formulae 2A to 2F may be different from each other.

25





In an implementation, the heterocyclic compound represented by Formula 1 may be one of the following Compounds 1 to 72 and 74 to 138.



26

-continued

5




6

7


8

27
-continued


9




10



11




12



28
-continued

13


5



15

14


20



30

15


35



45

16









55



65

29
-continued

30
-continued




21

22

23

31
-continued


24



32
-continued

27

5



10


15

20

28

25


25



30

35

40




45

26

29


50




55

60

65




33
-continued



30

34
-continued



33


5

10

15

20

34




31

30

35


40



32

45

35




50


55

60


65



35
-continued




36
-continued



37
-continued

43



38
-continued


46

5


10

15

20



44 25




47

30

35

40



45


45

50


55

60

65




48




39
-continued


49



50



51




65

40
-continued

52

5



10

15


20

25

30

53

35




40

45

50

54

55



60


65

41
-continued

42
-continued

59

55




5

10

15

56




20

25

30

57




35

40

45

50

58



55

60

65

5

10

15

20

25

30

35

40

45

50

55





60

65

60

61


62




43

-continued

63




64 20



25


30



65 35

40

45



66 50

55


60

65

44

-continued

67



5

10

15

64 20

25

30

65 35

40

45

50


66 50

55


60

65

68




69




45
-continued


70




71



72



74




65

46
-continued

75


5



10

15

71




20

25

30

72




40

45

74

50

55



60

65


76

77

78


47

-continued



48


-continued



49

-continued

89



50


-continued

93

5

10


15



90


20

94



25


30



91

35


95



40

45


50



92


55

96




60

65



51
-continued

97




5

10

15

20


98



25

30

99




40

45

50

100




55

60


65

52
-continued


101




102



103



104




53
-continued


54
-continued

105

109



5




10


15

110

106



20




25


30

111

107



35




40

45


50

108

112







55







60

65

55
-continued



56
-continued




57
-continued


58
-continued

121


125




122




123




124




125



126




127



59
-continued

128



5


10

15


20

60
-continued

131



129 25




132

30

35

40

45



133



50

130

55


60

65



61
-continued

134



62
-continued


137

5

10

15

20





138

135

25

30



35



The heterocyclic compound represented by Formula 1 may include a core structure or moiety represented by Formula 1a, below, and a glass transition temperature (T_g) or a melting point thereof may be high. Accordingly, the heterocyclic compound represented by Formula 1 may have excellent heat resistance, stability to charges, and charge transport capability. An electronic device including the heterocyclic compound represented by Formula 1, e.g., an organic light-emitting device, may have high efficiency, a low driving voltage, high luminance, and a long lifespan.

50
136

<Formula 1a>



60

Also, the core structure may include a dibenzofuran moiety or a dibenzothiophene moiety (as F₁) in a structure, wherein the dibenzofuran moiety and the dibenzothiophene moiety includes an oxygen atom or a sulfur atom as an atom having high electronegativity, and intermolecular attraction becomes strong, thereby improving heat resistance. An intermolecular distance may be shortened, and molecular

packing may be increased. As a result, it may be expected to improve a charge hopping rate due to an improved transfer integral. Therefore, heat resistance, stability to charges, and charge transport capability of the heterocyclic compound represented by Formula 1 may be further improved.

Synthesis methods of the heterocyclic compound represented by Formula 1 may include suitable synthetic methods.

At least one of the heterocyclic compounds represented by Formula 1 may be used or included between a pair of electrodes of an organic light-emitting device. For example, the heterocyclic compound may be included in at least one of a hole transport region and/or an emission layer.

Thus, another aspect provides an organic light-emitting device that includes: a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode and includes an emission layer. The organic layer may include at least one heterocyclic compound represented by Formula 1.

The expression that “(an organic layer) includes at least one heterocyclic compound” used herein may include a case in which “(an organic layer) includes an identical heterocyclic compound represented by Formula 1 and a case in which (an organic layer) includes two or more different heterocyclic compounds represented by Formula 1”.

For example, the organic layer may include only Compound 1 as the heterocyclic compound. In this regard, Compound 1 may be only in the electron transport layer of the organic light-emitting device. In an implementation, the organic layer may include Compound 1 and Compound 2 as the heterocyclic compound. In this regard, Compound 1 and Compound 2 may be in an identical layer (e.g., Compound 1 and Compound 2 both may be in an electron transport layer), or may be in different layers (e.g., Compound 1 may be in an electron transport layer, and Compound 2 may be in an electron injection layer).

In an implementation, the first electrode of the organic light-emitting device may be an anode.

In an implementation, the second electrode of the organic light-emitting device may be a cathode.

In an implementation, the organic layer may include i) a hole transport region between the first electrode and the emission layer, the hole transport region including at least one selected from a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer, and ii) an electron transport region between the emission layer and the second electrode, the electron transport region including at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer. For example, the electron transport region may include a single layer or layers.

In an implementation, at least one of the electron transport region and the emission layer may include at least one of the heterocyclic compounds. For example, the electron transport region of the organic light-emitting device may include an electron transport layer or an electron injection layer, and the electron transport layer or the electron injection layer may include at least one of the heterocyclic compounds. In an implementation, the emission layer of the organic light-emitting device may include a pyrene-based compound, an arylamine-based compound, or a styryl-based compound. The heterocyclic compound included in the emission layer may act as a host, and in this case, the emission layer may further include a dopant. The dopant may be a phosphorous dopant or a fluorescent dopant.

The term “organic layer” used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of the organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

FIG. 1 illustrates a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 may include a first electrode 110, an organic layer 150, and a second electrode 190.

Hereinafter, the structure of the organic light-emitting device 10 according to an embodiment and a method of manufacturing the organic light-emitting device 10 will be described in connection with FIG. 1.

In an implementation, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be a glass substrate or a plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water-resistance.

The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode 110 on the substrate. When the first electrode 110 is an anode, the material for a first electrode may be selected from materials with a high work function to facilitate hole injection.

The first electrode 110 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode 110 is a transmissible electrode, a material for forming a first electrode may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), zinc oxide (ZnO), and any combinations thereof. In an implementation, when the first electrode 110 is a semi-transmissible electrode or a reflective electrode, a material for forming a first electrode may be selected from magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and any combinations thereof.

The first electrode 110 may have a single-layered structure, or a multi-layered structure including two or more layers. In an implementation, the first electrode 110 may have a three-layered structure of ITO/Ag/ITO.

The organic layer 150 is disposed on the first electrode 110. The organic layer 150 may include an emission layer.

The organic layer 150 may further include a hole transport region between the first electrode 110 and the emission layer, and an electron transport region between the emission layer and the second electrode 190.



The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

The hole transport region may include at least one layer selected from a hole injection layer (HIL), a hole transport layer (HTL), an emission auxiliary layer, and an electron blocking layer (EBL).

For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a structure of hole injection layer/hole transport layer, hole injection layer/hole transport layer/emission auxiliary layer, hole injection layer/emission auxiliary layer, hole transport layer/emission auxiliary layer or hole injection layer/hole transport layer/electron blocking layer, wherein for each structure, constituting layers are sequentially stacked from the first electrode 110 in this stated order.





65

The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPB), β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), polyaniline/poly(4-styrenesulfonate) (Pani/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:



66


-continued




65

67


-continued




Spiro-TPD




Spiro-NPB



methylated NPB



TAPC




HMTPD

68


-continued

<Formula 201>



5

<Formula 202>



10

In Formulae 201 and 202,

L_{201} to L_{204} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

L_{205} may be selected from $^*-\text{O}-^*$, $^*-\text{S}-^*$, $^*-\text{N}(\text{Q}_{201})-^*$, a substituted or unsubstituted C_1 - C_{20} alkenylene group, a substituted or unsubstituted C_2 - C_{20} alkenylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

$\text{xa}1$ to $\text{xa}4$ may each independently be an integer selected from 0 to 3,

$\text{xa}5$ may be an integer selected from 1 to 10, and

R_{201} to R_{204} and Q_{201} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

For example, in Formula 202, R_{201} and R_{202} may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group, and R_{203} and R_{204} may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group.

In one or more embodiments, in Formulae 201 and 202,

L_{201} to L_{205} may each independently be selected from a phenylene group, a pentalenyne group, an indenylene group, a naphthylyne group, an azulenylyne group, a heptalenylyne group, an indacenylene group, an acenaphthylyne group, a fluorenylyne group, a spiro-bifluorenylyne group, a benzofluorenylyne group, a dibenzofluorenylyne group, a phenalenylyne group, a phenanthrenylene group, an anthracenylyne group, a fluoranthenylyne group, a triphenylynylene group, a pyrenylene group, a chrysenylene group, a

naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group; and

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an indacenylenylene group, an acenaphthylenylene group, a fluorenylenylene group, a spiro-bifluorenylenylene group, a benzofluorenylenylene group, a dibenzofluorenylenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenylyl group, a picenyl group, a perylenyl group, a pentaphenylyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenylyl group, an ovalenylyl group, a thiophenylyl group, a furanylyl group, a carbazolylyl group, an indolylyl group, an isoindolylyl group, a benzofuranylyl group, a benzothiophenylyl group, a dibenzofuranylyl group, a dibenzothiophenylyl group, a benzocarbazolylyl group, a dibenzocarbazolylyl group, a dibenzosilolylyl group, a pyridinylyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, xa1 to xa4 may each independently be 0, 1, or 2.

In one or more embodiments, xa5 may be 1, 2, 3, or 4.

In one or more embodiments, R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl

group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenylyl group, a picenyl group, a perylenyl group, a pentaphenylyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenylyl group, an ovalenylyl group, a thiophenylyl group, a furanylyl group, a carbazolylyl group, an indolylyl group, an isoindolylyl group, a benzofuranylyl group, a benzothiophenylyl group, a dibenzofuranylyl group, a dibenzothiophenylyl group, a benzocarbazolylyl group, a dibenzocarbazolylyl group, a dibenzosilolylyl group, and a pyridinylyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenylyl group, a picenyl group, a perylenyl group, a pentaphenylyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenylyl group, an ovalenylyl group, a thiophenylyl group, a furanylyl group, a carbazolylyl group, an indolylyl group, an isoindolylyl group, a benzofuranylyl group, a benzothiophenylyl group, a dibenzofuranylyl group, a dibenzothiophenylyl group, a benzocarbazolylyl group, a dibenzocarbazolylyl group, a dibenzosilolylyl group, and a pyridinylyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenylyl group, a picenyl group, a perylenyl group, a pentaphenylyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenylyl group, an ovalenylyl group, a thiophenylyl group, a furanylyl group, a carbazolylyl group, an indolylyl group, an isoindolylyl group, a benzofuranylyl group, a benzothiophenylyl group, a dibenzofuranylyl group, a dibenzothiophenylyl group, a benzocarbazolylyl group, a dibenzocarbazolylyl group, a dibenzosilolylyl group, a pyridinylyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

wherein Q₃₁ to Q₃₃ are the same as described above.

In one or more embodiments, at least one selected from R₂₀₁ to R₂₀₃ in Formula 201 may each independently be selected from:

a fluorenyl group, a spiro-bifluorenyl group, a carbazolylyl group, a dibenzofuranylyl group, and a dibenzothiophenylyl group; and

a fluorenyl group, a spiro-bifluorenyl group, a carbazolylyl group, a dibenzofuranylyl group, and a dibenzothiophenylyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl

71

group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.

In one or more embodiments, in Formula 202, i) R_{201} and R_{202} may be linked via a single bond, and/or ii) R_{203} and R_{204} may be linked via a single bond.


In one or more embodiments, at least one selected from R_{201} to R_{204} in Formula 202 may be selected from:

a carbazolyl group; and

a carbazolyl group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.


The compound represented by Formula 201 may be represented by Formula 201A:

<Formula 201A>



In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A(1):


<Formula 201A(1)>



In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A-1:


72

<Formula 201A-1>




In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A:

<Formula 202A>



In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A-1:

<Formula 202A-1>



In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1, L_{201} to L_{203} , $xa1$ to $xa3$, $xa5$, and R_{202} to R_{204} are the same as described above,

R_{211} and R_{212} are the same as described in connection with R_{203} ,

R_{213} to R_{217} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group,

73


a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzo-

74


furanyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

5 The hole transport region may include at least one compound selected from Compounds HT1 to HT39:


HT1




HT2



HT3



HT4





75

-continued
HT5



76

HT6



HT7

HT8





77

-continued
HT9



78

HT10



HT11

HT12





79

80

-continued



HT13

HT14





HT15

HT16



HT17

HT18





81

82

-continued



HT19

HT20




HT21

HT22




HT23




83

-continued
HT24





84

HT25





HT26

HT27



HT28

HT29





85

86

-continued



HT30

HT31





HT32

HT33




HT34

HT35




87

-continued
HT36



88

HT37



HT38

HT39



A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 9,000 Å, e.g., about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, e.g., about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer, and the electron blocking layer may block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include the materials as described above.

The hole transport region may further include, in addition to these materials, a charge-generation material for improv-

ing conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

50 The charge-generation material may be, for example, a p-dopant.

In one or more embodiments, a lowest unoccupied molecular orbital (LUMO) level of the p-dopant may be -3.5 eV or less.

55 The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound.


For example, the p-dopant may include at least one selected from:

60 a quinone derivative, such as tetracyanoquinodimethane (TCNQ) or 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);

a metal oxide, such as a tungsten oxide or a molybdenum oxide;

65 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and

a compound represented by Formula 221 below:



<Formula 221>

In Formula 221,

R_{221} to R_{223} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, wherein at least one selected from R_{221} to R_{223} has at least one substituent selected from a cyano group, —F, —Cl, —Br, —I, a C_1 - C_{20} alkyl group substituted with —F, a C_1 - C_{20} alkyl group substituted with —Cl, a C_1 - C_{20} alkyl group substituted with —Br, and a C_1 - C_{20} alkyl group substituted with —I.


When the organic light-emitting device **10** is a full-color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In an implementation, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers contact each other or are separated from each other. In an implementation, the emission layer may include two or more materials selected from a red-light emission material, a green-light emission material, and a blue-light emission material, in which the two or more materials are mixed with each other in a single layer to emit white light.

The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant and a fluorescent dopant.

An amount of the dopant in the emission layer may be, in general, in a range of about 0.01 to about 15 parts by weight based on 100 parts by weight of the host.

A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

In an implementation, the host may further include, the heterocyclic compound represented by Formula 1, a compound represented by Formula 301 below:



In Formula 301,

Ar_{301} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

$xb11$ may be 1, 2, or 3,

L_{301} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

$xb1$ may be an integer selected from 0 to 5,

R_{301} may be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_{301})(Q_{302})(Q_{303}), —N(Q_{301})(Q_{302}), —B(Q_{301})(Q_{302}), —C(=O)(Q_{301}), —S(=O)₂(Q_{301}), and —P(=O)(Q_{301})(Q_{302}),

$xb21$ may be an integer selected from 1 to 5,

Q_{301} to Q_{303} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In an implementation, Ar_{301} in Formula 301 may be selected from:

a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group; and


a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene

group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),


wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In Formula 301, when xb11 is two or more, two or more Ar₃₀₁(s) may be linked via a single bond.

In one or more embodiments, the compound represented by Formula 301 may be represented by Formula 301-1 or 301-2:



<Formula 301-1>



<Formula 301-2>

wherein, in Formulae 301-1 to 301-2,

A₃₀₁ to A₃₀₄ may each independently be selected from a benzene, a naphthalene, a phenanthrene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a pyridine, a pyrimidine, an indene, a fluorene, a spiro-bifluorene, a benzofluorene, a dibenzofluorene, an indole, a carbazole, benzocarbazole, a dibenzocarbazole, a furan, a benzofuran, a dibenzofuran, a naphthofuran, a benzonaphthofuran, a dinaphthofuran, a thiophene, a benzothiophene, a dibenzothiophene, a naphthothiophene, benzonaphthothiophene, and a dinaphthothiophene,

X₃₀₁ may be O, S, or N-[(L₃₀₄)xb4-R₃₀₄],

R₃₁₁ to R₃₁₄ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

xb22 and xb23 may each independently be 0, 1, or 2,

L₃₀₁, xb1, R₃₀₁, and Q₃₁ to Q₃₃ are the same as described above,

L₃₀₂ to L₃₀₄ are each independently the same as described in connection with L₃₀₁,

xb2 to xb4 are each independently the same as described in connection with xb1, and

R₃₀₂ to R₃₀₄ are each independently the same as described in connection with R₃₀₁.

For example, L₃₀₁ to L₃₀₄ in Formulae 301, 301-1, and 301-2 may each independently be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group,

an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazininylenylene group, a quinolininylenylene group, an isoquinolininylenylene group, a benzoquinolininylenylene group, a phthalazininylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolininylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarba-

zolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, a pyridinylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a thiadiazolylene group, an oxadiazolylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolyene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

wherein Q₃₁ to Q₃₃ are the same as described above.

In one or more embodiments, R₃₀₁ to R₃₀₄ in Formulas 301, 301-1, and 301-2 may each independently be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinoliny group, an

isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and


a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

wherein Q₃₁ to Q₃₃ are the same as described above.

95


In an implementation, the host may include an alkaline-earth metal complex. For example, the host may be selected from a Be complex (for example, Compound H55), an Mg complex, and a Zn complex.

The host may include, in addition to the heterocyclic compound represented by Formula 1 at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalene-2-yl)anthracene (MADN), 9,10-di-(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55:




96

-continued




97

-continued



5


10



15


20

25



30


35



40

45

50




55

60


65

98


-continued




H17




H18



H19




H20



H21

99

-continued



H22

100

-continued

5

H25

10

15

20

H23

25

30

35

40

45





H24

50




55

60

65



101
-continued



102
-continued

H27

5

H30

10

15

20

25

H28

30

35

40

45






50

H29

55

60

65




103
-continued


104
-continued

H35

H39




5



10


H36



15


20

H40



25


30



H37


35

40



45


H41



50


55

H38







60

65



105
-continued




106
-continued

H42

5

10

15



H46


H43

20

25

30

35




H47

H48

H44

40

45



H49


H45

50

55


60

65







H50

H51




107

-continued



The phosphorescent dopant may include an organometallic complex represented by Formula 401:




<Formula 401>

108

-continued

<Formula 402>



15

In Formulae 401 and 402,

M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

L_{401} may be selected from ligands represented by Formula 402, and xc1 may be 1, 2, or 3, wherein, when xc1 is two or more, two or more $L_{401}(s)$ may be identical to or different from each other,

L_{402} may be an organic ligand, and xc2 may be an integer selected from 0 to 4, wherein, when xc2 is two or more, two or more $L_{402}(s)$ may be identical to or different from each other,

X_{401} to X_{404} may each independently be nitrogen or carbon,

X_{401} and X_{403} may be linked via a single bond or a double bond, and X_{402} and X_{404} may be linked via a single bond or a double bond,

A_{401} and A_{402} may each independently be a C_5 - C_{60} cyclic group or a C_1 - C_{60} heterocyclic group,

X_{405} may be a single bond, $*-O-*$, $*-S-*$, $*-C(=O)-*$, $*-N(Q_{411})-*$, $*-C(Q_{411})(Q_{412})-*$, $*-C(Q_{411})=C(Q_{412})-*$, $*-C(Q_{411})=*$, or $*=C(Q_{411})=*$, wherein Q_{411} and Q_{412} may be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

X_{406} may be a single bond, O, or S,

R_{401} and R_{402} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_1 - C_{20} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group

$-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})(Q_{402})$, $-B(Q_{401})(Q_{402})$, $-C(=O)(Q_{401})$, $-S(=O)_2(Q_{401})$, and $-P(=O)(Q_{401})(Q_{402})$; wherein Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a C_6 - C_{20} aryl group, and a C_1 - C_{20} heteroaryl group,

xc11 and xc12 may each independently be an integer selected from 0 to 10, and

* and *' in Formula 402 each indicate a binding site to M in Formula 401.

In one or more embodiments, A_{401} and A_{402} in Formula 402 may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

In one or more embodiments, in Formula 402, i) X_{401} may be nitrogen and X_{402} may be carbon, or ii) X_{401} and X_{402} may each be nitrogen at the same time.

In one or more embodiments, R_{401} and R_{402} in Formula 402 may each independently be selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuran group, and a dibenzothiophenyl group;

a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuran group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuran group, and a dibenzothiophenyl group; and

—Si(Q_{401})(Q_{402})(Q_{403}), —N(Q_{401})(Q_{402}), —B(Q_{401})(Q_{402}), —C(=O)(Q_{401}), —S(=O)₂(Q_{401}), and —P(=O)(Q_{401})(Q_{402}),


wherein Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group.

In one or more embodiments, when xc1 in Formula 401 is two or more, two A_{401} (s) in two or more L_{401} (s) may optionally be linked via X_{407} , which is a linking group, or two A_{402} (s) in two or more L_{402} (s) may optionally be linked via X_{408} , which is a linking group (see Compounds PD1 to PD4 and PD7). X_{407} and X_{408} may each independently be a single bond, *—O—*, *—S—*, *—C(=O)—*, *—N(Q_{413})—*, *—C(Q_{413})(Q_{414})—*, or *—C(Q_{413})=C(Q_{414})—* (wherein Q_{413} and Q_{414} may each independently be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group).


L_{402} in Formula 401 may be a monovalent, divalent, or trivalent organic ligand. For example, L_{402} may be selected from a halogen, diketone (for example, acetylacetonate), a carboxylic acid (for example, picolinate), —C(=O), an isonitrile, —CN, and phosphorus (for example, phosphine or phosphite).

In one or more embodiments, the phosphorescent dopant may be selected from, for example, Compounds PD1 to PD25:


PD1









PD2



PD3




111
-continued



112
-continued

PD4

5




10

PD5

20

25



PD6

30

35

PD7


40

45

PD8

50


55



PD9

60

65



PD10






PD11

PD12

PD13


PD14

113
-continued




114
-continued


PD15




PD16




PD17



PD18



PD19



PD20


PD21

PD22


PD23

115

-continued




PD24



PD25

The fluorescent dopant may include an arylamine compound or a styrylamine compound.

The fluorescent dopant may include a compound represented by Formula 501:



<Formula 501> 40

In Formula 501,

Ar_{501} may be a substituted or unsubstituted $\text{C}_5\text{-C}_{60}$ carbocyclic group or a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heterocyclic group,

L_{501} to L_{503} may each independently be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkylene group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenylene group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

$\text{xd}1$ to $\text{xd}3$ may each independently be an integer selected from 0 to 3,

R_{501} and R_{502} may each independently be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a

116

substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryloxy group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylthio group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

$\text{xd}4$ may be an integer selected from 1 to 6.

10 In one or more embodiments, Ar_{501} in Formula 501 may be selected from:

a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group; and

20 a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{20}$ alkyl group, a $\text{C}_1\text{-C}_{20}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, L_{501} to L_{503} in Formula 501 may each independently be selected from:

35 a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylene group, a carbazolyene group, an indolyene group, an isoindolyene group, a benzofuranylene group, a benzothiophenylenylene group, a dibenzofuranylene group, a dibenzothiophenylenylene group, a benzocarbazolyene group, a dibenzocarbazolyene group, a dibenzosilolylene group, and a pyridinylenylene group; and

50 a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylene group, a carbazolyene group, an indolyene group, an isoindolyene group, a benzofuranylene group, a benzothiophenylenylene group, a dibenzofuranylene group, a dibenzothiophenylenylene group, a benzocarbazolyene group, a dibenzocarbazolyene group, a dibenzosilolylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{20}$ alkyl group, a $\text{C}_1\text{-C}_{20}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenan-

117

threnyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

In one or more embodiments, R_{501} and R_{501} in Formula 502 may each independently be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and


a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, and —Si(Q_{31})(Q_{32})(Q_{33}),

wherein, in Formula 501, Q_{31} to Q_{33} may be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.


In one or more embodiments, xd4 in Formula 501 may be 3.

For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:


118



FD1




FD2



FD3

119
-continued



FD4

5


10

15

20

25

120
-continued



FD7


FD5

30

35

40

45



FD8


FD6

50


55

60

65











FD9




121
-continued


122
-continued




123
-continued




124
-continued



In one or more embodiments, the fluorescent dopant may be selected from the following compounds.




DPVBi




DPAVBi

125




-continued


126




DCM



DCJTb



Coumarin 6



C545T

The electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer.

For example, the electron transport region may have a structure of electron transport layer/electron injection layer, a structure of hole blocking layer/electron transport layer/electron injection layer, a structure of electron control layer/electron transport layer/electron injection layer, or a structure of buffer layer/electron transport layer/electron injection layer, wherein for each structure, constituting layers are sequentially stacked from an emission layer.

The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include a metal-free compound containing at least one π electron-depleted nitrogen-containing ring.

The " π electron-depleted nitrogen-containing ring" indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

40

For example, the " π electron-depleted nitrogen-containing ring" may be i) a 5-membered to 7-membered hetero monocyclic group having at least one $*-N=*$ moiety, ii) a heteropoly cyclic group in which two or more 5-membered to 7-membered hetero monocyclic groups each having at least one $*-N=*$ moiety are condensed with each other, or iii) a heteropoly cyclic group in which at least one of 5-membered to 7-membered hetero monocyclic groups, each having at least one $*-N=*$ moiety, is condensed with at least one C_5 - C_{60} carbocyclic group.

Examples of the π electron-depleted nitrogen-containing ring include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, thiadiazol, an imidazopyridine, an imidazopyrimidine, and an azacarbazole.

In one or more embodiments, the electron transport region may further include, in addition to the heterocyclic compound represented by Formula 1, a compound represented by Formula 601.

In Formula 601,

Ar₆₀₁ may be a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₁-C₆₀ heterocyclic group,

xe11 may be 1, 2, or 3,

L₆₀₁ is selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

xe1 may be an integer selected from 0 to 5,

R₆₀₁ may be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₆₀₁)(Q₆₀₂)(Q₆₀₃), —C(=O)(Q₆₀₁), —S(=O)₂(Q₆₀₁), and —P(=O)(Q₆₀₁)(Q₆₀₂),

Q₆₀₁ to Q₆₀₃ may each independently be a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

xe21 may be an integer selected from 1 to 5.

In one or more embodiments, at least one selected from Ar₆₀₁(s) in the number of xe11 and R₆₀₁(s) in the number of xe21 may include the π electron-depleted nitrogen-containing ring described above.

In one or more embodiments, ring Ar₆₀₁ in Formula 601 may be selected from:

a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthacene group, a perylene group, a perylene group, a pentaphene

group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),


wherein, in Formula 601, Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

When xe11 in Formula 601 is two or more, two or more Ar₆₀₁(s) may be linked via a single bond.

In one or more embodiments, Ar₆₀₁ in Formula 601 may be an anthracene group.

In one or more embodiments, a compound represented by Formula 601 may be represented by Formula 601-1:

<Formula 601-1>



In Formula 601-1,

X₆₁₄ may be N or C(R₆₁₄), X₆₁₅ may be N or C(R₆₁₅), X₆₁₆ may be N or C(R₆₁₆), and at least one selected from X₆₁₄ to X₆₁₆ may be N,

L₆₁₁ to L₆₁₃ may each independently be substantially the same as described in connection with L₆₀₁,

xe611 to xe613 may each independently be substantially the same as described in connection with xe1,

R₆₁₁ to R₆₁₃ may each independently be substantially the same as described in connection with R₆₀₁, and

R₆₁₄ to R₆₁₆ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, L₆₀₁ and L₆₁₁ to L₆₁₃ in Formulae 601 and 601-1 may each independently be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene





131

azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and


—S(=O)₂(Q₆₀₁) and —P(=O)(Q₆₀₁)(Q₆₀₂),

wherein Q₆₀₁ and Q₆₀₂ are the same as described above.





The electron transport region may include at least one compound selected from Compounds ET1 to ET36:

132
-continued


133
-continued



134
-continued



135
-continued



ET10

5


10

15

20

25

136
-continued



ET13



ET11

30

35

40

45



ET14



ET12

50

55




60

65




ET15

137
-continued




138
-continued

ET16




15
lp;2.4p

ET17



ET18




ET19

ET20

ET21

139
-continued



ET22


5

10

15

20

140
-continued




ET25

25

ET23

ET26




30

35


40

45



ET24 50


ET27




55

60

65



141
-continued



ET28


5

10

15

20

ET29



25


30

35

40

45

ET30




50

55


60

65


142
-continued



ET31






ET32



ET33

143


-continued




In one or more embodiments, the electron transport region may include at least one selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq₃, BAlq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ.

144


ET34




15



ET35



35



45

ET36

A thickness of the buffer layer, the hole blocking layer, or the electron control layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.


A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.


The metal-containing material may include at least one selected from alkali metal complex and alkaline earth-metal

complex. The alkali metal complex may include a metal ion selected from an Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion, and the alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, an Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenylan oxazole, a hydroxy phenylthiazole, a hydroxy diphenylan oxadiazole, a hydroxy diphenylthiadiazol, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene.

For example, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.



ET-D1



ET-D2

The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode **190**. The electron injection layer may directly contact the second electrode **190**.

The electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

The electron injection layer may include alkali metal, alkaline earth metal, rare-earth metal, alkali metal compound, alkaline earth-metal compound, rare-earth metal compound, alkali metal complex, alkaline earth-metal complex, rare-earth metal complex, or any combinations thereof.

The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one embodiment, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs.

The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

The rare-earth metal may be selected from Sc, Y, Ce, Yb, Gd, and Tb.

The alkali metal compound, the alkaline earth-metal compound, and the rare-earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodines) of the alkali metal, the alkaline earth-metal and rare-earth metal.

The alkali metal compound may be selected from alkali metal oxides, such as Li₂O, Cs₂O, or K₂O, and alkali metal halides, such as LiF, NaF, CsF, KF, LiI, NaI, CsI, KI, or RbI. In one embodiment, the alkali metal compound may be selected from LiF, Li₂O, NaF, LiI, NaI, CsI, and KI.

The alkaline earth-metal compound may be selected from alkaline earth-metal compounds, such as BaO, SrO, CaO, Ba_xSr_{1-x}O (0<x<1), Ba_xCa_{1-x}O (0<x<1). In one embodiment, the alkaline earth-metal compound may be selected from BaO, SrO, and CaO.

The rare-earth metal compound may be selected from YbF₃, ScF₃, ScO₃, Y₂O₃, Ce₂O₃, GdF₃, and TbF₃. In one embodiment, the rare-earth metal compound may be selected from YbF₃, ScF₃, TbF₃, YbI₃, ScI₃, and TbI₃.

The alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare-earth metal as described above, and a ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may each independently be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenylan oxazole, hydroxy phenylthiazole, hydroxy diphenylan oxadiazole, hydroxy diphenylthiadiazol, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene.

The electron injection layer may consist of alkali metal, alkaline earth metal, rare-earth metal, alkali metal compound, alkaline earth-metal compound, rare-earth metal compound, alkali metal complex, alkaline earth-metal complex, rare-earth metal complex, or any combinations thereof, as described above. In one or more embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, alkali metal, alkaline earth metal, rare-earth metal, alkali metal compound, alkaline earth-metal compound, rare-earth metal compound, alkali metal complex, alkaline earth-metal complex, rare-earth metal complex, or any combinations thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, e.g., about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

The second electrode **190** may be disposed on the organic layer **150** having such a structure. The second electrode **190** may be a cathode which is an electron injection electrode, and in this regard, a material for forming the second electrode **190** may be selected from metal, an alloy, an electrically conductive compound, and a mixture thereof, which have a relatively low work function.

The second electrode **190** may include at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO. The second electrode **190** may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

The second electrode **190** may have a single-layered structure, or a multi-layered structure including two or more layers.

An organic light-emitting device **20** of FIG. **2** may include a first capping layer **210**, a first electrode **110**, an organic

147

layer 150, and a second electrode 190 which are sequentially stacked in this stated order, an organic light-emitting device 30 of FIG. 3 may include a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220 which are sequentially stacked in this stated order, and an organic light-emitting device 40 of FIG. 4 may include a first capping layer 210, a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220.

Regarding FIGS. 2 to 4, the first electrode 110, the organic layer 150, and the second electrode 190 may be understood by referring to the description presented in connection with FIG. 1.

In the organic layer 150 of each of the organic light-emitting devices 20 and 40, light generated in an emission layer may pass through the first electrode 110, which is a semi-transmissive electrode or a transmissive electrode, and the first capping layer 210 toward the outside, and in the organic layer 150 of each of the organic light-emitting devices 30 and 40, light generated in an emission layer may pass through the second electrode 190, which is a semi-transmissive electrode or a transmissive electrode, and the second capping layer 220 toward the outside.


The first capping layer 210 and the second capping layer 220 may increase external luminescent efficiency according to the principle of constructive interference.

The first capping layer 210 and the second capping layer 220 may each independently be an organic capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.

At least one selected from the first capping layer 210 and the second capping layer 220 may each independently include at least one material selected from carbocyclic compounds, heterocyclic compounds, amine-based compounds, porphyrine derivatives, phthalocyanine derivatives, naphthalocyanine derivatives, alkali metal complexes, and alkaline earth-based complexes. The carbocyclic compound, the heterocyclic compound, and the amine-based compound may be optionally substituted with a substituent containing at least one element selected from O, N, S, Se, Si, F, Cl, Br, and I. In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include an amine-based compound.


In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include the compound represented by Formula 201 or the compound represented by Formula 202.

In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include a compound selected from Compounds HT28 to HT33 and Compounds CP1 to CP5.



148

-continued



Hereinbefore, the organic light-emitting device according to an embodiment has been described in connection with FIGS. 1 through 4.

Layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region may be formed in a certain region by using one or more suitable methods selected from vacuum deposition, spin coating, casting, langmuir-blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of about 100 to about 500° C., at a vacuum degree of about 10⁻⁸ to about 10⁻³ torr, and at a deposition rate of about 0.01 to about 100 Å/sec by taking into account a compound to be included in a layer to be formed and the structure of a layer to be formed.

When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to 200° C. by taking into account a compound to be included in a layer to be formed and the structure of a layer to be formed.

The term "C₁-C₆₀ alkyl group" as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term "C₁-C₆₀ alkylene group" as used herein refers to a divalent group having the same structure as the C₁-C₆₀ alkyl group.

The term "C₂-C₆₀ alkenyl group" as used herein refers to a hydrocarbon group having at least one carbon-carbon double bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term "C₂-C₆₀ alkenylene group" as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkenyl group.

The term "C₂-C₆₀ alkynyl group" as used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethynyl group and a propynyl group. The term "C₂-C₆₀ alkynylene group" as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

The term "C₁-C₆₀ alkoxy group" as used herein refers to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is the C₁-C₆₀ alkyl group), and examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

The term "C₃-C₁₀ cycloalkyl group" as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term "C₃-C₁₀ cycloalkylene group" as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

The term "C₁-C₁₀ heterocycloalkyl group" as used herein refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuranlyl group, and a tetrahydrothiophenyl group. The term "C₁-C₁₀ heterocycloalkylene group," used herein, refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

The term "C₃-C₁₀ cycloalkenyl group" as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and does not have aromaticity, and examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a

cycloheptenyl group. The term "C₃-C₁₀ cycloalkenylene group" used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

The term "C₁-C₁₀ heterocycloalkenyl group" as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Examples of the C₁-C₁₀ heterocycloalkenyl group are a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranlyl group and a 2,3-dihydrothiophenyl group. The term "C₁-C₁₀ heterocycloalkenylene group," used herein, refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkenyl group.

The term "C₆-C₆₀ aryl group" as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C₆-C₆₀ arylene group used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Examples of the C₆-C₆₀ aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be fused to each other.

The term "C₁-C₆₀ heteroaryl group" as used herein refers to a monovalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. A C₁-C₆₀ heteroarylene group used herein refers to a divalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Examples of the C₁-C₆₀ heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, and an isoquinolyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be fused to each other.

The term "C₆-C₆₀ aryloxy group" as used herein indicates —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and a C₆-C₆₀ arylthio group indicates —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

The term "monovalent non-aromatic condensed polycyclic group" as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed with each other, only carbon atoms as a ring-forming atom, and non-aromaticity in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term "divalent non-aromatic condensed polycyclic group," used herein, refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

The term "monovalent non-aromatic condensed heteropolycyclic group" as used herein refers to a monovalent group (for example, having 1 to 60 carbon atoms) that has two or more rings condensed to each other, has at least one heteroatom selected from N, O, Si, P, and S, other than carbon atoms, as a ring-forming atom, and has non-aromaticity in the entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term "divalent non-aromatic condensed heteropolycyclic group," used herein, refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

The term "C₅-C₆₀ carbocyclic group" as used herein refers to a monocyclic or polycyclic group having 5 to 60

carbon atoms in which a ring-forming atom is a carbon atom only. The C₅-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₅-C₆₀ carbocyclic group may be a ring, such as a benzene group, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In one or more embodiments, depending on the number of substituents connected to the C₅-C₆₀ carbocyclic group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

The term "C₁-C₆₀ heterocyclic group" used herein refers to a group having the same structure as the C₅-C₆₀ carbocyclic group, except that as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used in addition to carbon (the number of carbon atoms may be in a range of 1 to 60).

At least one substituent of the substituted C₅-C₆₀ carbocyclic group, the substituted C₁-C₆₀ heterocyclic group, the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

deuterium (—D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and

a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

—Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂).

Unless otherwise defined herein, Q₁ to Q₆, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

The term "Ph" used herein refers to a phenyl group, the term "Me" used herein refers to a methyl group, the term "Et" used herein refers to an ethyl group, the term "ter-Bu" or "Bu^t" used herein refers to a tert-butyl, and the term "OMe" used herein refers to a methoxy group.

The term "biphenyl group" as used herein refers to "a phenyl group substituted with a phenyl group." In other words, the "biphenyl group" is a substituted phenyl group having a C₆-C₆₀ aryl group as a substituent.

The term "terphenyl group" as used herein refers to "a phenyl group substituted with a biphenyl group." In other words, the "terphenyl group" is a substituted phenyl group having, as a substituent, a C₆-C₆₀ aryl group substituted with a C₆-C₆₀ aryl group.

* and *^t used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.

Hereinafter, a compound according to embodiments and an organic light-emitting device according to embodiments will be described in detail with reference to Synthesis Examples and Examples. The wording "B was used instead of A" used in describing Synthesis Examples refers to that an identical molar equivalent of B was used in place of A.

The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples.


153

EXAMPLE

Synthesis Example

Synthesis Example 1

Synthesis of Compound 1




Synthesis of Compound 1

1.34 g (10 mmol) of 2-aminobenzofuran, 1.83 g (10 mmol) of 2-coumaranone, 3.32 g (10 mmol) of 10-(naphthalene-1-yl)anthracene-9-carbaldehyde, and 0.013 g (0.5 mmol) of diiodine were dissolved in 150 mL of THF and were then stirred at a temperature of 80° C. for 24 hours. The reacting solution was cooled to ambient temperature. Then, an organic layer was extracted three times by using 60 mL of water and 60 mL of diethyl ether. The organic layer was dried by using magnesium sulfate and the solvent was evaporated. The obtained residue was separated and purified by silica gel column chromatography, thereby completing the preparation of 3.67 g (60%) of Compound 1. Compound 1 was confirmed through LC-MS and ¹H NMR. C₄₅H₂₅N₂O₂; M+1 611.72


Synthesis Example 2

Synthesis of Compound 20



154

-continued





Synthesis of Intermediate 20-1

3 g (65%) of an Intermediate Compound 20-1 was prepared by following the same steps as in Synthesis Example 1, except that 3-bromobenzaldehyde was used instead of 10-(naphthalene-1-yl)anthracene-9-carbaldehyde. Intermediate Compound 20-1 was confirmed through LC-MS. C₂₇H₁₄BrNO₂; M+1 464.31


Synthesis of Compound 20

4.64 g (10 mmol) of Intermediate Compound 20-1, 3.45 g (10 mmol) of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile, 0.58 g (0.5 mmol) of Pd(PPh₃)₄, and 4.15 g (30.0 mmol) of K₂CO₃ were dissolved in 150 mL of a mixed solution of THF/H₂O (2/1) and were then stirred at a temperature of 80° C. for 16 hours. The reacting solution was cooled to ambient temperature. Then, an organic layer was extracted three times by using 60 mL of water and 60 mL of diethyl ether. The organic layer was dried by using magnesium sulfate and the solvent was evaporated. The obtained residue was separated and purified by silica gel column chromatography, thereby completing the preparation of 4.2 g (70%) of Compound 20. Compound 20 was confirmed through LC-MS and ¹H NMR. C₄₃H₂₆N₂O₂; M+1 602.71

Synthesis of Compound 23



23-1



compound 23

Synthesis of Intermediate 23-1

2.78 g (60%) of Intermediate Compound 23-1 was prepared in the same manner as in the synthesis of Intermediate 20-1, except that 4-bromobenzaldehyde was used instead of 3-bromobenzaldehyde. Intermediate Compound 23-1 was confirmed through LC-MS. $C_{27}H_{14}BrNO_2$: M+1 464.30


Synthesis of Compound 23

3.27 g (50%) of Compound 23 was prepared in the same manner as in the synthesis of Compound 20, except that Intermediate Compound 23-1 was used instead of Intermediate Compound 20-1, and 4-(9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-yl)pyridine was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-

carbonitrile. Compound 23 was confirmed through LC-MS and 1H NMR. $C_{47}H_{30}N_2O_2$: M+1 654.75


Synthesis Example 4

Synthesis of Compound 26




157


-continued

**158**


-continued



compound 26



20-1





Synthesis of Compound 26


4.93 g (80%) of Compound 26 was prepared in the same manner as in the synthesis of Compound 20, except that 2,4-diphenyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,5-triazine was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. Compound 26 was confirmed through LC-MS and ¹H NMR. C₄₂H₂₄N₄O₂; M+1 616.66

Synthesis Example 5

Synthesis of Compound 64




64-1



159

160

-continued



compound 64

Synthesis of Intermediate 64-1

1.4 g (30%) of Intermediate Compound 64-1 was prepared in the same manner as in the synthesis of Intermediate 20-1, except that dibenzo[b,d]furan-2-amine was used instead of dibenzo[b,d]furan-4-amine, and 4-bromobenzaldehyde was used instead of 3-bromobenzaldehyde. Intermediate Compound 64-1 was confirmed through LC-MS. $C_{27}H_{14}BrNO_2$: M+1 464.33


Synthesis of Compound 64

4.3 g (70%) of Compound 64 was prepared in the same manner as in the synthesis of Compound 20, except that Intermediate Compound 64-1 was used instead of Interme-

20 diate Compound 20-1, and 3,3'-(5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3-phenylene)dipyridine was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. Compound 64 was confirmed through LC-MS and 1H NMR. $C_{43}H_{25}N_3O_2$: M+1 616.68

Synthesis Example 6

Synthesis of Compound 94



compound 94

161

Synthesis of Intermediate 94-1


1.2 g (25%) of Intermediate Compound 94-1 was prepared in the same manner as in the synthesis of Intermediate 20-1, except that dibenzo[b,d]furan-2-amine was used instead of dibenzo[b,d]furan-4-amine, and 4-bromobenzaldehyde was used instead of 3-bromobenzaldehyde. Intermediate Compound 94-1 was confirmed through LC-MS. $C_{27}H_{14}BrNO_2$: M+1 464.32

Synthesis of Compound 94

3.91 g (60%) of Compound 94 was prepared in the same manner as in the synthesis of Compound 20, except that Intermediate Compound 94-1 was used instead of Intermediate Compound 20-1, and 9-phenyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-carbazole-2-carbonitrile was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. Compound 94 was confirmed through LC-MS and 1H NMR. $C_{46}H_{27}N_3O_2$: M+1 653.73

Synthesis Example 7

Synthesis of Compound 42



Synthesis of Compound 42


4.64 g (10 mmol) of Intermediate 20-1 was dissolved in 100 mL of THF, and 4 mL (2.5 M in hexane) of n-butyllithium was added thereto at a temperature of $-78^\circ C$. 2.20 g (10 mmol) of chlorodiphenylphosphine was slowly added dropwise after 1 hour. Then, the solution was stirred for 3 hours and then heated to ambient temperature. Then, water was added thereto and cleaning was performed three times by using ethyl acetate (30 mL). A cleaned ethyl acetate layer was dried with $MgSO_4$ and evaporated under reduced pressure, and the resultant was dissolved in 40 mL of dichloromethane. 4 mL of peroxide was added to the obtained

162

solution and then stirred at ambient temperature for 20 hours. 20 mL of water was added thereto. Then, an organic layer was extracted three times by using 20 mL of dichloromethane. The organic layer was dried by using magnesium sulfate and the solvent was evaporated. The obtained residue was separated and purified by silica gel column chromatography, thereby completing the preparation of 4.1 g (70%) of Compound 42. Compound 42 was confirmed through LC-MS and 1H NMR. $C_{39}H_{24}NO_3P$: M+1 585.58

Synthesis Example 8

Synthesis of Compound 114



Synthesis of Intermediate 114-1

3.3 g (70%) of Intermediate Compound 114-1 was prepared in the same manner as in the synthesis of Intermediate 20-1, except that dibenzo[b,d]thiophen-4-amine was used instead of dibenzo[b,d]furan-4-amine, and 4-bromoaldehyde was used instead of 3-bromoaldehyde. Intermediate Compound 114-1 was confirmed through LC-MS. $C_{27}H_{14}BrNOS$: M+1 480.40

Synthesis of Compound 114

4.2 g (70%) of Compound 114 was prepared in the same manner as in the synthesis of Compound 42, except that Intermediate Compound 114-1 was used instead of Intermediate Compound 20-1. Compound 114 was confirmed through LC-MS and 1H NMR. $C_{39}H_{24}NO_2PS$: M+1 601.67

TABLE 1

Compound	¹ H NMR (CDCl ₃ , 400 MHz)	MS/FAB	
		found	calc.
1	8.75-8.70 (m, 2H), 8.07-8.04 (m, 2H), 7.84-7.75 (m, 3H), 7.71-7.65 (m, 4H), 7.60-7.57 (m, 2H), 7.54-7.50 (m, 4H), 7.47-7.40 (m, 4H), 7.35-7.34 (m, 1H), 7.15-7.13 (m, 2H), 6.97-6.95 (m, 1H)	611.72	611.7
20	8.81-8.79 (m, 1H), 8.72-8.70 (m, 1H), 8.42-8.40 (m, 1H), 8.07-8.00 (m, 3H), 7.89-7.80 (m, 2H), 7.76-7.72 (m, 2H), 7.67-7.63 (m, 3H), 7.60-7.58 (m, 2H), 7.52-7.50 (m, 2H), 7.46-7.45 (m, 2H), 7.40-7.39 (m, 1H), 1.61-1.57 (m, 6H)	602.71	602.69
23	8.82-8.79 (m, 1H), 8.72-8.70 (m, 1H), 8.66-8.64 (m, 2H), 8.22-8.20 (m, 2H), 8.06-8.02 (m, 2H), 7.78-7.72 (m, 6H), 7.65-7.57 (m, 3H), 7.55-7.50 (m, 2H), 7.47-7.38 (m, 4H), 7.31-7.29 (m, 1H), 1.57-1.54 (m, 6H)	654.75	654.77
26	9.16-9.13 (m, 1H), 8.82-8.79 (m, 5H), 8.76-8.70 (m, 2H), 8.23-8.03 (m, 3H), 7.76-7.67 (m, 3H), 7.62-7.56 (m, 6H), 7.52-7.40 (m, 4H)	616.66	616.68
64	8.95-8.90 (m, 3H), 8.82-8.65 (m, 3H), 8.23-8.02 (m, 6H), 7.89-7.81 (m, 6H), 7.74-7.58 (m, 2H), 7.56-7.44 (m, 5H)	616.68	615.69
94	8.25-8.17 (m, 3H), 8.08-8.02 (m, 3H), 7.91-7.82 (m, 3H), 7.74-7.71 (m, 2H), 7.63-7.60 (m, 3H), 7.57-7.50 (m, 5H), 7.32-7.19 (m, 2H), 6.05-5.95 (m, 2H), 5.85-5.82 (m, 1H)	653.73	653.74
42	5.70-5.67 (m, 1H), 5.35-5.31 (m, 1H), 4.89-4.91 (m, 1H), 8.81-8.70 (m, 2H), 8.52-8.50 (m, 1H), 8.08-8.04 (m, 2H), 7.91-7.90 (m, 1H), 7.77-7.72 (m, 3H), 7.66-7.57 (m, 7H), 7.53-7.44 (m, 4H), 7.45-7.41 (m, 4H)	585.58	585.6
114	8.71-8.68 (m, 2H), 8.52-8.50 (m, 1H), 8.05-8.00 (m, 4H), 7.91-7.90 (m, 2H), 7.74-7.67 (m, 5H), 7.66-7.57 (m, 2H), 7.53-7.48 (m, 4H), 7.45-7.41 (m, 4H)	601.67	601.66

Example 1

As a substrate and an anode, a Corning 15 Ω/cm² (1,200 Å) ITO glass substrate was cut to a size of 50 mm×50 mm×0.7 mm, and then, sonicated with isopropyl alcohol and pure water each for 5 minutes, and then washed by irradiation of ultraviolet ray for 30 minutes and ozone, and the resultant glass substrate was provided to a vacuum deposition apparatus.

2-TNATA was vacuum-deposited on the ITO glass substrate (anode) to form a hole injection layer having a thickness of 600 Å, and 4,4'-bis[N-(1-naphthyl)-N-phenylamino]biphenyl (NPB) was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å.

9,10-di-naphthalene-2-yl-anthracene (ADN) (blue fluorescent host) and 4,4'-bis[2-(4-(N,N-diphenylamino)phenyl)vinyl]biphenyl (DPAVB) (blue fluorescent dopant) were co-deposited on the hole transport layer at a weight ratio of 98:2 to form an emission layer having a thickness of 300 Å.

Then, Compound 1 was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å. LiF was deposited on the electron transport layer to form

an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited on the electron injection layer to form an LiF/Al cathode having a thickness of 3,000 Å, thereby completing the manufacture of an organic light-emitting device.

Examples 2 to 8 and Comparative Example 1

Organic light-emitting devices of Examples 2 to 8 and Comparative Example 1 were manufactured in the same manner as in Example 1, except that Compounds shown in Table 2, below, were each used instead of Compound 1 to form an electron transport layer.

Evaluation Example 1


The driving voltage, current density, luminance, efficiency, emission color, and half lifespan of the organic light-emitting devices manufactured in Examples 1 to 8 and Comparative Example 1 were evaluated using Keithley SMU 236 and PR650 luminance meter. Results thereof are shown in Table 2. The half lifespan was obtained by measuring a time period that has lapsed until the luminance (@100 mA/cm²) was reduced to 50% of the initial luminance after the driving of the organic light-emitting device.

TABLE 2


	Electron transport material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
Example 1	Compound 1	4.75	50	2880	5.76	Blue	3115 hr
Example 2	Compound 20	4.91	50	3104	6.208	Blue	302 hr
Example 3	Compound 23	4.89	50	3088	6.176	Blue	335 hr
Example 4	Compound 26	5.13	50	2963	5.926	Blue	298 hr
Example 5	Compound 64	4.97	50	2870	5.74	Blue	301 hr
Example 6	Compound 94	5.45	50	2894	5.788	Blue	335 hr

TABLE 2-continued


	Electron transport material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
Example 7	Compound 42	5.66	50	2667	5.334	Blue	360 hr
Example 8	Compound 114	5.49	50	2695	5.39	Blue	349 hr
Comparative Example 1	Alq ₃	7.12	50	2039	4.078	Blue	162 hr



1




20




23

TABLE 2-continued


Electron transport material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
-----------------------------	---------------------	---------------------------------------	--------------------------------	-------------------	----------------	--




26



42





64



94

TABLE 2-continued

Electron transport material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
						
114						
						
Alq ₃						

Referring to Table 2, it may be seen that the organic light-emitting devices of Examples 1 to 8 had excellent efficiency and lifespan characteristics, compared to Comparative Example 1.

According to one or more embodiments, the organic light-emitting device including the heterocyclic compound may have high efficiency and a long lifespan.

Example embodiments have been disclosed herein, and although specific terms are employed, they are used and are to be interpreted in a generic and descriptive sense only and not for purpose of limitation. In some instances, as would be apparent to one of ordinary skill in the art as of the filing of the present application, features, characteristics, and/or elements described in connection with a particular embodiment may be used singly or in combination with features, characteristics, and/or elements described in connection with other embodiments unless otherwise specifically indicated. Accordingly, it will be understood by those of skill in the art that various changes in form and details may be made without departing from the spirit and scope of the present invention as set forth in the following claims.

What is claimed is:

1. A heterocyclic compound represented by Formula 1:





wherein, in Formula 1,

A₁ is a group represented by Formula 2,

A₂ is a group represented by Formula 3, and

a21 is an integer of 1 to 11, wherein, when a21 is 2 to 11, two or more of A₂ are identical to or different from each other:



wherein F₁ in Formula 2 is a dibenzofuran moiety or a dibenzothiophene moiety,

R₁ to R₅ and R₁₂ in Formula 2 are each independently selected from a binding site to A₂, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubsti-

tuted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_1)(\text{Q}_2)(\text{Q}_3)$, and $-\text{S}(=\text{O})_2(\text{Q}_1)$, b6 in Formula 2 is an integer of 0 to 6, wherein, when b6 is 2 to 6, two or more of R_6 are identical to or different from each other,

X_1 in Formula 2 is O or S,

at least one of R_1 to R_5 and R_{12} is a binding site to A_2 , a number of R_1 to R_5 and R_{12} being the binding site to A_2 being the same as a21,

L in Formula 3 is selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkylene group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenylene group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryloxy group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

n in Formula 3 is an integer of 0 to 7, wherein, when n is 2 to 7, two or more of L are identical to or different from each other,

Ar in Formula 3 is selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryloxy group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylthio group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_4)(\text{Q}_5)(\text{Q}_6)$, $-\text{C}(=\text{O})(\text{Q}_4)$, $-\text{S}(=\text{O})_2(\text{Q}_4)$, and $-\text{P}(=\text{O})(\text{Q}_4)(\text{Q}_5)$,

m in Formula 3 is an integer of 1 to 7, wherein, when m is 2 to 7, two or more of Ar are identical to or different from each other, and

at least one substituent of the substituted $\text{C}_3\text{-C}_{10}$ cycloalkylene group, the substituted $\text{C}_1\text{-C}_{10}$ heterocycloalkylene group, the substituted $\text{C}_3\text{-C}_{10}$ cycloalkenylene group, the substituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenylene group, the substituted $\text{C}_6\text{-C}_{60}$ arylene group, the substituted $\text{C}_1\text{-C}_{60}$ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted $\text{C}_1\text{-C}_{60}$ alkyl group, the substituted $\text{C}_2\text{-C}_{10}$ alkenyl group, the substituted $\text{C}_2\text{-C}_{60}$ alkynyl group, the substituted $\text{C}_1\text{-C}_{60}$ alkoxy group, the substituted $\text{C}_3\text{-C}_{10}$ cycloalkyl group, the substituted $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, the substituted $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, the substituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, the substituted $\text{C}_6\text{-C}_{60}$ aryl group, the substituted $\text{C}_6\text{-C}_{60}$ aryloxy group, the substituted $\text{C}_6\text{-C}_{60}$ arylthio group, the substituted $\text{C}_1\text{-C}_{60}$ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ alkyl group, a substituted or unsubstituted $\text{C}_2\text{-C}_{60}$ alkenyl group, a substituted or

unsubstituted $\text{C}_2\text{-C}_{60}$ alkynyl group, and a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ alkoxy group;

a $\text{C}_1\text{-C}_{60}$ alkyl group, a $\text{C}_2\text{-C}_{60}$ alkenyl group, a $\text{C}_2\text{-C}_{60}$ alkynyl group, and a $\text{C}_1\text{-C}_{60}$ alkoxy group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_{11})(\text{Q}_{12})(\text{Q}_{13})$, $-\text{N}(\text{Q}_{11})(\text{Q}_{12})$, $-\text{B}(\text{Q}_{11})(\text{Q}_{12})$, $-\text{C}(=\text{O})(\text{Q}_{11})$, $-\text{S}(=\text{O})_2(\text{Q}_{11})$, and $-\text{P}(=\text{O})(\text{Q}_{11})(\text{Q}_{12})$;

a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_1\text{-C}_{60}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{60}$ alkyl group, a $\text{C}_2\text{-C}_{60}$ alkenyl group, a $\text{C}_2\text{-C}_{60}$ alkynyl group, a $\text{C}_1\text{-C}_{60}$ alkoxy group, a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_{21})(\text{Q}_{22})(\text{Q}_{23})$, $-\text{N}(\text{Q}_{21})(\text{Q}_{22})$, $-\text{B}(\text{Q}_{21})(\text{Q}_{22})$, $-\text{C}(=\text{O})(\text{Q}_{21})$, $-\text{S}(=\text{O})_2(\text{Q}_{21})$, and $-\text{P}(=\text{O})(\text{Q}_{21})(\text{Q}_{22})$; and

$-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,

wherein Q_1 to Q_6 , Q_{11} to Q_{13} , Q_{21} to Q_{23} , and Q_{31} to Q_{33} are each independently selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{60}$ alkyl group, a $\text{C}_2\text{-C}_{60}$ alkenyl group, a $\text{C}_2\text{-C}_{60}$ alkynyl group, a $\text{C}_1\text{-C}_{60}$ alkoxy group, a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group.

2. The heterocyclic compound as claimed in claim 1, wherein:

X_1 is O, and

F_1 is a dibenzofuran moiety.

3. The heterocyclic compound as claimed in claim 1, wherein a21 is an integer of 1 to 6.

4. The heterocyclic compound as claimed in claim 1, wherein L in Formula 3 is selected from:


a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthroquinolylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthroquinolylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl


group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranlyl group, a benzothiophenyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthroquinolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃),

wherein Q₃₁ to Q₃₃ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group.


5. The heterocyclic compound as claimed in claim 1, wherein L in Formula 3 is a group represented by one of the following Formulae 5-1 to 5-34:



Formula 5-1










Formula 5-2




Formula 5-3

175
-continued




176
-continued


Formula 5-4





Formula 5-5





Formula 5-6





Formula 5-7




Formula 5-8



Formula 5-9



Formula 5-10



Formula 5-11

Formula 5-12

Formula 5-13

Formula 5-14

Formula 5-15

Formula 5-16










Formula 5-17

Formula 5-18

Formula 5-19


Formula 5-20

177
-continued




178
-continued


Formula 5-21




Formula 5-22




Formula 5-23




Formula 5-24




Formula 5-25




Formula 5-26




Formula 5-27



Formula 5-28



Formula 5-29



Formula 5-30

Formula 5-31

Formula 5-32

Formula 5-33

Formula 5-34

wherein, in Formulae 5-1 to 5-34,
 Y_1 is O, S, $C(Z_3)(Z_4)$, $N(Z_5)$, or $Si(Z_6)(Z_7)$,
 Z_1 to Z_7 are each independently selected from deuterium,
 $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group,
a nitro group, an amidino group, a hydrazino group,
a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20}
alkoxy group, a cyclopentyl group, a cyclohexyl group,
a cycloheptyl group, a cyclopentenyl group, a cyclo-
hexenyl group, a phenyl group, a biphenyl group, a
terphenyl group, a pentalenyl group, an indenyl group,
a naphthyl group, an azulenyl group, an indacenyl
group, an acenaphthyl group, a fluorenyl group, a
spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl
group, a benzofluorenyl group, a dibenzofluorenyl

group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazolinyl group, a benzoquinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indenocarbazolyl group, an indolocarbazolyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, in which Q_{31} to Q_{33} are each independently selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group,

d2 is an integer of 0 to 2,

d3 is an integer of 0 to 3,

d4 is an integer of 0 to 4,

d5 is an integer of 0 to 5,

d6 is an integer of 0 to 6,

d8 is an integer of 0 to 8, and

* and *' each indicate a binding site to a neighboring atom.

6. The heterocyclic compound as claimed in claim 5, wherein:

L is a group represented by one of Formula 5-1, 5-2, or 5-13, and

Z_1 in Formula 5-1, 5-2, or 5-13 is selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group.

7. The heterocyclic compound as claimed in claim 1, wherein n in Formula 3 is 0, 1, or 2.

8. The heterocyclic compound as claimed in claim 1, wherein Ar in Formula 3 is selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzo-

181

phenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiofenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiofenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiofenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiofenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiofenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and


Si(Q₄)(Q₅)(Q₆), —C(=O)(Q₄), —S(=O)₂(Q₄), and —P(=O)(Q₄)(Q₅),

wherein Q₄ to Q₆ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a


182

C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group.


9. The heterocyclic compound as claimed in claim 1, wherein Ar in Formula 3 is a group represented by one of the following Formulae 6-1 to 6-37:




Formula 6-1




Formula 6-2




Formula 6-3




Formula 6-4




Formula 6-5



Formula 6-6










Formula 6-7



Formula 6-8

185

-continued




186

-continued


Formula 6-23

5




Formula 6-24

15




Formula 6-25

20




Formula 6-26

30



Formula 6-27

40



Formula 6-28

50

Formula 6-29

60

65

Formula 6-30

Formula 6-31

Formula 6-32

Formula 6-33

Formula 6-34

Formula 6-35

Formula 6-36

Formula 6-37

wherein, in Formulae 6-1 to 6-37,

Y_{31} , Y_{33} and Y_{37} , are each independently O, S, C(Z_{35}) (Z_{36}), N(Z_{37}), or Si(Z_{38})(Z_{39}),

Y_{32} , Y_{34} , Y_{36} , and Y_{38} are each independently C(Z_{35}) or N,

Z_{31} to Z_{39} are each independently selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thi-

azolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazolinyl group, a benzoquinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothio-phenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indenocarbazolyl group, an indolocarbazolyl group, a benzoacridinyl group, a benzophenanthrenyl group, a di benzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, in which Q_{31} to Q_{33} are each independently selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group,

e2 is an integer of 0 to 2,

e3 is an integer of 0 to 3,

e4 is an integer of 0 to 4,


e5 is an integer of 0 to 5,

e6 is an integer of 0 to 6, and


* indicates a binding site to a neighboring atom.

10. The heterocyclic compound as claimed in claim 9, wherein Ar is a group represented by one of Formulae 6-1 to 6-6, 6-13 to 6-29, 6-31, 6-32, and 6-37.


11. The heterocyclic compound as claimed in claim 1, wherein Ar in Formula 3 is a group represented by one of the following Formulae 7-1 to 7-25:




Formula 7-1




Formula 7-2




Formula 7-3




Formula 7-4




Formula 7-5




Formula 7-6




Formula 7-7



Formula 7-8



Formula 7-9



Formula 7-10

5

10

15

20

25

30

35

40

45








50

55

60

65


189
-continued



190
-continued


Formula 7-11

5




Formula 7-12

15



Formula 7-13


20



Formula 7-14

25

30




Formula 7-15

35

40

Formula 7-16

45




50

55

Formula 7-17

60



65

Formula 7-18

Formula 7-19



Formula 7-20

Formula 7-21

Formula 7-22

Formula 7-23

191
-continued




wherein * in Formulae 7-1 to 7-25 indicates a binding site to a neighboring atom.

12. The heterocyclic compound as claimed in claim 1, wherein A₁ is a group represented by one of the following Formulae 2A to 2F:

Formula 7-24

5




10

15

Formula 7-25

20



25

30


35

40


45

50

Formula 2A



Formula 2B



192
-continued

Formula 2C

Formula 2D

Formula 2E

Formula 2F

wherein, in Formulae 2A to 2F,
X₁ and X₂ are each independently O or S,
R₁ to R₁₁ are each independently selected from a binding site to A₂, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalk-



193

enyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_1)(\text{Q}_2)(\text{Q}_3)$, and $-\text{S}(=\text{O})_2(\text{Q}_1)$, in which Q_1 to Q_3 are each independently selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group.

13. The heterocyclic compound as claimed in claim 12, wherein R_5 in Formulae 2A to 2F is a binding site to A_2 .





14. The heterocyclic compound as claimed in claim 12, wherein X_1 and X_2 in Formula 2A to 2F are O.

15. The heterocyclic compound as claimed in claim 1, wherein the heterocyclic compound represented by Formula 1 is one of the following Compounds 1 to 72 and 74 to 138:











194

-continued



195
-continued

196
-continued



30

35

40

45

50

55

11


12

13


14

197
-continued

15



16



20

25

17


35

40

45

50

18




65

198
-continued

19

5



15

16

20

25

30

17

35


40

45

50

18

55




65


21

199
-continued


22



25




24




200
-continued


25



26




27



201
-continued

28



202
-continued


31

5

10

15

20



32

25
29


30

35

40

45

30





33

50

55


60

65




203
-continued

34




204
-continued

37




35

38



45


36



65


50

39






205
-continued

40




41



206
-continued

44

5




10

15

20

25



35

40


45

50

55


60

65




207
-continued


47



48



49



65

208
-continued

50

5

10

15

20

25

30

35

40

45


49

50


55

60


65



51





52




209
-continued






210
-continued






211
-continued



212
-continued




213
-continued



214
-continued


67

70




15

71




68

72







69

74








65

215
-continued




216
-continued







217

-continued

84







85



218

-continued


89



219

-continued

93



220


-continued

97

5


10

15




94

20




25

30



95


35



40


45

50




96

55



60


65




221

-continued


101




102



103



104




222

-continued

105

5




10

15

106

20

25




30

107

35

40

45




50

108

55


60

65




223
-continued


109




110



111




112 50



224
-continued

113

5




10

15

114

20




25

30

111

35




115

40

45

112 50

55



60

65


116

225
-continued


226
-continued

117

121





118





119

122



119



123



120

120


124



121

227
-continued

125



5


10

15

20

25

126




30

35

40

127 45



50


55

60


65

228
-continued


128



129



130




229
-continued

230
-continued

131


134



20

25

135




133 45

40

50

136



65

60

55

5

10

15

20

25

30

35

40


45

50

55

60

65



30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30

35

40

45

50

55

60

65

5

10

15

20

25

30


35

40

45

231

-continued



232

16. An organic light-emitting device, comprising:
 a first electrode;
 a second electrode facing the first electrode; and
 an organic layer between the first electrode and the second
 electrode and comprises an emission layer,
 wherein the organic layer includes the heterocyclic com-
 pound as claimed in claim 1.

17. The organic light-emitting device as claimed in claim
 16, wherein:

the first electrode is an anode,
 the second electrode is a cathode, and
 the organic layer includes:

a hole transport region between the first electrode and
 the emission layer, the hole transport region includ-
 ing at least one selected from a hole injection layer,
 a hole transport layer, a buffer layer, and an electron
 blocking layer, and

an electron transport region between the emission layer
 and the second electrode, the electron transport
 region including at least one selected from a hole
 blocking layer, an electron transport layer, and an
 electron injection layer.

18. The organic light-emitting device as claimed in claim
 17, wherein at least one of the electron transport region and
 the emission layer includes the heterocyclic compound.

19. The organic light-emitting device as claimed in claim
 17, wherein:

the electron transport region includes the electron trans-
 port layer or the electron injection layer, and
 the electron transport layer or the electron injection layer
 includes heterocyclic compound.

20. The organic light-emitting device as claimed in claim
 18, wherein the emission layer further includes a pyrene-
 based compound, an arylamine compound, or a styryl-based
 compound.

* * * * *

专利名称(译)	杂环化合物和包括其的有机发光装置		
公开(公告)号	US10305046	公开(公告)日	2019-05-28
申请号	US15/496120	申请日	2017-04-25
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	SIM MUNKI KIM YOUNGKOOK PARK JUNHA LEE HYOYOUNG JEONG EUNJAE HWANG SEOKHWAN		
发明人	SIM, MUNKI KIM, YOUNGKOOK PARK, JUNHA LEE, HYOYOUNG JEONG, EUNJAE HWANG, SEOKHWAN		
IPC分类号	H01L51/00 H01L27/32 C07D491/153 H01L51/52 C07F9/6561 H01L51/50		
CPC分类号	H01L51/0071 C07D491/153 C07F9/6561 H01L27/3211 H01L51/0052 H01L51/0058 H01L51/0072 H01L51/5218 H01L51/0067 H01L51/5096 H01L51/5056 H01L51/5072 H01L51/5088 H01L51/5092		
审查员(译)	杨, JAY		
优先权	1020160061439 2016-05-19 KR		
其他公开文献	US20170338419A1		
外部链接	Espacenet		

摘要(译)

杂环化合物和包含其的有机发光装置，该化合物由式1表示：A 1 - (A 2) a₂₁，20032003 & #x3c; Formula 1> 其中，在公式1中，A 1 是表示的组通过式2，A 2 是由式3表示的基团，并且a₂₁是1至11的整数，其中，当a₂₁是2至11时，A 2 彼此相同或不同：

10

190

150

110