



US 20160043333A1

(19) **United States**

(12) **Patent Application Publication**
KIM et al.

(10) **Pub. No.: US 2016/0043333 A1**
(43) **Pub. Date: Feb. 11, 2016**

(54) **CONDENSED CYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

(52) **U.S. CL.**
CPC **H01L 51/0094** (2013.01); **C07F 7/0807** (2013.01); **H01L 51/5028** (2013.01)

(71) Applicants: **SAMSUNG DISPLAY CO., LTD.**,
Yongin-City (KR); **Gyeongsang National University Industry-Academic Cooperation Foundation**, Jinju (KR)

(57) **ABSTRACT**

A condensed cyclic compound and an organic light-emitting device, the condensed cyclic compound being represented by Formula 1:

(72) Inventors: **Mikyung KIM**, Yongin-City (KR); **Yunhi KIM**, Jinju (KR); **Jaeyoung HWANG**, Jinju (KR)

(21) Appl. No.: **14/618,835**

(22) Filed: **Feb. 10, 2015**

(30) **Foreign Application Priority Data**

Aug. 5, 2014 (KR) 10-2014-0100696

Publication Classification

(51) **Int. Cl.**
H01L 51/00 (2006.01)
H01L 51/50 (2006.01)
C07F 7/08 (2006.01)




FIG. 1

10

190
150
110

FIG. 2





FIG. 3



**CONDENSED CYCLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**

CROSS-REFERENCE TO RELATED
APPLICATION

[0001] Korean Patent Application No. 10-2014-0100696, filed on Aug. 5, 2014, in the Korean Intellectual Property Office, and entitled: "Condensed Cyclic Compound and Organic Light-Emitting Device Comprising the Same," is incorporated by reference herein in its entirety.

BACKGROUND

[0002] 1. Field

[0003] Embodiments relate to a condensed cyclic compound and an organic light-emitting device including the same.

[0004] 2. Description of the Related Art


[0005] Organic light-emitting devices (OLEDs), which are self-emitting devices, may have advantages such as wide viewing angles, excellent contrast, quick response, high brightness, excellent driving voltage characteristics, and can provide multicolored images.

[0006] An organic light-emitting device may have a structure in which a first electrode, a hole transport region, an emission layer, an electron transport region, and a second electrode are sequentially disposed in this order on a substrate. Holes injected from the first electrode may move to the emission layer via the hole transport region, while electrons injected from the second electrode may move to the emission layer via the electron transport region. Carriers such as the holes and electrons may recombine in the emission layer to generate excitons. When the excitons drop from an excited state to a ground state, light may be emitted.

SUMMARY

[0007] Embodiments are directed to a condensed cyclic compound and an organic light-emitting device including the same.

[0008] According to one or more embodiments, there is provided a condensed cyclic compound represented by Formula 1:



[0009] wherein, in Formula 1,

[0010] A₁ and A₂ are each independently a C₆-C₂₀ aromatic ring or a C₂-C₂₀ heteroaromatic ring, wherein A₁ and A₂ are not benzene at the same time;

[0011] L₁ to L₆ are each independently selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene

group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

[0012] a₁ to a₆ are each independently selected from 0, 1, 2, and 3;

[0013] R₁ to R₆ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —B(Q₃)(Q₄), and —Si(Q₅)(Q₆)(Q₇);

[0014] b₁ to b₆ are each independently selected from 0, 1, 2, and 3;

[0015] c₁ and c₂ are each independently selected from 0, 1, 2, and 3;

[0016] wherein at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from

[0017] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0018] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I,

a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —B(Q₁₃)(Q₁₄), and —Si(Q₁₅)(Q₁₆)(Q₁₇);

[0019] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0020] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —B(Q₂₃)(Q₂₄), and —Si(Q₂₅)(Q₂₆)(Q₂₇); and

[0021] —N(Q₃₁)(Q₃₂), —B(Q₃₃)(Q₃₄), and —Si(Q₃₅)(Q₃₆)(Q₃₇),

[0022] wherein Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent aromatic condensed heteropolycyclic group.

[0023] According to one or more embodiments, an organic light-emitting device includes: a first electrode; a second electrode disposed opposite to the first electrode; and an organic layer disposed between the first electrode and the second electrode and including an emission layer, wherein the organic layer includes at least one of the above-described condensed cyclic compounds of Formula 1.

BRIEF DESCRIPTION OF THE DRAWINGS

[0024] Features will be apparent to those of skill in the art by describing in detail exemplary embodiments with reference to the attached drawings in which:

[0025] FIG. 1 illustrates a schematic view of a structure of an organic light-emitting device according to an embodiment;

[0026] FIG. 2 illustrates UV absorption and photoluminescence (PL) spectra of Compound 5 obtained in Synthesis Example 1, and low-temperature PL spectrum thereof; and

[0027] FIG. 3 illustrates a cyclic voltammetry curve of Compound 5 of Synthesis Example 1.

DETAILED DESCRIPTION


[0028] Example embodiments will now be described more fully hereinafter with reference to the accompanying drawings; however, they may be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey exemplary implementations to those skilled in the art.

[0029] In the drawing figures, the dimensions of layers and regions may be exaggerated for clarity of illustration. Like reference numerals refer to like elements throughout.

[0030] As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

[0031] According to an embodiment, there is provided a condensed cyclic compound represented by Formula 1:

<Formula 1>



[0032] In Formula 1,

[0033] At and A₂ may be each independently a C₆-C₂₀ aromatic ring or a C₁-C₂₀ heteroaromatic ring. In an implementation, A₁ and A₂ may not both be a benzene at the same time.

[0034] In Formula 1, A₁ and A₂ may be condensed rings each sharing carbon with an adjacent 6-membered ring (e.g., the 6 membered ring that includes silicon).

[0035] For example, A₁ and A₂ may be the same or differ from each other. When A₁ and A₂ are the same, A₁ and A₂ may not both be benzene.

[0036] In an implementation, the C₆-C₂₀ aromatic ring may be a benzene, a naphthalene, or an anthracene; and the C₁-C₂₀ heteroaromatic ring may be a pyridine, a pyridazine, a pyrimidine, a pyrazine, a triazine, a quinoline, an isoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a benzothiofene, a benzofuran, a dibenzothiofene, or a dibenzofuran.

[0037] In an implementation, A_1 may be a benzene, a naphthalene, or an anthracene; and A_2 may be a naphthalene or an anthracene.

[0038] In an implementation, A_1 and A_2 may be each independently a pyridine, a pyridazine, a pyrimidine, a pyrazine, a triazine, a quinoline, an isoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a benzothiophene, a benzofuran, a dibenzothiophene, or a dibenzofuran.

[0039] In an implementation, A_1 may be a benzene, a naphthalene, an anthracene, a pyridine, a pyridazine, a pyrimidine, a pyrazine, a triazine, a quinoline, an isoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, or a cinnoline; and A_2 may be a naphthalene, an anthracene, a pyridine, a pyridazine, a pyrimidine, a pyrazine, a triazine, a quinoline, an isoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, or a cinnoline. However, embodiments of the present disclosure are not limited thereto.

[0040] In Formula 1, L_1 to L_6 may be each independently selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.


[0041] In an implementation, L_1 to L_6 in Formula 1 may be each independently selected from:

[0042] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isooxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzoimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzooxazolylene group, an isobenzooxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group; and


[0043] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isooxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzoimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzooxazolylene group, an isobenzooxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, an a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl

group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.


[0044] In an implementation, L_1 to L_6 in Formula 1 may be each independently a group represented by one of Formulae 3-1 to 3-32:




Formula 3-1




Formula 3-2




Formula 3-3




Formula 3-4



Formula 3-5




Formula 3-6




Formula 3-7


-continued




Formula 3-8




Formula 3-9




Formula 3-10




Formula 3-11




Formula 3-12




Formula 3-13




Formula 3-14



Formula 3-15













Formula 3-16




Formula 3-17

-continued




-continued

Formula 3-18




Formula 3-19




Formula 3-20

Formula 3-21




Formula 3-22

Formula 3-23



Formula 3-24



Formula 3-25

Formula 3-26

Formula 3-27

Formula 3-28

Formula 3-29

Formula 3-30

Formula 3-31

Formula 3-32

[0045] In Formulae 3-1 to 3-32,

[0046] Y_1 may be O, S, $C(Z_3)(Z_4)$, $N(Z_5)$, or $Si(Z_6)(Z_7)$;


[0047] Z_1 to Z_7 may be each independently selected from a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino groups, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;

[0048] d_1 may be an integer selected from 1 to 4;


[0049] d_2 may be an integer selected from 1 to 3;

[0050] d_3 may be an integer selected from 1 to 6;


- [0051] d4 may be an integer selected from 1 to 8;
 [0052] d5 may be 1 or 2;
 [0053] d6 may be an integer selected from 1 to 5; and
 [0054] * and *' may be binding sites with adjacent atoms.
 [0055] In an implementation, L₁ to L₆ in Formula 1 may be each independently a group represented by one of Formulae 4-1 to 4-23.




Formula 4-1




Formula 4-2




Formula 4-3




Formula 4-4




Formula 4-5




Formula 4-6




Formula 4-7



Formula 4-8




Formula 4-9




-continued


Formula 4-10




Formula 4-11




Formula 4-12




Formula 4-13



Formula 4-14




Formula 4-15




Formula 4-16


-continued




Formula 4-17




Formula 4-18




Formula 4-19




Formula 4-20



Formula 4-21



Formula 4-22



Formula 4-23

[0056] In Formulae 4-1 and 4-23, * and *' may be binding sites with adjacent atoms.

[0057] In Formula 1, a₁, which indicates the number of L₁s, may be selected from 0, 1, 2, and 3. In an implementation, a₁ may be 0 or 1. When a₁ is 0, *(L₁)_{a₁}*' may be a single bond. When a₁ is 2 or greater, two or more L₁s may be the same or differ from each other. In Formula 1, a₂, a₃, a₄, a₅, and a₆ may be understood based on the description of a₁ and the structure of Formula 1.

[0058] In an implementation, a₁ to a₆ in Formula 1 may be each independently 0 or 1.

[0059] In Formula 1, R₁ to R₆ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —B(Q₃)(Q₄), and —Si(Q₅)(Q₆)(Q₇), wherein Q₁ to Q₇ may be the same as defined herein.

[0060] In an implementation, R₁ to R₆ in Formula 1 may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C₁-C₂₀ alkyl group, a substituted or unsubstituted C₁-C₂₀ alkoxy group, a substituted or unsubstituted C₆-C₂₀ aryl group, a substituted or unsubstituted C₁-C₂₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and —Si(Q₅)(Q₆)(Q₇), wherein Q₅ to Q₇ may be the same as defined herein.

[0061] In an implementation, R₁ to R₆ in Formula 1 may be each independently selected from:

[0062] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0063] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinoxalinyl group, a cinnolinyl group, a carbazolyl group, a

phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0064] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group,

a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

[0065] —Si(Q₅)(Q₆)(Q₇), wherein Q₅ to Q₇ may be the same as defined herein.

[0066] In an implementation, R₁ to R₆ in Formula 1 may be each independently selected from:


[0067] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0068] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and


[0069] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-

fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.


[0070] In an implementation, R_1 to R_6 in Formula 1 may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, and a group represented by one of Formulae 6-1 to 6-49.




Formula 6-1




Formula 6-2




Formula 6-3




Formula 6-4




Formula 6-5




Formula 6-6




Formula 6-7




Formula 6-8



Formula 6-9












Formula 6-10






Formula 6-11

-continued


-continued




Formula 6-12




Formula 6-13




Formula 6-14




Formula 6-15




Formula 6-16




Formula 6-17



Formula 6-18



Formula 6-19



Formula 6-20

-continued

Formula 6-21

Formula 6-22

Formula 6-23

Formula 6-24

Formula 6-25

Formula 6-26

Formula 6-27









Formula 6-28

Formula 6-29

Formula 6-30

Formula 6-31

-continued



Formula 6-32

Formula 6-33

Formula 6-34

Formula 6-35






Formula 6-36

Formula 6-37

Formula 6-38

Formula 6-39

-continued



Formula 6-40


Formula 6-41

Formula 6-42


Formula 6-43

Formula 6-44


-continued




Formula 6-45




Formula 6-46



Formula 6-47



Formula 6-48



Formula 6-49

[0071] In Formulae 6-1 to 6-49, * may be a binding site with an adjacent atom.


[0072] In Formula 1, b_1 , which indicates the number of R_1 s, may be selected from 0, 1, 2 and 3. In an implementation, b_1 may be 1 or 2. When b_1 is 2 or greater, two or more R_1 s may be the same or differ from each other. In Formula 1, b_2 , b_3 , b_4 , b_5 , and b_6 may be understood based on the description of b_1 and the structure of Formula 1.

[0073] In Formula 1, c_1 , which indicates the number of $*(L_1)_{a1}-(R_1)_{b1}$ s, may be selected from 0, 1, 2, and 3. In an implementation, c_1 may be 0 or 1. When c_1 is 0, A_1 may be an unsubstituted ring. When c_1 is 2 or greater, two or more $*(L_1)_{a1}-(R_1)_{b1}$ may be the same or differ from each other. In Formula 1, c_2 may be understood based on the description of c_1 and the structure of Formula 1.


[0074] In an implementation, in Formula 1, c_1 and c_2 may be 0, a_3 to a_6 may be 0, and b_3 to b_6 may be 1.

[0075] In an implementation, in Formula 1, when c_1 and c_2 are 0, a_3 to a_6 are 0, and b_3 to b_6 are 1, R_3 to R_6 may be the same or differ from each other.


[0076] In an implementation, the condensed cyclic compound represented by Formula 1 may be represented by one of Formulae 1A to 1E.




<Formula 1A>




<Formula 1B>



<Formula 1C>



<Formula 1D>



<Formula 1E>

[0077] In Formulae 1A to 1E, L_1 to L_6 , a_1 to a_6 , R_1 to R_6 , b_1 to b_6 , c_1 , and c_2 may be the same as those defined above herein.


[0078] In an implementation, in Formulae 1A to 1E, L_1 to L_6 may be each independently a group represented by one of Formulae 4-1 to 4-23; and a_1 to a_6 may be each independently 0 or 1.

[0079] In an implementation, in Formulae 1A to 1E, R_1 to R_6 may be each independently selected from a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, and a group represented by one of Formulae 6-1 to 6-49; and b_1 to b_6 may be 1.


[0080] In an implementation, the condensed cyclic compound of Formula 1 may be represented by one of Formulae 1A(1) to 1A(5), Formulae 2A(1) to 2A(5), Formula 3A(1), and Formula 4A(1).

-continued


<Formula 1A(3)>




<Formula 1A(4)>




<Formula 1A(5)>



<Formula 1A(1)>




<Formula 1A(2)>




-continued

<Formula 2A(1)>




-continued


<Formula 3A(1)>




<Formula 2A(2)>



<Formula 4A(1)>




<Formula 2A(5)>



[0081] In Formulae 1A(1) to 1A(5), Formulae 2A(1) to 2A(5), Formula 3A(1), and Formula 4A(1), R₁, R₂, b1, and b2 may be the same as those defined herein, R_{3a} to R_{3g} may be each independently selected from the same groups as those defined for R₃ herein, R_{4a} to R_{4g} may be each independently selected from the same groups as those defined for R₄ herein, R_{5a} to R_{5g} may be each independently selected from the same groups as those defined for R₅ herein, and R_{6a} to R_{6g} may be each independently selected from the same groups as those defined for R₆ herein.


[0082] In an implementation, the condensed cyclic compound represented by Formula 1 may be one of Compounds 1 to 7.



-continued


1

6



2

7




3

4

[0083] The condensed cyclic compound represented by Formula 1 may include a core represented by Formula 1', and thus may have a high triplet energy level (T1) and may be suitable as a material for organic light-emitting devices, e.g., as a phosphorescent host.

<Formula 1'>



5

[0084] In Formula 1, A₁ and A₂ may not be benzene at the same time. For example, if a condensed cyclic compound of Formula 1 were to include both A₁ and A₂ as benzene at the same time, the compound (e.g., Compound A of Comparative Example 1, below) may have a relatively high glass transition temperature (T_g), a relatively high triplet energy level (T₁), and poor energy transfer characteristics, and thus may not be suitable for use as a material for organic light-emitting devices, e.g., as a phosphorescent host.

[0085] The condensed cyclic compound of Formula 1 may be synthesized using a suitable organic synthesis method. Methods of synthesizing the condensed cyclic compound of Formula 1 may be understood, e.g., based on the examples that will be described below.

[0086] The condensed cyclic compound of Formula 1 may be used or included between a pair of electrodes of an organic light-emitting device. In an implementation, the condensed cyclic compound of Formula 1 may be included in an electron transport region, e.g., in an electron transport layer.

[0087] According to another embodiment, an organic light-emitting device may include a first electrode, a second electrode opposite to the first electrode, and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer. The organic layer may include at least one condensed cyclic compound represented by Formula 1 described above.

[0088] As used herein, “(for example, the organic layer) including at least one condensed cyclic compound means that “(the organic layer) including one of the condensed cyclic compounds of Formula 1, or at least two different condensed cyclic compounds of Formula 1.”

[0089] In an implementation, the organic layer may include only Compound 1 as the condensed cyclic compound. In this regard, Compound 1 may be present in the emission layer of the organic light-emitting device. In an implementation, the organic layer may include Compounds 1 and 2 as the condensed cyclic compounds. In this regard, Compounds 1 and 2 may be present both in the same layer (for example, in the emission layer) or may be present in different layers (for example, in the emission layer and the electron transport layer, respectively).

[0090] The organic layer may include, e.g., i) a hole transport region between the first electrode (anode) and the emission layer and including at least one of a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer; and ii) an electron transport region between the emission layer and the second electrode (cathode) and including at least one of a hole blocking layer, an electron transport layer, and an electron injection layer. The emission layer may include the condensed cyclic compound of Formula 1.

[0091] As used herein, the term “organic layer” refers to a single layer and/or a plurality of layers disposed between the first and second electrodes of the organic light-emitting device. A material in the “organic layer” is not limited to an organic material.

[0092] FIG. 1 illustrates a schematic sectional view of an organic light-emitting device 10 according to an embodiment. Referring to FIG. 1, the organic light-emitting device 10 may include a first electrode 110, an organic layer 150, and a second electrode 190.

[0093] Hereinafter, a structure of an organic light-emitting device according to an embodiment and a method of manufacturing the same will now be described with reference to FIG. 1.

[0094] A substrate (not shown) may be disposed under the first electrode 110 or on the second electrode 190 in FIG. 1. The substrate may be a glass or transparent plastic substrate with good mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

[0095] For example, the first electrode 110 may be formed by depositing or sputtering a first electrode-forming material on the substrate. When the first electrode 110 is an anode, a material having a high work function may be used as the first electrode-forming material to facilitate hole injection. The first electrode 110 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. Transparent and conductive materials such as indium oxide (ITO),

indium zinc oxide (IZO), tin oxide (SnO₂), and zinc oxide (ZnO) may be used to form the first electrode. When the first electrode 110 is a semi-transmissive electrode or a reflective electrode, at least one material selected from magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag) may be used as a material for forming the first electrode.

[0096] The first electrode 110 may have a single-layer structure or a multi-layer structure including a plurality of layers. For example, the first electrode 110 may have a three-layered structure of ITO/Ag/ITO.

[0097] The organic layer 150 may be disposed on the first electrode 110. The organic layer 150 may include an emission layer (EML).

[0098] The organic layer 150 may include a hole transport region between the first electrode and the EML, and an electron transport region between the EML and the second electrode.

[0099] For example, the hole transport region may include at least one of a hole injection layer (HIL), a hole transport layer (HTL), a buffer layer, and an electron blocking layer (EBL).

[0100] For example, the electron transport region may include at least one of a hole blocking layer (HBL), an electron transport layer (ETL), and an electron injection layer (EIL).

[0101] The hole transport region may have a single-layered structure including a single material, a single-layered structure including a plurality of different materials, or a multi-layered structure including a plurality of layers including different materials.

[0102] In an implementation, the hole transport region may have a single-layered structure including a plurality of different materials, or a multi-layered structure of HIL/HTL, HIL/HTL/buffer layer, HIL/buffer layer, HTL/buffer layer, or HIL/HTL/EBL, wherein these layers forming a multi-layered structure are sequentially disposed on the first electrode 110 in the order stated above.

[0103] When the hole transport region includes a HIL, the HIL may be formed on the first electrode 110 by using any of a variety of methods, e.g., by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like.


[0104] When the HIL is formed using vacuum deposition, the deposition conditions may vary depending on a compound that is used to form the HIL and the structure of the HIL. For example, the deposition conditions may include a deposition temperature of about 100° C. to about 500° C., a degree of vacuum of about 10⁻⁸ to about 10⁻³ torr, and a deposition rate of about 0.01 to about 100 Å/sec.

[0105] When the HIL is formed using spin coating, the coating conditions may vary depending on a compound that is used to form the HIL and the structure of the HIL. For example, the coating conditions may include a coating rate of about 2,000 rpm to about 5,000 rpm and a heat treatment temperature of about 80° C. to about 200° C.


[0106] When the hole transport region includes a HTL, the HTL may be formed on the first electrode 110 or the HIL by using any of a variety of methods, e.g., by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like. When the HTL is formed

using vacuum deposition or spin coating, the conditions for deposition and coating for forming the HTL may be similar to the above-described deposition and coating conditions for forming the HIL, and accordingly will not be described in detail.


[0107] In an implementation, the hole transport region may include at least one of m-MTDATA, TDATA, 2-TNATA, NPB, β -NPB, TPD, Spiro-TPD, Spiro-NPB, α -NPB, TAPC, HMTPD, DNTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzene sulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate)(PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201 below, and a compound represented by Formula 202 below.




m-MTDATA





TDATA



2-TNATA



NPB

 β -NPB

TPD

quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylene group, a carbazolylene group, and a triazinylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, a quinolinylnyl group, an isoquinolinylnyl group, a quinoxalinylnyl group, a quinazolinylnyl group, a carbazolyl group, and a triazinyl group.

[0117] xa1 to xa4 may be each independently 0, 1, or 2;


[0118] xa5 may be 1, 2, or 3;

[0119] R₂₀₁ to R₂₀₄ may be each independently selected from:


[0120] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinylnyl group, an isoquinolinylnyl group, a quinoxalinylnyl group, a quinazolinylnyl group, a carbazolyl group, and a triazinyl group; and

[0121] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinylnyl group, an isoquinolinylnyl group, a quinoxalinylnyl group, a quinazolinylnyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an azulenylnyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinylnyl group, an isoquinolinylnyl group, a quinoxalinylnyl group, a quinazolinylnyl group, a carbazolyl group, and a triazinyl group. However, embodiments of the present disclosure are not limited thereto.


[0122] The compound of Formula 201 may be represented by Formula 201A.



[0123] For example, the compound of Formula 201 may be represented by Formula 201A-1:



[0124] The compound of Formula 202 may be represented by Formula 202A:



[0125] In Formulae 201A, 201A-1, and 202A,

[0126] L₂₀₁ to L₂₀₃, xa1 to xa3, xa5, and R₂₀₂ to R₂₀₄ may be defined as described in conjunction with Formula 201;

[0127] R₂₁₁ may be defined as described in conjunction with R₂₀₃ in Formula 201;

[0128] R₂₁₃ to R₂₁₆ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkoxy group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0129] In an implementation, in Formulae 201A, 201A-1, and 202A,

dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group,


[0145] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, and a triazinyl group, and

[0146] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group; and

[0147] xa5 may be 1 or 2.


[0148] In Formulae 201A and 201A-1, R₂₁₃ and R₂₁₄ may be linked to each other to form a saturated or unsaturated ring.

[0149] In an implementation, the compound represented by Formula 201 and/or the compound represented by Formula 202 may be compounds HT1 to HT20 illustrated below.




-continued

HT2




HT3




-continued

HT4




-continued


HT6



HT5




HT7




-continued

HT8




-continued


HT10



HT9




HT11




-continued

HT12




-continued


HT16




HT13




HT14




HT15




HT17




HT18



HT19




-continued




[0150] A thickness of the hole transport region may be about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å. When the hole transport region includes a HIL and a HTL, a thickness of the HIL may be about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å, and a thickness of the HTL may be about 50 Å to about 2,000 Å, e.g., about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the HIL, and the HTL are within these ranges, satisfactory hole transport characteristics may be obtained without a substantial increase in driving voltage.

[0151] The hole transport region may further include a charge-generating material to help improve conductivity, in addition to the materials as described above. The charge-generating material may be homogeneously or inhomogeneously dispersed in the hole transport region.

[0152] The charge-generating material may be, e.g., a p-dopant. The p-dopant may include one of quinone derivatives, metal oxides, and cyano group-containing compounds. Non-limiting examples of the p-dopant may include quinone derivatives such as tetracyanoquinonodimethane (TCNQ), 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonodimethane (F4-TCNQ), and the like; metal oxides such as tungsten oxide, molybdenum oxide, and the like; and Compound HT-D1 below.




-continued



[0153] The hole transport region may further include at least one of a buffer layer and an EBL, in addition to the HIL and HTL described above. The buffer layer may help compensate for an optical resonance distance of light according to a wavelength of the light emitted from the EML, and thus may help improve light-emission efficiency. A material in the buffer layer may be a suitable material used in the hole transport region. The EBL may help block injection of electrons from the electron transport region.

[0154] For example, a material for forming the EBL may include mCP, below.









[0155] The EML may be formed on the first electrode 110 or the hole transport region by using any of a variety of methods, e.g., by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like. When the EML is formed using vacuum deposition or spin coating, the deposition and coating conditions for forming the EML may be similar to the above-described deposition and coating conditions for forming the HIL, and accordingly may not be described in detail.

[0156] When the organic light-emitting device 10 is a full color organic light-emitting device, the EML may be patterned into a red emission layer, a green emission layer, and a blue emission layer to correspond to individual subpixels, respectively. In an implementation, the EML may have a structure in which a red emission layer, a green emission layer, and a blue emission layer are stacked upon one another, or a structure including a mixture of a red light-emitting material, a green light-emitting material, and a blue light-emitting material, and thus may emit white light.

[0157] The EML may include the condensed compound represented by Formula 1.

[0158] The EML may include a host and a dopant. The host may include the condensed cyclic compound represented by Formula 1.


[0159] The host may further include at least one of TPBi, TBADN, ADN (also referred to as "DNA"), CBP, CDBP, and TCP, in addition to the condensed cyclic compound of Formula 1.



[0160] The dopant may include at least one of a fluorescent dopant and a phosphorescent dopant.

[0161] The phosphorescent dopant may include an organo-metallic complex represented by Formula 401.

<Formula 401>



[0162] In Formula 401,

[0163] M may be selected from iridium (Ir), platinum (Pt), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), and thulium (Tm);

[0164] X_{401} to X_{404} may be each independently a nitrogen or a carbon;

[0165] rings A_{401} and A_{402} may be each independently selected from a substituted or unsubstituted benzene, a substituted or unsubstituted naphthalene, a substituted or unsubstituted fluorene, a substituted or unsubstituted spiro-fluorene, a substituted or unsubstituted indene, a substituted or unsubstituted pyrrole, a substituted or unsubstituted thiophene, a substituted or unsubstituted furan, a substituted or unsubstituted imidazole, a substituted or unsubstituted pyrazole, a substituted or unsubstituted thiazole, a substituted or unsubstituted isothiazole, a substituted or unsubstituted oxazole, a substituted or unsubstituted isoxazole, a substituted or unsubstituted pyridine, a substituted or unsubstituted pyrazine, a substituted or unsubstituted pyrimidine, a substituted or unsubstituted pyridazine, a substituted or unsubstituted quinoline, a substituted or unsubstituted isoquinoline, a substituted or unsubstituted benzoquinoline, a substituted or unsubstituted quinoxaline, a substituted or unsubstituted quiazoline, a substituted or unsubstituted carbazole, a substituted or unsubstituted benzimidazole, a substituted or unsubstituted benzofuran, a substituted or unsubstituted benzothiophene, a substituted or unsubstituted isobenzothiophene, a substituted or unsubstituted benzoxazole, a

substituted or unsubstituted isobenzoxazole, a substituted or unsubstituted triazole, a substituted or unsubstituted oxadiazole, a substituted or unsubstituted triazine, a substituted or unsubstituted dibenzofuran, and a substituted or unsubstituted dibenzothiophene;

[0166] at least one substituent of the substituted benzene, the substituted naphthalene, the substituted fluorene, the substituted spiro-fluorene, the substituted indene, the substituted pyrrole, the substituted thiophene, the substituted furan, the substituted imidazole, the substituted pyrazole, the substituted thiazole, the substituted isothiazole, the substituted oxazole, the substituted isoxazole, the substituted pyridine, the substituted pyrazine, the substituted pyrimidine, the substituted pyridazine, the substituted quinoline, the substituted isoquinoline, the substituted benzoquinoline, the substituted quinoxaline, the substituted quinazoline, the substituted carbazole, the substituted benzoimidazole, the substituted benzofuran, the substituted benzothiophene, the substituted isobenzothiophene, the substituted benzoxazole, the substituted isobenzoxazole, the substituted triazole, the substituted oxadiazole, the substituted triazine, the substituted dibenzofuran, and the substituted dibenzothiophene may be selected from:

[0167] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0168] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₄₀₁)(Q₄₀₂), —Si(Q₄₀₃)(Q₄₀₄)(Q₄₀₅), and —B(Q₄₀₆)(Q₄₀₇),

[0169] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group,

[0170] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group

or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₄₁₁)(Q₄₁₂), —Si(Q₄₁₃)(Q₄₁₄)(Q₄₁₅), and —B(Q₄₁₆)(Q₄₁₇), and

[0171] —N(Q₄₂₁)(Q₄₂₂), —Si(Q₄₂₃)(Q₄₂₄)(Q₄₂₅), and —B(Q₄₂₆)(Q₄₂₇);

[0172] L₄₀₁ may be an organic ligand;

[0173] xc1 may be 1, 2, or 3; and

[0174] xc2 may be 0, 1, 2, or 3,


[0175] wherein Q₄₀₁ to Q₄₀₇, Q₄₁₁ to Q₄₁₇ and Q₄₂₁ to Q₄₂₇ may be each independently selected from a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₆-C₆₀ aryl group, and a C₁-C₆₀ heteroaryl group.

[0176] For example, in Formula 401, L₄₀₁ may be a monovalent, divalent, or trivalent organic ligand. For example, L₄₀₁ in Formula 401 may be selected from a halogen ligand (for example, Cl or F), a diketone ligand (for example, acetylacetonate, 1,3-diphenyl-1,3-propanedionate, 2,2,6,6-tetramethyl-3,5-heptanedionate, or hexafluoroacetone), a carboxylic acid ligand (for example, picolinate, dimethyl-3-pyrazolecarboxylate, or benzoate), a carbon monoxide ligand, an isonitrile ligand, a cyano ligand, and a phosphorous ligand (for example, phosphine or phosphite). However, embodiments of the present disclosure are not limited thereto.

[0177] When A₄₀₁ in Formula 401 has at least two substituent groups, the at least two substituent groups of A₄₀₁ may be linked to each other to form a saturated or unsaturated ring.

[0178] When A₄₀₂ in Formula 401 has at least two substituent groups, the at least two substituent groups of A₄₀₂ may be linked to each other to form a saturated or unsaturated ring.


[0179] When xc1 in Formula 401 is 2 or greater, a plurality of ligands




in Formula 401 may be identical to or different from each other. When xc1 in Formula 1 is 2 or greater, A₄₀₁ and A₄₀₂ may be linked to A₄₀₁ and A₄₀₂ of another adjacent ligand, respectively, directly or via a linking group (for example, a C₁-C₅ alkylene group, —N(R')— (where R' is a C₁-C₁₀ alkyl group or a C₆-C₂₀ aryl group), or C(=O)—).

[0180] In an implementation, the phosphorescent dopant may include at least one of Compounds PD1 to PD74.


-continued




PD1




PD6




PD2




PD7




PD3




PD8




PD4



PD9








PD5




PD10

-continued




-continued

PD11




PD16

PD12




PD17

PD13




PD18

PD14








PD19

PD15




PD20

-continued




-continued

PD21




PD26

PD22




PD27

PD23




PD28

PD24








PD29

PD25




PD30

-continued




-continued

PD31




PD36

PD32




PD37

PD33




PD38

PD34








PD39

PD35




PD40

-continued




-continued

PD41




PD46

PD42




PD47

PD43




PD48

PD44




PD49

PD45




PD50



PD51


-continued

PD52





-continued

PD57




PD58


PD53




PD54




PD59




PD55




PD60








PD56



PD61




-continued




-continued

PD62




PD67

PD63




PD68

PD64




PD69

PD65




PD70

PD66




PD71


-continued



PD72




PtOEP




PD73

[0182] In an implementation, the phosphorescent dopant may include Flr6 below.




PD74



Flr6


[0181] In an implementation, the phosphorescent dopant may include PtOEP below.

[0183] In an implementation, the fluorescent dopant may further include at least one of DPAVBi, BDAVBi, TBPe, DCM, DCJTb, Coumarin 6, and C545T.




DPAVBi

-continued




DPAVBi



TBPe

DCM


DCJTb



Coumarin 6

C545T

[0184] In an implementation, the fluorescent dopant may include a compound represented by Formula 501.



<Formula 501>

[0185] In Formula 501,

[0186] Ar_{501} may be selected from:

[0187] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene,

a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene,

[0188] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio

group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, and —Si(Q₅₀₁)(Q₅₀₂)(Q₅₀₃), wherein Q₅₀₁ to Q₅₀₃ may be each independently selected from a hydrogen, C₁-C₆₀alkyl group, a C₂-C₆₀alkenyl group, a C₆-C₆₀ aryl group, and a C₁-C₆₀ heteroaryl group;

[0189] L₅₀₁ to L₅₀₃ may be the same as those for L₂₀₁ defined herein;

[0190] R₅₀₁ and R₅₀₂ may be each independently selected from:

[0191] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, and

[0192] a phenyl group, a naphthyl group, a fluorenyl group, spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, and a dibenzothio group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazolinyl group, a carbazolyl group, a triazinyl group, and a dibenzofuranyl group, and a dibenzothiophenyl group;

[0193] xd1 to xd3 may be each independently selected from 0, 1, 2, and 3; and

[0194] xb4 may be selected from 1, 2, 3, and 4.

[0195] The dopant may be included in the EML in an amount of, e.g., about 0.01 part by weight to about 15 parts by weight, based on 100 parts by weight of the host.

[0196] The thickness of the EML may be about 100 Å to about 1,000 Å, e.g., about 200 Å to about 600 Å. When the thickness of the EML is within these ranges, the EML may have good light emitting ability without a substantial increase in driving voltage.

[0197] Next, the electron transport region may be disposed on the EML.

[0198] The electron transport region may include, e.g., at least one of a HBL, an ETL, and an EIL.

[0199] In an implementation, the electron transport region may have a structure including an ETL/EIL, or a HBL/ETL/EIL, wherein the layers forming a structure of the electron transport region may be sequentially stacked on the EML in the order stated above.


[0200] In an implementation, the organic layer 150 of the organic light-emitting device 10 may include the electron

transport region between the EML and the second electrode 190, and the condensed cyclic compound of Formula 1 may be in the electron transport region.

[0201] The electron transport region may include a HBL. When the EML includes a phosphorescent dopant, the HBL may help reduce and/or help prevent diffusion of triplet excitons or holes into the ETL from the EML.

[0202] When the electron transport region includes a HBL, the HBL may be formed on the EML by using any of a variety of methods, e.g., by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like. When the HBL is formed using vacuum deposition or spin coating, the deposition and coating conditions for forming the HBL may be similar to the above-described deposition and coating conditions for forming the HIL, and accordingly may not be described in detail.

[0203] In an implementation, the HBL may include at least one of BCP and Bphen. However, embodiments of the present disclosure are not limited thereto.





[0204] A thickness of the HBL may be about 20Å to about 1,000Å, e.g., about 30Å to about 300Å. When the thickness of the HBL is within these ranges, the HBL may have improved hole blocking ability without a substantial increase in driving voltage.


[0205] The electron transport region may include an ETL. The ETL may be formed on the EML or the HBL by using any of a variety of methods, e.g., by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like. When the ETL is formed using vacuum deposition or spin coating, the deposition and coating conditions for forming the ETL may be similar to the above-described deposition and coating conditions for forming the HIL, and accordingly may not be described in detail.

[0206] In an implementation, the organic layer 150 of the organic light-emitting device may include an electron transport region between the EML and the second electrode 190. The electron transport region may include at least one of the ETL and the EIL.


[0207] In an implementation, the ETL may include at least one of BCP, Bphen, Alq₃, Balq, TAZ, NTAZ, TSP01, and TmPyPb.

Alq₃


BALq




TAZ



NTAZ



TSP01



TmPyPb

[0208] In an implementation, the ETL may include at least one selected from a compound represented by Formula 601 and a compound represented by Formula 602.



[0209] In Formula 601, Ar₆₀₁ may be selected from:

[0210] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene; and

[0211] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, and —Si(Q₃₀₁)(Q₃₀₂)(Q₃₀₃) (wherein Q₃₀₁ to Q₃₀₃ may be each independently selected from a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₆-C₆₀ aryl group, and a C₁-C₆₀ heteroaryl group);

[0212] L₆₀₁ may be the same as defined for L₂₀₁ herein;

[0213] E₆₀₁ may be selected from:

[0214] a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenan-

throlinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and


[0215] a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coroneryl group, an ovaerenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a ben-

zothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0216] xe1 may be selected from 0, 1, 2, and 3; and

[0217] xe2 may be selected from 1, 2, 3, and 4.

<Formula 602>



[0218] In Formula 602,

[0219] X₆₁₁ may be N or C-(L₆₁₁)_{xe611}-R₆₁₁; X₆₁₂ may be N or C-(L₆₁₂)_{xe612}-R₆₁₂; X₆₁₃ may be N or C-(L₆₁₃)_{xe613}-R₆₁₃; and at least one of X₆₁₁ to X₆₁₃ may be N;

[0220] L₆₁₁ to L₆₁₆ may be the same as defined for L₂₀₁ herein;

[0221] R₆₁₁ to R₆₁₆ may be each independently selected from:

[0222] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and


[0223] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀alkyl group, a C₁-C₂₀alkoxy group, a phenyl group, a naphthyl group, an azulenyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and

[0224] xe611 to xe616 may be each independently selected from 0, 1, 2, and 3.


[0225] The compound of Formula 601 and the compound of Formula 602 may be each independently selected from Compounds ET1 to ET15:

-continued


ET1




ET4




ET2




ET5



ET3




ET6




-continued


ET7



ET8




ET9




-continued


ET10



ET11





ET12




-continued

ET13


ET-D1




ET-D2



ET14



ET15



[0226] A thickness of the ETL may be about 100Å to about 1,000 Å, e.g., about 150 Å to about 500 Å. When the thickness of the ETL is within these ranges, the ETL may have satisfactory electron transporting ability without a substantial increase in driving voltage.

[0227] In an implementation, the ETL may further include a metal-containing material, in addition to the above-described materials.

[0228] The metal-containing material may include a lithium (Li) complex. Non-limiting examples of the Li complex may include compound ET-D1 (lithium quinolate (LiQ)), or compound ET-D2.

[0229] The electron transport region may include an EIL that may facilitate injection of electrons from the second electrode **190**.

[0230] The EIL may be formed on the ETL by using any of a variety of methods, e.g., by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like. When the EIL is formed using vacuum deposition or spin coating, the deposition and coating conditions for forming the EIL may be similar to the above-described deposition and coating conditions for forming the HIL, and accordingly may not be described in detail.

[0231] The EIL may include at least one selected from LiF, NaCl, CsF, Li₂O, BaO, and LiQ.

[0232] A thickness of the EIL may be about 1Å to about 100 Å, e.g., about 3Å to about 90Å. When the thickness of the EIL is within these ranges, the EIL may have satisfactory electron injection ability without a substantial increase in driving voltage.

[0233] The second electrode **190** may be disposed on the organic layer **150**, as described above. The second electrode **190** may be a cathode as an electron injecting electrode. A material for forming the second electrode **190** may be, e.g., a metal, an alloy, an electrically conductive compound, which have a low-work function, or a mixture thereof. Non-limiting examples of materials for forming the second electrode **190** may include lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag). In an implementation, a material for forming the second electrode **190** may be ITO or IZO. The second electrode **190** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode.

[0234] Although the organic light-emitting device of FIG. 1 is described above, embodiments are not limited thereto.

[0235] As used herein, a C₁-C₆₀ alkyl group refers to a monovalent linear or branched aliphatic hydrocarbon group having 1 to 60 carbon atoms. Non-limiting examples of the C₁-C₆₀ alkyl group are a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. A C₁-C₆₀ alkylene group refers to a divalent group having the same structure as the C₁-C₆₀ alkyl group.

[0236] As used herein, a C_1 - C_{60} alkoxy group refers to a monovalent group represented by $-OA_{101}$ (where A_{101} is a C_1 - C_{60} alkyl group as described above). Non-limiting examples of the C_1 - C_{60} alkoxy group are a methoxy group, an ethoxy group, and an isopropoxy group.

[0237] As used herein, a C_2 - C_{60} alkenyl group refers to a hydrocarbon group including at least one carbon double bond in the middle or terminal of the C_2 - C_{60} alkyl group. Non-limiting examples of the C_2 - C_{60} alkenyl group are an ethenyl group, a propenyl group, and a butenyl group. A C_2 - C_{60} alkylene group refers to a divalent group having the same structure as the C_2 - C_{60} alkenyl group.

[0238] As used herein, a C_2 - C_{60} alkynyl group refers to a hydrocarbon group including at least one carbon triple bond in the middle or terminal of the C_2 - C_{60} alkyl group. Non-limiting examples of the C_2 - C_{60} alkynyl group are an ethynyl group and a propynyl group. A C_2 - C_{60} alkynylene group refers to a divalent group having the same structure as the C_2 - C_{60} alkynyl group.

[0239] As used herein, a C_3 - C_{10} cycloalkyl group refers to a monovalent, monocyclic saturated hydrocarbon group having 3 to 10 carbon atoms. Non-limiting examples of the C_3 - C_{10} cycloalkyl group are a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. A C_3 - C_{10} cycloalkylene group refers to a divalent group having the same structure as the C_3 - C_{10} cycloalkyl group.

[0240] As used herein, a C_1 - C_{10} heterocycloalkyl group refers to a monovalent monocyclic group having 1 to 10 carbon atoms in which at least one hetero atom selected from N, O, P, and S is included as a ring-forming atom. Non-limiting examples of the C_1 - C_{10} heterocycloalkyl group are a tetrahydrofuranyl group and a tetrahydrothiophenyl group. A C_1 - C_{10} heterocycloalkylene group refers to a divalent group having the same structure as the C_1 - C_{10} heterocycloalkyl group.

[0241] As used herein, a C_3 - C_{10} cycloalkenyl group refers to a monovalent monocyclic group having 3 to 10 carbon atoms that includes at least one double bond in the ring but does not have aromaticity. Non-limiting examples of the C_3 - C_{10} cycloalkenyl group are a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. A C_3 - C_{10} cycloalkenylene group refers to a divalent group having the same structure as the C_3 - C_{10} cycloalkenyl group.

[0242] As used herein, a C_1 - C_{10} heterocycloalkenyl group refers to a monovalent monocyclic group having 1 to 10 carbon atoms that includes at least one double bond in the ring and in which at least one hetero atom selected from N, O, P, and S is included as a ring-forming atom. Non-limiting examples of the C_1 - C_{10} heterocycloalkenyl group are a 2,3-dihydrofuranyl group and a 2,3-dihydrothiophenyl group. A C_1 - C_{10} heterocycloalkenylene group refers to a divalent group having the same structure as the C_1 - C_{10} heterocycloalkenyl group.

[0243] As used herein, a C_6 - C_{60} aryl group refers to a monovalent, aromatic carbocyclic group having 6 to 60 carbon atoms, and a C_6 - C_{60} arylene group refers to a divalent, aromatic carbocyclic group having 6 to 60 carbon atoms. Non-limiting examples of the C_6 - C_{60} aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C_6 - C_{60} aryl group and the C_6 - C_{60} arylene group include at least two rings, the rings may be fused to each other.

[0244] As used herein, a C_1 - C_{60} heteroaryl group refers to a monovalent, aromatic carbocyclic group having 1 to 60 carbon atoms in which at least one hetero atom selected from N, O, P, and S is included as a ring-forming atom, and 1 to 60 carbon atoms. A C_1 - C_{60} heteroarylene group refers to a divalent, aromatic carbocyclic group having 1 to 60 carbon atoms in which at least one hetero atom selected from N, O, P, and S is included as a ring-forming atom. Non-limiting examples of the C_1 - C_{60} heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C_1 - C_{60} heteroaryl group and the C_1 - C_{60} heteroarylene group include at least two rings, the rings may be fused to each other.

[0245] As used herein, a C_6 - C_{60} aryloxy group indicates $-OA_{102}$ (where A_{102} is a C_6 - C_{60} aryl group as described above), and a C_6 - C_{60} arylthio group indicates $-SA_{103}$ (where A_{103} is a C_6 - C_{60} aryl group as described above).

[0246] As used herein, a monovalent non-aromatic condensed polycyclic group refers to a monovalent group having at least two rings condensed to each other, in which only carbon atoms (for example, 8 to 60 carbon atoms) are exclusively included as ring-forming atoms and the entire molecule represents non-aromaticity. A non-limiting example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. A divalent non-aromatic condensed polycyclic group refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0247] As used herein, a monovalent non-aromatic condensed heteropolycyclic group refers to a monovalent group having at least two rings condensed to each other, in which carbon atoms (for example, 2 to 60 carbon atoms) and a hetero atom selected from N, O, P, and S are ring-forming atoms and the entire molecule represents non-aromaticity. A non-limiting example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. A divalent non-aromatic condensed heteropolycyclic group refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0248] As used herein, at least one substituent of the substituted C_3 - C_{10} cycloalkylene group, the substituted C_1 - C_{10} heterocycloalkylene group, the substituted C_3 - C_{10} cycloalkenylene group, the substituted C_1 - C_{10} heterocycloalkenylene group, the substituted C_6 - C_{60} arylene group, the substituted C_1 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_1 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0249] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt

thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group,

[0250] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —B(Q₁₃)(Q₁₄), and —Si(Q₁₅)(Q₁₆)(Q₁₇),

[0251] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group,

[0252] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —B(Q₂₃)(Q₂₄), and —Si(Q₂₅)(Q₂₆)(Q₂₇), and

[0253] —N(Q₃₁)(Q₃₂), —B(Q₃₃)(Q₃₄), and —Si(Q₃₅)(Q₃₆)(Q₃₇),

[0254] wherein Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0255] For example, at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted

C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0256] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group,

[0257] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spirofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovarenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quiazolinyl group, a cinolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —N(Q₁₁)(Q₁₂), —B(Q₁₃)(Q₁₄), and —Si(Q₁₅)(Q₁₆)(Q₁₇),

[0258] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naph-

thyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovarenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group,

[0259] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovarenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopen-

tenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovarenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —N(Q₂₁)(Q₂₂), —B(Q₂₃)(Q₂₄), and —Si(Q₂₅)(Q₂₆)(Q₂₇), and

[0260] —N(Q₃₁)(Q₃₂), —B(Q₃₃)(Q₃₄), and —Si(Q₃₅)(Q₃₆)(Q₃₇),

[0261] wherein Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovarenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group,

a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.

[0262] The acronym “Ph” used herein refers to a phenyl group, the acronym “Me” used herein refers to a methyl group, the acronym “Et” used herein refers to an ethyl group, and the acronym “ter-Bu” or “Bu” used herein refers to a tert-butyl group.

[0263] One or more embodiments, which include condensed cyclic compounds, and organic light-emitting devices including the same, will now be described in detail with reference to the following examples. In the following synthesis example(s), the expression that “‘B’ instead of ‘A’ was used” means that the amounts of ‘B’ and ‘A’ were the same in equivalent amounts.


[0264] The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples.

EXAMPLES


Synthesis Example 1

Synthesis of Compound 5 (TPDSiN)

[0265]



-continued



Synthesis of Compound (A)

[0266] 5 g (17.5 mmol) of 2,3-dibromonaphthalene and 150 mL of a mixed solvent of anhydrous THF and Et₂O (1:1) were put into a 500-mL 3-necked flask, and then cooled down to about -110° C. in a cooling bath (EtOH, Et₂O/Liq-N₂ bath.), followed by slowly adding 7.2 mL (2.88 mmol) of n-BuLi (2.5 M) thereto. The resulting reaction mixture was stirred under a nitrogen atmosphere for about 30 minutes, and then a dilute solution of 2.2 g (8.75 mmol) of dichlorodiphenylsilane in 80 mL of a mixed solvent of anhydrous THF and Et₂O (1:1) was slowly dropwise added to the reaction mixture, followed by stirring at the same temperature for about 30 minutes. After slowly raising temperature to ambient temperature, the reaction mixture was stirred for 12 hours, followed by adding distilled water to terminate the reaction, extracting with diethyl ether three times (100 mL each time), drying with anhydrous magnesium sulfate (MgSO₄), and removing the solvent with a rotary evaporator. The residue was separated by column chromatography (n-hexane/dichloromethane, 15:1) to obtain 1.15 g of Compound (A) as a white solid (Yield: 22%).

[0267] ¹H NMR (300 MHz, DMSO-d₆) δ 7.42-7.51 (m, 14H), 7.86-7.84 (m, 6H), 8.63 (s, 2H).

Synthesis of Compound 5 (TPDSiN)

[0268] 1.15 g (1.94 mmol) of Compound (A) and 50 mL of anhydrous THF were put into a 100-mL 3-necked flask, and then cooled down to about -78° C. in a cooling bath (EtOAc/Liq-N₂ bath.), followed by adding 1.6 mL (0.64 mmol) of n-BuLi (2.5 M) thereto. The resulting reaction mixture was stirred under a nitrogen atmosphere for about 1 hour, and then a dilute solution of 0.98 g (3.88 mmol) of dichlorodiphenylsilane in 20 mL of anhydrous THF was slowly dropwise added to the reaction mixture, followed by stirring at the same temperature for about 30 minutes. After slowly raising temperature to ambient temperature, the reaction mixture was stirred for 24 hours, followed by adding 30 mL of distilled water to terminate the reaction, extracting with Et₂O three times (50 mL each time), drying with anhydrous magnesium sulfate (MgSO₄), and removing the solvent with a rotary evaporator. The residue was separated by column chromatography (n-hexane/ethyl acetate, 20:1) to obtain 0.61 g of Compound 5 (TPDSiN) as a white solid (Yield: 51%).

[0269] ^1H NMR (300 MHz, DMSO- d_6) δ 7.55-7.65 (m, 14H), 7.88-7.96 (m, 14H), 8.21 (d, 4H); HRMS (FAB+) m/z 616.21.

Evaluation Example 1

Characteristics Evaluation of Compound 5

[0270] Evaluation of Spectroscopic Characteristics of Compound 5

[0271] UV absorption and photoluminescence (PL) spectra of Compound 5, and low-temperature PL spectrum thereof were measured using the methods described in Table 1. The results are shown in FIG. 2.

TABLE 1

“UV_solution” spectrum	Compound 5 was dissolved in CHCl_3 to a concentration of 1×10^{-5} M, and then UV absorption spectrum thereof were measured at ambient temperature using a Shimadzu UV-350 Spectrometer.
“PL_solution” spectrum	A solution of Compound 5 in CHCl_3 to a concentration of 1×10^{-5} M was subjected to PL spectrum measurement at ambient temperature using an ISC PC1 Spectrofluorometer equipped with a Xenon lamp.
“PL_low temperature” spectrum	A solution of Compound 5 in CHCl_3 to a concentration of 1×10^{-5} M was subjected to PL spectrum measurement at a low temperature (77 K) using an ISC PC1 Spectrofluorometer equipped with a Xenon lamp.

[0272] Referring to FIG. 2, Compound 5 exhibited suitable spectroscopic characteristics for use as a material for organic light-emitting devices.

[0273] Evaluation of Electrochemical Characteristics of Compound 5

[0274] Electrochemical characteristics of Compound 5 were measured using cyclic voltammetry (CV) (Electrolyte: 0.1 M Bu_4NClO_4 /Solvent: CH_2Cl_2 /Electrode: 3-electrode system (working electrode: GC, reference electrode: Ag/AgCl, auxiliary electrode; Pt)). The results are shown in FIG. 3.

[0275] Referring to FIG. 3, Compound 5 exhibited suitable electrical electrochemical characteristics for use as a material for organic light-emitting devices.

[0276] Evaluation of HOMO and LUMO Energy Levels of Compound 5

[0277] A lowest unoccupied molecular orbital (LUMO) energy level of Compound 5 was calculated using a reduction onset in FIG. 3. A highest occupied molecular orbital (HOMO) energy level of Compound 5 was calculated using an optical band gap (E_g) based on the UV absorption edge in FIG. 2. The results are shown in Table 2. For comparison, data obtained from the UV absorption and PL spectra are also shown in Table 2. The HOMO and LUMO energy levels represent absolute values.

TABLE 2

UV (nm)	PL (nm)	E_T (eV)	E_g (eV)	HOMO (eV)	LUMO (eV)
250, 273, 327, 341	364	3.01	3.52	5.81	2.29


[0278] Referring to Table 2, Compound 5 was found to have a HOMO-LUMO energy gap that is suitable for used as a material, e.g., a host material, for organic light-emitting devices.

Example 1

[0279] A $15 \Omega/\text{cm}^2$ (1,000 Å) ITO glass substrate (available from Corning Co., Ltd) was cut to a size of 50 mm×50 mm×0.7 mm and then sonicated in isopropyl alcohol and pure water each for 5 minutes, and then cleaned by irradiation of ultraviolet rays for 30 minutes and exposure to ozone. The resulting ITO glass substrate was mounted into a vacuum deposition apparatus.


[0280] After DNTPD was deposited on the ITO anode to form an HIL having a thickness of about 600Å, NPB was deposited on the HIL to form a HTL having a thickness of about 100Å, mCP was deposited on the HTL to form an electron blocking layer (EBL) having a thickness of about 200Å, and then Compound 5 (host) and Fir6 (dopant) were co-deposited in a weight ratio of 90:10 to form an EML having a thickness of about 400Å on the EBL.

[0281] Next, TSPO1 was deposited on the EML to form a first ETL having a thickness of about 300Å, and then TmPyPb was deposited on the first ETL to form a second ETL having a thickness of about 250Å. Then, LiF was deposited on the second ETL to form an EIL having a thickness of about 10Å, and Al was then deposited on the EIL to form a cathode having a thickness of about 1000Å, thereby manufacturing an organic light-emitting device.




DNTPD


-continued




NPB




mCP



FIr6



TSP01



TmPyPb


Example 2

[0282] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 1, instead of Compound 5, was used as a host to form the EML.

Comparative Example 1


[0283] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound A, below, instead of Compound 5, was used as a host to form the EML.

Compound A



Comparative Example 2

[0284] An organic light-emitting device was manufactured in the same manner as in Example 1, except that CBP, below, instead of Compound 5, was used as a dopant to form the EML.



CBP

Evaluation Example 2

[0285] Driving voltages, current densities, luminances, efficiencies, and half-lifetimes of the organic light-emitting devices of Examples 1 and 2 and Comparative Examples 1 and 2 were evaluated using a Kethley Source-Measure Unit (SMU 236) and a PR650 (Spectroscan) Source Measurement Unit. (available from Photo Research, Inc.). The results are shown in Table 3.

TABLE 3

Example	Host of EML	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	CIE x
Example 1	Compound 5	6.5	4.16	500	12	0.18

TABLE 3-continued

Example	Host of EML	Driving voltage (V)	Current density (mA/cm ²)	Luminescence (cd/m ²)	Efficiency (cd/A)	CIE x
Example 2	Compound 1	7.0	4.54	500	11	0.17
Comparative Example 1	Compound A	7.8	7.14	500	7	0.18
Comparative Example 2	CBP	7.5	8.33	500	6	0.19


[0286] Referring to Table 3, it may be seen that the organic light-emitting devices of Examples 1 and 2 had lower driving voltages and improved efficiency characteristics, as compared to the organic light-emitting devices of Comparative Examples 1 and 2.

[0287] As described above, according to the one or more of the above embodiments, an organic light-emitting device including the condensed cyclic compounds of Formula 1 may have a low driving voltage, a high efficiency and a high luminance.

[0288] Example embodiments have been disclosed herein, and although specific terms are employed, they are used and are to be interpreted in a generic and descriptive sense only and not for purpose of limitation. In some instances, as would be apparent to one of ordinary skill in the art as of the filing of the present application, features, characteristics, and/or elements described in connection with a particular embodiment may be used singly or in combination with features, characteristics, and/or elements described in connection with other embodiments unless otherwise specifically indicated. Accordingly, it will be understood by those of skill in the art that various changes in form and details may be made without departing from the spirit and scope of the present invention as set forth in the following claims.

What is claimed is:

1. A condensed cyclic compound represented by Formula 1:



wherein, in Formula 1,

A₁ and A₂ are each independently a C₆-C₂₀ aromatic ring or a C₂-C₂₀ heteroaromatic ring, provided that A₁ and A₂ are not both a benzene ring at the same time;

L₁ to L₆ are each independently selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or

unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

a₁ to a₆ are each independently selected from 0, 1, 2, and 3;

R₁ to R₆ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —B(Q₃)(Q₄), and —Si(Q₅)(Q₆)(Q₇);

b₁ to b₆ are each independently selected from 0, 1, 2 and 3;

c₁ and c₂ are each independently selected from 0, 1, 2, and 3;

wherein at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt

thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —B(Q₁₃)(Q₁₄), and —Si(Q₁₅)(Q₁₆)(Q₁₇);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —B(Q₂₃)(Q₂₄), and —Si(Q₂₅)(Q₂₆)(Q₂₇); and
—N(Q₃₁)(Q₃₂), —B(Q₃₃)(Q₃₄), and —Si(Q₃₅)(Q₃₆)(Q₃₇);

wherein Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent aromatic condensed heteropolycyclic group.

2. The condensed cyclic compound as claimed in claim 1, wherein:

the C₆-C₂₀ aromatic ring is a benzene, a naphthalene, or an anthracene; and

the C₁-C₂₀ heteroaromatic ring is a pyridine, a pyridazine, a pyrimidine, a pyrazine, a triazine, a quinoline, an isoquinoline, a phthalazine, a naphthyridine, a quinox-

alaine, a quinazoline, a cinnoline, a benzothiophene, a benzofuran, a dibenzothiophene, or a dibenzofuran.

3. The condensed cyclic compound as claimed in claim 1, wherein:

A₁ is a benzene, a naphthalene, or an anthracene; and

A₂ is a naphthalene or an anthracene.

4. The condensed cyclic compound as claimed in claim 1, wherein A₁ and A₂ are each independently a pyridine, a pyridazine, a pyrimidine, a pyrazine, a triazine, a quinoline, an isoquinoline, a phthalazine, a naphthyridine, a quinoxalaine, a quinazoline, a cinnoline, a benzothiophene, a benzofuran, a dibenzothiophene, or a dibenzofuran.

5. The condensed cyclic compound as claimed in claim 1, wherein:

A₁ is a benzene, a naphthalene, an anthracene, a pyridine, a pyridazine, a pyrimidine, a pyrazine, a triazine, a quinoline, an isoquinoline, a phthalazine, a naphthyridine, a quinoxalaine, a quinazoline, or a cinnoline; and

A₂ is a naphthalene, an anthracene, a pyridine, a pyridazine, a pyrimidine, a pyrazine, a triazine, a quinoline, an isoquinoline, a phthalazine, a naphthyridine, a quinoxalaine, a quinazoline, or a cinnoline.

6. The condensed cyclic compound as claimed in claim 1, wherein L₁ to L₆ in Formula 1 are each independently selected from:


a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spirofluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isooxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzoimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzooxazolylene group, an isobenzooxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a diazopyridinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group;

and
a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an


acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylene group, a thiophenylenylene group, a furanylene group, an imidazolylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylene group, an isooxazolylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylenylene group, an indolylenylene group, an indazolylene group, a purinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a carbazolylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzoimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylenylene group, a benzooxazolylene group, an isobenzooxazolylene group, a triazolylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, and an imidazopyrimidinylenylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, an a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group,

an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.


7. The condensed cyclic compound as claimed in claim 1, wherein L₁ to L₆ in Formula 1 are each independently a group represented by one of Formulae 3-1 to 3-32:




Formula 3-1




Formula 3-2




Formula 3-3




Formula 3-4



Formula 3-5













Formula 3-6




Formula 3-7


-continued




Formula 3-8




Formula 3-9




Formula 3-10




Formula 3-11




Formula 3-12




Formula 3-13




Formula 3-14




Formula 3-15



Formula 3-16



Formula 3-17



-continued

Formula 3-18

Formula 3-19

Formula 3-20

Formula 3-21

Formula 3-22

Formula 3-23


Formula 3-24

Formula 3-25


Formula 3-26

Formula 3-27


-continued




Formula 3-28




Formula 3-29



Formula 3-30



Formula 3-31



Formula 3-32

d3 is an integer selected from 1 to 6;


d4 is an integer selected from 1 to 8;

d5 is 1 or 2;


d6 is an integer selected from 1 to 5; and

* and *' are binding sites with adjacent atoms.


8. The condensed cyclic compound as claimed in claim 1, wherein L₁ to L₆ in Formula 1 are each independently a group represented by one of Formulae 4-1 to 4-23:




Formula 4-1




Formula 4-2




Formula 4-3




Formula 4-4




Formula 4-5



Formula 4-6



Formula 4-7



Formula 4-8

wherein, in Formulae 3-1 to 3-32,








Y₁ is O, S, C(Z₃)(Z₄), N(Z₅), or Si(Z₆)(Z₇);

Z₁ to Z₇ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino groups, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group;

d1 is an integer selected from 1 to 4;

d2 is an integer selected from 1 to 3;

-continued



Formula 4-9

Formula 4-10

Formula 4-11









Formula 4-12

Formula 4-13

Formula 4-14

Formula 4-15

-continued



Formula 4-16

Formula 4-17

Formula 4-18

Formula 4-19

Formula 4-20

Formula 4-21

Formula 4-22

Formula 4-23

wherein, in Formulae 4-1 and 4-23, * and *' are binding sites with adjacent atoms.

9. The condensed cyclic compound as claimed in claim 1, wherein a1 to a6 in Formula 1 are each independently 0 or 1.

10. The condensed cyclic compound as claimed in claim 1, wherein R₁ to R₆ in Formula 1 are each independently

selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C₁-C₂₀ alkyl group, a substituted or unsubstituted C₁-C₂₀ alkoxy group, a substituted or unsubstituted C₆-C₂₀ aryl group, a substituted or unsubstituted C₁-C₂₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and —Si(Q₅)(Q₆)(Q₇).

11. The condensed cyclic compound as claimed in claim 1, wherein R₁ to R₆ in Formula 1 are each independently selected from:

a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a benzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a benzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

nyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a benzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a benzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

—Si(Q₅)(Q₆)(Q₇).

12. The condensed cyclic compound as claimed in claim 1, wherein R₁ to R₆ in Formula 1 are each independently selected from:


a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, a phenanthrolyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothienophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and


a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, a phenanthrolyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothienophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, a phenanthrolyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothienophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.

quinazoliny group, a carbazolyl group, a phenanthrolyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothienophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.


13. The condensed cyclic compound as claimed in claim 1, wherein R₁ to R₆ in Formula 1 are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, and a group represented by one of Formulae 6-1 to 6-49:




Formula 6-1




Formula 6-2



Formula 6-3









Formula 6-4



Formula 6-5

-continued



Formula 6-6

Formula 6-7










Formula 6-8

Formula 6-9

Formula 6-10

Formula 6-11

-continued



Formula 6-12

Formula 6-13

Formula 6-14

Formula 6-15

Formula 6-16












Formula 6-17

Formula 6-18

Formula 6-19

Formula 6-20

-continued



Formula 6-21

Formula 6-22

Formula 6-23

Formula 6-24

Formula 6-25

Formula 6-26

Formula 6-27









Formula 6-28

Formula 6-29

Formula 6-30

Formula 6-31

-continued



Formula 6-32

Formula 6-33

Formula 6-34

Formula 6-35






Formula 6-36

Formula 6-37

Formula 6-38

Formula 6-39

-continued



Formula 6-40






Formula 6-41

Formula 6-42

Formula 6-43

Formula 6-44

-continued



Formula 6-45

Formula 6-46

Formula 6-47


Formula 6-48

Formula 6-49





wherein, in Formulae 6-1 to 6-49, * is a binding site with an adjacent atom.

14. The condensed cyclic compound as claimed in claim 1, wherein the condensed cyclic compound represented by Formula 1 is represented by one of Formulae 1A to 1E:


<Formula 1A>



-continued



<Formula 1B>



<Formula 1C>

<Formula 1D>

<Formula 1E>


<Formula 1A(1)>

<Formula 1A(2)>

<Formula 1A(3)>


wherein, in Formulae 1A to 1E, L₁ to L₆, a₁ to a₆, R₁ to R₆, b₁ to b₆, c₁, and c₂ are the same as those defined with respect to Formula 1.

15. The condensed cyclic compound as claimed in claim 1, wherein the condensed cyclic compound represented by Formula 1 is represented by one of Formulae 1A(1) to 1A(5), Formulae 2A(1) to 2A(5), Formula 3A(1), and Formula 4A(1):




-continued

<Formula 1A(4)>




-continued


<Formula 2A(2)>



<Formula 1A(5)>





<Formula 2A(3)>




<Formula 2A(4)>

<Formula 2A(1)>




-continued


<Formula 2A(5)>




<Formula 3A(1)>




<Formula 4A(1)>




16. The condensed cyclic compound as claimed in claim 1, wherein the condensed cyclic compound represented by Formula 1 is one of Compounds 1 to 7:




1




2



3




4



5

wherein, in Formulae 1A(1) to 1A(5), Formulae 2A(1) to 2A(5), Formula 3A(1), and Formula 4A(1), R₁, R₂, b1, and b2 are the same as those defined with respect to Formula 1, R_{3a} to R_{3g} are each independently selected from the same groups as those defined for R₃ in Formula 1, R_{4a} to R_{4g} are each independently selected from the same groups as those defined for R₄ in Formula 1, R_{5a} to R_{5g} are each independently selected from the same groups as those defined for R₅ in Formula 1, and R_{6a} to R_{6g} are each independently selected from the same groups as those defined for R₆ in Formula 1.

-continued



- 6 **17.** An organic light-emitting device comprising:
 a first electrode;
 a second electrode opposite to the first electrode; and
 an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes at least one condensed cyclic compound as claimed in claim 1.
- 18.** The organic light-emitting device as claimed in claim 17, wherein:
 the first electrode is an anode,
 the second electrode is a cathode, and
 the organic layer includes:
- 7 a hole transport region between the first electrode and the emission layer, the hole transport region including at least one of a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer, and
 an electron transport region between the emission layer and the second electrode, the electron transport region including at least one of a hole blocking layer, an electron transport layer, and an electron injection layer.
- 19.** The organic light-emitting device as claimed in claim 17, wherein the condensed cyclic compound is included in the emission layer.
- 20.** The organic light-emitting device as claimed in claim 19, wherein the emission layer further includes a phosphorescent dopant, and the condensed cyclic compound is a host.

* * * * *

专利名称(译)	缩合环状化合物和包括其的有机发光器件		
公开(公告)号	US20160043333A1	公开(公告)日	2016-02-11
申请号	US14/618835	申请日	2015-02-10
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO., LTD. 庆尚大学产学合作基础		
当前申请(专利权)人(译)	三星DISPLAY CO., LTD. 产学合作基础庆尚大学		
[标]发明人	KIM MIKYUNG KIM YUNHI HWANG JAEYOUNG		
发明人	KIM, MIKYUNG KIM, YUNHI HWANG, JAEYOUNG		
IPC分类号	H01L51/00 C07F7/08 H01L51/50		
CPC分类号	H01L51/0094 H01L51/5028 C07F7/0807 H01L51/0067 H01L51/0085 H01L51/5016		
优先权	1020140100696 2014-08-05 KR		
其他公开文献	US9818960		
外部链接	Espacenet USPTO		

摘要(译)

缩合环状化合物和有机发光器件，缩合环状化合物由式1表示：

