





US 20160172603A1

(19) **United States**(12) **Patent Application Publication**
SHIN et al.(10) **Pub. No.: US 2016/0172603 A1**(43) **Pub. Date: Jun. 16, 2016**(54) **CONDENSED-CYCLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**(71) Applicant: **SAMSUNG DISPLAY CO., LTD.,**
Yongin-City (KR)(72) Inventors: **Dongwoo SHIN**, Yongin-City (KR);
Seulong KIM, Yongin-City (KR);
Younsun KIM, Yongin-City (KR);
Jungsub LEE, Yongin-City (KR);
Naoyuki ITO, Yongin-City (KR)(21) Appl. No.: **14/731,011**(22) Filed: **Jun. 4, 2015**(30) **Foreign Application Priority Data**

Dec. 10, 2014 (KR) 10-2014-0177821

Publication Classification(51) **Int. Cl.****H01L 51/00** (2006.01)**C09K 11/02** (2006.01)**C07D 403/04** (2006.01)**C07D 401/02** (2006.01)**C07D 223/14** (2006.01)**C09K 11/06** (2006.01)(52) **U.S. Cl.**CPC **H01L 51/0072** (2013.01); **C07D 223/14**
(2013.01); **C09K 11/06** (2013.01); **C07D**
403/04 (2013.01); **C07D 401/02** (2013.01);
C09K 11/025 (2013.01); **H01L 51/0054**
(2013.01); **H01L 51/0085** (2013.01); **H01L**
51/0058 (2013.01); **H01L 51/0067** (2013.01);
H01L 51/0056 (2013.01)(57) **ABSTRACT**

A condensed cyclic compound and an organic light-emitting device, the compound being represented by Formula 1:

 $(A_1)_{a1}-(L_1)_{b1}-(A_2)_{a2}$ <Formula 1>wherein, in Formula 1, A_1 is a group represented by Formula 2-1, below, a_1 is 1, 2, or 3, and, when a_1 is 2 or greater, two or more A_1 s are identical to or different from each other, A_2 is a group represented by Formula 2-2, below, a_2 is 1, 2, or 3, and, when a_2 is 2 or greater, two or more A_2 s are identical to or different from each other,10

190
150
110

FIG. 1

10

190
150
110

CONDENSED-CYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

[0001] Korean Patent Application No. 10-2014-0177821, filed on Dec. 10, 2014, in the Korean Intellectual Property Office, and entitled: “Condensed-Cyclic Compound and Organic Light-Emitting Device Including the Same,” is incorporated by reference herein in its entirety.

BACKGROUND

[0002] 1. Field

[0003] Embodiments relate to a condensed cyclic compound and an organic light-emitting device including the condensed cyclic compound.

[0004] 2. Description of the Related Art


[0005] Organic light-emitting devices (OLEDs) are self-emission devices that have wide viewing angles, high contrast ratios, and short response times. In addition, the OLEDs exhibit excellent luminance, driving voltage, and response speed characteristics and produce full-color images.

[0006] An organic light-emitting device may include a first electrode on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. The holes and the electrons may be recombined in the emission layer to produce excitons. These excitons change from an excited state to a ground state, thereby generating light.

SUMMARY

[0007] Embodiments are directed to a condensed cyclic compound and an organic light-emitting device including the condensed cyclic compound.


[0008] According to one or more exemplary embodiments, a condensed cyclic compound is represented by Formula 1:



<Formula 2-1>

-continued

<Formula 2-2>



[0009] wherein, in Formulae 1, 2-1, and 2-2,

[0010] A_1 is a group represented by Formula 2-1, a_1 is 1, 2, or 3, and, when a_1 is 2 or greater, two or more A_1 s are identical to or different from each other,

[0011] A_2 is a group represented by Formula 2-2, a_2 is 1, 2, or 3, and, when a_2 is 2 or greater, two or more A_2 s are identical to or different from each other,

[0012] L_1 and L_2 are each independently selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group, where L_2 is not a substituted or unsubstituted triphenylenylene group,

[0013] b_1 and b_2 are each independently selected from 0, 1, 2, and 3; when b_1 or b_2 is 0, $*(L_1)_{b1}-*$ or $*(L_2)_{b2}-*$ is a single bond, when b_1 is 2 or greater, two or more L_1 s are identical to or different from each other, and when b_2 is 2 or greater, two or more L_2 s are identical to or different from each other,

[0014] Ar_1 is selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, where Ar_1 is not a substituted or unsubstituted triphenylenyl group,

[0015] R_1 to R_{12} and R_{21} to R_{32} are each independently selected from a binding site to one of A_1 , L_1 , and A_2 in Formula 1, a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10}

heterocycloalkenyl group, a substituted or unsubstituted C_6-C_{60} aryl group, a substituted or unsubstituted C_6-C_{60} aryloxy group, a substituted or unsubstituted C_6-C_{60} arylthio group, a substituted or unsubstituted C_1-C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_1)(Q_2)(Q_3)$, $-B(Q_4)(Q_5)$, and $N(Q_6)(Q_7)$, where a1 of R_1 to R_{12} are each independently a binding site to L_1 or A_2 in Formula 1, and a2 of R_{21} to R_{32} are each independently a binding site to L_1 or A_2 in Formula 1;

[0016] at least one substituent of the substituted C_3-C_{10} cycloalkylene group, substituted C_1-C_{10} heterocycloalkylene group, substituted C_3-C_{10} cycloalkenylene group, substituted C_1-C_{10} heterocycloalkenylene group, substituted C_6-C_{60} arylene group, substituted C_1-C_{60} heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C_1-C_{60} alkyl group, substituted C_2-C_{60} alkenyl group, substituted C_2-C_{60} alkynyl group, substituted C_1-C_{60} alkoxy group, substituted C_3-C_{10} cycloalkyl group, substituted C_1-C_{10} heterocycloalkyl group, substituted C_3-C_{10} cycloalkenyl group, substituted C_1-C_{10} heterocycloalkenyl group, substituted C_6-C_{60} aryl group, substituted C_6-C_{60} aryloxy group, substituted C_6-C_{60} arylthio group, substituted C_1-C_{60} heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from

[0017] a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, and a C_1-C_{60} alkoxy group;

[0018] a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, and a C_1-C_{60} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{11})(Q_{12})(Q_{13})$, $-B(Q_{14})(Q_{15})$, and $-N(Q_{16})(Q_{17})$;

[0019] a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0020] a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a

deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_1-C_{60} alkoxy group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{21})(Q_{22})(Q_{23})$, $-B(Q_{24})(Q_{25})$, and $-N(Q_{26})(Q_{27})$; and

[0021] $-Si(Q_{31})(Q_{32})(Q_{33})$, $-B(Q_{34})(Q_{35})$, and $-N(Q_{36})(Q_{37})$;

[0022] wherein Q_1 to Q_7 , Q_{11} to Q_{17} , Q_{21} to Q_{27} , and Q_{31} to Q_{37} are each independently selected from a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_1-C_{60} alkoxy group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0023] According to one or more exemplary embodiments, an organic light-emitting device includes a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode and includes an emission layer, wherein the organic layer includes at least one type of the condensed cyclic compound.

BRIEF DESCRIPTION OF THE DRAWING

[0024] Features will be apparent to those of skill in the art by describing in detail exemplary embodiments with reference to the attached drawing in which:

[0025] FIG. 1 illustrates a schematic view of a structure of an organic light-emitting device according to an exemplary embodiment.

DETAILED DESCRIPTION

[0026] Example embodiments will now be described more fully hereinafter with reference to the accompanying drawing; however, they may be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey exemplary implementations to those skilled in the art.


[0027] In the drawing FIGURE, the dimensions of layers and regions may be exaggerated for clarity of illustration. Like reference numerals refer to like elements throughout.

[0028] As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.


[0029] According to one or more exemplary embodiments, a condensed cyclic compound may be represented by Formula 1.



[0030] In Formula 1, A_1 may be a group represented by Formula 2-1, and A_2 may be a group represented by Formula 2-2.



<Formula 2-1>



<Formula 2-2>

[0031] In Formula 1, a_1 denotes the number of A_1 s, and a_2 denotes the number of A_2 s, where a_1 and a_2 may each independently be selected from 1, 2, and 3, and when a_1 is 2 or greater, two or more A_1 s may be identical to or different from each other, and when a_2 is 2 or greater, two or more A_2 s may be identical to or different from each other.

[0032] In some embodiments, in Formula 1, a_1 and a_2 may each independently be 1 or 2.

[0033] In some embodiments, in Formula 1, a_1 and a_2 may each independently be 1 or 2, and $a_1 + a_2$ may be 3.

[0034] In some embodiments, in Formula 1, a_1 and a_2 each be 1.

[0035] In Formulae 1 and 2-1, L_1 and L_2 may each independently be selected from or include, e.g., a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group. In an implementation, L_2 may not be a substituted or unsubstituted triphenylenylene group.

[0036] In some embodiments, in Formulae 1 and 2-1, L_1 and L_2 may each independently be selected from, e.g.,


[0037] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a

heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylenylene group, a spiro-fluorenylenylene group, a benzofluorenylenylene group, a dibenzofluorenylenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylenylene group, an indolylenylene group, an indazolylenylene group, a purinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzoimidazolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, and an imidazopyrimidinylenylene group; and


[0038] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylenylene group, a spiro-fluorenylenylene group, a benzofluorenylenylene group, a dibenzofluorenylenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylenylene group, an indolylenylene group, an indazolylenylene group, a purinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzoimidazolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, and an imidazopyrimidinylenylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino

group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenaziny group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzo-carbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.


[0039] In one embodiment, L_1 and L_2 may each independently be a group represented by one of the following Formulae 3-1 to 3-38.




Formula 3-1




Formula 3-2



Formula 3-3




Formula 3-4




-continued


Formula 3-5




Formula 3-6




Formula 3-7




Formula 3-8




Formula 3-9




Formula 3-10




Formula 3-11



Formula 3-12















Formula 3-13




Formula 3-14

-continued




-continued

Formula 3-15




Formula 3-16


Formula 3-17



Formula 3-18




Formula 3-19




Formula 3-20

Formula 3-21




Formula 3-22


Formula 3-23



Formula 3-24



Formula 3-25



Formula 3-26

Formula 3-27

Formula 3-28

Formula 3-29


Formula 3-30

Formula 3-31


Formula 3-32

Formula 3-33


-continued




Formula 3-34




Formula 3-35



Formula 3-36



Formula 3-37



Formula 3-38

[0040] In Formulae 3-1 to 3-38,

[0041] Y_1 may be O, S, $C(Z_3)(Z_4)$, $N(Z_5)$, or $Si(Z_6)(Z_7)$;

[0042] Z_1 to Z_7 may each independently be selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazolinyl group, a carbazolyl group, and a triazinyl group,

[0043] d_1 may be an integer selected from 1 to 4, d_2 may be an integer selected from 1 to 3, d_3 may be an integer selected from 1 to 6, d_4 may be an integer selected from 1 to 8, d_5 may be an integer of 1 or 2, d_6 may be an integer selected from 1 to 5, and * and *' are each independently a binding site to a neighboring atom.

[0044] In one embodiment, L_1 and L_2 may each independently be selected from, e.g.,

[0045] a phenylene group, a naphthylene group, a pyridinylene group, a pyrimidinylene group, a quinolinylene group, a quinoxaliny group, and a triazinylene group; and

[0046] a phenylene group, a naphthylene group, a pyridinylene group, a pyrimidinylene group, a quinolinylene group, a quinoxaliny group, and a triazinylene group, each substituted with at least one selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, but embodiments are not limited thereto.

[0047] In Formulae 1 and 2-1, b_1 denotes the number of L_1 s, and b_2 denotes the number of L_2 s, and b_1 and b_2 may each independently be selected from 0, 1, 2, and 3.

[0048] When b_1 or b_2 is 0, $*(L_1)_{b_1}-'$ or $*(L_2)_{b_2}-*$ is a single bond. When b_1 is 2 or greater, two or more L_1 s may be identical to or different from each other, and when b_2 is 2 or greater, two or more L_2 s may be identical to or different from each other.

[0049] In one embodiment, b_1 and b_2 may each independently be selected from 0, 1, and 2.

[0050] In Formula 2-1, Ar_1 may be selected from or include, e.g., a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group. In an implementation, Ar_1 may not be a substituted or unsubstituted triphenylenyl group.

[0051] In one embodiment, Ar_1 may be selected from, e.g.,

[0052] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an

acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

[0053] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzox-


azolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃).

[0054] In one embodiment, Ar₁ may be selected from, e.g.,


[0055] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group; and

[0056] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃).


[0057] In one embodiment, Ar₁ may be a group represented by one of the following Formulae 5-1 to 5-15.



Formula 5-1














Formula 5-2




Formula 5-3

-continued



-continued

Formula 5-4



Formula 5-15

Formula 5-5

[0058] In Formulae 5-1 to 5-15,

[0059] Y_{31} may be $C(Z_{33})(Z_{34})$ or $N(Z_{35})$;[0060] Z_{31} to Z_{35} may each independently be selected from:[0061] a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

Formula 5-6

[0062] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, and a phosphoric acid or a salt thereof;

Formula 5-7

Formula 5-8

[0063] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group;

Formula 5-9

Formula 5-10

Formula 5-11


Formula 5-12

Formula 5-13











Formula 5-14

[0064] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group and a naphthyl group; and[0065] $-Si(Q_{31})(Q_{32})(Q_{33})$,[0066] e_1 may be an integer selected from 1 to 5, e_2 may be an integer selected from 1 to 7, e_3 may be an integer selected from 1 to 3, e_4 may be an integer selected from 1 to 4, e_5 may be an integer of 1 or 2, and $*$ is a binding site to a neighboring atom.[0067] In some embodiments, Ar_1 may be a group represented by one of Formulae 6-1 to 6-24.

Formula 6-1



-continued



Formula 6-2

Formula 6-3

Formula 6-4

Formula 6-5

Formula 6-6

Formula 6-7








Formula 6-8

Formula 6-9

Formula 6-10

Formula 6-11

-continued



Formula 6-12

Formula 6-13

Formula 6-14


Formula 6-15

Formula 6-16


Formula 6-17

Formula 6-18


-continued




Formula 6-19




Formula 6-20




Formula 6-21



Formula 6-22



Formula 6-23



Formula 6-24

[0068] In Formulae 6-1 to 6-24, * is a binding site to a neighboring atom.

[0069] In Formulae 2-1 and 2-2, R_1 to R_{12} and R_{21} to R_{32} may each independently be selected from or include, e.g., a binding site (to one of A_1 , L_1 , and A_2 in Formula 1), a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted

C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_1)(Q_2)(Q_3)$, $-B(Q_4)(Q_5)$, and $N(Q_6)(Q_7)$.

[0070] For example, at least one of R_1 to R_{12} is a binding site to L_1 or A_1 in Formula 1, and at least one of R_{21} to R_{32} is a binding site to L_1 or A_1 in Formula 1.

[0071] In one embodiment, R_1 to R_{12} and R_{21} to R_{32} may each independently be selected from, e.g.,

[0072] a binding site to one of A_1 , L_1 , and A_2 in Formula 1;

[0073] a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0074] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

[0075] a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0076] a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl

group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolynyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyrimidinyl group, an imidazopyrimidinyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃); and

[0077] —Si(Q₁)(Q₂)(Q₃),

[0078] where one of R₁ to R₁₂ is a binding site to L₁ or A₁ in Formula 1, and one of R₂₁ to R₃₂ is a binding site to L₁ or A₁ in Formula 1.

[0079] In one embodiment, R₁ to R₁₂ and R₂₁ to R₃₂ may each independently be selected from, e.g.,

[0080] a binding site to one of A₁, L₁, and A₂ in Formula 1,

[0081] a hydrogen, —F, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decanyl group, an iso-decanyl group, a sec-decanyl group, a tert-decanyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, and a triazinyl group;

[0082] a methyl group, an ethyl group, a propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decanyl group, an iso-decanyl group, a sec-decanyl group, a


tert-decanyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, and a triazinyl group, each substituted with at least one selected from —F, a cyano group, a nitro group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃); and

[0083] —Si(Q₁)(Q₂)(Q₃), and


[0084] one of R₁ to R₁₂ is a binding site to L₁ or A₂ in Formula 1, and one of R₂₁ to R₃₂ is a binding site to L₁ or A₁ in Formula 1.

[0085] In one embodiment, the condensed cyclic compound may be, e.g., represented by one of the following Formulae 1A to 1L.


<Formula 1A>



<Formula 1B>




<Formula 1C>




-continued


<Formula 1D>




<Formula 1E>



<Formula 1F>




<Formula 1G>




-continued


<Formula 1H>




<Formula 1I>



<Formula 1J>




<Formula 1K>




-continued

<Formula 1L>




-continued


Formula 4-7



Formula 4-8




Formula 4-9




[0086] In Formulae 1A to 1L,

[0087] L_1 , L_2 , b_1 , b_2 , Ar_1 , R_1 to R_{12} , and R_{21} to R_{32} may be the same as described above with respect to Formulae 1, 2-1, and/or 2-2.[0088] In one embodiment, in Formulae 1A to 1L, L_1 and L_2 may each independently be a group represented by one of Formula 4-1 to 4-20, and b_1 and b_2 may each independently be selected from 0, 1, and 2.


Formula 4-1




Formula 4-2




Formula 4-3




Formula 4-4




Formula 4-5




Formula 4-6




Formula 4-10




Formula 4-11




Formula 4-12




Formula 4-13




Formula 4-14




Formula 4-15




-continued




Formula 4-16




Formula 4-17



Formula 4-18



Formula 4-19




Formula 4-20

[0089] In Formulae 4-1 to 4-20, * and *' are a binding site to a neighboring atom.


[0090] In some embodiments, in Formulae 1A to 1L, L_1 and L_2 may each independently be a group represented by one of Formulae 4-1 to 4-3, and b_1 and b_2 may each independently be 0 or 1.

[0091] In one embodiment, in Formulae 1A to 1L, Ar_1 may be a group represented by one of Formulae 6-1 to 6-19, and R_1 to R_{12} and R_{21} to R_{32} may each independently be a group represented by one of Formulae 7-1 to 7-6.




Formula 6-1


-continued




Formula 6-2




Formula 6-3




Formula 6-4




Formula 6-5




Formula 6-6




Formula 6-7




Formula 6-8



Formula 6-9




Formula 6-10



Formula 6-11


-continued




Formula 6-12

-continued

Formula 6-19




Formula 6-13




[0092] In Formulae 6-1 to 6-19 and 7-1 to 7-6, * is a binding site to a neighboring atom.


[0093] In an implementation, the compound represented by Formula 1 may be one of the following Compounds 1 to 75:




Formula 6-14




1




Formula 6-15




2




Formula 6-16



Formula 6-17




3




Formula 6-18

-continued


4




5



6




7




-continued


8




9




10




11



12




-continued




13

-continued




17


14




15




18




16



19




20




-continued


21




22




23



24




25




-continued


26




27







28







29



-continued




-continued




-continued


38




39



40




41




-continued


42




43



44




45




-continued

46




-continued

50




51


47




48




49




52



53




54




-continued

55




-continued


58




56




59



57




60




-continued

61




-continued

64





65


62




63



66




67




-continued


68




69



70




71




-continued


72




73



74



75



[0094] In the condensed cyclic compound represented by Formula 1, A_1 may be a group represented by Formula 2-1, and A_2 may be a group represented by Formula 2-2. For example, the condensed cyclic compound represented by Formula 1 may include both the group represented by Formula 2-1 and the group represented by Formula 2-2. In this regard, the electrical stability of the condensed cyclic compound represented by Formula 1 may increase. Thus, an electrical device including the condensed cyclic compound, e.g., an organic light-emitting device, may have an increased lifespan.

[0095] In an implementation, in Formula 2-1, L_2 may not be “a substituted or unsubstituted triphenylenylene group” and/or Ar_1 may not be “a substituted or unsubstituted triphenylenylene group”. A number of R_1 to R_{12} in Formula 2-1 that are a binding site to L_1 or A_2 in Formula 1 may be equal to a1. For example, if a1 is 1, one of R_1 to R_{12} is a binding site, if a1 is 2, two of R_1 to R_{12} is a binding site, etc. In this regard, the condensed cyclic compound represented by Formula 1 may have high triplet energy. The condensed cyclic compound may be synthesized by using a suitable organic synthesis method. The synthesis method of the condensed cyclic compound may be recognized by referring to the Examples in the specification.

[0096] At least one of the condensed cyclic compounds represented by Formula 1 may be included between a pair of electrodes of an organic light-emitting device. For example, the condensed cyclic compound may be included in an emission layer. For example, an organic light-emitting device may include a first electrode, a second electrode facing the first electrode, and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer. The emission layer of the organic layer may include at least one condensed cyclic compound represented by Formula 1.

[0097] As used herein, the expression the “(organic layer) includes at least one condensed cyclic compound” may be construed as meaning the “(organic layer) may include one condensed cyclic compound in a range of Formula 1 or two different condensed cyclic compounds in a range of Formula 1”.

[0098] In some embodiments, the organic layer may include only Compound 1 as the condensed cyclic compound. Here, Compound 1 may be included in the emission layer of the organic light-emitting device.

[0099] The organic layer may include, e.g., i) a hole transport region that is between the first electrode (an anode) and the emission layer and that includes at least one of a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer and ii) an electron transport region that is between the emission layer and the second electrode (a cathode) and that includes at least one of a hole blocking layer, an electron transport layer, and an electron injection layer. The emission layer may include the condensed cyclic compound represented by Formula 1.

[0100] As used herein, the term the “organic layer” refers to a single and/or a plurality of layers disposed between the first electrode and the second electrode in an organic light-emitting

device. A material included in the “organic layer” is not limited to an organic material.

[0101] FIG. 1 illustrates a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 may include a first electrode 110, an organic layer 150, and a second electrode 190.

[0102] Hereinafter, a structure and a method of manufacturing an organic light-emitting device according to an embodiment will be described with reference to FIG. 1.

[0103] Referring to FIG. 1, a substrate may be additionally disposed under the first electrode 110 or on the second electrode 190. The substrate may be a glass substrate or transparent plastic substrate, each with excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

[0104] The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode 110 on the substrate. When the first electrode 110 is an anode, the material for the first electrode 110 may be selected from materials with a high work function to facilitate hole injection. The first electrode 110 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. The material for the first electrode 110 may be a transparent and highly conductive material, and examples of such a material may include indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO_2), and zinc oxide (ZnO). When the first electrode 110 is a semi-transmissive electrode or a reflective electrode, as a material for forming the first electrode, at least one selected from magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag) may be used.

[0105] The first electrode 110 may have a single-layer structure, or a multi-layer structure including a plurality of layers. For example, the first electrode 110 may have a triple-layer structure of ITO/Ag/ITO.

[0106] The organic layer 150 may be disposed on the first electrode 110. The organic layer 150 may include an emission layer.

[0107] The organic layer 150 may further include a hole transport region between the first electrode 110 and the emission layer, and an electron transport region between the emission layer and the second electrode 190.

[0108] The hole transport region may include at least one of a hole injection layer (HIL), a hole transport layer (HTL), a buffer layer, and an electron blocking layer (EBL). The electron transport region may include at least one of a hole blocking layer (HBL), an electron transport layer (ETL), and an electron injection layer (EIL).

[0109] The hole transport region may have a single-layered structure formed of a single material, a single-layered structure formed of a plurality of different materials, or a multi-layered structure having a plurality of layers formed of a plurality of different materials.

[0110] For example, the hole transport region may have a single-layered structure formed of a plurality of different materials or a structure of hole injection layer/hole transport layer, a structure of hole injection layer/hole transport layer/buffer layer, a structure of hole injection layer/buffer layer, a structure of hole transport layer/buffer layer, or a structure of hole injection layer/hole transport layer/electron blocking layer, each structure sequentially stacked on the first electrode 110.


[0111] When the hole transport region includes a hole injection layer, the hole injection layer may be formed on the

first electrode **110** by using various methods, e.g., vacuum deposition, spin coating, casting, a Langmuir-Blodgett (LB) method, inkjet printing, laser printing, or laser-induced thermal imaging (LITI).




[0112] When a hole injection layer is formed by vacuum deposition, e.g., the vacuum deposition may be performed at a deposition temperature in a range of about 100° C. to about 500° C., at a vacuum degree in a range of about 10^{-8} torr to about 10^{-3} torr, and/or at a deposition rate in a range of about 0.01 Å/sec to about 100 Å/sec, though the conditions may vary depending on a compound for a hole injection material and a structure of a desired hole injection layer.

[0113] When a hole injection layer is formed by spin coating, the spin coating may be performed at a coating rate of about 2,000 rpm to about 5,000 rpm and/or at a temperature of about 80° C. to about 200° C. by taking into account a compound for a hole injection layer to be deposited and the structure of a hole injection layer to be formed.


[0114] When the hole transport region includes a hole transport layer, the hole transport layer may be formed on the first electrode **110** or the hole injection layer by using various methods, e.g., vacuum deposition, spin coating, casting, LB method, inkjet printing, laser printing, or LITI. When the hole transport layer is formed by vacuum deposition or spin coating, conditions for vacuum deposition and coating may be similar to the above-described vacuum deposition and coating conditions for forming the hole injection layer. The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB, β -NPB, TPD, a spiro-TPD, a spiro-NPB, α -NPB, TAPC, HMTDP, 4,4',4"-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), (polyaniline)/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201 below, and a compound represented by Formula 202 below.




-continued




-continued

 β -NPB


-continued




TAPC




TPD




HMTPD




Spiro-TPD




<Formula 201>



<Formula 202>



Spiro-NPB



methylated NPB

[0115] In Formulae 201 and 202,**[0116]** L_{201} to L_{205} may each independently be the same as defined in connection with L_1 to L_3 ;**[0117]** $xa1$ to $xa4$ may each independently be selected from 0, 1, 2, and 3;**[0118]** $xa5$ may be selected from 1, 2, 3, 4, and 5;**[0119]** R_{201} to R_{204} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.**[0120]** For example, in Formulae 201 and 202,**[0121]** L_{201} to L_{205} may each independently be selected from:**[0122]** a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorene group, a dibenzofluorene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene

group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylene group, a carbazolyene group, and a triazinylene group; and

[0123] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a pyrenylene group, a chrysenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylenylene group, a quinazolinylene group, a carbazolyene group, and a triazinylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;

[0124] xa1 to xa4 may each independently be 0, 1, or 2;


[0125] xa5 may be 1, 2, or 3;

[0126] R₂₀₁ to R₂₀₄ may each independently be selected from:


[0127] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and

[0128] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an azulenyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, but embodiments are not limited thereto.


[0129] The compound represented by Formula 201 may be represented by Formula 201A:



[0130] For example, the compound represented by Formula 201 may be represented by Formula 201A-1.



[0131] The compound represented by Formula 202 may be represented by Formula 202A.



[0132] In Formula 201A, 201A-1, and 202A, L₂₀₁ to L₂₀₃, xa1 to xa3, xa5, and R₂₀₂ to R₂₀₄ may be the same as disclosed in the present specification, R₂₁₁ and R₂₁₂ may each independently be the same as defined in connection with R₂₀₃, and R₂₁₃ to R₂₁₆ may each independently be selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a

[0148] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a

pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;

[0149] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, and a triazinyl group; and

[0150] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;


[0151] xa5 may be 1 or 2.

[0152] In an implementation, in Formulae 201A and 201A-1, R₂₁₃ and R₂₁₄ may be linked to each other and form a saturated or unsaturated ring.


[0153] The compound represented by Formula 201 and the compound represented by Formula 202 may include, e.g., one of Compounds HT1 to HT20 below.

-continued


HT2



HT3




HT1




-continued

-continued


HT4




HT6



HT5




HT7




-continued

-continued


HT8




HT10



HT9




HT11




-continued

HT12




-continued


HT16




HT13




HT17




HT14




HT18



HT15




HT19



-continued

HT20




[0154] A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1000 Å. When the hole transport region includes both a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, e.g., about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0155] The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.


[0156] The charge-generation material may be, e.g., a p-dopant. The p-dopant may include one of a quinone derivative, a metal oxide, and a cyano group-containing compound. Examples of the p-dopant may include a quinone derivative, such as tetracyanoquinonedimethane (TCNQ) or 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonedimethane (F4-TCNQ); a metal oxide, such as a tungsten oxide or a molybdenum oxide; and Compound HT-D1 below.

<Compound HT-D1>



-continued

<F4-TCNQ>



[0157] The hole transport region may further include at least one of a buffer layer and an electron blocking layer, in addition to the hole injection layer and the hole transport layer. The buffer layer may compensate for an optical resonance distance according to a wavelength of light emitted from the emission layer, and the light-emission efficiency of a formed organic light-emitting device may be improved. A material included in the buffer layer may be a material included in the hole transport region. The electron blocking layer may help prevent injection of electrons from the electron transport region.

[0158] An emission layer may be formed on the first electrode 110 or the hole transport region by using various methods, e.g., vacuum deposition, spin coating, casting, LB method, inkjet printing, laser printing, or LITI. When the emission layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the emission layer may be determined by referring to the deposition and coating conditions for the hole injection layer.

[0159] When the organic light-emitting device 10 is a full-color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In an implementation, the emission layer may have a stacked structure of a red emission layer, a green emission layer, and a blue emission layer, or may include a red-light emission material, a green-light emission material, and a blue-light emission material, which are mixed with each other in a single layer, to emit white light.


[0160] The emission layer may include the condensed cyclic compound represented by Formula 1.

[0161] The emission layer may include a host and a dopant. In an implementation, the host may include the condensed cyclic compound represented by Formula 1.

[0162] The dopant may include, e.g., at least one of a fluorescent dopant and a phosphorescent dopant.

[0163] The phosphorescent dopant may include, e.g., an organometallic complex represented by Formula 401 below.

<Formula 401>



[0164] In Formula 401,

[0165] M may be selected from, e.g., iridium (Ir), platinum (Pt), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), and thulium (Tm);

[0166] X_{401} to X_{404} may each independently be, e.g., nitrogen or carbon;

[0167] rings A_{401} and A_{402} may each independently be selected from or include, e.g., a substituted or unsubstituted benzene, a substituted or unsubstituted naphthalene, a substituted or unsubstituted fluorene, a substituted or unsubstituted spiro-fluorene, a substituted or unsubstituted indene, a substituted or unsubstituted pyrrole, a substituted or unsubstituted thiophene, a substituted or unsubstituted furan, a substituted or unsubstituted imidazole, a substituted or unsubstituted pyrazole, a substituted or unsubstituted thiazole, a substituted or unsubstituted isothiazole, a substituted or unsubstituted oxazole, a substituted or unsubstituted isoxazole, a substituted or unsubstituted pyridine, a substituted or unsubstituted pyrazine, a substituted or unsubstituted pyrimidine, a substituted or unsubstituted pyridazine, a substituted or unsubstituted quinoline, a substituted or unsubstituted isoquinoline, a substituted or unsubstituted benzoquinoline, a substituted or unsubstituted quinoxaline, a substituted or unsubstituted quinazoline, a substituted or unsubstituted carbazole, a substituted or unsubstituted benzoimidazole, a substituted or unsubstituted benzofuran, a substituted or unsubstituted benzothiophene, a substituted or unsubstituted isobenzothiophene, a substituted or unsubstituted benzoxazole, a substituted or unsubstituted isobenzoxazole, a substituted or unsubstituted triazole, a substituted or unsubstituted oxadiazole, a substituted or unsubstituted triazine, a substituted or unsubstituted dibenzofuran, and a substituted or unsubstituted dibenzothiophene;

[0168] at least one substituent of the substituted benzene, substituted naphthalene, substituted fluorene, substituted spiro-fluorene, substituted indene, substituted pyrrole, substituted thiophene, substituted furan, substituted imidazole, substituted pyrazole, substituted thiazole, substituted isothiazole, substituted oxazole, substituted isoxazole, substituted pyridine, substituted pyrazine, substituted pyrimidine, substituted pyridazine, substituted quinoline, substituted isoquinoline, substituted benzoquinoline, substituted quinoxaline, substituted quinazoline, substituted carbazole, substituted benzoimidazole, substituted benzofuran, substituted benzothiophene, substituted isobenzothiophene, substituted benzoxazole, substituted isobenzoxazole, substituted triazole, substituted oxadiazole, substituted triazine, substituted dibenzofuran, and substituted dibenzothiophene may be selected from, e.g.,

[0169] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0170] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl

group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q_{401})(Q_{402}), —Si(Q_{403})(Q_{404})(Q_{405}), and —B(Q_{406})(Q_{407});

[0171] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0172] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q_{411})(Q_{412}), —Si(Q_{413})(Q_{414})(Q_{415}), and —B(Q_{416})(Q_{417}); and

[0173] —N(Q_{421})(Q_{422}), —Si(Q_{423})(Q_{424})(Q_{425}), and —B(Q_{426})(Q_{427});

[0174] L_{401} may be an organic ligand;

[0175] xc1 may be 1, 2, or 3; and


[0176] xc2 may be 0, 1, 2, or 3.

[0177] The descriptions of Q_{401} to Q_{407} , Q_{411} to Q_{417} , and Q_{421} to Q_{427} may be the same as defined in connection with Q_i herein.

[0178] L_{401} may be a monovalent, divalent, or trivalent organic ligand. In an implementation, L_{401} may be selected from, e.g., a halogen ligand (for example, Cl or F), a diketone ligand (for example, acetylacetonate, 1,3-diphenyl-1,3-propanedione, 2,2,6,6-tetramethyl-3,5-heptanedione, or hexafluoroacetonate), a carboxylic acid ligand (for example, picolinate, dimethyl-3-pyrazolecarboxylate, or benzoate), a carbon monoxide ligand, an isonitrile ligand, a cyano ligand, and a phosphorous ligand (for example, phosphine, or phosphite).


[0179] In an implementation, in Formula 401, when ring A_{401} has two or more substituents, the two or more substituents of ring A_{401} may link to each other and form a saturated or unsaturated ring.

[0180] In an implementation, in Formula 401, when ring A_{402} has two or more substituents, the two or more substituents of ring A_{402} may link to each other and form a saturated or unsaturated ring.




-continued

PD4




[0181] In Formula 401, when xc1 is 2 or greater, a plurality of ligands in Formula 401 may be identical to or different from each other. In Formula 401, when xc1 is 2 or greater, A₄₀₁ and A₄₀₂ may be respectively and directly linked to A₄₀₁ and A₄₀₂ of a different neighboring ligand or may link to A₄₀₁ and A₄₀₂ of a different neighboring ligand via a linking group (e.g. a C₁-C₅ alkylene group, —N(R')— (where, R' is a C₁-C₁₀ alkyl group or a C₆-C₂₀ aryl group), or —C(=O)—) therebetween.

PD5




[0182] The phosphorescent dopant may include, e.g., at least one of the following Compounds PD1 to PD74.


PD1




PD6




PD2




PD7




PD3



PD8




-continued




PD9


-continued




PD14




PD10




PD15




PD11




PD16




PD12



PD17








PD13



PD18


-continued




-continued

PD19

PD24




PD20



PD25


PD21

PD26




PD22

PD27








PD23

PD28




-continued




-continued

PD29




PD34

PD30




PD35

PD31




PD36

PD32




PD37

PD33




PD38


-continued




PD39




PD40



PD41




PD42




PD43


-continued




PD44




PD45




PD46



PD47




PD48




PD49


-continued




PD50




PD51




PD52



PD53




PD54




PD55


-continued




PD56




PD57



PD58




PD59




PD60

-continued




PD61

-continued




PD66


PD62




PD67




PD63




PD68




PD64




PD69




PD65



PD70




-continued




PD71

-continued




PD74




PD72

[0183] In some embodiments, the phosphorescent dopant may include PtOEP:




PD73


PtOEP




[0184] The fluorescent dopant may include, e.g., at least one selected from DPAVB_i, BDAVB_i, TBPe, DCM, DCJT_B, Coumarin 6, and C545T.

DPVB_i


-continued




DPAVBi




TBPc




DCM



DCJTb




Coumarin 6



C545T

[0185] In some embodiments, the fluorescent dopant may include, e.g., a compound represented by Formula 501:



[0186] In Formula 501,

[0187] Ar_{501} may be selected from:

[0188] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene; and

[0189] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, and $-\text{Si}(\text{Q}_{501})(\text{Q}_{502})(\text{Q}_{503})$ (where, Q_{501} to Q_{503} are each independently selected from a hydrogen, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_6 - C_{60} aryl group, and a C_1 - C_{60} heteroaryl group);

[0190] descriptions of L_{501} to L_{503} may be the same as defined in connection with the description of L_{201} herein;

[0191] R_{501} and R_{502} may each independently be selected from:

[0192] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazole group, a triazinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0193] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group,

a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0194] xd1 to xd3 may each independently be selected from 0, 1, 2, and 3; and

[0195] xd4 may be selected from 1, 2, 3, and 4.

[0196] In the emission layer, the amount of the dopant may be generally selected within a range of, e.g., about 0.01 parts to about 15 parts by weight, based on about 100 parts by weight of the host.

[0197] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0198] Then, an electron transport region may be disposed on the emission layer.

[0199] The electron transport region may include at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.


[0200] For example, the electron transport region may have a structure of electron transport layer/electron injection layer or a structure of hole blocking layer/electron transport layer/electron injection layer, wherein layers of each structure are sequentially stacked from the emission layer in the stated order.

[0201] According to an embodiment, the organic layer 150 of the organic light-emitting device may include an electron transport region between the emission layer and the second electrode 190.


[0202] The electron transport region may include a hole blocking layer. The hole blocking layer may be formed, when the emission layer includes a phosphorescent dopant, to help prevent diffusion of excitons or holes into an electron transport layer.

[0203] When the electron transport region includes a hole blocking layer, the hole blocking layer may be formed on the emission layer by using various methods, e.g., vacuum deposition, spin coating, casting, an LB method, inkjet printing, laser printing, or LITI. When the hole blocking layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the hole blocking layer may be determined by referring to the deposition and coating conditions for the hole injection layer.

[0204] The hole blocking layer may include, e.g., at least one selected from BCP and Bphen.



BCP




Bphen


[0205] A thickness of the hole blocking layer may be in a range of about 20 Å to about 1,000 Å, e.g., about 30 Å to about 300 Å. When the thickness of the hole blocking layer is within this range, excellent hole blocking characteristics may be obtained without a substantial increase in driving voltage.

[0206] The electron transport region may include an electron transport layer. The electron transport layer may be formed on the emission layer or the hole blocking layer by using various methods, e.g., vacuum deposition, spin coating, casting, an LB method, inkjet printing, laser printing, or LITI. When the electron transport layer is formed by using vacuum deposition or spin coating, vacuum deposition and coating conditions for the electron transport layer may be determined by referring to the vacuum deposition and coating conditions for the hole injection layer.


[0207] The electron transport layer may include, e.g., at least one selected from BCP, Bphen, Alq₃, BALq, TAZ, and NTAZ.

Alq₃


-continued



BALq




TAZ



NTAZ

[0208] In some embodiments, the electron transport layer may include, e.g., at least one compound selected from a compound represented by Formula 601 and a compound represented by Formula 602 illustrated below.



Formula 601>

[0209] In Formula 601,

[0210] Ar₆₀₁ may be selected from:

[0211] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene;

[0212] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀

heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, and —Si(Q₃₀₁)(Q₃₀₂)(Q₃₀₃) (where, Q₃₀₁ to Q₃₀₃ may be each independently selected from a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₆-C₆₀ aryl group, and a C₁-C₆₀ heteroaryl group);

[0213] the descriptions for L₆₀₁ may be the same as defined in connection with L₂₀₁ herein;

[0214] E₆₀₁ may be selected from:

[0215] a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and


[0216] a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl

group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0217] xe1 may be selected from 0, 1, 2, and 3; and

[0218] xe2 may be selected from 1, 2, 3, and 4.

<Formula 602>



[0219] In Formula 602,

[0220] X₆₁₁ may be N or C-(L₆₁₁)_{xe611}-R₆₁₁, X₆₁₂ may be N or C-(L₆₁₂)_{xe612}-R₆₁₂, X₆₁₃ may be N or C-(L₆₁₃)_{xe613}-R₆₁₃, and at least one selected from X₆₁₁ to X₆₁₃ may be N;

[0221] L₆₁₁ to L₆₁₆ may be each the same as defined in connection with L₂₀₁ provided herein;

[0222] R₆₁₁ to R₆₁₆ may each independently be selected from:






[0223] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinoliny group, an isoquinoliny group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group; and

[0224] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinoliny group, an isoquinoliny group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl

group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;


[0225] xe611 to xe616 may each independently be selected from 0, 1, 2, and 3.

[0226] The compound represented by Formula 601 and/or the compound represented by Formula 602 may each independently be one of Compounds ET1 to ET15 illustrated below.




-continued

ET6




-continued


ET9




ET7



ET10




ET8




-continued

ET11





-continued

ET14




ET15

ET12



ET13




[0227] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 150 Å to about 500 Å. When the thickness of the electron transport layer is within this range, excellent electron transport characteristics may be obtained without a substantial increase in driving voltage.

[0228] The electron transport layer may further include a metal-containing material in addition to the materials described above.


[0229] The metal-containing material may include a Li complex. The Li complex may include, e.g., Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.

ET-D1



-continued

ET-D2



[0230] The electron transport region may include an electron injection layer that facilitates electron injection from the second electrode **190**.

[0231] The electron injection layer may be formed on the electron transport layer by using various methods, e.g., vacuum deposition, spin coating, casting, an LB method, inkjet printing, laser printing, or LITI. When the electron injection layer is formed by vacuum deposition or spin coating, vacuum deposition and coating conditions for the electron injection layer may be determined by referring to the vacuum deposition and coating conditions for the hole injection layer.

[0232] The electron injection layer may include, e.g., at least one selected from LiF, NaCl, CsF, Li₂O, BaO, and LiQ.

[0233] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, e.g., about 3 Å to about 90 Å. When the thickness of the electron injection layer is within this range, excellent electron injection characteristics may be obtained without a substantial increase in driving voltage.

[0234] The second electrode **190** may be disposed on the organic layer **150**. The second electrode **190** may be a cathode that is an electron injection electrode, and in this regard, a material for forming the second electrode **190** may be a material having a low work function, and such a material may include metal, an alloy, an electrically conductive compound, or a mixture thereof. Examples of the material for the second electrode **190** may include lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag). In some embodiments, the material for forming the second electrode **190** may be ITO or IZO. The second electrode **190** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode.

[0235] Hereinbefore, an organic light-emitting device has been described with reference to FIG. 1, but it is not limited thereto.

[0236] A C₁-C₆₀ alkyl group used herein refers to a linear or branched aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and detailed examples thereof are a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, or a hexyl group. A C₁-C₆₀ alkylene group used herein refers to a divalent group having the same structure as a C₁-C₆₀ alkyl group.

[0237] A C₁-C₆₀ alkoxy group used herein refers to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is the C₁-C₆₀ alkyl group), and detailed examples thereof are a methoxy group, an ethoxy group, or an isopropoxy group.

[0238] A C₂-C₆₀ alkenyl group used herein refers to a hydrocarbon group formed by substituting at least one carbon double bond in the middle or terminal of the C₂-C₆₀ alkyl group, and detailed examples thereof are an ethenyl group, a propenyl group, or a butenyl group. A C₂-C₆₀ alkenylene group used herein refers to a divalent group having the same structure as a C₂-C₆₀ alkenyl group.

[0239] A C₂-C₆₀ alkynyl group used herein refers to a hydrocarbon group formed by substituting at least one carbon triple bond in the middle or terminal of the C₂-C₆₀ alkyl group, and detailed examples thereof are an ethynyl group or a propynyl group. A C₂-C₆₀ alkynylene group used herein refers to a divalent group having the same structure as a C₂-C₆₀ alkynyl group.

[0240] A C₃-C₁₀ cycloalkyl group used herein refers to a monovalent monocyclic saturated hydrocarbon group including 3 to 10 carbon atoms, and detailed examples thereof are a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, or a cycloheptyl group. A C₃-C₁₀ cycloalkylene group used herein refers to a divalent group having the same structure as a C₃-C₁₀ cycloalkyl group.

[0241] A C₁-C₁₀ heterocycloalkyl group used herein refers to a monovalent monocyclic group including at least one heteroatom selected from N, O, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and detailed examples thereof are a tetrahydrofuranyl group, or a tetrahydrothiophenyl group. A C₁-C₁₀ heterocycloalkylene group used herein refers to a divalent group having the same structure as a C₁-C₁₀ heterocycloalkyl group.

[0242] A C₃-C₁₀ cycloalkenyl group used herein refers to a monovalent monocyclic group including 3 to 10 carbon atoms and at least one double bond in the ring thereof and does not have aromaticity, and detailed examples thereof are a cyclopentenyl group, a cyclohexenyl group, or a cycloheptenyl group. A C₃-C₁₀ cycloalkenylene group used herein refers to a divalent group having the same structure as a C₃-C₁₀ cycloalkenyl group.

[0243] As used herein, a C₁-C₁₀ heterocycloalkenyl group refers to a monovalent monocyclic group that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one double bond in its ring. Examples of the C₁-C₁₀ heterocycloalkenyl group include a 2,3-dihydrofuranyl group and a 2,3-dihydrothiophenyl group. As used herein, a C₁-C₁₀ heterocycloalkenylene group refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkenyl group.

[0244] A C₆-C₆₀ aryl group used herein refers to a monovalent group including a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C₆-C₆₀ arylenylene group used herein refers to a divalent group including a carbocyclic aromatic system having 6 to 60 carbon atoms. Detailed examples of the C₆-C₆₀ aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, or a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylenylene group each include two or more rings, the rings may be fused to each other.

[0245] A C₁-C₆₀ heteroaryl group used herein refers to a monovalent group having a carbocyclic aromatic system including at least one heteroatom selected from N, O, P, and S as a ring-forming atom and 1 to 60 carbon atoms. A C₁-C₆₀ heteroarylenylene group used herein refers to a divalent group having a carbocyclic aromatic system including at least one heteroatom selected from N, O, P, and S as a ring-forming atom and 1 to 60 carbon atoms. Detailed examples of the

C₁-C₆₀ heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, or an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be fused to each other.

[0246] A C₆-C₆₀ aryloxy group used herein indicates —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and a C₆-C₆₀ arylthio group used herein indicates —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

[0247] A monovalent non-aromatic condensed polycyclic group used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed to each other and only carbon atoms as a ring forming atoms, wherein the molecular structure as a whole is non-aromatic in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. A divalent non-aromatic condensed polycyclic group used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0248] As used herein, a monovalent non-aromatic condensed polycyclic group refers to a monovalent non-aromatic condensed polycyclic group that has two or more rings condensed to each other, only carbon atoms (for example, the number of carbon atoms may be in a range of 8 to 60) as a ring forming atom, and non-aromaticity in the entire molecular structure. Examples of the monovalent non-aromatic condensed polycyclic group include a fluorenyl group. As used herein, a divalent non-aromatic condensed polycyclic group refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0249] As used herein, at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, substituted C₁-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₁-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₁-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from

[0250] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —B(Q₁₄)(Q₁₅), and —N(Q₁₆)(Q₁₇);

[0251] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0252] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —B(Q₂₄)(Q₂₅), and —N(Q₂₆)(Q₂₇); and

[0253] —Si(Q₃₁)(Q₃₂)(Q₃₃), —B(Q₃₄)(Q₃₅), and —N(Q₃₆)(Q₃₇);

[0254] wherein Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0255] In some embodiments, at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, substituted C₁-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₁-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₁-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from

[0256] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a

throlinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, $-\text{N}(\text{Q}_{21})$ (Q_{22}), $-\text{B}(\text{Q}_{23})(\text{Q}_{24})$, and $-\text{Si}(\text{Q}_{25})(\text{Q}_{26})(\text{Q}_{27})$; and

[0260] $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{33})(\text{Q}_{34})$, and $-\text{Si}(\text{Q}_{35})(\text{Q}_{36})(\text{Q}_{37})$;

[0261] wherein Q_{11} to Q_{17} , Q_{21} to Q_{27} , and Q_{31} to Q_{37} may be each independently selected from a hydrogen, a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, but embodiments are not limited thereto.

[0262] Hereinafter, an organic light-emitting device according to exemplary embodiments are described in detail with reference to Synthesis Examples and Examples. However, the organic light-emitting device is not limited thereto.


[0263] The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples.

EXAMPLES

Synthesis Example 1

Synthesis of Compound 1

[0264]




[0265] 1.4 g (4 mmol) of Intermediate A, 1.6 g (4 mmol) of Intermediate B, 231 mg (0.2 mmol) of $\text{Pd}(\text{PPh}_3)_4$, and 1.1 g (12 mmol) of sodium t-butoxide were added to 100 ml of toluene and 25 ml of ethanol in a nitrogen atmosphere and refluxed for 24 hours. When the reaction was completed, the solvent was evaporated, and 800 ml of methylene chloride and 800 ml of water were added thereto to wash the mixture. Then, an organic layer obtained therefrom was dried with anhydrous magnesium sulfate. Subsequently, the resultant was recrystallized and purified through silica gel chromatography, and thus, 1.6 g of Compound 1 (yield: 40%) was obtained.

[0266] MS (MALDI-TOF) m/z : 545 $[\text{M}]^+$.

Synthesis Example 2
Synthesis of Compound 4

[0267]





[0268] 1.2 g of Compound 4 (yield: 43%) was synthesized in the same manner as in the synthesis method of Compound 1, except that Intermediate C was used instead of Intermediate A.

[0269] MS (MALDI-TOF) m/z: 621 [M]⁺.

Synthesis Example 3
Synthesis of Compound 7

[0270]





[0271] 1.6 g of Compound 7 (yield: 45%) was synthesized in the same manner as in the synthesis method of Compound 1, except that Intermediate D was used instead of Intermediate A.


[0272] MS (MALDI-TOF) m/z: 621 [M]⁺.

Synthesis Example 4

Synthesis of Compound 26

[0273]






[0274] 0.9 g of Compound 26 (yield: 33%) was synthesized in the same manner as in the synthesis method of Compound 1, except that Intermediate E was used instead of Intermediate B.

[0275] MS (MALDI-TOF) m/z: 545 [M]⁺.

Synthesis Example 5

Synthesis of Compound 28

[0276]




[0277] 1.2 g of Compound 28 (yield: 30%) was synthesized in the same manner as in the synthesis method of Compound 1, except that Intermediate F was used instead of Intermediate A.

[0278] MS (MALDI-TOF) m/z: 671 [M]⁺.

Synthesis Example 6

Synthesis of Compound 32

[0279]




[0280] 1.1 g of Compound 32 (yield: 29%) was synthesized in the same manner as in the synthesis method of Compound 1, except that Intermediate F was used instead of Intermediate A, and Intermediate E was used instead of Intermediate B.

[0281] MS (MALDI-TOF) m/z: 671 [M]⁺.

Synthesis Example 7

Synthesis of Compound 36

[0282]




[0283] 1.4 g of Compound 36 (yield: 45%) was synthesized in the same manner as in the synthesis method of Compound 1, except that Intermediate C was used instead of Intermediate A, and Intermediate E was used instead of Intermediate B.

[0284] MS (MALDI-TOF) m/z : 621 [M]⁺.

Synthesis Example 8

Synthesis of Compound 37

[0285]



[0286] 1.5 g of Compound 37 (yield: 46%) was synthesized in the same manner as in the synthesis method of Compound 1, except that Intermediate D was used instead of Intermediate A, and Intermediate E was used instead of Intermediate B.


[0287] MS (MALDI-TOF) m/z : 621 [M]⁺.

Example 1

[0288] A indium tin oxide (ITO) glass substrate (available from Corning) having a sheet resistance and thickness of 15 Ω/cm^2 and 1,200 Å was cut to a size of 50 mm×50 mm×0.5 mm, sonicated in isopropyl alcohol and pure water for 15 minutes each, and then cleaned with UV and ozone for 30 minutes. The ITO glass substrate was then mounted on a vacuum depositor.

[0289] Compound HT-3 was deposited on the ITO glass substrate to form a hole injection layer having a thickness of 600 Å, NPB was deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å, and then Compound 1 (a host) and PD1 (a dopant) were co-deposited at a weight ratio of 0.15:1 on the hole transport layer to form an emission layer having a thickness of 400 Å.

[0290] BCP was deposited on the emission layer to form an electron transport layer having a thickness of 100 Å, LiF was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was deposited on the electrode injection layer to form a cathode having a thickness of 2,000 Å, thereby completing manufacture of an organic light-emitting device.



Example 2

[0291] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 4 was used instead of Compound 1 as a host in the formation of the emission layer.

Example 3

[0292] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 7 was used instead of Compound 1 as a host in the formation of the emission layer.

Example 4

[0293] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 26 was used instead of Compound 1 as a host in the formation of the emission layer.

Example 5

[0294] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 28 was used instead of Compound 1 as a host in the formation of the emission layer.

Example 6




[0295] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 32 was used instead of Compound 1 as a host in the formation of the emission layer.

Example 7

[0296] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 36 was used instead of Compound 1 as a host in the formation of the emission layer.


Example 8

[0297] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 37 was used instead of Compound 1 as a host in the formation of the emission layer.




Comparative Example 1

[0298] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound A, below, was used instead of Compound 1 as a host in the formation of the emission layer:




Comparative Example 2

[0299] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound B, below, was used instead of Compound 1 as a host in the formation of the emission layer:




Comparative Example 3

[0300] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound C, below, was used instead of Compound 1 as a host in the formation of the emission layer:



Comparative Example 4

[0301] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound D, below, was used instead of Compound 1 as a host in the formation of the emission layer:



Evaluation Example 1

[0302] Driving voltages, luminances, efficiencies, and lifespans (T_{90}) data of the organic light-emitting devices prepared in Examples 1 to 8 and Comparative Examples 1 and 4 were measured by using Kethley SMU 236 and a luminance meter PR650, and the results are shown in Table 1. The lifespan (T_{90}) is defined as the time for the luminance of an organic light-emitting device to decline to 90% of its initial luminance.

TABLE 1

	Emission layer (host)	Driving voltage (V)	Luminance (cd/m ²)	Efficiency (cd/A)	Color of emitted light	Lifespan (T_{90}) (hr) (@3000 cd/m ²)
Example 1	Compound 1	5.2	3000	52	green	125
Example 2	Compound 4	5.3	3000	55	green	105
Example 3	Compound 7	5.4	3000	59	green	109

TABLE 1-continued

	Emission layer (host)	Driving voltage (V)	Luminance (cd/m ²)	Efficiency (cd/A)	Color of emitted light	Lifespan (T ₉₀) (hr) (@3000 cd/m ²)
Example 4	Compound 26	5.3	3000	54	green	112
Example 5	Compound 28	5.4	3000	53	green	124
Example 6	Compound 32	5.2	3000	57	green	109
Example 7	Compound 36	5.3	3000	59	green	111
Example 8	Compound 37	5.2	3000	53	green	131
Comparative Example 1	Compound A	5.8	3000	28	green	60
Comparative Example 2	Compound B	5.9	3000	27	green	52
Comparative Example 3	Compound C	5.7	3000	12	green	20
Comparative Example 4	Compound D	5.8	3000	30	green	40

[0303] As shown in Table 1, driving voltages, luminances, efficiencies, and lifespans (T₉₀) of the organic light-emitting devices prepared in Examples 1 to 8 were excellent, compared to those of the organic light-emitting device prepared in Comparative Examples 1 and 4.

[0304] As described above, according to the one or more of the above embodiments, an organic light-emitting device including the condensed cyclic compound may have a low driving voltage, high efficiency, high luminance, and a long lifespan.

[0305] Example embodiments have been disclosed herein, and although specific terms are employed, they are used and are to be interpreted in a generic and descriptive sense only and not for purpose of limitation. In some instances, as would be apparent to one of ordinary skill in the art as of the filing of the present application, features, characteristics, and/or elements described in connection with a particular embodiment may be used singly or in combination with features, characteristics, and/or elements described in connection with other embodiments unless otherwise specifically indicated. Accordingly, it will be understood by those of skill in the art that various changes in form and details may be made without departing from the spirit and scope of the present invention as set forth in the following claims.

What is claimed is:

1. A condensed cyclic compound represented by Formula 1:




wherein, in Formula 1,


A₁ is a group represented by Formula 2-1, below, a₁ is 1, 2, or 3, and, when a₁ is 2 or greater, two or more A₁s are identical to or different from each other,

A₂ is a group represented by Formula 2-2, below, a₂ is 1, 2, or 3, and, when a₂ is 2 or greater, two or more A₂s are identical to or different from each other,

<Formula 2-1>



<Formula 2-2>



wherein, in Formulae 1, 2-1, and 2-2,

L₁ and L₂ are each independently selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group, provided that L₂ is not a substituted or unsubstituted triphenylene group,

b₁ and b₂ are each independently selected from 0, 1, 2, and 3; when b₁ or b₂ is 0, *-(L₁)_{b1}-* or *-(L₂)_{b2}-* is a single

bond, when b1 is 2 or greater, two or more L₁s are identical to or different from each other, and when b2 is 2 or greater, two or more L₂s are identical to or different from each other,

Ar₁ is selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, provided that Ar₁ is not a substituted or unsubstituted triphenylenyl group,

R₁ to R₁₂ and R₂₁ to R₃₂ are each independently selected from a binding site to one of A₁, L₁, and A₂ in Formula 1, a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —B(Q₄)(Q₅), and N(Q₆)(Q₇),

a number of R₁ to R₁₂ that are a binding site to L₁ or A₂ in Formula 1 is equal to a1, and a number of R₂₁ to R₃₂ that are a binding site to L₁ or A₂ in Formula 1 is equal to a2;

at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, substituted C₁-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₁-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₁-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino

group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —B(Q₁₄)(Q₁₅), and —N(Q₁₆)(Q₁₇);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —B(Q₂₄)(Q₂₅), and —N(Q₂₆)(Q₂₇); and

—Si(Q₃₁)(Q₃₂)(Q₃₃), —B(Q₃₄)(Q₃₅), and —N(Q₃₆)(Q₃₇);

wherein Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic

condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

2. The condensed cyclic compound as claimed in claim 1, wherein a₁ and a₂ are each independently 1 or 2.


3. The condensed cyclic compound as claimed in claim 1, wherein L₁ and L₂ are each independently selected from:

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzoimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group; and

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzoimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group.










group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzoimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group, each substituted with at least one selected from a deuterium, —F, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, C₁-C₂₀ alkyl group, C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.

4. The condensed cyclic compound as claimed in claim 1, wherein L₁ and L₂ are each independently a group represented by one of the following Formulae 3-1 to 3-38:



Formula 3-1

-continued



Formula 3-2

Formula 3-3

Formula 3-4

Formula 3-5

Formula 3-6













Formula 3-7

Formula 3-8

Formula 3-9

Formula 3-10

-continued



Formula 3-11

Formula 3-12

Formula 3-13

Formula 3-14

Formula 3-15

Formula 3-16

Formula 3-17

Formula 3-18










Formula 3-19

Formula 3-20

Formula 3-21

Formula 3-22

-continued



Formula 3-23

Formula 3-24

Formula 3-25

Formula 3-26

Formula 3-27








Formula 3-28

Formula 3-29

Formula 3-30

Formula 3-31

-continued



Formula 3-32

Formula 3-33

Formula 3-34

Formula 3-35

Formula 3-36

Formula 3-37

Formula 3-38

wherein, in Formulae 3-1 to 3-38,

Y_1 is O, S, $C(Z_3)(Z_4)$, $N(Z_5)$, or $Si(Z_6)(Z_7)$;

Z_1 to Z_7 are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxy-

lic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;

d1 is an integer selected from 1 to 4, d2 is an integer selected from 1 to 3, d3 is an integer selected from 1 to 6, d4 is an integer selected from 1 to 8, d5 is an integer of 1 or 2, d6 is an integer selected from 1 to 5, and * and *' are each independently a binding site to a neighboring atom.

5. The condensed cyclic compound as claimed in claim 1, wherein L₁ and L₂ are each independently selected from:

a phenylene group, a naphthylene group, a pyridinylene group, a pyrimidinylene group, a quinolinylene group, a quinoxalinylene group, and a triazinylene group; and

a phenylene group, a naphthylene group, a pyridinylene group, a pyrimidinylene group, a quinolinylene group, a quinoxalinylene group, and a triazinylene group, each substituted with at least one selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group.

6. The condensed cyclic compound as claimed in claim 1, wherein b1 and b2 are each independently selected from 0, 1, and 2.

7. The condensed cyclic compound as claimed in claim 1, wherein Ar₁ is selected from:

a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a

cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group,

a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$,

wherein Q_{31} to Q_{33} are each independently selected from a hydrogen, a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazolinyl group, a carbazolyl group, and a triazinyl group.


8. The condensed cyclic compound as claimed in claim 1, wherein Ar_1 is selected from:

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group; and


a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group, each substituted with at least one selected from a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$,

wherein Q_{31} to Q_{33} are each independently selected from a hydrogen, a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazolinyl group, a carbazolyl group, and a triazinyl group.


9. The condensed cyclic compound as claimed in claim 1, wherein Ar_1 is a group represented by one of the following Formulae 5-1 to 5-15:




Formula 5-1




Formula 5-2




Formula 5-3




Formula 5-4



Formula 5-5




Formula 5-6




Formula 5-7


-continued




Formula 5-8




Formula 5-9




Formula 5-10




Formula 5-11




Formula 5-12



Formula 5-13



Formula 5-14



Formula 5-15

wherein, in Formulae 5-1 to 5-15,

Y_{31} is $C(Z_{33})(Z_{34})$ or $N(Z_{35})$;

Z_{31} to Z_{35} are each independently selected from:

a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, and a phosphoric acid or a salt thereof;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a diben-


zofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, and a naphthyl group; and —Si(Q_{31})(Q_{32})(Q_{33}),


wherein Q_{31} to Q_{33} are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;

$e1$ is an integer selected from 1 to 5, $e2$ is an integer selected from 1 to 7, $e3$ is an integer selected from 1 to 3, $e4$ is an integer selected from 1 to 4, $e5$ is an integer of 1 or 2, and * is a binding site to a neighboring atom.


10. The condensed cyclic compound as claimed in claim 1, wherein Ar_1 is a group represented by one of the following Formulae 6-1 to 6-24:



Formula 6-1












Formula 6-2



Formula 6-3

-continued



Formula 6-4

Formula 6-5

Formula 6-6

Formula 6-7

Formula 6-8








Formula 6-9

Formula 6-10

Formula 6-11

Formula 6-12

-continued



Formula 6-13

Formula 6-14

Formula 6-15

Formula 6-16


Formula 6-17

Formula 6-18


Formula 6-19

-continued


Formula 6-20




Formula 6-21




Formula 6-22



Formula 6-23



Formula 6-24



wherein, in Formulae 6-1 to 6-24, * is a binding site to a neighboring atom.

11. The condensed cyclic compound as claimed in claim 1, wherein R_1 to R_{12} and R_{21} to R_{32} are each independently selected from:

a binding site to one of A_1 , L_1 , and A_2 in Formula 1;

a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl

group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazoyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$; and

$-\text{Si}(\text{Q}_1)(\text{Q}_2)(\text{Q}_3)$,

wherein one of R_1 to R_{12} is a binding site to L_1 or A_2 in Formula 1, and one of R_{21} to R_{32} is a binding site to L_1 or A_1 in Formula 1.

12. The condensed cyclic compound as claimed in claim 1, wherein R_1 to R_{12} and R_{21} to R_{32} are each independently selected from:

a binding site to one of A_1 , L_1 , and A_2 in Formula 1;

a hydrogen, $-\text{F}$, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decanyl group, an isodecanyl group, a sec-decanyl group, a tert-decanyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, and a triazinyl group;


a methyl group, an ethyl group, a propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decanyl group, an isodecanyl group, a sec-decanyl group, a tert-decanyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, and a triazinyl group, each substituted with at least one selected from $-\text{F}$, a cyano group, a nitro group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$; and

$-\text{Si}(\text{Q}_1)(\text{Q}_2)(\text{Q}_3)$,


wherein one of R_1 to R_{12} is a binding site to L_1 or A_2 in Formula 1, and one of R_{21} to R_{32} is a binding site to L_1 or A_1 in Formula 1.

13. The condensed cyclic compound as claimed in claim 1, wherein the compound represented by Formula 1 is represented by one of the following Formulae 1A to 1L:


<Formula 1A>




<Formula 1B>




<Formula 1C>



<Formula 1D>




-continued




-continued


<Formula 1H>




<Formula 1I>




<Formula 1J>




<Formula 1F>



<Formula 1G>




<Formula 1K>



-continued

<Formula 1L>




wherein, in Formulae 1A to 1L, the descriptions of L_1 , L_2 , b_1 , b_2 , Ar_1 , R_1 to R_{12} , and R_{21} to R_{32} are the same as L_1 , L_2 , b_1 , b_2 , Ar_1 , R_1 to R_{12} , and R_{21} to R_{32} of Formulae 1, 2-1, and 2-2.


14. The condensed cyclic compound as claimed in claim 13, wherein, in Formulae 1A to 1L,

b_1 and b_2 are each independently selected from 0, 1, and 2, and


L_1 and L_2 are each independently a group represented by one of the following Formulae 4-1 to 4-20,




Formula 4-1




Formula 4-2




Formula 4-3




Formula 4-4




Formula 4-5




Formula 4-6




Formula 4-7




Formula 4-8




Formula 4-9




Formula 4-10




Formula 4-11



Formula 4-12




Formula 4-13




Formula 4-14


-continued




Formula 4-15




Formula 4-16




Formula 4-17



Formula 4-18



Formula 4-19




Formula 4-20

wherein, in Formulae 4-1 to 4-20, * and *' are a binding site to a neighboring atom.

15. The condensed cyclic compound as claimed in claim 13, wherein:


Ar₁ is a group represented by one of the following Formulae 6-1 to 6-19, and

R₁ to R₁₂ and R₂₁ to R₃₂ are each independently a group represented by one of the following Formulae 7-1 to 7-6:




Formula 6-1


-continued




Formula 6-2




Formula 6-3




Formula 6-4




Formula 6-5




Formula 6-6




Formula 6-7




Formula 6-8



Formula 6-9










Formula 6-10




Formula 6-11

-continued





-continued


Formula 6-12




Formula 6-13




Formula 6-14




Formula 6-15



Formula 6-16



Formula 6-17



Formula 6-18

Formula 6-20

Formula 7-1

Formula 7-2

Formula 7-3

Formula 7-4









Formula 7-5

Formula 7-6

wherein, in Formulae 6-1 to 6-19 and 7-1 to 7-6, * is a binding site to a neighboring atom.


16. The condensed cyclic compound as claimed in claim 1, wherein the compound represented by Formula 1 is one of the following Compounds 1 to 75:

-continued




-continued


9




10




11



12




13




-continued


14




15



16




-continued




17


-continued




21




18




22




23




19



24




20



25


-continued

26




-continued


30



27




31





32

28




33

29




-continued


34




35



36




37




-continued


38



39




40




-continued

41




-continued


45




42



46





43




47

44




-continued




48


-continued




52




49




53




54



50




55



51


-continued

56




-continued


59




57




60



58




61




-continued


62



63




64




-continued


65




66



67




68




-continued


69




70



71




72




-continued


73



74



75



17. An organic light-emitting device, comprising:
a first electrode;

a second electrode facing the first electrode; and

an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes the condensed cyclic compound as claimed in claim 1.

18. The organic light-emitting device as claimed in claim 17, wherein:

the first electrode is an anode,

the second electrode is a cathode, and

the organic layer includes:

a hole transport region between the first electrode and the emission layer, the hole transport region including

at least one of a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer, and

an electron transport region between the emission layer and the second electrode, the electron transport region including at least one of a hole blocking layer, an electron transport layer, and an electron injection layer.

19. The organic light-emitting device as claimed in claim 17, wherein the emission layer includes the condensed cyclic compound.

20. The organic light-emitting device as claimed in claim 19, wherein the emission layer further includes a dopant.

* * * * *

专利名称(译)	缩合环状化合物和包括其的有机发光器件		
公开(公告)号	US20160172603A1	公开(公告)日	2016-06-16
申请号	US14/731011	申请日	2015-06-04
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	SHIN DONGWOO KIM SEULONG KIM YOUNSUN LEE JUNG SUB ITO NAOYUKI		
发明人	SHIN, DONGWOO KIM, SEULONG KIM, YOUNSUN LEE, JUNG SUB ITO, NAOYUKI		
IPC分类号	H01L51/00 C09K11/02 C07D403/04 C07D401/02 C07D223/14 C09K11/06		
CPC分类号	H01L51/0072 C09K2211/1007 C09K11/06 C07D403/04 C07D401/02 C09K11/025 H01L51/0054 H01L51/0085 H01L51/0058 H01L51/0067 H01L51/5056 H01L51/5012 H01L51/5088 H01L51/5008 H01L51/5096 H01L51/5072 H01L51/5092 C09K2211/185 C09K2211/1029 C07D223/14 C09B1/00 C09B57/00 C09B57/008 C09B57/10 C09K2211/1011 C09K2211/1044 H01L51/006 H01L51/5016 H05B33/14		
优先权	1020140177821 2014-12-10 KR		
其他公开文献	US9793496		
外部链接	Espacenet USPTO		

摘要(译)

缩合环状化合物和有机发光器件，该化合物由式1表示：(A₁)A₁-(L₁)B₁-(A₂)a₂ & #x3c; Formula 1> 其中，在式1中，A₁是由下面的式2-1表示的组，a₁是1,2或3，并且当a₁时为2或更大，两个或更多个A₁s彼此相同或不同，A₂是由式2-2表示的基团，下面，a₂是1,2或3，当a₂为2或更大时，两个或多个A₂彼此相同或不同，

