



(19) **United States**

(12) **Patent Application Publication**
SIM et al.

(10) **Pub. No.: US 2017/0338419 A1**

(43) **Pub. Date: Nov. 23, 2017**

(54) **HETEROCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

H01L 51/0052 (2013.01); *H01L 51/0058* (2013.01); *C07D 491/153* (2013.01); *H01L 51/0067* (2013.01); *H01L 51/5056* (2013.01); *H01L 51/5072* (2013.01); *H01L 51/5088* (2013.01); *H01L 51/5092* (2013.01); *H01L 51/5096* (2013.01)

(71) Applicant: **SAMSUNG DISPLAY CO., LTD.**,
Yongin-si (KR)

(72) Inventors: **Munki SIM**, Yongin-si (KR);
Youngkook KIM, Yongin-si (KR);
Junha PARK, Yongin-si (KR);
Hyoyoung LEE, Yongin-si (KR);
Eunjae JEONG, Yongin-si (KR);
Seokhwan HWANG, Yongin-si (KR)

(57) **ABSTRACT**

A heterocyclic compound and an organic light-emitting device including the same, the compound being represented by Formula 1:

$A_1-(A_2)_{a21}$ <Formula 1>

wherein, in Formula 1, A_1 is a group represented by Formula 2, A_2 is a group represented by Formula 3, and $a21$ is an integer of 1 to 11, wherein, when $a21$ is 2 to 11, two or more of A_2 are identical to or different from each other:

(21) Appl. No.: **15/496,120**

(22) Filed: **Apr. 25, 2017**


(30) **Foreign Application Priority Data**

May 19, 2016 (KR) 10-2016-0061439

Publication Classification

(51) **Int. Cl.**
H01L 51/00 (2006.01)
H01L 27/32 (2006.01)
C07D 491/153 (2006.01)
H01L 51/52 (2006.01)
C07F 9/6561 (2006.01)
H01L 51/50 (2006.01)

(52) **U.S. Cl.**
CPC *H01L 51/0071* (2013.01); *H01L 51/5218* (2013.01); *H01L 51/0072* (2013.01); *C07F 9/6561* (2013.01); *H01L 27/3211* (2013.01);



40

220
190
150
110
210

FIG. 1

10

190
150
110

FIG. 2

20

190
150
110
210

FIG. 3

30

220
190
150
110

FIG. 4

40

220
190
150
110
210

**HETEROCYCLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**

-continued

<Formula 3>

* —(L)_n—(Ar)_m,

CROSS-REFERENCE TO RELATED
APPLICATION

[0001] Korean Patent Application No. 10-2016-0061439, filed on May 19, 2016, in the Korean Intellectual Property Office, and entitled: "Heterocyclic Compound and Organic Light-Emitting Device Including the Same," is incorporated by reference herein in its entirety.

BACKGROUND

1. Field

[0002] Embodiments relate to a heterocyclic compound and an organic light-emitting device including the same.

2. Description of the Related Art

[0003] Organic light-emitting devices are self-emission devices and have wide viewing angles, high contrast ratios, short response times, and excellent brightness, driving voltage, and response speed characteristics, compared to devices in the art, and produce full-color images.

[0004] An example of such organic light-emitting devices may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode in this stated order. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, may recombine in the emission layer to produce excitons. These excitons transition from an excited state to a ground state, thereby generating light.

SUMMARY


[0005] Embodiments are directed to a heterocyclic compound and an organic light-emitting device including the same.

[0006] The embodiments may be realized by providing a heterocyclic compound represented by Formula 1:



[0007] wherein, in Formula 1, A₁ is a group represented by Formula 2, A₂ is a group represented by Formula 3, and a₂₁ is an integer of 1 to 11, wherein, when a₂₁ is 2 to 11, two or more of A₂ are identical to or different from each other:

<Formula 2>



[0008] wherein F₁ in Formula 2 is a dibenzofuran moiety or a dibenzothiophene moiety, R₁ to R₅ and R₁₂ in Formula 2 are each independently selected from a binding site to A₂, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), and —S(=O)₂(Q₁), b₆ in Formula 2 is an integer of 0 to 6, wherein, when b₆ is 2 to 6, two or more of R₆ are identical to or different from each other, X₁ in Formula 2 is O or S, at least one of R₁ to R₅ and R₁₂ is a binding site to A₂, a number of R₁ to R₅ and R₁₂ being the binding site to A₂ being the same as a₂₁, L in Formula 3 is selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group, n in Formula 3 is an integer of 0 to 7, wherein, when n is 2 to 7, two or more of L are identical to or different from each other, Ar in Formula 3 is selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₄)(Q₅)(Q₆), —C(=O)(Q₄), —S(=O)₂(Q₄), and —P(=O)(Q₄)(Q₅), m in Formula 3 is an integer of 1 to 7, wherein, when m is 2 to 7, two or more of Ar are identical to or different from each other, and at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed het-

eropolycyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_1 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

[0009] deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, and a substituted or unsubstituted C_1 - C_{60} alkoxy group; a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂); a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), wherein Q₁ to Q₆, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a naphthyl group.

cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0010] The embodiments may be realized by providing an organic light-emitting device including a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode and comprises an emission layer, wherein the organic layer includes the heterocyclic compound according to an embodiment

BRIEF DESCRIPTION OF THE DRAWINGS

[0011] Features will be apparent to those of skill in the art by describing in detail exemplary embodiments with reference to the attached drawings in which:

[0012] FIG. 1 illustrates a schematic cross-sectional diagram of an organic light-emitting device according to an embodiment;

[0013] FIG. 2 illustrates a schematic cross-sectional diagram of an organic light-emitting device according to an embodiment;

[0014] FIG. 3 illustrates a schematic cross-sectional diagram of an organic light-emitting device according to an embodiment; and

[0015] FIG. 4 illustrates a schematic cross-sectional diagram of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION


[0016] Example embodiments will now be described more fully hereinafter with reference to the accompanying drawings; however, they may be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey exemplary implementations to those skilled in the art.

[0017] In the drawing figures, the dimensions of layers and regions may be exaggerated for clarity of illustration. It will also be understood that when a layer or element is referred to as being “on” another layer or element, it can be directly on the other layer or element, or intervening layers may also be present. In addition, it will also be understood that when a layer is referred to as being “between” two layers, it can be the only layer between the two layers, or one or more intervening layers may also be present. Like reference numerals refer to like elements throughout.


[0018] A heterocyclic compound according to an embodiment may be represented by Formula 1.



[0019] In Formula 1, A₁ may be a group represented by Formula 2; A₂ may be a group represented by Formula 3; and a₂₁ may be an integer of 1 to 11. In an implementation, when a₂₁ is 2 to 11, two or more of A₂ may be identical to or different from each other:



<Formula 2>



<Formula 3>

[0020] F_1 , b_6 , R_1 to R_5 , R_{12} , X_1 , L , Ar , n , and m in Formulae 2 and 3 are described below.

[0021] In an implementation, F_1 in Formula 2 may be, e.g., a dibenzofuran moiety or a dibenzothiophene moiety. For example, F_1 may be a dibenzofuran.

[0022] R_1 to R_5 and R_{12} in Formula 2 may each independently be selected from or include, e.g., a binding site to A_2 , hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_1)(Q_2)(Q_3)$, and $-S(=O)_2(Q_1)$.

[0023] b_6 in Formula 2 may be an integer of 0 to 6. In an implementation, when b_6 is 2 to 6, two or more of R_6 may be identical to or different from each other.

[0024] X_1 in Formula 2 may be, e.g., O or S. In an implementation, X_1 may be O. In an implementation, X_1 may be S.

[0025] In an implementation, X_1 in Formula 2 may be O, and F_1 in Formula 2 may be a dibenzofuran moiety. In an implementation, X_1 in Formula 2 may be O, and F_1 in Formula 2 may be a dibenzothiophene moiety. In an implementation, X_1 in Formula 2 may be S, and F_1 in Formula 2 may be a dibenzofuran moiety. In an implementation, X_1 in Formula 2 may be S, and F_1 in Formula 2 may be a dibenzothiophene moiety.

[0026] In an implementation, at least one selected from R_1 to R_5 and R_{12} may be a binding site to A_2 Formula 1. For example, a number of R_1 to R_5 and R_{12} that are the binding site to A_2 may be the same number as the value of a_{21} . For example, when a_{21} is 2, two of R_1 to R_5 and R_{12} are binding sites to A_2 .

[0027] a_{21} in Formula 1 may be an integer of 1 to 6. In an implementation, a_{21} may be an integer of 1 to 3. In an implementation, a_{21} may be 1.

[0028] L in Formula 3 may be selected from or include, e.g., a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10}

cycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.


[0029] In an implementation, L in Formula 3 may be selected from:

[0030] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazololylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinylylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinnolinylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthrolinylylene group, a phenazinylylene group, a benzimidazololylylene group, an isobenzothiazolylylene group, a benzoxazololylylene group, an isobenzoxazololylylene group, a triazololylylene group, a tetrazolylylene group, an imidazopyridinylylene group, an imidazopyrimidinylylene group, and an azacarbazolylylene group; and


[0031] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazololylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinylylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinnolinylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthrolinylylene group, a phenazinylylene group, a benzimidazololylylene group, an isobenzothiazolylylene group, a benzoxazololylylene group, an isobenzoxazololylylene group, a triazololylylene group, a tetrazolylylene group, an imidazopyridi-

nylene group, an imidazopyrimidinylene group, and an azacarbazolylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenaziny group, a benzoimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃).


[0032] In an implementation, L in Formula 3 may be a group represented by one of the following Formulae 5-1 to 5-34:




Formula 5-1




Formula 5-2



Formula 5-3




Formula 5-4




-continued


Formula 5-5




Formula 5-6




Formula 5-7




Formula 5-8



Formula 5-9












Formula 5-10




Formula 5-11

-continued




-continued


Formula 5-12




Formula 5-13




Formula 5-14




Formula 5-15




Formula 5-16



Formula 5-17




Formula 5-18



Formula 5-19

Formula 5-20



Formula 5-21

Formula 5-22

Formula 5-23

Formula 5-24


Formula 5-25

Formula 5-26


Formula 5-27

Formula 5-28


-continued




Formula 5-29




Formula 5-30




Formula 5-31



Formula 5-32



Formula 5-33



Formula 5-34

[0033] In Formulae 5-1 to 5-34,

[0034] Y_1 may be O, S, $C(Z_3)(Z_4)$, $N(Z_5)$, or $Si(Z_6)(Z_7)$,

[0035] Z_1 to Z_7 may each independently be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano

group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, an isoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazoliny group, a benzoquinazoliny group, a cinnoliny group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indenocarbazolyl group, an indolocarbazolyl group, and —Si(Q_{31})(Q_{32})(Q_{33}).

[0036] d_2 may be an integer selected from 0 to 2,

[0037] d_3 may be an integer selected from 0 to 3,

[0038] d_4 may be an integer selected from 0 to 4,

[0039] d_5 may be an integer selected from 0 to 5,

[0040] d_6 may be an integer selected from 0 to 6,

[0041] d_8 may be an integer selected from 0 to 8, and

[0042] * and *' each indicate a binding site to a neighboring atom.

[0043] In an implementation, L may be a group represented by one of Formula 5-1, 5-2, or 5-13, and Z_1 in Formula 5-1, 5-2, or 5-13 may be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group.

[0044] n in Formula 3 may be an integer selected from 0 to 7. In an implementation, n in Formula 3 may be 0, 1, or 2.

[0045] In an implementation, n may be 1. When n is two or more, two or more of L may be identical to or different from each other, and when n is zero, “-(L)_n” indicates a single bond.

[0046] Ar in Formula 3 may be selected from or include, e.g., a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₄)(Q₅)(Q₆), —C(=O)(Q₄), —S(=O)₂(Q₄), and —P(=O)(Q₄)(Q₅).

[0047] In an implementation, Ar in Formula 3 may be selected from:

[0048] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;


[0049] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an

isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;


azolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

[0050] —Si(Q₄)(Q₅)(Q₆), —C(=O)(Q₄), —S(=O)₂(Q₄), and —P(=O)(Q₄)(Q₅).


[0051] In an implementation, Ar in Formula 3 may be a group represented by one of the following Formulae 6-1 to 6-37:




Formula 6-1




Formula 6-2




Formula 6-3




Formula 6-4




Formula 6-5



Formula 6-6




Formula 6-7




Formula 6-8


-continued




Formula 6-9




Formula 6-10




Formula 6-11




Formula 6-12




Formula 6-13



Formula 6-14




Formula 6-15




Formula 6-16


-continued




Formula 6-17




Formula 6-18




Formula 6-19



Formula 6-20




Formula 6-21




Formula 6-22


-continued




Formula 6-23




Formula 6-24




Formula 6-25




Formula 6-26



Formula 6-27




Formula 6-28




Formula 6-29


-continued




Formula 6-30




Formula 6-31




Formula 6-32




Formula 6-33




Formula 6-34



Formula 6-35



Formula 6-36



Formula 6-37

[0052] In Formulae 6-1 to 6-37,

[0053] Y_{31} , Y_{33} and Y_{37} may each independently be O, S, $C(Z_{35})(Z_{36})$, $N(Z_{37})$, or $Si(Z_{38})(Z_{39})$,

[0054] Y_{32} , Y_{34} , Y_{36} , and Y_{38} may each independently be $C(Z_{35})$ or N,

[0055] Z_{31} to Z_{39} may each independently be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an

imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazolinylnyl group, a benzoquinazolinylnyl group, a cinnolinylnyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafuorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indenocarbazolyl group, an indolocarbazolyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, and $-Si(Q_{31})(Q_{32})(Q_{33})$.

[0056] e2 may be an integer selected from 0 to 2,

[0057] e3 may be an integer selected from 0 to 3,

[0058] e4 may be an integer selected from 0 to 4,


[0059] e5 may be an integer selected from 0 to 5,

[0060] e6 may be an integer selected from 0 to 6, and

[0061] * indicates a binding site to a neighboring atom.










[0062] In an implementation, Ar in Formula 3 may a group represented by one of Formulae 6-1 to 6-6, 6-13 to 6-29, 6-31, 6-32, and 6-37.

[0063] In an implementation, Ar in Formula 3 may be a group represented by one of the following Formulae 7-1 to 7-25:




Formula 7-1

-continued




-continued

Formula 7-2




Formula 7-3




Formula 7-4

Formula 7-5




Formula 7-6




Formula 7-7


Formula 7-8



Formula 7-9



Formula 7-10



Formula 7-11

Formula 7-12

Formula 7-13


Formula 7-14

Formula 7-15


Formula 7-16

Formula 7-17


-continued




Formula 7-18




Formula 7-19




Formula 7-20



Formula 7-21




Formula 7-22




Formula 7-23

-continued



Formula 7-24




Formula 7-25

[0064] * in Formulae 7-1 to 7-25 indicates a binding site to a neighboring atom.


[0065] In an implementation, m in Formula 3 may be an integer selected from 1 to 7. In an implementation, m in Formula 3 may be 1, 2, or 3. When m is 2 to 7, two or more of Ar may be identical to or different from each other.

[0066] In an implementation, m in Formula 3 may be 1 or 2.

[0067] In an implementation, A₁ in Formula 1 may be a group represented by one of the following Formulae 2A to 2F:




Formula 2A




Formula 2B

-continued


Formula 2C




Formula 2D



Formula 2E



Formula 2F




[0068] In Formulae 2A to 2F, X_1 and X_2 may each independently be, e.g., O or S. R_1 to R_{11} may each independently be selected from or include, e.g., a binding site to A_2 in Formula 1, hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino

group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_1)(Q_2)(Q_3)$, and $-S(=O)_2(Q_1)$.


[0069] In an implementation, in Formulae 2A to 2F, X_1 and X_2 may be O, and R_5 may be a binding site to A_2 .

[0070] In an implementation, both X_1 and X_2 in Formulae 2A to 2F may be O. In an implementation, X_1 and X_2 in Formulae 2A to 2F may be different from each other.

[0071] In an implementation, the heterocyclic compound represented by Formula 1 may be one of the following Compounds 1 to 72 and 74 to 138.




1



2


-continued

3





-continued

7





4

8





5

9




6

10




-continued


11




12



13




14




-continued


15




16




17



18




-continued





-continued

22





20

23



24





-continued

-continued



28

25





26

29



27

30





-continued

-continued



31

34





32

35



33

36





-continued

-continued



37

40





38

41



39



42



-continued



-continued

46





44

47




45

48




-continued


-continued




49




52




50



53










51



54

-continued

-continued





-continued

-continued



62

65





63

66









64

67




-continued

-continued




-continued

75




-continued


79




76




80




77




81



78











82



-continued









-continued



-continued


-continued

95




-continued

99





-continued

103





104

100





105

101




106

102




-continued

107





-continued

111





108

112





109

113



110

114





-continued

-continued



119

115





116

120





117

121



118

122





-continued

-continued



123

126





124

127



125

128





-continued

-continued



129

132





133

130



131

134





-continued

-continued


138

135




136

[0072] The heterocyclic compound represented by Formula 1 may include a core structure or moiety represented by Formula 1a, below, and a glass transition temperature (T_g) or a melting point thereof may be high. Accordingly, the heterocyclic compound represented by Formula 1 may have excellent heat resistance, stability to charges, and charge transport capability. An electronic device including the heterocyclic compound represented by Formula 1, e.g., an organic light-emitting device, may have high efficiency, a low driving voltage, high luminance, and a long lifespan.



137

<Formula 1a>




[0073] Also, the core structure may include a dibenzofuran moiety or a dibenzothiophene moiety (as F₁) in a structure, wherein the dibenzofuran moiety and the dibenzothiophene moiety includes an oxygen atom or a sulfur atom as an atom having high electronegativity, and intermolecular attraction becomes strong, thereby improving heat resistance. An intermolecular distance may be shortened, and molecular packing may be increased. As a result, it may be expected to improve a charge hopping rate due to an improved transfer integral. Therefore, heat resistance, stability to charges, and charge transport capability of the heterocyclic compound represented by Formula 1 may be further improved.

[0074] Synthesis methods of the heterocyclic compound represented by Formula 1 may include suitable synthetic methods.

[0075] At least one of the heterocyclic compounds represented by Formula 1 may be used or included between a pair of electrodes of an organic light-emitting device. For example, the heterocyclic compound may be included in at least one of a hole transport region and/or an emission layer.

[0076] Thus, another aspect provides an organic light-emitting device that includes: a first electrode; a second



electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode and includes an emission layer. The organic layer may include at least one heterocyclic compound represented by Formula 1.

[0077] The expression that “(an organic layer) includes at least one heterocyclic compound” used herein may include a case in which “(an organic layer) includes an identical heterocyclic compound represented by Formula 1 and a case in which (an organic layer) includes two or more different heterocyclic compounds represented by Formula 1”.

[0078] For example, the organic layer may include only Compound 1 as the heterocyclic compound. In this regard, Compound 1 may be only in the electron transport layer of the organic light-emitting device. In an implementation, the organic layer may include Compound 1 and Compound 2 as the heterocyclic compound. In this regard, Compound 1 and Compound 2 may be in an identical layer (e.g., Compound 1 and Compound 2 both may be in an electron transport layer), or may be in different layers (e.g., Compound 1 may be in an electron transport layer, and Compound 2 may be in an electron injection layer).

[0079] In an implementation, the first electrode of the organic light-emitting device may be an anode.

[0080] In an implementation, the second electrode of the organic light-emitting device may be a cathode.

[0081] In an implementation, the organic layer may include i) a hole transport region between the first electrode and the emission layer, the hole transport region including at least one selected from a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer, and ii) an electron transport region between the emission layer and the second electrode, the electron transport region including at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer. For example, the electron transport region may include a single layer or layers.

[0082] In an implementation, at least one of the electron transport region and the emission layer may include at least one of the heterocyclic compounds. For example, the electron transport region of the organic light-emitting device may include an electron transport layer or an electron injection layer, and the electron transport layer or the electron injection layer may include at least one of the heterocyclic compounds. In an implementation, the emission layer of the organic light-emitting device may include a pyrene-based compound, an arylamine-based compound, or a styryl-based compound. The heterocyclic compound included in the emission layer may act as a host, and in this case, the emission layer may further include a dopant. The dopant may be a phosphorous dopant or a fluorescent dopant.

[0083] The term “organic layer” used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of the organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

[0084] FIG. 1 illustrates a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 may include a first electrode 110, an organic layer 150, and a second electrode 190.

[0085] Hereinafter, the structure of the organic light-emitting device 10 according to an embodiment and a method of

manufacturing the organic light-emitting device 10 will be described in connection with FIG. 1.

[0086] In an implementation, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be a glass substrate or a plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water-resistance.

[0087] The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode 110 on the substrate. When the first electrode 110 is an anode, the material for a first electrode may be selected from materials with a high work function to facilitate hole injection.

[0088] The first electrode 110 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode 110 is a transmissible electrode, a material for forming a first electrode may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), zinc oxide (ZnO), and any combinations thereof. In an implementation, when the first electrode 110 is a semi-transmissible electrode or a reflective electrode, a material for forming a first electrode may be selected from magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and any combinations thereof.

[0089] The first electrode 110 may have a single-layered structure, or a multi-layered structure including two or more layers. In an implementation, the first electrode 110 may have a three-layered structure of ITO/Ag/ITO.

[0090] The organic layer 150 is disposed on the first electrode 110. The organic layer 150 may include an emission layer.


[0091] The organic layer 150 may further include a hole transport region between the first electrode 110 and the emission layer, and an electron transport region between the emission layer and the second electrode 190.

[0092] The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.


[0093] The hole transport region may include at least one layer selected from a hole injection layer (HIL), a hole transport layer (HTL), an emission auxiliary layer, and an electron blocking layer (EBL).

[0094] For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a structure of hole injection layer/hole transport layer, hole injection layer/hole transport layer/emission auxiliary layer, hole injection layer/emission auxiliary layer, hole transport layer/emission auxiliary layer or hole injection layer/hole transport layer/electron blocking layer, wherein for each structure, constituting layers are sequentially stacked from the first electrode 110 in this stated order.


[0095] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPD), β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), polyaniline/poly(4-styrenesulfonate) (Pani/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:




m-MTDATA





2-TNATA



NPB



TDATA

 β -NPB

TPD

ylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylylene group, a phenanthrenylylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylylylene group, a pyrenylylene group, a chrysenylylene group, a naphthacenylylene group, a picenylylene group, a perylenylylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group; and

[0106] a phenylene group, a pentalenylylene group, an indenylene group, a naphthylene group, an azulenylylene group, a heptalenylylene group, an indacenylylene group, an acenaphthylene group, a fluorenylylene group, a spiro-bifluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenalenylylene group, a phenanthrenylylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylylylene group, a pyrenylylene group, a chrysenylylene group, a naphthacenylylene group, a picenylylene group, a perylenylylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a thiophenyl group, a furanyl group, a carbazolylylene group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

[0107] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0108] In one or more embodiments, xa1 to xa4 may each independently be 0, 1, or 2.

[0109] In one or more embodiments, xa5 may be 1, 2, 3, or 4.

[0110] In one or more embodiments, R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from:

[0111] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a thiophenyl group, a furanyl group, a carbazolylylene group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0112] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a thiophenyl group, a furanyl group, a carbazolylylene group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a thiophenyl group, a furanyl group, a carbazolylylene group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

[0113] wherein Q₃₁ to Q₃₃ are the same as described above.

[0114] In one or more embodiments, at least one selected from R₂₀₁ to R₂₀₃ in Formula 201 may each independently be selected from:

[0115] a fluorenyl group, a spiro-bifluorenyl group, a carbazolylylene group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0116] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.

[0117] In one or more embodiments, in Formula 202, i) R₂₀₁ and R₂₀₂ may be linked via a single bond, and/or ii) R₂₀₃ and R₂₀₄ may be linked via a single bond.


[0118] In one or more embodiments, at least one selected from R₂₀₁ to R₂₀₄ in Formula 202 may be selected from:

[0119] a carbazolyl group; and

[0120] a carbazolyl group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.


[0121] The compound represented by Formula 201 may be represented by Formula 201A:

<Formula 201A>




[0122] In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A(1):

<Formula 201A(1)>




[0123] In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A-1:

<Formula 201A-1>




[0124] In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A:

<Formula 202A>



[0125] In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A-1:

<Formula 202A-1>



[0126] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,

[0127] L₂₀₁ to L₂₀₃, xa1 to xa3, xa5, and R₂₀₂ to R₂₀₄ are the same as described above,

[0128] R₂₁₁ and R₂₁₂ are the same as described in connection with R₂₀₃,


[0129] R₂₁₃ to R₂₁₇ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl

group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a


perylene group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0130] The hole transport region may include at least one compound selected from Compounds HT1 to HT39:


HT1




HT2



HT3




HT4




-continued


HT5




HT6



HT7




HT8




-continued


HT9




HT10



HT11





HT12



-continued



HT13

HT14





HT15

HT16




HT17


HT18




-continued




HT19



HT20




HT21



HT22



-continued

HT23





HT24

HT25




HT26

HT27




-continued


HT28




HT29




HT30




HT31



HT32




HT33




-continued


HT34




HT35




HT36




HT37



HT38



HT39



[0131] A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 9,000 Å, e.g., about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, e.g., about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0132] The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer, and the electron blocking layer may block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include the materials as described above.

[0133] The hole transport region may further include, in addition to these materials, a charge-generation material for improving conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0134] The charge-generation material may be, for example, a p-dopant.

[0135] In one or more embodiments, a lowest unoccupied molecular orbital (LUMO) level of the p-dopant may be -3.5 eV or less.

[0136] The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound.



[0137] For example, the p-dopant may include at least one selected from:

[0138] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) or 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);


[0139] a metal oxide, such as a tungsten oxide or a molybdenum oxide;

[0140] 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and

[0141] a compound represented by Formula 221 below:



<Formula 221>



[0142] In Formula 221,

[0143] R_{221} to R_{223} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, wherein at least one selected from R_{221} to R_{223} has at least one substituent selected from a cyano group, $-F$, $-Cl$, $-Br$, $-I$, a C_1 - C_{20} alkyl group substituted with $-F$, a C_1 - C_{20} alkyl group substituted with $-Cl$, a C_1 - C_{20} alkyl group substituted with $-Br$, and a C_1 - C_{20} alkyl group substituted with $-I$.


[0144] When the organic light-emitting device 10 is a full-color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In an implementation, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers contact each other or are separated from each other. In an implementation, the emission layer may include two or more materials selected from a red-light emission material, a green-light emission material, and a blue-light emission material, in which the two or more materials are mixed with each other in a single layer to emit white light.

[0145] The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant and a fluorescent dopant.

[0146] An amount of the dopant in the emission layer may be, in general, in a range of about 0.01 to about 15 parts by weight based on 100 parts by weight of the host.

[0147] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0148] In an implementation, the host may further include, the heterocyclic compound represented by Formula 1, a compound represented by Formula 301 below:



<Formula 301>

[0149] In Formula 301,

[0150] Ar_{301} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0151] $xb11$ may be 1, 2, or 3,

[0152] L_{301} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0153] $xb1$ may be an integer selected from 0 to 5,

[0154] R_{301} may be selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{301})(Q_{302})(Q_{303})$, $-N(Q_{301})(Q_{302})$, $-B(Q_{301})(Q_{302})$, $-C(=O)(Q_{301})$, $-S(=O)_2(Q_{301})$, and $-P(=O)(Q_{301})(Q_{302})$,

[0155] $xb21$ may be an integer selected from 1 to 5,

[0156] Q_{301} to Q_{303} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0157] In an implementation, Ar_{301} in Formula 301 may be selected from:

[0158] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group; and


[0159] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})(Q_{32})$,

[0160] wherein Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{100} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.


[0161] In Formula 301, when $xb11$ is two or more, two or more $Ar_{301}(s)$ may be linked via a single bond.

[0162] In one or more embodiments, the compound represented by Formula 301 may be represented by Formula 301-1 or 301-2:

<Formula 301-1>



<Formula 301-2>



[0163] wherein, in Formulae 301-1 to 301-2,

[0164] A_{301} to A_{304} may each independently be selected from a benzene, a naphthalene, a phenanthrene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a pyridine, a pyrimidine, an indene, a fluorene, a spiro-bifluorene, a benzofluorene, a dibenzofluorene, an indole, a carbazole, benzocarbazole, a dibenzocarbazole, a furan, a benzofuran, a dibenzofuran, a naphthofuran, a benzonaphthofuran, a dinaphthofuran, a thiophene, a benzothiophene, a dibenzothiophene, a naphthothiophene, benzonaphthothiophene, and a dinaphthothiophene,

[0165] X_{301} may be O, S, or N- $[(L_{304})_{xb4}-R_{304}]$,

[0166] R_{311} to R_{314} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),

[0167] $xb22$ and $xb23$ may each independently be 0, 1, or 2,

[0168] L_{301} , $xb1$, R_{301} , and Q_{31} to Q_{33} are the same as described above,

[0169] L_{302} to L_{304} are each independently the same as described in connection with L_{301} ,

[0170] $xb2$ to $xb4$ are each independently the same as described in connection with $xb1$, and

[0171] R_{302} to R_{304} are each independently the same as described in connection with R_{301} .

[0172] For example, L_{301} to L_{304} in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0173] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazolylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a thiadiazolylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinolylylene group, a quinolinolylylene group, an isoquinolinolylylene group, a benzoquinolinolylylene group, a phthalazinolylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinnolinolylylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthrolinylylene group, a phenazinolylylene group, a benzimidazolylylene group, an isobenzothiazolylylene group, a benzoxazolylylene group, an isobenzoxazolylylene group, a triazolylylene group, a tetrazolylylene group, an imidazopyridinylylene group, an imidazopyrimidinylylene group, and an azacarbazolylylene group; and

[0174] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene

group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazolylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinolylylene group, a quinolinolylylene group, an isoquinolinolylylene group, a benzoquinolinolylylene group, a phthalazinolylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinnolinolylylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthrolinylylene group, a phenazinolylylene group, a benzimidazolylylene group, an isobenzothiazolylylene group, a benzoxazolylylene group, an isobenzoxazolylylene group, a triazolylylene group, a tetrazolylylene group, an imidazopyridinylylene group, an imidazopyrimidinylylene group, and an azacarbazolylylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylyl group, a quinazolinylyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),

[0175] wherein Q_{31} to Q_{33} are the same as described above.

[0176] In one or more embodiments, R_{301} to R_{304} in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0177] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group,

a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, and an azacarbazolyl group; and


[0178] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group,

a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),


[0179] wherein Q₃₁ to Q₃₃ are the same as described above.

[0180] In an implementation, the host may include an alkaline-earth metal complex. For example, the host may be selected from a Be complex (for example, Compound H55), an Mg complex, and a Zn complex.


[0181] The host may include, in addition to the heterocyclic compound represented by Formula 1 at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalene-2-yl)anthracene (MADN), 9,10-di-(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolyl-benzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55:



H1




H2




H3






-continued







-continued



-continued




-continued




-continued


H24



H25




H26




-continued


H27



H28




H29




-continued


H30




H31




H32



H33




H34




-continued


H35




H36




H37



H38




-continued




H39


-continued




H42




H40



H43




H41




H44


-continued




H45




H46



H47




H48




H49


-continued




H50




H51



H52




H53




H54

-continued



H55

[0182] The phosphorescent dopant may include an organometallic complex represented by Formula 401:



[0183] In Formulae 401 and 402,

[0184] M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

[0185] L₄₀₁ may be selected from ligands represented by Formula 402, and xc1 may be 1, 2, or 3, wherein, when xc1 is two or more, two or more L₄₀₁(s) may be identical to or different from each other,

[0186] L₄₀₂ may be an organic ligand, and xc2 may be an integer selected from 0 to 4, wherein, when xc2 is two or more, two or more L₄₀₂(s) may be identical to or different from each other,

[0187] X₄₀₁ to X₄₀₄ may each independently be nitrogen or carbon,

[0188] X₄₀₁ and X₄₀₃ may be linked via a single bond or a double bond, and X₄₀₂ and X₄₀₄ may be linked via a single bond or a double bond,

[0189] A₄₀₁ and A₄₀₂ may each independently be a C₅-C₆₀ cyclic group or a C₁-C₆₀ heterocyclic group,

[0190] X₄₀₅ may be a single bond, *—O—*, *—S—*, *—C(=O)—*, *—N(Q₄₁₁)—*, *—C(Q₄₁₁)(Q₄₁₂)—*, *—C(Q₄₁₁)=C(Q₄₁₂)—*, *—C(Q₄₁₁)=*, or *=C(Q₄₁₁)=*, wherein Q₄₁₁ and Q₄₁₂ may be hydrogen, deuterium, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

[0191] X₄₀₆ may be a single bond, O, or S,

[0192] R₄₀₁ and R₄₀₂ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl

group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₂₀ alkyl group, a substituted or unsubstituted C₁-C₂₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group —Si(Q₄₀₁)(Q₄₀₂)(Q₄₀₃), —N(Q₄₀₁)(Q₄₀₂), —B(Q₄₀₁)(Q₄₀₂), —C(=O)(Q₄₀₁), —S(=O)₂(Q₄₀₁), and —P(=O)(Q₄₀₁)(Q₄₀₂), wherein Q₄₀₁ to Q₄₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a C₆-C₂₀ aryl group, and a C₁-C₂₀ heteroaryl group,

[0193] xc11 and xc12 may each independently be an integer selected from 0 to 10, and

[0194] * and *' in Formula 402 each indicate a binding site to M in Formula 401.

[0195] In one or more embodiments, A₄₀₁ and A₄₀₂ in Formula 402 may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiothiophene group, an isobenzothiothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiothiophene group.

[0196] In one or more embodiments, in Formula 402, i) X₄₀₁ may be nitrogen and X₄₀₂ may be carbon, or ii) X₄₀₁ and X₄₀₂ may each be nitrogen at the same time.

[0197] In one or more embodiments, R₄₀₁ and R₄₀₂ in Formula 402 may each independently be selected from:

[0198] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0199] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

[0200] a cyclopentyl group, a cyclohexyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuran group, and a dibenzothiothiophenyl group;

[0201] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranlyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranlyl group, and a dibenzothiophenyl group; and


[0202] —Si(Q₄₀₁)(Q₄₀₂)(Q₄₀₃), —N(Q₄₀₁)(Q₄₀₂), —B(Q₄₀₁)(Q₄₀₂), —C(=O)(Q₄₀₁), —S(=O)₂(Q₄₀₁), and —P(=O)(Q₄₀₁)(Q₄₀₂),

[0203] wherein Q₄₀₁ to Q₄₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group.

[0204] In one or more embodiments, when xc1 in Formula 401 is two or more, two A₄₀₁(s) in two or more L₄₀₁(s) may optionally be linked via X₄₀₇, which is a linking group, or two A₄₀₂(s) in two or more L₄₀₂(s) may optionally be linked via X₄₀₈, which is a linking group (see Compounds PD1 to PD4 and PD7). X₄₀₇ and X₄₀₈ may each independently be a single bond, *—O—*, *—S—*, *—C(=O)—*, *—N(Q₄₁₃)—*, *C(Q₄₁₃)(Q₄₁₄)—*, or *—C(Q₄₁₃)=C(Q₄₁₄)—* (wherein Q₄₁₃ and Q₄₁₄ may each independently be hydrogen, deuterium, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group).


[0205] L₄₀₂ in Formula 401 may be a monovalent, divalent, or trivalent organic ligand. For example, L₄₀₂ may be selected from a halogen, diketone (for example, acetylacetonate), a carboxylic acid (for example, picolinate), —C(=O), an isonitrile, —CN, and phosphorus (for example, phosphine or phosphite).

[0206] In one or more embodiments, the phosphorescent dopant may be selected from, for example, Compounds PD1 to PD25:




PD1


-continued




PD2




PD3



PD4








PD5




PD6

-continued




-continued

PD7




PD12

PD8




PD13

PD9




PD14

PD10








PD15

PD11




PD16

-continued




-continued

PD17




PD22

PD18



PD23


PD19



PD24

PD20


PD21



PD25

[0207] The fluorescent dopant may include an arylamine compound or a styrylamine compound.

[0208] The fluorescent dopant may include a compound represented by Formula 501:



[0209] In Formula 501,

[0210] Ar_{501} may be a substituted or unsubstituted $\text{C}_5\text{-C}_{60}$ carbocyclic group or a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heterocyclic group,

[0211] L_{501} to L_{503} may each independently be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkylene group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenylene group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0212] $xd1$ to $xd3$ may each independently be an integer selected from 0 to 3,

[0213] R_{501} and R_{502} may each independently be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryloxy group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylthio group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

[0214] $xd4$ may be an integer selected from 1 to 6.

[0215] In one or more embodiments, Ar_{501} in Formula 501 may be selected from:

[0216] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group; and

[0217] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{20}$ alkyl group, a $\text{C}_1\text{-C}_{20}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0218] In one or more embodiments, L_{501} to L_{503} in Formula 501 may each independently be selected from:

[0219] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylene group, an indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, and a pyridinylene group; and

[0220] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylene group, an indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, and a pyridinylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{20}$ alkyl group, a $\text{C}_1\text{-C}_{20}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0221] In one or more embodiments, R_{501} and R_{502} in Formula 502 may each independently be selected from:

[0222] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0223] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl

group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃).


[0224] wherein, in Formula 501, Q₃₁ to Q₃₃ may be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0225] In one or more embodiments, xd4 in Formula 501 may be 3.


[0226] For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:

-continued


FD3




FD1



FD4




FD2




-continued

FD5




-continued


FD8




FD6




FD9




FD7










FD10



-continued




-continued




-continued

FD19



-continued



[0228] The electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0229] The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer.

[0230] For example, the electron transport region may have a structure of electron transport layer/electron injection layer, a structure of hole blocking layer/electron transport layer/electron injection layer, a structure of electron control layer/electron transport layer/electron injection layer, or a structure of buffer layer/electron transport layer/electron injection layer, wherein for each structure, constituting layers are sequentially stacked from an emission layer.

[0231] The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include a metal-free compound containing at least one π electron-depleted nitrogen-containing ring.

[0232] The “ π electron-depleted nitrogen-containing ring” indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

[0233] For example, the “ π electron-depleted nitrogen-containing ring” may be i) a 5-membered to 7-membered hetero monocyclic group having at least one $*-N=*$ moiety, ii) a heteropoly cyclic group in which two or more 5-membered to 7-membered hetero monocyclic groups each having at least one $*-N=*$ moiety are condensed with each other, or iii) a heteropoly cyclic group in which at least one of 5-membered to 7-membered hetero monocyclic groups, each having at least one $*-N=*$ moiety, is condensed with at least one C_5 - C_{60} carbocyclic group.

[0234] Examples of the π electron-depleted nitrogen-containing ring include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoxaline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, thiadiazol, an imidazopyridine, an imidazopyrimidine, and an azacarbazole.

[0235] In one or more embodiments, the electron transport region may further include, in addition to the heterocyclic compound represented by Formula 1, a compound represented by Formula 601.



[0236] In Formula 601,

[0237] Ar_{601} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0238] $\text{xe}11$ may be 1, 2, or 3,

[0239] L_{601} is selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0240] $\text{xe}1$ may be an integer selected from 0 to 5,

[0241] R_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_{601})(\text{Q}_{602})(\text{Q}_{603})$, $-\text{C}(=\text{O})(\text{Q}_{601})$, $-\text{S}(=\text{O})_2(\text{Q}_{601})$, and $-\text{P}(=\text{O})(\text{Q}_{601})(\text{Q}_{602})$,

[0242] Q_{601} to Q_{603} may each independently be a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

[0243] $\text{xe}21$ may be an integer selected from 1 to 5.

[0244] In one or more embodiments, at least one selected from $\text{Ar}_{601}(\text{s})$ in the number of $\text{xe}11$ and $\text{R}_{601}(\text{s})$ in the number of $\text{xe}21$ may include the π electron-depleted nitrogen-containing ring described above.

[0245] In one or more embodiments, ring Ar_{601} in Formula 601 may be selected from:

[0246] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinox-

line group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

[0247] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,


[0248] wherein, in Formula 601, Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0249] When $\text{xe}11$ in Formula 601 is two or more, two or more $\text{Ar}_{601}(\text{s})$ may be linked via a single bond.

[0250] In one or more embodiments, Ar_{601} in Formula 601 may be an anthracene group.

[0251] In one or more embodiments, a compound represented by Formula 601 may be represented by Formula 601-1:

<Formula 601-1>



[0252] In Formula 601-1,

[0253] X_{614} may be N or C(R_{614}), X_{615} may be N or C(R_{615}), X_{616} may be N or C(R_{616}), and at least one selected from X_{614} to X_{616} may be N,

[0254] L_{611} to L_{613} may each independently be substantially the same as described in connection with L_{601} ,

[0255] $xe611$ to $xe613$ may each independently be substantially the same as described in connection with $xe1$,

[0256] R_{611} to R_{613} may each independently be substantially the same as described in connection with R_{601} , and

[0257] R_{614} to R_{616} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0258] In one or more embodiments, L_{601} and L_{611} to L_{613} in Formulae 601 and 601-1 may each independently be selected from:

[0259] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

[0260] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a

dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenylyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinylyl group, an isoquinolinylyl group, a benzoquinolinylyl group, a phthalazinyl group, a naphthyridinylyl group, a quinoxalinylyl group, a quinazolinylyl group, a cinnolinylyl group, a phenanthridinylyl group, an acridinylyl group, a phenanthrolinylyl group, a phenazinylyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group.

[0261] In one or more embodiments, $xe1$ and $xe611$ to $xe613$ in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

[0262] In one or more embodiments, R_{601} and R_{611} to R_{613} in Formula 601 and 601-1 may each independently be selected from:

[0263] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenylyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an

isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;

[0264] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl


group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

[0265] —S(=O)₂(Q₆₀₁) and —P(=O)(Q₆₀₁)(Q₆₀₂),


[0266] wherein Q₆₀₁ and Q₆₀₂ are the same as described above.

[0267] The electron transport region may include at least one compound selected from Compounds ET1 to ET36:


ET1








ET2




-continued



-continued




-continued




ET9

-continued





ET11

ET12






ET10

ET13




-continued




-continued


ET14



ET15



ET16




ET17

ET18


ET19

-continued




ET20


-continued




ET23




ET21



ET24




ET22




ET25

-continued




ET26

-continued





ET29


ET27



ET28




ET30




ET31


-continued



ET32




ET33




ET34

-continued




ET35




ET36


[0268] In one or more embodiments, the electron transport region may include at least one selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq₃, BALq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ.



Alq₃



BAlq




[0269] A thickness of the buffer layer, the hole blocking layer, or the electron control layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.

[0270] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

[0271] The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.

[0272] The metal-containing material may include at least one selected from alkali metal complex and alkaline earth-metal complex. The alkali metal complex may include a metal ion selected from an Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion, and the alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, an Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenylan oxazole, a hydroxy phenylthiazole, a hydroxy diphenylan oxadiazole, a hydroxy diphenylthiadiazol, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene.


[0273] For example, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.



ET-D1

-continued

ET-D2



[0274] The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode 190. The electron injection layer may directly contact the second electrode 190.

[0275] The electron injection layer may have i) a single-layered structure including a single layer including a plurality of different materials, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0276] The electron injection layer may include alkali metal, alkaline earth metal, rare-earth metal, alkali metal compound, alkaline earth-metal compound, rare-earth metal compound, alkali metal complex, alkaline earth-metal complex, rare-earth metal complex, or any combinations thereof.

[0277] The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one embodiment, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs.

[0278] The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

[0279] The rare-earth metal may be selected from Sc, Y, Ce, Yb, Gd, and Tb.

[0280] The alkali metal compound, the alkaline earth-metal compound, and the rare-earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal and rare-earth metal.

[0281] The alkali metal compound may be selected from alkali metal oxides, such as Li₂O, Cs₂O, or K₂O, and alkali metal halides, such as LiF, NaF, CsF, KF, LiI, NaI, CsI, KI, or RbI. In one embodiment, the alkali metal compound may be selected from LiF, Li₂O, NaF, LiI, NaI, CsI, and KI.

[0282] The alkaline earth-metal compound may be selected from alkaline earth-metal compounds, such as BaO, SrO, CaO, Ba_xSr_{1-x}O (0<x<1), Ba_xCa_{1-x}O (0<x<1). In one embodiment, the alkaline earth-metal compound may be selected from BaO, SrO, and CaO.

[0283] The rare-earth metal compound may be selected from YbF₃, ScF₃, ScO₃, Y₂O₃, Ce₂O₃, GdF₃, and TbF₃. In one embodiment, the rare-earth metal compound may be selected from YbF₃, ScF₃, TbF₃, YbI₃, ScI₃, and TbI₃.

[0284] The alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare-earth metal as described above, and a ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may each independently be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenylan oxazole, hydroxy phenylthiazole, hydroxy diphenylan oxadiazole, hydroxy diphenylthiadiazol, hydroxy phenylpyridine,

hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene.

[0285] The electron injection layer may consist of alkali metal, alkaline earth metal, rare-earth metal, alkali metal compound, alkaline earth-metal compound, rare-earth metal compound, alkali metal complex, alkaline earth-metal complex, rare-earth metal complex, or any combinations thereof, as described above. In one or more embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, alkali metal, alkaline earth metal, rare-earth metal, alkali metal compound, alkaline earth-metal compound, rare-earth metal compound, alkali metal complex, alkaline earth-metal complex, rare-earth metal complex, or any combinations thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

[0286] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, e.g., about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

[0287] The second electrode 190 may be disposed on the organic layer 150 having such a structure. The second electrode 190 may be a cathode which is an electron injection electrode, and in this regard, a material for forming the second electrode 190 may be selected from metal, an alloy, an electrically conductive compound, and a mixture thereof, which have a relatively low work function.

[0288] The second electrode 190 may include at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO. The second electrode 190 may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

[0289] The second electrode 190 may have a single-layered structure, or a multi-layered structure including two or more layers.

[0290] An organic light-emitting device 20 of FIG. 2 may include a first capping layer 210, a first electrode 110, an organic layer 150, and a second electrode 190 which are sequentially stacked in this stated order, an organic light-emitting device 30 of FIG. 3 may include a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220 which are sequentially stacked in this stated order, and an organic light-emitting device 40 of FIG. 4 may include a first capping layer 210, a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220.

[0291] Regarding FIGS. 2 to 4, the first electrode 110, the organic layer 150, and the second electrode 190 may be understood by referring to the description presented in connection with FIG. 1.

[0292] In the organic layer 150 of each of the organic light-emitting devices 20 and 40, light generated in an emission layer may pass through the first electrode 110, which is a semi-transmissive electrode or a transmissive electrode, and the first capping layer 210 toward the outside, and in the organic layer 150 of each of the organic light-emitting devices 30 and 40, light generated in an emission layer may pass through the second electrode 190, which is a semi-transmissive electrode or a transmissive electrode, and the second capping layer 220 toward the outside.


[0293] The first capping layer 210 and the second capping layer 220 may increase external luminescent efficiency according to the principle of constructive interference.

[0294] The first capping layer 210 and the second capping layer 220 may each independently be an organic capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.


[0295] At least one selected from the first capping layer 210 and the second capping layer 220 may each independently include at least one material selected from carbocyclic compounds, heterocyclic compounds, amine-based compounds, porphyrine derivatives, phthalocyanine derivatives, naphthalocyanine derivatives, alkali metal complexes, and alkaline earth-based complexes. The carbocyclic compound, the heterocyclic compound, and the amine-based compound may be optionally substituted with a substituent containing at least one element selected from O, N, S, Se, Si, F, Cl, Br, and I. In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include an amine-based compound.

[0296] In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include the compound represented by Formula 201 or the compound represented by Formula 202.

[0297] In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include a compound selected from Compounds HT28 to HT33 and Compounds CP1 to CP5.



-continued



[0298] Hereinbefore, the organic light-emitting device according to an embodiment has been described in connection with FIGS. 1 through 4.

[0299] Layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region may be formed in a certain region by using one or more suitable methods selected from vacuum deposition, spin coating, casting, langmuir-blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

[0300] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of about 100 to about 500° C., at a vacuum degree of about 10⁻⁸ to about 10⁻³ torr, and at a deposition rate of about 0.01 to about 100 Å/sec by taking into account a compound to be included in a layer to be formed and the structure of a layer to be formed.

[0301] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to 200° C. by taking into account a compound to be included in a layer to be formed and the structure of a layer to be formed.

[0302] The term “C₁-C₆₀ alkyl group” as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “C₁-C₆₀ alkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₆₀ alkyl group.

[0303] The term “C₂-C₆₀ alkenyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon double bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C₂-C₆₀ alkenylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkenyl group.

[0304] The term “C₂-C₆₀ alkynyl group” used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethynyl group and a propynyl group. The term “C₂-C₆₀ alkynylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

[0305] The term “C₁-C₆₀ alkoxy group” as used herein refers to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is the C₁-C₆₀ alkyl group), and examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

[0306] The term “C₃-C₁₀ cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

[0307] The term “C₁-C₁₀ heterocycloalkyl group” used herein refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuranlyl group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group,” used herein, refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

[0308] The term “C₃-C₁₀ cycloalkenyl group” used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and does not have aromaticity, and examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group” used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

[0309] The term “C₁-C₁₀ heterocycloalkenyl group” used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Examples of the C₁-C₁₀ heterocycloalkenyl group are a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranlyl group and a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group,” used herein, refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkenyl group.

[0310] The term “C₆-C₆₀ aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C₆-C₆₀ arylene group used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Examples of the C₆-C₆₀ aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be fused to each other.

[0311] The term “C₁-C₆₀ heteroaryl group” used herein refers to a monovalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. A C₁-C₆₀ heteroarylene group used herein refers to a divalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Examples of the C₁-C₆₀ heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be fused to each other.

[0312] The term “C₆-C₆₀ aryloxy group” used herein indicates —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and a C₆-C₆₀ arylthio group indicates —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

[0313] The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed with each other, only carbon atoms as a ring-forming atom, and non-aromaticity in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group,” used herein, refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0314] The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group (for example, having 1 to 60 carbon atoms) that has two or more rings condensed to each other, has at least one heteroatom selected from N, O, Si, P, and S, other than carbon atoms, as a ring-forming atom, and has non-aromaticity in the entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group,” used herein, refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0315] The term “C₅-C₆₀ carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which a ring-forming atom is a carbon atom only. The C₅-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₅-C₆₀ carbocyclic group may be a ring, such as a benzene group, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In one or more embodiments, depending on the number of substituents connected to the C₅-C₆₀ carbocyclic group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

[0316] The term “C₁-C₆₀ heterocyclic group” used herein refers to a group having the same structure as the C₅-C₆₀ carbocyclic group, except that as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used in addition to carbon (the number of carbon atoms may be in a range of 1 to 60).

[0317] At least one substituent of the substituted C₅-C₆₀ carbocyclic group, the substituted C₁-C₆₀ heterocyclic group, the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the sub-

stituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0318] deuterium (—D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0319] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

[0320] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0321] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

[0322] —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂).

[0323] Unless otherwise defined herein, Q₁ to Q₆, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic con-

condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

[0324] The term “Ph” used herein refers to a phenyl group, the term “Me” used herein refers to a methyl group, the term “Et” used herein refers to an ethyl group, the term “ter-Bu” or “Bu” used herein refers to a tert-butyl, and the term “OMe” used herein refers to a methoxy group.

[0325] The term “biphenyl group” as used herein refers to “a phenyl group substituted with a phenyl group.” In other words, the “biphenyl group” is a substituted phenyl group having a C₆-C₆₀ aryl group as a substituent.

[0326] The term “terphenyl group” as used herein refers to “a phenyl group substituted with a biphenyl group.” In other words, the “terphenyl group” is a substituted phenyl group having, as a substituent, a C₆-C₆₀ aryl group substituted with a C₆-C₆₀ aryl group.

[0327] * and *¹ used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.

[0328] Hereinafter, a compound according to embodiments and an organic light-emitting device according to embodiments will be described in detail with reference to Synthesis Examples and Examples. The wording “B was used instead of A” used in describing Synthesis Examples refers to that an identical molar equivalent of B was used in place of A.


[0329] The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples.

EXAMPLE


Synthesis Example

Synthesis Example 1: Synthesis of Compound 1

[0330]



-continued




[0331] Synthesis of Compound 1


[0332] 1.34 g (10 mmol) of 2-coumaranone, 1.83 g (10 mmol) of dibenzo[b,d]furan-4-amine, 3.32 g (10 mmol) of 10-(naphthalene-1-yl)anthracene-9-carbaldehyde, and 0.013 g (0.5 mmol) of diiodine were dissolved in 150 mL of THF and were then stirred at a temperature of 80° C. for 24 hours. The reacting solution was cooled to ambient temperature. Then, an organic layer was extracted three times by using 60 mL of water and 60 mL of diethyl ether. The organic layer was dried by using magnesium sulfate and the solvent was evaporated. The obtained residue was separated and purified by silica gel column chromatography, thereby completing the preparation of 3.67 g (60%) of Compound 1. Compound 1 was confirmed through LC-MS and ¹H NMR. C₄₅H₂₅NO₂: M+1 611.72

Synthesis Example 2: Synthesis of Compound 20

[0333]



-continued



compound 20

Example 1, except that 3-bromobenzaldehyde was used instead of 10-(naphthalene-1-yl)anthracene-9-carbaldehyde. Intermediate Compound 20-1 was confirmed through LC-MS. $C_{27}H_{14}BrNO_2$: M+1 464.31


[0336] Synthesis of Compound 20

[0337] 4.64 g (10 mmol) of Intermediate Compound 20-1, 3.45 g (10 mmol) of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile, 0.58 g (0.5 mmol) of $Pd(PPh_3)_4$, and 4.15 g (30.0 mmol) of K_2CO_3 were dissolved in 150 mL of a mixed solution of THF/ H_2O (2/1) and were then stirred at a temperature of 80° C. for 16 hours. The reacting solution was cooled to ambient temperature. Then, an organic layer was extracted three times by using 60 mL of water and 60 mL of diethyl ether. The organic layer was dried by using magnesium sulfate and the solvent was evaporated. The obtained residue was separated and purified by silica gel column chromatography, thereby completing the preparation of 4.2 g (70%) of Compound 20. Compound 20 was confirmed through LC-MS and 1H NMR. $C_{43}H_{26}N_2O_2$: M+1 602.71


[0334] Synthesis of Intermediate 20-1

[0335] 3 g (65%) of an Intermediate Compound 20-1 was prepared by following the same steps as in Synthesis

Synthesis Example 3: Synthesis of Compound 23

[0338]

23-1



compound 23






[0339] Synthesis of Intermediate 23-1

[0340] 2.78 g (60%) of Intermediate Compound 23-1 was prepared in the same manner as in the synthesis of Intermediate 20-1, except that 4-bromobenzaldehyde was used instead of 3-bromobenzaldehyde. Intermediate Compound 23-1 was confirmed through LC-MS. $C_{27}H_{14}BrNO_2$: M+1 464.30

[0341] Synthesis of Compound 23

[0342] 3.27 g (50%) of Compound 23 was prepared in the same manner as in the synthesis of Compound 20, except that Intermediate Compound 23-1 was used instead of Intermediate Compound 20-1, and 4-(9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluoren-2-yl)pyridine was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. Compound 23 was confirmed through LC-MS and 1H NMR. $C_{47}H_{30}N_2O_2$: M+1 654.75

Synthesis Example 4: Synthesis of Compound 26



[0343]**[0344]** Synthesis of Compound 26

[0345] 4.93 g (80%) of Compound 26 was prepared in the same manner as in the synthesis of Compound 20, except that 2,4-diphenyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,5-triazine was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. Compound 26 was confirmed through LC-MS and 1H NMR. $C_{42}H_{24}N_4O_2$: M+1 616.66

Synthesis Example 5: Synthesis of Compound 64

[0346]

-continued



compound 64

[0347] Synthesis of Intermediate 64-1

[0348] 1.4 g (30%) of Intermediate Compound 64-1 was prepared in the same manner as in the synthesis of Intermediate 20-1, except that dibenzo[b,d]furan-2-amine was used instead of dibenzo[b,d]furan-4-amine, and 4-bromobenzaldehyde was used instead of 3-bromobenzaldehyde. Intermediate Compound 64-1 was confirmed through LC-MS. $C_{27}H_{14}BrNO_2$; M+1 464.33


[0349] Synthesis of Compound 64


[0350] 4.3 g (70%) of Compound 64 was prepared in the same manner as in the synthesis of Compound 20, except

that Intermediate Compound 64-1 was used instead of Intermediate Compound 20-1, and 3,3'-(5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3-phenylene)dipyridine was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. Compound 64 was confirmed through LC-MS and 1H NMR. $C_{43}H_{25}N_3O_2$; M+1 616.68

Synthesis Example 6: Synthesis of Compound 94

[0351]





[0352] Synthesis of Intermediate 94-1


[0353] 1.2 g (25%) of Intermediate Compound 94-1 was prepared in the same manner as in the synthesis of Intermediate 20-1, except that dibenzo[b,d]furan-2-amine was used instead of dibenzo[b,d]furan-4-amine, and 4-bromobenzaldehyde was used instead of 3-bromobenzaldehyde. Intermediate Compound 94-1 was confirmed through LC-MS. $C_{27}H_{14}BrNO_2$; M+1 464.32

[0354] Synthesis of Compound 94

[0355] 3.91 g (60%) of Compound 94 was prepared in the same manner as in the synthesis of Compound 20, except that Intermediate Compound 94-1 was used instead of Intermediate Compound 20-1, and 9-phenyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-carbazole-2-carbonitrile was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. Compound 94 was confirmed through LC-MS and 1H NMR. $C_{46}H_{27}N_3O_2$; M+1 653.73

Synthesis Example 7: Synthesis of Compound 42

[0356]




[0357] Synthesis of Compound 42

[0358] 4.64 g (10 mmol) of Intermediate 20-1 was dissolved in 100 mL of THF, and 4 mL (2.5 M in hexane) of n-butyllithium was added thereto at a temperature of $-78^\circ C$. 2.20 g (10 mmol) of chlorodiphenylphosphine was slowly added dropwise after 1 hour. Then, the solution was stirred for 3 hours and then heated to ambient temperature. Then, water was added thereto and cleaning was performed three times by using ethyl acetate (30 mL). A cleaned ethyl acetate layer was dried with $MgSO_4$ and evaporated under reduced pressure, and the resultant was dissolved in 40 mL of dichloromethane. 4 mL of peroxide was added to the obtained solution and then stirred at ambient temperature for 20 hours. 20 mL of water was added thereto. Then, an organic layer was extracted three times by using 20 mL of dichloromethane. The organic layer was dried by using magnesium sulfate and the solvent was evaporated. The obtained residue was separated and purified by silica gel column chromatography, thereby completing the prepara-


tion of 4.1 g (70%) of Compound 42. Compound 42 was confirmed through LC-MS and ^1H NMR. $\text{C}_{39}\text{H}_{24}\text{NO}_3\text{P}$: M+1 585.58

Synthesis Example 8: Synthesis of Compound 114

[0359]



-continued



[0360] Synthesis of Intermediate 114-1

[0361] 3.3 g (70%) of Intermediate Compound 114-1 was prepared in the same manner as in the synthesis of Intermediate 20-1, except that dibenzo[b,d]thiophen-4-amine was used instead of dibenzo[b,d]furan-4-amine, and 4-bromoaldehyde was used instead of 3-bromoaldehyde. Intermediate Compound 114-1 was confirmed through LC-MS. $\text{C}_{27}\text{H}_{14}\text{BrNOS}$: M+1 480.40

[0362] Synthesis of Compound 114

[0363] 4.2 g (70%) of Compound 114 was prepared in the same manner as in the synthesis of Compound 42, except that Intermediate Compound 114-1 was used instead of Intermediate Compound 20-1. Compound 114 was confirmed through LC-MS and ^1H NMR. $\text{C}_{39}\text{H}_{24}\text{NO}_2\text{PS}$: M+1 601.67

TABLE 1

Compound	^1H NMR (CDCl_3 , 400 MHz)	MS/FAB	
		found	calc.
1	8.75-8.70 (m, 2H), 8.07-8.04 -, 7.84-7.75 (m, 3H), 7.71-7.65 (m, 4H), 7.60-7.57 (m, 2H), 7.54-7.50 (m, 4H), 7.47-7.40 (m, 4H), 7.35-7.34 (m, 1H), 7.15-7.13 (m, 2H), 6.97-6.95 (m, 1H), 8.81-8.79 (m, 1H), 8.72-8.70 (m, 1H), 8.42-8.40 (m, 1H), 8.07-8.00 (m, 3H), 7.89-7.80 (m, 2H), 7.76-7.72 (m, 2H), 7.67-7.63 (m, 3H), 7.60-7.58 (m, 2H), 7.52-7.50 (m, 2H), 7.46-7.45 (m, 2H), 7.40-7.39 (m, 1H), 1.61-1.57 (m, 6H)	611.72	611.7
20	9.16-9.13 (m, 1H), 8.82-8.79 (m, 5H), 8.76-8.70 (m, 2H), 8.23-8.03 (m, 3H), 7.76-7.67 (m, 3H), 7.62-7.56 (m, 6H), 7.52-7.40 (m, 4H), 8.95-8.90 (m, 3H), 8.82-8.65 (m, 3H), 8.23-8.02 (m, 6H), 7.89-7.81 (m, 6H), 7.74-7.58 (m, 2H), 7.56-7.44 (m, 5H)	602.71	602.69
23	8.25-8.17 (m, 3H), 8.08-8.02 (m, 3H), 7.91-7.82 (m, 3H), 7.74-7.71 (m, 2H), 7.63-7.60 (m, 3H), 7.57-7.50 (m, 5H), 7.32-7.19 (m, 2H), 6.05-5.95 (m, 2H), 5.85-5.82 (m, 1H)	654.75	654.77
26	5.70-5.67 (m, 1H), 5.35-5.31 (m, 1H), 4.89-4.91 (m, 1H)	616.66	616.68
64	8.81-8.70 (m, 2H), 8.52-8.50 (m, 1H), 8.08-8.04 (m, 2H), 7.91-7.90 (m, 1H), 7.77-7.72 (m, 3H), 7.66-7.57 (m, 7H), 7.53-7.44 (m, 4H), 7.45-7.41 (m, 4H)	616.68	615.69
94	8.71-8.68 (m, 2H), 8.52-8.50 (m, 1H), 8.05-8.00 (m, 4H), 7.91-7.90 (m, 2H), 7.74-7.67 (m, 5H), 7.66-7.57 (m, 2H), 7.53-7.48 (m, 4H), 7.45-7.41 (m, 4H)	653.73	653.74
42	8.71-8.68 (m, 2H), 8.52-8.50 (m, 1H), 8.05-8.00 (m, 4H), 7.91-7.90 (m, 2H), 7.74-7.67 (m, 5H), 7.66-7.57 (m, 2H), 7.53-7.48 (m, 4H), 7.45-7.41 (m, 4H)	585.58	585.6
114	8.71-8.68 (m, 2H), 8.52-8.50 (m, 1H), 8.05-8.00 (m, 4H), 7.91-7.90 (m, 2H), 7.74-7.67 (m, 5H), 7.66-7.57 (m, 2H), 7.53-7.48 (m, 4H), 7.45-7.41 (m, 4H)	601.67	601.66

Example 1

[0364] As a substrate and an anode, a Corning 15 Ω/cm^2 (1,200 Å) ITO glass substrate was cut to a size of 50 mm \times 50 mm \times 0.7 mm, and then, sonicated with isopropyl alcohol and pure water each for 5 minutes, and then washed by irradiation of ultraviolet ray for 30 minutes and ozone, and the resultant glass substrate was provided to a vacuum deposition apparatus.

[0365] 2-TNATA was vacuum-deposited on the ITO glass substrate (anode) to form a hole injection layer having a thickness of 600 Å, and 4,4'-bis[N-(1-naphthyl)-N-phenylamino]biphenyl (NPB) was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å.

[0366] 9,10-di-naphthalene-2-yl-anthracene (ADN) (blue fluorescent host) and 4,4'-bis[2-(4-(N,N-diphenylamino)phenyl)vinyl]biphenyl (DPAVBi) (blue fluorescent dopant) were co-deposited on the hole transport layer at a weight ratio of 98:2 to form an emission layer having a thickness of 300 Å.

[0367] Then, Compound 1 was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å. LiF was deposited on the electron transport layer

to form an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited on the electron injection layer to form an LiF/Al cathode having a thickness of 3,000 Å, thereby completing the manufacture of an organic light-emitting device.

Examples 2 to 8 and Comparative Example 1

[0368] Organic light-emitting devices of Examples 2 to 8 and Comparative Example 1 were manufactured in the same manner as in Example 1, except that Compounds shown in Table 2, below, were each used instead of Compound 1 to form an electron transport layer.

Evaluation Example 1

[0369] The driving voltage, current density, luminance, efficiency, emission color, and half lifespan of the organic light-emitting devices manufactured in Examples 1 to 8 and Comparative Example 1 were evaluated using Keithley SMU 236 and PR650 luminance meter. Results thereof are shown in Table 2. The half lifespan was obtained by measuring a time period that has lapsed until the luminance (@100 mA/cm^2) was reduced to 50% of the initial luminance after the driving of the organic light-emitting device.

TABLE 2

	Electron transport material	Driving voltage (V)	Current density (mA/cm^2)	Luminance (cd/m^2)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm^2)
Example 1	Compound 1	4.75	50	2880	5.76	Blue	311.5 hr
Example 2	Compound 20	4.91	50	3104	6.208	Blue	302 hr
Example 3	Compound 23	4.89	50	3088	6.176	Blue	335 hr
Example 4	Compound 26	5.13	50	2963	5.926	Blue	298 hr
Example 5	Compound 64	4.97	50	2870	5.74	Blue	301 hr
Example 6	Compound 94	5.45	50	2894	5.788	Blue	335 hr
Example 7	Compound 42	5.66	50	2667	5.334	Blue	360 hr
Example 8	Compound 114	5.49	50	2695	5.39	Blue	349 hr
Comparative Example 1	Alq ₃	7.12	50	2039	4.078	Blue	162 hr







TABLE 2-continued


Electron transport material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
-----------------------------	---------------------	---------------------------------------	--------------------------------	-------------------	----------------	--



42

Alq₃


20




64

TABLE 2-continued


Electron transport material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
-----------------------------	---------------------	---------------------------------------	--------------------------------	-------------------	----------------	--




23



94



26



114

[0370] Referring to Table 2, it may be seen that the organic light-emitting devices of Examples 1 to 8 had excellent efficiency and lifespan characteristics, compared to Comparative Example 1.

[0371] According to one or more embodiments, the organic light-emitting device including the heterocyclic compound may have high efficiency and a long lifespan.

[0372] Example embodiments have been disclosed herein, and although specific terms are employed, they are used and are to be interpreted in a generic and descriptive sense only and not for purpose of limitation. In some instances, as would be apparent to one of ordinary skill in the art as of the filing of the present application, features, characteristics, and/or elements described in connection with a particular embodiment may be used singly or in combination with features, characteristics, and/or elements described in connection with other embodiments unless otherwise specifically indicated. Accordingly, it will be understood by those of skill in the art that various changes in form and details may be made without departing from the spirit and scope of the present invention as set forth in the following claims.

What is claimed is:

1. A heterocyclic compound represented by Formula 1:




wherein, in Formula 1,

A_1 is a group represented by Formula 2,

A_2 is a group represented by Formula 3, and

$a21$ is an integer of 1 to 11, wherein, when $a21$ is 2 to 11, two or more of A_2 are identical to or different from each other:



<Formula 2>



wherein F_1 in Formula 2 is a dibenzofuran moiety or a dibenzothiophene moiety,

R_1 to R_5 and R_{12} in Formula 2 are each independently selected from a binding site to A_2 , hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a

substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(Q_1)(Q_2)(Q_3)$, and $-\text{S}(=\text{O})_2(Q_1)$,

$b6$ in Formula 2 is an integer of 0 to 6, wherein, when $b6$ is 2 to 6, two or more of R_6 are identical to or different from each other,

X_1 in Formula 2 is O or S,

at least one of R_1 to R_5 and R_{12} is a binding site to A_2 , a number of R_1 to R_5 and R_{12} being the binding site to A_2 being the same as $a21$,

L in Formula 3 is selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

n in Formula 3 is an integer of 0 to 7, wherein, when n is 2 to 7, two or more of L are identical to or different from each other,

Ar in Formula 3 is selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(Q_4)(Q_5)(Q_6)$, $-\text{C}(=\text{O})(Q_4)$, $-\text{S}(=\text{O})_2(Q_4)$, and $-\text{P}(=\text{O})(Q_4)(Q_5)$,

m in Formula 3 is an integer of 1 to 7, wherein, when m is 2 to 7, two or more of Ar are identical to or different from each other, and

at least one substituent of the substituted C_3 - C_{10} cycloalkylene group, the substituted C_1 - C_{10} heterocycloalkylene group, the substituted C_3 - C_{10} cycloalkenylene group, the substituted C_1 - C_{10} heterocycloalkenylene group, the substituted C_6 - C_{60} arylene group, the substituted C_1 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_1 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, and a substituted or unsubstituted C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

—Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂).

wherein Q₁ to Q₆, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a fluorenylene group, a naphthylene group, a biphenyl group, a terphenyl group, and a naphthyl group.

lent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group.

2. The heterocyclic compound as claimed in claim 1, wherein:

X₁ is O, and

F₁ is a dibenzofuran moiety.

3. The heterocyclic compound as claimed in claim 1, wherein a21 is an integer of 1 to 6.

4. The heterocyclic compound as claimed in claim 1, wherein L in Formula 3 is selected from:


a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinoxalinylenylene group, a cinolinylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a

quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylyl group, a quinazolinylyl group, a cinnolinylyl group, a phenanthridinylyl group, an acridinylyl group, a phenanthrolinylyl group, a phenazinylyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃),


wherein Q₃₁ to Q₃₃ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group.

5. The heterocyclic compound as claimed in claim 1, wherein L in Formula 3 is a group represented by one of the following Formulae 5-1 to 5-34:




Formula 5-1


-continued




Formula 5-2




Formula 5-3




Formula 5-4




Formula 5-5




Formula 5-6



Formula 5-7












Formula 5-8




Formula 5-9


-continued




Formula 5-10




Formula 5-11




Formula 5-12




Formula 5-13




Formula 5-14




Formula 5-15




Formula 5-16



Formula 5-17



Formula 5-18



-continued

Formula 5-19

Formula 5-20

Formula 5-21

Formula 5-22

Formula 5-23


Formula 5-24

Formula 5-25


Formula 5-26

Formula 5-27


-continued




Formula 5-28




Formula 5-29




Formula 5-30



Formula 5-31




Formula 5-32



Formula 5-33

-continued

Formula 5-34



wherein, in Formulae 5-1 to 5-34,

 Y_1 is O, S, C(Z_3)(Z_4), N(Z_5), or Si(Z_6)(Z_7),

Z_1 to Z_7 are each independently selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazolinyl group, a benzoquinazolinyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafuorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indenocarbazolyl group, an indolocarbazolyl group, and —Si(Q_{31})(Q_{32})(Q_{33}), in which Q_{31} to Q_{33} are each independently selected from

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group,

d2 is an integer of 0 to 2,

d3 is an integer of 0 to 3,

d4 is an integer of 0 to 4,

d5 is an integer of 0 to 5,

d6 is an integer of 0 to 6,

d8 is an integer of 0 to 8, and

* and *' each indicate a binding site to a neighboring atom.

6. The heterocyclic compound as claimed in claim 5, wherein:

L is a group represented by one of Formula 5-1, 5-2, or 5-13, and

Z₁ in Formula 5-1, 5-2, or 5-13 is selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group.

7. The heterocyclic compound as claimed in claim 1, wherein n in Formula 3 is 0, 1, or 2.

8. The heterocyclic compound as claimed in claim 1, wherein Ar in Formula 3 is selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl

group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;


a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl

group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiazazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and


Si(Q₄)(Q₅)(Q₆), —C(=O)(Q₄), —S(=O)₂(Q₄), and —P(=O)(Q₄)(Q₅),

wherein Q₄ to Q₆ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group.


9. The heterocyclic compound as claimed in claim 1, wherein Ar in Formula 3 is a group represented by one of the following Formulae 6-1 to 6-37:



Formula 6-1




Formula 6-2




Formula 6-3


-continued




Formula 6-4




Formula 6-5




Formula 6-6




Formula 6-7



Formula 6-8










Formula 6-9




Formula 6-10

-continued




-continued

Formula 6-11




Formula 6-18

Formula 6-12




Formula 6-19

Formula 6-13




Formula 6-20

Formula 6-14




Formula 6-21

Formula 6-15




Formula 6-22

Formula 6-16




Formula 6-17




Formula 6-23


-continued




Formula 6-24




Formula 6-25




Formula 6-26




Formula 6-27




Formula 6-28



Formula 6-29




Formula 6-30




Formula 6-31


-continued




Formula 6-32




Formula 6-33




Formula 6-34



Formula 6-35



Formula 6-36



Formula 6-37

wherein, in Formulae 6-1 to 6-37,

Y₃₁, Y₃₃ and Y₃₇ are each independently O, S, C(Z₃₅) (Z₃₆), N(Z₃₇), or Si(Z₃₈)(Z₃₉),

Y₃₂, Y₃₄, Y₃₆, and Y₃₈ are each independently C(Z₃₅) or N,

Z₃₁ to Z₃₉ are each independently selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a qui-

nazolanyl group, a benzoquinazolanyl group, a cinnolanyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothio-phenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphtho-furanyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indenocarbazolyl group, an indolocarbazolyl group, a benzoacridinyl group, a benzophenanthrenyl group, a dibenzophenanthrenyl group, a benzophenanthridinyl group, a dibenzophenanthridinyl group, an indenophenanthrenyl group, an indolophenanthrenyl group, a benzofuranophenanthrenyl group, a benzothiophenanthrenyl group, a benzosilolephenanthrenyl group, an indenophenanthridinyl group, an indolophenanthridinyl group, a benzofuranophenanthridinyl group, a benzothiophenanthridinyl group, a benzosilolephenanthridinyl group, a dinaphthopyrrolyl group, a benzoanthracenyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, in which Q_{31} to Q_{33} are each independently selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{60}$ alkyl group, a $\text{C}_2\text{-C}_{60}$ alkenyl group, a $\text{C}_2\text{-C}_{60}$ alkynyl group, a $\text{C}_1\text{-C}_{60}$ alkoxy group, a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group,

e2 is an integer of 0 to 2,

e3 is an integer of 0 to 3,

e4 is an integer of 0 to 4,


e5 is an integer of 0 to 5,

e6 is an integer of 0 to 6, and


* indicates a binding site to a neighboring atom.

10. The heterocyclic compound as claimed in claim 7, wherein Ar is a group represented by one of Formulae 6-1 to 6-6, 6-13 to 6-29, 6-31, 6-32, and 6-37.


11. The heterocyclic compound as claimed in claim 1, wherein Ar in Formula 3 is a group represented by one of the following Formulae 7-1 to 7-25:




Formula 7-1




Formula 7-2




Formula 7-3




Formula 7-4




Formula 7-5




Formula 7-6




Formula 7-7



Formula 7-8




Formula 7-9




Formula 7-10


-continued




Formula 7-11




Formula 7-12




Formula 7-13




Formula 7-14



Formula 7-15




Formula 7-16




Formula 7-17


-continued




Formula 7-18




Formula 7-19




Formula 7-20



Formula 7-21




Formula 7-22




Formula 7-23

-continued




Formula 7-24




Formula 7-25

wherein * in Formulae 7-1 to 7-25 indicates a binding site to a neighboring atom.

12. The heterocyclic compound as claimed in claim 1, wherein A_1 is a group represented by one of the following Formulae 2A to 2F:




Formula 2A




Formula 2B

-continued


Formula 2C



Formula 2D



Formula 2E



Formula 2F

wherein, in Formulae 2A to 2F,

X_1 and X_2 are each independently O or S,

R_1 to R_{11} are each independently selected from a binding site to A_2 , hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a

substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), and —S(=O)₂(Q₁), in which Q₁ to Q₃ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, and a naphthyl group.


13. The heterocyclic compound as claimed in claim 12, wherein R₅ in Formulae 2A to 2F is a binding site to A₂.

14. The heterocyclic compound as claimed in claim 12, wherein X₁ and X₂ in Formula 2A to 2F are 0.


15. The heterocyclic compound as claimed in claim 1, wherein the heterocyclic compound represented by Formula 1 is one of the following Compounds 1 to 72 and 74 to 138:

-continued


3




4




5




6



1




2




-continued


7




8



9




10




-continued


11




12



13




14




-continued

-continued


15




19




16




20




17



21




18




-continued

-continued


22




25




26




23



24




27




-continued

-continued

28




31




32



29



30



33




-continued

-continued


34

37







35

38









36

39









-continued

-continued










-continued

-continued



-continued


-continued




-continued

-continued


59




62




63




60



61



64





-continued

-continued



65

68




69


66



67



70





-continued

-continued



75

71





76

72




77

74





78



-continued

-continued





-continued

-continued



87

91





88

92





89

93











90

94




-continued

-continued




-continued




103


-continued




107




104




108




105



109




106




110

-continued


111




112



113




114




-continued


115




116



117



118





-continued

-continued



119

123





120

124




121

125









122




-continued

-continued




-continued



-continued




133




134



136




137



-continued

138



16. An organic light-emitting device, comprising:
 a first electrode;
 a second electrode facing the first electrode; and
 an organic layer between the first electrode and the second electrode and comprises an emission layer,
 wherein the organic layer includes the heterocyclic compound as claimed in claim 1.

17. The organic light-emitting device as claimed in claim 16, wherein:

the first electrode is an anode,
 the second electrode is a cathode, and
 the organic layer includes:

a hole transport region between the first electrode and the emission layer, the hole transport region including at least one selected from a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer, and

an electron transport region between the emission layer and the second electrode, the electron transport region including at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

18. The organic light-emitting device as claimed in claim 17, wherein at least one of the electron transport region and the emission layer includes the heterocyclic compound.

19. The organic light-emitting device as claimed in claim 17, wherein:

the electron transport region includes the electron transport layer or the electron injection layer, and
 the electron transport layer or the electron injection layer includes heterocyclic compound.

20. The organic light-emitting device as claimed in claim 18, wherein the emission layer further includes a pyrene-based compound, an arylamine compound, or a styryl-based compound.

* * * * *

专利名称(译)	杂环化合物和包括其的有机发光装置		
公开(公告)号	US20170338419A1	公开(公告)日	2017-11-23
申请号	US15/496120	申请日	2017-04-25
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	SIM MUNKI KIM YOUNGKOOK PARK JUNHA LEE HYOYOUNG JEONG EUNJAE HWANG SEOKHWAN		
发明人	SIM, MUNKI KIM, YOUNGKOOK PARK, JUNHA LEE, HYOYOUNG JEONG, EUNJAE HWANG, SEOKHWAN		
IPC分类号	H01L51/00 H01L27/32 C07D491/153 H01L51/52 C07F9/6561 H01L51/50		
CPC分类号	H01L51/0071 H01L51/5218 H01L27/3211 C07D491/153 H01L51/0058 H01L51/0052 H01L51/5072 H01L51/0072 C07F9/6561 H01L51/5056 H01L51/5088 H01L51/5096 H01L51/5092 H01L51/0067		
优先权	1020160061439 2016-05-19 KR		
其他公开文献	US10305046		
外部链接	Espacenet USPTO		

摘要(译)

杂环化合物和包含其的有机发光装置，该化合物由式1表示：A 1 - (A 2) a₂₁ ， & #x3c;公式1> 其中，在公式1中，A 1 是表示的组通过式 2，A 2 是由式3表示的基团，并且a₂₁是1至11的整数，其中，当a₂₁是2 至11时，A 2 彼此相同或不同：

40

220
190
150
110
210