

US009288869B2

(12) **United States Patent**  
**Han et al.**

(10) **Patent No.:** **US 9,288,869 B2**  
(45) **Date of Patent:** **\*Mar. 15, 2016**

(54) **STYRYL-BASED COMPOUND,  
COMPOSITION CONTAINING  
STYRYL-BASED COMPOUND, AND  
ORGANIC LIGHT EMITTING DIODE  
INCLUDING STYRYL-BASED COMPOUND**

2211/1014; C09K 2211/1029; C09K  
2211/1088; C09K 2211/1092; C09K 11/06;  
H01L 51/0058; H01L 51/0059; H01L  
51/0061; H01L 51/0074; H01L 51/54; H05B  
33/14

See application file for complete search history.

(75) Inventors: **Sang-Hyun Han**, Yongin (KR);  
**Seok-Hwan Hwang**, Yongin (KR);  
**Young-Kook Kim**, Yongin (KR);  
**Hye-Jin Jung**, Yongin (KR); **Jin-O Lim**,  
Yongin (KR); **Soo-Yon Kim**, Yongin  
(KR); **Dae-Yup Shin**, Yongin (KR);  
**Jong-Hyuk Lee**, Yongin (KR)

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,635,308 A 6/1997 Inoue et al.  
5,972,247 A 10/1999 Shi et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CN 1854119 A 11/2006  
JP 8-12600 A 1/1996

(Continued)

OTHER PUBLICATIONS

Chinese OA issued by the Chinese Patent Office on Jun. 26, 2015 in  
the examination of the Chinese Patent Application No.  
201210443012.2, which corresponds to KR 10-2011-0117161.  
Request for Entry of the Accompanying Office Action is attached  
herewith.

*Primary Examiner* — Dawn L. Garrett

(74) *Attorney, Agent, or Firm* — Robert E. Bushnell, Esq.

(73) Assignee: **Samsung Display Co., Ltd.**,  
Samsung-ro, Giheung-Gu, Yongin-si,  
Gyeonggi-Do (KR)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 714 days.

This patent is subject to a terminal dis-  
claimer.

(21) Appl. No.: **13/609,633**

(22) Filed: **Sep. 11, 2012**

(65) **Prior Publication Data**

US 2013/0119355 A1 May 16, 2013

(30) **Foreign Application Priority Data**

Nov. 10, 2011 (KR) ..... 10-2011-0117161

(51) **Int. Cl.**  
**H01L 51/54** (2006.01)  
**C07C 211/58** (2006.01)


(Continued)

(52) **U.S. Cl.**  
CPC ..... **H05B 33/14** (2013.01); **C07B 59/00**  
(2013.01); **C07C 211/54** (2013.01);  
(Continued)

(58) **Field of Classification Search**  
CPC ..... C07C 2103/18; C07C 2103/24; C07C  
2103/26; C07C 2103/42; C07C 2103/48;  
C07C 2103/50; C07C 2103/94; C07C 211/54;  
C07C 211/56; C07C 211/58; C07C 211/60;  
C07C 211/61; C07C 255/58; C07D 209/86;  
C07D 209/88; C07D 213/38; C07D 213/74;  
C07D 307/91; C07D 333/76; C07D 401/10;  
C07D 409/12; C09K 2211/1011; C09K

(57) **ABSTRACT**


A styryl-based compound represented by Formula 1, a com-  
position containing the styryl-based compound, and an  
organic light-emitting diode (OLED) including the styryl-  
based compound:




[Formula 1]

The styryl-based compound may exhibit high heat resistance  
and thus an OLED including the same may have low driving  
voltage, high brightness, high efficiency, and long lifetime.

**17 Claims, 1 Drawing Sheet**



- (51) **Int. Cl.**  
*C07C 211/61* (2006.01)  
*C07D 213/38* (2006.01)  
*C07D 213/74* (2006.01)  
*C09K 11/06* (2006.01)  
*C07D 209/88* (2006.01)  
*H05B 33/14* (2006.01)  
*C07C 255/58* (2006.01)  
*C07B 59/00* (2006.01)  
*C07C 211/54* (2006.01)  
*C07C 211/56* (2006.01)  
*C07C 211/60* (2006.01)  
*C07D 307/91* (2006.01)  
*C07D 333/76* (2006.01)  
*C07D 401/10* (2006.01)  
*C07D 409/12* (2006.01)  
*C07D 209/86* (2006.01)  
*H01L 51/00* (2006.01)
- (52) **U.S. Cl.**  
 CPC ..... *C07C 211/56* (2013.01); *C07C 211/58* (2013.01); *C07C 211/60* (2013.01); *C07C 211/61* (2013.01); *C07C 255/58* (2013.01); *C07D 209/86* (2013.01); *C07D 209/88* (2013.01); *C07D 213/38* (2013.01); *C07D 213/74* (2013.01); *C07D 307/91* (2013.01); *C07D 333/76* (2013.01); *C07D 401/10* (2013.01); *C07D 409/12* (2013.01); *C09K 11/06* (2013.01); *H01L 51/0059* (2013.01); *C07B 2200/05* (2013.01); *C07C 2103/18* (2013.01); *C07C 2103/24* (2013.01); *C07C 2103/26* (2013.01); *C07C 2103/42* (2013.01); *C07C 2103/48* (2013.01); *C07C 2103/50* (2013.01); *C07C 2103/94* (2013.01); *C09K 2211/1011* (2013.01); *C09K 2211/1014* (2013.01); *C09K 2211/1029* (2013.01); *C09K 2211/1088* (2013.01); *C09K 2211/1092* (2013.01); *H01L 51/0058* (2013.01); *H01L 51/0061* (2013.01); *H01L 51/0074* (2013.01)
- (56) **References Cited**  
 U.S. PATENT DOCUMENTS
- |              |      |         |             |         |
|--------------|------|---------|-------------|---------|
| 6,465,115    | B2   | 10/2002 | Shi et al.  |         |
| 6,596,415    | B2   | 7/2003  | Shi et al.  |         |
| 8,679,648    | B2 * | 3/2014  | Han et al.  | 428/690 |
| 8,828,559    | B2 * | 9/2014  | Jung et al. | 428/690 |
| 2006/0246317 | A1   | 11/2006 | Lyu et al.  |         |
| 2010/0081846 | A1   | 4/2010  | Kim et al.  |         |
| 2013/0112948 | A1 * | 5/2013  | Jung et al. | 257/40  |
| 2014/0014916 | A1 * | 1/2014  | Han et al.  | 257/40  |
| 2014/0027721 | A1 * | 1/2014  | Kim et al.  | 257/40  |
| 2014/0239260 | A1 * | 8/2014  | Lee et al.  | 257/40  |
| 2014/0332772 | A1 * | 11/2014 | Han et al.  | 257/40  |
- FOREIGN PATENT DOCUMENTS
- |    |                 |    |         |
|----|-----------------|----|---------|
| JP | 2000-3782       | A  | 1/2000  |
| JP | 2006-273737     | A  | 10/2006 |
| KR | 10-2006-0111048 | A  | 10/2006 |
| KR | 10-0835601      | B1 | 5/2008  |
| KR | 10-2010-0069216 | A  | 6/2010  |
| KR | 10-2010-0118258 | A  | 11/2010 |
| WO | WO 2010/126233  | *  | 11/2010 |
- \* cited by examiner



1

**STYRYL-BASED COMPOUND,  
COMPOSITION CONTAINING  
STYRYL-BASED COMPOUND, AND  
ORGANIC LIGHT EMITTING DIODE  
INCLUDING STYRYL-BASED COMPOUND**

CROSS-REFERENCE TO RELATED PATENT  
APPLICATION

This application claims the benefit of Korean Patent Application No. 10-2011-0117161, filed on Nov. 10, 2011, in the Korean Intellectual Property Office, the disclosure of which is incorporated herein in its entirety by reference.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a styryl-based compound and an organic light-emitting diode (OLED) including the same.

2. Description of the Related Art

Organic light-emitting diodes (OLEDs) are self-emitting devices, have advantages such as a wide viewing angle, excellent contrast, quick response, high brightness, and excellent driving voltage, and can provide multicolored images.

A general OLED has a structure including a substrate, and an anode, a hole transport layer (HTL), an emission layer (EML), an electron transport layer (ETL), and a cathode which are sequentially stacked on the substrate. In this regard, the HTL, the EML, and the ETL are organic layers formed of organic compounds.


An operating principle of an OLED having the above-described structure is as follows.

When a voltage is applied between the anode and the cathode, holes injected from the anode move to the EML via the HTL, and electrons injected from the cathode move to the EML via the ETL. The holes and electrons recombine in the EML to generate excitons. When the excitons drop from an excited state to a ground state, light is emitted.

SUMMARY OF THE INVENTION

The present invention provides a styryl-based compound having a novel structure, a composition containing the styryl-based compound, and an organic light-emitting diode (OLED) including the styryl-based compound or the composition.

According to an aspect of the present invention, there is provided a styryl-based compound represented by Formula 1 below:



Formula 1

wherein  $\text{Ar}_3$  is a substituted or unsubstituted  $\text{C}_8\text{-C}_{20}$  aryl group having two or more rings fused with each other, or a substituted or unsubstituted  $\text{C}_2\text{-C}_{20}$  heteroaryl group having two or more rings fused with each other;


$\text{Ar}_1$  and  $\text{Ar}_2$  are each independently a substituted or unsubstituted  $\text{C}_5\text{-C}_{60}$  aryl group or a substituted or unsubstituted  $\text{C}_2\text{-C}_{60}$  heteroaryl group;

2

$\text{L}_1$  and  $\text{L}_2$  are each independently a substituted or unsubstituted  $\text{C}_5\text{-C}_{60}$  arylene group or a substituted or unsubstituted  $\text{C}_2\text{-C}_{60}$  heteroarylene group; and  $a$  and  $b$  are each independently an integer of 0 to 5.


According to another aspect of the present invention, there is provided a composition containing a styryl-based compound, including a styryl-based compound represented by Formula 1 below; and at least one of a styryl-based compound represented by Formula 1-1H-1 below, a styryl-based compound represented by Formula 1-1H-2 below, and a styryl-based compound represented by Formula 1-2H below:

Formula 1




15

Formula 1-1H-1




20

Formula 1-1H-2



25

Formula 1-2H



30

35

wherein substituents of Formulae 1-1H-1, 1-1H-2, and 1-2H are defined in a detailed description of Formula 1.

According to another aspect of the present invention, there is provided an organic light-emitting diode (OLED) including a first electrode; a second electrode facing the first electrode; and an organic layer interposed between the first electrode and the second electrode, wherein the organic layer includes at least one of the styryl-based compound, or at least one of the composition containing a styryl-based compound.

BRIEF DESCRIPTION OF THE DRAWINGS

The above and other features and advantages of the present invention will become more apparent by describing in detail exemplary embodiments thereof with reference to the attached drawings in which:


FIG. 1 is a schematic diagram illustrating an organic light-emitting diode (OLED) according to an embodiment of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

Hereinafter, the present embodiments will be described more fully with reference to the accompanying drawing, in which exemplary embodiments are shown. As used herein, the term "and/or" includes any and all combinations of one or more of the associated listed items. Expressions such as "at least one of," when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

## 3

A styryl-based compound according to an embodiment of the present invention is represented by Formula 1 below:




Formula 1


wherein Ar<sub>3</sub> may be a substituted or unsubstituted C<sub>8</sub>-C<sub>20</sub> aryl group having two or more rings fused with each other, or a substituted or unsubstituted C<sub>2</sub>-C<sub>20</sub> heteroaryl group having two or more rings fused with each other.

For example, in Formula 1, Ar<sub>3</sub> may be a substituted or unsubstituted naphthyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted dibenzofuranyl group, or a substituted or unsubstituted dibenzothiophenyl group. In this case, at least one substituent of the substituted naphthyl group, phenanthrenyl group, anthryl group, pyrenyl group, chrysenyl group, fluorenyl group, carbazolyl group, dibenzofuranyl group, and dibenzothiophenyl group may be selected from deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a C<sub>1</sub>-C<sub>10</sub> alkyl group; a C<sub>1</sub>-C<sub>10</sub> alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolyl group; an imidazopyridinyl group; an imidazopyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; an indolyl group; and a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>10</sub> alkyl group, and a C<sub>1</sub>-C<sub>10</sub> alkoxy group.

According to Ar<sub>3</sub> in Formula 1, the styryl-based compound represented by Formula 1 may be represented by any one of Formulae 1A through 1I below:



Formula 1A




65

## 4


-continued

Formula 1B




10

Formula 1C




15

Formula 1D




20

Formula 1E




25

Formula 1F



30

Formula 1G



35


Formula 1H

60

5

-continued

Formula II



In Formulae 1A through 1I,  $Z_1$  through  $Z_3$  may be each independently hydrogen; deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a  $C_1$ - $C_{10}$ alkyl group; a  $C_1$ - $C_{10}$ alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolanyl group; an imidazopyridinyl group; an imidazopyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; an indolyl group; an and phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazolanyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted by at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{10}$ alkyl group, and a  $C_1$ - $C_{10}$ alkoxy group. In this case, two or more neighboring substituents from among  $Z_1$  through  $Z_3$  may be optionally fused with each other or may be optionally connected by a single bond.

For example,  $Z_1$  through  $Z_3$  may be each independently hydrogen, deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a hexyl group, a heptyl group, an octyl group, a phenyl group, a naphthyl group, a fluorenyl group, a deuterated phenyl group, or a dimethylfluorenyl group. In this case, when  $Z_1$  and  $Z_2$  of Formula 1G are each a phenyl group,  $Z_1$  and  $Z_2$  may be connected by a single bond.

In Formulae 1A through 1I,  $T_1$  may be O or S.

In Formulae 1A through 1I,  $p$  may be an integer of 1 to 7, and  $q$  may be an integer of 1 to 4. When  $p$  is 2 or more, at least two  $Z_1$  may be identical to or different from each other and at least two  $Z_3$  may be identical to or different from each other. When  $q$  is 2 or more, at least two  $Z_2$  may be identical to or different from each other.

In Formula 1,  $Ar_1$  and  $Ar_2$  may be each independently a substituted or unsubstituted  $C_5$ - $C_{60}$ aryl group or a substituted or unsubstituted  $C_2$ - $C_{60}$ heteroaryl group. In this case, at least one substituent of the substituted  $C_5$ - $C_{60}$  aryl group and the substituted  $C_2$ - $C_{60}$  heteroaryl group may be selected from deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid

6

group or a salt thereof; a phosphoric acid or a salt thereof; a  $C_1$ - $C_{10}$ alkyl group; a  $C_1$ - $C_{10}$ alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolanyl group; an imidazopyridinyl group; an imidazopyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; an indolyl group; and a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazolanyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted by at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{10}$ alkyl group, and a  $C_1$ - $C_{10}$ alkoxy group.


For example, in Formula 1,  $Ar_1$  and  $Ar_2$  may be each independently, but are not limited to, a substituted or unsubstituted phenyl group, a substituted or unsubstituted pentalenyl group, a substituted or unsubstituted indenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted azulenyl group, a substituted or unsubstituted heptalenyl group, a substituted or unsubstituted indacenyl group, a substituted or unsubstituted acenaphthyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted spiro-fluorenyl group, a substituted or unsubstituted phenalenyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluoranthrenyl group, a substituted or unsubstituted triphenylenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted naphthacenyl group, a substituted or unsubstituted picenyl group, a substituted or unsubstituted perylenyl group, a substituted or unsubstituted pentaphenyl group, a substituted or unsubstituted hexacenyl group, a substituted or unsubstituted pyrrolyl group, a substituted or unsubstituted imidazolyl group, a substituted or unsubstituted pyrazolyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted pyridazinyl, a substituted or unsubstituted isoindolyl group, a substituted or unsubstituted indolyl group, a substituted or unsubstituted indazolyl group, a substituted or unsubstituted purinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted benzoquinolinyl group, a substituted or unsubstituted phthalazinyl group, a substituted or unsubstituted naphthyridinyl group, a substituted or unsubstituted quinoxalinyl group, a substituted or unsubstituted quinazolinyl group, a substituted or unsubstituted cinnolinyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted phenanthridinyl group, a substituted or unsubstituted acridinyl group, a substituted or unsubstituted phenanthrolinyl group, a substituted or unsubstituted phenazolinyl group, a substituted or unsubstituted benzoxazolyl group, a substituted or unsubstituted benzoimidazolyl group, a substituted or unsubstituted furanyl group, a substituted or unsubstituted benzofuranyl group, a substituted or unsubstituted thiophenyl group, a substituted or unsubstituted benzothiophenyl group, a substituted or unsubstituted thiazolyl group, a substituted or unsubstituted isothiazolyl group, a substituted or unsubstituted benzothiazolyl group, a substi-

7


tuted or unsubstituted isoxazolyl group, a substituted or unsubstituted oxazolyl group, a substituted or unsubstituted triazolyl group, a substituted or unsubstituted tetrazolyl group, a substituted or unsubstituted oxadiazolyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted benzoxazolyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted dibenzothiophenyl group, or a substituted or unsubstituted benzocarbazolyl group.

For example, in Formula 1 above, Ar<sub>1</sub> and Ar<sub>2</sub> may be each independently, but are not limited to, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted dibenzothiophenyl group, or a substituted or unsubstituted benzocarbazolyl group.


Ar<sub>1</sub> and Ar<sub>2</sub> may be each independently represented by any one of Formulae 2A through 2J below:




Formula 2A




Formula 2B




Formula 2C



Formula 2D



Formula 2E




Formula 2F


8

-continued


Formula 2G




Formula 2H



Formula 2I



Formula 2J



30









In Formulae 2A through 2J above, Z<sub>11</sub> through Z<sub>15</sub> may be each independently selected from hydrogen; deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a C<sub>1</sub>-C<sub>10</sub>alkyl group; a C<sub>1</sub>-C<sub>10</sub>alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolyl group; an imidazopyridinyl group; a pyridinyl group; a pyrimidinyl group; a pyrazinyl group; an indolyl group; and a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>10</sub>alkyl group, and a C<sub>1</sub>-C<sub>10</sub>alkoxy group. In this case, two or more neighboring substituents from among Z<sub>11</sub> through Z<sub>15</sub> may be optionally fused with each other or may be optionally connected by a single bond.

In Formulae 2A through 2J above, T<sub>11</sub> may be O or S.

In Formulae 2A and 2J above, r may be an integer of 1 to 7 and s may be an integer of 1 to 5. When r is 2 or more, at least two Z<sub>11</sub> may be identical to or different from each other and at least two Z<sub>13</sub> may be identical to or different from each other. When s is 2 or more, at least two Z<sub>12</sub> may be identical to or different from each other.

9


For example, Ar<sub>1</sub> and Ar<sub>2</sub> may be each independently represented by any one of Formulae 3A through 3O below:




10

-continued


Formula 3A 5




Formula 3B 10




Formula 3C 15




Formula 3D 20




Formula 3E 25




Formula 3F 30




Formula 3G 35



Formula 3H 40



Formula 3I 45



Formula 3J

Formula 3K

Formula 3L

Formula 3M

Formula 3N

Formula 3O

wherein a detailed description of Z<sub>11</sub> through Z<sub>15</sub> is already provided above (however, Formulae 3A through 3O above, Z<sub>11</sub> through Z<sub>15</sub> are not hydrogen). For example, in Formulae 3A through 3O above, Z<sub>11</sub> through Z<sub>15</sub> may be each independently, but are not limited to, deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a hexyl group, a heptyl group, an octyl group, a phenyl group, a naphthyl group, or a fluorenyl group. In Formulae 3A through 3O above, T<sub>11</sub> may be O or S.

In Formula 1 above, L<sub>1</sub> and L<sub>2</sub> may be each independently a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> arylene group or a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heteroarylene group. At least


one substituent of the substituted  $C_5$ - $C_{60}$  arylene group and the substituted  $C_2$ - $C_{60}$  heteroarylene group may be selected from deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a  $C_1$ - $C_{10}$  alkyl group; a  $C_1$ - $C_{10}$  alkoxy group; a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazoliny group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group; and a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazoliny group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{10}$  alkyl group, and a  $C_1$ - $C_{10}$  alkoxy group.

For example,  $L_1$  and  $L_2$  may be each independently a substituted or unsubstituted phenylene group, a substituted or unsubstituted pentalenylene group, a substituted or unsubstituted indenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted azulenylenylene group, a substituted or unsubstituted heptalenylene group, a substituted or unsubstituted indacenylene group, a substituted or unsubstituted acenaphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted phenalenylene group, a substituted or unsubstituted phenanthrenylene group, a substituted or unsubstituted fluoranthrylene group, a substituted or unsubstituted fluoranthrylene group, a substituted or unsubstituted triphenylenylene group, a substituted or unsubstituted pyrenylene group, a substituted or unsubstituted chrysenylene group, a substituted or unsubstituted naphthacenylenylene group, a substituted or unsubstituted picenylene group, a substituted or unsubstituted perylenylene group, a substituted or unsubstituted pentaphenylenylene group, a substituted or unsubstituted hexacenylenylene group, a substituted or unsubstituted pyrrolylene group, a substituted or unsubstituted pyrazolylenylene group, a substituted or unsubstituted imidazolylene group, a substituted or unsubstituted imidazolinylenylene group, a substituted or unsubstituted imidazopyridinylenylene group, a substituted or unsubstituted imidazopyrimidinylene group, a substituted or unsubstituted pyridinylenylene group, a substituted or unsubstituted pyrazinylenylene group, a substituted or unsubstituted pyrimidinylene group, a substituted or unsubstituted indolylenylene group, a substituted or unsubstituted purinylenylene group, a substituted or unsubstituted quinolylenylene group, a substituted or unsubstituted phthalazinylenylene group, a substituted or unsubstituted indolizinylenylene group, a substituted or unsubstituted naphthyridinylenylene group, a substituted or unsubstituted quinazolinylenylene group, a substituted or unsubstituted cinnolinylenylene group, a substituted or unsubstituted indazolylene group, a substituted or unsubstituted carbazolylene group, a substituted or unsubstituted phenazinylenylene group, a substituted or unsubstituted phenanthridinylenylene group, a substituted or unsubstituted pyranylene group, a substituted or unsubstituted chromenylenylene group, a substituted or unsubstituted furanyl group, a substituted or unsubstituted benzofuranylenylene group, a substituted or unsubstituted thiophenylenylene group, a substituted or unsubstituted benzothiophenylene group,


a substituted or unsubstituted isothiazolylenylene group, a substituted or unsubstituted benzoimidazolylene group, a substituted or unsubstituted isoxazolylene group, a substituted or unsubstituted dibenzothiophenylene group, a substituted or unsubstituted dibenzofuranylenylene group, a substituted or unsubstituted triazinylene group, or a substituted or unsubstituted oxadiazolylenylene group.

For example, in Formula 1 above,  $L_1$  and  $L_2$  may be each independently, but are not limited to, a substituted or unsubstituted phenylenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted phenanthrenylene group, a substituted or unsubstituted pyridinylenylene group, or a substituted or unsubstituted pyrazinylenylene group.


In Formula 1 above,  $L_1$  and  $L_2$  may be each independently represented by any one of Formulae 5A through 5K below:




Formula 5A




Formula 5B




Formula 5C




Formula 5D




Formula 5E



Formula 5F






Formula 5G



Formula 5H

13

-continued



In Formulae 5A through 5K above,  $Z_{31}$  through  $Z_{34}$  may be each independently selected from hydrogen; deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a  $C_1$ - $C_{10}$  alkyl group; a  $C_1$ - $C_{10}$  alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolyl group; an imidazopyridinyl group; an imidazopyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; an indolyl group; and a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{10}$  alkyl group, and a  $C_1$ - $C_{10}$  alkoxy group. In this case, two or more neighboring substituents from among  $Z_{31}$  through  $Z_{34}$  may be optionally fused with each other or may be optionally connected by a single bond.

For example,  $Z_{31}$  through  $Z_{34}$  may be each independently, but are not limited to, hydrogen, deuterium, a halogen atom, a hydroxyl group, a cyano group, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a hexyl group, or a heptyl group.

In Formulae 5A through 5K above,  $t$  may be an integer of 1 to 6 and  $u$  may be an integer of 1 to 3. When  $t$  is 2 or more, at least two  $Z_{31}$  may be identical to or different from each other. When  $u$  is 2 or more, at least two  $Z_{32}$  may be identical to or different from each other.

In Formula 1 above,  $a$  and  $b$  may be each independently an integer of 0 to 5.


For example, in Formula 1 above,  $a$  may be, but is not limited to, 0 or 1, and  $b$  may be, but is limited to, 1 or 2.

According to an embodiment of the present invention, the styryl-based compound of Formula 1 above may be represented by any one of Formula 1A-1 through 1I-3 below:

14

Formula 5I


5



Formula 1A-1

Formula 5J


10



Formula 1A-2


Formula 5K

15




Formula 1B-1

20



Formula 1B-2


30



Formula 1C-1

35


40



Formula 1C-2

45

50



Formula 1D-1


55

60


65

15


-continued




Formula 1E-1




Formula 1F-1




Formula 1G-1



Formula 1G-2




Formula 1H-1



Formula 1H-2

16


-continued



5

10


Formula 1I-1



15

20

Formula 1I-2



25

Formula 1I-3

In Formula 1A-1 through 1I-3 above, detailed descriptions for  $Z_1$ ,  $Z_2$ ,  $T_1$ ,  $Ar_1$ ,  $Ar_2$ ,  $L_1$ ,  $L_2$ ,  $a$ , and  $b$  are already provided above.

For example, in Formula 1A-1 through 1I-3 above,  $Z_1$  and  $Z_2$  may be each independently selected from hydrogen; deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a  $C_1$ - $C_{10}$  alkyl group; a  $C_1$ - $C_{10}$  alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolynyl group; an imidazopyridinyl group; an imidazopyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; an indolyl group; and a phenyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazolynyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{10}$  alkyl group, and a  $C_1$ - $C_{10}$  alkoxy group;  $T_1$  may be O or S;  $Ar_1$  and  $Ar_2$  may be each independently a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted dibenzothiophenyl group, or a substituted or unsubstituted benzocarbazolyl group;  $L_1$  and  $L_2$  may be each independently a substituted or unsubstituted phenylenyl group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted phenanthre-

17









nylene group, a substituted or unsubstituted pyridinylene group, or a substituted or unsubstituted pyrazinylene group; a is 0 or 1; and b is 1 or 2, but the present embodiment is not limited thereto.

For example, in Formulae 1A-1 through 1I-3 above,  $Z_1$  and  $Z_2$  may be each independently hydrogen, deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a

18

hexyl group, a heptyl group, an octyl group, a phenyl group, a naphthyl group, a fluorenyl group, a deuterated phenyl group, or a dimethylfluorenyl group;  $T_1$  may be O or S;  $Ar_1$  and  $Ar_2$  may be each independently represented by any one of Formulae 2A through 2J above;  $L_1$  and  $L_2$  may be each independently represented by any one of Formulae 5A through 5K above; a may be 0 or 1; and b may be 1 or 2, but the present embodiment is not limited thereto.

The styryl-based compound represented by Formula 1 above may be, but is not limited to, any one of Compounds 1 through 95 and 101 through 182 below:





19

20

-continued



9

10





11

12





13

14



15

16





21

22



-continued  
17

18





19

20





21

22



23

24





23

24



-continued  
25

26





27

28





29

30



31

32





25

26



-continued  
33

34





35

36





37

38



39

40





27

28

-continued



41

42





43

44





45

46



47

48





29

30

-continued



49

50





51

52





53

54





55

56



57

58





31

32

-continued



59

60





61

62





63

64





65

66



67

68





33

34

-continued



69

70





71

72





73

74





75

76





77

78



79

80





35

36

-continued



81

82




83

84





85




86


87



88



89




37

38



-continued

90




91


92




93



94



95





39

40

-continued



101

102





103

104





105

106





107

108



109

110





41

42

-continued



111

112





113

114





115

116



117

118





43

44

-continued



119

120





121

122





123

124





125

126



127

128





45

46



-continued  
129

130





131

132





133

134



135

136





47

48



-continued  
137

138





139

140





141

142



143

144





49

50



-continued  
145

146





147

148





149

150





151

152



153

154





51

52

-continued



155

156




157

158





159





160

161





162

163



164

165





53

54

-continued



166

167





168

169





170

171




172


173



174



175




55


56

-continued


176




177




178



179



180




57


58

-continued

181




182




When the styryl-based compound represented by Formula 1 above is synthesized, 100% deuteration of a styryl group may not be performed and thus a compound in which one of two carbon atoms of the styryl group is not substituted with deuterium and/or a compound in which both two carbon atoms of the styryl group are not substituted with deuterium may be simultaneously synthesized. Thus, a composition containing a styryl-based compound according to an embodiment of the present invention may include the styryl-based compound represented by Formula 1 above and may also include at least one of a styryl-based compound represented by Formula 1-1H-1 below, a styryl-based compound represented by Formula 1-1H-2 below, and a styryl-based compound represented by Formula 1-2H below:


30 For example, when Compound 3 is synthesized, i) only Compound 3 may be obtained, or ii) a composition including Compound 3 and at least one of Compounds 3-1H-1, 3-1H-2, and 3-2H may be obtained:



Formula 1-1H-1




Formula 1-H-2



Formula 1-2H


Compound 3



45

50

Compound 3-1H-1



55


60

65

A detailed description of substituents of Formula 1-1H-1, 1-1H-2, and 1-2H above is already provided above.

59

-continued



A deuterium substitution rate of the composition may be defined according to Equation 1 below and may be 70% or more:

$$\text{Deuterium substitution rate(\%)} = N_D / N_{D+H} \times 100 \quad (1)$$

In Equation 1 above,  $N_D$  is the number of deuterium atoms bonded to carbon atoms of the styryl group in Formulae 1, 1-1H-1, and 1-1H-2 above; and  $N_{D+H}$  is the total number of deuterium atoms and hydrogen atoms that are bonded to carbon atoms of the styryl group in Formulae 1, 1-1H-1, 1-1H-2, and 1-2H above.

When the deuterium substitution rate of the composition is 70% or more, an organic light-emitting diode (OLED) including an organic layer including the composition may exhibit low driving voltage, high efficiency, high brightness, long lifetime, and the like due to the styryl-based compound of Formula 1 above, which will be described later.


For example, the deuterium substitution rate of the composition may be, but is not limited to, 75% or more, 80% or more, 85% or more, 90% or more, or 95% or more.

For example, the deuterium substitution rate of the composition may be, but is not limited to, 96% or more, 97% or more, 98% or more, or 99% or more.

When an OLED including a compound having a styryl group (that is,  $-\text{CH}=\text{CH}-$ ) having a carbon-carbon double bond is stored and/or operated, the styryl group may have low resistance to Joule's heat or an electric field that is generated between organic layers, inside the organic layers, and/or between one of the organic layers and one of the electrodes and may have low resistance to heat provided during sublimation refinement or vapor deposition of a compound. Thus, a single bond of carbon-hydrogen, which is connected to a double bond in a styryl group exposed to heat and/or an electric field, may be weakened and may be disconnected in the form of radical. However, since a single bond of carbon-deuterium is about 7 times greater than a single bond of

60

carbon-hydrogen, a styryl group having a single bond of carbon-deuterium may have a stable structure. Since carbon is connected to deuterium by a single bond (refer to an ellipse indicated by dotted lines of Formula 1' below) in the styryl group of the styryl-based compound represented by Formula 1 above, although the styryl group is exposed to heat and/or an electric field, a single bond of carbon-deuterium may be effectively maintained according to a kinetic isotope effect:



Thus, although the styryl-based compound represented by Formula 1 above is exposed to heat and/or an electric field when an OLED including the same, since radicalization of a fully deuterated styryl group (i.e.,  $-\text{DC}=\text{CD}-$ ) of Formula 1 above is remarkably delayed compared to radicalization of a styryl group without deuterium (i.e.,  $-\text{CH}=\text{CH}-$ ), a structure of the styryl-based compound represented by Formula 1 may be maintained. Thus, an OLED including the styryl-based compound represented by Formula 1 may have low driving voltage, high efficiency high brightness, and long lifetime.

The styryl-based compound of Formula 1 and a composition including the same may be synthesized using a known organic synthesis method. The synthesis method of the styryl-based compound of Formula 1 may be easily understood by one of ordinary skill in the art with reference to Examples, which will be described later. The composition including the styryl-based compound represented by Formula 1 and at least one of the compounds represented by Formula 1-1H-1, Formula 1-1H-2, and Formula 1-2H may be obtained when a styryl group is imperfectly deuterated in the synthesis of the compound represented by Formula 1 above rather than being obtained by adding at least one of the compound represented by Formula 1-1H-1 above, the compound represented by Formula 1-1H-2 above, and the compound represented by Formula 1-2H above.

The styryl-based compound of Formula 1 or the composition containing the same may be used between a pair of electrodes of an OLED. For example, the styryl-based compound of Formula 1 or the composition containing the same may be used in an emission layer (EML) and/or between an anode and the EML (e.g., a hole injection layer (HIL), a hole transport layer (HTL), or a functional layer having hole injection and transport abilities).

According to another embodiment of the present invention, there is provided an OLED including a first electrode, a second electrode facing the first electrode, and an organic layer interposed between the first electrode and the second electrode, wherein the organic layer includes at least one of the styryl-based compound of Formula 1 or at least of the composition containing the same as described above.

The expression "the organic layer may include at least one of the styryl-based compound of Formula 1" as used herein means that the organic layer includes one of the styryl-based compounds represented by Formula 1 above or at least two different compounds selected from the styryl-based compounds represented by Formula 1 above.

For example, the organic layer may include only Compound 3 as the styryl-based compound. In this regard, Compound 3 may be included in an EML of the OLED. Also, the



61

organic layer may include Compound 3 and Compound 19 as the styryl-based compound. In this regard, Compounds 3 and 19 may be included in the same layer (e.g., in an EML) or in different layers (e.g., Compound 3 may be included in an EML and Compound 19 may be included in a HTL).

Throughout this specification, the expression “the organic layer may include at least one of the composition” means that “the organic layer includes one of the composition or at least two different compounds selected from the compositions.



For example, the organic layer may include, as the composition, a composition containing Compound 3 above and at least one of Compound 3-1H-1, Compound 3-1H-2, and Compound 3-2H. In this case, the composition may be included in an EML of the OLED.

In addition, the organic layer may include, as the composition, a first composition including Compound 3 above and at least one of Compound 3-1H-1, Compound 3-1H-2, and Compound 3-2H, and a second composition including Compound 19 above and at least one of Compound 19-1H-1, Compound 19-1H-2, and Compound 19-2H. In this case, the first composition and the second composition may be included in the same layer (e.g., the first composition and the second composition may be included in an EML) or in different layers (e.g., the first composition is included in an EML, and the second composition is included in a HTL).



62

-continued



The organic layer may include at least one of a HIL, a HTL, a functional layer having hole injection and transport abilities (hereinafter, referred to as “H-functional layer”), a buffer layer, an electron blocking layer (EBL), a hole blocking layer (HBL), an electron transport layer (ETL), an electron injection layer (EIL), and a functional layer having electron transport and injection abilities (hereinafter, referred to as “E-functional layer”).

The term “organic layer” used herein refers to a single layer or multiple layers interposed between the first electrode and the second electrode.

The organic layer may include an EML and the EML may include the styryl-based compound or the composition containing the styryl-based compound.

The styryl-based compound included in the EML may act as a fluorescent dopant. For example, the styryl-based compound may act as a blue fluorescent dopant that emits blue light.

63

FIG. 1 is a schematic cross-sectional view of an OLED 10 according to an embodiment of the present invention. Hereinafter, a structure and a manufacturing method of an OLED will be described in more detail with reference to FIG. 1.

A substrate 11 may be a substrate used in a general OLED, and may be a glass substrate or a transparent plastic substrate having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and waterproofness.

A first electrode 13 may be formed by applying a first electrode material on the substrate 11 by deposition or sputtering. When the first electrode 13 is an anode, the first electrode material may be selected from materials having a high work function so as to facilitate hole injection. The first electrode 13 may be a reflective electrode or a transparent electrode. Examples of the first electrode material may include indium-tin oxide (ITO), Indium-zinc-oxide (IZO), tin oxide ( $\text{SnO}_2$ ), and zinc oxide (ZnO). Also, when magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag) is used as the first electrode material, the first electrode 13 may be formed as a reflective electrode.

The first electrode 13 may be formed as a single layer or have a multi-layered structure having at least two layers. For example, the first electrode 13 may have a three-layered structure, e.g., ITO/Ag/ITO, but is not limited thereto.

An organic layer 15 is formed on the first electrode 13.

The organic layer 15 may include at least one of a HIL, a HTL, and a H-functional layer; an EML; an ETL; and an EIL.


The HIL may be formed on the first electrode 13 by using various methods such as vacuum deposition, spin coating, casting, or Langmuir-Blodgett (LB) deposition.

When the HIL is formed by vacuum deposition, the deposition conditions may vary according to a compound used as a material for forming the HIL, a structure of a desired HIL, and thermal characteristics. For example, the deposition condition may be, but is not limited to, a deposition temperature of about 100 to about 500° C., a degree of vacuum of about  $10^{-8}$  to about  $10^{-3}$  torr, and a deposition speed of about 0.01 to about 100 Å/sec.

When the HIL is formed by spin coating, the coating condition may vary according to a compound used as a material for forming the HIL, a structure of a desired HIL, and thermal characteristics. For example, the coating condition may be, but is not limited to, a coating speed of about 2,000 to about 5,000 rpm and a heat treatment temperature for removing a solvent after coating of about 80 to about 200° C.

The material for forming the HIL may be a known hole injection material. Examples of the known hole injection material include, but are limited to, N,N'-diphenyl-N,N'-bis-[4-(phenyl-m-tolyl-amino)-phenyl]-biphenyl-4,4'-diamine (DNTPD), a phthalocyanine compound such as copper phthalocyanine, 4,4',4''-tris(3-methylphenylphenylamino)triphenylamine (m-MTDATA), N,N'-di(1-naphthyl)-N,N'-diphenylbenzidine (NPB), TDATA, 2-TNATA, polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), and polyaniline/poly(4-styrenesulfonate) (PANI/PSS):

64




65

The thickness of the HIL may be in the range of about 100 Å to about 10,000 Å, for example, in the range of about 100 Å to about 1,000 Å. When the thickness of the HIL is within this range, satisfactory hole injection properties may be obtained without a substantial increase in driving voltage.

Next, an HTL may be formed on the HIL by using various methods such as vacuum deposition, spin coating, casting, or LB deposition. When the HTL is formed by vacuum deposition or spin coating, the deposition and coating conditions vary according to a used compound. However, in general, the condition may be almost the same as the condition for forming the HIL.

As a material for forming the HTL, at least one of a styryl-based compound, a composition containing a styryl-based compound, and a known hole transporting material. Examples of the known hole transporting material include, but are not limited to, carbazole derivatives such as N-phenylcarbazole and polyvinylcarbazole, N,N'-bis(3-methylphenyl)-N,N'-diphenyl-[1,1'-biphenyl]-4,4'-diamine (TPD), 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), and N,N'-di(1-naphthyl)-N,N'-diphenylbenzidine (NPB).




The thickness of the HTL may be in the range of about 50 Å to about 2,000 Å, for example, in the range of about 100 Å to about 1,500 Å. When the thickness of the HTL is within this range, satisfactory hole transport properties may be obtained without a substantial increase in driving voltage.

At least one of the hole injection material and the hole transporting material as described above may be included in the H-functional layer. The thickness of the H-functional layer may be in the range of about 500 Å to about 10,000 Å, for example, in the range of about 100 Å to about 1,000 Å. When the thickness of the H-functional layer is within this range, satisfactory hole injection and transport properties may be obtained without a substantial increase in driving voltage.


At least one of the HIL, the HTL, and the H-functional layer may include at least one of a compound represented by Formula 300 below and a compound represented by Formula 350 below:

66

Formula 300



Formula 350



wherein Ar<sub>11</sub>, Ar<sub>12</sub>, Ar<sub>21</sub>, and Ar<sub>22</sub> may be each independently a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> arylene group. Detailed descriptions for Ar<sub>11</sub>, Ar<sub>12</sub>, Ar<sub>21</sub>, and Ar<sub>22</sub> refer to the detailed description for L<sub>1</sub> above.

In Formula 300 above, e and f may be each independently an integer of 0 to 5, for example, 0, 1, or 2. For example, e may be 1, and f may be 0, but e and f are not limited to the above example.


In Formulae 300 and 350 above, R<sub>51</sub> through R<sub>58</sub>, R<sub>61</sub> through R<sub>69</sub>, and R<sub>71</sub> and R<sub>72</sub> may be each independently hydrogen, deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>60</sub> cycloalkyl group, a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> aryloxy group, or a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> arylthio group. For example, R<sub>51</sub> through R<sub>58</sub>, R<sub>61</sub> through R<sub>69</sub>, and R<sub>71</sub> and R<sub>72</sub> may be each independently one of hydrogen; deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a C<sub>1</sub>-C<sub>10</sub> alkyl group (e.g., methyl, ethyl, propyl, butyl, pentyl, hexyl, and the like), a C<sub>1</sub>-C<sub>10</sub> alkoxy group (e.g., methoxy, ethoxy, propoxy, butoxy, pentoxy, and the like); a C<sub>1</sub>-C<sub>10</sub> alkyl group and a C<sub>1</sub>-C<sub>10</sub> alkoxy group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl

67

group or a salt thereof, a sulfonic acid group or a salt thereof, and a phosphoric acid or a salt thereof; a phenyl group; a naphthyl group; an anthryl group; a fluorenyl group; a pyrenyl group; and a phenyl group, a naphthyl group, an anthryl group, a fluorenyl group, and a pyrenyl group that substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>10</sub> alkyl group, and a C<sub>1</sub>-C<sub>10</sub> alkoxy group, but are not limited thereto.



In Formula 300 above, R<sub>59</sub> may be one of a phenyl group; a naphthyl group; an anthryl group; a biphenyl group; a pyridinyl group; and a phenyl group, a naphthyl group, an anthryl group, a biphenyl group, and a pyridinyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> alkyl group, and a substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> alkoxy group.

The compound of Formula 300 may be represented by Formula 300A below, but is not limited thereto:



wherein a detailed description of R<sub>51</sub>, R<sub>60</sub>, R<sub>61</sub>, and R<sub>59</sub> is already provided above.


For example, at least one of the HIL, the HTL, and the H-functional layer may include at least one of Compounds 301 through 320 below, but is not limited thereto:




68

-continued


302



303




304



305

69  
-continued

306



70  
-continued


309

5

10

15

20



310


25

307

30

35

40



45



50

308

55


60

65



71  
-continued

312



72  
-continued

314

5

10

15

20

25

30

35

40

313

315



45

50

55


60

65



73  
-continued

316



74  
-continued

318

5

10

15

20

25

30

35

40

317




45

50

55


60

65




75

-continued




At least one of the HIL, the HTL, and the H-functional layer may further include a charge-generating material so as to increase the conductive of the layers, in addition to the known hole injection material, the known hole transporting material, the material for forming the H-functional layer having hole injection and transport abilities, the styryl-based compound and/or the composition containing the styryl-based compound.

The charge-generating material may be, for example, a p-dopant. The p-dopant may be one of a quinine derivative, a metal oxide, and a cyano-containing compound, but is not limited thereto. Examples of the p-dopant may include, but are not limited to, quinone derivatives such as tetra-cyanoquinodimethane (TCNQ) and 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinodimethane (F4-CTNQ); metal oxides such as an tungsten oxide and a molybdenum oxide; and cyano-containing compounds such as Compound 200 below and the like.



&lt;Compound 200&gt;



&lt;F4-CTNQ&gt;

When the HIL, the HTL, or the H-functional layer further includes the charge-generating material, the charge-generating material may be homogeneously or inhomogeneously dispersed in the HIL, the HTL, or the H-functional layer.

76

320

A buffer layer may be interposed between the EML and at least one of the HIL, the HTL, and the H-functional layer. The buffer layer increases efficiency by compensating for an optical resonance distance according to the wavelength of light emitted from the EML. The buffer layer may include a known hole injection material and a known hole transporting material. Also, the buffer layer may include the same material as one of the materials included in the HIL, the HTL, and the H-functional layer.

10

An EML may be formed on the HTL, the H-functional layer, or the buffer layer by vacuum deposition, spin coating, casting, or LB deposition. When the EML is formed by vacuum deposition or spin coating, the deposition and coating conditions vary according to a used compound. However, in general, the condition may be almost the same as the condition for forming the HIL.

15

20

The EML may include at least one of the styryl-based compound or at least one of the composition containing the styryl-based compound.

The EML may further include a host.

Examples of the host may include, but are not limited to, Alq<sub>3</sub>, 4,4'-N,N'-dicarbazole-biphenyl (CBP), poly(n-vinyl-carbazole) (PVK), 9,10-di(naphthalene-2-yl)anthracene (ADN), TCTA, 1,3,5-tris(N-phenylbenzimidazole-2-yl)benzene (TPBI), 3-tert-butyl-9,10-di(naphth-2-yl)anthracene (TBADN), E3, and distyrylarylene (DSA), dmCBP (refer to Formula below), and Compounds 501 through 509 below.

30

35

40


45

50


55

60


65



TPBI









TBADN



E3

77


-continued



78

-continued

5



10

15

20

25

30

35

40

501

45


50

502

55

60

65







503

504

505

79

-continued



80


Also, the host may be an anthracene-based compound represented by Formula 400 below:

506

5

Formula 400

10



15

507

20

wherein  $Ar_{111}$  and  $Ar_{112}$  may be each independently a substituted or unsubstituted  $C_5-C_{60}$  arylene group;  $Ar_{113}$  through  $Ar_{116}$  may be each independently a substituted or unsubstituted  $C_1-C_{10}$  alkyl group or a substituted or unsubstituted  $C_5-C_{60}$  aryl group; and  $g, h, i,$  and  $j$  may be each independently an integer of 0 to 4.

For example, in Formula 400 above,  $Ar_{111}$  and  $Ar_{112}$  may be each independently one of a phenylene group; a naphthylene group; a phenanthrenylene group; a pyrenylene group; and a phenylene group, a naphthylene group, a phenanthrenylene group, a fluorenyl group, and a pyrenylene group that are substituted with at least one of a phenyl group, a naphthyl group, and an anthryl group.

In Formula 400 above,  $g, h, i,$  and  $j$  may be each independently 0, 1, or 2.

508

35


509

40

In Formula 400 above,  $Ar_{113}$  through  $Ar_{116}$  may be each independently, but are not limited to, a  $C_1-C_{10}$  alkyl group that is substituted with at least one of a phenyl group, a naphthyl group, and an anthryl group; a phenyl group; a naphthyl group; an anthryl group; a pyrenyl group; a phenanthrenyl group; a fluorenyl group; a phenyl group, a naphthyl group, an anthryl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1-C_{60}$  alkyl group, a  $C_2-C_{60}$  alkenyl group, a  $C_2-C_{60}$  alkynyl group, a  $C_1-C_{60}$  alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group; and

55


60



65


For example, the anthracene-based compound of Formula 400 may be, but is not limited to, one of the compounds below:

81




82

-continued




83

-continued








84

-continued




85

-continued




86


-continued




10




20



30



40



50




60




87

-continued



88

-continued



5

10


15

Also, an anthracene-based compound represented by Formula 401 below may be used as the host:

<Formula 401>

25

30



35

wherein a detailed description of  $Ar_{122}$  through  $Ar_{125}$  is the same as the description of  $Ar_{113}$  of Formula 400 above.

In Formula 401 above,  $Ar_{126}$  and  $Ar_{127}$  may be each independently a  $C_1$ - $C_{10}$  alkyl group (e.g., a methyl group, an ethyl group, or a propyl group).

40

In Formula 401 above,  $k$  and  $l$  may be each independently an integer of 0 to 4. For example,  $k$  and  $l$  may be each independently 0, 1, or 2.

45




For example, the anthracene-based compound of Formula 401 may be, but is not limited to, one of the following compounds:

50



55

60


65



89  
-continued




90  
-continued



5  
10  
15  
20  
25  
If the OLED is a full-color OLED, the EML may be patterned as a red EML, a green EML, and a blue EML. In this regard, the above-described styryl-based compound or the composition containing the styryl-based compound may be included in the blue EML as a blue fluorescent dopant.

30  
35  
At least one of the red EML, the green EML, and the blue EML may include the following dopants (ppy=phenylpyridine).

For example, compounds described below may be used as a blue dopant, but are not limited thereto.



F<sub>2</sub>Irpic





(F<sub>2</sub>ppy)<sub>2</sub>Ir(tmd)

Ir(dfpp)<sub>3</sub>


91

92

-continued




DPVBi



DPAVBi

93




TBPe


94

-continued


For example, compounds described below may be used as a red dopant, but are not limited thereto. In addition, DCM or DCJTb below may be used as the red dopant.




PtOEP



Ir(piq)<sub>3</sub>



Btp<sub>2</sub>Ir(acac)



25

30

35

40

45


Ir(pq)<sub>2</sub>(acac)

50


55

60

65







Ir(2-phq)<sub>3</sub>



Ir(BT)<sub>2</sub>(acac)




95

-continued





For example, compounds described below may be used as a green dopant, but are not limited thereto. C545T below may be used as the green dopant.

96









Examples of the dopant included in the EML include Pt-complexes below, but are not limited thereto:



97

-continued



98

-continued

D3

5

10

D4 15

20

D5 25

30

D6 35

40






D7 45

50

55

D8 60

65



D9






D10

D11

D12

D13


**99**  
-continued



**100**  
-continued

D14


5



10

D15

15




20

25

D16

30




35

40

D17

45




50

55

D18

60



65

D19







D20

D21

D22

D23

**101**  
-continued




**102**  
-continued

D24

5

10




D30

D25

15

20




D31

D26

25

30




D32

D27

35

40




D33

D28

45

50

55




D34

D29


60

65



D35

**103**  
-continued




D36

5

10

D37


15



20

25


30



D38

35


40



D39

45

50




D40


55

**104**  
-continued


D41




D42




D43



D44



D45








60

65


105

-continued




106

D46




Os(fppz)<sub>2</sub>(CO)<sub>2</sub>

D47




Os(fppz)<sub>2</sub>(PPh<sub>2</sub>Me)<sub>2</sub>

D48




Os(bppz)<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>

D49



Os(fptz)<sub>2</sub>(PPh<sub>2</sub>Me)<sub>2</sub>

D50



Os(hptz)<sub>2</sub>(PPh<sub>2</sub>Me)<sub>2</sub>




Also, examples of the dopant included in the EML include, but are not limited to, Os-complexes:

When the EML includes a host and a dopant, the amount of the dopant in the EML may be generally in the range of about 0.01 to about 15 parts by weight based on 100 parts by weight of the host, but is not limited thereto.

107

The thickness of the EML may be in the range of about 100 Å to about 1,000 Å, for example, in the range of about 200 Å to about 600 Å. When the thickness of the EML is within this range, excellent luminescent properties may be obtained without a substantial increase in driving voltage.



Next, an ETL may be formed using various methods such as vacuum deposition, spin coating, or casting. When the ETL is formed by vacuum deposition or spin coating, the deposition and coating conditions vary according to a used compound. However, in general, the condition may be almost the same as the condition for forming the HIL. A material for forming the ETL may be a known electron transporting material that stably transports electrons injected from a cathode. Examples of the known electron transporting material may include, but are not limited to, a quinoline derivative such as tris(8-quinolinolate)aluminum (Alq<sub>3</sub>), TAZ, Balq, beryllium bis(benzoquinoline-10-olate (Bebq<sub>2</sub>), ADN, and known materials such as Compound 201 and Compound 202 below.



108

-continued


Compound 202



The thickness of the ETL may be in the range of about 100 Å to about 1,000 Å, for example, in the range of about 150 Å to about 500 Å. When the thickness of the ETL is within this range, satisfactory electron transport properties may be obtained without a substantial increase in driving voltage.

In addition, the ETL may further include a metal-containing material, in addition to a known electron transporting organic compound.

The metal-containing material may include a Li-complex. Examples of the Li-complex may include lithium quinolate (LiQ) and Compound 203 below:



Also, an EIL, which facilitates electron injection from a cathode, may be formed on the ETL, and a material for forming the EIL is not particularly limited.

The material for forming the EIL may include a known material for forming an EIL, such as LiF, NaCl, CsF, Li<sub>2</sub>O, or BaO. The deposition condition of the EIL may vary according to a used compound. However, in general, the condition may be almost the same as the condition for forming the HIL.


The thickness of the EIL may be in the range of about 1 Å to about 100 Å, for example, in the range of about 3 Å to about

90 Å. When the thickness of the EIL is within this range, satisfactory electron injection properties may be obtained without a substantial increase in driving voltage.

A second electrode **17** is formed on the organic layer **15**. The second electrode **17** may be a cathode, which is an electron injection electrode. In this regard, a metal for forming the second electrode **17** may include a metal having low work function, such as metal, an alloy, an electric conducting compound, and mixtures thereof. In particular, the second electrode **17** may be formed as a thin film by using lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag), thus being transparent. In order to obtain a top-emission type OLED, the second electrode **17** may be formed as a transparent electrode by using ITO or IZO.

The OLED has been described with reference to FIG. 1, but is not limited thereto.

Also, when a phosphorescent dopant is included in the EML, a HBL may be formed between the ETL and the EML or between the H-functional layer and the EML by vacuum deposition, spin coating, casting or LB deposition so as to prevent triplet excitons or holes from being diffused to the ETL. When the HBL is formed by vacuum deposition or spin coating, the conditions thereof may vary according to a used compound. However, in general, the conditions may be almost the same as the condition for forming the HIL. The HBL may include a known hole blocking material. Examples of the known hole blocking material may include an oxadiazole derivative, a triazole derivative, and a phenanthroline derivative. For example, BCP may be used as a hole blocking material.



The thickness of the HBL may be in the range of about 20 Å to about 1,000 Å, for example, in the range of about 30 Å to about 300 Å. When the thickness of the HBL is within this range, excellent hole blocking properties may be obtained without a substantial increase in driving voltage.

Examples of the unsubstituted  $C_1$ - $C_{60}$  alkyl group (or  $C_1$ - $C_{60}$  alkyl group) include  $C_1$ - $C_{60}$  linear or branched alkyl groups such as methyl, ethyl, propyl, isobutyl, sec-butyl, pentyl, iso-amyl, hexyl, and the like. The substituted  $C_1$ - $C_{60}$  alkyl group may be a group in which at least one hydrogen of the unsubstituted  $C_1$ - $C_{60}$  alkyl group is substituted with deuterium, a halogen atom, a hydroxyl group, a nitro group, a cyano group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{60}$  alkyl group, a  $C_2$ - $C_{60}$  alkenyl group, a  $C_2$ - $C_{60}$  alkynyl group, a  $C_6$ - $C_{60}$  aryl group, a  $C_2$ - $C_{60}$  heteroaryl group,  $-N(Q_{11})(Q_{12})$ , and  $-Si(Q_{13})(Q_{14})(Q_{15})$  (wherein  $Q_{11}$  through  $Q_{15}$  may be each independently selected from hydro-

gen, a  $C_1$ - $C_{60}$  alkyl group, a  $C_2$ - $C_{60}$  alkenyl group, a  $C_2$ - $C_{60}$  alkynyl group, a  $C_5$ - $C_{60}$  aryl group, and a  $C_2$ - $C_{60}$  heteroaryl group).

The unsubstituted  $C_1$ - $C_{60}$  alkoxy group (or  $C_1$ - $C_{60}$  alkoxy group) has a formula of  $-OA$  (in this regard, A is the unsubstituted  $C_1$ - $C_{60}$  alkyl group as described above) and examples thereof include methoxy, ethoxy, isopropoxy, and the like. At least one hydrogen atom of the unsubstituted  $C_1$ - $C_{60}$  alkoxy group may be substituted with the same substituent as in the substituted  $C_1$ - $C_{60}$  alkyl group described above.

The unsubstituted  $C_2$ - $C_{60}$  alkenyl group (or  $C_2$ - $C_{60}$  alkenyl group) is interpreted to contain at least one carbon-carbon double bond in the middle or at a terminal of the unsubstituted  $C_2$ - $C_{60}$  alkyl group. Examples of the unsubstituted  $C_2$ - $C_{60}$  alkenyl group include ethenyl, propenyl, butenyl, and the like. At least one hydrogen atom of the unsubstituted  $C_2$ - $C_{60}$  alkenyl group may be substituted with the same substituent as in the substituted  $C_1$ - $C_{60}$  alkyl group described above.

The unsubstituted  $C_2$ - $C_{60}$  alkynyl group (or  $C_2$ - $C_{60}$  alkynyl group) is interpreted to contain at least one carbon-carbon triple bond in the middle or at a terminal of the  $C_2$ - $C_{60}$  alkyl group defined above. Examples of the unsubstituted  $C_2$ - $C_{60}$  alkynyl group include ethynyl, propynyl, and the like. At least one hydrogen atom of the unsubstituted  $C_2$ - $C_{60}$  alkynyl group may be substituted with the same substituent as in the substituted  $C_1$ - $C_{60}$  alkyl group described above.

The unsubstituted  $C_5$ - $C_{60}$  aryl group indicates a monovalent group having an aromatic carbocyclic system that has 5 to 60 carbon atoms and at least one aromatic ring and the unsubstituted  $C_5$ - $C_{60}$  arylene group indicates a divalent group having an aromatic carbocyclic system that has 5 to 60 carbon atoms and at least one aromatic ring. If the  $C_5$ - $C_{60}$  aryl group and the  $C_5$ - $C_{60}$  arylene group each independently have two or more aromatic rings, the rings may be fused with each other. At least one hydrogen atom of each of the unsubstituted  $C_5$ - $C_{60}$  aryl group and the unsubstituted  $C_5$ - $C_{60}$  arylene group may be substituted with the same substituent as in the substituted  $C_1$ - $C_{60}$  alkyl group described above.

Examples of the substituted or unsubstituted  $C_5$ - $C_{60}$  aryl group include, but are not limited to, a phenyl group, a  $C_1$ - $C_{10}$  alkylphenyl group (e.g., an ethylphenyl group), a  $C_1$ - $C_{10}$  alkylbiphenyl group (e.g., an ethylbiphenyl group), a halophenyl group (e.g., an o-, m- and p-fluorophenyl group, and a dichlorophenyl group), a dicyanophenyl group, a trifluoromethoxyphenyl group, an o-, m-, and p-tolyl group, an o-, m- and p-cumenyl group, a mesityl group, a phenoxyphe-nyl group, an ( $\alpha,\alpha$ -dimethylbenzene)phenyl group, a ( $N,N'$ -dimethyl)aminophenyl group, a ( $N,N'$ -diphenyl)aminophenyl group, a pentalenyl group, an indenyl group, a naphthyl group, a halonaphthyl group (e.g., a fluoronaphthyl group), a  $C_1$ - $C_{10}$  alkyl-naphthyl group (e.g., a methyl-naphthyl group), a  $C_1$ - $C_{10}$  alkoxy-naphthyl group (e.g., a methoxy-naphthyl group), an anthracenyl group, an azulenyl group, a heptalenyl group, an acenaphthylenyl group, a phenalenyl group, a fluorenyl group, an anthraquinolyl group, a methylanthryl group, a phenanthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an ethyl-chrysenyl group, a picenyl group, a perylenyl group, a chloroperlylenyl group, a pentaphenyl group, a pentalenyl group, a tetraphenylenyl group, a hexaphenyl group, a hexacenyl group, a rubicenyl group, a coronenyl group, a trinaphthylenyl group, a heptaphenyl

111

group, a heptacenyl, a pyranthrenyl group, and an ovalenyl group. Examples of the substituted  $C_5$ - $C_{60}$  aryl group may be easily understood with reference to the examples of the unsubstituted  $C_5$ - $C_{60}$  aryl group described above and the substituents of the substituted  $C_1$ - $C_{60}$  alkyl group. Examples of the substituted or unsubstituted  $C_5$ - $C_{60}$  arylene group may be easily understood with reference to the substituted or unsubstituted  $C_5$ - $C_{60}$  aryl group described above.

The unsubstituted  $C_2$ - $C_{60}$  heteroaryl group indicates a monovalent group having at least one aromatic ring system including carbon rings and at least one hetero atom selected from the group consisting of N, O, P, and S, and the unsubstituted  $C_2$ - $C_{60}$  heteroarylene group indicates a divalent group having at least one aromatic ring system including carbon rings and at least one hetero atom selected from the group consisting of N, O, P, and S. In this regard, if the  $C_2$ - $C_{60}$  heteroaryl group and the  $C_2$ - $C_{60}$  heteroarylene group each independently have two or more aromatic rings, the rings may be fused with each other. At least one hydrogen atom of each of the unsubstituted  $C_2$ - $C_{60}$  heteroaryl group and the unsubstituted  $C_2$ - $C_{60}$  heteroarylene group may be substituted with the same substituents as in the  $C_1$ - $C_{60}$  alkyl group described above.

Examples of the unsubstituted  $C_2$ - $C_{60}$  heteroaryl group include, but are not limited to, a pyrazolyl group, an imidazolyl group, an oxazolyl group, a thiazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a pyridinyl group, a pyridazinyl group, a pyrimidinyl group, a triazinyl group, a carbazolyl group, an indolyl group, a quinolinyl group, an isoquinolinyl group, a benzoimidazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group. Examples of the unsubstituted  $C_2$ - $C_{60}$  heteroarylene group may be easily understood with reference to the examples of the substituted or unsubstituted  $C_2$ - $C_{60}$  arylene group.

The substituted or unsubstituted  $C_5$ - $C_{60}$  aryloxy group has a formula of  $-OA_2$  wherein  $A_2$  is the substituted or unsubstituted  $C_5$ - $C_{60}$  aryl group as described above, and the substituted or unsubstituted  $C_5$ - $C_{60}$  arylthio group has a formula of  $-SA_3$  wherein  $A_3$  is the substituted or unsubstituted  $C_5$ - $C_{60}$  aryl group described above.


An OLED according to an embodiment of the present invention will now be described in detail with reference to the following Examples. These Examples are for illustrative purposes only and are not intended to limit the scope of the invention.

## EXAMPLE

## Synthesis Example 1


## Synthesis of Compound 3

Compound 3 was synthesized according to Reaction Scheme 1 below:



112

-continued



## Synthesis of Intermediate 3-A

5.14 g (20.0 mmol) of 9-bromophenanthrene, 2.8 g (30.0 mmol) of aniline, 0.366 g (0.4 mmol) of tris(dibenzylideneacetone)dipalladium(0): $Pd_2(dba)_3$ , 0.081 g (0.4 mmol) of tri-tert-butylphosphine: $P(t-Bu)_3$ , and 2.88 g (30.0 mmol) of  $KOtBu$  were dissolved in 60 mL of toluene and stirred at 85° C. for 4 hours. The obtained reaction solution was cooled down to room temperature and then extracted three times with 50 ml of water and 50 ml of diethylether. The obtained organic layer was dried with magnesium sulfate and a solvent was evaporated therefrom to obtain a crude product. The crude product was purified with silica gel column chroma-

113

tography to obtain 4.63 g of Intermediate 3-A (yield 86%). The obtained compound was confirmed by MS/FAB.

$C_{20}H_{15}N$ : cal. 269.12, found 270.13.

#### Synthesis of Intermediate 3-B

2.69 g (10.0 mmol) of Intermediate 3-A, 2.75 g (15.0 mmol) of 2-bromo-4-vinylbenzene, 0.183 g (0.2 mmol) of  $Pd_2(dba)_3$ , 0.041 g (0.2 mmol) of  $P(t-Bu)_3$ , and 1.44 g (15.0 mmol) of  $KOtBu$  were dissolved in 30 mL of toluene and stirred at 85° C. for 4 hours. The obtained reaction solution was cooled down to room temperature and then extracted three times with 30 ml of water and 30 ml of diethylether. The obtained organic layer was dried with magnesium sulfate and a solvent was evaporated therefrom to obtain a crude product. The crude product was purified with silica gel column chromatography to obtain 2.30 g of Intermediate 3-B (yield 62%). The obtained compound was confirmed by MS/FAB.

$C_{28}H_{21}N$ : cal. 371.17, found 372.16.

#### Synthesis of Compound 3-2H

1.86 g (5.0 mmol) of Intermediate 3-B, 1.04 g (5.0 mmol) of 2-bromonaphthalene, 0.056 g (0.25 mmol) of palladium (II) acetate:  $Pd(OAc)_2$ , 0.76 g (0.25 mmol) of tri(o-tolyl)phosphine ( $(p-tolyl)_3P$ ), and 1.019 g (10.0 mmol) of  $Et_3N$  were dissolved in 30 ml of dimethylacetamide (DMAc) and stirred at 100° C. for 4 hours. The obtained reaction solution was cooled down to room temperature and then extracted three times with 30 ml of water and 30 ml of diethylether. The obtained organic layer was dried with magnesium sulfate and a solvent was evaporated therefrom to obtain a crude product. The crude product was purified with silica gel column chromatography to obtain 1.82 g of Compound 3-2H (yield 73%). The obtained compound was confirmed by MS/FAB and  $^1H$  nuclear magnetic resonance (NMR).

$C_{38}H_{27}N$ : cal. 497.21, found 498.22.

$^1H$  NMR ( $CDCl_3$ , 400 MHz)  $\delta$  (ppm) 8.59 (d, 1H), 8.20-8.17 (m, 1H), 8.00-7.92 (m, 3H), 7.85-7.82 (m, 3H), 7.71-7.67 (m, 2H), 7.60-7.39 (m, 7H), 7.22-7.12 (m, 2H), 7.08-7.02 (m, 3H), 6.97-6.91 (m, 3H), 6.85-6.83 (m, 2H)

#### Synthesis of Compound 3

1.49 g (3.0 mmol) of Compound 3-2H, 0.086 g (0.09 mmol) of carbonylchlorohydridotris(triphenylphosphine) ruthenium(II) ( $[(Ph_3P)_3Ru(CO)(Cl)H]$ ), and 0.54 mL (30.0 mmol) of  $D_2O$  were dissolved in 30 ml of 1,4-dioxane and stirred at 80° C. for 12 hours. The obtained reaction solution was cooled down to room temperature to remove a solvent and then extracted three times with 30 ml of water and 30 ml of dichloromethane. The obtained organic layer was dried with magnesium sulfate and a solvent was evaporated therefrom to obtain a crude product. The crude product was purified with silica gel column chromatography to obtain 1.33 g of Compound 3 (yield 89%). The obtained compound was confirmed by MS/FAB and  $^1H$  NMR.

$C_{38}H_{25}D_2N$ : cal. 499.23, found 450.25.

$^1H$  NMR ( $CDCl_3$ , 400 MHz)  $\delta$  (ppm) 8.59 (d, 1H), 8.20-8.17 (m, 1H), 7.95-7.93 (m, 1H), 7.85-7.83 (m, 4H), 7.71-7.63 (m, 3H), 7.59-7.50 (m, 3H), 7.47-7.39 (m, 4H), 7.12-7.06 (m, 3H), 6.98-6.93 (m, 3H), 6.83-6.80 (m, 2H), deuterium substitution rate: 92.5%


#### Synthesis Example 2

#### Synthesis of Compound 33

Compound 33 was synthesized according to Reaction Scheme 2 below:

114

Reaction scheme 2



Intermediate 33-A

3.14 mL (20.0 mmol) of vinylboronic acid pinacol ester, 5.58 g (30.0 mmol) of 2,6-dibromonaphthalene, 1.15 g (1.0

## 115

mmol) of Pd(PPh<sub>3</sub>)<sub>4</sub>, and 8.29 g (60.0 mmol) of K<sub>2</sub>CO<sub>3</sub> were dissolved in 60 mL of a mixture solution of THF/H<sub>2</sub>O (volume ratio of 2/1) and stirred at 75° C. for 5 hours. The obtained reaction solution was cooled down to room temperature and then extracted three times with 30 ml of water and 50 ml of diethylether. The obtained organic layer was dried with magnesium sulfate and a solvent was evaporated therefrom to obtain a crude product. The crude product was purified with silica gel column chromatography to obtain 2.98 g of Intermediate 33-A (yield 64%). The obtained compound was confirmed by MS/FAB.

C<sub>12</sub>H<sub>9</sub>Br: cal. 231.99. found 233.01.

## Synthesis of Intermediate 33-B

2.84 g of Intermediate 33-B (yield 65%) was prepared in the same manner as in the synthesis of Intermediate 3-B of Synthesis Example 1, except that 9,9'-dimethylfluoren-2-ylphenylamine was used instead of Intermediate 3-A and Intermediate 33-A was used instead of 1-bromo-4-vinylbenzene. The obtained compound was confirmed by MS/FAB.

C<sub>33</sub>H<sub>27</sub>N: cal. 437.21. found 438.24.

## Synthesis of Compound 33-2H

2.27 g of Compound 33-2H (yield 74%) was prepared in the same manner as in the synthesis of Compound 3-2H of Synthesis Example 1, except that Intermediate 33-B was used instead of Intermediate 3-B and 9-bromophenanthrene was used instead of 2-bromonaphthalene. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

C<sub>47</sub>H<sub>35</sub>N: cal. 613.28. found 614.29.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.66 (d, 1H), 8.46 (d, 1H), 8.15-8.06 (m, 3H), 7.92 (s, 1H), 7.83-7.58 (m, 9H), 7.47 (dt, 1H), 7.36-7.27 (m, 3H), 7.14-7.02 (m, 5H), 6.95-6.92 (m, 2H), 6.87 (s, 1H), 6.82-6.80 (m, 2H), 1.67 (s, 6H)

## Synthesis of Compound 33

Compound 33 (yield 85%) was prepared in the same manner as in the synthesis of Compound 3 of Synthesis Example 1, except that Compound 33-2H was used instead of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

C<sub>47</sub>H<sub>33</sub>D<sub>2</sub>N: cal. 615.29. found 616.30.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.66 (d, 1H), 8.46 (d, 1H), 8.11 (d, 1H), 7.93-7.91 (m, 2H), 7.83-7.62 (m, 8H), 7.58 (d, 1H), 7.48 (dt, 1H), 7.36-7.30 (m, 2H), 7.14-7.05 (m, 4H), 6.98 (dt, 1H), 6.93-6.90 (m, 2H), 6.87 (s, 1H), 6.81-6.79 (m, 2H), 1.67 (s, 6H), deuterium substitution rate: 84%

## Synthesis Example 3

## Synthesis of Compound 1

Compound 1 was prepared in the same manner as in Synthesis Example 1, except that N-phenylnaphthalen-2-amine was used instead of Intermediate 3-A. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

C<sub>34</sub>H<sub>23</sub>D<sub>2</sub>N: cal. 449.21. found 450.22.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 7.86-7.82 (m, 4H), 7.79-7.76 (m, 1H), 7.68-7.64 (m, 2H), 7.57-7.38 (m, 9H),

## 116

7.13-7.06 (m, 3H), 6.99-6.96 (m, 2H), 6.86-6.83 (m, 1H), 6.45-6.42 (m, 2H), deuterium substitution rate: 91.8%

## Synthesis Example 4


## Synthesis of Compound 5

Compound 5 was prepared in the same manner as in Synthesis Example 1, except that 2-bromonaphthalene was used instead of 9-bromophenanthrene, and Intermediate 5-A-(1) was used instead of aniline in the synthesis of Intermediate 3-A. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

C<sub>46</sub>H<sub>30</sub>D<sub>2</sub>FN: cal. 619.26. found 620.24.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 7.87-7.83 (m, 4H), 7.78-7.76 (m, 1H), 7.74-7.69 (m, 2H), 7.67-7.60 (m, 5H), 7.56-7.38 (m, 14H), 7.14-7.09 (m, 2H), 6.93-6.89 (m, 2H)

5-A-(1)



## Synthesis Example 5

## Synthesis of Compound 6

Compound 6 was prepared in the same manner as in Synthesis Example 1, except that 4-bromobenzonitrile was used instead of 9-bromophenanthrene and naphthalene-2-amine was used instead of aniline in the synthesis of Intermediate 3-A. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

C<sub>35</sub>H<sub>22</sub>D<sub>2</sub>N<sub>2</sub>: cal. 474.21. found 445.22.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 7.81-7.76 (m, 4H), 7.71-7.64 (m, 2H), 7.59-7.53 (m, 4H), 7.48-7.44 (m, 2H), 7.42-7.31 (m, 4H), 7.13-7.09 (m, 1H), 7.02-6.99 (m, 1H), 6.94-6.88 (m, 4H), deuterium substitution rate: 93.4%

## Synthesis Example 6

## Synthesis of Compound 9

Compound 9 was prepared in the same manner as in Synthesis Example 2, except that 2-bromonaphthalene was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

C<sub>43</sub>H<sub>31</sub>D<sub>2</sub>N: cal. 565.27. found 566.29.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 7.90-7.82 (m, 5H), 7.78-7.62 (m, 6H), 7.58-7.43 (m, 3H), 7.35-7.30 (m, 2H), 7.13-7.06 (m, 4H), 6.90-6.83 (m, 2H), 6.65-6.63 (m, 1H), 6.42-6.40 (m, 2H), 1.68 (s, 6H) deuterium substitution rate: 86.4%

117

## Synthesis Example 7

## Synthesis of Compound 12

Compound 12 was prepared in the same manner as in Synthesis Example 2, except that 2,7-dibromo-9,9-dimethyl-9H-fluorene was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, N-phenyl-naphthalen-2-amine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B, and 2-bromonaphthalene was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{43}\text{H}_{31}\text{D}_2\text{N}$ : cal. 565.27. found 566.28.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 7.89-7.83 (m, 4H), 7.78-7.64 (m, 6H), 7.58-7.51 (m, 4H), 7.46-7.33 (m, 3H), 7.20-7.15 (m, 3H), 6.91-6.83 (m, 2H), 6.73-6.71 (m, 1H), 6.42-6.40 (m, 2H), 1.67 (s, 6H) deuterium substitution rate: 84.4%

## Synthesis Example 8

## Synthesis of Compound 18

Compound 18 was prepared in the same manner as in Synthesis Example 1, except that 4-bromobenzonitrile was used instead of 9-bromophenanthrene and naphthalene-2-amine was used instead of aniline in the synthesis of Intermediate 3-A, and 9-bromophenanthrene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR

$\text{C}_{39}\text{H}_{24}\text{D}_2\text{N}_2$ : cal. 524.22. found 525.24.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.67-8.64 (m, 1H), 8.46-8.44 (m, 1H), 8.10-8.08 (m, 1H), 7.85-7.76 (m, 3H), 7.69-7.63 (m, 4H), 7.57-7.36 (m, 8H), 7.18-6.98 (m, 6H), deuterium substitution rate: 91.3%

## Synthesis Example 9

## Synthesis of Compound 19

Compound 19 was prepared in the same manner as in Synthesis Example 1, except that 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine was used instead of Intermediate 3-A, and 9-bromophenanthrene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{43}\text{H}_{31}\text{D}_2\text{N}$ : cal. 565.27. found 566.26.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.67-8.64 (m, 1H), 8.46-8.44 (m, 1H), 8.10-8.08 (m, 1H), 7.85-7.81 (m, 2H), 7.78-7.76 (m, 1H), 7.69-7.64 (m, 2H), 7.56-7.46 (m, 4H), 7.36-7.30 (m, 1H), 7.14-7.04 (m, 5H), 6.96-6.83 (m, 4H), 6.59-6.58 (m, 1H), 6.43-6.40 (m, 2H), 1.67 (s, 6H), deuterium substitution rate: 85.8%

## Synthesis Example 10

## Synthesis of Compound 25

Compound 25 was prepared in the same manner as in Synthesis Example 1, except that 3-iodo-9-phenyl-9H-carbazole was used instead of 9-bromophenanthrene in the synthesis of Intermediate 3-A, and 9-bromophenanthrene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

118

$\text{C}_{46}\text{H}_{30}\text{D}_2\text{N}_2$ : cal. 614.27. found 615.28.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.67-8.64 (m, 1H), 8.46-8.44 (m, 1H), 8.22-8.19 (m, 1H), 8.11-8.09 (m, 1H), 7.85-7.81 (m, 2H), 7.69-7.64 (m, 2H), 7.52-7.22 (m, 13H), 7.18-7.14 (m, 2H), 7.06-7.04 (m, 1H), 6.97-6.94 (m, 3H), 6.86-6.83 (m, 1H), 6.52-6.50 (m, 2H), deuterium substitution rate: 89.4%

## Synthesis Example 11


## Synthesis of Compound 29

Compound 29 was prepared in the same manner as in Synthesis Example 1, except that Intermediate A-29-(1) below was used instead of 9-bromophenanthrene in the synthesis of Intermediate 3-A, and 9-bromophenanthrene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{40}\text{H}_{25}\text{D}_2\text{NS}$ : cal. 555.20. found 556.23.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.67-8.64 (m, 1H), 8.46-8.44 (m, 1H), 8.12-8.04 (m, 2H), 7.85-7.81 (m, 3H), 7.72-7.58 (m, 5H), 7.50-7.42 (m, 4H), 7.18-7.14 (m, 2H), 7.06-7.04 (m, 1H), 6.97-6.94 (m, 3H), 6.86-6.83 (m, 1H), 6.51-6.49 (m, 2H), deuterium substitution rate: 86.9%

29-A-(1)



## Synthesis Example 12


## Synthesis of Compound 30

Compound 30 was prepared in the same manner as in Synthesis Example 1, except that Intermediate 30-A-(1) below was used instead of 9-bromophenanthrene and naphthalene-2-amine was used instead of aniline in the synthesis of Intermediate 3-A, and 9-bromophenanthrene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{44}\text{H}_{27}\text{D}_2\text{NO}$ : cal. 589.24. found 590.27.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.67-8.64 (m, 1H), 8.46-8.44 (m, 1H), 8.12-8.09 (m, 1H), 7.85-7.36 (m, 19H), 7.28-7.08 (m, 5H)

30-A-(1)



## Synthesis Example 13

## Synthesis of Compound 34

Compound 34 was prepared in the same manner as in Synthesis Example 2, except that 1,4-dibromonaphthalene

119

was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, and bis(9,9-dimethyl-9H-fluoren-2-yl) amine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{56}\text{H}_{41}\text{D}_2\text{N}$ : cal. 731.35. found 732.36.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.66 (d, 1H), 8.46 (d, 1H), 8.11 (d, 1H), 7.93-7.76 (m, 5H), 7.69-7.63 (m, 3H), 7.55-7.46 (m, 4H), 7.36-7.24 (m, 3H), 7.14-7.08 (m, 5H), 6.98-6.92 (m, 2H), 6.86-6.84 (m, 2H), 6.61-6.59 (m, 2H), 1.67 (s, 12H), deuterium substitution rate: 82%

## Synthesis Example 14

## Synthesis of Compound 36

Compound 36 was prepared in the same manner as in Synthesis Example 1, except that 9,10-di-tert-butyl-2-bromoanthracene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{50}\text{H}_{43}\text{D}_2\text{N}$ : cal. 661.37. found 662.35.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.58 (d, 1H), 8.43 (d, 1H), 8.20-8.07 (d, 4H), 7.95-7.93 (m, 1H), 7.71-7.68 (m, 2H), 7.61-7.56 (m, 3H), 7.48-7.39 (m, 5H), 7.07-7.03 (m, 3H), 6.97-6.91 (m, 3H), 6.52-6.49 (m, 2H), 1.72 (s, 9H), 1.71 (s, 9H), deuterium substitution rate: 84.6%


## Synthesis Example 15

## Synthesis of Compound 44

Compound 44 was prepared in the same manner as in Synthesis Example 1, except that Intermediate 44-A-(1) below was used instead of 9-bromophenanthrene in the synthesis of Intermediate 3-A, and 9,10-di-tert-butyl-2-bromoanthracene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{47}\text{H}_{42}\text{D}_2\text{N}_2$ : cal. 638.36. found 639.37.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.68 (d, 1H), 8.43 (d, 1H), 8.20-8.07 (d, 4H), 7.60 (dd, 1H), 7.50-7.41 (m, 7H), 7.17 (t, 1H), 7.10-7.04 (m, 3H), 6.99-6.96 (m, 2H), 6.86-6.83 (m, 1H), 6.54-6.48 (m, 3H), 1.72 (s, 9H), 1.71 (s, 9H), deuterium substitution rate: 82.6%



## Synthesis Example 16

## Synthesis of Compound 45

Compound 45 was prepared in the same manner as in Synthesis Example 1, except that bromobenzene was used instead of 9-bromophenanthrene, and Intermediate 5-A-(1) below was used instead of aniline in the synthesis of Intermediate 3-A, and 9,10-di-tert-butyl-2-bromoanthracene was used

120

instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{54}\text{H}_{46}\text{D}_2\text{FN}$ : cal. 731.38. found 732.37.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.43 (t, 1H), 8.19 (d, 1H), 8.14-8.06 (d, 2H), 7.72-7.39 (m, 16H), 7.14-7.04 (m, 3H), 6.97-6.90 (m, 3H), 6.64-6.62 (m, 2H), 1.71 (s, 9H), 1.70 (s, 9H)

## Synthesis Example 17


## Synthesis of Compound 47

Compound 47 was prepared in the same manner as in Synthesis Example 1, except that diphenylamine was used instead of Intermediate 3-A, and Intermediate 47-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{46}\text{H}_{31}\text{D}_2\text{N}$ : cal. 601.27. found 602.24.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.22 (d, 1H), 8.01 (d, 1H), 7.82-7.65 (m, 6H), 7.57-7.28 (m, 11H), 7.08-7.04 (m, 4H), 6.99-6.93 (m, 4H), 6.56-6.53 (m, 4H)

47-2H-(1)



## Synthesis Example 18


## Synthesis of Compound 52

Compound 52 was prepared in the same manner as in Synthesis Example 1, except that diphenylamine was used instead of Intermediate 3-A, and Intermediate 52-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{40}\text{H}_{22}\text{D}_7\text{N}$ : cal. 530.27. found 531.24.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.03 (dd, 2H), 7.93 (dd, 2H), 7.46-7.44 (m, 2H), 7.35-7.31 (m, 2H), 7.08-7.03 (m, 4H), 6.95-6.91 (m, 2H), 6.89-6.86 (m, 2H), 6.76-6.73 (m, 2H), 6.56-6.53 (m, 4H), deuterium substitution rate: 90.7%

52-2H-(1)



## 121


## Synthesis Example 19

## Synthesis of Compound 54

Compound 54 was prepared in the same manner as in Synthesis Example 1, except that diphenylamine was used instead of Intermediate 3-A, and Intermediate 54-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{44}\text{H}_{35}\text{D}_2\text{N}$ : cal. 581.30. found 582.31.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.18 (dd, 2H), 7.96 (dd, 2H), 7.70-7.63 (m, 4H), 7.52-7.49 (m, 2H), 7.44-7.40 (m, 2H), 7.06-7.02 (m, 6H), 6.89-6.83 (m, 4H), 6.56-6.52 (m, 4H), 1.68 (s, 9H), deuterium substitution rate: 83.2%




## Synthesis Example 20

## Synthesis of Compound 56

Compound 56 was prepared in the same manner as in Synthesis Example 1, except that diphenylamine was used instead of Intermediate 3-A, and Intermediate 56-2H-(1) was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{44}\text{H}_{29}\text{D}_2\text{N}$ : cal. 575.25. found 576.27.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.02 (dd, 2H), 7.83 (d, 1H), 7.75-7.64 (m, 5H), 7.46-7.26 (m, 6H), 7.08-6.94 (m, 7H), 6.89-6.86 (m, 2H), 6.76-6.73 (m, 2H), 6.56-6.52 (m, 4H), deuterium substitution rate: 94.6%



## Synthesis Example 21

## Synthesis of Compound 60

Compound 60 was prepared in the same manner as in Synthesis Example 1, except that bis(4-fluorophenyl)amine was used instead of Intermediate 3-A, and 1-bromopyrene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

## 122

$\text{C}_{36}\text{H}_{21}\text{D}_2\text{F}_2\text{N}$ : cal. 509.19. found 510.21.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.37 (d, 1H), 8.17 (d, 2H), 8.11-8.10 (m, 2H), 8.04-7.95 (m, 3H), 7.86 (d, 1H), 7.45-7.41 (m, 2H), 7.22-7.03 (m, 6H), 6.86-6.80 (m, 4H), deuterium substitution rate: 90.1%

## Synthesis Example 22

## Synthesis of Compound 61

Compound 61 was prepared in the same manner as in Synthesis Example 1, except that N-phenylnaphthalen-2-amine was used instead of Intermediate 3-A, and 1-bromopyrene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{40}\text{H}_{25}\text{D}_2\text{N}$ : cal. 523.22. found 524.21.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.37 (d, 1H), 8.17 (d, 2H), 8.04-7.95 (m, 3H), 7.86 (d, 1H), 7.79-7.77 (m, 1H), 7.65 (d, 1H), 7.58-7.38 (m, 6H), 7.13-7.04 (m, 3H), 6.86-6.83 (m, 1H), 6.54-6.52 (m, 2H)

## Synthesis Example 23

## Synthesis of Compound 64

Compound 64 was prepared in the same manner as in Synthesis Example 2, except that 5-bromo-2-fluoropyridine was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, diphenylamine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B, and 1-bromopyrene was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{35}\text{H}_{22}\text{D}_2\text{N}_2$ : cal. 474.20. found 475.21.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.45 (d, 1H), 8.29 (d, 1H), 8.17 (d, 2H), 8.10-7.88 (m, 6H), 7.35-7.33 (m, 1H), 7.18-7.08 (m, 5H), 7.06-7.03 (m, 2H), 6.73-6.71 (m, 4H), deuterium substitution rate: 91.5%

## Synthesis Example 24


## Synthesis of Compound 69

Compound 69 was prepared in the same manner as in Synthesis Example 1, except that 2,7-dibromo-9,9-dimethyl-9H-fluorene was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, diphenylamine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B, and Intermediate 69-2H-(1) below was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{53}\text{H}_{37}\text{D}_2\text{N}$ : cal. 691.32. found 692.29.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.99 (s, 1H), 8.69 (s, 1H), 8.33-8.31 (m, 2H), 8.13-8.05 (m, 4H), 7.75-7.61 (m, 6H), 7.50-7.35 (m, 4H), 7.18-7.03 (m, 5H), 6.85-6.83 (m, 3H), 6.77-6.75 (m, 1H), 6.55-6.52 (m, 4H), 1.68 (s, 6H)

123



Synthesis Example 25

## Synthesis of Compound 71

Compound 71 was prepared in the same manner as in Synthesis Example 1, except that 4-bromobenzonitrile was used instead of 9-bromophenanthrene in the synthesis of Intermediate 3-A and Intermediate 69-2H-(1) was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{45}H_{28}D_2N_2$ : cal. 600.25. found 601.24.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.80 (s, 1H), 8.69 (s, 1H), 8.33-8.31 (m, 2H), 8.13-8.05 (m, 3H), 7.71-7.63 (m, 3H), 7.50-7.35 (m, 7H), 7.18-7.03 (m, 4H), 6.92-6.73 (m, 5H), 6.52-6.50 (m, 2H), deuterium substitution rate: 90.6%

Synthesis Example 26

## Compound 74

Compound 74 was prepared in the same manner as in Synthesis Example 1, except that diphenylamine was used instead of Intermediate 3-A, and 2-bromotriphenylene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{38}H_{25}D_2N$ : cal. 499.22. found 450.21.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.86-8.81 (m, 5H), 8.50-8.49 (m, 1H), 7.70-7.60 (m, 5H), 7.47-7.44 (m, 2H), 7.08-7.04 (m, 4H), 6.89-6.83 (m, 4H), 6.56-6.52 (m, 4H), deuterium substitution rate: 91%

Synthesis Example 27

## Synthesis of Compound 80

Compound 80 was prepared in the same manner as in Synthesis Example 1, except that diphenylamine was used instead of Intermediate 3-A, and Intermediate 80-2H-(1) was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{49}H_{35}D_2N$ : cal. 641.30. found 642.31.


<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.64 (d, 1H), 8.42 (d, 1H), 7.89-7.86 (m, 2H), 7.79-7.77 (m, 1H), 7.68-7.63 (m, 6H), 7.54-7.34 (m, 5H), 7.15-7.03 (m, 5H), 6.89-6.83 (m, 4H), 6.56-6.54 (m, 4H), 1.69 (s, 6H)

124

62-2H-(1)

80-2H-(1)

5



10

Synthesis Example 28

15

## Synthesis of Compound 83

Compound 83 was prepared in the same manner as in Synthesis Example 1, except that diphenylamine was used instead of Intermediate 3-A, and Intermediate 83-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.


$C_{42}H_{27}D_2N$ : cal. 549.24. found 550.21.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.16-7.98 (m, 7H), 7.85 (d, 1H), 7.73-7.70 (m, 2H), 7.63-7.54 (m, 3H), 7.44-7.40 (m, 2H), 7.08-7.04 (m, 4H), 6.99-6.93 (m, 4H), 6.56-6.54 (m, 4H), deuterium substitution rate: 90.8%

25

30

35



83-2H-(1)

Synthesis Example 29

## Synthesis of Compound 86

Compound 86 was prepared in the same manner as in Synthesis Example 1, except that N-phenylnaphthalen-2-amine was used instead of Intermediate 3-A, and Intermediate 86-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.


$C_{52}H_{45}D_2N$ : cal. 687.38. found 688.37.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.22-8.21 (m, 1H), 8.13-8.07 (m, 2H), 7.78-7.73 (m, 3H), 7.66-7.39 (m, 13H), 7.13-7.05 (m, 3H), 6.99-6.95 (m, 2H), 6.86-6.83 (m, 1H), 6.64-6.62 (m, 2H), 1.74 (s, 9H), 1.71 (s, 9H)

55

60

65



86-2H-(1)

## 125


## Synthesis Example 30

## Synthesis of Compound 88

Compound 88 was prepared in the same manner as in Synthesis Example 1, except that diphenylamine was used instead of Intermediate 3-A, and Intermediate 88-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{45}\text{H}_{33}\text{D}_2\text{N}$ : cal. 591.28. found 592.31.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.06-8.04 (m, 1H), 7.93-7.86 (m, 3H), 7.79-7.75 (m, 2H), 7.65-7.43 (m, 8H), 7.36-7.34 (m, 1H), 7.08-7.03 (m, 4H), 6.99-6.93 (m, 4H), 6.66-6.64 (m, 4H), 1.69 (s, 6H), deuterium substitution rate: 82%




## Synthesis Example 31

## Synthesis of Compound 91

Compound 91 was prepared in the same manner as in Synthesis Example 2, except that 2,7-dibromophenanthrene was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, diphenylamine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B, and Intermediate 91-2H-(1) below was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{44}\text{H}_{29}\text{D}_2\text{N}$ : cal. 575.25. found 576.24.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.48-8.44 (m, 2H), 8.15-8.14 (m, 1H), 7.97-7.96 (m, 1H), 7.91-7.81 (m, 5H), 7.75-7.49 (m, 9H), 7.08-7.03 (m, 4H), 6.99-6.97 (m, 1H), 6.86-6.83 (m, 2H), 6.68-6.66 (m, 4H), deuterium substitution rate: 92.5%



## Synthesis Example 32

## Synthesis of Compound 93

Compound 93 was prepared in the same manner as in Synthesis Example 2, except that 2,7-dibromo-9,9-dimethyl-9H-fluorene was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, diphenylamine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B, and Intermediate 88-2H-(1)

## 126

was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{54}\text{H}_{41}\text{D}_2\text{N}$ : cal. 707.35. found 708.36.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.05-8.04 (m, 1H), 7.96-7.93 (m, 3H), 7.79-7.73 (m, 3H), 7.68-7.49 (m, 8H), 7.36-7.33 (m, 2H), 7.08-7.03 (m, 4H), 6.97-6.93 (m, 3H), 6.86-6.83 (m, 1H), 6.45-6.43 (m, 4H), 1.69 (s, 6H), 1.67 (s, 6H)

## Synthesis Example 33

## Synthesis of Compound 101

Compound 101 was prepared in the same manner as in Synthesis Example 1, except that diphenylamine was used instead of Intermediate 3-A, and 2-bromo-9,9-dimethyl-9H-fluorene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{35}\text{H}_{27}\text{D}_2\text{N}$ : cal. 465.24. found 466.25.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 7.85 (dd, 1H), 7.74 (dd, 1H), 7.46-7.44 (m, 2H), 7.38-7.33 (m, 2H), 7.28-7.27 (m, 1H), 7.20-7.04 (m, 6H), 6.99-6.93 (m, 4H), 6.66-6.63 (m, 4H), 1.67 (s, 6H), deuterium substitution rate: 85.6%

## Synthesis Example 34

## Synthesis of Compound 106

Compound 106 was prepared in the same manner as in Synthesis Example 1, except that 4-bromobenzonitrile was used instead of 9-bromophenanthrene and naphthalene-2-amine was used instead of aniline in the synthesis of Intermediate 3-A, and 2-bromo-9,9-dimethyl-9H-fluorene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{43}\text{H}_{28}\text{D}_2\text{N}_2$ : cal. 540.25. found 541.26.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 7.87-7.83 (m, 1H), 7.78-7.76 (m, 1H), 7.72 (d, 1H), 7.65 (d, 1H), 7.58-7.53 (m, 3H), 7.48-7.45 (m, 2H), 7.41-7.27 (m, 6H), 7.20-7.18 (m, 1H), 7.13-7.09 (m, 1H), 7.02-6.99 (m, 1H), 6.93-6.88 (m, 4H), 1.68 (s, 6H), deuterium substitution rate: 87.4%

## Synthesis Example 35

## Synthesis of Compound 110

Compound 110 was prepared in the same manner as in Synthesis Example 1, except that 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine was used instead of Intermediate, and 2-bromo-9,9-dimethyl-9H-fluorene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{44}\text{H}_{35}\text{D}_2\text{N}$ : cal. 581.30. found 582.31.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 7.86-7.84 (m, 1H), 7.78-7.76 (m, 1H), 7.72 (d, 1H), 7.72 (d, 1H), 7.55 (d, 1H), 7.48-7.45 (m, 2H), 7.36-7.31 (m, 3H), 7.28-7.27 (m, 1H), 7.14-7.04 (m, 5H), 6.93-6.83 (m, 4H), 6.69-6.67 (m, 1H), 6.53-6.51 (m, 2H), 1.68 (s, 6H), 1.65 (s, 6H), deuterium substitution rate: 82.1%

## Synthesis Example 36

## Synthesis of Compound 112

Compound 112 was prepared in the same manner as in Synthesis Example 1, except that 3-iodo-9-phenyl-9H-carba-

127

zole was used instead of 9-bromophenanthrene in the synthesis of Intermediate 3-A, and 2-bromo-9,9-dimethyl-9H-fluorene was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{47}H_{34}D_2N_2$ : cal. 630.30. found 631.32.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.22-8.20 (m, 1H), 7.86-7.84 (m, 1H), 7.73-7.71 (m, 1H), 7.54-7.04 (m, 19H), 6.87-6.82 (m, 3H), 6.76-6.73 (m, 1H), 6.61-6.59 (m, 2H), 1.68 (s, 6H)


## Synthesis Example 37

## Synthesis of Compound 118

Compound 118 was prepared in the same manner as in Synthesis Example 1, except that N-phenylnaphthalen-2-amine was used instead of Intermediate 3-A, and Intermediate 118-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{49}H_{31}D_2N$ : cal. 637.27. found 638.24.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.11-8.10 (m, 1H), 7.96 (d, 1H), 7.92-7.90 (m, 1H), 7.78-7.76 (m, 1H), 7.73-7.71 (m, 1H), 7.65 (d, 1H), 7.58-7.34 (m, 10H), 7.24-7.04 (m, 6H), 6.93-6.86 (m, 6H), 6.80-6.78 (m, 1H), 6.54-6.52 (m, 2H)



118-2H-(1) 30

## Synthesis Example 38

## Synthesis of Compound 121

Compound 121 was prepared in the same manner as in Synthesis Example 1, except that 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine was used instead of Intermediate 3-A, and 3-iodo-9-phenyl-9H-carbazole was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{47}H_{34}D_2N_2$ : cal. 449.21. found 450.22.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.28-8.24 (m, 1H), 8.02-8.00 (m, 1H), 7.77 (dd, 1H), 7.55 (d, 1H), 7.51-7.26 (m, 13H), 7.14-7.05 (m, 4H), 6.93-6.83 (m, 4H), 6.79-6.77 (m, 1H), 6.63-6.61 (m, 2H), 1.68 (s, 6H)

## Synthesis Example 39

## Synthesis of Compound 125

Compound 125 was prepared in the same manner as in Synthesis Example 1, except that 4-bromobenzonitrile was used instead of 9-bromophenanthrene and naphthalene-2-amine was used instead of aniline in the synthesis of Intermediate 3-A, and 3-iodo-9-phenyl-9H-carbazole was used instead of

128

2-bromonaphthalene during synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{43}H_{27}D_2N_3$ : cal. 589.24. found 590.24.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.26-8.24 (m, 1H), 8.02-8.00 (m, 1H), 7.78-7.76 (m, 1H), 7.65 (d, 1H), 7.57-7.52 (m, 3H), 7.50-7.25 (m, 15H), 7.12-7.10 (m, 1H), 6.93-6.88 (m, 4H), deuterium substitution rate: 88.1%

## Synthesis Example 40

## Synthesis of Compound 127

Compound 127 was prepared in the same manner as in Synthesis Example 1, except that 3-iodo-9-phenyl-9H-carbazole was used instead of 9-bromophenanthrene in the synthesis of Intermediate 3-A, and 3-iodo-9-phenyl-9H-carbazole was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{50}H_{33}D_2N_3$ : cal. 679.29. found 680.28.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.26-8.20 (m, 2H), 8.02-8.01 (m, 1H), 7.54-7.23 (m, 22H), 7.08-7.04 (m, 2H), 6.97-6.91 (m, 3H), 6.86-6.83 (m, 1H), 6.61-6.59 (m, 2H), deuterium substitution rate: 90.5%

## Synthesis Example 41

## Synthesis of Compound 128

Compound 128 was prepared in the same manner as in Synthesis Example 1, except that N,9,9-triphenyl-9H-fluoren-2-amine was used instead of Intermediate 3-A, and 3-iodo-9-phenyl-9H-carbazole was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{57}H_{38}D_2N_2$ : cal. 754.33. found 755.34.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.26-8.24 (m, 1H), 8.02-8.00 (m, 1H), 7.85 (dd, 1H), 7.57 (d, 1H), 7.51-7.26 (m, 13H), 7.19-7.04 (m, 13H), 6.91-6.89 (m, 1H), 6.83-6.73 (m, 4H), 6.68-6.67 (m, 1H), 6.53-6.51 (m, 2H)

## Synthesis Example 42


## Synthesis of Compound 133

Compound 133 was prepared in the same manner as in Synthesis Example 1, except that 9,9-dimethyl-N-(naphthalen-3-yl)-9H-fluoren-2-amine was used instead of Intermediate 3-A, and Intermediate 133-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{51}H_{35}D_2FN_2$ : cal. 698.30. found 699.31.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.14-8.12 (m, 1H), 7.93-7.91 (m, 1H), 7.79-7.76 (m, 2H), 7.69 (d, 1H), 7.58-7.54 (m, 4H), 7.49-7.23 (m, 1H), 7.16-7.05 (m, 5H), 6.99-6.91 (m, 3H), 6.79-6.78 (m, 1H), 1.67 (s, 6H)

129




## Synthesis Example 43

## Synthesis of Compound 136

Compound 136 was prepared in the same manner as in Synthesis Example 1, except that 1-bromo-4-fluorobenzene was used instead of 9-bromophenanthrene and naphthalene-2-amine was used instead of aniline in the synthesis of Intermediate 3-A, and Intermediate 136-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{36}\text{H}_{22}\text{D}_2\text{FNS}$ : cal. 523.17. found 524.27.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.29 (d, 1H), 8.10-8.04 (m, 2H), 7.80-7.76 (m, 2H), 7.65 (d, 1H), 7.58-7.54 (m, 4H), 7.48-7.35 (m, 5H), 7.25-7.20 (m, 3H), 7.04-7.01 (m, 2H), 6.84-6.81 (m, 2H), deuterium substitution rate: 91.3%



## Synthesis Example 44

## Synthesis of Compound 138

Compound 138 was prepared in the same manner as in Synthesis Example 1, except that Intermediate 29-A-(1) above was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{40}\text{H}_{25}\text{D}_2\text{NS}$ : cal. 555.19. found 556.17.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.58 (d, 1H), 8.27-8.26 (m, 1H), 8.21-8.11 (m, 2H), 7.95-7.92 (m, 1H), 7.86 (d, 1H), 7.79 (d, 1H), 7.72-7.54 (m, 6H), 7.46-7.39 (m, 4H), 7.07-7.03 (m, 3H), 6.97-6.91 (m, 3H), 6.52-6.49 (m, 2H), deuterium substitution rate: 90%

## Synthesis Example 45

## Synthesis of Compound 139

Compound 139 was prepared in the same manner as in Synthesis Example 1, except that Intermediate 118-2H-(1) above was used instead of 9-bromophenanthrene and naphthalene-2-amine was used instead of aniline in the synthesis of Intermediate 3-A, and Intermediate 29-A-(1) above was used

130

instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{51}\text{H}_{31}\text{D}_2\text{NS}$ : cal. 693.24. found 694.21.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.27-8.26 (m, 1H), 8.12 (d, 1H), 7.93-7.85 (m, 4H), 7.79 (d, 1H), 7.64-7.41 (m, 9H), 7.21-7.15 (m, 3H), 7.09-7.04 (m, 2H), 6.93-6.73 (m, 7H), 6.67-6.66 (m, 1H), 6.53-6.51 (m, 2H)

Synthesis Example 46

## Synthesis of Compound 142

Compound 142 was prepared in the same manner as in Synthesis Example 1, except that bromobenzene was used instead of 9-bromophenanthrene, and Intermediate 5-A-(1) above was used instead of aniline in the synthesis of Intermediate 3-A, and Intermediate 29-A-(1) above was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{44}\text{H}_{28}\text{D}_2\text{FNS}$ : cal. 625.22. found 626.23.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.27-8.26 (m, 1H), 8.12 (d, 1H), 7.86 (d, 1H), 7.79 (d, 1H), 7.72-7.70 (m, 2H), 7.66-7.40 (m, 14H), 7.13-7.03 (m, 3H), 6.98-6.90 (m, 3H), 6.54-6.52 (m, 2H), deuterium substitution rate: 92.8%


## Synthesis Example 47

## Synthesis of Compound 147

Compound 147 was prepared in the same manner as in Synthesis Example 1, except that 9,9-dimethyl-N-(naphthalen-3-yl)-9H-fluoren-2-amine was used instead of Intermediate 3-A, and Intermediate 147-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{45}\text{H}_{31}\text{D}_2\text{NO}$ : cal. 605.26. found 606.24.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 7.96-7.94 (m, 1H), 7.78-7.68 (m, 5H), 7.57-7.30 (m, 12H), 7.16-7.10 (m, 4H), 6.99-6.97 (m, 2H), 6.93-6.91 (m, 1H), 6.69-6.68 (m, 1H), 1.67 (s, 6H), deuterium substitution rate: 93%



## Synthesis Example 48

## Synthesis of Compound 150

Compound 150 was prepared in the same manner as in Synthesis Example 1, except that N,9,9-triphenyl-9H-fluoren-2-amine was used instead of Intermediate 3-A, and Intermediate 30-A-(1) above was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{51}\text{H}_{33}\text{D}_2\text{NO}$ : cal. 679.28. found 680.29.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 7.99-7.97 (m, 1H), 7.93-7.92 (m, 1H), 7.86-7.84 (m, 1H), 7.72 (d, 1H), 7.58-7.35

## 131

(m, 8H), 7.19-7.04 (m, 13H), 6.90 (dd, 1H), 6.83-6.73 (m, 4H), 6.67-6.66 (m, 1H), 6.53-6.51 (m, 2H)

## Synthesis Example 49

## Synthesis of Compound 157

Compound 157 was prepared in the same manner as in Synthesis Example 2, except that 2,7-dibromo-9,9-dimethyl-9H-fluorene was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, N-phenylnaphthalen-1-amine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B, and 2-bromo-9,9-dimethyl-9H-fluorene was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{48}H_{37}D_2N$ : cal. 631.32. found 632.33.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.13-8.10 (m, 1H), 7.87-7.84 (m, 2H), 7.73-7.64 (m, 5H), 7.49-7.43 (m, 3H), 7.36-7.02 (m, 8H), 6.93 (d, 1H), 6.85-6.78 (m, 2H), 6.55-6.54 (m, 1H), 6.46-6.44 (m, 2H), 1.69 (s, 6H), 1.67 (s, 6H)

## Synthesis Example 50

## Synthesis of Compound 161

Compound 161 was prepared in the same manner as in Synthesis Example 2, except that diphenylamine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B and 3-iodo-9-phenyl-9H-carbazole was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{42}H_{28}D_2N_2$ : cal. 564.25. found 565.24.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.26-8.24 (m, 1H), 8.01-8.00 (m, 1H), 7.88-7.87 (m, 1H), 7.76-7.67 (m, 3H), 7.63-7.62 (m, 1H), 7.52-7.26 (m, 10H), 7.09-7.04 (m, 4H), 6.97-6.95 (m, 1H), 6.86-6.83 (m, 2H), 6.58-6.56 (m, 4H), deuterium substitution rate: 90%

## Synthesis Example 51

## Synthesis of Compound 163

Compound 163 was prepared in the same manner as in Synthesis Example 2, except that 2,7-dibromo-9,9-dimethyl-9H-fluorene was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, N-phenylnaphthalen-2-amine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B, and 3-iodo-9-phenyl-9H-carbazole was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{51}H_{36}D_2N_2$ : cal. 680.31. found 681.34.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.26-8.24 (m, 1H), 8.01-8.00 (m, 1H), 7.78-7.64 (m, 5H), 7.58-7.26 (m, 15H), 7.10-7.05 (m, 3H), 6.91-6.83 (m, 2H), 6.74-6.73 (m, 1H), 6.52-6.50 (m, 2H), 1.68 (s, 6H)

## Synthesis Example 52

## Synthesis of Compound 164

Compound 164 was prepared in the same manner as in Synthesis Example 2, except that 5-bromo-2-iodopyridine


## 132

was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, diphenylamine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B, and Intermediate 164-2H-(1) below was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{49}H_{29}D_2N_3$ : cal. 591.26. found 592.27.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.14-8.12 (m, 1H), 8.08-8.07 (m, 1H), 7.92-7.89 (m, 2H), 7.82-7.80 (m, 2H), 7.70-7.67 (m, 2H), 7.54-7.48 (m, 4H), 7.39-7.26 (m, 4H), 7.21-7.19 (m, 2H), 7.14-7.08 (m, 5H), 6.96-6.93 (m, 2H), 6.73-6.71 (m, 4H), deuterium substitution rate: 93.5%

164-2H-(1)



## Synthesis Example 53


## Synthesis of Compound 167

Compound 167 was prepared in the same manner as in Synthesis Example 1, except that diphenylamine was used instead of Intermediate 3-A, and Intermediate 167-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

$C_{42}H_{27}D_2NS$ : cal. 581.21. found 582.24.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.36 (d, 1H), 8.16-8.06 (m, 2H), 7.98-7.97 (m, 1H), 7.89-7.78 (m, 3H), 7.63-7.58 (m, 2H), 7.46-7.35 (m, 4H), 7.12-6.94 (m, 6H), 6.85-6.73 (m, 4H), 6.66-6.63 (m, 4H)

167-2H-(1)



## Synthesis Example 54

## Synthesis of Compound 169

Compound 169 was prepared in the same manner as in Synthesis Example 2, except that 2,7-dibromo-9,9-dimethyl-9H-fluorene was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, N-phenylnaphthalen-2-amine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B, and Intermediate 30-A-(1) was used instead of 9-bromophenanthrene in

## 133

the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{45}\text{H}_{31}\text{D}_2\text{NO}$ : cal. 605.26. found 606.27.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 7.99-7.97 (m, 1H), 7.93-7.92 (m, 1H), 7.78-7.64 (m, 6H), 7.58-7.51 (m, 6H), 7.42-7.34 (m, 3H), 7.10-7.06 (m, 3H), 6.91-6.87 (m, 2H), 6.74-6.73 (m, 1H), 6.52-6.50 (m, 2H), 1.68 (s, 6H)

## Synthesis Example 55


## Synthesis of Compound 176

Compound 176 was prepared in the same manner as in Synthesis Example 1, except that N-phenyl-naphthalen-2-amine was used instead of Intermediate 3-A, and Intermediate 176-2H-(1) below was used instead of 2-bromonaphthalene in the synthesis of Compound 3-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{57}\text{H}_{40}\text{D}_2\text{N}_2$ : cal. 756.34. found 757.34.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 8.24-8.22 (m, 1H), 8.00-7.99 (m, 1H), 7.90-7.85 (m, 2H), 7.80-7.76 (m, 2H), 7.66-7.64 (m, 3H), 7.59-7.04 (m, 20H), 6.99-6.96 (m, 2H), 6.86-6.83 (m, 1H), 6.54-6.52 (m, 2H), 1.67 (s, 6H)

176-2H-(1)



## Synthesis Example 56


## Synthesis of Compound 179

Compound 179 was prepared in the same manner as in Synthesis Example 2, except that diphenylamine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine during synthesis of Intermediate 33-B, and Intermediate 179-2H-(1) below was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{45}\text{H}_{33}\text{D}_2\text{N}$ : cal. 591.28. found 592.29.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 7.83-7.79 (m, 2H), 7.75-7.67 (m, 6H), 7.63-7.62 (m, 1H), 7.57-7.55 (m, 1H), 7.50-7.46 (m, 2H), 7.35-7.31 (m, 1H), 7.28-7.27 (m, 1H), 7.15-7.04 (m, 6H), 6.96 (dd, 1H), 6.86-6.83 (m, 2H), 6.68-6.66 (m, 4H), 1.64 (s, 6H), deuterium substitution rate: 82%

179-2H-(1)



## 134

## Synthesis Example 57


## Synthesis of Compound 180

Compound 180 was prepared in the same manner as in Synthesis Example 2, except that 2,7-dibromo-9,9-dimethyl-9H-fluorene was used instead of 2,6-dibromonaphthalene in the synthesis of Intermediate 33-A, N-phenyl-naphthalen-2-amine was used instead of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine in the synthesis of Intermediate 33-B, Intermediate 180-2H-(1) below was used instead of 9-bromophenanthrene in the synthesis of Compound 33-2H. The obtained compound was confirmed by MS/FAB and  $^1\text{H}$  NMR.

$\text{C}_{64}\text{H}_{45}\text{D}_2\text{N}$ : cal. 831.38. found 832.40.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  (ppm) 7.90-7.88 (m, 1H), 7.85-7.64 (m, 8H), 7.58-7.33 (m, 10H), 7.22-7.05 (m, 14H), 6.94-6.92 (m, 1H), 6.81-6.73 (m, 2H), 6.73-6.72 (m, 1H), 6.53-6.50 (m, 2H), 1.67 (s, 6H)


180-2H-(1)



## Comparative Synthesis Example A


Compound A was synthesized according to Reaction Scheme A below:

## Reaction Scheme A



135

-continued



## Synthesis of Intermediate I-5

Intermediate I-5 was prepared in the same manner as in the synthesis of Intermediate 3-A of Synthesis Example 1, except that bromobenzene-d<sub>5</sub> was used instead of 9-bromophenanthrene, and 9,9-dimethyl-9H-fluorene-2-amine was used instead of aniline in the synthesis of Intermediate 3-A

## Synthesis of Intermediate I-6

Intermediate I-6 was prepared in the same manner as in the synthesis of Intermediate 33-B of Synthesis Example 2, except that Intermediate I-5 was used instead of 9,9-dimethyl-N-phenyl-9H-fluorene-2-amine.

## Synthesis of Compound A

Compound A was prepared in the same manner as in the synthesis of Compound 33-2H of Synthesis Example 2, except that Intermediate I-6 was used instead of Intermediate 33-B. The obtained compound was confirmed by MS/FAB and <sup>1</sup>H NMR.

C<sub>47</sub>H<sub>30</sub>D<sub>5</sub>N: cal. 618.31. found 619.32.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ (ppm) 8.66 (d, 1H), 8.46 (d, 1H), 8.15-8.06 (m, 3H), 7.92 (s, 1H), 7.83-7.56 (m, 9H), 7.48 (dt, 1H), 7.36-7.27 (m, 3H), 7.14-7.08 (m, 2H), 7.02 (dt, 1H), 6.97-6.95 (m, 1H), 6.89 (s, 1H), 1.66 (s, 6H)

## Example 1

As an anode, a 15 Ω/cm<sup>2</sup> (1200 Å) Corning ITO glass substrate was cut to a size of 50 mm×50 mm×0.7 mm, washed with ultrasonic waves in isopropyl alcohol and pure water for 5 minutes each, and then cleaned with UV and ozone for 30 minutes. The ITO glass substrate was mounted on a vacuum depositor.

2-TNATA was deposited on the ITO glass substrate to form a HIL having a thickness of 600 Å and 4,4'-bis[N-(1-naphthyl)-N-phenyl amino]biphenyl (NPB) was then deposited on the HIL to form a HTL having a thickness of 300 Å.

Then, 9,10-di(naphthalene-2-yl)anthracene (ADN) and Compound 3 were co-deposited on the HTL at a weight ratio of 98:2 to form an EML having a thickness of 300 Å.

Then, Alq<sub>3</sub> was deposited on the EML to form an ETL having a thickness of 300 Å, LiF was deposited on the ETL to form an EIL having a thickness of 10 Å, and Al was deposited on the EIL to form a second electrode (cathode) having a thickness of 3,000 Å, thereby completing the manufacture of an OLED.

136

Example 2

An OLED was manufactured in the same manner as in Example 1, except that Compound 19 was used instead of Compound 3 in the formation of the EML.

Example 3

An OLED was manufactured in the same manner as in Example 1, except that Compound 33 was used instead of Compound 3 in the formation of the EML.

Example 4

An OLED was manufactured in the same manner as in Example 1, except that Compound 45 was used instead of Compound 3 in the formation of the EML.

Example 5

An OLED was manufactured in the same manner as in Example 1, except that Compound 47 was used instead of Compound 3 in the formation of the EML.

Example 6

An OLED was manufactured in the same manner as in Example 1, except that Compound 54 was used instead of Compound 3 in the formation of the EML.

Example 7

An OLED was manufactured in the same manner as in Example 1, except that Compound 69 was used instead of Compound 3 in the formation of the EML.

Example 8

An OLED was manufactured in the same manner as in Example 1, except that Compound 74 was used instead of Compound 3 in the formation of the EML.

Example 9

An OLED was manufactured in the same manner as in Example 1, except that Compound 80 was used instead of Compound 3 in the formation of the EML.

Example 10

An OLED was manufactured in the same manner as in Example 1, except that Compound 92 was used instead of Compound 3 in the formation of the EML.

Example 11

An OLED was manufactured in the same manner as in Example 1, except that Compound 106 was used instead of Compound 3 in the formation of the EML.

Example 12

An OLED was manufactured in the same manner as in Example 1, except that Compound 118 was used instead of Compound 3 in the formation of the EML.

## 137

## Example 13

An OLED was manufactured in the same manner as in Example 1, except that Compound 128 was used instead of Compound 3 in the formation of the EML.

## Example 14

An OLED was manufactured in the same manner as in Example 1, except that Compound 136 was used instead of Compound 3 in the formation of the EML.

## Example 15

An OLED was manufactured in the same manner as in Example 1, except that Compound 161 was used instead of Compound 3 in the formation of the EML.

## Example 16

An OLED was manufactured in the same manner as in Example 1, except that Compound 169 was used instead of Compound 3 in the formation of the EML.

## Example 17

An OLED was manufactured in the same manner as in Example 1, except that Compound 180 was used instead of Compound 3 in the formation of the EML.

## 138

## Example 18

An OLED was manufactured in the same manner as in Example 1, except that Compound 3 was used instead of NPB in the formation of the HTL and Compound DPAVB<sub>i</sub> was used instead of Compound 3 in the formation of the EML.

## Example 19


An OLED was manufactured in the same manner as in Example 1, except that Compound 136 was used instead of NPB in the formation of the HTL and Compound DPAVB<sub>i</sub> was used instead of Compound 3 in the formation of the EML.

## Example 20

An OLED was manufactured in the same manner as in Example 1, except that Compound 161 was used instead of NPB in the formation of the HTL and Compound DPAVB<sub>i</sub> was used instead of Compound 3 in the formation of the EML.

## Comparative Example 1

An OLED was manufactured in the same manner as in Example 1, except that 1,4-bis-(2,2-diphenylvinyl)biphenyl (DPAVB<sub>i</sub>) was used instead of Compound 3 in the formation of the EML.

DPAVB<sub>i</sub>

45

## Comparative Example 2

An OLED was manufactured in the same manner as in Example 1, except that Compound A was used instead of Compound 3 in the formation of the EML.

50

## Evaluation Example 1

Driving voltage, current density, brightness, emission color, efficiency, and half lifetime (@100 mA/cm<sup>2</sup>) of each of the OLEDs of Examples 1 through 20 and Comparative Examples 1 and 2 were evaluated using PR650 Spectroscan Source Measurement Unit (available from PhotoResearch), and the results are shown in Table 1 below.

TABLE 1

	Hole transport layer	Dopant of Emission layer	Driving voltage (V)	Current density (mA/cm <sup>2</sup> )	Brightness (cd/m <sup>2</sup> )	efficiency (cd/A)	Emission color	Half lifetime (hr)
Example 1	NPB	Compound 3	6.39	50	3,075	6.15	Blue	259
Example 2	NPB	Compound 19	6.41	50	3,045	6.09	Blue	263
Example 3	NPB	Compound 33	6.42	50	3,085	6.17	Blue	269
Example 4	NPB	Compound 45	6.36	50	3,205	6.41	Blue	287

TABLE 1-continued

	Hole transport layer	Dopant of Emission layer	Driving voltage (V)	Current density (mA/cm <sup>2</sup> )	Brightness (cd/m <sup>2</sup> )	efficiency (cd/A)	Emission color	Half lifetime (hr)
Example 5	NPB	Compound 47	6.38	50	3,260	6.52	Blue	322
Example 6	NPB	Compound 54	6.39	50	3,120	6.24	Blue	271
Example 7	NPB	Compound 69	6.37	50	3,055	6.11	Blue	254
Example 8	NPB	Compound 74	6.36	50	2,985	5.97	Blue	322
Example 9	NPB	Compound 80	6.40	50	3,235	6.47	Blue	307
Example 10	NPB	Compound 92	6.38	50	3,035	6.07	Blue	297
Example 11	NPB	Compound 106	6.37	50	3,105	6.21	Blue	264
Example 12	NPB	Compound 118	6.41	50	3,290	6.58	Blue	326
Example 13	NPB	Compound 128	6.28	50	3,160	6.32	Blue	302
Example 14	NPB	Compound 136	6.31	50	3,165	6.33	Blue	273
Example 15	NPB	Compound 161	6.29	50	3,140	6.28	Blue	267
Example 16	NPB	Compound 169	6.28	50	3,305	6.61	Blue	279
Example 17	NPB	Compound 180	6.39	50	3,330	6.66	Blue	286
Example 18	Compound 3	DPVABi	5.72	50	2,680	5.36	Blue	209
Example 19	Compound 136	DPVABi	5.53	50	2,710	5.42	Blue	219
Example 20	Compound 161	DPVABi	5.54	50	2,880	5.76	Blue	226
Comparative Example 1	NPB	DPVABi	7.35	50	2,065	4.13	Blue	145
Comparative Example 2	NPB	Compound A	6.43	50	3,050	6.10	Blue	152

From the results shown in Table 1, it is confirmed that the OLEDs of Examples 1 through 20 each generally have excellent driving voltage, higher luminance, higher efficiency, higher color purity, and longer lifetime, as compared to the OLEDs of Comparative Examples 1 and 2.

In detail, it may be confirmed that the OLED of Example 3 using Compound 33 as a dopant of an EML has long lifetime compared to the OLED of Example 2 using Compound A as a dopant of an EML. The Compound A is different from Compound 33 in that a phenyl group bonded to nitrogen instead of a styryl group is substituted with deuterium. Thus, it may be confirmed that the lifetime properties of an OLED is increased by substituting carbon of the styryl group with deuterium styryl group.






It may be confirmed that the OLEDs of Examples 18 through 20 including Compounds 3, 136, and 161 as a hole transport material each have low driving voltage while having long lifetime compared to the OLED of Comparative Example 1.

As described above, according to the one or more embodiments of the present invention, a styryl-based compound may exhibit high heat resistance and thus an OLED including the same may have low driving voltage, high brightness, high efficiency, and long lifetime.

While the present invention has been particularly shown and described with reference to exemplary embodiments thereof, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope of the present invention as defined by the following claims.

What is claimed is:


1. A styryl-based compound represented by any one of Formulae 1A through 1E below:




141

-continued


Formula 1F




Formula 1G



Formula 1H



Formula 1I



wherein  $Z_1$  through  $Z_3$  are each independently selected from the group consisting of hydrogen; deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a  $C_1$ - $C_{10}$  alkyl group; a  $C_1$ - $C_{10}$  alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolynyl group; an imidazopyridinyl group; an imidazopyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; an indolyl group; and a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazolynyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a

142

salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{10}$  alkyl group, and a  $C_1$ - $C_{10}$  alkoxy group; two or more neighboring substituents from among  $Z_1$  through  $Z_3$  are optionally fused with each other or are optionally connected by a single bond;

$T_1$  is O or S;

$p$  is an integer of 1 to 7; and

$q$  is an integer of 1 to 4;

$Ar_1$  and  $Ar_2$  are each independently a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, or a substituted or unsubstituted  $C_2$ - $C_{60}$  heteroaryl group;

$L_1$  and  $L_2$  are each independently a substituted or unsubstituted  $C_6$ - $C_{60}$  arylene group, or a substituted or unsubstituted  $C_2$ - $C_{60}$  heteroarylene group; and

$a$  and  $b$  are each independently an integer of 0 to 5.


2. The styryl-based compound of claim 1, wherein  $Ar_1$  and  $Ar_2$  are each independently a substituted or unsubstituted phenyl group, a substituted or unsubstituted pentalenyl group, a substituted or unsubstituted indenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted azulenyl group, a substituted or unsubstituted heptalenyl group, a substituted, or unsubstituted indacenyl group, a substituted or unsubstituted acenaphthyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted spiro-fluorenyl group, a substituted or unsubstituted phenalenyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluoranthenyl group, a substituted or unsubstituted triphenylenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted naphthacenyl group, a substituted or unsubstituted picenyl group, a substituted or unsubstituted perylenyl group, a substituted or unsubstituted pentaphenyl group, a substituted or unsubstituted hexacenyl group, a substituted or unsubstituted pyrrolyl group, a substituted or unsubstituted imidazolyl group, a substituted or unsubstituted pyrazolyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted pyridazinyl group, a substituted or unsubstituted isoindolyl group, a substituted or unsubstituted indolyl group, a substituted or unsubstituted indazolyl group, a substituted or unsubstituted purinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted benzoquinolinyl group, a substituted or unsubstituted phthalazinyl group, a substituted or unsubstituted naphthyridinyl group, a substituted or unsubstituted quinoxalinyl group, a substituted or unsubstituted quinazolinyl group, a substituted or unsubstituted cinnolinyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted phenanthridinyl group, a substituted or unsubstituted acridinyl group, a substituted or unsubstituted phenanthrolinyl group, a substituted or unsubstituted phenazinyl group, a substituted or unsubstituted benzoxazolyl group, a substituted or unsubstituted benzimidazolyl group, a substituted or unsubstituted furanyl group, a substituted or unsubstituted benzofuranyl group, a substituted or unsubstituted thiophenyl group, a substituted or unsubstituted benzothiofenyl group, a substituted or unsubstituted thiazolyl group, a substituted or unsubstituted isothiazolyl group, a substituted or unsubstituted benzothiazolyl group, a substituted or unsubstituted isoxazolyl group, a substituted or unsubstituted oxazolyl group, a substituted or unsubstituted triazolyl group, a substituted or unsubstituted tetrazolyl group, a substituted or unsubstituted oxadiazolyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted benzoxazolyl group, a substituted or unsubsti-

143


tuted dibenzofuranyl group, a substituted or unsubstituted dibenzothiophenyl group, or a substituted or unsubstituted benzocarbazolyl group.

3. The styryl-based compound of claim 1, wherein Ar<sub>1</sub> and Ar<sub>2</sub> are each independently a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted dibenzothiophenyl group, or a substituted or unsubstituted benzocarbazolyl group.


4. The styryl-based compound of claim 1, wherein Ar<sub>1</sub> and Ar<sub>2</sub> are each independently represented by any one of Formulae 2A through 2J below:




Formula 2A




Formula 2B




Formula 2C




Formula 2D



Formula 2E



Formula 2F




Formula 2G

144


-continued

Formula 2H



5


Formula 2I



10

15

Formula 2J



20

25

wherein Z<sub>11</sub> through Z<sub>15</sub> are each independently selected from the group consisting of hydrogen; deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a C<sub>1</sub>-C<sub>10</sub> alkyl group; a C<sub>1</sub>-C<sub>10</sub> alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolynyl group; an imidazopyridinyl group; an imidazopyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; an indolyl group; and a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazolynyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>10</sub> alkyl group, and a C<sub>1</sub>-C<sub>10</sub> alkoxy group;

two or more neighboring substituents from among Z<sub>11</sub> through Z<sub>15</sub> are optionally fused with each other or are optionally connected by a single bond;

55

T<sub>11</sub> is O or S;  
r is an integer of 1 to 7; and  
s is an integer of 1 to 5.

5. The styryl-based compound of claim 1, wherein L<sub>1</sub> and L<sub>2</sub> are each independently a substituted or unsubstituted phenylenyl group, a substituted or unsubstituted pentalenylene group, a substituted or unsubstituted indenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted azulenylenylene group, a substituted or unsubstituted heptalenylene group, a substituted or unsubstituted indacenylene group, a substituted or unsubstituted acenaphthylenylene group, a substituted or unsubstituted fluorenylenylene group, a substituted or unsubstituted phenalenylene group, a

60


65

145


substituted or unsubstituted phenanthrenylene group, a substituted or unsubstituted anthrylene group, a substituted or unsubstituted fluoranthrenylene group, a substituted or unsubstituted triphenylenylene group, a substituted or unsubstituted pyrenylene group, a substituted or unsubstituted chrysenylene group, a substituted or unsubstituted naphthacenylenylene group, a substituted or unsubstituted picenylenylene group, a substituted or unsubstituted perylenylene group, a substituted or unsubstituted pentaphenylenylene group, a substituted or unsubstituted hexacenylenylene group, a substituted or unsubstituted pyrrolylene group, a substituted or unsubstituted pyrazolylenylene group, a substituted or unsubstituted imidazolinylenylene group, a substituted or unsubstituted imidazopyridinylenylene group, a substituted or unsubstituted imidazopyrimidinylenylene group, a substituted or unsubstituted pyridinylenylene group, a substituted or unsubstituted pyrazinylenylene group, a substituted or unsubstituted pyrimidinylenylene group, a substituted or unsubstituted indolylenylene group, a substituted or unsubstituted purinylenylene group, a substituted or unsubstituted quinolinylenylene group, a substituted or unsubstituted phthalazinylenylene group, a substituted or unsubstituted indolizinylenylene group, a substituted or unsubstituted naphthyridinylenylene group, a substituted or unsubstituted quinazolinylenylene group, a substituted or unsubstituted cinnolinylenylene group, a substituted or unsubstituted indazolinylenylene group, a substituted or unsubstituted carbazolinylenylene group, a substituted or unsubstituted phenazinylenylene group, a substituted or unsubstituted phenanthridinylenylene group, a substituted or unsubstituted pyranylene group, a substituted or unsubstituted chromenylenylene group, a substituted or unsubstituted furanyl group, a substituted or unsubstituted benzofuranylenylene group, a substituted or unsubstituted thiophenylenylene group, a substituted or unsubstituted benzothiophenyl group, a substituted or unsubstituted isothiazolylenylene group, a substituted or unsubstituted benzoimidazolinylenylene group, a substituted or unsubstituted isoxazolinylenylene group, a substituted or unsubstituted dibenzothiophenylenylene group, a substituted or unsubstituted dibenzofuranylenylene group, a substituted or unsubstituted triazininylenylene group, or a substituted or unsubstituted oxadiazolylenylene group.

6. The styryl-based compound of claim 1, wherein  $L_1$  and  $L_2$  are each independently a substituted or unsubstituted phenylenyl group, a substituted or unsubstituted naphthylenylene group, a substituted or unsubstituted fluorenylenylene group, a substituted or unsubstituted phenanthrenylene group, a substituted or unsubstituted pyridinylenylene group, or a substituted or unsubstituted pyrazinylenylene group.

7. The styryl-based compound of claim 1, wherein  $L_1$  and  $L_2$  are each independently represented by any one of Formulae 5A through 5K:



Formula 5A




Formula 5B

146


-continued

5 Formula 5C




5

10 Formula 5D




10

15 Formula 5E




15

20 Formula 5F




20

25 Formula 5G




25

30 Formula 5H




30

35 Formula 5I




35

40 Formula 5J




40

45 Formula 5K



45

50 Formula 5K



50

55

wherein  $Z_{31}$  through  $Z_{34}$  are each independently selected from the group consisting of hydrogen; deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a  $C_1$ - $C_{10}$  alkyl group; a  $C_1$ - $C_{10}$  alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolinylyl group; an imidazopyridinylyl

60

65

147


group; an imidazopyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; an indolyl group; and a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazolyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>10</sub> alkyl group, and a C<sub>1</sub>-C<sub>10</sub> alkoxy group;

two or more neighboring substituents from among Z<sub>31</sub> through Z<sub>34</sub> are optionally fused with each other or are optionally connected by a single bond;


t is an integer of 1 to 6; and

u is an integer of 1 to 3.


8. The styryl-based compound of claim 1, wherein the styryl-based compound is represented by any one of Formulae 1A-1 through 1I-3 below:




Formula 1A-1




Formula 1A-2




Formula 1B-1




Formula 1B-2




Formula 1C-1




Formula 1C-2




Formula 1D-1



Formula 1E-1



Formula 1F-1



Formula 1G-1

148

-continued

5

10

15

20

25

30

35

40

45

50

55


60

65


149

-continued


Formula 1G-2&gt;




Formula 1H-1




Formula 1H-2




Formula 1I-1



Formula 1I-2



Formula 1I-3



wherein  $Z_1$  and  $Z_2$  are each independently selected from  
 the group consisting of hydrogen; deuterium; a halogen  
 atom; a hydroxyl group; a cyano group; a nitro group; an  
 amino group; an amidino group; hydrazine; hydrazone;  
 a carboxyl group or a salt thereof; a sulfonic acid group  
 or a salt thereof; a phosphoric acid or a salt thereof; a  
 $C_1$ - $C_{10}$  alkyl group; a  $C_1$ - $C_{10}$  alkoxy group; a phenyl  
 group; a naphthyl group; a fluorenyl group; a phenan-

150

threnyl group; an anthryl group; a triphenylenyl group; a  
 pyrenyl group; a chrysenyl group; an imidazolyl group;  
 an imidazolynyl group; an imidazopyridinyl group; an  
 imidazopyrimidinyl group; a pyridinyl group; a pyrazi-  
 nyl group; a pyrimidinyl group; an indolyl group; and a  
 phenyl group, a naphthyl group, a fluorenyl group, a  
 phenanthrenyl group, an anthryl group, a triphenylenyl  
 group, a pyrenyl group, a chrysenyl group, an imidazolyl  
 group, an imidazolynyl group, an imidazopyridinyl  
 group, an imidazopyrimidinyl group, a pyridinyl group,  
 a pyrazinyl group, a pyrimidinyl group, and an indolyl  
 group that are substituted with at least one of deuterium,  
 a halogen atom, a hydroxyl group, a cyano group, a nitro  
 group, an amino group, an amidino group, hydrazine,  
 hydrazone, a carboxyl group or a salt thereof, as sulfonic  
 acid group or a salt thereof, a phosphoric acid or a salt  
 thereof, a  $C_1$ - $C_{10}$  alkyl group, and a  $C_1$ - $C_{10}$  alkoxy  
 group;

$T_1$  is O or S;

$Ar_1$  and  $Ar_2$  are each independently a substituted or unsub-  
 stituted phenyl group, a substituted or unsubstituted  
 naphthyl group, a substituted or unsubstituted phenan-  
 threnyl group, a substituted or unsubstituted pyrenyl  
 group, a substituted or unsubstituted carbazolyl group, a  
 substituted or unsubstituted fluorenyl group, a substi-  
 tuted or unsubstituted dibenzofuranyl group, a substi-  
 tuted or unsubstituted dibenzothiophenyl group, or a  
 substituted or unsubstituted benzocarbazolyl group;


$L_1$  and  $L_2$  are each independently a substituted or unsub-  
 stituted phenylenyl group, as substituted or unsubsti-  
 tuted naphthylene group, a substituted or unsubstituted  
 fluorenylene group, a substituted or unsubstituted  
 phenanthrenylene group, a substituted or unsubstituted  
 pyridinylene group, or a substituted or unsubstituted  
 pyrazinylene group;

$a$  is 0 or 1; and


$b$  is 1 or 2.

9. The styryl-based compound of claim 8, wherein  $Ar_1$  and  
 $Ar_2$  are each independently represented by any one of For-  
 mulae 2A through 2J below:


Formula 2A



Formula 2B










Formula 2C



151

-continued



wherein  $Z_{11}$  through  $Z_{15}$  are each independently hydrogen; deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a  $C_1$ - $C_{10}$  alkyl group; a  $C_1$ - $C_{10}$  alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolynyl group; an imidazopyridinyl group; an imidazopyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; an indolyl group; and a phenyl group, a naphthyl group, a fluorenyl group, as phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chryse-

152

Formula 2D

5

Formula 2E

10

Formula 2F

15

$T_{11}$  is O or S;  
 $r$  is an integer of 1 to 7; and  
 $s$  is an integer of 1 to 5.

10. The styryl-based compound of claim 8, wherein  $L_1$  and  $L_2$  are each independently represented by any one of Formulae 5A through 5K below:

Formula 2G

25

Formula 2H

30

Formula 2I

35

Formula 2J


45

50


55

60


65




Formula 5A




Formula 5B




Formula 5C




Formula 5D



Formula 5E




Formula 5F




Formula 5G

153


-continued




Formula 5H



Formula 5I



Formula 5J



Formula 5K

wherein  $Z_{31}$  through  $Z_{34}$  are each independently hydrogen; deuterium; a halogen atom; a hydroxyl group; a cyano group; a nitro group; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid

154


or a salt thereof; a  $C_1$ - $C_{10}$  alkyl group; a  $C_1$ - $C_{10}$  alkoxy group; a phenyl group; a naphthyl group; a fluorenyl group; a phenanthrenyl group; an anthryl group; a triphenylenyl group; a pyrenyl group; a chrysenyl group; an imidazolyl group; an imidazolynyl group; an imidazopyridinyl group; an imidazopyrimidinyl group; a pyridinyl group; a pyrazinyl group; a pyrimidinyl group; an indolyl group; and a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an imidazolyl group, an imidazolynyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, and an indolyl group that are substituted with at least one of deuterium, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{10}$  alkyl group, and a  $C_1$ - $C_{10}$  alkoxy group;

two or more neighboring substituents from among  $Z_{31}$  through  $Z_{34}$  are optionally fused with each other or are optionally connected by a single bond;


$t$  is an integer of 1 to 6; and

$u$  is an integer of 1 to 3.


11. The styryl-based compound of claim 1, wherein the styryl-based compound is any one of Compounds 1 through 95 and 101 through 182 below:




1




2




3



4



5



6



155

156

-continued



7

8





9

10





11

12





13

14



15

16





157

158

-continued



17

18





19

20





21

22



23

24





159

160

-continued



25

26





27

28





29

30



31

32





161

162

-continued



33

34





35

36





37

38



39

40





163

164



-continued  
41

42





43

44





45

46



47

48





165

166

-continued



49

50





51

52





53

54





55

56



57

58





167

168

-continued



59

60





61

62





63

64





65

66



67

68





169


170

-continued


69




70




71




72




73




74



75



76




77


171

172


-continued




79




80




81




82




83




84



85



86




87

173


174

-continued


88



89





90




91

92



93




175


176

-continued

94





101





102

103





104

105



106

107





177

178

-continued



108

109





110

111





112

113



114

115





179

180



-continued  
116

117





118

119





120

121





122

123



124

125





181

182



-continued  
126

127





128

129





130

131





132

133



134

135













183

184

-continued  
136

137





185

186

-continued



146

147





148

149





150

151





152

153



154

155





187

188

-continued



156

157





158

159





160

161





162

163





164

165



166

167





189

190

-continued



168

169





170

171




172


173




174



175



176




191


192

-continued


177




178




179



180




181




-continued

182




12. A composition, comprising:  
a styryl-based compound represented by Formula 1:




[Formula 1]

and


at least one of a styryl-based compound represented by  
Formula 1-1H-1, a styryl-based compound represented  
by Formula 1-1H-2, and a styryl-based compound rep-  
resented by Formula 1-2H:



[Formula 1-1H-1]



[Formula 1-1H-2]



[Formula 1-2H]

wherein  $Ar_3$  is a substituted or unsubstituted  $C_8$ - $C_{20}$  aryl  
group having two or more rings fused with each other,  
or a substituted, or unsubstituted  $C_2$ - $C_{20}$  heteroaryl  
group having two or more rings fused with each other;  
 $Ar_1$  and  $Ar_2$  are each independently a substituted or unsub-  
stituted  $C_5$ - $C_{60}$  aryl group or a substituted or unsub-  
stituted  $C_2$ - $C_{60}$  heteroaryl group;  
 $L_1$  and  $L_2$  are each independently a substituted or unsub-  
stituted  $C_5$ - $C_{60}$  arylene group or a substituted or unsub-  
stituted  $C_2$ - $C_{60}$  heteroarylene group; and  
 $a$  and  $b$  are each independently an integer of 0 to 5, wherein  
a deuterium substitution rate represented by Equation 1  
below is 70% or more:

$$\text{deuterium substitution rate(\%)} = \frac{N_D}{N_{D+H}} \times 100 \quad (1)$$

wherein  $N_D$  is the number of deuterium atoms bonded to  
carbon atoms of the styryl group in Formulae 1, 1-1H-1,  
and 1-1H-2 above; and

$N_{D+H}$  is the total number of deuterium atoms and hydrogen  
atoms that are bonded to carbon atoms of the styryl  
group in Formulae 1, 1-1H-1, 1-1H-2, and 1-2H above.


13. An organic light-emitting diode (OLED) comprising:  
a first electrode;  
a second electrode facing the first electrode; and  
an organic layer interposed between the first electrode and  
the second electrode, the organic layer comprising at  
least one of the styryl-based compound of claim 1 or a  
composition containing the styryl-based compound of  
claim 1.

14. The OLED of claim 13, wherein the organic layer  
comprises at least one selected from the group consisting of a  
hole injection layer, a hole transport layer, a functional layer  
having hole injection and transport abilities, a buffer layer, an  
electron blocking layer, an emission layer, a hole blocking  
layer, an electron transport layer, an electron injection layer,  
and a functional layer having electron injection and transport  
abilities.


15. The OLED of claim 14, wherein the organic layer  
comprises the emission layer, and the emission layer com-  
prises the styryl-based compound or the composition contain-  
ing the styryl-based compound.

16. The OLED of claim 15, wherein the emission layer  
further comprises at least one selected from the group con-  
sisting of an anthracene-based compound represented by For-  
mula 400 below and an anthracene-based compound rep-  
resented by Formula 401 below as a host:

Formula 400



Formula 401



wherein  $Ar_{111}$  and  $Ar_{112}$  are each independently a substi-  
tuted or unsubstituted  $C_5$ - $C_{60}$  arylene group;

Ar<sub>113</sub> through Ar<sub>116</sub> and Ar<sub>122</sub> through Ar<sub>125</sub> are each independently a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> alkyl group or a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> aryl group;

Ar<sub>126</sub> and Ar<sub>127</sub> are each independently a C<sub>1</sub>-C<sub>10</sub> alkyl group; and

g, h, i, j, k, and l are each independently an integer of 0 to 4.

17. The LED of claim 14, wherein the organic layer comprises at least one selected from the group consisting of a hole injection layer, a hole transport layer, and a functional layer having hole transport and injection abilities, and at least one selected from the hole injection layer, the hole transport layer, and the functional layer having hole transport and injection abilities comprises the styryl-based compound or the composition containing the styryl-based compound.

\* \* \* \* \*

专利名称(译)	基于苯乙烯基的化合物，含有基于苯乙烯基的化合物的组合物，和包含基于苯乙烯基的化合物的有机发光二极管		
公开(公告)号	<a href="#">US9288869</a>	公开(公告)日	2016-03-15
申请号	US13/609633	申请日	2012-09-11
[标]申请(专利权)人(译)	韩相HYUN HWANG SEOK HWAN 金荣KOOK JUNG HYE JIN LIM JINØ 金秀YON SHIN DAE YUP 李钟赫		
申请(专利权)人(译)	HAN , SANG-HYUN HWANG , SEOK-HWAN KIM , YOUNG-KOOK JUNG , HYE-JIN LIM , JIN-O KIM , SOO-YON SHIN , DAE-YUP LEE , JONG赫		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	HAN SANG HYUN HWANG SEOK HWAN KIM YOUNG KOOK JUNG HYE JIN LIM JIN O KIM SOO YON SHIN DAE YUP LEE JONG HYUK		
发明人	HAN, SANG-HYUN HWANG, SEOK-HWAN KIM, YOUNG-KOOK JUNG, HYE-JIN LIM, JIN-O KIM, SOO-YON SHIN, DAE-YUP LEE, JONG-HYUK		
IPC分类号	H01L51/54 C07D213/74 H05B33/14 C07C255/58 C07D213/38 C07C211/61 C07C211/58 C07B59/00 C07C211/54 C07C211/56 C07C211/60 C07D307/91 C07D333/76 C07D401/10 C07D409/12 C07D209 /86 C09K11/06 C07D209/88 H01L51/00		
CPC分类号	H05B33/14 C07B59/00 C07C211/54 C07C211/56 C07C211/58 C07C211/60 C07C211/61 C07C255/58 C07D209/86 C07D209/88 C07D213/38 C07D213/74 C07D307/91 C07D333/76 C07D401/10 C07D409 /12 C09K11/06 H01L51/0059 C09K2211/1011 C09K2211/1014 C09K2211/1029 C09K2211/1088 H01L51/0058 H01L51/0061 H01L51/0074 C09K2211/1092 C07B2200/05 C07C2103/18 C07C2103/24 C07C2103/26 C07C2103/42 C07C2103/48 C07C2103/50 C07C2103/94 C07C2603/18 C07C2603/24 C07C2603/26 C07C2603/42 C07C2603/48 C07C2603/50 C07C2603/94		

优先权 1020110117161 2011-11-10 KR

其他公开文献 US20130119355A1

外部链接 [Espacenet](#) [USPTO](#)

摘要(译)

由式1表示的基于苯乙烯基的化合物，包含基于苯乙烯基的化合物的组合  
物，和包含基于苯乙烯基的化合物的有机发光二极管（OLED）：基于  
苯乙烯基的化合物可以表现出高耐热性，因此包含其的OLED可以具有低  
驱动电压，高亮度，高效率 and 长寿命。

