



US 20180026206A1

(19) **United States**(12) **Patent Application Publication**
AHN et al.(10) **Pub. No.: US 2018/0026206 A1**(43) **Pub. Date: Jan. 25, 2018**(54) **HETEROCYCLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME****Publication Classification**(51) **Int. Cl.****H01L 51/00** (2006.01)**C09K 11/02** (2006.01)**C07F 9/64** (2006.01)(52) **U.S. Cl.****CPC** **H01L 51/0072** (2013.01); **C07F 9/64**(2013.01); **H01L 51/0067** (2013.01); **C09K****11/025** (2013.01); **H01L 51/5206** (2013.01)

(57)

ABSTRACT

A heterocyclic compound and an organic light emitting device, the compound being represented by Formula 1:

(21) Appl. No.: **15/634,392**

Formula 1

(22) Filed: **Jun. 27, 2017**(30) **Foreign Application Priority Data**

Jul. 25, 2016 (KR) 10-2016-0094145


**10****190****150****110**

FIG. 1

10

190
150
110

FIG. 2

20

190
150
110
210

FIG. 3

30

220
190
150
110

FIG. 4

40

220
190
150
110
210

HETEROCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

[0001] Korean Patent Application No. 10-2016-0094145, filed on Jul. 25, 2016, in the Korean Intellectual Property Office, and entitled: "Heterocyclic Compound and Organic Light-Emitting Device Including the Same," is incorporated by reference herein in its entirety.

BACKGROUND

1. Field

[0002] Embodiments relate to a heterocyclic compound and an organic light-emitting device including the same.

2. Description of the Related Art


[0003] Organic light-emitting devices are self-emission devices that have relatively wide viewing angles, relatively high contrast ratios, relatively short response times, and increased luminance, driving voltage, and response speed characteristics. Organic light-emitting devices may produce full-color images.

[0004] Organic light-emitting devices may include a first electrode on a substrate. Organic light-emitting devices may include a hole transport region, an emission layer, an electron transport region, and a second electrode sequentially on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region. Electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, may recombine in the emission layer to produce excitons. The excitons may transition from an excited state to a ground state, thereby generating light.

SUMMARY

[0005] Embodiments are directed to a heterocyclic compound and an organic light-emitting device including the same.

[0006] The embodiments may be realized by providing a heterocyclic compound represented by Formula 1:




Formula 1

[0007] wherein, in Formula 1, Ar₁ to Ar₃ are each independently a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group, two adjacent groups of Ar₁ to Ar₃ are separate or are bound to form a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group, R₁ to R₃ are each independently represented by the following Formula 2, b₁ to b₃ are each independently

an integer of 0 to 3, b₁ to b₃ satisfy b₁+b₂+b₃≥1; or when Ar₁ to Ar₃ are each a substituted or unsubstituted benzene group, b₁+b₂+b₃≥2,

Formula 2



[0008] wherein, in Formula 2, A₁ and A₂ are each independently a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, or a pyrimidine group, L₁₁ is a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group, a₁₁ is an integer of 0 to 3; when a₁₁ is 2 or 3, the 2 or 3 L₁₁(s) are identical to or different from each other, Y is C(R₁₇)(R₁₈), sulfur (S), Si(R₁₇)(R₁₈), S(=O), S(=O)₂, or P(=O)(R₁₇), R₁₇ to R₂₀ are each independently hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —S(=O)(Q₁), or —P(=O)(Q₁)(Q₂), b₁₉ and b₂₀ are each independently an integer of 0 to 8, wherein, in Formulae 1 and 2, at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, substituted C₁-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₁-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₁-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted

C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group; a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), or —P(=O)(Q₁₁)(Q₁₂); a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group; a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), or —P(=O)(Q₂₁)(Q₂₂); or —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), wherein Q₁, Q₂, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ are each independently hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, or a terphenyl group, and * indicates a binding site to an adjacent atom.

[0009] The embodiments may be realized by providing an organic light emitting device including a first electrode; a second electrode facing the first electrode; and an organic

layer between the first electrode and the second electrode, wherein the organic layer includes an emission layer, and wherein the organic layer includes the heterocyclic compound according to an embodiment.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] Features will be apparent to those of skill in the art by describing in detail exemplary embodiments with reference to the attached drawings in which:

[0011] FIG. 1 illustrates a schematic cross-sectional view of an organic light-emitting device according to an embodiment;

[0012] FIG. 2 illustrates a schematic cross-sectional view of an organic light-emitting device according to an embodiment;

[0013] FIG. 3 illustrates a schematic cross-sectional view of an organic light-emitting device according to an embodiment; and

[0014] FIG. 4 illustrates a schematic cross-sectional view of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION


[0015] Example embodiments will now be described more fully hereinafter with reference to the accompanying drawings; however, they may be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey exemplary implementations to those skilled in the art.

[0016] In the drawing figures, the dimensions of layers and regions may be exaggerated for clarity of illustration. It will also be understood that when a layer or element is referred to as being “on” another layer or element, it can be directly on the other layer or element, or intervening layers may also be present. In addition, it will also be understood that when a layer is referred to as being “between” two layers, it can be the only layer between the two layers, or one or more intervening layers may also be present. Like reference numerals refer to like elements throughout.

[0017] As used herein, the terms “or” and “and/or” are not exclusive, and include any and all combinations of one or more of the associated listed items. Expressions such as “at least one of” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

[0018] A heterocyclic compound may be represented by Formula 1.

Formula 1



[0019] In Formula 1, Ar₁ to Ar₃ may each independently be or include, e.g., a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group. In an implementation, in Formula 1, at

least two adjacent groups of Ar₁ to Ar₃ may be separate or may be bound to form a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group.

[0020] In an implementation, in Formula 1, Ar₁ to Ar₃ may each independently be selected from or include, e.g.,

[0021] a benzene group, a naphthalene group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a triazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, and a quinazoline group; and

[0022] a benzene group, a naphthalene group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a triazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, and a quinazoline group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂).

[0023] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0024] In an implementation, in Formula 1, Ar₁ to Ar₃ may each independently be selected from, e.g.,

[0025] a benzene group and a pyridine group; and

[0026] a benzene group and a pyridine group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl

group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂).

[0027] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0028] In an implementation, in Formula 1, at least one selected from Ar₁ to Ar₃ may be selected from, e.g., a substituted or unsubstituted π electron-depleted nitrogen-containing C₂-C₆₀ heterocyclic group.

[0029] In an implementation, in Formula 1, at least one selected from Ar₁ to Ar₃ may be selected from, e.g.,

[0030] a pyridine group, a pyrimidine group, a triazine group, a quinoline group, an isoquinoline group, a quinazoline group, and a quinoxaline group; and

[0031] a pyridine group, a pyrimidine group, a triazine group, a quinoline group, an isoquinoline group, a quinazoline group, and a quinoxaline group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂).


[0032] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0033] In Formula 1, b1 to b3 may each independently be, e.g., an integer of 0 to 3. In an implementation, b1 to b3 may each independently be, e.g., 0, 1, or 2.

[0034] In Formula 1, b1 to b3 may satisfy b1+b2+b3 $\geq$ 1. In an implementation, when Ar₁ to Ar₃ are each a substituted or unsubstituted benzene group, b1+b2+b3 $\geq$ 2.

[0035] In an implementation, when Ar₁ to Ar₃ are each a benzene group, b1 may be 2, b2 may be 0, and b3 may be 0. In an implementation, b1 may be 1, b2 may be 1, and b3 may be 0.

[0036] In Formula 1, R_1 to R_3 may each independently be a group represented by Formula 2.



Formula 2

[0037] In Formula 2, A_1 and A_2 may each independently be selected from, e.g., a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, and a pyrimidine group.

[0038] In an implementation, in Formula 2, A_1 and A_2 may each independently be a benzene group.

[0039] In Formula 2, L_{11} may be selected from or include, e.g., a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.

[0040] In an implementation, in Formula 2, L_{11} may be selected from, e.g.,

[0041] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylenylene group, a spiro-bifluorenylenylene group, a spiro-benzofluorene-fluorenylenylene group, a benzofluorenylenylene group, a dibenzofluorenylenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentacenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, a silolylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an indolylenylene group, an isoindolylenylene group, an indazolylenylene group, a purinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a benzoisoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolienylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a benzosilolylenylene group, a benzothiazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylene group, a carbazolylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a dibenzosilolylenylene group, a benzocarbazolylenylene group, a naphthobenzofuranylenylene group, a naphthobenzosilolylenylene group, a dibenzocarbazolylenylene group, a dinaphthofuranylenylene group, a dinaphthothiophenylenylene group, a dinaphthosilolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, an oxazolopyridinylenylene group, a thiazolopyridinylenylene group, a benzonaphthyridinylenylene group, an azafluorenylenylene group, an azaspiro-bifluorenylenylene group, an azacarbazolylenylene group, an azadibenzofuranylenylene group, an azadibenzothiophenylenylene group, an azadibenzosilolylenylene group, an indenopyrrolylenylene group, an indolopyrrolylenylene group, an indenocarbazolylenylene group, and an indolocarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{20} aryl group, a C_1 - C_{20} heteroaryl group, a monovalent non-aromatic condensed

group, a naphthobenzothiophenylenylene group, a naphthobenzosilolylenylene group, a dibenzocarbazolylenylene group, a dinaphthofuranylenylene group, a dinaphthothiophenylenylene group, a dinaphthosilolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, an oxazolopyridinylenylene group, a thiazolopyridinylenylene group, a benzonaphthyridinylenylene group, an azafluorenylenylene group, an azaspiro-bifluorenylenylene group, an azacarbazolylenylene group, an azadibenzofuranylenylene group, an azadibenzothiophenylenylene group, an azadibenzosilolylenylene group, an indenopyrrolylenylene group, an indolopyrrolylenylene group, an indenocarbazolylenylene group, and an indolocarbazolylenylene group; and

[0042] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylenylene group, a spiro-bifluorenylenylene group, a spiro-benzofluorene-fluorenylenylene group, a benzofluorenylenylene group, a dibenzofluorenylenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentacenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, a silolylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an indolylenylene group, an isoindolylenylene group, an indazolylenylene group, a purinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a benzoisoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolienylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a benzosilolylenylene group, a benzothiazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylene group, a carbazolylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a dibenzosilolylenylene group, a benzocarbazolylenylene group, a naphthobenzofuranylenylene group, a naphthobenzosilolylenylene group, a dibenzocarbazolylenylene group, a dinaphthofuranylenylene group, a dinaphthothiophenylenylene group, a dinaphthosilolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, an oxazolopyridinylenylene group, a thiazolopyridinylenylene group, a benzonaphthyridinylenylene group, an azafluorenylenylene group, an azaspiro-bifluorenylenylene group, an azacarbazolylenylene group, an azadibenzofuranylenylene group, an azadibenzothiophenylenylene group, an azadibenzosilolylenylene group, an indenopyrrolylenylene group, an indolopyrrolylenylene group, an indenocarbazolylenylene group, and an indolocarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{20} aryl group, a C_1 - C_{20} heteroaryl group, a monovalent non-aromatic condensed


polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a methylphenyl group, a biphenyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$.

[0043] wherein Q_{31} to Q_{33} may each independently be selected from, e.g.,


[0044] a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group; and

[0045] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group, each substituted with at least one selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, and a phenyl group.


[0046] In an implementation, in Formula 2, L_{11} may be, e.g., a group represented by one of the following Formulae 3-1 to 3-46.




Formula 3-1




Formula 3-2




Formula 3-3



Formula 3-4




Formula 3-5




Formula 3-6


-continued




Formula 3-7




Formula 3-8




Formula 3-9




Formula 3-10




Formula 3-11




Formula 3-12




Formula 3-13



Formula 3-14














Formula 3-15



Formula 3-16

-continued



Formula 3-17

Formula 3-18

Formula 3-19

Formula 3-20

Formula 3-21

Formula 3-22

Formula 3-23








Formula 3-24

Formula 3-25

Formula 3-26

Formula 3-27

-continued



Formula 3-28

Formula 3-29

Formula 3-30






Formula 3-31

Formula 3-32

Formula 3-33

Formula 3-34

-continued



Formula 3-35







Formula 3-36

Formula 3-37

Formula 3-38

Formula 3-39

-continued



Formula 3-40

Formula 3-41

Formula 3-42


Formula 3-43

Formula 3-44

Formula 3-45

-continued

Formula 3-46



[0047] In Formulae 3-1 to 3-46,

[0048] Y_1 may be selected from, e.g., oxygen (O), sulfur (S), $C(Z_3)(Z_4)$, $N(Z_5)$, and $Si(Z_6)(Z_7)$.

[0049] Z_1 to Z_7 may each independently be selected from, e.g., hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazolinyl group, a benzoquinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indenocarbazolyl group, and an indolocarbazolyl group.

[0050] d_2 may be 1 or 2,

[0051] d_3 may be an integer of 1 to 3,

[0052] d_4 may be an integer of 1 to 4,

[0053] d_5 may be an integer of 1 to 5,

[0054] d_6 may be an integer of 1 to 6, and

[0055] d_8 may be an integer of 1 to 8.

[0056] In an implementation, L_{11} may be a group represented by Formula 3-2, Formula 3-12, or Formula 3-31.

[0057] In Formula 2, a_{11} may be, e.g., an integer of 0 to 3. In an implementation, a_{11} may be 0, 1, or 2. In an implementation, a_{11} may be 0 or 1. When a_{11} is 2 or 3, the 2 or 3 $L_{11}(s)$ may be identical to or different from each other.

[0058] In Formula 2, Y may be selected from, e.g., $C(R_{17})(R_{18})$, S , $Si(R_{17})(R_{18})$, $S(=O)$, $S(=O)_2$, and $P(=O)(R_{17})$.

[0059] In an implementation, in Formula 2, Y may be $C(R_{17})(R_{18})$ or $Si(R_{17})(R_{18})$.

[0060] wherein R_{17} and R_{18} may each independently be selected from, e.g., hydrogen, deuterium, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, and a tert-butyl group.

[0061] In an implementation, in Formula 2, Y may be $C(R_{17})(R_{18})$, and

[0062] R_{17} and R_{18} may each be, e.g., a methyl group.

[0063] In an implementation, in Formula 2, R_{17} to R_{20} may each independently be selected from or include, e.g.,

[0064] hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-S(=O)(Q_1)$, and $-P(=O)(Q_1)(Q_2)$.

[0065] In an implementation, in Formula 2, R_{19} and R_{20} may each independently be selected from, e.g.,

[0066] hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0067] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group;

[0068] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl

group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂); and

[0069] —P(=O)(Q₁)(Q₂) and —S(=O)(Q₁),

[0070] wherein Q₁ to Q₂ and Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0071] In an implementation, in Formula 2, R₁₉ and R₂₀ may each independently be selected from, e.g.,

[0072] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0073] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, and a hydrazono group;

[0074] a phenyl group, a biphenyl group, a terphenyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group; and

[0075] a phenyl group, a biphenyl group, a terphenyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl

group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group; and

[0076] —P(=O)(Q₁)(Q₂) and —S(=O)(Q₁),

[0077] wherein Q₁ to Q₂ and Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0078] In an implementation, in Formula 2, R₁₉ and R₂₀ may each independently be selected from, e.g.,


[0079] hydrogen, a cyano group, a C₁-C₂₀ alkyl group, a phenyl group, a carbazole group, —P(=O)(Q₁)(Q₂), and —S(=O)(Q₁), and

[0080] Q₁ and Q₂ may each be a phenyl group.


[0081] In an implementation, in Formula 2, b19 and b20 may each independently be, e.g., an integer of 0 to 8. In an implementation, b19 and b20 may each independently be, e.g., an integer of 0 to 4. In an implementation, b19 and b20 may each independently be, e.g., an integer of 0 to 2.

[0082] In an implementation, the heterocyclic compound represented by Formula 1 may be represented by Formula 1-1, and groups represented by Formula 2 may be represented by Formula 2-1.

Formula 1-1



Formula 2-1



[0083] In Formula 1-1,

[0084] X₁₁ may be nitrogen (N) or C(R₁₁), X₁₂ may be N or C(R₁₂), X₁₃ may be N or C(R₁₃), X₁₄ may be N or C(R₁₄), X₁₅ may be N or C(R₁₅), X₁₆ may be N or C(R₁₆), X₂₁ may be N or C(R₂₁), X₂₂ may be N or C(R₂₂), X₂₃ may be N or C(R₂₃), X₂₄ may be N or C(R₂₄), X₂₅ may be N or C(R₂₅), X₂₆ may be N or C(R₂₆), X₃₁ may be N or C(R₃₁), X₃₂ may be N or C(R₃₂), X₃₃ may be N or C(R₃₃),

[0085] R₁₁ to R₁₆, R₂₁ to R₂₆, and R₃₁ to R₃₃ may each independently be selected from, e.g., a group represented by Formula 2-1, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino

group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group,

[0086] when X₁₁ is C(R₁₁), X₁₂ is C(R₁₂), X₁₃ is C(R₁₃), X₁₄ is C(R₁₄), X₁₅ is C(R₁₅), X₁₆ is C(R₁₆), X₂₁ is C(R₂₁), X₂₂ is C(R₂₂), X₂₃ is C(R₂₃), X₂₄ is C(R₂₄), X₂₅ is C(R₂₅), X₂₆ is C(R₂₆), X₃₁ is C(R₃₁), X₃₂ is C(R₃₂), and X₃₃ is C(R₃₃), at least two selected from R₁₁, R₂₁, R₃₁, R₂₂, and R₁₂ may be a group represented by Formula 2-1, or at least one selected from R₁₂ and R₁₃, R₂₃, R₃₂, R₂₄, and R₁₄ may be a group represented by Formula 2-1,

[0087] wherein R₁₂ and R₁₃, or R₁₄ and R₁₅ may be bound to form a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group, and

[0088] in Formula 2-1, L, n, Y, R₁₉, R₂₀, b19, and b20 may be the same as those described above.

[0089] In an implementation, in Formula 1-1, at least one selected from X₁₁ to X₁₆ may be N,

[0090] X₂₁ may be C(R₂₁), X₂₂ may be C(R₂₂), X₂₃ may be C(R₂₃), X₂₄ may be C(R₂₄), X₂₅ may be C(R₂₅), X₂₆ may be C(R₂₆), and

[0091] at least one selected from X₂₁ to X₂₆ may be a group represented by Formula 2-1.

[0092] In an implementation, in Formula 1-1, at least one selected from X₁₁ to X₁₆ may be N,

[0093] X₃₁ may be C(R₃₁), X₃₂ may be C(R₃₂), X₃₃ may be C(R₃₃), and

[0094] at least one selected from R₃₁ to R₃₃ may be a group represented by Formula 2-1.

[0095] In an implementation, in Formula 1-1, at least one selected from X₁₁ to X₁₆ may be N,

[0096] X₂₁ may be C(R₂₁), X₂₂ may be C(R₂₂), X₂₃ may be C(R₂₃), X₂₄ may be C(R₂₄), X₂₅ may be C(R₂₅), X₂₆ may be C(R₂₆),

[0097] X₃₁ may be C(R₃₁), X₃₂ may be C(R₃₂), X₃₃ may be C(R₃₃), and

[0098] at least one selected from R₂₁ to R₂₆ and R₃₁ to R₃₃ may be a group represented by Formula 2-1.

[0099] In an implementation, in Formula 1-1, at least one selected from X₂₁ to X₂₆ may be N,

[0100] X₃₁ may be C(R₃₁), X₃₂ may be C(R₃₂), X₃₃ may be C(R₃₃), and

[0101] at least one selected from R₃₁ to R₃₃ may be a group represented by Formula 2-1.

[0102] In an implementation, at least one selected from X₂₁ to X₂₆ may be N,

[0103] X₁₁ may be C(R₁₁), X₁₂ may be C(R₁₂), X₁₃ may be C(R₁₃), X₁₄ may be C(R₁₄), X₁₅ may be C(R₁₅), X₁₆ may be C(R₁₆),

[0104] X₃₁ may be C(R₃₁), X₃₂ may be C(R₃₂), X₃₃ may be C(R₃₃), and

[0105] at least one selected from R₁₁ to R₁₆ and R₃₁ to R₃₃ may be a group represented by Formula 2-1.

[0106] In an implementation, in Formula 1-1, X₂₂ may be N,

[0107] X₂₁ may be C(R₂₁), X₂₅ may be C(R₂₅), X₂₆ may be C(R₂₆), and

[0108] at least one selected from R₂₁, R₂₅, and R₂₆ may be a group represented by Formula 2-1.

[0109] In an implementation, in Formula 1-1, at least one selected from X₃₁ to X₃₃ may be N,

[0110] X₂₁ may be C(R₂₁), X₂₂ may be C(R₂₂), X₂₃ may be C(R₂₃), X₂₄ may be C(R₂₄), X₂₅ may be C(R₂₅), X₂₆ may be C(R₂₆), and

[0111] at least one selected from R₂₁ to R₂₆ may be a group represented by Formula 2-1.




[0112] In an implementation, in Formula 1-1, at least one selected from X₃₁ to X₃₃ may be N,

[0113] X₁₁ may be C(R₁₁), X₁₂ may be C(R₁₂), X₁₃ may be C(R₁₃), X₁₄ may be C(R₁₄), X₁₅ may be C(R₁₅), X₁₆ may be C(R₁₆),


[0114] X₂₁ may be C(R₂₁), X₂₂ may be C(R₂₂), X₂₃ may be C(R₂₃), X₂₄ may be C(R₂₄), X₂₅ may be C(R₂₅), X₂₆ may be C(R₂₆), and

[0115] at least one selected from R₁₁ to R₁₆ and R₂₁ to R₂₆ may be a group represented by Formula 2-1.


[0116] In an implementation, the heterocyclic compound represented by Formula 1 may be one of the following Compounds 1 to 77.




-continued




4



5




6




7


-continued



8




9




10

-continued




11


-continued




14




12




15



13




16




17

-continued




18


-continued




22




19




23




20



24




21



25


-continued

26




-continued

29




30

27




31




32


28



33




-continued




34


-continued




37




35



38




36




39

-continued




40


-continued




43




41




44



42




45




46

-continued




47


-continued




51




48




52




49



53




50









54

-continued




-continued




-continued

62




-continued


65




63




66




67




64



68




-continued




69


-continued




73




70




74



71




72



75

-continued



[0117] The heterocyclic compound represented by Formula 1 may have a heterocyclic structure having a phosphine oxide substituent, and thus may be advantageous in terms of luminous efficiency and energy transfer.

[0118] For example, in the heterocyclic compound, at least two substituents represented by Formula 2 may be substituted at a triaryl phosphine oxide, and thus the heterocyclic compound may have improved hole mobility as compared with a case in which the heterocyclic compound has only one substituent. Performance of a device may be varied by adjusting the position of the substituents to control the polarity of molecules.

[0119] In an implementation, in the heterocyclic compound, the substituent represented by Formula 2 may be bound to the phosphine oxide group via at least one π electron-depleted nitrogen-containing C_2-C_{60} heterocyclic group (e.g., a pyridine group), thus facilitating control of polarity of the heterocyclic compound molecule and enabling suitable control of hole transportability or electron transportability in a device. Consequently, the heterocyclic compound may help improve device characteristics.

[0120] For example, the substituent represented by Formula 2 may help prevent an electron conjugation effect, as compared with a case having "O" instead of Y, thus increasing an energy gap of a material. Further, in the heterocyclic compound, when a "N" position in the π electron-depleted nitrogen-containing C_2-C_{60} heterocyclic group bound to the phosphine oxide group is at an ortho-, meta-, or para-position with respect to a binding position to the phosphine oxide, the polarity of a molecule may increase in general, or intermolecular force may increase, resulting in an increase in

electron transportability of a material. Consequently, the heterocyclic compound may help improve device characteristics.

[0121] In an implementation, in the heterocyclic compound, when the substituent represented by Formula 2 bound to the π electron-depleted nitrogen-containing C_2-C_{60} heterocyclic group is at a meta- or para-position with respect to a binding position to the phosphine oxide, an energy level of a material may be easily controlled by adjusting electron donating characteristics, and device characteristics may improve by appropriately controlling electron and hole transportability of the material.

[0122] At least one heterocyclic compound represented by Formula 1 may be included between a pair of electrodes in an organic light-emitting device. In an implementation, the heterocyclic compound may be included in an emission layer and/or an electron transport region. In an implementation, the heterocyclic compound represented by Formula 1 may be used as a material for forming a capping layer, which is disposed on outer sides of a pair of electrodes in an organic light-emitting device.

[0123] Accordingly, there is provided an organic light-emitting device including a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, wherein the organic layer includes an emission layer. The organic layer may include at least one heterocyclic compound represented by Formula 1.

[0124] As used herein, "(for example, the organic layer) including at least one heterocyclic compound" means that "(the organic layer) including a heterocyclic compound represented by Formula 1, or at least two different heterocyclic compounds represented by Formula 1".

[0125] For example, the organic layer may include Compound 1 only as the heterocyclic compound. In this case, Compound 1 may be included in the emission layer of the organic light-emitting device. In an implementation, the organic layer may include Compounds 1 and 2 as the heterocyclic compounds. In this regard, Compounds 1 and 2 may be present in the same layer (e.g., Compounds 1 and 2 may be both present in an emission layer), or in different layers (e.g., Compound 1 may be present in an emission layer and Compound 2 may be present in a hole transport layer).

[0126] The organic layer may include i) a hole transport region between the first electrode (anode) and the emission layer that includes at least one selected from a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer, and ii) an electron transport region between the emission layer and the second electrode (cathode) that includes at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer. At least one selected from the hole transport region and the emission layer may include at least one heterocyclic compound represented by Formula 1. In an implementation, the organic light-emitting device may include an emission layer, and the emission layer may include at least one heterocyclic compound represented by Formula 1. The emission layer may include a host and a dopant, and at least one selected from the host and the dopant may include at least one heterocyclic compound represented by Formula 1. In an implementation, the heterocyclic compound represented by Formula 1 included in the emission layer may serve as a host, and in this case, the emission layer may further include a dopant described herein. In an implementation, the heterocyclic compound represented by Formula 1 included in

the emission layer may serve as a dopant, and in this case, the emission layer may further include a host described herein.

[0127] The organic light-emitting device may further include at least one selected from a first capping layer disposed in a path of light extracted or emitted from the emission layer, allowing the light to pass through to the outside after passing the first electrode; and a second capping layer disposed in a path of light extracted or emitted from the emission layer, allowing the light to pass through to the outside after passing the second electrode, wherein at least one selected from the first capping layer and the second capping layer may include at least one heterocyclic compound.

[0128] In an implementation, the organic light-emitting device may have a structure of i) first electrode, organic layer, second electrode, and second capping layer, ii) first capping layer, first electrode, organic layer, and second electrode, or iii) first capping layer, first electrode, organic layer, second electrode, and second capping layer, wherein layers of each structure are sequentially stacked in the stated order. At least one selected from the first capping layer and the second capping layer may include the heterocyclic compound.

[0129] The term “organic layer” as used herein refers to a single and/or a plurality of layers between the first electrode and the second electrode in an organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

[0130] FIG. 1 illustrates a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 may include the first electrode 110, the organic layer 150, and the second electrode 190.

[0131] Hereinafter, the structure of the organic light-emitting device 10 according to an embodiment and a method of manufacturing an organic light-emitting device according to an embodiment will be described in connection with FIG. 1.

[0132] In an implementation, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be a glass substrate or a plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

[0133] The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode 110 over the substrate. When the first electrode 110 is an anode, the material for forming the first electrode 110 may be selected from materials with a high work function that facilitate hole injection.

[0134] The first electrode 110 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode 110 is a transmissive electrode, a material for forming the first electrode 110 may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), zinc oxide (ZnO), and any combination thereof. When the first electrode 110 is a semi-transmissive electrode or a reflective electrode, as a material for forming the first electrode 110, at least one selected from magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and any combination thereof may be used.

[0135] The first electrode 110 may have a single-layered structure, or a multi-layered structure including two or more layers. In some embodiments, the first electrode 110 may have a triple-layer structure of ITO/Ag/ITO.

[0136] The organic layer 150 may be on the first electrode 110. The organic layer 150 may include an emission layer.


[0137] The organic layer 150 may further include a hole transport region between the first electrode 110 and the emission layer and an electron transport region between the emission layer and the second electrode 190.

[0138] The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.


[0139] The hole transport region may include at least one selected from a hole injection layer, an emission auxiliary layer, and an electron blocking layer.

[0140] For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials or a multi-layered structure, e.g., a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure, wherein layers of each structure are sequentially stacked on the first electrode 110 in each stated order.

[0141] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPD), β -NPB, TPD, a spiro-TPD, a spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), (polyaniline)/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:





-continued




TDATA


-continued

 β -NPB


TPD




2-TNATA




Spiro-TPD



Spiro-NPB



NPB



methylated NPB

Formula 202

[0144] L₂₀₅ may be selected from *—O—*, *—S—*, *—N(Q₂₀₁)—*, a substituted or unsubstituted C₁–C₂₀ alkylene group, a substituted or unsubstituted C₂–C₂₀ alkenylene group, a substituted or unsubstituted C₃–C₁₀ cycloalkylene group, a substituted or unsubstituted C₁–C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃–C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁–C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆–C₆₀ arylene group, a substituted or unsubstituted C₁–C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.

[0152] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a

cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

[0153] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0154] In one or more embodiments, xa1 to xa4 may each independently be 0, 1, or 2.

[0155] According to another embodiment, xa5 may be 1, 2, 3, or 4.

[0156] According to some embodiments, R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0157] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group,

an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂).

[0158] wherein Q₃₁ to Q₃₃ may be the same as those described herein.

[0159] According to some embodiments, at least one selected from R₂₀₁ to R₂₀₃ in Formula 201 may be selected from

[0160] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0161] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.


[0162] According to some embodiments, in Formula 202, i) R₂₀₁ and R₂₀₂ may be bound via a single bond, and/or ii) R₂₀₃ and R₂₀₄ may be bound via a single bond.

[0163] According to some embodiments, at least one selected from R₂₀₁ to R₂₀₄ in Formula 202 may be selected from

[0164] a carbazolyl group; and


[0165] a carbazolyl group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.

[0166] The compound represented by Formula 201 may be represented by Formula 201A:




Formula 201A

[0167] In some embodiments, the compound represented by Formula 201 may be represented by Formula 201A(1).




Formula 201A(1)

[0168] In some embodiments, the compound represented by Formula 201 may be represented by Formula 201A-1:




Formula 201A-1

[0169] In some embodiments, the compound represented by Formula 202 may be represented by Formula 202A:



Formula 202A

[0170] In some embodiments, the compound represented by Formula 202 may be represented by Formula 202A-1:



Formula 202A-1

[0171] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,


[0172] L_{201} to L_{203} , $xa1$ to $xa3$, $xa5$, and R_{202} to R_{204} may each independently be the same as those described herein,

[0173] R_{211} and R_{212} may each independently be substantially the same as the those described herein with reference to R_{203} , and


[0174] R_{213} to R_{217} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-F$, a pentalenyl

group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.


[0175] The hole transport region may include at least one compound selected from Compounds HT1 to HT39:




HT1



HT2




HT3




HT4

-continued


HT5




HT6



HT7




HT8




-continued


HT9




HT10




HT11




HT12




-continued
HT13




HT14




HT15




HT16



HT17




HT18




-continued


HT19




HT20



HT21




HT22




-continued

HT23





HT24




HT25

HT26




-continued


HT27




HT28




HT29




HT30




HT31




HT32



HT33




HT34




-continued


HT35




HT36




HT37



HT38



HT39



[0176] The thickness of the hole transport region may be in a range of about 100 (Angstroms) Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å. When the hole transport region includes at least one selected from a hole injection layer and a hole transport layer, the thickness of the hole injection layer may be in a range of about 100 Å to about 9,000 Å, e.g., about 100 Å to about 1,000 Å, and the thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, e.g., about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are within any of these ranges, excellent hole transport characteristics may be obtained without a substantial increase in driving voltage.

[0177] The emission auxiliary layer may increase light emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer. The electron blocking layer may reduce or eliminate the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include materials as described herein.

[0178] The hole transport region may include a charge generating material as well as the aforementioned materials, to improve conductive properties of the hole transport region. The charge generating material may be substantially homogeneously or non-homogeneously dispersed in the hole transport region.

[0179] The charge generating material may include, e.g., a p-dopant.

[0180] According to an embodiment, a lowest unoccupied molecular orbital (LUMO) of the p-dopant may be about -3.5 electron Volts (eV) or less.

[0181] The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a compound including a cyano group.



[0182] In some embodiments, the p-dopant may include at least one selected from

[0183] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) or 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);


[0184] a metal oxide, such as tungsten oxide or molybdenum oxide;

[0185] 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and

[0186] a compound represented by Formula 221:



Formula 221



[0187] In Formula 221,

[0188] R_{221} to R_{223} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, provided that at least one selected from R_{221} to R_{223} may include at least one substituent selected from a cyano group, —F, —Cl, —Br, —I, a C_1 - C_{20} alkyl group substituted with —F, a C_1 - C_{20} alkyl group substituted with —Cl, a C_1 - C_{20} alkyl group substituted with —Br, and a C_1 - C_{20} alkyl group substituted with —I.

[0189] When the organic light-emitting device 10 is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In one or more embodiments, the emission layer may have a stacked structure. The stacked structure may include two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer. The two or more layers may be in direct contact with each other. Alternatively, the two or more layers may be separated from each other. In one or more embodiments, the emission layer may include two or more materials. The two or more materials may include a red light-emitting material, a green light-emitting material, or a blue light-emitting material. The two or more materials may be mixed with each other in a single layer. The two or more materials mixed with each other in the single layer may emit white light.

[0190] The emission layer may include a host and a dopant. The dopant may include at least one of a fluorescent dopant and a phosphorescent dopant.

[0191] The amount of the dopant in the emission layer may be, in general, in a range of about 0.01 parts to about 15 parts by weight based on 100 parts by weight of the host.

[0192] The thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, and in some embodiments, about 200 Å to about 600 Å. When the thickness of the emission layer is within any of these ranges, increased luminous characteristics may be obtained without a substantial increase in driving voltage.

[0193] The host may further include, in addition to the heterocyclic compound represented by Formula 1, a compound represented by Formula 301:



[0194] In Formula 301,

[0195] Ar_{301} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0196] $xb11$ may be 1, 2, or 3,

[0197] L_{301} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0198] $xb1$ may be an integer from 0 to 5,

[0199] R_{301} may be selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{301})(Q_{302})(Q_{303})$, $-N(Q_{301})$

(Q_{302}) , $-B(Q_{301})(Q_{302})$, $-C(=O)(Q_{301})$, $-S(=O)_2(Q_{301})$, and $-P(=O)(Q_{301})(Q_{302})$, and

[0200] $xb21$ may be an integer from 1 to 5,

[0201] wherein Q_{301} to Q_{303} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0202] In some embodiments, Ar_{301} in Formula 301 may be selected from

[0203] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group; and


[0204] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})(Q_{32})$,

[0205] wherein Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.


[0206] When $xb11$ in Formula 301 is 2 or greater, at least two $Ar_{301}(s)$ may be bound via a single bond.

[0207] In one or more embodiments, the compound represented by Formula 301 may be represented by Formula 301-1 or 301-2:

Formula 301-1



Formula 301-2



[0208] In Formulae 301-1 to 301-2,

[0209] A_{301} to A_{304} may each independently be selected from a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, an indene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, an indole group, a carbazole group, a benzocarbazole group, a dibenzocarbazole group, a furan group, a benzofuran group, a dibenzofuran group, a naphthofuran group, a benzonaphthofuran group, a dinaphthofuran group, a thiophene group, a benzothiophene group, a dibenzothiophene group, a naphthothiophene group, a benzonaphthothiophene group, and a dinaphthothiophene group,

[0210] X_{301} may be O, S, or N- $[(L_{304})_{xb4}-R_{304}]$,

[0211] R_{311} to R_{314} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),

[0212] $xb22$ and $xb23$ may each independently be 0, 1, or 2,

[0213] L_{301} , $xb1$, R_{301} , and Q_{31} to Q_{33} may each independently be the same as those described herein,

[0214] L_{302} to L_{304} may each independently be substantially the same as those described with reference to L_{301} ,

[0215] $xb2$ to $xb4$ may each independently be substantially the same as those described with reference to $xb1$, and

[0216] R_{302} to R_{304} may each independently be substantially the same as those described herein with reference to R_{301} .

[0217] In some embodiments, in Formulae 301, 301-1, and 301-2, L_{301} to L_{304} may each independently be selected from

[0218] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinoxalinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a

triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

[0219] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinoxalinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a benzofuranyl group, a benzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalynyl group, a quinoxalynyl group, a cinnolynyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32})

[0220] wherein Q_{31} to Q_{33} may each independently be the same as those described herein.

[0221] In some embodiments, in Formulae 301, 301-1, and 301-2, R_{301} to R_{304} may each independently be selected from

[0222] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and


[0223] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl

group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, $-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})(Q_{32})$,


[0224] wherein Q_{31} to Q_{33} may each independently be the same as those described herein.

[0225] In some embodiments, the host may include an alkaline earth metal complex. For example, the host may include a beryllium (Be) complex, for example, Compound H55, a magnesium (Mg) complex, or a zinc (Zn) complex.


[0226] The host may include at least, in addition to the heterocyclic compound of represented by Formula 1, one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di-(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolyl-benzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55:



H1




H2




H3


-continued




H4




H5




H6



H7




H8




H9


-continued




H10




H11



H12




H13



H14


-continued

H15




-continued


H20




H16




H21




H17




H22




H18




H23



H19




-continued




H24

-continued




H26

H25




H27




-continued

H28





-continued

H32





H33

H29





H34

H30




H31

H35




-continued




H36

-continued




H40



H37




H38



H39






H41




H42

-continued




-continued

H43




H46

H44




H47

H48




H49


H45



H50




H51




-continued


H52




H53



H54




H55




[0227] The dopant may further include, in addition to the heterocyclic compound represented by Formula 1, a phosphorescent dopant or a fluorescent dopant.

[0228] The phosphorescent dopant may include an organometallic complex represented by Formula 401:

Formula 401



Formula 402



[0229] In Formulae 401 and 402,

[0230] M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

[0231] L_{401} may be selected from ligands represented by Formula 402. $xc1$ may be an integer selected from 1, 2, and 3; when $xc1$ is 2 or greater, at least two $L_{401}(s)$ may be identical to or different from each other,

[0232] L_{402} may be an organic ligand. $xc2$ may be an integer from 0 to 4; when $xc2$ is 2 or greater, at least two $L_{402}(s)$ may be identical to or different from each other,

[0233] X_{401} to X_{404} may each independently be nitrogen (N) or carbon (C),

[0234] X_{401} and X_{403} may be linked via a single bond or a double bond, X_{402} and X_{404} may be linked via a single bond or a double bond,

[0235] A_{401} and A_{402} may each independently be a C_5 - C_{60} carbocyclic group or a C_1 - C_{60} heterocyclic group,

[0236] X_{405} may be a single bond, $^*O^*$, $^*S^*$, $^*C(=O)^*$, $^*N(Q_{411})^*$, $^*C(Q_{411})(Q_{412})^*$, $^*C(Q_{411})=C(Q_{412})^*$, $^*C(Q_{411})=^*$, or $^*C(Q_{411})=^*$, wherein Q_{411} and Q_{412} may each independently be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

[0237] X_{406} may be a single bond, O, or S.

[0238] R_{401} and R_{402} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_1 - C_{20} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})(Q_{402})$, $-B(Q_{401})(Q_{402})$, $-C(=O)(Q_{401})$, $-S(=O)_2(Q_{401})$, and $-P(=O)(Q_{401})(Q_{402})$, wherein Q_{401} to Q_{403}

may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a C₆-C₂₀ aryl group, and a C₁-C₂₀ heteroaryl group,

[0239] xc11 and xc12 may each independently be an integer from 0 to 10, and

[0240] * and * in Formula 402 each indicate a binding site to M in Formula 401.

[0241] In an embodiment, in Formula 402, A₄₀₁ and A₄₀₂ may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

[0242] In one or more embodiments, in Formula 402, i) X₄₀₁ may be nitrogen, and X₄₀₂ may be carbon, or ii) X₄₀₁ and X₄₀₂ may each be nitrogen.

[0243] According to some embodiments, R₄₀₁ and R₄₀₂ in Formula 402 may each independently be selected from

[0244] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0245] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantyl group, a norbornanyl group, and a norbornenyl group;

[0246] a cyclopentyl group, a cyclohexyl group, an adamantyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0247] a cyclopentyl group, a cyclohexyl group, an adamantyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a

pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0248] —Si(Q₄₀₁)(Q₄₀₂)(Q₄₀₃), —N(Q₄₀₁)(Q₄₀₂), —B(Q₄₀₁)(Q₄₀₂), —C(=O)(Q₄₀₁), —S(=O)₂(Q₄₀₁), and —P(=O)(Q₄₀₁)(Q₄₀₂),


[0249] wherein Q₄₀₁ to Q₄₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group.

[0250] In one or more embodiments, when xc1 in Formula 401 is 2 or greater, two A₄₀₁(s) in at least two L₄₀₁(s) may optionally be linked via X₄₀₇ as a linking group, and two A₄₀₂(s) may optionally be linked via X₄₀₈ as a linking group (see e.g., Compounds PD1 to PD4 and PD7). X₄₀₇ and X₄₀₈ may each independently be selected from a single bond, *—O—*, *—S—*, *—C(=O)—*, *—N(Q₄₁₃)—, *—C(Q₄₁₃)(Q₄₁₄)—*, and *—C(Q₄₁₃)=C(Q₄₁₄)—*, wherein Q₄₁₃ and Q₄₁₄ may each independently be hydrogen, deuterium, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group.


[0251] L₄₀₂ in Formula 401 may be any suitable monovalent, divalent, or trivalent organic ligand. In some embodiments, L₄₀₂ may be selected from halogen, diketone (e.g., acetylacetonate), a carboxylic acid (e.g., picolinate), —C(=O), isonitrile, —CN, and phosphorus (e.g., phosphine or phosphite).

[0252] In some embodiments, the phosphorescent dopant may include, for example, at least one selected from Compounds PD1 to PD25:







PD1



PD2




-continued




-continued

PD3




PD9

PD4




PD10

PD5




PD11

PD6



PD12






PD7




PD13

PD8


-continued




PD14




PD15



PD16



PD17



PD18

-continued


PD19

PD20


PD21

PD22


-continued



PD23




PD24



PD25

[0253] The fluorescent dopant may include an arylamine compound or a styrylamine compound.

[0254] In some embodiments, the fluorescent dopant may include a compound represented by Formula 501:



Formula 501

[0255] In Formula 501,

[0256] Ar₅₀₁ may be a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₁-C₆₀ heterocyclic group,

[0257] L₅₀₁ to L₅₀₃ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

cloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0258] xd1 to xd3 may each independently be an integer from 0 to 3,

[0259] R₅₀₁ and R₅₀₂ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

[0260] xd4 may be an integer from 1 to 6.

[0261] In some embodiments, Ar₅₀₁ in Formula 501 may be selected from

[0262] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group; and

[0263] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0264] In one or more embodiments, L₅₀₁ to L₅₀₃ in Formula 501 may each independently be selected from

[0265] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacylenylene group, a pentacylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group; and

[0266] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0267] According to some embodiments, R_{501} and R_{502} in Formula 501 may each independently be selected from

[0268] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0269] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a


C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, and —Si(Q_{31})(Q_{32})(Q_{33}),

[0270] wherein Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.


[0271] In one or more embodiments, xd4 in Formula 501 may be 2.

[0272] In some embodiments, the fluorescent dopant may be selected from Compounds FD1 to FD22:


FD1



FD2




-continued




FD3

-continued

FD5





FD6




FD4

FD7




-continued

FD8




-continued


FD11




FD12




FD9




FD13







FD10



FD14




-continued



-continued


FD15

FD19




FD16

FD20




FD17

FD21




FD18


FD22




[0273] In some embodiments, the fluorescent dopant may be selected from the following compounds:




DPVBi




DPAVBi




TBPe




DCM



DCJTb



Coumarin 6



C545T

[0274] The electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure each having a plurality of layers, each having a plurality of different materials.

[0275] The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer.

[0276] In some embodiments, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, wherein layers of each structure are sequentially stacked on the emission layer in each stated order.

[0277] The electron transport region, e.g., a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region, may include a metal-free compound. The metal-free compound may include at least one π electron-depleted nitrogen-containing ring.

[0278] The “ π electron-depleted nitrogen-containing ring” may indicate a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

[0279] For example, the “ π electron-depleted nitrogen-containing ring” may be i) a 5-membered to 7-membered heteromonocyclic group having at least one $*-N=*$ moiety, ii) a heteropolycyclic group in which two or more 5-membered to 7-membered heteromonocyclic groups, each having at least one $*-N=*$ moiety are condensed, or iii) a heteropolycyclic group in which at least one of 5-membered to 7-membered heteromonocyclic group, each having at least one $*-N=*$ moiety, is condensed with at least one C_5 - C_{60} carbocyclic group.

[0280] Examples of the π electron-depleted nitrogen-containing ring may include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an iso-benzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, a thiadiazole, an imidazopyridine, an imidazopyrimidine, and an azacarbazole.

[0281] For example, the electron transport region may include, in addition to the heterocyclic compound represented by Formula 1, a compound represented by Formula 601:



[0282] In Formula 601,

[0283] Ar_{601} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0284] $xe11$ may be 1, 2, or 3,

[0285] L_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a sub-

stituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0286] $xe1$ may be an integer from 0 to 5,

[0287] R_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_{601})(\text{Q}_{602})(\text{Q}_{603})$, $-\text{C}(=\text{O})(\text{Q}_{601})$, $-\text{S}(=\text{O})_2(\text{Q}_{601})$, and $-\text{P}(=\text{O})(\text{Q}_{601})(\text{Q}_{602})$,

[0288] wherein Q_{601} to Q_{603} may each independently be a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

[0289] $xe21$ may be an integer from 1 to 5.

[0290] In some embodiments, at least one selected from the $xe11$ $Ar_{601}(s)$ and the $xe21$ $R_{601}(s)$ may include a π electron-depleted nitrogen-containing ring.

[0291] In some embodiments, in Formula 601, Ar_{601} may be selected from

[0292] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

[0293] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline

group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂).


[0294] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0295] When xe11 in Formula 601 is 2 or greater, at least two Ar_{601(s)} may be bound via a single bond.

[0296] In one embodiment, in Formula 601, Ar₆₀₁ may be an anthracene group.

[0297] In some embodiments, the compound represented by Formula 601 may be represented by Formula 601-1:

Formula 601-1



[0298] In Formula 601-1,

[0299] X₆₁₄ may be N or C(R₆₁₄), X₆₁₅ may be N or C(R₆₁₅), X₆₁₆ may be N or C(R₆₁₆), and at least one selected from X₆₁₄ to X₆₁₆ may be N,

[0300] L₆₁₁ to L₆₁₃ may each independently be substantially the same as those described herein with reference to L₆₀₁,

[0301] xe611 to xe613 may each independently be substantially the same as those described herein with reference to xe1,

[0302] R₆₁₁ to R₆₁₃ may each independently be substantially the same as those described herein with reference to R₆₀₁, and

[0303] R₆₁₄ to R₆₁₆ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0304] In one embodiment, in Formulae 601 and 601-1, L₆₀₁ and L₆₀₁ to L₆₁₃ may each be independently selected from

[0305] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

[0306] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocar-

bazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group.

[0307] In one or more embodiments, xe1 and xe611 to xe613 in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

[0308] In some embodiments, R₆₀₁ to R₆₁₁ and R₆₁₃ in Formulae 601 and 601-1 may each independently be selected from

[0309] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;

[0310] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl


group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

[0311] —S(=O)₂(Q₆₀₁) and —P(=O)(Q₆₀₁)(Q₆₀₂),

[0312] wherein Q₆₀₁ and Q₆₀₂ may each independently be substantially the same as those described herein.


[0313] The electron transport region may include at least one compound selected from Compounds ET1 to ET36:

ET1




-continued

ET2




-continued


ET5




ET3




ET6




ET4



ET7




-continued




ET8


-continued




ET10



ET9




ET11



ET12


-continued

ET13





-continued

ET16





ET17

ET14




ET18

ET15




-continued



ET19



-continued

ET22





ET20

ET23




ET21

ET24




-continued

ET25





-continued

ET28




ET29


ET26




ET27



ET30




-continued




ET31


-continued




ET34




ET32



ET35





ET33




ET36


[0314] In one or more embodiments, the electron transport region may include at least one selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq₃, BAlq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ:

Alq₃

BAlq



TAZ



NTAZ


[0315] The thickness of the buffer layer, the hole blocking layer, or the electron control layer may each independently be in a range of about 20 Å to about 1,000 Å, e.g., about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer or the electron control layer are within any of these ranges, excellent hole blocking characteristics or excellent electron controlling characteristics may be obtained without a substantial increase in driving voltage.

[0316] The thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 150 Å to about 500 Å. When the thickness of the electron transport layer is within any of these ranges, excellent electron transport characteristics may be obtained without a substantial increase in driving voltage.


[0317] The electron transport region (e.g., the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a material including metal.

[0318] The material including metal may include at least one selected from an alkali metal complex and an alkaline earth metal complex. The alkali metal complex may include a metal ion selected from a lithium (Li) ion, a sodium (Na) ion, a potassium (K) ion, a rubidium (Rb) ion, and a cesium (Cs) ion. The alkaline earth metal complex may include a metal ion selected from a beryllium (Be) ion, a magnesium (Mg) ion, a calcium (Ca) ion, an strontium (Sr) ion, and a barium (Ba) ion. Each ligand coordinated with the metal ion of the alkali metal complex and the alkaline earth metal complex may independently be selected from a hydroxyquinoline, a hydroxyisoquinoline, a hydroxybenzoquinoline, a hydroxyacridine, a hydroxyphenanthridine, a hydroxyphenyloxazole, a hydroxyphenylthiazole, a hydroxydiphenyl oxadiazole, a hydroxydiphenyl thiadiazole, a hydroxyphenyl pyridine, a hydroxyphenyl benzimidazole, a hydroxyphenyl benzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene.

[0319] For example, the material including metal may include a Li complex. The Li complex may include, e.g., Compound ET-D1 (lithium quinolate, LiQ) or ET-D2:



ET-D1



ET-D2

[0320] The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode 190. The electron injection layer may be in direct contact with the second electrode 190.

[0321] The electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers, each including a plurality of different materials.

[0322] The electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth metal complex, a rare earth metal complex, or a combination thereof.

[0323] The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one embodiment, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs.

[0324] The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

[0325] The rare earth metal may be selected from Sc, Y, Ce, Yb, Gd, and Tb.

[0326] The alkali metal compound, the alkaline earth metal compound, and the rare earth metal compound may each independently be selected from oxides and halides (e.g., fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth metal, and the rare earth metal, respectively.

[0327] The alkali metal compound may be selected from alkali metal oxides, such as Li_2O , Cs_2O , or K_2O , and alkali metal halides, such as LiF, NaF, CsF, KF, LiI, NaI, CsI, KI, or RbI. In one embodiment, the alkali metal compound may be selected from LiF, Li_2O , NaF, LiI, NaI, CsI, and KI.

[0328] The alkaline earth metal compound may be selected from alkaline earth metal compounds, such as BaO, SrO, CaO, $\text{Ba}_{1-x}\text{Sr}_x\text{O}$ (wherein $0 < x < 1$), and $\text{Ba}_x\text{Ca}_{1-x}\text{O}$ (wherein $0 < x < 1$). In one embodiment, the alkaline earth metal compound may be selected from BaO, SrO, and CaO.

[0329] The rare earth metal compound may be selected from YbF_3 , ScF_3 , ScO_3 , Y_2O_3 , Ce_2O_3 , GdF_3 , and TbF_3 . In one embodiment, the rare earth metal compound may be selected from YbF_3 , ScF_3 , TbF_3 , YbI_3 , ScI_3 , and TbI_3 .

[0330] The alkali metal complex, the alkaline earth metal complex, and the rare earth metal complex may each include ions of the above-described alkali metal, alkaline earth metal, and rare earth metal. Each ligand coordinated with the metal ion of the alkali metal complex, the alkaline earth metal complex, and the rare earth metal complex may independently be selected from a hydroxyquinoline, a hydroxyisoquinoline, a hydroxybenzoquinoline, a hydroxy-acridine, a hydroxyphenanthridine, a hydroxyphenyl oxazole, a hydroxyphenyl thiazole, a hydroxydiphenyl oxadiazole, a hydroxydiphenyl thiadiazole, a hydroxyphenyl pyridine, a hydroxyphenyl benzimidazole, a hydroxyphenyl benzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene.

[0331] The electron injection layer may consist of an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth metal complex, a rare earth metal complex, or a combination thereof, as described above. In some embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, the alkali metal, the alkaline earth metal, the rare earth metal, the alkali metal compound, the alkaline earth metal compound, the rare earth metal compound, the alkali metal complex, the alkaline earth metal complex, the rare earth metal complex, or a combination thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

[0332] The thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, e.g., about 3 Å to about 90 Å. When the thickness of the electron injection layer is within any of these ranges, excellent electron injection characteristics may be obtained without a substantial increase in driving voltage.

[0333] The second electrode 190 may be disposed on the organic layer 150. The second electrode 190 may be a cathode that is an electron injection electrode. A material for forming the second electrode 190 may be a material with a low work function, for example, a metal, an alloy, an electrically conductive compound, or a mixture thereof.

[0334] The second electrode 190 may include at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO. The second electrode 190 may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

[0335] The second electrode 190 may have a single-layered structure, or a multi-layered structure including two or more layers.

[0336] Referring to FIG. 2, an organic light-emitting device 20 has a first capping layer 210, the first electrode 110, the organic layer 150, and the second electrode 190 structure, wherein the layers are sequentially stacked in this stated order. Referring to FIG. 3, an organic light-emitting device 30 has the first electrode 110, the organic layer 150, the second electrode 190, and a second capping layer 220 structure, wherein the layers are sequentially stacked in this stated order. Referring to FIG. 4, an organic light-emitting device 40 has the first capping layer 210, the first electrode 110, the organic layer 150, the second electrode 190, and the second capping layer 220 structure, wherein the layers are stacked in this stated order.

[0337] The first electrode 110, the organic layer 150, and the second electrode 190 illustrated in FIGS. 2 to 4 may be substantially the same as those illustrated in FIG. 1.

[0338] In the organic light-emitting devices 20 and 40, light emitted from the emission layer in the organic layer 150 may pass through the first electrode 110 (which may be a semi-transmissive electrode or a transmissive electrode) and through the first capping layer 210 to the outside. In the organic light-emitting devices 30 and 40, light emitted from the emission layer in the organic layer 150 may pass through the second electrode 190 (which may be a semi-transmissive electrode or a transmissive electrode) and through the second capping layer 220 to the outside.


[0339] The first capping layer 210 and the second capping layer 220 may improve the external luminous efficiency based on the principle of constructive interference.

[0340] The first capping layer 210 and the second capping layer 220 may each independently be a capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.

[0341] At least one selected from the first capping layer 210 and the second capping layer 220 may each independently include at least one material selected from carbocyclic compounds, heterocyclic compounds, amine-based compounds, porphine derivatives, phthalocyanine derivatives, naphthalocyanine derivatives, alkali metal complexes, and alkaline earth metal complexes. The carbocyclic compound, the heterocyclic compound, and the amine-based compound may optionally be substituted with a substituent containing at least one element selected from O, N, S, Se, Si, F, Cl, Br, and I. In one embodiment, at least one of the first capping layer 210 and the second capping layer 220 may each independently include an amine-based compound.


[0342] In one or more embodiments, at least one of the first capping layer 210 and the second capping layer 220 may each independently include a compound represented by Formula 201 or 202.

[0343] In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include a compound selected from Compounds HT28 to HT33 and Compound CP1 to CP5:



-continued

CP5



[0344] Layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region may be formed in a certain region by using one or more suitable methods such as vacuum deposition, spin coating, casting, Langmuir-blodgett (LB) deposition, ink-jet printing, laser printing, and laser-induced thermal imaging.

[0345] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are each independently formed by vacuum deposition, the vacuum deposition may be performed at a temperature in a range of about 100° C. to about 500° C. at a vacuum degree in a range of about 10⁻⁸ torr to about 10⁻³ torr, and/or at a rate in a range of about 0.01 Angstroms per second (Å/sec) to about 100 Å/sec, depending on the compound to be included in each layer and the structure of each layer to be formed.

[0346] When layers constituting the hole transport region, the emission layer, and layers constituting the electron transport region are each independently formed by spin coating, the spin coating may be performed at a coating rate of about 2,000 revolutions per minute (rpm) to about 5,000 rpm, and at a heat treatment temperature of about 80 to about 200° C., depending on the compound to be included in each layer and the structure of each layer to be formed.

[0347] The term “C₁-C₆₀ alkyl group” as used herein refers to a linear or branched aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms. Examples thereof include a methyl group, an ethyl group, a propyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “C₁-C₆₀ alkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₆₀ alkyl group.

[0348] The term “C₂-C₆₀ alkenyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon double bond in the middle or at the terminus of the C₂-C₆₀ alkyl group. Examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C₂-C₆₀ alkenylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkenyl group.

[0349] The term “C₂-C₆₀ alkynyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the middle or at the terminus of the C₂-C₆₀ alkyl group. Examples thereof include an ethynyl group and a propynyl group. The term “C₂-C₆₀ alkynylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

[0350] The term “C₁-C₆₀ alkoxy group” as used herein refers to a monovalent group represented by —OA₁₀₁ (where A₁₀₁ is a C₁-C₆₀ alkyl group). Examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

[0351] The term “C₃-C₁₀ cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group including 3 to 10 carbon atoms. Examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

[0352] The term “C₁-C₁₀ heterocycloalkyl group” as used herein refers to a monovalent monocyclic group including at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms. Examples thereof include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuranyl group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

[0353] The term “C₃-C₁₀ cycloalkenyl group” as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one double bond in its ring, and is not aromatic. Examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

[0354] The term “C₁-C₁₀ heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group including at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one double bond in its ring. Examples of the C₁-C₁₀ heterocycloalkenyl group include a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranyl group, and a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

[0355] The term “C₆-C₆₀ aryl group” as used herein refers to an aromatic monovalent group having 6 to 60 carbon atoms. The term “C₆-C₆₀ arylene group” as used herein refers to an aromatic divalent group having 6 to 60 carbon atoms. Examples of the C₆-C₆₀ aryl group include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each independently include two or more rings, the respective rings may be fused.

[0356] The term “C₁-C₆₀ heteroaryl group” as used herein refers to a monovalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 60 carbon atoms. The term “C₁-C₆₀ heteroarylene group” as used herein refers to a divalent group having a heterocyclic aromatic system including at least one heteroatom selected from N, O, P, and S as a ring-forming atom and 1 to 60 carbon atoms. Examples of the C₁-C₆₀ heteroaryl group include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each independently include two or more rings, the respective rings may be fused.

[0357] The term “C₆-C₆₀ aryloxy group” as used herein refers to a group represented by —OA₁₀₂ (where A₁₀₂ is a C₆-C₆₀ aryl group). The term “C₆-C₆₀ arylthio group” as used herein refers to a group represented by —SA₁₀₃ (where A₁₀₃ is a C₆-C₆₀ aryl group).

[0358] The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group that has two or more rings condensed and only carbon atoms (e.g., 8 to 60 carbon atoms) as ring forming atoms, wherein the entire molecular structure is non-aromatic. An example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0359] The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group that has two or more condensed rings and at least one heteroatom selected from N, O, Si, P, and S, in addition to carbon atoms (for example, 1 to 60 carbon atoms), as a ring-forming atom, wherein the entire molecular structure is non-aromatic. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0360] The term “C₅-C₆₀ carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms only as ring-forming atoms. The term “C₅-C₆₀ carbocyclic group” as used herein refers to an aromatic carbocyclic group or a non-aromatic carbocyclic group. The term “C₅-C₆₀ carbocyclic group” as used herein refers to a ring (such as a benzene group), a monovalent group (such as a phenyl group), or a divalent group (such as a phenylene group). In one or more embodiments, depending on the number of substituents connected to the C₅-C₆₀ carbocyclic group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

[0361] The term “C₁-C₆₀ heterocyclic group” as used herein refers to a group having the same structure as the C₁-C₆₀ carbocyclic group, except that at least one heteroatom selected from N, O, Si, P, and S is used as a ring-forming atom, in addition to carbon atoms (e.g., 1 to 60 carbon atoms).

[0362] In the present specification, at least one substituent of the substituted C₅-C₆₀ carbocyclic group, substituted C₁-C₆₀ heterocyclic group, substituted C₃-C₁₀ cycloalkylene group, substituted C₁-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₁-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₁-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, a substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent

non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0363] deuterium (-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0364] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

[0365] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0366] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

[0367] —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂).

[0368] wherein Q₁ to Q₆, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

[0369] The term “Ph” as used herein refers to a phenyl group. The term “Me” as used herein refers to a methyl group. The term “Et” as used herein refers to an ethyl group. The term “ter-Bu” or “Bu” as used herein refers to a tert-butyl group. The term “OMe” as used herein refers to a methoxy group.

[0370] The term “biphenyl group” as used herein refers to a phenyl group substituted with a phenyl group. In other words, the “biphenyl group” may be a substituted phenyl group having a C₆-C₆₀ aryl group as a substituent.

[0371] The term “terphenyl group” as used herein refers to a phenyl group substituted with a biphenyl group. In other words, the “terphenyl group” may be a substituted phenyl group having a C₆-C₆₀ aryl group substituted with a C₆-C₆₀ aryl group as a substituent.

[0372] The symbols * and *' as used herein, unless defined otherwise, refer to a binding site to an adjacent atom in a corresponding formula.

[0373] Hereinafter a compound and an organic light-emitting device according to one or more embodiments will be described in detail with reference to Synthesis Examples and Examples. The expression “B was used instead of A” used in describing Synthesis Examples means that an identical molar equivalent of B was used in place of A.


[0374] The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples.



EXAMPLES

Synthesis Examples

Synthesis Example 1: Synthesis of Compound 1

[0375]





Synthesis of Intermediate 1-1

[0376] 2,5-dibromopyridine was dissolved in tetrahydrofuran (THF). Isopropyl magnesium chloride was added to

the solution, and the solution was stirred to obtain a Grignard reagent. Dichlorophenyl phosphine (PPhCl_2) was added to the Grignard reagent, and then the mixture was stirred to obtain Intermediate 1-1. Intermediate 1-1 was identified by liquid chromatography-mass spectrometry (LC-MS).

[0377] $\text{C}_{16}\text{H}_{11}\text{Br}_2\text{N}_2\text{P}$: M+1 420.8

Synthesis of Intermediate 1-2

[0378] Intermediate 1-1 was oxidized using a hydrogen peroxide solution to obtain Intermediate 1-2. Intermediate 1-2 was identified by LC-MS.

[0379] $\text{C}_{16}\text{H}_{11}\text{Br}_2\text{N}_2\text{OP}$: M+1 436.8

Synthesis of Compound 1


[0380] 5.0 g of Intermediate 1-2, 5.0 g of 9,9-dimethyl-9,10-dihydroacridine (CAS No. 6267-02-3), 0.10 g of palladium (II) acetate, 0.24 g of triphenylphosphine, and 1.65 g of NaOtBu were dissolved in 50 mL of toluene, and then the resulting solution was refluxed for 3 hours. The obtained reaction solution was cooled to ambient temperature, an organic layer was extracted using ethyl acetate, and then the organic layer was washed with water. The obtained organic layer was dried by using magnesium sulfate (MgSO_4), a solvent was removed therefrom by evaporation, and the obtained residue was separated and purified through silica gel chromatography to thereby obtain 4.9 g of Compound 1 (yield: 62%). Compound 1 was identified by LC-MS and ^1H -NMR.

[0381] $\text{C}_{46}\text{H}_{39}\text{N}_4\text{OP}$: M+1 695.3

[0382] ^1H NMR (500 MHz, CDCl_3) δ =7.94 (d, 2H), 7.80 (s, 2H), 7.72 (d, 2H), 7.51-7.42 (m, 3H), 7.21 (d, 2H), 7.21-7.11 (m, 12H), 6.99 (t, 4H), 1.62 (s, 12).

Synthesis Example 2: Synthesis of Compound 4

[0383]



Synthesis of Intermediate 4-1

[0384] 2,5-dibromopyridine was dissolved in THF, isopropyl magnesium chloride was added to the solution, and the solution was stirred to obtain a Grignard reagent. Dichlorophenyl phosphine was added to the Grignard reagent, 4-bromophenyl lithium was added to the mixture, and then the mixture was stirred continuously to obtain Intermediate 4-1. Intermediate 4-1 was identified by LC-MS.

[0385] $C_{17}H_{13}BrNP$: M+1 341.8

Synthesis of Intermediate 4-2

[0386] Intermediate 4-1 was oxidized using a hydrogen peroxide solution to obtain Intermediate 4-2. Intermediate 4-2 was identified by LC-MS.

[0387] $C_{17}H_{13}BrNOP$: M+1 357.9

Synthesis of Compound 4




[0388] 5.1 g of Compound 4 was obtained (yield: 52%) in substantially the same manner as in Synthesis of Compound 1, except that Intermediate 4-2 was used instead of Intermediate 1-2. Compound 4 was identified by LC-MS and 1H -NMR.

[0389] $C_{32}H_{27}N_2OP$: M+1 487.19


[0390] 1H NMR (500 MHz, $CDCl_3$) δ =9.14 (s, 1H), 8.77 (d, 1H), 8.28 (d, 1H), 7.75-7.70 (m, 2H), 7.52 (d, 1H), 7.21-7.11 (m, 6H), 6.93 (t, 2H), 1.67 (s, 6H).

Synthesis Example 3: Synthesis of Compound 5


[0391]



5-2



5-3



5

Synthesis of Intermediate 5-1

[0392] 2-bromo-9,9-dimethyl-9,10-dihydroacridine (CAS No. 1443680-94-1) was reacted with tosyl chloride (TsCl) under basic conditions to obtain Intermediate 5-1. Intermediate 5-1 was identified by LC-MS.

[0393] $C_{22}H_{20}BrNO_2S$: M+1 442.1

Synthesis of Intermediate 5-2

[0394] Intermediate 5-1 and carbazole was reacted using a copper catalyst to obtain Intermediate 5-2. Intermediate 5-2 was identified by LC-MS.

[0395] $C_{34}H_{28}N_2O_2S$: M+1 529.3

Synthesis of Intermediate 5-3

[0396] Under an acidic condition, the protective groups of Compound 5-2 were removed therefrom to obtain Intermediate 5-3. Intermediate 5-3 was identified by LC-MS.

[0397] $C_{27}H_{22}N_2$: M+1 375.2

Synthesis of Compound 5


[0398] 3.9 g of Intermediate 5-3, 3.7 g of Intermediate 4-2, 0.09 g of palladium (II) acetate, 0.22 g of triphenyl phosphine, and 1.00 g of NaOtBu were dissolved in 50 mL of toluene, and then the resulting solution was refluxed for 3 hours. The obtained reaction solution was cooled to ambient temperature, an organic layer was extracted using ethyl acetate, and then the organic layer was washed with water. The obtained organic layer was dried by using $MgSO_4$, a solvent was removed therefrom by evaporation, and the obtained residue was separated and purified through silica gel chromatography to thereby obtain 3.9 g of Compound 5 (yield: 58%). Compound 5 was identified by LC-MS and 1H -NMR.

[0399] $C_{44}H_{34}N_3OP$: M+1 652.3


[0400] 1H NMR (500 MHz, $CDCl_3$) δ =9.17 (s, 1H), 8.77 (d, 1H), 8.54 (d, 2H), 8.24 (d, 1H), 7.93 (d, 2H), 7.75-7.70 (m, 3H), 7.56-7.49 (m, 3H), 7.38-7.31 (m, 5H), 7.19-7.15 (m, 4H), 6.89 (t, 1H).

Synthesis Example 4: Synthesis of Compound 11

[0401]



-continued



Synthesis of Intermediate 11-1

[0402] Intermediate 11-1 was obtained in substantially the same manner as in Synthesis of Intermediate 1-1, except that 3,6-dibromo-2-methyl pyridine was used instead of 2,5-dibromopyridine. Intermediate 11-1 was identified by LC-MS.

[0403] $C_{18}H_{15}Br_2N_2P$: M+1 448.9

Synthesis of Intermediate 11-2

[0404] Intermediate 11-1 was oxidized using a hydrogen peroxide solution to obtain Intermediate 11-2. Intermediate 11-2 was identified by LC-MS.

[0405] $C_{18}H_{15}Br_2N_2OP$: M+1 464.9

Synthesis of Compound 11


[0406] 5.2 g of Compound 11 was obtained (yield: 63%) in substantially the same manner as in Synthesis of Compound 1, except that Intermediate 11-2 was used instead of Intermediate 1-2. Compound 11 was identified by LC-MS and 1H -NMR.

[0407] $C_{48}H_{43}N_4OP$: M+1 723.3


[0408] 1H NMR (500 MHz, $CDCl_3$) δ =7.78 (d, 4H), 7.52-7.49 (m, 3H), 7.27 (d, 2H), 7.21-7.11 (m, 12H), 6.92 (t, 4H), 2.51 (s, 3H), 1.69 (s, 12H).

Synthesis Example 5: Synthesis of Compound 16

[0409]



-continued



Synthesis of Intermediate 16-1

[0410] Intermediate 16-1 was obtained in substantially the same manner as in Synthesis of Intermediate 4-1, except that 3,6-dibromo-2-methyl pyridine was used instead of 2,5-dibromopyridine. Intermediate 16-1 was identified by LC-MS.

[0411] $C_{18}H_{15}BrNOP$: M+1 372.2

Synthesis of Compound 16


[0412] 5.7 g of Compound 16 was obtained (yield: 71%) in substantially the same manner as in Synthesis of Compound 4, except that Intermediate 16-1 was used instead of Intermediate 4-1. Compound 16 was identified by LC-MS and 1H -NMR.

[0413] $C_{48}H_{42}N_3OP$: M+1 708.7


[0414] 1H NMR (500 MHz, $CDCl_3$) δ =7.78-7.72 (m, 5H), 7.52-7.49 (m, 3H), 7.37-7.14 (m, 15H), 6.94 (t, 4H), 2.51 (s, 3H), 1.67 (s, 12H).

Synthesis Example 6: Synthesis of Compound 20

[0415]



-continued



Synthesis of Intermediate 20-1

[0416] 3,6-dibromo-2-methyl pyridine was dissolved in THF, isopropyl magnesium chloride (iPrMgCl) was added dropwise thereto, and the mixture was stirred to obtain a Grignard reagent. Diphenyl chlorophosphine (PPh_2Cl) was added to the Grignard reagent, and then the mixture was stirred to obtain Intermediate 20-1. Intermediate 20-1 was identified by LC-MS.

[0417] $C_{18}H_{15}BrNP$: M+1 356.1

Synthesis of Intermediate 20-2

[0418] Intermediate 20-1 was oxidized using a hydrogen peroxide solution to obtain Intermediate 20-2. Intermediate 20-2 was identified by LC-MS.

[0419] $C_{18}H_{15}BrNOP$: M+1 372.3

Synthesis of Compound 20


[0420] 2.2 g of Intermediate 20-2, 1.2 g of 9,9-dimethyl-9,10-dihydroacridine, 0.053 g of palladium (II) acetate, 0.12 g of triphenyl phosphine, and 0.85 g of NaOtBu were dissolved in 30 mL of toluene, and then the resulting solution was refluxed for 3 hours. The obtained reaction solution was cooled to ambient temperature, an organic layer was extracted using ethyl acetate, and then the organic layer was washed with water. The obtained organic layer was dried by using $MgSO_4$, a solvent was removed therefrom by evaporation, and the obtained residue was separated and purified through silica gel chromatography to thereby obtain 2.3 g of Compound 20 (yield: 77%). Compound 20 was identified by LC-MS and 1H -NMR.

[0421] $C_{33}H_{29}N_2OP$: M+1 501.5

[0422] 1H NMR (500 MHz, $CDCl_3$) δ =7.80-7.71 (m, 6H), 7.51-7.49 (m, 6H), 7.28 (d, 2H), 7.21-7.13 (m, 6H), 6.94 (t, 2H), 2.54 (s, 3H), 1.62 (s, 6H).

Synthesis Example 7: Synthesis of Compound 27

[0423]



Synthesis of Compound 27


[0424] 5.2 g of Intermediate 4-1, 5.5 g of Intermediate 27-1 (CAS No. 1821228-18-5), 0.13 g of palladium (II) acetate, 0.31 g of triphenyl phosphine, and 1.4 g of NaOtBu were dissolved in 80 mL of toluene, and then the resulting solution was refluxed for 3 hours. The obtained reaction solution was cooled to ambient temperature, an organic layer was extracted using ethyl acetate, and then the organic layer was washed with water. The obtained organic layer was dried by using MgSO_4 , a solvent was removed therefrom by evaporation, and the obtained residue was separated and purified through silica gel chromatography to thereby obtain 4.7 g of Compound 27 (yield: 49%). Compound 27 was identified by LC-MS and ^1H -NMR.

[0425] $\text{C}_{44}\text{H}_{34}\text{N}_3\text{OP}$: M+1 652.3

[0426] ^1H NMR (500 MHz, CDCl_3) δ =9.13 (s, 1H), 8.75 (d, 1H), 8.54 (d, 1H), 8.29 (d, 1H), 7.98-7.90 (m, 9H), 7.58-7.50 (m, 2H), 7.38-7.31 (m, 3H), 7.21-7.11 (m, 7H), 6.94 (t, 2H), 1.71 (s, 6H).

Synthesis Example 8: Synthesis of Compound 28

[0427]



Synthesis of Intermediate 28-1

[0428] 3-bromopyridine was dissolved in THF, isopropyl magnesium chloride ($i\text{PrMgCl}$) was added dropwise thereto, and the mixture was stirred to obtain a Grignard reagent. The Grignard reagent was allowed to react with phenyl dichlorophosphine, a (4-bromo-3-methylphenyl) lithium solution

(CAS No. 17989-26-3) was added dropwise thereto, and then the mixture was stirred to obtain Intermediate 28-1. Intermediate 28-1 was identified by LC-MS.

[0429] $C_{18}H_{15}BrNP$ M+1: 356.1

Synthesis of Intermediate 28-2

[0430] Intermediate 28-1 was oxidized using a hydrogen peroxide solution to obtain Intermediate 28-2. Intermediate 28-2 was identified by LC-MS.

[0431] $C_{18}H_{15}BrNOP$ M+1 372.1

Synthesis of Compound 28

[0432] 3.2 g of Compound 28 was obtained (yield: 43%) in substantially the same manner as in Synthesis of Com-



pound 4, except that Intermediate 28-2 was used instead of Intermediate 4-2. Compound 28 was identified by LC-MS and 1H -NMR.

[0433] $C_{45}H_{36}N_3OP$ M+1 666.4

[0434] 1H NMR (500 MHz, $CDCl_3$) δ =8.74 (d, 1H), 8.54 (d, 1H), 8.26 (d, 1H), 7.94 (d, 1H), 7.89-7.72 (m, 6H), 7.55-7.49 (m, 5H), 7.38-7.12 (m, 10H), 6.92 (t, 2H), 1.89 (s, 3H), 1.69 (s, 6H).

Synthesis Example 10: Synthesis of Compound 55

[0435]



Synthesis of Intermediate 55-2

[0436] 10 g of 2,6-dibromopyridine, 4.4 g of 9,9-dimethyl-9,10-dihydroacridine, 0.14 g of palladium (II) acetate, 0.33 g of triphenyl phosphine, and 3.0 g of NaOtBu were dissolved in 200 mL of toluene, and then the resulting solution was refluxed for 3 hours. The obtained reaction solution was cooled to ambient temperature, an organic layer was extracted using ethyl acetate, and then the organic layer was washed with water. The obtained organic layer was dried by using MgSO_4 , a solvent was removed therefrom by evaporation, and the obtained residue was separated and purified through silica gel chromatography to thereby obtain 3.2 g of Intermediate 55-2 (yield: 42%). Intermediate 55-2 was identified by LC-MS.

[0437] $\text{C}_{20}\text{H}_{17}\text{BrN}_2$; M+1 365.1

Synthesis of Compound 55

[0438] 3.0 g of Intermediate 55-2, 3.7 g of Intermediate 55-1 (CAS No. 328917-29-9), 0.38 g of $\text{Pd}(\text{PPh}_3)_4$, 2.8 g of


potassium carbonate, 40 mL of toluene, 10 mL of ethanol, and 10 mL of water were added to a round-bottom flask, and then the mixture was refluxed overnight. The obtained reaction solution was cooled to ambient temperature, an organic layer was extracted using ethyl acetate, and then the organic layer was washed with water. The obtained organic layer was dried by using MgSO_4 , a solvent was removed therefrom by evaporation, and the obtained residue was separated and purified through silica gel chromatography to thereby obtain 2.9 g of Compound 55 (yield: 62%). Compound 55 was identified by LC-MS and ^1H -NMR.


[0439] $\text{C}_{38}\text{H}_{31}\text{N}_2\text{OP}$; M+1 563.2

[0440] ^1H NMR (500 MHz, CDCl_3) δ =8.53 (s, 1H), 8.49 (d, 1H), 7.88 (d, 1H), 7.79-7.72 (m, 5H), 7.52-7.49 (m, 6H), 7.21-7.02 (m, 7H), 6.92 (t, 2H), 6.36 (d, 1H), 6.21 (d, 1H), 1.67 (s, 6H).

Synthesis Example 11: Synthesis of Compound 61

[0441]





Synthesis of Intermediate 61-1

[0442] 9-(3,5-dibromophenyl)-9H-carbazole (CAS No. 750573-26-3) was reacted with 9,9-dimethyl-9,10-dihydroacridine using a palladium catalyst to obtain Intermediate 61-1. Intermediate 61-1 was identified by LC-MS.

[0443] C₃₃H₂₅BrN₂ M+1: 529.1

Synthesis of Intermediate 61-2

[0444] Intermediate 61-1 was reacted with n-BuLi, and B(OMe)₃ was added dropwise thereto. Then, a purification process was performed thereon to obtain Intermediate 61-2. Intermediate 61-2 was identified by LC-MS.

[0445] C₃₃H₂₇BN₂O₂; M+1 495.3

Synthesis of Intermediate 61-3

[0446] 2,6-dibromopyridine was dissolved in THF, isopropyl magnesium chloride (iPrMgCl) was added dropwise thereto, and the mixture was stirred to obtain a Grignard reagent. The Grignard reagent was reacted with chlorodiphenylphosphine to obtain Intermediate 61-3. Intermediate 61-3 was identified by LC-MS.

[0447] C₁₇H₁₃BrNP M+1: 342.4

Synthesis of Intermediate 61-4

[0448] Intermediate 61-3 was oxidized using a hydrogen peroxide solution to obtain Intermediate 61-4. Intermediate 61-4 was identified by LC-MS.

[0449] C₁₇H₁₃BNOP; M+1 357.9

Synthesis of Compound 61

[0450] 3.7 g of Intermediate 61-4, 3.7 g of Intermediate 61-2 (CAS No. 328917-29-9), 0.17 g of Pd(PPh₃)₄, 3.6 g of potassium carbonate, 45 mL of toluene, 15 mL of ethanol, and 15 mL of water were added to a round-bottom flask, and then the mixture was refluxed overnight. The obtained reaction solution was cooled to ambient temperature, an organic layer was extracted using ethyl acetate, and then the organic layer was washed with water. The obtained organic layer was dried by using MgSO₄, a solvent was removed therefrom by evaporation, and the obtained residue was separated and purified through silica gel chromatography to thereby obtain 4.1 g of Compound 61 (yield: 55%). Compound 61 was identified by LC-MS and ¹H-NMR.



[0451] C₅₀H₃₈N₃OP; M+1 728.31

[0452] ¹H NMR (500 MHz, CDCl₃) δ=8.57 (d, 2H), 8.22 (s, 1H), 7.92 (d, 2H), 7.78-7.71 (m, 4H), 7.61 (s, 1H), 7.58

(t, 11H), 7.53-7.48 (m, 7H), 7.41-7.24 (m, 4H), 7.21-7.11 (m, 8H), 6.92 (t, 2H), 1.68 (s, 6H).

Synthesis Example 12: Synthesis of Compound 71

[0453]



Synthesis of Intermediate 71-1

[0454] 1,3,5-tribromobenzene was reacted with n-BuLi, diphenylchlorophosphine was added dropwise thereto, and

then the mixture was stirred to obtain Intermediate 71-1. Intermediate 71-1 was identified by LC-MS.

[0455] $C_{18}H_{13}Br_2OP$: M+1 433.91

Synthesis of Compound 71



[0456] 1.9 g of Intermediate 71-1, 1.8 g of 9,9-dimethyl-9,10-dihydroacridine (CAS No. 6267-02-3), 0.04 g of palladium (II) acetate, 0.09 g of triphenyl phosphine, and 0.63 g of NaOtBu were dissolved in 30 mL of toluene, and then the resulting solution was refluxed for 3 hours. The obtained reaction solution was cooled to ambient temperature, an organic layer was extracted using ethyl acetate, and then the organic layer was washed with water. The obtained organic layer was dried by using $MgSO_4$, a solvent was removed therefrom by evaporation, and the obtained residue was separated and purified through silica gel chromatography to thereby obtain 2.5 g of Compound 71 (yield: 82%). Compound 71 was identified by LC-MS and 1H -NMR.

[0457] $C_{48}H_{41}N_2OP$: M+1 693.41


[0458] 1H NMR (500 MHz, $CDCl_3$) δ =7.79-7.72 (m, 4H), 7.53-7.42 (m, 6H), 7.23 (s, 2H), 7.19-7.08 (m, 15H), 6.92 (t, 4H), 1.65 (s, 12H).

Synthesis Example 13: Synthesis of Compound 72

[0459]



-continued



Synthesis of Intermediate 72-1

[0460] 1,4-tribromobenzene was reacted with n-BuLi, dichlorophenylphosphine was added dropwise thereto, and then the mixture was stirred to obtain Intermediate 71-1. Intermediate 71-1 was identified by LC-MS.

[0461] $C_{18}H_{13}Br_2P$: M+1 418.8

Synthesis of Intermediate 72-2

[0462] Intermediate 72-1 was oxidized using a hydrogen peroxide solution to obtain Intermediate 72-2.

[0463] $C_{18}H_{13}Br_2OP$: M+1 434.7

Synthesis of Compound 72

[0464] 2.44 g of Compound 72 was obtained (yield: 81%) in substantially the same manner as in Synthesis of Compound 71, except that Intermediate 72-2 was used instead of Intermediate 71-1.

[0465] $C_{48}H_{41}N_2OP$: M+1 693.4

[0466] 1H NMR (500 MHz, $CDCl_3$) δ =7.81-7.70 (m, 6H), 7.53-7.48 (m, 3H), 7.41-7.34 (m, 4H), 7.21-7.11 (m, 12H), 6.95 (t, 4H), 1.71 (s, 12H).

Example 1


[0467] A Corning 15 ohms per square centimeter (Ω/cm^2) (1,200 Å thickness) ITO glass substrate was cut to a size of 50 millimeters (mm)×50 mm×0.7 mm, sonicated in isopropyl alcohol and pure water for 5 minutes each, and cleaned by exposure to ultraviolet rays and ozone, so that the resulting ITO glass substrate could be used as a substrate and an anode. Then, the glass substrate was mounted on a vacuum deposition apparatus.

[0468] NPD was vacuum-deposited on the ITO anode formed on the substrate to form a hole injection layer having a thickness of 300 Å, and mCP was vacuum-deposited as a hole transporting compound on the hole injection layer to form a hole transport layer having a thickness of 200 Å.


[0469] Compound 1 and FIrpic were co-deposited in a weight ratio of 92:8 on the hole transport layer to form an emission layer having a thickness of 250 Å.

[0470] Next, TAZ was deposited on the emission layer to form an electron transport layer having a thickness of 200 Å, LiF was deposited on the electron transport layer to form an


electron injection layer having a thickness of 10 Å, and aluminum was vacuum-deposited on the electron injection layer to form a LiF/Al cathode having a thickness of 100 Å, thereby completing the manufacture of an organic light-emitting device.




NPB




mCP



BCPO



Flrpic



TAZ

Examples 2 to 7 and Comparative Examples 1 to 3


[0471] Organic light-emitting devices were manufactured in substantially the same manner as in Example 1, except that the compounds shown in Table 1 were used instead of Compound 1 in forming each emission layer.

Evaluation Example 1



[0472] The driving voltage, current density, efficiency, and emission color of the organic light-emitting devices manufactured in Examples 1 to 7 and Comparative Examples 1 to 3 were measured. The results thereof are shown in Table 1.

TABLE 1

	Emission layer	Driving voltage (V)	Current density (mA/cm ²)	Maximum quantum efficiency (%)	Emission color
Example 1	1	3.0	1	24.4	blue
Example 2	5	2.7	1	28.5	blue
Example 3	27	3.1	1	20.2	blue
Example 4	55	2.7	1	27.3	blue
Example 5	61	2.9	1	28.8	blue
Example 6	71	3.2	1	23.9	blue
Example 7	72	3.1	1	22.7	blue
Comparative Example 1	BCPO	3.5	1	19.2	blue
Comparative Example 2	A	3.4	1	20.1	blue
Comparative Example 3	B	3.1	1	19.9	blue



BCPO



B

[0473] Referring to Table 1, the organic light-emitting devices of Examples 1 to 7 were found to have low driving voltage and excellent efficiency, as compared with the organic light-emitting devices of Comparative Examples 1 to 3.






Example 8

[0474] A Corning 15 Ω/cm^2 (1,200 Å thickness) ITO glass substrate was cut to a size of 50 mm×50 mm×0.7 mm, sonicated in isopropyl alcohol and pure water for 5 minutes each, and cleaned by exposure to ultraviolet rays and ozone, so that the resulting ITO glass substrate could be used as a substrate and an anode. Then, the glass substrate was mounted on a vacuum deposition apparatus.

[0475] NPD was vacuum-deposited on the ITO anode formed on the substrate to form a hole injection layer having a thickness of 300 Å, and TCTA was vacuum-deposited as a hole transporting compound on the hole injection layer to form a hole transport layer having a thickness of 200 Å.


[0476] A hole transport layer compound CzSi was vacuum-deposited on the hole transport layer to a thickness of 100 Å, and then DPEPO and Compound 4 were co-deposited in a weight ratio of 90:10 to form an emission layer having a thickness of 200 Å.

[0477] Next, DPEPO was deposited on the emission layer to form an electron transport layer having a thickness of 200 Å, TPBI was deposited on the electron transport layer to a thickness of 300 Å, LiF was deposited thereon to a thickness of 10 Å to form an electron injection layer, and aluminum was vacuum-deposited on the electron injection layer to form a LiF/Al cathode having a thickness of 3,000 Å, thereby completing the manufacture of an organic light-emitting device.



What is claimed is:

1. A heterocyclic compound represented by Formula 1:



Formula 1


wherein, in Formula 1,

Ar₁ to Ar₃ are each independently selected from a substituted or unsubstituted C₅-C₆₀ carbocyclic group and a substituted or unsubstituted C₂-C₆₀ heterocyclic group, two adjacent groups of Ar₁ to Ar₃ are separate or are bound to form a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group,

R₁ to R₃ are each independently represented by the following Formula 2,

b1 to b3 are each independently an integer of 0 to 3,

b1 to b3 satisfy b1+b2+b3≥1; or when Ar₁ to Ar₃ are each a substituted or unsubstituted benzene group, b1+b2+b3≥2,



Formula 2

wherein, in Formula 2,

A₁ and A₂ are each independently selected from a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, and a pyrimidine group,

L₁₁ is selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

a11 is an integer of 0 to 3; when a11 is 2 or 3, the 2 or 3 L₁₁(s) are identical to or different from each other,

Y is selected from C(R₁₇)(R₁₈), sulfur (S), Si(R₁₇)(R₁₈), S(=O), S(=O)₂, and P(=O)(R₁₇),

R₁₇ to R₂₀ are each independently selected from hydrogen,

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl

group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —S(=O)(Q₁), and —P(=O)(Q₁)(Q₂), and

b19 and b20 are each independently an integer of 0 to 8,

wherein, in Formulae 1 and 2, at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, substituted C₁-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₁-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₁-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀

aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

wherein Q₁, Q₂, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

* indicates a binding site to an adjacent atom.

2. The heterocyclic compound as claimed in claim 1, wherein Ar₁ to Ar₃ are each independently selected from:

a benzene group, a naphthalene group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a triazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, and a quinazoline group; and

a benzene group, a naphthalene group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a triazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, and a quinazoline group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a

pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphe-nyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

wherein Q₃₁ to Q₃₃ are each independently selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

3. The heterocyclic compound as claimed in claim 1, wherein at least one of Ar₁ to Ar₃ is selected from a substituted or unsubstituted π electron-depleted nitrogen-containing C₂-C₆₀ heterocyclic group.

4. The heterocyclic compound as claimed in claim 1, wherein A₁ and A₂ are each a benzene group.

5. The heterocyclic compound as claimed in claim 1, wherein L₁₁ is selected from:

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylene group, a spiro-bifluorenylene group, a spiro-benzofluorene-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentacenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, a silolylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an indolylene group, an isoindolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a benzoisoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylene group, a benzothiophenylene group, a benzosilolylene group, a benzothiazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a carbazolylenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a dibenzosilolylene group, a benzocarbazolylene group, a naphthobenzofuranylene group, a naphthobenzothiophenylene group, a naphthobenzosilolylene group, a dibenzocarbazolylene group, a dinaphthofuranylene group, a dinaphthothiophenylene group, a dinaphthosilolylene group, a thiadiazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, an oxazolopyridinylene group, a thiazolopyridinylene group, a benzonaphthyridinylene group, an aza-

fluorenylene group, an azaspiro-bifluorenylene group, an azacarbazolylene group, an azadibenzofuranylene group, an azadibenzothiophenylylene group, an azadibenzosilololylylene group, an indenopyrrolylene group, an indolopyrrolylene group, an indenocarbazolylylene group, and an indolocarbazolylylene group; and

a phenylene group, a pentalenylylene group, an indenylene group, a naphthylylene group, an azulenylylene group, an indacenylene group, an acenaphthylylene group, a fluorenylylene group, a spiro-bifluorenylylene group, a spiro-benzofluorene-fluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenalenylylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylylylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentacenylene group, a pyrrolylene group, a thiophenylylene group, a furanylylene group, a silolylylene group, an imidazolylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a pyridinylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, an indolylylene group, an isoindolylylene group, an indazolylylene group, a purinylylene group, a quinolinylylene group, an isoquinolinylylene group, a benzoquinolinylylene group, a benzoisoquinolinylylene group, a phthalazinylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinnolinylylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthrolinylylene group, a phenazinylylene group, a benzimidazolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a benzosilololylylene group, a benzothiazolylylene group, an isobenzothiazolylylene group, a benzoxazolylylene group, an isobenzoxazolylylene group, a triazolylylene group, a tetrazolylylene group, an oxadiazolylylene group, a diazinylylene group, a carbazolylylene group, a benzofuranylylene group, a dibenzothiophenylylene group, a dibenzosilololylylene group, a benzocarbazolylylene group, a naphthobenzofuranylylene group, a naphthobenzothiophenylylene group, a naphthobenzosilololylylene group, a dibenzocarbazolylylene group, a dinaphthofuranylylene group, a dinaphthothiophenylylene group, a dinaphthosilololylylene group, a thiadiazolylylene group, an imidazopyridinylylene group, an imidazopyrimidinylylene group, an oxazolopyridinylylene group, a thiazolopyridinylylene group, a benzonaphthyridinylylene group, an azafluorenylylene group, an azaspiro-bifluorenylylene group, an azacarbazolylylene group, an azadibenzofuranylylene group, an azadibenzothiophenylylene group, an azadibenzosilololylylene group, an indenopyrrolylene group, an indolopyrrolylene group, an indenocarbazolylylene group, and an indolocarbazolylylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{20} aryl group, a C_1 - C_{20} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monova-


lent non-aromatic condensed heteropolycyclic group, a methylphenyl group, a biphenyl group, and $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$,

wherein Q_{31} to Q_{33} are each independently selected from:


a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinylylene group, an isoquinolinylylene group, a quinoxalinylylene group, and a quinazolinylylene group; and

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinylylene group, an isoquinolinylylene group, a quinoxalinylylene group, and a quinazolinylylene group, each substituted with at least one selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, and a phenyl group.


6. The heterocyclic compound as claimed in claim 1, wherein L_{11} is selected from groups represented by Formula 3-1 to Formula 3-46:




Formula 3-1




Formula 3-2




Formula 3-3



Formula 3-4














Formula 3-5










Formula 3-6

-continued



-continued



Formula 3-28

Formula 3-29

Formula 3-30






Formula 3-31

Formula 3-32

Formula 3-33

Formula 3-34

-continued



Formula 3-35


Formula 3-36

Formula 3-37


Formula 3-38

Formula 3-39


-continued




Formula 3-40




Formula 3-41




Formula 3-42



Formula 3-43




Formula 3-44



Formula 3-45

-continued

Formula 3-46



wherein, in Formulae 3-1 to 3-46,

Y_1 is selected from oxygen (O), S, $C(Z_3)(Z_4)$, $N(Z_5)$, and $Si(Z_6)(Z_7)$,

Z_1 to Z_7 are each independently selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothio-phenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothio-phenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothio-phenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphtho-furanyl group, a dinaphthothio-phenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothio-phenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indenocarbazolyl group, and an indolocarbazolyl group,

d_2 is 1 or 2,

d_3 is an integer of 1 to 3,

d4 is an integer of 1 to 4,

d5 is an integer of 1 to 5,

d6 is an integer of 1 to 6,

d8 is an integer of 1 to 8, and

* and *¹ each indicate a binding site to an adjacent atom.

7. The heterocyclic compound as claimed in claim 1, wherein L_{11} is selected from groups represented by Formulae 3-2, 3-12, and 3-31.

8. The heterocyclic compound as claimed in claim 1, wherein a_{11} is 0, 1, or 2.

9. The heterocyclic compound as claimed in claim 1, wherein:

Y is selected from $C(R_{17})(R_{18})$ and $Si(R_{17})(R_{18})$, and

R_{17} and R_{18} are each independently selected from hydrogen, deuterium, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, and a tert-butyl group.

10. The heterocyclic compound as claimed in claim 1, wherein R_{19} and R_{20} are each independently selected from: hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group;

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group, each substituted with

at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group — $Si(Q_{31})(Q_{32})(Q_{33})$, and — $N(Q_{31})(Q_{32})$; and — $P(=O)(Q_1)(Q_2)$ and — $S(=O)(Q_1)$,

wherein Q_1 and Q_2 , and Q_{31} to Q_{33} are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.


11. The heterocyclic compound as claimed in claim 1, wherein b_{19} and b_{20} are each independently an integer of 0 to 4.

12. The heterocyclic compound as claimed in claim 1, wherein:


Formula 1 is represented by Formula 1-1, and

Formula 2 is represented by Formula 2-1,

Formula 1-1



Formula 2-1



wherein, in Formula 1-1,

X_{11} is nitrogen (N) or $C(R_{11})$, X_{12} is N or $C(R_{12})$, X_{13} is N or $C(R_{13})$, X_{14} is N or $C(R_{14})$, X_{15} is N or $C(R_{15})$, X_{16} is N or $C(R_{16})$, X_{21} is N or $C(R_{21})$, X_{22} is N or

C(R₂₂), X₂₃ is N or C(R₂₃), X₂₄ is N or C(R₂₄), X₂₅ is N or C(R₂₅), X₂₆ is N or C(R₂₆), X₃₁ is N or C(R₃₁), X₃₂ is N or C(R₃₂), X₃₃ is N or C(R₃₃),

R₁₁ to R₁₆, R₂₁ to R₂₆, and R₃₁ to R₃₃ are each independently selected from a group represented by Formula 2-1, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group,

when X₁₁ is C(R₁₁), X₁₂ is C(R₁₂), X₁₃ is C(R₁₃), X₁₄ is C(R₁₄), X₁₅ is C(R₁₅), X₁₆ is C(R₁₆), X₂₁ is C(R₂₁), X₂₂ is C(R₂₂), X₂₃ is C(R₂₃), X₂₄ is C(R₂₄), X₂₅ is C(R₂₅), X₂₆ is C(R₂₆), X₃₁ is C(R₃₁), X₃₂ is C(R₃₂), and X₃₃ is C(R₃₃); at least two of R₁₁, R₂₁, R₃₁, R₂₂, and R₁₂ are a group represented by Formula 2-1, or at least one of R₁, R₂₁, R₃₁, and R₂₂, and at least one of R₁₂, R₁₃, R₂₃, R₃₂, R₂₄, and R₁₄ is a group represented by Formula 2-1,

wherein R₁₂ and R₁₃, or R₁₄ and R₁₅ are bound to form a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group, and

in Formula 2-1, L, n, Y, R₁₉, R₂₀, b19, and b20 are defined the same as those of Formula 2.

13. The heterocyclic compound as claimed in claim 12, wherein

at least one selected from X₁₁ to X₁₆ is N, X₂₁ is C(R₂₁), X₂₂ is C(R₂₂), X₂₃ is C(R₂₃), X₂₄ is C(R₂₄), X₂₅ is C(R₂₅), X₂₆ is C(R₂₆), and at least one selected from X₂₁ to X₂₆ is selected from groups represented by Formula 2-1.

14. The heterocyclic compound as claimed in claim 12, wherein:

at least one selected from X₁₁ to X₁₆ is N, X₃₁ is C(R₃₁), X₃₂ is C(R₃₂), X₃₃ is C(R₃₃), and at least one selected from R₃₁ to R₃₃ is selected from groups represented by Formula 2-1.

15. The heterocyclic compound as claimed in claim 12, wherein:

at least one selected from X₂₁ to X₂₆ is N, X₃₁ is C(R₃₁), X₃₂ is C(R₃₂), X₃₃ is C(R₃₃), and at least one selected from R₃₁ to R₃₃ is selected from groups represented by Formula 2-1.





16. The heterocyclic compound as claimed in claim 12, wherein

X₂₂ is N, X₂₁ is C(R₂₁), X₂₅ is C(R₂₅), X₂₆ is C(R₂₆), and at least one selected from R₂₁, R₂₅, and R₂₆ is selected from groups represented by Formula 2-1.

17. The heterocyclic compound as claimed in claim 12, wherein


at least one selected from X₃₁ to X₃₃ is N, X₂₁ is C(R₂₁), X₂₂ is C(R₂₂), X₂₃ is C(R₂₃), X₂₄ is C(R₂₄), X₂₅ is C(R₂₅), X₂₆ is C(R₂₆), and at least one selected from R₂₁ to R₂₆ is selected from groups represented by Formula 2-1.

18. The heterocyclic compound as claimed in claim 1, wherein the compound is one of the following Compounds 1 to 77:




-continued

5




-continued


8




6




9




7



10




-continued




11


-continued




14




12




15



13




16




17

-continued




18


-continued




22




19




23




20



24




21



25


-continued

26




-continued

29




30

27




31




32


28



33




-continued




34


-continued




37




35



38




36




39

-continued




40


-continued




43




41




44



42




45




46

-continued




47


-continued




51




48




52




49



53




50




54

-continued




55


-continued




59




56




60



57




58



61


-continued

62




-continued


65




63




66




67




64




68




-continued




-continued




70




73





71




74



-continued



-continued



19. An organic light emitting device, comprising:
a first electrode;
a second electrode facing the first electrode; and
an organic layer between the first electrode and the second electrode,

wherein the organic layer includes an emission layer, and
wherein the organic layer includes the heterocyclic compound as claimed in claim 1.

20. The organic light-emitting device as claimed in claim 19, wherein the emission layer includes a host and a dopant, and at least one selected from the host and the dopant includes the heterocyclic compound.

* * * * *

专利名称(译)	异环化合物和有机发光装置，包括相同的装置		
公开(公告)号	US20180026206A1	公开(公告)日	2018-01-25
申请号	US15/634392	申请日	2017-06-27
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	AHN HEECHOON KO SOOBYUNG JUN MIEUN KIM YOUNGKOOK HWANG SEOKHWAN		
发明人	AHN, HEECHOON KO, SOOBYUNG JUN, MIEUN KIM, YOUNGKOOK HWANG, SEOKHWAN		
IPC分类号	H01L51/00 C09K11/02 C07F9/64		
CPC分类号	H01L51/0072 C07F9/64 H01L51/0067 H01L51/5016 H01L51/5206 H01L51/5221 H01L51/5012 C09K11/025 C07F9/65583 C07F9/65685 H01L51/0059		
优先权	1020160094145 2016-07-25 KR		
其他公开文献	US10388889		
外部链接	Espacenet USPTO		

(KR) 10-2016-0094145

摘要(译)

杂环化合物和有机发光器件，该化合物由式1表示：

10

190

150

110