



US 20180198074A1

(19) **United States**

(12) **Patent Application Publication**  
**LEE et al.**

(10) **Pub. No.: US 2018/0198074 A1**  
(43) **Pub. Date: Jul. 12, 2018**

(54) **HETEROCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

*C07F 9/6561* (2006.01)  
*C07D 495/04* (2006.01)  
*C07D 519/00* (2006.01)

(71) Applicant: **SAMSUNG DISPLAY CO., LTD.**,  
Yongin-si (KR)

(52) **U.S. Cl.**  
CPC ..... *H01L 51/0071* (2013.01); *C07D 491/056* (2013.01); *H01L 51/0067* (2013.01); *C07F 9/6561* (2013.01); *H01L 51/0052* (2013.01); *H01L 51/5206* (2013.01); *H01L 51/0056* (2013.01); *H01L 51/0072* (2013.01); *C07D 495/04* (2013.01); *C07D 519/00* (2013.01); *H01L 51/0058* (2013.01)

(72) Inventors: **Hyoyoung LEE**, Yongin-si (KR);  
**Junha PARK**, Yongin-si (KR); **Munki SIM**, Yongin-si (KR); **Eunjae JEONG**, Yongin-si (KR); **Youngkook KIM**, Yongin-si (KR); **Seokhwan HWANG**, Yongin-si (KR)

(57) **ABSTRACT**

(21) Appl. No.: **15/665,536**

Provided are a heterocyclic compound and an organic light-emitting device including the same. The heterocyclic compound is represented by Formula 1:

(22) Filed: **Aug. 1, 2017**


(30) **Foreign Application Priority Data**

Jan. 6, 2017 (KR) ..... 10-2017-0002556

**Publication Classification**

(51) **Int. Cl.**  
*H01L 51/00* (2006.01)  
*C07D 491/056* (2006.01)

<Formula 1>



**10**

<b>190</b>
<b>150</b>
<b>110</b>

**FIG. 1**

**10**

<b>190</b>
<b>150</b>
<b>110</b>

**FIG. 2**

**20**

<b>190</b>
<b>150</b>
<b>110</b>
<b>210</b>

**FIG. 3**

**30**

<b>220</b>
<b>190</b>
<b>150</b>
<b>110</b>

# FIG. 4

40

<b>220</b>
<b>190</b>
<b>150</b>
<b>110</b>
<b>210</b>

**HETEROCYCLIC COMPOUND AND  
ORGANIC LIGHT-EMITTING DEVICE  
INCLUDING THE SAME**

CROSS-REFERENCE TO RELATED  
APPLICATION

**[0001]** Korean Patent Application No. 10-2017-0002556, filed on Jan. 6, 2017, in the Korean Intellectual Property Office, and entitled: "Heterocyclic Compound and Organic Light-Emitting Device Including the Same," is incorporated by reference herein in its entirety.

BACKGROUND

1. Field

**[0002]** Embodiments relate to a heterocyclic compound and an organic light-emitting device including the same.


2. Description of the Related Art

**[0003]** Organic light-emitting devices are self-emission devices and produce full-color images, and also have wide viewing angles, high contrast ratios, short response times, as well as excellent characteristics in terms of brightness, driving voltage, and response speed.

**[0004]** An example of such organic light-emitting devices may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, may recombine in the emission layer to produce excitons. Excitons transiting from an excited state to a ground state may generate light.

SUMMARY

**[0005]** Embodiments are directed to a heterocyclic compound represented by Formula 1:



**[0006]** In Formula 1,

**[0007]** A<sub>1</sub> and A<sub>2</sub> are each independently selected from a C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a C<sub>2</sub>-C<sub>60</sub> heterocyclic group,

**[0008]** X and Y are each independently oxygen (O) or sulfur (S),

**[0009]** L<sub>1</sub> and L<sub>2</sub> are each independently selected from a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heterocyclic group, \*—Si(Q<sub>1</sub>)(Q<sub>2</sub>)-\*<sup>†</sup>, \*—N(Q<sub>1</sub>)-\*<sup>†</sup>, \*—B(Q<sub>1</sub>)-\*<sup>†</sup>, \*—C(=O)—\*<sup>†</sup>, \*—S(=O)<sub>2</sub>+\*<sup>†</sup>, and \*—P(=O)(Q<sub>1</sub>)-\*<sup>†</sup>,

**[0010]** a<sub>1</sub> and a<sub>2</sub> are each independently an integer selected from 0 to 5,

**[0011]** when a<sub>1</sub> is equal to 0, \*(L<sub>1</sub>)<sub>a1</sub>-\*<sup>†</sup> is a single bond, and when a<sub>2</sub> is equal to 0, \*(L<sub>2</sub>)<sub>a2</sub>-\*<sup>†</sup> is a single bond,

**[0012]** Ar<sub>1</sub> and Ar<sub>2</sub> are each independently selected from a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —N(Q<sub>1</sub>)(Q<sub>2</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —S(=O)<sub>2</sub>(Q<sub>1</sub>), and —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>),

**[0013]** b<sub>1</sub> and b<sub>2</sub> are each independently an integer selected from 1 to 5,

**[0014]** R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —C(=O)(Q<sub>1</sub>), —S(=O)<sub>2</sub>(Q<sub>1</sub>), and —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>),

**[0015]** c<sub>1</sub> and c<sub>2</sub> are each independently an integer selected from 0 to 7,

**[0016]** m and n are each independently an integer selected from 0 to 8, and the sum of m and n is equal to or greater than 1,

**[0017]** at least one substituent selected from the substituted C<sub>3</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

**[0018]** deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group;

**[0019]** a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —N(Q<sub>11</sub>)(Q<sub>12</sub>), —B(Q<sub>1</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

**[0020]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

**[0021]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —N(Q<sub>21</sub>)(Q<sub>22</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>), and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and

**[0022]** —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>), and

**[0023]** Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

**[0024]** Embodiments are also directed to an organic light-emitting device including: a first electrode; a second electrode facing the first electrode; and an organic layer disposed between the first electrode and the second electrode and including an emission layer, wherein the organic layer includes a heterocyclic compound according to an embodiment.

## BRIEF DESCRIPTION OF THE DRAWINGS

**[0025]** Features will become apparent to those of skill in the art by describing in detail exemplary embodiments with reference to the attached drawings in which:

**[0026]** FIGS. 1 to 4 each illustrate a schematic view of an organic light-emitting device according to an embodiment.


## DETAILED DESCRIPTION

**[0027]** Example embodiments will now be described more fully hereinafter with reference to the accompanying drawings; however, they may be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey exemplary implementations to those skilled in the art. Like reference numerals refer to like elements throughout.

**[0028]** As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

**[0029]** According to an aspect of embodiments, a heterocyclic compound is represented by Formula 1:

<Formula 1>



**[0030]** In Formula 1, A<sub>1</sub> and A<sub>2</sub> may each independently be a C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a C<sub>2</sub>-C<sub>60</sub> heterocyclic group.

**[0031]** In an embodiment, A<sub>1</sub> and A<sub>2</sub> in Formula 1 may each independently be selected from a benzene group, a naphthalene group, an anthracene group, and a phenanthrene group.

**[0032]** For example, A<sub>1</sub> and A<sub>2</sub> in Formula 1 may each independently be a benzene group and a naphthalene group.

**[0033]** In Formula 1, X and Y may each independently be oxygen (O) or sulfur (S).

**[0034]** In an embodiment, X and Y in Formula 1 may be identical to or different from each other. For example, in Formula 1, X may be O and Y may be S; X may be S and Y may be O; or both X and Y may be O or S.

**[0035]** In Formula 1, L<sub>1</sub> and L<sub>2</sub> may each independently be a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heterocyclic group, \*—Si(Q<sub>1</sub>)(Q<sub>2</sub>)-\*<sup>\*</sup>, \*—N(Q<sub>1</sub>)-\*<sup>\*</sup>, \*—B(Q<sub>1</sub>)-\*<sup>\*</sup>, \*—C(=O)-\*<sup>\*</sup>, \*—S(=O)<sub>2</sub>-\*<sup>\*</sup>, and \*—P(=O)(Q<sub>1</sub>)-\*<sup>\*</sup>.

**[0036]** In one or more embodiments, L<sub>1</sub> and L<sub>2</sub> in Formula 1 may each independently be a divalent group based on a group selected from:

**[0037]** a benzene group, a pentalene group, an indene group, a naphthalene group, an azulene group, an indacene group, an acenaphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a perylene group, a penta-

cene group, a pyrrole group, a thiophene group, a furan group, a silole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indole group, an iso-indole group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a benzoisoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, a benzofuran group, a benzothiophene group, a benzosilole group, a benzothiazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a carbazole group, a dibenzofuran group, a dibenzothiophene group, a dibenzosilole group, a benzocarbazole group, a naphtho benzofuran group, a naphtho benzothiophene group, a naphthobenzosilole group, a dibenzocarbazole group, a dinaphthofuran group, a dinaphthothiophene group, a dinaphthosilole group, a thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, an oxazolpyridine group, a thiazolpyridine group, a benzonaphthyridine group, an azafluorene group, an azaspiro-bifluorene group, an azacarbazole group, an azadibenzofuran group, an azadibenzothiophene group, an azadibenzosilole group, an indeno pyrrole group, an indolopyrrole group, an indeno carbazole group, and an indolocarbazole group;

[0038] a benzene group, a pentalene group, an indene group, a naphthalene group, an azulene group, an indacene group, an acenaphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a perylene group, a pentacene group, a pyrrole group, a thiophene group, a furan group, a silole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indole group, an iso-indole group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a benzoisoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, a benzofuran group, a benzothiophene group, a benzosilole group, a benzothiazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a carbazole group, a dibenzofuran group, a dibenzothiophene group, a dibenzosilole group, a benzocarbazole group, a naphtho benzofuran group, a naphtho benzothiophene group, a naphthobenzosilole group, a dibenzocarbazole group, a dinaphthofuran group, a dinaphthothiophene group, a dinaphthosilole group, a thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, an oxazolpyridine group, a thiazolpyridine group, a benzonaphthyridine group, an azafluorene group, an azaspiro-bifluorene group, an azacarbazole group, an azadibenzofuran group, an azadibenzothiophene group, an azadibenzosilole group, an indeno pyrrole group, an indolopyrrole group, an indeno carbazole group, and an indolo-

carbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>20</sub> aryl group, a C<sub>1</sub>-C<sub>20</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a methylphenyl group, a biphenyl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>); and


[0039] \*—P(=O)(Q<sub>1</sub>)-\*<sup>1</sup>, and

[0040] Q<sub>1</sub> and Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from:


[0041] a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group; and

[0042] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group, each substituted with at least one selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, and a phenyl group.


[0043] In one or more embodiments, in Formula 1, L<sub>1</sub> and L<sub>2</sub> may each independently be selected from groups represented by Formulae 3-1 to 3-47:




Formula 3-1




Formula 3-2



Formula 3-3












Formula 3-4




Formula 3-5


-continued




Formula 3-6




Formula 3-7




Formula 3-8




Formula 3-9




Formula 3-10




Formula 3-11





Formula 3-12



Formula 3-13



Formula 3-14



-continued

Formula 3-15

Formula 3-16

Formula 3-17

Formula 3-18

Formula 3-19

Formula 3-20

Formula 3-21









Formula 3-22

Formula 3-23

Formula 3-24


Formula 3-25

-continued




-continued

Formula 3-26




Formula 3-27

Formula 3-28




Formula 3-29

Formula 3-30




Formula 3-31



Formula 3-32

Formula 3-33



Formula 3-34







Formula 3-35

Formula 3-36


Formula 3-37

Formula 3-38


-continued




Formula 3-39



Formula 3-40



Formula 3-41



Formula 3-42

Formula 3-43

Formula 3-44

-continued

Formula 3-45

Formula 3-46

Formula 3-47

[0044] In Formulae 3-1 to 3-47,

[0045]  $Y_1$  may be O, S,  $C(Z_3)(Z_4)$ ,  $N(Z_5)$ , or  $Si(Z_6)(Z_7)$ ,

[0046]  $Z_1$  to  $Z_7$  may each independently be selected from:

[0047] hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazolinyl group, a benzoquinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imida-

zopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indeno carbazolyl group, and an indolocarbazolyl group.

**[0048]** d2 may be an integer selected from 0 to 2,

**[0049]** d3 may be an integer selected from 0 to 3,

**[0050]** d4 may be an integer selected from 0 to 4,

**[0051]** d5 may be an integer selected from 0 to 5,

**[0052]** d6 may be an integer selected from 0 to 6,

**[0053]** d8 may be an integer selected from 0 to 8, and

**[0054]** \* and \*' each indicate a binding site to a neighboring atom.

**[0055]** For example, in Formula 1,  $L_1$  and  $L_2$  may each independently be selected from groups represented by Formulae 3-1, 3-2, 3-6, 3-8, 3-12, 3-24, 3-32, and 3-47.

**[0056]** In Formula 1, a1 and a2 may each independently be an integer selected from 0 to 5.

**[0057]** In an embodiment, in Formula 1, a1 and a2 may each independently be 0, 1, or 2.

**[0058]** For example, when a1 is equal to 0,  $*(L_1)_{a1}-*$  may be a single bond, and when a2 is equal to 0,  $*(L_2)_{a2}-*$  may be a single bond. For example, when a1 is an integer of two or more, two or more  $L_1(s)$  may be identical to or different from each other. For example, when a2 is an integer of two or more, two or more  $L_2(s)$  may be identical to or different from each other.

**[0059]** In Formula 1,  $Ar_1$  and  $Ar_2$  may each independently be selected from a substituted or unsubstituted  $C_3-C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_1-C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3-C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_1-C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6-C_{60}$  aryl group, a substituted or unsubstituted  $C_6-C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6-C_{60}$  arylthio group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,  $-Si(Q_1)(Q_2)(Q_3)$ ,  $-N(Q_1)(Q_2)$ ,  $-B(Q_1)(Q_2)$ ,  $-C(=O)(Q_1)$ ,  $-S(=O)_2(Q_1)$ , and  $-P(=O)(Q_1)(Q_2)$ .

**[0060]** In an embodiment, in Formula 1,  $A_1$  and  $Ar_2$  may each independently be selected from:

**[0061]** a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a

quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

**[0062]** a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1-C_{20}$  alkyl group, a  $C_1-C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, and  $-Si(Q_{31})(Q_{32})(Q_{33})$ ; and

**[0063]**  $-Si(Q_1)(Q_2)(Q_3)$ ,  $-C(=O)(Q_1)$ ,  $-S(=O)_2(Q_1)$ , and  $-P(=O)(Q_1)(Q_2)$ , and

[0064] Q<sub>1</sub> to Q<sub>3</sub> and Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from:


[0065] a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group; and

[0066] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group, each substituted with at least one selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, and a phenyl group.


[0067] In one or more embodiments, in Formula 1, Ar<sub>1</sub> and Ar<sub>2</sub> may each independently be selected from groups represented by Formulae 5-1 to 5-84:

-continued


Formula 5-8




Formula 5-9




Formula 5-1




Formula 5-10




Formula 5-2




Formula 5-3




Formula 5-4




Formula 5-5




Formula 5-6




Formula 5-7




Formula 5-11




Formula 5-12











Formula 5-13




Formula 5-14




-continued




Formula 5-15




Formula 5-16




Formula 5-17




Formula 5-18




Formula 5-19







Formula 5-20



Formula 5-21



Formula 5-22



-continued

Formula 5-23

Formula 5-24

Formula 5-25

Formula 5-26

Formula 5-27

Formula 5-28

Formula 5-29














Formula 5-30

Formula 5-31


Formula 5-32

Formula 5-33


-continued




Formula 5-34




Formula 5-35




Formula 5-36




Formula 5-37




Formula 5-38




Formula 5-39




Formula 5-40




Formula 5-41




Formula 5-42




Formula 5-43




Formula 5-44



Formula 5-45



Formula 5-46



-continued

Formula 5-47

Formula 5-48

Formula 5-49

Formula 5-50

Formula 5-51

Formula 5-52

Formula 5-53

Formula 5-54











Formula 5-55

Formula 5-56

Formula 5-57


Formula 5-58

-continued




-continued


Formula 5-59




Formula 5-60




Formula 5-61




Formula 5-62




Formula 5-63




Formula 5-64




Formula 5-65




Formula 5-66



Formula 5-67



Formula 5-68



Formula 5-69

Formula 5-70

Formula 5-71

Formula 5-72

Formula 5-73

Formula 5-74


Formula 5-75

Formula 5-76


Formula 5-77

Formula 5-78


-continued




Formula 5-79




Formula 5-80




Formula 5-81



Formula 5-82



Formula 5-83



Formula 5-84

[0068] In Formulae 5-1 to 5-84,

[0069]  $Y_{31}$  may be O, S,  $C(Z_{35})(Z_{36})$ ,  $N(Z_{37})$ , or  $Si(Z_{38})(Z_{39})$ .[0070]  $Z_{31}$  to  $Z_{39}$  may each independently be selected from:

[0071] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl


group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazolinyl group, a benzoquinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolopyridinyl group, a thiazolopyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolopyrrolyl group, an indeno carbazolyl group, and an indolocarbazolyl group.

[0072]  $e_2$  may be an integer selected from 0 to 2,[0073]  $e_3$  may be an integer selected from 0 to 3,[0074]  $e_4$  may be an integer selected from 0 to 4,[0075]  $e_5$  may be an integer selected from 0 to 5,[0076]  $e_6$  may be an integer selected from 0 to 6,[0077]  $e_7$  may be an integer selected from 0 to 7,[0078]  $e_9$  may be an integer selected from 0 to 9, and


[0079] \* indicates a binding site to a neighboring atom.

[0080] For example, in Formula 1,  $Ar_1$  and  $Ar_2$  may each independently be selected from groups represented by Formulae 5-1 to 5-4, 5-7, 5-13, 5-14, 5-21 to 5-24, 5-28, 5-76, and 5-80 to 5-83.

[0081] In one or more embodiments, in Formula 1,  $Ar_1$  and  $Ar_2$  may each independently be selected from groups represented by Formulae 6-1 to 6-31:












Formula 6-1




Formula 6-2


-continued



Formula 6-3




Formula 6-4




Formula 6-5


Formula 6-6




Formula 6-7




Formula 6-8




Formula 6-9



Formula 6-10



Formula 6-11



-continued

Formula 6-12

Formula 6-13

Formula 6-14

Formula 6-15


Formula 6-16

Formula 6-17


Formula 6-18

Formula 6-19


-continued




Formula 6-20




Formula 6-21



Formula 6-22




Formula 6-23




Formula 6-24


-continued




Formula 6-25




Formula 6-26




Formula 6-27




Formula 6-28



Formula 6-29



Formula 6-30



Formula 6-31

[0082] In Formulae 6-1 to 6-31, \* indicates a binding site to a neighboring atom.

**[0083]** In Formula 1, b1 and b2 may each independently be an integer selected from 1 to 5. For example, in Formula 1, b1 and b2 may each independently be 1 or 2.

**[0084]** In Formula 1, R<sub>1</sub> and R<sub>2</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>60</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —C(=O)(Q<sub>1</sub>), —S(=O)<sub>2</sub>(Q<sub>1</sub>), and —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>).

**[0085]** In an embodiment, in Formula 1, R<sub>1</sub> and R<sub>2</sub> may each independently be selected from:

**[0086]** hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

**[0087]** a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group;

**[0088]** a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a

C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>); and

**[0089]** —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), and

**[0090]** Q<sub>1</sub> to Q<sub>3</sub> and Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from:

**[0091]** a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazoliny group; and

**[0092]** a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazoliny group, each substituted with at least one selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, and a phenyl group.

**[0093]** For example, in Formula 1, R<sub>1</sub> and R<sub>2</sub> may each independently be selected from:

**[0094]** hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

**[0095]** a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group;

**[0096]** a cyclopentyl group, a cyclohexyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, and a terphenyl group;

**[0097]** a cyclopentyl group, a cyclohexyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, and a terphenyl group; and

**[0098]** —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), and

**[0099]** Q<sub>1</sub> to Q<sub>3</sub> may each independently be selected from a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, and a naphthyl group.

**[0100]** In Formula 1, c1 and c2 may each independently be an integer selected from 0 to 7. When c1 is two or more, two

or more  $R_1(s)$  may be identical to or different from each other. When  $c_2$  is two or more, two or more  $R_2(s)$  may be identical to or different from each other.

[0101] In Formula 1,  $m$  and  $n$  may each independently be an integer selected from 0 to 8, wherein the sum of  $m$  and  $n$  is equal to or greater than 1.

[0102] In an embodiment, in Formula 1, the sum of  $m$  and  $n$  is equal to 1 or 2.

[0103] In one or more embodiments, in Formula 1,  $R_1$  and  $R_2$  may each independently be selected from:

[0104] hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group; and

[0105] a  $C_1$ - $C_{20}$  alkyl group and a  $C_1$ - $C_{20}$  alkoxy group, each substituted with at least one selected from deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group, and

[0106]  $m$  and  $n$  may each independently satisfy a relationship of  $m=0$  and  $n=1$ ;  $m=1$  and  $n=0$ ;  $m=0$  and  $n=2$ ;  $m=1$  and  $n=1$ ; or  $m=2$  and  $n=0$ .


[0107] For example, in Formula 1,  $R_1$  and  $R_2$  may each independently be hydrogen, and

[0108]  $m$  and  $n$  may each independently satisfy a relationship of  $m=0$  and  $n=1$ ;  $m=1$  and  $n=0$ ;  $m=0$  and  $n=2$ ;  $m=1$  and  $n=1$ ; or  $m=2$  and  $n=0$ .


[0109] In an embodiment, Formula 1 may be represented by one selected from Formulae 1-1 to 1-9:

-continued


Formula 1-4




Formula 1-5




Formula 1-6




Formula 1-7




Formula 1-8




Formula 1-1




Formula 1-2




Formula 1-3



-continued










-continued



[0110] In Formulae 1-1 to 1-9,


[0111] descriptions of X, Y, L<sub>1</sub>, L<sub>2</sub>, a<sub>1</sub>, a<sub>2</sub>, Ar<sub>1</sub>, Ar<sub>2</sub>, b<sub>1</sub>, b<sub>2</sub>, R<sub>1</sub>, R<sub>2</sub>, c<sub>1</sub>, c<sub>2</sub>, m, and n are each independently the same as defined above.

[0112] In one or more embodiments, Formula 1 may be represented by one selected from Formulae 1-11 to 1-19:



-continued

Formula 1-19



[0113] wherein, in Formulae 1-11 to 1-19,

[0114] descriptions of  $R_{11}$  to  $R_{18}$  may be the same as defined in connection with  $R_1$ , and descriptions of  $R_{21}$  to  $R_{28}$  may be the same as defined in connection with  $R_2$ ,

[0115] at least one selected from  $R_{11}$  to  $R_{14}$  may be a binding site to a group represented by Formula 2-1, or at least one selected from  $R_{21}$  to  $R_{24}$  may be a binding site to a group represented by Formula 2-2, and

[0116] descriptions of X, Y,  $L_1$ ,  $L_2$ , a1, a2,  $Ar_1$ ,  $Ar_2$ , b1, and b2 may be the same as defined above.

[0117] In an embodiment, Formula 1 may be represented by Formula 1-11,

[0118] wherein, in Formula 1-11,

[0119] (i)  $R_{11}$  to  $R_{14}$  may each independently be selected from:

[0120] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group; and

[0121] a  $C_1$ - $C_{20}$  alkyl group and a  $C_1$ - $C_{20}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group,

[0122]  $R_{21}$  to  $R_{24}$  may each independently be selected from:

[0123] a group represented by Formula 2-2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group; and

[0124] a  $C_1$ - $C_{20}$  alkyl group and a  $C_1$ - $C_{20}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group, and

[0125] at least one selected from  $R_{21}$  to  $R_{24}$  may be a group represented by Formula 2-2, or

[0126] (ii)  $R_{11}$  to  $R_{14}$  may each independently be selected from:

[0127] a group represented by Formula 2-1, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano

group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group; and

[0128] a  $C_1$ - $C_{20}$  alkyl group and a  $C_1$ - $C_{20}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group,

[0129] at least one selected from  $R_{11}$  to  $R_{14}$  may be a group represented by Formula 2-1, and

[0130]  $R_{21}$  to  $R_{24}$  may each independently be selected from:

[0131] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group; and

[0132] a  $C_1$ - $C_{20}$  alkyl group and a  $C_1$ - $C_{20}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group.

[0133] In one or more embodiments, Formula 1 may be represented by Formula 1-11,

[0134] wherein, in Formula 1-11,

[0135]  $R_{11}$  to  $R_{14}$  may each independently be selected from:

[0136] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group; and

[0137] a  $C_1$ - $C_{20}$  alkyl group and a  $C_1$ - $C_{20}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group,


[0138]  $R_{21}$  and  $R_{24}$  may each independently be selected from:

[0139] a group represented by Formula 2-2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group; and


[0140] a  $C_1$ - $C_{20}$  alkyl group and a  $C_1$ - $C_{20}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group,

[0141] at least one selected from  $R_{22}$  and  $R_{23}$  may be a group represented by Formula 2-2.


[0142] In an embodiment, the heterocyclic compound represented by Formula 1 may be one selected from Compounds 1 to 110:



-continued



-continued




-continued


15




16




17



18




19




-continued


20




21




22




23




-continued




-continued




25




29




26




30









27










31



-continued




-continued




-continued


45




46




47




48



49




50




-continued


51




52




53







54







55



-continued




-continued




-continued

64




-continued


68



65





69





70

66




71





67






72



-continued




-continued




-continued


80



81




82




-continued


83




84



85




86




-continued


87



88




89




-continued


90




91




92



93




94




-continued


95




96



97




98




-continued


99




100




101




102




-continued




103




104



105




106




107


-continued



108




109



110

**[0143]** The heterocyclic compound represented by Formula 1 may have a core having a structure shown below, and accordingly, condensed rings may be formed with many heteroatoms, which may form a pi-conjugated system with increased intermolecular packing. In this regard, the heterocyclic compound represented by Formula 1 may have advantages of a high glass transition temperature, a high melting point, and good charge transporting capability. Therefore, when the heterocyclic compound represented by Formula 1 is applied to an electronic device, such as an organic light-emitting device, the electronic device may have high efficiency and a long lifespan:



[0144] In addition, in the heterocyclic compound represented by Formula 1, heteroatoms (for example, X and Y) are located at a para-position with respect to each other, thereby having a molecular structure that may more easily form hydrogen bonds between molecules. Due to such hydrogen bonds, molecules may be uniformly piled up to increase intermolecular packing, which may help increase a glass transition temperature and a melting point, and enhance charge transporting capability.

[0145] In addition, in the heterocyclic compound represented by Formula 1, the sum of m and n in Formula 1 is equal to or greater than 1, and that is, a substituent other than hydrogen is present. In an implementation, the substituent may increase a conjugation length, which may enhance charge transporting capability.

[0146] A synthesis method of the heterocyclic compound represented by Formula 1 may be understood by one of skill in the art by referring to Examples below.

[0147] A heterocyclic compound represented by Formula 1 may be used between a pair of electrodes of an organic light-emitting device. For example, the heterocyclic compound represented by Formula 1 may be included in at least one layer selected from a hole transport region, an electron transport region, and an emission layer. In one or more embodiments, the heterocyclic compound represented by Formula 1 may be used as a material for a capping layer located outside a pair of electrodes of an organic light-emitting device.

[0148] According to another aspect of embodiments, there is provided an organic light-emitting device including: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode and including an emission layer, wherein the organic layer comprises at least one heterocyclic compound according to an embodiment.

[0149] The expression “(an organic layer) includes at least one heterocyclic compound” used herein includes a case in which “(an organic layer) includes identical heterocyclic compounds represented by Formula 1” or a case in which “(an organic layer) includes two or more different heterocyclic compounds”.

[0150] In an example embodiment, the first electrode may be an anode, the second electrode may be a cathode, and the organic layer may further include a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode, wherein the hole transport region may include a hole injection layer, a hole transport layer, an emission auxiliary layer, an electron blocking layer, or a combination thereof, and the electron transport region may include a buffer layer, a hole blocking layer, an electron

control layer, an electron transport layer, an electron injection layer, or a combination thereof.

[0151] In an embodiment, the electron transport region may include the at least one heterocyclic compound represented by Formula 1.

[0152] In an embodiment, the electron transport region may include an electron transport layer, and the electron transport layer may include the at least one heterocyclic compound represented by Formula 1.

[0153] In one or more embodiments, the electron transport region may include an emission auxiliary layer, and the emission auxiliary layer may include the at least one heterocyclic compound represented by Formula 1.

[0154] For example, the hole transport region may include a p-dopant, wherein a lowest unoccupied molecular orbital (LUMO) energy level of the p-dopant is equal to or less than about -3.5 eV. The p-dopant may include, for example, a quinone derivative, a metal oxide, a cyano group-containing compound, or a combination thereof.

[0155] For example, the electron transport region may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth metal complex, a rare earth metal complex, or a combination thereof.

[0156] For example, the emission layer may include a dopant and a host, wherein the host may include at least one selected from an anthracene-based compound, a pyrene-based compound, and a spiro-bifluorene-based compound.

[0157] In the organic light-emitting device, the emission layer may include a first color light emission layer,

[0158] the emission layer in the organic light-emitting device may further include i) at least one second color light emission layer or ii) at least one second color light emission layer and at least one third color light emission layer, between the first electrode and the second electrode,

[0159] wherein a maximum emission wavelength of the first color light emission layer, a maximum emission wavelength of the second color light emission layer, and a maximum emission wavelength of the third color light emission layer may be identical to or different from each other, and

[0160] the first color light and the second color light may be emitted as mixed light, or the first color light, the second color light, and the third color light may be emitted as mixed light.

[0161] The organic light-emitting device may further include at least one selected from a first capping layer disposed in a pathway along which light generated in the emission layer proceeds toward the outside through the first electrode and a second capping layer disposed in a pathway along which light generated in the emission layer proceeds toward the outside through the second electrode, wherein at least one of the first capping layer and the second capping layer may include the at least one heterocyclic compound represented by Formula 1.

[0162] For example, the organic light-emitting device may have i) a stack structure including a first electrode, an organic layer, a second electrode, and a second capping layer which are sequentially stacked in this stated order, ii) a stack structure including a first capping layer, a first electrode, an organic layer, and a second electrode which are sequentially stacked in this stated order, or iii) a stack structure including a first capping layer, a first electrode, an organic layer, a

second electrode, and a second capping layer which are sequentially stacked in this stated order, wherein, in each structure, at least one of the first capping layer and the second capping layer may include the at least one heterocyclic compound represented by Formula 1.

[0163] The term “organic layer” used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of the organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

[0164] [Description of FIG. 1]

[0165] FIG. 1 is a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 includes a first electrode 110, an organic layer 150, and a second electrode 190.

[0166] Hereinafter, the structure of the organic light-emitting device 10 according to an embodiment and a method of manufacturing the organic light-emitting device 10 will be described in connection with FIG. 1.

[0167] [First Electrode 110]

[0168] In FIG. 1, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be, for example, a glass substrate or a plastic substrate, which may have excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

[0169] The first electrode 110 may be formed by, for example, depositing or sputtering a material for forming the first electrode 110 on the substrate. When the first electrode 110 is an anode, the material for forming the first electrode 110 may be selected from, for example, materials with a high work function to facilitate hole injection.

[0170] The first electrode 110 may be, for example, a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode 110 is a transmissive electrode, the material for forming the first electrode 110 may be selected from, for example, indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO<sub>2</sub>), zinc oxide (ZnO), and a combination thereof. In one or more embodiments, when the first electrode 110 is a semi-transmissive electrode or a reflective electrode, the material for forming the first electrode 110 may be selected from, for example, magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and a combination thereof.

[0171] The first electrode 110 may have, for example, a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode 110 may have a three-layered structure of ITO/Ag/ITO.

[0172] [Organic Layer 150]

[0173] The organic layer 150 may be disposed on the first electrode 110. The organic layer 150 may include an emission layer.

[0174] The organic layer 150 may further include a hole transport region between the first electrode 110 and the emission layer, and an electron transport region between the emission layer and the second electrode 190.


[0175] [Hole Transport Region in Organic Layer 150]

[0176] The hole transport region may have, for example, i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0177] The hole transport region may include at least one layer selected from a hole injection layer, a hole transport layer, an emission auxiliary layer, and an electron blocking layer.


[0178] For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure, wherein, for each structure, constituting layers are sequentially stacked from the first electrode 110 in this stated order, but the structure of the hole transport region is not limited thereto.

[0179] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPD), β-NPB, TPD, spiro-TPD, spiro-NPB, methylated-NPB, TAPC, HMTDP, 4,4',4"-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:




m-MTDATA

-continued




TDATA


-continued




$\beta$ -NPB




TPD




2-TNATA




Spiro-TPD






Spiro-NPB



NPB



methylated NPB



[0180] In Formulae 201 and 202,

[0181]  $L_{201}$  to  $L_{204}$  may each independently be selected from a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkylene group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenylene group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylene group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

[0182]  $L_{205}$  may be selected from  $*-O-*$ ,  $*-S-*$ ,  $*-N(Q_{201})-*$ , a substituted or unsubstituted  $C_1$ - $C_{20}$  alkylene group, a substituted or unsubstituted  $C_2$ - $C_{20}$  alkenylene group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkylene group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenylene group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylene group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0183]  $xa1$  to  $xa4$  may each independently be an integer of 0 to 3,

[0184]  $xa5$  may be an integer of 1 to 10, and

[0185]  $R_{201}$  to  $R_{204}$  and  $Q_{201}$  may each independently be selected from a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0186] In an embodiment, in Formula 202,  $R_{201}$  and  $R_{202}$  may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group, and  $R_{203}$  and  $R_{204}$  may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group.

[0187] In one or more embodiments, in Formulae 201 and 202,

[0188]  $L_{201}$  to  $L_{205}$  may each independently be selected from:

[0189] a phenylene group, a pentalenylene group, an indenylene group, a naphthalenylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylene group; and

[0190] a phenylene group, a pentalenylene group, an indenylene group, a naphthalenylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylene group, each substituted with at least one selected from deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono

group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>) and —N(Q<sub>31</sub>)(Q<sub>32</sub>), and

**[0191]** Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

**[0192]** In one or more embodiments, xa1 to xa4 may each independently be 0, 1, or 2.

**[0193]** In one or more embodiments, xa5 may be 1, 2, 3, or 4.

**[0194]** In one or more embodiments, R<sub>201</sub> to R<sub>204</sub> and Q<sub>201</sub> may each independently be selected from a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

**[0195]** a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl,

—Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), and —N(Q<sub>31</sub>)(Q<sub>32</sub>), and

**[0196]** descriptions of Q<sub>31</sub> to Q<sub>33</sub> are the same as defined above.

**[0197]** In one or more embodiments, in Formula 201, at least one selected from R<sub>201</sub> to R<sub>203</sub> may each independently be selected from:

**[0198]** a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

**[0199]** a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.

**[0200]** In one or more embodiments, in Formula 202, i) R<sub>201</sub> and R<sub>202</sub> may be linked via a single bond and/or ii) R<sub>203</sub> and R<sub>204</sub> may be linked via a single bond.


**[0201]** In one or more embodiments, in Formula 202, at least one selected from R<sub>201</sub> to R<sub>204</sub> may be selected from:

**[0202]** a carbazolyl group; and

**[0203]** a carbazolyl group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.


[0204] The compound represented by Formula 201 may be represented by Formula 201A:

<Formula 201A>




[0205] In an embodiment, the compound represented by Formula 201 may be represented by Formula 201A(1) below:

<Formula 201A(1)>




[0206] In an embodiment, the compound represented by Formula 201 may be represented by Formula 201A-1 below:

<Formula 201A-1>




[0207] In an embodiment, the compound represented by Formula 202 may be represented by Formula 202A:

<Formula 202A>



[0208] In an embodiment, the compound represented by Formula 202 may be represented by Formula 202A-1:

<Formula 202A-1>



[0209] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,


[0210] descriptions of  $L_{201}$  to  $L_{203}$ ,  $xa1$  to  $xa3$ ,  $xa5$ , and  $R_{202}$  to  $R_{204}$  are each the same as defined above,

[0211] descriptions of  $R_{211}$  and  $R_{212}$  are each the same as defined in connection with  $R_{203}$ ,

[0212]  $R_{213}$  to  $R_{217}$  may each independently be selected from hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a  $C_1$ - $C_{10}$  alkyl group, a phenyl group substituted with  $-F$ , a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0213] The hole transport region may include at least one compound selected from Compounds HT1 to HT39:

HT1





-continued

-continued



HT2

HT4



HT3

HT5





-continued

-continued



HT6

HT8



HT7


HT9




-continued

-continued

HT10





HT12




HT13

HT11




-continued

HT14




-continued


HT17




HT15




HT18



HT16




HT19






-continued

HT26




-continued


HT29




HT27




HT30




HT31



HT28




HT32




-continued

HT33




-continued


HT36




HT34




HT37



HT35




HT38



-continued

HT39



[0214] A thickness of the hole transport region may be in a range of, for example, about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of, for example, about 100 Å to about 9,000 Å, for example, about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of, for example, about 50 Å to about 2,000 Å, for example about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0215] The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer, and the electron blocking layer may block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include the materials as described above.

[0216] [P-Dopant]

[0217] The hole transport region may further include, in addition to these materials, a charge-generation material for improving conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0218] The charge-generation material may be, for example, a p-dopant.

[0219] In an embodiment, a LUMO energy level of the p-dopant may be  $-3.5$  eV or less.

[0220] The p-dopant may include, for example, at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound.




[0221] For example, the p-dopant may include at least one selected from

[0222] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) or 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);

[0223] a metal oxide, such as tungsten oxide or molybdenum oxide;

[0224] 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and

[0225] a compound represented by Formula 221 below:



&lt;Formula 221&gt;

[0226] In Formula 221,

[0227]  $R_{221}$  to  $R_{223}$  may each independently be selected from a substituted or unsubstituted  $C_3$ - $C_{60}$  cycloalkyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, wherein at least one selected from  $R_{221}$  to  $R_{223}$  may have at least one substituent selected from a cyano group,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a  $C_1$ - $C_{20}$  alkyl group substituted with  $-F$ , a  $C_1$ - $C_{20}$  alkyl group substituted with  $-Cl$ , a  $C_1$ - $C_{20}$  alkyl group substituted with  $-Br$  and a  $C_1$ - $C_{20}$  alkyl group substituted with  $-I$ .

[0228] [Emission Layer in Organic Layer 150]

[0229] When the organic light-emitting device 10 is a full-color organic light-emitting device, the emission layer may be patterned into, for example, a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In one or more embodiments, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers contact each other or are separated from each other. In one or more embodiments, the emission layer may be patterned into a red

light-emitting material, a green-light emitting material, and a blue-light emitting material, in which two or more materials are mixed with each other in a single layer to emit white color.

[0230] The emission layer may include, for example, a host and a dopant. The dopant may include, for example, at least one of a phosphorescent dopant and a fluorescent dopant.

[0231] An amount of the dopant in the emission layer may be, for example, in a range of about 0.01 parts by weight to about 15 parts by weight based on 100 parts by weight of the host.

[0232] A thickness of the emission layer may be in a range of, for example, about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0233] [Host in Emission Layer]

[0234] In one or more embodiments, the host may include a compound represented by Formula 301 below.



[0235] In Formula 301,

[0236]  $\text{Ar}_{301}$  may be a substituted or unsubstituted  $\text{C}_5\text{-C}_{60}$  carbocyclic group or a substituted or unsubstituted  $\text{C}_1\text{-C}_{60}$  heterocyclic group,

[0237]  $\text{xb}11$  may be 1, 2, or 3,

[0238]  $\text{L}_{301}$  may be selected from a substituted or unsubstituted  $\text{C}_3\text{-C}_{10}$  cycloalkylene group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{10}$  heterocycloalkylene group, a substituted or unsubstituted  $\text{C}_3\text{-C}_{10}$  cycloalkenylene group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{10}$  heterocycloalkenylene group, a substituted or unsubstituted  $\text{C}_6\text{-C}_{60}$  arylene group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0239]  $\text{xb}1$  may be an integer of 0 to 5,

[0240]  $\text{R}_{301}$  may be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{60}$  alkyl group, a substituted or unsubstituted  $\text{C}_2\text{-C}_{60}$  alkenyl group, a substituted or unsubstituted  $\text{C}_2\text{-C}_{60}$  alkynyl group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{60}$  alkoxy group, a substituted or unsubstituted  $\text{C}_3\text{-C}_{10}$  cycloalkyl group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $\text{C}_3\text{-C}_{10}$  cycloalkenyl group, a substituted or unsubstituted

$\text{C}_1\text{-C}_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $\text{C}_6\text{-C}_{60}$  aryl group, a substituted or unsubstituted  $\text{C}_6\text{-C}_{60}$  aryloxy group, a substituted or unsubstituted  $\text{C}_6\text{-C}_{60}$  arylthio group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si( $\text{Q}_{301}$ )( $\text{Q}_{302}$ )( $\text{Q}_{303}$ ), —N( $\text{Q}_{301}$ )( $\text{Q}_{302}$ ), —B( $\text{Q}_{301}$ )( $\text{Q}_{302}$ ), —C(=O)( $\text{Q}_{301}$ ), —S(=O)<sub>2</sub>( $\text{Q}_{301}$ ), and —P(=O)( $\text{Q}_{301}$ )( $\text{Q}_{302}$ ),

[0241]  $\text{xb}21$  may be an integer from 1 to 5, and

[0242]  $\text{Q}_{301}$  to  $\text{Q}_{303}$  may each independently be selected from a  $\text{C}_1\text{-C}_{10}$  alkyl group, a  $\text{C}_1\text{-C}_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0243] In an embodiment,  $\text{Ar}_{301}$  in Formula 301 may be selected from:

[0244] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group; and


[0245] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $\text{C}_1\text{-C}_{20}$  alkyl group, a  $\text{C}_1\text{-C}_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si( $\text{Q}_{31}$ )( $\text{Q}_{32}$ )( $\text{Q}_{33}$ ), —N( $\text{Q}_{31}$ )( $\text{Q}_{32}$ ), —B( $\text{Q}_{31}$ )( $\text{Q}_{32}$ ), —C(=O)( $\text{Q}_{31}$ ), —S(=O)<sub>2</sub>( $\text{Q}_{31}$ ), and —P(=O)( $\text{Q}_{31}$ )( $\text{Q}_{32}$ ), and

[0246]  $\text{Q}_{31}$  to  $\text{Q}_{33}$  may each independently be selected from a  $\text{C}_1\text{-C}_{10}$  alkyl group, a  $\text{C}_1\text{-C}_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0247] When  $\text{xb}11$  in Formula 301 is two or more, two or more  $\text{Ar}_{301}(\text{s})$  may be linked via a single bond.


[0248] In one or more embodiments, the compound represented by Formula 301 may be represented by Formula 301-1 or 301-2:

<Formula 301-1>



-continued

&lt;Formula 301-2&gt;



[0249] wherein, in Formulae 301-1 and 301-2,

[0250]  $A_{301}$  to  $A_{304}$  may each independently be selected from a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, an indene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, an indole group, a carbazole group, a benzocarbazole group, a dibenzocarbazole group, a furan group, a benzofuran group, a dibenzofuran group, a naphthofuran group, a benzonaphthofuran group, a dinaphthofuran group, a thiophene group, a benzothiophene group, a dibenzothiophene group, a naphthothiophene group, a benzonaphthothiophene group, and a dinaphthothiophene group.

[0251]  $X_{301}$  may be O, S, or N- $[(L_{304})_{xb4}-R_{304}]$ .

[0252]  $R_{311}$  to  $R_{314}$  may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group —Si( $Q_{31}$ )( $Q_{32}$ )( $Q_{33}$ ), —N( $Q_{31}$ )( $Q_{32}$ ), —B( $Q_{31}$ )( $Q_{32}$ ), —C(=O)( $Q_{31}$ ), —S(=O)<sub>2</sub>( $Q_{31}$ ), and —P(=O)( $Q_{31}$ )( $Q_{32}$ ).

[0253]  $xb_{22}$  and  $xb_{23}$  may each independently be 0, 1, or 2,

[0254] descriptions of  $L_{301}$ ,  $xb_1$ ,  $R_{301}$ , and  $Q_{31}$  to  $Q_{33}$  are the same as defined above,

[0255] descriptions of  $L_{302}$  to  $L_{304}$  are the same as defined in connection with  $L_{301}$ ,

[0256] descriptions of  $xb_2$  to  $xb_4$  are the same as defined in connection with  $xb_1$ , and

[0257] descriptions of  $R_{302}$  to  $R_{304}$  are the same as defined in connection with  $R_{301}$ .

[0258] For example, in Formulae 301, 301-1, and 301-2,  $L_{301}$  to  $L_{304}$  may each independently be selected from:

[0259] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazolylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isox-

azolylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinylylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinnolinylylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthroliinylylene group, a phenazinylylene group, a benzimidazolylylene group, an isobenzothiazolylylene group, a benzoxazolylylene group, an isobenzoxazolylylene group, a triazolylylene group, a tetrazolylylene group, an imidazopyridinylylene group, an imidazopyrimidinylylene group, and an azacarbazolylylene group; and

[0260] a phenylene group, a naphthylene group, a fluorenylylene group, a spiro-bifluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazolylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinylylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinnolinylylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthroliinylylene group, a phenazinylylene group, a benzimidazolylylene group, an isobenzothiazolylylene group, a benzoxazolylylene group, an isobenzoxazolylylene group, a triazolylylene group, a tetrazolylylene group, an imidazopyridinylylene group, an imidazopyrimidinylylene group, and an azacarbazolylylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenylyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a penta-

phenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group,  $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$ ,  $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{C}(=\text{O})(\text{Q}_{31})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{31})$ , and  $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$ , and

**[0261]** descriptions of  $\text{Q}_{31}$  to  $\text{Q}_{33}$  are the same as defined above.

**[0262]** In an embodiment, in Formulae 301, 301-1, and 301-2,  $\text{R}_{301}$  to  $\text{R}_{304}$  may each independently be selected from:

**[0263]** a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

**[0264]** a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl


group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $\text{C}_1$ - $\text{C}_{20}$  alkyl group, a  $\text{C}_1$ - $\text{C}_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group,  $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$ ,  $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{C}(=\text{O})(\text{Q}_{31})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{31})$ , and  $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$ , and

**[0265]** descriptions of  $\text{Q}_{31}$  to  $\text{Q}_{33}$  are the same as defined above.

**[0266]** In one or more embodiments, the host may include, for example, an alkaline earth metal complex. For example, the host may be selected from a beryllium (Be) complex (for example, Compound H55), an Mg complex, and a Zn complex.



**[0267]** The host may include, for example, at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di-(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolyl-benzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55:

-continued





H1

H7





H2

H8





H3

H9





H4

H10




H5

H11




H6

H12




-continued




H13


-continued




H18




H14




H19




H15




H20




H16



H21




H17



H22


-continued

H23




-continued


H26




H24



H25




H27




-continued

H28




-continued


H32




H29




H33




H30




H34




H31



H35




-continued




H36

-continued




H40


H37




H38




H41






H39



H42




-continued




-continued

H43




H46

H44




H47

H48




H49

H45




H50


H51




-continued




H52




H53



H54




H55



&lt;Formula 401&gt;

&lt;Formula 402&gt;



[0270] In Formulae 401 and 402,

[0271] M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

[0272]  $L_{401}$  may be selected from ligands represented by Formula 402, and  $xc1$  may be 1, 2, or 3, wherein, when  $xc1$  is two or more, two or more  $L_{401}(s)$  may be identical to or different from each other,

[0273]  $L_{402}$  may be an organic ligand, and  $xc2$  may be an integer of 0 to 4, wherein, when  $xc2$  is two or more, two or more  $L_{402}(s)$  may be identical to or different from each other,

[0274]  $X_{401}$  to  $X_{404}$  may each independently be nitrogen or carbon,

[0275]  $X_{401}$  and  $X_{403}$  may be linked via a single bond or a double bond, and  $X_{402}$  and  $X_{404}$  may be linked via a single bond or a double bond,

[0276]  $A_{401}$  and  $A_{402}$  may each independently be a  $C_5$ - $C_{60}$  carbocyclic group or a  $C_1$ - $C_{60}$  heterocyclic group,

[0277]  $X_{405}$  may be a single bond,  $*-O-*'$ ,  $*-S-*'$ ,  $*-C(=O)-*'$ ,  $*-N(Q_{411})-*'$ ,  $*-C(Q_{411})(Q_{412})-*'$ ,  $*-C(Q_{411})=C(Q_{412})-*'$ ,  $*-C(Q_{411})-*'$ , or  $*=C=*'$ , wherein  $Q_{411}$  and  $Q_{412}$  may each independently be hydrogen, deuterium, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group.

[0278]  $X_{406}$  may be a single bond, O, or S.

[0279]  $R_{401}$  and  $R_{402}$  may each independently be selected from hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{20}$  alkyl group, a substituted or unsubstituted  $C_1$ - $C_{20}$  alkoxy group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,  $-Si(Q_{401})(Q_{402})(Q_{403})$ ,  $-N(Q_{401})$

[0268] [Phosphorescent Dopant Included in Emission Layer in Organic Layer 150]

[0269] The phosphorescent dopant may include, for example, an organometallic complex represented by Formula 401 below:

(Q<sub>402</sub>), —B(Q<sub>401</sub>)(Q<sub>402</sub>), —C(=O)(Q<sub>401</sub>), —S(=O)<sub>2</sub>(Q<sub>401</sub>), and —P(=O)(Q<sub>401</sub>)(Q<sub>402</sub>),

[0280] Q<sub>401</sub> to Q<sub>403</sub> may each independently be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>6</sub>-C<sub>20</sub> aryl group, and a C<sub>1</sub>-C<sub>20</sub> heteroaryl group,

[0281] xc11 and xc12 may each independently be an integer of 0 to 10, and

[0282] in Formula 402, \* and \*' each indicate a binding site to M in Formula 401.

[0283] In an embodiment, in Formula 402, A<sub>401</sub> and A<sub>402</sub> may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

[0284] In one or more embodiments, in Formula 402, i) X<sub>401</sub> may be nitrogen, and X<sub>402</sub> may be carbon or ii) X<sub>401</sub> and X<sub>402</sub> may each be nitrogen at the same time.

[0285] In one or more embodiments, R<sub>401</sub> and R<sub>402</sub> in Formula 402 may each independently be selected from

[0286] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

[0287] a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

[0288] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0289] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naph-

thyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0290] —Si(Q<sub>401</sub>)(Q<sub>402</sub>)(Q<sub>403</sub>), —N(Q<sub>401</sub>)(Q<sub>402</sub>), —B(Q<sub>401</sub>)(Q<sub>402</sub>), —C(=O)(Q<sub>401</sub>), —S(=O)<sub>2</sub>(Q<sub>401</sub>), and —P(=O)(Q<sub>401</sub>)(Q<sub>402</sub>), and


[0291] Q<sub>401</sub> to Q<sub>403</sub> may each independently be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group.

[0292] In one or more embodiments, when xc1 in Formula 401 is two or more, two A<sub>401</sub>(s) in two or more L<sub>401</sub>(s) may optionally be linked via X<sub>407</sub>, which is a linking group, or two A<sub>402</sub>(S) in two or more L<sub>401</sub>(s) may optionally be linked via X<sub>408</sub>, which is a linking group (see Compounds PD1 to PD4 and PD7). In one or more embodiments, X<sub>407</sub> and X<sub>408</sub> may each independently be a single bond, \*—O—\*<sup>\*</sup>, \*—S—\*<sup>\*</sup>, \*—C(=O)—\*<sup>\*</sup>, \*—N(Q<sub>413</sub>)-\*<sup>\*</sup>, \*—C(Q<sub>413</sub>)(Q<sub>414</sub>)-\*<sup>\*</sup> or \*—C(Q<sub>413</sub>)=C(Q<sub>414</sub>)-\*<sup>\*</sup> (wherein Q<sub>413</sub> and Q<sub>414</sub> may each independently be hydrogen, deuterium, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group).


[0293] L<sub>402</sub> in Formula 401 may be a monovalent, divalent, or trivalent organic ligand. For example, L<sub>402</sub> in Formula 401 may be selected from halogen, diketone (for example, acetylacetonate), carboxylic acid (for example, picolinate), —C(=O), isonitrile, —CN, and phosphorus (for example, phosphine or phosphite).

[0294] In one or more embodiments, the phosphorescent dopant may be, for example, selected from Compounds PD1 to PD25:







PD1



PD2




-continued




-continued

PD3




PD9

PD4




PD10

PD5




PD11

PD6



PD12






PD7



PD13


PD8

-continued




-continued


PD14




PD15



PD16



PD17



PD'8


PD19

PD20


PD21

PD22


-continued




PD23



PD24



PD25

**[0295]** [Fluorescent Dopant in Emission Layer]**[0296]** The fluorescent dopant may include, for example, an arylamine compound or a styrylamine compound.**[0297]** The fluorescent dopant may include, for example, a compound represented by Formula 501 below.**[0298]** In Formula 501,**[0299]** Ar<sub>501</sub> may be a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heterocyclic group,**[0300]** L<sub>501</sub> to L<sub>503</sub> may each independently be selected from a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkylene group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

cloalkylene group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenylene group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylene group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

**[0301]** x<sub>d1</sub> to x<sub>d3</sub> may each independently be an integer of 0 to 3,**[0302]** R<sub>501</sub> and R<sub>502</sub> may each independently be selected from a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and**[0303]** x<sub>d4</sub> may be an integer of 1 to 6.**[0304]** In an embodiment, Ar<sub>501</sub> in Formula 501 may be selected from:**[0305]** a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group; and**[0306]** a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.**[0307]** In one or more embodiments, L<sub>501</sub> to L<sub>503</sub> in Formula 501 may each independently be selected from:**[0308]** a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group; and**[0309]** a phenylene group, a naphthylene group, a fluorenylylene group, a spiro-bifluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthre-

nylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylynylene group, a pyrenylylene group, a chrysenylylene group, a perylynylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0310] In one or more embodiments,  $R_{501}$  and  $R_{502}$  in Formula 502 may each independently be selected from:

[0311] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and


[0312] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl

group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, and —Si( $Q_{31}$ )( $Q_{32}$ )( $Q_{33}$ ), and


[0313]  $Q_{31}$  to  $Q_{33}$  may each independently be selected from a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0314] In one or more embodiments, xd4 in Formula 501 may be 2.

[0315] For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:




FD1



FD2


-continued



FD3


-continued

FD5





FD6

FD4




FD7




-continued

FD8




-continued


FD11




FD9




FD12




FD13







FD10



FD14




-continued



-continued


FD15

FD19




FD16

FD20




FD17

FD21




FD18

FD22



[0316] In one or more embodiments, the fluorescent dopant may be selected from the following compounds.



[0317] [Electron Transport Region in Organic Layer 150]

[0318] The electron transport region may have, for example, i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0319] The electron transport region may include, for example, at least one selected from a buffer layer, a hole

blocking layer, an electron control layer, an electron transport layer, and an electron injection layer.

[0320] For example, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, wherein, for each structure, constituting layers are sequentially stacked from an emission layer.

However, embodiments of the structure of the electron transport region are not limited thereto.

**[0321]** The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include a metal-free compound containing at least one  $\pi$  electron-depleted nitrogen-containing ring.

**[0322]** The “ $\pi$  electron-depleted nitrogen-containing ring” indicates a  $C_1$ - $C_{60}$  heterocyclic group having at least one  $*-N=*$  moiety as a ring-forming moiety.

**[0323]** For example, the “ $\pi$  electron-depleted nitrogen-containing ring” may be i) a 5-membered to 7-membered heteromonocyclic group having at least one  $*-N=*$  moiety, ii) a heteropolycyclic group in which two or more 5-membered to 7-membered heteromonocyclic groups each having at least one  $*-N=*$  moiety are condensed with each other, or iii) a heteropolycyclic group in which at least one of 5-membered to 7-membered heteromonocyclic groups, each having at least one  $*-N=*$  moiety, is condensed with at least one  $C_5$ - $C_{60}$  carbocyclic group.

**[0324]** Examples of the  $\pi$  electron-depleted nitrogen-containing ring include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, thiadiazol, an imidazopyridine, an imidazopyrimidine, and an azacarbazole, but are not limited thereto.

**[0325]** For example, the electron transport region may include, in addition to the heterocyclic compound represented by Formula 1, a compound represented by Formula 601.



**[0326]** In Formula 601,

**[0327]**  $Ar_{601}$  may be a substituted or unsubstituted  $C_5$ - $C_{60}$  carbocyclic group or a substituted or unsubstituted  $C_1$ - $C_{60}$  heterocyclic group.

**[0328]**  $xe11$  may be 1, 2, or 3.

**[0329]**  $L_{601}$  may be selected from a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkylene group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenylene group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylene group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.

**[0330]**  $xe1$  may be an integer of 0 to 5,

**[0331]**  $R_{601}$  may be selected from a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed

heteropolycyclic group,  $-\text{Si}(Q_{601})(Q_{602})(Q_{603})$ ,  $-\text{C}(=\text{O})(Q_{601})$ ,  $-\text{S}(=\text{O})_2(Q_{601})$ , and  $-\text{P}(=\text{O})(Q_{601})(Q_{602})$ ,

**[0332]**  $Q_{601}$  to  $Q_{603}$  may each independently be a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

**[0333]**  $xe21$  may be an integer from 1 to 5.

**[0334]** In an embodiment, at least one selected from  $xe11$   $Ar_{601}(s)$  and  $xe21$   $R_{601}(s)$  may include the  $\pi$  electron-depleted nitrogen-containing ring.

**[0335]** In an embodiment, in Formula 601, ring  $Ar_{601}$  may be selected from:

**[0336]** a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a penta-phenene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and


**[0337]** a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a penta-phenene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group,  $-\text{Si}(Q_{31})(Q_{32})(Q_{33})$ ,  $-\text{S}(=\text{O})_2(Q_{31})$ , and  $-\text{P}(=\text{O})(Q_{31})(Q_{32})$ , and

[0338]  $Q_{31}$  to  $Q_{33}$  may each independently be selected from a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0339] When xe11 in Formula 601 is two or more, two or more  $Ar_{601}(s)$  may be linked via a single bond.

[0340] In one or more embodiments,  $Ar_{601}$  in Formula 601 may be an anthracene group.

[0341] In one or more embodiments, a compound represented by Formula 601 may be represented by Formula 601-1:



[0342] In Formula 601-1,

[0343]  $X_{614}$  may be N or C( $R_{614}$ ),  $X_{615}$  may be N or C( $R_{615}$ ), and  $X_{616}$  may be N or C( $R_{616}$ ), wherein at least one selected from  $X_{614}$  to  $X_{616}$  may be N,

[0344]  $L_{611}$  to  $L_{613}$  may each independently be the same as defined in connection with  $L_{601}$ ,

[0345] xe611 to xe613 may each independently be the same as defined in connection with xe1,

[0346]  $R_{611}$  to  $R_{613}$  may each independently be the same as defined in connection with  $R_{601}$ , and

[0347]  $R_{614}$  to  $R_{616}$  may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0348] In an embodiment,  $L_{601}$  and  $L_{611}$  to  $L_{613}$  in Formulae 601 and 601-1 may each independently be selected from:

[0349] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylene group, a dibenzofuranylenylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

[0350] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylene group, a dibenzofuranylenylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinylyl group, an isoquinolinylyl group, a benzoquinolinylyl group, a phthalazinyl group, a naphthyridinyl

group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridiny group, an acridiny group, a phenanthroliny group, a phenaziny group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridiny group, an imidazopyrimidinyl group, and an azacarbazolyl group.

**[0351]** In one or more embodiments, xe1 and xe611 to xe613 in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

**[0352]** In one or more embodiments, R<sub>601</sub> and R<sub>611</sub> to R<sub>613</sub> in Formula 601 and 601-1 may each independently be selected from:

**[0353]** a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridiny group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyraziny group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridiny group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridiny group, an acridiny group, a phenanthroliny group, a phenaziny group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridiny group, an imidazopyrimidinyl group, and an azacarbazolyl group;

**[0354]** a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridiny group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyraziny group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridiny group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridiny group, an acridiny group, a phenanthroliny group, a phenaziny group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridiny group, an imidazopyrimidinyl group, and an azacarbazolyl group;


azolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridiny group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridiny group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyraziny group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridiny group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridiny group, an acridiny group, a phenanthroliny group, a phenaziny group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridiny group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

**[0355]** —S(=O)<sub>2</sub>(Q<sub>601</sub>), and —P(=O)(Q<sub>601</sub>)(Q<sub>602</sub>), and




**[0356]** descriptions of Q<sub>601</sub> and Q<sub>602</sub> are the same as defined above.

**[0357]** The electron transport region may include, for example, at least one compound selected from Compounds ET1 to ET36:

ET1




-continued




ET2

-continued




ET5

ET3



ET6

ET7





-continued

-continued



ET10

ET8




ET11

ET9




ET12




-continued

ET13




-continued


ET16




ET14






ET17



ET15




-continued




-continued

ET18




ET21

ET19



ET22


ET20



ET23


-continued

ET24




-continued


ET27




ET25




ET28




ET26



ET29




-continued




ET30


-continued




ET33




ET31



ET34




ET32





ET35

-continued




ET36

[0358] In one or more embodiments, the electron transport region may include, for example, at least one selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq<sub>3</sub>, BAlq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ.


Alq<sub>3</sub>

BAlq



TAZ

-continued



NTAZ


[0359] A thickness of the buffer layer, the hole blocking layer, or the electron control layer may be in a range of, for example, about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.

[0360] A thickness of the electron transport layer may be, for example, in a range of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

[0361] The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.

[0362] The metal-containing material may include, for example, at least one of an alkali metal complex and an alkaline earth-metal complex. The alkali metal complex may include a metal ion selected from a Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion, and the alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, a Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenyloxazole, a hydroxy phenylthiazole, a hydroxy diphenyloxadiazole, a hydroxy diphenyl thiadiazol, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene.


[0363] For example, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.



ET-D1

-continued

ET-D2



**[0364]** The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode **190**. The electron injection layer may directly contact the second electrode **190**.

**[0365]** The electron injection layer may have, for example, i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

**[0366]** The electron injection layer may include, for example, an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or a combination thereof.

**[0367]** The alkali metal may be selected from Li, Na, K, Rb, and Cs. In an embodiment, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs.

**[0368]** The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

**[0369]** The rare earth metal may be selected from Sc, Y, Ce, Tb, Yb, and Gd.

**[0370]** The alkali metal compound, the alkaline earth-metal compound, and the rare earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal, and the rare earth metal.

**[0371]** The alkali metal compound may be selected from alkali metal oxides, such as  $\text{Li}_2\text{O}$ ,  $\text{Cs}_2\text{O}$ , or  $\text{K}_2\text{O}$ , and alkali metal halides, such as  $\text{LiF}$ ,  $\text{NaF}$ ,  $\text{CsF}$ ,  $\text{KF}$ ,  $\text{LiI}$ ,  $\text{NaI}$ ,  $\text{CsI}$ , or  $\text{KI}$ . In an embodiment, the alkali metal compound may be selected from  $\text{LiF}$ ,  $\text{Li}_2\text{O}$ ,  $\text{NaF}$ ,  $\text{LiI}$ ,  $\text{NaI}$ ,  $\text{CsI}$ , and  $\text{KI}$ .

**[0372]** The alkaline earth-metal compound may be selected from alkaline earth-metal oxides, such as  $\text{BaO}$ ,  $\text{SrO}$ ,  $\text{CaO}$ ,  $\text{Ba}_x\text{Sr}_{1-x}\text{O}$  ( $0 < x < 1$ ), and  $\text{Ba}_x\text{Ca}_{1-x}\text{O}$  ( $0 < x < 1$ ). In an embodiment, the alkaline earth-metal compound may be selected from  $\text{BaO}$ ,  $\text{SrO}$ , and  $\text{CaO}$ .

**[0373]** The rare earth metal compound may be selected from  $\text{YbF}_3$ ,  $\text{ScF}_3$ ,  $\text{ScO}_3$ ,  $\text{Y}_2\text{O}_3$ ,  $\text{Ce}_2\text{O}_3$ ,  $\text{GdF}_3$ , and  $\text{TbF}_3$ . In an embodiment, the rare earth metal compound may be selected from  $\text{YbF}_3$ ,  $\text{ScF}_3$ ,  $\text{TbF}_3$ ,  $\text{YbI}_3$ ,  $\text{ScI}_3$ , and  $\text{TbI}_3$ .

**[0374]** The alkali metal complex, the alkaline earth-metal complex, and the rare earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare earth metal as described above, and a ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, or the rare earth metal complex may be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy

benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenyloxazole, hydroxy phenylthiazole, hydroxy diphenyloxadiazole, hydroxy diphenylthiadiazole, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene.

**[0375]** The electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or a combination thereof, as described above. In one or more embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or a combination thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

**[0376]** A thickness of the electron injection layer may be in a range of, for example, about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

**[0377]** [Second Electrode **190**]

**[0378]** The second electrode **190** may be disposed on the organic layer **150** having such a structure. The second electrode **190** may be a cathode which is an electron injection electrode, and in this regard, a material for forming the second electrode **190** may be selected from, for example, a metal, an alloy, an electrically conductive compound, and a combination thereof, which may have a relatively low work function.

**[0379]** The second electrode **190** may include, for example, at least one selected from Li, Ag, Mg, Al, Al—Li, Ca, Mg—In, Mg—Ag, ITO, and IZO. The second electrode **190** may be, for example, a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

**[0380]** The second electrode **190** may have, for example, a single-layered structure, or a multi-layered structure including two or more layers.

**[0381]** [Description of FIGS. 2 to 4]

**[0382]** An organic light-emitting device **20** of FIG. 2 includes a first capping layer **210**, a first electrode **110**, an organic layer **150**, and a second electrode **190** which are sequentially stacked in this stated order, an organic light-emitting device **30** of FIG. 3 includes a first electrode **110**, an organic layer **150**, a second electrode **190**, and a second capping layer **220** which are sequentially stacked in this stated order, and an organic light-emitting device **40** of FIG. 4 includes a first capping layer **210**, a first electrode **110**, an organic layer **150**, a second electrode **190**, and a second capping layer **220** which are sequentially stacked in this stated order.

**[0383]** Regarding FIGS. 2 to 4, the first electrode **110**, the organic layer **150**, and the second electrode **190** may be understood by referring to the description presented in connection with FIG. 1.

[0384] In the organic layer 150 of each of the organic light-emitting devices 20 and 40, light generated in an emission layer may pass through the first electrode 110, which may be, for example, a semi-transmissive electrode or a transmissive electrode, and the first capping layer 210 toward the outside, and in the organic layer 150 of each of the organic light-emitting devices 30 and 40, light generated in an emission layer may pass through the second electrode 190, which may be, for example, a semi-transmissive electrode or a transmissive electrode, and the second capping layer 220 toward the outside.



[0385] The first capping layer 210 and the second capping layer 220 may increase external luminescent efficiency according to the principle of constructive interference.

[0386] The first capping layer 210 and the second capping layer 220 may each independently be, for example, an organic capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.




[0387] At least one selected from the first capping layer 210 and the second capping layer 220 may each independently include, for example, at least one material selected from carbocyclic compounds, heterocyclic compounds, amine-based compounds, porphrine derivatives, phthalocyanine derivatives, a naphthalocyanine derivatives, alkali metal complexes, and alkaline earth-based complexes. The carbocyclic compound, the heterocyclic compound, and the amine-based compound may be optionally substituted with a substituent containing at least one element selected from O, N, S, Se, Si, F, Cl, Br, and I. In an embodiment, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include an amine-based compound.

[0388] In an embodiment, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include, for example, the compound represented by Formula 201 or the compound represented by Formula 202.

[0389] In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include, for example, a compound selected from Compounds HT28 to HT33 and Compounds CP1 to CP5.



-continued



[0390] Layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region may be formed in a certain region by using one or more suitable methods selected from, for example, vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

[0391] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by vacuum deposition, the vacuum deposition may be performed, for example, at a deposition temperature of about 100° C. to about 500° C., at a vacuum degree of about 10<sup>-8</sup> torr to about 10<sup>-3</sup> torr, and at a deposition rate of about 0.01 Å/sec to about 100 Å/sec by taking into account a material to be included in a layer to be formed and the structure of a layer to be formed.

**[0392]** When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by spin coating, the spin coating may be performed, for example, at a coating speed of about 2000 rpm to about 5000 rpm and at a heat treatment temperature of about 80° C. to 200° C. by taking into account a material to be included in a layer to be formed and the structure of a layer to be formed.

**[0393]** [General Definition of Substituents]

**[0394]** The term “C<sub>1</sub>-C<sub>60</sub> alkyl group” as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “C<sub>1</sub>-C<sub>60</sub> alkylene group” as used herein refers to a divalent group having the same structure as the C<sub>1</sub>-C<sub>60</sub> alkyl group.

**[0395]** The term “C<sub>2</sub>-C<sub>60</sub> alkenyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon double bond in the middle or at the terminus of the C<sub>2</sub>-C<sub>60</sub> alkyl group, and examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C<sub>2</sub>-C<sub>60</sub> alkenylene group” as used herein refers to a divalent group having the same structure as the C<sub>2</sub>-C<sub>60</sub> alkenyl group.

**[0396]** The term “C<sub>2</sub>-C<sub>60</sub> alkynyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the middle or at the terminus of the C<sub>2</sub>-C<sub>60</sub> alkyl group, and examples thereof include an ethynyl group and a propynyl group. The term “C<sub>2</sub>-C<sub>60</sub> alkynylene group” as used herein refers to a divalent group having the same structure as the C<sub>2</sub>-C<sub>60</sub> alkynyl group.

**[0397]** The term “C<sub>1</sub>-C<sub>60</sub> alkoxy group” as used herein refers to a monovalent group represented by —OA<sub>101</sub> (wherein A<sub>101</sub> is the C<sub>1</sub>-C<sub>60</sub> alkyl group), and examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

**[0398]** The term “C<sub>3</sub>-C<sub>10</sub> cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C<sub>3</sub>-C<sub>10</sub> cycloalkylene group” as used herein refers to a divalent group having the same structure as the C<sub>3</sub>-C<sub>10</sub> cycloalkyl group.

**[0399]** The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group” as used herein refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a 1,2,3,4-oxatriazolidiny group, a tetrahydrofuranly group, and a tetrahydrothiophenyl group. The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkylene group” used herein refers to a divalent group having the same structure as the C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group.

**[0400]** The term “C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group” as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and no aromaticity, and examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group” as used herein refers to a divalent group having the same structure as the C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group.

**[0401]** The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Examples of the C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group include a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranly group, and a 2,3-dihydrothiophenyl group. The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkenylene group” as used herein refers to a divalent group having the same structure as the C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group.

**[0402]** The term “C<sub>6</sub>-C<sub>60</sub> aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C<sub>6</sub>-C<sub>60</sub> arylylene group used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Examples of the C<sub>6</sub>-C<sub>60</sub> aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C<sub>6</sub>-C<sub>60</sub> aryl group and the C<sub>6</sub>-C<sub>60</sub> arylylene group each include two or more rings, the rings may be fused to each other.

**[0403]** The term “C<sub>1</sub>-C<sub>60</sub> heteroaryl group” as used herein refers to a monovalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C<sub>1</sub>-C<sub>60</sub> heteroarylylene group” as used herein refers to a divalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Examples of the C<sub>1</sub>-C<sub>60</sub> heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinoxinyl group, and an isoquinolinyl group. When the C<sub>1</sub>-C<sub>60</sub> heteroaryl group and the C<sub>1</sub>-C<sub>60</sub> heteroarylylene group each include two or more rings, the rings may be fused to each other.

**[0404]** The term “C<sub>6</sub>-C<sub>60</sub> aryloxy group” as used herein indicates —OA<sub>102</sub> (wherein A<sub>102</sub> is the C<sub>6</sub>-C<sub>60</sub> aryl group), and the term “C<sub>6</sub>-C<sub>60</sub> arylythio group” as used herein indicates —SA<sub>103</sub> (wherein A<sub>103</sub> is the C<sub>6</sub>-C<sub>60</sub> aryl group).

**[0405]** The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group that has two or more rings condensed and only carbon atoms (e.g., 8 to 60 carbon atoms) as ring-forming atoms, wherein the entire molecular structure is non-aromatic. Examples of the monovalent non-aromatic condensed polycyclic group may include a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

**[0406]** The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group that has two or more rings condensed, at least one heteroatom selected from N, O, Si, P, and S, in addition to carbon atoms (e.g., 1 to 60 carbon atoms), as ring-forming atoms, wherein the entire molecular structure is non-aromatic. Examples of the monovalent non-aromatic condensed heteropolycyclic group may include a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

**[0407]** The term “C<sub>5</sub>-C<sub>60</sub> carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which a ring-forming atom is a carbon atom only. The C<sub>5</sub>-C<sub>60</sub> carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C<sub>5</sub>-C<sub>60</sub> carbocyclic group may be a ring, such as a benzene group, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In one or more embodiments, depending on the number of substituents connected to the C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the C<sub>5</sub>-C<sub>60</sub> carbocyclic group may be a trivalent group or a quadrivalent group.

**[0408]** The term “C<sub>1</sub>-C<sub>60</sub> heterocyclic group” as used herein refers to a group having the same structure as the C<sub>5</sub>-C<sub>60</sub> carbocyclic group, except that as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used in addition to carbon (the number of carbon atoms may be in a range of 1 to 60).

**[0409]** In the present specification, at least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkylene group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenylene group, the substituted C<sub>6</sub>-C<sub>60</sub> arylene group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>60</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

**[0410]** deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

**[0411]** a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —N(Q<sub>11</sub>)(Q<sub>12</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>).

**[0412]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

**[0413]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>50</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —N(Q<sub>21</sub>)(Q<sub>22</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>), and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and

**[0414]** —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>), and

**[0415]** Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

**[0416]** The term “Ph” as used herein refers to a phenyl group, the term “Me” as used herein refers to a methyl group, the term “Et” as used herein refers to an ethyl group, the term “ter-Bu” or “But” as used herein refers to a tert-butyl group, and the term “OMe” as used herein refers to a methoxy group.

**[0417]** The term “biphenyl group” as used herein refers to “a phenyl group substituted with a phenyl group.” In other words, the term “biphenyl group” is a substituted phenyl group having a C<sub>6</sub>-C<sub>60</sub> aryl group as a substituent.

**[0418]** The term “terphenyl group” as used herein refers to “a phenyl group substituted with a biphenyl group.” In other words, the “terphenyl group” is a phenyl group having, as a substituent, a C<sub>6</sub>-C<sub>60</sub> aryl group substituted with a C<sub>6</sub>-C<sub>60</sub> aryl group.

[0419] The symbols \* and \*\* as used herein unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.



[0420] Hereinafter, a compound according to embodiments and an organic light-emitting device according to embodiments will be described in detail with reference to Synthesis Examples and Examples. The wording "B was used instead of A" used in describing Synthesis Examples refers to that an identical molar equivalent of B was used in place of A.

[0421] The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples.

#### EXAMPLES

##### Synthesis Example 1: Synthesis of Compound 13

[0422]



13

##### Synthesis of Intermediate I-1

[0423] 1.99 g (10 mmol) of 2,3-dichloroquinoxaline, 2.27 g (12 mmol) of 4-bromobenzene-1,2-diol, and 4.15 g (30 mmol) of potassium carbonate were dissolved in 100 ml of DMF, and the mixed solution was stirred at room temperature for 24 hours. Distilled water was added to the reaction solution, and an extraction process was performed thereon

three times using 60 ml of ethyl acetate. An organic layer collected therefrom was dried using magnesium sulfate ( $\text{MgSO}_4$ ), and then, the residual obtained by evaporating a solvent therefrom was separation-purified by gel chromatography to obtain 2.68 g (yield: 85%) of Intermediate I-1. The obtained compound was confirmed by LC-MS.

[0424]  $\text{C}_{14}\text{H}_7\text{BrN}_2\text{O}_2$ ;  $M^+$  313.97

##### Synthesis of Compound 13

[0425] 3.15 g (10 mmol) of Intermediate I-1, 4.73 g (11 mmol) of 4,4,5,5-tetramethyl-2-(10-(naphthalen-1-yl)anthracen-9-yl)-1,3,2-dioxaborolane, 0.57 g (5 mol %) of  $\text{Pd}(\text{PPh}_3)_4$ , 4.15 g (3 equiv) of  $\text{K}_2\text{CO}_3$  were dissolved in 100 ml of a solution containing THF and  $\text{H}_2\text{O}$  at a volume ratio of 2:1, and then, the mixed solution was stirred at a temperature of  $80^\circ\text{C}$ . for 24 hours. After the reaction solution was cooled to room temperature, an extraction process was performed thereon three times using 60 ml of ethyl acetate. An organic layer collected therefrom was dried using  $\text{MgSO}_4$ , and then, the residual obtained by evaporating a solvent therefrom was separation-purified by gel chromatography to obtain 4.15 g (yield: 77%) of Compound 13. The obtained compound was confirmed by LC-MS.

[0426]  $\text{C}_{38}\text{H}_{22}\text{N}_2\text{O}_2$ ;  $M^+$  538.17

##### Synthesis Example 2: Synthesis of Compound 15

[0427] 3.08 g (yield: 63%) of Compound 15 was synthesized in the same manner as in Synthesis of Compound 13, except that 4-(10-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)pyridine was used instead of 4,4,5,5-tetramethyl-2-(10-(naphthalen-1-yl)anthracen-9-yl)-1,3,2-dioxaborolane. The obtained compound was identified by MS/FAB and  $^1\text{H}$  NMR.

[0428]  $\text{C}_{33}\text{H}_{19}\text{N}_3\text{O}_2$ ;  $M^+$  489.15

##### Synthesis Example 3: Synthesis of Compound 16

[0429] 3.70 g (yield: 72%) of Compound 16 was synthesized in the same manner as in the Synthesis of Compound 13, except that 4-(10-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)benzotrile was used instead of 4,4,5,5-tetramethyl-2-(10-(naphthalen-1-yl)anthracen-9-yl)-1,3,2-dioxaborolane. The obtained compound was identified by MS/FAB and  $^1\text{H}$  NMR.

[0430]  $\text{C}_{35}\text{H}_{19}\text{N}_3\text{O}_2$ ;  $M^+$  513.15

##### Synthesis Example 4: Synthesis of Compound 20

[0431] 3.92 g (yield: 61%) of Compound 20 was synthesized in the same manner as in the Synthesis of Compound 13, except that 3,3'-(5-(10-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)-1,3-phenylene)dipyridine was used instead of 4,4,5,5-tetramethyl-2-(10-(naphthalen-1-yl)anthracen-9-yl)-1,3,2-dioxaborolane. The obtained compound was identified by MS/FAB and  $^1\text{H}$  NMR.

[0432]  $\text{C}_{44}\text{H}_{26}\text{N}_4\text{O}_2$ ;  $M^+$  643.20

## Synthesis Example 5: Synthesis of Compound 21

[0433] 4.44 g (yield: 69%) of Compound 21 was synthesized in the same manner as in the Synthesis of Compound 13, except that 2,4-diphenyl-6-(10-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)-1,3,5-triazine was used instead of 4,4,5,5-tetramethyl-2-(10-(naphthalen-1-yl)anthracen-9-yl)-1,3,2-dioxaborolane. The obtained compound was identified by MS/FAB and <sup>1</sup>H NMR.

[0434] C<sub>43</sub>H<sub>25</sub>N<sub>5</sub>O<sub>2</sub>; M<sup>+</sup> 643.20

## Synthesis Example 6: Synthesis of Compound 27

[0435] 4.47 g (yield: 71%) of Compound 27 was synthesized in the same manner as in the Synthesis of Compound 13, except that 9,9-dimethyl-7-(10-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)-9H-fluorene-2-carbonitrile was used instead of 4,4,5,5-tetramethyl-2-(10-(naphthalen-1-yl)anthracen-9-yl)-1,3,2-dioxaborolane. The obtained compound was identified by MS/FAB and <sup>1</sup>H NMR.

[0436] C<sub>44</sub>H<sub>27</sub>N<sub>3</sub>O<sub>2</sub>; M<sup>+</sup> 629.21

benzonitrile was used instead of 4,4,5,5-tetramethyl-2-(10-(naphthalen-1-yl)anthracen-9-yl)-1,3,2-dioxaborolane. The obtained compound was identified by MS/FAB and <sup>1</sup>H NMR.

[0438] C<sub>35</sub>H<sub>19</sub>N<sub>3</sub>S<sub>2</sub>; M<sup>+</sup> 545.10

## Synthesis Example 8: Synthesis of Compound 59

[0439] 4.20 g (yield: 73%) of Compound 59 was synthesized in the same manner as in the Synthesis of Compound 52, except that 2,4-diphenyl-6-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1,3,5-triazine was used instead of 4-(10-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)benzonitrile. The obtained compound was identified by MS/FAB and <sup>1</sup>H NMR.

[0440] C<sub>35</sub>H<sub>21</sub>N<sub>5</sub>S<sub>2</sub>; M<sup>+</sup> 575.12

[0441] The <sup>1</sup>H NMR and MS/FAB results of the compounds synthesized according to Synthesis Examples 1 to 8 are shown in Table 1.

[0442] Methods of synthesizing compounds other than the compound shown in Table 1 are recognizable by one of ordinary skill in the art by referring to the synthesis path and source materials described above.

TABLE 1


Compound	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500 MHz)	HR-EIMS	
		found	calc.
13	δ = 7.96-7.92 (m, 2H), 7.83-7.75 (m, 5H), 7.71-7.69 (m, 2H), 7.67-7.64 (m, 1H), 7.55-7.49 (m, 3H), 7.46-7.44 (m, 1H), 7.39-7.30 (m, 7H), 6.97-6.95 (m, 1H)	538.15	538.17
15	δ = 8.81-7.78 (m, 2H), 7.97-7.87 (m, 4H), 7.76-7.71 (m, 4H), 7.66-7.64 (m, 1H), 7.55-7.49 (m, 2H), 7.38-7.34 (m, 2H), 7.32-7.29 (m, 4H)	489.16	489.15
16	δ = 7.96-7.94 (m, 2H), 7.86-7.80 (m, 4H), 7.75-7.72 (m, 2H), 7.69-7.64 (m, 3H), 7.53-7.49 (m, 2H), 7.38-7.31 (m, 6H)	513.13	513.15
20	δ = 8.95-8.92 (m, 2H), 8.69-8.66 (m, 2H), 8.11-8.07 (m, 4H), 7.95-7.93 (m, 2H), 7.89-7.83 (m, 3H), 7.70-7.63 (m, 3H), 7.53-7.45 (m, 4H), 7.39-7.31 (m, 6H)	642.22	642.21
21	δ = 8.87-8.83 (m, 4H), 8.28-8.26 (m, 2H), 7.95-7.93 (m, 2H), 7.88-7.85 (m, 2H), 7.66-7.60 (m, 5H), 7.58-7.56 (m, 2H), 7.53-7.50 (m, 4H), 7.41-7.33 (m, 4H)	643.19	643.20
27	δ = 8.01-7.92 (m, 5H), 7.81-7.79 (m, 2H), 7.69-7.64 (m, 4H), 7.53-7.49 (m, 3H), 7.46-7.44 (m, 1H), 7.39-7.31 (m, 6H), 1.64 (s, 6H)	629.22	629.21
52	δ = 8.09-8.07 (m, 1H), 7.92-7.79 (m, 6H), 7.77-7.67 (m, 4H), 7.60-7.57 (m, 4H), 7.37-7.29 (m, 4H)	545.09	545.10
59	δ = 8.81-8.78 (m, 4H), 8.58-8.53 (m, 2H), 8.02-8.00 (m, 1H), 7.83-7.75 (m, 5H), 7.62-7.51 (m, 7H), 7.41-7.39 (m, 2H)	575.13	575.12

## Synthesis Example 7: Synthesis of Compound 52

[0437] 3.44 g (yield: 68%) of Compound 52 was synthesized in the same manner as in the Synthesis of Intermediate I-1, except that 4-bromobenzene-1,2-dithiol was used instead of 4-bromobenzene-1,2-diol, and in the same manner as in the Synthesis of Compound 13, except that 4-(10-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)

## Example 1

[0443] An anode was prepared by cutting a Corning 15 Ω/cm<sup>2</sup> (1.200 Å) ITO glass substrate to a size of 50 mm×50 mm×0.7 mm, sonicating the glass substrate by using isopropyl alcohol and pure water for 5 minutes each, and then irradiating UV light for 30 minutes to ozone to clean the glass substrate. Then, the anode was loaded onto a vacuum deposition apparatus.



[0444] Then, 2-TNATA, which is a material for forming a hole injection layer, was vacuum-deposited on the substrate to form a hole injection layer having a thickness of 600 Å, and then, 4,4'-bis[N-(1-naphthyl)-N-phenylamino]biphenyl (hereinafter, referred to as NPB), which is a known hole transporting compound, was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å. 9,10-di-naphthalene-2-yl-anthracene (hereinafter, referred to as ADN), which is a blue fluorescent host, and 4,4'-bis[2-(4-(N,N-diphenylamino)phenyl)vinyl]biphenyl (hereinafter, referred to as DPAVBi), which is a blue fluorescent dopant, were co-deposited on the hole

transport layer at a weight ratio of 98:2 to form an emission layer having a thickness of 300 Å.

[0445] Then, Compound 13 synthesized according to the embodiments of the present disclosure was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å, and then, LiF, which is a halogenated alkali metal, was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited thereon to a thickness of 3,000 Å (i.e., a cathode electrode), thereby forming an LiF/Al electrode and completing the manufacture of an organic light-emitting device.


Examples 2 to 8 and Comparative Examples 1 and

3


[0446] Organic light-emitting devices were manufactured in the same manner as in Example 1, except that in forming an electron transport layer, compounds shown in Table 2 were used instead of Compound 1.

TABLE 2


	Material	Driving voltage (V)	Current density (mA/cm <sup>2</sup> )	Brightness (cd/m <sup>2</sup> )	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm <sup>2</sup> )
Example 1	Compound 13	4.65	50	3,625	7.23	Blue	361
Example 2	Compound 15	4.61	50	3,845	7.69	Blue	378
Example 3	Compound 16	4.53	50	3,625	7.25	Blue	352
Example 4	Compound 20	4.49	50	3,945	7.89	Blue	368
Example 5	Compound 21	4.45	50	3,615	7.23	Blue	370
Example 6	Compound 27	4.70	50	3,525	7.05	Blue	421
Example 7	Compound 52	5.00	50	3,375	6.75	Blue	341
Example 8	Compound 59	4.95	50	3,375	6.75	Blue	352
Comparative Example 1	Compound 200	5.06	50	3,010	6.52	Blue	325
Comparative Example 7	Compound A	5.12	50	3,165	6.33	Blue	307
Comparative Example 3	Compound B	5.08	50	3,225	6.45	Blue	331



200



Compound A



Compound B


[0447] Referring to Table 2, it was confirmed that the organic light-emitting devices of Examples 1 to 8 had a low driving voltage, high efficiency, high brightness, and long lifespan characteristics, compared to those of the organic light-emitting devices of Comparative Examples 1 to 3.

[0448] As described above, an organic light-emitting device including a heterocyclic compound may have a low driving voltage, high efficiency, high luminance, and a long lifespan.

[0449] Example embodiments have been disclosed herein, and although specific terms are employed, they are used and are to be interpreted in a generic and descriptive sense only and not for purpose of limitation. In some instances, as would be apparent to one of ordinary skill in the art as of the filing of the present application, features, characteristics, and/or elements described in connection with a particular embodiment may be used singly or in combination with features, characteristics, and/or elements described in connection with other embodiments unless otherwise specifically indicated. Accordingly, it will be understood by those of skill in the art that various changes in form and details may be made without departing from the spirit and scope of the present invention as set forth in the following claims.

What is claimed is:

1. A heterocyclic compound represented by Formula 1:



wherein, in Formula 1,

A<sub>1</sub> and A<sub>2</sub> are each independently a C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a C<sub>2</sub>-C<sub>60</sub> heterocyclic group,

X and Y are each independently oxygen (O) or sulfur (S),

L<sub>1</sub> and L<sub>2</sub> are each independently selected from a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heterocyclic group, \*—Si(Q<sub>1</sub>)(Q<sub>2</sub>)-\*, \*—N(Q<sub>1</sub>)-\*, \*—B(Q<sub>1</sub>)-\*, \*—C(=O)-\*, \*—S(=O)<sub>2</sub>-\*, and \*—P(=O)(Q<sub>1</sub>)-\*,

a<sub>1</sub> and a<sub>2</sub> are each independently an integer selected from 0 to 5,

when a<sub>1</sub> is 0, \*-(L<sub>1</sub>)<sub>a1</sub>-\* is a single bond, and when a<sub>2</sub> is 0, \*-(L<sub>2</sub>)<sub>a2</sub>-\* is a single bond,

Ar<sub>1</sub> and Ar<sub>2</sub> are each independently selected from a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —N(Q<sub>1</sub>)(Q<sub>2</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —S(=O)<sub>2</sub>(Q<sub>1</sub>), and —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>),

b<sub>1</sub> and b<sub>2</sub> are each independently an integer selected from 1 to 5,

R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —C(=O)(Q<sub>1</sub>), —S(=O)<sub>2</sub>(Q<sub>1</sub>), and —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>),

c<sub>1</sub> and c<sub>2</sub> are each independently an integer selected from 1 to 7,

m and n are each independently an integer selected from 0 to 8, provided that m+n≥1,

at least one substituent of the substituted C<sub>3</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group;

a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —N(Q<sub>11</sub>)(Q<sub>12</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub>

aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

- a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —N(Q<sub>21</sub>)(Q<sub>22</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>), and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>).

Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

\* and \*' each indicate a binding site to a neighboring atom.

2. The heterocyclic compound as claimed in claim 1, wherein A<sub>1</sub> and A<sub>2</sub> are each independently selected from a benzene group, a naphthalene group, an anthracene group, and a phenanthrene group.

3. The heterocyclic compound as claimed in claim 1, wherein X and Y are identical to each other.

4. The heterocyclic compound as claimed in claim 1, wherein L<sub>1</sub> and L<sub>2</sub> are each independently a divalent group based on a group selected from:

a benzene group, a pentalene group, an indene group, a naphthalene group, an azulene group, an indacene group, an acenaphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a perylene group, a pentacene group, a pyrrole group, a thiophene group, a furan group, a silole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine

group, a pyridazine group, an indole group, an isoindole group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a benzoisoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, a benzofuran group, a benzothiophene group, a benzosilole group, a benzothiazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a carbazole group, a dibenzofuran group, a dibenzothiophene group, a dibenzosilole group, a benzocarbazole group, a naphthobenzofuran group, a naphthobenzothiophene group, a naphthobenzosilole group, a dibenzocarbazole group, a dinaphthofuran group, a dinaphthothiophene group, a dinaphthosilole group, a thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, an oxazolpyridine group, a thiazolpyridine group, a benzonaphthyridine group, an azafluorene group, an azaspiro-bifluorene group, an azacarbazole group, an azadibenzofuran group, an azadibenzothiophene group, an azadibenzosilole group, an indenopyrrole group, an indolpyrrole group, an indenocarbazole group, and an indolcarbazole group;

a benzene group, a pentalene group, an indene group, a naphthalene group, an azulene group, an indacene group, an acenaphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a perylene group, a pentacene group, a pyrrole group, a thiophene group, a furan group, a silole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indole group, an isoindole group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a benzoisoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, a benzofuran group, a benzothiophene group, a benzosilole group, a benzothiazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a carbazole group, a dibenzofuran group, a dibenzothiophene group, a dibenzosilole group, a benzocarbazole group, a naphthobenzofuran group, a naphthobenzothiophene group, a naphthobenzosilole group, a dibenzocarbazole group, a dinaphthofuran group, a dinaphthothiophene group, a dinaphthosilole group, a thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, an oxazolpyridine group, a thiazolpyridine group, a benzonaphthyridine group, an azafluorene group, an azaspiro-bifluorene group, an azacarbazole group, an azadibenzofuran group, an azadibenzothiophene group, an azadibenzosilole group, an indenopyrrole group, an indolpyrrole group,

an indenocarbazole group, and an indolcarbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>20</sub> aryl group, a C<sub>1</sub>-C<sub>20</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a methylphenyl group, a biphenyl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>); and

\*—P(=O)(Q<sub>1</sub>)\*,

Q<sub>1</sub> and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from:

a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group; and


a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group, each substituted with at least one selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, and a phenyl group, and

\* and \*' each indicate a binding site to a neighboring atom.


5. The heterocyclic compound as claimed in claim 1, wherein:

at least one of a1 and a2 is nonzero, and


L<sub>1</sub> and L<sub>2</sub> are each independently a group represented by Formulae 3-1 to 3-47:




Formula 3-1



Formula 3-2




Formula 3-3




Formula 3-4


-continued




Formula 3-5




Formula 3-6




Formula 3-7




Formula 3-8




Formula 3-9




Formula 3-10




Formula 3-11



Formula 3-12














Formula 3-13




Formula 3-14


-continued



Formula 3-15




Formula 3-16




Formula 3-17

Formula 3-18




Formula 3-19

Formula 3-20




Formula 3-21




Formula 3-22


Formula 3-23



Formula 3-24



Formula 3-25



-continued

Formula 3-26

Formula 3-27

Formula 3-28

Formula 3-29






Formula 3-30

Formula 3-31

Formula 3-32


Formula 3-33

-continued




-continued


Formula 3-34




Formula 3-35





Formula 3-36



Formula 3-37



Formula 3-38



Formula 3-39

Formula 3-40


Formula 3-41

Formula 3-42


Formula 3-43

Formula 3-44


-continued



Formula 3-45



Formula 3-46



Formula 3-47

wherein, in Formulae 3-1 to 3-47,

$Y_1$  is O, S,  $C(Z_3)(Z_4)$ ,  $N(Z_5)$ , or  $Si(Z_6)(Z_7)$ ,

$Z_1$  to  $Z_7$  are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a quinazolinyl group, a benzoquinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothio-phenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolpyridinyl group, a thiazolpyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolpyrrolyl group, an indenocarbazolyl group, and an indolcarbazolyl group,

lityl group, a dibenzocarbazolyl group, a dinaphthofuranyl group, a dinaphthothiophenyl group, a dinaphthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolpyridinyl group, a thiazolpyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolpyrrolyl group, an indenocarbazolyl group, and an indolcarbazolyl group,

$d_2$  is an integer selected from 0 to 2,

$d_3$  is an integer selected from 0 to 3,

$d_4$  is an integer selected from 0 to 4,

$d_5$  is an integer selected from 0 to 5,

$d_6$  is an integer selected from 0 to 6,

$d_8$  is an integer selected from 0 to 8, and

\* and \*' each indicate a binding site to a neighboring atom.

6. The heterocyclic compound as claimed in claim 1, wherein  $a_1$  and  $a_2$  are each independently 0, 1, 2, or 3.

7. The heterocyclic compound as claimed in claim 1, wherein  $Ar_1$  and  $Ar_2$  are each independently selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothi-


azolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoanthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>); and —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —C(=O)(Q<sub>1</sub>), —S(=O)<sub>2</sub>(Q<sub>1</sub>), and —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and

Q<sub>1</sub> to Q<sub>3</sub> and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from:


a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinyl group, each substituted with at least one selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, and a phenyl group.


8. The heterocyclic compound as claimed in claim 1, wherein Ar<sub>1</sub> and Ar<sub>2</sub> are each independently selected from groups represented by Formulae 5-1 to 5-84:




Formula 5-1




Formula 5-2




Formula 5-3




Formula 5-4




Formula 5-5




Formula 5-6



Formula 5-7










Formula 5-8




Formula 5-9

-continued




-continued


Formula 5-10




Formula 5-11




Formula 5-12




Formula 5-13






Formula 5-14



Formula 5-15



Formula 5-16



Formula 5-17

Formula 5-18

Formula 5-19

Formula 5-20

Formula 5-21












Formula 5-22

Formula 5-23


Formula 5-24

Formula 5-25


-continued




Formula 5-26




Formula 5-27




Formula 5-28




Formula 5-29




Formula 5-30




Formula 5-31




Formula 5-32




Formula 5-33






Formula 5-34



Formula 5-35



Formula 5-36



-continued

Formula 5-37

Formula 5-38

Formula 5-39

Formula 5-40

Formula 5-41

Formula 5-42

Formula 5-43

Formula 5-44

Formula 5-45

Formula 5-46










Formula 5-47

Formula 5-48

Formula 5-49



-continued



Formula 5-72

Formula 5-73

Formula 5-74

Formula 5-75

Formula 5-76





Formula 5-77

Formula 5-78

Formula 5-79

Formula 5-80

-continued



Formula 5-81

Formula 5-82

Formula 5-83

Formula 5-84

wherein, in Formulae 5-1 to 5-84,

Y<sub>31</sub> is O, S, C(Z<sub>35</sub>)(Z<sub>36</sub>), N(Z<sub>37</sub>), or Si(Z<sub>38</sub>)(Z<sub>39</sub>),Z<sub>31</sub> to Z<sub>39</sub> are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-benzofluorene-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentacenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, an isoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a benzoquinoxalinyl group, a qui-

nazolanyl group, a benzoquinazolanyl group, a cinnolanyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranlyl group, a benzothio-phenyl group, a benzosilolyl group, a benzothiazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a carbazolyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a naphthobenzofuranlyl group, a naphthobenzothiophenyl group, a naphthobenzosilolyl group, a dibenzocarbazolyl group, a dinaphtho-furanlyl group, a dinaphthothiophenyl group, a dina-phthosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an oxazolpyridinyl group, a thiazolpyridinyl group, a benzonaphthyridinyl group, an azafluorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranlyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group, an indenopyrrolyl group, an indolpyrrolyl group, an indenocarbazolyl group, and an indolcarbazolyl group,

e2 is an integer selected from 0 to 2,

e3 is an integer selected from 0 to 3,

e4 is an integer selected from 0 to 4,

e5 is an integer selected from 0 to 5,

e6 is an integer selected from 0 to 6,

e7 is an integer selected from 0 to 7,

e9 is an integer selected from 0 to 9, and

\* indicates a binding site to a neighboring atom.

9. The heterocyclic compound as claimed in claim 1, wherein b1 and b2 are each independently an integer selected from 1 to 4.

10. The heterocyclic compound as claimed in claim 1, wherein R<sub>1</sub> and R<sub>2</sub> are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenan-threnyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chry-senyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenyl group, a corone-nyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranlyl group, a benzothiophenyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group;

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenan-

threnyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chry-senyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenyl group, a corone-nyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranlyl group, a benzothiophenyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphe-nyl group, a terphenyl group, a phenyl group substi-tuted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substi-tuted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenan-threnyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thio-phenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranlyl group, a benzothiophenyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>); and —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), and

Q<sub>1</sub> to Q<sub>3</sub> and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from:

a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazi-nyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinylyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazolinylyl group, each substituted with at least one selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, and a phenyl group.

11. The heterocyclic compound as claimed in claim 1, wherein the sum of m and n is 1 or 2.

12. The heterocyclic compound as claimed in claim 1, wherein R<sub>1</sub> and R<sub>2</sub> are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and C<sub>1</sub>-C<sub>20</sub> alkoxy group; and

a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium,

—F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group, and

m and n are selected from:

m is 0 and n is 1;






m is 1 and n is 0;

m is 0 and n is 2;

m is 1 and n is 1; or


m is 2 and n is 0.

13. The heterocyclic compound as claimed in claim 1, wherein the heterocyclic compound is selected from groups represented by Formulae 1-1 to 1-9:




-continued


Formula 1-6




Formula 1-7



Formula 1-8




Formula 1-9



wherein, in Formulae 1-1 to 1-9, descriptions of X, Y, L<sub>1</sub>, L<sub>2</sub>, a<sub>1</sub>, a<sub>2</sub>, Ar<sub>1</sub>, Ar<sub>2</sub>, b<sub>1</sub>, b<sub>2</sub>, R<sub>1</sub>, R<sub>2</sub>, c<sub>1</sub>, c<sub>2</sub>, m, and n are the same as defined in claim 1.


14. The heterocyclic compound as claimed in claim 1, wherein the heterocyclic compound is represented by Formulae 1-11 to 1-19:

Formula 1-11




-continued

Formula 1-12





-continued

Formula 1-17





Formula 1-18

Formula 1-13



Formula 1-19

Formula 1-14




Formula 1-15

\* —(L<sub>1</sub>)<sub>a1</sub>—(Ar<sub>1</sub>)<sub>b1</sub>


&lt;Formula 2-1&gt;

\* —(L<sub>2</sub>)<sub>a2</sub>—(Ar<sub>2</sub>)<sub>b2</sub>

&lt;Formula 2-2&gt;



Formula 1-16



wherein, in Formulae 1-11 to 1-19,

descriptions of R<sub>11</sub> to R<sub>18</sub> are the same as defined in connection with R<sub>1</sub> and descriptions of R<sub>21</sub> to R<sub>28</sub> are the same as defined in connection with R<sub>2</sub>,at least one selected from R<sub>11</sub> to R<sub>14</sub> is a binding site to the group represented by Formula 2-1, or at least one selected from R<sub>21</sub> to R<sub>24</sub> is a binding site to the group represented by Formula 2-2, anddescriptions of X, Y, L<sub>1</sub>, L<sub>2</sub>, a<sub>1</sub>, a<sub>2</sub>, Ar<sub>1</sub>, Ar<sub>2</sub>, b<sub>1</sub>, and b<sub>2</sub> are the same as claimed in claim 1.

**15.** The heterocyclic group as claimed in claim 14, wherein Formula 1 is represented by Formula 1-11, wherein, in Formula 1-11,

(i) R<sub>11</sub> to R<sub>14</sub> are each independently selected from:hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group; and

a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group,

R<sub>21</sub> to R<sub>24</sub> are each independently selected from:

the group represented by Formula 2-2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group; and

a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group, and

at least one selected from R<sub>21</sub> to R<sub>24</sub> is the group represented by Formula 2-2, or

(ii) R<sub>11</sub> to R<sub>14</sub> are each independently selected from:

the group represented by Formula 2-1, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group; and

a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group,

at least one selected from R<sub>11</sub> to R<sub>14</sub> is the group represented by Formula 2-1, and

R<sub>21</sub> to R<sub>24</sub> are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group; and

a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group.

**16.** The heterocyclic compound as claimed in claim **14**, wherein Formula 1 is represented by Formula 1-11, wherein, in Formula 1-11,

R<sub>11</sub> to R<sub>14</sub> are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group; and

a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group,

R<sub>21</sub> and R<sub>24</sub> are each independently selected from:


the group represented by Formula 2-2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group; and

a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium,

—F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, and a hydrazono group, and


at least one selected from R<sub>22</sub> and R<sub>23</sub> is the group represented by Formula 2-2.

**17.** The heterocyclic compound as claimed in claim **1**, wherein the heterocyclic compound is a compound selected from Compounds 1 to 110:




-continued


7




8




9




10



11




12




-continued


13




14




15



16




17




-continued


18




19




20



21




22




-continued


23




24



25




26




-continued


27




28




29



30




31




-continued


32




33




34




35




36




37



-continued




-continued




-continued


51




52




53



54




55




-continued


56




57



58




59






-continued

68




-continued


73




69




74




70



75





71




76

72




-continued


77



78




79




-continued


80



81




82




-continued


83




84



85




86




-continued


87



88




89




-continued


90




91




92



93




94




-continued


95




96



97




98




-continued


99




100



101




102




-continued


103




104




105




106



107




-continued




108

-continued

110



109



- 18.** An organic light-emitting device comprising:  
 a first electrode;  
 a second electrode facing the first electrode; and  
 an organic layer between the first electrode and the second electrode and comprising an emission layer,  
 wherein the organic layer comprises at least one heterocyclic compound of claim 1.
- 19.** The organic light-emitting device as claimed in claim 18, wherein the first electrode is an anode, the second electrode is a cathode, and the organic layer comprises a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode,  
 wherein the hole transport region comprises a hole injection layer, a hole transport layer, an emission auxiliary layer, an electron blocking layer, or a combination thereof, and  
 the electron transport region comprises a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, an electron injection layer, or a combination thereof.
- 20.** The organic light-emitting device as claimed in claim 18, wherein the organic layer comprises an electron transport region, wherein the electron transport region comprises the at least one heterocyclic compound.

\* \* \* \* \*

专利名称(译)	杂环化合物和包括其的有机发光装置		
公开(公告)号	<a href="#">US20180198074A1</a>	公开(公告)日	2018-07-12
申请号	US15/665536	申请日	2017-08-01
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	LEE HYOYOUNG PARK JUNHA SIM MUNKI JEONG EUNJAE KIM YOUNGKOOK HWANG SEOKHWAN		
发明人	LEE, HYOYOUNG PARK, JUNHA SIM, MUNKI JEONG, EUNJAE KIM, YOUNGKOOK HWANG, SEOKHWAN		
IPC分类号	H01L51/00 C07D491/056 C07F9/6561 C07D495/04 C07D519/00		
CPC分类号	H01L51/0071 H01L51/5092 H01L51/0067 C07F9/6561 H01L51/0052 H01L51/0058 H01L51/0056 H01L51/0072 C07D495/04 C07D519/00 H01L51/5206 H01L51/5221 H01L51/5012 H01L51/5088 H01L51/5056 H01L51/5096 H01L51/5072 C07D491/056 C09K11/06 C09K2211/1007 C09K2211/1011 C09K2211/1029 C09K2211/1059 C09K2211/1088 C09K2211/1092 C09K2211/1096 H01L51/441		
优先权	1020170002556 2017-01-06 KR		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

本发明提供杂环化合物和包含该杂环化合物的有机发光装置。杂环化合物由式1表示：

<Formula 1>

