

(10) **Patent No.:** **US 9,818,952 B2**
(45) **Date of Patent:** **Nov. 14, 2017**

- | | | | | |
|--------------|------|---------|--------------------|------------------------|
| 2005/0156164 | A1 | 7/2005 | Sotoyama | |
| 2005/0221124 | A1 * | 10/2005 | Hwang | C07F 9/5728
428/690 |
| 2008/0193797 | A1 | 8/2008 | Heil et al. | |
| 2010/0013381 | A1 | 1/2010 | Stoessel et al. | |
| 2010/0032658 | A1 | 2/2010 | Lee et al. | |
| 2011/0284831 | A1 * | 11/2011 | Kaiser | H01L 51/0056
257/40 |
| 2012/0228584 | A1 * | 9/2012 | Wigglesworth | C07D 493/06
257/40 |

CN	103497758	A	*	1/2014
EP	2100941	A2	*	12/2008
JP	2009-033067	A		2/2009
JP	2010-034451	A		2/2010
JP	2010-109360	A	*	5/2010
JP	2013-161860	A	*	8/2013
KR	10-2006-0006760	A		1/2006
KR	10-2008-0049770	A		6/2008
KR	10-2009-0033493	A		4/2009
KR	10-2010-0007780	A		1/2010
WO	WO-2012/070226	A1		5/2012

Machine translation of JP 2013-161860 A (publication date Aug. 2013).*

Machine translation of JP 2010-109360 A (publication date May 2010).*


Machine translation of CN 103497758 A (publication date Jan. 2014).*

* cited by examiner

Primary Examiner — Dawn Garrett

(74) *Attorney, Agent, or Firm* — Lee & Morse, P.C.

A condensed cyclic compound and an organic light-emitting device, the compound being represented by Formula 1:

$$\langle \text{Formula 1} \rangle$$


- (56) **References Cited**
- U.S. PATENT DOCUMENTS

- | | | | | |
|--------------|------|--------|---------------|-------------------------|
| 7,053,255 | B2 | 5/2006 | Ikeda et al. | |
| 7,233,019 | B2 | 6/2007 | Ionkin et al. | |
| 2004/0185300 | A1 * | 9/2004 | Hatwar | H01L 51/5064
428/690 |

10

190
150
110

CONDENSED CYCLIC COMPOUND AND ORGANIC LIGHT EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

Korean Patent Application No. 10-2013-0126105, filed on Oct. 22, 2013, in the Korean Intellectual Property Office, and entitled: "Condensed Cyclic Compound And Organic Light Emitting Device Comprising The Same," is incorporated by reference herein in its entirety.

BACKGROUND

1. Field

Embodiments relate to a condensed cyclic compound and an organic light-emitting device including the same.

2. Description of the Related Art


Organic light-emitting devices (OLEDs), which are self-emitting devices, may have advantages such as wide viewing angles, excellent contrast, quick response, high brightness, excellent driving voltage characteristics, and may provide multicolored images.

An organic light-emitting device may have a structure in which a first electrode, a hole transport region, an emission layer, an electron transport region, and a second electrode are sequentially disposed in this order on a substrate. Holes injected from the first electrode may move to the emission layer via the hole transport region, while electrons injected from the second electrode may move to the emission layer via the electron transport region. Carriers, such as the holes and electrons, may recombine in the emission layer to generate excitons. When the excitons drop from an excited state to a ground state, light is emitted.

SUMMARY

Embodiments are directed to a condensed cyclic compound and an organic light-emitting device including the same.

According to one or more embodiments, there is provided a condensed cyclic compound represented by Formula 1:



wherein, in Formula 1,

L_1 and L_2 are each independently a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_2 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic

group, or a substituted or unsubstituted divalent non-aromatic hetero-condensed polycyclic group,

a_1 and a_2 are each independently an integer of 0 to 5,

R_1 to R_4 are each independently a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group.

R_{11} to R_{13} are each independently a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, $-S(Q_1)(Q_2)(Q_3)$, or $-B(Q_4)(Q_5)$,

b_1 and b_2 are each independently an integer of 1 to 3, and b_3 is an integer of 1 to 4,

wherein at least one substituent of the substituted C_3 - C_{10} cycloalkylene group, the substituted C_2 - C_{10} heterocycloalkylene group, the substituted C_3 - C_{10} cycloalkenylene group, the substituted C_2 - C_{10} heterocycloalkenylene group, the substituted C_6 - C_{60} arylene group, the substituted C_2 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_2 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_2 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_2 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic hetero-condensed polycyclic group, the substituted C_2 - C_{60} alkynyl group, and the substituted C_1 - C_{60} alkoxy group is

a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, or a C_1 - C_{60} alkoxy group;

a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_3 - C_{60} alkynyl group, or a C_1 - C_{60} alkoxy group, each substituted with at least one of a deuterium $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof,

3

thereof, a C₃-C₁₀ cyclo alkyl group, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₁₃)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

a C₃-C₁₀ cyclo alkyl group, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic hetero-condensed polycyclic group;

a C₃-C₁₀ cyclo alkyl condensed, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic hetero-condensed polycyclic group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cyclo alkyl group, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇); or

—Si(Q₃₁)(Q₃₂)(Q₃₃), or —B(Q₃₄)(Q₃₅),

wherein Q₁ to Q₅, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₅ are each independently a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cyclo alkyl group, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic hetero-condensed polycyclic group.

According to one or more embodiments, an organic light-emitting device includes: a first electrode; a second electrode disposed opposite to the first electrode; and an organic layer disposed between the first electrode and the second electrode and including an emission layer, wherein the organic layer includes at least one of the condensed cyclic compounds of Formula 1 above.

BRIEF DESCRIPTION OF THE DRAWING

Features will be apparent to those of skill in the art by describing in detail exemplary embodiments with reference to the attached drawing in which:

The FIGURE illustrates a schematic view of a structure of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION

Example embodiments will now be described more fully hereinafter with reference to the accompanying drawing;

4


however, they may be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey exemplary implementations to those skilled in the art.

In the drawing FIGURE, the dimensions of layers and regions may be exaggerated for clarity of illustration. Like reference numerals refer to like elements throughout.

As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

According to an embodiment, there may be provided a condensed cyclic compound represented by Formula 1.

<Formula 1>




In formula 1, L₁ and L₂ may be each independently:

a phenylene group, a pentalenylene group, an indeylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indazenylene group, an acenaphthylenylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a pentalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, a ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzoimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzooxazolylene group, an isobenzooxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, or a dibenzocarbazolylene group; or

a phenylene group, a pentalenylene group, an indeylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indazenylene group, an acenaphthylenylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a













phenalenylene group, a phenanthrenylene group, an anthrazenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylene group, a cinnolinylenylene group, a carbazolylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylene group, a benzoimidazolylene group, a benzofuranylenylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzooxazolylene group, an isobenzooxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylenylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, or a dibenzocarbazolylenylene, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, an C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indeyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluorantenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a pycenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and Si(Q₃₁)(Q₃₂)(Q₃₃) (where Q₃₁ to Q₃₃ are each independently a hydrogen, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, or a naphthyl group).

For example, L1 and L2 in Formula 1 may be each independently represented by one of Formulae 3-1 to 3-32.



7

-continued




8

-continued


Formula 3-10

5




Formula 3-11

10




Formula 3-12

15




Formula 3-13

20




Formula 3-14

25




Formula 3-15

30




Formula 3-16

35



Formula 3-17

40




Formula 3-18

45

Formula 3-19

50




Formula 3-20

55

Formula 3-21

60



65

Formula 3-22

Formula 3-23

Formula 3-24

Formula 3-25

Formula 3-26

Formula 3-27

Formula 3-28


Formula 3-29

Formula 3-30

Formula 3-31

9

-continued



In formulae 3-1 to 3-32,

Z_1 and Z_2 may be each independently a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, or —Si(Q_{31})(Q_{32})(Q_{33}) (where Q_{31} to Q_{33} may be each independently, a hydrogen, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, or a naphthyl group;

d_1 may be an integer of 1 to 4;

d_2 may be an integer of 1 to 3;

d_3 may be an integer of 1 to 6;





d_4 may be an integer of 1 to 8;

d_5 may be 1 or 2; and

d_6 may be an integer of 1 to 5.

In Formulae 3-1 to 3-32, * and *' are binding sites. For example, * may be a binding site to a core, L_1 , or L_2 in Formula 1, and *' may be a binding site to L_1 , L_2 , or N in Formula 1.

For example, L_1 and L_2 in Formula 1 may be each independently represented by one of Formulae Formula 4-1 to 4-23:



Formula 3-32

5

10

30

35

40

Formula 4-1

Formula 4-2

Formula 4-3

Formula 4-4

45

50









55

60

65

10

-continued



Formula 4-5

Formula 4-6

Formula 4-7

Formula 4-8

Formula 4-9









Formula 4-10

Formula 4-11

Formula 4-12

11


-continued





12

-continued

Formula 4-13



Formula 4-14



Formula 4-15

In Formula 1, a₁, which indicates the number of L₁s, may be an integer of 0 to 5, e.g., may be 0, 1, or 2. When a₁ is 0, L₁ may be a single bond. When a₁ is 2 or greater, a plurality of L₁s may be identical to or different from each other.

Formula 4-16

In Formula 1, a₂, which indicates the number of L₂s, may be an integer of 0 to 5, e.g., may be 0, 1, or 2. When a₂ is 0, L₂ may be a single bond. When a₂ is 2 or greater, a plurality of L₂s may be identical to or different from each other.

Formula 4-17

In Formula 1, R₁ to R₄ may be each independently: a phenyl group, a pentalenyl group, an indeyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluorantenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a pycenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, a ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinoxalinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group; or

Formula 4-18

a phenyl group, a pentalenyl group, an indeyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluorantenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a pycenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, a ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinoxalinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group; or

Formula 4-19

a phenyl group, a pentalenyl group, an indeyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluorantenyl group, a triphenylenyl

Formula 4-21

Formula 4-22

Formula 4-23

13

group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a pycenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, a ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, or a dibenzocarbazolyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclohexyl group, a cyclopentyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indeyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a pycenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, a ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and $Si(Q_{31})(Q_{32})(Q_{33})$ (where Q_{31} to Q_{33} are each independently a hydrogen, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, or a naphthyl group).

For example, R_1 and R_4 in Formula 1 may be each independently;


a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a

14


triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, or a dibenzocarbazolyl group; or

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, or a dibenzocarbazolyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and $-Si(Q_{31})(Q_{32})(Q_{33})$ (where Q_{31} to Q_{33} may be each independently a hydrogen, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, or a naphthyl group).


For example, R_1 to R_4 in Formula 1 may be each independently represented by one of Formulae 5-1 to 5-14.




Formula 5-1




Formula 5-2



Formula 5-3












Formula 5-4



Formula 5-5

15

-continued



In Formulae 5-1 to 5-14,

Y_{31} may be O, S, C(Z_{33})(Z_{34}), or N(Z_{35});

Z_{31} to Z_{35} may be each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carba-

16

Formula 5-6

zoyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, or dibenzocarbazoyl group, or —Si(Q_{31})(Q_{32})(Q_{33}) (where Q_{31} to Q_{33} may be each independently, a hydrogen, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, or a naphthyl group);

Formula 5-7

e1 may be an integer of 1 to 5;

e2 may be an integer of 1 to 7;

e3 may be an integer of 1 to 3;

Formula 5-8

e4 may be an integer of 1 to 4;

e5 may be 1 or 2; and

* is a binding site to N in Formula 1.


Formula 5-9

In an implementation, R_1 to R_4 in Formula 1 may be each independently represented by one of Formulae 6-1 to 6-45.

Formula 5-9


20

Formula 5-10




25

Formula 5-11




30

Formula 5-12




35

Formula 5-13




40




Formula 5-14



45











50



17


-continued

**18**

-continued


Formula 6-10

5




Formula 6-11

10




Formula 6-12

15




Formula 6-13

20




Formula 6-14

30




Formula 6-15

40




45




Formula 6-16

50




55



Formula 6-17

60



65

Formula 6-18

Formula 6-19

Formula 6-20

Formula 6-21

Formula 6-22

Formula 6-23

Formula 6-24








Formula 6-25

Formula 6-26

Formula 6-27

19


-continued

**20**

-continued

Formula 6-28


5



10

Formula 6-29


15



20

Formula 6-30


25



30

formula 6-31


35



40

Formula 6-32


45



50

Formula 6-33


55



60

formula 6-34

65



Formula 6-35

Formula 6-36

Formula 6-37

Formula 6-38

Formula 6-39




Formula 6-40

Formula 6-41

Formula 6-42

21

-continued



In Formula 1 above, $R_1=R_2=R_3=R_4$; $R_1=R_3$, R_1-R_2 , and $R_2=R_4$; $R_2=R_3$, $R_1 \neq R_2$, and $R_1=R_4$; $R_1=R_2$, $R_1 \neq R_3$, and $R_3=R_4$; $R_2=R_3=R_4$, and $R_1 \neq R_2$; $R_3=R_4$, and $R_1 \neq R_2 \neq R_3$; or $R_1 \neq R_2 \neq R_3 \neq R_4$. For example, in some implementations, some of R_1 to R_4 may be the same, some of R_1 to R_4 may be different, all of R_1 to R_4 may be same, or all of R_1 to R_4 may be different.

In Formula 1, R_{11} to R_{13} may be each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuran-yl group, a dibenzothiophenyl group, a benzocarbazolyl group, or a dibenzocarbazolyl group, or —S(Q_1)(Q_2)(Q_3), (where Q_1 to Q_3 may be each independently selected from a hydrogen, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, or naphthyl group).

In an implementation, R_{11} to R_{13} in Formula 1 may be each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, or a C_1 - C_{20} alkoxy group, but are not limited thereto.

In an implementation, R_{11} to R_{13} in Formula 1 may be hydrogen.

22

In an implementation, the condensed cyclic compound of Formula 1 may be represented by one of Formulae 1A to 1D.

Formula 6-43

5

<Formula 1A>

Formula 6-44

10

<Formula 1B>

Formula 6-45

20

<Formula 1C>

25

30

<Formula 1D>

35

40





45

50

55

60

65



Substituents in Formulae 1A to 1D may be the same as described above with respect to Formula 1.

For example, in Formulae 1A to 1D,

L_1 and L_2 may be each independently represented by one of Formulae 3-1 to 3-32 above (e.g., one of Formulae 4-1 to 4-23 above);

$a1$ and $a2$ may be each independently 0, 1, or 2;

R_1 to R_4 may be each independently represented by one of Formulae 5-1 to 5-14 above (e.g., one of Formulae 6-1 to 6-45 above);

R_{11} to R_{13} may be each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group,

23

a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazoyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, or —Si(Q₁)(Q₂)(Q₃) (where Q₁ to Q₃


24

may be each independently selected from a hydrogen, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, or a naphthyl group); and


b1 to b3 may be each independently 1 or 2.

In an implementation, the condensed cyclic compound represented by Formula 1 may be represented by Formula 1A above.


In an implementation, the condensed cyclic compound represented by Formula 1 may include one of Compounds 1 to 96, below.




1




2




3




4




5



6



7



8



25

26


-continued

9


10




11



12





13



14

15





27

28

-continued



16

17





18

19





20

21





22

23



24

25





29

30

-continued



26

27





28

29





30

31





32

33



34

35





31

32

-continued



36

37





38

39





40

41



42

43





33

34

-continued



44

45





46

47





48

49




50

51




35



-continued


52

36




53


54




55




56




57




58




59



60



61





37

38

-continued



62

63




64


65



66



67




39

40



-continued

68





69

70




71

72



73




41


42

-continued


74



75



76





43

44


-continued

77


78




79



80



81





45

46


-continued

82

83




84




85

86





87

88



89

90





47

48


-continued

91


92




93



94




95




-continued

96



The condensed cyclic compound of Formula 1 may include a core represented by



with oxygens in symmetrical positions. Unshared electron pairs of the oxygens may provide extra electrons, so that the condensed cyclic compound has a molecular structure that is rich in π -electrons. This may help increase a transition probability, so that the condensed cyclic compound of Formula 1 may help improve emission efficiency.

An organic light-emitting device including the condensed cyclic compound represented by Formula 1, above, may have a low driving voltage, a high luminance, a high efficiency, and/or a long lifetime.

The condensed cyclic compound of Formula 1 may be synthesized using a suitable organic synthesis method. Methods of synthesizing the condensed cyclic compounds of Formula 1 may be understood based on the examples that will be described below.

The condensed cyclic compound of Formula 1 may be included between a pair of electrodes of an organic light-emitting device. In an implementation, the condensed cyclic compound of Formula 1 may be in an electron transport region, e.g., in an electron transport layer.

According to another embodiment, an organic light-emitting device may include a first electrode, a second electrode opposite to the first electrode, and an organic layer between the first electrode and the second electrode and including an emission layer. The organic layer may include at least one condensed cyclic compound represented by Formula 1, above.

As used herein, “(for example, the organic layer) including at least one condensed cyclic compound means that “(the organic layer) including one of the condensed cyclic compounds of Formula 1 above, or at least two different condensed cyclic compounds of Formula 1 above”.

In an implementation, the organic layer may include only Compound 1 above as the condensed cyclic compound. In this regard, Compound 1 may be present in the electron transport layer of the organic light-emitting device. In some embodiments, the organic layer may include Compounds 1 and 2 as the condensed cyclic compounds. In this regard, Compounds 1 and 2 may be present both in the same layer (for example, in the electron transport layer) or may be present in different layers (for example, in the emission layer and the electron transport layer, respectively).

The organic layer may include i) a hole transport region between the first electrode and the emission layer and including at least one of a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer; and ii) an electron transport region between the emission layer and the second electrode and including at least one of a hole blocking layer, an electron transport layer, and an electron injection layer. The electron transport region may include the condensed cyclic compound represented by Formula 1 above. For example, the electron transport region may include the electron transport layer, the electron transport layer may include the condensed cyclic compound represented by Formula 1 above.

As used herein, the term “organic layer” refers to a single layer and/or a plurality of layers disposed between the first and second electrodes of the organic light-emitting device. A material in the “organic layer” is not limited to an organic material.

Hereinafter, a structure of an organic light-emitting device according to an embodiment and a method of manufacturing the same will now be described with reference to the FIGURE.

The FIGURE illustrates a schematic sectional view of an organic light-emitting device **10** according to an embodiment. Referring to the FIGURE, the organic light-emitting device **10** may include a first electrode **110**, an organic layer **150**, and a second electrode **190**.

A substrate (not shown) may be disposed under the first electrode **110** or on the second electrode **190** in the FIGURE. The substrate may be a glass or transparent plastic substrate with good mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

51

For example, the first electrode **110** may be formed by depositing or sputtering a first electrode-forming material on the substrate **11**. When the first electrode **110** is an anode, a material having a high work function may be used as the first electrode-forming material to facilitate hole injection. The first electrode **110** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. Transparent and conductive materials such as ITO, IZO, SnO₂, and ZnO may be used to form the first electrode. The first electrode **110** as a semi-transmissive electrode or a reflective electrode may be formed of at least one material selected from magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag).

The first electrode **110** may have a single-layer structure or a multi-layer structure including a plurality of layers. For example, the first electrode **110** may have a three-layered structure of ITO/Ag/ITO.

The organic layer **150** may be disposed on the first electrode **110**. The organic layer **150** may include an emission layer (EML).

The organic layer **150** may include a hole transport region between the first electrode and the EML, and an electron transport region between the EML and the second electrode.

For example, the hole transport region may include at least one of a hole injection layer (HIL), a hole transport layer (HTL), a buffer layer, and an electron blocking layer (EBL). For example, the electron transport layer may include at least one of a hole blocking layer (HBL), an electron transport layer (ETL), and an electron injection layer (EIL).

The hole transport region may have a single-layered structure including a single material, a single-layered structure including a plurality of materials, or a multi-layered structure including a plurality of layers including different materials.

In an implementation, the electron transport region may have a single-layered structure including a plurality of materials, or a multi-layered structure of HIL/HTL, HIL/HTL/buffer layer, HIL/buffer layer, HTL/buffer layer, or HIL/HTL/EBL, wherein these layers forming a multi-layered structure are sequentially disposed on the first electrode **110** in the order stated above.

When the hole transport region includes a HIL, the HIL may be formed on the first electrode **110** by using any of a variety of methods, for example, by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like.

When the HIL is formed using vacuum deposition, the deposition conditions may vary depending on the material that is used to form the HIL and the structure of the HIL. For example, the deposition conditions may be selected from the following conditions: a deposition temperature of about 100° C. to about 500° C., a degree of vacuum of about 10⁻⁸ to about 10⁻³ torr, and a deposition rate of about 0.01 to 100 Å/sec.


When the HIL is formed using spin coating, the coating conditions may vary depending on the material that is used to form the HIL and the structure of the HIL. For example, the coating conditions may be selected from the following conditions: a coating rate of about 2,000 rpm to about 5,000 rpm and a heat treatment temperature of about 800° C. to about 200° C.

52


When the hole transport region includes a HTL, the HTL may be formed on the first electrode **110** or the HIL by using any of a variety of methods, for example, by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like. When the HTL is formed using vacuum deposition or spin coating, the conditions for deposition and coating may be similar to the above-described deposition and coating conditions for forming the HIL, and accordingly will not be described in detail.

In an implementation, the hole transport region may include at least one of m-MTDATA, TDATA, 2-TNATA, NPB, β-NPB, TPD, Spiro-TPD, Spiro-NPB, α-NPB, TAPC, HMTPD, 4,4',4"-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzene sulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201 below, and a compound represented by Formula 202 below.

m-MTDATA




TDATA



53

-continued

2-TNATA



5


10

15

20


25

NPB



30

35


 β -NPB

40

45

50

TPD



55


60

65

54

-continued

Spiro-TPD



5


10

Spiro-NPB

15

20

25

 α -NPB

30

35

TAPC

40

45


50

HMTPD


55

60

65




<Formula 201>



55

-continued



In Formulae 201 and 202

L_{201} to L_{205} may be each independently defined as described above in conjunction with L_1 in Formula 1;

$xa1$ to $xa4$ are each independently selected from 0, 1, 2, and 3;

$xa5$ may be selected from 1, 2, 3, 4, and 5;

R_{201} to R_{205} may be each independently defined as described above in conjunction with R_1 in Formula 1.

For example, in Formulae 201 and 202,

L_{201} to L_{205} may be each independently:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorene group, a dibenzofluorene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylene group, a carbazolyne group, or a triazinylene; or

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, or a triazinylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group,

$xa1$ to $xa4$ may be each independently 0, 1, or 2;

$xa5$ may be 1, 2, or 3;


R_{201} to R_{205} may be each independently:

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, or a triazinyl group; or


56

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an azulenyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group.


In an implementation, the compound of Formula 201 may be represented by Formula 201A.



For example, the compound of Formula 201 may be represented by Formula 201A-1.



In an implementation, the compound of Formula 201 may be represented by Formula 201A.



In Formulae 201A, 201A-1, and 202A,

L_{201} to L_{203} , $xa1$ to $xa3$, $xa5$, and R_{202} to R_{204} may be defined as described in conjunction with Formula 201;

R_{211} may be defined as described in conjunction with R_{203} in Formula 201;

R_{213} to R_{216} may be each independently a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group, or a salt thereof, a sulfonic acid group, or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cyclo alkyl group, a C_2 - C_{10} heterocyclo alkyl group, a C_3 - C_{10} cyclo alkenyl group, a C_2 - C_{10} heterocyclo alkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic hetero-condensed polycyclic group.

In an implementation, in Formulae 201A, 201A-1, and 202A,

L_{201} to L_{203} may be each independently:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a pyrenylene group, a chrysénylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a carbazolylenylene group, or a triazinylene group; or

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a pyrenylene group, a chrysénylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a carbazolylenylene group, or a triazinylene group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl

group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, a quinoliny group, an isoquinoliny group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group,

$xa1$ to $xa3$ may be each independently 0 or 1,

R_{203} , R_{211} , and R_{212} may be each independently:

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinoliny group, an isoquinoliny group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, or a triazinyl group; or

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinoliny group, an isoquinoliny group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, a triazinyl group, each substituted with at least one of a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinoliny group, an isoquinoliny group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group.

R_{213} and R_{214} may be each independently:

a C_1 - C_{20} alkyl group or a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group or a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinoliny group, an isoquinoliny group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinoliny group, an isoquinoliny group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, or a triazinyl group; or

59

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group.

R₂₁₅ and R₂₁₆ may be each independently:

a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, or a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group or a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, or a triazinyl group; and

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or

60


a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, and

xa5 may be 1 or 2.


In an implementation, in Formulae 201A and 201A-1, R₂₁₃ and R₂₁₄ may be linked to each other to form a saturated or unsaturated ring.

The compound represented by Formula 201, and the compound represented by Formula 202 may include compounds HT1 to HT20 illustrated below.

HT1




HT2



61


-continued

HT3

**62**

-continued

HT5




25

30

35

40

HT4



65

HT6

45

50


55

60

63

-continued

HT7



5

10

15

20

25

30


35

40

64

-continued

HT9



HT8


45

50


55

60

65




HT10



65

-continued

HT11



5

10

15


20

HT12

25

30

35



40

45


50

HT13

55


60

65


**66**

-continued


HT14




HT15



HT16




HT17



67

-continued




A thickness of the hole transport region may be from about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1000 Å. When the hole transport region includes a HIL and a HTL, a thickness of the HIL may be from about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å, and a thickness of the HTL may be from about 50 Å to about 2,000 Å, e.g., about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the HIL, and the HTL are within these ranges, satisfactory hole transport characteristics may be obtained without a substantial increase in driving voltage.

The hole transport region may further include a charge-generating material to help improve conductivity, in addition to the materials as described above. The charge-generating material may be homogeneously or inhomogeneously dispersed in the hole transport region.

The charge-generating material may be, e.g., a p-dopant. The p-dopant may be one of quinone derivatives, metal

68

oxides, and compounds with a cyano group. Non-limiting examples of the p dopant may include quinone derivatives such as tetracyanoquinonodimethane (TCNQ), 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonodimethane (F4-TCNQ), and the like; metal oxides such as tungsten oxide, molybdenum oxide, and the like; and Compound HT-D1 below.



The hole transport region may further include at least one of a buffer layer and an EBL, in addition to the HIL and HTL described above. The buffer layer may help compensate for an optical resonance distance of light according to a wavelength of the light emitted from the EML, and thus may improve light-emission efficiency. A material in the buffer layer may be a suitable material used in the hole transport region. The EBL may block migration of electrons from the electron transport region into the EML.


The EML may be formed on the first electrode 110 or the hole transport region by using any of a variety of methods, e.g., by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like. When the EML is formed using vacuum deposition or spin coating, the deposition and coating conditions for forming the EML may be similar to the above-described deposition and coating conditions for forming the HIL, and accordingly will not be described in detail.

When the organic light-emitting device 10 is a full color organic light-emitting device, the EML may be patterned into a red emission layer, a green emission layer, and a blue emission layer to correspond to individual subpixels, respectively. In an implementation, the EML may have a structure in which a red emission layer, a green emission layer and a blue emission layer are stacked upon one another, or a structure including a mixture of a red light-emitting material, a green light-emitting material, and a blue light-emitting material, and thus may emit white light.

The EML may include a host and a dopant.


For example, the host may include at least one of TPBi, TBADN, ADN (also referred to as "DNA"), CBP, CDBP, and TCP.

69




70

-continued



In an implementation, the host may include a compound represented by Formula 301.



<Formula 301>

In Formula 301,

AR₃₀₁ may be:





a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, or an indenoanthracene; and

a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, or an indenoanthracene, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cyclo alkyl group, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a non-valent non-aromatic condensed polycyclic group, a non-valent non-aromatic hetero-condensed polycyclic group, and —Si(Q₃₀₁)(Q₃₀₂)(Q₃₀₃) (where Q₃₀₁ to Q₃₀₃ may be each independently, a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₆-C₆₀ aryl group, or a C₂-C₆₀ heteroaryl group),

L₃₀₁ may be defined as described above in conjunction with L₂₀₁ in Formula 201,

R₃₀₁ may be:

a C₁-C₂₀ alkyl group, or a C₁-C₂₀ alkoxy group;
 a C₁-C₂₀ alkyl group, or a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl



71

group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, or a triazinyl group; or

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group,

xb1 may be selected from 0, 1, 2, and 3;

xb2 may be selected from 1, 2, 3, and 4.

For example, in Formula 301,

L₃₀₁ may be:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a pyrenylene group, or a chrysenylene group; or

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a pyrenylene group, or a chrysenylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, and

R₃₀₁ may be:

a C₁-C₂₀ alkyl group, or a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group or a C₁-C₂₀ alkoxy group, each substituted at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl

72


group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, or a chrysenyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, or a chrysenyl group; each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group.

For example, the host may include a compound represented by formula 301A.


<Formula 301A>




Substituents in Formula 301A may be defined as described above in conjunction with other formulae herein.

The compound of Formula 301 may include at least one of Compounds H1 to 42.

H1









H2



73

-continued

**74**

-continued

H3

5

10

H4

15

20

25

H5

30

35

H6

40

45

H7

50

55

H8

60

65

H9







H10

H11

H12


H13

H14




75

-continued





**76**

-continued



77

-continued



78

-continued

H25

5

10

15

H26

20

25

30

H27 35

40

H28 45

50

H29 55

60

65






H30

H31

H32




H33

H34



79

-continued



80

-continued

H35

H38

5

10

15

H36

20

25

30

35

H37



45

50

55


60

65




H39

H40



81

-continued



H41

5


10

15

20

82

-continued



H44


H42 25

30


35

40

45



H45




In an implementation, the host may include at least one of Compounds H43 to H49.

H43 50


55

60

65






H46



83

-continued



The dopant may include a condensed cyclic compound of Formula 1.

In an implementation, an amount of the dopant in the EML may be from about 0.01 parts to about 15 parts by weight, based on 100 parts by weight of the host.

The thickness of the EML may be about 100 Å to about 1,000 Å, e.g., about 200 Å to about 600 Å. When the thickness of the EML is within these ranges, the EML may have good light emitting ability without a substantial increase in driving voltage.

Next, the electron transport region may be disposed on the EML.

The electron transport region may include at least one of a HBL, an ETL, and an EIL.

84


In an implementation, the electron transport region may have a structure including an ETL, an ETL/EIL, or a HBL/ETL/EIL, wherein the layers forming a structure of the electron transport region may be sequentially stacked on the EML in the order stated above.

In an implementation, the organic layer 150 of the organic light-emitting device 10 may include the electron transport region between the EML and the second electrode 190.


The electron transport region may include a HBL. When the EML includes a phosphorescent dopant, the HBL may prevent diffusion of triplet excitons or holes into the ETL from the EML.

When the electron transport region includes a HBL, the HBL may be formed on the EML by using any of a variety of methods, for example, by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like. When the HBL is formed using vacuum deposition or spin coating, the deposition and coating conditions for forming the HBL may be similar to the above-described deposition and coating conditions for forming the HIL, and accordingly will not be described in detail.

For example, the HBL may include at least one of BCP below and Bphen, below.



BCP




Bphen

A thickness of the HBL may be from about 20 Å to about 1,000 Å, e.g., about 30 Å to about 300 Å. When the thickness of the HBL is within these ranges, the HBL may have improved hole blocking ability without a substantial increase in driving voltage.


The electron transport region may include an ETL. The ETL may be formed on the EML or the HBL by using any of a variety of methods, for example, by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like. When the ETL is formed using vacuum deposition or spin coating, the deposition and coating conditions for forming the ETL may be similar to the above-described deposition and coating conditions for forming the HIL, and accordingly will not be described in detail.

85


The ETL may further include at least one of Alq₃, Balq, TAZ, and NTAZ below, in addition to BCP and Bphen described above.




Alq3



BAIq



TAZ.



NTAZ.

In an implementation, the ETL may further include, at least one of compounds represented by Formula 601 below, in addition to the condensed cyclic compound represented by Formula 1.

$$\text{AR}_{601} = [(\text{L}_{601})_{xe1} - \text{E}_{601}]_{xe2}$$

<Formula 601>

86

In Formula 601,

AR₆₀₁ may be defined as described in conjunction with AR₃₀₁ in Formula 301,

L_{601} may be defined as described above in conjunction
5 with L_{201} in Formula 301,

E₆₀₁ may be selected from:

a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthroli-
n-1-yl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, or a dibenzocarbazolyl group; or

a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, a oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthroli-
nyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzo-
thiazolyl group, a benzooxazolyl group, an isobenzo-
oxazolyl group, a triazolyl group, a tetrazolyl group, an oxadi-
azolyl group, a triazinyl group, a dibenzofuranyl group, a
dibenzothiophenyl group, a benzocarbazolyl group, or a
dibenzocarbazolyl group, each substituted with at least one
selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl
group, a cyano group, a nitro group, an amino group, an
amidino group, a hydrazine group, a hydrazone group, a
carboxylic group or a salt thereof, a sulfonic acid group or
a salt thereof, a phosphoric acid group or a salt thereof, a
C₁–C₂₀ alkyl group, a C₁–C₂₀ alkoxy group, a phenyl group,
a pentalenyl group, an indeyl group, a naphthyl group, an
azulenyl group, a heptalenyl group, an indacenyl group, an
acenaphthyl group, a fluorenyl group, a spiro-fluorenyl
group, a benzofluorenyl group, a dibenzofluorenyl group, a
phenalenyl group, a phenanthrenyl group, an anthracenyl
group, a fluorantenyl group, a triphenylenyl group, a pyrenyl
group, a chrysenyl group, a naphthacenyl group, a pycenyl
group, a perylenyl group, a pentaphenyl group, a hexacenyl
group, a pentacenyl group, a rubicenyl group, a coronenyl
group, a ovalenyl group, a pyrrolyl group, a thiophenyl
group, a furanyl group, an imidazolyl group, a pyrazolyl
group, a thiazolyl group, an isothiazolyl group, a oxazolyl
group, an isooxazolyl group, a pyridinyl group, a pyrazinyl
group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl
group, a quinolinyl group, an isoquinolinyl group, a benzo-
quinolinyl group, a phthalazinyl group, a naphthyridinyl
group, a quinoxalinyl group, a quinazolinyl group, a cinnol-


87

linyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothienophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group,

xe1 may be selected from 0, 1, 2, and 3, and

xe2 may be selected from 1, 2, 3, and 4.

In an implementation, the electron transport layer may include at least one of compounds represented by Formula 602:



In Formula 602,

X_{611} may be N or C-(L_{611})_{xe611}- R_{611} , X_{612} may be N or C-(L_{612})_{xe612}- R_{612} , X_{613} may be N or C-(L_{613})_{xe613}- R_{613} , at least one of X_{611} to X_{613} may be N,

L_{611} to L_{616} may be each independently defined as described above in conjunction L_{201} in other formulae herein,

R_{611} to R_{616} may be each independently:

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, or a triazinyl group; or


a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an azulenyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, or a triazinyl group, and

xe611 to xe616 may be each independently selected from, 0, 1, 2, and 3.


The compound represented by Formula 601 and the compound represented by Formula 602 may include at least one of Compounds ET1 to ET15 illustrated below.

88


ET1



ET2




ET3








89

-continued




**90**

-continued



91

-continued




92


-continued

ET10


ET13



ET14




ET15



55 A thickness of the ETL may be from about 100 Å to about 1,000 Å, e.g., about 150 Å to about 500 Å. When the thickness of the ETL is within these ranges, the ETL may have satisfactory electron transporting ability without a substantial increase in driving voltage.

60 In an implementation, the ETL may further include a metal-containing material, in addition to the above-described materials.

The metal-containing material may include a lithium (Li) 65 complex. Non-limiting examples of the Li complex may include compound ET-D1 below (lithium quinolate (LiQ)), or compound ET-D2 below.



The electron transport region may include an EIL that may facilitate injection of electrons from the second electrode **190**.

The EIL may be formed on the ETL by using any of a variety of methods, e.g., by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, inkjet printing, laser printing, laser induced thermal imaging (LITI), or the like. When the EIL is formed using vacuum deposition or spin coating, the deposition and coating conditions for forming the EIL may be similar to the above-described deposition and coating conditions for forming the HIL, and accordingly will not be described in detail.

The EIL may include at least one selected from LiF, NaCl, CsF, Li₂O, BaO, and LiQ.

A thickness of the EIL may be from about 1 Å to about 100 Å, e.g., about 3 Å to about 90 Å. When the thickness of the EIL is within these ranges, the EIL may have satisfactory electron injection ability without a substantial increase in driving voltage.

The second electrode **190** may be disposed on the organic layer **150**, as described above. The second electrode **190** may be a cathode as an electron injecting electrode. A material for forming the second electrode **190** may be a metal, an alloy, an electrically conductive compound, which have a low-work function, or a mixture thereof. Non-limiting examples of materials for forming the second electrode **190** may include lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag). In some embodiments, a material for forming the second electrode **190** may be ITO or IZO. The second electrode **190** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode.

As used herein, a C₁-C₆₀ alkyl group refers to a linear or branched aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms. Non-limiting examples of the C₁-C₆₀ alkyl group are a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. A C₁-C₆₀ alkylene group refers to a divalent group having the same structure as the C₁-C₆₀ alkyl.

As used herein, a C₁-C₆₀ alkoxy group refers to a monovalent group represented by —OA₁₀₁ (where A₁₀₁ is a C₁-C₆₀ alkyl group as described above. Non-limiting examples of the C₁-C₆₀ alkoxy group as described above.

Non-limiting examples of the C₁-C₆₀ alkoxy group are a methoxy group, an ethoxy group, and an isopropoxy group.

As used herein, a C₂-C₆₀ alkenyl group refers to a structure including at least one carbon double bond in the middle or terminal of the C₂-C₆₀ alkyl group. Non-limiting examples of the C₂-C₆₀ alkenyl group are an ethenyl group, a propenyl group, and a butenyl group. A C₂-C₆₀ alkylene group refers to a divalent group having the same structure as the C₂-C₆₀ alkenyl group.

As used herein, a C₂-C₆₀ alkynyl group refers to a structure including at least one carbon triple bond in the middle or terminal of the C₂-C₆₀ alkyl group. Non-limiting examples of the C₂-C₆₀ alkynyl group are an ethynyl group and a propynyl group. A C₂-C₆₀ alkynylene group used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

As used herein, a C₃-C₁₀ cycloalkyl group refers to a monovalent, monocyclic hydrocarbon group having 3 to 10 carbon atoms. Non-limiting examples of the C₃-C₁₀ cycloalkyl group are a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. A C₃-C₁₀ cycloalkylene group refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

As used herein, a C₂-C₁₀ heterocycloalkyl group refers to a monovalent monocyclic group having 2 to 10 carbon atoms in which at least one hetero atom selected from N, O, P, and S is included as a ring-forming atom. Non-limiting examples of the C₂-C₁₀ heterocycloalkyl group are a tetrahydrofuranyl group and a tetrahydrothiophenyl group. A C₂-C₁₀ heterocycloalkylene group refers to a divalent group having the same structure as the C₂-C₁₀ heterocycloalkyl group.

As used herein, a C₃-C₁₀ cycloalkenyl group refers to a monovalent monocyclic group having 3 to 10 carbon atoms that includes at least one double bond in the ring but does not have aromaticity. Non-limiting examples of the C₃-C₁₀ cycloalkenyl group are a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. A C₃-C₁₀ cycloalkenylene group refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

As used herein, a C₂-C₁₀ heterocycloalkenyl group used herein refers to a monovalent monocyclic group having 2 to 10 carbon atoms that includes at least one double bond in the ring and in which at least one hetero atom selected from N, O, P, and S is included as a ring-forming atom. Non-limiting examples of the C₂-C₁₀ heterocycloalkenyl group are a 2,3-dihydrofuranyl group and a 2,3-dihydrothiophenyl group. A C₂-C₁₀ heterocycloalkenylene group used herein refers to a divalent group having the same structure as the C₂-C₁₀ heterocycloalkenyl group.

As used herein, a C₆-C₆₀ aryl group refers to a monovalent, aromatic carbocyclic aromatic group having 6 to 60 carbon atoms, and a C₆-C₆₀ arylene group refers to a divalent, aromatic carbocyclic group having 6 to 60 carbon atoms. Non-limiting examples of the C₆-C₆₀ aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group include at least two rings, the rings may be fused to each other.

As used herein, a C₂-C₆₀ heteroaryl group refers to a monovalent, aromatic carbocyclic aromatic group having 2 to 60 carbon atoms in which at least one hetero atom selected from N, O, P, and S is included as a ring-forming atom, and 2 to 60 carbon atoms. A C₂-C₆₀ heteroarylene group refers to a divalent, aromatic carbocyclic group hav-

ing 2 to 60 carbon atoms in which at least one hetero atom selected from N, O, P, and S is included as a ring-forming atom. Non-limiting examples of the C₂-C₆₀ heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₂-C₆₀ heteroaryl and the C₂-C₆₀ heteroarylene include at least two rings, the rings may be fused to each other.

As used herein, a C₆-C₆₀ aryloxy group indicates —OA₁₀₂ (where A₁₀₂ is a C₆-C₆₀ aryl group as described above), and a C₆-C₆₀ arylthio group indicates —SA₁₀₃ (wherein A₁₀₃ is a C₆-C₆₀ aryl group as described above).

As used herein, a monovalent non-aromatic condensed polycyclic group refers to a monovalent group having at least two rings condensed to each other, in which only carbon atoms (for example, 8 to 60 carbon atoms) are exclusively included as ring-forming atoms and the entire molecule has non-aromaticity. A non-limiting example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. A divalent non-aromatic condensed polycyclic group refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

As used herein, a monovalent non-aromatic condensed heteropolycyclic group refers to a monovalent group having at least two rings condensed to each other, in which carbon atoms (for example, 2 to 60 carbon atoms) and a hetero atom selected from N, O, P, and S are as ring-forming atoms and the entire molecule has non-aromaticity. A non-limiting example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazoyl group. A divalent non-aromatic condensed heteropolycyclic group refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

As used herein, at least one of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₂-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₂-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₂-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed cyclic group, the substituted divalent non-aromatic hetero-condensed cyclic group, the substituted C₃-C₁₀ cyclo alkyl group, the substituted C₂-C₁₀ heterocyclo alkyl group, the substituted C₃-C₁₀ cyclo alkenyl group, the substituted C₂-C₁₀ heterocyclo alkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₂-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic hetero-condensed polycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, and the substituted C₁-C₆₀ alkoxy group may be selected from

a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt

thereof, a cyclopentenyl group, a cyclohexyl group, a cycloheptyl, a cyclopentenyl, a cycloheptenyl, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coroneryl group, an obarenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazoyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, an imidazopyridinyl group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

a cyclopentenyl group, a cyclohexyl group, a cycloheptyl, a cyclopentenyl, a cycloheptenyl, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coroneryl group, an obarenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazoyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, and an imidazopyridinyl group;

a cyclopentenyl group, a cyclohexyl group, a cycloheptyl, a cyclopentenyl, a cycloheptenyl, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group,

a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coroneryl group, an obarenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a phenyl group, a naphthyl group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇); and —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇).

wherein Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ may be each independently selected from a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a cyclopentenyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl, a cycloheptenyl, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a

pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coroneryl group, an obarenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group.


The acronym “Ph” used herein refers to phenyl, the acronym “Me” used herein refers to methyl, the acronym “Et” used herein refers to ethyl, and the acronym “ter-Bu” or “Bu” used herein refers to tert-butyl.

One or more embodiments, which may include condensed cyclic compounds, and organic light-emitting devices including the same, will now be described in detail with reference to the following examples. In the following synthesis example, the expression that “‘B’ instead of ‘A’ was used” means that the amounts of ‘B’ and ‘A’ were the same in equivalent amounts.

The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples.

EXAMPLES


Synthesis Example 1: Synthesis of Compound 5



99

100

-continued



5

Synthesis of Intermediate 5-1

5.2 g of 2-bromo-5-chlorodimethylfluorene (23.6 mmol) was dissolved in 100 mL of tetrahydrofuran (THF), and 10 mL of normal butyl lithium (25.0 mmol, 2.5 M in hexane) was slowly dropwise added thereto at added thereto -78°C . After the reaction solution was stirred at the same temperature for about 1 hour, 9.3 mL (50.0 mmol) of 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane was slowly dropwise added to the reaction solution, which was then stirred

at about -78°C . for about 1 hour, and further at room or ambient temperature for about 24 hours. After the reaction was completed, 50 mL of a 10% HCl aqueous solution and 50 mL of H_2O were added thereto, followed by extracting three times with 80 mL of diethyl ether. An organic layer was collected, and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 5.83 g of Intermediate 5-1 (Yield: 92%). This compound was identi-

101

fied using liquid chromatography-mass spectroscopy (LC-MS).

$C_{13}H_{18}BClO_3$; M^+ 268.1

Synthesis of Intermediate 5-2

5.90 g (22.0 mmol) of Intermediate 1-6, 4.18 g (11.0 mmol) of 1,4-(4-bromophenyl)-1-phenyl-1-benzoimidazole, 2.54 g (2.2 mmol) of tetrakis(triphenylphosphine)palladium ($Pd(PPh_3)_4$), and 9.00 g (66 mmol) of K_2CO_3 were dissolved in 40 mL of a mixed tetrahydrofuran (THF) and H_2O (2:1 by volume) solution to obtain a solution, which was then stirred at about 70° C. for about 5 hours. The reaction solution was cooled down to ambient temperature, and 60 mL of water was added thereto, followed by extracting three times with 60 mL of ethyl ether. The organic phase was collected, and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 3.42 g of Intermediate 5-2 (Yield: 76%). This compound was identified using liquid chromatography-mass spectroscopy (LC-MS).

$C_{24}H_{18}Cl_2O_2$; M^+ 408.1

Synthesis of Intermediate 5-3

8.18 g (20.0 mmol) of Intermediate 5-2, 9.66 g (40.0 mmol) of Intermediate 5-A, 0.37 g (0.4 mmol) of $Pd_2(dba)_3$, 0.08 g (0.4 mmol) of $PtBu_3$, and 5.76 g (60.0 mmol) of $KOtBu$ were dissolved in 90 mL of toluene to obtain a mixture, which was then stirred at about 85° C. for 4 hours. The reaction solution was cooled down to ambient temperature, followed by extracting three times with 50 mL of water and 50 mL of diethyl ether. The organic phase was collected, and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 13.6 g of Intermediate 5-3 (Yield: 83%). This compound was identified using LC-MS.

$C_{54}H_{54}N_2O_2Si_2$ M^+ 818.4

102

Synthesis of Intermediate 5-4

1.64 g (2.00 mmol) of Intermediate 5-3 was dissolved in 10 mL of methyl chloroform (MC) to obtain a reaction solution, and 0.33 mL (3.5 mmol) of BBR_3 was slowly dropwise added thereto at about -78° C. After raising the temperature of the reaction solution to ambient temperature, the reaction solution was stirred at ambient temperature for about 24 hours. After the reaction was completed, 5 mL of MeOH and 10 mL of H_2O were added thereto, followed by extracting three times with 10 mL of MC. The organic phase was collected, and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 1.19 g of Intermediate 5-4 (Yield: 75%). This compound was identified using LC-MS.


$C_{52}H_{50}N_2O_2Si_2$; M^+ 790.3

Synthesis of Compound 5

1.58 g (2.00 mmol) of Intermediate 5-4 was dissolved in 10 mL of dimethylformamide (DMF), and 0.48 g (6.0 mmol) of CuO was dropwise added thereto at ambient temperature. The reaction mixture was stirred at about 140° C. for about 48 hours. After the reaction was completed, the reaction product was filtered using Celite to obtain an organic layer. 10 mL of H_2O was added to the organic layer, followed by extracting three times with 10 mL of ethyl acetate. The resulting organic layer was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 1.04 g of Compound 5 (Yield: 66%). This compound was identified using liquid chromatography-mass spectroscopy (LC-MS) and nuclear magnetic resonance (NMR).

$C_{52}H_{46}N_2O_2Si_2$; M^+ found 786.32, calc. 786.31.


Synthesis Example 2: Synthesis of Compound 23



23-3(1)

103

104



Synthesis of Intermediate 23-3

Intermediate 23-3(1) was obtained in the same manner as in the synthesis of Intermediate 5-3 in Synthesis Example 1, except that Intermediate 23-A (20.0 mmol) instead of Intermediate 5-A (40.0 mmol) was used.

Synthesis of Intermediate 23-3

Intermediate 23-3 was obtained in the same manner as in the synthesis of Intermediate 5-3 in Synthesis Example 1, except that Intermediate 23-3(1) (20.0 mmol) and Intermediate 23-B (20.0 mmol) instead of Intermediate 5-2 (20.0 mmol) and Intermediate 5-A (40.0 mmol) was used.

Synthesis of Intermediate 23-4

Intermediate 23-4 was obtained in the same manner as in the synthesis of Intermediate 5-4 in Synthesis Example 1, except that Intermediate 23-3 instead of Intermediate 5-3 was used.


Synthesis of Compound 23

2.30 g of Compound 23 (Yield 71%) was synthesized in the same manner as in the synthesis of Compound 5 in Synthesis Example 1, except that Intermediate 23-4 instead of Intermediate 5-4 was used. This compound was identified using LC-MS and NMR.

$C_{58}H_{37}FN_2O_2$; M^+ found 848.39, Calc. 848.36

105


Synthesis Example 3: Synthesis of Compound 25



3.5 g of Compound 25 (Yield 68%) was synthesized in the same manner as in Synthesis Example 2, except that Intermediate 25-A instead of Intermediate 23-A was used. This compound was identified using LC-MS and NMR.

$C_{28}H_{24}N_2O$: M^+ found 380.18, Calc. 380.18

Synthesis Example 4: Synthesis of Compound 31



25-A 5

10

15

20

31-A


35

106

2.2 g of Compound 31 (Yield 65%) was synthesized in the same manner as in Synthesis Example 2, except that Intermediate 31-A instead of Intermediate 23-A was used. This compound was identified using LC-MS and NMR.

$C_{56}H_{34}N_2O_3$: M^+ found 782.27, Calc. 782.26

Synthesis Example 5: Synthesis of Compound 50




50-A

3.2 g of Compound 50 (Yield 70%) was synthesized in the same manner as in Synthesis Example 2, except that Intermediate 50-An instead of Intermediate 23-A was used. This compound was identified using LC-MS and NMR.

$C_{25}H_{17}N_3O$: M^+ found 391.13, Calc. 391.13



Synthesis Example 6: Synthesis of Compound 63



107

108

-continued



Synthesis of Intermediate 63-3(1)

Intermediate 63-3(1) was obtained in the same manner as in the synthesis of Intermediate 5-3 in Synthesis Example 1, except that Intermediate 63-A (20.0 mmol) instead of Intermediate 5-A (40.0 mmol) was used.

Synthesis of Intermediate 63-3

Intermediate 63-3 was obtained in the same manner as in the synthesis of Intermediate 5-3 in Synthesis Example 1, except that Intermediate 63-3(1) (20.0 mmol) and Intermediate 63-B (20.0 mmol) instead of Intermediate 5-2 (20.0 mmol) and Intermediate 5-A (40.0 mmol) were used.

Synthesis of Intermediate 63-4

Intermediate 63-4 was obtained in the same manner as in the synthesis of Intermediate 5-4 in Synthesis Example 1, except that Intermediate 63-3 instead of Intermediate 5-3 was used.


Synthesis of Intermediate 63

2.6 g of Compound 63 (Yield 75%) was synthesized in the same manner as in the synthesis of Compound 5 in Synthesis Example 1, except that Intermediate 63-4 instead of Intermediate 5-4 was used. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

$C_{61}H_{42}N_2O_2$: M^+ found 834.33, Calc. 834.32

109


Synthesis Example 7: Synthesis of Compound 70



1.9 g of Compound 70 (Yield 61%) was synthesized in the same manner as in Synthesis Example 6, except that Intermediate 70-A instead of Intermediate 63-A was used. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

$C_{20}H_{18}N$; M^+ found 818.30, Calc. 818.29


Synthesis Example 8: Synthesis of Compound 73



2.3 g of Compound 73 (Yield 73%) was synthesized in the same manner as in Synthesis Example 6, except that Intermediate 73-B instead of Intermediate 63-B was used.

$C_{20}H_{18}N$; M^+ found 848.31, Calc. 848.30

Synthesis Example 9: Synthesis of Compound 91



1.8 g of Compound 91 (Yield 60%) was synthesized in the same manner as in Synthesis Example 6, except that Intermediate 91-A instead of Intermediate 63-A was used. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

$C_{20}H_{18}N$; M^+ found 769.28, Calc. 769.27

Synthesis Example 10: Synthesis of Compound 1

2.6 g of Compound 1 (Yield 65%) was synthesized in the same manner as in Synthesis Example 1, except that Intermediate 23-B of Synthesis Example 2 instead of Interme-

110

diolate 5-A was used to synthesize Intermediate 5-3. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

70-A 5 Synthesis Example 11: Synthesis of Compound 7


2.8 g of Compound 7 (Yield 66%) was synthesized in the same manner as in Synthesis Example 1, except that Intermediate 63-B of Synthesis Example 6 instead of Intermediate 5-A was used to synthesize Intermediate 5-3. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 12: Synthesis of Compound 11

2.5 g of Compound 11 (Yield 71%) was synthesized in the same manner as in Synthesis Example 1, except that Intermediate 73-B of Synthesis Example 6 instead of Intermediate 5-A was used to synthesize Intermediate 5-3. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 13: Synthesis of Compound 14

73-B 25



14-A


3.0 g of Compound 14 (Yield 70%) was synthesized in the same manner as in Synthesis Example 1, except that Intermediate 14-A instead of Intermediate 5-A was used to synthesize Intermediate 5-3. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 14: Synthesis of Compound 24

3.2 g of Compound 24 (Yield 72%) was synthesized in the same manner as in Synthesis Example 3, except that Intermediate 73-B of Synthesis Example 8 instead of Intermediate 23-A was used to synthesize Intermediate 23-3(1). This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 15: Synthesis of Compound 29

91-A 55




29-A

60

111


2.8 g of Compound 29 (Yield 65%) was synthesized in the same manner as in Synthesis Example 2, except that Intermediate 29-A instead of Intermediate 23-A was used to synthesize Intermediate 23-3(1). This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 16: Synthesis of Compound 35




2.2 g of Compound 35 (Yield 63%) was synthesized in the same manner as in Synthesis Example 2, except that Intermediate 35-A instead of Intermediate 23-A was used to synthesize Intermediate 23-3(1). This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 17: Synthesis of Compound 40



3.2 g of Compound 40 (Yield 71%) was synthesized in the same manner as in Synthesis Example 2, except that Intermediate 40-A instead of Intermediate 23-A was used to synthesize Intermediate 23-3(1). This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 18: Synthesis of Compound 53



112

2.2 g of Compound 53 (Yield 62%) was synthesized in the same manner as in Synthesis Example 2, except that Intermediate 53-A instead of Intermediate 23-A was used to synthesize Intermediate 23-3(1). This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 19: Synthesis of Compound 57

10


35-A

15

20

25

30



57-A

2.0 g of Compound 57 (Yield 64%) was synthesized in the same manner as in Synthesis Example 2, except that Intermediate 57-A instead of Intermediate 23-A was used to synthesize Intermediate 23-3(1). This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 20: Synthesis of Compound 64

40-A

35

40

2.9 g of Compound 64 (Yield 69%) was synthesized in the same manner as in Synthesis Example 6, except that Intermediate 23-A of Synthesis Example 2 instead of Intermediate 63-A was used. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 21: Synthesis of Compound 65

45

50

55




2.7 g of Compound 65 (Yield 73%) was synthesized in the same manner as in Synthesis Example 6, except that Intermediate 73-B of Synthesis Example 8 instead of Intermediate 63-A was used to synthesize Intermediate 63-3(1). This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 22: Synthesis of Compound 69

60


65

3.0 g of Compound 69 (Yield 66%) was synthesized in the same manner as in Synthesis Example 6, except that Intermediate 70-A of Synthesis Example 7 instead of Intermediate 63-A was used to synthesize Intermediate 63-3(1). This compound was identified using LC-MS and nuclear magnetic resonance (NMR).



115

116



Synthesis of Intermediate 78-3(1)

Intermediate 78-3(1) was obtained in the same manner as in the synthesis of Intermediate 5-3 in Synthesis Example 1, except that the amount of Intermediate 5-A was changed from 40.0 to 20.0 mmol.

Synthesis of Intermediate 78-3

Intermediate 78-3 was obtained in the same manner as in the synthesis of Intermediate 5-3 in Synthesis Example 1, except that Intermediate 78-3(1) (20.0 mmol) and Intermediate 14-A (20.0 mmol) instead of Intermediate 5-2 (20.0 mmol) and Intermediate 5-A (40.0 mmol) were used.

Synthesis of Intermediate 78-4

Intermediate 78-4 was obtained in the same manner as in the synthesis of Intermediate 5-4 in Synthesis Example 1, except that Intermediate 78-3 instead of Intermediate 5-3 was used.

Synthesis of Intermediate 78

3.2 g of Compound 78 (Yield 65%) was synthesized in the same manner as in the synthesis of Compound 5 in Synthesis Example 1, except that Intermediate 78-4 instead of Intermediate 5-4 was used. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).


Synthesis Example 24: Synthesis of Compound 82

3.0 g of Compound 82 (Yield 68%) was synthesized in the same manner as in Synthesis Example 23, except that Intermediate 23-A of Synthesis Example 2 instead of Intermediate 5-A was used to synthesize Intermediate 78-3(1). This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

Synthesis Example 25: Synthesis of Compound 85

2.5 g of Compound 85 (Yield 62%) was synthesized in the same manner as in Synthesis Example 1, except that Intermediate 85-A instead of Intermediate 5-A was used to synthesize Intermediate 5-3. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).




Synthesis Example 26: Synthesis of Compound 95



117

118

-continued



Synthesis of Intermediate 95-1

Intermediate 95-1 was obtained in the same manner as in the synthesis of Intermediate 5-3 in Synthesis Example 1, except that the amount of Intermediate 5-A was changed from 40.0 to 20.0 mmol.

Synthesis of Intermediate 95-2

11.9 g (22.0 mmol) of Intermediate 95-1, 10.1 g (22.0 mmol) of Intermediate 95-A, 1.27 g (1.1 mmol) of tetrakis (triphenylphosphine)palladium ($\text{Pd}(\text{PPh}_3)_4$), and 4.50 g (33 mmol) of K_2CO_3 were dissolved in 60 mL of a mixed THF and H_2O (2:1 by volume) solution to obtain a solution,

which was then stirred at about 70° C. for about 5 hours. The reaction mixture was cooled down to ambient temperature, and 60 mL of water was then added thereto, followed by extracting three times with 60 mL of ethyl ether. The organic phase was collected, and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 11.8 g of Intermediate 95-2 (Yield: 64%). This compound was identified using LC-MS.

Synthesis of Intermediate 95-3

Intermediate 95-3 was obtained in the same manner as in the same manner as in the synthesis of Intermediate 5-4 in

Synthesis Example 1, except that Intermediate 95-2 instead of Intermediate 5-3 was used.

Synthesis of Compound 95

2.8 g of Compound 95 (Yield 69%) was synthesized in the same manner as in the synthesis of Compound 5 in Synthesis Example 1, except that Intermediate 95-3 instead of Inter-

mediate 5-4 was used. This compound was identified using LC-MS and nuclear magnetic resonance (NMR).

The compounds obtained in the above-described synthesis examples were identified using ¹H NMR and mass spectroscopy/fast atom bombardment (MS/FAB). The results are shown in Table 1, below. The compounds represented by Formula 1 above may be synthesized with reference to Synthesis Examples 1 to 26.

TABLE 1

Compound	¹ H NMR (CDCl ₃ , 400 MHz)	MS/FAB	
		found	calc.
1	δ = 7.90-7.85 (m, 2H), 7.65-7.62 (m, 2H), 7.31-7.29 (m, 2H), 7.09-7.04 (m, 8H), 6.76-6.74 (m, 2H), 6.66-6.62 (m, 4H), 6.50-6.48 (m, 2H), 6.32-6.27 (m, 8H),	642.24	642.23
5	δ = 7.92-7.87 (m, 2H), 7.65-7.62 (m, 2H), 7.40-7.36 (m, 4H), 7.31-7.29 (m, 2H), 7.09-7.04 (m, 4H), 6.76-6.52 (m, 8H), 6.52-6.50 (m, 2H), 6.37-6.34 (m, 4H), 0.24 (s, 18H),	786.31	786.31
7	δ = 7.90-7.87 (m, 2H), 7.60-7.54 (m, 6H), 7.49-7.44 (m, 6H), 7.31-7.29 (m, 2H), 7.21-7.15 (m, 4H), 7.07-6.96 (m, 6H), 6.82-6.80 (m, 2H), 6.65-6.61 (m, 4H), 6.43-6.40 (m, 2H), 6.19-6.16 (m, 4H),	794.30	794.29
11	δ = 7.90-7.87 (m, 2H), 7.84-07.82 (m, 2H), 7.73-7.66 (m, 4H), 7.56-7.53 (m, 2H), 7.48-7.40 (m, 4H), 7.31-7.29 (m, 2H), 7.08-6.94 (m, 8H), 6.72-6.70 (m, 2H), 6.65-6.61 (m, 2H), 6.49-6.47 (m, 2H), 6.36-6.33 (m, 4H),	822.26	822.25
14	δ = 7.95-7.93 (m, 2H), 7.78-7.76 (m, 2H), 7.70-7.67 (m, 4H), 7.60-7.53 (m, 6H), 7.41-7.38 (m, 2H), 7.31-7.29 (m, 2H), 7.10-7.07 (m, 4H), 6.69-6.66 (m, 2H), 6.76-6.74 (m, 2H), 6.66-6.62 (m, 2H), 6.54-6.51 (m, 2H), 6.36-6.32 (m, 4H),	742.27	742.26
23	δ = 8.10-8.06 (m, 2H), 7.74-7.69 (m, 2H), 7.65-7.60 (m, 4H), 7.57-7.48 (m, 6H), 7.44-7.40 (m, 1H), 7.31-7.29 (m, 2H), 7.14-7.03 (m, 7H), 6.75-6.73 (m, 1H), 6.67-6.59 (m, 3H), 6.53-6.48 (m, 2H), 6.39-6.37 (m, 1H), 6.30-6.22 (m, 6H),	812.29	812.28
24	δ = 8.10-8.06 (m, 2H), 7.84-7.82 (m, 1H), 7.73-7.69 (m, 2H), 7.64 (d, 1H), 7.54 (d, 1H), 7.48-7.40 (m, 2H), 7.31-7.29 (m, 2H), 7.09-7.04 (m, 7H), 7.00 (d, 1H), 6.96-6.94 (m, 1H), 6.74-6.61 (m, 5H), 6.50-6.48 (m, 2H), 6.36-6.27 (m, 5H),	732.24	732.24
25	δ = 8.10-8.06 (m, 2H), 7.84-7.82 (m, 1H), 7.72-7.69 (m, 2H), 7.65-7.61 (m, 3H), 7.56-7.38 (m, 8H), 7.31-7.29 (m, 2H), 7.09-6.94 (m, 6H), 6.74-6.62 (m, 6H), 6.50-6.48 (m, 2H), 6.30-6.27 (m, 4H)	808.28	808.27
29	δ = 8.10-8.06 (m, 2H), 7.89 (dd, 1H), 7.76 (dd, 1H), 7.72-7.69 (m, 3H), 7.65-7.63 (m, 2H), 7.42-7.35 (m, 3H), 7.31-7.29 (m, 2H), 7.22 (t, 1H), 7.09-7.03 (m, 6H), 6.97 (d, 1H), 6.91 (t, 1H), 6.74-6.71 (m, 2H), 6.65-6.61 (m, 3H), 6.50-6.47 (m, 2H), 6.36-6.33 (m, 2H), 6.30-6.27 (m, 4H)	808.27	808.27
31	δ = 8.06-8.04 (m, 2H), 7.83 (d, 1H), 7.78-7.71 (m, 4H), 7.67-7.62 (m, 2H), 7.58-7.50 (m, 5H), 7.44-7.38 (m, 2H), 7.31-7.29 (m, 2H), 7.09-7.04 (m, 5H), 7.00 (dd, 1H), 6.66-6.62 (m, 2H), 6.79-6.74 (m, 2H), 6.55 (dd, 1H), 6.49 (dd, 1H), 6.31-6.27 (m, 4H)	782.27	782.26
35	δ = 8.10-8.06 (m, 2H), 7.83 (d, 1H), 7.74-7.62 (m, 3H), 7.59-7.51 (m, 3H), 7.44-7.40 (m, 1H), 7.31-7.29 (m, 2H), 7.09-7.05 (m, 5H), 6.87-6.81 (m, 2H), 6.75-6.73 (m, 1H), 6.68-6.62 (m, 4H), 6.52-6.48 (m, 2H), 6.30-6.27 (m, 4H)	768.83	768.82
40	δ = 8.58-8.57 (m, 1H), 8.43-8.41 (m, 2H), 8.20-8.16 (m, 1H), 8.00-7.95 (m, 1H), 7.78-7.76 (m, 1H), 7.71-7.63 (m, 4H), 7.58-7.54 (m, 3H), 7.43-7.30 (m, 1H), 7.35-7.29 (m, 4H), 7.14-7.04 (m, 6H), 6.77-6.74 (m, 2H), 6.66-6.60 (m, 6H), 6.30-6.27 (m, 4H), 1.36 (s, 6H)	858.33	858.32
50	δ = 8.59-8.57 (m, 1H), 8.43-8.41 (m, 2H), 7.96-7.92 (m, 1H), 7.72-7.68 (m, 2H), 7.65-7.63 (m, 2H), 7.58-7.53 (m, 2H), 7.43-7.36 (m, 4H), 7.31-7.29 (m, 2H), 7.09-7.04 (m, 4H), 6.74-6.71 (m, 3H), 6.67-6.62 (m, 3H), 6.50-6.45 (m, 3H), 6.30-6.27 (m, 4H)	767.27	767.26
53	δ = 8.60-8.57 (m, 1H), 8.43-8.41 (m, 2H), 8.19-8.15 (m, 1H), 7.84-7.80 (m, 1H), 7.74-7.54 (m, 12H), 7.50-7.40 (m, 6H), 7.35-7.29 (m, 3H), 7.09-7.04 (m, 4H), 6.97 (d, 1H), 6.92-6.90 (m, 1H), 6.76-6.74 (m, 1H), 6.65-6.63 (m, 2H), 6.49 (dd, 1H), 6.39-6.37 (m, 1H), 6.31-6.27 (m, 5H)	894.32	894.32
57	δ = 8.59-8.57 (m, 1H), 8.43-8.41 (m, 2H), 8.20-8.17 (m, 1H), 8.13-8.11 (m, 1H), 8.00-7.96 (m, 1H), 7.82 (d, 1H), 7.76 (d, 1H), 7.72-7.62 (m, 4H), 7.56-7.53 (m, 2H), 7.46-7.35 (m, 4H), 7.31-7.29 (m, 2H), 7.09-7.04 (m, 5H), 6.90 (dd, 1H), 6.76-6.74 (m, 1H), 6.66-6.60 (m, 3H), 6.50-6.47 (m, 2H), 6.30-6.27 (m, 4H)	848.26	848.25

TABLE 1-continued

Compound	¹ H NMR (CDCL ₃ , 400 MHz)	MS/FAB	
		found	calc.
63	δ = 7.85-7.82 (m, 2H), 7.77 (dd, 1H), 7.68 (dd, 1H), 7.59-7.54 (m, 4H), 7.50-7.44 (m, 3H), 7.37-7.29 (m, 3H), 7.21-6.97 (m, 9H), 6.86 (dd, 1H), 6.81 (d, 1H), 6.69-6.60 (m, 4H), (6.50-6.46 (m, 2H), 6.41 (m, 1H), 6.34-6.31 (m, 2H), 6.19-6.16 (m, 6H), 1.61 (s, 6H)	834.33	834.32
64	δ = 8.10-8.07 (m, 2H), 8.73-8.69 (m, 2H), 7.65-7.40 (m, 16H), 7.31-7.29 (m, 2H), 7.21-6.96 (m, 8H), 6.81 (d, 1H), 6.65-6.59 (m, 3H), 6.53-6.51 (m, 1H), 6.43-6.37 (m, 2H), 6.26-6.23 (m, 2H), 6.19-6.16 (m, 2H)	888.33	888.32
65	δ = 8.10-8.07 (m, 2H), 7.84-7.82 (m, 1H), 7.73-7.68 (m, 2H), 7.59-7.54 (m, 4H), 7.50-7.40 (m, 5H), 7.31-7.29 (m, 2H), 7.21-7.15 (m, 2H), 7.05-6.94 (m, 7H), 6.82-6.80 (m, 1H), 6.72-6.70 (m, 1H), 6.65-6.61 (m, 3H), 6.48 (dd, 1H), 6.41 (dd, 1H), 6.37-6.35 (m, 2H), 6.19-6.15 (m, 2H)	808.27	808.27
69	δ = 8.59-8.57 (m, 1H), 8.43-8.41 (m, 2H), 8.20-8.18 (m, 1H), 7.97-7.94 (m, 1H), 7.71-7.54 (m, 8H), 7.94-7.39 (m, 4H), 7.31-7.29 (m, 2H), 7.21-7.15 (m, 3H), 7.08-6.96 (m, 5H), 6.81 (d, 1H), 6.65-6.61 (m, 4H), 6.45-6.40 (m, 2H), 6.27-6.16 (m, 4H)	818.30	818.29
70	δ = 8.35-8.30 (m, 1H), 8.22-8.20 (m, 2H), 8.11 (d, 1H), 7.93-7.90 (m, 1H), 7.76-7.54 (m, 8H), 7.50-7.44 (m, 4H), 7.31-7.29 (m, 2H), 7.21-7.15 (m, 2H), 7.11-6.96 (m, 6H), 6.82-6.80 (m, 1H), 6.76-6.75 (m, 1H), 6.65-6.60 (m, 3H), 6.53 (dd, 1H), 6.42 (dd, 1H), 6.36-6.33 (m, 2H), 6.19-6.16 (m, 2H)	818.30	818.29
73	δ = 8.12-8.10 (m, 2H), 7.84-7.82 (m, 1H), 7.78-7.76 (m, 1H), 7.72-7.68 (m, 3H), 7.58-7.54 (m, 2H), 7.48-7.40 (m, 2H), 7.38-7.29 (m, 3H), 7.14-6.99 (m, 7H), 6.96-6.94 (m, 1H), 6.86 (dd, 1H), 6.71-6.61 (m, 4H), 6.50-6.46 (m, 3H), 6.36-6.31 (m, 4H), 1.61 (s, 6H)	848.31	848.30
78	δ = 8.10-8.08 (m, 2H), 7.78-7.77 (m, 1H), 7.70-7.54 (m, 6H), 7.41-7.36 (m, 3H), 7.31-7.29 (m, 2H), 7.10-7.04 (m, 4H), 6.89-6.86 (m, 1H), 6.77-6.75 (m, 2H), 6.71-6.62 (m, 4H), 6.54-6.50 (m, 2H), 6.37-6.33 (m, 4H), 0.24 (s, 9H)	764.30	764.29
82	δ = 8.03-8.00 (m, 2H), 7.78-7.76 (m, 1H), 7.73-7.49 (m, 16H), 7.44-7.38 (m, 2H), 7.31-7.29 (m, 2H), 7.14-7.03 (m, 5H), 6.89-6.86 (m, 1H), 6.76-6.74 (m, 1H), 6.67-6.59 (m, 2H), 6.54-6.51 (m, 2H), 6.39-6.33 (m, 3H), 6.26-6.22 (m, 2H)	862.30	862.30
85	δ = 8.43-8.40 (m, 2H), 8.19-8.16 (m, 2H), 7.66-7.63 (m, 2H), 7.45-7.41 (m, 2H), 7.31-7.21 (m, 8H), 6.95-6.88 (m, 4H), 6.79-6.71 (m, 6H), 6.64-6.62 (m, 2H)	644.23	644.22
91	δ = 8.33-8.31 (m, 2H), 8.10-8.06 (m, 1H), 7.78-7.73 (m, 2H), 7.70-7.66 (m, 1H), 7.61-7.54 (m, 6H), 7.50-7.38 (m, 5H), 7.31-7.29 (m, 2H), 7.21-7.15 (m, 3H), 7.07-6.96 (m, 4H), 6.92-6.90 (m, 1H), 6.82-6.76 (m, 2H), 6.67-6.60 (m, 3H), 6.41 (dd, 1H), 6.19-6.16 (m, 2H)	769.28	769.27
95	δ = 8.30 (dd, 1H), 8.26 (dd, 1H), 7.94-7.92 (m, 1H), 7.84-7.79 (m, 2H), 7.73-7.71 (m, 1H), 7.83 (dd, 1H), 7.57-7.52 (m, 1H), 7.48-7.39 (m, 5H), 7.30 (d, 1H), 7.18 (d, 1H), 7.09-6.96 (m, 8H), 6.76-6.74 (m, 1H), 6.66-6.61 (m, 3H), 6.51-6.47 (m, 3H), 6.30-6.22 (m, 6H)	808.27	808.27

Example 1

45

Acorning 15 Ω/cm² (1,200 Å) ITO glass substrate was cut to a size of 50 mm×50 mm×0.7 mm and then sonicated in isopropyl alcohol and pure water each for five minutes, and then cleaned by irradiation of ultraviolet rays for 30 minutes and exposure to ozone. The resulting ITO glass substrate was mounted into a vacuum deposition device.

After 2-TNATA was vacuum-deposited on the ITO anode of the ITO glass substrate to form an HIL having a thickness of 600 Å, NPB was deposited on the HIL to form a HTL having a thickness of about 300 Å, and then ADN (host) and Compound 5 (dopant) were co-deposited in a weight ratio of 98:2 on the HTL to form an EML having a thickness of about 300 Å.

Then, Alq₃ was deposited on the EML to form an ETL having a thickness of about 300 Å, and then LiF was deposited on the ETL to form an EIL having a thickness of about 10 Å. Then, Al was deposited on the EIL to form a cathode having a thickness of about 3,000 Å, thereby completing the manufacture of an organic light-emitting device.

Example 2

An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 23 instead of Compound 5 was used to form the EML.

Example 3

An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 25 instead of Compound 5 was used to form the EML.

Example 4

An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 31 instead of Compound 5 was used to form the EML.

Example 5

An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 50 instead of Compound 5 was used to form the EML.

123

Example 6

An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 63 instead of Compound 5 was used to form the EML.

Example 7

An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 170 instead of Compound 5 was used to form the EML.

Example 8


An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 73 instead of Compound 5 was used to form the EML.

Example 9

An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 91 instead of Compound 5 was used to form the EML.

Comparative Example 1

An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound A instead of Compound 5 was used to form the EML.




Comparative Example 2

An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound B instead of Compound 5 was used to form the EML.

124


<Compound B>



Comparative Example 3

An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound C instead of Compound 5 was used to form the EML.

<Compound C>



Evaluation Example 1

Driving voltages, current densities, luminances, efficiencies, and half-lifetimes of the organic light-emitting devices of Examples 1 to 9 and Comparative Examples 1 to 3 were evaluated using a Keithley Source-Measure Unit (SMU 236) and a PR650 (Spectroscan) Source Measurement Unit. (available from Photo Research, Inc.). The results are shown in Table 2, below. A half-lifetime was measured as the time taken until a measured initial luminosity (assumed as 100%) is reduced to 50%.)

TABLE 2

Example	Dopant in EML	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half-lifetime (hr @100 mA/cm ²)
Example 1	Compound 5	6.22	50	3230	6.46	Blue	362
Example 2	Compound 23	6.20	50	3560	7.12	Blue	345
Example 3	Compound 25	6.15	50	3611	7.22	Blue	351
Example 4	Compound 31	6.30	50	3575	7.15	Blue	359
Example 5	Compound 50	6.20	50	3150	6.30	Blue	370

TABLE 2-continued

Example	Dopant in EML	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half-lifetime (hr @100 mA/cm ²)
Example 6	Compound 63	6.20	50	3550	7.10	Blue	346
Example 7	Compound 70	6.18	50	3520	7.04	Blue	350
Example 8	Compound 73	6.22	50	3605	7.21	Blue	360
Example 9	Compound 91	6.20	50	3220	6.44	Blue	315
Comparative Example 1	Compound A	7.01	50	2645	5.29	Blue	258
Comparative Example 2	Compound B	6.75	50	2420	4.84	Blue	260
Comparative Example 3	Compound C	6.89	50	2608	5.21	Blue	245


Referring to Table 1, the organic light-emitting devices of Examples 1 to 9 exhibited improved driving voltages, improved current densities, improved luminances, improved efficiencies, and improved half-lifetimes, compared to those of the organic light-emitting devices of Comparative Examples 1 to 3.

As described above, according to the one or more of the above embodiments, an organic light-emitting device including a condensed cyclic compound of Formula 1 above may have a low driving voltage, a high efficiency, a high luminance, and a long lifetime.

Example embodiments have been disclosed herein, and although specific terms are employed, they are used and are to be interpreted in a generic and descriptive sense only and not for purpose of limitation. In some instances, as would be apparent to one of ordinary skill in the art as of the filing of the present application, features, characteristics, and/or elements described in connection with a particular embodiment may be used singly or in combination with features, characteristics, and/or elements described in connection with other embodiments unless otherwise specifically indicated. Accordingly, it will be understood by those of skill in the art that various changes in form and details may be made without departing from the spirit and scope of the present invention as set forth in the following claims.

What is claimed is:

1. A condensed cyclic compound represented by Formula 1:



wherein, in Formula 1,

L_1 and L_2 are each independently a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_2 - C_{10} heterocycloalk-

enylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_2 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted divalent non-aromatic hetero-condensed polycyclic group,

$a1$ and $a2$ are each independently an integer of 0 to 5,

R_1 to R_4 are each independently a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group,

R_{11} to R_{13} are each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, —Si(Q_1)(Q_2)(Q_3), or —B(Q_4)(Q_5),

$b1$ and $b2$ are each independently an integer of 1 to 3, and $b3$ is an integer of 1 to 4,

wherein at least one substituent of the substituted C_3 - C_{10} cycloalkylene group, the substituted C_2 - C_{10} heterocycloalkylene group, the substituted C_3 - C_{10} cycloalkenylene group, the substituted C_2 - C_{10} heterocycloalkenylene group, the substituted C_6 - C_{60} arylene group,

the substituted C₂-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₂-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₂-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic hetero-condensed polycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, and the substituted C₁-C₆₀ alkoxy group is:

a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₃-C₁₀ cyclo alkyl group, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), or —B(Q₁₆)(Q₁₇);

a C₃-C₁₀ cyclo alkyl group, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic hetero-condensed polycyclic group;

a C₃-C₁₀ cyclo alkyl group, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cyclo alkyl group, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), or —B(Q₂₆)(Q₂₇); or —Si(Q₃₁)(Q₃₂)(Q₃₃), or —B(Q₃₄)(Q₃₅),

wherein Q₁ to Q₅, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₅ are each independently a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cyclo alkyl group, a C₂-C₁₀ heterocyclo alkyl group, a C₃-C₁₀ cyclo alkenyl group, a C₂-C₁₀ heterocyclo alkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic hetero-condensed polycyclic group.

2. The condensed cyclic compound as claimed in claim 1, wherein L₁ and L₂ are each independently:


a phenylene group, a pentalenylene group, an indeylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, a ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isooxazolylenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylenylene group, an indolylenylene group, an indazolylenylene group, a purinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylene group, a benzoimidazolylenylene group, a benzofuranylenylene group, a benzothienophenylene group, an isobenzothiazolylenylene group, a benzooxazolylenylene group, an isobenzooxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylene group, a dibenzofuranylenylene group, a dibenzothienophenylene group, a benzocarbazolylenylene group, or a dibenzocarbazolylenylene group;

a phenylene group, a pentalenylene group, an indeylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a Spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, a ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isooxazolylenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a

129

pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylene group, a cinnolinylenylene group, a carbazolylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzoimidazolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, an isobenzothiazolylenylene group, a benzooxazolylenylene group, an isobenzooxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, or a dibenzocarbazolylenylene, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, an a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indeyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluorantenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a pycenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, a ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, a oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃), in which Q₃₁ to Q₃₃ are each independently a hydrogen, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, or a naphthyl group.

3. The condensed cyclic compound as claimed in claim 1, wherein L₁ and L₂ are each independently represented by one of Formulae 3-1 to 3-32:




Formula 3-1

65


130

-continued


Formula 3-2




Formula 3-3




Formula 3-4




Formula 3-5




Formula 3-6




Formula 3-7




Formula 3-8



Formula 3-9















Formula 3-10



131

-continued




132

-continued

Formula 3-11

5




Formula 3-12

10


Formula 3-13

15




Formula 3-14

20




Formula 3-15

25




Formula 3-16

30




Formula 3-17

35



Formula 3-18

40




Formula 3-19

45

Formula 3-20

50




Formula 3-21

55

Formula 3-22

60



65

Formula 3-23

Formula 3-24

Formula 3-25

Formula 3-26

Formula 3-27

Formula 3-28


Formula 3-29

Formula 3-30

Formula 3-31

133

-continued



wherein, in Formulae 3-1 to 3-32,

Z_1 and Z_2 are each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spirofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, or —Si(Q_{31})(Q_{32})(Q_{33}), in which Q_{31} to Q_{33} are each independently a hydrogen, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, or a naphthyl group,

d_1 is an integer of 1 to 4,

d_2 is an integer of 1 to 3,

d_3 is an integer of 1 to 6,





d_4 is an integer of 1 to 8,

d_5 is 1 or 2,

d_6 is an integer of 1 to 5, and

* and *1 are binding sites to adjacent atoms.

4. The condensed cyclic compound as claimed in claim 1, wherein L_1 and L_2 are each independently represented by one of Formulae 4-1 to 4-23, in which * and *1 are binding sites to adjacent atoms:



Formula 3-32

5

10

15

20

25

30

35

40

Formula 4-1









Formula 4-2

Formula 4-3

Formula 4-4

134

-continued



Formula 4-5

Formula 4-6

Formula 4-7

Formula 4-8

Formula 4-9

Formula 4-10









Formula 4-11

Formula 4-12

65

135


-continued





136

-continued

Formula 4-13



Formula 4-14



Formula 4-15

5. The condensed cyclic compound as claimed in claim 1, wherein a1 and a2 are each independently 0, 1, or 2.

6. The condensed cyclic compound as claimed in claim 1, wherein R₁ to R₄ are each independently:

a phenyl group, a pentalenyl group, an indeyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluo-ranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, a ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzo-quinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, or a dibenzocarbazolyl group; or

a phenyl group, a pentalenyl group, an indeyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluo-ranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a pycenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl

Formula 4-21

Formula 4-22

Formula 4-23

Formula 4-16

Formula 4-17

Formula 4-18

Formula 4-19

Formula 4-20

137

group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, a oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, or a dibenzocarbazolyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluorantenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a pycenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, a ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and Si(Q₃₁)(Q₃₂)(Q₃₃), in which Q₃₁ to Q₃₃ are each independently a hydrogen, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, or a naphthyl group.

7. The condensed cyclic compound as claimed in claim 1, wherein R₁ to R₄ are each independently:


a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an

138


isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, or a dibenzocarbazolyl group; or

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, or a dibenzocarbazolyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃), in which Q₃₁ to Q₃₃ are each independently a hydrogen, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, or a naphthyl group.


8. The condensed cyclic compound as claimed in claim 1, wherein R₁ to R₄ are each independently represented by one of Formulae 5-1 to 5-14:




Formula 5-1



Formula 5-2













Formula 5-3



Formula 5-4

139

-continued



wherein, in Formulae 5-1 to 5-14,
 Y_{31} is O, S, $C(Z_{33})(Z_{34})$, or $N(Z_{35})$,

Z_{31} to Z_{35} are each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl

140

Formula 5-5

5

Formula 5-6

10

Formula 5-7

15

Formula 5-8

20

Formula 5-9

25

Formula 5-10

30

Formula 5-11

35

Formula 5-12

40

Formula 5-13

45

Formula 5-14

50

55

60


group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, or $-\text{Si}(Q_{31})(Q_{32})(Q_{33})$, in which Q_{31} to Q_{33} are each independently a hydrogen, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, or a naphthyl group,

e1 is an integer of 1 to 5,
 e2 is an integer of 1 to 7,
 e3 is an integer of 1 to 3,
 e4 is an integer of 1 to 4,
 e5 is 1 or 2, and


* is a binding site to an adjacent atom.

9. The condensed cyclic compound as claimed in claim 1, wherein R_1 to R_4 are each independently represented by one of Formulae 6-1 to 6-45, in which * is a binding site to an adjacent atom:


Formula 6-1




Formula 6-2




Formula 6-3




Formula 6-4




Formula 6-5




Formula 6-6



Formula 6-7











Formula 6-8



141


-continued

**142**

-continued

Formula 6-9

5




Formula 6-10

10


Formula 6-11

15




Formula 6-12

25




Formula 6-13

30




Formula 6-14

40




Formula 6-15

50




55




Formula 6-16

60



65



Formula 6-17

Formula 6-18

Formula 6-19

Formula 6-20

Formula 6-21

Formula 6-22









Formula 6-23

Formula 6-24

Formula 6-25

143


-continued

**144**

-continued

Formula 6-26

5




Formula 6-27

10


Formula 6-28

15




Formula 6-29

25




Formula 6-30

35




Formula 6-31

40




Formula 6-32

50




Formula 6-33

60



65



Formula 6-34

Formula 6-35

Formula 6-36

Formula 6-37

Formula 6-38





Formula 6-39

Formula 6-40

Formula 6-41

145

-continued



10. The condensed cyclic compound as claimed in claim 1, wherein:

$R_1=R_2=R_3=R_4$;

$R_1=R_3$, $R_1 \neq R_2$, and $R_2=R_4$;

$R_2=R_3$, $R_1 \neq R_2$, and $R_1=R_4$;

$R_1=R_2$, $R_1 \neq R_3$, and $R_3=R_4$;

$R_2=R_3=R_4$, and $R_1 \neq R_2$;

$R_3=R_4$, and $R_1 \neq R_2 \neq R_3$; or

$R_1 \neq R_2 \neq R_3 \neq R_4$.


11. The condensed cyclic compound as claimed in claim 1, wherein R_{11} to R_{13} are each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, or —Si(Q_1)(Q_2)(Q_3), in which Q_1 to Q_3 are each independently a hydrogen, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, or a naphthyl group.

12. The condensed cyclic compound as claimed in claim 1, wherein the condensed cyclic compound represented by Formula 1 is represented by one of Formulae 1A to 1D:

146

Formula 6-42

<Formula 1A>



Formula 6-43

10

<Formula 1B>

Formula 6-44

15

Formula 6-45

20

25

<Formula 1C>

30

35

<Formula 1D>

40

45

50

55

60

65

wherein R_1 to R_4 , R_{11} to R_{13} , L_1 , L_2 , a_1 , a_2 , b_1 , and b_2 in Formulae 1A to 1D are the same as those defined with respect to Formula 1.

13. The condensed cyclic compound as claimed in claim 12, wherein:

the condensed cyclic compound represented by Formula 1 is represented by Formula 1A,

L_1 and L_2 are each independently represented by one of Formulae 4-1 to 4-23, below, in

which * and *' are binding sites to adjacent atoms, a_1 and a_2 are each independently 0, 1, or 2,









R_1 to R_4 are each independently represented by one of Formulae 6-1 to 6-45, below, in which * is a binding site to an adjacent atom,

R_{11} to R_{13} are each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano

147

group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spirofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a benzothienophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, or —Si(Q₁)(Q₂)(Q₃), in which Q₁ to Q₃ are each independently a hydrogen, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, or a naphthyl group, and

b1 to b3 are each independently 1 or 2:



Formula 4-1

Formula 4-2

Formula 4-3

Formula 4-4

Formula 4-5









Formula 4-6

Formula 4-7

Formula 4-8

148

-continued



Formula 4-9

Formula 4-10

Formula 4-11

Formula 4-12

Formula 4-13











Formula 4-14

Formula 4-15











Formula 4-16

149

-continued

**150**

-continued



Formula 4-17

5

Formula 4-18

10

Formula 4-19

20

Formula 4-20

25

Formula 4-21

30

Formula 4-22

35

Formula 4-23

40

Formula 4-24

45

Formula 4-25

50

Formula 6-1

55

Formula 6-2

60

Formula 6-3

65

Formula 6-4

Formula 6-5

Formula 6-6

Formula 6-7

Formula 6-8

Formula 6-9

Formula 6-10








Formula 6-11

Formula 6-12

Formula 6-13

151


-continued

**152**

-continued


Formula 6-14

5



10

Formula 6-15 15




20

25

Formula 6-16


30



35

Formula 6-17

40




45

Formula 6-18

50

Formula 6-19


55



Formula 6-20

60

65



Formula 6-21

Formula 6-22

Formula 6-23

Formula 6-24

Formula 6-25









Formula 6-26

Formula 6-27

Formula 6-28

Formula 6-29


153
-continued



154
-continued

Formula 6-30


5



Formula 6-38

Formula 6-31

10




Formula 6-39

15

Formula 6-32

20




Formula 6-40

25

Formula 6-33


30



Formula 6-41

Formula 6-34

35




Formula 6-42

40

Formula 6-35

45




Formula 6-43

50

Formula 6-36

55



Formula 6-44






60

Formula 6-37






65

14. The condensed cyclic compound as claimed in claim 1, wherein the condensed cyclic compound represented by Formula 1 includes one of Compounds 1 to 96:

155



156




157


158

-continued


11




12




13




14




15



16



17





159

160

-continued



18

19





20

21





22

23





24

25



26

27





161

162

-continued



28

29





30

31





32

33





34

35



36

37





163

164



-continued
38

39





40

41





42

43



44

45





165

166

-continued



46

47





48

49





50

51



52

53





167

168

-continued



54

55





56

57





58

59





60

61



62

63





169

170


-continued

64


65




66



67



68





171

172

-continued



69

70




71


72



73



74




173


174

-continued

75




76




77

78



79




175


176

-continued

80





81




82

83












84



177

178

-continued




179


180

-continued


94



95



96



15. An organic light-emitting device, comprising:
a first electrode;

a second electrode opposite to the first electrode; and
an organic layer between the first electrode and the second
electrode, the organic layer including an emission layer,
wherein the organic layer includes at least one condensed
cyclic compound as claimed in claim 1.

16. The organic light-emitting device as claimed in claim
15, wherein the organic layer includes:

a hole transport region between the first electrode and the
emission layer, the hole transport region including at
least one of a hole injection layer, a hole transport layer,
a buffer layer, and an electron blocking layer, and
an electron transport region between the emission layer
and the second electrode, the electron transport region
including at least one of a hole injection layer, an
electron transport layer, and an electron blocking layer.


17. The organic light-emitting device as claimed in claim
16, wherein the emission layer includes the at least one
condensed cyclic compound.

18. The organic light-emitting device as claimed in claim
17, wherein:

the emission layer further includes a host,
the at least condensed cyclic compound is a dopant, and

the host includes a compound represented by Formula
301A:

<Formula 301A>



wherein, in Formula 301A,

L₃₀₁ is:

a phenylene group, a naphthylene group, a fluorenylene
group, a spiro-fluorenylene group, a benzofluorenylene
group, a dibenzofluorenylene group, a phenanthrene-
nylene group, an anthracenylenylene group, a pyrenylene
group, or a chrysénylene group; or

a phenylene group, a naphthylene group, a fluorenylene
group, a spiro-fluorenylene group, a benzofluorenylene

181


group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, or a chrysenylene group, each substituted with a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a car-

182


xb1 iso, 1, 2, or 3, and
xb2 is 1, 2, 3, or 4.

19. The organic light-emitting device as claimed in claim 16, wherein the hole transport region includes at least one of a compound represented by Formula 201A and a compound represented by Formula 202A:

<Formula 201A>



<Formula 202A>



boxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, or a chrysenyl group,

R₃₀₁ is:

a C₁-C₂₀ alkyl group, or a C₁-C₂₀ alkoxy group;
a C₁-C₂₀ alkyl group or a C₁-C₂₀ alkoxy group, each substituted with a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, or a chrysenyl group;
a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, or a chrysenyl group;
a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, or a chrysenyl group, each substituted with a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, or a chrysenyl group,

wherein, in Formulae 201A and 202A,

L₂₀₁ to L₂₀₃ are each independently:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylene group, a carbazolinylene group, or a triazinylene group; or

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylene group, a carbazolinylene group, or a triazinylene group, each substituted with a deuterium, —F, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolinyl group, or a triazinyl group,


xa1 to xa3 are each independently 0 or 1,

185

group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, or a triazinyl group, and


xa5 is 1 or 2.

20. The organic light-emitting device as claimed in claim 16, wherein the electron transport layer includes at least one of a compound represented by Formula 601 and a compound represented by Formula 602:



<Formula 601>

<Formula 602>



wherein, in Formulae 601 and 602,

Ar_{601} is:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a pyrenylene group, or a chrysenylene group; or

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a pyrenylene group, or a chrysenylene group, each substituted with a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, or a chrysenyl group,

L_{611} to L_{616} are each independently:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a pyrenylene group, a chrysenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a carbazolylene group, or a triazinylene group; or

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a pyrenylene group, a chrysenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a carbazolylene group, or a triazinylene group, each substituted with a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic

186

group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, or a triazinyl group,

E_{601} is:

a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzo-carbazolyl group, or a dibenzocarbazolyl group; or

a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzo-carbazolyl group, or a dibenzocarbazolyl group, each substituted with a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coroneryl group, an obarenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl

187

group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzo-carbazolyl group, or a dibenzocarbazolyl group,

xe1 is 0, 1, 2, or 3,
 xe2 is 1, 2, 3, or 4,
 X_{611} is N or $C-(L_{611})_{xe611}-R_{611}$,
 X_{612} is N or $C-(L_{612})_{xe612}-R_{612}$,
 X_{613} is N or $C-(L_{613})_{xe613}-R_{613}$, at least one of X_{611} to X_{613} being N,
 R_{611} to R_{616} are each independently:
 a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl

188

group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, or a triazinyl group; or
 a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, or a triazinyl group, each substituted with a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an azulenyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, or a triazinyl group, and
 xe611 to xe616 are each independently 0, 1, 2, or 3.

* * * * *

专利名称(译)	缩合环状化合物和包括其的有机发光器件		
公开(公告)号	US9818952	公开(公告)日	2017-11-14
申请号	US14/300540	申请日	2014-06-10
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	JUNG HYE JIN LIM JIN O HAN SANG HYUN		
发明人	JUNG, HYE-JIN LIM, JIN-O HAN, SANG-HYUN		
IPC分类号	H01L51/00 C09K11/06 H01L51/50 C07D493/06		
CPC分类号	H01L51/006 C07D493/06 C09K11/06 H01L51/0061 H01L51/0073 H01L51/5072 H01L51/0081 H01L51/5012 H01L51/5056		
优先权	1020130126105 2013-10-22 KR		
其他公开文献	US20150108440A1		
外部链接	Espacenet USPTO		

摘要(译)

<Formula 1>

缩合环状化合物和有机发光器件，该化合物由式1表示：

