




US 20170084850A1

(19) **United States**(12) **Patent Application Publication**
HWANG et al.(10) **Pub. No.: US 2017/0084850 A1**(43) **Pub. Date: Mar. 23, 2017**(54) **ORGANOMETALLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**(71) Applicant: **SAMSUNG ELECTRONICS CO.,
LTD.**, Suwon-si (KR)(72) Inventors: **Kyuyoung HWANG**, Anyang-si (KR);
Ohyun KWON, Yongin-si (KR);
Seungyeon KWAK, Yongin-si (KR);
Kum Hee LEE, Suwon-si (KR); **Aram
JEON**, Suwon-si (KR); **Hyeonho
CHOI**, Seoul (KR); **Byoungki CHOI**,
Hwaseong-si (KR)(21) Appl. No.: **15/263,850**(22) Filed: **Sep. 13, 2016**(30) **Foreign Application Priority Data**

Sep. 22, 2015 (KR) 10-2015-0134134

Publication Classification(51) **Int. Cl.****H01L 51/00** (2006.01)**C09K 11/02** (2006.01)**C07F 15/00** (2006.01)**C09K 11/06** (2006.01)(52) **U.S. Cl.**CPC **H01L 51/0085** (2013.01); **C09K 11/06**
(2013.01); **C09K 11/025** (2013.01); **C07F**
15/004 (2013.01); **H01L 51/5206** (2013.01)(57) **ABSTRACT**

An organometallic compound represented by Formula 1:



Formula 1

wherein in Formula 1, M in Formula 1 is selected from iridium (Ir), platinum (Pt), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), thulium (Tm), and rhodium (Rh), L_1 in Formula 1 is selected from a ligand represented by Formula 2A or 2B as they are defined in the specification, and L_2 in Formula 1 is selected from a mono-anionic organic ligand.





FIG. 1



ORGANOMETALLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application claims priority to Korean Patent Application No. 10-2015-0134134, filed on Sep. 22, 2015, in the Korean Intellectual Property Office, and all the benefits accruing therefrom under 35 U.S.C. §119, the content of which is incorporated herein in its entirety by reference.

BACKGROUND

[0002] 1. Field

[0003] One or more embodiments relate to an organometallic compound and an organic light-emitting device including the same.

[0004] 2. Description of the Related Art

[0005] Organic light emitting devices (OLEDs) are self-emission devices that have wide viewing angles, high contrast ratios, and short response times. In addition, OLEDs exhibit excellent brightness, driving voltage, and response speed characteristics, and produce full-color images.

[0006] In an example, an organic light-emitting device includes an anode, a cathode, and an organic layer that is disposed between the anode and the cathode, wherein the organic layer includes an emission layer. A hole transport region may be disposed between the anode and the emission layer, and an electron transport region may be disposed between the emission layer and the cathode. Holes provided from the anode may move toward the emission layer through the hole transport region, and electrons provided from the cathode may move toward the emission layer through the electron transport region. The holes and the electrons recombine in the emission layer to produce excitons. These excitons change from an excited state to a ground state, thereby generating light.


[0007] Various types of organic light emitting devices are known. However, there still remains a need in OLEDs having low driving voltage, high efficiency, high brightness, and long lifespan.

SUMMARY

[0008] One or more embodiments relate to a novel organometallic compound and an organic light-emitting device including the same.


[0009] Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented exemplary embodiments.

[0010] An aspect provides an organometallic compound represented by Formula 1 below:




Formula 1

Formula 2A



Formula 2B



[0011] wherein

[0012] M in Formula 1 is selected from iridium (Ir), platinum (Pt), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), thulium (Tm), and rhodium (Rh),

[0013] L₁ in Formula 1 is a ligand represented by Formula 2A or 2B,

[0014] L₂ in Formula 1 is selected from a mono-anionic organic ligand,

[0015] * and *' in Formulae 2A and 2B each indicate a binding site to M in Formula 1,

[0016] n₁ in Formula 1 is 1, 2, or 3, and when n₁ is 2 or more, 2 or more groups L₁ are identical to or different from each other,

[0017] n₂ in Formula 1 is 0, 1, 2, or 3, and when n₂ is 2 or more, 2 or more groups L₂ are identical to or different from each other,

[0018] Y₁ and Y₃ in Formulae 2A and 2B are each independently carbon or nitrogen,

[0019] Y₂ in Formulae 2A and 2B is carbon,

[0020] a bond between Y₂ and Y₃ in Formula 2B is a double bond,

[0021] CY₁ in Formulae 2A and 2B is a C₂-C₃₀ heterocyclic group,

[0022] in Formulae 2A and 2B, X₁ is N or C(R₁), X₂ is N or C(R₂), X₃ is N or C(R₃), X₄ is N or C(R₄), X₅ is N or C(R₅), X₆ is N or C(R₆), X₇ is N or C(R₇),

[0023] T₁ in Formula 2B is selected from a single bond, *—O—*, *—S—*, *—N(R₃₁)—*, *—C(R₃₁)(R₃₂)—*, *—C(R₃₁)=C(R₃₂)—*, and *—Si(R₃₁)(R₃₂)—*, and R₃₁ and R₃₂ are optionally bound to each other to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group or a substituted or unsubstituted C₂-C₃₀ heterocyclic group,

[0024] R_1 to R_7 , R_{10} , R_{31} , and R_{32} are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, —SF₅, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q_1)(Q_2), —Si(Q_3)(Q_4)(Q_5), —B(Q_6)(Q_7), and —P(=O)(Q_8)(Q_9),

[0025] a_1 is an integer selected from 0 to 5,

[0026] two or more among R_1 to R_3 in Formulae 2A and 2B are optionally bound to each other to form a substituted or unsubstituted C_6 - C_{30} carbocyclic group or a substituted or unsubstituted C_2 - C_{30} heterocyclic group,

[0027] two or more among R_4 to R_7 in Formula 2A are optionally bound to each other to form a substituted or unsubstituted C_5 - C_{30} carbocyclic group or a substituted or unsubstituted C_2 - C_{30} heterocyclic group,

[0028] two or more among R_4 to R_6 in Formula 2B are optionally bound to each other to form a substituted or unsubstituted C_5 - C_{30} carbocyclic group or a substituted or unsubstituted C_2 - C_{30} heterocyclic group,

[0029] two or more among groups R_{10} in the number of a_1 in Formulae 2A and 2B are optionally bound to each other to form a substituted or unsubstituted C_5 - C_{30} carbocyclic group or a substituted or unsubstituted C_2 - C_{30} heterocyclic group, and

[0030] at least one of substituents of the substituted C_5 - C_{30} carbocyclic group, substituted C_2 - C_{30} heterocyclic group, substituted C_1 - C_{60} alkyl group, substituted C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group, substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10} cycloalkyl group, substituted C_1 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substituted C_1 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_1 - C_{60} heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

[0031] deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0032] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, —F,

—Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q_{11})(Q_{12}), —Si(Q_{13})(Q_{14})(Q_{15}), —B(Q_{16})(Q_{17}), and —P(=O)(Q_{18})(Q_{19}),

[0033] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0034] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q_{21})(Q_{22}), —Si(Q_{23})(Q_{24})(Q_{25}), —B(Q_{26})(Q_{27}), and —P(=O)(Q_{28})(Q_{29}); and

[0035] —N(Q_{31})(Q_{32}), —Si(Q_{33})(Q_{34})(Q_{35}), —B(Q_{36})(Q_{37}) and —P(=O)(Q_{38})(Q_{39}),

[0036] wherein Q_1 to Q_9 , Q_{11} to Q_{19} , Q_{21} to Q_{29} , and Q_{31} to Q_{39} are each independently selected from a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} alkyl group, and a C_6 - C_{60} aryl group.

[0037] According to another aspect of an exemplary embodiment, an organic light-emitting device includes:

[0038] a first electrode,

[0039] a second electrode, and

[0040] an organic layer disposed between the first electrode and the second electrode,

[0041] wherein the organic layer includes an emission layer and at least one of the organometallic compounds described above.

[0042] The organometallic compounds may act as a dopant in the emission layer.

BRIEF DESCRIPTION OF THE DRAWING

[0043] These and/or other aspects will become apparent and more readily appreciated from the following description of the exemplary embodiments, taken in conjunction with FIG. 1 which is a schematic view of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION

[0044] Reference will now be made in detail to exemplary embodiments, examples of which are illustrated in the accompanying drawings, wherein like reference numerals refer to like elements throughout. In this regard, the present exemplary embodiments may have different forms and should not be construed as being limited to the descriptions set forth herein. Accordingly, the exemplary embodiments are merely described below, by referring to the figures, to explain aspects. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

[0045] It will be understood that when an element is referred to as being “on” another element, it can be directly in contact with the other element or intervening elements may be present therebetween. In contrast, when an element is referred to as being “directly on” another element, there are no intervening elements present.

[0046] It will be understood that, although the terms first, second, third etc. may be used herein to describe various elements, components, regions, layers, and/or sections, these elements, components, regions, layers, and/or sections should not be limited by these terms. These terms are only used to distinguish one element, component, region, layer, or section from another element, component, region, layer, or section. Thus, a first element, component, region, layer, or section discussed below could be termed a second element, component, region, layer, or section without departing from the teachings of the present embodiments.

[0047] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. As used herein, the singular forms “a,” “an,” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.

[0048] The term “or” means “and/or.” It will be further understood that the terms “comprises” and/or “comprising,” or “includes” and/or “including” when used in this specification, specify the presence of stated features, regions, integers, steps, operations, elements, and/or components, but do not preclude the presence or addition of one or more other


features, regions, integers, steps, operations, elements, components, and/or groups thereof.

[0049] Unless otherwise defined, all terms (including technical and scientific terms) used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this general inventive concept belongs. It will be further understood that terms, such as those defined in commonly used dictionaries, should be interpreted as having a meaning that is consistent with their meaning in the context of the relevant art and the present disclosure, and will not be interpreted in an idealized or overly formal sense unless expressly so defined herein.

[0050] Exemplary embodiments are described herein with reference to cross section illustrations that are schematic illustrations of idealized embodiments. As such, variations from the shapes of the illustrations as a result, for example, of manufacturing techniques and/or tolerances, are to be expected. Thus, embodiments described herein should not be construed as limited to the particular shapes of regions as illustrated herein but are to include deviations in shapes that result, for example, from manufacturing. For example, a region illustrated or described as flat may, typically, have rough and/or nonlinear features. Moreover, sharp angles that are illustrated may be rounded. Thus, the regions illustrated in the figures are schematic in nature and their shapes are not intended to illustrate the precise shape of a region and are not intended to limit the scope of the present claims.

[0051] “About” or “approximately” as used herein is inclusive of the stated value and means within an acceptable range of deviation for the particular value as determined by one of ordinary skill in the art, considering the measurement in question and the error associated with measurement of the particular quantity (i.e., the limitations of the measurement system). For example, “about” can mean within one or more standard deviations, or within $\pm 30\%$, 20% , 10% , 5% of the stated value.

[0052] An organometallic compound according to an embodiment is represented by Formula 1 below:



Formula 1

[0053] M in Formula 1 may be selected from iridium (Ir), platinum (Pt), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), thulium (Tm), and rhodium (Rh).

[0054] For example, M in Formula 1 may be selected from iridium, platinum, and osmium, but is not limited thereto.

[0055] L_1 in Formula 1 may be a group represented by Formula 2A or 2B. Formulae 2A and 2B are the same as described below.

[0056] L_2 in Formula 1 may be selected from mono-anionic organic ligands.

[0057] * and *' in Formulae 2A and 2B may each be a binding site to M in Formula 1.

[0058] n_1 in Formula 1 may be 1, 2, or 3, and when n_1 is 2 or more, 2 or more groups L_1 may be identical to or different from each other.

[0059] n_2 in Formula 1 may be 0, 1, 2, or 3, and when n_2 is 2 or more, 2 or more groups L_2 may be identical to or different from each other.

[0060] For example, in Formula 1,

[0061] M may be Ir or Os and the sum of n_1 and n_2 may be 3 or 4; or

[0062] M may be Pt and the sum of n_1 and n_2 may be 2, but M is not limited thereto.

[0063] In an embodiment, in Formula 1,
 [0064] M may be Ir, and when n₁ may be 2, n₂ may be 1;
 [0065] M may be Os, and when n₁ may be 2, and n₂ may be 2; or
 [0066] M may be Pt, and when n₁ may be 2, and n₂ may be 0, but M is not limited thereto.
 [0067] Y₁ and Y₃ in Formulae 2A and 2B may each independently be carbon or nitrogen.
 [0068] For example, Y₃ in Formulae 2A and 2B may be carbon.
 [0069] Y₂ in Formulae 2A and 2B may be carbon, and the bond between Y₂ and Y₃ in Formula 2B may be a double bond.
 [0070] CY₁ in Formulae 2A and 2B may be a C₂-C₃₀ heterocyclic group.

[0071] For example, CY₁ in Formulae 2A and 2B may include a five-membered ring containing at least one nitrogen atom as a ring-forming atom.

[0072] In an embodiment, CY₁ in Formulae 2A and 2B may be selected from a pyrrole ring, an isoindole ring, an indole ring, an indazole ring, a pyrazole ring, an imidazole ring, a triazole ring, an oxazole ring, an isoxazole ring, an oxadiazole ring, a thiazole ring, an isothiazole ring, a thiadiazole ring, a purine ring, a benzoimidazole ring, an isobenzothiazole ring, a benzoxazole ring, an isobenzoxazole ring, an imidazopyridine ring, an imidazopyrimidine ring, and a pyrrolopyridine ring, but is not limited thereto.
 [0073] In some embodiments, CY₁ in Formulae 2A and 2B may each independently be a pyrrole ring, an isoindole ring, an indole ring, a pyrrolopyridine ring, a pyrazole ring, or a benzoimidazole ring.

[0074] In Formulae 2A and 2B, X₁ may be N or C(R₁), X₂ may be N or C(R₂), X₃ may be N or C(R₃), X₄ may be N or C(R₄), X₅ may be N or C(R₅), X₆ may be N or C(R₆), and X₇ may be N or C(R₇).


[0075] For example, in Formulae 2A and 2B, X₁ may be C(R₁), X₂ may be C(R₂), X₃ may be C(R₃), X₄ may be C(R₄), X₅ may be C(R₅), X₆ may be C(R₆), and X₇ may be C(R₇), but they are not limited thereto.

[0076] In Formula 2B, T₁ may be selected from a single bond, *—O—*, *—S—*, *—N(R₃₁)—*, *—C(R₃₁)(R₃₂)—*, *—C(R₃₁)=C(R₃₂)—*, and *—Si(R₃₁)(R₃₂)—*, and R₃₁ and R₃₂ are optionally bound to each other to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group, or a substituted or unsubstituted C₂-C₃₀ heterocyclic group (for example, a substituted or unsubstituted cyclopropane, a substituted or unsubstituted cyclobutane, a substituted or unsubstituted cyclopentane, a substituted or unsubstituted cyclohexane, a substituted or unsubstituted benzene).

[0077] For example, T₁ in Formula 2B may be selected from *—O—*, *—S—*, *—C(R₃₁)(R₃₂)—* and groups represented by Formulae 11-1 to 11-4:



Formula 11-1




Formula 11-2

-continued

Formula 11-3



Formula 11-4



[0078] * and *' in Formulae 11-1 to 11-4 each indicate a binding site to a neighboring atom.

[0079] R₁ to R₇, R₁₀, R₃₁, and R₃₂ in Formula 1 may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, —SF₅, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), and —P(=O)(Q₈)(Q₉).

[0080] For example, R₁ to R₇, R₁₀, R₃₁, and R₃₂ in Formula 1 may each independently be selected from:

[0081] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, —SF₅, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0082] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl (adamantyl) group, a norbornanyl (norbornyl) group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

[0083] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a

phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a

benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and
[0084] $-\text{N}(\text{Q}_1)(\text{Q}_2)$, $-\text{Si}(\text{Q}_3)(\text{Q}_4)(\text{Q}_5)$, $-\text{B}(\text{Q}_6)(\text{Q}_7)$, and $-\text{P}(=\text{O})(\text{Q}_8)(\text{Q}_9)$.

[0085] Q_1 to Q_9 may each independently be selected from:

[0086] $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCH}_3$, $-\text{CHDCH}_2\text{H}$, $-\text{CHDCH}_2\text{H}$, $-\text{CHDCH}_3$, $-\text{CD}_2\text{CD}_2\text{H}$, and $-\text{CD}_2\text{CDH}_2$;

[0087] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0088] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C_1 - C_{10} alkyl group, and a phenyl group.

[0089] In some embodiments, R_1 to R_7 , R_{10} , R_{31} , and R_{32} in Formula 1 may each independently be

[0090] hydrogen, deuterium, $-\text{F}$, a cyano group, a nitro group, $-\text{SF}_5$, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0091] a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrim-

idinyll group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, $-F$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0092] $-N(Q_1)(Q_2)$, $-Si(Q_3)(Q_4)(Q_5)$, $-B(Q_5)(Q_7)-P(=O)(Q_8)(Q_9)$, and

[0093] Q_1 to Q_9 may each independently be selected from:

[0094] $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CH_2CH_3$, $-CH_2CD_3$, $-CH_2CD_2H$, $-CH_2CDH_2$, $-CHDC H_3$, $-CHDCD_2H$, $-CHDCDH_2$, $-CHDCD_3$, $-CD_2CD_3$, $-CD_2CD_2H$, and $-CD_2CDH_2$;

[0095] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0096] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C_1 - C_{10} alkyl group, and a phenyl group.


[0097] In some embodiments, R_1 to R_7 , R_{10} , R_{31} , and R_{32} in Formula 1 may each independently be selected from hydrogen, deuterium, $-F$, a cyano group, a nitro group, $-SF_5$, $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, groups represented by Formulae 9-1 to 9-19, groups represented by Formulae 10-1 to 10-36, and $-Si(Q_3)(Q_4)(Q_5)$, and

[0098] Q_3 to Q_5 may each independently be selected from:

[0099] $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CH_2CH_3$, $-CH_2CD_3$, $-CH_2CD_2H$, $-CH_2CDH_2$, $-CHDC H_3$, $-CHDCD_2H$, $-CHDCDH_2$, $-CHDCD_3$, $-CD_2CD_3$, $-CD_2CD_2H$, and $-CD_2CDH_2$;

[0100] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0101] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C_1 - C_{10} alkyl group, and a phenyl group:



Formula 9-1


Formula 9-2

Formula 9-3


-continued




Formula 9-4




Formula 9-5




Formula 9-6




Formula 9-7




Formula 9-8




Formula 9-9




Formula 9-10




Formula 9-11




Formula 9-12




Formula 9-13




Formula 9-14




Formula 9-15




Formula 9-16



Formula 9-17















Formula 9-18



Formula 9-19

-continued



Formula 10-1

Formula 10-2

Formula 10-3

Formula 10-4

Formula 10-5

Formula 10-6

Formula 10-7

Formula 10-8











Formula 10-9

Formula 10-10

Formula 10-11

Formula 10-12

-continued



Formula 10-14

Formula 10-16

Formula 10-18










Formula 10-19

Formula 10-20

Formula 10-21

Formula 10-22

-continued



Formula 10-23

Formula 10-24

Formula 10-25

Formula 10-26

Formula 10-27






Formula 10-28

Formula 10-29

Formula 10-30

Formula 10-31

-continued



Formula 10-32

Formula 10-33

Formula 10-34

Formula 10-35

Formula 10-36

[0102] * in Formulae 9-1 to 9-19 and 10-1 to 10-36 indicates a binding site to a neighboring atom.

[0103] a₁ in Formulae 2A and 2B indicates the number of groups R₁₀, and may be an integer selected from 0 to 5. When a₁ is 2 or more, 2 or more groups R₁ may be identical to or different from each other.

[0104] Two neighboring substituents among R₁ to R₃ in Formulae 2A and 2B may optionally be bound to each other to form a substituted or unsubstituted C₅-C₃₀ carbocyclic ring or a substituted or unsubstituted C₂-C₃₀ heterocyclic ring (for example, a substituted or unsubstituted cyclopentane ring, a substituted or unsubstituted cyclopentene ring, a substituted or unsubstituted cyclohexane ring, a substituted or unsubstituted cyclohexene ring, a substituted or unsubstituted adamantane ring, a substituted or unsubstituted bicycloheptane ring, a substituted or unsubstituted bicycloheptene ring, a substituted or unsubstituted benzene ring, a substituted or unsubstituted naphthalene ring, a substituted or unsubstituted pyridine ring, a substituted or unsubstituted pyrimidine ring, or the like).

[0105] Two neighboring substituents among R₄ to R₇ in Formula 2A may optionally be bound to each other to form a substituted or unsubstituted C₅-C₃₀ carbocyclic ring or a

substituted or unsubstituted C_2 - C_{30} heterocyclic ring (for example, a substituted or unsubstituted cyclopentane ring, a substituted or unsubstituted cyclopentene ring, a substituted or unsubstituted cyclohexane ring, a substituted or unsubstituted cyclohexene ring, a substituted or unsubstituted adamantane ring, a substituted or unsubstituted bicycloheptane ring, a substituted or unsubstituted bicycloheptene ring, a substituted or unsubstituted benzene ring, a substituted or unsubstituted naphthalene ring, a substituted or unsubstituted pyridine ring, a substituted or unsubstituted pyrimidine ring, or the like).


[0106] Two neighboring substituents in R_4 to R_6 in Formula 2B may optionally be bound to each other to form a substituted or unsubstituted C_5 - C_{30} carbocyclic ring or a substituted or unsubstituted C_2 - C_{30} heterocyclic ring (for example, a substituted or unsubstituted cyclopentane ring, a substituted or unsubstituted cyclopentene ring, a substituted or unsubstituted cyclohexane ring, a substituted or unsubstituted cyclohexene ring, a substituted or unsubstituted adamantane ring, a substituted or unsubstituted bicycloheptane ring, a substituted or unsubstituted bicycloheptene ring, a substituted or unsubstituted benzene ring, a substituted or unsubstituted naphthalene ring, a substituted or unsubstituted pyridine ring, a substituted or unsubstituted pyrimidine ring, or the like).

[0107] Two or more of groups R_{10} in the number of a1 in Formulae 2A and 2B may optionally be bound to each other to form a substituted or unsubstituted $_{05-030}$ carbocyclic ring or a substituted or unsubstituted C_2 - C_{30} heterocyclic ring (for example, a substituted or unsubstituted cyclopentane ring, a substituted or unsubstituted cyclopentene ring, a substituted or unsubstituted cyclohexane ring, a substituted or unsubstituted cyclohexene ring, a substituted or unsubstituted adamantane ring, a substituted or unsubstituted bicycloheptane ring, a substituted or unsubstituted bicycloheptene ring, a substituted or unsubstituted benzene ring, a substituted or unsubstituted naphthalene ring, a substituted or unsubstituted pyridine ring, a substituted or unsubstituted pyrimidine ring, or the like).

[0108] At least one of X_1 to X_7 in Formulae 2A and 2B may be C(CN). Accordingly, the organometallic compound represented by Formula 1 may have high color purity and high heat resistance.


[0109] In an embodiment, X_5 in Formulae 2A and 2B may be C(CN), but is not limited thereto.

[0110] In an embodiment, ligand L_1 in Formula 1 may be selected from ligands represented by Formulae 2A-1 to 2A-4 and 2B-1 to 2B-3:




Formula 2A-1


-continued




Formula 2A-2




Formula 2A-3



Formula 2A-4




Formula 2B-1



Formula 2B-2

-continued

Formula 2B-3



[0111] In Formulae 2A-1 to 2A-4 and 2B-1 to 2B-3,

[0112] descriptions of X_1 to X_7 and T_1 are the same as described herein,

[0113] X_{11} is N or C(R_{11}), X_{12} is N or C(R_{12}), X_{13} is N or C(R_{13}), X_{14} is N or C(R_{14}), and X_{15} is N or C(R_{15}),

[0114] X_{21} is O, S, N(R_{21}), or C(R_{22})(R_{23}),

[0115] X_{21} is O, S, N(R_{24}), or C(R_{25})(R_{26}),

[0116] descriptions of R_{11} to R_{15} and R_{21} to R_{26} are the same as described in connection with R_{10} , and

[0117] * and *' each indicate a binding site to M in Formula 1.

[0118] For example, in Formulae 2A-1 to 2A-4 and 2B-1 to 2B-3,

[0119] X_1 is C(R_1), X_2 is C(R_2), X_3 is C(R_3), X_4 is C(R_4), X_5 is C(R_5), X_6 is C(R_6), and X_7 is C(R_7),

[0120] T_1 is selected from $*-O-*'$, $*-S-*'$, $*-C(R_{31})(R_{32})-*'$, and groups represented by Formulae 11-1 to 11-4,

[0121] R_1 to R_7 , R_{11} to R_{15} , R_{21} to R_{26} , R_{31} , and R_{32} are each independently selected from: hydrogen, deuterium, —F, a cyano group, a nitro group, —SF₅, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0122] a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a

sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0123] —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), —P(=O)(Q₈)(Q₉); and

[0124] Q₁ to Q₉ are each independently selected from:

[0125] —CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCH₂CD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;

[0126] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0127] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C₁-C₁₀ alkyl group, and a phenyl group,

[0128] two neighboring substituents among R_1 to R_3 may optionally be bound to each other to form a substituted or unsubstituted C₅-C₁₀ carbocyclic ring,

[0129] two neighboring substituents among R_4 to R_7 may optionally be bound to each other to form a substituted or unsubstituted C₅-C₁₀ carbocyclic ring, and

[0130] two neighboring substituents among R_{11} to R_{15} and R_{21} to R_{26} may optionally be bound to each other to form a substituted or unsubstituted C₅-C₁₀ carbocyclic ring (for example, a substituted or unsubstituted cyclopentane ring, a substituted or unsubstituted cyclopentene ring, a substituted or unsubstituted cyclohexane ring, a substituted or unsubstituted cyclohexene ring, a substituted or unsubstituted adamantane ring, a substituted or unsubstituted bicycloheptane ring, a substituted or unsubstituted bicycloheptene ring, a substituted or unsubstituted benzene ring, or a substituted or unsubstituted naphthalene ring), but is not limited thereto.

[0131] In some embodiments, in Formulae 2A-1 to 2A-4 and 2B-1 to 2B-3,

[0132] X_1 is C(R_1), X_2 is C(R_2), X_3 is C(R_3), X_4 is C(R_4), X_5 is C(R_5), X_6 is C(R_6), and X_7 is C(R_7),

[0133] T_1 is selected from $*-O-*'$, $*-S-*'$, $*-C(R_{31})(R_{32})-*'$, and groups represented by Formulae 11-1 to 11-4,

[0134] R_1 to R_7 , R_{11} to R_{15} , R_{21} to R_{26} , R_{31} , and R_{32} are each independently selected from hydrogen, deuterium, —F, a cyano group, a nitro group, —SF₅, —CH₃, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, groups rep-

resented by Formulae 9-1 to 9-19, groups represented by Formulae 10-1 to 10-36, and $-\text{Si}(\text{Q}_3)(\text{Q}_4)(\text{Q}_5)$,

[0135] Q_3 to Q_5 are each independently selected from:


[0136] $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCCH}_3$, $-\text{CHDCD}_2\text{H}$, $-\text{CHDCDH}_2$, $-\text{CHDCD}_3$, $-\text{CD}_2\text{CD}_3$, $-\text{CD}_2\text{CD}_2\text{H}$, and $-\text{CD}_2\text{CDH}_2$;

[0137] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and


[0138] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C_1 - C_{10} alkyl group, and a phenyl group, but is not limited thereto.

[0139] In an embodiment, at least one selected from X_1 to X_7 in Formulae 2A-1 to 2A-4 and 2B-1 to 2B-3 may be $\text{C}(\text{CN})$. In some embodiments, in Formulae 2A-1 to 2A-4, and 2B-1 to 2B-3, X_5 may be $\text{C}(\text{CN})$ or X_{12} may be $\text{C}(\text{CN})$. Accordingly, the organometallic compound represented by Formula 1 may have a high color purity and a high heat resistance.

[0140] In some embodiments, ligand L_1 in Formula 1 may be selected from ligands represented by Formulae 2A-1(1) to 2A-1(6):




Formula 2A-1(1)




Formula 2A-1(2)


-continued




Formula 2A-1(3)



Formula 2A-1(4)



Formula 2A-1(5)



Formula 2A-1(6)

[0141] In Formulae 2A-1(1) to 2A-1(6),

[0142] X_1 is N or $\text{C}(\text{R}_1)$, X_2 is N or $\text{C}(\text{R}_2)$, X_3 is N or $\text{C}(\text{R}_3)$, X_4 is N or $\text{C}(\text{R}_4)$, X_5 is N or $\text{C}(\text{R}_5)$, X_6 is N or $\text{C}(\text{R}_6)$, and X_7 is N or $\text{C}(\text{R}_7)$,

[0143] T_1 is selected from $*-O-*$, $*-S-*$, $*-C(R_{31})(R_{32})-*$, and groups represented by Formulae 11-1 to 11-4,

[0144] X_{11} is N or C(R_{11}),

[0145] R_1 to R_7 , R_{11} , R_{31} , R_{32} , and R_{41} to R_{48} are each independently selected from:

[0146] hydrogen, deuterium, $-F$, a cyano group, a nitro group, $-SF_5$, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0147] a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, $-F$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0148] $-N(Q_1)(Q_2)$, $-Si(Q_3)(Q_4)(Q_5)$, $-B(Q_6)(Q_7)$ and $-P(=O)(Q_8)(Q_9)$,

[0149] Q_1 to Q_9 are each independently selected from:

[0150] $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CH_2CH_3$, $-CH_2CD_3$, $-CH_2CD_2H$, $-CH_2CDH_2$, $-CHDCH_3$, $-CHDCD_2H$, $-CHDCDH_2$, $-CHDCD_3$, $-CD_2CD_3$, $-CD_2CD_2H$, and $-CD_2CDH_2$;

[0151] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl

group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0152] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C_1 - C_{10} alkyl group, and a phenyl group;

[0153] two neighboring substituents among R_1 to R_3 may optionally be bound to each other to form a substituted or unsubstituted C_5 - C_{10} carbocyclic ring (for example, a substituted or unsubstituted benzene ring, a substituted or unsubstituted naphthalene ring, or the like),

[0154] two neighboring substituents among R_4 to R_7 may optionally be bound to each other to form a substituted or unsubstituted C_5 - C_{10} carbocyclic ring (for example, a substituted or unsubstituted benzene ring, a substituted or unsubstituted naphthalene ring, or the like), and

[0155] $*$ and $*$ may each be a binding site to M in Formula 1.

[0156] In some embodiments, in Formulae 2A-1(1) to 2A-1(6),

[0157] X_1 is N or C(R_1), X_2 is N or C(R_2), X_3 is N or C(R_3), X_4 is N or C(R_4), X_5 is N or C(R_5), X_6 is N or C(R_6), and X_7 is N or C(R_7),

[0158] T_1 is selected from $*-O-*$, $*-S-*$, $*-C(R_{31})(R_{32})-*$, and groups represented by Formulae 11-1 to 11-4,

[0159] X_{11} is N or C(R_{11}),

[0160] R_1 to R_7 , R_{11} , R_{31} , R_{32} , and R_{41} to R_{48} are each independently selected from hydrogen, deuterium, $-F$, a cyano group, a nitro group, $-SF_5$, $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, groups represented by Formulae 9-1 to 9-19, groups represented by Formulae 10-1 to 10-36, and $-Si(Q_3)(Q_4)(Q_5)$,

[0161] Q_3 to Q_5 are each independently selected from:


[0162] $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CH_2CH_3$, $-CH_2CD_3$, $-CH_2CD_2H$, $-CH_2CDH_2$, $-CHDCH_3$, $-CHDCD_2H$, $-CHDCDH_2$, $-CHDCD_3$, $-CD_2CD_3$, $-CD_2CD_2H$ and $-CD_2CDH_2$;

[0163] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group and a naphthyl group; and


[0164] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C_1 - C_{10} alkyl group, and a phenyl group, but they are not limited thereto.

[0165] In an embodiment, at least one selected from X_1 to X_7 in Formulae 2A-1(1) to 2A-1(6) may be C(CN). In some embodiments, X_5 in Formulae 2A-1(1) to 2A-1(6) may be C(CN). Accordingly, the organometallic compound represented by Formula 1 may have high color purity and high heat resistance.


[0166] In some embodiments, ligand L_1 in Formula 1 may be selected from ligands represented by Formulae 4-1 to 4-7, 4-11 to 4-17, 4-21 to 4-27, 4-31 to 4-37, 4-41 to 4-44, 4-51 to 4-54, 4-61 to 4-64, and 4-71 to 4-74:




Formula 4-1




Formula 4-2




Formula 4-3




Formula 4-4




Formula 4-5



Formula 4-6




Formula 4-7




Formula 4-11

-continued




Formula 4-12


-continued




Formula 4-15




Formula 4-13




Formula 4-16



Formula 4-14




Formula 4-17




Formula 4-21

-continued


Formula 4-26




Formula 4-27




Formula 4-31



Formula 4-32




-continued




Formula 4-33


-continued




Formula 4-37




Formula 4-34




Formula 4-41




Formula 4-35



Formula 4-42




Formula 4-36




Formula 4-43

-continued




Formula 4-44


-continued




Formula 4-53




Formula 4-51




Formula 4-54



Formula 4-52




Formula 4-61




Formula 4-62

-continued




Formula 4-63

-continued




Formula 4-73


Formula 4-64




Formula 4-74



Formula 4-71



Formula 4-72



[0167] Regarding Formulae 4-1 to 4-7, 4-11 to 4-17, 4-21 to 4-27, 4-31 to 4-37, 4-41 to 4-44, 4-51 to 4-54, 4-61 to 4-64, and 4-71 to 4-74, descriptions of X_{11} , X_{14} , T_1 , R_1 to R_7 , R_{11} to R_{15} , and R_{41} to R_{48} are the same as described herein, descriptions of R_8 and R_9 are the same as described in connection with R_1 , and * and *' each indicates a binding site to M in Formula 1.

[0168] For example, in Formulae 4-1 to 4-7, 4-11 to 4-17, 4-21 to 4-27, 4-31 to 4-37, 4-41 to 4-44, 4-51 to 4-54, 4-61 to 4-64, and 4-71 to 4-74,

[0169] X_{11} is N or C(R_{11}),

[0170] X_{14} is N or C(R_{14}),

[0171] T_1 is selected from $^*O^*$, $^*S^*$, $^*C(R_{31})(R_{32})^*$, and groups represented by Formulae 11-1 to 11-4,

[0172] R_1 to R_9 , R_{11} to R_{15} , R_{31} , R_{32} , and R_{41} to R_{48} are each independently selected from hydrogen, deuterium, $-F$, a cyano group, a nitro group, $-SF_5$, $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, groups represented by Formulae 9-1 to 9-19, groups represented by Formulae 10-1 to 10-36, and $-Si(Q_3)(Q_4)(Q_5)$,

[0173] Q_3 to Q_5 are each independently selected from:

[0174] $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CH_2CH_3$, $-CH_2CD_3$, $-CH_2CD_2H$, $-CH_2CDH_2$, $-CHDCH_3$, $-CHDCD_2H$, $-CHDCDH_2$, $-CHDCD_3$, $-CD_2CD_3$, $-CD_2CD_2H$, and $-CD_2CDH_2$;

[0175] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0197] —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), and —P(=O)(Q₈)(Q₉),

[0198] Q₁ to Q₉ are each independently selected from:

[0199] —CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDC₂H₃, —CHDCD₂H, —CHDCDH₂, —CHDCD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;


[0200] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0201] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C₁-C₁₀ alkyl group, and a phenyl group.


[0202] In an embodiment, ligand L₂ in Formula 1 may be selected from ligands represented by Formulae 3-1 to 3-15, but is not limited thereto:

-continued


Formula 3-4




Formula 3-5




Formula 3-6




Formula 3-7




Formula 3-8




Formula 3-9




Formula 3-10




Formula 3-11




Formula 3-1




Formula 3-2




Formula 3-3




-continued




Formula 3-12



Formula 3-13



Formula 3-14



Formula 3-15

[0203] In Formulae 3-1 to 3-15,

[0204] X_{31} is N or C(Z_{1a}), X_{32} is N or C(Z_{1b}),

[0205] X_{41} is O, S, N(Z_{1a}), or C(Z_{1a})(Z_{1b}),

[0206] descriptions of Z_1 to Z_4 and Z_{11} to Z_{14} are the same as described herein,

[0207] descriptions of Z_{1a} and Z_{1b} are the same as described in connection with Z_1 ,

[0208] e2 is 1 or 2,

[0209] e3 is an integer selected from 1 to 3,

[0210] e4 is an integer selected from 1 to 4,

[0211] * and *' each indicate a binding site to M in Formula 1.

[0212] For example, Z_1 to Z_4 , Z_{1a} , Z_{1b} , and Z_{11} to Z_{14} in Formulae 3-1 to 3-15 may each independently be selected from:

[0213] hydrogen, deuterium, —F, a cyano group, a nitro group, —SF₅, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0214] a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0215] —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), and —P(=O)(Q₈)(Q₉),

[0216] Q₁ to Q₉ are each independently selected from:

[0217] —CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCH₃, —CHDCH₂H, —CHDCHD₂, —CHDCHD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;

[0218] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0219] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C₁-C₁₀ alkyl group, and a phenyl group, but they are not limited thereto.

[0220] In an embodiment, ligand L₁ in Formula 1 may be selected from ligands represented by Formulae 2A-1 to 2A-4 and 2B-1 to 2B-3, and ligand L₂ in Formula 1 may be selected from ligands represented by Formulae 3A to 3F (for example, ligands represented by Formulae 3-1 to 3-15).


[0221] In some embodiments, ligand L₁ in Formula 1 may be selected from ligands represented by Formulae 2A-1(1) to 2A-1(6), and ligand L₂ in Formula 1 may be selected from ligands represented by Formulae 3A to 3F (for example, ligands represented by Formulae 3-1 to 3-15).

[0222] In some embodiments, ligand L₁ in Formula 1 may be selected from ligands represented by Formulae 4-1 to 4-7, 4-11 to 4-17, 4-21 to 4-27, 4-31 to 4-37, 4-41 to 4-44, 4-51 to 4-54, 4-61 to 4-64, and 4-71 to 4-74, and ligand L₂ in Formula 1 may be selected from ligands represented by Formulae 3A to 3F (for example, ligands represented by Formulae 3-1 to 3-15).


[0223] The organometallic compound may be one of Compounds 1 to 68:

-continued


5




1





2




3




4







7







8




-continued



-continued




-continued




18


-continued




19



20




21




22




-continued




-continued




26




27




28




30




31




32



33




-continued




34


-continued




38




35




39




36




40



37








41








42

-continued



-continued



43

44

45

46

47

48


49

50

51


52

-continued




53


-continued




58




54




59




55




60




56



61








57




62

-continued



-continued



63

68

64

[0224] Regarding the organometallic compound represented by Formula 1, ligand L_1 may be selected from a ligand represented by Formula 2A or 2B.

[0225] Nitrogen of a carboline-based ring in Formulae 2A and 2B is bound to M in Formula 1, and ring CY_1 in Formulae 2A and 2B is bound to M in Formula 1 via nitrogen. Accordingly, the organometallic compound represented by Formula 1 may have high luminescent efficiency.


65

[0226] For example, HOMO, LUMO, singlet (S_1), and triplet (T_1) energy levels of some of the organometallic compounds were evaluated by using a DFT method of Gaussian program (structurally optimized at a level of B3LYP, 6-31G(d,p)), and results thereof are shown in Table 1 below.

TABLE 1

Compound No.	HOMO (eV)	LUMO (eV)	T_1 energy level (eV)
3	-4.208	-1.705	1.860
4	-4.508	-1.769	2.140
9	-4.941	-1.580	2.593
10	-5.084	-1.842	2.551
19	-4.938	-1.949	2.225
49	-4.831	-1.828	2.366
66	-4.508	-1.547	2.203
68	-5.246	-2.028	2.545
Comparative Compound 1	-4.691	-1.298	2.528

66



67

Comparative Compound 1

[0227] Referring to Table 1, it can be seen that the organometallic compound represented by Formula 1 has electronic characteristics that are suitable for a material for an electronic device, for example, a dopant for an organic light-emitting device.

[0228] Synthesis methods of the organometallic compound represented by Formula 1 may be recognizable by one of ordinary skill in the art by referring to Synthesis Examples provided below.

[0229] The organometallic compound represented by Formula 1 is suitable for use in an organic layer of an organic light-emitting device, for example, for use as a dopant in an emission layer of the organic layer. Thus, another aspect provides an organic light-emitting device that includes:

[0230] a first electrode;

[0231] a second electrode; and

[0232] an organic layer that is disposed between the first electrode and the second electrode,

[0233] wherein the organic layer includes an emission layer and at least one organometallic compound represented by Formula 1.

[0234] The organic light-emitting device may have, due to the inclusion of an organic layer including the organometallic compound represented by Formula 1, excellent driving voltage, efficiency, power, electroluminescent (EL) emission wavelength, and color purity characteristics.

[0235] The organometallic compound represented by Formula 1 may be used between a pair of electrodes of an organic light-emitting device. For example, the organometallic compound represented by Formula 1 may be included in the emission layer. In this regard, the organometallic compound may act as a dopant, and the emission layer may further include a host (that is, an amount of the organometallic compound represented by Formula 1 is smaller than an amount of the host).

[0236] The expression that “(an organic layer) includes at least one of organometallic compounds” as used herein may include an embodiment in which “(an organic layer) includes identical organometallic compounds represented by Formula 1 and an embodiment in which (an organic layer) includes two or more different organometallic compounds represented by Formula 1.

[0237] For example, the organic layer may include only Compound 1 as the organometallic compound. In this regard, Compound 1 may be included only in the emission layer of the organic light-emitting device. In some embodiments, the organic layer may include, as the organometallic compound, Compound 1 and Compound 2. In this regard, Compound 1 and Compound 2 may be included in an identical layer (for example, Compound 1 and Compound 2 all may be included in an emission layer).

[0238] The first electrode may be an anode, which is a hole injection electrode, and the second electrode may be a cathode, which is an electron injection electrode; or the first electrode may be a cathode, which is an electron injection electrode, and the second electrode may be an anode, which is a hole injection electrode.

[0239] For example, in the organic light-emitting device, the first electrode may be an anode, the second electrode may be a cathode, and the organic layer may further include a hole transport region disposed between the first electrode and the emission layer and an electron transport region disposed between the emission layer and the second electrode, wherein the hole transport region may include at least one selected from a hole injection layer, a hole transport layer, and an electron blocking layer, and wherein the electron transport region may include at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

[0240] The term “organic layer” as used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of the organic light-emitting device. The “organic layer” may include, in addition to an organic compound, an organometallic complex including metal.

[0241] FIG. 1 is a schematic cross-sectional view of an organic light-emitting device 10 according to an embodiment. Hereinafter, the structure of an organic light-emitting device according to an embodiment and a method of manufacturing an organic light-emitting device according to an embodiment will be described in connection with FIG. 1. The organic light-emitting device 10 includes a first electrode 11, an organic layer 15, and a second electrode 19, which are sequentially stacked.

[0242] A substrate may be additionally disposed under the first electrode 11 or above the second electrode 19. For use as the substrate, any substrate that is used in general organic light-emitting devices may be used, and the substrate may be a glass substrate having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water-resistance, or a transparent plastic substrate.

[0243] The first electrode 11 may be formed by depositing or sputtering a material for forming the first electrode 11 on the substrate. The first electrode 11 may be an anode. The material for the first electrode 11 may be selected from materials with a high work function to facilitate hole injection. The first electrode 11 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. The material for the first electrode may be, for example, indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), and zinc oxide (ZnO). In some embodiments, magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag) may be used as the material for the first electrode.

[0244] The first electrode 11 may have a single-layer structure or a multi-layer structure including two or more layers. For example, the first electrode 11 may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode 110 is not limited thereto.

[0245] The organic layer 15 is disposed on the first electrode 11.

[0246] The organic layer 15 may include a hole transport region, an emission layer, and an electron transport region.

[0247] The hole transport region may be disposed between the first electrode 11 and the emission layer.

[0248] The hole transport region may include at least one selected from a hole injection layer, a hole transport layer, an electron blocking layer, and a buffer layer.

[0249] The hole transport region may include only either a hole injection layer or a hole transport layer. In some embodiments, the hole transport region may have a structure of hole injection layer/hole transport layer or hole injection layer/hole transport layer/electron blocking layer, which are sequentially stacked in this stated order from the first electrode 11.


[0250] A hole injection layer may be formed on the first electrode 11 by using one or more methods selected from vacuum deposition, spin coating, casting, or Langmuir-Blodgett (LB) deposition.

[0251] When a hole injection layer is formed by vacuum deposition, the deposition conditions may vary according to a material that is used to form the hole injection layer, and the structure and thermal characteristics of the hole injection layer. For example, the deposition conditions may include a deposition temperature of about 100 to about 500° C., a vacuum pressure of about 10^{-8} to about 10^{-3} torr, and a deposition rate of about 0.01 to about 100 Angstroms per second (Å/sec). However, the deposition conditions are not limited thereto.




[0252] When the hole injection layer is formed using spin coating, coating conditions may vary according to the material used to form the hole injection layer, and the structure and thermal properties of the hole injection layer. For example, a coating speed may be from about 2,000 revolutions per minute (rpm) to about 5,000 rpm, and a temperature at which a heat treatment is performed to remove a solvent after coating may be from about 80° C. to about 200° C. However, the coating conditions are not limited thereto.

[0253] Conditions for forming a hole transport layer and an electron blocking layer may be understood by referring to conditions for forming the hole injection layer.



[0254] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB, β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4', 4"-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzene sulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), (polyaniline)/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201 below, and a compound represented by Formula 202 below:




-continued




-continued

 β -NPB


TPD



Spiro-TPD




Spiro-NPB




methylated NPB

-continued




TAPC




HMTPD

Formula 201



Formula 202



[0255] Ar_{101} and Ar_{102} in Formula 201 may be each independently selected from

[0256] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluore-

nylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, and a pentacenylene group; and

[0257] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, and a pentacenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0258] In Formula 201, xa and xb may each independently be an integer selected from 0 to 5, or 0, 1, or 2. For example, xa may be 1 and xb may be 0, but they are not limited thereto.

[0259] R_{101} to R_{108} , R_{111} to R_{110} , and R_{121} to R_{124} in Formulae 201 and 202 may be each independently selected from

[0260] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{10} alkyl group (for example, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a hexyl group, and so on), or a C_1 - C_{10} alkoxy group (for example, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, and so on);

[0261] a C_1 - C_{10} alkyl group or a C_1 - C_{10} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, and a phosphoric acid group or a salt thereof;

[0262] a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, and a pyrenyl group; and

[0263] a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, and a pyrenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{10} alkyl group, and a C_1 - C_{10} alkoxy group, but they are not limited thereto.


[0264] R_{109} in Formula 201 may be

[0265] a phenyl group, a naphthyl group, an anthracenyl group, and a pyridinyl group; and

[0266] a phenyl group, a naphthyl group, an anthracenyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, and a pyridinyl group.

[0267] According to an embodiment, the compound represented by Formula 201 may be represented by Formula 201A, but is not limited thereto:


Formula 201A



[0268] R_{101} , R_{111} , R_{112} , and R_{109} in Formula 201A may be understood by referring to the description provided herein.


[0269] For example, the compound represented by Formula 201, and the compound represented by Formula 202 may include compounds HT1 to HT20 illustrated below, but are not limited thereto.

HT1




-continued

HT2




-continued


HT4



HT3




HT5




-continued

HT6




-continued


HT8




HT9



HT7




HT10




-continued


HT11



HT12




HT13




-continued


HT14




HT15



HT16




HT17




-continued


HT18



HT19



HT20




[0270] A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes a hole injection layer and a hole transport layer, the thickness of the hole injection layer may be in a range of about 100 Å to about 10,000 Å, and for example, about 100 Å to about 1,000 Å, and the thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, and for example, about 100 Å to about 1,500 Å. While not wishing to be bound by theory, it is understood that when the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0271] The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-


generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0272] The charge-generation material may be, for example, a p-dopant. The p-dopant may be one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but embodiments are not limited thereto. Non-limiting examples of the p-dopant are a quinone derivative, such as tetracyanoquinonodimethane (TCNQ) or 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonodimethane (F4-TCNQ); a metal oxide, such as a tungsten oxide or a molybdenum oxide; and a cyano group-containing compound, such as Compound HT-D1 below, but are not limited thereto.

Compound HT-D1



F4-TCNQ



[0273] The hole transport region may include a buffer layer.


[0274] Also, the buffer layer may compensate for an optical resonance distance according to a wavelength of light emitted from the emission layer, and thus, efficiency of a formed organic light-emitting device may be improved.

[0275] Then, an emission layer (EML) may be formed on the hole transport region by vacuum deposition, spin coating, casting, LB deposition, or the like. When the emission layer is formed by vacuum deposition or spin coating, the deposition or coating conditions may be similar to those applied to form the hole injection layer although the deposition or coating conditions may vary according to the material that is used to form the emission layer.


[0276] Meanwhile, when the hole transport region includes an electron blocking layer, a material for the electron blocking layer may be selected from materials for the hole transport region described above and materials for a host to be explained later. However, the material for the electron blocking layer is not limited thereto. For example, when the hole transport region includes an electron blocking layer, a material for the electron blocking layer may be mCP, which will be explained later.

[0277] The emission layer may include a host and a dopant, and the dopant may include the organometallic compound represented by Formula 1.


[0278] The host may include at least one selected from TPBi, TBADN, ADN (also referred to as “DNA”), CBP, CDBP, TCP, mCP, Compound H50, and Compound H51:




TPBi




TBADN



ADN




CBP




CDBP

-continued




TCP




mCP

Compound H50




Compound H51



[0279] In some embodiments, the host may further include a compound represented by Formula 301 below.

Formula 301



[0280] Ar_{111} and Ar_{112} in Formula 301 may each independently be selected from

[0281] a phenylene group, a naphthylene group, a phenanthrenylene group, and a pyrenylene group; and

[0282] a phenylene group, a naphthylene group, a phenanthrenylene group, and a pyrenylene group, each substituted with at least one selected from a phenyl group, a naphthyl group, and an anthracenyl group.

[0283] Ar_{113} to Ar_{116} in Formula 301 may each independently be selected from

[0284] a C_1 - C_{10} alkyl group, a phenyl group, a naphthyl group, a phenanthrenyl group, and a pyrenyl group; and

[0285] a phenyl group, a naphthyl group, a phenanthrenyl group, and a pyrenyl group, each substituted with at least one selected from a phenyl group, a naphthyl group, and an anthracenyl group.


[0286] g, h, i, and j in Formula 301 may be each independently an integer selected from 0 to 4, and may be, for example, 0, 1, or 2.

[0287] Ar_{113} to Ar_{116} in Formula 301 may be each independently selected from

[0288] a C_1 - C_{10} alkyl group, substituted with at least one selected from a phenyl group, a naphthyl group, and an anthracenyl group;


[0289] a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group;

[0290] a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group; and



[0291] but embodiments are not limited thereto.

[0292] In some embodiments, the host may include a compound represented by Formula 302 below:








[0293] Ar_{122} to Ar_{125} in Formula 302 are the same as described in detail in connection with Ar_{113} in Formula 301.


[0294] Ar_{126} and Ar_{127} in Formula 302 may be each independently a C_1 - C_{10} alkyl group (for example, a methyl group, an ethyl group, or a propyl group).

[0295] k and l in Formula 302 may be each independently an integer of 0 to 4. For example, k and l may be 0, 1, or 2.


[0296] The compound represented by Formula 301 and the compound represented by Formula 302 may include Compounds H1 to H42 illustrated below, but are not limited thereto.




-continued




H6




H7



H8




H9




H10


-continued




H11




H12




H13



H14







H15








H16

-continued




-continued




-continued


H26




H27




H28




H29



H30




H31




-continued


H32




H33




H34



H35




-continued




H36

-continued





H38

H39




H37




H40

-continued

H41



H42



[0297] When the organic light-emitting device is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, and a blue emission layer. In some embodiments, due to a stack structure including a red emission layer, a green emission layer, and/or a blue emission layer, the emission layer may emit white light.

[0298] When the emission layer includes a host and a dopant, an amount of the dopant may be in a range of about 0.01 to about 15 parts by weight based on 100 parts by weight of the host, but is not limited thereto.

[0299] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. While not wishing to be bound by theory, it is understood that when the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.


[0300] Then, an electron transport region may be disposed on the emission layer.

[0301] The electron transport region may include at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.


[0302] For example, the electron transport region may have a structure of hole blocking layer/electron transport layer/electron injection layer or a structure of electron transport layer/electron injection layer, but the structure of the electron transport region is not limited thereto. The electron transport layer may have a single-layered structure or a multi-layer structure including two or more different materials.

[0303] Conditions for forming the hole blocking layer, the electron transport layer, and the electron injection layer which constitute the electron transport region may be understood by referring to the conditions for forming the hole injection layer.

[0304] When the electron transport region includes a hole blocking layer, the hole blocking layer may include, for example, at least one of BCP, Bphen, and BA1q but is not limited thereto.




BCP




Bphen

[0305] A thickness of the hole blocking layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. While not wishing to be bound by theory, it is understood that when the thickness of the hole blocking layer is within these ranges, the hole blocking layer may have improved hole blocking ability without a substantial increase in driving voltage.


[0306] The electron transport layer may further include at least one selected from BCP, Bphen, Alq₃, BA1q, TAZ, and NTAZ.

Alq₃


-continued



BALq




TAZ



NTAZ


[0307] In some embodiments, the electron transport layer may include at least one of ET1 and ET2, but are not limited thereto:

ET1



-continued

ET2




[0308] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. While not wishing to be bound by theory, it is understood that when the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.


[0309] Also, the electron transport layer may further include, in addition to the materials described above, a metal-containing material.

[0310] The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.

ET-D1



ET-D2



[0311] The electron transport region may include an electron injection layer (EIL) that promotes flow of electrons from the second electrode 19 thereto.

[0312] The electron injection layer may include at least one selected from, LiF, NaCl, CsF, Li₂O, BaO, and LiQ.

[0313] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. While not wishing to be bound by theory, it is understood that when the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

[0314] The second electrode **19** is disposed on the organic layer **15**. The second electrode **19** may be a cathode. A material for forming the second electrode **19** may be selected from metal, an alloy, an electrically conductive compound, and a combination thereof, which have a relatively low work function. For example, lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag) may be used as a material for forming the second electrode **19**. In some embodiments, to manufacture a top emission type light-emitting device, a transmissive electrode formed using ITO or IZO may be used as the second electrode **19**.

[0315] Hereinbefore, the organic light-emitting device has been described with reference to FIG. 1, but is not limited thereto.

[0316] A C_1 - C_{60} alkyl group as used herein refers to a linear or branched aliphatic saturated hydrocarbon monovalent group having 1 to 60 carbon atoms. Examples thereof are a methyl group, an ethyl group, a propyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. A C_1 - C_{60} alkylene group as used herein refers to a divalent group having the same structure as the C_1 - C_{60} alkyl group.

[0317] A C_1 - C_{60} alkoxy group as used herein refers to a monovalent group represented by $-OA_{101}$ (wherein A_{101} is the C_1 - C_{60} alkyl group). Examples thereof are a methoxy group, an ethoxy group, and an iso-propyloxy (iso-propoxy) group.

[0318] A C_2 - C_{60} alkenyl group as used herein refers to a hydrocarbon group formed by including at least one carbon-carbon double bond in the middle or at the terminal of the C_2 - C_{60} alkyl group. Examples thereof are an ethenyl group, a propenyl group, and a butenyl group. A C_2 - C_{60} alkenylene group as used herein refers to a divalent group having the same structure as the C_2 - C_{60} alkenyl group.

[0319] A C_2 - C_{60} alkynyl group as used herein refers to a hydrocarbon group formed by including at least one carbon-carbon triple bond in the middle or at the terminal of the C_2 - C_{60} alkyl group. Examples thereof are an ethynyl group, and a propynyl group. A C_2 - C_{60} alkynylene group as used herein refers to a divalent group having the same structure as the C_2 - C_{60} alkynyl group.

[0320] A C_3 - C_{10} cycloalkyl group as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms. Examples thereof are a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. A C_3 - C_{10} cycloalkylene group as used herein refers to a divalent group having the same structure as the C_3 - C_{10} cycloalkyl group.

[0321] A C_1 - C_{10} heterocycloalkyl group as used herein refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, P, and S as a ring-forming atom and 1 to 10 carbon atoms. Examples thereof are a tetrahydrofuranlyl group, and a tetrahydrothiophenyl group.

A C_1 - C_{10} heterocycloalkylene group as used herein refers to a divalent group having the same structure as the C_1 - C_{10} heterocycloalkyl group.

[0322] A C_3 - C_{10} cycloalkenyl group as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof, and which is not aromatic. Examples thereof are a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. A C_3 - C_{10} cycloalkenylene group as used herein refers to a divalent group having the same structure as the C_3 - C_{10} cycloalkenyl group.

[0323] A C_1 - C_{10} heterocycloalkenyl group as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one double bond in its ring. Examples of the C_1 - C_{10} heterocycloalkenyl group are a 2,3-dihydrofuranlyl group and a 2,3-dihydrothiophenyl group. A C_1 - C_{10} heterocycloalkenylene group as used herein refers to a divalent group having the same structure as the C_1 - C_{10} heterocycloalkenyl group.

[0324] A C_6 - C_{60} aryl group as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C_6 - C_{60} arylene group as used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Examples of the C_6 - C_{60} aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C_6 - C_{60} aryl group and the C_6 - C_{60} arylene group each include two or more rings, the rings may be fused to each other.

[0325] A C_1 - C_{60} heteroaryl group as used herein refers to a monovalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, P, Si, and S as a ring-forming atom, and 1 to 60 carbon atoms. A C_1 - C_{60} heteroarylene group as used herein refers to a divalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, P, Si, and S as a ring-forming atom, and 1 to 60 carbon atoms. Examples of the C_1 - C_{60} heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C_1 - C_{60} heteroaryl group and the C_1 - C_{60} heteroarylene group each include two or more rings, the rings may be fused to each other.

[0326] A C_6 - C_{60} aryloxy group as used herein indicates $-OA_{102}$ (wherein A_{102} is the C_6 - C_{60} aryl group), and a C_6 - C_{60} arylthio group as used herein indicates $-SA_{103}$ (wherein A_{103} is the C_6 - C_{60} aryl group).

[0327] A monovalent non-aromatic condensed polycyclic group as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed to each other, only carbon atoms as a ring forming atom, and which is non-aromatic in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. A divalent non-aromatic condensed polycyclic group as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0328] A monovalent non-aromatic condensed heteropolycyclic group as used herein refers to a monovalent group (for example, having 2 to 60 carbon atoms) that has two or more rings condensed to each other, has a heteroatom selected from N, O, P, Si, and S, other than carbon atoms, as a ring forming atom, and which is non-aromatic in the entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. A divalent non-aromatic condensed heteropolycyclic

group as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0329] At least one of substituents of the substituted C_5 - C_{30} carbocyclic group, substituted C_2 - C_{30} heterocyclic group, substituted C_1 - C_{60} alkyl group, substituted C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group, substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10} cycloalkyl group, substituted C_1 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substituted C_1 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_1 - C_{60} heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group as used herein may be selected from:

[0330] deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0331] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_{11})(Q_{12})$, $-Si(Q_{13})(Q_{14})(Q_{15})$, $-B(Q_{16})(Q_{17})$, and $-P(=O)(Q_{18})(Q_{19})$;

[0332] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0333] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a

C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_{21})(Q_{22})$, $-Si(Q_{23})(Q_{24})(Q_{25})$, $-B(Q_{26})(Q_{27})$, and $-P(=O)(Q_{28})(Q_{29})$; and [0334] $-N(Q_{31})(Q_{32})$, $-Si(Q_{33})(Q_{34})(Q_{35})$, $-B(Q_{36})(Q_{37})$, and $-P(=O)(Q_{38})(Q_{39})$,

[0335] wherein Q_{11} to Q_{19} , Q_{21} to Q_{29} , and Q_{31} to Q_{39} are each independently selected from a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} alkyl group, and a C_6 - C_{60} aryl group.

[0336] When a group containing a specified number of carbon atoms is substituted with any of the groups listed in the preceding paragraphs, the number of carbon atoms in the resulting "substituted" group is defined as the sum of the carbon atoms contained in the original (unsubstituted) group and the carbon atoms (if any) contained in the substituent. For example, when the term "substituted C_1 - C_{60} alkyl" refers to a C_1 - C_{60} alkyl group substituted with C_6 - C_{60} aryl group, the total number of carbon atoms in the resulting aryl substituted alkyl group is C_7 - C_{120} .

[0337] Hereinafter, a compound and an organic light-emitting device according to embodiments are described in detail with reference to Synthesis Example and Examples. However, the organic light-emitting device is not limited thereto. The wording "B was used instead of A" used in describing Synthesis Examples means that an amount of A in molar equivalents was identical to an amount of B in molar equivalents.


EXAMPLE

Synthesis Example 1


Synthesis of Compound 9

[0338]


Synthesis of Compound M1A



Compound A



[0339] 5.08 grams (g) (14.83 millimoles, mmol) of Compound A and 2.32 g (6.59 mmol) of iridium chloride were mixed with 60 milliliters (mL) of ethoxyethanol and 20 mL of distilled water. The resulting mixture was stirred under reflux for about 24 hours until the reaction was complete, and then the temperature was reduced to room temperature. A solid formed therefrom was separated by filtration. The solid was thoroughly washed with water, methanol, and hexane in the stated order, and dried in a vacuum oven, to obtain 3.8 g (63%) of Compound M1A.



[0340] 2.80 g (1.54 mmol) of Compound M1A, 1.54 g (15.40 mmol) of acetylacetone, and 2.13 g (15.40 mmol) of K_2CO_3 were mixed with 30 mL of ethoxyethanol. The


resulting mixture was stirred under reflux for about 12 hours until the reaction was complete, and then the temperature was reduced. The cooled reaction mixture was filtered to obtain a solid, which was then thoroughly washed with ethanol and hexane. The solid was purified by column chromatography using methylene chloride (MC) and hexane to obtain 0.35 g (12%) of Compound 9. The obtained compound was identified by using mass spectroscopy and high-performance liquid chromatography (HPLC) analysis. **[0341]** HRMS (MALDI) calcd for $\text{C}_{49}\text{H}_{49}\text{IrN}_8\text{O}_2$: m/z 974.3608, Found: 974.3601

Synthesis Example 2

Synthesis of Compound 19

[0342]

Synthesis of Compound M1B



[0343] 5.24 g (13.36 mmol) of Compound B and 2.09 g (5.94 mmol) of iridium chloride were mixed with 60 mL of ethoxyethanol and 20 mL of distilled water. The resulting mixture was stirred under reflux for about 24 hours until the




reaction was complete, and then the temperature was reduced to room temperature. A solid formed therefrom was separated by filtration. The solid was thoroughly washed with water, methanol, and hexane in the stated order, and dried in a vacuum oven, to obtain 4.5 g (75%) of Compound M1 B.

[0345] HRMS (MALDI) calcd for $C_{57}H_{53}IrN_8O_2$: m/z 1074.3921, Found: 1074.3926


Comparative Synthesis Example

Synthesis of Comparative Compound 1


[0346]




[0344] 2.82 g (1.40 mmol) of Compound M1B, 1.40 g (13.96 mmol) of acetylacetone, and 1.93 g (13.96 mmol) of K_2CO_3 were mixed with 30 mL of ethoxyethanol. The resulting mixture was stirred under reflux for about 12 hours until the reaction was complete, and then the temperature was reduced. The cooled mixture was then filtered to obtain a solid, which was thoroughly washed with ethanol and hexane. The solid was purified by column chromatography using MC and hexane to obtain 0.72 g (24%) of Compound 19. The obtained compound was identified by using mass spectroscopy and HPLC analysis.



[0347] 7.70 g (31.50 mmol) of Compound C and 4.94 g (14.00 mmol) of iridium chloride were mixed with 90 mL of ethoxyethanol and 30 mL of distilled water. The resulting mixture was stirred under reflux for about 24 hours until the reaction was complete, and then the temperature was reduced to room temperature. A solid formed therefrom was separated by filtration. The washed solid was thoroughly washed with water, methanol, and hexane in the stated order, and dried in a vacuum oven, to obtain 8.6 g (86%) of Compound M1C.





[0348] 4.13 g (2.89 mmol) of Compound M1C, 2.90 g (28.92 mmol) of acetylacetone, and 4.00 g (28.92 mmol) of K_2CO_3 were mixed with 50 mL of ethoxyethanol. The resulting mixture was stirred under reflux for about 12 hours until the reaction was complete. The formed solid was filtered and thoroughly washed with ethanol and hexane. Then, the solid was purified by column chromatography using MC and hexane to obtain 0.95 g (21%) of Comparative Compound 1. The obtained compound was identified by using mass spectroscopy and HPLC analysis.

[0349] HRMS (MALDI) calcd for $C_{39}H_{29}IrN_4O_2$: m/z 778.1920, Found: 778.1914

Example 1

[0350] An ITO glass substrate was cut to a size of 50 millimeters (mm)×50 mm×0.5 mm. Then the glass substrate was sonicated in acetone, iso-propyl alcohol, and pure water for about 15 minutes in each solvent, and cleaned by exposure to ultraviolet rays with ozone for 30 minutes.

[0351] Thereafter, m-MTDATA was deposited on an ITO electrode (anode) of the glass substrate at a deposition rate of about 1 Angstroms per second ($\text{\AA}/\text{sec}$) to form a hole injection layer having a thickness of about 600 Angstroms ($\text{\AA}$) and α -NPD was deposited on the hole injection layer at

a deposition rate of about 1 $\text{\AA}/\text{sec}$ to form a hole transport layer having a thickness of about 250 $\text{\AA}$.

[0352] Compound 9 (as a dopant) and CBP (as a host) were co-deposited on the hole transport layer at a deposition rate of about 0.1 $\text{\AA}/\text{sec}$ and 1 $\text{\AA}/\text{sec}$, respectively to form an emission layer having a thickness of about 400 $\text{\AA}$.

[0353] BAQ was deposited on the emission layer at a deposition rate of 1 $\text{\AA}/\text{sec}$ to form a hole blocking layer having a thickness of about 50 $\text{\AA}$. Then, Alq_3 was deposited on the hole blocking layer to form an electron transport layer having a thickness of about 300 $\text{\AA}$, and LiF was deposited on the electron transport layer to form an electron injection layer having a thickness of about 10 $\text{\AA}$. A second electrode (cathode) was formed on the electron injection layer by depositing Al to have a thickness of about 1,200 $\text{\AA}$. Accordingly, the manufacture of an organic light-emitting device was completed, in which the organic light-emitting device included a structure of ITO/m-MTDATA (600 $\text{\AA}$)/ α -NPD (250 $\text{\AA}$)/CBP+10% (Compound 9) (400 $\text{\AA}$)/BAQ (50 $\text{\AA}$)/ Alq_3 (300 $\text{\AA}$)/LiF (10 $\text{\AA}$)/Al (1,200 $\text{\AA}$).

Example 2 and Comparative Example 1

[0354] An organic light-emitting device was manufactured in the same manner as in Example 1, except that the compounds listed in Table 2 were used instead of Compound 9 as a dopant in the formation of the emission layer.


Evaluation Example 1

Evaluation of Characteristics of Organic Light-Emitting Device

[0355] Driving voltage, efficiency, power, EL emission wavelength, and color-coordination of the organic light-emitting devices manufactured in Examples 1 and 2 and Comparative Example 1 were evaluated at a selected luminance of about 1,000 candelas per square meter (cd/m^2). The results thereof are shown in Table 2. A Keithley 2400 current voltmeter and a luminance meter (Minolta Cs-1000A) were used in evaluation.



TABLE 2

	Dopant	Driving	Efficiency (cd/A)	Power (lm/W)	λ_{max}	CIE _x	CIE _y
		voltage (V)					
Example 1	Compound 9	7.0	10	4.5	525	0.28	0.69
Example 2	Compound 19	7.3	7.5	3.2	601.8	0.61	0.38
Comparative Example 1	Comparative Compound 1	8.0	6.0	2.4	535	0.40	0.58



Compound 9

TABLE 2-continued

Dopant	Driving voltage (V)	Efficiency (cd/A)	Power (lm/W)	λ_{max}	CIE _x	CIE _y
						
Compound 19						
						
Comparative Compound 1						

[0356] Referring to Table 2, it was found that the organic light-emitting devices manufactured in Examples 1 and 2 had excellent driving voltage, efficiency, power, EL emission wavelength, and color-coordination, as compared with the organic light-emitting device of Comparative Example 1.


[0357] The organometallic compound according to embodiments has excellent electric characteristics and thermal stability. Accordingly, an organic light-emitting device including the organometallic compound may have excellent driving voltage, current density, efficiency, power, color purity, and lifespan characteristics.

[0358] It should be understood that exemplary embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each exemplary embodiment should typically be considered as available for other similar features or aspects in other exemplary embodiments.

[0359] While one or more exemplary embodiments have been described with reference to the figures, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope of the present disclosure as defined by the following claims.


What is claimed is:

1. An organometallic compound represented by Formula 1:




Formula 1

Formula 2A



Formula 2B



wherein

M in Formula 1 is selected from iridium (Ir), platinum (Pt), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), thulium (Tm), and rhodium (Rh),

L₁ in Formula 1 is selected from a ligand represented by Formula 2A or 2B,

L₂ in Formula 1 is selected from a mono-anionic organic ligand,

* and *[†] in Formulae 2A and 2B each indicate a binding site to M in Formula 1,

n₁ in Formula 1 is 1, 2, or 3, and when n₁ is 2 or more, 2 or more groups L₁ are identical to or different from each other,

n₂ in Formula 1 is 0, 1, 2, or 3, and when n₂ is 2 or more, 2 or more groups L₂ are identical to or different from each other,

Y₁ and Y₃ in Formulae 2A and 2B are each independently carbon or nitrogen,

Y₂ in Formulae 2A and 2B is carbon,

a bond between Y₂ and Y₃ in Formula 2B is a double bond,

CY₁ in Formulae 2A and 2B is a C₂-C₃₀ heterocyclic group,

in Formulae 2A and 2B, X₁ is N or C(R₁), X₂ is N or C(R₂), X₃ is N or C(R₃), X₄ is N or C(R₄), X₅ is N or C(R₅), X₆ is N or C(R₆), X₇ is N or C(R₇),

T₁ in Formula 2B is selected from a single bond, *—O*, *—S—*, *—N(R₃₁)—*, *—C(R₃₁)(R₃₂)—*, *—C(R₃₁)=C(R₃₂)—', and *—Si(R₃₁)(R₃₂)—', and R₃₁ and R₃₂ are optionally bound to each other to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group or a substituted or unsubstituted C₂-C₃₀ heterocyclic group,

R₁ to R₇, R₁₀, R₃₁, and R₃₂ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, —SF₅, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), and —P(=O)(Q₈)(Q₉),

a₁ is an integer selected from 0 to 5,

two or more among R₁ to R₃ in Formulae 2A and 2B are optionally bound to each other to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group or a substituted or unsubstituted C₂-C₃₀ heterocyclic group,

two or more among R₄ to R₇ in Formula 2A are optionally bound to each other to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group or a substituted or unsubstituted C₂-C₃₀ heterocyclic group,

two or more among R₄ to R₆ in Formula 2B are optionally bound to each other to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group or a substituted or unsubstituted C₂-C₃₀ heterocyclic group,

two or more among groups R₁₀ in the number of a₁ in Formulae 2A and 2B are optionally bound to each other to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group or a substituted or unsubstituted C₂-C₃₀ heterocyclic group, and

at least one of substituents of the substituted C₅-C₃₀ carbocyclic group, substituted C₂-C₃₀ heterocyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), —B(Q₁₆)(Q₁₇), and —P(=O)(Q₁₈)(Q₁₉);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

eroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅) —B(Q₂₆)(Q₂₇), and —P(=O)(Q₂₈)(Q₂₉); and

—N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅) —B(Q₃₆)(Q₃₇) and —P(=O)(Q₃₈)(Q₃₉),

wherein Q₁ to Q₆, Q₁₁ to Q₁₉, Q₂₁ to Q₂₉, and Q₃₁ to Q₃₉ are each independently selected from a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ alkyl group, and a C₆-C₆₀ aryl group.

2. The organometallic compound of claim 1, wherein in Formula 1,

M is Ir or Os, and the sum of n₁ and n₂ is 3 or 4; or

M is Pt, and the sum of n₁ and n₂ is 2.

3. The organometallic compound of claim 1, wherein

CY₁ in Formulae 2A and 2B is selected from a pyrrole ring, an isoindole ring, an indole ring, an indazole ring, a pyrazole ring, an imidazole ring, a triazole ring, an oxazole ring, an isoxazole ring, an oxadiazole ring, a thiazole ring, an isothiazole ring, a thiadiazole ring, a purine ring, a benzimidazole ring, an isobenzothiazole ring, a benzoxazole ring, an isobenzoxazole ring, an imidazopyridine ring, an imidazopyrimidine ring, and a pyrrolopyridine ring.

4. The organometallic compound of claim 1, wherein

T₁ in Formula 2B is selected from *—O—*, *—S—*, *—C(R₃₁)(R₃₂)—*, and groups represented by Formulae 11-1 to 11-4 below:




Formula 11-1



Formula 11-2



Formula 11-3



Formula 11-4

wherein * and *' in Formulae 11-1 to 11-4 each indicate a binding site to a neighboring atom.

5. The organometallic compound of claim 1, wherein

R₁ to R₇, R₁₀, R₃₁, and R₃₂ are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, —SF₅, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl

group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a carbazolyl group, a phenanthroliny group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a carbazolyl group, a phenanthroliny group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a carbazolyl group, a phenanthroliny group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a diben-

zofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

—N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), and —P(=O)(Q₈)(Q₉),

wherein Q₁ to Q₉ are each independently selected from:

—CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCH₂, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C₁-C₁₀ alkyl group, and a phenyl group.

6. The organometallic compound of claim 1, wherein

R₁ to R₇, R₁₀, R₃₁, and R₃₂ are each independently selected from:

hydrogen, deuterium, —F, a cyano group, a nitro group, —SF₅, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group and a dibenzothiophenyl group; and

a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl

group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, a cyano group, a nitro group, a $\text{C}_1\text{-C}_{10}$ alkyl group, a $\text{C}_1\text{-C}_{10}$ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

$-\text{N}(\text{Q}_1)(\text{Q}_2)$, $-\text{Si}(\text{Q}_3)(\text{Q}_4)(\text{Q}_5)$, $-\text{B}(\text{Q}_6)(\text{Q}_7)$, and $-\text{P}(=\text{O})(\text{Q}_8)(\text{Q}_9)$;

wherein Q_1 to Q_9 are each independently selected from:

$-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCCH}_3$, $-\text{CHDCD}_2\text{H}$, $-\text{CHDCDH}_2$, $-\text{CHDCD}_3$, $-\text{CD}_2\text{CD}_3$, $-\text{CD}_2\text{CD}_2\text{H}$, and $-\text{CD}_2\text{CDH}_2$;

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a $\text{C}_1\text{-C}_{10}$ alkyl group, and a phenyl group.

7. The organometallic compound of claim 1, wherein


R_1 to R_7 , R_{10} , R_{31} , and R_{32} are each independently selected from hydrogen, deuterium, $-\text{F}$, a cyano group, a nitro group, $-\text{SF}_5$, $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, groups represented by Formulae 9-1 to 9-19, groups represented by Formulae 10-1 to 10-36, and $-\text{Si}(\text{Q}_3)(\text{Q}_4)(\text{Q}_5)$,

Q_3 to Q_5 are each independently selected from:


$-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCCH}_3$, $-\text{CHDCD}_2\text{H}$, $-\text{CHDCDH}_2$, $-\text{CHDCD}_3$, $-\text{CD}_2\text{CD}_3$, $-\text{CD}_2\text{CD}_2\text{H}$, and $-\text{CD}_2\text{CDH}_2$;

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a $\text{C}_1\text{-C}_{10}$ alkyl group, and a phenyl group:



Formula 9-1



Formula 9-2


-continued




Formula 9-3




Formula 9-4




Formula 9-5




Formula 9-6




Formula 9-7




Formula 9-8




Formula 9-9




Formula 9-10




Formula 9-11



Formula 9-12




Formula 9-13




Formula 9-14




Formula 9-15



Formula 9-16















Formula 9-17



Formula 9-18

-continued



Formula 9-19

Formula 10-1

Formula 10-2

Formula 10-3

Formula 10-4

Formula 10-5

Formula 10-6

Formula 10-7











Formula 10-8

Formula 10-9

Formula 10-10

Formula 10-11

-continued



Formula 10-12

Formula 10-13

Formula 10-14

Formula 10-15

Formula 10-16

Formula 10-17











Formula 10-18

Formula 10-19

Formula 10-20

Formula 10-21

-continued



Formula 10-22

Formula 10-23

Formula 10-24

Formula 10-25

Formula 10-26

Formula 10-27






Formula 10-28

Formula 10-29

Formula 10-30


Formula 10-31

-continued



wherein, * in Formulae 9-1 to 9-19 and 10-1 to 10-36 indicates a binding site to a neighboring atom.

8. The organometallic compound of claim 1, wherein ligand L_1 in Formula 1 is selected from groups represented by Formulae 2A-1 to 2A-4 and 2B-1 to 2B-3:



Formula 10-32

Formula 10-33


Formula 10-34

Formula 10-35


Formula 10-36

Formula 2A-1


-continued




Formula 2A-2




Formula 2A-3



Formula 2A-4




Formula 2B-1



Formula 2B-2

-continued



Formula 2B-3

wherein, in Formulae 2A-1 to 2A-4 and 2B-1 to 2B-3, X_1 to X_7 , and T_1 are the same as described in claim 1, X_{11} is N or C(R_{11}), X_{12} is N or C(R_{12}), X_{13} is N or C(R_{13}),

X_{14} is N or C(R_{14}), X_{15} is N or C(R_{15}),

X_{21} is O, S, N(R_{21}), or C(R_{22})(R_{23}),

X_{21} is O, S, N(R_{24}), or C(R_{25})(R_{26}),

R_{11} to R_{15} and R_{21} to R_{26} are the same as described in connection with R_{10} in claim 1, and

* and *' each indicate a binding site to M in Formula 1.

9. The organometallic compound of claim 8, wherein

X_1 is C(R_1), X_2 is C(R_2), X_3 is C(R_3), X_4 is C(R_4), X_5 is C(R_5), X_6 is C(R_6), X_7 is C(R_7),

T_1 is selected from $*-O-*'$, $*-S-*'$, $*-C(R_{31})(R_{32})-*'$, and groups represented by Formulae 11-1 to 11-4,

R_1 to R_7 , R_{11} to R_{15} , R_{21} to R_{26} , R_{31} , and R_{32} are each independently selected from:

hydrogen, deuterium, —F, a cyano group, a nitro group, —SF₅, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a

tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

—N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), and —P(=O)(Q₈)(Q₉),

wherein Q₁ to Q₉ are each independently selected from:

—CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C₁-C₁₀ alkyl group, and a phenyl group,

two neighboring substituents among R₁ to R₃ are optionally bound to each other to form a substituted or unsubstituted C₅-C₁₀ carbocyclic ring,

two neighboring substituents among R₄ to R₇ are optionally bound to each other to form a substituted or unsubstituted C₅-C₁₀ carbocyclic ring, and

two neighboring substituents among R₁₁ to R₁₅ and R₂₁ to R₂₆ are optionally bound to each other to form a substituted or unsubstituted C₅-C₁₀ carbocyclic ring;



Formula 11-1




Formula 11-2



Formula 11-3

-continued

Formula 11-4




wherein * and *' in Formulae 11-1 to 11-4 each indicate a binding site to a neighboring atom.


10. The organometallic compound of claim 8, wherein X₅ is C(CN) or X₁₂ is C(CN).

11. The organometallic compound of claim 1, wherein ligand L₁ in Formula 1 is selected from ligands represented by Formulae 2A-1(1) to 2A-1(6):


Formula 2A-1(1)



Formula 2A-1(2)




Formula 2A-1(3)




-continued


Formula 2A-1(4)



Formula 2A-1(5)



Formula 2A-1(6)



wherein, in Formulae 2A-1(1) to 2A-1(6),
 X_1 is N or C(R₁), X_2 is N or C(R₂), X_3 is N or C(R₃), X_4
 is N or C(R₄), X_5 is N or C(R₅), X_6 is N or C(R₆), X_7
 is N or C(R₇),

T₁ is selected from *—O—*, *—S—*, *—C(R₃₁)
(R₃₂)—*, and groups represented by Formulae 11-1 to
11-4.

X_{11} is $\mathbb{N}$ or $C(\mathbb{R}_{11})$,

R_1 to R_7 , R_{11} , R_{31} , R_{32} , and R_{41} to R_{48} are each independently selected from:

hydrogen, deuterium, —F, a cyano group, a nitro group, —SF₅, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group,

an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranlyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranlyl group, and a dibenzothiophenyl group; and

$$-\text{N}(\text{Q}_1)(\text{Q}_2), \quad -\text{Si}(\text{Q}_3)(\text{Q}_4)(\text{Q}_5), \quad -\text{B}(\text{Q}_6)(\text{Q}_7), \quad \text{and} \\ -\text{P}(=\text{O})(\text{Q}_8)(\text{Q}_9),$$

wherein Q_1 to Q_8 are each independently selected from:

$$\begin{array}{ccccccc} -\text{CH}_3, & -\text{CD}_3, & -\text{CD}_2\text{H}, & -\text{CDH}_2, & -\text{CH}_2\text{CH}_3, \\ -\text{CH}_2\text{CD}_3, & -\text{CH}_2\text{CD}_2\text{H}, & -\text{CH}_2\text{CDH}_2, & -\text{CHDCH}_3, \\ -\text{CHDCH}_2\text{H}, & -\text{CHDCH}_2\text{H}_2, & -\text{CHDCH}_3, \\ -\text{CD}_2\text{CD}_3, & -\text{CD}_2\text{CD}_2\text{H}, & \text{and} & -\text{CD}_2\text{CDH}_2; \end{array}$$

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C₁-C₁₀ alkyl group, and a phenyl group.

two neighboring substituents among R₁ to R₃ are optionally bound to each other to form a substituted or unsubstituted C₅-C₁₀ carbocyclic ring,

two neighboring substituents among R_4 to R_7 are optionally bound to each other to form a substituted or unsubstituted C_5 - C_{10} carbocyclic ring, and

* and *' each indicate a binding site to M in Formula 1:




Formula 11-1



Formula 11-2




Formula 11-3



Formula 11-4

wherein * and *' in Formulae 11-1 to 11-4 each indicate a binding site to a neighboring atom.


12. The organometallic compound of claim 1, wherein ligand L_1 in Formula 1 is selected from ligands represented by Formulae 4-1 to 4-7, 4-11 to 4-17, 4-21 to 4-27, 4-31 to 4-37, 4-41 to 4-44, 4-51 to 4-54, 4-61 to 4-64, and 4-71 to 4-74:




Formula 4-1

-continued


Formula 4-2




Formula 4-3




Formula 4-4



Formula 4-5




-continued




Formula 4-6

-continued




Formula 4-13


Formula 4-7




Formula 4-11



Formula 4-12




Formula 4-14




Formula 4-15

-continued




Formula 4-16

-continued




Formula 4-23


Formula 4-17




Formula 4-21




Formula 4-24




Formula 4-22



Formula 4-25




Formula 4-26




-continued


Formula 4-34




Formula 4-35




Formula 4-36



Formula 4-37




-continued




Formula 4-41

-continued




Formula 4-51




Formula 4-42

Formula 4-52




Formula 4-43




Formula 4-44

Formula 4-53




-continued




Formula 4-54


-continued




Formula 4-64




Formula 4-61




Formula 4-71




Formula 4-62




Formula 4-72



Formula 4-63



Formula 4-73



Formula 4-74

wherein, in Formulae 4-1 to 4-7, 4-11 to 4-17, 4-21 to 4-27, 4-31 to 4-37, 4-41 to 4-44, 4-51 to 4-54, 4-61 to 4-64, and 4-71 to 4-74,

X_{11} is N or C(R_{11}),

X_{14} is N or C(R_{14}),

T_1 is the same as described in claim 1,

R_1 to R_9 , R_{11} to R_{15} and R_{41} to R_{48} are each independently selected from hydrogen, deuterium, —F, a cyano group, a nitro group, —SF₅, —CH₃, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, groups represented by Formulae 9-1 to 9-19, groups represented by Formulae 10-1 to 10-36, and —Si(Q_3)(Q_4)(Q_5),

Q_3 to Q_5 are each independently selected from:

—CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;


an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C₁-C₁₀ alkyl group, and a phenyl group, and

* and *' each indicate a binding site to M in Formula 1:



Formula 9-1



Formula 9-2




Formula 9-3




Formula 9-4


-continued




Formula 9-5




Formula 9-6




Formula 9-7




Formula 9-8




Formula 9-9




Formula 9-10




Formula 9-11




Formula 9-12




Formula 9-13




Formula 9-14




Formula 9-15





Formula 9-16





Formula 9-17




Formula 9-18















Formula 9-19




Formula 10-1




-continued




Formula 10-2




Formula 10-3




Formula 10-4



Formula 10-5




Formula 10-6




Formula 10-7

Formula 10-8




Formula 10-9




Formula 10-10


Formula 10-11



Formula 10-12



Formula 10-13



-continued

Formula 10-14

Formula 10-15

Formula 10-16

Formula 10-17

Formula 10-18

Formula 10-19










Formula 10-20

Formula 10-21

Formula 10-22

Formula 10-23

-continued



Formula 10-24

Formula 10-25

Formula 10-26

Formula 10-27

Formula 10-28





Formula 10-29

Formula 10-30

Formula 10-31





Formula 10-32

-continued



wherein * in Formulae 9-1 to 9-19 and 10-1 to 10-36 indicates a binding site to a neighboring atom.

13. The organometallic compound of claim 1, wherein ligand L_2 in Formula 1 is selected from ligands represented by Formula 3A to 3F:



Formula 10-33

Formula 10-34

Formula 10-35

Formula 10-36


Formula 3A

Formula 3B


Formula 3C

Formula 3D

-continued



Formula 3E



Formula 3F

wherein, in Formulae 3A to 3F

Y_{11} is selected from O, N, $N(Z_1)$, $P(Z_1)(Z_2)$, and $As(Z_1)(Z_2)$,

Y_{12} is selected from O, N, $N(Z_3)$, $P(Z_3)(Z_4)$, and $As(Z_3)(Z_4)$,

CY_{11} is a C_2 - C_{30} heterocyclic group,

T_{11} are each independently selected from a single bond, a double bond, $*—C(Z_{11})(Z_{12})—*$, $*—C(Z_{11})=C(Z_{12})—*$, $*=C(Z_{11})—*$, $*—C(Z_{11})=*$, $*=C(Z_{11})—C(Z_{12})=C(Z_{13})—*$, $*—C(Z_{11})=C(Z_{12})—C(Z_{13})=*$, $*—N(Z_{11})—*$, and a substituted or unsubstituted C_6 - C_{30} arylene group,

a_{11} is an integer selected from 1 to 10,

Y_{13} to Y_{16} are each independently carbon (C) or nitrogen (N), Y_{13} and Y_{14} are bound to each other via a single bond or a double bond, Y_{15} and Y_{16} are bound to each other via a single bond or a double bond,

CY_{12} and CY_{13} are each independently selected from a C_5 - C_{60} cyclic group and a C_2 - C_{60} heterocyclic group,

A_7 is P or As,

Z_1 to Z_4 and Z_{11} to Z_{13} are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, — $N(Q_1)(Q_2)$, — $Si(Q_3)(Q_4)(Q_5)$, — $B(Q_6)(Q_7)$, and — $P(=O)(Q_8)(Q_9)$,

d_1 and d_2 are each independently an integer selected from 0 to 10, and

* and *' each indicate a binding site to M in Formula 1.

14. The organometallic compound of claim 1, wherein CY_{11} is selected from a pyridine ring, a pyrimidine ring, a triazine ring, a pyrrole ring, a pyrazole ring, an imidazole ring, and a triazole ring,

CY_{12} and CY_{13} are each independently selected from a benzene ring, a naphthalene ring, a pyridine ring, a pyrimidine ring, a triazine ring, a pyrrole ring, a pyrazole ring, an imidazole ring, and a triazole ring, and

Z_1 to Z_4 and Z_{11} to Z_{13} are each independently selected from:

hydrogen, deuterium, —F, a cyano group, a nitro group, — SF_5 , a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, — CD_3 , — CD_2H , — CDH_2 , — CF_3 , — CF_2H , — CFH_2 , a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

— $N(Q_1)(Q_2)$, — $Si(Q_3)(Q_4)(Q_5)$, — $B(Q_6)(Q_7)$ and — $P(=O)(Q_8)(Q_9)$,

Q_1 to Q_9 are each independently selected from:


— CH_3 , — CD_3 , — CD_2H , — CDH_2 , — CH_2CH_3 , — CH_2CD_3 , — CH_2CD_2H , — CH_2CDH_2 , — $CHDCH_3$, — $CHDCD_2H$, — $CHDCDH_2$, — $CHDCD_3$, — CD_2CD_3 , — CD_2CD_2H , and — CD_2CDH_2 ;

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl


group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C₁-C₁₀ alkyl group, and a phenyl group.


15. The organometallic compound of claim 1, wherein ligand L₂ in Formula 1 is selected from ligands represented by Formulae 3-1 to 3-15:




Formula 3-1




Formula 3-2




Formula 3-3



Formula 3-4




Formula 3-5




Formula 3-6


-continued




Formula 3-7




Formula 3-8




Formula 3-9




Formula 3-10




Formula 3-11



Formula 3-12




Formula 3-13



Formula 3-14

-continued

Formula 3-15



wherein, in Formulae 3-1 to 3-15,

X_{31} is N or C(Z_{1a}), and X_{32} is N or C(Z_{1b}),

X_{41} is O, S, N(Z_{1a}), or C(Z_{1a})(Z_{1b}),

Z_1 to Z_4 , Z_{1a} , Z_{1b} , and Z_{11} to Z_{14} are each independently selected from:

hydrogen, deuterium, —F, a cyano group, a nitro group, —SF₅, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group and a dibenzothiophenyl group;

a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

—N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), and —P(=O)(Q₈)(Q₉);

Q₁ to Q₉ are each independently selected from:

—CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from deuterium, a C₁-C₁₀ alkyl group, and a phenyl group,




e2 is an integer selected from 0 to 2,

e3 is an integer selected from 0 to 3,





e4 is an integer selected from 0 to 4, and

* and *' each indicate a binding site to M in Formula 1.





16. The organometallic compound of claim 1, wherein the organometallic compound is one of Compounds 1 to 68:







-continued







-continued




-continued










-continued







-continued








-continued








-continued








-continued




-continued



-continued












-continued




-continued


47




-continued




57




58



59




60




61


-continued




62




63



64





65



66

-continued



67

17. An organic light-emitting device comprising:

a first electrode;

a second electrode; and

an organic layer disposed between the first electrode and the second electrode,

wherein the organic layer comprises an emission layer and at least one organometallic compound of claim 1.

18. The organic light-emitting device of claim 17, wherein the first electrode is an anode,

the second electrode is a cathode,

the organic layer comprises a hole transport region disposed between the first electrode and the emission layer and an electron transport region disposed between the emission layer and the second electrode,

68

wherein the hole transport region comprises a hole injection layer, a hole transport layer, a buffer layer, an intermediate layer, an electron blocking layer, or any combination thereof and

wherein the electron transport region comprises a hole blocking layer, an electron transport layer, an electron injection layer, or any combination thereof.

19. The organic light-emitting device of claim 17, wherein the emission layer comprises the organometallic compound.


20. The organic light-emitting device of claim 19, wherein the emission layer further comprises a host.

* * * * *


专利名称(译)	有机金属化合物和包括其的有机发光器件		
公开(公告)号	US20170084850A1	公开(公告)日	2017-03-23
申请号	US15/263850	申请日	2016-09-13
[标]申请(专利权)人(译)	三星电子株式会社		
申请(专利权)人(译)	SAMSUNG ELECTRONICS CO. , LTD.		
当前申请(专利权)人(译)	SAMSUNG ELECTRONICS CO. , LTD.		
[标]发明人	HWANG KYUYOUNG KWON OHYUN KWAK SEUNGYEON LEE KUM HEE JEON ARAM CHOI HYEONHO CHOI BYOUNGKI		
发明人	HWANG, KYUYOUNG KWON, OHYUN KWAK, SEUNGYEON LEE, KUM HEE JEON, ARAM CHOI, HYEONHO CHOI, BYOUNGKI		
IPC分类号	H01L51/00 C09K11/02 C07F15/00 C09K11/06		
CPC分类号	H01L51/0085 C09K2211/185 C09K11/025 C07F15/004 H01L51/5206 H01L51/5221 H01L51/5024 H01L51/5056 H01L51/5072 H01L51/5088 H01L51/5008 H01L51/5096 H01L51/5092 H01L51/56 C09K2211/1044 C09K11/06 C07F15/0026 C07F15/0033 H01L51/5016		
优先权	1020150134134 2015-09-22 KR		
其他公开文献	US10164200		
外部链接	Espacenet USPTO		

摘要(译)

由式1表示的有机金属化合物： $M(L_1)_{n1}(L_2)_{n2}$ 公式1 其中在式1中，式1中的M选自铱(Ir)，铂(Pt)，锇(Os)，钛(Ti)，锆(Zr)，铪(Hf)，镱(Eu)，铽(Tb)，钍(Tm)和铑(Rh)，L₁选自式2A或2B表示的配体，因为它们在说明书中定义，并且式1中的L₂选自单阴离子有机物配位体。



Formula 2A



Formula 2B