



US010069085B2

(12) **United States Patent**
Lee et al.

(10) **Patent No.: US 10,069,085 B2**
 (45) **Date of Patent: Sep. 4, 2018**

(54) **HETEROCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

(71) Applicant: **SAMSUNG DISPLAY CO., LTD.**,
 Yongin-si, Gyeonggi-do (KR)

(72) Inventors: **Hyoyoung Lee**, Yongin-si (KR);
Youngkook Kim, Yongin-si (KR);
Junha Park, Yongin-si (KR); **Munki Sim**,
 Yongin-si (KR); **Eunjae Jeong**, Yongin-si (KR);
Seokhwan Hwang, Yongin-si (KR)

(73) Assignee: **Samsung Display Co., Ltd.**, Yongin-si,
 Gyeonggi-do (KR)

(*) Notice: Subject to any disclaimer, the term of this
 patent is extended or adjusted under 35
 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/425,351**

(22) Filed: **Feb. 6, 2017**

(65) **Prior Publication Data**
 US 2018/0019406 A1 Jan. 18, 2018

(30) **Foreign Application Priority Data**
 Jul. 18, 2016 (KR) 10-2016-0090775

(51) **Int. Cl.**
H01L 51/50 (2006.01)
H01L 51/00 (2006.01)
 (Continued)

(52) **U.S. Cl.**
 CPC **H01L 51/0072** (2013.01); **C07D 471/14**
 (2013.01); **C07D 487/04** (2013.01);
 (Continued)

(58) **Field of Classification Search**
 None
 See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,645,948 A 7/1997 Shi et al.
 6,465,115 B2 10/2002 Shi et al.
 (Continued)

FOREIGN PATENT DOCUMENTS

JP 10-017860 A 1/1998
 JP 11-087067 A 3/1999
 (Continued)

OTHER PUBLICATIONS

Tang, et al. "Organic Electroluminescent Diodes" Appl. Phys. Lett.,
 vol. 51 (1987) pp. 913-915.

(Continued)


Primary Examiner — Gregory D Clark

(74) Attorney, Agent, or Firm — Lee & Morse, P.C.

(57) **ABSTRACT**

A heterocyclic compound is represented by Formula 1:

<Formula 1>



wherein, in Formula 1, R₁ to R₄, R₁₁, ring A₁, L₁, Ar₁, c₁₁,
 a₁, and b₁ are as defined in the specification.

20 Claims, 4 Drawing Sheets

10

190

150

110

(51) **Int. Cl.**

C07D 471/14 (2006.01)
C07D 487/04 (2006.01)
C07D 491/147 (2006.01)
C07F 9/6561 (2006.01)

KR 10-2014-0049861 A 4/2014
 KR 10-2015-0036402 A 4/2015
 KR 10-2015-0065157 A 6/2015
 KR 10-2015-0075378 A 7/2015
 WO 2014/009317 A1 1/2014
 WO 2015/150234 A1 10/2015
 WO 2015/150234 A8 10/2015

(52) **U.S. Cl.**

CPC **C07D 491/147** (2013.01); **C07F 9/6561**
 (2013.01); **H01L 51/0052** (2013.01); **H01L**
51/0067 (2013.01); **H01L 51/0071** (2013.01);
H01L 51/0073 (2013.01); **H01L 51/0077**
 (2013.01); **H01L 51/5004** (2013.01); **H01L**
51/506 (2013.01); **H01L 51/5072** (2013.01);
H01L 2251/552 (2013.01)

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,596,415 B2 7/2003 Shi et al.
 6,878,469 B2 4/2005 Yoon et al.
 2015/0207083 A1* 7/2015 Schaefer C07D 487/04
 430/79

FOREIGN PATENT DOCUMENTS

KR 10-0691543 B1 3/2007
 KR 10-2010-0097797 A 9/2010
 KR 10-2014-0006708 A 1/2014

OTHER PUBLICATIONS

Adachi, et al. "Confinement of Charge Carriers and Molecular Excitons within 5nm-thick Emitter Layer in Organic Electroluminescent Devices with a Double Heterostructure" Appl. Phys. Lett., vol. 57 (1990) pp. 531-533.

Johansson, et al. "Solid-State Amplified Spontaneous Emission in Some Spiro-Type Molecules: A New Concept for the Design of Solid-State Lasing Molecules" Advanced Materials vol. 10 (1998) pp. 1136-1141.

Tao, et al. "Sharp Green Electroluminescence from 1H-pyrazolo[3,4-b]quinolone-based Light-Emitting Diodes" Appl. Phys. Lett., vol. 77 (2000) pp. 1575-1577.

Sakamoto, et al. "Synthesis, Characterization, and Electron-Transport Property of Perfluorinated Phenylene Dendrimers" J. Am. Chem. Soc., vol. 122 (2000) pp. 1832-1833.

Yamaguchi, et al. "Diphenylamino-Substituted 2,5-Diarylsiloles for Single-Layer Organic Electroluminescent Devices" Chemistry Letters (2001) pp. 98-99.

* cited by examiner

FIG. 1

10

190
150
110

FIG. 2**20**

190
150
110
210

FIG. 3

30

220
190
150
110

FIG. 4

40

220
190
150
110
210

1

HETEROCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

Korean Patent Application No. 10-2016-0090775, filed on Jul. 18, 2016, in the Korean Intellectual Property Office, and entitled: "Heterocyclic Compound and Organic Light-Emitting Device Including the Same," is incorporated by reference herein in its entirety.

BACKGROUND

1. Field

Embodiments relate to a compound for an organic light-emitting device and an organic light-emitting device including the same.


2. Description of the Related Art

Compared to devices in the art, organic light-emitting devices are self-emission devices, have wide viewing angles, high contrast ratios, short response times, and excellent brightness, driving voltage, and response speed characteristics, and produce full-color images.

An example of such organic light-emitting devices may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, recombine in the emission layer to produce excitons. These excitons transition from an excited state to a ground state, thereby generating light.

SUMMARY

Embodiments are directed to a heterocyclic compound represented by Formula 1:



wherein, in Formula 1,

ring A₁ may be a C₁₀-C₆₀ carbocyclic group or a C₆-C₆₀ heterocyclic group,

L₁ may be selected from a substituted or unsubstituted C₃-C₆₀ carbocyclic group, a substituted or unsubstituted C₁-C₆₀ heterocyclic group, *—S(=O)—*, *—S(=O)₂—*, *—P(=O)(Q₄)—*, and *—P(=S)(Q₄)—*,

2

a₁ may be an integer selected from 0 to 5, wherein, when a₁ is two or more, two or more L₁(s) may be identical to or different from each other,

Ar₁ may be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₁-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —S(=O)(Q₁), —S(=O)₂(Q₁), —P(=O)(Q₁)(Q₂), and —P(=S)(Q₁)(Q₂),

b₁ may be an integer selected from 1 to 5, wherein, when b₁ is two or more, two or more Ar₁(s) may be identical to or different from each other,

R₁ to R₄ and R₁₁ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂),

c₁₁ may be an integer selected from 0 to 6, wherein, when c₁₁ is two or more, two or more R₁₁(s) may be identical to or different from each other,

at least one substituent selected from a substituent(s) of the substituted C₁₀-C₆₀ carbocyclic group, the substituted C₆-C₆₀ heterocyclic group, the substituted C₃-C₆₀ carbocyclic group, the substituted C₁-C₆₀ heterocyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an

amino group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

—Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂);

Q₁ to Q₄, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and * and *' each indicate a binding site to a neighboring atom.

According to one or more embodiments, an organic light-emitting device includes: a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode and includes an emission layer, wherein the organic layer includes at least one of the heterocyclic compound provided above.

BRIEF DESCRIPTION OF THE DRAWINGS

Features will become apparent to those of skill in the art by describing in detail exemplary embodiments with reference to the attached drawings in which:

FIG. 1 illustrates a schematic view of an organic light-emitting device according to an embodiment;

FIG. 2 illustrates a schematic view of an organic light-emitting device according to an embodiment;

FIG. 3 illustrates a schematic view of an organic light-emitting device according to an embodiment; and


FIG. 4 illustrates a schematic view of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION

Example embodiments will now be described more fully hereinafter with reference to the accompanying drawings; however, they may be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey exemplary implementations to those skilled in the art.

In the drawing figures, the dimensions of layers and regions may be exaggerated for clarity of illustration. It will also be understood that when a layer or element is referred to as being “on” another layer or substrate, it can be directly on the other layer or substrate, or intervening layers may also be present. In addition, it will also be understood that when a layer is referred to as being “between” two layers, it can be the only layer between the two layers, or one or more intervening layers may also be present. Like reference numerals refer to like elements throughout.

A heterocyclic compound according to an embodiment may be represented by Formula 1:



wherein, in Formula 1,

ring A₁ may be a C₁₀-C₆₀ carbocyclic group or a C₆-C₆₀ heterocyclic group.

In one or more embodiments, ring A₁ may be a C₁₀-C₆₀ carbocyclic group with two or more rings condensed with each other or a C₆-C₆₀ heterocyclic group with two or more rings condensed with each other.

In one or more embodiments, ring A₁ may be a naphthalene, a fluorene, a benzofluorene, a phenanthrene, an anthracene, a quinoline, an isoquinoline, a benzoquinoline, a carbazole, a dibenzofuran, a dibenzothiophene, or a benzo-carbazolyl.

In one or more embodiments, ring A₁ may be a naphthalene, a phenanthrene, a quinoline, or a dibenzofuran, as examples.

L₁ in Formula 1 may be selected from a substituted or unsubstituted C₃-C₆₀ carbocyclic group, a substituted or unsubstituted C₁-C₆₀ heterocyclic group, *—S(=O)—*, *—S(=O)₂—*, *—P(=O)(Q₄)—*, and *—P(=S)(Q₄)—*.

5

In one or more embodiments, L_1 may be selected from:

a benzene group, a pentalene group, an indene group, a naphthalene group, an azulene group, a heptalene group, an indacene group, an acenaphthalene group, a fluorene group, a spiro-bifluorene group, a spiro-benzofluorene-fluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pyrrole group, a thiophene group, a furan group, a silole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a triazine group, a benzofuran group, a benzothiophene group, a benzosilole group, a dibenzofuran group, a dibenzothiophene group, a dibenzosilole group, a carbazole group, a benzocarbazole group, a dibenzocarbazole group, a benzimidazole group, and an imidazopyridine group;

a benzene group, a pentalene group, an indene group, a naphthalene group, an azulene group, a heptalene group, an indacene group, an acenaphthalene group, a fluorene group, a spiro-bifluorene group, a spiro-benzofluorene-fluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pyrrole group, a thiophene group, a furan group, a silole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a triazine group, a benzofuran group, a benzothiophene group, a benzosilole group, a dibenzofuran group, a dibenzothiophene group, a dibenzosilole group, a carbazole group, a benzocarbazole group, a dibenzocarbazole group, a benzimidazole group, and an imidazopyridine group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a carbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), and —B(Q_{31})(Q_{32}); and


—S(=O)—, *—S(=O)₂—*, *—P(=O)(Q_4)—*, and *—P(=S)(Q_4)—*,

Q_4 and Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, and


* and *' each indicate a binding site to a neighboring atom.

6


In one or more embodiments, L_1 may be selected from groups represented by Formulae 3-1 to 3-51, as examples:




Formula 3-1




Formula 3-2




Formula 3-3




Formula 3-4




Formula 3-5




Formula 3-6



Formula 3-7











Formula 3-8



Formula 3-9

7

-continued




8

-continued

Formula 3-10

5



Formula 3-11 10

Formula 3-12 15

Formula 3-13 20

Formula 3-14 30

Formula 3-15 35

Formula 3-16 45

Formula 3-17 55

25

30

35

40





45

50

55

60

65



Formula 3-18










Formula 3-19

Formula 3-20

Formula 3-21

Formula 3-22


-continued




-continued

Formula 3-23

5




10




Formula 3-24

15




Formula 3-25

20




25




Formula 3-26

30




35




Formula 3-27

40




Formula 3-28

45




Formula 3-29

50



Formula 3-30

55



Formula 3-31

60

65

Formula 3-32

Formula 3-33

Formula 3-34

Formula 3-35

Formula 3-36

Formula 3-37

Formula 3-38

Formula 3-39









Formula 3-40

Formula 3-41

Formula 3-42

11


-continued



12

-continued

Formula 3-43



Formula 3-44

wherein, in Formulae 3-1 to 3-51,

Y₁ may be O, S, C(Z₃)(Z₄), N(Z₃), or Si(Z₃)(Z₄);

Z₁ to Z₄ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkenyl group, a C₁-C₂₀ alkynyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

Formula 3-45

wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

d₂ may be 1 or 2,d₃ may be an integer selected from 1 to 3,d₄ may be an integer selected from 1 to 4,d₅ may be an integer selected from 1 to 5,d₆ may be an integer selected from 1 to 6,d₈ may be an integer selected from 1 to 8, and

Formula 3-47

* and *' each indicate a binding site to a neighboring atom.

In one or more embodiments, L₁ may be selected from:

a benzene group, a naphthalene group, a fluorene group,

a phenanthrene group, an anthracene group, a pyridine group,

a pyrazine group, a pyrimidine group, a pyridazine group,

a triazine group, a benzofuran group, a benzothiophene group,

a benzosilole group, a dibenzofuran group, a dibenzothiophene group,

a dibenzosilole group, and a carbazole group;

Formula 3-48

a benzene group, a naphthalene group, a fluorene group,

a phenanthrene group, an anthracene group, a pyridine group,

a pyrazine group, a pyrimidine group, a pyridazine group,

a triazine group, a benzofuran group, a benzothiophene group,

a benzosilole group, a dibenzofuran group, a dibenzothiophene group,

a dibenzosilole group, and a carbazole group, each substituted with at least one selected

from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group,

a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group,a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), and—B(Q₃₁)(Q₃₂); and*—S(=O)—*', *—S(=O)₂—*', *—P(=O)(Q₄)—*',and *—P(=S)(Q₄)—*',

Formula 3-50

wherein Q₄ and Q₃₁ to Q₃₃ may each independently beselected from a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group,

a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, and

* and *' each indicate a binding site to a neighboring atom.

In one or more embodiments, L_1 may be selected from:

a benzene group, a fluorene group, an anthracene group, a pyridine group, a triazine group, and a carbazole group; a benzene group, a fluorene group, an anthracene group, a pyridine group, a triazine group, and a carbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), and —B(Q_{31})(Q_{32}); and

—S(=O)—, *—S(=O)₂—*, *—P(=O)(Q_4)—*, and *—P(=S)(Q_4)—*,

wherein Q_4 and Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, and

* and *' each indicate a binding site to a neighboring atom.

a_1 in Formula 1 may be an integer selected from 0 to 5. a_1 indicates the number of $L_1(s)$ in Formula 1, wherein, when a_1 is two or more, two or more $L_1(s)$ may be identical to or different from each other. For example, a_1 may be 0, 1, 2, or 3.

Ar_1 in Formula 1 may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —S(=O)(Q_1), —S(=O)₂(Q_1), —P(=O)(Q_1)(Q_2), and —P(=S)(Q_1)(Q_2).

In one or more embodiments, Ar_1 may be selected from:


a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a thiophenyl group, a furanyl group, a carbazolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzonaphthofuranyl group, a dinaphthofuranyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a benzonaphthosilolyl group, a dinaphthosilolyl group, a benzimidazolyl group, a benzoimidazonaphthoimidazolyl group, and an imidazopyridinyl group;

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrene-


nyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a thiophenyl group, a furanyl group, a carbazolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzonaphthofuranyl group, a dinaphthofuranyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a benzonaphthosilolyl group, a dinaphthosilolyl group, a benzimidazolyl group, a benzoimidazonaphthoimidazolyl group, and an imidazopyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a thiophenyl group, a furanyl group, a carbazolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, —N(Q_{31})(Q_{32}), and —Si(Q_{31})(Q_{32})(Q_{33}); and —S(=O)(Q_1), —S(=O)₂(Q_1), —P(=O)(Q_1)(Q_2), and —P(=S)(Q_1)(Q_2),

wherein Q_1 , Q_2 , and Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, as examples.


In one or more embodiments, Ar_1 may be selected from groups represented by Formulae 5-1 to 5-29, —S(=O)(Q_1), —S(=O)₂(Q_1), —P(=O)(Q_1)(Q_2), and —P(=S)(Q_1)(Q_2):




Formula 5-1



Formula 5-2













Formula 5-3



Formula 5-4

15

-continued




16

-continued


Formula 5-5

5




Formula 5-6

10



Formula 5-7

15




Formula 5-8

20

Formula 5-9

25




Formula 5-10

30

Formula 5-11

35




Formula 5-12

40

Formula 5-13

50




45

55

Formula 5-14

60



65

Formula 5-15

Formula 5-16

Formula 5-17

Formula 5-18

Formula 5-19







Formula 5-20

Formula 5-21

Formula 5-22

17

-continued




18

-continued

Formula 5-29

Formula 5-23



Formula 5-24

wherein, in Formulae 5-1 to 5-29,
 15 Y_{31} may be O, S, C(Z_{35})(Z_{36}), N(Z_{35}), or Si(Z_{35})(Z_{36}),
 Z_{31} to Z_{36} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group,
 20 a C_1 - C_{20} alkenyl group, a C_1 - C_{20} alkynyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group,
 25 —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),
 30 wherein Q_1 , Q_2 , and Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

Formula 5-25

Formula 5-26

Formula 5-27

Formula 5-28

Formula 5-29

Formula 5-30

Formula 5-31

Formula 5-32

Formula 5-33

Formula 5-34

Formula 5-35

Formula 5-36

Formula 5-37

Formula 5-38

Formula 5-39

Formula 5-40

Formula 5-41

Formula 5-42

Formula 5-43

Formula 5-44

Formula 5-45

Formula 5-46

Formula 5-47

Formula 5-48

Formula 5-49

Formula 5-50

Formula 5-51

Formula 5-52

Formula 5-53

Formula 5-54

Formula 5-55

Formula 5-56

Formula 5-57

Formula 5-58

Formula 5-59

Formula 5-60

Formula 5-61

Formula 5-62

Formula 5-63

Formula 5-64

Formula 5-65

Formula 5-66

Formula 5-67

Formula 5-68

Formula 5-69

Formula 5-70

Formula 5-71

Formula 5-72

Formula 5-73

Formula 5-74

Formula 5-75

Formula 5-76

Formula 5-77

Formula 5-78

Formula 5-79

Formula 5-80

Formula 5-81

Formula 5-82

Formula 5-83

Formula 5-84

Formula 5-85

Formula 5-86

Formula 5-87

Formula 5-88

Formula 5-89

Formula 5-90

Formula 5-91

Formula 5-92

Formula 5-93

Formula 5-94

Formula 5-95

Formula 5-96

Formula 5-97

Formula 5-98

Formula 5-99

Formula 5-100

Formula 5-101

Formula 5-102

Formula 5-103

Formula 5-104

Formula 5-105

Formula 5-106

Formula 5-107

Formula 5-108

Formula 5-109

Formula 5-110

Formula 5-111

Formula 5-112

Formula 5-113

Formula 5-114

Formula 5-115

Formula 5-116

Formula 5-117

Formula 5-118

Formula 5-119

Formula 5-120

Formula 5-121

Formula 5-122

Formula 5-123

Formula 5-124

Formula 5-125

Formula 5-126

Formula 5-127

Formula 5-128

Formula 5-129

Formula 5-130

Formula 5-131

Formula 5-132

Formula 5-133

Formula 5-134

Formula 5-135

Formula 5-136

Formula 5-137

Formula 5-138

Formula 5-139

Formula 5-140

Formula 5-141

Formula 5-142

Formula 5-143

Formula 5-144

Formula 5-145

Formula 5-146

Formula 5-147

Formula 5-148

Formula 5-149

Formula 5-150

Formula 5-151

Formula 5-152

Formula 5-153

Formula 5-154

Formula 5-155

Formula 5-156

Formula 5-157

Formula 5-158

Formula 5-159

Formula 5-160

Formula 5-161

Formula 5-162

Formula 5-163

Formula 5-164

Formula 5-165

Formula 5-166

Formula 5-167

Formula 5-168

Formula 5-169

Formula 5-170

Formula 5-171

Formula 5-172

Formula 5-173

Formula 5-174

Formula 5-175

Formula 5-176

Formula 5-177

Formula 5-178

Formula 5-179

Formula 5-180

Formula 5-181

Formula 5-182

Formula 5-183

Formula 5-184

Formula 5-185

Formula 5-186

Formula 5-187

Formula 5-188

Formula 5-189

Formula 5-190

Formula 5-191

Formula 5-192

Formula 5-193

Formula 5-194

Formula 5-195

Formula 5-196

Formula 5-197

Formula 5-198

Formula 5-199

Formula 5-200

Formula 5-201

Formula 5-202

Formula 5-203

Formula 5-204

Formula 5-205

Formula 5-206

Formula 5-207

Formula 5-208

Formula 5-209

Formula 5-210

Formula 5-211

Formula 5-212

Formula 5-213

Formula 5-214

Formula 5-215

Formula 5-216

Formula 5-217

Formula 5-218

Formula 5-219

Formula 5-220

Formula 5-221

Formula 5-222

Formula 5-223

Formula 5-224

Formula 5-225

Formula 5-226

Formula 5-227

Formula 5-228

Formula 5-229

Formula 5-230

Formula 5-231

Formula 5-232

Formula 5-233

Formula 5-234

Formula 5-235

Formula 5-236

Formula 5-237

Formula 5-238

Formula 5-239

Formula 5-240

Formula 5-241

Formula 5-242

Formula 5-243

Formula 5-244

Formula 5-245

Formula 5-246

Formula 5-247

Formula 5-248

Formula 5-249

Formula 5-250

Formula 5-251

Formula 5-252

Formula 5-253

Formula 5-254

Formula 5-255

Formula 5-256

Formula 5-257

Formula 5-258

Formula 5-259

Formula 5-260

Formula 5-261

Formula 5-262

Formula 5-263

Formula 5-264

Formula 5-265

Formula 5-266

Formula 5-267

Formula 5-268

Formula 5-269

Formula 5-270

Formula 5-271

Formula 5-272

Formula 5-273

Formula 5-274

Formula 5-275

Formula 5-276

Formula 5-277

Formula 5-278

Formula 5-279

Formula 5-280

Formula 5-281

Formula 5-282

Formula 5-283

Formula 5-284

Formula 5-285

Formula 5-286

Formula 5-287

Formula 5-288

Formula 5-289

Formula 5-290

Formula 5-291

Formula 5-292

Formula 5-293

Formula 5-294

Formula 5-295

Formula 5-296

Formula 5-297

Formula 5-298

Formula 5-299

Formula 5-300

Formula 5-301

Formula 5-302

Formula 5-303

Formula 5-304

Formula 5-305

Formula 5-306

Formula 5-307

Formula 5-308











Formula 5-309

Formula 5-310

Formula 5-311

19

-continued




20

-continued


Formula 6-5

5




Formula 6-6

10




Formula 6-7

15




Formula 6-8

20




Formula 6-9

25




Formula 6-10

30




Formula 6-11

40




Formula 6-12

45




Formula 6-13

50



Formula 6-14

60



65

Formula 6-15

Formula 6-16

Formula 6-17

Formula 6-18

Formula 6-19

Formula 6-20

Formula 6-21







Formula 6-22

Formula 6-23

Formula 6-24

21

-continued




22

-continued

Formula 6-25


5



10

Formula 6-26


15



20

Formula 6-27


25



30

Formula 6-28


35




40

Formula 6-29

45




50



55

Formula 6-30

60



65

Formula 6-31

Formula 6-32

Formula 6-33

Formula 6-34








Formula 6-35

Formula 6-36

Formula 6-37

23


-continued

**24**

-continued

Formula 6-38

5



Formula 6-39

10

Formula 6-40

15

Formula 6-41

25

Formula 6-42

30

Formula 6-43




45

Formula 6-44

55

60

65



Formula 6-45

Formula 6-46

Formula 6-47






Formula 6-48

Formula 6-49

Formula 6-50

25

-continued



wherein * indicates a binding site to a neighboring atom.

In one or more embodiments, Ar₁ may be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, a pyridinyl group, a dinaphthofuranyl group, and a benzoimidazodaphthoimidazolyl group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, a pyridinyl group, a dinaphthofuranyl group, and a benzoimidazodaphthoimidazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, —N(Q₃₁)(Q₃₂), and —Si(Q₃₁)(Q₃₂)(Q₃₃); and

—S(=O)(Q₁), —S(=O)₂(Q₁), —P(=O)(Q₁)(Q₂), and —P(=S)(Q₁)(Q₂),

wherein Q₁, Q₂, and Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

b1 in Formula 1 may be an integer selected from 1 to 5. b1 indicates the number of Ar₁(s) in Formula 1, wherein,

26

Formula 6-51

when b1 is two or more, two or more Ar₁(s) may be identical to or different from each other. For example, b1 may be 1 or 2.

Formula 6-52

R₁ to R₄ and R₁₁ in Formula 1 may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂).

Formula 6-53

Formula 6-54

In one or more embodiments, R₁ to R₄ may each independently be selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, and a nitro group;

—Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂); and groups represented by Formulae 5-1 to 5-29, and

Formula 6-55


R₁₁ may be selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;


a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, and a nitro group; and

—Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂);


Formula 5-1




Formula 5-2



Formula 5-3













Formula 5-4



27

-continued




28

-continued


Formula 5-5

5




Formula 5-6

10



Formula 5-7

15




Formula 5-8

20

Formula 5-9

25




Formula 5-10

30

Formula 5-11

35




Formula 5-12

40

Formula 5-13

50




45

55

Formula 5-14

60



65

Formula 5-15

Formula 5-16

Formula 5-17

Formula 5-18

Formula 5-19







Formula 5-20

Formula 5-21

Formula 5-22

29


-continued



30

-continued

Formula 5-29



Formula 5-23

Formula 5-24

Formula 5-25

Formula 5-26

Formula 5-27

Formula 5-28

In Formulae 5-1 to 5-29,

- 15 Y_{31} may be O, S, C(Z_{35})(Z_{36}), N(Z_{35}), or Si(Z_{35})(Z_{36}),
 Z_{31} to Z_{36} may each independently be selected from
hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl
group, a cyano group, a nitro group, an amidino group, a
hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group,
20 a C_1 - C_{20} alkenyl group, a C_1 - C_{20} alkynyl group, a C_1 - C_{20}
alkoxy group, a phenyl group, a biphenyl group, a terphenyl
group, a naphthyl group, a fluorenyl group, a spiro-bifluo-
renyl group, a benzofluorenyl group, a dibenzofluorenyl
group, a phenanthrenyl group, an anthracenyl group, a
25 pyrenyl group, a chrysenyl group, a pyridinyl group, a
pyrazinyl group, a pyrimidinyl group, a pyridazinyl group,
a quinolinyl group, an isoquinolinyl group, a quinoxalinyl
group, a quinazolinyl group, a carbazolyl group, a triazinyl
group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})
30 (Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})
(Q_{32}),

wherein Q_1 , Q_2 , and Q_{31} to Q_{33} may each independently
be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy
group, a phenyl group, a biphenyl group, a terphenyl group,
35 and a naphthyl group,

e_2 may be 1 or 2,

e_3 may be an integer selected from 1 to 3,

e_4 may be an integer selected from 1 to 4,

e_5 may be an integer selected from 1 to 5,

40 e_6 may be an integer selected from 1 to 6,

e_7 may be an integer selected from 1 to 7,

e_9 may be an integer selected from 1 to 9, and

* and *' each indicate a binding site to a neighboring
atom.

- 45 In one or more embodiments, R_1 to R_4 may each inde-
pendently be selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl
group, a cyano group, a nitro group, a C_1 - C_{20} alkyl group,
and a C_1 - C_{20} alkoxy group;

- 50 a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each
substituted with at least one selected from deuterium, —F,
—Cl, —Br, —I, a hydroxyl group, a cyano group, and a
nitro group;

- a phenyl group, a biphenyl group, a terphenyl group, a
55 fluorenyl group, a phenanthrenyl group, an anthracenyl
group, a pyridinyl group, a carbazolyl group, a dibenzofura-
nyl group, and a dibenzothiophenyl group;

- a phenyl group, a biphenyl group, a terphenyl group, a
fluorenyl group, a phenanthrenyl group, an anthracenyl
60 group, a pyridinyl group, a carbazolyl group, a dibenzofura-
nyl group, and a dibenzothiophenyl group, each substituted
with at least one selected from deuterium, —F, —Cl, —Br,
—I, a hydroxyl group, a cyano group, a nitro group, an
amidino group, a hydrazino group, a hydrazono group, a
65 C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group,
a biphenyl group, a terphenyl group, —N(Q_{31})(Q_{32}), and
—Si(Q_{31})(Q_{32})(Q_{33}); and

31

—Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂),
—C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂), and

R₁₁ may be selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl
group, a cyano group, a nitro group, a C₁-C₂₀ alkyl group,
and a C₁-C₂₀ alkoxy group;


a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each
substituted with at least one selected from deuterium, —F,
—Cl, —Br, —I, a hydroxyl group, a cyano group, and a
nitro group; and

—Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂),
—C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂),

wherein Q₁ to Q₃ and Q₃₁ to Q₃₃ may each independently
be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy
group, a phenyl group, a biphenyl group, a terphenyl group,
and a naphthyl group, as examples.






c11 in Formula 1 may be an integer selected from 0 to 6.
c11 indicates the number of R₁₁(s) in Formula 1, wherein,
when c11 is two or more, two or more R₁₁(s) may be
identical to or different from each other. For example, c11
may be 0, 1, or 2.

In one or more embodiments, the compound represented
by Formula 1 may be represented by one selected from
Formulae 1A to 1J:




32

-continued



33

-continued



In Formulae 1A to 1J, L_1 , a_1 , Ar_1 , b_1 , and R_1 to R_4 are the same as described above,

Y_{51} may be O, S, C(Z_{53})(Z_{54}), N(Z_{53}), or Si(Z_{53})(Z_{54}),

Y_{52} to Y_{57} may each independently be C(Z_{55}) or N,

Z_{51} to Z_{55} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkenyl group, a C_1 - C_{20} alkynyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

f_2 may be 1 or 2,

f_4 may be an integer selected from 1 to 4,

f_6 may be an integer selected from 1 to 6, and

f_8 may be an integer selected from 1 to 8.

In Formulae 1A to 1J,

L_1 may be selected from:

a benzene group, a fluorene group, an anthracene group, a pyridine group, a triazine group, and a carbazole group;

a benzene group, a fluorene group, an anthracene group, a pyridine group, a triazine group, and a carbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), and —B(Q_{31})(Q_{32}); and

—S(=O)—, *—S(=O)₂—*, *—P(=O)(Q_4)—*, and *—P(=S)(Q_4)—*,

a_1 may be an integer selected from 0 to 3,

Ar_1 may be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, a pyridinyl group, a dinaphthofuranyl group, and a benzoimidazonaphthoimidazolyl group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, a pyridinyl group, a dinaphthofuranyl group, and a benzoimidazonaphthoimidazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino

34

group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, —N(Q_{31})(Q_{32}), and —Si(Q_{31})(Q_{32})(Q_{33}); and

5 —S(=O)(Q_1), —S(=O)₂(Q_1), —P(=O)(Q_1)(Q_2), and —P(=S)(Q_1)(Q_2),

b_1 may be 1 or 2, and

R_1 to R_4 may each independently be selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, and a nitro group;

a phenyl group, a biphenyl group, a terphenyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyridinyl group, a carbazolyl group, a dibenzofuran group, and a dibenzothiophenyl group;

a phenyl group, a biphenyl group, a terphenyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyridinyl group, a carbazolyl group, a dibenzofuran group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a

C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, —N(Q_{31})(Q_{32}), and —Si(Q_{31})(Q_{32})(Q_{33}); and

—Si(Q_1)(Q_2)(Q_3), —N(Q_1)(Q_2), B(Q_1)(Q_2), —C(=O)(Q_1), —S(=O)₂(Q_1), and —P(=O)(Q_1)(Q_2),

wherein Q_1 to Q_4 and Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, as examples.

In one or more embodiments, in Formulae 1A to 1J,

R_1 , R_2 , and R_4 may each independently be selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, and a nitro group; and

—Si(Q_1)(Q_2)(Q_3), —N(Q_1)(Q_2), —B(Q_1)(Q_2), —C(=O)(Q_1), —S(=O)₂(Q_1), and —P(=O)(Q_1)(Q_2), and R_3 may be selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, and a nitro group;

a phenyl group, a biphenyl group, a terphenyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyridinyl group, a carbazolyl group, a dibenzofuran group, and a dibenzothiophenyl group;

a phenyl group, a biphenyl group, a terphenyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyridinyl group, a carbazolyl group, a dibenzofuran group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a

35

C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, $-N(Q_{31})(Q_{32})$, and $-Si(Q_{31})(Q_{32})(Q_{33})$; and









$-Si(Q_1)(Q_2)(Q_3)$, $-N(Q_1)(Q_2)$, $-B(Q_1)(Q_2)$, $-C(=O)(Q_1)$, $-S(=O)_2(Q_1)$, and $-P(=O)(Q_1)(Q_2)$,

wherein Q_1 to Q_3 and Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, as examples.

In one or more embodiments, the heterocyclic compound represented by Formula 1 may be one selected from Compounds 1 to 56:





36

-continued








37

-continued






**38**

-continued



39

-continued

**40**


-continued

18

23

5

10




15

24

19

20

25




20

30

35

40




21

26

45

50




22

55

27





60

65



41

-continued




42

-continued

28

32

5




10

15

33

29


20



25

34

30




35

30

40

35

45




50

36

31

55







60

65

43

-continued



44

-continued

37

5

10

38 15

20

25

30

39

35

40

45





50

40

55

60

65



41


42

43

44

45

-continued



45

5

10

15

20

46

25

30

35

47

40

45

50

48


55

60

65

46

-continued







49

50

51


47

-continued



48

-continued



The heterocyclic compound represented by Formula 1 may include a benzimidazole and has a heterocyclic core with five or more rings condensed with one another. For example, due to the inclusion of three or more heteroatoms and a condensed ring, the heterocyclic compound represented by Formula 1 may form a π -conjugated system. Thus, packing between molecules may be increased, and a glass transition temperature and a melting point may be increased. Also, a charge transport capability may also be increased. Therefore, an organic light-emitting device including the heterocyclic compound represented by Formula 1 may have high luminance and a long lifespan.

The compound represented by Formula 1 may have relatively high charge (hole or electron) transport capability. Accordingly, it may be possible to improve an exciton-forming ratio in an emission layer in the organic light-emitting device including the heterocyclic compound represented by Formula 1. Thus, the organic light-emitting device may have a low driving voltage and high efficiency.

The synthesis methods of the heterocyclic compound represented by Formula 1 may be recognizable by one of ordinary skill in the art by referring to Examples provided below.

At least one of the heterocyclic compounds represented by Formula 1 may be used between a pair of electrodes of an organic light-emitting device. For example, the heterocyclic compound may be included in at least one of a hole transport region, an electron transport region, and an emission layer. In one or more embodiments, the heterocyclic compound represented by Formula 1 may be used as a material for a capping layer located outside a pair of electrodes of an organic light-emitting device.

According to one or more embodiments, an organic light-emitting device may include a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode. The organic layer includes an emission layer. The organic layer may include at least one of the heterocyclic compounds represented by Formula 1.

The expression “(an organic layer) includes at least one of the heterocyclic compounds” used herein may include a case in which “(an organic layer) includes identical heterocyclic compounds represented by Formula 1” and a case in which “(an organic layer) includes two or more different heterocyclic compounds represented by Formulae 1A to 1J.”

In one or more embodiments, the first electrode may be an anode, the second electrode may be a cathode, the organic layer may further include a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second

electrode. The hole transport region may include a hole injection layer, a hole transport layer, an emission auxiliary layer, an electron blocking layer, or any combination thereof. The electron transport region may include a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, an electron injection layer, or any combination thereof.

The organic light-emitting device may further include at least one selected from a first capping layer disposed in a pathway along which light generated in an emission layer proceeds toward outside through the first electrode and a second capping layer disposed in a pathway along which light generated in an emission layer proceeds toward outside through the second electrode. The at least one selected from the first capping layer and the second capping layer may include at least one of the heterocyclic compounds represented by Formula 1.

In one or more embodiments, the organic light-emitting device may have i) a stack structure including a first electrode, an organic layer, a second electrode, and a second capping layer which are sequentially stacked in this stated order, ii) a stack structure including a first capping layer, a first electrode, an organic layer, and a second electrode which are sequentially stacked in this stated order, or iii) a stack structure including a first capping layer, a first electrode, an organic layer, a second electrode, and a second capping layer which are sequentially stacked in this stated order. At least one selected from the first capping layer and the second capping layer may include the heterocyclic compound.

The term "organic layer" as used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode in the organic light-emitting device. The organic layer may include other materials besides an organic material.

Description of FIG. 1

FIG. 1 illustrates a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 may include a first electrode 110, an organic layer 150, and a second electrode 190.

Hereinafter, the structure of the organic light-emitting device 10 according to an embodiment and a method of manufacturing the organic light-emitting device 10 will be described in connection with FIG. 1.

First Electrode 110

In FIG. 1, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be a substrate having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water-resistance. For example, the substrate may be glass substrate or a plastic substrate.

The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode 110 on the substrate. When the first electrode 110 is an anode, the material for forming the first electrode 110 may be selected from materials with a high work function to facilitate hole injection.

The first electrode 110 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode 110 is a transmissive electrode, a material for forming the first electrode 110 may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), zinc oxide (ZnO), and any combinations thereof, as examples. When the first electrode 110 is a semi-transmissive electrode or a reflective electrode, as a material for forming the first electrode 110, magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), or any combination thereof may be used, as examples.

The first electrode 110 may have a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode 110 may have a three-layered structure of ITO/Ag/ITO, as an example.

Organic Layer 150

The organic layer 150 may be disposed on the first electrode 110. The organic layer 150 may include an emission layer.

The organic layer 150 may further include a hole transport region between the first electrode 110 and the emission layer, and an electron transport region between the emission layer and the second electrode 190.


Hole Transport Region in Organic Layer 150

The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

The hole transport region may include at least one layer selected from a hole injection layer, a hole transport layer, an emission auxiliary layer, and an electron blocking layer.

For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure, wherein, for each structure, constituting layers are sequentially stacked from the first electrode 110 in the stated order.


The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPB), β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTDP, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), PEDOT/PSS (poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate)), polyaniline/camphor sulfonic acid (Pani/CSA), polyaniline/poly(4-styrenesulfonate) (Pani/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:



m-MTDATA

51


-continued




TDATA

52


-continued



TPD




2-TNATA




Spiro-TPD

55





NPB

65



methylated NPB

-continued



In Formulae 201 and 202,

L₂₀₁ to L₂₀₄ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

L_{205} may be selected from $*-O-*$, $*-S-*$, $*-N$ (Q₂₀₁)-*, a substituted or unsubstituted C₁-C₂₀ alkylene group, a substituted or unsubstituted C₂-C₂₀ alkenylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

54

xa1 to xa4 may each independently be an integer selected from 0 to 3,

xa5 may be an integer selected from 1 to 10, and

R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

For example, in Formula 202, R₂₀₁ and R₂₀₂ may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group, and R₂₀₃ and R₂₀₄ may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group.

In one or more embodiments, in Formulae 201 and 202,

L₂₀₁ to L₂₀₅ may each independently be selected from:

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, dibenzosilolylene group, and a pyridinylene group; and

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, dibenzosilolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁–C₂₀ alkyl group, a C₁–C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁–C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenylene group, an indenyl group, a naphthyl group, an azulenyl

55

group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, and $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$,

wherein Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, xa1 to xa4 may each independently be an integer selected from 0, 1, and 2.

In one or more embodiments, xa5 may be 1, 2, 3, or 4.

In one or more embodiments, R_{201} to R_{204} and Q_{201} may each independently be selected from a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a Spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cycloheptenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-\text{F}$, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-

56

bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, and $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$,

wherein Q_{31} to Q_{33} are the same as described above.

In one or more embodiments, at least one selected from R_{201} to R_{203} in Formula 201 may each independently be selected from:

a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-\text{F}$, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, as examples.

In one or more embodiments, in Formula 202, i) R_{201} and R_{202} may be linked via a single bond and/or ii) R_{203} and R_{204} may be linked via a single bond.


In one or more embodiments, at least one selected from R_{201} to R_{204} in Formula 202 may be selected from:

a carbazolyl group; and

a carbazolyl group, substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-\text{F}$, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, as examples.


The compound represented by Formula 201 may be represented by Formula 201A:

<Formula 201A>




57


In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A(1) below, as an example:



In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A-1 below, as an example:




In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A:



58

In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A-1:



In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1, L₂₀₁ to L₂₀₃, xa1 to xa3, xa5, and R₂₀₂ to R₂₀₄ are the same as described above, and


R₂₁₁ and R₂₁₂ may be the same as described in connection with R₂₀₃.

R₂₁₃ to R₂₁₇ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

The hole transport region may include at least one compound selected from Compounds HT1 to HT39, as examples:


59

HT1




60


HT2



HT3



HT4





61

62

-continued



HT5

HT6




HT7

HT8




63

-continued
HT9




64


HT10



HT11




HT12




65

-continued


HT13



HT15




HT17




66


HT14



HT16




HT18




67

-continued
HT19





68

HT20




HT21

HT22



HT23




69

70



-continued

HT24





HT25

HT26



HT27

HT28





71

72

-continued



HT29

HT30





HT31

HT32





HT33

HT34




HT35


HT36




73

-continued
HT37

74



HT38



HT39

45

A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, or, for example, about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer and a hole transport layer, the thickness of the hole injection layer may be in a range of about 100 Å to about 9,000 Å, or, for example, about 100 Å to about 1,000 Å, and the thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, or, for example, about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer.

The electron blocking layer may block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include the materials as described above.

p-dopant

The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

The charge-generation material may be, for example, a p-dopant.

In one or more embodiments, a lowest unoccupied molecular orbital (LUMO) level of the p-dopant may be -3.5 eV or less.

The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, as examples.

For example, the p-dopant may include at least one selected from:


a quinone derivative, such as tetracyanoquinodimethane (TCNQ) or 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);

a metal oxide, such as tungsten oxide or molybdenum oxide;

65

75

1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and
a compound represented by Formula 221, as examples:



In Formula 221,

R_{221} to R_{223} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, provided that at least one selected from R_{221} to R_{223} has at least one substituent selected from a cyano group, —F, —Cl, —Br, —I, a C_1 - C_{20} alkyl group substituted with —F, a C_1 - C_{20} alkyl group substituted with —Cl, a C_1 - C_{20} alkyl group substituted with —Br, and a C_1 - C_{20} alkyl group substituted with —I.

Emission Layer in Organic Layer 150

When the organic light-emitting device **10** is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In one or more embodiments, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers contact each other or are separated from each other. In one or more embodiments, the emission layer may include two or more materials selected from a red-light emission material, a green-light emission material, and a blue-light emission material, in which the two or more materials are mixed with each other in a single layer to emit white light.

76


The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant and a fluorescent dopant.

An amount of the dopant in the emission layer may be, for example, in a range of about 0.01 to about 15 parts by weight based on 100 parts by weight of the host.

A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, or, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

Host in Emission Layer

In one or more embodiments, the host may include a compound represented by Formula 301 below:



In Formula 301,

Ar_{301} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

$xb11$ may be 1, 2, or 3,

L_{301} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

$xb1$ may be an integer selected from 0 to 5,

R_{301} may be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_{301})(Q_{302})(Q_{303}), —N(Q_{301})(Q_{302}), —B(Q_{301})(Q_{302}), —C(=O)(Q_{301}), —S(=O)₂(Q_{301}), and —P(=O)(Q_{301})(Q_{302}), and

$xb21$ may be an integer selected from 1 to 5,

wherein Q_{301} to Q_{303} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, as examples.

In one or more embodiments, Ar_{301} in Formula 301 may be selected from:

a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene

77


group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group and a dibenzothio-
phene group; and

a naphthalene group, a fluorene group, a spiro-bifluorene
group, a benzofluorene group, a dibenzofluorene group, a
phenalene group, a phenanthrene group, an anthracene
group, a fluoranthene group, a triphenylene group, a pyrene
group, a chrysene group, a naphthacene group, a picene
group, a perylene group, a pentaphene group, an indenoan-
thracene group, a dibenzofuran group, and a dibenzothio-
phene group, each substituted with at least one selected from
deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano
group, a nitro group, an amidino group, a hydrazino group,
a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy
group, a phenyl group, a biphenyl group, a terphenyl group,
a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂),
—B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and
—P(=O)(Q₃₁)(Q₃₂),


wherein Q₃₁ to Q₃₃ may each independently be selected
from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl
group, a biphenyl group, a terphenyl group, and a naphthyl
group, as examples.

When xb11 in Formula 301 is two or more, two or more
Ar₃₀₁ (s) may be linked via a single bond.

In one or more embodiments, the compound represented
by Formula 301 may be represented by Formula 301-1 or
301-2:



<Formula 301-1>



<Formula 301-2>

50

In Formulae 301-1 to 301-2,

A₃₀₁ to A₃₀₄ may each independently be selected from a
benzene, a naphthalene, a phenanthrene, a fluoranthene, a
triphenylene, a pyrene, a chrysene, a pyridine, a pyrimidine,
an indene, a fluorene, a spiro-bifluorene, a benzofluorene, a
dibenzofluorene, an indole, a carbazole, benzocarbazole,
dibenzocarbazole, a furan, a benzonaphthofuran, a dibenzofuran,
a naphthofuran, a benzonaphthofuran, a dinaphthofuran, a
thiophene, a benzothiophene, a dibenzothiophene, a naph-
thothiophene, a benzonaphthothiophene, and dinaphtho-
thiophene,

X₃₀₁ may be O, S, or N—[(L₃₀₄)xb4-R₃₀₄],

R₃₁₁ to R₃₁₄ may each independently be selected from
hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl
group, a cyano group, a nitro group, an amidino group, a
hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group,

78

a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a
terphenyl group, a naphthyl group —Si(Q₃₁)(Q₃₂)(Q₃₃),
—N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂
(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

xb22 and xb23 may each independently be 0, 1, or 2,

L₃₀₁, xb1, R₃₀₁, and Q₃₁ to Q₃₃ are the same as described
above,

L₃₀₂ to L₃₀₄ are each independently the same as described
in connection with L₃₀₁,

xb2 to xb4 are each independently the same as described
in connection with xb1, and

R₃₀₂ to R₃₀₄ are each independently the same as described
in connection with R₃₀₁.

For example, L₃₀₁ to L₃₀₄ in Formulae 301, 301-1, and
301-2 may each independently be selected from:

a phenylene group, a naphthylene group, a fluorenylene
group, a spiro-bifluorenylene group, a benzofluorenylene
group, a dibenzofluorenylene group, a phenanthrenylene
group, an anthracenylenylene group, a fluoranthenylenylene
group, a triphenylenylene group, a pyrenylene group, a chrysenylene
group, a perylenylene group, a pentaphenylenylene group, a
hexacenylenylene group, a pentacenylenylene group, a thiophe-
nylenylene group, a furanylenylene group, a carbazolylenylene group, an
indolylenylene group, an isoindolylenylene group, a benzofura-
nylenylene group, a benzothiophenylenylene group, a dibenzofura-
nylenylene group, a dibenzothiophenylenylene group, a benzocaba-
zolylenylene group, a dibenzocarbazolylenylene group, a

55

60

dibenzosilolylenylene group, a pyridinylenylene group, an imida-
zolylenylene group, a pyrazolylenylene group, a thiazolylenylene group,
an isothiazolylenylene group, an oxazolylenylene group, an isox-
azolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene
group, a pyrazinylenylene group, a pyrimidinylenylene group, a
pyridazinylenylene group, a triazininylenylene group, a quinolinylenylene
group, an isoquinolinylenylene group, a benzoquinolinylenylene group,
a phthalazinylenylene group, a naphthyridinylenylene group, a
quinoxalinylenylene group, a quinazolinylenylene group, a cinnol-
inylenylene group, a phenanthridinylenylene group, an acridinylenylene
group, a phenanthrolinylenylene group, a phenazinylenylene group,
a benzimidazolylenylene group, an isobenzothiazolylenylene group,
a benzoxazolylenylene group, an isobenzoxazolylenylene group, a
triazolylenylene group, a tetrazolylenylene group, an imidazopyrimi-
dinylenylene group, an imidazopyrimidinylenylene group, and an
azacarbazolylenylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylenylene group, a pentacenylenylene group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),

wherein Q_{31} to Q_{33} may be the same as described above.

In one or more embodiments, R_{301} to R_{304} in Formulae 301, 301-1, and 301-2 may each independently be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group,

a perylenyl group, a pentaphenyl group, a hexacenylenylene group, a pentacenylenylene group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylenylene group, a pentacenylenylene group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylenylene group, a pentacenylenylene group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),


81

linyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,


wherein Q_{31} to Q_{33} may be the same as described above.

In one or more embodiments, the host may include an alkaline-earth metal complex. For example, the host may be selected from a Be complex (for example, Compound H55), an Mg complex, and a Zn complex.


The host may include at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di(2-naphthyl)-2-t-butyl-anthracene (TBADN), a 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55, as examples:




H1 25



H2 35




H3 45




H4 60

82


-continued




H5




H6




H7



H8




H9



H10

83


-continued



H11


5

H12



15

H13




25

30

35

H14




40

45

50

55

H15




60


65

84


-continued




H16




H17



H18



H19




H20

85

-continued

H21



5


10

15

86

-continued

H24




H25

20

25


H22



30

35

40




H26

45

H23


50



55




60

65



87

-continued



88

-continued

H27

H30

5

10

15

20

25

H28

30

35

40

45

50

H29

55

60






65

H31

H32


H33

H34



89

-continued




H35

5

10

H36 15



20


25

30

35

H37

40




45

50

55

H38


60




65

90


-continued




H39



H40







H41



91

-continued

**92**

-continued

H42

H46

5

10

15

H43

20

H47

25

30

H48

35

H44

40

H49

45

H50

H45







50

55

H51





60

65



93

-continued



Phosphorescent Dopant Included in Emission Layer in Organic Layer 150

94

The phosphorescent dopant may include an organometallic complex represented by Formula 401 below:

H52

5

$$M(L_{401})_{xc1}(L_{402})_{xc2}, \text{ and}$$

<Formula 401>


10

<Formula 402>

H53

15

20



In Formulae 401 and 402,

M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

L_{401} may be selected from ligands represented by Formula 402, $xc1$ may be 1, 2, or 3, wherein, when $xc1$ is two or more, two or more L_{401} (s) may be identical to or different from each other,

H54

30

L_{402} may be an organic ligand, and $xc2$ may be an integer selected from 0 to 4, wherein, when $xc2$ is two or more, two or more L_{402} (s) may be identical to or different from each other,

35

X_{401} to X_{404} may each independently be nitrogen or carbon,

40

X_{401} and X_{403} may be linked via a single bond or a double bond, and X_{402} and X_{404} may be linked via a single bond or a double bond,

45

A_{401} and A_{402} may each independently be selected from a C_5 - C_{60} carbocyclic group or a C_1 - C_{60} heterocyclic group,

50

X_{405} may be a single bond, $*-O-*$, $*-S-*$, $*-C(=O)-*$, $*-N(Q_{411})-*$, $*-C(Q_{411})(Q_{412})-*$, $*-C(Q_{411})=C(Q_{412})-*$, $*-C(Q_{411})=*$, or $*=C(Q_{411})=*$, wherein Q_{411} and Q_{412} may be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

55

X_{406} may be a single bond, O, or S,

H55

60

R_{401} and R_{402} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_1 - C_{20} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})(Q_{402})$, $-B(Q_{401})(Q_{402})$, $-C(=O)(Q_{401})$, $-S(=O)_2$

95

(Q₄₀₁), and —P(=O)(Q₄₀₁)(Q₄₀₂), wherein Q₄₀₁ to Q₄₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a C₆-C₂₀ aryl group, and a C₁-C₂₀ heteroaryl group,

xc11 and xc12 may each independently be an integer selected from 0 to 10, and

* and *' in Formula 402 each indicate a binding site to M in Formula 401.

In one or more embodiments, A₄₀₁ and A₄₀₂ in Formula 402 may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

In one or more embodiments, in Formula 402, i) X₄₀₁ may be nitrogen, and X₄₀₂ may be carbon, or ii) X₄₀₁ and X₄₀₂ may each be nitrogen at the same time.

In one or more embodiments, R₄₀₁ and R₄₀₂ in Formula 402 may each independently be selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

96

—Si(Q₄₀₁)(Q₄₀₂)(Q₄₀₃), —N(Q₄₀₁)(Q₄₀₂), —B(Q₄₀₁)(Q₄₀₂), —C(=O)(Q₄₀₁), —S(=O)₂(Q₄₀₁), and —P(=O)(Q₄₀₁)(Q₄₀₂),


wherein Q₄₀₁ to Q₄₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group, as examples.

In one or more embodiments, when xc1 in Formula 401 is two or more, two A₄₀₁(s) in two or more L₄₀₁(s) may be optionally linked to each other via X₄₀₇, which is a linking group, or two A₄₀₂(s) in two or more L₄₀₁(s) may be optionally linked to each other via X₄₀₈, which is a linking group (see Compounds PD1 to PD4 and PD7). X₄₀₇ and X₄₀₈ may each independently be a single bond, *—O—*, *—S—*, *—C(=O)—*, *—N(Q₄₁₃)—*, *—C(Q₄₁₃)(Q₄₁₄)—*, or *—C(Q₄₁₃)=C(Q₄₁₄)—* (wherein Q₄₁₃ and Q₄₁₄ may each independently be hydrogen, deuterium, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group), as examples.


L₄₀₂ in Formula 401 may be a monovalent, divalent, or trivalent organic ligand. For example, L₄₀₂ may be selected from halogen, diketone (for example, acetylacetonate), carboxylic acid (for example, picolinate), —C(=O), isonitrile, —CN, and phosphorus (for example, phosphine or phosphite), as examples.

In one or more embodiments, the phosphorescent dopant may be selected from, for example, Compounds PD1 to PD25, as examples:


PD1



PD2









PD3



97

-continued




98

-continued

PD4

5


10



PD5

20

25




PD6

30

35

PD7

40




PD8

45

50


55



PD9

60

65



PD10

PD11






PD12

PD13

PD14

99

-continued


**100**

-continued

PD15

5

10



PD16


15

20

25

PD17

30




35

40

PD18

45

50




PD19

55

60

65



PD20



PD21

PD22

PD23

101


-continued



Fluorescent Dopant in Emission Layer

The fluorescent dopant may include an arylamine compound or a styrylamine compound.

The fluorescent dopant may include a compound represented by Formula 501 below.



In Formula 501,

Ar₅₀₁ may be a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₁-C₆₀ heterocyclic group,

L₅₀₁ to L₅₀₃ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

xd1 to xd3 may each independently be an integer of 0 to 3;

102

PD24

R₅₀₁ and R₅₀₂ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

xd4 may be an integer of 1 to 6.

In one or more embodiments, Ar₅₀₁ in Formula 501 may be selected from:

a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group; and

a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, L₅₀₁ to L₅₀₃ in Formula 501 may each independently be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, dibenzosilolylenylene group, and a pyridinylenylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, dibenzosilolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono

103

group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

In one or more embodiments, R₅₀₁ and R₅₀₂ in Formula 501 may each independently be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and


a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃),

wherein Q₃₁ to Q₃₃ may each be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.


In one or more embodiments, xd4 in Formula 501 may be 2, as an example.

104


For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:



FD1




FD2



FD3

105

-continued



FD4

5

10

15

20

25

FD5

30

35

40

45


FD6

50


55

60

65

**106**

-continued



FD7

5


10

15

20

25

FD8




30

35

40






45

FD9







107


-continued

**108**


-continued




109
-continued




110
-continued



In one or more embodiments, the fluorescent dopant may be selected from the following compounds, as examples.



DPVBi




DPAVB

111

112

-continued



40

Electron Transport Region in Organic Layer 150

The electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer, as examples.

For example, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, wherein, for each structure, constituting layers are sequentially stacked from an emission layer.

The electron transport region may include at least one of the heterocyclic compounds.

In one or more embodiments, the electron transport region may include an electron transport layer. The electron transport layer may include at least one of the heterocyclic compounds, as examples.

The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an

electron transport layer in the electron transport region) may include a metal-free compound containing at least one π electron-depleted nitrogen-containing ring.

The term “ π electron-depleted nitrogen-containing ring” indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

For example, the π electron-depleted nitrogen-containing ring may be i) a 5-membered to 7-membered hetero monocyclic group having at least one $*-N=*$ moiety, ii) a heteropoly cyclic group in which two or more 5-membered to 7-membered hetero monocyclic groups each having at least one $*-N=*$ moiety are condensed with each other, or iii) a heteropoly cyclic group in which at least one of 5-membered to 7-membered hetero monocyclic groups, each having at least one $*-N=*$ moiety, is condensed with at least one C_5 - C_{60} carbocyclic group.

Examples of the π electron-depleted nitrogen-containing ring include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, thiadiazol, an imidazopyridine, an imidazopyrimidine, and an azacabazole.

For example, the electron transport region may include a compound represented by Formula 601:



In Formula 601,

Ar_{601} may be a substituted or unsubstituted $\text{C}_5\text{-C}_{60}$ carbocyclic group or a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heterocyclic group,

xe11 may be 1, 2, or 3,

L_{601} may be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkylene group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenylene group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

xe1 may be an integer selected from 0 to 5,

R_{601} may be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryloxy group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylthio group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_{601})(\text{Q}_{602})(\text{Q}_{603})$, $-\text{C}(=\text{O})(\text{Q}_{601})$, $-\text{S}(=\text{O})_2(\text{Q}_{601})$, and $-\text{P}(=\text{O})(\text{Q}_{601})(\text{Q}_{602})$,

Q_{601} to Q_{603} may each independently be a $\text{C}_1\text{-C}_{10}$ alkyl group, a $\text{C}_1\text{-C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

xe21 may be an integer selected from 1 to 5.

In one or more embodiments, at least one of $\text{Ar}_{601}(\text{s})$ in the number of xe11 and/or at least one of $\text{R}_{601}(\text{s})$ in the number of xe21 may include the π electron-depleted nitrogen-containing ring.

In one or more embodiments, ring Ar_{601} in Formula 601 may be selected from:

a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzo-

fluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{20}$ alkyl group, a $\text{C}_1\text{-C}_{20}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,


wherein Q_{31} to Q_{33} may each independently be selected from a $\text{C}_1\text{-C}_{10}$ alkyl group, a $\text{C}_1\text{-C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

When xe11 in Formula 601 is two or more, two or more $\text{Ar}_{601}(\text{s})$ may be linked to each other via a single bond.

In one or more embodiments, Ar_{601} in Formula 601 may be an anthracene group.

In one or more embodiments, a compound represented by Formula 601 may be represented by Formula 601-1:

<Formula 601-1>



In Formula 601-1,

X_{614} may be N or $\text{C}(\text{R}_{614})$, X_{615} may be N or $\text{C}(\text{R}_{615})$, X_{616} may be N or $\text{C}(\text{R}_{616})$, and at least one selected from X_{614} to X_{616} may be N,

L_{611} to L_{613} may be each independently substantially the same as described in connection with L_{601} ,

xe611 to xe613 may be each independently substantially the same as described in connection with xe1 ,

R_{611} to R_{613} may be each independently substantially the same as described in connection with R_{601} ,

R_{614} to R_{616} may each independently be selected from hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{20}$ alkyl group, a $\text{C}_1\text{-C}_{20}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, L_{601} and L_{611} to L_{613} in Formulae 601 and 601-1 may each independently be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolininylenylene group, an isoquinolininylenylene group, a benzoquinolininylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinoxalinylenylene group, a cinnolininylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 – C_{20} alkyl group, a C_1 – C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an

indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, as examples.

In one or more embodiments, xel and xe611 to xe613 in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

In one or more embodiments, R₆₀₁ and R₆₁₁ to R₆₁₃ in Formula 601 and 601-1 may each independently be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl


117

group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and


—S(=O)₂(Q₆₀₁), and —P(=O)(Q₆₀₁)(Q₆₀₂),

wherein Q₆₀₁ and Q₆₀₂ are substantially the same as described above.

The electron transport region may include at least one compound selected from Compounds ET1 to ET36, as examples:




ET1




118

-continued

ET2






ET3





ET4

119
-continued




120
-continued



121

-continued



ET10

5

10

15

20

25


ET11

30

35

40

45




ET12

50

55


60

65


**122**

-continued


ET13



ET14






ET15



123

-continued

**124**

-continued


ET16

5

10

15

20




ET17

25

30

35

40



ET18


45

50

55

60

65




ET19

ET20

ET21

125
-continued

ET22



5


10

15

20


126
-continued

ET25



25

ET23




30

35

40


45

ET26



ET24

50




55


60

65

ET27



127
-continued



ET28


5

10

15

20

ET29




25

30

35

40

45



ET30


50

55


60

65


128
-continued



ET31






ET32



ET33

129



-continued





In one or more embodiments, the electron transport region may include at least one selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq₃, BAlq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ:

130

ET34



ET35



NTAZ

ET36

The thickness of the buffer layer, the hole blocking layer, or the electron control layer may each independently be in a range of about 20 Å to about 1,000 Å, or, for example, about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.


A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, or, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.


131

The metal-containing material may include at least one selected from an alkali metal complex and an alkaline earth-metal complex. The alkali metal complex may include a metal ion selected from an Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion. The alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, an Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenylan oxazole, a hydroxy phenylthiazole, a hydroxy diphenylan oxadiazole, a hydroxy diphenylthiadiazol, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene, as examples.

For example, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2:



ET-D1



ET-D2

The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode **190**. The electron injection layer may directly contact the second electrode **190**.

The electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

The electron injection layer may include alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof.

The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one or more embodiments, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs, as examples.

The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

The rare-earth metal may be selected from Sc, Y, Ce, Tb, Yb, and Gd.

The alkali metal compound, the alkaline earth-metal compound, and the rare-earth metal compound may be selected

132

from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal and rare-earth metal.

The alkali metal compound may be selected from alkali metal oxides, such as Li_2O , Cs_2O , or K_2O , and alkali metal halides, such as LiF, NaF, CsF, KF, LiI, NaI, CsI, or KI. In one or more embodiments, the alkali metal compound may be selected from LiF, Li_2O , NaF, LiI, NaI, CsI, and KI, as examples.

The alkaline earth-metal compound may be selected from alkaline earth-metal compounds, such as BaO, SrO, CaO, $\text{Ba}_x\text{Sr}_{1-x}\text{O}$ ($0 < x < 1$), or $\text{Ba}_x\text{Ca}_{1-x}\text{O}$ ($0 < x < 1$). In one or more embodiments, the alkaline earth-metal compound may be selected from BaO, SrO, and CaO, as examples.

The rare-earth metal compound may be selected from YbF_3 , ScF_3 , ScO_3 , Y_2O_3 , Ce_2O_3 , GdF_3 , and TbF_3 . In one or more embodiments, the rare-earth metal compound may be selected from YbF_3 , ScF_3 , TbF_3 , YbI_3 , ScI_3 , and TbI_3 , as examples.

The alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare-earth metal as described above, and a ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may each independently be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenylan oxazole, hydroxy phenylthiazole, hydroxy diphenylan oxadiazole, hydroxy diphenylthiadiazol, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene, as examples.

The electron injection layer may consist of alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof, as described above. In one or more embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

Second Electrode **190**

The second electrode **190** may be disposed on the organic layer **150** having such a structure. The second electrode **190** may be a cathode that is an electron injection electrode, and in this regard, a material for forming the second electrode **190** may be a material having a low work function, and such a material may be metal, alloy, an electrically conductive compound, or a combination thereof.

The second electrode **190** may include at least one selected from lithium (Li), silver (Si), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO, as examples. The second electrode **190**

133

may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

The second electrode 190 may have a single-layered structure, or a multi-layered structure including two or more layers.

Description of FIGS. 2 to 4

An organic light-emitting device 20 of FIG. 2 may include a first capping layer 210, a first electrode 110, an organic layer 150, and a second electrode 190 which are sequentially stacked in this stated order. An organic light-emitting device 30 of FIG. 3 may include a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220 which are sequentially stacked in this stated order. An organic light-emitting device 40 of FIG. 4 may include a first capping layer 210, a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220.

Regarding FIGS. 2 to 4, the first electrode 110, the organic layer 150, and the second electrode 190 may be understood by referring to the description presented in connection with FIG. 1.

In the organic layer 150 of each of the organic light-emitting devices 20 and 40, light generated in an emission layer may pass through the first electrode 110, which may be a semi-transmissive electrode or a transmissive electrode, and the first capping layer 210 toward the outside. In the organic layer 150 of each of the organic light-emitting devices 30 and 40, light generated in an emission layer may pass through the second electrode 190, which is a semi-transmissive electrode or a transmissive electrode, and the second capping layer 220 toward the outside.

The first capping layer 210 and the second capping layer 220 may increase external luminescent efficiency according to the principle of constructive interference.


The first capping layer 210 and the second capping layer 220 may each independently be an organic capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.

At least one selected from the first capping layer 210 and the second capping layer 220 may each independently include at least one material selected from a carbocyclic compound, a heterocyclic compound, an amine-based compound, a porphyrine derivative, a phthalocyanine derivative, a naphthalocyanine derivative, an alkali metal complex, and an alkaline earth-based complex. The carbocyclic compound, the heterocyclic compound, and the amine-based compound may be optionally substituted with a substituent containing at least one element selected from O, N, S, Se, Si, F, Cl, Br, and I. In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may independently include an amine-based compound.

In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may independently include the compound represented by Formula 201 or the compound represented by Formula 202.

In one or more embodiments, at least one selected from the first capping layer 210 and the second capping layer 220 may independently include a compound selected from Compounds HT28 to HT33 and Compounds CP1 to CP5, as examples.

134



Hereinbefore, the organic light-emitting device according to an embodiment has been described in connection with FIGS. 1-4. The organic light-emitting device may have other configurations.

Respective layers constituting the hole transport region, an emission layer, and respective layers constituting the electron transport region may be formed in a certain region by using one or more suitable methods selected from vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

When the respective layers of the hole transport region, the emission layer, and the respective layers of the electron transport region are formed by deposition, the deposition may be performed at a deposition temperature of about 100 to about 500° C., at a vacuum degree of about 10⁻⁸ to about 10⁻³ torr, and at a deposition rate of about 0.01 to about 100 Å/sec by taking into account a material for forming a layer to be deposited and the structure of a layer to be formed.

When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to 200° C., depending on a material to be included in a layer and the structure of each layer to be formed.

General Definition of Substituents

The term “C₁-C₆₀ alkyl group” as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and non-limiting examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “C₁-C₆₀ alkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₆₀ alkyl group.

The term “C₂-C₆₀ alkenyl group” as used herein refers to a hydrocarbon group formed by substituting at least one carbon-carbon double bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and non-limiting examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C₂-C₆₀ alkenylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkenyl group.

The term “C₂-C₆₀ alkynyl group” as used herein refers to a hydrocarbon group formed by substituting at least one carbon-carbon triple bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and non-limiting examples thereof include an ethynyl group and a propynyl group. The term “C₂-C₆₀ alkynylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

The term “C₁-C₆₀ alkoxy group” as used herein refers to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is the C₁-C₆₀ alkyl group), and non-limiting examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

The term “C₃-C₁₀ cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and non-limiting examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

The term “C₁₀-C₁₀ heterocycloalkyl group” as used herein refers to a monovalent saturated monocyclic group

having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and non-limiting examples thereof include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuranyl group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

The term “C₃-C₁₀ cycloalkenyl group” as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and does not have aromaticity, and non-limiting examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

The term “C₁-C₁₀ heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Non-limiting examples of the C₁-C₁₀ heterocycloalkenyl group include a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranyl group, and a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkenyl group.

The term “C₆-C₆₀ aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and the term “C₆-C₆₀ arylene group” as used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Non-limiting examples of the C₆-C₆₀ aryl group include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be fused to each other.

The term “C₁-C₆₀ heteroaryl group” as used herein refers to a monovalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C₁-C₆₀ heteroarylene group” as used herein refers to a divalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Non-limiting examples of the C₁-C₆₀ heteroaryl group include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be fused to each other.

The term “C₆-C₆₀ aryloxy group” as used herein refers to —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and a C₆-C₆₀ arylthio group used herein indicates —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed with each other, only carbon atoms as a ring-forming atom, and non-aromaticity in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term “divalent non-aromatic condensed polycy-

clic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group (for example, having 1 to 60 carbon atoms) that has two or more rings condensed to each other, has at least one heteroatom selected from N, O, Si, P, and S, other than carbon atoms, as a ring-forming atom, and has non-aromaticity in the entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

The term “C₅-C₆₀ carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which a ring-forming atom is a carbon atom only. The C₅-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₅-C₆₀ carbocyclic group may be a ring, such as a benzene, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In one or more embodiments, depending on the number of substituents connected to the C₅-C₆₀ carbocyclic group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

The term “C₁-C₆₀ heterocyclic group” as used herein refers to a group having the same structure as the C₁-C₆₀ carbocyclic group, except that as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used in addition to carbon (the number of carbon atoms may be in a range of 1 to 60).

At least one of substituents of the substituted C₅-C₆₀ carbocyclic group, substituted C₁-C₆₀ heterocyclic group, substituted C₃-C₁₀ cycloalkylene group, substituted C₁-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₁-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₁-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, a substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

deuterium (-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group,

—Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

—Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

wherein Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

The term “Ph”, as used herein, may refer to a phenyl group; the term “Me”, as used herein, may refer to a methyl group; the term “Et”, as used herein, may refer to an ethyl group; the term “ter-Bu” or “But”, as used herein, may refer to a tert-butyl group; and the term “OMe” as used herein may refer to a methoxy group.

The term “biphenyl group” as used therein refers to “a phenyl group substituted with a phenyl group.” In other words, a “biphenyl group” is a substituted phenyl group having a C₆-C₆₀ aryl group as a substituent.

The term “terphenyl group” as used herein refers to “a phenyl group substituted with a biphenyl group.” In other words, a “terphenyl group” is a substituted phenyl group having a C₆-C₆₀ aryl group substituted with a C₆-C₆₀ aryl group as a substituent

* and * used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.


The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the

139

scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples. The wording "B was used instead of A" used in describing Synthesis Examples indicates that an identical molar equivalent of B was used in place of A.

EXAMPLE

Synthesis Example 1: Synthesis of Compound 1



Synthesis of Intermediate I-1

1.08 g (10 mmol) of o-phenylene diamine and 1.06 g (10 mmol) of cyanogen bromide were dissolved in 100 mL of MeOH and were then stirred at room temperature for 24 hours. An ammonia solution was added thereto, and an extraction process was performed thereon three times by using 60 mL of ethyl acetate. An organic layer collected therefrom was dried by using MgSO_4 and a solvent was removed therefrom by evaporation. The residue obtained

140

therefrom was separated and purified by silica gel column chromatography, thereby completing the preparation of 1.06 g (80%) of Intermediate I-1. The obtained compound was identified by LC-MS.

$\text{C}_7\text{H}_7\text{N}_3$; M^+ 133.06

Synthesis of Intermediate I-2

1.33 g (10 mmol) of Intermediate I-1, 2.23 g (10 mmol) of 1-bromonaphthalene, 2.89 g (1.5 equiv) of CsCO_3 , 20 mg (10 mol %) of CuI , and 200 mg (15 mol %) of 8-quinolinol were dissolved in 100 mL of t-BuOH and were then stirred at a temperature of 90° C. for 18 hours. The obtained result was cooled to room temperature, and an extraction process was performed thereon three times by using 60 mL of ethyl acetate. An organic layer obtained therefrom was dried by using MgSO_4 and a solvent was removed therefrom by evaporation. The residue obtained therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 2.02 g (78%) Intermediate I-2. The obtained compound was identified by LC-MS.

$\text{C}_{17}\text{H}_{13}\text{N}_3$; M^+ 259.10

Synthesis of Intermediate I-3

2.59 g (10 mmol) of Intermediate I-2, 1.73 g (11 mmol) of bromobenzene, 0.1 mol % of $\text{Pd}_2(\text{dba})_3$, and 3.18 g (1.5 equiv) of K_3PO_4 were dissolved in 100 mL of t-BuOH and were then stirred at a temperature of 120° C. for 5 hours. The obtained result was cooled to room temperature, and an extraction process was performed thereon three times by using 60 mL of ethyl acetate. An organic layer collected therefrom was dried by using MgSO_4 and a solvent was removed therefrom by evaporation. The residue obtained therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 2.58 g (77%) of Intermediate I-3. The obtained compound was identified by LC-MS.

$\text{C}_{23}\text{H}_{17}\text{N}_3$; M^+ 335.15

Synthesis of Compound 1

3.35 g (10 mmol) of Intermediate I-3, $\text{Cu}(\text{OAc})_2$, PPh_3 , 1,10-phenanthroline, and NaOAc were dissolved and stirred at a temperature of 80° C. for 12 hours. The obtained residue was cooled to room temperature, and an extraction process was performed thereon three times by using 30 mL of water and 30 mL of ethyl acetate. An organic layer obtained therefrom was dried by using magnesium sulfate and a solvent was removed therefrom by evaporation. Then, the residue obtained therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 2.33 g (70%) of Compound 1. The obtained compound was identified by MS/FAB and ^1H NMR.

Synthesis Example 2: Synthesis of Compound 4

2.09 g (63%) of Compound 4 was obtained in the same manner as in Synthesis of Compound 1, except that 3-bromopyridine was used instead of bromobenzene in synthesizing Intermediate I-3. The obtained compound was identified by MS/FAB and ^1H NMR.

Synthesis Example 3: Synthesis of Compound 12

2.14 g (58%) of Compound 12 was obtained in the same manner as in Synthesis of Compound 1, except that

141

3-bromo-2,6-difluoropyridine was used instead of bromobenzene in synthesizing Intermediate I-3. The obtained compound was identified by MS/FAB and ^1H NMR.

Synthesis Example 4: Synthesis of Compound 15

4.19 g (70%) of Compound 15 was obtained in the same manner as in Synthesis of Compound 1, except that 6-(3-bromophenyl)dinaphtho[2,1-b:1',2'-d]furan was used instead of bromobenzene in synthesizing Intermediate I-3. The obtained compound was identified by MS/FAB and ^1H NMR.

Synthesis Example 5: Synthesis of Compound 30

4.19 g (70%) of Compound 30 was obtained in the same manner as in Synthesis of Compound 1, except that 3-(3-bromophenyl)-2,6-difluoropyridine was used instead of bromobenzene in synthesizing Intermediate I-3. The obtained compound was identified by MS/FAB and ^1H NMR.

Synthesis Example 6: Synthesis of Compound 31

2.37 g (71%) of Compound 31 was obtained in the same manner as in Synthesis of Compound 1, except that 5-bromoquinoline was used instead of 1-bromonaphthalene in synthesizing Intermediate I-2. The obtained compound was identified by MS/FAB and ^1H NMR.

Synthesis Example 7: Synthesis of Compound 37

2.24 g (68%) Compound 37 was obtained in the same manner as in Synthesis of Compound 1, except that 4-bromoquinoline was used instead of 1-bromonaphthalene in synthesizing Intermediate I-2. The obtained compound was identified by MS/FAB and ^1H NMR.

142

Synthesis Example 8: Synthesis of Compound 40

3.67 g (64%) of Compound 2 was obtained in the same manner as in Synthesis of Compound 1, except that 4-bromodibenzo[b,d]furan was used instead of 1-bromonaphthalene in synthesizing Intermediate I-2, and (3-bromophenyl)diphenylphosphine oxide was used instead of bromobenzene in synthesizing Intermediate I-3. The obtained compound was identified by MS/FAB and ^1H NMR.

Synthesis Example 9: Synthesis of Compound 41

3.17 g (71%) of Compound 41 was obtained in the same as in Synthesis of Compound 1, except that 4-bromoquinoline was used instead of 1-bromonaphthalene in synthesizing Intermediate I-2, and 3-(4-bromophenyl)-2,6-difluoropyridine was used instead of bromobenzene in synthesizing Intermediate I-3. The obtained compound was identified by MS/FAB and ^1H NMR.

Synthesis Example 10: Synthesis of Compound 50

3.60 g (61%) of Compound 50 was obtained in the same manner as in Synthesis of Compound 1, except that 4-bromodibenzo[b,d]furan was used instead of 1-bromonaphthalene in synthesizing Intermediate I-2, and 7-(6-bromopyridin-2-yl)-9,9-dimethyl-9H-fluorene-2-carbonitrile was used instead of bromobenzene in synthesizing Intermediate I-3. The obtained compound was identified by MS/FAB and ^1H NMR.

^1H NMR and MS/FAB of Compounds synthesized in Synthesis Examples 1 to 10 are shown in Table 1 below.

Methods of synthesizing compounds other than Compounds shown in Table 1 are recognizable by one of ordinary skill in the art by referring to the synthesis paths and source materials described above.

TABLE 1


Compound	^1H NMR (CDCl_3 , 500 MHz)	HR-EIMS	
		found	calc.
1	δ = 8.32-8.38 (m, 1H), 8.08-7.97 (m, 2H), 7.60-7.35 (m, 11H)	333.15	333.13
4	δ = 8.66-8.60 (m, 2H), 8.35-8.24 (m, 2H), 8.06-8.02 (m, 2H), 7.94-7.92 (d, 1H), 7.53-7.45 (m, 6H), 7.30-7.35 (m, 1H)	334.14	334.12
12	δ = 8.35-8.31 (d, 1H), 8.06-8.00 (m, 4H), 7.55-7.33 (m, 6H), 7.94-7.92 (d, 1H), 7.53-7.45 (m, 6H), 6.94-6.92 (d, 1H)	370.11	370.10
15	δ = 8.84-8.82 (m, 2H), 8.45-8.35 (m, 2H), 8.06-8.01 (m, 2H), 7.94-7.88 (m, 5H), 7.73-7.71 (m, 2H), 7.62-7.45 (m, 10H), 7.35-7.29 (d, 1H)	599.21	599.20
30	δ = 8.64-8.56 (m, 2H), 8.42-8.38 (m, 3H), 7.79 (s, 1H), 7.70-7.69 (d, 2H), 7.53-7.51 (m, 3H), 7.50-7.35 (m, 6H), 6.68 (s, 1H)	496.17	496.15
31	δ = 9.29 (s, 1H), 8.24 (s, 1H), 7.95 (s, 1H), 7.77 (s, 1H), 7.63-7.52 (m, 6H), 7.48-7.40 (m, 4H)	334.11	334.12
37	δ = 9.18 (s, 1H), 8.34 (s, 1H), 8.23 (s, 1H), 7.82 (s, 1H), 7.69-7.60 (m, 6H), 7.47-7.35 (m, 4H)	334.11	334.12
40	δ = 8.15-8.17 (d, 1H), 7.96-7.98 (d, 1H), 7.87-7.84 (d, 1H), 7.82-7.80 (m, 1H), 7.73-7.69 (m, 6H), 7.63-7.55 (m, 3H), 7.49-7.38 (m, 11H)	573.17	573.16
41	δ = 9.24 (s, 1H), 8.37-8.34 (m, 2H), 8.24-8.22 (d, 1H), 7.84-7.80 (m, 1H), 7.68-7.67 (m, 3H), 7.60-7.58 (d, 1H), 7.50-7.45 (m, 4H), 7.37-7.30 (m, 1H), 6.76-6.73 (m, 12H)	447.15	447.13
50	δ = 8.33-8.31 (d, 1H), 8.19-8.16 (d, 2H), 7.97-7.95 (m, 3H), 7.88-7.78 (m, 5H), 7.66-7.58 (m, 3H), 7.50-7.38 (m, 6H), 1.65 (s, 6H)	591.20	591.21

143


Example 1

An anode was prepared by cutting an ITO glass substrate (manufactured by Corning), on which an ITO layer was deposited to a thickness of $15 \Omega/\text{cm}^2$ (1,200 Å), to a size of 50 mm×50 mm×0.7 mm, ultrasonically cleaning the ITO glass substrate (anode) using isopropyl alcohol and pure water each for 5 minutes, and then, exposing the ITO glass substrate (anode) to UV irradiation for 30 minutes and ozone to clean the ITO glass substrate. Then, the ITO glass substrate (anode) was loaded into a vacuum deposition apparatus.


2-TNATA was vacuum-deposited on the ITO glass substrate (anode) to form a hole injection layer having a



2-TNATA



ADN




DPAVBi

144

thickness of 600 Å, and NPB was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å, thereby forming a hole transport region.

AND (host) and DPAVBi (dopant) were co-deposited on the hole transport layer at a weight ratio of 98:2 to form an emission layer having a thickness of 300 Å.

Then, Compound 1 was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å, LiF was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited on the electron injection layer to form a cathode having a thickness of 3,000 Å, thereby completing the manufacture of an organic light-emitting device.







NPB

145

Examples 2 to 10 and Comparative Examples 1 to






4

Organic light-emitting devices of Examples 2 to 10 and Comparative Examples 1 to 4 were manufactured in the same manner as in Example 1, except that Compounds shown in Table 2 were each used in forming an electron transport layer.




146

-continued



147

-continued



50


5

10

15

20

A




25

30

35

40

B




45

50


55

148

-continued



C



D

TABLE 2

	Material for electron transport layer	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
Example 1	Compound 1	4.33	50	3,025	6.05	Blue	242 hr
Example 2	Compound 4	4.21	50	3,115	6.23	Blue	265 hr
Example 3	Compound 12	4.03	50	3,285	6.57	Blue	223 hr

TABLE 2-continued

	Material for electron transport layer	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
Example 4	Compound 15	4.19	50	3,165	6.33	Blue	277 hr
Example 5	Compound 30	4.03	50	3,330	6.66	Blue	215 hr
Example 6	Compound 31	4.14	50	3,205	6.41	Blue	223 hr
Example 7	Compound 37	4.11	50	3,155	6.31	Blue	285 hr
Example 8	Compound 40	4.07	50	3,375	6.75	Blue	220 hr
Example 9	Compound 41	4.02	50	3,315	6.63	Blue	205 hr
Example 10	Compound 50	4.15	50	3,070	6.14	Blue	210 hr
Comparative Example 1	A	4.50	50	2,325	4.65	Blue	70 hr
Comparative Example 2	B	4.39	50	2,685	5.37	Blue	150 hr
Comparative Example 3	C	4.94	50	2,275	4.55	Blue	57 hr
Comparative Example 4	D	4.59	50	2,425	4.85	Blue	83 hr


Referring to Table 2, it can be seen that the organic light-emitting devices of Examples 1 to 10 had a high driving voltage, high brightness, high efficiency, and a long half lifespan, compared to the devices of Comparative Examples 1 to 4.

According to one or more embodiments, an organic light-emitting device including the heterocyclic compound may have a low driving voltage, high luminance, high efficiency, and a long lifespan.

Example embodiments have been disclosed herein, and although specific terms are employed, they are used and are to be interpreted in a generic and descriptive sense only and not for purpose of limitation. In some instances, as would be apparent to one of ordinary skill in the art as of the filing of the present application, features, characteristics, and/or elements described in connection with a particular embodiment may be used singly or in combination with features, characteristics, and/or elements described in connection with other embodiments unless otherwise specifically indicated. Accordingly, it will be understood by those of skill in the art that various changes in form and details may be made without departing from the spirit and scope of the present invention as set forth in the following claims.

What is claimed is:

1. A heterocyclic compound represented by Formula 1:



wherein, in Formula 1,

ring A₁ is a C₁₀-C₆₀ carbocyclic group or a C₆-C₆₀ heterocyclic group,

L₁ is selected from a substituted or unsubstituted C₃-C₆₀ carbocyclic group, a substituted or unsubstituted C₁-C₆₀ heterocyclic group, *—S(=O)—*, *—S(=O)₂—*, *—P(=O)(Q₄)—*, and *—P(=S)(Q₄)—*,

a₁ is an integer selected from 0 to 5, wherein, when a₁ is two or more, two or more L₁(s) are identical to or different from each other,

Ar₁ is selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —S(=O)(Q₁), —S(=O)₂(Q₁), —P(=O)(Q₁)(Q₂), and —P(=S)(Q₁)(Q₂),

b₁ is an integer selected from 1 to 5, wherein, when b₁ is two or more, two or more Ar₁(s) are identical to or different from each other,

R₁ to R₄ and R₁₁ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂),

c₁₁ is an integer selected from 0 to 6, wherein, when c₁₁ is two or more, two or more R₁₁(s) are identical to or different from each other,

151

at least one substituent selected from a substituent(s) of the substituted C₁₀-C₆₀ carbocyclic group, the substituted C₆-C₆₀ heterocyclic group, the substituted C₃-C₆₀ carbocyclic group, the substituted C₁-C₆₀ heterocyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl

152

group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

—Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

Q₁ to Q₄, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

* and *' each indicate a binding site to a neighboring atom.

2. The heterocyclic compound as claimed in claim 1, wherein:


ring A₁ is a C₁₀-C₆₀ carbocyclic group with two or more rings condensed with each other, or a C₆-C₆₀ heterocyclic group with two or more rings condensed with each other.

3. The heterocyclic compound as claimed in claim 1, wherein:


ring A₁ is a naphthalene, a fluorene, a benzofluorene, a phenanthrene, an anthracene, a quinoline, an isoquinoline, a benzoquinoline, a carbazole, a dibenzofuran, a dibenzothiophene, or a benzocarbazole.

4. The heterocyclic compound as claimed in claim 1, wherein:


L₁ is selected from groups represented by Formulae 3-1 to 3-51:




Formula 3-1



Formula 3-2












Formula 3-3



Formula 3-4

153


-continued




154

-continued


Formula 3-5




Formula 3-6




Formula 3-7




Formula 3-8




Formula 3-9



Formula 3-10




Formula 3-11



Formula 3-12



Formula 3-13



65

Formula 3-14

Formula 3-15

Formula 3-16







Formula 3-17

Formula 3-18

Formula 3-19

155


-continued




156

-continued


Formula 3-20




Formula 3-21




Formula 3-22





Formula 3-23



Formula 3-24



Formula 3-25



Formula 3-26

Formula 3-27

Formula 3-28

Formula 3-29

Formula 3-30

Formula 3-31

Formula 3-32

Formula 3-33










Formula 3-34

Formula 3-35

Formula 3-36

157


-continued



158

-continued


Formula 3-37



Formula 3-38


10 Formula 3-47

Formula 3-39




Formula 3-40

15 Formula 3-48




Formula 3-41

20 Formula 3-49




Formula 3-42

25 Formula 3-50



Formula 3-43

30 Formula 3-51



Formula 3-44

45 wherein, in Formulae 3-1 to 3-51,
Y₁ is O, S, C(Z₃)(Z₄), N(Z₃), or Si(Z₃)(Z₄),
Z₁ to Z₄ are each independently selected from hydrogen,

50 deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkenyl group, a C₁-C₂₀ alkynyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

Formula 3-45

55 wherein Q₃₁ to Q₃₃ are each independently selected from
60 a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

159

d2 is 1 or 2,

d3 is an integer selected from 1 to 3,

d4 is an integer selected from 1 to 4,

d5 is an integer selected from 1 to 5,

d6 is an integer selected from 1 to 6,

d8 is an integer selected from 1 to 8, and

* and *' each indicate a binding site to a neighboring atom.

5. The heterocyclic compound as claimed in claim 1, wherein:

L₁ is selected from:

a benzene group, a fluorene group, an anthracene group, a pyridine group, a triazine group, and a carbazole group;

a benzene group, a fluorene group, an anthracene group, a pyridine group, a triazine group, and a carbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), and —B(Q₃₁)(Q₃₂); and

—S(=O)—', *—S(=O)₂—*', *—P(=O)(Q₄)—*', and *—P(=S)(Q₄)—*'

Q₄ and Q₃₁ to Q₃₃ are each independently selected from a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, and


* and *' each indicate a binding site to a neighboring atom.

6. The heterocyclic compound as claimed in claim 1, wherein:


a1 is an integer selected from 0 to 3.

7. The heterocyclic compound as claimed in claim 1, wherein:


Ar₁ is selected from groups represented by Formulae 5-1 to 5-29, —S(=O)(Q₁), —S(=O)₂(Q₁), —P(=O)(Q₁)(Q₂), and —P(=S)(Q₁)(Q₂):




Formula 5-1




Formula 5-2



Formula 5-3



Formula 5-4




Formula 5-5


160

-continued


Formula 5-6




Formula 5-7




Formula 5-8




Formula 5-9




Formula 5-10




Formula 5-11




Formula 5-12




Formula 5-13



Formula 5-14










Formula 5-15



161


-continued

**162**

-continued

Formula 5-16

5



Formula 5-17

10

Formula 5-18

15

20

25

Formula 5-19

30

Formula 5-20

35

40

Formula 5-21

50

55

Formula 5-22

60

65

Formula 5-23






Formula 5-24

Formula 5-25

Formula 5-26


Formula 5-27

Formula 5-28



163

-continued



wherein, in Formulae 5-1 to 5-29,

Y₃₁ is O, S, C(Z₃₅)(Z₃₆), N(Z₃₅), or Si(Z₃₅)(Z₃₆),

Z₃₁ to Z₃₆ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkenyl group, a C₁-C₂₀ alkynyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

wherein Q₁, Q₂, and Q₃₁ to Q₃₃ are each independently selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

e2 is 1 or 2,

e3 is an integer selected from 1 to 3,

e4 is an integer selected from 1 to 4,

e5 is an integer selected from 1 to 5,

e6 is an integer selected from 1 to 6,

e7 is an integer selected from 1 to 7,

e9 is an integer selected from 1 to 9, and

* and *' each indicate a binding site to a neighboring atom.

8. The heterocyclic compound as claimed in claim 1, wherein:

Ar₁ is selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, a pyridinyl group, a dinaphthofuranyl group, and a benzoimidazonaphthoimidazolyl group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, a pyridinyl group, a dinaphthofuranyl group, and a benzoimidazonaphthoimidazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀

164

alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, —N(Q₃₁)(Q₃₂), and —Si(Q₃₁)(Q₃₂)(Q₃₃); and

5 —S(=O)(Q₁), —S(=O)₂(Q₁), —P(=O)(Q₁)(Q₂), and —P(=S)(Q₁)(Q₂),

wherein Q₁, Q₂, and Q₃₁ to Q₃₃ are each independently selected from a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

9. The heterocyclic compound as claimed in claim 1, wherein:

15 b1 is 1 or 2.

10. The heterocyclic compound as claimed in claim 1, wherein:

R₁ to R₄ are each independently selected from:

20 hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, and a nitro group;

a phenyl group, a biphenyl group, a terphenyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyridinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

a phenyl group, a biphenyl group, a terphenyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyridinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, —N(Q₃₁)(Q₃₂), and —Si(Q₃₁)(Q₃₂)(Q₃₃); and

—Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂),

50 R₁₁ is selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, and a nitro group; and


60 —Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂),

wherein Q₁ to Q₃ and Q₃₁ to Q₃₃ are each independently selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

165

11. The heterocyclic compound as claimed in claim 1, wherein:


the compound represented by Formula 1 is represented by one selected from Formulae 1A to 1J:




166

-continued


1F




1G




1H



1I



1J



wherein, in Formulae 1A to 1J,

L_1 , a_1 , Ar_1 , b_1 , and R_1 to R_4 are the same as described in claim 1,

Y_{51} is O, S, $C(Z_{53})(Z_{54})$, $N(Z_{53})$, or $Si(Z_{53})(Z_{54})$,

Y_{52} to Y_{57} are each independently $C(Z_{55})$ or N,

Z_{51} to Z_{55} are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a

167

cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkenyl group, a C₁-C₂₀ alkynyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, 5

f2 is 1 or 2,
 f4 is an integer selected from 1 to 4,
 f6 is an integer selected from 1 to 6, and
 f8 is an integer selected from 1 to 8.

12. The heterocyclic compound as claimed in claim 11, 10
 wherein,
 in Formulae 1A and 1J,
 L₁ is selected from:
 a benzene group, a fluorene group, an anthracene group, 15
 a pyridine group, a triazine group, and a carbazole group;
 a benzene group, a fluorene group, an anthracene group,
 a pyridine group, a triazine group, and a carbazole 20
 group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), 25
 —N(Q₃₁)(Q₃₂), and —B(Q₃₁)(Q₃₂); and
 —S(=O)—[†], *—S(=O)₂—*[†], *—P(=O)(Q₄)—*[†],
 and *—P(=S)(Q₄)—*[†],
 a1 is an integer selected from 0 to 3,
 Ar₁ is selected from:
 a phenyl group, a biphenyl group, a terphenyl group, a 30
 naphthyl group, a fluorenyl group, a phenanthrenyl group, a pyridinyl group, dinaphthofuranyl group, and a benzoimidazonaphthoimidazolyl group;
 a phenyl group, a biphenyl group, a terphenyl group, a 35
 naphthyl group, a fluorenyl group, a phenanthrenyl group, a pyridinyl group, a dinaphthofuranyl group, and a benzoimidazonaphthoimidazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ 40
 alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, —N(Q₃₁)(Q₃₂), and —Si(Q₃₁)(Q₃₂)(Q₃₃); and
 —S(=O)(Q₁), —S(=O)₂(Q₁), —P(=O)(Q₁)(Q₂), and 45
 —P(=S)(Q₁)(Q₂),
 b1 is 1 or 2,
 R₁ to R₄ are each independently selected from:
 hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group; 50
 a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, and a nitro group;
 a phenyl group, a biphenyl group, a terphenyl group, a 55
 fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyridinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;
 a phenyl group, a biphenyl group, a terphenyl group, a 60
 fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyridinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group,

168

a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, —N(Q₃₁)(Q₃₂), and —Si(Q₃₁)(Q₃₂)(Q₃₃); and
 —Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂),
 —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂),
 wherein Q₁ to Q₄ and Q₃₁ to Q₃₃ are each independently selected from a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

13. An organic light-emitting device, comprising:
 a first electrode;
 a second electrode facing the first electrode; and
 an organic layer between the first electrode and the second electrode, the organic layer comprising an emission layer,
 wherein the organic layer comprises at least one of the heterocyclic compounds as claimed in claim 1.

14. The organic light-emitting device as claimed in claim 13, wherein
 the first electrode is an anode,
 the second electrode is a cathode, and
 the organic layer further comprises a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode,
 the hole transport region comprises a hole injection layer, a hole transport layer, an emission auxiliary layer, an electron blocking layer, or any combination thereof, and
 the electron transport region comprises a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, an electron injection layer, or any combination thereof.

15. The organic light-emitting device as claimed in claim 14, wherein:
 the electron transport region comprises the at least one of the heterocyclic compounds.

16. The organic light-emitting device as claimed in claim 14, wherein:
 the electron transport region comprises an electron transport layer, and
 the electron transport layer comprises the at least one of the heterocyclic compounds.

17. The organic light-emitting device as claimed in claim 16, wherein
 the electron transport layer further comprises a metal-containing material.

18. The organic light-emitting device as claimed in claim 17, wherein the metal-containing material comprises a Li complex.

19. The organic light-emitting device as claimed in claim 14, wherein
 the hole transport region comprises a p-dopant, and
 a lowest unoccupied molecular orbital (LUMO) level of the p-dopant is about -3.5 eV or less.

20. The organic light-emitting device as claimed in claim 19, wherein
 the p-dopant comprises a quinone derivative, a metal oxide, a cyano group-containing compound, or any combination thereof.

* * * * *

专利名称(译)	杂环化合物和包括其的有机发光装置		
公开(公告)号	US10069085	公开(公告)日	2018-09-04
申请号	US15/425351	申请日	2017-02-06
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	LEE HYOYOUNG KIM YOUNGKOOK PARK JUNHA SIM MUNKI JEONG EUNJAE HWANG SEOKHWAN		
发明人	LEE, HYOYOUNG KIM, YOUNGKOOK PARK, JUNHA SIM, MUNKI JEONG, EUNJAE HWANG, SEOKHWAN		
IPC分类号	H01L51/50 C07F9/6561 H01L51/00 C07D471/14 C07D487/04 C07D491/147		
CPC分类号	H01L51/0072 C07D471/14 C07D487/04 C07D491/147 C07F9/6561 H01L51/0052 H01L51/0071 H01L51/0073 H01L51/0077 H01L51/5004 H01L51/0067 H01L2251/552 H01L51/506 H01L51/5072		
审查员(译)	CLARK , GREGORYð		
优先权	1020160090775 2016-07-18 KR		
其他公开文献	US20180019406A1		
外部链接	Espacenet		

摘要(译)

杂环化合物由式1表示：其中，在式1中，R₁至R₄，R₁₁，环A₁，L₁，Ar₁，c₁₁，a₁和b₁如说明书中所定义。

