



US 20150333273A1

(19) **United States**

(12) **Patent Application Publication**
LEE et al.

(10) **Pub. No.: US 2015/0333273 A1**

(43) **Pub. Date: Nov. 19, 2015**

(54) **CONDENSED CYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

(30) **Foreign Application Priority Data**

May 15, 2014 (KR) 10-2014-0058395

(71) Applicants: **Samsung Electronics Co., Ltd.**,
Suwon-si (KR); **Samsung SDI Co., Ltd.**,
Yongin-si (KR)

Publication Classification

(51) **Int. Cl.**
H01L 51/00 (2006.01)
C07D 519/00 (2006.01)
C07D 487/06 (2006.01)

(72) Inventors: **Sunyoung LEE**, Seoul (KR); **Soyeon KIM**, Anyang-si (KR); **Yoonhyun KWAK**, Seoul (KR); **Sangdong KIM**, Hwaseong-si (KR); **Sangyeob LEE**, Hwaseong-si (KR)

(52) **U.S. Cl.**
CPC *H01L 51/0072* (2013.01); *C07D 487/06* (2013.01); *C07D 519/00* (2013.01); *H01L 51/0067* (2013.01); *H01L 51/5012* (2013.01)

(21) Appl. No.: **14/599,724**

(57) **ABSTRACT**

A condensed cyclic compound represented by one of Formulae 1A to 1C, wherein Formulae 1A to 1C are disclosed in the detailed description.

(22) Filed: **Jan. 19, 2015**





FIG. 1



**CONDENSED CYCLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**

CROSS-REFERENCE TO RELATED
APPLICATION

[0001] This application claims priority to Korean Patent Application No. 10-2014-0058395, filed on May 15, 2014, in the Korean Intellectual Property Office, and all the benefits accruing therefrom under 35 U.S.C. §119, the content of which is incorporated herein in its entirety by reference.

BACKGROUND

[0002] 1. Field

[0003] One or more embodiments relate to condensed cyclic compounds and organic light-emitting devices including the same.

[0004] 2. Description of the Related Art

[0005] Organic light-emitting devices (OLEDs) are self-emission devices that have wide viewing angles, high contrast ratios, and short response times. In addition, the OLEDs exhibit excellent brightness, driving voltage, and response speed characteristics, and produce full-color images.

[0006] A typical organic light-emitting device includes an anode, a cathode, and an organic layer that is disposed between the anode and the cathode and includes an emission layer. A hole transport region may be disposed between the anode and the emission layer, and an electron transport region may be disposed between the emission layer and the cathode. Holes provided from the anode may move toward the emission layer through the hole transport region, and electrons provided from the cathode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, are recombined in the emission layer to produce excitons. These excitons change from an excited state to a ground state, thereby generating light.


[0007] Various types of organic light emitting devices are known. However, there still remains a need in OLEDs having low driving voltage, high efficiency, high brightness, and long lifespan.

SUMMARY

[0008] Provided are novel condensed cyclic compounds and organic light-emitting devices including the same.






[0009] Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments.

[0010] According to an aspect, a condensed cyclic compound is represented by one of Formulae 1A to 1C:




-continued

Formula 1B




-continued

Formula 2E



Formula 2F



[0011] wherein in the formulae,

[0012] Ring A is a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, a group represented by Formula 2A, or a group represented by Formula 2B;

[0013] Ring B₁ is a C₅-C₂₀ carbocyclic group or a C₂-C₂₀ heterocyclic group that does not include a nitrogen atom as a ring-forming atom;

[0014] Ring B₂, Ring B₃, Ring C₁, and Ring C₂ are each independently a C₅-C₂₀ carbocyclic group, or a C₂-C₂₀ heterocyclic group, provided that at least one selected from Ring B₂ and Ring B₃ is a C₂-C₂₀ heterocyclic group including a nitrogen atom as a ring-forming atom, and at least one selected from Ring C₁ and Ring C₂ is a C₂-C₂₀ heterocyclic group including a nitrogen atom as a ring-forming atom;

[0015] Pc₁ and Pc₂ in Formula 1A and 1B are each independently a group represented by Formulae 2A or a group represented by Formula 2B;

[0016] Cz₁, Cz₂, and Cz₃ in Formula 1C are each independently a group represented by Formula 2C, a group represented by Formula 2D, a group represented by Formula 2E, or a group represented by Formula 2F;

[0017] c₁, c₂, c₁₁, c₁₂, c₁₃, p₁, p₂, p₁₁, p₁₂, and p₁₃ are each independently an integer selected from 0 to 3;

[0018] when c₁ and c₂ in Formula 1A and 1B both are 0, Ring A is a group represented by Formula 2A or a group represented by Formula 2B;

[0019] when Ring A in Formula 1A and 1B is neither a group represented by Formula 2A nor a group represented by Formula 2B;

$$c_1 + c_2 \neq 0;$$

[0020] at least one selected from Ring A in Formula 1A, all R₁ in Formula 1A, and all R₂ in Formula 1A is a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, or a substituted or

unsubstituted monovalent non-aromatic condensed heteropolycyclic group, each including at least one Moiety 1 as a ring-forming moiety;

[0021] the sum of c₁₁, c₁₂, and c₁₃ in Formula 1C is not 0;

[0022] when each of Cz₁, Cz₂, and Cz₃ in Formula 1C is independently a group represented by Formula 2C or a group represented by Formula 2D, at least one selected from all R₁₁ in Formula 1C, all R₁₂ in Formula 1C, and all R₁₃ in Formula 1C is a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, or a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, each including at least one Moiety 1 as a ring-forming moiety;

[0023] L₁ to L₃, L₁₁ to L₁₃, L₂₁ to L₂₄, and L₃₁ to L₃₃ are each independently selected from a substituted or unsubstituted C₁-C₆₀ alkylene group, a substituted or unsubstituted C₂-C₆₀ alkenylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₂-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

[0024] a₁ to a₃, a₁₁ to a₁₃, a₂₁ to a₂₄, and a₃₁ to a₃₃ are each independently an integer selected from 0 to 6;

[0025] R₁, R₂, R₁₁ to R₁₃, R₂₁ to R₂₃, and R₃₁ to R₃₃ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), and —B(Q₆)(Q₇);

[0026] b₁, b₂, b₁₁ to b₁₃, b₂₁ to b₂₃, and b₃₁ to b₃₃ are each independently an integer selected from 0 to 4;

[0027] at least one of substituents of the substituted C₁-C₆₀ alkylene group, substituted C₂-C₆₀ alkenylene group, substituted C₃-C₁₀ cycloalkylene group, substituted C₂-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₂-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₂-C₁₀ heterocy-

cloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₂-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₂-C₆₀ heteroaryl group, a substituted monovalent non-aromatic condensed polycyclic group, and a substituted monovalent non-aromatic condensed hetero-polycyclic group is selected from

[0028] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0029] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed hetero-polycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

[0030] a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed hetero-polycyclic group;

[0031] a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇); and

[0032] —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇);

[0033] wherein Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ may be each independently selected from a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed hetero-condensed polycyclic group.

=N=*

Moiety 1

[0034] in Moiety 1, ** and *** each indicates a binding site to a ring-forming atom included in a ring including Moiety 1.

[0035] According to another aspect, an organic light-emitting device includes:

[0036] a first electrode;

[0037] a second electrode; and

[0038] an organic layer disposed between the first electrode and the second electrode,

[0039] wherein the organic layer includes an emission layer, and further includes at least one condensed cyclic compound represented by Formula 1.

BRIEF DESCRIPTION OF THE DRAWINGS

[0040] These and/or other aspects will become apparent and more readily appreciated from the following description of the embodiments, taken in conjunction with FIG. 1 which is a schematic view of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION

[0041] Reference will now be made in detail to embodiments, examples of which are illustrated in the accompanying drawings, wherein like reference numerals refer to like elements throughout. In this regard, the present embodiments may have different forms and should not be construed as being limited to the descriptions set forth herein. Accordingly, the embodiments are merely described below, by referring to the figures, to explain aspects. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. The term “or” means “and/or.” Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

[0042] It will be understood that when an element is referred to as being “on” another element, it can be directly in contact with the other element or intervening elements may be present therebetween. In contrast, when an element is referred to as being “directly on” another element, there are no intervening elements present.

[0043] It will be understood that, although the terms first, second, third etc. may be used herein to describe various elements, components, regions, layers, and/or sections, these elements, components, regions, layers, and/or sections should not be limited by these terms. These terms are only used to distinguish one element, component, region, layer, or section from another element, component, region, layer, or section. Thus, a first element, component, region, layer, or section discussed below could be termed a second element, component, region, layer, or section without departing from the teachings of the present embodiments.

[0044] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. As used herein, the singular forms “a,” “an,” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.

[0045] It will be further understood that the terms “comprises” and/or “comprising,” or “includes” and/or “including” when used in this specification, specify the presence of stated features, regions, integers, steps, operations, elements, and/or components, but do not preclude the presence or addition of one or more other features, regions, integers, steps, operations, elements, components, and/or groups thereof.

[0046] Spatially relative terms, such as “beneath,” “below,” “lower,” “above,” “upper” and the like, may be used herein for ease of description to describe one element or feature’s relationship to another element(s) or feature(s) as illustrated in the figures. It will be understood that the spatially relative terms are intended to encompass different orientations of the


device in use or operation in addition to the orientation depicted in the figures. For example, if the device in the figures is turned over, elements described as “below” or “beneath” other elements or features would then be oriented “above” the other elements or features. Thus, the exemplary term “below” can encompass both an orientation of above and below. The device may be otherwise oriented (rotated 90 degrees or at other orientations) and the spatially relative descriptors used herein interpreted accordingly.

[0047] “About” or “approximately” as used herein is inclusive of the stated value and means within an acceptable range of deviation for the particular value as determined by one of ordinary skill in the art, considering the measurement in question and the error associated with measurement of the particular quantity (i.e., the limitations of the measurement system). For example, “about” can mean within one or more standard deviations, or within $\pm 30\%$, 20% , 10% , 5% of the stated value.

[0048] Unless otherwise defined, all terms (including technical and scientific terms) used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. It will be further understood that terms, such as those defined in commonly used dictionaries, should be interpreted as having a meaning that is consistent with their meaning in the context of the relevant art and the present disclosure, and will not be interpreted in an idealized or overly formal sense unless expressly so defined herein.


[0049] Exemplary embodiments are described herein with reference to cross section illustrations that are schematic illustrations of idealized embodiments. As such, variations from the shapes of the illustrations as a result, for example, of manufacturing techniques and/or tolerances, are to be expected. Thus, embodiments described herein should not be construed as limited to the particular shapes of regions as illustrated herein but are to include deviations in shapes that result, for example, from manufacturing. For example, a region illustrated or described as flat may, typically, have rough and/or nonlinear features. Moreover, sharp angles that are illustrated may be rounded. Thus, the regions illustrated in the figures are schematic in nature and their shapes are not intended to illustrate the precise shape of a region and are not intended to limit the scope of the present claims.

[0050] A condensed cyclic compound according to an embodiment is represented by one of Formulae 1A to 1C:




-continued


Formula 1B




Formula 1C




Formula 2A




Formula 2B



Formula 2C




Formula 2D




-continued

Formula 2E



Formula 2F



[0051] The relationships between the groups represented by Formulae 2A to 2F and the condensed cyclic compound represented by one of Formulae 1A to 1C will be described below in detail.

[0052] Ring A in Formula 1A may be a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, a group represented by Formula 2A, or a group represented by Formula 2B.

[0053] According to an embodiment, Ring A in Formula 1A may be selected from


[0054] a phenyl group, a naphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, phenanthridinyl, phenanthrolinyl, phenazinyl, a benzoimidazolyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzofuopyrazinyl group, a benzofuopyrimidinyl group, a benzofuopyridinyl group, a benzothienopyrazinyl group, a benzothienopyrimidinyl group, a benzothienopyridinyl group, a group represented by Formula 2A, and a group represented by Formula 2B; and

[0055] a phenyl group, a naphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, phenanthridinyl, phenanthrolinyl, phenazinyl, a benzoimidazolyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzofuopyrazinyl group, a benzofuopyrimidinyl group, a benzofuopyridinyl group, a benzothienopyrazinyl group, a benzothienopyrimidinyl group, a benzothienopyridinyl group, a group represented by Formula 2A, and a group represented by Formula 2B; and


an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, phenanthridinyl, phenanthrolinyl, phenazinyl, a benzoimidazolyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzofuopyrazinyl group, a benzofuopyrimidinyl group, a benzofuopyridinyl group, a benzothienopyrazinyl group, a benzothienopyrimidinyl group, and a benzothienopyridinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a phenyl group substituted with a C₁-C₂₀ alkyl group, a naphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a biphenyl group, and —Si(Q₃₃)(Q₃₄)(Q₃₅),

[0056] wherein Q₃₃ to Q₃₅ may be each independently selected from a hydrogen, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group.


[0057] In some embodiments, Ring A in Formula 1A may be represented by one of Formulae 4-1 to 4-51, Formulae 6-1 to 6-5 below, a group represented Formula 2A, and a group represented by Formula 2B, but a formula representing Ring A is not limited thereto:




Formula 4-1



Formula 4-2















Formula 4-3




Formula 4-4

-continued




-continued


Formula 4-5




Formula 4-6




Formula 4-7




Formula 4-8




Formula 4-9




Formula 4-10




Formula 4-11




Formula 4-12




Formula 4-13




Formula 4-14



Formula 4-15



Formula 4-16



Formula 4-17

Formula 4-18

Formula 4-19

Formula 4-20

Formula 4-21

Formula 4-22

Formula 4-23

Formula 4-24


Formula 4-25

Formula 4-26


Formula 4-27

Formula 4-28


-continued




Formula 4-29




Formula 4-30




Formula 4-31




Formula 4-32




Formula 4-33




Formula 4-34



Formula 4-35




Formula 4-36




Formula 4-37


-continued




Formula 4-38




Formula 4-39




Formula 4-40




Formula 4-41




Formula 4-42




Formula 4-43




Formula 4-44



Formula 4-45



Formula 4-46




Formula 4-47




Formula 4-48


-continued




Formula 4-49




Formula 4-50




Formula 4-51




Formula 6-1




Formula 6-2



Formula 6-2



Formula 6-4



Formula 6-5

[0058] In Formulae 4-1 to 4-51 and 6-1 to 6-5,

[0059] Y_{31} is O, S, C(Z_{33})(Z_{34}), N(Z_{35}), or Si(Z_{36})(Z_{37});

[0060] Z_{31} to Z_{37} may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group substituted

with a C_1 - C_{20} alkyl group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a quinoxaliny group, a biphenyl group, and —Si(Q_{33})(Q_{34})(Q_{35}); wherein

[0061] Q_{33} to Q_{35} are each independently selected from a hydrogen, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, and a quinoxaliny group;

[0062] e_1 is an integer of 1 to 5;

[0063] e_2 is an integer of 1 to 7;

[0064] e_3 is an integer of 1 to 3;

[0065] e_4 is an integer of 1 to 4;

[0066] e_5 is an integer of 1 or 2;

[0067] e_6 is an integer of 1 to 6; and

[0068] * indicates a binding site to a neighboring atom.

[0069] Ring B_1 in Formula 1A is a C_5 - C_{20} carbocyclic group or a C_2 - C_{20} heterocyclic group that does not include a nitrogen atom as a ring-forming atom.

[0070] For example, Ring B_1 in Formula 1A may be a benzene, a naphthalene, an anthracene, a phenanthrene, a benzofuran, or a benzothiophene.

[0071] According to an embodiment, Ring B_1 in Formula 1A may be a benzene, a naphthalene, an anthracene, a phenanthrene, or a benzofuran.

[0072] Ring B_2 , Ring B_3 , Ring C_1 , and Ring C_2 in Formulae 1B and 1C are each independently a C_5 - C_{20} carbocyclic group or a C_2 - C_{20} heterocyclic group, provided that at least one selected from Ring B_2 and Ring B_3 is a C_2 - C_{20} heterocyclic group including a nitrogen atom as a ring-forming atom, and at least one selected from Ring C_1 and Ring C_2 is a C_2 - C_{20} heterocyclic group including a nitrogen atom as a ring-forming atom.

[0073] In some embodiments, Ring B_2 , Ring B_3 , Ring C_1 , and Ring C_2 may be each independently a benzene, a naphthalene, an anthracene, a phenanthrene, a benzofuran, a benzothiophene, a pyridine, a pyrimidine, a pyrazine, a benzofuropyrazine, a benzofuopyrimidine, benzofuopyridine, a benzothienopyrazine, a benzothienopyrimidine, or a benzothienopyridine;

[0074] at least one selected from Ring B_2 and Ring B_3 is a pyridine, a pyrimidine, a pyrazine, a benzofuropyrazine, a benzofuopyrimidine, a benzofuopyridine, a benzothienopyrazine, a benzothienopyrimidine, or a benzothienopyridine, and

[0075] at least one selected from Ring C_1 and Ring C_2 is a pyridine, a pyrimidine, a pyrazine, a benzofuropyrazine, a benzofuopyrimidine, a benzofuopyridine, a benzothienopyrazine, a benzothienopyrimidine, or a benzothienopyridine.

[0076] In some embodiments, Ring B_2 , Ring B_3 , Ring C_1 , and Ring C_2 may be each independently a benzene, a naphthalene, a pyridine, a pyrimidine, or a pyrazine;

[0077] at least one of Ring B₂ and Ring B₃ may be a pyridine, a pyrimidine, or a pyrazine; and

[0078] at least one of Ring C₁ and Ring C₂ may be a pyridine, a pyrimidine, or a pyrazine.

[0079] Pc₁ and Pc₂ in Formula 1A and 1B are each independently a group represented by Formulae 2A or a group represented by Formula 2B.

[0080] Each of Cz₁, Cz₂, and Cz₃ in Formula 1C is independently a group represented by Formula 2C, a group represented by Formula 2D, a group represented by Formula 2E, or a group represented by Formula 2F.

[0081] c₁, c₂, c₁₁, c₁₂, c₁₃, p₁, p₂, p₁₁, p₁₂, and p₁₃ in Formulae 1A to 1C are each independently an integer selected from 0 to 3.

[0082] When each of c₁ and c₂ in Formulae 1A and 1B is 0, Ring A is a group represented by Formula 2A or a group represented by Formula 2B, and when Ring A in Formulae 1A and 1B is neither a group represented by Formula 2A nor a group represented by Formula 2B, c₁+c₂≠0.

[0083] That is, Formulae 1A and 1B necessarily include one selected from the group represented by Formula 2A and the group represented by Formula 2B.

[0084] At least one selected from Ring A in Formula 1A, all R₁ in Formula 1A, and all R₂ in Formula 1A is an electron transport moiety.

[0085] The electron transport moiety may be a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, or a substituted or unsubstituted monovalent non-aromatic condensed hetero-polycyclic group, each including Moiety 1 below as a ring-forming moiety. Herein, a carbazole-based group that does not include Moiety 1 is not an electron transport moiety, due to the absence of Moiety 1.

=N—*

Moiety 1

[0086] in Moiety 1, ** and *** each indicates a binding site to a ring-forming atom in a ring including Moiety 1.

[0087] Examples of the electron transport moiety are groups represented by Formulae 4-1 to 4-51, but are not limited thereto.

[0088] The sum of c₁₁, c₁₂, and c₁₃ in Formula 1C is not 0.

[0089] That is, Formula 1C necessarily includes at least one of groups represented by Formulae 2C to 2F.

[0090] When Cz₁, Cz₂, and Cz₃ in Formula 1C is a group represented by Formula 2C or a group represented by Formula 2D, at least one selected from R₁₁ in Formula 1C, R₁₂ in Formula 1C, and R₁₃ in Formula 1C is an electron transport moiety. Herein, the electron transport moiety has been described above and will not be repeated.

[0091] According to an embodiment, in Formulae 1A and 1B,

[0092] c₁=0, c₂=0, and Ring A is a group represented by Formula 2A or a group represented by Formula 2B;

[0093] c₁=1 and c₂=0; or

[0094] c₁=1 and c₂=1.

[0095] In some embodiments, in Formula 1C,

[0096] c₁₁=1, c₁₂=0, and c₁₃=0;

[0097] c₁₁=0, c₁₂=0, and c₁₃=1;

[0098] c₁₁=1, c₁₂=1, and c₁₃=0; or

[0099] c₁₁=1, c₁₂=0, and c₁₃=1.

[0100] L₁ to L₃, L₁₁ to L₁₃, L₂₁ to L₂₄, and L₃₁ to L₃₃ in Formulae 1A to 1C and 2A to 2F may be each independently

selected from a substituted or unsubstituted C₁-C₆₀ alkylene group, a substituted or unsubstituted C₂-C₆₀ alkenylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylenylene group, a substituted or unsubstituted C₂-C₆₀ heteroarylenylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed hetero-polycyclic group.

[0101] For example, L₁ to L₃, L₁₁ to L₁₃, L₂₁ to L₂₄ and L₃₁ to L₃₃ in Formulae 1A to 1C and 2A to 2F may be each independently selected from


[0102] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pyrrolylene group, an imidazolylene group, a pyrazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzoxazolylene group, a benzoimidazolylene group, a furanylene group, a benzofuranylene group, a thiophenylene group, a benzothiophenylene group, a thiazolylene group, an isothiazolylene group, a benzothiazolylene group, an isoxazolylene group, an oxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, an imidazopyrimidinylene group, and an imidazopyridinylene group; and

[0103] a phenylene group, a pentalenylene group, an indenylene group, naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pyrrolylene group, an imidazolylene group, a pyrazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzooxazolylene group, a benzoimidazolylene group, a


furanylene group, a benzofuranylene group, a thiophenylylene group, a benzothiophenylylene group, a thiazolylylene group, an isothiazolylylene group, a benzothiazolylylene group, an isoxazolylylene group, an oxazolylylene group, a triazolylylene group, a tetrazolylylene group, an oxadiazolylylene group, a triazinylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, an imidazopyrimidinylene group, and an imidazopyridinylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₆-C₂₀ aryl group, a C₂-C₂₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, and —Si(Q₃₃)(Q₃₄)(Q₃₅);

[0104] wherein Q₃₃ to Q₃₅ are each independently selected from a hydrogen, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a phthalazinyl group, a quinoxalinyl group, a cinnolinyl group, and a quinazolinyl group.


[0105] According to an embodiment, L₁ to L₃, L₁₁ to L₁₃, L₂₁ to L₂₄ and L₃₁ to L₃₃ in Formulae 1A to 1C and 2A to 2F may be each independently one selected from Formulae 2-1 to 2-37:




Formula 2-1



Formula 2-2




Formula 2-3




Formula 2-4


-continued




Formula 2-5




Formula 2-6




Formula 2-7




Formula 2-8




Formula 2-9




Formula 2-10




Formula 2-11



Formula 2-12














Formula 2-13




Formula 2-14

-continued




-continued

Formula 2-15




Formula 2-16




Formula 2-17

Formula 2-18




Formula 2-19


Formula 2-10




Formula 2-21



Formula 2-22




Formula 2-23



Formula 2-24

Formula 2-25



Formula 2-26

Formula 2-27

Formula 2-28

Formula 2-29


Formula 2-30

Formula 2-31


Formula 2-32

Formula 2-33


-continued




Formula 2-34



Formula 2-35



Formula 2-36



Formula 2-37

[0106] wherein in Formulae 2-1 to 2-37,

[0107] Y_1 may be O, S, S(=O), S(=O)₂, C(Z₃)(Z₄), N(Z₅), or Si(Z₆)(Z₇);

[0108] Z₁ to Z₇ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, a quinoxalinyl group, a biphenyl group, and —Si(Q₃₃)(Q₃₄)(Q₄₅), wherein

[0109] Q₃₃ to Q₄₅ are each independently selected from a hydrogen, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl

group, an isoquinolinyl group, a phthalazinyl group, a quinoxalinyl group, a cinnolinyl group, and a quinazolinyl group;

[0110] d₁ is an integer of 1 to 4;

[0111] d₂ is an integer of 1 to 3;

[0112] d₃ is an integer of 1 to 6;

[0113] d₄ is an integer of 1 to 8;

[0114] d₅ is an integer of 1 or 2;

[0115] d₆ is an integer of 1 to 5; and


[0116] * indicates a binding site to a neighboring atom.

[0117] In some embodiments, L₁ to L₃, L₁₁ to L₁₃, L₂₁ to L₂₄ and L₃₁ to L₃₃ in Formulae 1A to 1C and 2A to 2F may be each independently selected from


[0118] a phenylene group, a naphthylene group, a phenanthrenylene group, an anthracenylene group, a triphenylenylene group, and a carbazolylene group; and

[0119] a phenylene group, a naphthylene group, a phenanthrenylene group, an anthracenylene group, a triphenylenylene group, and a carbazolylene group each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a phenanthrenyl group, and a biphenyl group, but they are not limited thereto.


[0120] In some embodiments, L₁ to L₃, L₁₁ to L₁₃, L₂₁ to L₂₄ and L₃₁ to L₃₃ in Formulae 1A to 1C and 2A to 2F may be each independently one of Formulae 3-1 to 3-33, but are not limited thereto:




Formula 3-1




Formula 3-2




Formula 3-3



Formula 3-4











Formula 3-5



Formula 3-6

-continued



Formula 3-7

Formula 3-8

Formula 3-9

Formula 3-10








Formula 3-11

Formula 3-12

Formula 3-13

Formula 3-14

-continued



Formula 3-15

Formula 3-16

Formula 3-17


Formula 3-18

Formula 3-19


Formula 3-20

Formula 3-21


-continued




Formula 3-22




Formula 3-23




Formula 3-24




Formula 3-25



Formula 3-26




Formula 3-27




Formula 3-28


-continued




Formula 3-29




Formula 3-30



Formula 3-31



Formula 3-32



Formula 3-33

[0121] wherein * and *¹ in Formulae 3-1 to 3-33 each independently indicates a binding site to a neighboring atom.

[0122] a₁ in Formulae 1A to 1C indicates the number of L₁, and may be 0, 1, 2, 3, 4 or 6, for example, 0, 1 or 2, or for example, 0 or 1. When a₁ is 0, L₁ is a single bond. When a₁ is 2 or more, groups L₁ may be identical or different. a₂ to a₃, a₁₁ to a₁₃, a₂₁ to a₂₄ and a₃₁ to a₃₃ may be understood by referring to the description presented in connection with a₁ and structures of Formulae 1A to 1C and 2A to 2F.

[0123] According to an embodiment, a₁ to a₃, a₁₁ to a₁₃, a₂₁ to a₂₄ and a₃₁ to a₃₃ in Formula 1 may be each independently 0, 1, or 2, for example, 0 or 1, but they are not limited thereto.

[0124] R₁, R₂, R₁₁ to R₁₃, R₂₁ to R₂₃, and R₃₁ to R₃₃ in Formulae 1A to 1C and 2A to 2F may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a

salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkyne group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed hetero-polycyclic group, a group represented by Formula 2A, a group represented by Formula 2B, $-N(Q_1)(Q_2)$, $-Si(Q_3)(Q_4)(Q_5)$, and $-B(Q_6)(Q_7)$.

[0125] For example, R_1 , R_2 , R_{11} to R_{13} , R_{21} to R_{23} , and R_{31} to R_{33} may be each independently selected from a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0126] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, and a phosphoric acid or a salt thereof;

[0127] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0128] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl

group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, phenanthridinyl, acridinyl, phenanthrolinyl, phenazinyl, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a phenyl group substituted with a C_1 - C_{20} alkyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a biphenyl group, and $-Si(Q_{33})(Q_{34})(Q_{35})$; and

[0129] $-Si(Q_3)(Q_4)(Q_5)$;

[0130] wherein Q_3 to Q_5 and Q_{33} to Q_{35} are each independently selected from a hydrogen, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a

fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzozofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group.

[0131] In some embodiments, R_1 , R_2 , R_{11} to R_{13} , R_{21} to R_{23} , and R_{31} to R_{33} may be each independently selected from


[0132] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0133] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, and a phosphoric acid or a salt thereof;


[0134] one selected from Formulae 4-1 to 4-51, 5-1 to 5-6, and 6-1 to 6-5; and

[0135] —Si(Q_3)(Q_4)(Q_5),


[0136] wherein Q_3 to Q_5 may be each independently selected from a hydrogen, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzozofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group, but they are not limited thereto.




Formula 4-1




Formula 4-2



Formula 4-3




Formula 4-4




Formula 4-5


-continued




Formula 4-6




Formula 4-7




Formula 4-8




Formula 4-9




Formula 4-10




Formula 4-11




Formula 4-12




Formula 4-13




Formula 4-14




Formula 4-15



Formula 4-16














Formula 4-17




Formula 4-18

-continued




-continued

Formula 4-19




Formula 4-20




Formula 4-21

Formula 4-22




Formula 4-23




Formula 4-24


Formula 4-25




Formula 4-26




Formula 4-27



Formula 4-28



Formula 4-29



Formula 4-30

Formula 4-31

Formula 4-32

Formula 4-33

Formula 4-34












Formula 4-35

Formula 4-36

Formula 4-37


Formula 4-38

-continued




-continued

Formula 4-39




Formula 4-40

Formula 4-41




Formula 4-42

Formula 4-43




Formula 4-44

Formula 4-45




Formula 4-46




Formula 4-47



Formula 4-48



Formula 4-49



Formula 4-50

Formula 4-51

Formula 5-1

Formula 5-2

Formula 5-3


Formula 5-4

Formula 5-5


Formula 5-6

Formula 6-1


-continued




Formula 6-2



Formula 6-3



Formula 6-4



Formula 6-5

[0137] wherein in Formulae 4-1 to 4-51, 5-1 to 5-6, and 6-1 to 6-5,

[0138] Y_{31} is O, S, C(Z_{33})(Z_{34}), N(Z_{35}), or Si(Z_{36})(Z_{37});

[0139] Z_{31} to Z_{37} may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group substituted with a C_1 - C_{20} alkyl group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, a quinoxalinyl group, a biphenyl group, and —Si(Q_{33})(Q_{34})(Q_{35}); wherein

[0140] Q_{33} to Q_{35} are each independently selected from a hydrogen, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group;

[0141] e_1 is an integer of 1 to 5;

[0142] e_2 is an integer of 1 to 7;

[0143] e_3 is an integer of 1 to 3;

[0144] e_4 is an integer of 1 to 4;

[0145] e_5 is an integer of 1 or 2;


[0146] e_6 is an integer of 1 to 6; and

[0147] * indicates a binding site to a neighboring atom.


[0148] b_1 , b_2 , b_{11} to b_{13} , b_{21} to b_{23} , and b_{31} to b_{33} in the formulae above may be each independently an integer selected from 0 to 4. For example, b_1 , b_2 , b_{11} to b_{13} , b_{21} to b_{23} and b_{31} to b_{33} may be each independently 0 or 1, but they are not limited thereto.

[0149] Formula 2A may be represented by one of Formulae 2A(1) to 2A(5) below, Formula 2B may be represented by one of Formulae 2B(1) and 2B(2) below, Formula 2D may be represented by one of Formulae 2D(1) to 2D(5) below, and Formula 2F may be represented by one of Formulae 2F(1) and 2F(2) below:


Formula 2A(1)




Formula 2A(2)




Formula 2A(3)




Formula 2A(4)




Formula 2A(5)




-continued




Formula 2B(2)




Formula 2E(1)




Formula 2E(2)




Formula 2E(3)




Formula 2E(4)




-continued



Formula 2F(1)



Formula 2F(2)




[0150] L₂₁ to L₂₄, a₂₁ to a₂₄, R₂₁ to R₂₃, b₂₁ to b₂₃, L₃₁ to L₃₃, a₃₁ to a₃₃, R₃₁ to R₃₃, b₃₁ to b₃₃, c₂₁ to c₂₃, c₃₁, and c₃₂ in Formulae 2A (1) to 2A (5), 2B(1), 2B(2), 2E(1) to 2E(5), 2F(1), and 2F(2) are as described above.


[0151] For example, a₂₄ in Formulae 2A(1) to 2A(5), 2B(1), and 2B(2) is 0, 1, or 2, and L₂₄ is selected from Formulae 2-1 to 2-37, and c₁₁, c₁₂, and c₁₃ may each be a hydrogen, but they are not limited thereto.

[0152] The condensed cyclic compound represented by Formula 1 may be represented by one of Formulae 1A(1) to 1A(8) below:

Formula 1A(1)




-continued




Formula 1A(2)


-continued




Formula 1A(6)




Formula 1A(3)




Formula 1A(7)



Formula 1A(4)



Formula 1A(8)



Formula 1A(5)


[0153] In Formulae 1A(1) and 1A(8),

[0154] Ring A is a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, or a substituted or unsubstituted monovalent non-aromatic condensed hetero-polycyclic group, each including Moiety 1 as a ring-forming moiety, (for example, a group represented by one of Formulae 4-1 to 4-55),


[0155] Pc₂, Pc₂, L₃, a₃, R₁, R₂, p₁, and p₂ are as described above, and

[0156] R₃ to R₄ are as described above.


[0157] In some embodiments, the condensed cyclic compound represented by Formula 1 may be represented by one of Formulae 1B(1) to 1B(10) below:




Formula 1B(1)




Formula 1B(2)




Formula 1B(3)




Formula 1B(4)




Formula 1B(5)




Formula 1B(6)



Formula 1B(7)




Formula 1B(8)



Formula 1B(9)


-continued




Formula 1B(10)

[0158] Ring A, P_{c1}, P_{c2}, L₃, a₃, R₁, R₂, p₁, and p₂ in Formulae 1B(1) to 1B(10) are as described above.

[0159] In some embodiments, the condensed cyclic compound may be represented by Formulae 1C(1) or 1C(2) below:




Formula 1C(1)



Formula 1C(2)

[0160] C_{z1}, C_{z2}, C_{z3}, R₁₁, R₁₂, R₁₃, p₁₁, p₁₂, and p₁₃ in Formulae 1C(1) and 1C(2) are as described above.

[0161] In some embodiments, the condensed cyclic compound may be represented by Formulae 1C(3) below:




Formula 1C(3)

[0162] wherein in Formula 1C(3),


[0163] C_{z3}, L₁₁, and a₁₁ are as described above; R₁₁ may be a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalk-

enyl group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, or a substituted or unsubstituted monovalent non-aromatic condensed hetero-polycyclic group, each including Moiety 1 as a ring-forming moiety, (for example, a group represented by one of Formulae 4-1 to 4-55); and b₁₁ is 1 or 2.

[0164] In some embodiments, the condensed cyclic compound may be represented by Formulae 1C(4) or 1C(5) below:




Formula 1C(4)



Formula 1C(5)


[0165] wherein in Formulae 1C(4) and 1C(5), C_{z1} and C_{z3} are each a group represented by Formula 2E or a group represented by Formula 2F.

[0166] The condensed cyclic compound may be one of Compounds 1 to 76 below, but is not limited thereto:




-continued

2




-continued


4



3




5




-continued


6



7




8




-continued


9



10




11




-continued


12



13




14




-continued


15



16




17




-continued

18




-continued


21




19



20




22




-continued

23




-continued


25



24




26




-continued


27



28




29




-continued

30




31




-continued

32




-continued


34




33



35




36




-continued

37




-continued


39



38




40




-continued

41





-continued

43




44


42



45




-continued




46


-continued




49




47



50




48



51

-continued




52

53

54

-continued




55

56


57

-continued


58



59




60




-continued

61




62




-continued

63




-continued


66




64




67




65



68




-continued




69


-continued



71




70




72

-continued





-continued



75

76



[0167] Formulae 1A and 1B necessarily include at least one group represented by one of Formulae 2A and 2B, and Formula 1C has a core that is substantially similar to Formulae 2A and 2B. Accordingly, excellent thermal stability may be obtained.

[0168] In some embodiments, Formula 1A necessarily includes the electron transport moiety (for example, see Formulae 4-1 to 4-55) used herein, at least one selected from Ring B₁ and Ring B₂ in Formula 1B necessarily includes at least one nitrogen as a ring-forming atom, and Formula 1C necessarily includes either the electron transport moiety used herein in its molecular structure or a group represented by at least one of Formulae 2E and 2F. Accordingly, the condensed cyclic compound represented by one of Formulae 1A to 1C may have excellent bipolar characteristics and high T₁ energy levels, and due to these characteristics, the condensed cyclic compound is appropriate for use as a material for forming an electric device, for example, an organic light-emitting device. For example, the condensed cyclic compound represented by one of Formulae 1A to 1C may be appropriate for use as a host for an organic light-emitting device.

[0169] For example, the highest occupied molecular orbital (HOMO), lowest unoccupied molecular orbital (LUMO), T₁

energy level, and S_1 energy level of Compounds 2 to 19 and 21 to 76 and Compounds A and B were simulated by using Gaussian, and simulation evaluation results are shown in Table 1 below:


TABLE 1

	HOMO (eV)	LUMO (eV)	T_1 (eV)	S_1 (eV)
Compound A	-5.113	-1.25	2.933	3.224
Compound B	-5.118	-1.589	2.856	3.039
Compound 2	-5.344	-1.916	2.912	3.02
Compound 3	-5.133	-1.991	2.627	2.731
Compound 4	-5.196	-1.798	2.781	2.947
Compound 5	-5.083	-1.792	2.862	2.928
Compound 6	-5.32	-1.891	2.74	2.924
Compound 7	-5.167	-1.973	2.424	2.767
Compound 8	-5.167	-2.019	2.399	2.728
Compound 9	-5.282	-1.664	2.924	3.04
Compound 11	-5.337	-1.933	2.886	3.01
Compound 12	-5.273	-1.953	2.773	2.919
Compound 13	-5.308	-1.84	2.733	2.999
Compound 14	-5.259	-1.992	2.47	2.835
Compound 15	-5.100	-1.793	2.878	3.040
Compound 16	-5.279	-1.691	2.901	3.085
Compound 17	-5.487	-1.957	2.865	3.126
Compound 18	-5.291	-2.015	2.455	2.845
Compound 19	-5.284	-1.925	2.508	2.924
Compound 26	-5.343	-1.927	2.802	3.023
Compound 21	-5.430	-1.997	2.836	3.012
Compound 22	-5.246	-2.021	2.688	2.804
Compound 23	-5.139	-2.004	2.643	2.743
Compound 24	-5.269	-2.015	2.727	2.843
Compound 25	-5.252	-2.018	2.7	2.816
Compound 26	-5.045	-1.996	2.554	2.649
Compound 27	-5.255	-1.934	2.811	2.846
Compound 28	-5.266	-2.011	2.737	2.87
Compound 29	-5.267	-1.985	2.713	2.837
Compound 30	-5.419	-1.992	2.697	2.965
Compound 31	-5.243	-1.972	2.697	2.821
Compound 32	-5.06	-2.01	2.564	2.656
Compound 33	-5.155	-2.007	2.657	2.764
Compound 34	-5.224	-2.004	2.523	2.843
Compound 35	-5.213	-2.012	2.538	2.804
Compound 36	-5.256	-1.997	2.693	2.819
Compound 37	-5.292	-2.004	2.704	2.837
Compound 38	-5.282	-2.001	2.559	2.833
Compound 39	-5.082	-1.974	2.586	2.696
Compound 40	-5.201	-1.986	2.64	2.769
Compound 41	-5.111	-1.939	2.576	2.703
Compound 42	-5.17	-2.035	2.403	2.726
Compound 43	-5.227	-1.279	2.917	3.191
Compound 44	-5.384	-1.78	2.849	3.121
Compound 45	-5.214	-1.222	2.92	3.129
Compound 46	-5.295	-1.327	2.918	3.108
Compound 47	-5.269	-1.302	2.915	3.158
Compound 48	-5.236	-1.387	2.894	3.057
Compound 49	-5.589	-1.653	2.986	3.32
Compound 50	-5.361	-1.704	2.892	3.24
Compound 51	-5.603	-1.914	2.907	3.102
Compound 52	-5.252	-1.675	2.927	3.062
Compound 53	-5.509	-1.857	2.499	3.107
Compound 54	-5.895	-1.767	2.992	3.297
Compound 55	-5.87	-1.693	2.959	3.345
Compound 56	-5.709	-1.642	2.532	3.188
Compound 57	-5.819	-1.894	2.667	3.124
Compound 58	-5.436	-2.124	2.924	3.016
Compound 59	-5.666	-1.853	2.865	3.096
Compound 60	-5.501	-1.492	2.713	3.119
Compound 61	-5.089	-1.762	2.745	3.022
Compound 62	-5.331	-2.008	2.827	2.925
Compound 63	-5.097	-1.815	2.053	2.901
Compound 64	-5.314	-2.015	2.587	2.804
Compound 65	-5.177	-1.964	2.571	2.816
Compound 66	-5.248	-1.655	2.662	3.094
Compound 67	-5.369	-1.413	2.879	3.118
Compound 68	-5.153	-1.984	2.785	2.839
Compound 69	-5.094	-1.804	2.734	2.94


TABLE 1-continued

	HOMO (eV)	LUMO (eV)	T_1 (eV)	S_1 (eV)
Compound 70	-5.426	-2.123	2.73	2.977
Compound 71	-5.334	-1.938	2.868	2.964
Compound 72	-5.254	-1.915	2.441	2.887
Compound 73	-5.321	-1.918	2.894	3.017
Compound 74	-5.276	-1.926	2.305	2.971
Compound 75	-5.016	-1.580	2.186	2.964

Compound A



Compound B



[0170] From Table 1, it is confirmed that Compounds 2 to 19 and 21 to 76 have lower LUMO levels (that is, larger LUMO absolute values) than Compounds A and B. Accordingly, it is confirmed that the condensed cyclic compound represented by one of Formulae 1A to 1C is appropriate for use as a material for an organic layer, for example, an auxiliary layer or an emission layer, of an organic light-emitting device.

[0171] A method of synthesizing the condensed cyclic compound represented by one of Formulae 1A to 1C may be apparent to one of ordinary skill in the art by referring to Synthesis Examples used herein.

[0172] Accordingly, since the condensed cyclic compound represented by one of Formulae 1A to 1C is appropriate for use in an organic layer of an organic light-emitting device, for

example, for use as a host in an emission layer of the organic layer, another aspect provides an organic light-emitting device including:

[0173] a first electrode;

[0174] a second electrode; and

[0175] an organic layer disposed between the first electrode and the second electrode,

[0176] wherein the organic layer includes an emission layer and at least one condensed cyclic compound represented by one of Formulae 1A to 1C.

[0177] Due to the inclusion of the organic layer including the condensed cyclic compound represented by one of Formulae 1A to 1C, the organic light-emitting device may have low driving voltage, high efficiency, high brightness, and long lifespan.

[0178] The condensed cyclic compound represented by one of Formulae 1A to 1C may be used between a pair of electrodes of an organic light-emitting device. For example, the condensed cyclic compound may be included in at least one selected from an emission layer, a hole transport region (including, for example, at least one of a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer) that is disposed between the first electrode and the emission layer, and an electron transport region (including, for example, at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer) that is disposed between the emission layer and the second electrode. For example, the condensed cyclic compound represented by one of Formulae 1A to 1C may be included in the emission layer. In this regard, the emission layer may further include a dopant, and the condensed cyclic compound included in the emission layer may act as a host. The emission layer may be a green emission layer emitting green light or a blue emission layer emitting blue light, and the dopant may be a phosphorescent dopant.

[0179] The expression “(an organic layer) includes at least one condensed cyclic compound” as used herein may include a case in which “(an organic layer) includes identical condensed cyclic compounds represented by Formulae 1A to 1C and a case in which (an organic layer) includes two or more different condensed cyclic compounds represented by Formulae 1A to 1C.

[0180] For example, the organic layer may include, as the condensed cyclic compound, only Compound 1. In this regard, Compound 1 may be situated in an emission layer of the organic light-emitting device. In another embodiment, the organic layer may include, as the condensed cyclic compound, Compound 1 and Compound 2. In this regard, Compound 1 and Compound 2 may be situated in either an identical layer (for example, Compound 1 and Compound 2 all may be situated in an emission layer), or different layers.

[0181] The first electrode may be an anode, which is a hole injection electrode, and the second electrode may be a cathode, which is an electron injection electrode, or the first electrode may be a cathode, which is an electron injection electrode, or the second electrode may be an anode, which is a hole injection electrode.

[0182] For example, the first electrode is an anode, and the second electrode is a cathode, and the organic layer includes;

[0183] i) a hole transport region disposed between the first electrode and the emission layer, including at least one of a hole injection layer, a hole transport layer, and an electron blocking layer, and

[0184] ii) an electron transport region disposed between the emission layer and the second electrode, including at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

[0185] The term “organic layer” as used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of an organic light-emitting device. The “organic layer” may include, in addition to an organic compound, an organometallic complex including metal.

[0186] FIG. 1 is a schematic view of an organic light-emitting device 10 according to an embodiment. Hereinafter, the structure of an organic light-emitting device according to an embodiment and a method of manufacturing an organic light-emitting device according to an embodiment will be described in connection with FIG. 1. The organic light-emitting device 10 includes a first electrode 11, an organic layer 15, and a second electrode 19, which are sequentially stacked.

[0187] In FIG. 1, a substrate may be additionally disposed under the first electrode 11 or above the second electrode 19. For use as the substrate, any substrate that is used in general organic light-emitting devices may be used, and the substrate may be a glass substrate or transparent plastic substrate, each with excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and waterproofness.

[0188] The first electrode 11 may be formed by depositing or sputtering a material for forming the first electrode on the substrate. The first electrode 11 may be an anode. The material for the first electrode 11 may be selected from materials with a high work function to allow holes to be easily provided. The first electrode 11 may be a reflective electrode or a transmissive electrode. The material for the first electrode 11 may be an indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), or zinc oxide (ZnO). In some embodiments, the material for the first electrode 11 may be metal, such as magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag).

[0189] The first electrode 11 may have a single-layer structure or a multi-layer structure including two or more layers. For example, the first electrode 11 may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode 110 is not limited thereto.

[0190] An organic layer 15 is disposed on the first electrode 11.

[0191] The organic layer 15 may include a hole transport region, an emission layer, and an electron transport region.

[0192] The hole transport region may be disposed between the first electrode 11 and the emission layer.

[0193] The hole transport region may include at least one of a hole injection layer, a hole transport layer, an electron blocking layer, and a buffer layer.

[0194] The hole transport region may include only either a hole injection layer or a hole transport layer. According to another embodiment, the hole transport region may have a structure of hole injection layer/hole transport layer or hole injection layer/hole transport layer/electron blocking layer, which are sequentially stacked in this stated order in the direction from the first electrode 11.

[0195] When the hole transport region includes a hole injection layer (HIL), the hole injection layer may be formed on the first electrode 11 by using any one of various methods,


for example, vacuum deposition, spin coating, casting, or Langmuir-Blodgett (LB) deposition.

[0196] When a hole injection layer is formed by vacuum deposition, the deposition conditions may vary according to a material that is used to form the hole injection layer, and the structure and thermal characteristics of the hole injection layer. For example, the deposition conditions may include a deposition temperature of about 100 to about 500° C., a vacuum pressure of about 10^{-8} to about 10^{-3} torr, and a deposition rate of about 0.01 to about 100 Å/sec. However, the deposition conditions are not limited thereto.




[0197] When the hole injection layer is formed using spin coating, coating conditions may vary according to the material used to form the hole injection layer, and the structure and thermal properties of the hole injection layer. For example, a coating speed may be from about 2,000 revolutions per minute (rpm) to about 5,000 rpm, and a temperature at which a heat treatment is performed to remove a solvent after coating may be from about 80° C. to about 200° C. However, the coating conditions are not limited thereto.

[0198] Conditions for a hole transport layer and an electron blocking layer may be understood by referring to conditions for forming the hole injection layer.


[0199] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB, β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), (polyaniline)/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201 below, and a compound represented by Formula 202 below:




-continued




-continued




β -NPB




TPD



Spiro-TPD




Spiro-NPB




methylated NPB

-continued




TAPC




HMTPD

Formula 201



Formula 202



[0200] Ar_{101} and Ar_{102} in Formula 201 may be each independently selected from

[0201] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, and a pentacenylene group; and

[0202] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, and a pentacenylene group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed hetero-polycyclic group.

[0203] x_a and x_b in Formula 201 may be each independently an integer of 0 to 5, or 0, 1 or 2. For example, x_a may be 1 and x_b may be 0, but x_a and x_b are not limited thereto.

[0204] R_{101} to R_{108} , R_{111} to R_{119} , and R_{121} to R_{124} in Formulae 201 and 202 may be each independently selected from

[0205] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{10} alkyl group (for example, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a hexyl group, and so on), and a C_1 - C_{10} alkoxy group (for example, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, and so on);

[0206] a C_1 - C_{10} alkyl group and a C_1 - C_{10} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, and a phosphoric acid or a salt thereof;

[0207] a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, and a pyrenyl group; and

[0208] a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, and a pyrenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{10} alkyl group, or a C_1 - C_{10} alkoxy group, but they are not limited thereto.


[0209] R_{109} in Formula 201 may be selected from

[0210] a phenyl group, a naphthyl group, an anthracenyl group, and a pyridinyl group; or

[0211] a phenyl group, a naphthyl group, an anthracenyl group, and a pyridinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group.

[0212] According to an embodiment, the compound represented by Formula 201 may be represented by Formula 201A below, but is not limited thereto:


Formula 201A



[0213] R_{101} , R_{111} , R_{112} , and R_{109} in Formula 201A may be understood by referring to the description provided herein.


[0214] For example, the compound represented by Formula 201, and the compound represented by Formula 202 may include compounds HT1 to HT20 illustrated below, but are not limited thereto.

HT1




-continued

HT2




-continued


HT4



HT3




HT5




-continued

HT6





-continued

HT8




HT9

HT7




-continued

HT10




-continued


HT12




HT13




HT11




HT14



HT15




-continued




HT16


-continued




HT20



HT17



HT18




HT19


[0215] A thickness of the hole transport region may be in a range of about 100 Angstroms (Å) to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes both a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, for example about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0216] The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0217] The charge-generation material may be, for example, a p-dopant. The p-dopant may be one of a quinone derivative, a metal oxide, and a cyano group-containing compound, but is not limited thereto. Non-limiting examples of the p-dopant are a quinone derivative, such as tetracyanoquinonodimethane (TCNQ) or 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonodimethane (F4-TCNQ); a metal oxide, such as a tungsten oxide or a molybdenum oxide; and a cyano group-containing compound, such as Compound HT-D1 below, but are not limited thereto.



Compound HT-D1



[0218] The hole transport region may include a buffer layer.

[0219] Also, the buffer layer may compensate for an optical resonance distance according to a wavelength of light emitted from the emission layer, and thus, efficiency of a formed organic light-emitting device may be improved.


[0222] The emission layer may include at least one condensed cyclic compound represented by one of Formulae 1A to 1C. The emission layer may include a dopant. The dopant may be at least one selected from a phosphorescent dopant and a fluorescent dopant.

[0223] The emission layer includes a host and a dopant, and the host may include at least one condensed cyclic compound represented by one of Formulae 1A to 1C.


[0224] According to an embodiment, the emission layer includes a first host and a second host.

[0225] The condensed cyclic compound represented by one of Formulae 1A to 1C may be included in the emission layer. The first host may include the condensed cyclic compound represented by one of Formulae 1A to 1C, and the second host may include at least one selected from a first compound represented by Formula 41 and a second compound represented by Formula 61:


Formula 41



Formula 61



Formula 61A



Formula 61B



[0220] Then, an emission layer (EML) may be formed on the hole transport region by vacuum deposition, spin coating, casting, LB deposition, or the like. When the emission layer is formed by vacuum deposition or spin coating, the deposition or coating conditions may be similar to those applied to form the hole injection layer although the deposition or coating conditions may vary according to the material that is used to form the emission layer.

[0221] When the organic light-emitting device is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, and a blue emission layer. According to another embodiment, due to a stacking structure including a red emission layer, a green emission layer, and/or a blue emission layer, the emission layer may emit white light.

[0226] wherein in the formulae above,

[0227] X_{41} is $N-[(L_{42})_{a42}-(R_{42})_{b42}]$, S, O, $S(=O)$, $S(=O)_2$, $C(=O)$, $C(R_{43})(R_{44})$, $Si(R_{43})(R_{44})$, $P(R_{43})$, $P(=O)(R_{43})$, or $C=N(R_{43})$;

[0228] Ring A_{61} in Formula 61 is represented by Formula 61A;

[0229] Ring A_{62} in Formula 61 is represented by Formula 61B;

[0230] X_{61} is $N-[(L_{62})_{a62}-(R_{62})_{b62}]$, S, O, $S(=O)$, $S(=O)_2$, $C(=O)$, $C(R_{63})(R_{64})$, $Si(R_{63})(R_{64})$, $P(R_{63})$, $P(=O)(R_{63})$, or $C=N(R_{63})$;

[0231] X_{71} is $C(R_{71})$ or N;

[0232] X_{72} is $C(R_{72})$ or N;

[0233] X_{73} is $C(R_{73})$ or N;

[0234] X_{74} is $C(R_{74})$ or N;

[0235] X_{75} is C(R_{75}) or N;
[0236] X_{76} is C(R_{76}) or N;
[0237] X_{77} is C(R_{77}) or N;
[0238] X_{78} is C(R_{78}) or N;
[0239] Ar_{41} , L_{41} , L_{42} , L_{61} , and L_{62} are each independently selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_2 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed hetero-polycyclic group;
[0240] n_1 and n_2 are each independently an integer selected from 0 to 3;
[0241] a_{41} , a_{42} , a_{61} , and a_{62} are each independently an integer selected from 0 to 3;
[0242] R_{41} to R_{44} , R_{51} to R_{54} , R_{61} to R_{64} , and R_{71} to R_{79} are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed hetero-polycyclic group, —N(Q_1)(Q_2), —Si(Q_3)(Q_4)(Q_5), and —B(Q_6)(Q_7);
[0243] b_{41} , b_{42} , b_{51} to b_{54} , b_{61} , b_{62} , and b_{79} are each independently an integer selected from 1 to 3;
[0244] at least one of substituents of the substituted C_1 - C_{60} alkylene group, substituted C_2 - C_{60} alkenylene group, substituted C_3 - C_{10} cycloalkylene group, substituted C_2 - C_{10} heterocycloalkylene group, substituted C_3 - C_{10} cycloalkenylene group, substituted C_2 - C_{10} heterocycloalkenylene group, substituted C_6 - C_{60} arylene group, substituted C_2 - C_{60} heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed hetero-polycyclic group, substituted C_1 - C_{60} alkyl group, substituted C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group, substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10} cycloalkyl group, substituted C_2 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substituted C_2 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_2 - C_{60} heteroaryl group, a substituted monovalent non-aromatic condensed polycyclic group, and a substituted monovalent non-aromatic condensed hetero-polycyclic group may be selected from
[0245] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic

acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0246] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed hetero-polycyclic group, —N(Q_{11})(Q_{12}), —Si(Q_{13})(Q_{14})(Q_{15}), and —B(Q_{16})(Q_{17});

[0247] a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed hetero-polycyclic group;

[0248] a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q_{21})(Q_{22}), —Si(Q_{23})(Q_{24})(Q_{25}), and —B(Q_{26})(Q_{27}); and

[0249] —N(Q_{31})(Q_{32}), —Si(Q_{33})(Q_{34})(Q_{35}), and —B(Q_{36})(Q_{37});

[0250] wherein Q_1 to Q_7 , Q_{11} to Q_{17} , Q_{21} to Q_{27} , and Q_{31} to Q_{37} may be each independently selected from a hydrogen, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed hetero-condensed polycyclic group.

[0251] According to an embodiment, R_{41} to R_{44} , R_{51} to R_{54} , R_{61} to R_{64} , and R_{71} to R_{79} may be each independently selected from

[0252] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a

salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0253] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, and a phosphoric acid or a salt thereof;

[0254] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furanyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl, an acridinyl, a phenanthrolinyl, a phenazinyl, a benzoimidazolyl group, a benzofuranyl group, a benzothiofenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiofenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group, each substituted with at least one selected from a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furanyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl, an acridinyl, a phenanthrolinyl, a phenazinyl, a benzoimidazolyl group, a benzofuranyl group, a benzothiofenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiofenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and a biphenyl group; and


[0255] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiofenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiofenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, carboxylic acid or a salt thereof, sulfonic acid or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, —Si(Q₃₃)(Q₃₄)(Q₃₅), a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiofenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiofenyl group, a benzocarbazolyl group, a dibenzocarbazolyl

group, an imidazopyridinyl group, an imidazopyrimidinyl group, and a biphenyl group; and


[0256] —Si(Q₃)(Q₄)(Q₅),

[0257] wherein Q₃ to Q₅ and Q₃₃ to Q₃₅ are each independently selected from a hydrogen, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group, but are not limited thereto.


[0258] For example, the first compound may be represented by one of Formulae 41-1 to 41-12 below, and the second compound may be represented by one of Formulae 61-1 to 61-6:



Formula 41-1




Formula 41-2




Formula 41-3

-continued


Formula 41-4




Formula 41-5



Formula 41-6




Formula 41-7




Formula 41-8


-continued




Formula 41-9




Formula 41-10




Formula 41-11



Formula 41-12




Formula 61-1




Formula 61-2


-continued




Formula 61-3



Formula 61-4



Formula 61-5




Formula 61-6

[0259] X_{41} , X_{61} , L_{41} , L_{42} , a_{41} , a_{42} , L_{61} , L_{62} , a_{61} , a_{62} , R_{41} to R_{44} , b_{41} , b_{42} , R_{61} to R_{64} , b_{61} , b_{62} , R_{71} to, R_{76} and b_{79} in Formulae 41-1 to 41-12 and 61-1 to 61-6 are as described above.


[0260] For example, when the second host includes the first compound represented by Formula 41, at least one selected from R_{41} , R_{42} , and R_{52} of the first compound, and when the second host includes the second compound represented by Formula 61, at least one selected from R_{61} and R_{62} of the second compound may be each independently selected from

[0261] a thiophenyl group, a furanyl group, a carbazolyl group, an acridinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group; and



A1

[0262] a thiophenyl group, a furanyl group, a carbazolyl group, acridinyl, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, carboxylic acid or a salt thereof, sulfonic acid or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a phenyl group substituted with C₁-C₂₀ alkyl group, a naphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a biphenyl group, and —Si(Q₃₃)(Q₃₄)(Q₃₅),




A2

[0263] wherein Q₃₃ to Q₃₅ may be each independently selected from a hydrogen, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group.

[0264] In some embodiments, when the second host includes the first compound represented by Formula 41, at least one selected from R₄₁, R₄₂, and R₅₂ of the first compound, and when the second host includes the second compound represented by Formula 61, at least one selected from R₆₁ and R₆₂ of the second compound may be represented by one of Formulae 5-1 to 5-6 and 6-1 to 6-5.


[0265] According to an embodiment, the first compound represented by Formula 41 may include one of Compounds A1 to A83 below, and the second compound represented by Formula 61 may include one of Compounds B1 to B20, but they are not limited thereto:

-continued




A3


-continued



A5




A4




A6

-continued




A7

-continued




A10





A11


A8



A9




A12




A13

-continued




A14


-continued




A17




A15



A18




A16




A19

-continued


A20




A21



A22




A23




-continued


A24



A25




A26




-continued


A27



A28




A29




-continued


A30



A31




-continued




A32

-continued




A34




A35

A33




A36




A37

-continued


A38



A39




A40




-continued


A41



A42




A43




-continued

A44




-continued


A47




A45




A48



A46




A49




-continued


A50




A51



A52




A53




-continued

A54




A55




-continued


A56



A57




A58




-continued


A59




A60



A61




-continued




A62


-continued




A65




A63



A66




A64




A67

-continued




A68


-continued




A71




A69




A72



A70




A73




A74

-continued




A75


-continued




A78




A76



A79




A77




A80

-continued




A81

-continued




B1




B2


A82




B3




B4




A83



B5




-continued




B6


-continued




B10




B7




B11




B8



B12








B9





B13

-continued



-continued




[0266] When the first host and the second host all are included in an emission layer, holes and electrons provided to the emission layer may be effectively controlled. Accordingly, an organic light-emitting device with high luminescent efficiency and long lifespan may be obtained.

[0267] A weight ratio of the first host to the second host in the emission layer may be each independently in a range of 1:99 to 99:1, for example, 10:90 to 90:10. When the weight ratio of the first host and the second host is within these ranges, holes and electrons provided to the emission layer may be effectively controlled.

[0268] A dopant in the emission layer may be a fluorescent dopant that emits light according to a fluorescent emission mechanism or a phosphorescent dopant that emits light according to a phosphorescent emission mechanism.

[0269] According to an embodiment, the dopant in the emission layer may be a phosphorescent dopant, and the phosphorescent dopant may include an organometallic compound represented by Formula 81 below:

Formula 81



[0270] wherein in Formula 81,
 [0271] M may be selected from iridium (Ir), platinum (Pt), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), and thulium (Tm);

[0272] Y_1 to Y_4 are each independently carbon (C) or nitrogen (N);

[0273] Y_1 and Y_2 are linked via a single bond or a double bond, and Y_3 and Y_4 are linked via a single bond or a double bond;

[0274] CY_1 and CY_2 are each independently selected from a benzene, a naphthalene, a fluorene, a spiro-fluorene, an indene, a pyrrole, a thiophene, a furan, an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isooxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, a quinoline, an isoquinoline, a benzoquinoline, a quinoxaline, a quinoxaline, a carbazole, a benzoimidazole, a benzofuran, a benzothiophene, an isobenzothiophene, a benzooxazole, an isobenzooxazole, a triazole, a tetrazole, an oxadiazole, a triazine, a dibenzofuran, and a dibenzothiophene, and CY_1 and CY_2 are optionally linked to each other through a single bond or an organic linking group;

[0275] R_{81} to R_{82} may be each independently a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, $-SF_5$, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_1)(Q_2)$, $-Si(Q_3)(Q_4)(Q_5)$, or $-B(Q_6)(Q_7)$;

[0276] a_{81} and a_{82} are each independently an integer of 1 to 5;


[0277] n_{81} is an integer of 0 to 4;

[0278] n_{82} is 1, 2, or 3; and






[0279] L_{81} is a monovalent organic ligand, a divalent organic ligand, or a trivalent organic ligand.

[0280] R_{81} and R_{82} may be understood by referring to the description provided herein in connection with R_{41} .






[0281] The phosphorescent dopant may include at least one of Compounds PD1 to PD8 below, but is not limited thereto:



-continued




-continued




-continued

PD7




PD12

PD8




PD13

PD9




PD14

PD10







PD15

PD11




PD16

-continued




-continued

PD17




PD21

PD18




PD22

PD19




PD23

PD20








PD24

PD26




-continued




-continued

PD26




PD31

PD27




PD32

PD28




PD33

PD29








PD34

PD30




PD35

-continued




-continued

PD36




PD41

PD37




PD42

PD38




PD43

PD39









PD44

PD40




PD45

-continued




-continued

PD46




PD52

PD47




PD53

PD48




PD54

PD49




PD55

PD50




PD51



PD56


-continued




PD57

-continued


PD62




PD58



PD59




PD63




PD60

PD64





PD61






PD65



PD66




-continued




-continued

PD67




PD72

PD68




PD73

PD69




PD74

PD70





PD75

PD71



PD76


-continued



PD77

[0282] According to another embodiment, the phosphorescent dopant may include PtOEP illustrated below:


PD78




PtOEP


[0283] The fluorescent dopant may include at least one selected from DPAVBi, BDAVBi, TBPe, DCM, DCJTb, Coumarin 6, and C545T.

DPAVBi



BDAVBi





[0284] When the emission layer includes a host and a dopant, an amount of the dopant may be in a range of about 0.01 to about 20 parts by weight based on 100 parts by weight of the host, but is not limited thereto.

[0285] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0286] Then, an electron transport region may be disposed on the emission layer.


[0287] The electron transport region may include at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.



[0288] For example, the electron transport region may have a structure of hole blocking layer/electron transport layer/electron injection layer or a structure of electron transport layer/electron injection layer, but the structure of the electron transport region is not limited thereto. The electron transport layer may have a single-layered structure or a multi-layer structure including two or more different materials.

[0289] Conditions for forming the hole blocking layer, the electron transport layer, and the electron injection layer

which constitute the electron transport region may be understood by referring to the conditions for forming the hole injection layer.

[0290] When the electron transport layer includes a hole blocking layer, the hole blocking layer may include, for example, at least one of BCP and Bphen, but may also include other materials.






[0291] A thickness of the hole blocking layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thickness of the hole blocking layer is within these ranges, the hole blocking layer may have excellent hole blocking characteristics without a substantial increase in driving voltage.




[0292] The electron transport layer may further include, in addition to the organometallic compound represented by Formula 1, at least one selected from BCP, Bphen, Alq₃, Balq, TAZ, and NTAZ.

[0293] According to another embodiment, the electron transport layer may include at least one of ET1 and ET2, but are not limited thereto:


ET1



ET2

Alq₃

BAIq




TAZ


[0294] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

[0295] Also, the electron transport layer may further include, in addition to the materials described above, a metal-containing material.

[0296] The metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.



ET-D1



ET-D2

[0297] The electron transport region may include an electron injection layer (EIL) that allows electrons to be easily provided from a second electrode **19**.

[0298] The electron injection layer may include at least one selected from, LiF, NaCl, CsF, Li₂O, BaO, and LiQ.

[0299] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

[0300] The second electrode **19** is disposed on the organic layer **15**. The second electrode **19** may be a cathode. A material for forming the second electrode **19** may be metal, an alloy, an electrically conductive compound, and a combination thereof, which have a relatively low work function. For example, lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag) may be used as the material for forming the second electrode **19**. To manufacture a top emission type light-emitting device, a transmissive electrode formed using ITO or IZO may be used as the second electrode **19**.

[0301] Hereinbefore, the organic light-emitting device has been described with reference to FIG. **1**, but is not limited thereto.

[0302] A C₁-C₆₀ alkyl group used herein refers to a linear or branched aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms. Detailed examples thereof are a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an isomethyl group, and a hexyl group. A C₁-C₆₀ alkylene group used herein refers to a divalent group having the same structure as the C₁-C₆₀ alkyl group.

[0303] A C₁-C₆₀ alkoxy group used herein refers to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is

the C₁-C₆₀ alkyl group). Detailed examples thereof are a methoxy group, an ethoxy group, and an isopropoxy group.

[0304] A C₂-C₆₀ alkenyl group used herein refers to a hydrocarbon group formed by substituting at least one carbon double bond in the middle or at the terminal of the C₂-C₆₀ alkyl group. Detailed examples thereof are an ethenyl group, a propenyl group, and a butenyl group. A C₂-C₆₀ alkenylene group used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkenyl group.

[0305] A C₂-C₆₀ alkynyl group used herein refers to a hydrocarbon group formed by substituting at least one carbon triple bond in the middle or at the terminal of the C₂-C₆₀ alkyl group. Detailed examples thereof are an ethynyl group, and a propynyl group. A C₂-C₆₀ alkynylene group used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

[0306] A C₃-C₁₀ cycloalkyl group used herein refers to a monovalent hydrocarbon monocyclic group having 3 to 10 carbon atoms. Detailed examples thereof are a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. A C₃-C₁₀ cycloalkylene group used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

[0307] A C₂-C₁₀ heterocycloalkyl group used herein refers to a monovalent monocyclic group having at least one hetero atom selected from N, O, P, and S as a ring-forming atom and 2 to 10 carbon atoms. Detailed examples thereof are a tetrahydrofuran group, and a tetrahydrothiophenyl group. A C₂-C₁₀ heterocycloalkylene group used herein refers to a divalent group having the same structure as the C₂-C₁₀ heterocycloalkyl group.

[0308] A C₃-C₁₀ cycloalkenyl group used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one double bond in the ring thereof and does not have aromaticity. Detailed examples thereof are a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. A C₃-C₁₀ cycloalkenylene group used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

[0309] A C₂-C₁₀ heterocycloalkenyl group used herein refers to a monovalent monocyclic group that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, 2 to 10 carbon atoms, and at least one double bond in its ring. Detailed examples of the C₂-C₁₀ heterocycloalkenyl group are a 2,3-dihydrofuran group and a 2,3-dihydrothiophenyl group. A C₂-C₁₀ heterocycloalkenylene group used herein refers to a divalent group having the same structure as the C₂-C₁₀ heterocycloalkenyl group.

[0310] A C₆-C₆₀ aryl group used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C₆-C₆₀ arylene group used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Detailed examples of the C₆-C₆₀ aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be fused to each other.

[0311] A C₂-C₆₀ heteroaryl group used herein refers to a monovalent group having a carbocyclic aromatic system that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, and 2 to 60 carbon atoms. A C₂-C₆₀ heteroarylene group used herein refers to a divalent group having a carbocyclic aromatic system that has at least one

hetero atom selected from N, O, P, and S as a ring-forming atom, and 2 to 60 carbon atoms. Examples of the C₂-C₆₀ heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₂-C₆₀ heteroaryl group and the C₂-C₆₀ heteroarylene group each include two or more rings, the rings may be fused to each other.

[0312] A C₆-C₆₀ aryloxy group used herein indicates —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and a C₆-C₆₀ arylthio group indicates —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

[0313] A monovalent non-aromatic condensed polycyclic group used herein refers to a monovalent group that has two or more rings condensed to each other, only carbon atoms (for example, the number of carbon atoms may be in a range of 8 to 60) as ring forming atoms, wherein the molecular structure as a whole is non-aromatic in the entire molecular structure. An example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. A divalent non-aromatic condensed polycyclic group used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0314] A monovalent non-aromatic condensed hetero-polycyclic group used herein refers to a monovalent group that has two or more rings condensed to each other, has a heteroatom selected from N, O, P, and S, other than carbon atoms (for example, the number of carbon atoms may be in a range of 2 to 60), as ring-forming atoms, wherein the molecular structure as a whole is non-aromatic in the entire molecular structure. An example of the monovalent non-aromatic condensed hetero-polycyclic group is a carbazolyl group. A divalent non-aromatic condensed hetero-polycyclic group used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed hetero-polycyclic group.

[0315] In the present specification, at least one of substituents of substituents of the substituted C₁-C₆₀ alkylene group, substituted C₂-C₆₀ alkenylene group, substituted C₂-C₆₀ alkynylene group, substituted C₃-C₁₀ cycloalkylene group, substituted C₂-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₂-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₂-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed hetero-polycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₃-C₁₀ cycloalkyl group, substituted C₂-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₂-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₂-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed hetero-polycyclic group may be selected from

[0316] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0317] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed hetero-polycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

[0318] a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed hetero-polycyclic group;

[0319] a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed hetero-polycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed hetero-polycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇); and

[0320] —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇), wherein

[0321] Q₁ to Q₇, Q₁₁ to Q₁₇, and Q₃₁ to Q₃₇ used herein may be each independently selected from a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed hetero-polycyclic group.

[0322] In the present specification, at least one of substituents of the substituted C₁-C₆₀ alkylene group, substituted C₂-C₆₀ alkenylene group, substituted C₂-C₆₀ alkynylene group, substituted C₃-C₁₀ cycloalkylene group, substituted C₂-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₂-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₂-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed hetero-polycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₃-C₁₀ cycloalkyl

group, substituted C_2 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substituted C_2 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_2 - C_{60} heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from

[0323] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0324] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q_{11})(Q_{12}), —Si(Q_{13})(Q_{14})(Q_{16}), and —B(Q_{16})(Q_{17});

[0325] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pyrrolyl group, an imidazolyl group, a pyrazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzooxazolyl group, a benzoimidazolyl group, a furanyl group, a benzofuranyl group, a thiophenyl group, a benzothiophenyl group, a thiazolyl group, an isothiazolyl group, a benzothiazolyl group, an isooxazolyl group, an oxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyrimidinyl group, and an imidazopyridinyl group, each substituted with at least one selected from a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, cyclooctyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a phthalazinyl group, a quinoxalinyl group, a cinnolinyl group, and a quinazolinyl group;

[0326] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pyrrolyl group, an imidazolyl group, a pyrazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furanyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzooxazolyl group, a benzoimidazolyl group, a furanyl group, a benzofuranyl group, a thiophenyl group, a benzothiophenyl group, a thiazolyl group, an isothiazolyl group, a benzothiazolyl group, an isooxazolyl group, an oxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyrimidinyl group, and an imidazopyridinyl group, each substituted with at least one selected from a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a phthalazinyl group, a quinoxalinyl group, a cinnolinyl group, and a quinazolinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, monovalent non-aromatic hetero-condensed polycyclic group, —N(Q_{21})(Q_{22}), —Si(Q_{23})(Q_{24})(Q_{25}), and —B(Q_{26})(Q_{27}); and

[0327] —N(Q_{31})(Q_{32}), —Si(Q_{33})(Q_{34})(Q_{35}), and —B(Q_{36})(Q_{37}),

[0328] wherein Q_1 to Q_7 , Q_{11} to Q_{17} , and Q_{31} to Q_{37} may be each independently a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pyrrolyl group, an imidazolyl group, a pyrazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furanyl group, a quinolinyl group, an isoquinolinyl

group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzooxazolyl group, a benzoimidazolyl group, a furanyl group, a benzofuranyl group, a thiophenyl group, a benzothiophenyl group, a thiazolyl group, an isothiazolyl group, a benzothiazolyl group, an isooxazolyl group, an oxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyrimidinyl group, and an imidazopyridinyl group, each substituted with at least one selected from a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a phthalazinyl group, a quinoxalinyl group, a cinnolinyl group, or a quinazolinyl group, but they are not limited thereto.

[0329] The term “biphenyl group” as used therein refers to “a phenyl group substituted with a phenyl group.”



[0330] Hereinafter, a compound and an organic light-emitting device according to embodiments are described in detail with reference to Synthesis Example and Examples. However, the organic light-emitting device is not limited thereto. The wording “B was used instead of A” used in describing Synthesis Examples means that a molar equivalent of A was identical to a molar equivalent of B.


EXAMPLES

Synthesis Example 1

Synthesis of Compound 3

[0331]





Synthesis of Intermediate 3-5

[0332] Intermediate 3-6 (10 g, 59.8 mmol) and Intermediate A (8.44 g, 59.8 mmol) were dissolved in 200 mL of DMSO, and then, Cs_2CO_3 (23.4 g, 71.8 mmol) was added to the resultant and stirred for 15 hours. 500 mL of water was added to the reaction product, and the resultant was filtered and a solid was collected. The solid was dissolved in 300 mL of methylene chloride, and 300 mL of water was added thereto, and then the resultant was extracted. An organic layer obtained therefrom was dried by using MgSO_4 , and the residual obtained by evaporating the solvent was separation-purified by silica gel column chromatography to obtain Intermediate 3-5 (12.93 g, yield of 75%). The obtained compound was identified by LC-MS.

[0333] $\text{C}_{18}\text{H}_{12}\text{N}_2\text{O}_2$: M+ 288.09

Synthesis of Intermediate 3-4

[0334] Intermediate 3-5 (10 g, 34.7 mmol) and $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (25.16 g, 111 mmol) was added to EtOH and then, the mixture was heated to 70° C. for 8 hours or more. When the reaction stopped, the reaction product was cooled to room temperature, and then, concentrated, and an organic layer obtained by adding 300 mL of methylene chloride and 300 mL of NaHCO_3 (aq) thereto was dried by using MgSO_4 , and the residual obtained by evaporating the solvent was separation-purified by silica gel column chromatography to obtain Intermediate 3-4 (6.27 g, yield of 70%). The obtained compound was identified by LC-MS.

[0335] $\text{C}_{18}\text{H}_{14}\text{N}_2$: M+ 258.12

Synthesis of Intermediate 3-3

[0336] Intermediate 3-4 (6 g, 23.23 mmol), 60 mL of acetic acid, 5 mL of conc. H_2SO_4 in a reaction container were

deposited to an ice bath to decrease the temperature to 10° C. Then, NaNO_2 (1.6 g, 1 eq) was dissolved in 150 mL of water, and the resultant was added dropwise thereto for 20 minutes, and stirred for 10 minutes, and then, the temperature of the mixture was raised to 130° C., and then stirred for 20 minutes. When the reaction stopped, the temperature was decreased to room temperature, and then, 300 mL of water was added thereto, and filtered to isolate a precipitated solid. The residual was separation-purified by silica gel column chromatography to obtain Intermediate 3-3 (4.03 g, 72%). The obtained compound was identified by LC-MS.

[0337] $\text{C}_{18}\text{H}_{11}\text{N}$: M+ 241.09

Synthesis of Intermediate 3-2

[0338] Intermediate 3-3 (3 g, 12.43 mmol) and NBS (2.21 g, 12.43 mmol) were dissolved in 100 mL of methylene chloride, and then, the mixture was stirred at room temperature for 15 hours. When the reaction stopped, the reaction product was cooled to room temperature, and 200 mL of methylene chloride was added thereto, and then 200 mL of water was added dropwise thereto, and an obtained organic layer was dried by using MgSO_4 , and the residual obtained by evaporating the solvent was separation-purified by silica gel column chromatography to obtain Intermediate 3-2 (2.03 g, 51%). The obtained compound was identified by LC-MS.

[0339] $\text{C}_{18}\text{H}_{10}\text{BrN}$: M+ 319.00

Synthesis of Intermediate 3-1

[0340] Intermediate 3-2 (1.5 g, 4.68 mmol), Intermediate B (1.19 g, 5.62 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.27 g, 0.234 mmol), and K_2CO_3 (3.24 g, 23.4 mmol) were mixed with 100 mL of THF and 30 mL of distilled water. The mixture was heated to a temperature of 120° C. and stirred for 24 hours while refluxing, and then, cooled to room temperature, and extracted using 200 mL of water and 200 mL of methylene chloride. An organic layer obtained therefrom was dried by using MgSO_4 , and the residual obtained by evaporating the solvent was separation-purified by silica gel column chromatography to obtain Intermediate 3-1 (1.5 g, 79%). The obtained compound was identified by LC-MS.

[0341] $\text{C}_{30}\text{H}_{18}\text{N}_2$: M+ 406.15

Synthesis of Compound 3

[0342] Intermediate 3-1 (1 g, 2.46 mmol) was dissolved in 30 mL of DMF, and then, sodium hydride was added thereto, and the result was stirred for 1 hour. Intermediate C (0.8 g, 2.95 mmol) dissolved in 10 mL of DMF was added thereto, and then, stirred for 12 hours at room temperature. Resulting precipitate was washed with methylene chloride, and then added to dichlorobenzene, and the result was heated, and then, while cooling to room temperature, recrystallized and separated to obtain Compound 3 (1.14 g, 65%). The obtained compound was identified by LC-MS and NMR.


[0343] $\text{C}_{51}\text{H}_{31}\text{N}_5$: M+ 713.26

[0344] $^1\text{H NMR}$ (CDCl_3 , 300 MHz) δ (ppm) 8.36-8.19 (m, 11H), 7.89 (s, 1H), 7.68-7.50 (m, 16H), 7.20 (m, 3H)

Synthesis Example 2

Synthesis of Compound 2

[0345]



[0346] Compound 2 (68%) was synthesized in the same manner as in Synthesis Example 1, except that in synthesizing Compound 3, Intermediate D was used instead of Intermediate C. The obtained compound was identified by LC-MS and NMR.


[0347] $C_{45}H_{27}N_5$: M+ 637.75

[0348] 1H NMR ($CDCl_3$, 300 MHz) δ (ppm) 7.36-8.30 (m, 5H), 8.19-8.13 (m, 4H), 7.89 (s, 1H), 7.65-7.50 (m, 14H), 7.20 (m, 3H)

Synthesis Example 3

Synthesis of Compound 4

[0349]



[0350] Compound 4 (70%) was synthesized in the same manner as in Synthesis Example 1, except that in synthesizing Compound 3, Intermediate E was used instead of Intermediate C. The obtained compound was identified by LC-MS and NMR.


[0351] $C_{46}H_{28}N_4$: M+ 636.23

[0352] 1H NMR ($CDCl_3$, 300 MHz) δ (ppm) 8.9 (s, 1H), 8.30-8.13 (m, 5H), 7.94-7.89 (m, 5H), 7.65-7.50 (m, 14H), 7.20 (m, 3H)

Synthesis Example 4

Synthesis of Compound 7

[0353]



[0354] Compound 7 (62%) was synthesized in the same manner as in Synthesis Example 1, except that in synthesizing Compound 3, Intermediate F was used instead of Intermediate C. The obtained compound was identified by LC-MS and NMR.


[0355] $C_{44}H_{26}N_4$: M+ 610.22

[0356] 1H NMR ($CDCl_3$, 300 MHz) δ (ppm) 8.30-8.13 (m, 6H), 7.89-7.80 (m, 5H), 7.65-7.50 (m, 12H), 7.20 (m, 3H)

Synthesis Example 5

Synthesis of Compound 8

[0357]



[0358] Compound 8 (62%) was synthesized in the same manner as in Synthesis Example 1, except that in synthesizing Compound 3, Intermediate G was used instead of Intermediate C. The obtained compound was identified by LC-MS and NMR.


[0359] $C_{50}H_{30}N_4$: M+ 686.25

[0360] 1H NMR ($CDCl_3$, 300 MHz) δ (ppm) 8.30-8.13 (m, 8H), 7.89-7.75 (m, 7H), 7.65-7.41 (m, 12H), 7.20 (m, 3H)


Synthesis Example 6

Synthesis of Compound 76


[0361]




Intermediate 76-3



Intermediate 76-1



Intermediate D



Compound 76

Synthesis of Intermediate 76-1

[0362] Intermediate 76-1 was prepared in the same manner as used in synthesizing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 76-2 and Intermediate 76-3 were respectively used instead of Intermediate 3-2 and Intermediate B.

Synthesis of Compound 76

[0363] Compound 76 (yield of 74%) was prepared in the same manner as used in synthesizing Intermediate 3 of Synthesis Example 1, except that in synthesizing Compound 3, Intermediate 76-1 and Intermediate D were respectively used instead of Intermediate 3-1 and Intermediate C. The obtained compound was identified by LC-MS and NMR.


[0364] $\text{C}_{45}\text{H}_{27}\text{N}_5$; M^+ 686.25

[0365] $^1\text{H NMR}$ (CDCl_3 , 300 MHz) δ (ppm) 8.62 (d, 1H), 8.36 (m, 4H), 8.22-8.19 (m, 4H), 7.74 (s, 1H), 7.65-7.50 (m, 14H), 7.20 (m, 3H)


Synthesis Example 7

Synthesis of Compound 60


[0366]



Intermediate 60-3



Intermediate 60-2



Synthesis of Intermediate 60-2

[0367] Intermediate 60-2 was synthesized in the same manner as in synthesizing Intermediate 3-2 of Synthesis Example 1, except that Intermediate 60-3 was used instead of Intermediate 3-3.

Synthesis of Intermediate 60-1


[0368] Intermediate 60-1 was prepared in the same manner as used in synthesizing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 60-2 and Intermediate H were respectively used instead of Intermediate 3-2 and Intermediate B.

Synthesis of Compound 60

[0369] Compound 60 (yield of 51%) was prepared in the same manner as used in synthesizing Intermediate 3 of Synthesis Example 1, except that Intermediate 60-1 and Intermediate I were respectively used instead of Intermediate 3-1 and Intermediate C.

[0370] $C_{38}H_{22}N_4$: M+ 534.18

[0371] 1H NMR ($CDCl_3$, 300 MHz) δ (ppm) 8.87 (d, 1H), 8.51-8.01 (m, 8H), 7.89 (s, 1H), 7.70-7.43 (m, 8H), 7.20-7.15 (m, 4H)



Evaluation Example 1

Evaluation on HOMO, LUMO, and Triplets (T1) Energy Levels

[0372] HOMO, LUMO and T1 energy levels of Compounds 2, 3, and 7 were evaluated according to the method indicated in Table 2, and results thereof are shown in Table 3.

TABLE 2

HOMO energy level evaluation method	A potential (V)-current (A) graph of each compound was obtained by using cyclic voltammetry (CV) (electrolyte: 0.1M Bu_4NClO_4 /solvent: CH_2Cl_2 /electrode: 3 electrode system (working electrode: GC, reference electrode: Ag/AgCl, auxiliary electrode: Pt)), and then, from reduction onset of the graph, a HOMO energy level of the compound was calculated.
LUMO energy level evaluation method	Each compound was diluted at a concentration of 1×10^{-5} M in $CHCl_3$, and an UV absorption spectrum thereof was measured at room temperature by using a Shimadzu UV-350 spectrometer, and a LUMO energy level thereof was calculated by using an optical band gap (Eg) from an edge of the absorption spectrum.
T1 energy level evaluation method	A mixture (each compound was dissolved in an amount of 1 mg in 3 cc of toluene) of toluene and each compound was loaded into a quartz cell, and then, the resultant quartz cell was loaded into liquid nitrogen (77 K) and a photoluminescence spectrum thereof was measured by using a device for measuring photoluminescence, and the obtained spectrum was compared with a photoluminescence spectrum measured at room temperature, and peaks observed only at low temperature were analyzed to calculate T1 energy levels.

TABLE 3

Compound No.	HOMO (eV)	LUMO (eV)	T1 energy level (eV)
	(calc.)	(calc.)	
2	-5.515	-2.376	2.63
3	-5.303	-2.298	2.66
7	-5.587	-2.385	2.47

[0373] From Table 3, it is confirmed that the compounds above have electric characteristics that are suitable for use as a material for forming an organic light-emitting device.

Evaluation Example 2


Thermal Characteristics Evaluation

[0374] Thermal analysis (N_2 atmosphere, temperature range: from room temperature to $800^\circ C$. ($10^\circ C./min$)-TGA, from room temperature to $400^\circ C$.-DSC, Pan Type: Pt Pan in disposable Al Pan (TGA), and disposable Al pan (DSC)) was performed on Compounds 7 and 8 and Compounds C and D by using thermo gravimetric analysis (TGA) and differential scanning calorimetry (DSC), and results thereof are shown in Table 4. As shown in Table 4, it was confirmed that the synthesized compounds had excellent thermal stability.

TABLE 4


Compound No.	Tg ($^\circ C$.)	Td (1%, $^\circ C$.)	Td (5%, $^\circ C$.)
Compound 7	149.45	447.8	486.4
Compound 8	167.33	482.5	521.5
Compound C	128.8	427.3	473.9
Compound D	144.2	462.5	502.1

Compound 7




-continued

Compound 8




Compound C



-continued

Compound D



Example 1

[0375] A glass substrate with ITO/Ag/ITO (70/1,000/70 Å) anode thereon was cut to a size of 50 mm×50 mm×0.5 mm, and then, ultrasonically cleaned with isopropyl alcohol, acetone, and pure water for 5 minutes each, and then exposed to ultraviolet (UV) light for 30 minutes and then ozone. Then, the glass substrate was loaded into a vacuum deposition apparatus.

[0376] 2-TNATA was deposited on the anode on the glass substrate to form a hole injection layer having a thickness of 600 Å, and then NPB was deposited on the hole injection layer to form a hole transport layer having a thickness of 1,000 Å to form a hole transport region.

[0377] Subsequently, Compound 3 (host) and Ir(ppy)₃ (dopant, referred to as Compound PD1 in this application) were co-deposited on the hole transport layer to form an emission layer having a thickness of 250 Å.

[0378] BCP was deposited on the emission layer to form a hole blocking layer having a thickness of 50 Å, and Alq₃ was deposited on the hole blocking layer to form an electron transport layer having a thickness of 350 Å, and LiF was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å to form an electron transport region.

[0379] Mg and Ag were deposited on the electron injection layer at a weight ratio of 90:10 to form a cathode having a thickness of 120 Å, thereby completing the manufacture of an organic light-emitting device.

Example 2

[0380] An organic light-emitting device was manufactured in the same manner as in Example 1, except that in forming an emission layer, as a host, Compound 2 was used instead of Compound 3.

Example 3

[0381] An organic light-emitting device was manufactured in the same manner as in Example 1, except that in forming an emission layer, as a host, Compound 4 was used instead of Compound 3.

Example 4

[0382] An organic light-emitting device was manufactured in the same manner as in Example 1, except that in forming an emission layer, as a host, Compound 76 was used instead of Compound 3.

Example 5

[0383] An organic light-emitting device was manufactured in the same manner as in Example 1, except that a thickness of the hole transport layer was 1,350 Å, and in forming an emission layer, Compound 7 was used as the host, PtOEP was used as the dopant, and a thickness of the emission layer was 400 Å.


Example 6

[0384] An organic light-emitting device was manufactured in the same manner as in Example 4, except that in forming an emission layer, as a host, Compound 8 was used instead of Compound 7.

Comparative Example 1

[0385] An organic light-emitting device was manufactured in the same manner as in Example 1, except that in forming an emission layer, as a host, Compound A was used instead of Compound 3.


Compound A



Comparative Example 2

[0386] An organic light-emitting device was manufactured in the same manner as in Example 1, except that in forming an emission layer, as a host, Compound B was used instead of Compound 3.

Compound B



Comparative Example 3

[0387] An organic light-emitting device was manufactured in the same manner as in Example 5, except that in forming an emission layer, as a host, Compound A was used instead of Compound 7.

Comparative Example 3

[0388] An organic light-emitting device was manufactured in the same manner as in Example 5, except that in forming an emission layer, as a host, Compound B was used instead of Compound 7.

Evaluation Example 4

Evaluation on Characteristics of an Organic Light-Emitting Device

[0389] Driving voltage, current density, brightness, luminescence color, efficiency, and LT_{97} (@10 milliamperes per square centimeter, mA/cm^2) of the organic light-emitting devices of Examples 1 to 6 and Comparative Examples 1 to 4 were evaluated by using PR650 Spectroscan Source Measurement Unit. (a product of PhotoResearch Company). Results thereof are shown in Table 6 below.

TABLE 6

	EML		Driving	Current	Brightness	Efficiency	Emission	
	Host	Dopant	Voltage	Density				
			(V)	(mA/cm^2)	(cd/m^2)	(cd/A)	color	LT_{97} (hr)
Example 1	Compound 3	$\text{Ir}(\text{ppy})_3$	5.4	10	6,829	68.3	Green	75
Example 2	Compound 2	$\text{Ir}(\text{ppy})_3$	5.3	10	6,724	67.2	Green	79
Example 3	Compound 4	$\text{Ir}(\text{ppy})_3$	5.9	10	6,709	67.1	Green	76
Example 4	Compound 76	$\text{Ir}(\text{ppy})_3$	5.6	10	6,171	61.7	Green	70
Example 5	Compound 7	PtOEP	5.8	10	3,422	34.2	Red	110
Example 6	Compound 8	PtOEP	6.0	10	3,117	31.2	Red	114
Comparative Example 1	Compound A	$\text{Ir}(\text{ppy})_3$	8.2	10	4,564	45.6	Green	32
Comparative Example 2	Compound B	$\text{Ir}(\text{ppy})_3$	7.3	10	4,276	42.8	Green	35
Comparative Example 3	Compound A	PtOEP	7.9	10	1,810	18.1	Red	52
Comparative Example 4	Compound B	PtOEP	7.7	10	1,921	19.2	Red	50

[0390] From Table 6, it was confirmed that the organic light-emitting devices of Examples 1 to 4 had lower driving voltage, higher efficiency, and longer lifespan of the organic light-emitting devices of Comparative Examples 1 and 2, and the organic light-emitting devices of Examples 5 and 6 had lower driving voltage, higher efficiency, and longer lifespan than the organic light-emitting devices of Comparative Examples 3 and 4.


[0391] The condensed cyclic compounds according to embodiments have excellent electric characteristics and thermal stability. Accordingly, organic light-emitting devices including the condensed cyclic compounds may have a low driving voltage, high efficiency, high brightness, and a long lifespan.

[0392] It should be understood that the exemplary embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as available for other similar features or aspects in other embodiments.

What is claimed is:


1. A condensed cyclic compound represented by one of Formulae 1A to 1C:

Formula 1A




-continued


Formula 1B




Formula 1C




Formula 2A




Formula 2B



Formula 2C




Formula 2D




-continued

Formula 2E



Formula 2F



wherein in the Formulae 1A to 1C and 2A to 2F,

Ring A is selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, a group represented by Formula 2A, and a group represented by Formula 2B;

Ring B₁ is a C₅-C₂₀ carbocyclic group or a C₂-C₂₀ heterocyclic group that does not comprise a nitrogen atom as a ring-forming atom;

Ring B₂, Ring B₃, Ring C₁, and Ring C₂ are each independently a C₅-C₂₀ carbocyclic group, or a C₂-C₂₀ heterocyclic group, provided that at least one selected from Ring B₂ and Ring B₃ is a C₂-C₂₀ heterocyclic group comprising a nitrogen atom as a ring-forming atom, and at least one selected from Ring C₁ and Ring C₂ is a C₂-C₂₀ heterocyclic group comprising a nitrogen atom as a ring-forming atom;

Pc₁ and Pc₂ in Formula 1A and 1B are each independently a group represented by Formulae 2A or a group represented by Formula 2B;

Cz₁, Cz₂, and Cz₃ in Formula 1C are each independently selected from a group represented by Formula 2C, a group represented by Formula 2D, a group represented by Formula 2E, and a group represented by Formula 2F;

c₁, c₂, c₁₁, c₁₂, c₁₃, p₁, p₂, p₁₁, p₁₂, and p₁₃ are each independently an integer selected from 0 to 3;

when c₁ and c₂ in Formula 1A and 1B both are 0, Ring A is a group represented by Formula 2A or a group represented by Formula 2B;

when Ring A in Formula 1A and 1B is neither a group represented by Formula 2A nor a group represented by Formula 2B;

$$c_1 + c_2 \neq 0;$$

at least one selected from Ring A in Formula 1A, all R₁ in Formula 1A, and all R₂ in Formula 1A is a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group,

a substituted or unsubstituted C_2 - C_{60} heteroaryl group, or a substituted or unsubstituted monovalent non-aromatic condensed hetero-polycyclic group, each comprising at least one Moiety 1 as a ring-forming moiety; the sum of c11, c12, and c13 in Formula 1C is not 0;

when each of Cz_1 , Cz_2 , and Cz_3 in Formula 1C is independently a group represented by Formula 2C or a group represented by Formula 2D, at least one selected from all R_{11} in Formula 1C, all R_{12} in Formula 1C, and all R_{13} in Formula 1C is a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, or a substituted or unsubstituted monovalent non-aromatic condensed hetero-polycyclic group, each comprising at least one Moiety 1 as a ring-forming moiety;

L_1 to L_3 , L_{11} to L_{13} , L_{21} to L_{24} , and L_{31} to L_{33} are each independently selected from a substituted or unsubstituted C_1 - C_{60} alkylene group, a substituted or unsubstituted C_2 - C_{60} alkenylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_2 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed hetero-polycyclic group;

a1 to a3, a11 to a13, a21 to a24, and a31 to a33 are each independently an integer selected from 0 to 6;

R_1 , R_2 , R_{11} to R_{13} , R_{21} to R_{23} , and R_{31} to R_{33} are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed hetero-polycyclic group, —N(Q_1)(Q_2), —Si(Q_3)(Q_4)(Q_5), and —B(Q_6)(Q_7);

b1, b2, b11 to b13, b21 to b23, and b31 to b33 are each independently an integer selected from 0 to 4;

at least one of substituents of the substituted C_1 - C_{60} alkylene group, substituted C_2 - C_{60} alkenylene group, substituted C_3 - C_{10} cycloalkylene group, substituted C_2 - C_{10} heterocycloalkylene group, substituted C_3 - C_{10} cycloalkenylene group, substituted C_2 - C_{10} heterocycloalkenylene group, substituted C_6 - C_{60} arylene group,

substituted C_2 - C_{60} heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed hetero-polycyclic group, substituted C_1 - C_{60} alkyl group, substituted C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group, substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10} cycloalkyl group, substituted C_2 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substituted C_2 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_2 - C_{60} heteroaryl group, a substituted monovalent non-aromatic condensed polycyclic group, and a substituted monovalent non-aromatic condensed hetero-polycyclic group is selected from

- a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;
- a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed hetero-polycyclic group, —N(Q_{11})(Q_{12}), —Si(Q_{13})(Q_{14})(Q_{15}), and —B(Q_{16})(Q_{17});
- a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed hetero-polycyclic group;
- a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group,

a monovalent non-aromatic condensed heteropolycyclic group, $-\text{N}(\text{Q}_{21})(\text{Q}_{22})$, $-\text{Si}(\text{Q}_{23})(\text{Q}_{24})(\text{Q}_{25})$, and $-\text{B}(\text{Q}_{26})(\text{Q}_{27})$; and $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{Si}(\text{Q}_{33})(\text{Q}_{34})(\text{Q}_{35})$, and $-\text{B}(\text{Q}_{36})(\text{Q}_{37})$;

wherein Q_1 to Q_7 , Q_{11} to Q_{17} , Q_{21} to Q_{27} , and Q_{31} to Q_{37} are each independently selected from a hydrogen, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group:

=N=*

Moiety 1

wherein in Moiety 1, ** and *** each indicates a binding site to a ring-forming atom included in a ring comprising Moiety 1.

2. The condensed cyclic compound of claim 1, wherein Ring A in Formula 1A is selected from

a phenyl group, a naphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, phenanthridinyl, phenanthrolinyl, phenazinyl, a benzoimidazolyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzofuopyrazinyl group, a benzofuopyrimidinyl group, a benzothienopyrazinyl group, a benzothienopyrimidinyl group, a benzothienopyridinyl group, a group represented by Formula 2A, and a group represented by Formula 2B; and


a phenyl group, a naphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, phenanthridinyl, phenanthrolinyl, phenazinyl, a benzoimidazolyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzofuopyrazinyl group, a benzofuopyrimidinyl group, a benzothienopyrazinyl group, a benzothienopyrimidinyl group, and a

benzothienopyridinyl group, each substituted with one selected from a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a phenyl group substituted with a C_1 - C_{20} alkyl group, a naphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a biphenyl group, and $-\text{Si}(\text{Q}_{33})(\text{Q}_{34})(\text{Q}_{35})$,


wherein Q_{33} to Q_{35} are each independently selected from a hydrogen, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group.

3. The condensed cyclic compound of claim 1, wherein


Ring A in Formula 1A is represented by one selected from Formulae 4-1 to 4-51, Formulae 6-1 to 6-5, a group represented Formula 2A, and a group represented by Formula 2B:




Formula 4-1




Formula 4-2




Formula 4-3



Formula 4-4
















Formula 4-5



Formula 4-6

-continued



Formula 4-7

Formula 4-8

Formula 4-9

Formula 4-10

Formula 4-11

Formula 4-12

Formula 4-13

Formula 4-14

Formula 4-15












Formula 4-16

Formula 4-17

Formula 4-18

Formula 4-19

-continued



Formula 4-20

Formula 4-21

Formula 4-22

Formula 4-23

Formula 4-24

Formula 4-25

Formula 4-26











Formula 4-27

Formula 4-28

Formula 4-29


Formula 4-30

-continued




-continued


Formula 4-31




Formula 4-32




Formula 4-33




Formula 4-34




Formula 4-35




Formula 4-36




Formula 4-37




Formula 4-38



Formula 4-39



Formula 4-40



Formula 4-41

Formula 4-42

Formula 4-43

Formula 4-44

Formula 4-45

Formula 4-46


Formula 4-47

Formula 4-48


Formula 4-49

Formula 4-50


-continued




Formula 4-51




Formula 6-1




Formula 6-2



Formula 6-3



Formula 6-4



Formula 6-5

wherein in Formulae 4-1 to 4-51 and 6-1 to 6-5:

Y_{31} is O, S, C(Z_{33})(Z_{34}), N(Z_{35}), or Si(Z_{36})(Z_{37});

Z_{31} to Z_{37} are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group substituted with a C_1 - C_{20} alkyl group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, a quinoxalinyl group, a biphenyl group, and —Si(Q_{33})(Q_{34})(Q_{35}), wherein

Q_{33} to Q_{35} are each independently selected from a hydrogen, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a

triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group;

e_1 is an integer of 1 to 5;

e_2 is an integer of 1 to 7;

e_3 is an integer of 1 to 3;

e_4 is an integer of 1 to 4;

e_5 is an integer of 1 or 2;

e_6 is an integer of 1 to 6; and

* indicates a binding site to a neighboring atom.

4. The condensed cyclic compound of claim 1, wherein Ring B_1 is a benzene, a naphthalene, an anthracene, a phenanthrene, a benzofuran, or a benzothiophene.

5. The condensed cyclic compound of claim 1, wherein Ring B_2 , Ring B_3 , Ring C_1 , and Ring C_2 are each independently a benzene, a naphthalene, an anthracene, a phenanthrene, a benzofuran, a benzothiophene, a pyridine, a pyrimidine, a pyrazine, a benzofuopyrazine, a benzofuopyrimidine, benzofuopyridine, a benzothienopyrazine, a benzothienopyrimidine, or a benzothienopyridine;

at least one selected from Ring B_2 and Ring B_3 is a pyridine, a pyrimidine, a pyrazine, a benzofuopyrazine, a benzofuopyrimidine, a benzofuopyridine, a benzothienopyrazine, a benzothienopyrimidine, or a benzothienopyridine; and

at least one selected from Ring C_1 and Ring C_2 is a pyridine, a pyrimidine, a pyrazine, a benzofuopyrazine, a benzofuopyrimidine, a benzofuopyridine, a benzothienopyrazine, a benzothienopyrimidine, or a benzothienopyridine.

6. The condensed cyclic compound of claim 1, wherein $c_1=0$, $c_2=0$, and Ring A is a group represented by Formula 2A or a group represented by Formula 2B;

$c_1=1$ and $c_2=0$;

$c_1=1$, and $c_2=1$;

$c_{11}=1$, $c_{12}=0$, and $c_{13}=0$;

$c_{11}=0$, $c_{12}=0$, and $c_{13}=1$;

$c_{11}=1$, $c_{12}=1$, and $c_{13}=0$; or

$c_{11}=1$, $c_{12}=0$, and $c_{13}=1$.

7. The condensed cyclic compound of claim 1, wherein L_1 to L_3 , L_{11} to L_{13} , L_{21} to L_{24} , and L_{31} to L_{33} are each independently selected from

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylene group, a spirofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pyrrolylene group, an imidazolylene group, a pyrazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a

quinoxalinylene group, a quinazolinylene group, a cinolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzoxazolylene group, a benzoimidazolylene group, a furanylene group, a benzofuranylene group, a thiophenylylene group, a benzothiophenylylene group, a thiazolylylene group, an isothiazolylylene group, a benzothiazolylylene group, an isoxazolylene group, an oxazolylene group, a triazolylene group, a tetrazolylylene group, an oxadiazolylylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, an imidazopyrimidinylene group, and an imidazopyridinylylene group; and


a phenylene group, a pentalenylylene group, an indenylene group, a naphthylene group, an azulenylylene group, a heptalenylylene group, an indacenylene group, an acenaphthylene group, a fluorenylylene group, a spiro-fluorenylylene group, a phenalenylylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthrenylene group, a triphenylylylene group, a pyrenylene group, a chrysenylene group, a naphthacenylylene group, a picenylylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pyrrolylylene group, an imidazolylene group, a pyrazolylylene group, a pyridinylylene group, a pyrazinylylene group, a pyrimidinylene group, a pyridazinylylene group, an isoindolylylene group, an indolylylene group, an indazolylene group, a purinylylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinolinylene group, a carbazolylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthrolinylylene group, a phenazinylylene group, a benzoxazolylene group, a benzoimidazolylene group, a furanylylene group, a benzofuranylylene group, a thiophenylylene group, a benzothiophenylylene group, a thiazolylylene group, an isothiazolylylene group, a benzothiazolylylene group, an isoxazolylene group, an oxazolylene group, a triazolylene group, a tetrazolylylene group, an oxadiazolylylene group, a triazinylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, an imidazopyrimidinylene group, or an imidazopyridinylylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₆-C₂₀ aryl group, a C₂-C₂₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed hetero-polycyclic group, and —Si(Q₃₃)(Q₃₄)(Q₃₅),

wherein Q₃₃ to Q₃₅ are each independently selected from a hydrogen, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a


triazinyl group, a quinolinyl group, an isoquinolinyl group, a phthalazinyl group, a quinoxalinylylene group, a cinolinyl group, and a quinazolinylylene group.

8. The condensed cyclic compound of claim 1, wherein


L₁ to L₃, L₁₁ to L₁₃, L₂₁ to L₂₄, and L₃₁ to L₃₃ are each independently represented by one of Formulae 2-1 to 2-37:




Formula 2-1




Formula 2-2




Formula 2-3




Formula 2-4




Formula 2-5



Formula 2-6














Formula 2-7




Formula 2-8


-continued




Formula 2-9




Formula 2-10




Formula 2-11




Formula 2-12




Formula 2-13




Formula 2-14




Formula 2-15




Formula 2-16



Formula 2-17



Formula 2-18



Formula 2-19

-continued

Formula 2-20

Formula 2-21

Formula 2-22

Formula 2-23

Formula 2-24

Formula 2-25


Formula 2-26

Formula 2-27


Formula 2-28

Formula 2-29


-continued




Formula 2-30




Formula 2-31




Formula 2-32




Formula 2-33




Formula 2-34



Formula 2-35



Formula 2-36



Formula 2-37

wherein in Formulae 2-1 to 2-37,

Y₁ may be O, S, S(=O), S(=O)₂, C(Z₃)(Z₄), N(Z₅), or Si(Z₆)(Z₇);

Z₁ to Z₇ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a quinoxaliny group, a biphenyl group, and —Si(Q₃₃)(Q₃₄)(Q₃₅), wherein

Q₃₃ to Q₃₅ are each independently selected from a hydrogen, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a phthalazinyl group, a quinoxaliny group, a cinolinyl group, and a quinazoliny group;

d₁ is an integer of 1 to 4;d₂ is an integer of 1 to 3;d₃ is an integer of 1 to 6;d₄ is an integer of 1 to 8;d₅ is an integer of 1 or 2;d₆ is an integer of 1 to 5; and


* indicates a binding site to a neighboring atom.

9. The condensed cyclic compound of claim 1, whereinL₁ to L₃, L₁₁ to L₁₃, L₂₁ to L₂₄, and L₃₁ to L₃₃ are each independently selected from

a phenylene group, a naphthylene group, a phenanthrenylene group, an anthracenylene group, a triphenylenylene group, and a carbazolylene group; and











a phenylene group, naphthylene group, a phenanthrenylene group, an anthracenylene group, a triphenylenylene group, and a carbazolylene group each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a phenanthrenyl group, and a biphenyl group.

10. The condensed cyclic compound of claim 1, wherein L₁ to L₃, L₁₁ to L₁₃, L₂₁ to L₂₄, and L₃₁ to L₃₃ are each independently represented by one of Formulae 3-1 to 3-33:




Formula 3-1

-continued




Formula 3-2




Formula 3-3

Formula 3-4




Formula 3-5

Formula 3-6




Formula 3-7


Formula 3-8




Formula 3-9



Formula 3-10



Formula 3-11



-continued

Formula 3-12

Formula 3-13

Formula 3-14








Formula 3-15

Formula 3-16

Formula 3-17


Formula 3-18

-continued




-continued


Formula 3-19




Formula 3-20




Formula 3-21




Formula 3-22




Formula 3-23



Formula 3-24



Formula 3-25



Formula 3-26

Formula 3-27

Formula 3-28

Formula 3-29


Formula 3-30

Formula 3-31

Formula 3-32

-continued

Formula 3-33



wherein * and *¹ in Formulae 3-1 to 3-33 each independently indicates a binding site to a neighboring atom.

11. The condensed cyclic compound of claim 1, wherein a₁ to a₃, a₁₁ to a₁₃, a₂₁ to a₂₄, and a₃₁ to a₃₃ are each independently 0, 1, or 2.

12. The condensed cyclic compound of claim 1, wherein R₁, R₂, R₁₁ to R₁₃, R₂₁ to R₂₃, and R₃₁ to R₃₃ are each independently selected from

a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazine group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, and a phosphoric acid or a salt thereof;

a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinalinyl group, a quinazolinalinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiofenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiofenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group,

an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinalinyl group, a quinazolinalinyl group, a cinnolinyl group, a carbazolyl group, phenanthridinyl, acridinyl, phenanthrolinyl, phenazinyl, a benzoimidazolyl group, a benzofuranyl group, a benzothiofenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiofenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazine group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a phenyl group substituted with a C₁-C₂₀ alkyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinalinyl group, a quinazolinalinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiofenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiofenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.

zopyridinyl group, an imidazopyrimidinyl group, a biphenyl group, and $-\text{Si}(\text{Q}_{33})(\text{Q}_{34})(\text{Q}_{35})$; and $-\text{Si}(\text{Q}_3)(\text{Q}_4)(\text{Q}_5)$,

wherein Q_3 to Q_5 and Q_{33} to Q_{35} are each independently selected from a hydrogen, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group.


13. The condensed cyclic compound of claim 1, wherein R_1 , R_2 , R_{11} to R_{13} , R_{21} to R_{23} , and R_{31} to R_{33} are each independently selected from

a hydrogen, a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;


a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, and a phosphoric acid or a salt thereof;

one of Formulae 4-1 to 4-51, 5-1 to 5-6, and 6-1 to 6-5; and $-\text{Si}(\text{Q}_3)(\text{Q}_4)(\text{Q}_5)$,


wherein Q_3 to Q_5 are each independently selected from a hydrogen, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group:




Formula 4-1




Formula 4-2



Formula 4-3




Formula 4-4




-continued


Formula 4-5




Formula 4-6




Formula 4-7




Formula 4-8




Formula 4-9




Formula 4-10




Formula 4-11




Formula 4-12




Formula 4-13



Formula 4-14














Formula 4-15




Formula 4-16

-continued




-continued



Formula 4-38

Formula 4-49


Formula 4-39



Formula 4-50

Formula 4-40


Formula 4-41



Formula 4-51


Formula 4-42

Formula 4-43



Formula 5-1


Formula 4-44



Formula 5-2


Formula 4-45

Formula 4-46




Formula 5-3

Formula 4-47




Formula 5-4

Formula 4-48




Formula 5-5


-continued




Formula 5-6




Formula 6-1




Formula 6-2



Formula 6-3



Formula 6-4



Formula 6-5

wherein in Formulae 4-1 to 4-51, 5-1 to 5-6, and 6-1 to 6-5, Y_{31} is O, S, C(Z_{33})(Z_{34}), N(Z_{35}), or Si(Z_{36})(Z_{37});

Z_{31} to Z_{37} are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group substituted with a C_1 - C_{20} alkyl group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a quinoxaliny group, a biphenyl group, and —Si(Q_{33})(Q_{34})(Q_{35}), wherein

Q_{33} to Q_{35} are each independently selected from a hydrogen, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl

group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, and a quinoxaliny group;

e_1 is an integer of 1 to 5;

e_2 is an integer of 1 to 7;

e_3 is an integer of 1 to 3;

e_4 is an integer of 1 to 4;

e_5 is an integer of 1 or 2;


e_6 is an integer of 1 to 6; and

* indicates a binding site to a neighboring atom.


14. The condensed cyclic compound of claim 1, wherein

Formula 2A is represented by one of Formulae 2A(1) to 2A(5), Formula 2B is represented by one of Formulae 2B(1) and 2B(2), Formula 2D is represented by one of Formulae 2D(1) to 2D(5), and Formula 2F is represented by one of Formulae 2F(1) and 2F(2):


Formula 2A(1)




Formula 2A(2)




Formula 2A(3)




Formula 2A(4)




-continued




Formula 2B(1)




Formula 2B(2)




Formula 2E(1)




Formula 2E(2)




Formula 2E(3)




Formula 2E(4)




-continued



Formula 2F(1)




Formula 2F(2)




wherein L_{21} to L_{24} , a_{21} to a_{24} , R_{21} to R_{23} , b_{21} to b_{23} , L_{31} to L_{33} , a_{31} to a_{33} , R_{31} to R_{33} , b_{31} to b_{33} , c_{21} to c_{23} , c_{31} and c_{32} in Formulae 2A (1) to 2A (5), 2B(1), 2B(2), 2E(1) to 2E(5), 2F(1), and 2F(2) are the same as recited in claim 1.

15. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound is represented by one of Formulae 1A(1) to 1A(8):


Formula 1A(1)




-continued




Formula 1A(2)



Formula 1A(3)




Formula 1A(4)




Formula 1A(5)


-continued



Formula 1A(6)



Formula 1A(7)



Formula 1A(8)


wherein, in Formulae 1A(1) and 1A(8),

Ring A is a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, or a substituted or unsubstituted monovalent non-aromatic condensed hetero-polycyclic group, each comprising Moiety 1 as a ring-forming moiety;


Pc_1 , Pc_2 , L_3 , a_3 , R_1 , R_2 , p_1 , and p_2 are the same as recited in claim 1; and

R_3 and R_4 are the same as recited in claim 1.


16. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound is represented by one selected from Formulae 1B(1) to 1B(10):




Formula 1B(1)




Formula 1B(2)




Formula 1B(3)




Formula 1B(4)




Formula 1B(5)



Formula 1B(6)




Formula 1B(7)




Formula 1B(8)

-continued




Formula 1B(9)




Formula 1B(10)

wherein Ring A, Pc₂, Pc₂, L₃, a₃, R₁, R₂, p₁, and p₂ in Formulae 1B(1) to 1B(10) are the same as recited in claim 1.

17. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound is represented by Formulae 1C(1) or 1C(2):




Formula 1C(1)



Formula 1C(2)

wherein Cz₁, Cz₂, Cz₃, R₁₁, R₁₂, R₁₃, p₁₁, p₁₂, and p₁₃ in Formulae 1C(1) and 1C(2) are the same as recited in claim 1.

18. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound is represented by Formula 1C(3):



Formula 1C(3)


wherein in Formula 1C(3),

Cz₃, L₁₁, and a₁₁ are the same as recited in claim 1;


R₁₁ is a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, or a substituted or unsubstituted monovalent non-aromatic condensed hetero-polycyclic group, each comprising Moiety 1 as a ring-forming moiety; and

b₁₁ is 1 or 2.

19. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound represented by Formulae 1C(4) or 1C(5):







Formula 1C(4)



Formula 1C(5)

wherein in Formulae 1C(4) and 1C(5), Cz₁ and Cz₃ are each a group represented by Formula 2E or a group represented by Formula 2F.



20. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound is one of Compounds 1 to 76:



-continued



5

6




7

8




-continued


9




10




11




12



13




14




-continued


21




22



23





24



-continued



25

26



27



28



-continued



29

30



31



32



-continued



33

34




35

36




-continued

37




38




-continued

39





40



-continued



41

42




43

44




-continued


45




46



47





48



-continued



49

50



51



52



-continued



53

54





55

56





-continued
57

58





59

60



61



62



-continued



63

64




65

66




-continued


67




68




69




70





71





72



-continued
73

75

76



21. An organic light-emitting device comprising:
a first electrode;
a second electrode; and
an organic layer disposed between the first electrode and the second electrode,


wherein the organic layer comprises an emission layer and at least one condensed cyclic compound represented by Formula 1 of claim 1.

22. The organic light-emitting device of claim 21, wherein the emission layer comprises the condensed cyclic compound represented by Formula 1 of claim 1.


23. The organic light-emitting device of claim 21, wherein the emission layer comprises a first host and a second host, the condensed cyclic compound represented by one of Formulae 1A to 1C exists in the emission layer, the first host comprises the condensed cyclic compound represented by one of Formulae 1A to 1C, and

the second host comprises at least one selected from a first compound represented by Formula 41 and a second compound represented by Formula 61:


Formula 41



-continued



Formula 61



Formula 61A



Formula 61B

wherein in the formulae 41, 61, 61A, and 61B,
 X_{41} is $N-[(L_{42})_{a42}-(R_{42})_{b42}]$, S, O, $S(=O)$, $S(=O)_2$,
 $C(=O)$, $C(R_{43})(R_{44})$, $Si(R_{43})(R_{44})$, $P(R_{43})$, $P(=O)$
 (R_{43}) , or $C=N(R_{43})$;

Ring A_{61} in Formula 61 is represented by Formula 61A;

Ring A_{62} in Formula 61 is represented by Formula 61B;

X_{61} is $N-[(L_{62})_{a62}-(R_{62})_{b62}]$, S, O, $S(=O)$, $S(=O)_2$,
 $C(=O)$, $C(R_{63})(R_{64})$, $Si(R_{63})(R_{64})$, $P(R_{63})$, $P(=O)$
 (R_{63}) , or $C=N(R_{63})$;

X_{71} is $C(R_{71})$ or N;

X_{72} is $C(R_{72})$ or N;

X_{73} is $C(R_{73})$ or N;

X_{74} is $C(R_{74})$ or N;

X_{75} is $C(R_{75})$ or N;

X_{76} is $C(R_{76})$ or N;

X_{77} is $C(R_{77})$ or N;

X_{78} is $C(R_{78})$ or N;

Ar_{41} , L_{41} , L_{42} , L_{61} , and L_{62} are each independently
 selected from a substituted or unsubstituted C_3 - C_{10}
 cycloalkylene group, a substituted or unsubstituted
 C_2 - C_{10} heterocycloalkylene group, a substituted or
 unsubstituted C_3 - C_{10} cycloalkenylene group, a substi-
 tuted or unsubstituted C_2 - C_{10} heterocycloalkenylene
 group, a substituted or unsubstituted C_6 - C_{60} arylene
 group, a substituted or unsubstituted C_2 - C_{60} hetero-
 arylylene group, a substituted or unsubstituted divalent
 non-aromatic condensed polycyclic group, or a substi-
 tuted and unsubstituted divalent non-aromatic con-
 densed hetero-polycyclic group;

$n1$ and $n2$ are each independently an integer selected from
 0 to 3;

$a41$, $a42$, $a61$, and $a62$ are each independently an integer
 selected from 0 to 3;

R_{41} to R_{43} , R_{51} to R_{54} , R_{61} to R_{64} , and R_{71} to R_{79} are each
 independently selected from a hydrogen, a deuterium,
 $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a
 nitro group, an amino group, an amidino group, a hydra-
 zine group, a hydrazone group, a carboxylic acid or a salt
 thereof, a sulfonic acid or a salt thereof, a phosphoric
 acid or a salt thereof, a substituted or unsubstituted
 C_1 - C_{60} alkyl group, a substituted or unsubstituted
 C_2 - C_{60} alkenyl group, a substituted or unsubstituted
 C_2 - C_{60} alkynyl group, a substituted or unsubstituted
 C_1 - C_{60} alkoxy group, a substituted or unsubstituted

C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted
 C_2 - C_{10} heterocycloalkyl group, a substituted or unsub-
 stituted C_3 - C_{10} cycloalkenyl group, a substituted or unsub-
 stituted C_2 - C_{10} heterocycloalkenyl group, a substi-
 tuted or unsubstituted C_6 - C_{60} aryl group, a substituted
 or unsubstituted C_6 - C_{60} aryloxy group, a substituted or
 unsubstituted C_6 - C_{60} arylthio group, a substituted or
 unsubstituted C_2 - C_{60} heteroaryl group, a substituted or
 unsubstituted monovalent non-aromatic condensed
 polycyclic group, a substituted or unsubstituted
 monovalent non-aromatic condensed hetero-polycyclic
 group, $-N(Q_1)(Q_2)$, $-Si(Q_3)(Q_4)(Q_5)$, and $-B(Q_6)$
 (Q_7) ;

$b41$, $b42$, $b51$ to $b54$, $b61$, $b62$, and $b79$ are each indepen-
 dently an integer selected from 1 to 3;

at least one of substituents of the substituted C_1 - C_{60} alky-
 lene group, substituted C_2 - C_{60} alkenylene group, substi-
 tuted C_3 - C_{10} cycloalkylene group, substituted C_2 - C_{10}
 heterocycloalkylene group, substituted C_3 - C_{10}
 cycloalkenylene group, substituted C_2 - C_{10} heterocy-
 cloalkenylene group, substituted C_6 - C_{60} arylene group,
 substituted C_2 - C_{60} heteroarylene group, substituted
 divalent non-aromatic condensed polycyclic group, substi-
 tuted divalent non-aromatic condensed hetero-poly-
 cyclic group, substituted C_1 - C_{60} alkyl group, substituted
 C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group,
 substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10}
 cycloalkyl group, substituted C_2 - C_{10} heterocycloalkyl
 group, substituted C_3 - C_{10} cycloalkenyl group, substi-
 tuted C_2 - C_{10} heterocycloalkenyl group, substituted
 C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group,
 substituted C_6 - C_{60} arylthio group, substituted C_2 - C_{60}
 heteroaryl group, a substituted monovalent non-aro-
 matic condensed polycyclic group, and a substituted
 monovalent non-aromatic condensed hetero-polycyclic
 group is selected from

a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a
 cyano group, a nitro group, an amino group, an amidino
 group, a hydrazine group, a hydrazone group, a carboxy-
 lic acid group or a salt thereof, a sulfonic acid group or a
 salt thereof, a phosphoric acid group or a salt thereof, a
 C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60}
 alkynyl group, and a C_1 - C_{60} alkoxy group;

a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60}
 alkynyl group, and a C_1 - C_{60} alkoxy group, each substi-
 tuted with at least one selected from a deuterium, $-F$,
 $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro
 group, an amino group, an amidino group, a hydrazine
 group, a hydrazone group, a carboxylic acid or a salt
 thereof, a sulfonic acid or a salt thereof, a phosphoric
 acid or a salt thereof, a C_3 - C_{10} cycloalkyl group, a
 C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl
 group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl
 group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group,
 a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic
 condensed polycyclic group, a monovalent non-aro-
 matic condensed hetero-polycyclic group, $-N(Q_{11})$
 (Q_{12}) , $-Si(Q_{13})(Q_{14})(Q_{15})$, and $-B(Q_{16})(Q_{17})$;

a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl
 group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocy-
 cloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy
 group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl

group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed hetero-polycyclic group;

- a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group,

a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇); and —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇);

wherein Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ are each independently selected from a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group.

24. The organic light-emitting device of claim **21**, wherein the emission layer further comprises a phosphorescent dopant.

25. The organic light-emitting device of claim **21**, wherein the electron transport region comprises at least one of the condensed cyclic compounds represented by Formula 1.

* * * * *

专利名称(译)	缩合环状化合物和包括其的有机发光器件		
公开(公告)号	US20150333273A1	公开(公告)日	2015-11-19
申请号	US14/599724	申请日	2015-01-19
[标]申请(专利权)人(译)	三星电子株式会社		
申请(专利权)人(译)	SAMSUNG ELECTRONICS CO. , LTD. 三星SDI CO. , LTD		
当前申请(专利权)人(译)	SAMSUNG ELECTRONICS CO. , LTD. 三星SDI CO. , LTD.		
[标]发明人	LEE SUNYOUNG KIM SOYEON KWAK YOONHYUN KIM SANGDONG LEE SANGYEOB		
发明人	LEE, SUNYOUNG KIM, SOYEON KWAK, YOONHYUN KIM, SANGDONG LEE, SANGYEOB		
IPC分类号	H01L51/00 C07D519/00 C07D487/06		
CPC分类号	H01L51/0072 C07D487/06 C07D519/00 H01L51/5072 H01L51/5012 H01L51/5024 H01L51/0067 H01L51/0085 H01L51/0087 H01L51/5016 H01L2251/5384		
优先权	1020140058395 2014-05-15 KR		
其他公开文献	US9771373		
外部链接	Espacenet USPTO		

摘要(译)
由式1A至1C之一表示的稠合环状化合物，其中式1A至1C在详述中公开。

