




US 20160141526A1

(19) **United States**(12) **Patent Application Publication**
LEE et al.(10) **Pub. No.: US 2016/0141526 A1**(43) **Pub. Date: May 19, 2016**(54) **ORGANOMETALLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**(52) **U.S. CL.**
CPC **H01L 51/0094** (2013.01); **C07F 15/0033**
(2013.01); **C09K 11/06** (2013.01); **C09K**
2211/1055 (2013.01); **H01L 51/5206** (2013.01)(71) Applicant: **Samsung Electronics Co., Ltd.**,
Suwon-si (KR)(57) **ABSTRACT**
An organometallic compound represented by Formula 1:(72) Inventors: **BangLin LEE**, Suwon-si (KR);
Hyeonho CHOI, Seoul (KR); **Youngjae**
PARK, Seoul (KR); **Kyuhyun IM**,
Yongin-si (KR)

Formula 1

(21) Appl. No.: **14/676,242**(22) Filed: **Apr. 1, 2015**(30) **Foreign Application Priority Data**

Nov. 14, 2014 (KR) 10-2014-0158906

Publication Classification(51) **Int. Cl.**
H01L 51/00 (2006.01)
C09K 11/06 (2006.01)
C07F 15/00 (2006.01)

wherein in Formula 1, groups and variables are the same as described in the specification.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6





FIG. 7



ORGANOMETALLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application claims priority to Korean Patent Application No. 10-2014-0158906, filed on Nov. 14, 2014, in the Korean Intellectual Property Office, and all the benefits accruing therefrom under 35 U.S.C. §119, the content of which is incorporated herein in its entirety by reference.

BACKGROUND

[0002] 1. Field

[0003] One or more embodiments relate to an organometallic compound and an organic light-emitting device including the same.

[0004] 2. Description of the Related Art

[0005] Organic light emitting devices (OLEDs) are self-emission devices that have wide viewing angles, high contrast ratios, and short response times. In addition, OLEDs exhibit excellent luminance, driving voltage, and response speed characteristics, and produce full-color images.

[0006] In an example, an organic light-emitting device includes an anode, a cathode, and an organic layer that is disposed between the anode and the cathode and includes an emission layer. A hole transport region may be disposed between the anode and the emission layer, and an electron transport region may be disposed between the emission layer and the cathode. Holes provided from the anode may move toward the emission layer through the hole transport region, and electrons provided from the cathode may move toward the emission layer through the electron transport region. The holes and the electrons are recombined in the emission layer to produce excitons. These excitons change from an excited state to a ground state, thereby generating light.

[0007] Various types of organic light emitting devices are known. However, there still remains a need in OLEDs having low driving voltage, high efficiency, high brightness, and long lifespan.


SUMMARY

[0008] One or more embodiments relate to a novel organometallic compound and an organic light-emitting device including the same.

[0009] Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented exemplary embodiments.

[0010] An aspect of exemplary embodiments provides an organometallic compound represented by Formula 1:

Formula 1



[0011] wherein in Formula 1,

[0012] CY_1 may be selected from a C_5 - C_{60} carbocyclic group and a C_1 - C_{60} heterocyclic group,

[0013] Q_1 to Q_3 may be each independently selected from

[0014] a C_1 - C_{20} alkyl group and a phenyl group; and

[0015] a C_1 - C_{20} alkyl group and a phenyl group, each substituted with at least one selected from a deuterium, a C_1 - C_{10} alkyl group, and a phenyl group,

[0016] R_1 to R_8 may be each independently selected from a hydrogen, a deuterium, $-CH_3$, $-CDH_2$, $-CD_2H$, and $-CD_3$,

[0017] R_{11} to R_{14} may be each independently selected from

[0018] a hydrogen, a deuterium, a C_1 - C_{10} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a phenyl group, and a naphthyl group; and

[0019] a C_1 - C_{10} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from a deuterium, a C_1 - C_{10} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a phenyl group, and a naphthyl group,

[0020] R_{21} may be selected from

[0021] a hydrogen, a deuterium, a hydroxyl group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0022] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group,

a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0023] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0024] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

[0025] —N(Q₁₁)(Q₁₂) and —P(=O)(Q₁₃)(Q₁₄),

[0026] a₁ and b₁ may be each independently an integer selected from 0 to 4, provided that when a₁ is 2 or more, two or more groups R₁₁ may be identical or different, and when b₁ is 2 or more, 2 or more groups R₂₁ may be identical or different, and

[0027] Q₁₁ to Q₁₄ may be each independently selected from a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group and a C₆-C₁₄ aryl group; a C₆-C₁₄ aryl group substituted with at least one selected from a C₁-C₂₀ alkyl group and a C₆-C₁₄ aryl group; and a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0028] Another aspect provides an organic light-emitting device including:

[0029] a first electrode;

[0030] a second electrode; and

[0031] an organic layer disposed between the first electrode and the second electrode, wherein the organic layer includes an emission layer and includes at least one organometallic compound represented by Formula 1.

[0032] The emission layer may include the organometallic compound represented by Formula 1, the organometallic compound represented by Formula 1 included in the emission layer may act as a dopant, and the emission layer may further include a host, wherein an amount of the organometallic compound represented by Formula 1 in the emission layer is smaller than an amount of the host.

BRIEF DESCRIPTION OF THE DRAWINGS

[0033] These and/or other aspects will become apparent and more readily appreciated from the following description

of the exemplary embodiments, taken in conjunction with the accompanying drawings in which:

[0034] FIG. 1 is a schematic view of an organic light-emitting device according to an embodiment;

[0035] FIG. 2 is a graph of weight (percent, %) versus temperature (degree Centigrade, ° C.) showing thermogravimetric analysis (TGA) data of Compound 3;

[0036] FIG. 3 is a graph of intensity (arbitrary unit, a. u.) versus wavelength (nanometer, nm) showing an ultraviolet (UV)-absorption spectrum and a photoluminescent (PL) of Compound 3 in solution;

[0037] FIG. 4 is a graph of intensity (arbitrary unit, a. u.) versus time (minute, min) showing high performance liquid chromatography (HPLC) data of Compound 3 to explain a purity change after a heat treatment of Compound 3;

[0038] FIG. 5 is a graph of intensity (arbitrary unit, a. u.) versus time (minute, min) showing HPLC data of Compound A to explain a purity change after a heat treatment of Compound A;

[0039] FIG. 6 is a graph of electroluminescent (EL) intensity (arbitrary unit, a. u.) versus wavelength (nanometer, nm) showing an electroluminescent (EL) spectrum of organic light-emitting devices manufactured according to Examples 1, 3, 4, and 5; and

[0040] FIG. 7 is a graph of luminance (percent, %) versus time (hour, hr) showing lifetime data (at 10 milliAmperes per square centimeter (mA/cm²)) of the organic light-emitting device manufactured according to Example 3.

DETAILED DESCRIPTION

[0041] Reference will now be made in detail to exemplary embodiments, examples of which are illustrated in the accompanying drawings, wherein like reference numerals refer to like elements throughout. In this regard, the present exemplary embodiments may have different forms and should not be construed as being limited to the descriptions set forth herein. Accordingly, the exemplary embodiments are merely described below, by referring to the figures, to explain aspects. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

[0042] It will be understood that when an element is referred to as being “on” another element, it can be directly in contact with the other element or intervening elements may be present therebetween. In contrast, when an element is referred to as being “directly on” another element, there are no intervening elements present.

[0043] It will be understood that when an element is referred to as being “on” another element, it can be directly in contact with the other element or intervening elements may be present therebetween. In contrast, when an element is referred to as being “directly on” another element, there are no intervening elements present.

[0044] It will be understood that, although the terms first, second, third etc. may be used herein to describe various elements, components, regions, layers, and/or sections, these elements, components, regions, layers, and/or sections should not be limited by these terms. These terms are only used to distinguish one element, component, region, layer, or section from another element, component, region, layer, or section. Thus, a first element, component, region, layer, or section discussed below could be termed a second element,

component, region, layer, or section without departing from the teachings of the present embodiments.

[0045] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. As used herein, the singular forms “a,” “an,” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.

[0046] It will be further understood that the terms “comprises” and/or “comprising,” or “includes” and/or “including” when used in this specification, specify the presence of stated features, regions, integers, steps, operations, elements, and/or components, but do not preclude the presence or addition of one or more other features, regions, integers, steps, operations, elements, components, and/or groups thereof.

[0047] Spatially relative terms, such as “beneath,” “below,” “lower,” “above,” “upper” and the like, may be used herein for ease of description to describe one element or feature’s relationship to another element(s) or feature(s) as illustrated in the figures. It will be understood that the spatially relative terms are intended to encompass different orientations of the device in use or operation in addition to the orientation depicted in the figures. For example, if the device in the figures is turned over, elements described as “below” or “beneath” other elements or features would then be oriented “above” the other elements or features. Thus, the exemplary term “below” can encompass both an orientation of above and below. The device may be otherwise oriented (rotated 90 degrees or at other orientations) and the spatially relative descriptors used herein interpreted accordingly.


[0048] “About” or “approximately” as used herein is inclusive of the stated value and means within an acceptable range of deviation for the particular value as determined by one of ordinary skill in the art, considering the measurement in question and the error associated with measurement of the particular quantity (i.e., the limitations of the measurement system). For example, “about” can mean within one or more standard deviations, or within $\pm 30\%$, 20% , 10% , 5% of the stated value.

[0049] Unless otherwise defined, all terms (including technical and scientific terms) used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. It will be further understood that terms, such as those defined in commonly used dictionaries, should be interpreted as having a meaning that is consistent with their meaning in the context of the relevant art and the present disclosure, and will not be interpreted in an idealized or overly formal sense unless expressly so defined herein.

[0050] Exemplary embodiments are described herein with reference to cross section illustrations that are schematic illustrations of idealized embodiments. As such, variations from the shapes of the illustrations as a result, for example, of manufacturing techniques and/or tolerances, are to be expected. Thus, embodiments described herein should not be construed as limited to the particular shapes of regions as illustrated herein but are to include deviations in shapes that result, for example, from manufacturing. For example, a region illustrated or described as flat may, typically, have rough and/or nonlinear features. Moreover, sharp angles that are illustrated may be rounded. Thus, the regions illustrated in the figures are schematic in nature and their shapes are not intended to illustrate the precise shape of a region and are not intended to limit the scope of the present claims.

[0051] An organometallic compound according to an embodiment is represented by Formula 1 below:

Formula 1



[0052] CY_1 in Formula 1 may be selected from a C_5 - C_{60} carbocyclic group and a C_1 - C_{60} heterocyclic group.

[0053] For example, CY_1 in Formula 1 may be selected from a benzene, a dibenzofuran, a dibenzothiophene, a fluorene, and a carbazole, but embodiments are not limited thereto.

[0054] Q_1 to Q_3 in Formula 1 may be each independently selected from

[0055] a C_1 - C_{20} alkyl group and a phenyl group; and

[0056] a C_1 - C_{20} alkyl group and a phenyl group, each substituted with at least one selected from a deuterium, a C_1 - C_{10} alkyl group, and a phenyl group.

[0057] For example, Q_1 to Q_3 in Formula 1 may be each independently selected from

[0058] a C_1 - C_{10} alkyl group and a phenyl group; and

[0059] a C_1 - C_{10} alkyl group and a phenyl group, each substituted with at least one selected from a deuterium and a C_1 - C_{10} alkyl group.

[0060] In some embodiments, Q_1 to Q_3 in Formula 1 may be each independently selected from

[0061] $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CH_2CH_3$, $-CH_2CD_3$, $-CH_2CD_2H$, $-CH_2CDH_2$, $-CHDCH_3$, $-CHDCD_2H$, $-CHDCDH_2$, $-CHDCD_3$, $-CD_2CD_3$, $-CD_2CD_2H$, and $-CD_2CDH_2$;

[0062] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group; and

[0063] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group, each substituted with at least one selected from a deuterium and a C_1 - C_{10} alkyl group, but embodiments are not limited thereto.

[0064] In some embodiments, Q_1 to Q_3 in Formula 1 may all be identical;

[0065] Q_1 and Q_3 may be identical and Q_2 and Q_1 may be different from each other; or

[0066] Q_1 to Q_3 may all be different from each other.

[0067] In some embodiments, Q_1 to Q_3 in Formula 1 may all be identical, and may be selected from

[0068] $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCD}_3$, $-\text{CHDCD}_2\text{H}$, $-\text{CHDCDH}_2$, $-\text{CHDCD}_3$, $-\text{CD}_2\text{CD}_3$, $-\text{CD}_2\text{CD}_2\text{H}$, and $-\text{CD}_2\text{CDH}_2$;

[0069] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group; and

[0070] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group, each substituted with at least one selected from a deuterium and a C_1 - C_{10} alkyl group, but they are not limited thereto.

[0071] In some embodiments, Q_1 and Q_3 in Formula 1 may be each independently selected from $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCD}_3$, $-\text{CHDCD}_2\text{H}$, $-\text{CHDCDH}_2$, $-\text{CHDCD}_3$, $-\text{CD}_2\text{CD}_3$, $-\text{CD}_2\text{CD}_2\text{H}$, and $-\text{CD}_2\text{CDH}_2$;

[0072] Q_2 may be selected from

[0073] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group; and

[0074] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group, each substituted with at least one selected from a deuterium and a C_1 - C_{10} alkyl group, but embodiments are not limited thereto.

[0075] R_1 to R_8 in Formula 1 may be each independently selected from a hydrogen, a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$.

[0076] For example, R_1 to R_8 in Formula 1 may be each independently selected from a hydrogen and a deuterium, but they are not limited thereto.

[0077] In some embodiments, in Formula 1,

[0078] R_1 to R_8 may all be a hydrogen;

[0079] R_1 , R_2 , and R_4 to R_3 may be a hydrogen and R_3 may be selected from a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$;

[0080] R_1 , R_2 , R_4 , R_5 , R_6 and R_8 may be a hydrogen, and R_3 and R_7 may be each independently selected from a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$;

[0081] R_1 to R_4 may be a hydrogen, and R_5 to R_8 may be each independently selected from a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$;

[0082] R_1 to R_5 may be a hydrogen, and R_6 to R_8 may be each independently selected from a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$;

[0083] R_1 to R_4 and R_8 may be a hydrogen, and R_5 to R_7 may be each independently selected from a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$; or

[0084] R_1 to R_8 may be each independently selected from a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$, but are not limited thereto.

[0085] R_{11} to R_{14} in Formula 1 may be each independently selected from

[0086] a hydrogen, a deuterium, a C_1 - C_{10} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a phenyl group, and a naphthyl group; and

[0087] a C_1 - C_{10} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from a deuterium, a C_1 - C_{10} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a phenyl group, and a naphthyl group.

[0088] For example, R_{11} to R_{14} in Formula 1 may be each independently selected from


[0089] a hydrogen, a deuterium, a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an isohexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an isoheptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an isooctyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an isodecyl group, a sec-decyl group, a tert-decyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, and a naphthyl group; and

[0090] a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an isohexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an isoheptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an isooctyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an isodecyl group, a sec-decyl group, a tert-decyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from a deuterium, a C_1 - C_{10} alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, and a naphthyl group.

[0091] In some embodiments, R_{11} to R_{14} in Formula 1 may be each independently selected from a hydrogen, a deuterium, $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, a group represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-12, but embodiments are not limited thereto:



Formula 9-1




Formula 9-2



Formula 9-3

-continued



Formula 9-4

Formula 9-5

Formula 9-6

Formula 9-7

Formula 9-8

Formula 9-9

Formula 9-10

Formula 9-11

Formula 9-12

Formula 9-13

Formula 9-14


Formula 9-15

Formula 9-16

Formula 9-17

Formula 10-1

-continued



Formula 10-2

Formula 10-3

Formula 10-4

Formula 10-5

Formula 10-6

Formula 10-7

Formula 10-8

Formula 10-9

Formula 10-10

Formula 10-11

Formula 10-12

[0092] a_1 in Formula 1 may be an integer selected from 0 to 4. When a_1 is 2 or more, two or more groups R_{11} may be identical or different. For example, a_1 in Formula 1 may be 0, 1, or 2.

[0093] R_1 in Formula 21 may be selected from

[0094] a hydrogen, a deuterium, a hydroxyl group, a nitro group, an amino group, an amidino group, a hydrazine group,

a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0095] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0096] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0097] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

[0098] —N(Q₁₁)(Q₁₂) and —P(=O)(Q₁₃)(Q₁₄),

[0099] wherein Q₁₁ to Q₁₄ are each independently selected from a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, and a C₆-C₁₄ aryl group; a C₆-C₁₄ aryl group substituted with at least one selected from a C₁-C₂₀ alkyl group and a C₆-C₁₄ aryl group; and a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0100] In some embodiments, R₂₁ in Formula 1 may be selected from

[0101] a hydrogen, a deuterium, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0102] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro

group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

[0103] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzo-carbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0104] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzo-carbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl

group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

[0105] $-\text{B}(\text{Q}_{11})(\text{Q}_{12})$ and $-\text{P}(=\text{O})(\text{Q}_{13})(\text{Q}_{14})$,

[0106] Q_{11} to Q_{14} may be each independently selected from

[0107] a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, and a carbazolyl group; and

[0108] a phenyl group and a naphthyl group, each substituted with at least one selected from a C_1 - C_{20} alkyl group, a phenyl group, and a naphthyl group.

[0109] In some embodiments, R_{21} in Formula 1 may be selected from

[0110] a hydrogen, a deuterium, a cyano group, a nitro group, a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an isohexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an isoheptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an isooctyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an isodecyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0111] a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an isohexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an isoheptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an isooctyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an isodecyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with

at least one selected from a deuterium, $-\text{F}$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

[0112] $-\text{B}(\text{Q}_{11})(\text{Q}_{12})$ and $-\text{P}(=\text{O})(\text{Q}_{13})(\text{Q}_{14})$,

[0113] Q_{11} to Q_{14} may be each independently selected from


[0114] a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, and a naphthyl group; and

[0115] a phenyl group and a naphthyl group, each substituted with at least one selected from a C_1 - C_{20} alkyl group, a phenyl group, and a naphthyl group, but they are not limited thereto.

[0116] In some embodiments, R_{21} in Formula 1 may be selected from a hydrogen, a deuterium, a cyano group, a nitro group, $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, groups represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-30, but embodiments are not limited thereto:



Formula 9-1



Formula 9-2



Formula 9-3




Formula 9-4



Formula 9-5




Formula 9-6



Formula 9-7




Formula 9-8




Formula 9-9




Formula 9-10
















Formula 9-11



Formula 9-12



-continued



Formula 9-13

Formula 9-14

Formula 9-15

Formula 9-16

Formula 9-17

Formula 10-1

Formula 10-2

Formula 10-3

Formula 10-4












Formula 10-5

Formula 10-6

Formula 10-7

Formula 10-8

-continued



Formula 10-9

Formula 10-10

Formula 10-11

Formula 10-12

Formula 10-13

Formula 10-14

Formula 10-15











Formula 10-16

Formula 10-17


Formula 10-18

Formula 10-19

-continued



-continued



Formula 10-20

Formula 10-30

Formula 10-21

Formula 10-22

Formula 10-23

Formula 10-24

Formula 10-25

Formula 10-26

Formula 10-27


Formula 10-28

Formula 10-29


[0117] b_1 in Formula 1 may be an integer selected from 0 to 4. When b_1 is 2 or more, two or more groups R_{21} may be identical or different.

[0118] In some embodiments, the organometallic compound represented by Formula 1 may be represented by one of Formulae 1-1 to 1-12:

Formula 1-1




Formula 1-2




-continued


Formula 1-3



Formula 1-4




Formula 1-5




-continued


Formula 1-6



Formula 1-7




Formula 1-8




-continued

Formula 1-9




-continued


Formula 1-12



Formula 1-10



Formula 1-11



[0119] In connection with Formulae 1-1 to 1-12, descriptions of CY_1 , Q_1 to Q_3 , R_1 to R_8 , R_{11} to R_{14} , R_{21} , and $b1$ are the same as described above, and descriptions of R_{11a} and R_{11b} are the same as those of R_{11} . In this embodiment, however, R_{11} , R_{11a} , and R_{11b} in Formulae 1-1 to 1-12 are not a hydrogen.

[0120] For example, in Formulae 1-1 and 1-12,

[0121] Q_1 to Q_3 may be each independently selected from

[0122] $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CH_2CH_3$, $-CH_2CD_3$, $-CH_2CD_2H$, $-CH_2CDH_2$, $-CHDCH_3$, $-CHDCD_2H$, $-CHDCDH_2$, $-CHDCD_3$, $-CD_2CD_3$, $-CD_2CD_2H$, and $-CD_2CDH_2$;

[0123] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group; and

[0124] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group, each substituted with at least one selected from a deuterium and a C_1 - C_{10} alkyl group,

[0125] R_1 to R_8 may all be a hydrogen;

[0126] R_1 , R_2 , and R_4 to R_8 may be a hydrogen, and R_3 may be selected from a deuterium, $-CH_3$, $-CDH_2$, $-CD_2H$, and $-CD_3$;

[0127] R_1 , R_2 , R_4 , R_5 , R_6 and R_8 may be a hydrogen, and R_3 and R_7 may be each independently selected from a deuterium, $-CH_3$, $-CDH_2$, $-CD_2H$, and $-CD_3$;

[0128] R_1 to R_4 may be a hydrogen, and R_5 to R_8 may be each independently selected from a deuterium, $-CH_3$, $-CDH_2$, $-CD_2H$, and $-CD_3$;

[0129] R_1 to R_5 may be a hydrogen, and R_6 to R_8 may be each independently selected from a deuterium, $-CH_3$, $-CDH_2$, $-CD_2H$, and $-CD_3$;

[0130] R_1 to R_4 and R_8 may be a hydrogen, and R_5 to R_7 may be each independently selected from a deuterium, $-CH_3$, $-CDH_2$, $-CD_2H$, and $-CD_3$; or

[0131] R_1 to R_8 may be each independently selected from a hydrogen, a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$,


[0132] R_{11} , R_{11a} , R_{11b} , R_{12} , R_{13} and R_{14} may be each independently selected from a hydrogen, a deuterium, $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, groups represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-12 (provided that R_{11} , R_{11a} and R_{11b} are not a hydrogen), and

[0133] R_{21} may be selected from a hydrogen, a deuterium, a cyano group, a nitro group, $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, groups represented by Formulae 9-1 to 9-17, and groups represented by Formulae 10-1 to 10-30.


[0134] In some embodiments, the organometallic compound represented by Formula 1 may be represented by one of Formulae 1(1) to 1(12):

-continued


Formula 1(3)




Formula 1(1)




Formula 1(4)



Formula 1(2)




Formula 1(5)




-continued

Formula 1(6)




-continued


Formula 1(9)




Formula 1(7)




Formula 1(10)



Formula 1(8)




Formula 1(11)



-continued

Formula 1(12)



[0135] In connection with Formulae 1(1) to 1(12), descriptions of Q₁ to Q₃, R₁ to R₈, R₁₁, a₁, R₁₃ and R₂₁ are the same as described above, X₁ may be N(R₂₁), O, or S, and descriptions of R₂₂ to R₂₄ are the same as those of R₂₁, provided that R₁₃ and R₂₁ to R₂₄ in Formulae 1(1) to 1(12) are not be a hydrogen.

[0136] For example, in Formulae 1(1) and 1(12),

[0137] Q₁ to Q₃ may be each independently selected from

[0138] —CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;

[0139] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group; and

[0140] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group, each substituted with at least one selected from a deuterium and a C₁-C₁₀ alkyl group,

[0141] R₁ to R₈ may all be a hydrogen;

[0142] R₁, R₂, and R₄ to R₈ may be a hydrogen, and R₃ may be selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃;

[0143] R₁, R₂, R₄, R₅, R₆ and R₃ may be a hydrogen, and R₇ and R₈ may be each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃;

[0144] R₁ to R₄ may be a hydrogen, and R₅ to R₈ may be each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃;

[0145] R₁ to R₅ may be a hydrogen, and R₆ to R₈ may be each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃;

[0146] R₁ to R₄ and R₈ may be a hydrogen, and R₅ to R₇ may be each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃; or

[0147] R₁ to R₈ may be each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃,


[0148] R₁₁ and R₁₃ may be each independently selected from a hydrogen, a deuterium, —CH₃, —CD₃, —CD₂H, —CDH₂, a group represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-12 (provided that R₁₃ is not a hydrogen),

[0149] R₂₁ to R₂₄ may be each independently selected from a deuterium, a cyano group, a nitro group, —CH₃, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, groups represented by Formulae 9-1 to 9-17, and groups represented by Formulae 10-1 to 10-30, and


[0150] a₁ may be 0, 1, or 2.

[0151] In some embodiments, the organometallic compound represented by Formula 2 may be represented by one of Formulae 2-1 to 2-72:

Formula 2-1




Formula 2-2




-continued


Formula 2-3



Formula 2-4




Formula 2-5




-continued


Formula 2-6



Formula 2-7




Formula 2-8




-continued

Formula 2-15




-continued


Formula 2-18




Formula 2-16




Formula 2-19



Formula 2-17




Formula 2-20




-continued

Formula 2-21





-continued



Formula 2-24



Formula 2-25




Formula 2-26




-continued


Formula 2-27



Formula 2-28




Formula 2-29




-continued


Formula 2-30



Formula 2-31




Formula 2-32




-continued


Formula 2-33



Formula 2-34




Formula 2-35




-continued


Formula 2-36



Formula 2-37




Formula 2-38




-continued


Formula 2-39



Formula 2-40




Formula 2-41




-continued


Formula 2-42



Formula 2-43




Formula 2-44




-continued


Formula 2-48



Formula 2-49




Formula 2-50




-continued


Formula 2-51



Formula 2-52




Formula 2-53




-continued


Formula 2-54



Formula 2-55




Formula 2-56




-continued


Formula 2-57



Formula 2-58




Formula 2-59




-continued


Formula 2-60



Formula 2-61




Formula 2-62




-continued


Formula 2-63



Formula 2-64




Formula 2-65




-continued


Formula 2-66



Formula 2-67




Formula 2-68





-continued


Formula 2-72



Formula 2-70



Formula 2-71



[0152] In connection with Formulae 2-1 to 2-72, descriptions of Q₁ to Q₃, R₁ to R₈, R₁₁, R₁₃, and R₂₁ are the same as described above, X₁ may be N(R₂₁), O, or S, descriptions of R_{11a} and R_{11b} are the same as descriptions of R₁₁, and descriptions of R₂₂ to R₂₄ are the same as descriptions of R₂₁. In this embodiment, however, R₁₁, R_{11a}, R_{11b}, R₁₃ and R₂₁ to R₂₄ in Formulae 2-1 to 2-72 are not a hydrogen.

[0153] For example, in Formulae 2-1 and 2-72,

[0154] Q_1 to Q_3 may be each independently selected from

[0155] $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$,
 $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCH}_3$,
 $-\text{CHDCH}_2\text{H}$, $-\text{CHDCH}_2\text{H}$, $-\text{CHDCH}_3$, $-\text{CD}_2\text{CD}_3$,
 $-\text{CD}_2\text{CD}_2\text{H}$, and $-\text{CD}_2\text{CDH}_2$;

[0156] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group; and

[0157] an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group, each substituted with at least one selected from a deuterium and a C₁-C₁₀ alkyl group,

[0158] R_1 to R_8 may all be a hydrogen;

[0159] R₁, R₂, and R₄ to R₃ may be a hydrogen, and R₃ may be selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃;

[0160] R₁, R₂, R₄, R₅, R₆ and R₈ may be a hydrogen, and R₃ and R₇ may be each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃;

[0161] R₁ to R₄ may be a hydrogen, and R₅ to R₈ may be each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃;

[0162] R_1 to R_5 may be a hydrogen, and R_6 to R_8 may be each independently selected from a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$;

[0163] R_1 to R_4 and R_8 may be a hydrogen, and R_5 to R_7 may be each independently selected from a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$; or

[0164] R_1 to R_8 may be each independently selected from a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$,


[0165] R_{11} , R_{11a} , R_{11b} , and R_{13} may be each independently selected from a deuterium, $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, a group represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-12,

[0166] R_{21} to R_{24} may be each independently selected from a deuterium, a cyano group, a nitro group, $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, a group represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-30.


[0167] In some embodiments, the organometallic compound represented by Formula 1 may be represented by one of Formulae 1 to 54 below:

-continued


3




4




1




2



5




-continued




6

-continued




7


9




8




10



11




-continued




12

-continued




15

13




16


14





-continued





-continued




18



19




-continued




23


-continued




26




24



27




25




28

-continued




29

-continued




32


30




33




31



34




-continued




35

-continued




38


36




39




37








40




-continued




-continued




-continued




-continued




48




50







51




54



52



53



[0168] R_1 to R_8 of Ligand A in Formula 1 may be each independently selected from a hydrogen, a deuterium, $-\text{CH}_3$, $-\text{CDH}_2$, $-\text{CD}_2\text{H}$, and $-\text{CD}_3$. Ligand B in Formula 1 does not include other silyl groups than a "silyl group A." An organometallic compound represented by Formula 1 having two Ligands A and one Ligand B has excellent thermal stability and process stability. Accordingly, when the organometallic compound represented by Formula 1 is used, a high-quality organic light-emitting device may be manufactured in bulk quantities.

[0169] In an example, the highest occupied molecular orbital (HOMO), the lowest unoccupied molecular orbital (LUMO), and triplet (T_1) energy levels of Compounds 1 to 7 were evaluated by using a DFT method of Gaussian program (structurally optimized at a level of B3LYP, 6-31G(d,p)). The evaluation results are shown in Table 1 below.

TABLE 1

Compound No.	HOMO (eV)	LUMO (eV)	T ₁ energy level (eV)
1	-4.852	-1.404	2.486
2	-4.847	-1.353	2.506
3	-4.811	-1.278	2.571
4	-4.806	-1.191	2.628
5	-4.846	-1.309	2.552
6	-4.840	-1.267	2.571
7	-4.814	-1.171	2.630

[0170] From Table 1, it is confirmed that the compound represented by Formula 1 has electric characteristics that are suitable for use as a material for manufacturing an electric device, for example, an organic light-emitting device.

[0171] Synthesis methods of the organometallic compounds represented by Formula 1 may be understood by one of ordinary skill in the art by referring to Synthesis Examples provided below.

[0172] The organometallic compound represented by Formula 1 is suitable for use in an organic layer of an organic light-emitting device, for example, for use as a dopant in an emission layer of the organic layer. Thus, another aspect provides an organic light-emitting device that includes:

[0173] a first electrode;

[0174] a second electrode; and

[0175] an organic layer that is disposed between the first electrode and the second electrode, wherein the organic layer includes an emission layer and at least one organometallic compound represented by Formula 1.

[0176] The organic light-emitting device may have, due to the inclusion of an organic layer including the organometallic compound represented by Formula 1, a low driving voltage, high efficiency, high luminance, and a long lifespan.

[0177] The organometallic compound of Formula 1 may be used between a pair of electrodes of an organic light-emitting device. For example, the organometallic compound represented by Formula 1 may be included in the emission layer. In this regard, the organometallic compound may act as a dopant, and the emission layer may further include a host (that is, an amount of the organometallic compound represented by Formula 1 is smaller than an amount of the host).

[0178] The expression “(an organic layer) includes at least one organometallic compounds” as used herein may include an embodiment in which “(an organic layer) includes identical organometallic compounds of Formula 1 and an embodiment in which (an organic layer) includes two or more different organometallic compounds of Formula 1.

[0179] For example, the organic layer may include, as the organometallic compound, only Compound 1. In this regard, Compound 1 may be included in an emission layer of the organic light-emitting device. In some embodiments, the organic layer may include, as the organometallic compound, Compound 1 and Compound 2. In this regard, Compound 1 and Compound 2 may be included in an identical layer (for example, Compound 1 and Compound 2 all may exist in an emission layer).

[0180] The first electrode may be an anode, which is a hole injection electrode, and the second electrode may be a cathode, which is an electron injection electrode; or the first electrode may be a cathode, which is an electron injection electrode, or the second electrode may be an anode, which is a hole injection electrode.

[0181] For example, the first electrode may be an anode, and the second electrode may be a cathode, and the organic layer includes:

[0182] i) a hole transport region that is disposed between the first electrode and the emission layer, wherein the hole transport region includes at least one selected from a hole injection layer, a hole transport layer, and an electron blocking layer, and

[0183] ii) an electron transport region that is disposed between the emission layer and the second electrode, wherein the electron transport region includes at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

[0184] The term “organic layer” as used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of an organic light-emitting device. The “organic layer” may include, in addition to an organic compound, an organometallic complex including metal.

[0185] FIG. 1 is a schematic view of an organic light-emitting device 10 according to an embodiment. Hereinafter, the structure of an organic light-emitting device according to an embodiment and a method of manufacturing an organic light-emitting device according to an embodiment will be described in connection with FIG. 1. The organic light-emitting device 10 includes a first electrode 11, an organic layer 15, and a second electrode 19, which are sequentially stacked.

[0186] In FIG. 1, a substrate may be additionally disposed under the first electrode 11 or above the second electrode 19. As the substrate, any substrate that is used in general organic light-emitting devices may be used. The substrate may be a glass substrate or transparent plastic substrate, each with excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water-resistance.

[0187] The first electrode 11 may be formed by depositing or sputtering a material for forming the first electrode on the substrate. The first electrode 11 may be an anode. The material for the first electrode 11 may be selected from materials with a high work function to readily provide holes. The first electrode 11 may be a reflective electrode or a transmissive electrode. The material for the first electrode may be, for example, indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), and zinc oxide (ZnO). In some embodiments, magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag) may be used as the material for the first electrode.

[0188] The first electrode 11 may have a single-layer structure or a multi-layer structure including two or more layers. For example, the first electrode 11 may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode 110 is not limited thereto.

[0189] An organic layer 15 is disposed on the first electrode 11.

[0190] The organic layer 15 may include a hole transport region, an emission layer, and an electron transport region.

[0191] The hole transport region may be disposed between the first electrode 11 and the emission layer.

[0192] The hole transport region may include at least one selected from a hole injection layer, a hole transport layer, an electron blocking layer, and a buffer layer.

[0193] The hole transport region may include only either a hole injection layer or a hole transport layer. In some embodiments, the hole transport region may have a structure of hole injection layer/hole transport layer or hole injection layer/hole transport layer/electron blocking layer, which are sequentially stacked in this stated order from the first electrode 11.


[0194] A hole injection layer hole injection layer may be formed on the first electrode **11** by using various methods, such as vacuum deposition, spin coating, casting, or Langmuir-Blodgett (LB).

[0195] When a hole injection layer is formed by vacuum deposition, the deposition conditions may vary depending on a material that is used to form the hole injection layer, and the structure and thermal characteristics of the hole injection layer. For example, the deposition conditions may include a deposition temperature of about 100 to about 500° C., a vacuum pressure of about 10^{-8} to about 10^{-3} torr, and a deposition rate of about 0.01 to about 100 Angstroms per second (Å/sec). However, the deposition conditions are not limited thereto.




[0196] When the hole injection layer is formed using spin coating, coating conditions may vary according to the material used to form the hole injection layer, and the structure and thermal properties of the hole injection layer. For example, a coating speed may be from about 2,000 revolutions per minute (rpm) to about 5,000 rpm, and a temperature at which a heat treatment is performed to remove a solvent after coating may be from about 80° C. to about 200° C. However, the coating conditions are not limited thereto.

[0197] Conditions for a hole transport layer and an electron blocking layer may be understood by referring to conditions for forming the hole injection layer.


[0198] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB, β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated NPB, TAPC, HMTPD, 4,4',4"-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), (polyaniline)/poly(4-styrenesulfonate) (Pani/PSS), a compound represented by Formula 201 below, and a compound represented by Formula 202 below:




-continued




-continued

 β -NPB


-continued



TAPC




TDP




HMTPD


Formula 201




Spiro-TPD




Formula 202



Spiro-NPB



methylated NPB



[0199] Ar_{101} and Ar_{102} in Formula 201 may be each independently selected from

[0200] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluore-

nylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, and a pentacenylene group; and

[0201] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, and a pentacenylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0202] In Formula 201, xa and xb may be each independently an integer selected from 0 to 5, for example, an integer selected from 0, 1, and 2. For example, xa is 1 and xb is 0, but xa and xb are not limited thereto.

[0203] R_{101} to R_{108} , R_{111} to R_{119} and R_{121} to R_{124} in Formulae 201 and 202 may be each independently selected from

[0204] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{10} alkyl group (for example, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a hexyl group, and so on), and a C_1 - C_{10} alkoxy group (for example, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, and so on);

[0205] a C_1 - C_{10} alkyl group or a C_1 - C_{10} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, and a phosphoric acid group or a salt thereof;

[0206] a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, and a pyrenyl group; and


[0207] a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, and a pyrenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{10} alkyl group, and a C_1 - C_{10} alkoxy group, but they are not limited thereto.

[0208] R_{109} in Formula 201 may be selected from

[0209] a phenyl group, a naphthyl group, an anthracenyl group and a pyridinyl group; and

[0210] a phenyl group, a naphthyl group, an anthracenyl group and a pyridinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, and a pyridinyl group.

[0211] According to an embodiment, the compound represented by Formula 201 may be represented by Formula 201A, but is not limited thereto:




Formula 201A

[0212] R_{101} , R_{111} , R_{112} , and R_{109} in Formula 201A may be understood by referring to the description provided herein.

[0213] For example, the compound represented by Formula 201, and the compound represented by Formula 202 may include compounds HT1 to HT20 illustrated below, but are not limited thereto.


HT1




-continued

-continued


HT2




HT4



HT3




HT5




-continued

-continued


HT6




HT8



HT7



HT9





-continued


-continued

HT12

HT10





HT13




HT14

HT11




HT15




-continued

HT16




-continued


HT20




HT17



HT18



HT19





[0214] A thickness of the hole transport region may be in a range of about 100 Angstroms (Å) to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes a hole injection layer and a hole transport layer, the thickness of the hole injection layer may be in a range of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å, and the thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, and for example, about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0215] The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0216] The charge-generation material may be, for example, a p-dopant. The p-dopant may be one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but embodiments are not limited thereto. Non-limiting examples of the p-dopant are a quinone derivative, such as tetracyanoquinonodimethane (TCNQ) or 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonodimethane (F4-TCNQ); a metal oxide, such as a tungsten oxide or a molybdenum oxide; and a cyano group-containing compound, such as Compound HT-D1 below, but are not limited thereto.

Compound HT-D1





[0217] The hole transport region may include a buffer layer.







[0218] The buffer layer may compensate for an optical resonance distance depending on a wavelength of light emitted from the emission layer, thus improving the efficiency of a formed organic light-emitting device.


[0219] Then, an emission layer (EML) may be formed on the hole transport region by vacuum deposition, spin coating, casting, LB deposition, or the like. When the emission layer is formed by vacuum deposition or spin coating, the deposition or coating conditions may be similar to those applied to form the hole injection layer. However, the deposition or coating conditions may vary according to the material that is used to form the emission layer.

[0220] Meanwhile, when the hole transport region includes an electron blocking layer, a material for the electron blocking layer may be selected from materials for the hole transport region described above and materials for a host, that will be discussed later. However, the material for the electron blocking layer is not limited thereto. For example, when the hole transport region includes an electron blocking layer, a material for the electron blocking layer may be mCP, which will also be discussed later.


[0221] The emission layer may include a host and a dopant, and the dopant may include the organometallic compound represented by Formula 1.

[0222] The host may include at least one selected from TPBi, TBADN, ADN (also referred to as "DNA"), CBP, CDBP, TCP, Mcp, Compound HSO, and Compound H51:







Compound H50



Compound H51



[0223] In some embodiments, the host may further include a compound represented by Formula 301 below.



[0224] Ar_{111} and Ar_{112} in Formula 301 may be each independently selected from

a phenylene group, a naphthylene group, a phenanthrenylene group, and a pyrenylene group; and

a phenylene group, a naphthylene group, a phenanthrenylene group, and a pyrenylene group, each substituted with at least one selected from a phenyl group, a naphthyl group, and an anthracenyl group.

[0225] Ar_{113} to Ar_{116} in Formula 301 may be each independently selected from

[0226] a C_1 - C_{10} alkyl group, a phenyl group, a naphthyl group, a phenanthrenyl group, and a pyrenyl group; and

[0227] a phenyl group, a naphthyl group, a phenanthrenyl group, and a pyrenyl group, each substituted with at least one selected from a phenyl group, a naphthyl group, and an anthracenyl group.


[0228] g , h , i , and j in Formula 301 may be each independently an integer selected from 0 to 4, for example, an integer selected from 0, 1, and 2.

[0229] Ar_{113} to Ar_{116} in Formula 301 may be each independently selected from

[0230] a C_1 - C_{10} alkyl group substituted with at least one selected from a phenyl group, a naphthyl group, and an anthracenyl group;


[0231] a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl, a phenanthrenyl group, and a fluorenyl group;

[0232] a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group; and



but embodiments are not limited thereto.

[0233] In some embodiments, the host may include a compound represented by Formula 302 below:




[0234] Ar_{122} to Ar_{125} in Formula 302 are the same as described in detail in connection with Ar_{113} in Formula 301.


[0235] Ar_{126} and Ar_{127} in Formula 302 may be each independently a C_1 - C_{10} alkyl group (for example, a methyl group, an ethyl group, or a propyl group).

[0236] k and l in Formula 302 may be each independently an integer selected from 0 to 4. For example, k and l may be 0, 1, or 2.


[0237] The compound represented by Formula 301 and the compound represented by Formula 302 may include Compounds H1 to H42 illustrated below, but are not limited thereto.




H1




H7




H2




H8




H3




H9




H4




H10



H5




H11




H6


-continued




H12




H13



H14




H15




H16


-continued




H17



H18




H19




H20

-continued




H21


-continued




H26




H22




H27




H23




H29




H24




H30



H25




H31




H28

-continued


H32




H33



H34




H35




-continued

H36




H37




-continued

H38




-continued


H41




H39



H42



H40



[0238] When the organic light-emitting device is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, and a blue emission layer. In some embodiments, due to a stack structure including a red emission layer, a green emission layer, and/or a blue emission layer, the emission layer may emit white light.

[0239] When the emission layer includes a host and a dopant, an amount of the dopant may be in a range of about 0.01 to about 15 parts by weight based on 100 parts by weight of the host, but is not limited thereto.

[0240] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0241] Then, an electron transport region may be disposed on the emission layer.


[0242] The electron transport region may include at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

[0243] For example, the electron transport region may have a structure of hole blocking layer/electron transport layer/electron injection layer or a structure of electron transport


layer/electron injection layer, but the structure of the electron transport region is not limited thereto. The electron transport layer may have a single-layered structure or a multi-layer structure including two or more different materials.

[0244] Conditions for forming the hole blocking layer, the electron transport layer, and the electron injection layer, which constitute the electron transport region, may be understood by referring to the conditions for forming the hole injection layer.

[0245] When the electron transport layer includes a hole blocking layer, the hole blocking layer may include, for example, at least one of BCP, Bphen, and Balq but is not limited thereto.





BCP




Bphen

[0246] A thickness of the hole blocking layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thickness of the hole blocking layer is within these ranges, the hole blocking layer may have improved hole blocking ability without a substantial increase in driving voltage.


[0247] The electron transport layer may further include at least one selected from BCP, Bphen, Alq₃, Balq, TAZ, and NTAZ.

Alq₃

Balq




TAZ



NTAZ


[0248] In some embodiments, the electron transport layer may include at least one of ET1 and ET2, but are not limited thereto:



ET1

-continued


ET2




[0249] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

[0250] Also, the electron transport layer may further include, in addition to the materials described above, a metal-containing material.

[0251] The metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.



ET-D1



ET-D2

[0252] The electron transport region may include an electron injection layer (EIL) that allows electrons to be easily provided from a second electrode 19.

[0253] The electron injection layer may include at least one selected from, LiF, NaCl, CsF, Li₂O, BaO, and LiQ.

[0254] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

[0255] The second electrode 19 is disposed on the organic layer 15. The second electrode 19 may be a cathode. A material for forming the second electrode 19 may be selected from metal, an alloy, an electrically conductive compound, and a combination thereof, which have a relatively low work function. For example, lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag) may be formed as a material for forming the second electrode 19. In some embodiments, to manufacture a top emission type light-emitting device, a transmissive electrode formed using ITO or IZO may be used as the second electrode 19.

[0256] Hereinbefore, the organic light-emitting device has been described with reference to FIG. 1, but is not limited thereto.

[0257] A C₁-C₆₀ alkyl group as used herein refers to a linear or branched aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms. Detailed examples thereof are a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an isomethyl group, and a hexyl group. A C₁-C₆₀ alkylene group as used herein refers to a divalent group having the same structure as the C₁-C₆₀ alkyl group.

[0258] A C₁-C₆₀ alkoxy group as used herein refers to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is the C₁-C₆₀ alkyl group). Detailed examples thereof are a methoxy group, an ethoxy group, and an isopropoxy group.

[0259] A C₂-C₆₀ alkenyl group as used herein refers to a hydrocarbon group having at least one carbon double bond in the middle or at the terminal of the C₂-C₆₀ alkyl group. Detailed examples thereof are an ethenyl group, a propenyl group, and a butenyl group. A C₂-C₆₀ alkenylene group as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkenyl group.

[0260] A C₂-C₆₀ alkynyl group as used herein refers to a hydrocarbon group having at least one carbon triple bond in the middle or at the terminal of the C₂-C₆₀ alkyl group. Detailed examples thereof are an ethynyl group, and a propynyl group. A C₂-C₆₀ alkynylene group as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

[0261] A C₃-C₁₀ cycloalkyl group as used herein refers to a monovalent hydrocarbon monocyclic group having 3 to 10 carbon atoms. Detailed examples thereof are a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. A C₃-C₁₀ cycloalkylene group as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

[0262] A C₁-C₁₀ heterocycloalkyl group as used herein refers to a monovalent monocyclic group having at least one hetero atom selected from N, O, P, and S as a ring-forming atom and 1 to 10 carbon atoms. Detailed examples thereof are a tetrahydrofuran group, and a tetrahydrothiophene group. A C₁-C₁₀ heterocycloalkylene group as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

[0263] A C₃-C₁₀ cycloalkenyl group as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one double bond in the ring thereof, and which is not aromatic. Detailed examples thereof are a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. A C₃-C₁₀ cycloalkenylene group as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

[0264] A C₁-C₁₀ heterocycloalkenyl group as used herein refers to a monovalent monocyclic group that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one double bond in its ring. Detailed examples of the C₁-C₁₀ heterocycloalkenyl group are a 2,3-dihydrofuranyl group and a 2,3-dihydrothiophenyl group. A C₁-C₁₀ heterocycloalkenylene group as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkenyl group.

[0265] A C₆-C₆₀ aryl group as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C₆-C₆₀ arylene group as used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Detailed examples of the C₆-C₆₀ aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be fused to each other.

[0266] A C₁-C₆₀ heteroaryl group as used herein refers to a monovalent group having a carbocyclic aromatic system that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, and 1 to 60 carbon atoms. A C₁-C₆₀ heteroarylene group as used herein refers to a divalent group having a carbocyclic aromatic system that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, and 1 to 60 carbon atoms. Examples of the C₁-C₆₀ heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be fused to each other.

[0267] A C₆-C₆₀ aryloxy group as used herein indicates —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and a C₆-C₆₀ arylthio group as used herein indicates —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

[0268] A monovalent non-aromatic condensed polycyclic group as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed to each other, only carbon atoms as a ring forming atom, and which is non-aromatic in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. A divalent non-aromatic condensed polycyclic group as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0269] A monovalent non-aromatic condensed heteropolycyclic group as used herein refers to a monovalent group (for example, having 2 to 60 carbon atoms) that has two or more rings condensed to each other, has a heteroatom selected from N, O, P, and S, other than carbon atoms, as a ring forming atom, and which is non-aromatic in the entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazoyl group. A diva-

lent non-aromatic condensed heteropolycyclic group as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0270] Hereinafter, a compound and an organic light-emitting device according to embodiments are described in detail with reference to Synthesis Example and Examples. However, the organic light-emitting device is not limited thereto. The wording “B was used instead of A” used in describing Synthesis Examples means that an amount (molar equivalents) of A used was identical to an amount of B used.


EXAMPLE

Synthesis Example 1

Synthesis of Compound 1

Synthesis of Intermediate (A)

[0271]




[0272] 2,5-dibromopyridine (2.35 grams (g), 10 millimoles (mmol)), a phenylboronic acid (1.34 g, 11 mmol), palladium (0) acetate (Pd(OAc)₂, (0.09 g, 0.4 mmol), triphenylphosphine (0.26 g, 1.0 mmol), and potassium carbonate (K₂CO₃, 3.4 g, 25 mmol) were placed into a two-neck flask. Acetonitrile (CH₃CN, 80 milliliters (mL)) and methanol (40 mL) were added thereto, the resultant mixture was purged with nitrogen, and stirred at a temperature of 50° C. for 24 hours. Once the reaction was complete, the reaction was cooled to room temperature and filtered to remove white solid therefrom. The obtained residual solution was concentrated under reduced pressure. The product was extracted with methylene chloride (MC)/H₂O. The organic layer was washed with water and brine, and dried with magnesium sulfate (MgSO₄). The crude product was purified by column chromatography (hexane/MC=1/1) to obtain Intermediate (A) (2.1 g, yield=92%) as white solid.

[0273] MS: m/z calcd 227.38, found 227.95.

Synthesis of Intermediate (B)

[0274]




[0275] Intermediate (A) (4.7 g, 20 mmol), 4-(trimethylsilyl)phenylboronic acid (4.6 g, 24 mmol), tetrakis(triphenylphosphine)palladium(0) ($\text{Pd(PPh}_3)_4$, 1.6 g, 1.4 mmol) and potassium carbonate (K_2CO_3 , 6.9 g, 50 mmol) were placed into a two-neck flask. 70 mL of tetrahydrofuran (THF) and 15 mL of H_2O were added thereto, and the resultant mixture was purged with nitrogen and refluxed for 12 hours. Thereafter, the resultant mixture was cooled to room temperature. The product was extracted with methylene chloride (MC) to obtain an organic layer, which was dried with magnesium sulfate (MgSO_4), concentrated, and purified with column chromatography (hexane/MC=1/2) to obtain Intermediate (B) (3.9 g, yield=65%).

[0276] MS: m/z calcd 302.47. found 303.35.

Synthesis of Intermediate (C)


[0277]



[0278] 2-phenylpyridine (12.4 g, 80 mmol), iridium chloride (12.5 g, 35 mmol), ethoxyethanol (150 mL), and water (50 mL) were placed into a round flask, and refluxed at a temperature of 120° C. for 12 hours. The temperature was decreased to room temperature, and the resultant mixture was added to water to precipitate a solid. The solid was filtered, and sequentially washed with water, methanol, ether, and hexane to obtain Intermediate (C) (13 g, yield=70%).

Synthesis of Compound 1

[0279]



[0280] Intermediate (C) (4.28 g, 4 mmol), silver (I) trifluoromethane sulfonate (AgOTf , 2.56 g, 8 mmol), methylene chloride (MC) (100 mL), and methanol (30 mL) were placed into a round flask, and refluxed for 12 hours. The temperature was decreased to room temperature, and the resultant mixture was filtered to remove white solid therefrom. The solvent was removed under reduced pressure to obtain a solid (5.17 g, 7.2 mmol). The solid and Intermediate (B) (3.27 g, 10.8 mmol) were dissolved in ethanol, and refluxed for 12 hours. Once the reaction was complete, the temperature was decreased to room temperature. The yellow solid obtained by filtration was purified by column chromatography (hexane/MC=1/1) to obtain Compound 1 (1.5 g, yield=23%).

[0281] MS: m/z calcd 803.06. found 804.00.

[0282] ^1H NMR (CDCl_3 , 300 MHz): 7.89~7.84 (m, 3H), 7.69~7.62 (m, 4H), 7.45~7.41 (m, 2H), 7.38~7.36 (m, 4H), 6.98~6.96 (d, 2H), 6.92~6.89 (m, 8H), 6.82~6.77 (m, 4H), 0.21 (s, 9H).

Synthesis Example 2


Synthesis of Compound 2

[0283] Compound 2 (1.2 g, yield=30%) was synthesized in the same manner as in Synthesis Example 1, except that in

synthesis of Intermediate (B), 3-(trimethylsilyl)phenylboronic acid was used instead of 4-(trimethylsilyl)phenylboronic acid.

[0284] MS: m/z calcd 817.08. found 817.94.

[0285] ^1H NMR (CDCl_3 , 300 MHz): 7.88~7.84 (m, 3H), 7.67~7.61 (m, 4H), 7.45~7.42 (m, 2H), 7.37~7.34 (m, 4H), 6.98~6.96 (d, 2H), 6.95~6.89 (m, 8H), 6.82~6.78 (m, 4H), 0.21 (s, 9H).




Synthesis Example 3

Synthesis of Compound 3

Synthesis of Intermediate (D)

[0286]




[0287] Intermediate (D) (yield=85%) was obtained in the same manner as Intermediate (A) in Synthesis Example 1, except that 2,5-dibromo-4-methylpyridine was used instead of 2,5-dibromopyridine.

[0288] MS: m/z calcd 248.12. found 249.01.

Synthesis of Intermediate (E)

[0289]




[0290] Intermediate (E) (yield of 62%) was synthesized in the same manner as Intermediate (B) in Synthesis Example 1, except that Intermediate (D) was used instead of Intermediate (A).

[0291] MS: m/z calcd 317.50. found 318.27.

Synthesis of Compound 3

[0292]



[0293] Compound 3 (yield=25%) was obtained in the same manner as Compound 1 in Synthesis Example 1, except that Intermediate (E) was used instead of Intermediate (B).

[0294] MS: m/z calcd 817.08. found 818.04.


[0295] ^1H NMR (CDCl_3 , 300 MHz): 7.78 (t, 2H), 7.67 (s, 1H), 7.64~7.50 (m, 4H), 7.47~7.42 (m, 2H), 7.41~7.39 (m, 4H), 7.35 (s, 1H), 7.02~7.00 (d, 2H), 6.89~6.81 (m, 8H), 6.81~6.77 (m, 2H), 2.32 (s, 3H), 0.20 (s, 9H).

Synthesis Example 4

Synthesis of Compound 4

Synthesis of Intermediate (F)

[0296]




[0297] Intermediate (E) (6.3 g, 20 mmol), lithium diisopropylamide (LDA, 2.2 equivalents (eq)) (4.7 g, 44 mmol), and THF (50 mL) were placed into a two-neck flask, and the mixture was stirred at a temperature of -70°C . for 30 minutes. 2-bromopropane (5.9 g, 48 mmol) was slowly added by drops thereto. The resultant mixture was subsequently stirred at a temperature of -70°C . for 24 hours. The temperature was slowly raised to room temperature. Water was added thereto to quench the reaction. The product was extracted with methylene chloride (MC). The combined organic layers were dried with magnesium sulfate (MgSO_4) and concentrated. The product was purified by column chromatography (hexane/MC=1/1) to obtain Intermediate (F) (5.2 g, yield=73%).


[0298] MS: m/z calcd 359.58. found 360.45.

Synthesis of Compound 4

[0299]



-continued



[0300] Compound 4 (yield=21%) was obtained in the same manner as Compound 1 in Synthesis Example 1, except that Intermediate (F) was used instead of Intermediate (B).

[0301] MS: m/z calcd 859.16. found 860.05.


[0302] ^1H NMR (CDCl_3 , 300 MHz): 7.79 (t, 2H), 7.65 (s, 1H), 7.64~7.50 (m, 4H), 7.47~7.43 (m, 2H), 7.42~7.39 (m, 4H), 7.35 (s, 1H), 7.02~7.00 (d, 2H), 6.89~6.81 (m, 8H), 6.81~6.77 (m, 2H), 2.52 (d, 2H), 1.90 (m, 1H), 0.93 (d, 6H), 0.23 (s, 9H).

Synthesis Example 5

Synthesis of Compound 9

Synthesis of Intermediate (G)

[0303]




[0304] Intermediate (G) (yield=82%) was obtained in the same manner as Intermediate (A) in Synthesis Example 1, except that [1,1'-biphenyl]-3-boronic acid was used instead of a phenylboronic acid.

[0305] MS: m/z calcd 324.21. found 325.15.

Synthesis of Intermediate (H)

[0306]




[0307] Intermediate (H) (yield of 78%) was synthesized in the same manner as Intermediate (B) of Synthesis Example 1, except that Intermediate (G) was used instead of Intermediate (A).

[0308] MS: m/z calcd 393.60. found 394.52.

Synthesis of Compound 9

[0309]



[0310] Compound 9 (yield=18%) was obtained in the same manner as Compound 1 in Synthesis Example 1, except that Intermediate (H) was used instead of Intermediate (B).

[0311] MS: m/z calcd 893.18. found 894.07.

[0312] ^1H NMR (CDCl_3 , 300 MHz): 7.80 (m, 2H), 7.64 (s, 1H), 7.62~7.51 (m, 4H), 7.47~7.43 (m, 2H), 7.41~7.38 (m, 4H), 7.34 (s, 1H), 7.02~7.00 (d, 4H), 6.89~6.80 (m, 12H), 2.32 (s, 3H), 0.21 (s, 9H)

Synthesis Example 6

Synthesis of Compound 5

[0313] Compound 5 was synthesized in the same manner as in Synthesis Example 1, except that in synthesis of Intermediate (B), 2-(methyl)-4-(trimethylsilyl)phenylboronic acid was used instead of 4-(trimethylsilyl)phenylboronic acid.

Synthesis Example 7

Synthesis of Compound 6

[0314] Compound 6 was synthesized in the same manner as in Synthesis Example 1, except that in synthesis of Intermediate (B), 2-(t-butyl)-4-(trimethylsilyl)phenylboronic acid was used instead of 4-(trimethylsilyl)phenylboronic acid.

Synthesis Example 8

Synthesis of Compound 7

[0315] Compound 7 was synthesized in the same manner as in Synthesis Example 3, except that in synthesis of Intermediate (B), 2-(methyl)-4-(trimethylsilyl)phenylboronic acid was used instead of 4-(trimethylsilyl)phenylboronic acid.

Evaluation Example 1

Evaluation of Thermal Characteristics

[0316] Compound 3 was subjected to thermal analysis (N_2 atmosphere, temperature range: room temperature to 800°C . (10 degrees Centigrade per minute ($^\circ\text{C}/\text{min}$))-TGA, room temperature to 450°C .-DSC, Pan Type: Pt Pan in disposable Al Pan (TGA), disposable Al pan (DSC)) using thermogravimetric analysis (TGA) and differential scanning calorimetry (DSC). The obtained results are shown in FIG. 2 and Table 2 below. Referring to FIG. 2 and Table 2, it was confirmed that Compound 3 had excellent thermal stability.

TABLE 2

	Td ($^\circ\text{C}$.) (1%, 5%)	Tg ($^\circ\text{C}$.)	Tm ($^\circ\text{C}$.)
Compound 3	374, 397	not detected.	not detected.

Evaluation Example 2

Evaluation on HOMO, LUMO, and T_1 Energy Levels

[0317] HOMO, LUMO and T_1 energy levels of Compound 3 were evaluated according to the method indicated in Table 3, and results thereof are shown in Table 4.

TABLE 3

HOMO energy level	Each compound was diluted at a concentration of 1×10^{-5} molar (M) in CHCl_3 , and an UV absorption spectrum thereof was measured at room temperature
-------------------	--

TABLE 3-continued

evaluation method	by using a Shimadzu UV-350 spectrometer. The HOMO energy level thereof was calculated by using an optical band gap (Eg) from an edge of the absorption spectrum.
LUMO energy level evaluation method	A potential (Volts, V)-current (Amperes, A) graph of each compound was obtained by using cyclic voltammetry (CV) (electrolyte: 0.1M Bu ₄ NClO ₄ /solvent: CH ₂ Cl ₂ /electrode: 3 electrode system (working electrode: GC, reference electrode: Ag/AgCl, auxiliary electrode: Pt)). Subsequently, from reduction onset of the graph, the LUMO energy level of the compound was calculated.
T1 energy level evaluation method	A mixture (each compound was dissolved in an amount of 1 milligram (mg) in 3 cubic centimeters (cc) of toluene) of toluene and each compound was loaded into a quartz cell, and then, the resultant quartz cell was loaded into liquid nitrogen (77 Kelvins (K)). A photoluminescence spectrum thereof was measured by using a device for measuring photoluminescence. The obtained spectrum was compared with a photoluminescence spectrum measured at room temperature, and peaks observed only at low temperature were analyzed to calculate T1 energy levels.

TABLE 4

Compound No.	HOMO (eV) (found)	LUMO (eV) (found)	T ₁ energy level (eV)
Compound 1	-5.049	-2.521	2.375
Compound 2	-5.043	-2.523	2.396
Compound 3	-5.016	-2.486	2.455
Compound 4	-5.007	-2.400	2.507
Compound 5	-5.045	-2.507	2.432
Compound 6	-5.040	-2.465	2.451
Compound 7	-5.013	-2.379	2.510

[0318] From Table 4, it was confirmed that Compounds 1 to 7 have electric characteristics that are suitable for use as a material for forming an organic light-emitting device.

Evaluation Example 3

Luminance Characteristics Evaluation

[0319] UV absorption spectrum and photoluminescence (PL) spectrum of Compound 3 were analyzed to evaluate luminance characteristics thereof. First, Compound 3 was diluted in toluene to a concentration of 0.2 millimolar (mM), and then analyzed by Shimadzu UV-350 Spectrometer to obtain an UV absorption spectrum. Compound 3 was diluted in toluene to a concentration of 10 mM, and then analyzed by Xenon-equipped ISC PC1 spectrofluorometer to obtain a PL spectrum. Results thereof are shown in Table 3.

[0320] Referring to FIG. 3, it was confirmed that Compound 3 has luminance characteristics that are suitable for use as a material for an organic light-emitting device.

Evaluation Example 4

Process Stability Evaluation

[0321] Heat-resistant stability and process stability of Compound 3 were evaluated by analyzing purify levels before and after Compound 3 and Compound A were heat treated. First, Compound 3 and Compound A were heat treated under a nitrogen atmosphere (180° C., 100 hours), and a change in purity of Compound 3 and Compound A was

measured by high performance liquid chromatography (HPLC). Results thereof are shown in FIG. 4 (Compound 3) and FIG. 5 (Compound A), and summarized in Table 5.

Compound A

[0322]




TABLE 5

Compound No.	Purity level before the heat treatment (%)	Purity level after the heat treatment (%)	Purity change (%)
Compound 3	99.47	99.46	0.01
Compound A	99.30	95.90	3.80

[0323] Referring to Table 5 and FIGS. 4 and 6, it was confirmed that Compound 3 had heat-resistant stability. Accordingly, it was confirmed that Compound 3 had excellent process stability that minimizes the difference in characteristics of a device depending on a process time during a deposition process that is required in manufacturing an organic light-emitting device.

Example 1

[0324] A glass substrate with ITO/Ag/ITO (70 Å/1,000 Å/70 Å) as an anode thereon was cut to a size of 50 millimeters (mm)×50 mm×0.5 mm, sonicated in isopropyl alcohol and pure water for about 5 minutes, and cleaned by irradiation of ultraviolet rays for 30 minutes and exposure to ozone. The resultant structure was subsequently mounted on a vacuum deposition apparatus.

[0325] 2-TNATA was deposited on the anode to form a hole injection layer having a thickness of 60 nanometers (nm), and NPB was deposited on the hole injection layer to form a hole transport layer having a thickness of 100 nm.

[0326] CBP (host) and Compound 1 (dopant) were co-deposited on the hole transport layer at a weight ratio of 90:10 to form an emission layer having a thickness of 25 nm. BCP was deposited on the emission layer to form a hole blocking layer having a thickness of 5 nm. Alq₃ was deposited on the

hole blocking layer to form an electron transport layer having a thickness of 35 nm. LiF was deposited on the electron transport layer to form an electron injection layer having a thickness of 1 nm. Mg and Ag were co-deposited on the electron injection layer at a weight ratio of 9:1 to form a cathode having a thickness of 12 nm, thereby completing manufacture of an organic light-emitting device (emission of green light).

Examples 2 to 5 and Comparative Example 1

[0327] An organic light-emitting device was manufactured in the same manner as in Example 1, except that in forming an emission layer, a dopant shown in Table 6 was used instead of Compound 1.

Evaluation Example 5

Evaluation of Characteristics of Organic Light-Emitting Devices

[0328] The driving voltage, efficiency, luminance, lifespan (T_{97}), and color purity of the organic light-emitting devices of Examples 1 to 5 and Comparative Example 1 were evaluated, and evaluation results are shown in Table 6. EL spectra of the organic light-emitting devices of Examples 1, 3, 4 and 5 are shown in Table 6. FIG. 7 shows a time-luminance (at 10 milliAmperes per square meter (mA/m^2)) graph of the organic light-emitted device of Example 3. The efficiency, luminance, and color coordinate of each of the organic light-emitting device were measured by using a luminance meter PR650 while power source was supplied thereto by a current-voltage meter (Kethley SMU 236). A lifespan was obtained by measuring an amount of time that lapsed when luminance was reduced from 100% (initial luminance) to 97% at the current density of 10 milliAmperes per square centimeter (mA/cm^2).

TABLE 6

	Dopant	Driving Voltage (V)	Efficiency (Cd/A)	Luminance (cd/m^2)	T_{97} (hr)		
					(at 10 mA/ cm^2)	Color coordinate	
						CIEx	CIy
Example 1	Compound 1	5.5	99.8	9000	123	0.37	0.59
Example 2	Compound 2	5.3	92.4	9000	100	0.34	0.60
Example 3	Compound 3	5.3	87.6	9000	230	0.34	0.61
Example 4	Compound 4	5.5	100.4	9000	250	0.35	0.60
Example 5	Compound 9	5.1	95.4	9000	114	0.36	0.59

TABLE 6-continued

	Dopant	Driving Voltage (V)	Efficiency (Cd/A)	Luminance (cd/m^2)	T_{97} (hr)		
					(at 10 mA/ cm^2)	Color coordinate	
						CIEx	CIy
Comparative Example 1	Ir(PPY) ₃	5.8	90.2	9000	35	0.33	0.61






TABLE 6-continued

Dopant	Driving Voltage (V)	Efficiency (Cd/A)	Luminance (cd/m ²)	T ₉₇ (hr) (at 10 mA/cm ²)	Color coordinate	
					CIEx	CIEy
						
						

[0329] Referring to Table 6, it was confirmed that the organic light-emitting devices manufactured according to Examples 1 to 5 have a lower driving voltage, higher efficiency, higher luminance, a longer lifespan, and a higher color purity than the organic light-emitting devices manufactured according to Comparative Example 1.

[0330] The organometallic compound according to embodiments has excellent electric characteristics and thermal stability. Accordingly, an organic light-emitting device including the organometallic compound may have a low driving voltage, high efficiency, and high color purity.

[0331] It should be understood that the exemplary embodiments described therein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each exemplary embodiment should typically be considered as available for other similar features or aspects in other exemplary embodiments.


[0332] While one or more exemplary embodiments have been described with reference to the figures, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing

from the spirit and scope of the present inventive concept as defined by the following claims.

What is claimed is:

1. An organometallic compound represented by Formula 1:

Formula 1



wherein in Formula 1,

CY₁ is selected from a C₅-C₆₀ carbocyclic group and a C₁-C₆₀ heterocyclic group,

Q₁ to Q₃ are each independently selected from

a C₁-C₂₀ alkyl group and a phenyl group; and

a C₁-C₂₀ alkyl group and a phenyl group, each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, and a phenyl group,

R₁ to R₈ are each independently selected from a hydrogen, a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃,

R₁₁ to R₁₄ are each independently selected from

a hydrogen, a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a

C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group; and

a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a

C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group, each substituted with at least one

selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a

C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,

each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a phenyl group, and a naphthyl group,


- C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;
- a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;
- a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and —N(Q₁₁)(Q₁₂) and —P(=O)(Q₁₃)(Q₁₄),
- a1 and b1 are each independently an integer selected from 0 to 4, provided that when a1 is 2 or more, 2 or more, groups R₁₁ are identical or different, and when b1 is 2 or more, 2 or more groups R₂₁ are identical or different, and Q₁₁ to Q₁₄ are each independently selected from a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, and a C₆-C₁₄ aryl group; a C₆-C₁₄ aryl group substituted with at least one selected from a C₁-C₂₀ alkyl group and a C₆-C₁₄ aryl group; a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.
2. The organometallic compound of claim 1, wherein CY₁ is selected from a benzene, a dibenzofuran, a dibenzothiophene, a fluorene, and a carbazole.
3. The organometallic compound of claim 1, wherein Q₁ to Q₃ are each independently selected from a C₁-C₁₀ alkyl group and a phenyl group; and a C₁-C₁₀ alkyl group and a phenyl group, each substituted with at least one selected from a deuterium and a C₁-C₁₀ alkyl group.
4. The organometallic compound of claim 1, wherein Q₁ to Q₃ are each independently selected from —CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCH₂, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂; an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group; and
5. The organometallic compound of claim 1, wherein Q₁ to Q₃ are all identical; Q₁ and Q₃ are identical, and Q₂ and Q₁ are different from each other; or Q₁ to Q₃ are all different from each other.
6. The organometallic compound of claim 1, wherein Q₁ and Q₃ are each independently selected from —CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCH₂, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂, Q₂ is selected from an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group; and an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, and a phenyl group, each substituted with at least one selected from a deuterium and a C₁-C₁₀ alkyl group.
7. The organometallic compound of claim 1, wherein R₁ to R₈ are all a hydrogen; R₁, R₂, and R₄ to R₈ are a hydrogen, and R₃ is selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃; R₁, R₂, R₄, R₅, R₆ and R₈ are a hydrogen, and R₃ and R₇ are each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃; R₁ to R₄ are a hydrogen, and R₅ to R₈ are each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃; R₁ to R₅ are a hydrogen, and R₆ to R₈ are each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃; R₁ to R₄ and R₈ are a hydrogen, and R₅ to R₇ are each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃; or R₁ to R₈ are each independently selected from a deuterium, —CH₃, —CDH₂, —CD₂H, and —CD₃.
8. The organometallic compound of claim 1, wherein R₁₁ to R₁₄ are each independently selected from a hydrogen, a deuterium, a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an isohexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an isoheptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an isooctyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an isodecyl group, a sec-decyl group, a tert-decyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a

norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, and a naphthyl group; and


a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an isohexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an isoheptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an isooctyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an isodecyl group, a sec-decyl group, a tert-decyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, and a naphthyl group.

9. The organometallic compound of claim 1, wherein

R₁₁ to R₁₄ are each independently selected from a hydrogen, a deuterium, —CH₃, —CD₃, —CD₂H, —CDH₂, a group represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-12:



Formula 9-1



Formula 9-2




Formula 9-3



Formula 9-4




Formula 9-5




Formula 9-6



Formula 9-7




Formula 9-8




Formula 9-9




-continued




Formula 9-10




Formula 9-11




Formula 9-12




Formula 9-13




Formula 9-14




Formula 9-15




Formula 9-16




Formula 9-17




Formula 10-1




Formula 10-2




Formula 10-3



Formula 10-4




Formula 10-5




Formula 10-6


-continued




Formula 10-7




Formula 10-8




Formula 10-9



Formula 10-10



Formula 10-11



Formula 10-12

10. The organometallic compound of claim 1, wherein R_{21} is selected from

a hydrogen, a deuterium, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{10} alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidi-

nyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthroli-nyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthroli-nyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthroli-nyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl

group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

$-\text{B}(\text{Q}_{11})(\text{Q}_{12})$ and $-\text{P}(=\text{O})(\text{Q}_{13})(\text{Q}_{14})$,

Q_{11} to Q_{14} are each independently selected from

a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, and a carbazolyl group; and a phenyl group and a naphthyl group, each substituted with at least one selected from a C_1 - C_{20} alkyl group, a phenyl group, and a naphthyl group.

11. The organometallic compound of claim 1, wherein

R_{21} is selected from

a hydrogen, a deuterium, a cyano group, a nitro group, a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an isohexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an isoheptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an isooctyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an isodecyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an isohexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an isoheptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an isooctyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an isodecyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from a deuterium, $-\text{F}$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantyl group, a norbornyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

$-\text{B}(\text{Q}_{11})(\text{Q}_{12})$ and $-\text{P}(=\text{O})(\text{Q}_{13})(\text{Q}_{14})$, and

Q_{11} to Q_{14} are each independently selected from

a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, and a naphthyl group; and


a phenyl group and a naphthyl group, each substituted with at least one selected from a C_1 - C_{10} alkyl group, a phenyl group, and a naphthyl group.

12. The organometallic compound of claim 1, wherein

R_{21} is selected from a hydrogen, a deuterium, a cyano group, a nitro group, $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, groups represented by Formulae 9-1 to 9-17, and groups represented by Formulae 10-1 to 10-30:



Formula 9-1



Formula 9-2




Formula 9-3




Formula 9-4




Formula 9-5




Formula 9-6




Formula 9-7




Formula 9-8




Formula 9-9




Formula 9-10




Formula 9-11




Formula 9-12





Formula 9-13















Formula 9-14




Formula 9-15




-continued




Formula 9-16




Formula 9-17




Formula 10-1




Formula 10-2




Formula 10-3




Formula 10-4



Formula 10-5




Formula 10-6




Formula 10-7

Formula 10-8



Formula 10-9

Formula 10-10



-continued

Formula 10-11

Formula 10-12

Formula 10-13

Formula 10-14

Formula 10-15

Formula 10-16










Formula 10-17

Formula 10-18


Formula 10-19

Formula 10-20

-continued



-continued



Formula 10-21

Formula 10-30

Formula 10-22

13. The organometallic compound of claim 1, wherein the organometallic compound is represented by one of Formulae 1-1 to 1-12:


Formula 10-23

Formula 1-1

Formula 10-24

Formula 10-25

Formula 10-26




Formula 10-27

Formula 1-2


Formula 10-28

Formula 10-29



-continued



Formula 1-3



Formula 1-4



-continued

Formula 1-6




Formula 1-5

Formula 1-7




Formula 1-8




-continued

Formula 1-9




-continued

Formula 1-12




Formula 1-10




wherein in Formulae 1-1 to 1-12, CY_1 , Q_1 to Q_3 , R_1 to R_8 , R_{11} to R_{14} , R_{21} , and $b1$ are the same as in claim 1, and R_{11a} and R_{11b} are the same as R_{11} , provided that each of R_{11} , R_{11a} , and R_{11b} in Formulae 1-1 to 1-12 is not a hydrogen.

14. The organometallic compound of claim 1, wherein the organometallic compound is represented by one of Formulae 1(1) to 1(12):

Formula 1(1)




Formula 1-11




-continued

Formula 1(2)




-continued


Formula 1(5)




Formula 1(3)




Formula 1(6)



Formula 1(4)




Formula 1(7)




-continued


Formula 1(8)



Formula 1(9)




Formula 1(10)




-continued

Formula 1(11)








Formula 1(12)




wherein in Formulae 1(1) to 1(12), Q₁ to Q₃, R₁ to R₈, R₁₁, a₁, R₁₃ and R₂₁ are the same as in claim 1, X₁ is N(R₂₁), O, or S, and R₂₂ to R₂₄ are the same as R₂₁, provided that each of R₁₃ and R₂₁ to R₂₄ in Formulae 1(1) to 1(12) is not a hydrogen.

15. The organometallic compound of claim 1, wherein the organometallic compound is represented by one of Formulae 2-1 to 2-72:




-continued


Formula 2-6



Formula 2-7




Formula 2-8




-continued


Formula 2-9



Formula 2-10




Formula 2-11




-continued

Formula 2-12




-continued


Formula 2-15




Formula 2-16



Formula 2-14




Formula 2-17




-continued

Formula 2-18




-continued


Formula 2-21




Formula 2-19




Formula 2-22



Formula 2-20




Formula 2-23




-continued


Formula 2-24



Formula 2-25




Formula 2-26




-continued


Formula 2-27



Formula 2-28




Formula 2-29




-continued


Formula 2-30



Formula 2-31




Formula 2-32




-continued


Formula 2-33



Formula 2-34




Formula 2-35




-continued


Formula 2-39



Formula 2-40




Formula 2-41




-continued


Formula 2-42



Formula 2-43




Formula 2-44




-continued


Formula 2-45



Formula 2-46




Formula 2-47




-continued


Formula 2-48



Formula 2-49




Formula 2-50




-continued


Formula 2-51



Formula 2-52




Formula 2-53




-continued


Formula 2-54



Formula 2-55




Formula 2-56




-continued


Formula 2-57



Formula 2-58




Formula 2-59




-continued


Formula 2-60



Formula 2-61




Formula 2-62




-continued


Formula 2-63



Formula 2-64




Formula 2-65




-continued


Formula 2-66



Formula 2-67




Formula 2-68




-continued


Formula 2-69



Formula 2-70





Formula 2-71



-continued

-continued

Formula 2-72




2

3



wherein in with Formulae 2-1 to 2-72, Q_1 to Q_3 , R_1 to R_8 , R_{11} , R_{13} , and R_{21} are the same as in claim 1, X_1 is $N(R_{21})$, O, or S, R_{11a} and R_{11b} are the same as R_{11} , and R_{22} to R_{24} are the same as R_{21} , provided that, each of R_{11} , R_{11a} , R_{11b} , R_{13} and R_{21} to R_{24} in Formulae 2-1 to 2-72 is not a hydrogen.

16. The organometallic compound of claim 1, wherein the organometallic compound is one of Compounds 1 to 54:




4

1




-continued




5


-continued




8




6



9




7




10


-continued




-continued




12




15




13




16





-continued





-continued



18




19




-continued

23




-continued

26




24



27




25




28





-continued





-continued




33




34




-continued




-continued




36




37




39








40




-continued






-continued




-continued



-continued




-continued




51


-continued



54



52



53

17. An organic light-emitting device comprising:

a first electrode;

a second electrode; and

an organic layer disposed between the first electrode and the second electrode,

wherein the organic layer comprises an emission layer, and at least one organometallic compound of claim 1.

18. The organic light-emitting device of claim 17, wherein

the first electrode is an anode,

the second electrode is a cathode, and

the organic layer comprises:

i) a hole transport region disposed between the first electrode and the emission layer, wherein the hole transport region comprises at least one of a hole injection layer, a hole transport layer, and an electron blocking layer, and

ii) an electron transport region disposed between the emission layer and the second electrode, wherein the electron transport region comprises at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

19. The organic light-emitting device of claim 17, wherein the emission layer comprises the organometallic compound of claim 1.

20. The organic light-emitting device of claim 19, wherein the emission layer further comprises a host, and wherein an amount of the organometallic compound of claim 1 in the emission layer is smaller than an amount of the host.

* * * * *

专利名称(译)	有机金属化合物和包括其的有机发光器件		
公开(公告)号	US20160141526A1	公开(公告)日	2016-05-19
申请号	US14/676242	申请日	2015-04-01
[标]申请(专利权)人(译)	三星电子株式会社		
申请(专利权)人(译)	SAMSUNG ELECTRONICS CO. , LTD.		
当前申请(专利权)人(译)	SAMSUNG ELECTRONICS CO. , LTD.		
[标]发明人	LEE BANGLIN CHOI HYEONHO PARK YOUNGJAE IM KYUHYUN		
发明人	LEE, BANGLIN CHOI, HYEONHO PARK, YOUNGJAE IM, KYUHYUN		
IPC分类号	H01L51/00 C09K11/06 C07F15/00		
CPC分类号	H01L51/0094 C07F15/0033 C09K11/06 C09K2211/1055 H01L51/5206 H01L51/5221 H01L2251/552 H01L51/5056 H01L51/5088 H01L51/5096 H01L51/5072 H01L51/5092 H01L51/5004 H01L51/5012 C09K2211/1007 C09K2211/1014 C09K2211/1029 C09K2211/185 H01L51/0081 H01L51/0085 H01L51/5016		
优先权	1020140158906 2014-11-14 KR		
其他公开文献	US9758719		
外部链接	Espacenet USPTO		

摘要(译)

由式1表示的有机金属化合物：其中在式1中，基团和变量与说明书中描述的相同。

Formula 1

