


(19) **United States**(12) **Patent Application Publication**  
**LEE et al.**(10) **Pub. No.: US 2019/0074451 A1**(43) **Pub. Date: Mar. 7, 2019**(54) **CONDENSED CYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**(71) Applicants: **Samsung Electronics Co., Ltd.**, Suwon-si (KR); **Samsung SDI Co., Ltd.**, Yongin-si (KR)(72) Inventors: **Seungjae LEE**, Suwon-si (KR); **Byungku KIM**, Suwon-si (KR); **Youngkwon KIM**, Suwon-si (KR); **Changwoo KIM**, Suwon-si (KR); **Hyungsun KIM**, Suwon-si (KR); **Changju SHIN**, Suwon-si (KR); **Eunsun YU**, Suwon-si (KR); **Byoungki CHOI**, Hwaseong-si (KR); **Kyu-Young HWANG**, Suwon-si (KR)**Publication Classification**(51) **Int. Cl.**  
*H01L 51/00* (2006.01)  
*C07D 405/14* (2006.01)  
*C09K 11/06* (2006.01)  
*C07D 491/048* (2006.01)  
(52) **U.S. Cl.**  
CPC ..... *H01L 51/0072* (2013.01); *C07D 405/14* (2013.01); *C09K 11/06* (2013.01); *H01L 51/5088* (2013.01); *H01L 51/0073* (2013.01); *C07D 491/048* (2013.01); *H01L 51/0071* (2013.01); *H01L 51/0067* (2013.01)(57) **ABSTRACT**  
A condensed-cyclic compound represented by Formula 1A:

Formula 1A

(21) Appl. No.: **16/177,914**(22) Filed: **Nov. 1, 2018****Related U.S. Application Data**

(63) Continuation-in-part of application No. 14/573,422, filed on Dec. 17, 2014, now Pat. No. 10,158,085.

(30) **Foreign Application Priority Data**

Dec. 17, 2013 (KR) ..... 10-2013-0157532

wherein in Formula 1A, groups, substituents, and variables are the same as defined in the specification.

**10****19****15****11**

# FIGURE

10

19
15
11

**CONDENSED CYCLIC COMPOUND AND  
ORGANIC LIGHT-EMITTING DEVICE  
INCLUDING THE SAME**

CROSS-REFERENCE TO RELATED  
APPLICATIONS

[0001] This application is a continuation-in-part of U.S. patent application Ser. No. 14/573,422, filed on Dec. 17, 2014, which claims priority to Korean Patent Application No. 10-2013-0157532, filed on Dec. 17, 2013, and all the benefits accruing therefrom under 35 U.S.C. § 119, the contents of which are incorporated herein in their entireties by reference.

BACKGROUND

1. Field

[0002] One or more embodiments of the present disclosure relate to condensed-cyclic compounds and organic light-emitting devices including the condensed-cyclic compounds.

2. Description of the Related Art

[0003] Organic light-emitting devices (OLEDs) are self-emitting devices that have advantages such as wide viewing angles, excellent contrast ratios, and quick response times. In addition, OLEDs exhibit excellent brightness, driving voltage, and response speed characteristics, and can provide multicolored images.

[0004] A typical OLED has a structure including an anode, a cathode, and an organic layer disposed between the anode and the cathode and including an emission layer. A hole transporting region may be disposed between the anode and the cathode, and an electron transporting region may be disposed between the emission layer and the cathode. Holes injected from the anode move to the EML via the hole transport region, and electrons injected from the cathode move to the EML via the electron transport region. Carriers such as holes and electrons recombine in the EML to generate excitons. When the excitons drop from an excited state to a ground state, light is emitted.

[0005] Different types of organic light emitting devices are known. However, there still remains a need in OLEDs having low driving voltage, high efficiency, high brightness, and long lifespan.


SUMMARY

[0006] One or more embodiments include novel condensed-cyclic compounds and organic light-emitting devices including the condensed-cyclic compounds.


[0007] Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments.

[0008] According to one or more embodiments, provided is a condensed-cyclic compound represented by Formula 1A or 1B:


Formula 1A




Formula 1B



Formula 1C



Formula 1D



[0009] In the Formulae above,

[0010] ring A<sub>1</sub> in Formulae 1A and 1B is represented by Formula 1C or 1D;

[0011] X<sub>1</sub> is N or C-[(L<sub>1</sub>)<sub>a1</sub>-(R<sub>1</sub>)<sub>b1</sub>],

[0012] X<sub>2</sub> is N or C-[(L<sub>2</sub>)<sub>a2</sub>-(R<sub>2</sub>)<sub>b2</sub>],

[0013] X<sub>3</sub> is N or C-[(L<sub>3</sub>)<sub>a3</sub>-(R<sub>3</sub>)<sub>b3</sub>], and

[0014] at least one of X<sub>1</sub> to X<sub>3</sub> is N;

[0015] X<sub>4</sub> is O or S;

[0016] X<sub>11</sub> is selected from N-[(L<sub>12</sub>)<sub>a12</sub>-(R<sub>12</sub>)<sub>b12</sub>], S, O, S(=O), S(=O)<sub>2</sub>, C(=O), C(R<sub>13</sub>)(R<sub>14</sub>), Si(R<sub>13</sub>)(R<sub>14</sub>), P(R<sub>13</sub>), P(=O)(R<sub>13</sub>), and C=N(R<sub>12</sub>);

[0017] L<sub>1</sub> to L<sub>4</sub>, L<sub>11</sub>, L<sub>12</sub> and L<sub>21</sub> may be each independently selected from a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkylene group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenylene group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylene group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic hetero-condensed polycyclic group;

[0018] a<sub>1</sub> to a<sub>4</sub>, a<sub>11</sub>, a<sub>12</sub>, and a<sub>21</sub> may be each independently selected from integers of 0 to 3;

[0019] R<sub>1</sub> to R<sub>3</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>11</sub> to R<sub>17</sub> may be each independently selected from a hydrogen, a deuterium, —F (a fluoro group), —Cl (a chloro group), —Br (a bromo group),

—I (an iodo group), a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted  $C_1-C_{60}$  alkyl group, a substituted or unsubstituted  $C_2-C_{60}$  alkenyl group, a substituted or unsubstituted  $C_2-C_{60}$  alkynyl group, a substituted or unsubstituted  $C_1-C_{60}$  alkoxy group, a substituted or unsubstituted  $C_3-C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_2-C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3-C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_2-C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6-C_{60}$  aryl group, a substituted or unsubstituted  $C_6-C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6-C_{60}$  arylthio group, a substituted or unsubstituted  $C_2-C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group,  $-N(Q_1)(Q_2)$ ,  $-Si(Q_3)(Q_4)(Q_5)$ , and  $-B(Q_6)(Q_7)$ ,

**[0020]**  $R_4$  is selected from a substituted or unsubstituted  $C_3-C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_2-C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3-C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_2-C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6-C_{60}$  aryl group, a substituted or unsubstituted  $C_6-C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6-C_{60}$  arylthio group, a substituted or unsubstituted  $C_2-C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group;

**[0021]** b1 to b6 and b11 to b17 may be each independently selected from integers of 1 to 3;

**[0022]** at least one substituent of the substituted  $C_3-C_{10}$  cycloalkylene group, substituted  $C_2-C_{10}$  heterocycloalkylene group, substituted  $C_3-C_{10}$  cycloalkenylene group, substituted  $C_2-C_{10}$  heterocycloalkenylene group, substituted  $C_6-C_{60}$  arylene group, substituted  $C_2-C_{60}$  heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic hetero-condensed polycyclic group, substituted  $C_1-C_{60}$  alkyl group, substituted  $C_2-C_{60}$  alkenyl group, substituted  $C_2-C_{60}$  alkynyl group, substituted  $C_1-C_{60}$  alkoxy group, substituted  $C_3-C_{10}$  cycloalkyl group, substituted  $C_2-C_{10}$  heterocycloalkyl group, substituted  $C_3-C_{10}$  cycloalkenyl group, substituted  $C_2-C_{10}$  heterocycloalkenyl group, substituted  $C_6-C_{60}$  aryl group, substituted  $C_6-C_{60}$  aryloxy group, substituted  $C_6-C_{60}$  arylthio group, substituted  $C_2-C_{60}$  heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic hetero-condensed polycyclic group may be selected from:

**[0023]** a deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1-C_{60}$  alkyl group, a  $C_2-C_{60}$  alkenyl group, a  $C_2-C_{60}$  alkynyl group, and a  $C_1-C_{60}$  alkoxy group;

**[0024]** a  $C_1-C_{60}$  alkyl group, a  $C_2-C_{60}$  alkenyl group, a  $C_2-C_{60}$  alkynyl group, and a  $C_1-C_{60}$  alkoxy group, each substituted with at least one of a deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a

hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a  $C_3-C_{10}$  cycloalkyl group, a  $C_2-C_{10}$  heterocycloalkyl group, a  $C_3-C_{10}$  cycloalkenyl group, a  $C_2-C_{10}$  heterocycloalkenyl group, a  $C_6-C_{60}$  aryl group, a  $C_6-C_{60}$  aryloxy group, a  $C_6-C_{60}$  arylthio group, a  $C_2-C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group,  $-N(Q_{11})(Q_{12})$ ,  $-Si(Q_{13})(Q_{14})(Q_{15})$  and  $-B(Q_{16})(Q_{17})$ ;

**[0025]** a  $C_3-C_{10}$  cycloalkyl group, a  $C_2-C_{10}$  heterocycloalkyl group, a  $C_3-C_{10}$  cycloalkenyl group, a  $C_2-C_{10}$  heterocycloalkenyl group, a  $C_6-C_{60}$  aryl group, a  $C_6-C_{60}$  aryloxy group, a  $C_6-C_{60}$  arylthio group, a  $C_2-C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group;


**[0026]** a  $C_3-C_{10}$  cycloalkyl group, a  $C_2-C_{10}$  heterocycloalkyl group, a  $C_3-C_{10}$  cycloalkenyl group, a  $C_2-C_{10}$  heterocycloalkenyl group, a  $C_6-C_{60}$  aryl group, a  $C_6-C_{60}$  aryloxy group, a  $C_6-C_{60}$  arylthio group, a  $C_2-C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, each substituted with at least one of a deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1-C_{60}$  alkyl group, a  $C_2-C_{60}$  alkenyl group, a  $C_2-C_{60}$  alkynyl group, a  $C_1-C_{60}$  alkoxy group, a  $C_3-C_{10}$  cycloalkyl group, a  $C_2-C_{10}$  heterocycloalkyl group, a  $C_3-C_{10}$  cycloalkenyl group, a  $C_2-C_{10}$  heterocycloalkenyl group, a  $C_6-C_{60}$  aryl group, a  $C_6-C_{60}$  aryloxy group, a  $C_6-C_{60}$  arylthio group, a  $C_2-C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group,  $-N(Q_{21})(Q_{22})$ ,  $-Si(Q_{23})(Q_{24})(Q_{25})$ , and  $-B(Q_{26})(Q_{27})$ ; and

**[0027]**  $-N(Q_{31})(Q_{32})$ ,  $-Si(Q_{33})(Q_{34})(Q_{35})$  and  $-B(Q_{36})(Q_{37})$ ;

**[0028]** wherein  $Q_1$  to  $Q_7$ ,  $Q_{11}$  to  $Q_{17}$ ,  $Q_{21}$  to  $Q_{27}$ , and  $Q_{31}$  to  $Q_{37}$  may be each independently selected from a hydrogen, a  $C_1-C_{60}$  alkyl group, a  $C_2-C_{60}$  alkenyl group, a  $C_2-C_{60}$  alkynyl group, a  $C_1-C_{60}$  alkoxy group, a  $C_3-C_{10}$  cycloalkyl group, a  $C_2-C_{10}$  heterocycloalkyl group, a  $C_3-C_{10}$  cycloalkenyl group, a  $C_2-C_{10}$  heterocycloalkenyl group, a  $C_6-C_{60}$  aryl group, a  $C_6-C_{60}$  aryloxy group, a  $C_6-C_{60}$  arylthio group, a  $C_2-C_{60}$  heteroaryl group, monovalent a non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group.


**[0029]** According to one or more other embodiments, provided is a condensed-cyclic compound represented by Formula 1A:

Formula 1A



[0030] wherein in Formula 1A,

[0031] ring A<sub>1</sub> is represented by Formula 1D;




Formula 1D


[0032] X<sub>1</sub> to X<sub>3</sub> is N;

[0033] X<sub>4</sub> is O or S;


[0034] L<sub>4</sub> and L<sub>21</sub> are each independently selected from groups represented by Formulae 2-1, 2-2 and 2-34;



Formula 2-1



Formula 2-2



Formula 2-34

[0035] wherein in Formulae 2-1, 2-2 and 2-34, Z<sub>1</sub> is selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group and a phenyl group and d<sub>1</sub> is selected from integers of 1 to 4;

[0036] a<sub>4</sub> is selected from integers 0 to 3 and a<sub>21</sub> is 0, 1, 2 or 3;

[0037] R<sub>5</sub>, R<sub>6</sub>, R<sub>15</sub> and R<sub>17</sub> are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

[0038] R<sub>4</sub> is selected from:

[0039] a phenyl group; and

[0040] a phenyl group, substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, and a biphenyl group; and

[0041] b<sub>4</sub> to b<sub>6</sub>, b<sub>15</sub> and b<sub>17</sub> are each independently selected from integers of 1 to 3.

[0042] According to one or more other embodiments, provided is an organic light-emitting device including

[0043] a first electrode;

[0044] a second electrode; and

[0045] an organic layer disposed between the first electrode and the second electrode,

[0046] wherein the organic layer includes an emission layer and at least one condensed-cyclic compound described above.

[0047] The condensed-cyclic compound may be included in the emission layer, wherein the emission layer further includes a dopant, and the condensed-cyclic compound included in the emission layer may act as a host.

#### BRIEF DESCRIPTION OF THE DRAWING

[0048] These and/or other aspects will become apparent and more readily appreciated from the following description of the embodiments, taken in conjunction with the accompanying drawing in which:

[0049] the FIGURE is a schematic view showing an organic light-emitting device according to an embodiment.

#### DETAILED DESCRIPTION

[0050] Reference will now be made in detail to embodiments, examples of which are illustrated in the accompanying drawings, wherein like reference numerals refer to the like elements throughout. In this regard, the present embodiments may have different forms and should not be construed as being limited to the descriptions set forth herein. Accordingly, the embodiments are merely described below, by referring to the FIGURES, to explain aspects of the present description. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

[0051] It will be understood that when an element is referred to as being “on” another element, it can be directly in contact with the other element or intervening elements may be present therebetween. In contrast, when an element is referred to as being “directly on” another element, there are no intervening elements present.

[0052] It will be understood that, although the terms first, second, third etc. may be used herein to describe various elements, components, regions, layers, and/or sections, these elements, components, regions, layers, and/or sections should not be limited by these terms. These terms are only used to distinguish one element, component, region, layer, or section from another element, component, region, layer, or section. Thus, a first element, component, region, layer, or section discussed below could be termed a second element, component, region, layer, or section without departing from the teachings of the present embodiments.

[0053] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. As used herein, the singular forms “a,” “an,” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.





[0054] The term “or” means “and/or.” It will be further understood that the terms “comprises” and/or “comprising,” or “includes” and/or “including” when used in this specification, specify the presence of stated features, regions, integers, steps, operations, elements, and/or components, but do not preclude the presence or addition of one or more other features, regions, integers, steps, operations, elements, components, and/or groups thereof.

[0055] Unless otherwise defined, all terms (including technical and scientific terms) used herein have the same meaning as commonly understood by one of ordinary skill in the


art to which this general inventive concept belongs. It will be further understood that terms, such as those defined in commonly used dictionaries, should be interpreted as having a meaning that is consistent with their meaning in the context of the relevant art and the present disclosure, and will not be interpreted in an idealized or overly formal sense unless expressly so defined herein.

**[0056]** Exemplary embodiments are described herein with reference to cross section illustrations that are schematic illustrations of idealized embodiments. As such, variations from the shapes of the illustrations as a result, for example, of manufacturing techniques and/or tolerances, are to be expected. Thus, embodiments described herein should not be construed as limited to the particular shapes of regions as illustrated herein but are to include deviations in shapes that result, for example, from manufacturing. For example, a region illustrated or described as flat may, typically, have rough and/or nonlinear features. Moreover, sharp angles that are illustrated may be rounded. Thus, the regions illustrated in the figures are schematic in nature and their shapes are not intended to illustrate the precise shape of a region and are not intended to limit the scope of the present claims.



**[0057]** A condensed-cyclic compound according to an embodiment may be represented by Formula 1A or 1B:



**[0058]** ring A<sub>1</sub> in Formula 1A and 1B may be represented by Formula 1C or 1D.




**[0059]** In Formulae 1A and 1B, ring A<sub>1</sub> may be fused with an adjacent 5-membered cyclic ring by sharing the carbon atoms disposed therebetween. Accordingly, the condensed-cyclic compound represented by Formula 1A or 1B may be represented by any one of Formulae 1A-1 to 1A-7 and 1B-1 to 1B-4:




-continued


Formula 1A-5




Formula 1A-6




Formula 1A-7




Formula 1B-1




Formula 1B-2



-continued



Formula 1B-3



Formula 1B-4


[0060] Descriptions of Formulae 1A-1 to 1A-7 and 1B-1 to 1B-4, X<sub>1</sub> to X<sub>4</sub>, X<sub>11</sub>, L<sub>1</sub> to L<sub>4</sub>, L<sub>11</sub>, L<sub>12</sub>, L<sub>21</sub>, a<sub>1</sub> to a<sub>4</sub>, a<sub>11</sub>, a<sub>12</sub>, a<sub>21</sub>, R<sub>1</sub> to R<sub>6</sub>, R<sub>11</sub> to R<sub>17</sub>, b<sub>1</sub> to b<sub>6</sub>, and b<sub>11</sub> to b<sub>17</sub> are given below.

[0061] According to an embodiment, the condensed-cyclic compound may be represented by Formula 1A-1, 1A-2, 1A-3, 1B-1, 1B-2, 1B-3, or 1B-4, but it is not limited thereto.


[0062] According to another embodiment, the condensed-cyclic compound may be represented by Formula 1A-1 or 1A-2, but it is not limited thereto.

[0063] Formulae 1A and 1B may be represented by any one of Formulae 1A(1) to 1A(4) and 1B(1) to 1B(4), but they are not limited thereto:


-continued



Formula 1A(2)




Formula 1A(1)







Formula 1A(3)

-continued



-continued



**[0064]** In Formulae 1A(1) to 1A(4) and 1B(1) to 1B(4), descriptions of ring A<sub>1</sub>, X<sub>1</sub> to X<sub>4</sub>, X<sub>11</sub>, L<sub>1</sub> to L<sub>4</sub>, L<sub>11</sub>, L<sub>12</sub>, L<sub>21</sub>, a<sub>1</sub> to a<sub>4</sub>, a<sub>11</sub>, a<sub>12</sub>, a<sub>21</sub>, R<sub>1</sub> to R<sub>6</sub>, R<sub>11</sub> to R<sub>17</sub>, b<sub>1</sub> to b<sub>6</sub>, and b<sub>11</sub> to b<sub>17</sub> are given below.

**[0065]** In Formula 1C, X<sub>11</sub> is selected from N-[(L<sub>12</sub>)<sub>a12</sub>-(R<sub>12</sub>)<sub>b12</sub>], S, O, S(=O), S(=O)<sub>2</sub>, C(=O), C(R<sub>13</sub>)(R<sub>14</sub>), Si(R<sub>13</sub>)(R<sub>14</sub>), P(R<sub>13</sub>), P(=O)(R<sub>13</sub>), and C=N(R<sub>12</sub>). For example, in Formula 1C, X<sub>11</sub> may be selected from N-[(L<sub>12</sub>)<sub>a12</sub>-(R<sub>12</sub>)<sub>b12</sub>], S, O, and C(R<sub>13</sub>)(R<sub>14</sub>), but it is not limited thereto. Descriptions of L<sub>12</sub>, a<sub>12</sub>, R<sub>12</sub> to R<sub>14</sub>, and b<sub>12</sub> may be understood by referring to the description below.

**[0066]** In Formula 1A and 1B, X<sub>1</sub> is N or C-[(L<sub>1</sub>)<sub>a1</sub>-(R<sub>1</sub>)<sub>b1</sub>], X<sub>2</sub> is N or C-[(L<sub>2</sub>)<sub>a2</sub>-(R<sub>2</sub>)<sub>b2</sub>], X<sub>3</sub> is N or C-[(L<sub>3</sub>)<sub>a3</sub>-(R<sub>3</sub>)<sub>b3</sub>], and at least one of X<sub>1</sub> to X<sub>3</sub> is N.

**[0067]** For example, in Formulae 1A and 1B,

**[0068]** X<sub>1</sub> to X<sub>3</sub> may be N;

**[0069]** X<sub>1</sub> may be C-[(L<sub>1</sub>)<sub>a1</sub>-(R<sub>1</sub>)<sub>b1</sub>], X<sub>2</sub> and X<sub>3</sub> may be N;

**[0070]** X<sub>1</sub> may be N, X<sub>2</sub> may be C-[(L<sub>2</sub>)<sub>a2</sub>-(R<sub>2</sub>)<sub>b2</sub>], and X<sub>3</sub> may be N;

**[0071]** X<sub>1</sub> and X<sub>2</sub> may be N, and X<sub>3</sub> may be C-[(L<sub>3</sub>)<sub>a3</sub>-(R<sub>3</sub>)<sub>b3</sub>];

**[0072]** X<sub>1</sub> may be C-[(L<sub>1</sub>)<sub>a1</sub>-(R<sub>1</sub>)<sub>b1</sub>], X<sub>2</sub> may be N, and X<sub>3</sub> may be C-[(L<sub>3</sub>)<sub>a3</sub>-(R<sub>3</sub>)<sub>b3</sub>];

**[0073]** X<sub>1</sub> may be C-[(L<sub>1</sub>)<sub>a1</sub>-(R<sub>1</sub>)<sub>b1</sub>], X<sub>2</sub> may be C-[(L<sub>2</sub>)<sub>a2</sub>-(R<sub>2</sub>)<sub>b2</sub>], and X<sub>3</sub> may be N; or

**[0074]** X<sub>1</sub> may be N, X<sub>2</sub> may be C-[(L<sub>2</sub>)<sub>a2</sub>-(R<sub>2</sub>)<sub>b2</sub>], and X<sub>3</sub> may be C-[(L<sub>3</sub>)<sub>a3</sub>-(R<sub>3</sub>)<sub>b3</sub>]

**[0075]** According to an embodiment,

**[0076]** X<sub>1</sub> may be C-[(L<sub>1</sub>)<sub>a1</sub>-(R<sub>1</sub>)<sub>b1</sub>], X<sub>2</sub> and X<sub>3</sub> may be N; or

**[0077]** X<sub>1</sub> and X<sub>2</sub> may be N, and X<sub>3</sub> may be C-[(L<sub>3</sub>)<sub>a3</sub>-(R<sub>3</sub>)<sub>b3</sub>], but they are not limited thereto.

**[0078]** Descriptions of L<sub>1</sub> to L<sub>3</sub>, a<sub>1</sub> to a<sub>3</sub>, R<sub>1</sub> to R<sub>3</sub>, and b<sub>1</sub> to b<sub>3</sub> may be understood by referring to the description below.

**[0079]** In Formulae 1A, 1B, and 1C, L<sub>1</sub> to L<sub>4</sub>, L<sub>11</sub>, L<sub>12</sub>, and L<sub>21</sub> may be each independently selected from a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkylene group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenylene group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylene group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic hetero-condensed polycyclic group.

**[0080]** For example, in Formulae 1A, 1B, and 1C,  $L_1$  to  $L_4$ ,  $L_{11}$ , Lie, and  $L_{21}$  may be each independently selected from:


**[0081]** a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pyrrolylene group, an imidazolylene group, a pyrazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzooxazolylene group, a benzimidazolylene group, a furanylene group, a benzofuranylene group, a thiophenylene group, a benzothiophenylene group, a thiazolylene group, a isothiazolylene group, a benzothiazolylene group, an isoxazolylene group, an oxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, an imidazopyrimidinylene group, and an imidazopyridinylene group; and

**[0082]** a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pyrrolylene group, an imidazolylene group, a pyrazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzooxazolylene group, a benzimidazolylene group, a furanylene group, a benzofuranylene group, a thiophenylene group, a benzothiophenylene group, a thiazolylene group, a isothiazolylene group, a benzothiazolylene group, an isoxazolylene group, an oxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, an imidazopyrimidinylene group, and an imidazopyridinylene group, each substituted with at least one of a deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a


hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a  $C_6$ - $C_{20}$  aryl group, a  $C_2$ - $C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed cyclic group, and  $-Si(Q_{33})(Q_{34})(Q_{35})$ ,

**[0083]**  $Q_{33}$  to  $Q_{35}$  may be each independently selected from a hydrogen, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a phthalazinyl group, a quinoxalinyl group, a cinnolinyl group, and a quinazolinyl group, but they are not limited thereto.


**[0084]** For example, in Formulae 1A, 1B, and 1C,  $L_1$  to  $L_4$ ,  $L_{11}$ , Lie, and  $L_{21}$  may be each independently selected from Formulae 2-1 to 2-34:




Formula 2-1




Formula 2-2




Formula 2-3



Formula 2-4













Formula 2-5




Formula 2-6


-continued




Formula 2-7




Formula 2-8




Formula 2-9




Formula 2-10




Formula 2-11




Formula 2-12




Formula 2-13




Formula 2-14



Formula 2-15



Formula 2-16



-continued

Formula 2-17

Formula 2-18

Formula 2-19

Formula 2-20

Formula 2-21

Formula 2-22


Formula 2-23

Formula 2-24


Formula 2-25

Formula 2-26

-continued




Formula 2-27




-continued


Formula 2-34




Formula 2-28




Formula 2-29




Formula 2-30



Formula 2-31



Formula 2-32



Formula 2-33

[0085] In Formula 2-1 to 2-34,

[0086]  $Y_1$  may be selected from O, S, S(=O), S(=O)<sub>2</sub>, C(Z<sub>3</sub>)(Z<sub>4</sub>), N(Z<sub>5</sub>), or Si(Z<sub>6</sub>)(Z<sub>7</sub>);

[0087] Z<sub>1</sub> to Z<sub>7</sub> may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a quinoxaliny group, and —Si(Q<sub>33</sub>)(Q<sub>34</sub>)(Q<sub>35</sub>);

[0088] wherein Q<sub>33</sub> to Q<sub>35</sub> may be each independently selected from a hydrogen, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, and a quinoxaliny group;

[0089] d<sub>1</sub> is selected from integers of 1 to 4,

[0090] d<sub>2</sub> is selected from integers of 1 to 3,

[0091] d<sub>3</sub> is selected from integers of 1 to 6,

[0092] d<sub>4</sub> is selected from integers of 1 to 8,

[0093] d<sub>5</sub> is 1 or 2,

[0094] d<sub>6</sub> is selected from integers of 1 to 5, and

[0095] each of \* and \*' indicates a bonding site to neighboring atoms.

[0096] For example, in Formulae 2-1 to 2-34, \*' indicates a bonding site to neighboring atoms of L<sub>1</sub> to L<sub>4</sub>, L<sub>11</sub>, L<sub>12</sub>, and L<sub>21</sub> or bonding site to each of R<sub>1</sub> to R<sub>4</sub>, R<sub>11</sub>, R<sub>12</sub>, and R<sub>21</sub>.

[0097] According to an embodiment, in Formulae 1A and 1B, L<sub>1</sub> to L<sub>4</sub>, L<sub>11</sub>, L<sub>12</sub>, and L<sub>21</sub> may be each independently selected from Formulae 2-1 to 2-5, 2-9 to 2-23, and 2-34, but they are not limited thereto.

[0098] According to another embodiment, in Formulae 1A and 1B, L<sub>1</sub> to L<sub>4</sub>, L<sub>11</sub>, L<sub>12</sub>, and L<sub>21</sub> may be each independently selected from:

[0099] a phenylene group and a naphthylene group; and

[0100] a phenylene group and a naphthylene group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt

thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, a quinoxalinyl group, a biphenyl group, and —Si(Q<sub>33</sub>)(Q<sub>34</sub>)(Q<sub>35</sub>) (wherein Q<sub>33</sub> to Q<sub>35</sub> may be each independently selected from a hydrogen, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, and a quinoxalinyl group, but they are not limited thereto.

**[0101]** In Formula 1A, a<sub>1</sub> represents the number of groups L<sub>1</sub>, which may be 0, 1, 2, or 3, for example, 0, 1, or 2, or may be 0 or 1. When a<sub>1</sub> is 0, -(L<sub>1</sub>)<sub>a<sub>1</sub></sub>- is a single bond. When a<sub>1</sub> is 2 or greater, two or more groups L<sub>1</sub> may be the same or different. Descriptions of a<sub>2</sub> to a<sub>4</sub>, a<sub>11</sub>, a<sub>12</sub>, and a<sub>21</sub> may be understood by referring to the description of a<sub>1</sub> and structures of Formulae 1A and 1B.

**[0102]** According to an embodiment, in Formulae above, a<sub>1</sub> to a<sub>4</sub>, a<sub>11</sub>, a<sub>12</sub>, and a<sub>21</sub> may be each independently, 0, 1, or 2.

**[0103]** According to another embodiment, in Formulae above, a<sub>21</sub> may be 1, but it is not limited thereto.

**[0104]** In Formulae 1A, 1B, 1C, and 1D, R<sub>1</sub> to R<sub>3</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>11</sub> to R<sub>17</sub> may be each independently selected from a hydrogen, a deuterium, —F (a fluoro group), —Cl (a chloro group), —Br (a bromo group), —I (an iodo group), a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, —N(Q<sub>1</sub>)(Q<sub>2</sub>), —Si(Q<sub>3</sub>)(Q<sub>4</sub>)(Q<sub>5</sub>), and —B(Q<sub>6</sub>)(Q<sub>7</sub>). Here, descriptions of Q<sub>1</sub> to Q<sub>7</sub> are as described below.

**[0105]** In Formulae 1A and 1B, R<sub>4</sub> is selected from a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or

unsubstituted C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group. For example, R<sub>4</sub> may be selected from a substituted or unsubstituted C<sub>6</sub>-C<sub>20</sub> aryl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>30</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, but it is not limited thereto.

**[0106]** According to an embodiment, in Formulae 1A and 1B, R<sub>11</sub> and R<sub>17</sub> may be each independently selected from a substituted or unsubstituted C<sub>6</sub>-C<sub>30</sub> aryl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>30</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group.

**[0107]** For example, in Formulae 1A, 1B, 1C, and 1D, R<sub>1</sub> to R<sub>3</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>13</sub> to R<sub>17</sub> may be each independently selected from:

**[0108]** a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

**[0109]** a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof and a phosphoric acid or a salt thereof;

**[0110]** a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthroline group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

**[0111]** a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluore-

nyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranly group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranly group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, —Si(Q<sub>33</sub>)(Q<sub>34</sub>)(Q<sub>35</sub>), a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranly group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranly group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and a biphenyl group; and

**[0112]** —Si(Q<sub>3</sub>)(Q<sub>4</sub>)(Q<sub>5</sub>) (provided that, R<sub>13</sub> and R<sub>14</sub> are not —Si(Q<sub>3</sub>)(Q<sub>4</sub>)(Q<sub>5</sub>));

**[0113]** wherein Q<sub>3</sub> to Q<sub>5</sub> and Q<sub>33</sub> to Q<sub>35</sub> may be each independently selected from a hydrogen, C<sub>1</sub>-C<sub>20</sub> alkyl group, C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a naphthyl

group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranly group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, and a quinoxalinyl group, but they are not limited thereto.

**[0114]** According to another embodiment, in Formulae 1A, 1B, 1C, and 1D, R<sub>1</sub> to R<sub>3</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>13</sub> to R<sub>17</sub> may be each independently selected from:

**[0115]** a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

**[0116]** a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, and a phosphoric acid or a salt thereof;

**[0117]** a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranly group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, and a quinoxalinyl group;

**[0118]** a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranly group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, and a quinoxalinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, —Si(Q<sub>33</sub>)(Q<sub>34</sub>)(Q<sub>35</sub>), a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranly group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, and a quinoxalinyl group; and


**[0119]** —Si(Q<sub>3</sub>)(Q<sub>4</sub>)(Q<sub>5</sub>), provided that R<sub>13</sub> and R<sub>14</sub> are not —Si(Q<sub>3</sub>)(Q<sub>4</sub>)(Q<sub>5</sub>);

**[0120]** wherein Q<sub>3</sub> to Q<sub>5</sub> and Q<sub>33</sub> to Q<sub>35</sub> may be each independently selected from a hydrogen, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranly group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl




[0131] wherein,  $R_4$ ,  $R_{11}$ , and  $R_{17}$  may be each independently selected from Formulae 4-1 to 4-31 (for example, Formulae 4-1 to 4-3 and 4-6 to 4-13), but they are not limited thereto.


-continued




Formula 4-1




Formula 4-2




Formula 4-3




Formula 4-4




Formula 4-5




Formula 4-6




Formula 4-7




Formula 4-8




Formula 4-9




Formula 4-10




Formula 4-11




Formula 4-12




Formula 4-13




Formula 4-14




Formula 4-15




Formula 4-16




Formula 4-17



Formula 4-18




Formula 4-19




Formula 4-20


-continued




Formula 4-21




Formula 4-22




Formula 4-23




Formula 4-24




Formula 4-25




Formula 4-26



Formula 4-27




Formula 4-28




Formula 4-29

-continued

Formula 4-30



Formula 4-31



[0132] In Formulae 4-1 to 4-31,

[0133]  $Y_{31}$  may be O, S, C( $Z_{33}$ )( $Z_{34}$ ), N( $Z_{35}$ ) or Si( $Z_{36}$ )( $Z_{37}$ ),[0134]  $Z_{31}$  to  $Z_{37}$  may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof,  $C_1$ - $C_{20}$  alkyl group,  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a quinoxaliny group, a biphenyl group, and —Si( $Q_{33}$ )( $Q_{34}$ )( $Q_{35}$ );[0135]  $Q_3$  to  $Q_5$  and  $Q_{33}$  to  $Q_{35}$  may be each independently selected from a hydrogen, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, and a quinoxaliny group;

[0136] e1 may be selected from integers of 1 to 5,

[0137] e2 is selected from integers of 1 to 7,

[0138] e3 is selected from integers of 1 to 3,

[0139] e4 is selected from integers of 1 to 4,

[0140] e5 is 1 or 2,

[0141] e6 is selected from integers of 1 to 6, and

[0142] \* indicates a bonding site to a neighboring atom.

[0143] According to another embodiment,  $R_1$  to  $R_3$ ,  $R_5$ ,  $R_6$ , and  $R_{13}$  to  $R_{17}$  may be each independently selected from:[0144] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group;[0145] a  $C_1$ - $C_{20}$  alkyl group and a  $C_1$ - $C_{20}$  alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group,

an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, and a phosphoric acid or a salt thereof;

[0146] a phenyl group and a naphthyl group;

[0147] a phenyl group and a naphthyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, and a naphthyl group; and

[0148] —Si(Q<sub>3</sub>)(Q<sub>4</sub>)(Q<sub>5</sub>) (provided that, R<sub>13</sub> and R<sub>14</sub> are not —Si(Q<sub>3</sub>)(Q<sub>4</sub>)(Q<sub>5</sub>) and Q<sub>3</sub> to Q<sub>5</sub> are each independently selected from a hydrogen, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, and a naphthyl group);

[0149] R<sub>4</sub>, R<sub>11</sub>, and R<sub>12</sub> are each independently selected from:

[0150] a phenyl group and a naphthyl group; and

[0151] a phenyl group and a naphthyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, and a naphthyl group; but, they are not limited thereto.

[0152] According to another embodiment, R<sub>5</sub>, R<sub>6</sub>, and R<sub>15</sub> to R<sub>17</sub> may all be hydrogen, but they are not limited thereto.

[0153] In Formulae 1A and 1B, b1 represents the number of groups R<sub>1</sub> and may be selected from integers of 1 to 3. For example, b1 may be 1 or 2. In greater detail, b1 may be 1. When b1 is 2 or greater, two or more of groups R<sub>1</sub> may be the same or different. Descriptions of b2 to b6 and b11 to b17 may be understood by referring to the description of b1 and structures of Formulae 1A, 1B, 1C, and 1D.

[0154] According to an embodiment, at least one substituent of the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkylene group, substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenylene group, substituted C<sub>6</sub>-C<sub>60</sub> arylene group, substituted C<sub>2</sub>-C<sub>60</sub> heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic hetero-condensed polycyclic group, substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, substituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, substituted C<sub>6</sub>-C<sub>60</sub> aryl group, substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, substituted C<sub>2</sub>-C<sub>60</sub> heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic hetero-condensed polycyclic group may be selected from:

[0155] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

[0156] a C<sub>1</sub>-C<sub>60</sub> alkyl group, C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q<sub>11</sub>)(Q<sub>12</sub>), —Si(Q<sub>13</sub>)(Q<sub>14</sub>)(Q<sub>15</sub>) and —B(Q<sub>16</sub>)(Q<sub>17</sub>);


[0157] a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group;

[0158] a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q<sub>21</sub>)(Q<sub>22</sub>), —Si(Q<sub>23</sub>)(Q<sub>24</sub>)(Q<sub>25</sub>), and —B(Q<sub>26</sub>)(Q<sub>27</sub>); and

[0159] —N(Q<sub>31</sub>)(Q<sub>32</sub>), —Si(Q<sub>33</sub>)(Q<sub>34</sub>)(Q<sub>35</sub>) and —B(Q<sub>36</sub>)(Q<sub>37</sub>);


[0160] wherein Q<sub>1</sub> to Q<sub>7</sub>, Q<sub>11</sub> to Q<sub>17</sub>, Q<sub>21</sub> to Q<sub>27</sub> and Q<sub>31</sub> to Q<sub>37</sub> may be each independently selected from a hydrogen, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group.

[0161] According to one or more embodiments, provided is a condensed-cyclic compound represented by Formula 1A:



[0162] wherein in Formula 1A,




[0163] ring A<sub>1</sub> is represented by Formula 1D;



[0164] X<sub>1</sub> to X<sub>3</sub> is N;

[0165] X<sub>4</sub> is O or S;

[0166] L<sub>4</sub> and L<sub>21</sub> are each independently selected from groups represented by Formulae 2-1, 2-2 and 2-34;



[0167] wherein in Formulae 2-1, 2-2 and 2-34, Z<sub>1</sub> is selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group and a phenyl group and d<sub>1</sub> is selected from integers of 1 to 4;

[0168] a<sub>4</sub> is selected from integers 0 to 3 and a<sub>21</sub> is 0, 1, 2 or 3 (or a<sub>21</sub> may be 1 or 2);

[0169] R<sub>5</sub>, R<sub>6</sub>, R<sub>15</sub> and R<sub>17</sub> are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or

a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

[0170] R<sub>4</sub> is selected from:




[0171] a phenyl group; and

[0172] a phenyl group, substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, and a biphenyl group; and

[0173] b<sub>4</sub> to b<sub>6</sub>, b<sub>15</sub> and b<sub>17</sub> are each independently selected from integers of 1 to 3.


[0174] For example, a condensed-cyclic compound represented by Formula 1A described above may be represented by one of Formulae 1A(1) to 1A(4) described herein.

[0175] The condensed-cyclic compound may be any one of Compounds 1 to 825, but it is not limited thereto:




-continued


4




5



6




7




-continued


8




9



10




11




-continued


12




13



15




15




-continued


16



17




18




20

-continued





19

-continued




22

20





23



24


21




25

-continued


26




27



28




29




-continued


30




31



32









33




-continued

-continued




-continued


40




41



42




43




-continued


44







45







46



-continued




-continued




-continued


55




56



57




58




-continued


59




60






61







62




-continued




-continued




-continued




70



71




72




73


-continued




74



75




76




77

-continued




78


-continued




81




79



82




80




83

-continued


-continued




84




87




85



88









86




89

-continued

-continued




-continued




96


-continued




99




97



100




98




101

-continued




102


-continued




105




103



106




104




107

-continued




108


-continued




111




109



112




110




113

-continued




114


-continued




117




115



118




116




119

-continued




119


-continued




122




120



123




121




124

-continued




125


-continued




128




126



129




127




130


-continued




-continued




132




135



133




136




-continued

137




-continued


140




138




141



139









142




-continued

-continued




-continued




149


-continued




152




150



153









151



154







-continued

-continued




-continued

-continued




-continued


167




168



169




170




-continued


171




172






173







174




-continued



-continued




-continued




182

-continued




185


183




186



184




187




-continued

188




-continued


191




189




193



190




192




-continued


194




195



196




197




-continued


198




199



200




201




-continued

202




-continued


205




203




206



204




207




-continued


208




209



210




211




-continued


212



213




214




-continued


215




216



217




218




-continued


219




220



221




222




-continued

223




-continued


227




224




228




225




229






226







230



-continued



-continued





-continued

-continued



242

238





243

239





244

240




245

241




-continued




246


-continued




249




247



250







248







251


-continued




-continued




-continued




-continued




261




265




262




266




263




-continued




-continued




268




271



269




272




-continued


273




274



275




276




-continued


277



278




279




-continued


280




281



282




283




-continued


284




285




286




287




-continued



288




289




290


-continued




291



292




293




294

-continued




295


-continued




298




296



299










297




300

-continued


-continued




-continued




308



309




310




311


-continued




312



313






314







315

-continued




-continued

319




-continued


323




324



325




326




-continued


327




328








329



-continued




-continued




-continued

-continued


336




339




337




340





338




341




-continued




343




346



344




347




-continued

348




-continued


351




349




352




350



353




-continued




354


-continued




357




355



358




356




359

-continued


-continued




360




363




361



364









362




365

-continued






-continued






-continued






-continued



-continued




-continued




-continued

-continued


384




387




385




388



386




389




-continued

390




-continued


393




391




394



392









395









-continued

-continued




-continued

-continued




-continued

408




-continued


411




409




412




410



413




-continued




414


-continued




417




415



418




416




419

-continued


420




421



422




423




-continued


424




425



426




427




-continued

428




-continued

432




433

429





434

430




435

431




-continued

436




-continued


439




437




440



438




441




-continued

442




-continued


445




443




446



444




447




-continued


448




449



450




451




-continued


452




453




454




455




-continued



456




457




458


-continued




459



460




461




462

-continued


463



464




465




-continued


466



467




468




-continued


469




470



471




472




-continued


473







474







475






-continued







-continued



-continued




-continued




-continued


491




492



493




494




-continued


495




496



497




-continued




498


-continued




501




499



502




500




503

-continued


504




505



506




507




-continued


508




509



510







511




-continued

-continued




-continued

516




-continued


519




520




517




521



518




522




-continued


523



524




525




-continued

526




527




-continued

528




-continued


531




532




529




533






530





534



-continued




-continued




-continued

540




-continued


543




544




541



545




542




-continued

546




-continued

550




551


547




552




548



549










553











-continued

-continued



-continued


-continued




-continued

-continued


569




572




573




570



574




571




-continued

575




-continued


578




576




579




577



580




-continued




581


-continued




584




582



585




583




586

-continued


587




588



589




590




-continued


591



592




593




-continued


594



595




596




-continued


597



598




599




-continued

600




-continued


603




601




604




602



605




-continued




606


-continued




609




607



610




608




611


-continued




-continued




613




616




614




617




-continued




-continued




621




619



622




620




623


-continued




-continued




625




628



626




629





-continued

-continued




630

633





631

634




632

635




-continued

636




-continued


639




637




640




638




641




-continued




-continued



643





646




647

644




-continued




648


-continued




651




649



652




650




653

-continued




-continued




655

657





656




658



659




-continued




-continued


663




664




665




666



-continued




-continued




669

667


670



668










671




-continued


-continued




-continued




-continued



680



682





-continued

-continued


686

683





687


684




688




685



689




-continued




690


-continued



692




691



693


-continued

694




-continued

697





698

695




699

696




-continued

700




-continued


704




705




701




706




702



703




707




-continued

-continued

708





711



712





709




713

710




714




-continued

715




-continued


719



716





720





721

717



722


718




-continued

-continued


723




726



727





724









728


725



-continued




-continued




735


-continued




738




736



739




737




740


-continued




741



742




743




744


-continued



745




746



747


-continued

748




-continued


751




749




752



750




753




-continued

754





-continued

757




758


755




756



759




-continued




760


-continued




763




761



764




762



765


-continued

766




-continued


769




767




770




768








771




-continued



-continued




-continued




778


-continued




781




779



782




780



783


-continued

784




-continued


787




785




788




786



789




-continued




790


-continued




793




791



794




792




795

-continued




796


-continued




799




797



800




798




801



-continued




-continued




803




804



807




-continued




808


-continued




811




809



812




810




813

-continued


814




815




816



817




818




-continued


819




820




821



822




823




-continued

824




825




-continued


169



253




337




[0176] Alternatively, the condensed-cyclic compound may be any one of Compounds 1, 85, 169, 253, 337, 421, 505, 506, 575, 659, 660, 729 and 825, but it is not limited thereto:


1



85



421




-continued


-continued

660


505





506




575




659



729



825



[0177] In the condensed-cyclic compound represented by Formula 1A or 1B,  $R_4$  is selected from a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_2$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group.

[0178] In other words, in Formula 1A and 1B,  $R_4$  necessarily includes a ring structure. As a result, the condensed-cyclic compound represented by Formula 1A or 1B is chemically and structurally stable and may actually have a spherical molecular structure. Accordingly, the condensed-cyclic compound represented by Formula 1A or 1B may have excellent thermal stability, which may increase deposition temperature. As a result, efficiency and lifespan of an organic light-emitting device including the condensed-cyclic compound may be improved to improve formability of the organic light-emitting device during the manufacturing process thereof.

[0179] Also, because both group "A" and the "carbazole-based" group in Formulae 1A and 1B are bound to a "nitrogen-containing 6-membered ring" (see Formulae 1A' and 1B'), hole injection and hole transport and electron injection and electron transport may occur thoroughly and Formulae 1A and 1B may each actually have a spherical molecular structure. Accordingly, the condensed-cyclic compound may simultaneously have excellent charge-transporting ability and thermal stability, such that the organic light-emitting device including the condensed-cyclic compound may have increased emission efficiency, reduced driving voltage, and a long lifespan.

[0182] a first electrode;



[0183] a second electrode; and

[0184] an organic layer disposed between the first electrode and the second electrode,

[0185] wherein the organic layer includes the EML, which includes at least one condensed-cyclic compound represented by Formula 1A or 1B.

[0186] The organic light-emitting device including an organic layer including the condensed-cyclic compound represented by Formula 1A or 1B has low driving voltage, high efficiency, high brightness, and a long lifespan.

[0187] The condensed-cyclic compound represented by Formula 1A or 1B may be used between a pair of electrodes in the organic light-emitting device. For example, the condensed-cyclic compound may be included in at least one of an EML, a hole-transport region disposed between the first electrode and the EML (for example, the hole transport region may include at least one of a hole-injecting layer (HIL), a hole-transporting layer (HTL), and an electron-blocking layer (EBL)), and an electron-transport region disposed between the EML and the second electrode (for example, the electron transport region may include at least one of a hole-blocking layer (HBL), an electron-transporting layer (ETL), and an electron-injecting layer (EIL)). For example, the condensed-cyclic compound represented by



[0180] A method of synthesizing the condensed-cyclic compound represented by Formula 1A or 1B may be understood by one of ordinary skill in the art by referring to the embodiments described below.

[0181] Accordingly, the condensed-cyclic compound represented by Formula 1A or 1B may be suitable as a material for an organic layer (for example, a host of an EML) in an organic light-emitting device. According to another embodiment, provided is an organic light-emitting device including

Formula 1A or 1B may be included in the EML. In this regard, the EML further includes a dopant and the condensed-cyclic compound included in the EML may act as a host. The EML may be a green EML emitting green light and the dopant may be a phosphorescent dopant.

[0188] As used herein, the term "(the organic layer) includes at least one condensed-cyclic compound" may be understood as "(the organic layer) may include at least one condensed-cyclic compound belonging to the group of For-

mula 1A or 1B or two different condensed-cyclic compounds belonging to the group of Formula 1A or 1B”.

[0189] For example, the organic layer may only include Compound 1 as the condensed-cyclic compound. In this regard, Compound 1 may be situated in the EML of the organic light-emitting device. Alternatively, the organic layer may include Compound 1 and Compound 2 as the condensed-cyclic compound. In this regard, Compound 1 and Compound 2 may be present on the same layer (for example, Compound 1 and Compound 2 may all be present on the EML) or on different layers.

[0190] The organic layer includes

[0191] i) a hole transport region that is disposed between the first electrode and the EML and includes at least one of an HIL, an HTL, a buffer layer, and an EBL, and

[0192] ii) an electron transport region that is disposed between the EML and the second electrode and includes at least one layer selected from a HBL, an ETL, and an EIL.

[0193] The expression “organic layer”, as used herein refers to a single layer and/or a plurality of layers disposed between the first and second electrodes of an organic light-emitting device. A material of the “organic layer” is not limited to an organic material and may include an organic metal complex including a metal.

[0194] The FIGURE is a schematic view of an organic light-emitting device 10 according to an embodiment. Hereinafter, a structure and a method of manufacturing the organic light-emitting device according to an embodiment will be described with reference to the FIGURE. The organic light-emitting device 10 includes a first electrode 11, an organic layer 15, and a second electrode 19, which are sequentially stacked in the stated order.

[0195] A substrate may be additionally disposed under the first electrode 11 or on the second electrode 19. The substrate may be a conventional glass substrate or a transparent plastic substrate, each with excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water repellency.

[0196] The first electrode 11 may be formed by depositing or sputtering a material for forming the first electrode 11 on the substrate. When the first electrode 11 is an anode, the material for the first electrode 11 may be selected from materials with a high work function for an easy hole injection. The first electrode 11 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. The material for the first electrode 110 may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO<sub>2</sub>), and zinc oxide (ZnO). Alternatively, a metal such as magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag).

[0197] The first electrode 11 may have a single-layer structure or a multi-layer structure including two or more layers.

[0198] The organic layer 15 is disposed on the first electrode 11.

[0199] The organic layer 15 may include a hole transport region; an EML; and an electron transport region.

[0200] The hole transport region may be disposed between the first electrode 11 and the EML.

[0201] The hole transport region may include at least one of the HIL, HTL, EBL, and buffer layer.

[0202] The hole transport region may only include the HIL or HTL. Alternatively, the hole transport region may have an

HIL/HTL structure or an HIL/HTL/EBL structure, wherein layers of each structure are sequentially stacked on the first electrode 11 in this stated order, but it is not limited thereto.


[0203] When the hole transport region includes an HIL, the HIL may be formed on the first electrode 11 by using various methods, such as vacuum deposition, spin coating, casting, a Langmuir-Blodgett (LB) method, or the like.

[0204] When an HIL is formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of about 100 to about 500° C., at a vacuum degree of about 10<sup>-8</sup> to about 10<sup>-3</sup> torr, and at a deposition rate of about 0.01 Angstrom per second (Å/sec) to about 100 Å/sec in consideration of a compound for an HIL to be deposited, and the structure of an HIL to be formed, but the conditions are not limited thereto.

[0205] When an HIL is formed by spin coating, the spin coating may be performed at a coating rate of about 2,000 revolutions per minute (rpm) to about 5,000 rpm, and at a temperature of about 80° C. to 200° C. for removing a solvent after the spin coating, in consideration of a compound for an HIL to be deposited, and the structure of an HIL to be formed, but the conditions are not limited thereto.


[0206] The conditions for forming the HTL and EBL may be inferred based on the conditions for forming the HIL.

[0207] The hole transport region may include at least one compound selected from m-MTDATA, TDATA, 2-TNATA, NPB, 8-NPB, TPD, Spiro-TPD, Spiro-NPB, methylated NPB, TAPC, HMTPD, 4,4',4"-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), (polyaniline)/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201 below, and a compound represented by Formula 202 below:




m-MTDATA


-continued



TDATA




2-TNATA




NPB


-continued




$\beta$ -NPB




TPD




Spiro-TPD




Spiro-NPB



methylated NPB



Formula 201



**[0208]** In Formula 201, Ar<sub>101</sub> and Ar<sub>102</sub> may be each independently selected from:

**[0209]** a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluore-

nylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, and a pentacenylene group; and

**[0210]** a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, and a pentacenylene group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group.

**[0211]** In Formula 201, xa and xb may be each independently integers of 0 to 5, or 0, 1, or 2. For example, xa may be 1 and xb may be 0, but they are not limited thereto.

**[0212]** In Formulae 201 and 202, R<sub>101</sub> to R<sub>108</sub>, R<sub>111</sub> to R<sub>119</sub>, and R<sub>121</sub> to R<sub>124</sub> may be each independently selected from:

**[0213]** a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>10</sub> alkyl group (for example, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, and a hexyl group) and a C<sub>1</sub>-C<sub>10</sub> alkoxy group (for example, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, and a pentoxy group);

**[0214]** a C<sub>1</sub>-C<sub>10</sub> alkyl group and a C<sub>1</sub>-C<sub>10</sub> alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, and a phosphoric acid group or a salt thereof;

**[0215]** a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, and a pyrenyl group; and


**[0216]** a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, and a pyrenyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>10</sub> alkyl group, and a C<sub>1</sub>-C<sub>10</sub> alkoxy group, but they are not limited thereto.

**[0217]** In Formula 201, R<sub>109</sub> may be any one of a phenyl group, a naphthyl group, an anthracenyl group and a pyridinyl group; a phenyl group, a naphthyl group, an anthracenyl

group, and a pyridinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group.


**[0218]** According to an embodiment, the compound represented by Formula 201 may be represented by Formula 201A, but it is not limited thereto:

Formula 201A



-continued

HT2





**[0219]** In Formula 201A, detailed descriptions of R<sub>101</sub>, R<sub>111</sub>, R<sub>112</sub>, and R<sub>109</sub> may be the same as described herein.

**[0220]** For example, the compound represented by Formula 201 and the compound represented by Formula 202 may include Compounds HT1 to HT20, but the compound is not limited thereto:

HT3

HT1





-continued

-continued



HT4

HT6




HT5

HT7




-continued


HT8



HT9




HT10




-continued


HT11







HT12



HT13




-continued




-continued


HT18



HT19





HT20




[0221] A thickness of the hole transport region may be in a range of about 100 Angstrom ( $\text{\AA}$ ) to about 10,000  $\text{\AA}$ , for example, about 100  $\text{\AA}$  to about 1,000  $\text{\AA}$ . When the hole transport region includes an HIL and a HTL, a thickness of the HIL may be in a range of about 100  $\text{\AA}$  to about 10,000  $\text{\AA}$ , for example, about 100  $\text{\AA}$  to about 1,000  $\text{\AA}$ , and a thickness of the HTL may be in a range of about 50  $\text{\AA}$  to about 2,000  $\text{\AA}$ , for example, about 100  $\text{\AA}$  to about 1,500  $\text{\AA}$ . When the thicknesses of the hole transport region, the HIL, and the HTL are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0222] The hole transport region may further include, in addition to the abovementioned materials, a charge-generating material for the improvement of conductive properties. The charge-generating material may be homogeneously or non-homogeneously dispersed throughout the hole transport region.


[0223] The charge-generating material may be, for example, a p-dopant. The p-dopant may be one of a quinone derivative, a metal oxide, and a cyano group-containing compound, but is not limited thereto. For example, non-limiting examples of the p-dopant are a quinone derivative, such as tetracyanoquinonedimethane (TCNQ) or 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonedimethane (F4-TCNQ); a metal oxide, such as a tungsten oxide or a molybdenum oxide; and Compound HT-D1 illustrated below, but are not limited thereto.




Compound HT-D1




TPBi




TBADN



ADN



CBP



CDBP


[0224] The hole transport region may further include a buffer layer.

[0225] The buffer layer may compensate for an optical resonance distance according to a wavelength of light emitted from the EML, and thus, efficiency of an organic light-emitting device may be improved.

[0226] An EML may be formed on the hole transport region by using various methods, such as vacuum deposition, spin coating, casting, or an LB method. When the EML is formed by vacuum deposition or spin coating, deposition and coating conditions for the EML may be determined by referring to the deposition and coating conditions for the HIL.

[0227] The EML may include a host and a dopant. The host may include at least one condensed-cyclic compound represented by Formula 1A or 1B.

[0228] The host may include at least one compound selected from TPBi, TBADN, ADN (also referred to as "DNA"), CBP, CDBP, and TCP:




[0229] When the organic light-emitting device **10** is a full color organic light-emitting device, the EML may be patterned into a red EML, a green EML, and a blue EML. In some embodiments, the EML may have a stacked structure of a red EML, a green EML, and/or a blue EML to emit white light. The host in the red EML, green EML, and blue EML may include the condensed-cyclic compound represented by Formula 1. According to an embodiment, the host in the green EML may include the condensed-cyclic compound represented by Formula 1A or 1B.

[0230] The EML may include a fluorescent dopant that emits light according to a fluorescent light emission mechanism or a phosphorescent dopant that emits light according to a phosphorescent light emission mechanism.

[0231] According to an embodiment, the EML may include a fluorescent and a phosphorescent dopant including the condensed-cyclic compound represented by Formula 1A or 1B. The phosphorescent dopant may include an organic metal complex including a transition metal (for example, iridium (Ir), platinum (Pt), osmium (Os), rhodium (Rh), or the like).

[0232] The phosphorescent compound may include an organometallic compound represented by Formula 81:



[0233] In Formula 81,

[0234] M may be selected from Iridium (Ir), platinum (Pt), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), and thulium (Tm);

[0235] Y<sub>1</sub> to Y<sub>4</sub> may be each independently carbon (C) or nitrogen (N);

[0236] Y<sub>1</sub> and Y<sub>2</sub> may be connected by a single bond or a double bond, and Y<sub>3</sub> and Y<sub>4</sub> may be connected by a single bond or a double bond;

[0237] CY<sub>1</sub> and CY<sub>2</sub> may be each independently selected from a benzene, a naphthalene, a fluorene, a spiro-fluorene, an indene, a pyrrole, a thiophene, a furan, an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isooxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, a quinoline, an isoquinoline, a benzoquinoline, a quinoxaline, a quinazoline, a carbazole, a benzoimidazole, benzofuran, a benzothiophene, an isobenzothiophene, a benzooxazole, an isobenzooxazole, a triazole, a tetrazole, an oxadiazole, a triazine, a dibenzofuran, and a dibenzothiophene, wherein CY<sub>1</sub> and CY<sub>2</sub> are optionally bound to each other via a single bond or an organic linking group;

[0238] R<sub>81</sub> and R<sub>82</sub> are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, —SF<sub>5</sub>, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, —N(Q<sub>1</sub>)(Q<sub>2</sub>), —Si(Q<sub>3</sub>)(Q<sub>4</sub>)(Q<sub>5</sub>), and —B(Q<sub>6</sub>)(Q<sub>7</sub>);

[0239] a81 and a82 are each independently selected from integers of 1 to 5;


[0240] n81 is selected from integers of 0 to 4;

[0241] n82 is 1, 2, or 3; and






[0242] L<sub>81</sub> is selected from a monovalent organic ligand, a divalent organic ligand, and a trivalent organic ligand.

[0243] Descriptions of R<sub>81</sub> and R<sub>82</sub> may be the same as the description of R<sub>5</sub>.

[0244] The phosphorescent dopant may include at least one of Compounds PD1 to PD74, but it is not limited thereto:




-continued




-continued

PD2




PD7

PD3




PD8

PD4




PD9

PD5








PD10

PD6




PD11

-continued




-continued

PD12




PD17

PD13




PD18

PD14



PD19


PD15




PD20

PD16


-continued




PD21




PD22



PD23




PD24




PD25


-continued




PD26




PD27



PD28








PD29




PD30

-continued




-continued

PD31




PD36

PD32




PD37

PD33




PD38

PD34








PD39

PD35




PD40

-continued




-continued

PD41




PD46

PD42




PD47

PD43




PD48

PD44




PD49

PD45




PD50



PD51


-continued

PD52




-continued


PD57




PD53




PD58




PD54




PD59




PD55




PD60




PD56




PD61




-continued




PD62




PD63



PD64




PD65




PD66


-continued




PD67




PD68



PD69






PD70




PD71

-continued



[0245] Alternatively, the phosphorescent dopant may include PtOEP or compound PhGD:


PD72



PD73


PtOEP

PD74




PhGD

[0246] The fluorescent dopant may include at least one of DPVBi, DPAVBi, TBPe, DCM, DCJTb, Coumarin 6, and C545T.




DPVBi




DPAVBi


-continued




TBPe




DCM



DCJTb



Coumarin 6



C545T

[0247] When the EML includes the host and the dopant, an amount of the dopant may be selected from a range of about 0.01 parts by weight to about 15 parts by weight based on 100 parts by weight of the host, but the amount is not limited thereto.

[0248] A thickness of the EML may be about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the EML is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0249] Then, an electron transport region may be disposed on the EML.


[0250] The electron transport region may include at least one layer selected from a HBL, an ETL, and an EIL, but is not limited thereto.

[0251] For example, the electron transport region may have an HBL/ETL/EIL structure or an ETL/EIL structure, wherein layers of each structure are sequentially stacked from the EML in the stated order, but is not limited thereto.



The ETL may have a single layer or a multi-layer structure including two or more different materials.

[0252] The conditions for forming the HBL, ETL, and EIL may be understood by referring to the conditions for forming the HIL.

[0253] When the electron transport region includes the HBL, the HBL may include, for example, at least one of BCP and Bphen, but it is not limited thereto.



BCP




**[0254]** A thickness of the HBL may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thickness of the HBL is within the range described above, the HBL may have excellent hole blocking characteristics without a substantial increase in driving voltage.



**[0255]** The ETL may include at least one of BCP and Bphen, and may further include at least one of Alq<sub>3</sub>, Balq, TAZ, and NTAZ.

**[0256]** Alternatively, the ETL may include at least one of Compound ET1 and ET2, but it is not limited thereto.

ET1




ET2




**[0257]** A thickness of the ETL may be in a range of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the ETL is within the range described above, the ETL may have satisfactory electron transportation characteristics without a substantial increase in driving voltage.

**[0258]** Also, the ETL may further include, in addition to the materials described above, a metal-containing material.

[0259] The metal-containing material may include a  $L_1$  complex. The  $L_1$  complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.



ET-D1



ET-D2

[0260] The electron transport region may include an EIL that allows electrons to be easily provided from the second electrode 19.

[0261] The EIL may include at least one compound selected from LiF, NaCl, CsF,  $Li_2O$ , and BaO.

[0262] A thickness of the EIL may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the EIL is within the range described above, the EIL may have satisfactory electron transportation characteristics without a substantial increase in driving voltage.

[0263] The second electrode 19 is disposed on the organic layer 15 having the structure described above. The second electrode 19 may be a cathode that is an electron injection electrode, and in this regard, a material for forming the second electrode 19 may be a material having a low work function, and such a material may be a metal, an alloy, an electrically conductive compound, or a mixture thereof. Detailed examples of the material for forming second electrode 19 are lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag). Alternatively, ITO or IZO may be used to form a transmissive second electrode 19 to manufacture a top emission light-emitting device.

[0264] Hereinbefore, the organic light-emitting device has been described with reference to the FIGURE, but is not limited thereto.

[0265] A  $C_1$ - $C_{60}$  alkyl group used herein refers to a linear or branched aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms. Detailed examples thereof are a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. A  $C_1$ - $C_{60}$  alkylene used herein refers to a divalent group having the same structure as the  $C_1$ - $C_{60}$  alkyl group.

[0266] A  $C_1$ - $C_{60}$  alkoxy group used herein refers to a monovalent group represented by  $-OA_{101}$  (wherein  $A_{101}$  is

the  $C_1$ - $C_{60}$  alkyl group). Detailed examples thereof are a methoxy group, an ethoxy group, and an isopropoxy group.

[0267] A  $C_2$ - $C_{60}$  alkenyl group used herein refers to a hydrocarbon group formed by substituting at least one carbon double bond in the middle or at the terminal of the  $C_2$ - $C_{60}$  alkyl group. Detailed examples thereof are an ethenyl group, a propenyl group, and a butenyl group. A  $C_2$ - $C_{60}$  alkenylene group used herein refers to a divalent group having the same structure as the  $C_2$ - $C_{60}$  alkenyl group.

[0268] A  $C_2$ - $C_{60}$  alkynyl group used herein refers to a hydrocarbon group formed by substituting at least one carbon triple bond in the middle or at the terminal of the  $C_2$ - $C_{60}$  alkyl group. Detailed examples thereof are an ethynyl group and a propynyl group. A  $C_2$ - $C_{60}$  alkynylene group used herein refers to a divalent group having the same structure as the  $C_2$ - $C_{60}$  alkynyl group.

[0269] A  $C_3$ - $C_{10}$  cycloalkyl group used herein refers to a monovalent hydrocarbon monocyclic group having 3 to 10 carbon atoms. Detailed examples thereof are a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. A  $C_3$ - $C_{10}$  cycloalkylene group used herein refers to a divalent group having the same structure as the  $C_3$ - $C_{10}$  cycloalkyl group.

[0270] A  $C_3$ - $C_{10}$  heterocycloalkyl group used herein refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, P, and S as a ring-forming atom and 3 to 10 carbon atoms. Detailed examples thereof are tetrahydrofuran and tetrahydrothiophene. A  $C_3$ - $C_{10}$  heterocycloalkylene group used herein refers to a divalent group having the same structure as the  $C_3$ - $C_{10}$  heterocycloalkyl group.

[0271] A  $C_3$ - $C_{10}$  cycloalkenyl group used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one double bond in the ring thereof and does not have aromaticity. Detailed examples thereof are a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. A  $C_3$ - $C_{10}$  cycloalkenylene group used herein refers to a divalent group having the same structure as the  $C_3$ - $C_{10}$  cycloalkenyl group.

[0272] A  $C_2$ - $C_{10}$  heterocycloalkenyl group used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, P, and S as a ring-forming atom, 2 to 10 carbon atoms, and at least one double bond in its ring. Detailed examples of the  $C_2$ - $C_{10}$  heterocycloalkenyl group are a 2,3-dihydrofuran group and a 2,3-dihydrothiophene group. A  $C_2$ - $C_{10}$  heterocycloalkenylene group used herein refers to a divalent group having the same structure as the  $C_2$ - $C_{10}$  heterocycloalkenyl group.

[0273] A  $C_6$ - $C_{60}$  aryl group used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a  $C_6$ - $C_{60}$  arylene group used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Detailed examples of the  $C_6$ - $C_{60}$  aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the  $C_6$ - $C_{60}$  aryl group and the  $C_6$ - $C_{60}$  arylene group each include two or more rings, the rings may be fused to each other.

[0274] A  $C_2$ - $C_{60}$  heteroaryl group used herein refers to a monovalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, P, and S as a ring-forming atom, and 2 to 60 carbon atoms. A  $C_2$ - $C_{60}$  heteroarylene group used herein refers to a divalent group

having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, P, and S as a ring-forming atom, and 2 to 60 carbon atoms. Detailed examples of the C<sub>2</sub>-C<sub>60</sub> heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C<sub>2</sub>-C<sub>60</sub> heteroaryl group and the C<sub>2</sub>-C<sub>60</sub> heteroarylene group each include two or more rings, the rings may be fused to each other.

[0275] A C<sub>6</sub>-C<sub>60</sub> aryloxy group used herein indicates —OA<sub>102</sub> (wherein A<sub>102</sub> is the C<sub>6</sub>-C<sub>60</sub> aryl) and a C<sub>6</sub>-C<sub>60</sub> arylthio group used herein indicates —SA<sub>103</sub> (wherein A<sub>103</sub> is the C<sub>6</sub>-C<sub>60</sub> aryl group).

[0276] A monovalent non-aromatic condensed polycyclic group (for example, having 8 to 60 carbon atoms) used herein refers to a monovalent group that has two or more rings condensed to each other, only carbon atoms as ring-forming atoms, wherein the molecular structure as a whole is non-aromatic. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. A divalent non-aromatic condensed polycyclic group used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0277] A monovalent non-aromatic condensed heteropolycyclic group (for example, having 2 to 60 carbon atoms) used herein refers to a monovalent group that has two or more rings condensed to each other, has a heteroatom selected from N, O, P, and S, other than carbon atoms, as a ring forming atom, wherein the molecular structure as a whole is non-aromatic. Detailed examples of the monovalent non-aromatic condensed heteropolycyclic group are a carbazolyl group. A divalent non-aromatic condensed heteropolycyclic group used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.


[0278] Hereinafter, an organic light-emitting device according to an embodiment will be described in detail with reference to Synthesis Examples and Examples. The wording "B was used instead of A" used in describing Synthesis Examples means that a molar equivalent of A was identical to a molar equivalent of B.

#### EXAMPLE

##### Synthesis Example 1: Synthesis of Compound 813


##### Synthesis of Intermediate B

[0279]



Molecular Weight: 196.20

Intermediate A




[0280] 44.0 g (224.3 mmol) of Intermediate A, 126.9 g (224.3 mmol) of acetophenone, and 9.0 g (224.3 mmol) of sodium hydroxide were added in 670 mL of ethanol in a 1,000 mL round bottom flask and then stirred in nitrogen atmosphere for 2 hours at room temperature to prepare a mixture. Crystallized solids in the mixture were filtered to obtain Intermediate B (59.2 g and yield 88.0%). Element analysis results of Intermediate B are as follows.

[0281] calcd. C<sub>21</sub>H<sub>14</sub>O<sub>2</sub>: C, 84.54; H, 4.73; O, 10.73; found: C, 84.51; H, 4.75; O, 10.71.


##### Synthesis of Intermediate C

[0282]





Molecular Weight: 296.33

Intermediate B



Molecular Weight: 235.51




**[0283]** 13.0 g (106.2 mmol) of Intermediate B, 30.0 g (127.4 mmol) of 3-bromobenzimidamide, and 8.5 g (212.3 mmol) of sodium hydroxide were added in 500 ml of ethanol in a 1,000 mL round bottom flask, heated in a nitrogen atmosphere for 15 hours to reflux the same to prepare a mixture. Crystallized solids in the mixture were stirred with water and then filtered. The crystallized solids were stirred again by using ethanol and then filtered to obtain Intermediate C (23.8 g, yield 47%). Element analysis results of the Intermediate C are as follows.

**[0284]** calcd.  $C_{28}H_{17}BrN_2O$ : C, 70.45; H, 3.59; Br, 16.74; N, 5.87; O, 3.35; found C, 70.43; H, 3.60; Br, 16.72; N, 5.85; O, 3.36.

#### Synthesis of Compound 813

**[0285]**




**[0286]** 23.8 g (49.9 mmol) of Intermediate C, 7.0 g (41.6 mmol) of 9H-carbazole, 8.0 g (83.1 mmol) of sodium t-butoxide, 3.8 g (4.2 mmol) of Pd(dba)<sub>2</sub>, and 4.2 mL (8.3 mmol) of tri-*t*-butyl phosphine (50% in toluene solution) were added to 166.2 mL of xylene in a 500 mL round bottom flask, and heated in a nitrogen atmosphere for 15 hours to reflux the same to prepare a mixture. The mixture was added to 1,000 mL of methanol to filter crystallized solids, dissolved in monochlorobenzene to filter the same by using a silica gel/celite, a suitable amount of organic solvent was removed therefrom, and then the same was recrystallized with methanol to obtain Compound 813 (11.7 g and yield 50%). Element analysis results and NMR analysis results of Compound 813 are as follows.



**[0287]** calcd.  $C_{40}H_{25}N_3O$ : C, 85.24; H, 4.47; N, 7.46; O, 2.84; found C, 85.22; H, 4.46; N, 7.47; O, 2.83.

**[0288]** 300 MHz (CDCl<sub>3</sub>, ppm): δ 8.960 (m, 1H), 8.865 (m, 2H), 8.657 (dd, 1H), 8.359 (dd, 2H), 8.196 (d, 2H), 8.081 (dd, 1H), 7.999 (d, 1H), 7.816 (t, 1H), 7.659-7.727 (m, 2H), 7.381-7.601 (m, 10H), 7.316 (t, 2H)

#### Synthesis Example 2: Synthesis of Compound 814

**[0289]**





**[0290]** 15.0 g (31.4 mmol) of Intermediate C, 9.7 g (37.7 mmol) of 5H-benzofuro[3,2-c]carbazole, 6.0 g (62.9 mmol) of sodium t-butoxide, 2.9 g (3.1 mmol) of Pd(dba)<sub>2</sub>, and 3.1 mL (6.3 mmol) of tri-t-butyl phosphine (50% in toluene solution) were added to 125.7 mL of xylene in a 500 mL round bottom flask, and heated in a nitrogen atmosphere for 15 hours to reflux the same to prepare a mixture. The mixture was added to 1,000 mL of methanol to filter crystallized solids, dissolved in dichlorobenzene to filter the same by using a silica gel/celite, a suitable amount of organic solvent was removed therefrom, and then the same was recrystallized with methanol to obtain Compound 814 (11.8 g, yield 57%). Element analysis results and NMR analysis results of Compound 814 are as follows.


**[0291]** calcd. C<sub>46</sub>H<sub>27</sub>N<sub>3</sub>O<sub>2</sub>: C, 84.51; H, 4.16; N, 6.43; O, 4.89; found C, 84.49; H, 4.17; N, 6.44; O, 4.87.

**[0292]** 300 MHz (CDCl<sub>3</sub>, ppm): δ 9.023 (t, 1H), 8.905-8.959 (m, 2H), 8.797 (dd, 1H), 8.666 (dd, 1H), 8.392 (m, 2H), 8.110 (dd, 1H), 8.012 (m, 3H), 7.684-7.884 (t, 4H), 7.362-7.627 (m, 12H)

#### Synthesis Example 3: Synthesis of Compound 815

##### Synthesis of Intermediate D

**[0293]**





**[0294]** 30.0 g (152.9 mmol) of Intermediate A, 30.4 g (152.9 mmol) of 1-(3-bromophenyl)ethanone, and 6.1 g (152.9 mmol) of sodium hydroxide were added to 458.7 mL of ethanol in a 1,000 mL round bottom flask and then stirred in a nitrogen current for 2 hours at room temperature to prepare a mixture. Crystallized solids in the mixture were filtered to obtain Intermediate D (46.2 g, yield 80.0%). Element analysis results of Intermediate D are as follows.

**[0295]** calcd. C<sub>21</sub>H<sub>13</sub>BrO<sub>2</sub>: C, 66.86; H, 3.47; Br, 21.18; O, 8.48; found C, 66.83; H, 3.45; Br, 21.17; O, 8.49.

##### Synthesis of Intermediate E

**[0296]**







**[0297]** 80.3 g (212.9 mmol) of Intermediate D, 40.0 g (255.4 mmol) of benzimidamide, and 17 g (425.7 mmol) of sodium hydroxide were added to 1,000 ml of ethanol in a 2,000 mL round bottom flask and then heated in a nitrogen atmosphere for 15 hours to reflux the same to prepare a mixture. Crystallized solids in the mixture were filtered and then stirred with water to filter the same. The crystallized solids were stirred again with ethanol and then filtered to obtain Intermediate E (45.8 g, yield 45%). Element analysis results of Intermediate E are as follows.

**[0298]** calcd.  $C_{28}H_{17}BrN_2O$ : C, 70.45; H, 3.59; Br, 16.74; N, 5.87; O, 3.35; found C, 70.44; H, 3.61; Br, 16.71; N, 5.84; O, 3.35.

#### Synthesis of Compound 815

**[0299]**




**[0300]** 34.3 g (71.8 mmol) of Intermediate E, 10.0 g (60.0 mmol) of 9H-carbazole, 11.5 g (119.6 mmol) of sodium t-butoxide, 5.5 g (6.0 mmol) of  $Pd(dba)_2$ , and 2.9 mL (12.0 mmol) of tri-t-butyl phosphine (50% in toluene solution) were added to 239.2 mL of xylene in a 500 mL round bottom flask and then heated in a nitrogen atmosphere for 15 hours to reflux the same to prepare a mixture. The mixture was added to 1,000 mL of methanol to filter crystallized solids, which were dissolved in monochlorobenzene to be filtered by using a silica gel/celite, a suitable amount of organic solvent was removed therefrom and then recrystallized with methanol to obtain Compound 815 (22.2 g, yield 66%). Element analysis results and NMR analysis results of Compound 815 are as follows.


**[0301]** calcd.  $C_{40}H_{25}N_3O$ : C, 85.24; H, 4.47; N, 7.46; O, 2.84; found C, 85.21; H, 4.47; N, 7.48; O, 2.86.

**[0302]** 300 MHz ( $CDCl_3$ , ppm):  $\delta$  8.905 (s, 1H), 8.745 (m, 3H), 8.635 (t, 1H), 8.473 (tt, 1H), 8.208 (d, 2H), 8.120 (dd, 1H), 8.016 (d, 1H), 7.802-7.885 (m, 2H), 7.317-7.600 (m, 13H)

#### Synthesis Example 4: Synthesis of Compound 816

**[0303]**







**[0304]** 19.0 g (56.0 mmol) of Intermediate E, 12.0 g (46.6 mmol) of 5H-benzofuro[3,2-c]carbazole, 6.7 g (70.0 mmol) of sodium t-butoxide, 1.3 g (1.4 mmol) of Pd(dba)<sub>2</sub>, and 2.1 mL (4.2 mmol) tri t-butylphosphine (50% in toluene) were added to 182.9 mL of xylene in a 500 mL round bottom flask and then heated in a nitrogen atmosphere for 15 hours to reflux the same to prepare a mixture. The mixture was added to 1,000 mL of methanol to filter crystallized solids, dissolved in dichlorobenzene to filter the same by using a silica gel/celite, a suitable amount of organic solvent was removed therefrom, and then the same was recrystallized with methanol to obtain Compound 816 (14.7 g, yield 48%). Element analysis results and NMR analysis results of Compound 816 are as follows.

**[0305]** calcd. C<sub>46</sub>H<sub>27</sub>N<sub>3</sub>O<sub>2</sub>: C, 84.51; H, 4.16; N, 6.43; O, 4.89; found C, 84.53; H, 4.18; N, 6.42; O, 4.87.

**[0306]** 300 MHz (CDCl<sub>3</sub>, ppm): δ 8.903 (s, 1H), 8.739 (d, 3H), 8.641 (t, 2H), 8.506 (d, 1H), 8.096 (d, 1H), 7.976 (m, 3H), 7.868 (m, 2H), 7.763 (d, 1H), 7.347-7.634 (m, 13H)

#### Synthesis Example 5: Synthesis of Compound 506

**[0307]**



#### a) Synthesis of Intermediate 506-1

**[0308]** 22.6 g (100 mmol) of 2,4-dichloro-6-phenyltriazine was added to 100 mL of tetrahydrofuran, 100 mL of toluene, and 100 mL of distilled water in a 500 mL round-bottom flask, and then, 0.9 equivalent weights of dibenzofuran-3-boronic acid, 0.03 equivalent weights of tetrakis(triphenyl)phosphine palladium, and 2 equivalent weights of potassium carbonate were added thereto, followed by heating while refluxing in a nitrogen atmosphere. After 6 hours, the reaction solution was cooled, the water layer was removed, and the organic layer was dried under reduced pressure. The resulting solid was washed with water and hexane, and the solid was recrystallized by using 200 mL of toluene to complete the preparation of Intermediate 506-1 at a yield of 60%.

#### b) Synthesis of Compound 506


**[0309]** 8.01 g (22.4 mmol) of Intermediate 506-1 synthesized above was added to 80 mL of tetrahydrofuran and 40 mL of distilled water in a 500 mL round-bottom flask, and 1.0 equivalent weight of 9-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-phenyl)-carbazole (CAS: 785051-54-9), 0.03 equivalent weights of tetrakis(triphenyl)phosphine palladium, and 2 equivalent weights of potassium carbonate were added thereto, followed by heating while refluxing in

a nitrogen atmosphere. After 18 hours, the reaction solution was cooled and the precipitated solid was filtered and washed with 500 mL of water. The solid was recrystallized by using 500 mL of monochlorobenzene to complete the preparation of 10.12 g of Compound 506.


[0310] LC/MS calculated for:  $C_{39}H_{24}N_4O$  Exact Mass: 564.1950 found for: 565.20.

Synthesis Example 6: Synthesis of Compound 825

[0311]



-continued





[0312] Compound 825 was synthesized in the same manner as in the same manner as in "b)" of Synthesis Example 5, except that 9-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-phenyl)-carbazole (CAS: 870119-58-7) was used in an equivalent weight of 1.0 instead of 9-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-phenyl)-carbazole.


[0313] LC/MS calculated for:  $C_{39}H_{24}N_4O$  Exact Mass: 564.1950 found for: 565.20.

Comparative Synthesis Example 1: Synthesis of Comparative Compound R1

[0314]



-continued



## a) Synthesis of Intermediate R1-1

[0315] Intermediate R1-1 was synthesized in the same manner as in “b)” of Synthesis Example 5, except that each of 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-dibenzofuran and 1-bromo-4-chloro-benzene was used in an equivalent weight of 1 instead of Intermediate 506-1 and 9-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-phenyl)-carbazole.

## b) Synthesis of Intermediate R1-2

[0316] One equivalent weight of Intermediate R1-1 synthesized above was added to 150 mL of xylene in a 500 mL round-bottom flask, and 0.05 equivalent weights of dichlorodiphenylphosphinoferrrocene palladium, 1.2 equivalent weights of bis(pinacolato)diboron, and 2 equivalent weights of potassium acetate were added thereto, followed by heating while refluxing for 18 hours in a nitrogen atmosphere. After cooling the reaction solution, the resultant solution was washed with water through extraction, the organic layer was treated with activated carbon and filtered through silica gel, and the filtrate was concentrated. The concentrated solid

was stirred with a small amount of hexane, and then the solid was filtered therefrom to complete the preparation of Intermediate R1-2 at a yield of 75%.

## c) Synthesis of Intermediate R1-3

[0317] Intermediate R1-3 was synthesized in the same manner as in “a)” of Synthesis Example 5, except that 0.9 equivalent weights of [4-(9H-carbazol-9-yl)phenyl]boronic acid was used instead of dibenzofuran-3-boronic acid.


## d) Synthesis of Comparative Compound R1

[0318] Comparative Compound R1 was synthesized in the same manner as in “a)” of Synthesis Example 5, except that Intermediate R1-2 and Intermediate R1-3 was used in an equivalent weight of 1 instead of 2,4-dichloro-6-phenyl-1,3,5-triazine and dibenzofuran-3-boronic acid.


[0319] LC/MS calculated for:  $C_{45}H_{28}N_4O$  Exact Mass: 640.2263 found for: 641.23.

## Comparative Synthesis Example 2: Synthesis of Comparative Compound R2

[0320]



-continued




## a) Synthesis of Intermediate R2-1

[0321] Intermediate R2-1 was synthesized in the same manner as in "a)" of Synthesis Example 5, except that 2,4-dichloro-6-phenylpyrimidine was used in an equivalent weight of 1 instead of 2,4-dichloro-6-phenyltriazine.

## b) Synthesis of Comparative Compound R2


[0322] Comparative Compound R2 was synthesized in the same manner as in "b)" of Synthesis Example 5, except that each of Intermediate R2-1 synthesized above was used in an equivalent weight of 1.0 instead of Intermediate 506-1.

[0323] LC/MS calculated for:  $C_{40}H_{25}N_3O$  Exact Mass: 563.1998 found for: 564.23.




## Comparative Synthesis Example 3: Synthesis of Comparative Compound R3

[0324]



-continued



R3

## a) Synthesis of Intermediate R3-1

**[0325]** In a nitrogen atmosphere, 7.86 g (323 mmol) of magnesium and 1.64 g (6.46 mmol) of iodine were added to 0.1 L of tetrahydrofuran (THF) and stirred for 30 minutes, and then, 80 g (323 mmol) of 3-bromodibenzofuran dissolved in 0.3 L of THF was slowly added dropwise thereto at a temperature of 0° C. for 30 minutes. The resulting mixed solution was slowly added dropwise to 29.5 g (160 mmol) of cyanuric chloride dissolved in 0.5 L of THF at a temperature of 0° C. for 30 minutes. The reaction temperature was raised to room temperature, followed by 1 hour of stirring. Then, in the refluxing condition, the resultant solution was additionally stirred for 12 hours. After cooling the reaction mixture, water was added slowly to terminate the reaction, and the organic solvent was concentrated under reduced pressure to obtain a solid. This was stirred with 200 mL of acetone and then filtered to complete the preparation of Intermediate R3-1 at a yield of 40%.

## b) Synthesis of Comparative Compound R3

**[0326]** Comparative Compound R3 was synthesized in the same manner as in “b)” of Synthesis Example 5, except that each of Intermediate R3-1 synthesized above was used in an equivalent weight of 1.0 instead of Intermediate 506-1.

**[0327]** LC/MS calculated for:  $C_{45}H_{26}N_4O_2$  Exact Mass: 654.2056 found for: 655.2089.

## Evaluation Example 1: Evaluation of HOMO, LUMO, and Triplet (T1) Energy Levels of Compounds

**[0328]** HOMO, LUMO, and triplet (T1) energy levels of Compounds 813 to 824 were evaluated by using a DFT method of a Gaussian program (optimized at B3LYP and 6-31G(d,p) levels) and the results obtained therefrom are shown in Table 1 below.

TABLE 1

Compound No.	HOMO (eV)	LUMO (eV)	T <sub>1</sub> energy level (eV)
813	-5.236	-1.880	2.994
815	-5.355	-1.802	3.096
817	-5.252	-1.877	3.002
819	-5.435	-1.894	3.089
814	-5.150	-1.894	2.9
816	-5.344	-1.909	3.02
820	-5.303	-1.925	2.959
818	-5.139	-1.885	2.904
821	-5.188	-1.893	2.945
822	-5.350	-1.934	2.997
823	-5.206	-1.897	2.951
824	-5.394	-1.913	3.063

**[0329]** Thereafter, HOMO, LUMO, S1 energy level, and T1 energy level of Compounds 813 to 816 obtained according to the method in Table 2 and the results thereof are shown in Table 3.

TABLE 2

An evaluation method of a HOMO energy level	Each compound was diluted in $CHCl_3$ at a concentration of $1 \times 10^{-5}M$ to measure UV absorption spectrum by using a Shimadzu UV-350 Spectrometer at room temperature and then a HOMO energy level was calculated by using an optical band gap (Eg) from an edge of the absorption spectrum.
An evaluation method of a LUMO energy level	Cyclic voltammetry (CV) (electrolyte: 0.1M $Bu_4NClO_4$ /solvent: $CH_2Cl_2$ /electrode: a 3-electrode system (working electrode: GC, standard electrode: Ag/AgCl, and supply electrode: Pt)) was used to obtain a voltage (V)-current (A) graph for each compound and a LUMO energy level was calculated from a reduction onset of the graph.
An evaluation method of eVT1	A mixture of toluene and each compound (1 mg of each compound was dissolved in 3 cc of toluene) was added to a quartz cell and liquid nitrogen (77 K)

TABLE 2-continued

energy level	was added thereto, a photoluminescence measurement device was used to measure photoluminescence spectrum, which was compared to a conventional room temperature photoluminescence spectrum to only analyze the peak observed only at low temperature and calculate a T1 energy level.
--------------	---

TABLE 3

Compound No.	HOMO (eV)	LUMO (eV)	T1 energy level (eV)
813	-5.432	-1.901	2.994
815	-5.665	-1.972	3.096
814	-5.451	-1.998	2.9
816	-5.643	-1.942	3.02

[0330] It may be concluded from Tables 1 and 3 that the condensed-cyclic compounds have suitable electrical properties to be used as a material for an organic light-emitting device.

#### Evaluation Example 2: Evaluation of Thermal Properties of Synthesized Compounds

[0331] compounds 813 to 816 were subjected to thermal analyses by using Thermo Gravimetric Analysis (TGA) and Differential Scanning calorimetry (DSC) (N<sub>2</sub> atmosphere and temperature range for TGA: room temperature ~800° C. (10° C./min), for DSC: from room temperature to 400° C., Pan Type: for TGA: Pt Pan in a disposable Al Pan, for DSC: a disposable Al pan) and results obtained therefrom are shown in Table 4. From Table 4, it may be concluded that Compounds 813 to 816 have excellent thermal stabilities.

TABLE 4

Compound No.	Tc (° C.)	Tm (° C.)	Tg (° C.)
813	—	—	101.21
815	218.72	241.72	102.57
814	—	312.38	—
816	—	—	139.38

#### Example 1

[0332] An ITO glass substrate was cut to a size of 50 mm×50 mm×0.5 mm, and the ITO glass substrate was ultrasonically washed using isopropyl alcohol and pure water for 15 minutes each, followed by irradiation of UV and exposure to ozone for cleaning for about 30 minutes.

[0333] m-MTDATA was vacuum deposited on the ITO glass substrate to form an HIL having a thickness of 600 Å, and α-NPB was vacuum deposited at a rate of 1 Å/sec on the HIL to form an EML having a thickness of 300 Å. Thereafter, Ir(ppy)<sub>3</sub> (dopant) and Compound 813 (host) were co-deposited at a rate of 0.1 Å/sec and 1 Å/sec, respectively, on the HTL to form an EML having a thickness of 400 Å. BAiq was vacuum deposited on the EML at a rate of 1 Å/sec to form an HBL having a thickness of 50 Å and then Alq<sub>3</sub> was vacuum deposited on the HBL to form an ETL having a thickness of 300 Å. Then, LiF 10 Å (EIL) and Al 2000 Å (cathode) were sequentially vacuum deposited on the ETL to manufacture an organic light-emitting device.

#### Example 2

[0334] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 815 was used as a host instead of Compound 813 when forming an EML.

#### Example 3


[0335] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 814 was used as a host instead of Compound 813 when forming an EML.

#### Example 4

[0336] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 816 was used as a host instead of Compound 813 when forming an EML.


#### Comparative Example 1

[0337] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound A was used as a host instead of Compound 813 when forming an EML.




#### Comparative Example 2

[0338] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound B was used as a host instead of Compound 813 when forming an EML.



## Comparative Example 3

[0339] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound C was used as a host instead of Compound 813 when forming an EML.



## Evaluation Example 2: Evaluation of Characteristics of Organic Light-Emitting Device

[0340] Changes in current density and brightness, and emission efficiency of each organic light-emitting device manufactured in Examples 1 to 4 and Comparative Examples 1 to 3 were measured. A detailed method of measurement is as described below and results obtained therefrom are shown in Table 5 below:

[0341] (1) Measurement of Changes in Current Density According to Changes in Voltage

[0342] For each organic light-emitting device, voltage was increased from 0 volts (V) to 10 V to measure current that flows through a unit cell therein by using a voltage-current meter (Keithley 2400) and the current was divided by surface area to obtain a current density.

[0343] (2) Measurement of Changes in Brightness According to Changes in Voltage

[0344] For each organic light-emitting device, brightness was measured while increasing voltage from 0 V to 10 V to by using Cs-1000A (a product of Minolta).

[0345] (3) Measurement of Emission Efficiency

[0346] Brightness, current density, and voltage measured from (1) and (2) were used to calculate current efficiency (candelas per ampere (cd/A)) at the same current density (10 milliamperes per square centimeter (mA/cm<sup>2</sup>)).

TABLE 5

	Host	Dopant	Driving voltage (V)	Current density (cd/A)	Brightness (cd/m <sup>2</sup> )
Example 1	Compound 813	Ir(ppy) <sub>3</sub>	3.7	40.3	3500
Example 2	Compound 815	Ir(ppy) <sub>3</sub>	3.4	46.2	3500
Example 3	Compound 814	Ir(ppy) <sub>3</sub>	4.1	36.1	3500
Example 4	Compound 816	Ir(ppy) <sub>3</sub>	3.8	33.8	3500

TABLE 5-continued

	Host	Dopant	Driving voltage (V)	Current density (cd/A)	Brightness (cd/m <sup>2</sup> )
Comparative Example 1	Compound A	Ir(ppy) <sub>3</sub>	4.4	32.6	3500
Comparative Example 2	Compound B	Ir(ppy) <sub>3</sub>	4.3	31.5	3500
Comparative Example 3	Compound C	Ir(ppy) <sub>3</sub>	4.2	32.9	3500

[0347] From Table 5, it may be concluded that the organic light-emitting devices of Examples 1 to 4 have lower driving voltage, high efficiency, and high brightness compared the organic light-emitting devices of Comparative Examples 1 to 3.

## Examples 5 and 6

[0348] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 506 and 825, respectively, was used as a host instead of Compound 813 when forming an EML.

## Comparative Examples R1 to R3

[0349] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound R1 to R3, respectively, was used as a host instead of Compound 813 when forming an EML.

## Evaluation Example 3

[0350] Driving voltage, luminous power efficiency and lifespan (T90@24K) of each organic light-emitting device manufactured in Example 5 and Comparative Example R1 were measured. A detailed method of measurement is as described below and results obtained therefrom are shown as a relative value (%) in Table 6 below. The relative value of the driving voltage, the luminous power efficiency and the lifespan of the organic light-emitting device of Comparative Example R1 were regarded as "100%."

[0351] (1) Measurement of Current Density Change Depending on Voltage Change

[0352] Current values flowing in the unit device of the manufactured organic light emitting diodes were measured for, while increasing the voltage from 0 V to 10 V using a current-voltage meter (Keithley 2400), and the measured current values were divided by an area to provide the results.

[0353] (2) Measurement of Luminance Change Depending on Voltage Change

[0354] Luminance of the manufactured organic light emitting diodes was measured for luminance, while increasing the voltage from 0 V to 10 V using a luminance meter (Minolta Cs-1000A).

[0355] (3) Measurement of Luminous Power Efficiency

[0356] Luminous Power efficiency (lm/W) at the same current density (10 mA/cm<sup>2</sup>) was calculated by using the luminance, current density, and voltages (V) from the items (1) and (2).

[0357] (4) Measurement of Driving Voltage



[0358] Driving voltages (V) of each device were measured at 15 mA/cm<sup>2</sup> using a current-voltage meter (Keithley 2400).

[0359] (5) Measurement of Lifespan (T<sub>90</sub>@24K)

[0360] T<sub>90</sub> was measured an amount of time that lapsed when 100% of the initial luminance (2400 cd/m<sup>2</sup>) was decreased to 90% at 24K using a Polanonix life-span measurement system.

TABLE 6

	Host	Driving voltage (relative value)	Luminous Power Efficiency (relative value)	T <sub>90</sub> @24K (relative value)
Example 5	Compound 506	91%	137%	288%
Comparative	Comparative	100%	100%	100%
Example R1	Compound R1			



[0361] From Table 6, it may be concluded that the organic light-emitting device of Example 5 has lower driving voltage, higher luminous power efficiency, and longer lifespan compared the organic light-emitting devices of Comparative Example R1.



#### Evaluation Example 4

[0362] Driving voltage and lifespan (T<sub>90</sub>@24K) of each organic light-emitting device manufactured in Example 6 and Comparative Example R2 were measured. A detailed method of measurement is as described in Evaluation Example 3 and results obtained therefrom are shown as a

relative value (%) in Table 7 below. The relative value of the driving voltage and the lifespan of the organic light-emitting device of Comparative Example R2 were regarded as "100%."

TABLE 7

	Host	Driving voltage (relative value)	T <sub>90</sub> @24K (relative value)
Example 6	Compound 825	96%	162%
Comparative	Comparative	100%	100%
Example R2	Compound R2		




[0363] From Table 7, it may be concluded that the organic light-emitting device of Example 6 has lower driving voltage, and longer lifespan compared the organic light-emitting devices of Comparative Example R2.

#### Evaluation Example 5


[0364] Driving voltage and lifespan (T<sub>90</sub>@24K) of each organic light-emitting device manufactured in Example 6 and Comparative Example R3 were measured. A detailed method of measurement is as described in Evaluation Example 3 and results obtained therefrom are shown as a relative value (%) in Table 8 below. The relative value of the driving voltage and the lifespan of the organic light-emitting device of Comparative Example R3 were regarded as "100%."

TABLE 8

	Host	Driving voltage (relative value)	T <sub>90</sub> @24K (relative value)
Example 6	Compound 825	105%	141%
Comparative	Comparative	100%	100%
Example R3	Compound R3		



825



R3

[0365] From Table 8, it may be concluded that the organic light-emitting device of Example 6 has longer lifespan compared the organic light-emitting devices of Comparative Example R2.


[0366] As described above, according to the one or more of the above embodiments, the condensed-cyclic compound have excellent electrical properties and thermal stability and thus, an organic light-emitting device including the condensed-cyclic compound may have low driving voltage, high efficiency, high brightness, and a long lifespan.

[0367] It should be understood that the exemplary embodiments described therein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as available for other similar features or aspects in other embodiments.

[0368] While one or more embodiments have been described with reference to the FIGURES, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope of the present disclosure as defined by the following claims.


What is claimed is:

1. A condensed-cyclic compound represented by Formula 1A:



Formula 1A


wherein in Formula 1A,  
ring A<sub>1</sub> is represented by Formula 1D;




Formula 1D

X<sub>1</sub> to X<sub>3</sub> is N;  
X<sub>4</sub> is O or S;


L<sub>4</sub> and L<sub>21</sub> are each independently selected from groups represented by Formulae 2-1, 2-2 and 2-34;



Formula 2-1



Formula 2-2



Formula 2-34

wherein in Formulae 2-1, 2-2 and 2-34, Z<sub>1</sub> is selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group and a phenyl group and d1 is selected from integers of 1 to 4;

a4 is selected from integers 0 to 3 and a21 is 0, 1, 2 or 3;  
R<sub>5</sub>, R<sub>6</sub>, R<sub>15</sub> and R<sub>17</sub> are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino

group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group;




$R_4$  is selected from:

a phenyl group; and

a phenyl group, substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, and a biphenyl group;


$b_4$  to  $b_6$ ,  $b_{15}$  and  $b_{17}$  are each independently selected from integers of 1 to 3.

2. The condensed-cyclic compound of claim 1, wherein the condensed-cyclic compound is represented by one of Formulae 1A(1) to 1A(4):



-continued



Formula 1A(4)




wherein in Formulae 1A(1) to 1A(4), descriptions of ring  $A_1$ ,  $X_1$  to  $X_4$ ,  $L_4$ ,  $L_{21}$ ,  $a_1$ ,  $a_{21}$ ,  $R_4$  to  $R_6$ ,  $R_{15}$ ,  $R_{17}$ ,  $b_4$  to  $b_6$ ,  $b_{15}$  and  $b_{17}$  are the same as described in claim 1.

3. The condensed-cyclic compound of claim 1, wherein  $a_{21}$  is 0, 1 or 2.


4. The condensed-cyclic compound of claim 1, wherein the condensed-cyclic compound is one of Compounds 1, 85, 169, 253, 337, 421, 505, 506, 575, 659, 660, 729 and 825:




-continued




169



253




337




421


-continued




505



506




575




659

-continued


660



729



825



5. An organic light-emitting device comprising  
a first electrode;  
a second electrode; and  
an organic layer disposed between the first electrode and the second electrode,  
wherein the organic layer comprises an emission layer  
and the condensed-cyclic compound of claim 1.

6. The organic light-emitting device of claim 5, wherein  
the first electrode is an anode and  
the second electrode is a cathode,  
wherein the organic layer comprises  
i) a hole transport region disposed between the first  
electrode and the emission layer,


wherein the hole transport region comprises at least one of  
a hole-injection layer, a hole-transporting layer, and an  
electron-blocking layer; and

ii) an electron transport region disposed between the  
emission layer and the second electrode,  
wherein the electron transport region comprises at least  
one layer selected from a hole-blocking layer, an elec-  
tron-transporting layer, and an electron-injecting layer.

7. The organic light-emitting device of claim 5, wherein  
the emission layer comprises the condensed-cyclic com-  
pound.

8. The organic light-emitting device of claim 7, wherein  
the emission layer further comprises an organometallic  
compound represented by Formula 81:

Formula 81



wherein in Formula 81,

M is selected from Ir, Pt, Os, Ti, Zr, Hf, Eu, Tb, and Tm;

Y<sub>1</sub> to Y<sub>4</sub> are each independently C or N;

Y<sub>1</sub> and Y<sub>2</sub> are connected by a single bond or a double  
bond and

Y<sub>3</sub> and Y<sub>4</sub> are connected by a single bond or a double  
bond;

CY<sub>1</sub> and CY<sub>2</sub> are each independently selected from a  
benzene, a naphthalene, a fluorene, a spiro-fluorene, an  
indene, a pyrrole, a thiophene, a furan, an imidazole, a  
pyrazole, a thiazole, an isothiazole, an oxazole, an  
isooxazole, a pyridine, a pyrazine, a pyrimidine, a  
pyridazine, a quinoline, an isoquinoline, a benzoqui-  
noline, a quinoxaline, a quinazoline, a carbazole, a  
benzimidazole, a benzofuran, a benzothiophene, an  
isobenzothiophene, a benzooxazole, an isobenzoox-  
azole, a triazole, a tetrazole, an oxadiazole, a triazine,  
a dibenzofuran, and a dibenzothiophene, wherein CY<sub>1</sub>  
and CY<sub>2</sub> are optionally bound to each other by a single  
bond or an organic linking group;

R<sub>81</sub> and R<sub>82</sub> are each independently selected from a  
hydrogen, a deuterium, —F, —Cl, —Br, —I, a  
hydroxyl group, a cyano group, a nitro group, an amino  
group, an amidino group, a hydrazine group, a hydra-  
zone group, a carboxylic acid or a salt thereof, a  
sulfonic acid or a salt thereof, a phosphoric acid or a  
salt thereof, —SF<sub>5</sub>, a substituted or unsubstituted  
C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted  
C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted  
C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted  
C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted  
C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted  
C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsub-  
stituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or

unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, —N(Q<sub>1</sub>)(Q<sub>2</sub>), —Si(Q<sub>3</sub>)(Q<sub>4</sub>)(Q<sub>5</sub>), and —B(Q<sub>6</sub>)(Q<sub>7</sub>);

a81 and a82 are each independently selected from integers of 1 to 5;

n81 is selected from integers of 0 to 4;

n82 is 1, 2, or 3; and

L<sub>81</sub> is selected from a monovalent organic ligand, a divalent organic ligand, and a trivalent organic ligand.

\* \* \* \* \*

专利名称(译)	缩合环状化合物和包括其的有机发光器件		
公开(公告)号	<a href="#">US20190074451A1</a>	公开(公告)日	2019-03-07
申请号	US16/177914	申请日	2018-11-01
[标]申请(专利权)人(译)	三星电子株式会社		
申请(专利权)人(译)	SAMSUNG ELECTRONICS CO. , LTD. 三星SDI CO. , LTD.		
当前申请(专利权)人(译)	SAMSUNG ELECTRONICS CO. , LTD. 三星SDI CO. , LTD.		
[标]发明人	LEE SEUNGJAE KIM BYUNGKU KIM YOUNGKWON KIM CHANGWOO KIM HYUNGSUN SHIN CHANGJU YU EUNSUN CHOI BYOUNGKI HWANG KYU YOUNG		
发明人	LEE, SEUNGJAE KIM, BYUNGKU KIM, YOUNGKWON KIM, CHANGWOO KIM, HYUNGSUN SHIN, CHANGJU YU, EUNSUN CHOI, BYOUNGKI HWANG, KYU-YOUNG		
IPC分类号	H01L51/00 C07D405/14 C09K11/06 C07D491/048		
CPC分类号	H01L51/0072 C07D405/14 C09K11/06 H01L51/0067 H01L51/0073 C07D491/048 H01L51/0071 H01L51/5088 H01L51/5056 H01L51/5096 H01L51/5012 H01L51/5072 H01L51/5092 C09K2211/1018 H01L51/0081 H01L51/0085 H01L51/5016		
优先权	1020130157532 2013-12-17 KR		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

#### 摘要(译)

由式1A表示的稠环化合物：其中在公式1A中，组，取代基和变量与规范中定义的相同。

**10**

<b>19</b>
<b>15</b>
<b>11</b>