



US 20160181530A1

(19) **United States**(12) **Patent Application Publication**
CHOI et al.(10) **Pub. No.: US 2016/0181530 A1**(43) **Pub. Date: Jun. 23, 2016**(54) **ORGANOMETALLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME****Publication Classification**(51) **Int. Cl.***H01L 51/00* (2006.01)*C09K 11/06* (2006.01)*C09K 11/02* (2006.01)*C07F 15/00* (2006.01)(52) **U.S. Cl.**CPC *H01L 51/009* (2013.01); *C07F 15/0033*(2013.01); *C09K 11/06* (2013.01); *C09K**11/025* (2013.01); *H01L 51/0072* (2013.01);*H01L 51/5004* (2013.01)(71) Applicant: **Samsung Electronics Co., Ltd.**,
Suwon-si (KR)(72) Inventors: **Whail CHOI**, Seoul (KR); **Seungyeon
KWAK**, Yongin-si (KR); **Yoonhyun
KWAK**, Seoul (KR); **Ohyun KWON**,
Yongin-si (KR); **Hyeonho CHOI**, Seoul
(KR); **Kyuyoung HWANG**, Anyang-si
(KR); **Sangdong KIM**, Seoul (KR);
Youngjae PARK, Seoul (KR)(21) Appl. No.: **14/967,594**(22) Filed: **Dec. 14, 2015**(30) **Foreign Application Priority Data**

Dec. 17, 2014 (KR) 10-2014-0182552

(57) **ABSTRACT**

An organometallic compound represented by Formula 1:

 $M(L_1)_{n1}(L_2)_{n2}$

Formula 1



wherein groups M, L_1 , L_2 , $n1$, and $n2$ are the same as they are
defined in the specification.

FIG. 1



**ORGANOMETALLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**

**CROSS-REFERENCE TO RELATED
APPLICATION**

[0001] This application claims priority to and the benefit of Korean Patent Application No. 10-2014-0182552, filed on Dec. 17, 2014, in the Korean Intellectual Property Office, the disclosure of which is incorporated herein in its entirety by reference.

BACKGROUND

[0002] 1. Field

[0003] The present disclosure relates to an organometallic compound and an organic light-emitting device including the same.

[0004] 2. Description of the Related Art

[0005] Organic light-emitting devices (OLEDs) are self-emission devices that have wide viewing angles, high contrast ratios, and short response times. In addition, the OLEDs exhibit excellent luminance, driving voltage, and response speed characteristics, and produce full-color images.

[0006] A typical organic light-emitting device includes an anode, a cathode, and an organic layer that is disposed between the anode and the cathode and includes an emission layer. A hole transport region may be disposed between the anode and the emission layer, and an electron transport region may be disposed between the emission layer and the cathode. Holes provided from the anode may move toward the emission layer through the hole transport region, and electrons provided from the cathode may move toward the emission layer through the electron transport region. The holes and electrons recombine in an emission layer and generate excitons. These excitons change from an excited state to a ground state to thereby generate light.


[0007] Various types of organic light emitting devices are known. However, there still remains a need in OLEDs having low driving voltage, high efficiency, high brightness, and long lifespan.

SUMMARY


[0008] Provided are an organometallic compound and an organic light-emitting device including the organometallic compound.

[0009] Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented exemplary embodiments.


[0010] According to an aspect of an exemplary embodiment, an organometallic compound is represented by Formula 1:




Formula 1



Formula 2A



Formula 2B



Formula 2C

[0011] wherein in Formula 1,

[0012] M is selected from Ir, Pt, Os, Ti, Zr, Hf, Eu, Tb, Tm, and Rh,

[0013] L₁ in Formula 1 is selected from ligands represented by Formula 2A,

[0014] L₂ in Formula 1 is selected from ligands represented by Formula 2B,

[0015] provided that L₁ and L₂ in Formula 1 are different from each other,

[0016] n₁ and n₂ in Formula 1 are each independently 1 or 2, and the sum of n₁ and n₂ is 2 or 3,

[0017] Y₁ and Y₂ in Formula 2A are each independently C or N, and Y₃ and Y₄ in Formula 2B are each independently C or N,

[0018] in Formulae 2A and 2B,

[0019] CY₁ and CY₃ are each independently a C₁-C₆₀ heterocyclic group,

[0020] CY₂ and CY₄ are each independently selected from a C₅-C₆₀ carbocyclic group and a C₁-C₆₀ heterocyclic group,

[0021] wherein CY₁ and CY₂ are optionally linked to each other through a first linking group, and, CY₃ and CY₄ are optionally linked to each other through a second linking group,

[0022] in Formulae 2A and 2B, Z₁ to Z₄ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, —SF₅, a hydroxyl group, a cyano group, a nitro

group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_1)(Q_2)$, $-B(Q_6)(Q_7)$, and $-P(=O)(Q_8)(Q_9)$.

[0023] a1 to a4 are each independently an integer selected from 0 to 4,

[0024] in Formula 2B, R_{11} and R_{12} are each independently selected from groups represented by Formula 2C,

[0025] b1 and b2 are each independently an integer selected from 0 to 3, and the sum of b1 and b2 is 1 or greater,

[0026] in Formula 2C, R_1 to R_3 are each independently selected from a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

[0027] when b1 is 1 or greater, at least one of R_1 to R_3 in Formula 2C is optionally bound to neighboring group Z_3 to form a saturated or unsaturated ring having two to ten carbons,

[0028] when b2 is 1 or greater, at least one of R_1 to R_3 in Formula 2C is optionally bound to neighboring group Z_4 to form a saturated or unsaturated ring having two to ten carbons,

[0029] * and *' in Formulae 2A and 2B are a binding site to M in Formula 1, and * in Formula 2C is a binding site to CY_3 or CY_4 in Formula 2B,

[0030] at least one of substituents of the substituted C_1 - C_{60} alkyl group, substituted C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group, substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10} cycloalkyl group, substituted C_1 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substituted C_1 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_1 - C_{60} heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from

[0031] a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0032] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_{11})(Q_{12})$, $-Si(Q_{13})(Q_{14})(Q_{15})$, $-B(Q_{16})(Q_{17})$, and $-P(=O)(Q_{18})(Q_{19})$;

[0033] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0034] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_{21})(Q_{22})$, $-Si(Q_{23})(Q_{24})(Q_{25})$, $-B(Q_{26})(Q_{27})$, and $-P(=O)(Q_{28})(Q_{29})$, and

[0035] $-N(Q_{31})(Q_{32})$, $-Si(Q_{33})(Q_{34})(Q_{35})$, $-B(Q_{36})(Q_{37})$, and $-P(=O)(Q_{38})(Q_{39})$,

[0036] wherein Q_1 , Q_2 , Q_6 to Q_9 , Q_{11} to Q_{19} , Q_{21} to Q_{29} , and Q_{31} to Q_{39} are each independently selected from a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted

C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0037] According to an aspect of another exemplary embodiment, an organic light-emitting device includes:

[0038] a first electrode;

[0039] a second electrode; and

[0040] an organic layer disposed between the first electrode and the second electrode,

[0041] wherein the organic layer includes an emission layer, and

[0042] wherein the organic layer includes at least one organometallic compound represented by Formula 1.

[0043] The organometallic compound may be included in the emission layer, the emission layer may further include a host, and an amount of the organometallic compound may be less than an amount of the host.

BRIEF DESCRIPTION OF THE DRAWINGS

[0044] These and/or other aspects will become apparent and more readily appreciated from the following description of the exemplary embodiments, taken in conjunction with the accompanying drawings in which:

[0045] FIG. 1 is a schematic cross-sectional view of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION

[0046] Reference will now be made in detail to exemplary embodiments, examples of which are illustrated in the accompanying drawings, wherein like reference numerals refer to like elements throughout. In this regard, the present exemplary embodiments may have different forms and should not be construed as being limited to the descriptions set forth herein. Accordingly, the exemplary embodiments are merely described below, by referring to the figures, to explain aspects. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

[0047] It will be understood that when an element is referred to as being “on” another element, it can be directly in contact with the other element or intervening elements may be present therebetween. In contrast, when an element is referred to as being “directly on” another element, there are no intervening elements present.

[0048] It will be understood that, although the terms first, second, third etc. may be used herein to describe various elements, components, regions, layers, and/or sections, these elements, components, regions, layers, and/or sections should not be limited by these terms. These terms are only used to distinguish one element, component, region, layer, or section from another element, component, region, layer, or

section. Thus, a first element, component, region, layer, or section discussed below could be termed a second element, component, region, layer, or section without departing from the teachings of the present embodiments.

[0049] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. As used herein, the singular forms “a,” “an,” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.

[0050] It will be further understood that the terms “comprises” and/or “comprising,” or “includes” and/or “including” when used in this specification, specify the presence of stated features, regions, integers, steps, operations, elements, and/or components, but do not preclude the presence or addition of one or more other features, regions, integers, steps, operations, elements, components, and/or groups thereof.

[0051] Spatially relative terms, such as “beneath,” “below,” “lower,” “above,” “upper” and the like, may be used herein for ease of description to describe one element or feature's relationship to another element(s) or feature(s) as illustrated in the figures. It will be understood that the spatially relative terms are intended to encompass different orientations of the device in use or operation in addition to the orientation depicted in the figures. For example, if the device in the figures is turned over, elements described as “below” or “beneath” other elements or features would then be oriented “above” the other elements or features. Thus, the exemplary term “below” can encompass both an orientation of above and below. The device may be otherwise oriented (rotated 90 degrees or at other orientations) and the spatially relative descriptors used herein interpreted accordingly.

[0052] “About” or “approximately” as used herein is inclusive of the stated value and means within an acceptable range of deviation for the particular value as determined by one of ordinary skill in the art, considering the measurement in question and the error associated with measurement of the particular quantity (i.e., the limitations of the measurement system). For example, “about” can mean within one or more standard deviations, or within $\pm 30\%$, 20%, 10%, 5% of the stated value.

[0053] Unless otherwise defined, all terms (including technical and scientific terms) used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. It will be further understood that terms, such as those defined in commonly used dictionaries, should be interpreted as having a meaning that is consistent with their meaning in the context of the relevant art and the present disclosure, and will not be interpreted in an idealized or overly formal sense unless expressly so defined herein.

[0054] Exemplary embodiments are described herein with reference to cross section illustrations that are schematic illustrations of idealized embodiments. As such, variations from the shapes of the illustrations as a result, for example, of manufacturing techniques and/or tolerances, are to be expected. Thus, embodiments described herein should not be construed as limited to the particular shapes of regions as illustrated herein but are to include deviations in shapes that result, for example, from manufacturing. For example, a region illustrated or described as flat may, typically, have rough and/or nonlinear features. Moreover, sharp angles that are illustrated may be rounded. Thus, the regions illustrated in the figures are schematic in nature and their shapes are not

intended to illustrate the precise shape of a region and are not intended to limit the scope of the present claims.


[0055] An organometallic compound represented by Formula 1 below is provided:




[0056] In Formula 1,

[0057] L_1 is selected from ligands that are each represented by Formula 2A, and


[0058] L_2 is selected from ligands that are each represented by Formula 2B:



Formula 2A



Formula 2B



Formula 2C

[0059] In Formulae 2A and 2B, * and *' are each a binding site to M in Formula 1.

[0060] In Formula 1, M may be selected from iridium (Ir), platinum (Pt), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), thulium (Tm), and rhodium (Rh).

[0061] For example, in Formula 1, M may be Ir or Pt.

[0062] In Formula 1, L_1 and L_2 are different from each other, n_1 and n_2 are each independently 1 or 2, and the sum of n_1 and n_2 is 2 or 3.

[0063] In Formula 2A, Y_1 and Y_2 may be each independently carbon (C) or nitrogen (N), and, in Formula 2B, Y_3 and Y_4 may be each independently C or N.

[0064] For example, in Formula 2A, Y_1 to Y_4 may be C, but embodiments are not limited thereto.

[0065] In Formulae 2A and 2B, CY_1 and CY_3 may be each independently a C_1 - C_{60} heterocyclic group; CY_2 and CY_4 may be each independently selected from a C_5 - C_{60} carbocyclic group and a C_1 - C_{60} heterocyclic group; CY_1 and CY_2 may be optionally bound to each other through a first linking group; and, CY_3 and CY_4 may be optionally bound to each other through a second linking group. The C_5 - C_{60} carbocyclic

group and C_1 - C_{60} heterocyclic group may be “a monocyclic group” or “a polycyclic group”.

[0066] In an embodiment, in Formulae 2A and 2B, CY_1 and CY_3 may be each independently selected from a pyridine ring, a pyrimidine ring, a pyrazine ring, a triazine ring, a quinoline ring, an isoquinoline ring, a quinazoline ring, a quinoxaline ring, a triazole ring, an imidazole ring, and a pyrazole ring; and CY_2 and CY_4 may be each independently selected from a benzene ring, a naphthalene ring, a pyridine ring, a pyrimidine ring, a pyrazine ring, a triazine ring, a quinoline ring, an isoquinoline ring, a quinazoline ring, a quinoxaline ring, a carbazole ring, a dibenzofuran ring, and a dibenzothiophene ring.


[0067] In another embodiment, in Formulae 2A and 2B, CY_1 and CY_3 may be each independently selected from a pyridine ring, a pyrimidine ring, a pyrazine ring, a triazine ring, a triazole ring, an imidazole ring, and a pyrazole ring; and CY_2 and CY_4 may be each independently selected from a benzene ring, a naphthalene ring, a pyridine ring, a pyrimidine ring, a pyrazine ring, a carbazole ring, a dibenzofuran ring, and a dibenzothiophene ring.

[0068] In some embodiments, in Formulae 2A and 2B, CY_1 and CY_3 may be each independently selected from a pyridine ring, a pyrimidine ring, a pyrazine ring, and a triazine ring; and CY_2 and CY_4 may be each independently selected from a benzene ring, a naphthalene ring, a carbazole ring, a dibenzofuran ring, and a dibenzothiophene ring, but embodiments are not limited thereto.

[0069] In Formula 2A, CY_1 and CY_2 may be optionally bound to each other through the first linking group; and, in Formula 2B, CY_3 and CY_4 may be optionally bound to each other through the second linking group. The first linking group and the second linking group may be each independently selected from linking groups represented by Formula 6:



[0070] In Formula 6, wherein Z_{31} is selected from $*-O-*'$, $*-S-*'$, $*-N(Q_{41})-*'$, $*-C(Q_{42})(Q_{43})-*'$, $*-C(Q_{44})=C(Q_{45})-*'$, and



[0071] Q_{41} to Q_{49} are each independently selected from

[0072] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0073] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

[0074] a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

[0075] b10 is an integer selected from 1 to 10, and when b10 is 2 or greater, groups Z_{31} are identical to or different from each other.


[0076] For example, in Formula 6, Q_{41} to Q_{49} are each independently selected from

[0077] a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0078] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

[0079] a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group, but embodiments are not limited therefrom.

[0080] For example, in Formula 2A, CY_1 and CY_2 are optionally bound to each other through the first linking group, and/or, in Formula 2B, CY_3 and CY_4 are optionally bound to each other through the second linking group, wherein the first linking group and the second linking group are each independently represented by $*-C(Q_{44})=C(Q_{45})-^{*}$ or



(i.e., b10=1 in Formula 6), wherein Q_{44} to Q_{49} are each independently selected from a hydrogen, a C_1 - C_{10} alkyl group, and a C_1 - C_{10} alkoxy group, embodiments are not limited thereto.

[0081] In Formulae 2A and 2B, Z_1 to Z_4 may be each independently selected from a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-SF_5$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or

unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_1)(Q_2)$, $-B(Q_6)(Q_7)$, and $-P(=O)(Q_8)(Q_9)$.

[0082] In an embodiment, in Formulae 2A and 2B, Z_1 to Z_4 are each independently

[0083] a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-SF_5$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0084] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{14} aryl group, a C_1 - C_{14} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0085] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{14} aryl group, a C_1 - C_{14} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0086] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{14} aryl group, a C_1 - C_{14} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{14} aryl group, a C_1 - C_{14} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

[0087] $-N(Q_1)(Q_2)$, $-B(Q_6)(Q_7)$, and $-P(=O)(Q_8)(Q_9)$;

[0088] wherein Q_1 , Q_2 and Q_6 to Q_9 may be each independently selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{14} aryl group, a C_6 - C_{14} aryl group substituted with at least one of a C_1 - C_{20} alkyl group and a C_6 - C_{14} aryl group, a C_1 - C_{14} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0089] In another embodiment, in Formulae 2A and 2B, Z_1 to Z_4 may be each independently selected from

[0090] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, —SF₅, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0091] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl (adamantyl) group, a norbornanyl (norbornyl) group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

[0092] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0093] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzo-

carbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

[0094] —B(Q₆)(Q₇) and —P(=O)(Q₈)(Q₉).

[0095] wherein Q₆ to Q₉ are each independently selected from

[0096] —CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;

[0097] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0098] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one of a deuterium, a C₁-C₁₀ alkyl group, and a phenyl group, embodiments are not limited thereto.

[0099] In some embodiments, in Formulae 2A and 2B, Z_1 to Z_4 are each independently selected from

[0100] a hydrogen, a deuterium, —F, a cyano group, a nitro group, —SF₅, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy

group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

[0101] a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group, each substituted with at least one selected from a deuterium, $-F$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

[0102] $-B(Q_6)(Q_7)$ and $-P(=O)(Q_8)(Q_9)$,

[0103] wherein Q_6 to Q_9 are each independently selected from

[0104] $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CH_2CH_3$, $-CH_2CD_3$, $-CH_2CD_2H$, $-CH_2CDH_2$, $-CHDCH_3$, $-CHDCD_2H$, $-CHDCDH_2$, $-CHDCD_3$, $-CD_2CD_3$, $-CD_2CD_2H$, and $-CD_2CDH_2$;

[0105] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0106] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from a deuterium, a C_1 - C_{10} alkyl group, and a phenyl group.

[0107] In some embodiments, in Formulae 2A and 2B, Z_1 to Z_4 may be each independently selected from a hydrogen, a deuterium, $-F$, a cyano group, a nitro group, $-SF_5$, $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a group represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-32:

-continued


Formula 9-3




Formula 9-4




Formula 9-5




Formula 9-6




Formula 9-7




Formula 9-8




Formula 9-9




Formula 9-10




Formula 9-11




Formula 9-12




Formula 9-13




Formula 9-14




Formula 9-15




Formula 9-16



Formula 9-17




Formula 10-1















Formula 9-1




Formula 9-2




-continued




Formula 10-2




Formula 10-3




Formula 10-4




Formula 10-5




Formula 10-6



Formula 10-7




Formula 10-8




Formula 10-9

Formula 10-10




Formula 10-11

Formula 10-12



Formula 10-13



-continued

Formula 10-14

Formula 10-15

Formula 10-16

Formula 10-17

Formula 10-18

Formula 10-19


Formula 10-20

Formula 10-21


Formula 10-22

Formula 10-23


-continued




Formula 10-24




Formula 10-25




Formula 10-26




Formula 10-27




Formula 10-28




Formula 10-29



Formula 10-30



Formula 10-31



Formula 10-32

[0108] In Formulae 2A and 2B, a₁ to a₄ may be each independently an integer selected from 0 to 4. When a₁ is 2 or greater, groups Z₁ may be identical to or different from each other; when a₂ is 2 or greater, groups Z₂ may be identical to or

different from each other; when a₃ is 2 or greater, groups Z₃ may be identical to or different from each other; and when a₄ is 2 or greater, groups Z₄ may be identical to or different from each other.

[0109] In Formula 2B, R₁₁ and R₁₂ are each independently a group represented by Formula 2C, b₁ and b₂ are each independently an integer selected from 0 to 3, and the sum of b₁ and b₂ is 1 or greater. That is, the ligand represented by Formula 2B includes at least one of groups represented by Formula 2C as a substituent.

[0110] In an embodiment, in Formula 2B, b₁ may be 1 or 2, and b₂ may be 0.

[0111] In another embodiment, in Formula 2B, b₁ may be 1, and b₂ may be 1.

[0112] In another embodiment, in Formula 2B, b₁ may be 1, and b₂ may be 0.

[0113] In Formula 2C, R₁ to R₃ may be each independently selected from a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0114] In an embodiment, in Formula 2C, R₁ to R₃ may be each independently selected from

[0115] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group;

[0116] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0117] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

[0118] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an

amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₁₄ aryl group, a C₁-C₁₄ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0119] In another embodiment, in Formula 2C, R₁ to R₃ may be each independently selected from

[0120] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group;

[0121] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

[0122] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

[0123] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a ben-

zoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.

[0124] In some embodiments, in Formula 2C, R₁ to R₃ may be each independently selected from

[0125] a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

[0126] a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group,

a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group, each substituted with at least one selected from a deuterium, $-F$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group, but embodiments are not limited thereto.

[0127] In some embodiments, in Formula 2C, R_1 to R_3 may be each independently selected from $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, the group represented by Formulae 9-1 to 9-17, and the group represented by Formulae 10-1 to 10-18.

[0128] In an embodiment, in Formula 2C, R_1 to R_3 may be identical to each other.

[0129] In another embodiment, in Formula 2C, R_1 and R_2 are identical to each other and R_3 is different from R_1 ; or R_1 to R_3 may be all different from one another.

[0130] In an embodiment, in Formula 2C, R_1 to R_3 may be all a methyl group, an ethyl group, or $-CD_3$.

[0131] In some embodiments, in Formula 2C, R_1 and R_2 may be a methyl group, and R_3 may be a phenyl group.


[0132] In Formula 2B, when b_1 is 1 or greater, at least one of R_1 to R_3 in Formula 2C may be optionally bound to a neighboring group Z_3 to form a saturated or unsaturated ring having two to ten carbons (e.g., see Formulae 2B(4) to 2B(10)).

[0133] When b_2 is 1 or greater, at least one of R_1 to R_3 in Formula 2C may be optionally bound to a neighboring group Z_4 to form a saturated or unsaturated ring having two to ten carbons.


[0134] In some embodiments, in Formula 1, L_1 may be selected from Formulae 3-1 to 3-110:

-continued


Formula 3-3




Formula 3-4




Formula 3-5




Formula 3-6




Formula 3-7




Formula 3-1




Formula 3-2




-continued




Formula 3-8




Formula 3-9



Formula 3-10




Formula 3-11




Formula 3-12


-continued




Formula 3-13




Formula 3-14



Formula 3-15




Formula 3-16




Formula 3-17


-continued




Formula 3-18




Formula 3-19



Formula 3-20




Formula 3-21




Formula 3-22


-continued




Formula 3-23




Formula 3-24



Formula 3-25




Formula 3-26




Formula 3-27


-continued




Formula 3-28




Formula 3-29



Formula 3-30




Formula 3-31




Formula 3-32


-continued




Formula 3-33




Formula 3-34



Formula 3-35








Formula 3-36



Formula 3-37

-continued



Formula 3-38






Formula 3-39

Formula 3-40

Formula 3-41

Formula 3-42

-continued



Formula 3-43


Formula 3-44

Formula 3-45


Formula 3-46

Formula 3-47


-continued




Formula 3-48




Formula 3-49



Formula 3-50




Formula 3-51




Formula 3-52


-continued




Formula 3-53




Formula 3-54



Formula 3-55








Formula 3-56



Formula 3-57

-continued



Formula 3-58






Formula 3-59

Formula 3-60

Formula 3-61

Formula 3-62

-continued



Formula 3-63


Formula 3-64

Formula 3-65


Formula 3-66

Formula 3-67


-continued




Formula 3-68




Formula 3-69



Formula 3-70




Formula 3-71




Formula 3-72


-continued




Formula 3-73




Formula 3-74



Formula 3-75




Formula 3-76




Formula 3-77


-continued




Formula 3-78




Formula 3-79



Formula 3-80




Formula 3-81




Formula 3-82


-continued




Formula 3-83




Formula 3-84



Formula 3-85




Formula 3-86




Formula 3-87

-continued


Formula 3-93




Formula 3-94




Formula 3-95




Formula 3-96




Formula 3-97




-continued




Formula 3-98




Formula 3-99



Formula 3-100




Formula 3-101




Formula 3-102


-continued




Formula 3-103




Formula 3-104



Formula 3-105




Formula 3-106




Formula 3-107

-continued


Formula 3-108



Formula 3-109



Formula 3-110



[0135] In Formulae 3-1 to 3-110, Z_1 and Z_2 are the same as defined herein;

[0136] descriptions of Z_{1a} and Z_{1b} are each independently the same as defined in connection with Z_1 ;

[0137] descriptions of Z_{2a} , Z_{2b} , and Z_{2c} each independently the same as defined in connection with Z_2 ;

[0138] aa2 and ab2 are each independently 1 or 2;

[0139] aa3 and ab3 are each independently an integer selected from 1 to 3;

[0140] aa4 and ab4 are each independently an integer selected from 1 to 4; and

[0141] * and *' are each independently a binding site to M.

[0142] In some embodiments, in Formulae 3-1 to 3-110,

[0143] Z_1 , Z_2 , Z_{1a} , Z_{1b} , Z_{2a} , Z_{2b} , and Z_{2c} may be each independently selected from

[0144] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, —SF₃, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0145] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃,

—CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

[0146] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0147] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cyclo-

heptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluo-ranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnoli-nyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranlyl group, a ben-zothiophenyl group, an isobenzothiazolyl group, a benzox-azolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzo-carbazolyl group, a dibenzocarbazolyl group, an imidazopy-ridinyl group, and an imidazopyrimidinyl group; and

[0148] $-\text{N}(\text{Q}_1)(\text{Q}_2)$, $-\text{B}(\text{Q}_6)(\text{Q}_7)$, and $-\text{P}(=\text{O})(\text{Q}_8)(\text{Q}_9)$,

[0149] wherein Q_1 , Q_2 and Q_6 to Q_9 are each independently selected from

[0150] $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCH}_3$, $-\text{CHDCD}_2\text{H}$, $-\text{CHDCDH}_2$, $-\text{CHDCD}_3$, $-\text{CD}_2\text{CD}_3$, $-\text{CD}_2\text{CD}_2\text{H}$, and $-\text{CD}_2\text{CDH}_2$;

[0151] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0152] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one of a deuterium and a C_1 - C_{10} alkyl group,


[0153] aa2 and ab2 are each independently 1 or 2,

[0154] aa3 and ab3 are each independently an integer selected from 1 to 3,

[0155] aa4 and ab4 are each independently an integer selected from 1 to 4, and


[0156] * and *' may be a binding site to M in Formula 1.

[0157] In some embodiments, in Formula 1, L_1 may be selected from ligands represented by Formulae 3-1(1) to 3-1(59):




Formula 3-1(1)


-continued




Formula 3-1(2)




Formula 3-1(3)




Formula 3-1(4)



Formula 3-1(5)









Formula 3-1(6)



Formula 3-1(7)

-continued



Formula 3-1(8)

Formula 3-1(9)







Formula 3-1(10)

Formula 3-1(11)

Formula 3-1(12)

Formula 3-1(13)

-continued



Formula 3-1(14)

Formula 3-1(15)






Formula 3-1(16)

Formula 3-1(17)

Formula 3-1(18)

Formula 3-1(19)

-continued



Formula 3-1(20)






Formula 3-1(21)

Formula 3-1(22)

Formula 3-1(23)

Formula 3-1(24)

-continued



Formula 3-1(25)







Formula 3-1(26)

Formula 3-1(27)

Formula 3-1(28)

Formula 3-1(29)

-continued



Formula 3-1(30)

Formula 3-1(31)






Formula 3-1(32)

Formula 3-1(33)

Formula 3-1(34)

Formula 3-1(35)

-continued



Formula 3-1(36)


Formula 3-1(37)

Formula 3-1(38)


Formula 3-1(39)

Formula 3-1(40)


-continued




Formula 3-1(41)




Formula 3-1(42)



Formula 3-1(43)




Formula 3-1(44)




Formula 3-1(45)


-continued




Formula 3-1(46)




Formula 3-1(47)



Formula 3-1(48)




Formula 3-1(49)




Formula 3-1(50)

-continued




Formula 3-1(51)

-continued




Formula 3-1(55)


Formula 3-1(52)




Formula 3-1(56)




Formula 3-1(53)




Formula 3-1(57)




Formula 3-1(54)



Formula 3-1(58)



-continued




Formula 3-1(59)


[0158] In Formulae 3-1(1) to 3-1(59),

[0159] Z_1 , Z_2 , Z_{1a} , Z_{1b} , Z_{1c} , Z_{1d} , Z_{2a} , Z_{2b} , Z_{2c} , and Z_{2d} may be each independently selected from a deuterium, $-F$, a cyano group, a nitro group, $-SF_5$, $-CH_3$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, the group represented by one of Formulae 9-1 to 9-17, and the group represented by one of Formulae 10-1 to 10-32.


[0160] In some embodiments, L_2 of Formula 1 may be selected from ligands represented by Formulae 2B(1) to 2B(10):




Formula 2B(1)




Formula 2B(2)




Formula 2B(3)




Formula 2B(4)




Formula 2B(5)



Formula 2B(6)




Formula 2B(7)




Formula 2B(8)

-continued




Formula 2B(9)




Formula 2B(10)

[0161] In Formulae 2B(1) to 2B(10), descriptions of Z₃, Z₄, a₃, a₄, and R₁ to R₃ are the same as defined herein, CY₄ is selected from a benzene, a naphthalene, a carbazole, a dibenzofuran, and a dibenzothiophene, description of Z₅ is the same as defined in connection with Z₃, and a₅ is an integer selected from 1 to 6.

[0162] In some embodiments, L₂ of Formula 1 may be selected from ligands represented by Formulae 2B-1 to 2B-60:




Formula 2B-1




Formula 2B-2


-continued




Formula 2B-3



Formula 2B-4




Formula 2B-5




Formula 2B-6


-continued




Formula 2B-7



Formula 2B-8




Formula 2B-9




Formula 2B-10


-continued




Formula 2B-11



Formula 2B-12




Formula 2B-13




Formula 2B-14

-continued




Formula 2B-15


-continued




Formula 2B-19




Formula 2B-16




Formula 2B-20




Formula 2B-17



Formula 2B-21




Formula 2B-18




Formula 2B-22

-continued




Formula 2B-23


-continued




Formula 2B-27




Formula 2B-24




Formula 2B-28



Formula 2B-25








Formula 2B-29




Formula 2B-26

-continued




Formula 2B-30

-continued




Formula 2B-35

Formula 2B-31




Formula 2B-36

Formula 2B-32



Formula 2B-37


Formula 2B-33




Formula 2B-38

Formula 2B-34


-continued




Formula 2B-39



Formula 2B-40




Formula 2B-41




Formula 2B-42


-continued




Formula 2B-43



Formula 2B-44




Formula 2B-45




Formula 2B-46


-continued




Formula 2B-47




Formula 2B-48



Formula 2B-49




Formula 2B-50




Formula 2B-51


-continued




Formula 2B-52



Formula 2B-53




Formula 2B-54




Formula 2B-55


-continued




Formula 2B-56



Formula 2B-57




Formula 2B-58



Formula 2B-59

-continued



Formula 2B-60

[0163] In Formulae 2B-1 to 2B-60,**[0164]** R_1 to R_3 may be each independently selected from

[0165] a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

[0166] a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group, each substituted with at least one selected from a deuterium, $-F$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group,

[0167] Z_3 , Z_{3a} , Z_{3b} , Z_4 , Z_{4a} , Z_{4b} , Z_{4c} , and Z_{4d} are each independently selected from

[0168] a deuterium, —F, a cyano group, a nitro group, —SF₅, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

[0169] a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

[0170] —B(Q₆)(Q₇) and —P(=O)(Q₈)(Q₉),

[0171] wherein Q₆ to Q₉ are each independently selected from

[0172] —CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;

[0173] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

[0174] an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, and a phenyl group, but embodiments are not limited thereto.

[0175] In another embodiment, in Formulae 2B-1 to 2B-60,

[0176] R₁ to R₃ may be each independently selected from —CH₃, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, the group represented by one of Formulae 9-1 to 9-17, and the group represented by one of Formulae 10-1 to 10-18,

[0177] Z_3 , Z_{3a} , Z_{3b} , Z_4 , Z_{4a} , Z_{4b} , Z_{4c} , and Z_{4d} may be each independently selected from a deuterium, —F, a cyano group, a nitro group, —SF₅, —CH₃, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, the group represented by one of Formulae 9-1 to 9-17, and the group represented by one of Formulae 10-1 to 10-32, but embodiments are not limited thereto.

[0178] In Formula 1, n₁ and n₂ are each independently 1 or 2, and n₁+n₂ is 2 or 3.

[0179] In some embodiments, M may be Ir, and n₁+n₂=3; or M may be Pt, and n₁+n₂=2, but embodiments are not limited thereto.

[0180] In another embodiment, in Formula 1, n₂ may be 1.

[0181] The organometallic compound represented by Formula 1 does not have a form of a salt that is formed of an ion pair, and the organometallic compound may be neutral.

[0182] In an embodiment, in Formula 1,

[0183] M is Ir, and n₁+n₂=3; or M is Pt, and n₁+n₂=2,

[0184] L₁ is a ligand represented by one of Formulae 3-1 and 3-110 (e.g., a ligand represented by one of Formula 3-1(1) to 3-1(59)),

[0185] L₂ is a ligand represented by one of Formulae 2B(1) to 2B(10),

[0186] the organometallic compound represented by Formula 1 may be neutral, but embodiments are not limited thereto.

[0187] In another embodiment, in Formula 1,


[0188] M is Ir, and n₁+n₂=3; or M is Pt, and n₁+n₂=2,

[0189] L₁ is a ligand represented by one of Formulae 3-1 and 3-110 (e.g., a ligand represented by one of Formula 3-1(1) to 3-1(59)),


[0190] L₂ is a ligand represented by one of Formulae 2B-1 to 2B-60, and

[0191] the organometallic compound represented by Formula 1 may be neutral, but embodiments are not limited thereto.

[0192] The organometallic compound may be one of Compounds 1 to 109, but embodiments are not limited thereto:




-continued




2

-continued




6


3




7




4




8




5










9




-continued




-continued




-continued




-continued




19




23




20




24




21



25




-continued




26


-continued




30




27




31




28



32




29




33

-continued




34


-continued




38




35




39




36



40




37




41

44


-continued




-continued




43




47




44




48




45



49




-continued




50


-continued




54




51




55




52



56







53







57


-continued




-continued




-continued




-continued



67





71





72

68




73








69




-continued



-continued




-continued




82


-continued




86




83




87




84



88




85




89

-continued




90


-continued




94




91




95




92



96




93




97


-continued




-continued




99




103




100




104




101



105



-continued



[0193] In Formula 1, n₁ and n₂ may be 1 or 2. That is, n₁ is not 0, and thus the organometallic compound represented by Formula 1 essentially includes the ligand represented by Formula 2A. Also, the ligand represented by Formula 2A is a

N—C bidentate ligand that is linked to a metal M of Formula 1 through carbon and nitrogen. Thus, the organometallic compound represented by Formula 1 may have excellent thermal stability.

[0194] Also, the organometallic compound represented by Formula 1 has the ligand represented by Formula 2B, and the ligand represented by Formula 2B essentially has a group represented by Formula 2C as a substituent. The group represented by Formula 2C includes “Ge”, and thus an electric device, for example, an organic light-emitting device, including the organometallic compound represented by Formula 1 may have a high efficiency.

[0195] For example, the results of HOMO, LUMO, and triplet (T₁) energy levels of some compounds among the organometallic compounds evaluated by using a DFT method of Gaussian program (geometry optimized at the B3LYP, 6-31G (d,p) level) are shown in Table 1.

TABLE 1

Compound No.	HOMO (eV)	LUMO (eV)	T ₁ Energy level (eV)
1	-4.822	-1.242	2.567
2	-4.799	-1.160	2.623
3	-4.822	-1.242	2.567
4	-4.760	-1.142	2.644
5	-4.799	-1.160	2.623
6	-4.793	-1.156	2.622

[0196] Referring to Table 1, it may be confirmed that the compound is suitable to be used as a material for an organic light-emitting device.

[0197] A method of synthesizing the organometallic compound represented by Formula 1 may be understood to one of ordinary skill in the art by referring to Synthesis Examples used herein.

[0198] Therefore, the organometallic compound represented by Formula 1 is suitable as, for example, a dopant in an emission layer in an organic layer of an organic light-emitting device, and, according to another aspect of an embodiment, provided is an organic light-emitting device including:

[0199] a first electrode;

[0200] a second electrode; and

[0201] an organic layer disposed between the first electrode and the second electrode,

[0202] wherein the organic layer includes at least one of the organometallic compounds represented by Formula 1.

[0203] The organic light-emitting device includes the organic layer including the organometallic compound represented by Formula 1 and thus may have low driving voltage, high efficiency, high electrical power, high quantum efficiency, long lifespan, and excellent color purity.

[0204] The organometallic compound represented by Formula 1 may be used between a pair of electrodes in an organic light-emitting device. For example, the organometallic compound represented by Formula 1 may be included in the emission layer. Here, the organometallic compound serves as a dopant, and the emission layer may further include a host (that is, an amount of the organometallic compound represented by Formula 1 is smaller than an amount of the host).

[0205] As used herein, the term “(organic layer) including at least one organometallic compound” denotes “(organic layer) including one of the organometallic compounds of Formula 1 above, or at least two different organometallic compounds of Formula 1 above.”

[0206] For example, the organic layer may include only Compound 1 as the organometallic compound. In this regard, Compound 1 may be included in the emission layer of the organic light-emitting device. Alternatively, the organic layer may include Compound 1 and Compound 2 as the organometallic compounds. In this regard, Compound 1 and Compound 2 may be included in the same layer (for example, both Compound 1 and Compound 2 may be included in the emission layer).

[0207] The first electrode may be an anode, which is a hole injection electrode, and the second electrode may be a cathode, which is an electron injection electrode. Alternatively, the first electrode may be a cathode, which is an electron injection electrode, and the second electrode may be an anode, which is a hole injection electrode.

[0208] For example, the first electrode may be an anode, the second electrode may be a cathode, and the organic layer may include:

[0209] i) a hole transport region disposed between the first electrode and the emission layer, wherein the hole transport region may include at least one selected from a hole injection layer, a hole transport layer, and an electron blocking layer; and

[0210] ii) an electron transport region disposed between the emission layer and the second electrode, wherein the electron transport region may include at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

[0211] As used herein, the term "organic layer" refers to a single and/or a plurality of layers disposed between the first electrode and the second electrode in an organic light-emitting device. The "organic layer" may include not only organic compounds but also organometallic complexes including metals.

[0212] FIG. 1 is a schematic view of an organic light-emitting device 10 according to an embodiment. Hereinafter, a structure and a method of manufacturing the organic light-emitting device according to an embodiment will be described with reference to FIG. 1. The organic light-emitting device 10 includes a first electrode 11, an organic layer 15, and a second electrode 19, which are sequentially layered in the stated order.

[0213] A substrate may be additionally disposed under the first electrode 11 or on the second electrode 19. The substrate may be a conventional substrate that is used in an organic light-emitting device, such as glass substrate or a transparent plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water repellency.

[0214] The first electrode 11 may be formed by vacuum-depositing or sputtering a material for forming the first electrode on the substrate. The first electrode 11 may be an anode. The material for the first electrode 11 may be selected from materials with a high work function for an easy hole injection. The first electrode 11 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. Examples of the material for the first electrode 11 may include indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), and zinc oxide (ZnO). Alternatively, a metal such as magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), cal-

cium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag) may be used.

[0215] The first electrode 11 may have a single layer structure or a multi-layer structure including a plurality of layers. For example, the first electrode 11 may have a triple-layer structure of ITO/Ag/ITO, but embodiments are not limited thereto.

[0216] The organic layer 15 is disposed on the first electrode 11.

[0217] The organic layer 15 may include a hole transport region, an emission layer, and an electron transport region.

[0218] The hole transport region may be disposed between the first electrode 11 and the emission layer.

[0219] The hole transport region may include at least one of a hole injection layer, a hole transport layer, an electron blocking layer, and a buffer layer.

[0220] The hole transport region may only include a hole injection layer or a hole transport layer. Alternatively, the hole transport region may include a structure in which a hole injection layer/a hole transport layer or a hole injection layer/a hole transport layer/an electron blocking layer are sequentially layered on the first electrode 11.

[0221] When the hole transport region includes a hole injection layer, the hole injection layer (HIL) may be formed on the first electrode 11 by using various methods such as vacuum-deposition, spin coating, casting, and Langmuir-Blodgett (LB) method.







[0222] When the hole injection layer is formed using a vacuum deposition, vacuum deposition conditions may vary according to the compound that is used to form the hole injection layer, and the desired structure and thermal properties of the hole injection layer to be formed. For example, vacuum deposition may be performed at a temperature in a range of about 100° C. to about 500° C., a vacuum degree in a range of about 10⁻⁸ torr to about 10⁻³ torr, and a deposition rate in a range of about 0.01 Angstroms per second (Å/sec) to about 100 Å/sec. However, the deposition conditions are not limited thereto.

[0223] When the hole injection layer is formed using spin coating, the coating conditions may vary according to the compound that is used to form the hole injection layer, and the desired structure and thermal properties of the hole injection layer to be formed. For example, the coating rate may be in a range of about 2,000 revolutions per minute (rpm) to about 5,000 rpm, and a temperature at which heat treatment is performed to remove a solvent after coating may be in a range of about 80° C. to about 200° C. However, the coating conditions are not limited thereto.


[0224] The conditions for forming a hole transport layer and an electron blocking layer may be inferred based on the conditions for forming the hole injection layer.

[0225] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB, β-NPB, TPD, a spiro-TPD, a spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/


DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), (polyaniline)/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201 below, and a compound represented by Formula 202 below:




-continued




Spiro-TPD




Spiro-NPB



methylated NPB




TAPC




HMTPD

-continued

Formula 201



Formula 202



[0226] In Formula 201, Ar₁₀₁ and Ar₁₀₂ may be each independently selected from

[0227] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an acenaphthylenylene group, a fluorenylenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, and a pentacenylene group; and

[0228] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an acenaphthylenylene group, a fluorenylenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, and a pentacenylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group

or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0229] In Formula 201, xa and xb may be each independently an integer selected from 0 to 5, or 0, 1, or 2. For example, xa may be 1, and xb may be 0, but embodiments are not limited thereto.

[0230] In Formulae 201 and 202, R₁₀₁ to R₁₀₈, R₁₁₁ to R₁₁₉, and R₁₂₁ to R₁₂₄ may be each independently selected from

[0231] a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group (e.g., a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, or a hexyl group), and a C₁-C₁₀ alkoxy group (e.g., a methoxy group, an ethoxy group, a propoxy group, a butoxy group, or a pentoxy group);

[0232] a C₁-C₁₀ alkyl group and a C₁-C₁₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, and a phosphoric acid group or a salt thereof;

[0233] a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, and a pyrenyl group; and

[0234] a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, and a pyrenyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, and a C₁-C₁₀ alkoxy group, but embodiments are not limited thereto.


[0235] In Formula 201, R₁₀₉ may be selected from

[0236] a phenyl group, a naphthyl group, an anthracenyl group and a pyridinyl group; and

[0237] a phenyl group, a naphthyl group, an anthracenyl group, and a pyridinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, and a pyridinyl group, but embodiments are not limited thereto.

[0238] In an embodiment, the compound represented by Formula 201 may be represented by Formula 201A, but embodiments are not limited thereto:


Formula 201A



[0239] In Formula 201A, descriptions of R₁₀₁, R₁₁₁, R₁₁₂, and R₁₀₉ are the same as provided herein.


[0240] For example, the compound represented by Formula 201 and the compound represented by Formula 202 may include Compounds HT1 to HT20, but they are not limited thereto:

HT1




-continued

HT2




-continued


HT4



HT3




HT5




-continued

HT6




-continued


HT8




HT9



HT7




HT10




-continued


HT11



HT12




HT13




-continued


HT14




HT15



HT16




HT17




-continued


HT18



HT19



HT20




[0241] A thickness of the hole transport region may be in a range of about 100 Angstroms (Å) to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes a hole injection layer and a hole transport layer, the thickness of the hole injection layer may be in a range of about 100 Å to about 10,000 Å, and for example, about 100 Å to about 1,000 Å, and the thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, and for example, about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are within these ranges, excellent hole transport characteristics may be obtained without a substantial increase in driving voltage.

[0242] The hole transport region may further include, in addition to the mentioned materials above, a charge-generating material to improve conductive properties. The charge-

generating material may be homogeneously or non-homogeneously dispersed throughout the hole transport region.

[0243] The charge-generating material may be, for example, a p-dopant. The p-dopant may be one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but embodiments are not limited thereto. For example, non-limiting examples of the p-dopant are a quinone derivative, such as tetracyanoquinonedimethane (TCNQ) or 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinodimethane (F4-TCNQ); a metal oxide, such as a tungsten oxide or a molybdenum oxide; and a compound containing a cyano group, such as Compound HT-D1 illustrated below, but they are not limited thereto.

Compound HT-D1F4-TCNQ



[0244] The hole transport region may further include a buffer layer.


[0245] The buffer layer may compensate for an optical resonance distance according to a wavelength of light emitted from the emission layer, and thus may increase efficiency.

[0246] The emission layer may be formed on the hole transport region by using various methods, such as vacuum-deposition, spin coating, casting, or an LB method. When the emission layer is formed by vacuum-deposition or spin coating, vacuum-deposition and coating conditions for the emission layer may be generally similar to the conditions for forming a hole injection layer, though the conditions may vary depending on the compound used.


[0247] When the hole transport region include an electron blocking layer, a material for the electron blocking layer may be selected from a material that may be used in the hole transport region and a material for a host, but embodiments are not limited thereto. For example, when the hole transport region includes the electron blocking layer, the material for the electron blocking layer may be mCP.

[0248] The emission layer may include a host and a dopant, and the dopant includes the organometallic compound represented by Formula 1.


[0249] The host may include at least one of TPBi, TBADN, ADN (also, referred to as "DNA"), CBP, CDBP, TCP, mCP, Compound H50, and Compound H51:




TPBi




TBADN



ADN




CBP




CDBP

-continued




TCP




mCP

Compound H50




Compound H51



[0250] Alternatively, the host may further include a compound represented by Formula 301:

Formula 301



[0251] In Formula 301, Ar_{111} and Ar_{112} may be each independently selected from

[0252] a phenylene group, a naphthylene group, a phenanthrenylene group, and a pyrenylene group; and

[0253] a phenylene group, a naphthylene group, a phenanthrenylene group, and a pyrenylene group, each substituted with at least one selected from a phenyl group, a naphthyl group, and an anthracenyl group.

[0254] In Formula 301, Ar_{113} to Ar_{116} may be each independently selected from

[0255] a C_1 - C_{10} alkyl group, a phenyl group, a naphthyl group, a phenanthrenyl group, and a pyrenyl group; and

[0256] a phenyl group, a naphthyl group, a phenanthrenyl group, and a pyrenyl group, each substituted with at least one selected from a phenyl group, a naphthyl group, and an anthracenyl group.


[0257] In Formula 301, g, h, i, and j may be each independently an integer selected from 0 to 4, for example, 0, 1, or 2.

[0258] In Formula 301, Ar_{113} to Ar_{116} may be each independently selected from

[0259] a C_1 - C_{10} alkyl group substituted with at least one of a phenyl group, a naphthyl group, and an anthracenyl group;


[0260] a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group;

[0261] a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group; and



but embodiments are not limited thereto.

[0262] Alternatively, the host may include a compound represented by Formula 302:




Formula 302

[0263] In Formula 302, descriptions of Ar_{122} to Ar_{125} are the same as defined in connection with Ar_{113} of Formula 301.


[0264] In Formula 302, Ar_{126} and Ar_{127} may be each independently a C_1 - C_{10} alkyl group (e.g., a methyl group, an ethyl group, or a propyl group).

[0265] In Formula 302, k and l may be each independently an integer selected from 0 to 4. For example, k and l may be 0, 1, or 2.


[0266] The compound represented by Formula 301 and the compound represented by Formula 302 may include Compounds H1 to H42, but embodiments are not limited thereto:




H1




H2



H3




H4




H5


-continued




H6




H7



H8




H9




H10


-continued




H11




H12




H13



H14







H15








H16

-continued




-continued




-continued


H26




H27




H28



H29




H30




-continued


H31




H32




H33



H34




-continued




H35

-continued




H38


H36




H39



H37




H40




-continued

H41



H42



[0267] When the organic light-emitting device is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, and a blue emission layer. Alternatively, the emission layer may have a structure in which the red emission layer, the green emission layer, and/or the blue emission layer are layered to emit white light or other various embodiments are possible.

[0268] When the emission layer includes a host and a dopant, an amount of the dopant may be in a range of about 0.01 parts to about 15 parts by weight based on 100 parts by weight of the host, but embodiments are not limited thereto.

[0269] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.


[0270] Then, an electron transport region may be disposed on the emission layer.

[0271] The electron transport region may include at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer, but is not limited thereto.


[0272] For example, the electron transport region may have a structure of a hole blocking layer/an electron transport layer/an electron injection layer or an electron transport layer/an electron injection layer, but it is not limited thereto. The electron transport layer may have a single layer structure or a multi-layer structure including two or more different materials.

[0273] The conditions for forming a hole blocking layer, an electron transport layer, and an electron injection layer may be inferred based on the conditions for forming the hole injection layer.

[0274] When the electron transport region includes a hole blocking layer, the hole blocking layer may include, for example, at least one of BCP, Bphen and Balq, but embodiments are not limited thereto:




BCP




Bphen

[0275] A thickness of the hole blocking layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thickness of the hole blocking layer is within this range, excellent hole blocking characteristics may be obtained without a substantial increase in driving voltage.


[0276] The electron transport layer may further include at least one of BCP and Bphen above and Alq₃, Balq, TAZ, and NTAZ below:

Alq₃


-continued



BAIq




TAZ



NTAZ


[0277] Alternatively, the electron transport layer may include at least one selected from Compounds ET1 and ET2, but it is not limited thereto.

ET1



-continued

ET2




[0278] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within this range, excellent electron transport characteristics may be obtained without a substantial increase in driving voltage.


[0279] The electron transport layer may further include a metal-containing material in addition to the materials described above.

[0280] The metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.

ET-D1



ET-D2



[0281] The electron transport region may include an electron injection layer (EIL) that facilitates electron injection from the second electrode 19.

[0282] The electron injection layer may include at least one selected from LiF, NaCl, CsF, Li₂O, and BaO.

[0283] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection

layer is within this range, excellent electron injection characteristics may be obtained without a substantial increase in driving voltage.

[0284] The second electrode **19** is disposed on the organic layer **15**. The second electrode **19** may be a cathode. A material for the second electrode **19** may be a material having a relatively low work function, such as a metal, an alloy, an electrically conductive compound, and a mixture thereof. Detailed examples of the material for forming the second electrode **19** are lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), and magnesium-silver (Mg—Ag). Alternatively, ITO or IZO may be used to form a transmissive second electrode **19** to manufacture a top emission light-emitting device, and such a variation may be possible.

[0285] Hereinbefore, the organic light-emitting device has been described with reference to FIG. 1, but embodiments are not limited thereto.

[0286] A C_1 - C_{60} alkyl group as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms. Detailed examples thereof are a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. A C_1 - C_{60} alkenylene group as used herein refers to a divalent group having the same structure as a C_1 - C_{60} alkyl group.

[0287] A C_1 - C_{60} alkoxy group as used herein refers to a monovalent group represented by $—OA_{101}$ (wherein A_{101} is the C_1 - C_{60} alkyl group). Detailed examples thereof are a methoxy group, an ethoxy group, and an isopropoxy group.

[0288] A C_2 - C_{60} alkenyl group as used herein refers to a hydrocarbon aliphatic group formed by including at least one carbon-carbon double bond in the middle or at the terminal of the C_2 - C_{60} alkyl group as defined above. Detailed examples thereof are an ethenyl group, a propenyl group, and a butenyl group. A C_2 - C_{60} alkenylene group as used herein refers to a divalent group having the same structure as a C_2 - C_{60} alkenyl group.

[0289] A C_2 - C_{60} alkynyl group as used herein refers to a hydrocarbon group formed by including at least one carbon-carbon triple bond in the middle or at the terminal of the C_2 - C_{60} alkyl group as defined above. Detailed examples thereof are an ethynyl group and a propynyl group. A C_2 - C_{60} alkynylene group as used herein refers to a divalent group having the same structure as a C_2 - C_{60} alkynyl group.

[0290] A C_3 - C_{10} cycloalkyl group as used herein refers to a monovalent monocyclic saturated hydrocarbon group including 3 to 10 carbon atoms. Detailed examples thereof are a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. A C_3 - C_{10} cycloalkylene group as used herein refers to a divalent group having the same structure as a C_3 - C_{10} cycloalkyl group.

[0291] A C_1 - C_{10} heterocycloalkyl group as used herein refers to a monovalent monocyclic saturated group including at least one hetero atom selected from N, O, P, and S as a ring-forming atom and 1 to 10 carbon atoms. Detailed examples thereof are a tetrahydrofuran group and a tetrahydrothiophenyl group. A C_1 - C_{10} heterocycloalkylene group as used herein refers to a divalent group having the same structure as a C_1 - C_{10} heterocycloalkyl group.

[0292] A C_3 - C_{10} cycloalkenyl group as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one double bond in its ring, and which is not aromatic. Detailed examples thereof are a cyclopentenyl

group, a cyclohexenyl group, and a cycloheptenyl group. A C_3 - C_{10} cycloalkenylene group as used herein refers to a divalent group having the same structure as a C_3 - C_{10} cycloalkenyl group.

[0293] A C_1 - C_{10} heterocycloalkenyl group as used herein refers to a monovalent monocyclic group including at least one hetero atom selected from N, O, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one double bond in its ring. Detailed examples of the C_1 - C_{10} heterocycloalkenyl group are a 2,3-dihydrofuran group and a 2,3-dihydrothiophenyl group. A C_1 - C_{10} heterocycloalkenylene group as used herein refers to a divalent group having the same structure as a C_1 - C_{10} heterocycloalkenyl group.

[0294] A C_6 - C_{60} aryl group as used herein refers to a monovalent group including a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C_6 - C_{60} arylene group as used herein refers to a divalent group including a carbocyclic aromatic system having 6 to 60 carbon atoms. Detailed examples of the C_6 - C_{60} aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C_6 - C_{60} aryl group and the C_6 - C_{60} arylene group each include two or more rings, the rings may be fused to each other.

[0295] A C_1 - C_{60} heteroaryl group as used herein refers to a monovalent group having an aromatic system including at least one hetero atom selected from N, O, P, and S as a ring-forming atom and 1 to 60 carbon atoms. A C_1 - C_{60} heteroarylene group as used herein refers to a divalent group having a carbocyclic aromatic system including at least one hetero atom selected from N, O, P, and S as a ring-forming atom and 1 to 60 carbon atoms. Detailed examples of the C_1 - C_{60} heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C_1 - C_{60} heteroaryl group and the C_1 - C_{60} heteroarylene group each include a plurality of rings, the rings may be fused to each other.

[0296] A C_6 - C_{60} aryloxy group as used herein indicates $—OA_{102}$ (wherein A_{102} is the C_6 - C_{60} aryl group), and a C_6 - C_{60} arylthio group as used herein indicates $—SA_{103}$ (wherein A_{103} is the C_6 - C_{60} aryl group).

[0297] A monovalent non-aromatic condensed polycyclic group as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed to each other, only carbon atoms as ring forming atoms, wherein the molecular structure as a whole is non-aromatic in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. A divalent non-aromatic condensed polycyclic group as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0298] A monovalent non-aromatic condensed heteropolycyclic group as used herein refers to a monovalent group (for example, having 1 to 60 carbon atoms) that has two or more rings condensed to each other, has a hetero atom selected from N, O, P, and S, other than carbon atoms, as a ring forming atom, wherein the molecular structure as a whole is non-aromatic in the entire molecular structure. The monovalent non-aromatic condensed heteropolycyclic group includes a carbazolyl group. A divalent non-aromatic condensed hetero-polycyclic group as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed hetero-polycyclic group.

[0299] At least one of substituents of the substituted C_1 - C_{60} alkyl group, substituted C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group, substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10} cycloalkyl group, substituted C_1 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substituted C_1 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_1 - C_{60} heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from

[0300] a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0301] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_{11})(Q_{12})$, $-Si(Q_{13})(Q_{14})(Q_{15})$, $-B(Q_{16})(Q_{17})$, and $-P(=O)(Q_{18})(Q_{19})$;

[0302] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0303] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, $-F$, $-Cl$, $-Br$, $-I$, $-CD_3$, $-CD_2H$, $-CDH_2$, $-CF_3$, $-CF_2H$, $-CFH_2$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-N(Q_{21})(Q_{22})$, $-Si(Q_{23})(Q_{24})(Q_{25})$, $-B(Q_{26})(Q_{27})$, and $-P(=O)(Q_{28})(Q_{29})$; and

[0304] $-N(Q_{31})(Q_{32})$, $-Si(Q_{33})(Q_{34})(Q_{35})$, $-B(Q_{36})(Q_{37})$, and $-P(=O)(Q_{38})(Q_{39})$,

[0305] wherein Q_1 , Q_2 , Q_6 to Q_9 , Q_{11} to Q_{19} , Q_{21} to Q_{29} , and Q_{31} to Q_{39} are each independently selected from a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group (for example, a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group).

[0306] The present inventive concept will be described in further detail with reference to the following examples. These examples are for illustrative purposes only and are not intended to limit the scope of the present inventive concept.


[0307] Hereinafter, the compound according to an embodiment and an organic light-emitting device will be described in detail by referring to Synthesis Examples and Examples, but embodiments are not limited thereto. The wording "B was used instead of A" used in describing Synthesis Examples means that an amount of B used was identical to an amount of A used based on molar equivalence.

EXAMPLE

Synthesis Example 1

Synthesis of Compound 2

[0308]




[0309] 10.0 grams (g) (42.22 millimoles (mmol)) of 2,5-dibromopyridine was added to 200 milliliters (ml) of diethyl ether. The mixture was cooled to -78°C ., 42.22 mmol of n-BuLi was slowly added thereto, and the resulting mixture was stirred for 1 hour at a temperature of -78°C .. Then, 5.2 ml (42.22 mmol) of chloro(trimethyl)germane was added thereto, and the mixture was reacted for 1 hour at a temperature of -78°C ., heated to room temperature, and then allowed to stir for 12 hours. The resultant was extracted with dichloromethane. The organic layer was dried using anhydrous magnesium sulfate and filtered. The solvent was removed from the resultant under reduced pressure, and the residue thus obtained was purified by column chromatography with ethyl acetate:hexane=1:15 as an eluent to obtain 6.3 g of Compound A2 (yield: 54%). The compound was identified using LCMS and ^1H NMR.

[0310] ^1H -NMR (CDCl_3) δ 8.36 (s, 1H), 7.58 (d, 1H), 7.44 (d, 2H), 0.42 (s, 9H)

[0311] MS: m/z 275.94 [(M+1) $^+$]

Synthesis of Compound A1

[0312]



[0313] 6.30 g (22.93 mmol) of Compound A2, 5.35 g (43.85 mmol) of biphenylboronic acid, 0.90 g (3.99 mmol) of $\text{Pd}(\text{PPh}_3)_4$, and 2.09 g (7.97 mmol) of K_2CO_3 were mixed with 100 ml of toluene and 50 ml of distilled water, and the mixture was stirred for 18 hours at 70°C ., cooled to room temperature, and filtered. The resultant was extracted with dichloromethane. The organic layer, was dried using anhydrous magnesium sulfate (MgSO_4) and filtered. The solvent was removed from the resultant under reduced pressure, and


the residue thus obtained was purified by column chromatography with ethyl acetate:hexane=1:25 to obtain 5.1 g of Compound A1 (yield: 64%). The compound was identified using LCMS and ^1H NMR.

[0314] ^1H -NMR (CDCl_3) δ 8.75 (s, 1H), 8.23 (s, 1H), 7.96 (d, 1H), 7.84 (dd, 1H), 7.75 (d, 1H), 7.65 (t, 3H), 7.54 (t, 1H), 7.45 (t, 2H), 7.38 (t, 1H), 0.46 (s, 9H)

[0315] MS: m/z 349.09 [(M+1) $^+$]

Synthesis of Compound M2A


[0316]





[0317] 10.0 g (64.43 mmol) of 2-phenylpyridine and 10.1 g (28.64 mmol) of iridium chloride trihydrate were mixed with 150 ml of ethoxyethanol and 50 ml of distilled water, and the mixture was stirred under reflux for 24 hours to allow a reaction to proceed. The mixture was cooled to room temperature. The solid thus obtained was filtered and separated, thoroughly washed with water, methanol, and hexane in the stated order. Then, the solid was dried in a vacuum oven to obtain 13.8 g of Compound M2A (yield: 90%).

Synthesis of Compound M1A

[0318]






[0319] 2.36 g (2.20 mmol) of Compound M2A was mixed with 60 ml of methylene chloride (MC), and then 1.13 g (4.40 mmol) of AgOTf was dissolved in 20 ml of a mixture of dichloromethane/methanol (3:1). Then, the flask was wrapped with aluminum foil to protect the contents from the action of light, and the mixture was stirred for 18 hours at room temperature. The solid thus produced was removed by using a celite filter. The solvent was removed from the resultant under reduced pressure. The solid (Compound M1A) thus obtained was used in the next step without additional purification.

Synthesis of Compound 2

[0320]



[0321] 3.06 g (4.29 mmol) of Compound M1A and 1.79 g (5.15 mmol) of Compound A1 was mixed with 20 ml of ethanol, and the mixture was stirred under reflux for 15 hours to allow the reaction to proceed, and cooled to room temperature. A solid obtained by filtering the mixture was thoroughly washed with ethanol and hexane, and the resultant was purified by column chromatography with ethyl acetate:hexane=1:6 as an eluent to obtain 2.1 g of Compound 2 (yield: 58%). The compound was identified using LCMS and ¹H NMR.

[0322] ¹H-NMR (CDCl₃) δ 7.89 (m, 4H), 7.61 (m, 8H), 7.50 (d, 1H), 7.40 (m, 3H), 7.24 (m, 1H), 7.13 (d, 1H), 6.89 (m, 9H), 0.19 (s, 9H)


[0323] MS: m/z 849.18 [(M+1)⁺]

Synthesis Example 2

Synthesis of Compound 7

Synthesis of Compound B3

[0324]




[0325] 20.0 g (79.71 mmol) of 2,5-dibromo-4-methylpyridine, 11.7 g (95.65 mmol) of phenylboronic acid, 0.90 g (3.99 mmol) of Pd(OAc)₂, 2.09 g (7.97 mmol) of PPh₃, and 22.0 g (159.4 mmol) of K₂CO₃ were mixed with 200 ml of acetonitrile and 100 ml of methanol, and the mixture was stirred at 50° C. for 18 hours, cooled to room temperature, and filtered. The resultant was extracted with dichloromethane to collect an organic layer, which was then dried using anhydrous magne-

sium sulfate (MgSO_4) and filtered. The solvent was removed from the resultant under reduced pressure, and the residue thus obtained was purified by column chromatography with ethyl acetate:hexane=1:2.5 as an eluent to obtain 9.6 g of Compound B3 (yield: 48%).

Synthesis of Compound B2

[0326]




[0327] 9.1 g of Compound B2 (yield: 82%) was obtained in the same manner as in the synthesis of Compound A2 of Synthesis Example 1, except that 9.6 g (38.61 mmol) of Compound B3 was used instead of 2,5-dibromopyridine. The compound was identified using LCMS and ^1H NMR.

[0328] ^1H -NMR (CDCl_3) δ 8.53 (s, 1H), 7.92 (d, 2H), 7.39 (m, 4H), 2.40 (s, 3H), 0.44 (s, 9H)


[0329] MS: m/z 287.07 $[(M+1)^+]$

Synthesis of Compound 7

[0330]



-continued



[0331] 1.3 g of Compound 7 (yield: 33%) was obtained in the same manner as in the synthesis of Compound 2 of Synthesis Example 1, except that 1.7 g (6.11 mmol) of Compound B2 instead of Compound A1 and 3.6 g (5.09 mmol) of Compound M1A were used. The compound was identified using LCMS and ^1H NMR.

[0332] ^1H -NMR (CDCl_3) δ 7.91 (m, 4H), 7.65 (m, 8H), 7.54 (s, 1H), 7.40 (m, 4H), 7.24 (s, 1H), 7.13 (d, 2H), 6.89 (m, 2H), 2.43 (s, 3H), 0.20 (s, 9H)


[0333] MS: m/z 787.16 $[(M+1)^+]$

Synthesis Example 3

Synthesis of Compound 8

Synthesis of Compound B1

[0334]



[0335] 2.30 g (8.04 mmol) of Compound B2 was mixed with 40 ml of THF and cooled to -78°C . Then, 7.24 ml (14.48



mmol) of lithium diisopropylamide (LDA) was slowly added thereto. The mixture was stirred at -78°C . for 1 hour to allow the reaction to proceed, then warmed to room temperature, and stirred at room temperature for additional 1.5 hour. Then, the temperature of the resultant mixture was lowered to -78°C ., 1.25 mL (13.27 mmol) of 2-bromopropane was slowly added thereto, and the mixture was warmed to room temperature, at which the reaction was allowed to stir for 12 hours. The resultant was extracted with dichloromethane to collect an organic layer, which was then dried using anhydrous magnesium sulfate (MgSO_4) and filtered. The solvent was removed from the resultant under reduced pressure, and the residue thus obtained was purified by column chromatography with ethyl acetate:hexane=4:96 as an eluent to obtain 2.0 g of Compound B1 (yield: 76%). The compound was identified using LCMS and ^1H NMR.

[0336] ^1H -NMR (CDCl_3) δ 8.55 (s, 1H), 7.89 (d, 2H), 7.40 (m, 4H), 3.25 (d, 2H), 1.85 (m, 1H), 0.90 (d, 6H), 0.45 (s, 9H)


[0337] MS: m/z 329.12 $[(\text{M}+1)^+]$

Synthesis of Compound 8

[0338]



-continued



[0339] 1.0 g of Compound 8 (yield: 27%) was obtained in the same manner as in the synthesis of Compound 2 in Synthesis Example 1, except that 1.7 g (5.26 mmol) of Compound B1 instead of Compound B2 and 3.1 g (4.39 mmol) of Compound M1A were used. The compound was identified using LCMS and ^1H NMR.

[0340] ^1H -NMR (CDCl_3) δ 7.89 (m, 4H), 7.63 (m, 8H), 7.53 (s, 1H), 7.39 (m, 4H), 7.25 (s, 1H), 7.15 (d, 2H), 6.90 (m, 2H), 3.31 (d, 2H), 1.89 (m, 1H), 0.91 (d, 6H), 0.18 (s, 9H)


[0341] MS: m/z 829.20 $[(\text{M}+1)^+]$

Synthesis Example 4

Synthesis of Compound 10

Synthesis of Compound C3



[0342]



[0343] 9.5 g of Compound C3 (yield: 78%) was obtained in the same manner as in the synthesis of Compound B3 in Synthesis Example 2, except that 8.0 g (40.34 mmol) of ([1,1'-biphenyl]-3-yl)boronic acid was used instead of phenylboronic acid.

Synthesis of Compound C2



[0344]




[0345] 2.5 g of Compound C2 (yield: 76%) was obtained in the same manner as in the synthesis of Compound A2 in Synthesis Example 1, except that 3.0 g (9.25 mmol) of Compound C3 was used instead of 2,5-dibromopyridine.

Synthesis of Compound C1

[0346]




-continued




[0347] 2.1 g of Compound C1 (yield: 75%) was obtained in the same manner as in the synthesis of Compound B1 in Synthesis Example 3, except that 2.5 g (7.03 mmol) of Compound C2 was used instead of Compound B2.

Synthesis of Compound 10

[0348]



-continued



Compound 10

[0349] 1.2 g of Compound 10 (yield: 34%) was obtained in the same manner as in the synthesis of Compound 2 in Synthesis Example 1, except that 1.9 g (4.65 mmol) of Compound C1 instead of Compound A1 and 2.8 g (3.87 mmol) of Compound M1A were used. The compound was identified using LCMS and ^1H NMR.

[0350] ^1H -NMR (CDCl_3) δ 7.89 (m, 4H), 7.59 (m, 8H), 7.49 (d, 1H), 7.38 (m, 3H), 7.21 (m, 1H), 7.12 (d, 1H), 6.87 (m, 8H), 3.30 (d, 2H), 1.88 (m, 1H), 0.90 (d, 6H), 0.19 (s, 9H)

[0351] MS: m/z 905.23 $[(M+1)^+]$

[0355] BA1q was deposited on the emission layer to form a hole blocking layer having a thickness of 50 Å, Alq₃ was deposited on the hole blocking layer to form an electron transport layer having a thickness of 300 Å, LiF was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was deposited on the electron injection layer to form a second electrode (a cathode) having a thickness of 1,200 Å, thereby manufacturing an organic light-emitting device having a structure of ITO/m-MTDATA (600 Å)/ α -NPD (250 Å)/CBP+10% (Compound 2) (400 Å)/BA1q (50 Å)/Alq₃ (300 Å)/LiF (10 Å)/Al (1200 Å).

Examples 2 to 4 and Comparative Examples 1 to 3

[0356] Organic light-emitting devices were manufactured in the same manner as in Example 1, except that Compounds shown in Table 2 were each used instead of Compound 2 as a dopant in the formation of the emission layer.

Evaluation Example 1

Evaluation of Organic Light-Emitting Device Characteristics

[0357] Driving voltage, efficiency, color purity, maximum efficiency, FWHM of EL spectrum, maximum emission wavelengths of the organic light-emitting devices manufactured in Examples 1 to 4 and Comparative Examples 1 to 3 were measured, and the results are shown in Table 2. The measurement was performed by using a current-voltmeter (available from Keithley 2400) and a luminance meter (available from Minolta Cs-1000A).

TABLE 2

	Dopant	Driving voltage (V)	Efficiency (Cd/A) (at driving voltage)	CIE x (at driving voltage)	Maximum efficiency (Cd/A)	FWHM (nm)	Maximum emission wavelength (nm)
Example 1	Compound 2	4.9	49.6	0.354	59.0	74.6	523
Example 2	Compound 7	4.9	49.3	0.330	60.5	75.9	517
Example 3	Compound 8	5.0	47.5	0.325	62.0	77	515
Example 4	Compound 10	4.9	50.0	0.322	60.0	79	518
Comparative Example 1	Compound R1	5.4	49.2	0.374	57.7	81.8	528
Comparative Example 2	Compound R2	5.2	42.0	0.331	55.0	77	520
Comparative Example 3	Compound R3	5.4	39.0	0.315	46.5	84	513


Example 1

[0352] An ITO glass substrate (an anode) was cut to a size of 50 millimeters (mm)×50 mm×0.5 mm, and then ultrasonicated in acetone, isopropyl alcohol, and pure water (for 15 minutes in each solvent), and then cleaned by UV ozone for 30 minutes.


[0353] Then, m-MTDATA was deposited on the glass substrate at a deposition rate of 1 Å/sec to form a hole injection layer having a thickness of 600 Å on the ITO electrode (anode), and α -NPD was deposited on the hole injection layer at a deposition rate of 1 Å/sec to form a hole transport layer having a thickness of 250 Å.

[0354] Compound 2 (a dopant) and CBP (a host) were co-deposited on the hole transport layer at deposition rates of 0.1 Å/sec and 1 Å/sec, respectively, to form an emission layer having a thickness of 400 Å.


Compound R1



-continued



Compound R2



Compound R3

[0358] Referring to Table 2, a coordinate of CIE x of the organic light-emitting devices prepared in Examples 1 to 4 was in a range of 0.322 to 0.354, but a coordinate of CIE x of the organic light-emitting device prepared in Comparative Example 1 was 0.374. Thus, it was confirmed that the organic light-emitting devices of Examples 1 to 4 had better color purities than that of the organic light-emitting device of Comparative Example 1. Also, it may be confirmed that efficiencies and maximum efficiencies of the organic light-emitting devices of Examples 1 to 4 are better than those of organic light-emitting devices of Comparative Examples 2 and 3.


[0359] As described above, according to the one or more of the above exemplary embodiments, the organometallic compound has excellent electric characteristics and thermal stability. Thus, an organic light-emitting device including the organometallic compound may have low driving voltage, high efficiency, and excellent color purity.

[0360] It should be understood that the exemplary embodiments described therein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each exemplary embodiment should typically be considered as available for other similar features or aspects in other exemplary embodiments.


[0361] While one or more exemplary embodiments have been described with reference to the figures, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope as defined by the following claims.

What is claimed is:


1. An organometallic compound represented by Formula 1:




Formula 1



Formula 2A



Formula 2B



Formula 2C

wherein in Formula 1,

M is selected from Ir, Pt, Os, Ti, Zr, Hf, Eu, Tb, Tm, and Rh, L₁ in Formula 1 is selected from ligands represented by Formula 2A,

L₂ in Formula 1 is selected from ligands represented by Formula 2B,

provided that L₁ and L₂ in Formula 1 are different from each other,

n₁ and n₂ in Formula 1 are each independently 1 or 2, and the sum of n₁ and n₂ is 2 or 3,

Y₁ and Y₂ in Formula 2A are each independently C or N, and Y₃ and Y₄ in Formula 2B are each independently C or N,

in Formulae 2A and 2B,

CY₁ and CY₃ are each independently a C₁-C₆₀ heterocyclic group,

CY₂ and CY₄ are each independently selected from a C₅-C₆₀ carbocyclic group and a C₁-C₆₀ heterocyclic group,

wherein CY₁ and CY₂ are optionally linked to each other through a first linking group, and,

CY₃ and CY₄ are optionally linked to each other through a second linking group, in Formulae 2A and 2B,

Z₁ to Z₄ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, —SF₅, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —B(Q₆)(Q₇), and —P(=O)(Q₈)(Q₉),

a1 to a4 are each independently an integer selected from 0 to 4,

in Formula 2B,

R₁₁ and R₁₂ are each independently selected from groups represented by Formula 2C,

b1 and b2 are each independently an integer selected from 0 to 3, and the sum of b1 and b2 is 1 or greater,

in Formula 2C,

R₁ to R₃ are each independently selected from a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

when b1 is 1 or greater, at least one of R₁ to R₃ in Formula 2C is optionally bound to neighboring group Z₃ to form a saturated or unsaturated ring having two to ten carbons,

when b2 is 1 or greater, at least one of R₁ to R₃ in Formula 2C is optionally bound to neighboring group Z₄ to form a saturated or unsaturated ring having two to ten carbons,

* and *' in Formulae 2A and 2B are a binding site to M in Formula 1, and * in Formula 2C is a binding site to CY₃ or CY₄ in Formula 2B,

at least one of substituents of the substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀

heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from

a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), —B(Q₁₆)(Q₁₇), and —P(=O)(Q₁₈)(Q₁₉);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), —B(Q₂₆)(Q₂₇), and —P(=O)(Q₂₈)(Q₂₉); and

—N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), —B(Q₃₆)(Q₃₇), and —P(=O)(Q₃₈)(Q₃₉),

wherein Q₁, Q₂, Q₆ to Q₉, Q₁₁ to Q₁₉, Q₂₁ to Q₂₉, and Q₃₁ to Q₃₉ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

2. The organometallic compound of claim 1, wherein in Formula 2A and 2B,

CY₁ and CY₃ are each independently selected from a pyridine ring, a pyrimidine ring, a pyrazine ring, a triazine ring, a quinoline ring, an isoquinoline ring, a quinazoline ring, a quinoxaline ring, a triazole ring, an imidazole ring, and a pyrazole ring, and

CY₂ and CY₄ are each independently selected from a benzene ring, a naphthalene ring, a pyridine ring, a pyrimidine ring, a pyrazine ring, a triazine ring, a quinoline ring, an isoquinoline ring, a quinazoline ring, a quinoxaline ring, a carbazole ring, a dibenzofuran ring, and a dibenzothiophene ring.

3. The organometallic compound of claim 1, wherein in Formula 2A and 2B,

CY₁ and CY₃ are each independently selected from a pyridine ring, a pyrimidine ring, a pyrazine ring, a triazine ring, a triazole ring, an imidazole ring, and a pyrazole ring, and

CY₂ and CY₄ are each independently selected from a benzene ring, a naphthalene ring, a pyridine ring, a pyrimidine ring, a pyrazine ring, a carbazole ring, a dibenzofuran ring, and a dibenzothiophene ring.

4. The organometallic compound of claim 1, wherein in Formula 2A and 2B,

Z₁ to Z₄ are each independently selected from

a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, —SF₅, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid

group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthroli-nyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthroli-nyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclo-

hexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzo-carbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

$-\text{B}(\text{Q}_6)(\text{Q}_7)$ and $-\text{P}(=\text{O})(\text{Q}_8)(\text{Q}_9)$,

wherein Q_6 to Q_9 are each independently selected from

$-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCD}_3$, $-\text{CHDCD}_2\text{H}$, $-\text{CHDCDH}_2$, $-\text{CHDCD}_3$, $-\text{CD}_2\text{CD}_3$, $-\text{CD}_2\text{CD}_2\text{H}$, and $-\text{CD}_2\text{CDH}_2$;

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from a deuterium, a C_1 - C_{10} alkyl group, and a phenyl group.

5. The organometallic compound of claim 1, wherein in Formula 2A and 2B,

Z_1 to Z_4 are each independently selected from

a hydrogen, a deuterium, $-\text{F}$, a cyano group, a nitro group, $-\text{SF}_5$, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group, each substituted with at least one selected from a deuterium, $-\text{F}$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, C_1 - C_{10} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

$-\text{B}(\text{Q}_6)(\text{Q}_7)$ and $-\text{P}(=\text{O})(\text{Q}_8)(\text{Q}_9)$,

wherein Q_6 to Q_9 are each independently selected from

$-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCD}_3$, $-\text{CHDCD}_2\text{H}$, $-\text{CHDCDH}_2$, $-\text{CHDCD}_3$, $-\text{CD}_2\text{CD}_3$, $-\text{CD}_2\text{CD}_2\text{H}$, and $-\text{CD}_2\text{CDH}_2$;

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and


an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from a deuterium, a C_1 - C_{10} alkyl group, and a phenyl group.

6. The organometallic compound of claim 1, wherein in Formula 2A and 2B,

Z_1 to Z_4 are each independently selected from a hydrogen, a deuterium, $-\text{F}$, a cyano group, a nitro group, $-\text{SF}_5$, $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, a group represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-32:



Formula 9-1




Formula 9-2



Formula 9-3

-continued



Formula 9-4

Formula 9-5

Formula 9-6

Formula 9-7

Formula 9-8

Formula 9-9

Formula 9-10

Formula 9-11

Formula 9-12

Formula 9-13

Formula 9-14


Formula 9-15

Formula 9-16

Formula 9-17

Formula 10-1

-continued



Formula 10-2

Formula 10-3

Formula 10-4

Formula 10-5

Formula 10-6

Formula 10-7

Formula 10-8

Formula 10-9











Formula 10-10

Formula 10-11

Formula 10-12

Formula 10-13

-continued



Formula 10-14

Formula 10-15

Formula 10-16

Formula 10-17

Formula 10-18

Formula 10-19










Formula 10-20

Formula 10-21

Formula 10-22

Formula 10-23

-continued



Formula 10-24

Formula 10-25

Formula 10-26

Formula 10-27

Formula 10-28

Formula 10-29

Formula 10-30

Formula 10-31

Formula 10-32

7. The organometallic compound of claim 1, wherein b1 is 1 or 2, and b2 is 0.

8. The organometallic compound of claim 1, wherein in Formula 2C,

R₁ to R₃ are each independently selected from

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with

at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.

9. The organometallic compound of claim 1, wherein in Formula 2C,

R₁ to R₃ are each independently selected from

a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and


a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a

butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group.

10. The organometallic compound of claim 1, wherein in Formula 2C, R₁ to R₃ are each independently selected from —CH₃, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a group represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-18:



Formula 9-1




Formula 9-2




Formula 9-3




Formula 9-4




Formula 9-5




Formula 9-6




Formula 9-7




Formula 9-8




Formula 9-9




Formula 9-10



Formula 9-11




Formula 9-12




Formula 9-13


-continued




Formula 9-14




Formula 9-15




Formula 9-16




Formula 9-17




Formula 10-1




Formula 10-2




Formula 10-3




Formula 10-4




Formula 10-5



Formula 10-6













Formula 10-7



Formula 10-8

-continued



11. The organometallic compound of claim 1, wherein in Formula 2C, R_1 to R_3 are identical to one another.

12. The organometallic compound of claim 1, wherein in Formula 2C, R_1 and R_2 are identical to each other and R_3 is different from R_1 ; or all of R_1 to R_3 are different from one another.

13. The organometallic compound of claim 1, wherein in Formula 1,

L_1 is selected from Formulae 3-1 to 3-110:

Formula 10-9

Formula 10-10

Formula 10-11

Formula 10-12

Formula 10-13






Formula 10-14

Formula 10-15

Formula 10-16

Formula 10-17

Formula 10-18



Formula 3-1






Formula 3-2

Formula 3-3

Formula 3-4

Formula 3-5

-continued



Formula 3-6






Formula 3-7

Formula 3-8

Formula 3-9

Formula 3-10

-continued



Formula 3-11






Formula 3-12

Formula 3-13


Formula 3-14

Formula 3-15


-continued




Formula 3-16




Formula 3-17




Formula 3-18



Formula 3-19



Formula 3-20



-continued

Formula 3-21


Formula 3-22

Formula 3-23


Formula 3-24

Formula 3-25


-continued




Formula 3-26




Formula 3-27



Formula 3-28




Formula 3-29




Formula 3-30


-continued




Formula 3-31




Formula 3-32



Formula 3-33








Formula 3-34




Formula 3-35


-continued




Formula 3-36




Formula 3-37




Formula 3-38



Formula 3-39



Formula 3-40



-continued

Formula 3-41


Formula 3-42

Formula 3-43


Formula 3-44

Formula 3-45


-continued




Formula 3-46




Formula 3-47



Formula 3-48




Formula 3-49




Formula 3-50


-continued




Formula 3-51




Formula 3-52



Formula 3-53




Formula 3-54




Formula 3-55


-continued




Formula 3-56




Formula 3-57



Formula 3-58




Formula 3-59




Formula 3-60


-continued




Formula 3-61




Formula 3-62



Formula 3-63




Formula 3-64




Formula 3-65


-continued




Formula 3-66




Formula 3-67



Formula 3-68




Formula 3-69




Formula 3-70


-continued




Formula 3-71




Formula 3-72



Formula 3-73




Formula 3-74




Formula 3-75

-continued




Formula 3-76




Formula 3-81


Formula 3-77




Formula 3-82




Formula 3-78




Formula 3-83




Formula 3-79




Formula 3-84




Formula 3-80




Formula 3-85




-continued




Formula 3-86




Formula 3-87



Formula 3-88




Formula 3-89




Formula 3-90


-continued




Formula 3-91




Formula 3-92



Formula 3-93




Formula 3-94



Formula 3-95


-continued



Formula 3-96


-continued

Formula 3-101




Formula 3-97

Formula 3-102




Formula 3-98

Formula 3-103





Formula 3-99

Formula 3-104




Formula 3-100

Formula 3-105



-continued



Formula 3-106

wherein in Formulae 3-1 to 3-110,


Z_1 , Z_2 , Z_{1a} , Z_{1b} , Z_{2a} , Z_{2b} , and Z_{2c} are each independently selected from

a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, —SF₅, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;


a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthroli-nyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;


a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthroli-




Formula 3-107



Formula 3-108



Formula 3-109



Formula 3-110

nyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

—N(Q₁)(Q₂), —B(Q₆)(Q₇), and —P(=O)(Q₈)(Q₉), wherein Q₁, Q₂ and Q₆ to Q₉ are each independently selected from

—CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCD₃, —CHDCD₂H, —CHDCDH₂, —CHDCD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from a deuterium and a C₁-C₁₀ alkyl group,

aa2 and ab2 are each independently 1 or 2,


aa3 and ab3 are each independently an integer selected from 1 to 3,

aa4 and ab4 are each independently an integer selected from 1 to 4, and


* and *' are a binding site to M of Formula 1.

14. The organometallic compound of claim 1, wherein in Formula 1, L₁ is selected from ligands represented by Formulae 3-1 (1) to 3-1(59):


Formula 3-1(1)




Formula 3-1(2)




Formula 3-1(3)




Formula 3-1(4)




Formula 3-1(5)




Formula 3-1(6)




-continued




Formula 3-1(7)




Formula 3-1(8)




Formula 3-1(9)



Formula 3-1(10)




Formula 3-1(11)




Formula 3-1(12)


-continued




Formula 3-1(13)




Formula 3-1(14)




Formula 3-1(15)



Formula 3-1(16)




Formula 3-1(17)




Formula 3-1(18)


-continued




Formula 3-1(19)




Formula 3-1(20)



Formula 3-1(21)




Formula 3-1(22)




Formula 3-1(23)


-continued




Formula 3-1(24)




Formula 3-1(25)



Formula 3-1(26)








Formula 3-1(27)




Formula 3-1(28)


-continued




Formula 3-1(29)




Formula 3-1(30)




Formula 3-1(31)



Formula 3-1(32)



Formula 3-1(33)



-continued

Formula 3-1(34)


Formula 3-1(35)

Formula 3-1(36)


Formula 3-1(37)

Formula 3-1(38)


-continued




Formula 3-1(39)




Formula 3-1(40)



Formula 3-1(41)




Formula 3-1(42)




Formula 3-1(43)


-continued




Formula 3-1(44)




Formula 3-1(45)



Formula 3-1(46)




Formula 3-1(47)




Formula 3-1(48)

-continued




Formula 3-1(49)


-continued




Formula 3-1(53)




Formula 3-1(50)




Formula 3-1(54)




Formula 3-1(51)



Formula 3-1(55)




Formula 3-1(52)




Formula 3-1(56)


-continued



Formula 3-1(57)



Formula 3-1(58)




Formula 3-1(59)

wherein in Formulae 3-1 (1) to 3-1(59),

Z₁, Z₂, Z_{1a}, Z_{1b}, Z_{1c}, Z_{1d}, Z_{2a}, Z_{2b}, Z_{2c}, and Z_{2d} are each independently selected from a deuterium, —F, a cyano group, a nitro group, —SF₅, —CH₃, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a group represented by Formulae 9-1 to 9-17, and a group represented by Formulae 10-1 to 10-32:



Formula 9-1



Formula 9-2




Formula 9-3




Formula 9-4


-continued




Formula 9-5




Formula 9-6



Formula 9-7



Formula 9-8




Formula 9-9




Formula 9-10




Formula 9-11




Formula 9-12




Formula 9-13




Formula 9-14




Formula 9-15




Formula 9-16



Formula 9-17















Formula 10-1



Formula 10-2

-continued



Formula 10-3

Formula 10-4

Formula 10-5

Formula 10-6

Formula 10-7

Formula 10-8

Formula 10-9

Formula 10-10











Formula 10-11

Formula 10-12

Formula 10-13

Formula 10-14

-continued



Formula 10-15

Formula 10-16

Formula 10-17

Formula 10-18

Formula 10-19

Formula 10-20









Formula 10-21

Formula 10-22

Formula 10-23

Formula 10-24

-continued



Formula 10-25

Formula 10-26

Formula 10-27






Formula 10-28

Formula 10-29

Formula 10-30

Formula 10-31

Formula 10-32



Formula 2B(1)

Formula 2B(2)






Formula 2B(3)

Formula 2B(4)

Formula 2B(5)

15. The organometallic compound of claim 1, wherein L_2 is selected from ligands represented by Formula 2B(1) to 2B(10):

-continued



wherein in Formulae 2B(1) to 2B(10), Z_3 , Z_4 , a_3 , a_4 , and R_1 to R_3 are the same as defined in claim 1, CY_4 is selected from a benzene, a naphthalene, a carbazole, a

dibenzofuran, and a dibenzothiophene, Z_5 is the same as defined in connection with Z_3 , and a_5 is an integer selected from 1 to 6.

16. The organometallic compound of claim 1, wherein L_2 is selected from ligands represented by Formulae 2B-1 to 2B-60:

Formula 2B(6)


Formula 2B(7)

Formula 2B(8)


Formula 2B(9)

Formula 2B(10)


Formula 2B-1




Formula 2B-2




Formula 2B-3




Formula 2B-4




-continued




Formula 2B-5



Formula 2B-6




Formula 2B-7




Formula 2B-8


-continued




Formula 2B-9



Formula 2B-10




Formula 2B-11




Formula 2B-12

-continued




Formula 2B-13


-continued




Formula 2B-16




Formula 2B-14




Formula 2B-17



Formula 2B-15




Formula 2B-18




Formula 2B-19


-continued




Formula 2B-20



Formula 2B-21




Formula 2B-22




Formula 2B-23


-continued




Formula 2B-24



Formula 2B-25




Formula 2B-26




Formula 2B-27

-continued




Formula 2B-28


-continued




Formula 2B-32




Formula 2B-29




Formula 2B-33




Formula 2B-30



Formula 2B-34




Formula 2B-31




Formula 2B-35

Formula 2B-36


-continued




Formula 2B-37



Formula 2B-38




Formula 2B-39




Formula 2B-40


-continued




Formula 2B-41



Formula 2B-42




Formula 2B-43




Formula 2B-44


-continued




Formula 2B-45




Formula 2B-50




Formula 2B-46




Formula 2B-51




Formula 2B-47




Formula 2B-52



Formula 2B-48




Formula 2B-53




Formula 2B-49


-continued




Formula 2B-54



Formula 2B-55




Formula 2B-56




Formula 2B-57


-continued



Formula 2B-58



Formula 2B-59



Formula 2B-60

wherein in Formulae 2B-1 to 2B-60,

 R_1 to R_3 are each independently selected from

a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a

cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

Z₃, Z_{3a}, Z_{3b}, Z₄, Z_{4a}, Z_{4b}, Z_{4c}, and Z_{4d} are each independently selected from

a deuterium, —F, a cyano group, a nitro group, —SF₅, a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group;

a methyl group, an ethyl group, an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an iso-hexyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an iso-heptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an iso-octyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an iso-nonyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an iso-decyl group, a sec-decyl group, a tert-decyl group, a

methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, and a pyrimidinyl group; and

—B(Q₆)(Q₇) and —P(=O)(Q₈)(Q₉),

wherein Q₆ to Q₉ are each independently selected from

—CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCD₃, —CHDCD₂H, —CHDCD₂H, —CHDCD₃, —CD₂CD₃, —CD₂CD₂H, and —CD₂CDH₂;

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group; and

an n-propyl group, an iso-propyl group, an n-butyl group, an iso-butyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an iso-pentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, and a naphthyl group, each substituted with at least one selected from a deuterium, a C₁-C₁₀ alkyl group, and a phenyl group.


17. The organometallic compound of claim 1, wherein in Formula 1,

M is Ir and the sum of n1 and n2 is 3, or


M is Pt and the sum of n1 and n2 is 2,

wherein the organometallic compound does not comprise an ionic pair.

18. The organometallic compound of claim 1, wherein the organometallic compound is represented by Compounds 1 to 109:




-continued




2

-continued




6


3




4




7




8




9




5




-continued




-continued




11




15




16




12




13




17




-continued




-continued





19




23




24




21




25




-continued




-continued



27





31




32


28




29




33




-continued




-continued




35




39




36




40




37










41







-continued







-continued




-continued




-continued




-continued




-continued




59




63




64




60




61




65




-continued




-continued



67





71




72


68




69










73




-continued



-continued




-continued




82


-continued




86




83




87




84



88




85




89

-continued




90


-continued




94




91




95




92



96




93




97

-continued




98


-continued




102




99




103



100




104




101

-continued




105

-continued




106

108



107

109



19. An organic light-emitting device comprising:
 a first electrode;
 a second electrode; and
 an organic layer disposed between the first electrode and the second electrode,
 wherein the organic layer comprises an emission layer, and
 wherein the organic layer comprises at least one organo-metallic compound of claim 1.


20. The organic light-emitting device of claim 19, wherein the emission layer comprises the organometallic compound of claim 1, wherein the emission layer further comprises a host, and wherein an amount of the organometallic compound is less than an amount of the host.

* * * * *


专利名称(译)	有机金属化合物和包括其的有机发光器件		
公开(公告)号	US20160181530A1	公开(公告)日	2016-06-23
申请号	US14/967594	申请日	2015-12-14
[标]申请(专利权)人(译)	三星电子株式会社		
申请(专利权)人(译)	SAMSUNG ELECTRONICS CO. , LTD.		
当前申请(专利权)人(译)	SAMSUNG ELECTRONICS CO. , LTD.		
[标]发明人	CHOI WHAIL KWAK SEUNGYEON KWAK YOONHYUN KWON OHYUN CHOI HYEONHO HWANG KYUYOUNG KIM SANGDONG PARK YOUNGJAE		
发明人	CHOI, WHAIL KWAK, SEUNGYEON KWAK, YOONHYUN KWON, OHYUN CHOI, HYEONHO HWANG, KYUYOUNG KIM, SANGDONG PARK, YOUNGJAE		
IPC分类号	H01L51/00 C09K11/06 C09K11/02 C07F15/00		
CPC分类号	H01L51/009 C07F15/0033 C09K11/06 C09K11/025 H01L51/0072 C09K2211/185 H01L2251/552 H01L51/5016 C09K2211/1007 C09K2211/1029 H01L51/5004		
优先权	1020140182552 2014-12-17 KR		
外部链接	Espacenet USPTO		

摘要(译)


由式1表示的有机金属化合物： $M(L_1)_{n1}(L_2)_{n2}$ 公式1 其中基团M， L_1 ， L_2 ， $n1$ 和 $n2$ 与它们在说明书中定义的相同。



Formula 2A



Formula 2B



Formula 2C