



US 20200227650A1

(19) **United States**(12) **Patent Application Publication**
KIM et al.(10) **Pub. No.: US 2020/0227650 A1**(43) **Pub. Date: Jul. 16, 2020**(54) **ORGANIC COMPOUND AND ORGANIC
ELECTROLUMINESCENT DEVICE
COMPRISING SAME****Publication Classification**(51) **Int. Cl.**

H01L 51/00 (2006.01)
C07D 403/10 (2006.01)
C09K 11/06 (2006.01)
C07D 405/14 (2006.01)
C07D 401/14 (2006.01)
C07D 405/10 (2006.01)
C07D 251/24 (2006.01)
C07D 409/10 (2006.01)

(52) **U.S. Cl.**

CPC *H01L 51/0072* (2013.01); *C07D 403/10*
(2013.01); *C09K 11/06* (2013.01); *H01L*
51/0067 (2013.01); *C07D 405/14* (2013.01);
H01L 51/5088 (2013.01); *C07D 401/14*
(2013.01); *C07D 405/10* (2013.01); *C07D*
251/24 (2013.01); *C07D 409/10* (2013.01);
H01L 51/0074 (2013.01); *H01L 51/0073*
(2013.01)

(71) Applicant: **DOOSAN CORPORATION**, Seoul
(KR)(72) Inventors: **Hong Suk KIM**, Yongin-si,
Gyeonggi-do (KR); **Young Bae KIM**,
Yongin-si (KR)(73) Assignee: **DOOSAN CORPORATION**, Seoul
(KR)(21) Appl. No.: **16/626,634**(22) PCT Filed: **May 2, 2018**(86) PCT No.: **PCT/KR2018/005079**

§ 371 (c)(1),


(2) Date: **Dec. 26, 2019**(30) **Foreign Application Priority Data**

Jun. 30, 2017 (KR) 10-2017-0083287


(57)

ABSTRACT


The present invention relates to a novel compound and an organic electroluminescent device including the same, and by using the compound according to the present invention in an organic material layer of an organic electroluminescent device, preferably a light emitting layer, luminous efficiency, driving voltage, lifetime and the like of the organic electroluminescent device may be enhanced.



【FIG. 1】



【FIG. 2】



ORGANIC COMPOUND AND ORGANIC ELECTROLUMINESCENT DEVICE COMPRISING SAME

TECHNICAL FIELD

[0001] The present invention relates to a novel organic compound capable of being used as a material for an organic electroluminescent device, and an organic electroluminescent device including the same.

BACKGROUND ART


[0002] With the observation of organic thin film light emission made by Bemanose in 1950s as a start, studies on organic electroluminescent (EL) devices have been continued leading to blue electroluminescence using a single anthracene crystal in 1965, and in 1987, an organic electroluminescent device having a laminated structure divided into functional layers of a hole layer and a light emitting layer has been proposed by Tang. After that, in order to manufacture organic electroluminescent devices with high efficiency and long lifetime, development has been made in the form of introducing each characteristic organic material layer into the device, which leads to the development of specialized materials used therein.

[0003] When a voltage is applied between the two electrodes in an organic electroluminescent device, holes and electrons are injected to an organic material layer from the anode and the cathode, respectively. When the injected holes and electrons meet, excitons are formed, and light emits when these excitons fall back to the ground state. Herein, materials used as the organic material layer may be divided into a light emitting material, a hole injection material, a hole transport material, an electron transport material, an electron injection material and the like depending on the function.



[0004] The light emitting material may be divided into, depending on the light emitting color, blue, green and red light emitting materials, and yellow and orange light emitting materials for obtaining better natural colors. In addition, in order to increase color purity and increase luminous efficiency through energy transfer, host/dopant series may be used as the light emitting material.

[0005] The dopant material may be divided into fluorescent dopants using organic materials and phosphorescent dopants using metal complex compounds including heavy atoms such as Ir or Pt. Herein, development of phosphorescent materials may enhance luminous efficiency up to 4 times compared to fluorescence theoretically, and therefore, studies on phosphorescent host materials have been widely progressed as well as on phosphorescent dopants.


[0006] So far, NPB, BCP, Alq₃ and the like have been widely known as materials of a hole injection layer, a hole transport layer, a hole blocking layer and an electron transport layer, and anthracene derivatives have been reported as a material of a light emitting layer. Particularly, among light emitting layer materials, metal complex compounds including Ir such as Firpic, Ir(ppy)₃ or (acac)Ir(btp)₂ having advantages in terms of efficiency enhancement have been used as blue, green and red phosphorescent dopant materials, and 4,4-dicarbazolylbiphenyl (CBP) has been used as a phosphorescent host material.




NPB

Alq₃

CBP




BCP



Firpic

Ir(ppy)₃



[0007] However, although being advantageous in terms of a light emission property, existing organic material layer materials have a low glass transition temperature and thereby have very unfavorable thermal stability, which is not satisfactory in terms of an organic electroluminescent device lifetime. Accordingly, development of organic material layer materials having superior performance has been required.

DISCLOSURE


Technical Problem

[0008] The present invention is directed to providing a novel organic compound capable of being used in an organic electroluminescent device, and having excellent hole and electron injection and transport abilities, a light emitting ability and the like.

[0009] The present invention is also directed to providing an organic electroluminescent device including the novel organic compound, and thereby exhibiting a low driving voltage, high luminous efficiency, and an enhanced lifetime.

Technical Solution

[0010] In view of the above, one embodiment of the present invention provides a compound represented by the following Chemical Formula 1:




[Chemical Formula 1]

[0011] in Chemical Formula 1,

[0012] X₁ is selected from the group consisting of S, O, N(R₁) and C(R₂)(R₃);


[0013] X₂ and X₃ are each independently N or C(Ar₁), and at least one thereof is C(Ar₁); and

[0014] ring A is represented by any one of the following Chemical Formulae 2 to 4;




[Chemical Formula 2]

-continued



[Chemical Formula 3]



[Chemical Formula 4]

[0015] in Chemical Formulae 1 to 4,

[0016] a dotted line means a part that is fused;


[0017] m is an integer of 0 to 4;

[0018] n is an integer of 0 to 6;

[0019] R₁ to R₄ are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a C₁~C₄₀ alkyl group, a C₂~C₄₀ alkenyl group, a C₂~C₄₀ alkynyl group, a C₃~C₄₀ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C₆~C₆₀ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a C₁~C₄₀ alkyloxy group, a C₆~C₆₀ aryloxy group, a C₃~C₄₀ alkylsilyl group, a C₆~C₆₀ arylsilyl group, a C₁~C₄₀ alkylboron group, a C₆~C₆₀ arylboron group, a C₆~C₆₀ arylphosphanyl group, a C₆~C₆₀ mono or diarylphosphinyl group and a C₆~C₆₀ arylamine group;

[0020] the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R₁ to R₄ are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a C₁~C₄₀ alkyl group, a C₂~C₄₀ alkenyl group, a C₂~C₄₀ alkynyl group, a C₆~C₆₀ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a C₆~C₆₀ aryloxy group, a C₁~C₄₀ alkyloxy group, a C₆~C₆₀ arylamine group, a C₃~C₄₀ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C₁~C₄₀ alkylsilyl group, a C₁~C₄₀ alkylboron group, a C₆~C₆₀ arylboron group, a C₆~C₆₀ arylphosphanyl group, a C₆~C₆₀ mono or diarylphosphinyl group and a C₆~C₆₀ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other; and

[0021] Ar₁ is a substituent represented by the following Chemical Formula 5, and when Ar₁ is present in plural numbers, these are the same as or different from each other;



[Chemical Formula 5]

[0022] in Chemical Formula 5,

[0023] * means a part where a bond is formed;

[0024] L_1 and L_2 are each independently selected from the group consisting of a direct bond, a $C_6\sim C_{18}$ arylene group and a heteroarylene group having 5 to 18 nuclear atoms;

[0025] Ar_2 is selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group; and

[0026] the arylene group and the heteroarylene group of L_1 and L_2 , and the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkoxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of Ar_2 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.

[0027] Another embodiment of the present invention provides an organic electroluminescent device including an anode, a cathode, and one or more organic material layers provided between the anode and the cathode, wherein at least one of the one or more organic material layers includes the compound of Chemical Formula 1.

[0028] The “alkyl” in the present invention is a monovalent substituent derived from linear or branched saturated hydrocarbon having 1 to 40 carbon atoms. Examples thereof may include methyl, ethyl, propyl, isobutyl, sec-butyl, pentyl, iso-amyl, hexyl and the like, but are not limited thereto.

[0029] The “alkenyl” in the present invention is a monovalent substituent derived from linear or branched unsaturated hydrocarbon having one or more carbon-carbon double bonds and having 2 to 40 carbon atoms. Examples thereof may include vinyl, allyl, isopropenyl, 2-butenyl and the like, but are not limited thereto.

[0030] The “alkynyl” in the present invention is a monovalent substituent derived from linear or branched unsaturated hydrocarbon having one or more carbon-carbon triple bonds and having 2 to 40 carbon atoms. Examples thereof may include ethynyl, 2-propynyl and the like, but are not limited thereto.

[0031] The “aryl” in the present invention means a monovalent substituent derived from aromatic hydrocarbon having a single ring or two or more rings combined and having 6 to 60 carbon atoms. In addition, a monovalent substituent having two or more rings fused with each other, including only carbon (for example, the number of carbon atoms may

be from 8 to 60) as a ring-forming atom, and with the whole molecule having non-aromaticity may also be included. Examples of such aryl may include phenyl, naphthyl, phenanthryl, anthryl, fluorenyl and the like, but are not limited thereto. The “heteroaryl” in the present invention means a monovalent substituent derived from monoheterocyclic or polyheterocyclic aromatic hydrocarbon having 5 to 60 nuclear atoms. Herein, one or more carbons, preferably 1 to 3 carbons, in the ring are substituted with a heteroatom selected from among N, O, P, S and Se. In addition, a monovalent group having two or more rings simply attached (pendant) or fused with each other, including a heteroatom selected from among N, O, P, S and Se as a ring-forming atom in addition to carbon, and with the whole molecule having non-aromaticity is interpreted to be included as well. Examples of such heteroaryl may include 6-membered monocyclic rings such as pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl or triazinyl; polycyclic rings such as phenoxathieryl, indolizyl, indolyl, purinyl, quinolyl, benzothiazole or carbazolyl; 2-furanyl, N-imidazolyl, 2-isoxazolyl, 2-pyridinyl, 2-pyrimidinyl and the like, but are not limited thereto.

[0032] The “aryloxy” in the present invention is a monovalent substituent represented by $RO-$, and R means aryl having 5 to 60 carbon atoms. Examples of such aryloxy may include phenyloxy, naphthyloxy, diphenyloxy and the like, but are not limited thereto.

[0033] The “alkyloxy” in the present invention is a monovalent substituent represented by $R'O-$, and R' means alkyl having 1 to 40 carbon atoms and is interpreted to include a linear, branched or cyclic structure. Examples of such alkyloxy may include methoxy, ethoxy, n-propoxy, 1-propoxy, t-butoxy, n-butoxy, pentoxy and the like, but are not limited thereto.

[0034] The “arylamine” in the present invention means amine substituted with aryl having 6 to 60 carbon atoms.

[0035] The “cycloalkyl” in the present invention means a monovalent substituent derived from monocyclic or polycyclic non-aromatic hydrocarbon having 3 to 40 carbon atoms. Examples of such cycloalkyl may include cyclopropyl, cyclopentyl, cyclohexyl, norbornyl, adamantyl and the like, but are not limited thereto.

[0036] The “heterocycloalkyl” in the present invention means a monovalent substituent derived from non-aromatic hydrocarbon having 3 to 40 nuclear atoms, and one or more carbons, preferably 1 to 3 carbons, in the ring are substituted with a heteroatom such as N, O, S or Se. Examples of such heterocycloalkyl may include morpholine, piperazine and the like, but are not limited thereto.

[0037] The “alkylsilyl” in the present invention means silyl substituted with alkyl having 1 to 40 carbon atoms, and the “aryl silyl” means silyl substituted with aryl having 5 to 60 carbon atoms.

[0038] The “fused ring” in the present invention means a fused aliphatic ring, a fused aromatic ring, a fused heteroaliphatic ring, a fused heteroaromatic ring, or a combined form thereof.

Advantageous Effects

[0039] A compound of the present invention has excellent thermal stability, carrier transport ability, light emitting ability and the like, and therefore, is useful as a material of an organic material layer of an organic electroluminescent device.

[0040] In addition, an organic electroluminescent device including a compound of the present invention in an organic material layer exhibits greatly enhanced properties in terms of light emitting performance, driving voltage, lifetime, efficiency and the like, and can be effectively used in a full color display panel and the like.

DESCRIPTION OF DRAWINGS

[0041] FIG. 1 is a sectional diagram illustrating an organic electroluminescent device according to one embodiment of the present invention.


[0042] FIG. 2 is a sectional diagram illustrating an organic electroluminescent device according to one embodiment of the present invention.

MODE FOR DISCLOSURE

[0043] Hereinafter, the present invention will be described in detail.

[0044] 1. Novel Organic Compound

[0045] A novel compound of the present invention may be represented by the following Chemical Formula 1:




[Chemical Formula 1]

[0046] in Chemical Formula 1,


[0047] X_1 is selected from the group consisting of S, O, $N(R_1)$ and $C(R_2)(R_3)$;

[0048] X_2 and X_3 are each independently N or $C(Ar_1)$, and at least one thereof is $C(Ar_1)$; and


[0049] ring A is represented by any one of the following Chemical Formulae 2 to 4;



[Chemical Formula 2]



[Chemical Formula 3]



[Chemical Formula 4]

[0050] in Chemical Formulae 1 to 4,

[0051] a dotted line means a part that is fused;

[0052] m is an integer of 0 to 4;


[0053] n is an integer of 0 to 6;

[0054] R_1 to R_4 are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano

group, a nitro group, a C_1 - C_{40} alkyl group, a C_2 - C_{40} alkenyl group, a C_2 - C_{40} alkynyl group, a C_3 - C_{40} cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C_6 - C_{60} aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a C_1 - C_{40} alkyloxy group, a C_6 - C_{60} aryloxy group, a C_3 - C_{40} alkylsilyl group, a C_6 - C_{60} arylsilyl group, a C_1 - C_{40} alkylboron group, a C_6 - C_{60} arylboron group, a C_6 - C_{60} arylphosphanyl group, a C_6 - C_{60} mono or diarylphosphinyl group and a C_6 - C_{60} arylamine group;

[0055] the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_1 to R_4 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a C_1 - C_{40} alkyl group, a C_2 - C_{40} alkenyl group, a C_2 - C_{40} alkynyl group, a C_3 - C_{40} cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C_1 - C_{40} alkylsilyl group, a C_1 - C_{40} alkylboron group, a C_6 - C_{60} arylboron group, a C_6 - C_{60} arylphosphanyl group, a C_6 - C_{60} mono or diarylphosphinyl group and a C_6 - C_{60} arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other; and

[0056] Ar_1 is a substituent represented by the following Chemical Formula 5, and when Ar_1 is present in plural numbers, these are the same as or different from each other;



[Chemical Formula 5]

[0057] in Chemical Formula 5,

[0058] * means a part where a bond is formed;

[0059] L_1 and L_2 are each independently selected from the group consisting of a direct bond, a C_6 - C_{18} arylene group and a heteroarylene group having 5 to 18 nuclear atoms;

[0060] Ar_2 is selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a C_1 - C_{40} alkyl group, a C_2 - C_{40} alkenyl group, a C_2 - C_{40} alkynyl group, a C_3 - C_{40} cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C_6 - C_{60} aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a C_1 - C_{40} alkyloxy group, a C_6 - C_{60} aryloxy group, a C_3 - C_{40} alkylsilyl group, a C_6 - C_{60} arylsilyl group, a C_1 - C_{40} alkylboron group, a C_6 - C_{60} arylboron group, a C_6 - C_{60} arylphosphanyl group, a C_6 - C_{60} mono or diarylphosphinyl group and a C_6 - C_{60} arylamine group; and


[0061] the arylene group and the heteroarylene group of L_1 and L_2 , and the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of Ar_2 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a

cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkyloxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.


[0062] More specifically, the compound represented by Chemical Formula 1 of the present invention has an EWG bonding to a 5-membered aromatic ring or a 5-membered aromatic heteroring such as indole, indazole, indene, benzofuran, benzothiophene or triazolo, and may be used as a host material since, by having a similar energy level with carbazole, the energy level may be adjusted to be higher than a dopant energy level. Particularly, benzofuran and benzothiophene moieties are electron abundant, and when used as an electron transport layer material of an organic electroluminescent device, mobility increases, and as a result, an increase in the luminous efficiency and a decrease in the driving voltage may be expected. In addition, the 5-membered aromatic ring or the 5-membered aromatic heteroring of the present invention has a smaller molecular weight compared to existing compounds, and therefore, may be deposited at a relatively lower deposition temperature when deposited compared to other materials, and favorable processability and enhanced thermal stability may be obtained.

[0063] Accordingly, the compound represented by Chemical Formula 1 of the present invention may be used as a material of an organic material layer of an organic electroluminescent device, and may be preferably used as a light emitting layer material (green phosphorescent host material), an electron transport layer/injection layer material, a light emitting auxiliary layer material or an electron transport auxiliary layer material, and more preferably as a light emitting layer material, an electron transport layer material or an electron transport auxiliary layer material. In addition, the organic electroluminescent device including the compound of Chemical Formula 1 may have performance and lifetime properties greatly enhanced, and a full color organic light emitting panel using such an organic electroluminescent device may have its performance maximized.

[0064] According to preferred one embodiment of the present invention, the compound may be represented by any one of the following Chemical Formulae 6 to 14:




[Chemical Formula 6]




[Chemical Formula 7]


-continued




[Chemical Formula 8]




[Chemical Formula 9]




[Chemical Formula 10]




[Chemical Formula 11]



[Chemical Formula 12]



[Chemical Formula 13]



[Chemical Formula 14]

[0065] in Chemical Formulae 6 to 14,

[0066] ring A, Ar_1 and R_1 to R_3 each have the same definitions as in Chemical Formula 1.

[0067] According to preferred one embodiment of the present invention, the compound is preferably represented by any one of Chemical Formulae 6 to 8 in terms of securing low driving voltage and high luminous efficiency, and may be more preferably represented by Chemical Formula 6 or 7, and even more preferably represented by Chemical Formula 8 or 10.

[0068] According to preferred one embodiment of the present invention, R_1 to R_4 are each independently selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and


[0069] the alkyl group, the aryl group and the heteroaryl group of R_1 to R_4 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these may be the same as or different from each other.

[0070] According to preferred one embodiment of the present invention, R_1 to R_4 are each independently selected


from the group consisting of a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthalenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a carbazolyl group, a fluorenyl group, a spirofluorenyl group and a dibenzodioxynyl group, and

[0071] the methyl group, the ethyl group, the propyl group, the butyl group, the pentyl group, the phenyl group, the biphenyl group, the terphenyl group, the naphthalenyl group, the pyridinyl group, the pyrimidinyl group, the triazinyl group, the dibenzofuranyl group, the dibenzothiophenyl group, the carbazolyl group, the fluorenyl group, the spirofluorenyl group and the dibenzodioxynyl group of R_1 to R_4 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a C_1 - C_{40} alkyl group, a C_6 - C_{60} alylamine group, a C_6 - C_{60} aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these may be the same as or different from each other.


[0072] According to preferred one embodiment of the present invention, L_1 and L_2 are each independently a direct bond, or a linker represented by any one of the following Chemical Formulae A-1 to A-7, but are not limited thereto:




A-1




A-2




A-3



A-4




A-5



A-6

-continued

A-7



[0073] in Chemical Formulae A-1 to A-7,

[0074] * means a part where a bond is formed;

[0075] Z_1 to Z_8 are each independently N or C(R_5);

[0076] any two of Z_1 to Z_6 forming a bond as a linker in Chemical Formula A-1 are C(R_5), and herein, R_5 is not present;

[0077] any one of Z_1 to Z_4 and any one of Z_5 to Z_8 forming a bond as a linker in Chemical Formula A-4 are C(R_5), and herein, R_5 is not present;

[0078] any two of Z_1 to Z_4 forming a bond as a linker in Chemical Formula A-5 are C(R_5), and herein, R_5 is not present;


[0079] X_4 and X_5 are each independently O, S, N(R_6) or C(R_7)(R_8);

[0080] X_6 is N or C(R_9);

[0081] R_5 to R_9 are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a C_1 - C_{40} alkyl group, a C_2 - C_{40} alkenyl group, a C_2 - C_{40} alkynyl group, a C_3 - C_{40} cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C_6 - C_{60} aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a C_1 - C_{40} alkyloxy group, a C_6 - C_{60} aryloxy group, a C_3 - C_{40} alkylsilyl group, a C_6 - C_{60} arylsilyl group, a C_1 - C_{40} alkylboron group, a C_6 - C_{60} arylboron group, a C_6 - C_{60} arylphosphanyl group, a C_6 - C_{60} mono or diarylphosphinyl group and a C_6 - C_{60} arylamine group, or adjacent groups bond to form a fused ring, and when R_5 to R_8 are each present in plural numbers, these are the same as or different from each other; and

[0082] the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_5 to R_9 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a C_1 - C_{40} alkyl group, a C_2 - C_{40} alkenyl group, a C_2 - C_{40} alkynyl group, a C_6 - C_{60} aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a C_6 - C_{60} aryloxy group, a C_1 - C_{40} alkyloxy group, a C_6 - C_{60} arylamine group, a C_3 - C_{40} cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C_1 - C_{40} alkylsilyl group, a C_1 - C_{40} alkylboron group, a C_6 - C_{60} arylboron group, a C_6 - C_{60} arylphosphanyl group, a C_6 - C_{60} mono or diarylphosphinyl group and a C_6 - C_{60} arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.

[0083] According to preferred one embodiment of the present invention, L_1 and L_2 are each independently a direct bond, or a linker represented by any one of the following Chemical Formulae B-1 to B-12, but are not limited thereto:



- continued
- B-11
- B-12
- B-2
- B-3
- B-4
- B-5
- B-6
- B-7
- B-8
- B-9
- B-10
- [0084]** in Chemical Formulae B-1 to B-12,
- [0085]** * means a part where a bond is formed;
- [0086]** R_6 to R_9 are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkyloxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group; and
- [0087]** the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_6 to R_9 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkyloxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.
- [0088]** According to preferred one embodiment of the present invention, in Chemical Formulae B-1 to B-12, R_6 to R_9 are each independently selected from the group consisting of hydrogen, a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms; and
- [0089]** the alkyl group, the aryl group and the heteroaryl group of R_6 to R_9 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and


when substituted with a plurality of the substituents, these are the same as or different from each other.

[0090] According to preferred one embodiment of the present invention, L_1 and L_2 are each independently a direct bond, or may be a linker selected from the group consisting of Chemical Formulae B-1 to B-3 and B-7 to B-12.

[0091] According to preferred one embodiment of the present invention, Ar_2 is a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, or an mylamine group, and

[0092] the aryl group, the heteroaryl group and the mylamine group of Ar_2 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ mylamine group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these may be the same as or different from each other.

[0093] According to preferred one embodiment of the present invention, Ar_2 may be a substituent represented by the following Chemical Formula 15:



[Chemical Formula 15]

[0094] in Chemical Formula 15,

[0095] * means a part where a bond is formed;


[0096] Y_1 to Y_5 are each independently N or C(R_{10});

[0097] R_{10} is selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group, and when R_{10} is present in plural numbers, these are the same as or different from each other; and

[0098] the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkoxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_{10} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and

when substituted with a plurality of the substituents, these are the same as or different from each other.

[0099] According to preferred one embodiment of the present invention, the substituent represented by Chemical Formula 15 may be a substituent represented by the following Chemical Formula 16:



[Chemical Formula 16]

[0100] in Chemical Formula 16,

[0101] * means a part where a bond is formed;

[0102] R_{11} and R_{12} are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group, or bond to adjacent groups to form a fused ring;


[0103] the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkoxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_{11} and R_{12} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other; and

[0104] Y_1 , Y_3 and Y_5 each have the same definitions as in Chemical Formula 15.


[0105] According to preferred one embodiment of the present invention, the substituent represented by Chemical Formula 15 may be a substituent represented by any one of the following

Chemical Formulae C-1 to C-5:


[0106]




C-1




C-2



C-3



C-4



C-5

[0107] in Chemical Formulae C-1 to C-5,

[0108] * means a part where a bond is formed,

[0109] R_{11} and R_{12} are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkyloxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group, or bond to adjacent groups to form a fused ring; and

[0110] the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_{11} and R_{12} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group,

a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkyloxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.

[0111] According to preferred one embodiment of the present invention, R_{11} and R_{12} are each independently selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and

[0112] the alkyl group, the aryl group and the heteroaryl group of R_{11} and R_{12} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these are the same as or different from each other.

[0113] According to preferred one embodiment of the present invention, R_{11} and R_{12} are each independently selected from the group consisting of a phenyl group, a biphenyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, a carbazolyl group, a fluorenyl group and a dibenzothiophenyl group, and


[0114] the phenyl group, the biphenyl group, the pyridinyl group, the pyrimidinyl group, the dibenzofuranyl group, the carbazolyl group, the fluorenyl group and the dibenzothiophenyl group of R_{11} and R_{12} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these are the same as or different from each other.

[0115] According to preferred one embodiment of the present invention, R_{11} and R_{12} are each independently selected from the group consisting of a phenyl group, a biphenyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, a carbazolyl group, a fluorenyl group and a dibenzothiophenyl group, and

[0116] the phenyl group, the biphenyl group, the pyridinyl group, the pyrimidinyl group, the dibenzofuranyl group, the carbazolyl group, the fluorenyl group and the dibenzothiophenyl group of R_{11} and R_{12} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a methyl group, an ethyl group, a propanyl group, a butyl group, a phenyl group, a biphenyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, a carbazolyl group, a fluorenyl group and a dibenzothiophenyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.

[0117] According to preferred one embodiment of the present invention, Ar_2 may be a substituent represented by the following Chemical Formula 17:

[Chemical Formula 17]



[0118] in Chemical Formula 17,

[0119] * means a part where a bond is formed;

[0120] R_{13} and R_{14} are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkyloxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group, or bond to adjacent groups to form a fused ring; and

[0121] the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_{13} and R_{14} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkyloxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.

[0122] According to preferred one embodiment of the present invention, R_{13} and R_{14} are each independently selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and

[0123] the alkyl group, the aryl group and the heteroaryl group of R_{13} and R_{14} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these are the same as or different from each other.

[0124] According to preferred one embodiment of the present invention, R_{13} and R_{14} are each independently selected from the group consisting of a phenyl group, a biphenyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, a carbazolyl group, a fluorenyl group and a dibenzothiophenyl group, and


[0125] the phenyl group, the biphenyl group, the pyridinyl group, the pyrimidinyl group, the dibenzofuranyl group, the carbazolyl group, the fluorenyl group and the dibenzothiophenyl group of R_{13} and R_{14} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these are the same as or different from each other.

[0126] According to preferred one embodiment of the present invention, R_{13} and R_{14} are each independently selected from the group consisting of a phenyl group, a biphenyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, a carbazolyl group, a fluorenyl group and a dibenzothiophenyl group, and


[0127] the phenyl group, the biphenyl group, the pyridinyl group, the pyrimidinyl group, the dibenzofuranyl group, the carbazolyl group, the fluorenyl group and the dibenzothiophenyl group of R_{13} and R_{14} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a methyl group, an ethyl group, a propyl group, a butyl group, a phenyl group, a biphenyl group, a pyridinyl group, a pyrimidinyl group, a dibenzofuranyl group, a carbazolyl group, a fluorenyl group and a dibenzothiophenyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.

[0128] The compound represented by Chemical Formula 1 of the present invention may be represented by the following compounds, but is not limited thereto:




R1



R2




-continued



-continued


R6

R3




R7

R4




R8


R5



R9




-continued




R10


-continued




R13




R11




R14



R15




R12








R16


-continued




-continued




-continued



R23




R24




R25


-continued




R26



R27




R28




R29

-continued




R30


-continued




R33




R31




R34



R35




R32




R36

-continued





-continued



R40




R41



R42




-continued




R43

-continued




R46

R47





R44

R48




R45

R49




-continued





R50

-continued




R53

R51





R54

R55




R56






R52






-continued



-continued




-continued



-continued


R66

R63




R67

R64




R68


R65



R69




-continued




R70


-continued




R73




R71




R74



R75




R72








R76


-continued



-continued




-continued




R103


-continued




R106




R104



R107




R105




R108


-continued



R109




R110




R111


-continued



R112




R113




R114

-continued




-continued

R116c1ccc(cc1)-c2ccc(cc2)-c3cc(ccc3N2C(=N1C(=N(C(=N1)C2)C4=CC=CC=C4)C5=CC=CC=C5)C6=CC=CC=C6)C7=CC=CC=C7R119c1ccc(cc1)-c2ccc(cc2)-c3cc(ccc3N2C(=N1C(=N(C(=N1)C2)C4=CC=CC=C4)C5=CC=CC=C5)C6=CC=CC=C6)C7=CC=CC=C7R117c1ccc(cc1)-c2ccc(cc2)-c3cc(ccc3N2C(=N1C(=N(C(=N1)C2)C4=CC=CC=C4)C5=CC=CC=C5)C6=CC=CC=C6)C7=CC=CC=C7R120c1ccc(cc1)-c2ccc(cc2)-c3cc(ccc3N2C(=N1C(=N(C(=N1)C2)C4=CC=CC=C4)C5=CC=CC=C5)C6=CC=CC=C6)C7=CC=CC=C7


-continued

R121




-continued


R124




R122




R125



R123




R126




-continued

R127




-continued


R130




R128




R131







R129






R132



-continued




-continued




-continued


R140



R141




R142




-continued


R143



R144




R145




-continued


R146




R147



R148




R149




-continued


R150







R151






R152




-continued




-continued




-continued




-continued




R161




R164



R162




R165




-continued


R166




R167



R168




R169




-continued


R170




R171




R172




-continued




R174




R175




R176




-continued



R178




R179




-continued


R180



R181




R182




-continued


R183



R184




R185




-continued


R186




R187



R188




R189




-continued


R190




R191



R192




-continued




-continued


R196




R194




R197




R195




R198




-continued




-continued




R200




R203




R201



R204




-continued




R212


-continued




R215




R213



R216




R214








R217

-continued




-continued




-continued

R224




-continued


R228




R225




R229




R226



R227




R230




-continued

R231




-continued


R234




R235




R236




R232








R233




-continued



-continued




-continued




R243


-continued




R245



R244




R246




R247



-continued




-continued




R252




R253




R254



-continued




-continued

R256c1ccc(cc1)-n2c3ccccc3cc2c4ccccc4-c5ccc(cc5)-c6ccc(cc6)-c7ccc(cc7)-c8ccccc8-c9nnnc9-c10ccccc10-c11ccccn11R259c1ccc(cc1)-n2c3ccccc3cc2c4ccccc4-c5ccc(cc5)-c6ccc(cc6)-c7ccc(cc7)-c8ccccc8-c9nnnc9-c10ccccc10-c11ccc(cc11)-c12ccccc12-s13ccccc13R257c1ccc(cc1)-n2c3ccccc3cc2c4ccccc4-c5ccc(cc5)-c6ccc(cc6)-c7ccc(cc7)-c8ccccc8-c9nnnc9-c10ccccc10-c11ccccn11R260c1ccc(cc1)-n2c3ccccc3cc2c4ccccc4-c5ccc(cc5)-c6ccc(cc6)-c7ccc(cc7)-c8ccccc8-c9nnnc9-c10ccccc10-c11ccc(cc11)-c12ccccc12-s13ccccc13


-continued

R261




-continued


R264



R262





R265




R266


R263




R267




-continued




-continued




R269




R272









R270



R273




-continued




-continued

R280




-continued


R283




R281




R284



R282




R285




-continued

R286




-continued


R289




R287




R290




R288



R291




-continued




R292

-continued





R295


R293



R294




R296




R297

-continued



-continued

R300



[0

129] The compound of Chemical Formula 1 of the present invention may be synthesized using general synthesis methods (refer to *Chem. Rev.*, 60:313 (1960); *J. Chem. SOC.* 4482 (1955); *Chem. Rev.* 95: 2457 (1995) or the like). Detailed synthesis processes of the compounds of the present invention will be specifically described in synthesis examples to be described later.

[0130] 2. Organic Electroluminescent Device

[0131] Meanwhile, another aspect of the present invention relates to an organic electroluminescent device (organic EL device) including the compound represented by Chemical Formula 1 according to the present invention.


[0132] Specifically, the present invention relates to an organic electroluminescent device including an anode, a cathode, and one or more organic material layers provided between the anode and the cathode, and at least one of the one or more organic material layers includes the compound represented by Chemical Formula 1. Herein, the compound may be used either alone or as a mixture of two or more.

[0133] The one or more organic material layers may be any one or more of a hole injection layer, a hole transport layer, a light emitting layer, a light emitting auxiliary layer, a lifetime improving layer, an electron transport layer, an electron transport auxiliary layer and an electron injection layer, and at least one organic material layer among these may include the compound represented by Chemical Formula 1.

[0134] The structure of the organic electroluminescent device according to the present invention described above is not particularly limited, but, when referring to FIG. 1 as one example, includes an anode (10) and a cathode (20) facing each other, and an organic layer (30) located between the anode (10) and the cathode (20). Herein, the organic layer (30) may include a hole transport layer (31), a light emitting layer (32) and an electron transport layer (34). In addition, a hole transport auxiliary layer (33) may be included between the hole transport layer (31) and the light emitting layer (32), and an electron transport auxiliary layer (35) may be included between the electron transport layer (34) and the light emitting layer (32).

[0135] When referring to FIG. 2 as another example of the present invention, the organic layer (30) may further include

R299



a hole injection layer (37) between the hole transport layer (31) and the anode (10), and may further include an electron injection layer (36) between the electron transport layer (34) and the cathode (20).

[0136] The hole injection layer (37) laminated between the hole transport layer (31) and the anode (10) in the present invention is a layer having a function of, as well as improving interfacial properties between ITO used as the anode and an organic material used as the hole transport layer (31), smoothing the ITO surface by being coated on the top of the ITO of which surface is not smooth, and those commonly used in the art may be used without particular limit, and for example, amine compounds may be used. However, the hole injection layer is not limited thereto.

[0137] In addition, the electron injection layer (36) is a layer laminated on the top of the electron transport layer and having a function of facilitating electron injection from the cathode and eventually improving power efficiency, and is not particularly limited as long as it is commonly used in the art. For example, materials such as LiF, Liq, NaCl, CsF, Li₂O or BaO may be used.

[0138] Although not shown in the drawings in the present invention, a light emitting auxiliary layer may be further included between the hole transport auxiliary layer (33) and the light emitting layer (32). The light emitting auxiliary layer may perform a role of adjusting a thickness of the organic layer (30) while performing a role of transporting holes to the light emitting layer (32). The light emitting auxiliary layer may include a hole transport material, and may be formed with the same material as the hole transport layer (31).

[0139] In addition, although not shown in the drawings in the present invention, a lifetime improving layer may be further included between the electron transport auxiliary layer (35) and the light emitting layer (32). Holes migrating to the light emitting layer (32) by getting on an ionization potential level in an organic light emitting device are not able to diffuse or migrate to the electron transport layer by being blocked by a high energy barrier of the lifetime improving layer, and consequently, the lifetime improving layer has a function of limiting the holes in the light emitting layer. Such a function of limiting the holes in the light emitting layer prevents the holes from diffusing to the electron transport layer migrating electrons by reduction, and therefore, suppresses a lifetime decrease phenomenon caused through an irreversible decomposition reaction by oxidation, and thereby contributes to improving a lifetime of the organic light emitting device.

[0140] The compound represented by Chemical Formula 1 in the present invention has an EWG bonding to a 5-membered aromatic ring or a 5-membered aromatic heteroring such as indole, indazole, indene, benzofuran, benzothiophene or triazolo, and may be used as a host material since, by having a similar energy level with carbazole, the energy level may be adjusted to be higher than a dopant energy level. Particularly, benzofuran and benzothiophene moieties are electron abundant, and when used as an electron transport layer material of an organic electroluminescent device, mobility increases, and as a result, an increase in the luminous efficiency and a decrease in the driving voltage may be expected. In addition, the 5-membered aromatic ring or the 5-membered aromatic heteroring of the present invention has a smaller molecular weight compared to existing compounds, and therefore, may be deposited at a relatively

lower deposition temperature when deposited compared to other materials, and favorable processability and enhanced thermal stability may be obtained.

[0141] Accordingly, the compound represented by Chemical Formula 1 of the present invention may be used as a material of any one of a hole injection layer, a hole transport layer, a light emitting layer, an electron transport layer and an electron injection layer, an organic material layer of an organic electroluminescent device, but may be preferably used as a material of any of a light emitting layer, an electron transport layer, and an electron transport auxiliary layer further laminated on the electron transport layer, and more preferably used as a material of an electron transport layer or an electron transport auxiliary layer.

[0142] In addition, when using the compound according to the present invention as a light emitting layer material, the compound represented by Chemical Formula 1 may be specifically used as a phosphorescent host, a fluorescent host or a dopant material of the light emitting layer, and may be preferably used as a phosphorescent host (blue, green and/or red phosphorescent host material).

[0143] In addition, the organic electroluminescent device in the present invention has, as described above, an anode, one or more organic material layers and a cathode consecutively laminated, and in addition thereto, may further include an insulating layer or an adhesive layer at an interface between the electrode and the organic material layer.

[0144] Except that at least one or more of the organic material layers (for example, electron transport auxiliary layer) are formed to include the compound represented by Chemical Formula 1, the organic electroluminescent device of the present invention may be manufactured by forming other organic material layers and electrodes using materials and methods known in the art.

[0145] The organic material layer may be formed using a vacuum deposition method or a solution coating method. Examples of the solution coating method may include spin coating, dip coating, doctor blading, inkjet printing, thermal transfer method or the like, but are not limited thereto.

[0146] A substrate capable of being used in the present invention is not particularly limited, and silicon wafers, quartz, glass plates, metal plates, plastic films, sheets and the like may be used.

[0147] The anode material may be prepared using, for example, a conductor having high work function so as to have smooth hole injection, and examples thereof may include metals such as vanadium, chromium, copper, zinc or gold, or alloys thereof; metal oxides such as zinc oxide, indium oxide, indium tin oxide (ITO) or indium zinc oxide (IZO); combinations of metals and oxides such as ZnO:Al or SnO₂:Sb; conductive polymers such as polythiophene, poly(3-methylthiophene), poly[3,4-(ethylene-1,2-dioxy)thiophene] (PEDT), polypyrrole or polyaniline; carbon black, and the like, but are not limited thereto.

[0148] The cathode material may be prepared using, for example, a conductor having low work function so as to have smooth electron injection, and examples thereof may include metals such as magnesium, calcium, sodium, potassium, titanium, indium, yttrium, lithium, gadolinium, aluminum, silver, tin or lead, or alloys thereof; and multilayer-structured materials such as LiF/Al or LiO₂/Al, but are not limited thereto.


[0149] Hereinafter, the present invention will be described in detail with reference to examples as follows. However,

the following examples are for illustrative purposes only, and the present invention is not limited to the following examples.

PREPARATION EXAMPLE 1

Synthesis of A1

[0150]



[0151] Under a nitrogen stream, 3-(3-bromophenyl)-1-phenyl-1H-indazole (8.5 g, 24.4 mmol), 4,4',4'',5,5',5'',8-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0152] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A1 (6.8 g, 17.1 mmol, yield 70%).


[0153] GC-Mass (theoretical value: 396.3 g/mol, measured value: 396 g/mol)

[0154] ¹H-NMR: δ 1.25 (s, 12H), 7.45~7.55 (m, 8H), 7.62 (s, 1H), 7.84~9.2 (m, 4H)

PREPARATION EXAMPLE 2

Synthesis of A2

[0155]



[0156] Under a nitrogen stream, 3-(3-bromophenyl)-1-(pyridin-4-yl)-1H-indazole (8.5 g, 24.4 mmol), 4,4',4'',5,5',5'',8-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0157] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A2 (7.3 g, 18.3 mmol, yield 75%).


[0158] GC-Mass (theoretical value: 397.3 g/mol, measured value: 397 g/mol)

[0159] ¹H-NMR: δ 1.25 (s, 12H), 7.45~7.55 (m, 5H), 7.62 (s, 1H), 7.84~9.2 (m, 4H), 8.11~12 (d, 2H)

PREPARATION EXAMPLE 3

Synthesis of A3

[0160]



[0161] Under a nitrogen stream, 3-(3-bromophenyl)-1,5-diphenyl-1H-indazole (10.4 g, 24.4 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0162] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A3 (8.6 g, 18.3 mmol, yield 75%).


[0163] GC-Mass (theoretical value: 472.4 g/mol, measured value: 472 g/mol)

[0164] ¹H-NMR: δ 1.25 (s, 12H), 7.44~7.59 (m, 14H), 7.62 (s, 1H), 8.23~25 (d, 2H)

PREPARATION EXAMPLE 4

Synthesis of A4

[0165]



[0166] Under a nitrogen stream, 1-(3-bromophenyl)-3-phenyl-3H-benzo[e]indazole (9.7 g, 24.4 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0167] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A4 (7.6 g, 17.1 mmol, yield 70%).


[0168] GC-Mass (theoretical value: 446.4 g/mol, measured value: 446 g/mol)

[0169] ¹H-NMR: δ 1.25 (s, 12H), 7.45~7.55 (m, 10H), 7.62 (s, 1H), 7.81~99 (m, 4H)

PREPARATION EXAMPLE 5

Synthesis of A5

[0170]



[0171] Under a nitrogen stream, 1-(3-bromophenyl)-3-(pyridin-4-yl)-3H-benzo[e]indazole (9.8 g, 24.4 mmol), 4,4',4'',5,5',5'',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0172] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A5 (7.6 g, 17.1 mmol, yield 70%).


[0173] GC-Mass (theoretical value: 447.4 g/mol, measured value: 447 g/mol)

[0174] ¹H-NMR: δ 1.25 (s, 12H), 7.45~7.55 (m, 9H), 7.62 (s, 1H), 7.81~9.9 (m, 2H), 8.21~23 (d, 2H)

PREPARATION EXAMPLE 6

Synthesis of A6

[0175]



[0176] Under a nitrogen stream, 3-(3-bromophenyl)-1-phenyl-1H-benzo[f]indazole (9.7 g, 24.4 mmol), 4,4',4'',5,5',5'',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0177] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A6 (7.6 g, 17.1 mmol, yield 70%).


[0178] GC-Mass (theoretical value: 446.4 g/mol, measured value: 446 g/mol)

[0179] ¹H-NMR: δ 1.25 (s, 12H), 7.45~7.55 (m, 10H), 7.58 (s, 1H), 7.61 (s, 1H), 7.78 (s, 1H), 7.81~8.6 (m, 2H)

PREPARATION EXAMPLE 7

Synthesis of A7

[0180]



[0181] Under a nitrogen stream, 3-(3-bromophenyl)-1-phenyl-1H-benzotriazole (9.7 g, 24.4 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0182] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A7 (7.6 g, 17.1 mmol, yield 70%).


[0183] GC-Mass (theoretical value: 446.36 g/mol, measured value: 446 g/mol)

[0184] ¹H-NMR: δ 1.25 (s, 12H), 7.45~7.54 (m, 10H), 7.62 (s, 1H), 7.81~88 (m, 4H)

PREPARATION EXAMPLE 8

Synthesis of A8

[0185]



[0186] Under a nitrogen stream, 3-(3-bromophenyl)-1-phenyl-1H-indole (8.5 g, 24.4 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0187] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A8 (6.7 g, 17.1 mmol, yield 70%).


[0188] GC-Mass (theoretical value: 395.3 g/mol, measured value: 395 g/mol)

[0189] ¹H-NMR: δ 1.25 (s, 12H), 7.43~7.55 (m, 9H), 7.62 (s, 1H), 7.84~92 (m, 4H)

PREPARATION EXAMPLE 9

Synthesis of A9

[0190]



[0191] Under a nitrogen stream, 3-(3-bromophenyl)-1-(pyridin-4-yl)-1H-indole (8.5 g 24.4 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0192] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A9 (6.8 g, 17.1 mmol, yield 70%).


[0193] GC-Mass (theoretical value: 396.3 g/mol, measured value: 396 g/mol)

[0194] ¹H-NMR: δ 1.25 (s, 12H), 7.41~7.55 (m, 6H), 7.62 (s, 1H), 7.84~9.2 (m, 4H), 8.11~12 (d, 2H)

PREPARATION EXAMPLE 10

Synthesis of A10

[0195]



[0196] Under a nitrogen stream, 3-(3-bromophenyl)-1,5-diphenyl-1H-indole (10.3 g, 24.4 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0197] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A10 (8.6 g, 18.3 mmol, yield 75%).


[0198] GC-Mass (theoretical value: 471.4 g/mol, measured value: 471 g/mol)

[0199] ¹H-NMR: δ 1.25 (s, 12H), 7.41~7.60 (m, 15H), 7.63 (s, 1H), 8.23~25 (d, 2H)

PREPARATION EXAMPLE 11

Synthesis of A11

[0200]



[0201] Under a nitrogen stream, 1-(3-bromophenyl)-3-phenyl-3H-benzo[e]indole (9.7 g, 24.4 mmol), 4,4',4'',5,5',5'',8-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0202] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A11 (7.6 g, 17.1 mmol, yield 70%).


[0203] GC-Mass (theoretical value: 445.4 g/mol, measured value: 445 g/mol)

[0204] ¹H-NMR: δ 1.25 (s, 12H), 7.41~7.55 (m, 11H), 7.62 (s, 1H), 7.81~9.9 (m, 4H)

PREPARATION EXAMPLE 12

Synthesis of A12

[0205]



[0206] Under a nitrogen stream, 1-(3-bromophenyl)-3-(pyridin-4-yl)-3H-benzo[e]indazole (9.7 g, 24.4 mmol), 4,4',4'',5,5',5'',8-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0207] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A12 (7.6 g, 17.1 mmol, yield 70%).


[0208] GC-Mass (theoretical value: 446.4 g/mol, measured value: 446 g/mol)

[0209] ¹H-NMR: δ 1.25 (s, 12H), 7.42~7.55 (m, 10H), 7.62 (s, 1H), 7.81~9.9 (m, 2H), 8.21~23 (d, 2H)

PREPARATION EXAMPLE 13

Synthesis of A13

[0210]



[0211] Under a nitrogen stream, 3-(3-bromophenyl)-1-phenyl-1H-benzo[f]indole (9.7 g, 24.4 mmol), 4,4,4',4',5,5',5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0212] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A13 (7.6 g, 17.1 mmol, yield 70%).


[0213] GC-Mass (theoretical value: 445.4 g/mol, measured value: 445 g/mol)

[0214] ¹H-NMR: δ 1.25 (s, 12H), 7.42~7.55 (m, 11H), 7.58 (s, 1H), 7.61 (s, 1H), 7.78 (s, 1H), 7.81~8.5 (m, 2H)

PREPARATION EXAMPLE 14

Synthesis of A14

[0215]



[0216] Under a nitrogen stream, 3-(3-bromophenyl)-1-phenyl-1H-benzo[g]indazole (9.7 g, 24.4 mmol), 4,4,4',4',5,5',5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0217] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A14 (7.6 g, 17.1 mmol, yield 70%).


[0218] GC-Mass (theoretical value: 445.37 g/mol, measured value: 445 g/mol)

[0219] ¹H-NMR: δ 1.25 (s, 12H), 7.42~7.54 (m, 11H), 7.62 (s, 1H), 7.82~8.9 (m, 4H)

PREPARATION EXAMPLE 15

Synthesis of A15

[0220]



[0221] Under a nitrogen stream, 2-(3-bromophenyl)benzofuran (6.7 g, 24.4 mmol), 4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0222] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A15 (5.5 g, 17.1 mmol, yield 70%).


[0223] GC-Mass (theoretical value: 320.2 g/mol, measured value: 320 g/mol)

[0224] ¹H NMR: δ 1.25 (s, 12H), 7.11 (s, 1H), 7.58 (s, 1H), 7.67~7.89 (m, 7H)


PREPARATION EXAMPLE 16

Synthesis of A16

[0225]



-continued



[0226] Under a nitrogen stream, 2-(3-bromophenyl)-1,1-dimethyl-1H-indene (7.3 g, 24.4 mmol), 4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ 0.6 g, 0.7 mmol, KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0227] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed using MgSO₄, and the result was purified using column chromatography to obtain target Compound A16 (5.9 g, 17.1 mmol, yield 70%).


[0228] GC-Mass (theoretical value: 346.28 g/mol, measured value: 346 g/mol)

[0229] ¹H-NMR: δ 1.25 (s, 12H), 1.69 (s, 6H), 7.01 (s, 1H), 7.45 (s, 1H), 7.57~7.77 (m, 7H)

PREPARATION EXAMPLE 17

Synthesis of A17

[0230]



[0231] Under a nitrogen stream, 2-(3-bromophenyl)benzo[b]thiophene (7.0 g, 24.4 mmol), 4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.4 g, 29.2 mmol), Pd(dppf)Cl₂ (0.6 g, 0.7 mmol), KOAc (7.2 g, 73.1 mmol) and 1,4-dioxane (200 ml) were mixed, and stirred for 6 hours at 130° C.

[0232] After the reaction was finished, the result was extracted with ethyl acetate, the moisture was removed

using MgSO_4 , and the result was purified using column chromatography to obtain target Compound A17 (6.1 g, 18.3 mmol, yield 75%).


[0233] GC-Mass (theoretical value: 336.26 g/mol, measured value: 336 g/mol)

[0234] $^1\text{H-NMR}$: δ 1.25 (s, 12H), 7.51 (s, 1H), 7.59 (s, 1H), 7.72~7.92 (m, 7H)

SYNTHESIS EXAMPLE 1

Synthesis of R1

[0235]



[0236] Under a nitrogen stream, A1 (6.8 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g 18.8 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/ H_2O /ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110°C .


[0237] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO_4 . After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R1 (8.4 g, 12.8 mmol, yield 75%).

[0238] GC-Mass (theoretical value: 653.8 g/mol, measured value: 653 g/mol)

SYNTHESIS EXAMPLE 2

Synthesis of R8

[0239]



[0240] Under a nitrogen stream, A1 (6.8 g 17.1 mmol), 2-bromo-4,6-bis(dibenzo[b,d]furan-4-yl)-1,3,5-triazine (9.2 g 18.8 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/ H_2O /ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110°C .


[0241] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO_4 . After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R8 (9.0 g, 13.1 mmol, yield 77%).

[0242] GC-Mass (theoretical value: 681.8 g/mol, measured value: 682 g/mol)

SYNTHESIS EXAMPLE 3

Synthesis of R21

[0243]



[0244] Under a nitrogen stream, A2 (6.8 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), $\text{Pd(PPh}_3)_4$ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/ H_2O /ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110°C .


[0245] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO_4 . After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R21 (7.8 g, 11.9 mmol, yield 70%).

[0246] GC-Mass (theoretical value: 654.8 g/mol, measured value: 655 g/mol)

SYNTHESIS EXAMPLE 4

Synthesis of R41

[0247]



[0248] Under a nitrogen stream, A3 (8.6 g, 18.3 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (9.3 g, 20.1 mmol), $\text{Pd(PPh}_3)_4$ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/ H_2O /ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110°C .


[0249] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO_4 . After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R41 (10.0 g, 13.7 mmol, yield 75%).

[0250] GC-Mass (theoretical value: 729.9 g/mol, measured value: 730 g/mol)

SYNTHESIS EXAMPLE 5

Synthesis of R61

[0251]



R61

[0252] Under a nitrogen stream, A4 (7.6 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/ H_2O /ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110°C .


[0253] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO_4 . After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R61 (9.0 g, 12.8 mmol, yield 75%).

[0254] GC-Mass (theoretical value: 703.9 g/mol, measured value: 703 g/mol)

SYNTHESIS EXAMPLE 6

Synthesis of R68

[0255]



R68

[0256] Under a nitrogen stream, A4 (7.6 g, 17.1 mmol), 2-bromo-4,6-bis(dibenzo[b,d]furan-4-yl)-1,3,5-triazine (9.2 g, 18.8 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.0 g, 5 mol %), potassium

carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0257] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R68 (9.4 g, 12.8 mmol, yield 75%).

[0258] GC-Mass (theoretical value: 731.8 g/mol, measured value: 731 g/mol)

SYNTHESIS EXAMPLE 7

Synthesis of R81

[0259]



[0260] Under a nitrogen stream, A5 (7.6 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0261] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R81 (8.9 g 12.6 mmol, yield 74%).

[0262] GC-Mass (theoretical value: 703.9 g/mol, measured value: 704 g/mol)

SYNTHESIS EXAMPLE 8

Synthesis of R101

[0263]



[0264] Under a nitrogen stream, A6 (7.6 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0265] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R101 (9.0 g, 12.8 mmol, yield 75%).

[0266] GC-Mass (theoretical value: 703.9 g/mol, measured value: 703 g/mol)

SYNTHESIS EXAMPLE 9

Synthesis of R121

[0267]



[0268] Under a nitrogen stream, A7 (7.6 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0269] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R121 (8.9 g, 12.0 mmol, yield 74%).

[0270] GC-Mass (theoretical value: 703.9 g/mol, measured value: 703 g/mol)

SYNTHESIS EXAMPLE 10

Synthesis of R141

[0271]



[0272] Under a nitrogen stream, A8 (6.7 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0273] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R141 (8.3 g, 12.8 mmol, yield 75%).

[0274] GC-Mass (theoretical value: 652.8 g/mol, measured value: 653 g/mol)

SYNTHESIS EXAMPLE 11

Synthesis of R148

[0275]



[0276] Under a nitrogen stream, A8 (6.7 g, 17.1 mmol), 2-bromo-4,6-bis(dibenzo[b,d]furan-4-yl)-1,3,5-triazine (9.2 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0277] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R148 (8.7 g, 12.8 mmol, yield 75%).

[0278] GC-Mass (theoretical value: 680.8 g/mol, measured value: 681 g/mol)

SYNTHESIS EXAMPLE 12

Synthesis of R161

[0279]



[0280] Under a nitrogen stream, A9 (6.8 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0281] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R161 (7.9 g, 11.1 mmol, yield 71%).

[0282] GC-Mass (theoretical value: 653.8 g/mol, measured value: 654 g/mol)

SYNTHESIS EXAMPLE 13

Synthesis of R181

[0283]



[0284] Under a nitrogen stream, A10 (8.6 g, 18.3 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (9.3 g, 20.1 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0285] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R181 (10.0 g, 13.7 mmol, yield 75%).

[0286] GC-Mass (theoretical value: 729.9 g/mol, measured value: 730 g/mol)

SYNTHESIS EXAMPLE 14

Synthesis of R201

[0287]



[0288] Under a nitrogen stream, A11 (7.6 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0289] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R201 (9.0 g, 12.8 mmol, yield 75%).

[0290] GC-Mass (theoretical value: 703.9 g/mol, measured value: 703 g/mol)

SYNTHESIS EXAMPLE 15

Synthesis of R208

[0291]



[0292] Under a nitrogen stream, A11 (7.6 g 17.1 mmol), 2-bromo-4,6-bis(dibenzo[b,d]furan-4-yl)-1,3,5-triazine (9.2 g 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0293] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R208 (9.4 g, 12.8 mmol, yield 75%).

[0294] GC-Mass (theoretical value: 731.8 g/mol, measured value: 731 g/mol)

SYNTHESIS EXAMPLE 16

Synthesis of R221

[0295]



[0296] Under a nitrogen stream, A12 (7.6 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0297] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R221 (8.9 g, 12.6 mmol, yield 74%).

[0298] GC-Mass (theoretical value: 703.9 g/mol, measured value: 704 g/mol)

SYNTHESIS EXAMPLE 17

Synthesis of R241

[0299]



[0300] Under a nitrogen stream, A13 (7.6 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0301] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R241 (9.0 g, 12.8 mmol, yield 75%).

[0302] GC-Mass (theoretical value: 702.9 g/mol, measured value: 703 g/mol)

SYNTHESIS EXAMPLE 18

Synthesis of R261

[0303]



[0304] Under a nitrogen stream, A14 (7.6 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0305] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R261 (8.9 g, 12.0 mmol, yield 74%).

[0306] GC-Mass (theoretical value: 702.9 g/mol, measured value: 703 g/mol)

SYNTHESIS EXAMPLE 19

Synthesis of R281

[0307]



[0308] Under a nitrogen stream, A15 (5.5 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium

carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0309] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO₄. After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R281 (9.9 g, 12.8 mmol, yield 75%).

[0310] GC-Mass (theoretical value: 577.7 g/mol, measured value: 578 g/mol)

SYNTHESIS EXAMPLE 20

Synthesis of R285

[0311]



[0312] Under a nitrogen stream, A15 (5.5 g, 17.1 mmol), 2-bromo-4,6-bis(dibenzo[b,d]furan-4-yl)-1,3,5-triazine (9.2 g, 18.8 mmol), Pd(PPh₃)₄ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/H₂O/ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110° C.


[0313] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO_4 . After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R285 (8.0 g, 13.1 mmol, yield 77%).

[0314] GC-Mass (theoretical value: 605.7 g/mol, measured value: 606 g/mol)

SYNTHESIS EXAMPLE 21

Synthesis of R286

[0315]



[0316] Under a nitrogen stream, A16 (5.9 g, 17.1 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (8.7 g, 18.8 mmol), $\text{Pd(PPh}_3)_4$ (1.0 g, 5 mol %), potassium carbonate (7.1 g, 51.2 mmol) and toluene/ H_2O /ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110°C .

[0317] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO_4 . After the solvent of the organic layer


was removed, the result was purified using column chromatography to obtain target Compound R286 (7.2 g, 11.9 mmol, yield 70%).

[0318] GC-Mass (theoretical value: 603.8 g/mol, measured value: 604 g/mol)

SYNTHESIS EXAMPLE 22

Synthesis of R291

[0319]



[0320] Under a nitrogen stream, A17 (6.1 g, 18.3 mmol), 2-(3'-bromobiphenyl-3-yl)-4,6-diphenyl-1,3,5-triazine (9.3 g, 20.1 mmol), $\text{Pd(PPh}_3)_4$ (1.0 g, 5 mol %), potassium carbonate (7.6 g, 53.1 mmol) and toluene/ H_2O /ethanol (80 ml/40 ml/40 ml) were introduced, and stirred for 3 hours at 110°C .

[0321] After the reaction was finished, the organic layer was separated using methylene chloride, and water was removed using MgSO_4 . After the solvent of the organic layer was removed, the result was purified using column chromatography to obtain target Compound R291 (9.2 g, 13.7 mmol, yield 75%).

[0322] GC-Mass (theoretical value: 669.9 g/mol, measured value: 670 g/mol)

EXAMPLES 1 TO 6


Manufacture of Green Organic Electroluminescent Device

[0323] After high purity sublimation purifying the compounds synthesized in the synthesis examples using commonly known methods, green organic electroluminescent devices were manufactured using the following procedure.


[0324] First, a glass substrate on which indium tin oxide (ITO) was coated as a thin film to a thickness of 1500 Å was ultrasonic cleaned using distilled water. After the cleaning with distilled water was finished, the substrate was ultrasonic cleaned with solvents of isopropyl alcohol, acetone, methanol and the like, dried, then transferred to a UV OZONE washer (Power sonic 405, manufactured by Hwashin Tech. Co., Ltd.), and then, after cleaning the substrate for 5 minutes using UV, the substrate was transferred to a vacuum deposition apparatus.

[0325] On the transparent ITO electrode prepared as above, m-MTDATA (60 nm)/TCTA (80 nm)/each compound of R1, R8, R141, R148, R201 and R208+10% Ir(ppy)₃ (300 nm)/BCP (10 nm)/Alq₃ (30 nm)/LiF (1 nm)/Al (200 nm) were laminated in this order to manufacture an organic electroluminescent device.

[0326] Structures of m-MTDATA, TCTA, Ir(ppy)₃, CBP and BCP are as follows.





m-MTDATA




TCTA

-continued

Ir(ppy)₃

CBP



BCP

COMPARATIVE EXAMPLE 1

Manufacture of Green Organic Electroluminescent Device

Device

[0327] A green organic electroluminescent device was manufactured in the same manner as in Example 1 except that CBP was used instead of Compound R141 as the light emitting host material when forming the light emitting layer.

EVALUATION EXAMPLE 1

[0328] For each of the green organic electroluminescent devices manufactured in Examples 1 to 6 and Comparative Example 2, driving voltage, current efficiency and light emission peak at current density of (10) mA/cm² were measured, and the results are shown in the following Table 1.

TABLE 1

Sample	Host	Driving Voltage (V)	EL Peak (nm)	Current Efficiency (cd/A)
Example 1	R1	5.2	515	12.5
Example 2	R8	4.3	515	11.4
Example 3	R141	5.1	515	12.3
Example 4	R148	4.2	515	11.1

TABLE 1-continued

Sample	Host	Driving Voltage (V)	EL Peak (nm)	Current Efficiency (cd/A)
Example 5	R201	5.4	515	12.2
Example 6	R208	4.4	515	10.8
Comparative Example 1	CBP	7.1	516	7.2

[0329] As shown in Table 1, it was seen that using the compound (R1, R8, R141, R148, R201, R208) according to the present invention in a light emitting layer of the green organic EL device (Examples 1 to 6) resulted in more superior performance in terms of efficiency and driving voltage compared to the green organic EL device using existing CBP (Comparative Example 1).

EXAMPLES 7 TO 28

Manufacture of Blue Organic Electroluminescent Device

[0330] After high purity sublimation purifying the compounds synthesized in the synthesis examples using commonly known methods, blue organic electroluminescent devices were manufactured using the following procedure.

[0331] First, a glass substrate on which indium tin oxide (ITO) was coated as a thin film to a thickness of 1500 Å was ultrasonic cleaned using distilled water. After the cleaning with distilled water was finished, the substrate was ultrasonic cleaned with solvents of isopropyl alcohol, acetone, methanol and the like, dried, then transferred to a UV OZONE washer (Power sonic 405, manufactured by Hwashin Tech. Co., Ltd), and then, after cleaning the substrate for 5 minutes using UV, the substrate was transferred to a vacuum deposition apparatus.



[0332] On the transparent ITO electrode prepared as above, DS-205 (Doosan Corporation Electro-Materials, 80 nm)/NPB (15 nm)/ADN+5% DS-405 (Doosan Corporation Electro-Materials, 30 nm)/each compound of R1, R8, R21, R41, R61, R68, R81, R101, R121, R141, R141, R148, R161, R181, R201, R208, R221, R241, R261, R281, R285, R286 and R291 (5 nm)/Alq₃ (25 nm)/LiF (1 nm)/Al (200 nm) were laminated in this order to manufacture an organic electroluminescent device.

COMPARATIVE EXAMPLE 2

Manufacture of Blue Organic Electroluminescent Device

[0333] A blue organic electroluminescent device was manufactured in the same manner as in Example 6 except that R1 was not used as the electron transport auxiliary layer material, and Alq₃, the electron transport layer material, was deposited to 30 nm instead of 25 nm.


[0334] Structures of NPB, AND and Alq₃ used in Examples 6 to 28 and Comparative Example 2 are as follows.



COMPARATIVE EXAMPLE 3

Manufacture of Blue Organic Electroluminescent Device

[0335] A blue organic electroluminescent device was manufactured in the same manner as in Example 6 except that A1 of the following structural formula was used instead of R1 as the electron transport auxiliary layer material.



EVALUATION EXAMPLE 2

[0336] For each of the organic electroluminescent devices manufactured in Examples 7 to 28 and Comparative Examples 2 and 3, driving voltage, light emission wave-

length and current efficiency at current density of 10 mA/cm² were measured, and the results are shown in the following Table 2.

TABLE 2

Sample	Electron Transport Auxiliary Layer	Driving Voltage (V)	Light Emission Peak (nm)	Current Efficiency (cd/A)
Example 7	R1	4.4	458	8.3
Example 8	R8	3.9	458	7.5
Example 9	R21	4.4	458	8.0
Example 10	R41	4.4	458	8.2
Example 11	R61	4.4	458	8.1
Example 12	R68	4.4	458	8.2
Example 13	R81	4.4	458	8.3
Example 14	R101	4.4	458	8.0
Example 15	R121	4.4	458	8.2
Example 16	R141	4.3	458	7.8
Example 17	R148	3.5	458	7.1
Example 18	R161	4.4	458	7.3
Example 19	R181	4.5	458	7.4
Example 20	R201	4.1	458	7.5
Example 21	R208	3.9	458	7.0
Example 22	R221	4.2	458	7.5
Example 23	R241	4.1	458	7.6
Example 24	R261	4.5	458	7.3
Example 25	R281	3.8	458	6.9
Example 26	R285	3.4	458	6.8
Example 27	R286	4.7	458	7.2
Example 28	R291	3.8	458	6.9
Comparative Example 2	Alq ₃	4.8	458	6.2
Comparative Example 3	Al	4.7	457	6.5

[0337] As shown in Table 2, it was seen that the blue organic electroluminescent devices using the compound of the present invention in an electron transport auxiliary layer (Examples 6 to 28) exhibited superior performance in terms of current efficiency, light emission peak and driving voltage compared to the blue organic electroluminescent devices without an electron transport auxiliary layer (Comparative Examples 2 and 3).

EXAMPLES 29 TO 32

Manufacture of Blue Organic Electroluminescent Device

[0338] After high purity sublimation purifying the compounds synthesized in the synthesis examples using commonly known methods, blue organic electroluminescent devices were manufactured using the following procedure.

[0339] First, a glass substrate on which indium tin oxide (ITO) was coated as a thin film to a thickness of 1500 Å was ultrasonic cleaned using distilled water. After the cleaning with distilled water was finished, the substrate was ultrasonic cleaned with solvents of isopropyl alcohol, acetone, methanol and the like, dried, then transferred to a UV OZONE washer (Power sonic 405, manufactured by Hwashin Tech. Co., Ltd.), and then, after cleaning the substrate for 5 minutes using UV, the substrate was transferred to a vacuum deposition apparatus.

[0340] On the transparent ITO electrode prepared as above, DS-205 (Doosan Corporation Electro-Materials, 80 nm)/NPB (15 nm)/ADN+5% DS-405 (Doosan Corporation Electro-Materials, 30 nm)/each compound of R1, R8, R21 and R41 (30 nm)/LiF (1 nm)/Al (200 nm) were laminated in this order to manufacture an organic electroluminescent device.

COMPARATIVE EXAMPLE 4

Manufacture of Blue Organic Electroluminescent Device

[0341] A blue organic electroluminescent device was manufactured in the same manner as in Example 29 except that Alq₃ was used instead of R1 as the electron transport layer material.

COMPARATIVE EXAMPLE 5

Manufacture of Blue Organic Electroluminescent Device

[0342] A blue organic electroluminescent device was manufactured in the same manner as in Example 29 except that R1 was not used as the electron transport layer material.

EVALUATION EXAMPLE 3

[0343] For each of the blue organic electroluminescent devices manufactured in Examples 29 to 32 and Comparative Examples 2 and 3, driving voltage, current efficiency and light emission wavelength at current density of 10 mA/cm² were measured, and the results are shown in the following Table 3.

TABLE 3

Sample	Electron Transport Layer	Driving Voltage (V)	Light Emission Peak (nm)	Current Efficiency (cd/A)
Example 29	R1	4.5	455	8.7
Example 30	R8	3.9	455	7.8
Example 31	R21	4.4	455	8.4
Example 32	R41	4.2	455	9.3
Comparative Example 4	Alq ₃	4.7	458	5.5
Comparative Example 5	—	4.8	460	6.2

[0344] As shown in Table 3, it was seen that the blue organic electroluminescent devices using the compound of the present invention in an electron transport layer (Examples 29 to 32) exhibited superior performance in terms of driving voltage, light emission peak and current efficiency compared to the blue organic electroluminescent device using existing Alq₃ in an electron transport layer (Comparative Example 4), and the blue organic electroluminescent device without an electron transport layer (Comparative Example 5).


REFERENCE NUMERAL

- [0345] 10: Anode
- [0346] 20: Cathode
- [0347] 30: Organic Layer
- [0348] 31: Hole Transport Layer
- [0349] 32: Light Emitting Layer
- [0350] 33: Hole Transport Auxiliary Layer
- [0351] 34: Electron Transport Layer
- [0352] 35: Electron Transport Auxiliary Layer

[0353] 36: Electron Injection Layer

[0354] 37: Hole Injection Layer

1. A compound represented by the following Chemical Formula 1:




[Chemical Formula 1]

wherein, in Chemical Formula 1,

X_1 is selected from the group consisting of S, O, $N(R_1)$ and $C(R_2)(R_3)$;

X_2 and X_3 are each independently N or $C(Ar_1)$, and at least one thereof is $C(Ar_1)$; and


ring A is represented by any one of the following Chemical Formulae 2 to 4;



[Chemical Formula 2]



[Chemical Formula 3]



[Chemical Formula 4]

in Chemical Formulae 1 to 4,

a dotted line means a part that is fused;

m is an integer of 0 to 4;


n is an integer of 0 to 6;

R_1 to R_4 are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group;

the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkoxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_1 to R_4 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro

group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other; and

Ar_1 is a substituent represented by the following Chemical Formula 5, and when Ar_1 is present in plural numbers, these are the same as or different from each other;



[Chemical Formula 5]

in Chemical Formula 5,


* means a part where a bond is formed;

L_1 and L_2 are each independently selected from the group consisting of a direct bond, a $C_6\sim C_{18}$ arylene group and a heteroarylene group having 5 to 18 nuclear atoms;


Ar_2 is selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group; and

the arylene group and the heteroarylene group of L_1 and L_2 , and the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkoxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of Ar_2 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.


2. The compound of claim 1, which is represented by any one of the following Chemical Formulae 6 to 14:




[Chemical Formula 6]




[Chemical Formula 7]




[Chemical Formula 8]




[Chemical Formula 9]




[Chemical Formula 10]




[Chemical Formula 11]



[Chemical Formula 12]



[Chemical Formula 13]



[Chemical Formula 14]

wherein, in Chemical Formulae 6 to 14,


ring A, Ar₁ and R₁ to R₃ each have the same definitions as in claim 1.

3. The compound of claim 1, wherein R₁ to R₄ are each independently selected from the group consisting of a


C₁~C₄₀ alkyl group, a C₆~C₆₀ aryl group and a heteroaryl group having 5 to 60 nuclear atoms; and

the alkyl group, the aryl group and the heteroaryl group of R₁ to R₄ are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a C₁~C₄₀ alkyl group, a C₆~C₆₀ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these are the same as or different from each other.


4. The compound of claim 1, wherein L₁ and L₂ are each independently a direct bond, or a linker represented by any one of the following Chemical Formulae A-1 to A-7:




A-1




A-2




A-3




A-4



A-5



A-6



A-7

in Chemical Formulae A-1 to A-7,

* means a part where a bond is formed;

Z₁ to Z₈ are each independently N or C(R₅);

any two of Z₁ to Z₆ forming a bond as a linker in Chemical Formula A-1 are C(R₅), and herein, R₅ is not present;

any one of Z₁ to Z₄ and any one of Z₅ to Z₈ forming a bond as a linker in Chemical Formula A-4 are C(R₅), and herein, R₅ is not present;

any two of Z_1 to Z_4 forming a bond as a linker in Chemical Formula A-5 are $C(R_5)$, and herein, R_5 is not present;

X_4 and X_5 are each independently O, S, $N(R_6)$ or $C(R_7)(R_8)$;


X_6 is N or $C(R_9)$;

R_5 to R_9 are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\text{--}C_{40}$ alkyl group, a $C_2\text{--}C_{40}$ alkenyl group, a $C_2\text{--}C_{40}$ alkynyl group, a $C_3\text{--}C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\text{--}C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\text{--}C_{40}$ alkyloxy group, a $C_6\text{--}C_{60}$ aryloxy group, a $C_3\text{--}C_{40}$ alkylsilyl group, a $C_6\text{--}C_{60}$ arylsilyl group, a $C_1\text{--}C_{40}$ alkylboron group, a $C_6\text{--}C_{60}$ arylboron group, a $C_6\text{--}C_{60}$ arylphosphanyl group, a $C_6\text{--}C_{60}$ mono or diarylphosphinyl group and a $C_6\text{--}C_{60}$ arylamine group, or adjacent groups bond to form a fused ring, and when R_5 to R_8 are each present in plural numbers, these are the same as or different from each other; and


the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_5 to R_9 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a $C_1\text{--}C_{40}$ alkyl group, a $C_2\text{--}C_{40}$ alkenyl group, a $C_2\text{--}C_{40}$ alkynyl group, a $C_6\text{--}C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\text{--}C_{60}$ aryloxy group, a $C_1\text{--}C_{40}$ alkyloxy group, a $C_6\text{--}C_{60}$ arylamine group, a $C_3\text{--}C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\text{--}C_{40}$ alkylsilyl group, a $C_1\text{--}C_{40}$ alkylboron group, a $C_6\text{--}C_{60}$ arylboron group, a $C_6\text{--}C_{60}$ arylphosphanyl group, a $C_6\text{--}C_{60}$ mono or diarylphosphinyl group and a $C_6\text{--}C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.

5. The compound of claim 1, wherein L_1 and L_2 are each independently a direct bond, or a linker represented by any one of the following Chemical Formulae B-1 to B-12:


-continued




B-3




B-4




B-5




B-6




B-7




B-8




B-9



B-10




B-1




B-2

-continued

B-11



B-12



in Chemical Formulae B-1 to B-12,

* means a part where a bond is formed;

R_6 to R_9 are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group; and


the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkoxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_6 to R_9 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.

6. The compound of claim 1, wherein Ar_2 is a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, or an arylamine group; and

the aryl group, the heteroaryl group and the arylamine group of Ar_2 are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ arylamine group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these are the same as or different from each other.

7. The compound of claim 1, wherein Ar_2 is a substituent represented by the following Chemical Formula 15:

[Chemical Formula 15]



in Chemical Formula 15,

* means a part where a bond is formed;


Y_1 to Y_5 are each independently N or C(R_{10});

R_{10} is selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group, and when R_{10} is present in plural numbers, these are the same as or different from each other; and

the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkoxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_{10} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkoxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.

8. The compound of claim 7, wherein the substituent represented by Chemical Formula 15 is a substituent represented by the following Chemical Formula 16:

[Chemical Formula 16]



in Chemical Formula 16,

* means a part where a bond is formed;


R_{11} and R_{12} are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a

C₂-C₄₀ alkenyl group, a C₂-C₄₀ alkynyl group, a C₃-C₄₀ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C₆-C₆₀ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a C₁-C₄₀ alkyloxy group, a C₆-C₆₀ aryloxy group, a C₃-C₄₀ alkylsilyl group, a C₆-C₆₀ arylsilyl group, a C₁-C₄₀ alkylboron group, a C₆-C₆₀ arylboron group, a C₆-C₆₀ arylphosphanyl group, a C₆-C₆₀ mono or diarylphosphinyl group and a C₆-C₆₀ arylamine group, or bond to adjacent groups to form a fused ring;


the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R₁₁ and R₁₂ are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a C₁-C₄₀ alkyl group, a C₂-C₄₀ alkenyl group, a C₂-C₄₀ alkynyl group, a C₃-C₄₀ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C₆-C₆₀ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a C₆-C₆₀ aryloxy group, a C₁-C₄₀ alkyloxy group, a C₆-C₆₀ arylamine group, a C₃-C₄₀ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C₁-C₄₀ alkylsilyl group, a C₁-C₄₀ alkylboron group, a C₆-C₆₀ arylphosphanyl group, a C₆-C₆₀ mono or diarylphosphinyl group and a C₆-C₆₀ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other; and

Y₁, Y₃ and Y₅ each have the same definitions as in claim 7.


9. The compound of claim 7, wherein the substituent represented by Chemical Formula 15 is represented by any one of the following Chemical Formulae C-1 to C-5:




C-1




C-2



C-3



C-4



C-5

in Chemical Formulae C-1 to C-5,

* means a part where a bond is formed;


R₁₁ and R₁₂ are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a C₁-C₄₀ alkyl group, a C₂-C₄₀ alkenyl group, a C₂-C₄₀ alkynyl group, a C₃-C₄₀ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C₆-C₆₀ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a C₁-C₄₀ alkyloxy group, a C₆-C₆₀ aryloxy group, a C₃-C₄₀ alkylsilyl group, a C₆-C₆₀ arylsilyl group, a C₁-C₄₀ alkylboron group, a C₆-C₆₀ arylboron group, a C₆-C₆₀ arylphosphanyl group, a C₆-C₆₀ mono or diarylphosphinyl group and a C₆-C₆₀ arylamine group, or bond to adjacent groups to form a fused ring; and

the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R₁₁ and R₁₂ are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro group, a C₁-C₄₀ alkyl group, a C₂-C₄₀ alkenyl group, a C₂-C₄₀ alkynyl group, a C₃-C₄₀ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C₆-C₆₀ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a C₆-C₆₀ aryloxy group, a C₁-C₄₀ alkyloxy group, a C₆-C₆₀ arylamine group, a C₃-C₄₀ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a C₁-C₄₀ alkylsilyl group, a C₁-C₄₀ alkylboron group, a C₆-C₆₀ arylphosphanyl group, a C₆-C₆₀ mono or diarylphosphinyl group and a C₆-C₆₀ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.

10. The compound of claim 9, wherein R₁₁ and R₁₂ are each independently selected from the group consisting of a C₁-C₄₀ alkyl group, a C₆-C₆₀ aryl group and a heteroaryl group having 5 to 60 nuclear atoms; and

the alkyl group, the aryl group and the heteroaryl group of R₁₁ and R₁₂ are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a C₁-C₄₀ alkyl group, a C₆-C₆₀ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these are the same as or different from each other.

11. The compound of claim 1, wherein Ar₂ is a substituent represented by the following Chemical Formula 17:



[Chemical Formula 17]

in Chemical Formula 17,

* means a part where a bond is formed;

R_{13} and R_{14} are each independently selected from the group consisting of hydrogen, deuterium, halogen, a cyano group, a nitro group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_1\sim C_{40}$ alkyloxy group, a $C_6\sim C_{60}$ aryloxy group, a $C_3\sim C_{40}$ alkylsilyl group, a $C_6\sim C_{60}$ arylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylamine group, or bond to adjacent groups to form a fused ring; and


the alkyl group, the alkenyl group, the alkynyl group, the aryl group, the heteroaryl group, the aryloxy group, the alkyloxy group, the cycloalkyl group, the heterocycloalkyl group, the arylamine group, the alkylsilyl group, the alkylboron group, the arylboron group, the arylphosphanyl group, the mono or diarylphosphinyl group and the arylsilyl group of R_{13} and R_{14} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of deuterium, halogen, a cyano group, a nitro

group, a $C_1\sim C_{40}$ alkyl group, a $C_2\sim C_{40}$ alkenyl group, a $C_2\sim C_{40}$ alkynyl group, a $C_6\sim C_{60}$ aryl group, a heteroaryl group having 5 to 60 nuclear atoms, a $C_6\sim C_{60}$ aryloxy group, a $C_1\sim C_{40}$ alkyloxy group, a $C_6\sim C_{60}$ arylamine group, a $C_3\sim C_{40}$ cycloalkyl group, a heterocycloalkyl group having 3 to 40 nuclear atoms, a $C_1\sim C_{40}$ alkylsilyl group, a $C_1\sim C_{40}$ alkylboron group, a $C_6\sim C_{60}$ arylboron group, a $C_6\sim C_{60}$ arylphosphanyl group, a $C_6\sim C_{60}$ mono or diarylphosphinyl group and a $C_6\sim C_{60}$ arylsilyl group, and when substituted with a plurality of the substituents, these are the same as or different from each other.


12. The compound of claim 11, wherein R_{13} and R_{14} are each independently selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms; and

the alkyl group, the aryl group and the heteroaryl group of R_{13} and R_{14} are each independently unsubstituted or substituted with one or more types of substituents selected from the group consisting of a $C_1\sim C_{40}$ alkyl group, a $C_6\sim C_{60}$ aryl group and a heteroaryl group having 5 to 60 nuclear atoms, and when substituted with a plurality of the substituents, these are the same as or different from each other.


13. The compound of claim 1, which is selected from the group consisting of the following compounds:




R1



R2




R3




R4

-continued


R5




R6



R7




R8




-continued


R9




R10




R11



R12




R13




-continued


R14




R15



R16




R17




-continued


R18



R19




R20




-continued


R21




R22




R23




R24




R25




-continued




R26




R27



R28




R29




R30


-continued




R31



R32




R33




R34


-continued




R35



R36




R37




R38

-continued


R39




R40



R41




R42




-continued


R43




R44



R45




R46




-continued


R47




R48




R49




R50



R51




R52




-continued


R53



R54




R55




-continued


R56




R57



R58




R59




-continued


R60




R61




R62



R63




R64




-continued


R65




R66




R67




R68




-continued




R69




R70



R71




R72




R73

-continued

R74





R75




R76


R77




-continued



R78




R79




R80

-continued


R81




R82




R83




R84




R85




-continued




R86




R87



R88




R89




R90


-continued




R91



R92




R93




R94


-continued




R95



R96




R97




R98

-continued

R99





R100




R101

R102




-continued


R103




R104



R105




R106




-continued


R107




R108




R109




R110



R111




R112




-continued


R113



R114




R115




-continued


R116




R117



R118




R119




-continued


R120




R121




R122



R123




R124




-continued


R125




R126



R127




R128




-continued


R129




R130




R131



R132




R133




-continued


R134



R135




R136




-continued


R137



R138




R139





-continued

R140





R141

R142




R143

R144




-continued


R145




R146



R147




R148




-continued


R149




R150




R151



R152




R153




-continued


R154




R155




R156




R157




-continued



R158








R159




R160

-continued




-continued


R166




R167




R168




R169




R170




-continued




R171



R172




R173




R174


-continued




R175



R176




R177




R178

-continued


R179




R180



R221




R222




-continued


R223




R224



R225




R226




-continued


R227




R228




R229




R230



R231




R232




-continued


R233



R234




R235




-continued


R236




R237



R238




R239



-continued

R240



14. An organic electroluminescent device comprising:
an anode;
a cathode; and
one or more organic material layers provided between the
anode and the cathode,
wherein at least one of the one or more organic material
layers includes the compound represented by Chemical
Formula 1 of claim 1.

15. The organic electroluminescent device of claim 14,
wherein the organic material layer includes one or more
layers selected from the group consisting of a hole injection
layer, a hole transport layer, a hole transport auxiliary layer,
an electron transport layer, an electron transport auxiliary
layer and a light emitting layer.

* * * * *

专利名称(译)	有机化合物和包括该化合物的有机电致发光器件		
公开(公告)号	US20200227650A1	公开(公告)日	2020-07-16
申请号	US16/626634	申请日	2018-05-02
[标]申请(专利权)人(译)	株式会社斗山		
申请(专利权)人(译)	斗山公司		
当前申请(专利权)人(译)	斗山公司		
[标]发明人	KIM HONG SUK KIM YOUNG BAE		
发明人	KIM, HONG SUK KIM, YOUNG BAE		
IPC分类号	H01L51/00 C07D403/10 C09K11/06 C07D405/14 C07D401/14 C07D405/10 C07D251/24 C07D409/10		
CPC分类号	C09K2211/1018 H01L51/0074 H01L51/5064 C07D405/14 H01L51/5012 C07D403/10 C07D401/14 H01L51/0073 H01L51/0067 C07D251/24 C07D405/10 C09K11/06 C07D409/10 H01L51/5088 H01L51/5016 H01L51/0072 H01L51/508 C07D231/56 C07D409/14 H01L51/00 H01L51/50		
优先权	1020170083287 2017-06-30 KR		
外部链接	Espacenet USPTO		

摘要(译)

本发明涉及一种新颖的化合物和包括该化合物的有机电致发光器件,并且通过在有机电致发光器件的有机材料层,优选发光层中使用根据本发明的化合物,发光效率,驱动电压,可以提高有机电致发光器件的寿命等。

