



US 20190296247A1

(19) **United States**(12) **Patent Application Publication**
SAKAMOTO(10) **Pub. No.: US 2019/0296247 A1**(43) **Pub. Date: Sep. 26, 2019**(54) **ORGANIC ELECTROLUMINESCENCE
DEVICE AND POLYCYCLIC COMPOUND
FOR ORGANIC ELECTROLUMINESCENCE
DEVICE**(71) Applicant: **Samsung Display Co., Ltd.**, Yongin-si
(KR)(72) Inventor: **Naoya SAKAMOTO**, Yokohama (JP)(21) Appl. No.: **16/270,108**(22) Filed: **Feb. 7, 2019**(30) **Foreign Application Priority Data**

Mar. 22, 2018 (KR) 10-2018-0033427

Publication Classification(51) **Int. Cl.****H01L 51/00** (2006.01)**C07D 401/14** (2006.01)**C09K 11/06** (2006.01)(52) **U.S. Cl.**CPC **H01L 51/0072** (2013.01); **C07D 401/14**(2013.01); **H01L 51/5012** (2013.01); **C09K****11/06** (2013.01); **H01L 51/0067** (2013.01)

(57)

ABSTRACT

An organic electroluminescence device includes a first electrode, a hole transport region disposed on the first electrode, an emission layer disposed on the hole transport region, an electron transport region disposed on the emission layer, and a second electrode disposed on the electron transport region, wherein the emission layer includes a polycyclic compound containing two electron donors and one electron acceptor, and the electron acceptor includes a benzonitrile part and a pyridine part, thereby showing high emission efficiency.

10




FIG. 1




FIG. 2





FIG. 3



ORGANIC ELECTROLUMINESCENCE DEVICE AND POLYCYCLIC COMPOUND FOR ORGANIC ELECTROLUMINESCENCE DEVICE

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application claims priority to and the benefit of Korean Patent Application No. 10-2018-0033427, filed on Mar. 22, 2018, the entire content of which is hereby incorporated by reference.

BACKGROUND

1. Field

[0002] The present disclosure herein relates to an organic electroluminescence device and a polycyclic compound utilized in the organic electroluminescence device.

2. Description of the Related Art

[0003] Recently, the development of an organic electroluminescence display device as an image display device is being actively conducted. Different from a liquid crystal display device, the organic electroluminescence display device is a so-called self-luminescent display device in which holes and electrons injected from a first electrode and a second electrode recombine in an emission layer, and a light emission material including an organic compound in the emission layer emits light to attain display (e.g., an image).

[0004] In the application of an organic electroluminescence device to a display device, the decrease of the driving voltage, and the increase of the light-emitting efficiency and the life (e.g., lifespan) of the organic electroluminescence device are required, and materials for an organic electroluminescence device that can stably attain (e.g., satisfy) the requirements are being continuously developed.

[0005] Particularly, recently, in order to accomplish an organic electroluminescence device with high efficiency, techniques on phosphorescence emission (which utilizes energy in a triplet state) or delayed fluorescence emission (which utilizes the generating phenomenon of singlet excitons by the collision of triplet excitons (triplet-triplet annihilation, TTA)) are being developed, and development on a material for thermally activated delayed fluorescence (TADF) utilizing delayed fluorescence phenomenon is being conducted.

SUMMARY

[0006] Aspects according to one or more embodiments of the present disclosure are directed toward an organic electroluminescence device having long lifespan and high efficiency, and a polycyclic compound utilized therein.

[0007] Aspects according to one or more embodiments of the present disclosure are directed toward an organic electroluminescence device including a material for emitting thermally activated delayed fluorescence, and a polycyclic compound utilized as a material for emitting thermally activated delayed fluorescence.

[0008] According to an embodiment of the inventive concept, an organic electroluminescence device includes a first electrode, a hole transport region on the first electrode, an emission layer on the hole transport region, an electron transport region on the emission layer, and a second electrode on the electron transport region, wherein the emission layer includes a polycyclic compound containing two electron donors (e.g., two electron donor groups) and one


electron acceptor (e.g., one electron acceptor group), and the electron acceptor includes a benzonitrile part (e.g., a benzonitrile group) and a pyridine part (e.g., a pyridine group). [0009] In an embodiment, the emission layer may be configured to emit delayed fluorescence.

[0010] In an embodiment, the emission layer may include a host and a dopant, and the dopant may be the polycyclic compound.

[0011] In an embodiment, the emission layer may be a thermally activated delayed fluorescence emission layer configured to emit blue light.

[0012] In an embodiment, the polycyclic compound may be represented by the following Formula 1:


Formula 1



[0013] In Formula 1, X_1 and X_2 are each independently a direct linkage, CR_7R_8 , SiR_9R_{10} , O, or S; L is a direct linkage, CO, SO_2 , $SiR_{11}R_{12}$, a substituted or unsubstituted arylene group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroarylene group having 2 to 30 carbon atoms for forming a ring; R_1 to R_4 are each independently a hydrogen atom, a deuterium atom, a halogen atom, $NR_{13}R_{14}$, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring, R_1 to R_4 may be each optionally independently combined with an adjacent group to form a ring; a, b, c and d are each independently an integer of 0 to 4; R_5 to R_{14} are each independently a hydrogen atom, a deuterium atom, a halogen atom, a cyano group, a substituted or unsubstituted amino group, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring, R_5 to R_{14} may be each optionally independently combined with an adjacent group to form a ring; and e and f are each independently an integer of 0 to 3.


[0014] In an embodiment, Formula 1 may be represented by one of the following Formula 1-1 to Formula 1-3:

Formula 1-1




-continued

Formula 1-2



Formula 1-3




[0015] In Formula 1-1 to Formula 1-3, X_1 and X_2 , L , R_1 to R_6 , and a to f are the same as respectively defined in association with Formula 1.


[0016] In an embodiment, X_1 and X_2 of Formula 1 may be the same.

[0017] In an embodiment, R_5 and R_6 of Formula 1 may be each independently represented by one of the following Formula 2-1 to Formula 2-4:


Formula 2-1



Formula 2-2




Formula 2-3




-continued

Formula 2-4




[0018] In an embodiment, R_1 to R_4 of Formula 1 may be each independently represented by one of the following Formula 3-1 to Formula 3-3:


Formula 3-1



Formula 3-2



Formula 3-3



[0019] In an embodiment of the inventive concept, an organic electroluminescence device includes a first electrode, a hole transport region on the first electrode, an emission layer on the hole transport region, an electron transport region on the emission layer, and a second electrode on the electron transport region, wherein the emission layer includes the polycyclic compound represented by Formula 1 above.

[0020] In an embodiment of the inventive concept, a polycyclic compound is represented by Formula 1 above.

BRIEF DESCRIPTION OF THE FIGURES

[0021] The accompanying drawings are included to provide a further understanding of the inventive concept, and are incorporated in and constitute a part of this specification. The drawings illustrate exemplary embodiments of the inventive concept and, together with the description, serve to explain principles of the inventive concept. In the drawings:

[0022] FIG. 1 is a cross-sectional view schematically illustrating an organic electroluminescence device according to an embodiment of the inventive concept;

[0023] FIG. 2 is a cross-sectional view schematically illustrating an organic electroluminescence device according to an embodiment of the inventive concept; and

[0024] FIG. 3 is a cross-sectional view schematically illustrating an organic electroluminescence device according to an embodiment of the inventive concept.

DETAILED DESCRIPTION

[0025] The inventive concept may have various modifications and may be embodied in different forms, and example embodiments will be explained in more detail with reference to the accompany drawings. The inventive concept may, however, be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, all modifications, equivalents, and substituents which are included in the spirit and technical scope of the inventive concept should be included in the inventive concept.

[0026] Like reference numerals refer to like elements throughout. In the drawings, the dimensions of structures may be exaggerated for clarity of illustration. It will be understood that, although the terms “first,” “second,” etc., may be used herein to describe various elements, these elements should not be limited by these terms. These terms are only used to distinguish one element from another element. Thus, a first element could be termed a second element without departing from the teachings of the present invention. Similarly, a second element could be termed a first element. As used herein, the singular forms are intended to include the plural forms as well, unless the context clearly indicates otherwise.

[0027] It will be further understood that the terms “comprises” and/or “comprising,” when used in this specification, specify the presence of stated features, numerals, steps, operations, elements, parts, or the combination thereof, but do not preclude the presence or addition of one or more other features, numerals, steps, operations, elements, parts, or the combination thereof. It will also be understood that when a layer, a film, a region, a plate, etc., is referred to as being “on” another part, it can be “directly on” the other part, or intervening layers may also be present.

[0028] In the description, “-” represents a connecting position.

[0029] In the description, the term “unsubstituted” corresponds to an unsubstituted functional group and the term “substituted” corresponds to a functional group substituted with at least one substituent selected from the group consisting of a deuterium atom, a halogen atom, a cyano group, a nitro group, an amino group, a silyl group, a boron group, a phosphine oxide group, a phosphine sulfide group, an alkyl group, an alkenyl group, an aryl group, and a heterocyclic group. In addition, each of the substituents may be substituted or unsubstituted. For example, a biphenyl group may be interpreted as an aryl group or a phenyl group substituted with a phenyl group.

[0030] In the description, the terms “forming a ring via the combination with an adjacent group” may refer to forming a substituted or unsubstituted hydrocarbon ring, or a substituted or unsubstituted heterocycle (e.g., heterocyclic ring) via the combination with an adjacent group. The hydrocarbon ring may include an aliphatic hydrocarbon ring and/or an aromatic hydrocarbon ring. The heterocycle may include an aliphatic heterocycle and/or an aromatic heterocycle. The hydrocarbon ring and the heterocycle may be monocyclic or polycyclic. In addition, the ring formed via the combination with an adjacent group may be combined with another ring to form a spiro structure.

[0031] In the description, the terms “an adjacent group” may refer to a substituent substituted for an atom which is directly combined with an atom substituted with a corresponding substituent, another substituent substituted for an atom which is substituted with a corresponding substituent, or a substituent sterically positioned at the nearest position to a corresponding substituent. For example, in 1,2-dimethylbenzene, two methyl groups may be interpreted as “adjacent groups” to each other, and in 1,1-diethylcyclopentene, two ethyl groups may be interpreted as “adjacent groups” to each other.

[0032] In the description, the halogen atom may be a fluorine atom, a chlorine atom, a bromine atom and/or an iodine atom.

[0033] In the description, the alkyl (e.g., the alkyl group) may be a linear, branched or cyclic group. The carbon number of the alkyl may be 1 to 50, 1 to 30, 1 to 20, 1 to 10, or 1 to 6. Examples of the alkyl may include methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl, t-butyl, i-butyl, 2-ethylbutyl, 3,3-dimethylbutyl, n-pentyl, i-pentyl, neopentyl, t-pentyl, cyclopentyl, 1-methylpentyl, 3-methylpentyl, 2-ethylpentyl, 4-methyl-2-pentyl, n-hexyl, 1-methylhexyl, 2-ethylhexyl, 2-butylhexyl, cyclohexyl, 4-methylcyclohexyl, 4-t-butylcyclohexyl, n-heptyl, 1-methylheptyl, 2,2-dimethylheptyl, 2-ethylheptyl, 2-butylheptyl, n-octyl, t-octyl, 2-ethyloctyl, 2-butyloctyl, 2-hexyloctyl, 3,7-dimethyloctyl, cyclooctyl, n-nonyl, n-decyl, adamantyl, 2-ethyldecyl, 2-butyldecyl, 2-hexyldecyl, 2-octyldecyl, n-undecyl, n-dodecyl, 2-ethyldodecyl, 2-butyldecyl, 2-hexyldodecyl, 2-octyldodecyl, n-tridecyl, n-tetradecyl, n-pentadecyl, n-hexadecyl, 2-ethylhexadecyl, 2-butylhexadecyl, 2-hexylhexadecyl, 2-octylhexadecyl, n-heptadecyl, n-octadecyl, n-nonadecyl, n-eicosyl, 2-ethyleicosyl, 2-butyleicosyl, 2-hexyleicosyl, 2-octyleicosyl, n-henicosyl, n-docosyl, n-tricosyl, n-tetracosyl, n-pentacosyl, n-hexacosyl, n-heptacosyl, n-octacosyl, n-nonacosyl, n-triacontyl, etc., without being limited thereto.

[0034] In the description, the term “aryl group” refers to an optional functional group or substituent derived from an aromatic hydrocarbon ring. The aryl group may be a monocyclic aryl group or a polycyclic aryl group. The carbon number for forming a ring in the aryl group may be, 6 to 30, 6 to 20, or 6 to 15. Examples of the aryl group may include phenyl, naphthyl, fluorenyl, anthracenyl, phenanthryl, biphenyl, terphenyl, quaterphenyl, quinphenyl, sexiphenyl, triphenylenyl, pyrenyl, benzo[fluoranthene], chrysenyl, etc., without being limited thereto.

[0035] In the description, the heteroaryl (e.g., the heteroaryl group) may be a heteroaryl including at least one of O, N, P, Si and S as a heteroatom. The carbon number for forming a ring of the heteroaryl may be 2 to 30, or 2 to 20. The heteroaryl may be monocyclic heteroaryl or polycyclic heteroaryl. Examples of the polycyclic heteroaryl may have a dicyclic or tricyclic structure. Examples of the heteroaryl may include thiophene, furan, pyrrole, imidazole, thiazole, oxazole, oxadiazole, triazole, pyridyl, bipyridyl, pyrimidyl, triazine, triazole, acridyl, pyridazine, pyrazinyl, quinolinyl, quinazoline, quinoxalyl, phenoxazyl, phthalaziny, pyrido pyrimidinyl, pyrido pyrazinyl, pyrazino pyrazinyl, isoquinoline, indole, carbazole, N-arylcarbazole, N-heteroarylcarbazole, N-alkylcarbazole, benzoxazole, benzimidazole, benzothiazole, benzocarbazole, benzothiophene, dibenzothiophene, thienothiophene, benzofuranyl, phenanthroline, thiazolyl, isooxazolyl, oxadiazolyl, thiadiazolyl, benzothi-

azolyl, phenothiazinyl, dibenzosilole, dibenzofuran, etc., without being limited thereto.

[0036] In the description, the silyl group may include an alkyl silyl group and/or an aryl silyl group. Examples of the silyl group may include trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, vinyl dimethylsilyl, propyl dimethylsilyl, triphenylsilyl, diphenylsilyl, phenylsilyl, etc. However, an embodiment of the inventive concept is not limited thereto.

[0037] In the description, the carbon number of the amine group is not specifically limited, but may be 1 to 30. The amine group may include an alkyl amine group and/or an aryl amine group. Examples of the amine group include a methylamine group, a dimethylamine group, a phenylamine group, a naphthylamine group, a 9-methyl-anthracenylamine group, a triphenylamine group, etc., without being limited thereto.

[0038] In the description, the explanation on the aryl group may be applied to the arylene group except that the arylene group is a divalent group.

[0039] In the description, the explanation on the heteroaryl group may be applied to the heteroarylene group except that the heteroarylene group is a divalent group.

[0040] Hereinafter, the organic electroluminescence device according to an embodiment of the inventive concept will be explained with reference to FIGS. 1 to 3.

[0041] Referring to FIGS. 1 to 3, an organic electroluminescence device 10 according to an embodiment may include a first electrode EL1, a hole transport region HTR, an emission layer EML, an electron transport region ETR and a second electrode EL2, laminated one by one (e.g., in the stated order).

[0042] The first electrode EL1 and the second electrode EL2 are oppositely disposed from each other, and a plurality of organic layers may be disposed between the first electrode EL1 and the second electrode EL2. The plurality of the organic layers may include a hole transport region HTR, an emission layer EML, and an electron transport region ETR. The organic electroluminescence device 10 of an embodiment may include the polycyclic compound of an embodiment in the emission layer EML.

[0043] When compared with FIG. 1, FIG. 2 shows the cross-sectional view of an organic electroluminescence device 10 of an embodiment, wherein a hole transport region HTR includes a hole injection layer HIL and a hole transport layer HTL, and an electron transport region ETR includes an electron injection layer EIL and an electron transport layer ETL. In addition, when compared with FIG. 1, FIG. 3 shows the cross-sectional view of an organic electroluminescence device 10 of an embodiment, wherein a hole transport region HTR includes a hole injection layer HIL, a hole transport layer HTL, and an electron blocking layer EBL, and an electron transport region ETR includes an electron injection layer EIL, an electron transport layer ETL, and a hole blocking layer HBL.

[0044] In the organic electroluminescence devices 10 of exemplary embodiments shown in FIGS. 1 to 3, the polycyclic compound of an embodiment, including two electron donors (e.g., two electron donor groups) and one electron acceptor (e.g., one electron acceptor group) may be included in at least one organic layer among the plurality of organic layers. In the polycyclic compound of an embodiment, the electron acceptor may include a benzonitrile part (e.g., a benzonitrile group) and a pyridine part (e.g., a pyridine group). For example, the polycyclic compound of an

embodiment may have an electron donor-electron acceptor-electron donor (D-A-D) structure.

[0045] The organic electroluminescence device 10 of an embodiment may include a polycyclic compound including two electron donors and one electron acceptor in an emission layer EML. In the polycyclic compound, the electron acceptor may include a benzonitrile part and a pyridine part.

[0046] In the organic electroluminescence devices 10 of an embodiment, the first electrode EL1 has conductivity. The first electrode EL1 may be formed utilizing a metal alloy or a conductive compound. The first electrode EL1 may be an anode.

[0047] The first electrode EL1 may be a transmissive electrode, a transfective electrode, or a reflective electrode. If the first electrode EL1 is the transmissive electrode, the first electrode EL1 may be formed utilizing a transparent metal oxide such as indium tin oxide (ITO), indium zinc oxide (IZO), zinc oxide (ZnO), and indium tin zinc oxide (ITZO). If the first electrode EL1 is the transfective electrode or the reflective electrode, the first electrode EL1 may include Ag, Mg, Cu, Al, Pt, Pd, Au, Ni, Nd, Ir, Cr, Li, Ca, LiF/Ca, LiF/Al, Mo, Ti, a compound thereof, or a mixture thereof (for example, a mixture of Ag and Mg). Also, the first electrode EL1 may have a structure including a plurality of layers including a reflective layer or a transfective layer formed utilizing the above materials, and a transmissive conductive layer formed utilizing ITO, IZO, ZnO, or ITZO. For example, the first electrode EL1 may include a plurality of layers, such as ITO/Ag/ITO.

[0048] The hole transport region HTR is provided on the first electrode EL1. The hole transport region HTR may include at least one of a hole injection layer HIL, a hole transport layer HTL, a hole buffer layer, or an electron blocking layer EBL.

[0049] The hole transport region HTR may have a single layer formed utilizing a single material, a single layer formed utilizing a plurality of different materials, or have a multilayer structure including a plurality of layers formed utilizing a plurality of different materials.

[0050] For example, the hole transport region HTR may have the structure of a single layer such as a hole injection layer HIL, or a hole transport layer HTL, and may have a structure of a single layer formed utilizing a hole injection material and a hole transport material. Alternatively, the hole transport region HTR may have a structure of a single layer formed utilizing a plurality of different materials, or a multilayer structure laminated (e.g., stacked) from the first electrode EL1 of hole injection layer HIL/hole transport layer HTL, hole injection layer HIL/hole transport layer HTL/hole buffer layer, hole injection layer HIL/hole buffer layer, hole transport layer HTL/hole buffer layer, or hole injection layer HIL/hole transport layer HTL/electron blocking layer, without being limited thereto.

[0051] The hole transport region HTR may be formed utilizing various suitable methods, such as a vacuum deposition method, a spin coating method, a cast method, a Langmuir-Blodgett (LB) method, an inkjet printing method, a laser printing method, and a laser induced thermal imaging (LITI) method.

[0052] The hole injection layer HIL of the organic electroluminescence device 10 of an embodiment may include a suitable (e.g., a known) hole injection material. For example, the hole injection layer HIL may include triphenylamine-containing polyetherketone (TPAPEK), 4-iso-

propyl-4'-methyldiphenyliodoniumtetrakis(pentafluorophenyl)borate (PPBI), N,N'-diphenyl-N,N'-bis-[4-(phenyl-methyl-amino)-phenyl]-phenyl-4,4'-diamine (DNTPD), a phthalocyanine compound such as copper phthalocyanine, 4,4',4''-tris(3-methyl phenyl phenylamino)triphenylamine (m-MTDATA), N,N'-di(1-naphthyl)-N,N'-diphenylbenzidine (NPB), N,N'-bis(1-naphthyl)-N,N'-diphenyl-4,4'-diamine (α -NPD), 4,4',4''-tris(N,N-diphenylamino)triphenylamine (TDATA), 4,4',4''-tris(N,N-2-naphthyl phenylamino)triphenylamine (2-TNATA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), or dipyrzino[2,3-f:2',3'-h]quinoxaline-2,3,6,7,10,11-hexacarbonitrile (HAT-CN). However, an embodiment of the inventive concept is not limited thereto.

[0053] The hole transport layer HTL of the organic electroluminescence device 10 of an embodiment may include a suitable (e.g., a known) hole transport material. For example, the hole transport layer HTL may include 1,1-bis[(di-4-trileamino)phenyl]cyclohexane (TAPC), carbazole derivatives such as N-phenyl carbazole and polyvinyl carbazole, N,N'-bis(3-methylphenyl)-N,N'-diphenyl-[1,1-biphenyl]-4,4'-diamine (TPD), 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), N,N'-di(1-naphthyl)-N,N'-diphenylbenzidine (NPB), N,N'-bis(1-naphthyl)-N,N'-diphenyl-4,4'-diamine (α -NPD), etc. However, an embodiment of the inventive concept is not limited thereto.

[0054] Meanwhile, the hole transport region HTR may further include an electron blocking layer EBL, and the electron blocking layer EBL may be disposed between a hole transport layer HTL and an emission layer EML. The electron blocking layer EBL may play the role of preventing or substantially preventing electron injection from an electron transport region ETR to a hole transport region HTR.

[0055] The electron blocking layer EBL may include a common material (e.g., a known material) in the art. The electron blocking layer EBL may include, for example, carbazole derivatives (such as N-phenylcarbazole, and polyvinyl carbazole), fluorine-based derivatives, N,N'-bis(3-methylphenyl)-N,N'-diphenyl-[1,1-biphenyl]-4,4'-diamine (TPD), triphenylamine-based derivatives such as 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), N,N'-di(naphthalene-1-yl)-N,N'-diphenyl-benzidine (NPD), 4,4'-cyclohexylidene bis[N,N-bis(4-methylphenyl)benzenamine] (TAPC), 4,4'-bis[N,N'-(3-tolyl)amino]-3,3'-dimethylbiphenyl (HMTDP), mCP, etc. In addition, as described above, the electron blocking layer EBL may include the polycyclic compound according to an embodiment of the inventive concept.

[0056] The thickness of the hole transport region HTR may be from about 100 Å to about 10,000 Å, for example, from about 100 Å to about 5,000 Å. The thickness of the hole injection layer HIL may be, for example, from about 30 Å to about 1,000 Å, and the thickness of the hole transport layer HTL may be from about 30 Å to about 1,000 Å. For example, the thickness of the electron blocking layer EBL may be from about 10 Å to about 1,000 Å. If the thicknesses of the hole transport region HTR, the hole injection layer

HIL, the hole transport layer HTL and the electron blocking layer EBL satisfy the above-described ranges, satisfactory hole transport properties may be achieved without substantial increase of a driving voltage.

[0057] The hole transport region HTR may further include a charge generating material in addition to the above-described materials to improve conductivity. The charge generating material may be dispersed uniformly or non-uniformly in the hole transport region HTR. The charge generating material may be, for example, a p-dopant. The p-dopant may be one of quinone derivatives, metal oxides, or cyano group-containing compounds, without being limited thereto. For example, non-limiting examples of the p-dopant may include quinone derivatives (such as tetracyanoquinodimethane (TCNQ) and 2,3,5,6-tetrafluoro-tetracyanoquinodimethane (F4-TCNQ)), and metal oxides (such as tungsten oxide and molybdenum oxide), without being limited thereto.

[0058] As described above, the hole transport region HTR may further include at least one of a hole buffer layer or an electron blocking layer EBL in addition to the hole injection layer HIL and the hole transport layer HTL. The hole buffer layer may compensate a resonance distance according to the wavelength of light emitted from the emission layer EML and increase light emission efficiency. Materials included in the hole transport region HTR may be utilized as materials included in the hole buffer layer.

[0059] The emission layer EML is provided on the hole transport region HTR. The emission layer EML may have a thickness of, for example, about 100 Å to about 600 Å. The emission layer EML may have a single layer formed utilizing a single material, a single layer formed utilizing a plurality of different materials, or have a multilayer structure having a plurality of layers formed utilizing a plurality of different materials.

[0060] The emission layer EML may emit one of red light, green light, blue light, white light, yellow light, and cyan light. The emission layer EML may include a fluorescence emitting material or a phosphorescence emitting material.


[0061] In an embodiment, the emission layer EML includes a polycyclic compound containing two electron donors and one electron acceptor, and the electron acceptor includes a benzonitrile part and a pyridine part.

[0062] In an embodiment, the emission layer EML may be a fluorescence emission layer. For example, a portion of the light emitted from the emission layer EML may be attributed to thermally activated delayed fluorescence. In an embodiment, the emission layer EML may include a light-emitting component that emits thermally activated delayed fluorescence. In an embodiment, the emission layer EML may be an emission layer emitting thermally activated delayed fluorescence that emits blue light. Light-emitting component emitting thermally activated delayed fluorescence may be a material having excellent electron accepting and electron donating properties and smooth charge transfer (CT) in a molecule.

[0063] In an embodiment, the emission layer EML may include a polycyclic compound containing two electron donors and one electron acceptor. In addition, in an embodiment, the emission layer EML may include a host and a dopant, and the dopant may include a polycyclic compound containing two electron donors and one electron acceptor.

[0064] The polycyclic compound may, for example, have a structure represented by the following Formula 1:

Formula 1



[0065] In Formula 1, X_1 and X_2 are each independently a direct linkage, CR_7R_8 , SiR_9R_{10} , O, or S; and L is a direct linkage, CO, SO_2 , $SiR_{11}R_{12}$, a substituted or unsubstituted arylene group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroarylene group having 2 to 30 carbon atoms for forming a ring.

[0066] In Formula 1, R_1 to R_4 are each independently a hydrogen atom, a deuterium atom, a halogen atom, $NR_{13}R_{14}$, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring, or combined with an adjacent group to form a ring; and a, b, c and d are each independently an integer of 0 to 4.


[0067] If each of a to d is an integer of 2 or more, a plurality of R_1 , R_2 , R_3 and R_4 may be respectively the same or different from each other. For example, when a, b, c or d is an integer of 2 or more, a plurality of R_1 , R_2 , R_3 or R_4 may be the same or different from each other.

[0068] In Formula 1, R_5 to R_{14} are each independently a hydrogen atom, a deuterium atom, a halogen atom, a cyano group, a substituted or unsubstituted amino group, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring, or combined with an adjacent group to form a ring; and e and f are each independently an integer of 0 to 3.


[0069] If each of e and f is an integer of 2 or more, a plurality of R_5 and R_6 may be respectively the same or different from each other. For example, when e or f is an integer of 2 or more, a plurality of R_5 or R_6 may be the same or different from each other.

[0070] e and f may each be 0. For example, if e is 0, a pyridine group in the polycyclic compound represented by Formula 1 may be an unsubstituted pyridylene group.

[0071] According to an embodiment, in the polycyclic compound represented by Formula 1,




and




Formula 1-1


-continued



parts may be electron donors, and



corresponding to a benzonitrile part and



corresponding to a pyridine part may be electron acceptors. However, an embodiment of the inventive concept is not limited thereto.


[0072] In Formula 1, if L (which corresponds to a linker) is a direct linkage, the benzonitrile part and the pyridine part of the electron donor are directly bonded, and if L (which corresponds to a linker) is CO, SO_2 , $SiR_{11}R_{12}$, a substituted or unsubstituted arylene group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroarylene group having 2 to 30 carbon atoms for forming a ring, the benzonitrile part and the pyridine part of the electron acceptor are bonded via L.

[0073] The electron donor of Formula 1 does not include a cyano group. If the electron donor includes a cyano group, the absolute value (ΔE_{ST}) of the difference between a singlet energy level (S1) and a triplet energy level (T1), which will be explained later, becomes relatively large, and is not applicable as a material for emitting thermally activated delayed fluorescence.


[0074] In an embodiment, Formula 1 may be represented by one of the following Formula 1-1 to Formula 1-3:

-continued

Formula 1-2



Formula 1-3




[0075] In Formula 1-1 to Formula 1-3, X_1 and X_2 , L , R_1 to R_6 , and a to f are the same as respectively defined in association with Formula 1. Formula 1-1 represents a case where X_1 , X_2 and L are direct linkages, Formula 1-2 represents a case where X_1 and X_2 are direct linkages, and Formula 1-3 represents a case where L is a direct linkage.


[0076] In an embodiment, X_1 and X_2 of Formula 1 may be the same. For example, if X_1 is a direct linkage, X_2 is also a direct linkage, and if X_1 is $C(CH_3)_2$, X_2 is also $C(CH_3)_2$.

[0077] In an embodiment, R_5 and R_6 of Formula 1 may be each independently one of the following Formula 2-1 to Formula 2-4:


Formula 2-1



Formula 2-2




Formula 2-3



-continued

Formula 2-4




[0078] That is, R_5 and R_6 are each independently a substituted or unsubstituted carbazole group, a substituted pyridyl group, or a substituted benzonitrile group. In an embodiment, R_5 and R_6 are each independently an unsubstituted carbazole group, a carbazole group substituted with an alkyl group, a pyridyl group substituted with a carbazole group, or a benzonitrile group substituted with a carbazole group.

[0079] In Formula 1, R_1 to R_4 are each independently a hydrogen atom, a halogen atom, $NR_{13}R_{14}$, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring, or may be combined with an adjacent group to form a ring.


[0080] In R_1 to R_4 of Formula 1, a cyano group is excluded from “substituted or unsubstituted” substituents. Particularly, the substituents in R_1 to R_4 may be each independently a hydrogen atom, a fluorine atom, a methyl group, an unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, or an unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring.

[0081] In an embodiment, R_1 to R_4 of Formula 1 may be each independently represented by one of the following Formula 3-1 to Formula 3-3:


Formula 3-1




Formula 3-2




Formula 3-3



[0082] In an embodiment, L of Formula 1 may be represented by one of the following Formulae 4-1 and 4-2:



Formula 4-1



Formula 4-2


[0083] According to an embodiment, the polycyclic compound represented by Formula 1 may be a material for emitting delayed fluorescence. The polycyclic compound of an embodiment may be a material for emitting thermally activated delayed fluorescence.

[0084] According to an embodiment, the polycyclic compound represented by Formula 1 may have an absolute value (ΔE_{ST}) of the difference between a singlet energy level (S1) and a triplet energy level (T1) of about 0.2 eV or less. For example, $S1-T1 \leq 0.2$ eV may be satisfied.

[0085] For example, the polycyclic compound represented by Formula 1 has a small difference between a singlet energy level (S1) and a triplet energy level (T1) and may be utilized as a material for emitting thermally activated delayed fluorescence. In an embodiment, the polycyclic compound represented by Formula 1 may be utilized as a material for emitting thermally activated delayed fluorescence, which emits blue light. However, an embodiment of the inventive concept is not limited thereto. The polycyclic compound of an embodiment may be a material for emitting thermally activated delayed fluorescence, which emits red light.


[0086] According to an embodiment, the polycyclic compound represented by Formula 1 may be any one among the compounds represented in the following Compound Group 1:

Compound Group 1




1


-continued




2



3




4




5

-continued




6

-continued




10


7




11




8




12




9



13




-continued




14


-continued




18




15




19




16




20



17




21




22

-continued


23




24




25



26




27




-continued


28




29




30



31




32




-continued

33




-continued


36




34




37




38




39




35




40




-continued




41




42



43




44




45


-continued



46




47




48

-continued




49


-continued




52




50




53



54




51




55

-continued




56


-continued




60




57




61




58



62




59




63

-continued




64


-continued




69




65




66




70



67




68



71


-continued

72




-continued


75




73




76



74




77




-continued

78




-continued

81




82

79




83

80




84




-continued


85




86




87



88




89




-continued


90




91




92



93




94




-continued


95




96




97



98




99




-continued


100




101



102




103




-continued


104




105




106



107




108




-continued


109



110




111




-continued


112




113



114




115




-continued


116




117



118




119




-continued

120




-continued


124




125




126




127




128




121



122




123




-continued


129




130




131



132




133




-continued


134




135




136








137








138




-continued



-continued




-continued




149


-continued




155




150




156




151




157




152




158



153




159












154


-continued



-continued




-continued




170


-continued




175




171




176




172




177




173



178




174




179

-continued




180


-continued




185




181




186




182




187




183



188




184




189

-continued




190


-continued




194




191




192



195




193



196


-continued

197





-continued

200





201

198




202

199




203




-continued

204





-continued

207





208

205




209


206



210




-continued




211


-continued




215




212




216



213




214




217

-continued




218


-continued




221




219




222



223




220




224

-continued




225

-continued




229


226




230




227



228




231




-continued

232




-continued


236




233




237




234




238




235



239




-continued




240


-continued




244




241




245




242



246




243



247


-continued

248




-continued


251




249




252



250




253




-continued

254




-continued

257




258


255



256




259




-continued

260




-continued


263




264



261



262



[0087] The polycyclic compound represented by Formula 1 may be utilized in an organic electroluminescence device **10** of an embodiment to improve the efficiency and life of the organic electroluminescence device. For example, the polycyclic compound represented by Formula 1 may be utilized in an emission layer EML of an organic electroluminescence device **10** of an embodiment and may improve the emission efficiency and life of the organic electroluminescence device.

[0088] In addition, the polycyclic compound of an embodiment includes two electron donors and one electron acceptor in one compound unit, in which the two electron donors each does not include a cyano group and the electron acceptor includes a benzonitrile part and a pyridine part. Accordingly, the difference between singlet energy and triplet energy is minimized or reduced and a triplet energy level is increased, and thus, the polycyclic compound may be utilized as a material for emitting blue light, which emits thermally activated delayed fluorescence. In addition, because the polycyclic compound has excellent electron accepting and electron donating properties, charge transfer (CT) in a molecule is smooth, and the compound may be utilized as a material for emitting thermally activated delayed fluorescence.

[0089] In an embodiment, the emission layer EML includes a host and a dopant, and the host may be a host for emitting delayed fluorescence and the dopant may be a dopant for emitting delayed fluorescence. According to an embodiment, the polycyclic compound represented by Formula 1 may be included as a dopant material of an emission

layer EML. For example, according to an embodiment, the polycyclic compound represented by Formula 1 may be utilized as a TADF dopant.

[0090] Meanwhile, in an embodiment, the emission layer EML may include a suitable (e.g., a known) host material. For example, in an embodiment, the emission layer EML may include, as a host material, tris(8-hydroxyquinolino)aluminum (Alq₃), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), poly(N-vinylcarbazole) (PVK), 9,10-di(naphthalene-2-yl)anthracene (ADN), 4,4',4''-tris(carbazol-9-yl)-triphenylamine (TCTA), 1,3,5-tris(N-phenylbenzimidazole-2-yl)benzene (TPBi), 3-tert-butyl-9,10-di(naphth-2-yl)anthracene (TBADN), distyrylarylene (DSA), 4,4'-bis(9-carbazolyl)-2,2'-dimethyl-biphenyl (CDBP), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), bis[2-(diphenylphosphino)phenyl]ether oxide (DPEPO), hexaphenyl cyclotriphosphazene (CP1), 1,4-bis(triphenylsilyl)benzene (UGH2), hexaphenylcyclotrisiloxane (DPSiO₃), octaphenylcyclotetrasiloxane (DPSiO₄), 2,8-bis(diphenylphosphoryl)dibenzofuran (PPF), etc. However, an embodiment of the inventive concept is not limited thereto. Any suitable (e.g., known) host materials for emitting delayed fluorescence other than the suggested host materials may be included.

[0091] In the organic electroluminescence device **10** of an embodiment, the emission layer EML may further include a suitable (e.g., a known) dopant material. In an embodiment, the emission layer EML may include, as a dopant, styryl derivatives (for example, 1,4-bis[2-(3-N-ethylcarbazoryl)vinyl]benzene (BCzVB), 4-(di-p-tolylamino)-4'-[(di-p-tolylamino)styryl]stilbene (DPAVB), and N-(4-((E)-2-(6-((E)-4-(diphenylamino)styryl)naphthalen-2-yl)vinyl)phenyl)-N-phenylbenzenamine (N-BDAVB), perylene and the derivatives thereof (for example, 2,5,8,11-tetra-*t*-butylperylene (TBP)), pyrene and the derivatives thereof (for example, 1,1-dipyrene, 1,4-dipyrenylbenzene, 1,4-bis(N,N-diphenylamino)pyrene), etc.

[0092] Referring to FIGS. **1** to **3** again, in the organic electroluminescence device **10** of an embodiment, the electron transport region ETR is provided on the emission layer EML. The electron transport region ETR may include at least one of an electron blocking layer, an electron transport layer ETL or an electron injection layer EIL. However, an embodiment of the inventive concept is not limited thereto.

[0093] The electron transport region ETR may have a single layer formed utilizing a single material, a single layer formed utilizing a plurality of different materials, or a multilayer structure having a plurality of layers formed utilizing a plurality of different materials.

[0094] For example, the electron transport region ETR may have a single layer structure of an electron injection layer EIL or an electron transport layer ETL, or a single layer structure formed utilizing an electron injection material and an electron transport material. Further, the electron transport region ETR may have a single layer structure having a plurality of different materials, or a structure laminated from the first electrode EL1 of electron transport layer ETL/electron injection layer EIL, or hole blocking layer/electron transport layer ETL/electron injection layer EIL, without being limited thereto. The thickness of the electron transport region ETR may be, for example, from about 100 Å to about 1,500 Å.

[0095] The electron transport region ETR may be formed utilizing various suitable methods, such as a vacuum depo-

sition method, a spin coating method, a cast method, a Langmuir-Blodgett (LB) method, an inkjet printing method, a laser printing method, and a laser induced thermal imaging (LITI) method.

[0096] If the electron transport region ETR includes an electron transport layer ETL, the electron transport region ETR may include, for example, tris(8-hydroxyquinolino)aluminum (Alq₃), 1,3,5-tri[(3-pyridyl)-phen-3-yl]benzene, 2,4,6-tris(3'-(pyridin-3-yl)biphenyl-3-yl)-1,3,5-triazine, 2-(4-(N-phenylbenzimidazolyl-1-ylphenyl)-9,10-dinaphthylanthracene, 1,3,5-tri(1-phenyl-1H-benzo[d]imidazol-2-yl)benzene (TPBi), 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), 3-(4-biphenyl)-4-phenyl-5-tert-butylphenyl-1,2,4-triazole (TAZ), 4-(naphthalen-1-yl)-3,5-diphenyl-4H-1,2,4-triazole (NTAZ), 2-(4-biphenyl)-5-(4-tert-butylphenyl)-1,3,4-oxadiazole (tBu-PBD), bis(2-methyl-8-quinolinolato-N1,O8)-(1,1'-biphenyl-4-olato)aluminum (BAIq), berylliumbis(benzoquinolin-10-olate) (Bebq2), 9,10-di(naphthalene-2-yl)anthracene (ADN), or a mixture thereof, without being limited thereto.

[0097] If the electron transport region ETR includes the electron transport layer ETL, the thickness of the electron transport layer ETL may be from about 100 Å to about 1,000 Å and may be, for example, from about 150 Å to about 500 Å. If the thickness of the electron transport layer ETL satisfies the above-described range, satisfactory electron transport properties may be obtained without substantial increase of a driving voltage.

[0098] If the electron transport region ETR includes the electron injection layer EIL, the electron transport region ETR may include, for example, LiF, lithium quinolate (LiQ), Li₂O, BaO, NaCl, CsF, a metal in lanthanoides (such as Yb), or a metal halide (such as RbCl, RbI and KI). However, an embodiment of the inventive concept is not limited thereto. The electron injection layer EIL may also be formed utilizing a mixture material of an electron transport material and an insulating organo metal salt. The organo metal salt may be a material having an energy band gap of about 4 eV or more. The organo metal salt may include, for example, metal acetates, metal benzoates, metal acetoacetates, metal acetylacetonates, or metal stearates.

[0099] If the electron transport region ETR includes the electron injection layer EIL, the thickness of the electron injection layer EIL may be from about 1 Å to about 100 Å, or from about 3 Å to about 90 Å. If the thickness of the electron injection layer EIL satisfies the above described ranges, satisfactory electron injection properties may be obtained without inducing substantial increase of a driving voltage.

[0100] The electron transport region ETR may include a hole blocking layer HBL as described above. The hole blocking layer may include, for example, at least one of 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), or 4,7-diphenyl-1,10-phenanthroline (Bphen). However, an embodiment of the inventive concept is not limited thereto.

[0101] The second electrode EL2 is provided on the electron transport region ETR. The second electrode EL2 has conductivity. The second electrode EL2 may be formed utilizing a metal alloy or a conductive compound. The second electrode EL2 may be a cathode. The second electrode EL2 may be a transmissive electrode, a transfective electrode or a reflective electrode. If the second electrode

EL2 is the transmissive electrode, the second electrode EL2 may include a transparent metal oxide, for example, ITO, IZO, ZnO, ITZO, etc.

[0102] If the second electrode EL2 is the transmissive electrode or the reflective electrode, the second electrode EL2 may include Ag, Mg, Cu, Al, Pt, Pd, Au, Ni, Nd, Ir, Cr, Li, Ca, LiF/Ca, LiF/Al, Mo, Ti, a compound thereof, or a mixture thereof (for example, a mixture of Ag and Mg). The second electrode EL2 may have a multilayered structure including a reflective layer or a transmissive layer formed utilizing the above-described materials and a transparent conductive layer formed utilizing ITO, IZO, ZnO, ITZO, etc.

[0103] Though not shown, the second electrode EL2 may be connected with an auxiliary electrode. If the second electrode EL2 is connected with the auxiliary electrode, the resistance of the second electrode EL2 may decrease.


[0104] In the organic electroluminescence device 10, according to the application of a voltage to each of the first electrode EL1 and second electrode EL2, holes injected from the first electrode EL1 may move via the hole transport region HTR to the emission layer EML, and electrons injected from the second electrode EL2 may move via the electron transport region ETR to the emission layer EML. The electrons and the holes are recombined in the emission layer EML to produce excitons, and the excitons may emit light via transition from an excited state to a ground state.

[0105] If the organic electroluminescence device 10 is a top emission device, the first electrode EL1 may be a reflective electrode and the second electrode EL2 may be a transmissive electrode or a transmissive electrode. If the organic electroluminescence device 10 is a bottom emission device, the first electrode EL1 may be a transmissive electrode or a transmissive electrode and the second electrode EL2 may be a reflective electrode.

[0106] The organic electroluminescence device of an embodiment of the inventive concept utilizes the polycyclic compound as a material for an emission layer and may have improved light-emitting efficiency and life characteristics.

[0107] An embodiment of the inventive concept provides a polycyclic compound represented by the following Formula 1:

Formula 1



[0108] In Formula 1, X_1 and X_2 are each independently a direct linkage, CR_7R_8 , SiR_9R_{10} , O, or S, and L is a direct linkage, CO, SO_2 , $SiR_{11}R_{12}$, a substituted or unsubstituted arylene group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroarylene group having 2 to 30 carbon atoms for forming a ring.

[0109] In Formula 1, R_1 to R_4 are each independently a hydrogen atom, a deuterium atom, a halogen atom, $NR_{13}R_{14}$, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring, or combined with an adjacent group to form a ring; a, b, c and d are each independently an integer of 0 to 4.

[0110] If each of a to d is an integer of 2 or more, a plurality of R_1 , R_2 , R_3 and R_4 may be respectively the same or different from each other.

[0111] In Formula 1, R_5 to R_{14} are each independently a hydrogen atom, a deuterium atom, a halogen atom, a cyano group, a substituted or unsubstituted amino group, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring, or combined with an adjacent group to form a ring; and e and f are each independently an integer of 0 to 3.

[0112] If each of e and f is an integer of 2 or more, a plurality of R_5 and R_6 may be respectively the same or different from each other.

[0113] The same explanation on the polycyclic compound in the organic electroluminescence device of an embodiment may be applied to the polycyclic compound of an embodiment, represented by Formula 1.

[0114] The polycyclic compound according to an embodiment may be any one selected from the compounds represented in Compound Group 1 above.

[0115] Hereinafter, the inventive concept will be more particularly explained referring to embodiments and comparative embodiments. The following embodiments are only illustrations to assist the understanding of the inventive concept, and the scope of the inventive concept is not limited thereto.


EXAMPLES

1. Synthesis of Polycyclic Compounds

[0116] First, the synthetic method of the polycyclic compounds according to exemplary embodiments of the inventive concept will be particularly explained referring to the synthetic methods of Compound 2, Compound 69, Compound 71, Compound 74, Compound 77, Compound 80, Compound 83, Compound 84, Compound 119, Compound 174, and Compound 212. In addition, the synthetic methods of the polycyclic compounds explained below are only embodiments, and the synthetic method of the polycyclic compound according to an embodiment of the inventive concept is not limited thereto.

Synthesis of Compound 2

[0117] Compound 2, which is a polycyclic compound according to an embodiment, may be synthesized, for example, by the following Reaction 1:




[0118] Under an argon atmosphere, to a 500 mL, three-neck flask, 5.00 g of 4-bromo-3-fluorobenzonitrile, 6.35 g of bis(pinacolato)diboron, 2.04 g of a [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) dichloromethane adduct, and 4.91 g of potassium acetate were added and dissolved in 130 mL of 1,4-dioxane, followed by stirring at about 90° C. for about 6 hours. After dissipating heat, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 5.25 g (yield 85%) of Intermediate A. The molecular weight of Intermediate A measured by FAB-MS was 247.

[0119] Under an argon atmosphere, in a 500 mL, three-neck flask, 5.00 g of Intermediate A, 2.66 g of 4-chloro-3-fluoropyridine, 0.23 g of palladium acetate ($\text{Pd}(\text{OAc})_2$), 0.83 g of 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (SPhos), and 8.59 g of tripotassium phosphate (K_3PO_4) were dissolved in 100 mL of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80° C. for about 10 hours. After cooling in the air, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 3.50 g (yield 80%) of Intermediate B. The molecular weight of Intermediate B measured by FAB-MS was 216.


[0120] Under an argon atmosphere, to a 200 mL, three-neck flask, 3.00 g of Intermediate B, 4.64 g of carbazole, and 9.59 g of potassium carbonate (K_2CO_3) were added and dissolved in 50 mL of anhydrous DMSO, followed by stirring at about 150° C. for about 6 hours. After cooling in the air, 300 mL of water was added to the reaction mixture, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 150 mL of CH_2Cl_2 . The resultant solution was dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 4.82 g (yield 68%) of Compound 2 was obtained as a target product. The molecular weight of the target product measured by FAB-MS was 510. From the result, the target product was identified as Compound 2.

Synthesis of Compound 69

[0121] Compound 69, which is a polycyclic compound according to an embodiment may be synthesized, for example, by the following Reaction 2:



-continued




[0122] Under an argon atmosphere, to a 500 mL, three-neck flask, 5.00 g of 2-fluoropyridine-3-boronic acid, 7.10 g of 3-bromo-4-fluorobenzonitrile, 0.40 g of $\text{Pd}(\text{OAc})_2$, 1.46 g of SPhos, and 15.06 g of K_3PO_4 were dissolved in 180 ml of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80° C. for about 10 hours. After finishing the reaction, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 5.98 g (yield 78%) of Intermediate C. The molecular weight of Intermediate C measured by FAB-MS was 216.

[0123] Under an argon atmosphere, to a 200 mL, three-neck flask, 3.00 g of Intermediate C, 5.42 g of dimethylcarbazole, and 9.59 g of K_2CO_3 were added and dissolved in 50 ml of anhydrous DMSO, followed by stirring at about 150° C. for about 6 hours. After cooling in the air, 300 ml of water was added to the reaction mixture, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 150 ml of CH_2Cl_2 . The resultant solution was dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 6.29 g (yield 80%) of Compound 69 was obtained as a target product. The molecular weight of the target product measured by FAB-MS was 566. From the result, the target product was identified as Compound 69.

Synthesis of Compound 71

[0124] Compound 71, which is a polycyclic compound according to an embodiment may be synthesized, for example, by the following Reaction 3:


Reaction 3



[0125] Under an argon atmosphere, to a 200 ml, three-neck flask, 3.00 g of Intermediate C, 8.86 g of 3,6-diphenylcarbazole, and 9.59 g of K_2CO_3 were added and dissolved in 50 ml of anhydrous DMSO, followed by stirring at about 150° C. for about 6 hours. After cooling in the air, 300 ml of water was added to the reaction mixture, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 150 ml of CH_2Cl_2 . The resultant solution was dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 8.14 g (yield 72%) of Compound 71 was obtained as a target product. The molecular weight of the target product measured by FAB-MS was 814. From the result, the target product was identified as Compound 71.

Synthesis of Compound 74

[0126] Compound 74, which is a polycyclic compound according to an embodiment may be synthesized, for example, by the following Reaction 4:



[0127] Under an argon atmosphere, to a 200 ml, three-neck flask, 5.00 g of 3-bromo-4-fluorobenzonitrile, 4.18 g of carbazole and 5.18 g of K_2CO_3 were added and dissolved in 50 ml of anhydrous DMSO, followed by stirring at about 110° C. for about 8 hours. After cooling in the air, 300 ml of water was added to the reaction solution, followed by


stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 150 ml of CH_2Cl_2 . The resultant solution was dried with $MgSO_4$, and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 6.08 g (yield 70%) of Intermediate D was obtained. The molecular weight of Intermediate D measured by FAB-MS was 347.


[0128] Under an argon atmosphere, in a 500 ml, three-neck flask, 6.00 g of Intermediate D, 2.44 g of (2-fluoropyridine-3-yl)boronic acid, 0.19 g of $Pd(OAc)_2$, 0.71 g of SPhos, and 7.34 g of K_3PO_4 were dissolved in 100 ml of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80° C. for about 16 hours. After finishing the reaction, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with $MgSO_4$, and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 3.08 g (yield 49%) of Intermediate E. The molecular weight of Intermediate E measured by FAB-MS was 363.

[0129] Under an argon atmosphere, to a 200 ml, three-neck flask, 3.00 g of Intermediate E, 2.74 g of 3,9'-bicarbazole, and 2.28 g of K_2CO_3 were added and dissolved in 30 ml of anhydrous DMSO, followed by stirring at about 150° C. for about 6 hours. After cooling in the air, 300 ml of water was added to the reaction mixture, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 150 ml of CH_2Cl_2 . The resultant solution was dried with $MgSO_4$, and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 3.07 g (yield 55%) of Compound 74 was obtained as a target product. The molecular weight of the target product measured by FAB-MS was 675. From the result, the target product was identified as Compound 74.

Synthesis of Compound 77

[0130] Compound 77, which is a polycyclic compound according to an embodiment may be synthesized, for example, by the following Reaction 5:






[0131] Under an argon atmosphere, to a 200 ml, three-neck flask, 5.00 g of 3-bromo-4-fluorobenzonitrile, 8.31 g of 3,9'-bicarbazole and 6.91 g of K_2CO_3 were added and dissolved in 50 ml of anhydrous DMSO, followed by stirring at about 110° C. for about 8 hours. After cooling in the air, 300 ml of water was added to the reaction solution, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 150 ml of CH_2Cl_2 . The resultant solution was dried with $MgSO_4$, and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 7.56 g (yield 59%) of Intermediate F was obtained. The molecular weight of Intermediate F measured by FAB-MS was 512.


[0132] Under an argon atmosphere, in a 500 ml, three-neck flask, 7.00 g of Intermediate F, 1.93 g of (2-fluoropyridine-3-yl)boronic acid, 0.15 g of $Pd(OAc)_2$, 0.56 g of SPhos, and 5.80 g of K_3PO_4 were dissolved in 100 ml of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80° C. for about 16 hours. After finishing the reaction, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with $MgSO_4$, and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 3.03 g (yield 42%) of Intermediate G. The molecular weight of Intermediate G measured by FAB-MS was 528.

[0133] Under an argon atmosphere, to a 200 ml, three-neck flask, 3.00 g of Intermediate G, 0.95 g of carbazole, and 1.56 g of K_2CO_3 were added and dissolved in 30 ml of anhydrous DMSO, followed by stirring at about 150° C. for about 6 hours. After cooling in the air, 300 ml of water was added to the reaction mixture, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 150 ml of CH_2Cl_2 . The resultant solution was dried with $MgSO_4$, and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 2.15 g (yield 56%) of Compound 77 was obtained as a target product. The molecular weight of the target product measured by FAB-MS was 675. From the result, the target product was identified as Compound 77.

Synthesis of Compound 80

[0134] Compound 80, which is a polycyclic compound according to an embodiment may be synthesized, for example, by the following Reaction 6:






[0135] Under an argon atmosphere, in a 500 ml, three-neck flask, 5.00 g of 2,6-difluoro-3-pyridineboronic acid, 5.46 g of 5-chloro-2,4-difluorobenzonitrile, 0.35 g of $\text{Pd}(\text{OAc})_2$, 1.29 g of SPhos, and 13.36 g of K_3PO_4 were dissolved in 160 ml of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80°C . for about 12 hours. After finishing the reaction, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 5.71 g (yield 72%) of Intermediate H. The molecular weight of Intermediate H measured by FAB-MS was 252.

[0136] Under an argon atmosphere, to a 200 ml, three-neck flask, 3.00 g of Intermediate H, 7.96 g of carbazole, and 16.44 g of K_2CO_3 were added and dissolved in 80 ml of anhydrous DMSO, followed by stirring at about 150°C . for about 8 hours. After cooling in the air, 500 ml of water was added to the reaction mixture, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 200 ml of CH_2Cl_2 . The resultant solution was dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 4.80 g (yield 48%) of Compound 80 was obtained as a target product. The molecular weight of the target product measured by FAB-MS was 840. From the result, the target product was identified as Compound 80.

Synthesis of Compound 83

[0137] Compound 83, which is a polycyclic compound according to an embodiment may be synthesized, for example, by the following Reaction 7:




[0138] Under an argon atmosphere, in a 500 ml, three-neck flask, 5.00 g of 2,6-difluoro-3-iodopyridine, 3.80 g of 3-cyano-2,4-difluorophenylboronic acid, 1.20 g of tetrakis (triphenylphosphine)palladium(0), 0.85 g of SPhos, and 5.74 g of K_2CO_3 were dissolved in 100 ml of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80°C . for about 10 hours. After finishing the reaction, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 3.98 g (yield 76%) of Intermediate J. The molecular weight of Intermediate J measured by FAB-MS was 252.

[0139] Under an argon atmosphere, to a 200 ml, three-neck flask, 3.00 g of Intermediate J, 7.96 g of carbazole, and 16.44 g of K_2CO_3 were added and dissolved in 80 ml of anhydrous DMSO, followed by stirring at about 150°C . for about 8 hours. After cooling in the air, 500 ml of water was added to the reaction mixture, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 200 ml of CH_2Cl_2 . The resultant

solution was dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 4.60 g (yield 46%) of Compound 83 was obtained as a target product. The molecular weight of the target product measured by FAB-MS was 840. From the result, the target product was identified as Compound 83.

Synthesis of Compound 84

[0140] Compound 84, which is a polycyclic compound according to an embodiment may be synthesized, for example, by the following Reaction 8:




[0141] Under an argon atmosphere, in a 500 ml, three-neck flask, 5.00 g of (2-fluoropyridin-3-yl)boronic acid, 6.99 g of 4,5-dichlorophthalonitrile, 0.40 g of $\text{Pd}(\text{OAc})_2$, 1.47 g of SPhos, and 15.06 g of K_3PO_4 were dissolved in 180 ml of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80°C . for about 8 hours. After finishing the reaction, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 4.85 g (yield 53%) of Intermediate K. The molecular weight of Intermediate K measured by FAB-MS was 257.

[0142] Under an argon atmosphere, to a 200 ml, three-neck flask, 3.00 g of Intermediate K, 3.89 g of carbazole, and 8.05 g of K_2CO_3 were added and dissolved in 40 ml of anhydrous DMSO, followed by stirring at about 150°C . for about 8 hours. After cooling in the air, 300 ml of water was added to the reaction mixture, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 150 ml of CH_2Cl_2 . The resultant solution was dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 4.37 g (yield 70%) of Compound 84 was obtained as a target product. The molecular weight of the target product measured by FAB-MS was 535. From the result, the target product was identified as Compound 84.

Synthesis of Compound 119

[0143] Compound 119, which is a polycyclic compound according to an embodiment may be synthesized, for example, by the following Reaction 9:





[0144] Under an argon atmosphere, in a 500 ml, three-neck flask, 5.00 g of (2,6-difluoropyridin-4-yl)boronic acid, 6.29 g of 3-bromo-4-fluorobenzonitrile, 0.35 g of $\text{Pd}(\text{OAc})_2$, 1.29 g of SPhos, and 13.36 g of K_3PO_4 were dissolved in 160 ml of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80° C. for about 8 hours. After finishing the reaction, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 4.79 g (yield 65%) of Intermediate L. The molecular weight of Intermediate L measured by FAB-MS was 234.

[0145] Under an argon atmosphere, to a 200 ml, three-neck flask, 3.00 g of Intermediate L, 6.43 g of carbazole, and 14.16 g of K_2CO_3 were added and dissolved in 50 ml of anhydrous DMSO, followed by stirring at about 150° C. for about 10 hours. After cooling in the air, 300 ml of water was added to the reaction mixture, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 150 ml of CH_2Cl_2 . The resultant solution was dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 6.06 g (yield 70%) of Compound 119 was obtained as a target product. The molecular weight of the target product measured by FAB-MS was 675. From the result, the target product was identified as Compound 119.

Synthesis of Compound 174

[0146] Compound 174, which is a polycyclic compound according to an embodiment may be synthesized, for example, by the following Reaction 10:



[0147] Under an argon atmosphere, to a 1 L, three-neck flask, 10.00 g of 2,6-dichlorobenzonitrile, 4.79 g of 5-cyano-2-fluorophenylboronic acid, 0.32 g of $\text{Pd}(\text{OAc})_2$, 1.19 g of SPhos, and 12.34 g of K_3PO_4 were dissolved in 290 ml of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80° C. for about 8 hours. After finishing the reaction, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 5.07 g (yield 68%) of Intermediate M. The molecular weight of Intermediate M measured by FAB-MS was 256.


[0148] Under an argon atmosphere, to a 500 ml, three-neck flask, 5.00 g of Intermediate M, 2.75 g of (2-fluoropyridin-3-yl)boronic acid, 0.22 g of $\text{Pd}(\text{OAc})_2$, 0.80 g of SPhos, and 8.27 g of K_3PO_4 were dissolved in 100 ml of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80° C. for about 8 hours. After finishing the reaction, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 3.96 g (yield 64%) of Intermediate N. The molecular weight of Intermediate N measured by FAB-MS was 317.

[0149] Under an argon atmosphere, to a 200 ml, three-neck flask, 3.00 g of Intermediate N, 3.16 g of carbazole, and 6.53 g of K_2CO_3 were added and dissolved in 40 ml of anhydrous DMSO, followed by stirring at about 150° C. for about 8 hours. After cooling in the air, 300 ml of water was added to the reaction mixture, followed by stirring. The precipitate thus obtained was recovered by filtering with suction and dissolved in 150 ml of CH_2Cl_2 . The resultant solution was dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 3.87 g (yield 67%) of Compound 174 was obtained as a target

product. The molecular weight of the target product measured by FAB-MS was 611. From the result, the target product was identified as Compound 174.

Synthesis of Compound 212

[0150] Compound 212, which is a polycyclic compound according to an embodiment may be synthesized, for example, by the following Reaction 11:



[0151] Under an argon atmosphere, to a 500 ml, three-neck flask, 5.00 g of (6-bromopyridin-3-yl)boronic acid, 7.63 g of 2-bromo-5-iodobenzonitrile, 0.28 g of $\text{Pd}(\text{OAc})_2$, 1.02 g of SPhos, and 10.52 g of K_3PO_4 were dissolved in 120 ml of a deaerated mixture solvent of toluene/ethanol/water (10:1:2), followed by stirring at about 80°C . for about 12 hours. After finishing the reaction, water was added, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by silica gel column chromatography to obtain 6.20 g (yield 74%) of Intermediate O. The molecular weight of Intermediate O measured by FAB-MS was 338.

[0152] Under an argon atmosphere, to a 200 ml, three-neck flask, 3.00 g of Intermediate O, 3.71 g of 9,9-dimethyl-9,10-dihydroacridine, 0.51 g of bis(dibenzylideneacetone) palladium(0), 1.78 ml of tri-tert-butylphosphine (2 M in toluene) and 1.71 g of sodium tert-butoxide were added and dissolved in 50 ml of anhydrous toluene, followed by heating and refluxing for about 8 hours. After finishing the reaction, water was added to the reaction mixture, and extraction with CH_2Cl_2 was conducted. An organic layer was separated and dried with MgSO_4 , and solvents were removed by distillation under a reduced pressure. The crude product thus obtained was separated by recrystallization, and 3.96 g (yield 75%) of Compound 212 was obtained as a target product. The molecular weight of the target product measured by FAB-MS was 594. From the result, the target product was identified as Compound 212.

2. Manufacture and Evaluation of Organic Electroluminescence Device Including a Polycyclic Compound

Manufacture of Organic Electroluminescence Devices


[0153] Organic electroluminescence devices of exemplary embodiments including the polycyclic compounds of exemplary embodiments in an emission layer were manufactured by a method described below. Organic electroluminescence devices of Examples 1 to 11 were manufactured utilizing the polycyclic compounds of Compound 2, Compound 69, Compound 71, Compound 74, Compound 77, Compound 80, Compound 83, Compound 84, Compound 119, Compound 174 and Compound 212 respectively as materials for an emission layer. Compounds utilized in the emission layer in Examples 1 to 11 and Comparative Examples 1 to 5 are shown below in Table

TABLE 1


Example Compounds	

TABLE 1-continued

69



71



74




TABLE 1-continued








TABLE 1-continued


84



119



174



212







TABLE 1-continued

Comparative
Compounds


X-1



X-2




X-3



X-4

TABLE 1-continued



[0154] The organic electroluminescence devices of the examples and the comparative examples were manufactured by a method described below.

[0155] On a glass substrate, ITO was patterned to a thickness of about 1,500 Å and washed with ultra-pure water, and a UV ozone treatment was conducted for about 10 minutes. Then, HAT-CN was deposited to a thickness of about 100 Å, α-NPD was deposited to a thickness of about 800 Å, and mCP was deposited to a thickness of about 50 Å to form a hole transport region.

[0156] Then, each of the polycyclic compounds of an embodiment and the Comparative Compound, and DPEPO were co-deposited in a ratio of 18:82 to form an emission layer to a thickness of about 200 Å. Then, a layer with a thickness of about 100 Å was formed utilizing DPEPO. That is, in order to form the emission layer by the co-deposition, each of Compounds 2, 69, 71, 74, 77, 80, 83, 84, 119, 174 and 212 was mixed with DPEPO and deposited in Examples 1 to 11, and each of Comparative Compounds X-1, X-2, X-3, X-4 and X-5 was mixed with DPEPO and deposited in Comparative Examples 1 to 5.

[0157] On the emission layer, a layer was formed utilizing TPBi to a thickness of about 300 Å, and a layer was formed utilizing LiF to a thickness of about 5 Å to form an electron transport region. Then, a second electrode was formed utilizing aluminum (Al) to a thickness of about 1,000 Å.

[0158] In the examples, the hole transport region, the emission layer, the electron transport region and the second electrode were formed by utilizing a vacuum deposition apparatus.

Energy Levels of Each Compound

[0159] In Table 2 below, the singlet (S1) energy levels and the triplet (T1) energy levels of Compounds 2, 69, 71, 74, 77, 80, 83, 84, 119, 174 and 212, which are example compounds, and Comparative Compounds X-1, X-2, X-3, X-4 and X-5 are shown. The energy level values in Table 2 were calculated by a non-empirical molecular orbital method. Particularly, the energy level values were calculated by B3LYP/6-31G(d) utilizing Gaussian 09 of Gaussian Co. ΔE_{ST} represents the difference between a singlet (S1) energy level and a triplet (T1) energy level.

TABLE 2

Compound	S1 energy level	T1 energy level	ΔE_{ST}
Compound 2	2.96	2.87	0.09
Compound 69	2.96	2.81	0.15
Compound 71	2.85	2.72	0.13
Compound 74	2.82	2.80	0.02

TABLE 2-continued

Compound	S1 energy level	T1 energy level	ΔE_{ST}
Compound 77	2.95	2.83	0.12
Compound 80	2.97	2.80	0.17
Compound 83	2.96	2.85	0.11
Compound 84	2.55	2.47	0.08
Compound 119	2.99	2.81	0.18
Compound 174	3.08	2.99	0.09
Compound 212	2.45	2.44	0.01
Comparative Compound X-1	3.20	2.85	0.35
Comparative Compound X-2	2.96	2.87	0.09
Comparative Compound X-3	3.28	3.07	0.21
Comparative Compound X-4	3.35	3.14	0.21
Comparative Compound X-5	3.24	3.03	0.21

[0160] Compounds 2, 69, 71, 74, 77, 80, 83, 84, 119, 174 and 212, which are example compounds, showed low ΔE_{ST} values of about 0.2 eV or less. In comparison, Comparative Compounds X-1, and X-3 to X-5 were found to show ΔE_{ST} values greater than about 0.2 eV.

Evaluation of Properties of Organic Electroluminescence Device

[0161] In order to evaluate the properties of the organic electroluminescence devices of the examples and the comparative examples, the maximum emission wavelength (nm) and external quantum yield (%) were measured. The measurement was conducted utilizing a luminous brightness measurement apparatus, C9920-11 of HAMAMATSU Photonics Co.

TABLE 3

	Emission layer dopant	Maximum emission wavelength (nm)	External quantum yield (%)
Example 1	Compound 2	492	10
Example 2	Compound 69	462	12
Example 3	Compound 71	460	11
Example 4	Compound 74	460	13
Example 5	Compound 77	448	10
Example 6	Compound 80	458	13
Example 7	Compound 83	456	14
Example 8	Compound 84	486	10
Example 9	Compound 119	455	10
Example 10	Compound 174	446	8
Example 11	Compound 212	490	12
Comparative Example 1	Comparative Compound X-1	445	1

TABLE 3-continued

	Emission layer dopant	Maximum emission wavelength (nm)	External quantum yield (%)
Comparative Example 2	Comparative Compound X-2	458	4
Comparative Example 3	Comparative Compound X-3	430	2
Comparative Example 4	Comparative Compound X-4	428	1
Comparative Example 5	Comparative Compound X-5	431	2

[0162] Referring to Table 3, it may be found that the organic electroluminescence devices of Examples 1 to 11, which utilized the polycyclic compounds of exemplary embodiments as dopant materials of an emission layer, showed excellent external quantum efficiency when compared to Comparative Examples 1 to 5. In addition, the organic electroluminescence devices of Examples 1 to 11, which utilized the polycyclic compounds of exemplary embodiments as the dopant materials of an emission layer, and the organic electroluminescence devices of Comparative Examples 1 to 5, which utilized Comparative Compounds X-1 to X-5, were found to have emission wavelength of about 495 nm or less and emit blue light.

[0163] Referring to the results of Tables 2 and 3, the Example Compounds had a small ΔE_{ST} and showed high emission efficiency in a blue emission region at the same time, thus might be utilized as materials for thermally activated delayed fluorescence with high efficiency.

[0164] The organic electroluminescence device of an embodiment includes the polycyclic compound of an embodiment in an emission layer and may accomplish deep blue light with relatively short wavelengths and may show high emission efficiency at the same time.

[0165] Also, the polycyclic compound of an embodiment has an electron donor-electron acceptor-electron donor (D-A-D) structure, and the electron donor does not include a cyano group but the electron acceptor includes a benzonitrile part and a pyridine part at the same time, and thus, the polycyclic compound shows a low ΔE_{ST} value and high emission efficiency.

[0166] In comparison to the examples, Comparative Compound X-1 had an electron donor-electron donor-electron acceptor (D-D-A) structure, and the electron acceptor included only a pyridine part. Thus, this compound showed a high ΔE_{ST} value and may not be utilized as a thermally activated delayed fluorescence material. In addition, the organic electroluminescence device of Comparative Example 1 showed low external quantum efficiency than those of the examples.

[0167] Comparative Compound X-2, which was utilized in Comparative Example 2, had a low ΔE_{ST} value, but included only a benzonitrile part in an electron acceptor, and thus, had low electron accepting properties. Accordingly, charge transfer (CT) in a molecule was not smooth and thermally activated delayed fluorescence was not shown. Therefore, the organic electroluminescence device of Comparative Example 2 was found to show low external quantum efficiency.

[0168] Comparative Compounds X-3 and X-4, which were utilized in Comparative Examples 3 and 4, showed relatively not high ΔE_{ST} values but included only a pyridine

part in an electron acceptor, and thus, the accepting properties of the electron acceptor was low. Accordingly, charge transfer (CT) in a molecule was not smooth and thermally activated delayed fluorescence was not shown. Therefore, the organic electroluminescence devices of Comparative Examples 3 and 4 were found to show low external quantum efficiency.

[0169] In addition, Comparative Compound X-5 utilized in Comparative Example 5 showed a relatively not high ΔE_{ST} value but included only a pyridine part in an electron acceptor, and thus, the accepting properties of the electron acceptor was low. Because the carbazole group of the electron donor included a cyano group, the donating properties of the electron donor was low and charge transfer (CT) in a molecule was not smooth, and thus, thermally activated delayed fluorescence was not shown. Therefore, the organic electroluminescence device of Comparative Example 5 was found to show low external quantum efficiency.

[0170] Referring to Table 3, the organic electroluminescence devices of exemplary embodiments, which utilized the polycyclic compounds of exemplary embodiments as luminescent materials of an emission layer, showed higher emission efficiency than the comparative examples. In addition, with regard to emission wavelengths, it was found that maximum emission wavelength of about 495 nm or less was shown and deep blue color was achieved.

[0171] The organic electroluminescence device of an embodiment has an electron donor-electron acceptor-electron donor (D-A-D) structure, which includes two electron donors and one electron acceptor, and utilizes a polycyclic compound including an electron acceptor having a benzonitrile part and a pyridine part as a material for an emission layer. Thus, high emission efficiency in a blue light wavelength region may be achieved.

[0172] The organic electroluminescence device according to an embodiment of the inventive concept may attain high efficiency and long life.

[0173] The polycyclic compound according to an embodiment of the inventive concept may improve the life and efficiency of an organic electroluminescence device.

[0174] Throughout the disclosure, the use of “may” when describing embodiments of the present invention refers to “one or more embodiments of the present invention.” Moreover, any numerical range recited herein is intended to include all sub-ranges of the same numerical precision subsumed within the recited range. For example, a range of “1.0 to 10.0” is intended to include all subranges between (and including) the recited minimum value of 1.0 and the recited maximum value of 10.0, that is, having a minimum value equal to or greater than 1.0 and a maximum value equal to or less than 10.0, such as, for example, 2.4 to 7.6. Any maximum numerical limitation recited herein is intended to include all lower numerical limitations subsumed therein and any minimum numerical limitation recited in this specification is intended to include all higher numerical limitations subsumed therein. Accordingly, Applicant reserves the right to amend this specification, including the claims, to expressly recite any sub-range subsumed within the ranges expressly recited herein. All such ranges are intended to be inherently described in this specification such that amending to expressly recite any such subranges


would comply with the requirements of 35 U.S.C. § 112, first paragraph, or 35 U.S.C. § 112(a), and 35 U.S.C. § 132(a).

[0175] Although the exemplary embodiments of the present invention have been described, it is understood that the present invention should not be limited to these exemplary embodiments but various changes and modifications can be made by one ordinary skilled in the art within the spirit and scope of the present invention as hereinafter claimed, and equivalents thereof.

What is claimed is:

1. An organic electroluminescence device, comprising:
a first electrode;
a hole transport region on the first electrode;
an emission layer on the hole transport region;
an electron transport region on the emission layer; and
a second electrode on the electron transport region,
wherein the emission layer comprises a polycyclic compound comprising two electron donors and one electron acceptor, and the electron acceptor comprises a benzonitrile part and a pyridine part.
2. The organic electroluminescence device of claim 1, wherein the emission layer is configured to emit delayed fluorescence.
3. The organic electroluminescence device of claim 1, wherein the emission layer is a delayed fluorescence emission layer comprising a host and a dopant, and the dopant is the polycyclic compound.
4. The organic electroluminescence device of claim 1, wherein the emission layer is a thermally activated delayed fluorescence emission layer configured to emit blue light.
5. The organic electroluminescence device of claim 1, wherein the polycyclic compound is represented by following Formula 1:

Formula 1



wherein in Formula 1,

X_1 and X_2 are each independently a direct linkage, CR_7R_8 , SiR_9R_{10} , O, or S,

L is a direct linkage, CO, SO_2 , $SiR_{11}R_{12}$, a substituted or unsubstituted arylene group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroarylene group having 2 to 30 carbon atoms for forming a ring,

R_1 to R_4 are each independently a hydrogen atom, a deuterium atom, a halogen atom, $NR_{13}R_{14}$, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring;

R_1 to R_4 are each optionally independently combined with an adjacent group to form a ring,


a, b, c and d are each independently an integer of 0 to 4, R_5 to R_{14} are each independently a hydrogen atom, a deuterium atom, a halogen atom, a cyano group, a substituted or unsubstituted amino group, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring;

R_5 to R_{14} are each optionally independently combined with an adjacent group to form a ring, and


e and f are each independently an integer of 0 to 3.

6. The organic electroluminescence device of claim 5, wherein the polycyclic compound represented by Formula 1 is represented by one of following Formula 1-1 to Formula 1-3:


Formula 1-1



Formula 1-2



Formula 1-3




wherein in Formula 1-1 to Formula 1-3, X_1 and X_2 , L, R_1 to R_6 , and a to f are the same as respectively defined in association with Formula 1.

7. The organic electroluminescence device of claim 5, wherein X_1 and X_2 of Formula 1 are the same.

8. The organic electroluminescence device of claim 5, wherein R_5 and R_6 of Formula 1 are each independently represented by one of following Formula 2-1 to Formula 2-4:


10. The organic electroluminescence device of claim 5, wherein the compound represented by Formula 1 is any one among compounds represented in following Compound Group 1:

Formula 2-1




[Compound Group 1]


Formula 2-2



Formula 2-3




Formula 2-4




9. The organic electroluminescence device of claim 5, wherein R_1 to R_4 of Formula 1 are each independently represented by one of following Formula 3-1 to Formula 3-3:



Formula 3-1




Formula 3-2




Formula 3-3




1



2




3




4

-continued




5

-continued




9


6




10




7




11




8



12




-continued




13


-continued




17




14




18




15




19



16




20




21

-continued




22


-continued




27




23




28




24




29




25



30




26




31

-continued




32


-continued




36




33




37




34




38



35




39




40


-continued




41




42



43




44




45


-continued



46




47




48

-continued




49


-continued



52





50



53


51



54


55

-continued




56


-continued




61




57




62




58




63




59



64




60



65


66

-continued




66

-continued




70

67





71

68




72

69




67

-continued




73

-continued




76


74




77



75




78




-continued

79




-continued

82




83

80




84

81




85




-continued


86




87




88



89




90




-continued


91




92




93



94




95




-continued


101




102



103




104




100

-continued


105




106




107



109




110




-continued


111







112






113



-continued




-continued




-continued

120




-continued


124




125




126




127




128




121



122




123




-continued


129




130




131



132




133




-continued


134




135




136




137



138




-continued




139


-continued




144




140




145




141




146




142



147









143








148






-continued









-continued




-continued



-continued




-continued




171

-continued




176


172




173




177




174





178



175




179




180

-continued




181


-continued




186




182




187




183




188




184



189




185




190

-continued


191




192



193




194




-continued


195



196




197




-continued

198




-continued


201




199




202




203




200









204




-continued



-continued




-continued




212


-continued




216




213




217



214




218




215


-continued





-continued




220



221




224




226

-continued




226


-continued




229




227



230




228




231

-continued




232


-continued




236




233




237




234



238




235




239

-continued




240


-continued




244




241




245




242



246




243




247

-continued




248

-continued




251


249




252



250




253




-continued

254




-continued


257




255




258



256




259




-continued

260




-continued


263



264





261



11. An organic electroluminescence device, comprising:
a first electrode;
a hole transport region on the first electrode;
an emission layer on the hole transport region;
an electron transport region on the emission layer; and
a second electrode on the electron transport region,
wherein the emission layer comprises a polycyclic compound represented by following Formula 1:

262

Formula 1



wherein in Formula 1,
 X_1 and X_2 are each independently a direct linkage,
 CR_7R_8 , SiR_9R_{10} , O, or S,
L is a direct linkage, CO, SO_2 , $SiR_{11}R_{12}$, a substituted or unsubstituted arylene group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroarylene group having 2 to 30 carbon atoms for forming a ring,

R_1 to R_4 are each independently a hydrogen atom, a deuterium atom, a halogen atom, $NR_{13}R_{14}$, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring;

R_1 to R_4 are each optionally independently combined with an adjacent group to form a ring,

a , b , c and d are each independently an integer of 0 to 4,


R_5 to R_{14} are each independently a hydrogen atom, a deuterium atom, a halogen atom, a cyano group, a substituted or unsubstituted amino group, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring;

R_5 to R_{14} are each optionally independently combined with an adjacent group to form a ring, and


e and f are each independently an integer of 0 to 3.

12. The organic electroluminescence device of claim 11, wherein the polycyclic compound represented by Formula 1 is represented by one of following Formula 1-1 to Formula 1-3:


Formula 1-1



Formula 1-2




Formula 1-3



wherein in Formula 1-1 to Formula 1-3, X_1 and X_2 , L , R_1 to R_6 , and a to f are the same as respectively defined in association with Formula 1.

13. A polycyclic compound represented by following Formula 1:

Formula 1



wherein in Formula 1,

X_1 and X_2 are each independently a direct linkage, CR_7R_8 , SiR_9R_{10} , O , or S ,

L is a direct linkage, CO , SO_2 , $SiR_{11}R_{12}$, a substituted or unsubstituted arylene group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroarylene group having 2 to 30 carbon atoms for forming a ring,

R_1 to R_4 are each independently a hydrogen atom, a deuterium atom, a halogen atom, $NR_{13}R_{14}$, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring;

R_1 to R_4 are each optionally independently combined with an adjacent group to form a ring,


a , b , c and d are each independently an integer of 0 to 4, R_5 to R_{14} are each independently a hydrogen atom, a deuterium atom, a halogen atom, a cyano group, a substituted or unsubstituted amino group, a substituted or unsubstituted alkyl group having 1 to 20 carbon atoms, a substituted or unsubstituted aryl group having 6 to 30 carbon atoms for forming a ring, or a substituted or unsubstituted heteroaryl group having 2 to 30 carbon atoms for forming a ring;

R_5 to R_{14} are each optionally independently combined with an adjacent group to form a ring, and

e and f are each independently an integer of 0 to 3.


14. The polycyclic compound of claim 13, wherein the polycyclic compound represented by Formula 1 is represented by one of following Formula 1-1 to Formula 1-3:

Formula 1-1




-continued

Formula 1-2



Formula 1-3



wherein in Formula 1-1 to Formula 1-3, X_1 and X_2 , L , R_1 to R_6 , and a to f are the same as respectively defined in association with Formula 1.


15. The polycyclic compound of claim 13, wherein the compound represented by Formula 1 is a material for emitting thermally activated delayed fluorescence.

16. The polycyclic compound of claim 13, wherein the compound represented by Formula 1 has an absolute value of a difference between a singlet energy level and a triplet energy level of about 0.2 eV or less.


17. The polycyclic compound of claim 13, wherein X_1 and X_2 of Formula 1 are the same.

18. The polycyclic compound of claim 13, wherein R_5 and R_6 of Formula 1 are each independently one of following Formula 2-1 to Formula 2-4:

Formula 2-1




Formula 2-2




-continued

Formula 2-3




Formula 2-4




19. The polycyclic compound of claim 13, wherein R_1 to R_4 of Formula 1 are each independently one of following Formula 3-1 to Formula 3-3:


Formula 3-1



Formula 3-2




Formula 3-3




20. The polycyclic compound of claim 13, wherein the compound represented by Formula 1 is any one among compounds represented in following Compound Group 1:

[Compound Group 1]




-continued




2


-continued




6




3




7




4



8




5




9

-continued




10


-continued




14




11




15




12



16




13




17

-continued


18




19




20



21




22




-continued


23




24




25




26




27




-continued




28




29



30




31




32


-continued



33




34



35


-continued

36




-continued


40




37




41




42




38




43



39




44




-continued

45




-continued


48




46




49




47



50




-continued




51


-continued




55




52




56




57




53



58




54



59


-continued

60




-continued

64




65


61




62




67




63



68




-continued




69


-continued




72




70



73




71



74


-continued

75




-continued


78




76




79



77




80




-continued

81




-continued

84




85

82





86

83




87




-continued


88




89




90



91




92




-continued


93




94




95



96




97




-continued


98




99




100



101




102




-continued


103




104




105




106



107




-continued




108


-continued




111




109



112




110




113

-continued


114




115



116




117




-continued


118




119



120




121




-continued


122




123




124



125




126




-continued


127




128




129




130



131




-continued




132


-continued




137




133




138




134




139




135



140




136




141


-continued




142




143



144




145




146


-continued




147




148




149



150




151




152

-continued


153




154




155



156




157




-continued


158




159




160




161




162




-continued




163




164




165



166




167




168


-continued




169




170




171




172




173




174




175



176




177




178


-continued




179




180




181



182








183







184

-continued




-continued




-continued

194




-continued


197




195




198



196




199




-continued

200




-continued


204




201



205





202







206





203




-continued



-continued




-continued




215


-continued




218




216



219




217




220

-continued




221


-continued




225




222




226




223



227




224



228


-continued

229




-continued


232




230




233




234



235




231




-continued

236




-continued


240




237




241




238




242



239




243




-continued

244




-continued


248




245




249




246



250




247




-continued

251




-continued


254




252




255



253




256




-continued

257




-continued


260



258





261




262


259



-continued



-continued



* * * * *

专利名称(译)	有机电致发光器件和用于有机电致发光器件的多环化合物		
公开(公告)号	US20190296247A1	公开(公告)日	2019-09-26
申请号	US16/270108	申请日	2019-02-07
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	SAKAMOTO NAOYA		
发明人	SAKAMOTO, NAOYA		
IPC分类号	H01L51/00 C07D401/14 C09K11/06		
CPC分类号	H01L2251/301 H01L51/001 C09K11/06 H01L51/56 H01L51/0067 H01L51/5206 C09K2211/1018 H01L2251/558 H01L51/5056 H01L2251/308 H01L51/5088 H01L51/5072 C07D401/14 H01L51/0072 H01L51/5012 H01L51/0021 H01L51/5092 H01L51/5221 C07D405/14 C07D413/14 C07D417/14 C07D519/00 C07F7/081 C09K2211/1007 C09K2211/1029 C09K2211/1033 C09K2211/1037 C09K2211/104 C09K2211/1088 H01L51/0071 H01L51/0094 H01L51/5024 C07D491/10 C07F7/0814 C07F7/0816		
优先权	1020180033427 2018-03-22 KR		
外部链接	Espacenet USPTO		

摘要(译)

有机电致发光器件包括第一电极，设置在第一电极上的空穴传输区，设置在空穴传输区上的发射层，设置在发射层上的电子传输区以及设置在电子传输区上的第二电极，其中，发射层包括含有两个电子供体和一个电子受体的多环化合物，并且电子受体包括苯膦部分和吡啶部分，从而显示出高发射效率。

