

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 August 2005 (18.08.2005)

PCT

(10) International Publication Number
WO 2005/074361 A2

- (51) International Patent Classification: **Not classified**
- (21) International Application Number:
PCT/IL2005/000113
- (22) International Filing Date: 31 January 2005 (31.01.2005)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/541,779 5 February 2004 (05.02.2004) US
- (71) Applicant (for all designated States except US): **EARLY-SENSE LTD.** [IL/IL]; 29 Kiryati Street, 52223 Ramat Gan (IL).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **LANGE, Daniel, H.** [IL/IL]; 10 Yarden Street, 25147 Kfar Vradim (IL). **GROSS, Yosef** [IL/IL]; 10 Hanotea Street, 73160 Moshav Mazor (IL). **HALPERIN, Avner** [IL/IL]; 29 Kiryati Street, 52223 Ramat Gan (IL).
- (74) Agents: **SANFORD T. COLB & CO.** et al.; P.O. Box 2273, 76122 Rehovot (IL).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 2005/074361 A2

(54) Title: TECHNIQUES FOR PREDICTION AND MONITORING OF RESPIRATION-MANIFESTED CLINICAL EPISODES

(57) Abstract: A method is provided for predicting an onset of a clinical episode, the method including sensing breathing of a subject, determining at least one breathing pattern of the subject responsively to the sensed breathing, comparing the breathing pattern with a baseline breathing pattern, and predicting the onset of the episode at least in part responsively to the comparison. Other embodiments are also described.

TECHNIQUES FOR PREDICTION AND MONITORING OF RESPIRATION-
MANIFESTED CLINICAL EPISODES

CROSS-REFERENCES TO RELATED APPLICATIONS

This application claims the benefit of US Provisional Patent Application
5 60/541,779, filed February 5, 2004, entitled, "Method and apparatus for prediction and
monitoring of respiration manifested clinical episodes," which is assigned to the assignee
of the present application and is incorporated herein by reference.

This application is related to a US regular patent application filed on even date
herewith, entitled, "Techniques for prediction and monitoring of respiration-manifested
10 clinical episodes," which is assigned to the assignee of the present patent application and
is incorporated herein by reference.

FIELD OF THE INVENTION

The present invention relates generally to predicting and monitoring abnormal
physiological conditions, and specifically to methods and apparatus for predicting and
15 monitoring abnormal physiological conditions by measuring and analyzing characteristics
of respiration.

BACKGROUND OF THE INVENTION

Chronic diseases are often expressed by episodic worsening of clinical symptoms.
Preventive treatment of chronic diseases reduces the overall dosage of required
20 medication and associated side effects. Generally, preventive treatment should be
initiated or intensified as soon as the earliest clinical symptoms are detected, in order to
prevent progression and worsening of the clinical episode and to stop and reverse the
pathophysiological process. Therefore, an ability to accurately monitor pre-episodic
indicators increases the effectiveness of preventive treatment of chronic diseases.

25 Many chronic diseases interfere with normal breathing patterns, through a variety
of physiological mechanisms. Common respiratory disorders, such as asthma, chronic
obstructive pulmonary disease (COPD), and cystic fibrosis (CF), are direct modifiers of
breathing patterns. Other chronic diseases, such as diabetes, epilepsy, and certain heart

diseases, are also known to modify breathing activity, because of pathophysiologies leading to abnormal sympathetic and parasympathetic neural activity.

Asthma is a chronic disease with no known cure. Substantial alleviation of asthma symptoms is possible via preventive therapy, such as the use of bronchodilators and anti-inflammatory agents. Asthma management is aimed at improving the quality of life of asthma patients. Asthma management presents a serious challenge to the patient and physician, as preventive therapies require constant monitoring of lung function and corresponding adaptation of medication type and dosage. However, monitoring of lung function is not simple, and requires sophisticated instrumentation and expertise, which are generally not available in the non-clinical or home environment.

Monitoring of lung function is viewed as a major factor in determining an appropriate treatment, as well as in patient follow-up. Preferred therapies are based on aerosol-type medications to minimize systemic side-effects. The efficacy of aerosol type therapy is highly dependent on patient compliance, which is difficult to assess, further contributing to the importance of lung-function monitoring.

Asthma episodes usually develop over a period of several days, although they may sometimes seem to appear unexpectedly. The gradual onset of the asthmatic episode provides an opportunity to start countermeasures to stop and reverse the inflammatory process. Early treatment at the pre-episode stage may reduce the clinical episode manifestation considerably, and may even prevent the transition from the pre-clinical stage to a clinical episode altogether.

Two techniques are generally used for asthma monitoring. The first technique, spirometry, evaluates lung function using a spirometer, an instrument that measures the volume of air inhaled and exhaled by the lungs. Airflow dynamics are measured during a forceful, coordinated inhalation and exhalation effort by the patient into a mouthpiece connected via a tube to the spirometer. A peak-flow meter is a simpler device that is similar to the spirometer, and is used in a similar manner. The second technique evaluates lung function by measuring nitric-oxide concentration using a dedicated nitric-oxide monitor. The patient breathes into a mouthpiece connected via a tube to the monitor.

Efficient asthma management requires daily monitoring of respiratory function, which is generally impractical, particularly in non-clinical or home environments. Peak-flow meters and nitric-oxide monitors provide a general indication of the status of lung

function. However, these monitoring devices do not possess predictive value, and are used as during-episode markers. In addition, peak-flow meters and nitric-oxide monitors require active participation of the patient, which is difficult to obtain from many children and substantially impossible to obtain from infants.

5 Congestive heart failure (CHF) is a condition in which the heart is weakened and unable to circulate blood to meet the body's needs. The subsequent buildup of fluids in the legs, kidneys, and lungs characterizes the condition as congestive. The weakening may be associated with either the left, right, or both sides of the heart, with different etiologies and treatments associated with each type. In most cases, it is the left side of the
10 heart which fails, so that it is unable to efficiently pump blood to the systemic circulation. The ensuing fluid congestion of the lungs results in changes in respiration, including alterations in rate and/or pattern, accompanied by increased difficulty in breathing and tachypnea.

 Quantification of such abnormal breathing provides a basis for assessing CHF
15 progression. For example, Cheyne-Stokes Respiration (CSR) is a breathing pattern characterized by rhythmic oscillation of tidal volume with regularly recurring periods of alternating apnea and hyperpnea. While CSR may be observed in a number of different pathologies (e.g., encephalitis, cerebral circulatory disturbances, and lesions of the bulbar center of respiration), it has also been recognized as an independent risk factor for
20 worsening heart failure and reduced survival in patients with CHF. In CHF, CSR is associated with frequent awakening that fragments sleep, and with concomitant sympathetic activation, both of which may worsen CHF. Other abnormal breathing patterns may involve prolonged expiration or inspiration, or gradual changes in respiration rate usually leading to tachypnea.

25 US Patent 5,853,005 to Scanlon, which is incorporated herein by reference, describes a transducer in communication with fluid in a pad. The pad is held in close contact against a sound or movement source, and monitors acoustic signals transferred into the fluid. The signal pattern is monitored aurally and/or compared to predetermined reference patterns, and optional control and stimulation means can be activated in
30 response to the comparison results. The sensed acoustic signal can be transmitted to a remote receiver or processed locally. Typically, the acoustic signal is representative of the heartbeat or breathing of a living organism. The monitoring system may be applied to

diverse situations including SIDS, apnea, home baby monitoring, medical transport devices, blood pressure cuffs, seats, combat casualty care and hand-held devices. An embodiment is described in which the system is attached to home or institution mattresses for health monitoring, recovery, research, or presence detection.

5 US 6,666,830 to Lehrman et al., which is incorporated herein by reference, describes a system for detecting the onset of an obstructive sleep apnea event before the obstructive sleep apnea event fully develops, and before the cessation of breathing occurs. The system includes one or more microphones capable of detecting breathing sounds within an airway of a person. The microphones generate signals representative of the
10 breathing sounds, and send the signals to a controller. The controller identifies at least one signal pattern that is associated with a breathing pattern of the person that occurs at the onset of an obstructive sleep apnea event. The controller may also identify at least one signal pattern that is associated with a partially-occluded breathing pattern of the person. The controller identifies the signal patterns by using digital signal processing
15 techniques to analyze the signals representative of breathing sounds. The method involves detecting breathing sounds within an airway of a person, generating signals representative of the breathing sounds, and identifying at least one signal pattern that is associated with a breathing pattern of the person that occurs at the onset of an obstructive sleep apnea event.

20 US 6,790,183 to Murphy, which is incorporated herein by reference, describes a lung sound diagnostic system for use in collecting, organizing and analyzing lung sounds associated with the inspiration(s) and expiration(s) of a patient. The system includes a plurality of transducers that may be placed at various sites around the patient's chest. The microphones are coupled to signal processing circuitry and A/D converters which digitize
25 the data and preferably provides the digital data to a computer station. The system may also include application programs for detecting and classifying abnormal sounds. The resulting information may be displayed in a variety of formats to facilitate diagnosis. Additionally, the system may include an analysis program for comparing selected criteria corresponding to the detected abnormal sounds with predefined thresholds in order to
30 provide a likely diagnosis. Also described are a system and method for differentiating between the crackles produced by an patient with interstitial pulmonary fibrosis (IPF) from the crackles produced by a CHF patient.

US Patent 6,168,568 to Gavriely, which is incorporated herein by reference, describes a phonopneumograph system for analyzing breath sounds. The system includes a plurality of breath-related sensors placed around the respiratory system of a patient for measuring breath-related activity, and a breath analyzer. The breath analyzer matches the
5 breath sound data produced by the breath-related sensors to a plurality of breath sound templates, each of which parameterizes one type of breath sound, and determines the presence of regular and/or adventitious breath sounds only when the breath sound data matches, within predetermined goodness of fit criteria, one or more of the breath sound templates.

10 US Patent 6,261,238 to Gavriely, which is incorporated herein by reference, describes a method for analyzing breath sounds produced by a respiratory system. The method includes measuring breath sounds produced by the respiratory system; tentatively identifying a signal as being caused by a breath sound of a given type if it meets a first criterion characteristic of the breath sound of the given type; and confirming the
15 identification if a tentatively identified signal meets a second criterion characteristic of the breath sound of the given type.

US Patent 5,738,102 to Lemelson, which is incorporated herein by reference, describes a system for monitoring and computer analyzing select physiological variables of a patient in real time in order to alert medical personnel to the need for medical
20 treatment or automatically administering such treatment under computer control. Such physiological variables monitored by the system may include lung sounds, respiratory rate and rhythm, heart rate and rhythm, heart sounds, and body temperature. Coded signals relating to the physiological variables are produced and compared with reference versions of same by a decision computer in order to evaluate the patient's condition. If the
25 evaluation indicates medical treatment is needed, the decision computer activates a local and/or a remote alarm to alert medical personnel and/or activates one or more actuators for administering a medical treatment such as the injection or infusion of a drug.

An article by Shochat M et al., entitled, "PedemaTOR: Innovative method for detecting pulmonary edema at the pre-clinical stage," undated, available at http://www.isrmed.info/rsmm_rabinovich/pedemator.htm, which is incorporated herein by
30 reference, describes an impedance monitor for pre-clinical detection of pulmonary edema. The impedance monitor measures "internal thoracic impedance" (ITI), which is roughly

equal to lung impedance, by automatically calculating skin-electrode impedance and subtracting it from the measured transthoracic impedance (TTI).

The following articles, which are incorporated herein by reference, may be of interest:

5 Bentur L et al., "Wheeze monitoring in children for assessment of nocturnal asthma and response to therapy," *Eur Respir J* 21(4):621-626 (2003).

Stegmaier-Stracca PA et al., "Cough detection using fuzzy classification," Symposium on Applied Computing, Proceedings of the 1995 ACM Symposium on Applied Computing, Nashville, Tennessee, United States, pp. 440 - 444 (1995).

10 Waris M et al., "A new method for automatic wheeze detection," *Technol Health Care* 6(1):33-40 (1998).

The inclusion of the foregoing references in this Background section does not imply that they constitute prior art or analogous art with respect to the invention disclosed herein.

15

SUMMARY OF THE INVENTION

In some embodiments of the present invention, a method for monitoring a chronic medical condition comprises non-invasively monitoring at least one breathing pattern of a subject, typically during sleep at night. The pattern is analyzed in order to (a) predict an approaching clinical episode, such as an asthma attack, and/or (b) monitor the severity and progression of a clinical episode as it occurs. Analyzing the pattern typically comprises comparing the pattern to a baseline pattern. Prediction of an approaching clinical episode facilitates early preventive treatment, which generally reduces the required dosage of medication.

In some embodiments of the present invention, the breathing pattern is monitored by continuously acquiring breathing-related body motion data of the subject during sleep. The motion data is processed to yield at least one periodic breathing-related movement pattern, from which the breathing pattern is extracted. For some applications, the motion data is acquired using a sensing device that does not come in contact with the subject or clothes the subject is wearing. For example, the sensing device may be a pressure gauge, which is typically adapted to be installed under a mattress upon which the subject sleeps.

30

Because the data acquisition is non-invasive (and typically not noticeable), it is generally suitable for monitoring both children and adults in a home environment.

The effectiveness of the techniques described herein is in part based on the observation that some chronic medical conditions interfere with normal breathing during sleep and while awake, resulting in condition-specific abnormal breathing patterns. Various direct and indirect physiological mechanisms modify breathing patterns, resulting in specific patterns related to the cause of modification. Respiratory diseases, such as asthma, chronic obstructive pulmonary disease (COPD), and cystic fibrosis (CF), directly modify breathing patterns, while physiological abnormalities associated with some conditions indirectly modify breathing patterns. For example, such indirect breathing pattern-modifying physiological abnormalities include: (a) congestive heart failure (CHF), which sometimes causes abnormal breathing patterns such as Cheyne-Stokes Respiration (CSR), (b) hypoglycemia, such as caused by diabetes, and (c) abnormal autonomic nervous system activity, such as caused by some neurological conditions.

In some embodiments of the present invention, a system for monitoring chronic medical conditions comprises a breathing-related motion acquisition module, a breathing pattern analysis module, and an output module.

There is therefore provided, in accordance with an embodiment of the present invention, a method for predicting an onset of a clinical episode, including:

- sensing breathing of a subject;
- determining at least one breathing pattern of the subject responsively to the sensed breathing;
- comparing the breathing pattern with a baseline breathing pattern; and
- predicting the onset of the episode at least in part responsively to the comparison.

For some applications, the breathing pattern includes a breathing rate pattern of the subject, the baseline breathing pattern includes a baseline breathing rate pattern, and comparing the breathing pattern with the baseline breathing pattern includes comparing the breathing rate pattern with the baseline breathing rate pattern.

For some applications, comparing includes determining the baseline breathing pattern by analyzing breathing of the subject during at least one non-symptomatic period.

For some applications, comparing includes setting the baseline breathing pattern responsively to a population average breathing pattern.

For some applications, predicting the onset includes predicting the onset responsively to a prolonged inspirium time of the subject, and/or to a prolonged expirium time of the subject. For some applications, the breathing pattern includes successive segments of inspirium and expirium, and predicting the onset includes predicting the onset responsively to a trend towards greater durations of at least one of: the inspirium segments and the expirium segments.

In an embodiment, the clinical episode includes an episode associated with a condition selected from the list consisting of: asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), CHF, diabetes, and epilepsy.

In an embodiment, the breathing pattern includes a breathing duty-cycle pattern, and predicting the onset includes predicting the onset responsively to an increase in a breathing duty-cycle of the subject.

For some applications, sensing breathing of the subject includes sensing at least one breathing-related sound selected from the list consisting of: a sound caused by wheezing, and a sound caused by coughing, and predicting the onset includes predicting the onset responsively to an aspect of the breathing-related sound.

For some applications, sensing breathing of the subject includes sensing at least one type of breathing-related mechanical vibrations selected from the list consisting of: mechanical vibrations caused by wheezing, and mechanical vibrations caused by coughing, and predicting the onset includes predicting the onset responsively to an aspect of the breathing-related mechanical vibrations.

In an embodiment, the breathing pattern includes a breathing rate variability pattern, the baseline breathing pattern includes a baseline breathing rate variability pattern, and predicting the onset includes predicting the onset responsively to a decrease in breathing rate variability over time compared to the baseline breathing rate variability pattern. For some applications, determining the at least one breathing pattern includes determining the breathing rate variability pattern and a slow trend breathing rate pattern, comparing the breathing pattern with the baseline breathing pattern includes comparing the breathing rate variability pattern with the baseline breathing rate variability pattern,

and comparing the slow trend breathing rate pattern with a baseline slow trend breathing rate pattern, and predicting the onset includes predicting the onset responsively to both comparisons. For some applications, sensing the breathing includes sensing at least one of: breathing sounds of the subject, and respiratory air-flow of the subject. For some applications, the clinical episode includes an asthma attack, and predicting the onset of the episode includes predicting the onset of the asthma attack.

In an embodiment, the breathing pattern and the baseline breathing pattern include respective slow trend breathing rate patterns, and comparing the breathing pattern with the baseline breathing pattern includes comparing the slow trend breathing rate pattern with the baseline slow trend breathing rate pattern. For some applications, the baseline slow trend breathing rate pattern includes a monotonic decline in breathing rate over at least 1 hour, and predicting the onset includes predicting the onset responsively to a difference between the slow trend breathing rate pattern and the monotonic decline in breathing rate.

In an embodiment, sensing the breathing includes acquiring breathing-related body motion data of the subject. For some applications, acquiring the body motion data includes acquiring the body motion data while the subject is sleeping. For some applications, determining the breathing pattern includes analyzing the body motion data to determine a breathing-related movement pattern, and determining the breathing pattern responsively to the breathing-related movement pattern. For some applications, determining the breathing pattern includes removing non-breathing-related motion data from the body motion data. For example, removing the non-breathing-related motion data from the body motion data may include applying analysis techniques such as frequency-domain spectral analysis or time-domain regression analysis.

In an embodiment, acquiring the body motion data includes acquiring the body motion data without contacting the subject or clothes the subject is wearing. For some applications, the clinical episode includes an asthma attack, and predicting the onset of the episode includes predicting the onset of the asthma attack. For some applications, acquiring the breathing-related body motion data includes measuring a pressure. For some applications, measuring the pressure includes measuring a pressure at a mattress upon which the subject lies. Alternatively or additionally, measuring the pressure includes measuring a pressure under a mattress upon which the subject lies. Further alternatively or additionally, measuring the pressure includes measuring a pressure under

a mattress covering upon which the subject lies, for example, a sheet, a mattress pad, or a mattress cover.

For some applications, the breathing pattern includes a breathing rate variability pattern, the baseline breathing pattern includes a baseline breathing rate variability pattern, and predicting the onset includes predicting the onset responsively to a decrease
5 in breathing rate variability over time compared to the baseline breathing rate variability pattern. For some applications, determining the at least one breathing pattern includes determining the breathing rate variability pattern and a slow trend breathing rate pattern; comparing the breathing pattern with the baseline breathing pattern includes comparing
10 the breathing rate variability pattern with the baseline breathing rate variability pattern, and comparing the slow trend breathing rate pattern with a baseline slow trend breathing rate pattern; and predicting the onset includes predicting the onset responsively to both comparisons.

There is also provided, in accordance with an embodiment of the present
15 invention, a method including:

sensing breathing of a subject during a clinical episode;

determining at least one breathing pattern of the subject responsively to the sensed
breathing;

comparing the breathing pattern with a baseline breathing pattern; and

20 assessing a progression of the episode at least in part responsively to the comparison.

For some applications, the breathing pattern includes a breathing rate pattern of the subject, the baseline breathing pattern includes a baseline breathing rate pattern, and comparing the breathing pattern with the baseline breathing pattern includes comparing
25 the breathing rate pattern with the baseline breathing rate pattern.

For some applications, assessing the progression includes assessing the progression responsively to a prolonged inspirium time of the subject, and/or to a prolonged expirium time of the subject.

For some applications, the breathing pattern includes successive segments of
30 inspirium and expirium, and assessing the progression includes assessing the progression

responsively to a trend towards greater durations of at least one of: the inspirium segments and the expirium segments.

In an embodiment, the clinical episode includes an episode associated with a condition selected from the list consisting of: asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), CHF, diabetes, and epilepsy.

In an embodiment, the breathing pattern includes a breathing duty-cycle pattern, and assessing the progression includes assessing the progression responsively to an increase in a breathing duty-cycle of the subject.

For some applications, sensing breathing of the subject includes sensing at least one breathing-related sound selected from the list consisting of: a sound caused by wheezing, and a sound caused by coughing, and assessing the progression includes assessing the progression responsively to an aspect of the breathing-related sound. For some applications, the clinical episode includes an asthma attack, and assessing the progression includes assessing the progression of the asthma attack responsively to the aspect.

For some applications, sensing breathing of the subject includes sensing at least one type of breathing-related mechanical vibrations selected from the list consisting of: mechanical vibrations caused by wheezing, and mechanical vibrations caused by coughing, and assessing the progression includes assessing the progression responsively to an aspect of the breathing-related mechanical vibrations. For some applications, the clinical episode includes an asthma attack, and assessing the progression includes assessing the progression of the asthma attack responsively to the aspect.

In an embodiment, the breathing pattern includes a breathing rate variability pattern, the baseline breathing pattern includes a baseline breathing rate variability pattern, and assessing the progression includes assessing the progression responsively to a decrease in breathing rate variability over time compared to the baseline breathing rate variability pattern. For some applications, determining the at least one breathing pattern includes determining the breathing rate variability pattern and a slow trend breathing rate pattern; comparing the breathing pattern with the baseline breathing pattern includes comparing the breathing rate variability pattern with the baseline breathing rate variability pattern, and comparing the slow trend breathing rate pattern with a baseline slow trend breathing rate pattern; and assessing the progression includes assessing the progression

responsively to both comparisons. For some applications, the clinical episode includes an asthma attack, and assessing the progression of the episode includes assessing a severity of the asthma attack.

5 In an embodiment, the breathing pattern and the baseline breathing pattern include respective slow trend breathing rate patterns, and comparing the breathing pattern with the baseline breathing pattern includes comparing the slow trend breathing rate pattern with the baseline slow trend breathing rate pattern. For some applications, the baseline slow trend breathing rate pattern includes a monotonic decline in breathing rate over at least 1 hour, and assessing the progression includes assessing the progression responsively to a
10 difference between the slow trend breathing rate pattern and the monotonic decline in breathing rate.

In an embodiment, sensing the breathing includes acquiring breathing-related body motion data of the subject. For some applications, determining the breathing pattern includes analyzing the body motion data to determine a breathing-related movement
15 pattern, and determining the breathing pattern responsively to the breathing-related movement pattern.

In an embodiment, acquiring the body motion data includes acquiring the body motion data without contacting the subject or clothes the subject is wearing. For some applications, the clinical episode includes an asthma attack, and assessing the progression
20 of the episode includes assessing a severity of the asthma attack. For some applications, acquiring the breathing-related body motion data includes measuring a pressure. For some applications, measuring the pressure includes measuring a pressure at a mattress upon which the subject lies. Alternatively or additionally, measuring the pressure includes measuring a pressure under a mattress upon which the subject lies. Further
25 alternatively or additionally, measuring the pressure includes measuring a pressure under a mattress covering upon which the subject lies, for example, a sheet, a mattress pad, or a mattress cover.

For some applications, the breathing pattern includes a breathing rate variability pattern, the baseline breathing pattern includes a baseline breathing rate variability
30 pattern, and assessing the progression includes assessing the progression responsively to a decrease in breathing rate variability over time compared to the baseline breathing rate variability pattern. For some applications, determining the at least one breathing pattern

includes determining the breathing rate variability pattern and a slow trend breathing rate pattern; comparing the breathing pattern with the baseline breathing pattern includes comparing the breathing rate variability pattern with the baseline breathing rate variability pattern, and comparing the slow trend breathing rate pattern with a baseline slow trend breathing rate pattern; and assessing the progression includes assessing the progression
5 responsively to both comparisons.

There is further provided, in accordance with an embodiment of the present invention, a method including:

sensing breathing of a subject;
10 determining at least one breathing pattern of the subject responsively to the sensed breathing;
comparing the breathing pattern with a baseline breathing pattern; and
detecting an abnormal breathing pattern associated with congestive heart failure (CHF), at least in part responsively to the comparison.

15 For some applications, determining the breathing pattern includes determining a breathing rate pattern of the subject, and comparing the breathing pattern with the baseline breathing pattern includes comparing the breathing rate pattern with a baseline breathing rate pattern.

For some applications, detecting the abnormal breathing pattern includes detecting
20 Cheyne-Stokes Respiration (CSR), and/or detecting tachypnea.

In an embodiment, sensing the breathing includes acquiring breathing-related body motion data of the subject. For some applications, acquiring the body motion data includes acquiring the body motion data while the subject is sleeping.

In an embodiment, acquiring the body motion data includes acquiring the body
25 motion data without contacting the subject or clothes the subject is wearing. For some applications, detecting the abnormal breathing pattern includes detecting Cheyne-Stokes Respiration (CSR) and/or tachypnea.

For some applications, acquiring the breathing-related body motion data includes measuring a pressure. For some applications, measuring the pressure includes measuring
30 a pressure at a mattress upon which the subject lies. Alternatively or additionally, measuring the pressure includes measuring a pressure under a mattress upon which the

subject lies. Further alternatively or additionally, measuring the pressure includes measuring a pressure under a mattress covering upon which the subject lies, for example, a sheet, a mattress pad, or a mattress cover.

5 There is further provided, in accordance with an embodiment of the present invention, a method including:

sensing breathing of a subject;

determining at least one breathing pattern of the subject responsively to the sensed breathing;

comparing the breathing pattern with a baseline breathing pattern; and

10 detecting an abnormal breathing pattern associated with a condition of the subject, at least in part responsively to the comparison, the condition selected from the list consisting of: chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), diabetes, and epilepsy.

For some applications, determining the breathing pattern includes determining a
15 breathing rate pattern of the subject, and comparing the breathing pattern with the baseline breathing pattern includes comparing the breathing rate pattern with a baseline breathing rate pattern.

In an embodiment, sensing the breathing includes acquiring breathing-related body motion data of the subject. For some applications, acquiring the body motion data
20 includes acquiring the body motion data while the subject is sleeping.

In an embodiment, acquiring the body motion data includes acquiring the body motion data without contacting the subject or clothes the subject is wearing.

For some applications, acquiring the breathing-related body motion data includes measuring a pressure. For some applications, measuring the pressure includes measuring
25 a pressure at a mattress upon which the subject lies. Alternatively or additionally, measuring the pressure includes measuring a pressure under a mattress upon which the subject lies. Further alternatively or additionally, measuring the pressure includes measuring a pressure under a mattress covering upon which the subject lies, for example, a sheet, a mattress pad, or a mattress cover.

30 There is still further provided, in accordance with an embodiment of the present invention, apparatus for predicting an onset of a clinical episode, including:

a breathing sensor, adapted to sense breathing of a subject, and to generate a signal responsively thereto; and

a control unit, adapted to:

receive the signal,

5 determine at least one breathing pattern of the subject responsive to the signal,

compare the breathing pattern with a baseline breathing pattern, and

predict the onset of the episode at least in part responsively to the comparison.

10 There is additionally provided, in accordance with an embodiment of the present invention, apparatus including:

a breathing sensor, adapted to sense breathing of a subject during a clinical episode, and to generate a signal responsively thereto; and

a control unit, adapted to:

15 receive the signal,

determine at least one breathing pattern of the subject responsive to the signal,

compare the breathing pattern with a baseline breathing pattern, and

assess a progression of the episode at least in part responsively to

20 the comparison.

There is still additionally provided, in accordance with an embodiment of the present invention, apparatus including:

a breathing sensor, adapted to sense breathing of a subject during a clinical episode, and to generate a signal responsively thereto; and

25 a control unit, adapted to:

receive the signal,

determine at least one breathing pattern of the subject responsive to the signal,

compare the breathing pattern with a baseline breathing pattern, and

30 detect an abnormal breathing pattern associated with congestive heart failure (CHF), at least in part responsively to the comparison.

There is additionally provided, in accordance with an embodiment of the present invention, apparatus including:

a breathing sensor, adapted to sense breathing of a subject during a clinical episode, and to generate a signal responsively thereto; and

5 a control unit, adapted to:

receive the signal,

determine at least one breathing pattern of the subject responsive to the signal,

compare the breathing pattern with a baseline breathing pattern, and

10 detect an abnormal breathing pattern associated with a condition of the subject, at least in part responsively to the comparison, the condition selected from the list consisting of: chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), diabetes, and epilepsy.

There is also provided, in accordance with an embodiment of the present invention, a method for clinical episode prediction and assessment, including:

15 measuring breathing rate variability patterns during night sleep;

comparing said breathing rate variability patterns to normal breathing rate variability patterns; and

20 determining a likelihood of a nearing clinical episode or a progression or severity of an ongoing episode.

For some applications, said measuring of breathing rate variability patterns is executed by means of measurement of a composite body movement signal and extraction of a periodic, breathing-related movement signal from said composite body movement signal. Alternatively, said measuring of breathing rate variability patterns is executed by

25 means of measurement of respiration airflow from a mouth and/or a nose. Further alternatively, said measuring of breathing rate variability patterns is executed by means of acoustic measurement of airway and lung sounds from a chest, a back, a neck, and/or a face.

For some applications, said normal breathing rate patterns are extracted from the

30 patient during non-symptomatic periods. For some applications, the normal breathing rate patterns are extracted from averaged patterns of normal, healthy subjects with similar character of age, height, weight, and/or gender.

For some applications, said breathing rate variability patterns include: (1) cyclic patterns, whose typical durations range from several seconds to several minutes, and/or (2) slow trends of segmented, monotonically declining breathing rate usually lasting several hours.

5 For some applications, said comparing is based on a calculation of a degree of deviation of said breathing rate variability patterns from said normal breathing rate variability patterns.

In an embodiment, said clinical episode is a clinical asthma episode.

For some applications, said clinical episode relates to any chronic disease affecting
10 breathing rate patterns, such as diabetes, a heart condition, a neurological disorder, or epilepsy.

There is further provided, in accordance with an embodiment of the present invention, apparatus for clinical episode assessment and prediction, including:

- 15 a breathing sensor which measures breathing;
- an amplifier which amplifies the output signal of the breathing sensor;
- an A/D card which digitizes the amplifier output;
- a processor, which extracts breathing rate patterns and compares said patterns to normal patterns; and
- 20 an output device presenting the result on a numerical, textual or graphical display, or transmitting the results to a clinical follow-up center.

For some applications, the breathing sensor is implemented as a motion-sensitive sensor installed under a bed mattress. Alternatively, the breathing sensor is implemented as an airflow detector aimed at a face of the subject. Further alternatively, the breathing sensor is implemented as an acoustic detector aimed or attached to a face, chest, or back
25 of the subject.

The present invention will be more fully understood from the following detailed description of embodiments thereof, taken together with the drawings, in which:

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a schematic block diagram illustrating a system for monitoring a chronic
30 medical condition, in accordance with an embodiment of the present invention;

Fig. 2 is a schematic block diagram illustrating a data acquisition module of the system of Fig. 1, in accordance with an embodiment of the present invention;

Fig. 3 is a schematic block diagram illustrating a pattern analysis module of the system of Fig. 1, in accordance with an embodiment of the present invention; and

5 Fig. 4 is a graph illustrating breathing rate patterns of a chronic asthma patient, measured during an experiment conducted in accordance with an embodiment of the present invention.

DETAILED DESCRIPTION OF EMBODIMENTS

10 Fig. 1 is a schematic block diagram illustrating a system 10 for monitoring a chronic medical condition, in accordance with an embodiment of the present invention. System 10 typically comprises a breathing-related data acquisition module 20, a breathing pattern analysis module 22, and an output module 24. For some applications, two or more of modules 20, 22, and 24 are packaged in a single housing. For other applications, the modules are packaged separately, such as to enable remote analysis by pattern analysis
15 module 22 of breathing signals acquired locally by breathing-related data acquisition module 20.

In an embodiment of the present invention, data acquisition module 20 is adapted to non-invasively monitor breathing patterns of a subject. Pattern analysis module 22 is adapted to analyze the patterns in order to (a) predict an approaching clinical episode,
20 such as an asthma attack, and/or (b) monitor the severity and progression of a clinical episode as it occurs. Output module 24 is adapted to notify the subject and/or a healthcare worker of the predicted or occurring episode. Prediction of an approaching clinical episode facilitates early preventive treatment, which generally reduces the required dosage of medication. When treating asthma, such a reduced dosage generally minimizes
25 the side-effects associated with high dosages typically required to reverse the inflammatory condition once the episode has begun.

Although system 10 may monitor breathing patterns at any time, for some conditions it is generally most effective to monitor such patterns during sleep at night. When the subject is awake, physical and mental activities unrelated to the monitored
30 condition often affect breathing patterns. Such unrelated activities generally have less influence during most night sleep. For some applications, system 10 monitors and records

patterns throughout all or a large portion of a night. The resulting data set generally encompasses typical long-term respiratory patterns, and facilitates comprehensive analysis. Additionally, such a large data set enables rejection of segments contaminated with movement or other artifacts, while retaining sufficient data for a statistically significant analysis.

Reference is now made to Fig. 2, which is a schematic block diagram illustrating data acquisition module 20, in accordance with an embodiment of the present invention. Data acquisition module 20 typically comprises a breathing sensor 30, and other circuitry as appropriate, such as at least one pre-amplifier 32, at least one band-pass filter 34, and an analog-to-digital (A/D) converter 36.

In an embodiment of the present invention, breathing sensor 30 comprises a pressure gauge, which is typically adapted to be installed in, on, or under a mattress upon which the subject sleeps, and to sense breathing-related motion of the subject. For some applications, breathing sensor 30 may be adapted to be installed under a mattress covering upon which the subject sleeps, such as under a sheet, a mattress pad, or a mattress cover. Pattern analysis module 22 is adapted to extract breathing patterns from the motion data, as described hereinbelow with reference to Fig. 3. Alternatively or additionally, breathing sensor 30 comprises another type of sensor, such as an acoustic or air-flow sensor, attached or directed at the subject's face, neck, chest and/or back.

Fig. 3 is a schematic block diagram illustrating pattern analysis module 22, in accordance with an embodiment of the present invention. Pattern analysis module 22 typically comprises a digital signal processor (DSP) 40, dual port RAM (DPR) 42, EEPROM 44, and an I/O port 46. Pattern analysis module 22 is adapted to extract breathing patterns from the raw data generated by data acquisition module 20, and to perform processing and classification of the breathing patterns. Pattern analysis module 22 analyzes changes in breathing patterns, typically during sleep. Responsively to the analysis, module 22 (a) predicts an approaching clinical episode, and/or (b) monitors episode severity and progression.

As mentioned above, in an embodiment, breathing sensor 30 comprises a pressure gauge adapted to be installed under a mattress, and to sense breathing-related motion of the subject. Motion of the subject during sleep includes regular breathing movements as well as other, unrelated body movements. In general, breathing-related motion is the

dominant contributor to body motion during sleep. Pattern analysis module 22 is adapted to substantially eliminate the portion of the motion signal received from the pressure gauge that represents motion unrelated to breathing. For example, the pattern analysis module may remove segments of the signal contaminated by non-breathing related motion. While breathing-related motion is periodic, non-breathing-related motion is generally random and non-predictable. For some applications, the pattern analysis module eliminates the non-breathing related motion using frequency-domain spectral analysis or time-domain regression analysis. Techniques for applying these analysis techniques will be evident to those skilled in art who have read the present application. For some applications, pattern analysis module 22 uses statistical methods, such as linear prediction or outlier analysis, to remove non-breathing-related motion from the signal. The pattern analysis module typically digitizes the motion data at a sampling rate of at least 10 Hz, although lower frequencies are suitable for some applications.

Pattern analysis module 22 is typically adapted to extract breathing patterns from a train of transient breathing pulses, each pulse including one inhalation-exhalation cycle. Breathing patterns during night sleep generally fall into one of several categories, including:

- relatively fast-changing, random breathing patterns, which occur mainly during REM sleep;
- cyclic breathing rate variability patterns, whose typical duration ranges from several seconds to several minutes;
- slow trends in breathing rates (typically, during normal sleep of a healthy subject, such slow trends include segmented, substantially monotonically declining breathing rates usually lasting several hours; for subjects suffering chronically from certain conditions, such as asthma, the monotonic decline may be less pronounced or absent, as discussed, for example, hereinbelow with reference to Fig. 4);
- interruptions in breathing patterns such as coughing and other sleep disturbances; and
- interruptions in breathing patterns caused by momentary waking.

These breathing patterns are associated with various physiological parameters, such as sleep-stage, anxiety, and body temperature. For example, REM sleep is usually accompanied by randomly variable breathing patterns, while deep sleep stages are usually accompanied by more regular and stable patterns. Abnormally high body temperature
5 may accelerate breathing rate, but usually maintains normal cyclic breathing rate variability patterns. Psychological variables such as anxiety are also modulators of breathing patterns during sleep, yet their effect is normally reduced with sleep progression. Interruptions in breathing patterns such as coughing or that caused by momentary waking may be normal, associated with asthma, or associated with other
10 unrelated pathology, and are assessed in context.

In an embodiment of the present invention, pattern analysis module 22 is configured to predict the onset of an asthma attack, and/or monitor its severity and progression. Module 22 typically analyzes changes in breathing rate and in breathing rate variability patterns in combination to predict the onset of an asthma attack. Although
15 breathing rate typically slightly increases prior to the onset of an attack, this increase alone is not always a specific marker of the onset of an attack. Therefore, in order to more accurately predict the onset of an attack, and monitor the severity and progression of an attack, module 22 typically additionally analyzes changes in breathing rate variability patterns. For some applications, module 22 compares one or more of the following
20 patterns to respective baseline patterns, and interprets a deviation from baseline as indicative of (a) the onset of an attack, and/or (b) the severity of an attack in progress:

- a slow trend breathing rate pattern. Module 22 interprets as indicative of an approaching or progressing attack an increase vs. baseline, for example, for generally healthy subjects, an attenuation of the typical segmented,
25 monotonic decline of breathing rate typically over at least 1 hour, e.g., over at least 2, 3, or 4 hours, or the transformation of this decline into an increasing breathing rate pattern, depending on the severity of the attack;
- a breathing rate variability pattern. Module 22 interprets as indicative of an approaching or progressing attack a decrease in breathing rate
30 variability. Such a decrease generally occurs as the onset of an episode approaches, and intensifies with the progression of shortness of breath during an attack;

- a breathing duty-cycle pattern. Module 22 interprets a substantial increase in the breathing duty-cycle as indicative of an approaching or progressing attack. Breathing duty-cycle patterns include, but are not limited to, inspirium time / total breath cycle time, expirium time / total breath cycle time, and (inspirium + expirium time) / total breath cycle time; and
- interruptions in breathing pattern such as caused by coughs, sleep disturbances, or waking. Module 22 quantifies these events, and determines their relevance to prediction of potential asthma attacks.

Pattern analysis module 22 typically determines baseline patterns by analyzing breathing patterns of the subject during non-symptomatic periods. Alternatively or additionally, module 22 is programmed with baseline patterns based on population averages. For some applications, such population averages are segmented by characteristic traits such as age, height, weight, and gender.

In an embodiment of the present invention, breathing cycles are divided into successive segments of inspirium and expirium. Module 22 interprets as indicative of an approaching or progressing attack a trend towards greater durations of the inspirium and/or expirium segments during sleep (typically night sleep).

In an embodiment of the present invention, breathing sensor 30 further comprises an acoustic sensor for measurement of breathing-related sounds such as those caused by wheezing or coughing. (For some applications, in which breathing sensor 30 comprises a pressure gauge, the acoustic sensor is integrated with the pressure gauge. Alternatively, the acoustic sensor is a separate component.) Pattern analysis module 22 processes such breathing sounds independently, or time-locked to expirium and/or inspirium, e.g., by using spectral averaging to enhance the signal-to-noise ratio of wheezing sounds. For some applications, the level of wheezing and its timing with respect to the timing of inspirium and expirium provides additional information for predicting an upcoming asthma attack and/or monitoring the severity and progression of an attack.

Wheezing and coughing can be attributed to specific parts of the breathing cycle (mainly inspirium and expirium), and thus provide a useful insight regarding the type of upcoming or progressing respiratory distress. In addition, wheezing can be filtered according to the periodicity of the breathing cycle, thus enhancing identification of breathing-related sounds of the obstructed airways. Periodic, breathing-cycle-related

wheezing can provide additional insight regarding the type of upcoming or progressing respiratory distress.

In an embodiment of the present invention, pattern analysis module 22 is configured to detect, typically during night sleep, an abnormal breathing pattern associated with congestive heart failure (CHF), such as tachypnea or Cheyne-Stokes Respiration (CSR). Because treatment of CHF appears to be beneficial, its early detection is important.

Reference is again made to Fig. 1. Output module 24 typically comprises a dedicated display unit, such as an LCD or CRT monitor. Alternatively or additionally, the output module comprises a wireless or wired communication port for relaying the acquired and processed data to a remote site for further analysis or interpretation.

Reference is made to Fig. 4, which is a graph illustrating breathing rate patterns of a chronic asthma patient, measured during an experiment conducted in accordance with an embodiment of the present invention. Breathing of the asthma patient was monitored during sleep on several nights. The patient's breathing rate was averaged for each hour of sleep (excluding periods of rapid eye movement (REM) sleep). During the first approximately two months that the patient was monitored, the patient did not experience any episodes of asthma. A line 100 is representative of a typical slow trend breathing pattern recorded during this non-episodic period, and thus represents a baseline slow trend breathing rate pattern for this patient. It should be noted that, unlike the monotonic decline in breathing rate typically observed in non-asthmatic patients, the baseline breathing rate pattern of the chronically asthmatic patient of the experiment reflects an initial decline in breathing rate during the first few hours of sleep, followed by a gradual increase in breathing rate throughout most of the rest of the night.

Line 102 and 104 were recorded on two successive nights at the conclusion of the approximately two-month period, line 102 on the first of these two nights, and line 104 on the second of these two nights. The patient experienced an episode of asthma during the second of these nights. Lines 102 and 104 thus represent a pre-episodic slow trend breathing rate pattern and an episodic slow trend breathing rate pattern, respectively. As can be seen in the graph, the patient's breathing rate was substantially elevated vs. baseline during all hours of the pre-episodic night, and even further elevated vs. baseline during the episodic night.

Using techniques described herein, the pattern of line 102 is compared with the baseline pattern of line 100, in order to predict that the patient may experience an asthmatic episode. The pattern of line 104 is compared with the baseline pattern of line 100 in order to assess a progression of the asthmatic episode.

5 Although some embodiments described herein relate specifically to asthmatic episodes or CHF, the principles of the present invention may be applied, *mutatis mutandis*, to predicting and monitoring other respiratory and non-respiratory conditions that affect normal breathing patterns, such as chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), diabetes, a neurological disorder (e.g., epilepsy), and certain
10 heart diseases in addition to CHF.

It will be appreciated by persons skilled in the art that the present invention is not limited to what has been particularly shown and described hereinabove. Rather, the scope of the present invention includes both combinations and subcombinations of the various features described hereinabove, as well as variations and modifications thereof that are
15 not in the prior art, which would occur to persons skilled in the art upon reading the foregoing description.

CLAIMS

1. A method for predicting an onset of a clinical episode, comprising:
sensing breathing of a subject;
determining at least one breathing pattern of the subject responsively to the sensed
5 breathing;
comparing the breathing pattern with a baseline breathing pattern; and
predicting the onset of the episode at least in part responsively to the comparison.
2. The method according to claim 1,
wherein the breathing pattern includes a breathing rate pattern of the subject,
10 wherein the baseline breathing pattern includes a baseline breathing rate pattern,
and
wherein comparing the breathing pattern with the baseline breathing pattern
comprises comparing the breathing rate pattern with the baseline breathing rate pattern.
3. The method according to claim 1, wherein comparing comprises determining the
15 baseline breathing pattern by analyzing breathing of the subject during at least one non-
symptomatic period.
4. The method according to claim 1, wherein comparing comprises setting the
baseline breathing pattern responsively to a population average breathing pattern.
5. The method according to claim 1, wherein predicting the onset comprises
20 predicting the onset responsively to a prolonged inspirium time of the subject.
6. The method according to claim 1, wherein predicting the onset comprises
predicting the onset responsively to a prolonged expirium time of the subject.
7. The method according to claim 1, wherein the breathing pattern includes
successive segments of inspirium and expirium, and wherein predicting the onset
25 comprises predicting the onset responsively to a trend towards greater durations of at least
one of: the inspirium segments and the expirium segments.
8. The method according to claim 1, wherein the clinical episode includes an episode
associated with a condition selected from the list consisting of: asthma, chronic
obstructive pulmonary disease (COPD), cystic fibrosis (CF), CHF, diabetes, and epilepsy.

9. The method according to claim 1, wherein the breathing pattern includes a breathing duty-cycle pattern, and wherein predicting the onset comprises predicting the onset responsively to an increase in a breathing duty-cycle of the subject.
10. The method according to claim 1, wherein sensing breathing of the subject
5 comprises sensing at least one breathing-related sound selected from the list consisting of: a sound caused by wheezing, and a sound caused by coughing, and wherein predicting the onset comprises predicting the onset responsively to an aspect of the breathing-related sound.
11. The method according to claim 1, wherein sensing breathing of the subject
10 comprises sensing at least one type of breathing-related mechanical vibrations selected from the list consisting of: mechanical vibrations caused by wheezing, and mechanical vibrations caused by coughing, and wherein predicting the onset comprises predicting the onset responsively to an aspect of the breathing-related mechanical vibrations.
12. The method according to any one of claims 1-11,
15 wherein the breathing pattern includes a breathing rate variability pattern, wherein the baseline breathing pattern includes a baseline breathing rate variability pattern, and wherein predicting the onset comprises predicting the onset responsively to a decrease in breathing rate variability over time compared to the baseline breathing rate
20 variability pattern.
13. The method according to claim 12, wherein determining the at least one breathing pattern comprises determining the breathing rate variability pattern and a slow trend breathing rate pattern, wherein comparing the breathing pattern with the baseline breathing pattern
25 comprises comparing the breathing rate variability pattern with the baseline breathing rate variability pattern, and comparing the slow trend breathing rate pattern with a baseline slow trend breathing rate pattern, and wherein predicting the onset comprises predicting the onset responsively to both comparisons.
- 30 14. The method according to claim 12, wherein sensing the breathing comprises sensing at least one of: breathing sounds of the subject, and respiratory air-flow of the subject.

15. The method according to claim 12, wherein the clinical episode includes an asthma attack, and wherein predicting the onset of the episode comprises predicting the onset of the asthma attack.
16. The method according to any one of claims 1-11, wherein the breathing pattern and the baseline breathing pattern include respective slow trend breathing rate patterns, and wherein comparing the breathing pattern with the baseline breathing pattern comprises comparing the slow trend breathing rate pattern with the baseline slow trend breathing rate pattern.
17. The method according to claim 16, wherein the baseline slow trend breathing rate pattern includes a monotonic decline in breathing rate over at least 1 hour, and wherein predicting the onset comprises predicting the onset responsively to a difference between the slow trend breathing rate pattern and the monotonic decline in breathing rate.
18. The method according to any one of claims 1-11, wherein sensing the breathing comprises acquiring breathing-related body motion data of the subject.
19. The method according to claim 18, wherein acquiring the body motion data comprises acquiring the body motion data while the subject is sleeping.
20. The method according to claim 18, wherein determining the breathing pattern comprises analyzing the body motion data to determine a breathing-related movement pattern, and determining the breathing pattern responsively to the breathing-related movement pattern.
21. The method according to claim 20, wherein determining the breathing pattern comprises removing non-breathing-related motion data from the body motion data.
22. The method according to claim 21, wherein removing the non-breathing-related motion data from the body motion data comprises applying at least one analysis technique selected from the list consisting of: frequency-domain spectral analysis, and time-domain regression analysis.
23. The method according to claim 18, wherein acquiring the body motion data comprises acquiring the body motion data without contacting the subject or clothes the subject is wearing.

24. The method according to claim 23, wherein the clinical episode includes an asthma attack, and wherein predicting the onset of the episode comprises predicting the onset of the asthma attack.
25. The method according to claim 23, wherein acquiring the breathing-related body motion data comprises measuring a pressure.
26. The method according to claim 25, wherein measuring the pressure comprises measuring a pressure at a mattress upon which the subject lies.
27. The method according to claim 25, wherein measuring the pressure comprises measuring a pressure under a mattress upon which the subject lies.
28. The method according to claim 25, wherein measuring the pressure comprises measuring a pressure under a mattress covering upon which the subject lies.
29. The method according to claim 23,
wherein the breathing pattern includes a breathing rate variability pattern,
wherein the baseline breathing pattern includes a baseline breathing rate variability pattern, and
wherein predicting the onset comprises predicting the onset responsively to a decrease in breathing rate variability over time compared to the baseline breathing rate variability pattern.
30. The method according to claim 29,
wherein determining the at least one breathing pattern comprises determining the breathing rate variability pattern and a slow trend breathing rate pattern,
wherein comparing the breathing pattern with the baseline breathing pattern comprises comparing the breathing rate variability pattern with the baseline breathing rate variability pattern, and comparing the slow trend breathing rate pattern with a baseline slow trend breathing rate pattern, and
wherein predicting the onset comprises predicting the onset responsively to both comparisons.
31. A method comprising:
sensing breathing of a subject during a clinical episode;
determining at least one breathing pattern of the subject responsively to the sensed breathing;

comparing the breathing pattern with a baseline breathing pattern; and
assessing a progression of the episode at least in part responsively to the
comparison.

32. The method according to claim 31,
5 wherein the breathing pattern includes a breathing rate pattern of the subject,
wherein the baseline breathing pattern includes a baseline breathing rate pattern,
and
wherein comparing the breathing pattern with the baseline breathing pattern
comprises comparing the breathing rate pattern with the baseline breathing rate pattern.
- 10 33. The method according to claim 31, wherein assessing the progression comprises
assessing the progression responsively to a prolonged inspirium time of the subject.
34. The method according to claim 31, wherein assessing the progression comprises
assessing the progression responsively to a prolonged expirium time of the subject.
35. The method according to claim 31, wherein the breathing pattern includes
15 successive segments of inspirium and expirium, and wherein assessing the progression
comprises assessing the progression responsively to a trend towards greater durations of at
least one of: the inspirium segments and the expirium segments.
36. The method according to claim 31, wherein the clinical episode includes an
episode associated with a condition selected from the list consisting of: asthma, chronic
20 obstructive pulmonary disease (COPD), cystic fibrosis (CF), CHF, diabetes, and epilepsy.
37. The method according to claim 31, wherein the breathing pattern includes a
breathing duty-cycle pattern, and wherein assessing the progression comprises assessing
the progression responsively to an increase in a breathing duty-cycle of the subject.
38. The method according to any one of claims 31-37, wherein sensing breathing of
25 the subject comprises sensing at least one type of breathing-related mechanical vibrations
selected from the list consisting of: a sound caused by wheezing, a sound caused by
coughing, mechanical vibrations caused by wheezing, and mechanical vibrations caused
by coughing, and wherein assessing the progression comprises assessing the progression
responsively to an aspect of the breathing-related mechanical vibrations.

39. The method according to claim 38, wherein the clinical episode includes an asthma attack, and wherein assessing the progression comprises assessing the progression of the asthma attack responsively to the aspect.
40. The method according to any one of claims 31-37,
5 wherein the breathing pattern includes a breathing rate variability pattern,
wherein the baseline breathing pattern includes a baseline breathing rate variability pattern, and
wherein assessing the progression comprises assessing the progression responsively to a decrease in breathing rate variability over time compared to the baseline
10 breathing rate variability pattern.
41. The method according to claim 40,
wherein determining the at least one breathing pattern comprises determining the breathing rate variability pattern and a slow trend breathing rate pattern,
wherein comparing the breathing pattern with the baseline breathing pattern
15 comprises comparing the breathing rate variability pattern with the baseline breathing rate variability pattern, and comparing the slow trend breathing rate pattern with a baseline slow trend breathing rate pattern, and
wherein assessing the progression comprises assessing the progression responsively to both comparisons.
- 20 42. The method according to claim 40, wherein the clinical episode includes an asthma attack, and wherein assessing the progression of the episode comprises assessing a severity of the asthma attack.
43. The method according to any one of claims 31-37, wherein the breathing pattern and the baseline breathing pattern include respective slow trend breathing rate patterns,
25 and wherein comparing the breathing pattern with the baseline breathing pattern comprises comparing the slow trend breathing rate pattern with the baseline slow trend breathing rate pattern.
44. The method according to claim 43, wherein the baseline slow trend breathing rate pattern includes a monotonic decline in breathing rate over at least 1 hour, and wherein
30 assessing the progression comprises assessing the progression responsively to a difference between the slow trend breathing rate pattern and the monotonic decline in breathing rate.

45. The method according to any one of claims 31-37, wherein sensing the breathing comprises acquiring breathing-related body motion data of the subject.
46. The method according to claim 45, wherein determining the breathing pattern comprises analyzing the body motion data to determine a breathing-related movement pattern, and determining the breathing pattern responsively to the breathing-related movement pattern.
47. The method according to claim 45, wherein acquiring the body motion data comprises acquiring the body motion data without contacting the subject or clothes the subject is wearing.
48. The method according to claim 47, wherein the clinical episode includes an asthma attack, and wherein assessing the progression of the episode comprises assessing a severity of the asthma attack.
49. The method according to claim 47, wherein acquiring the breathing-related body motion data comprises measuring a pressure.
50. The method according to claim 49, wherein measuring the pressure comprises measuring a pressure at a mattress upon which the subject lies.
51. The method according to claim 49, wherein measuring the pressure comprises measuring a pressure under a mattress upon which the subject lies.
52. The method according to claim 49, wherein measuring the pressure comprises measuring a pressure under a mattress covering upon which the subject lies.
53. The method according to claim 47,
wherein the breathing pattern includes a breathing rate variability pattern,
wherein the baseline breathing pattern includes a baseline breathing rate variability pattern, and
wherein assessing the progression comprises assessing the progression responsively to a decrease in breathing rate variability over time compared to the baseline breathing rate variability pattern.
54. The method according to claim 53,
wherein determining the at least one breathing pattern comprises determining the breathing rate variability pattern and a slow trend breathing rate pattern,

wherein comparing the breathing pattern with the baseline breathing pattern comprises comparing the breathing rate variability pattern with the baseline breathing rate variability pattern, and comparing the slow trend breathing rate pattern with a baseline slow trend breathing rate pattern, and

5 wherein assessing the progression comprises assessing the progression responsively to both comparisons.

55. A method comprising:

sensing breathing of a subject;

10 determining at least one breathing pattern of the subject responsively to the sensed breathing;

comparing the breathing pattern with a baseline breathing pattern; and

detecting an abnormal breathing pattern associated with congestive heart failure (CHF), at least in part responsively to the comparison.

56. The method according to claim 55, wherein determining the breathing pattern
15 comprises determining a breathing rate pattern of the subject, and wherein comparing the breathing pattern with the baseline breathing pattern comprises comparing the breathing rate pattern with a baseline breathing rate pattern.

57. The method according to claim 55, wherein detecting the abnormal breathing pattern comprises detecting Cheyne-Stokes Respiration (CSR).

20 58. The method according to claim 55, wherein detecting the abnormal breathing pattern comprises detecting tachypnea.

59. The method according to any one of claims 55-58, wherein sensing the breathing comprises acquiring breathing-related body motion data of the subject.

60. The method according to claim 59, wherein acquiring the body motion data
25 comprises acquiring the body motion data while the subject is sleeping.

61. The method according to claim 59, wherein acquiring the body motion data comprises acquiring the body motion data without contacting the subject or clothes the subject is wearing.

62. The method according to claim 61, wherein detecting the abnormal breathing
30 pattern comprises detecting a pattern selected from: Cheyne-Stokes Respiration (CSR), and tachypnea.

63. The method according to claim 61, wherein acquiring the breathing-related body motion data comprises measuring a pressure.
64. The method according to claim 63, wherein measuring the pressure comprises measuring a pressure at a mattress upon which the subject lies.
- 5 65. The method according to claim 63, wherein measuring the pressure comprises measuring a pressure under a mattress upon which the subject lies.
66. The method according to claim 63, wherein measuring the pressure comprises measuring a pressure under a mattress covering upon which the subject lies.
67. A method comprising:
- 10 sensing breathing of a subject;
determining at least one breathing pattern of the subject responsively to the sensed breathing;
comparing the breathing pattern with a baseline breathing pattern; and
detecting an abnormal breathing pattern associated with a condition of the subject,
- 15 at least in part responsively to the comparison, the condition selected from the list consisting of: chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), diabetes, and epilepsy.
68. The method according to claim 67, wherein determining the breathing pattern comprises determining a breathing rate pattern of the subject, and wherein comparing the
- 20 breathing pattern with the baseline breathing pattern comprises comparing the breathing rate pattern with a baseline breathing rate pattern.
69. The method according to any one of claims 67-68, wherein sensing the breathing comprises acquiring breathing-related body motion data of the subject.
70. The method according to claim 69, wherein acquiring the body motion data
- 25 comprises acquiring the body motion data while the subject is sleeping.
71. The method according to claim 69, wherein acquiring the body motion data comprises acquiring the body motion data without contacting the subject or clothes the subject is wearing.
72. The method according to claim 71, wherein acquiring the breathing-related body
- 30 motion data comprises measuring a pressure.

73. The method according to claim 72, wherein measuring the pressure comprises measuring a pressure at a mattress upon which the subject lies.
74. The method according to claim 72, wherein measuring the pressure comprises measuring a pressure under a mattress upon which the subject lies.
- 5 75. The method according to claim 72, wherein measuring the pressure comprises measuring a pressure under a mattress covering upon which the subject lies.
76. Apparatus for predicting an onset of a clinical episode, comprising:
a breathing sensor, adapted to sense breathing of a subject, and to generate a signal
responsively thereto; and
10 a control unit, adapted to:
receive the signal,
determine at least one breathing pattern of the subject responsive to
the signal,
compare the breathing pattern with a baseline breathing pattern, and
15 predict the onset of the episode at least in part responsively to the
comparison.
77. The apparatus according to claim 76,
wherein the breathing pattern includes a breathing rate pattern of the subject,
wherein the baseline breathing pattern includes a baseline breathing rate pattern,
20 and
wherein the control unit is adapted to compare the breathing rate pattern with the
baseline breathing rate pattern.
78. The apparatus according to claim 76, wherein the control unit is adapted to
determine the baseline breathing pattern by analyzing breathing of the subject during at
25 least one non-symptomatic period.
79. The apparatus according to claim 76, wherein the control unit is adapted to set the
baseline breathing pattern responsively to a population average breathing pattern.
80. The apparatus according to claim 76, wherein the control unit is adapted to predict
the onset responsively to a prolonged inspirium time of the subject.
- 30 81. The apparatus according to claim 76, wherein the control unit is adapted to predict
the onset responsively to a prolonged expirium time of the subject.

82. The apparatus according to claim 76, wherein the breathing pattern includes successive segments of inspirium and expirium, and wherein the control unit is adapted to predict the onset responsively to a trend towards greater durations of at least one of: the inspirium segments and the expirium segments.
- 5 83. The apparatus according to claim 76, wherein the clinical episode includes an episode associated with a condition selected from the list consisting of: asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), CHF, diabetes, and epilepsy.
84. The apparatus according to claim 76, wherein the breathing pattern includes a breathing duty-cycle pattern, and wherein the control unit is adapted to predict the onset
10 responsively to an increase in a breathing duty-cycle of the subject.
85. The apparatus according to claim 76, wherein the breathing sensor comprises an acoustic sensor, adapted to sense at least one type of breathing-related mechanical vibrations selected from the list consisting of: a sound caused by wheezing, a sound caused by coughing, mechanical vibrations caused by wheezing, and mechanical
15 vibrations caused by coughing, and wherein the control unit is adapted to predict the onset responsively to an aspect of the breathing-related mechanical vibrations.
86. The apparatus according to any one of claims 76-85,
wherein the breathing pattern includes a breathing rate variability pattern,
wherein the baseline breathing pattern includes a baseline breathing rate variability
20 pattern, and
wherein the control unit is adapted to predict the onset responsively to a decrease in breathing rate variability over time compared to the baseline breathing rate variability pattern.
87. The apparatus according to claim 86, wherein the control unit is adapted to:
25 determine the breathing rate variability pattern and a slow trend breathing rate pattern,
compare the breathing rate variability pattern with the baseline breathing rate variability pattern,
compare the slow trend breathing rate pattern with a baseline slow trend breathing
30 rate pattern, and
predict the onset responsively to both comparisons.

88. The apparatus according to claim 86, wherein the breathing sensor is adapted to sense the breathing by sensing at least one of: breathing sounds of the subject, and respiratory air-flow of the subject.
89. The apparatus according to claim 86, wherein the clinical episode includes an asthma attack, and wherein the control unit is adapted to predict the onset of the asthma attack.
90. The apparatus according to any one of claims 76-85, wherein the breathing pattern and the baseline breathing pattern include respective slow trend breathing rate patterns, and wherein the control unit is adapted to compare the slow trend breathing rate pattern with the baseline slow trend breathing rate pattern.
91. The apparatus according to claim 90, wherein the baseline slow trend breathing rate pattern includes a monotonic decline in breathing rate over at least 1 hour, and wherein the control unit is adapted to predict the onset responsively to a difference between the slow trend breathing rate pattern and the monotonic decline in breathing rate.
92. The apparatus according to any one of claims 76-85, wherein the breathing sensor comprises a motion sensor, adapted to sense breathing-related body motion of the subject indicative of the breathing.
93. The apparatus according to claim 92, wherein the control unit is adapted to determine the breathing pattern by:
- analyzing the body motion data to determine a breathing-related movement pattern, and
 - determining the breathing pattern responsively to the breathing-related movement pattern.
94. The apparatus according to claim 93, wherein the control unit is adapted to determine the breathing pattern by removing non-breathing-related motion data from the body motion data.
95. The apparatus according to claim 94, wherein the control unit is adapted to remove the non-breathing-related motion data from the body motion data by applying at least one analysis technique selected from the list consisting of: frequency-domain spectral analysis, and time-domain regression analysis.

96. The apparatus according to claim 92, wherein the breathing sensor is adapted to acquire the body motion data without contacting the subject or clothes the subject is wearing.
97. The apparatus according to claim 96, wherein the clinical episode includes an asthma attack, and wherein the control unit is adapted to predict the onset of the asthma attack.
98. The apparatus according to claim 96, wherein the breathing sensor comprises a pressure gauge.
99. The apparatus according to claim 98, wherein the pressure gauge is configured to measure a pressure at a mattress upon which the subject lies.
100. The apparatus according to claim 98, comprising a mattress, adapted to support the subject lying thereon, wherein the pressure gauge is configured to measure a pressure at the mattress.
101. The apparatus according to claim 98, wherein the pressure gauge is configured to measure a pressure under a mattress upon which the subject lies.
102. The apparatus according to claim 98, comprising a mattress, adapted to support the subject lying thereon, wherein the pressure gauge is configured to measure a pressure under the mattress.
103. The apparatus according to claim 98, wherein the pressure gauge is configured to measure a pressure under a mattress covering upon which the subject lies.
104. The apparatus according to claim 98, comprising a mattress covering, adapted for the subject to lie thereon, wherein the pressure gauge is configured to measure a pressure under the mattress covering.
105. The apparatus according to claim 96,
wherein the breathing pattern includes a breathing rate variability pattern,
wherein the baseline breathing pattern includes a baseline breathing rate variability pattern, and
wherein the control unit is adapted to predict the onset responsively to a decrease in breathing rate variability over time compared to the baseline breathing rate variability pattern.

106. The apparatus according to claim 105, wherein the control unit is adapted to:
determine the breathing rate variability pattern and a slow trend breathing rate
pattern,
compare the breathing rate variability pattern with the baseline breathing rate
5 variability pattern,
compare the slow trend breathing rate pattern with a baseline slow trend breathing
rate pattern, and
predict the onset responsively to both comparisons.
107. Apparatus comprising:
10 a breathing sensor, adapted to sense breathing of a subject during a clinical
episode, and to generate a signal responsively thereto; and
a control unit, adapted to:
receive the signal,
determine at least one breathing pattern of the subject responsive to
15 the signal,
compare the breathing pattern with a baseline breathing pattern, and
assess a progression of the episode at least in part responsively to
the comparison.
108. The apparatus according to claim 107,
20 wherein the breathing pattern includes a breathing rate pattern of the subject,
wherein the baseline breathing pattern includes a baseline breathing rate pattern,
and
wherein the control unit is adapted to compare the breathing rate pattern with the
baseline breathing rate pattern.
- 25 109. The apparatus according to claim 107, wherein the control unit is adapted to assess
the progression responsively to a prolonged inspirium time of the subject.
110. The apparatus according to claim 107, wherein the control unit is adapted to assess
the progression responsively to a prolonged expirium time of the subject.
111. The apparatus according to claim 107, wherein the breathing pattern includes
30 successive segments of inspirium and expirium, and wherein the control unit is adapted to

assess the progression responsively to a trend towards greater durations of at least one of: the inspirium segments and the expirium segments.

112. The apparatus according to claim 107, wherein the clinical episode includes an episode associated with a condition selected from the list consisting of: asthma, chronic
5 obstructive pulmonary disease (COPD), cystic fibrosis (CF), CHF, diabetes, and epilepsy.

113. The apparatus according to claim 107, wherein the breathing pattern includes a breathing duty-cycle pattern, and wherein the control unit is adapted to assess the progression responsively to an increase in a breathing duty-cycle of the subject.

114. The apparatus according to any one of claims 107-113, wherein the breathing
10 sensor comprises an acoustic sensor, adapted to sense at least one type of breathing-related mechanical vibrations selected from the list consisting of: a sound caused by wheezing, a sound caused by coughing, mechanical vibrations caused by wheezing, and mechanical vibrations caused by coughing, and wherein the control unit is adapted to assess the progression responsively to an aspect of the breathing-related mechanical
15 vibrations.

115. The apparatus according to claim 114, wherein the clinical episode includes an asthma attack, and wherein the control unit is adapted to assess the progression of the asthma attack responsively to the aspect.

116. The apparatus according to any one of claims 107-113,
20 wherein the breathing pattern includes a breathing rate variability pattern,
wherein the baseline breathing pattern includes a baseline breathing rate variability pattern, and

wherein the control unit is adapted to assess the progression responsively to a decrease in breathing rate variability over time compared to the baseline breathing rate
25 variability pattern.

117. The apparatus according to claim 116, the control unit is adapted to:
determine the breathing rate variability pattern and a slow trend breathing rate
pattern,
compare the breathing rate variability pattern with the baseline breathing rate
30 variability pattern,

compare the slow trend breathing rate pattern with a baseline slow trend breathing rate pattern, and

assess the progression responsively to both comparisons.

118. The apparatus according to claim 116, wherein the clinical episode includes an asthma attack, and wherein the control unit is adapted to assess a severity of the asthma attack.

119. The apparatus according to any one of claims 107-113, wherein the breathing pattern and the baseline breathing pattern include respective slow trend breathing rate patterns, and wherein the control unit is adapted to compare the slow trend breathing rate pattern with the baseline slow trend breathing rate pattern.

120. The apparatus according to claim 119, wherein the baseline slow trend breathing rate pattern includes a monotonic decline in breathing rate over at least 1 hour, and wherein the control unit is adapted to assess the progression responsively to a difference between the slow trend breathing rate pattern and the monotonic decline in breathing rate.

121. The apparatus according to any one of claims 107-113, wherein the breathing sensor comprises a motion sensor, adapted to sense breathing-related body motion of the subject indicative of the breathing.

122. The apparatus according to claim 121, wherein the control unit is adapted to determine the breathing pattern by:

analyzing the body motion data to determine a breathing-related movement pattern, and

determining the breathing pattern responsively to the breathing-related movement pattern.

123. The apparatus according to claim 121, wherein the breathing sensor is adapted to acquire the body motion data without contacting the subject or clothes the subject is wearing.

124. The apparatus according to claim 123, wherein the clinical episode includes an asthma attack, and wherein the control unit is adapted to the control unit is adapted to assess a severity of the asthma attack.

125. The apparatus according to claim 123, wherein the breathing sensor comprises a pressure gauge.

126. The apparatus according to claim 125, wherein the pressure gauge is configured to measure a pressure at a mattress upon which the subject lies.
127. The apparatus according to claim 125, comprising a mattress, adapted to support the subject lying thereon, wherein the pressure gauge is configured to measure a pressure at the mattress.
128. The apparatus according to claim 125, wherein the pressure gauge is configured to measure a pressure under a mattress upon which the subject lies.
129. The apparatus according to claim 125, comprising a mattress, adapted to support the subject lying thereon, wherein the pressure gauge is configured to measure a pressure under the mattress.
130. The apparatus according to claim 125, wherein the pressure gauge is configured to measure a pressure under a mattress covering upon which the subject lies.
131. The apparatus according to claim 125, comprising a mattress covering, adapted for the subject to lie thereon, wherein the pressure gauge is configured to measure a pressure under the mattress covering.
132. The apparatus according to claim 123,
wherein the breathing pattern includes a breathing rate variability pattern,
wherein the baseline breathing pattern includes a baseline breathing rate variability pattern, and
wherein the control unit is adapted to assess the progression responsively to a decrease in breathing rate variability over time compared to the baseline breathing rate variability pattern.
133. The apparatus according to claim 132, the control unit is adapted to:
determine the breathing rate variability pattern and a slow trend breathing rate pattern,
compare the breathing rate variability pattern with the baseline breathing rate variability pattern,
compare the slow trend breathing rate pattern with a baseline slow trend breathing rate pattern, and
assess the progression responsively to both comparisons.
134. Apparatus comprising:

a breathing sensor, adapted to sense breathing of a subject during a clinical episode, and to generate a signal responsively thereto; and

a control unit, adapted to:

receive the signal,

5 determine at least one breathing pattern of the subject responsive to the signal,

compare the breathing pattern with a baseline breathing pattern, and

detect an abnormal breathing pattern associated with congestive

heart failure (CHF), at least in part responsively to the comparison.

10 135. The apparatus according to claim 134, wherein the breathing pattern includes a breathing rate pattern of the subject, wherein the baseline breathing pattern includes a baseline breathing rate pattern, and wherein the control unit is adapted to compare the breathing rate pattern with the baseline breathing rate pattern.

136. The apparatus according to claim 134, wherein the abnormal breathing pattern
15 includes Cheyne-Stokes Respiration (CSR), and wherein the control unit is adapted to detect the CSR.

137. The apparatus according to claim 134, wherein the abnormal breathing pattern includes tachypnea, and wherein the control unit is adapted to detect the tachypnea.

138. The apparatus according to any one of claims 134-137, wherein the breathing
20 sensor comprises a motion sensor, adapted to sense breathing-related body motion of the subject indicative of the breathing.

139. The apparatus according to claim 138, wherein the breathing sensor is adapted to acquire the body motion data without contacting the subject or clothes the subject is wearing.

25 140. The apparatus according to claim 139, wherein the abnormal breathing pattern includes Cheyne-Stokes Respiration (CSR), and wherein the control unit is adapted to detect the CSR.

141. The apparatus according to claim 139, wherein the breathing sensor comprises a pressure gauge.

30 142. The apparatus according to claim 141, wherein the pressure gauge is configured to measure a pressure at a mattress upon which the subject lies.

143. The apparatus according to claim 141, comprising a mattress, adapted to support the subject lying thereon, wherein the pressure gauge is configured to measure a pressure at the mattress.

144. The apparatus according to claim 141, wherein the pressure gauge is configured to
5 measure a pressure under a mattress upon which the subject lies.

145. The apparatus according to claim 141, comprising a mattress, adapted to support the subject lying thereon, wherein the pressure gauge is configured to measure a pressure under the mattress.

146. The apparatus according to claim 141, wherein the pressure gauge is configured to
10 measure a pressure under a mattress covering upon which the subject lies.

147. The apparatus according to claim 141, comprising a mattress covering, adapted for the subject to lie thereon, wherein the pressure gauge is configured to measure a pressure under the mattress covering.

148. Apparatus comprising:
15 a breathing sensor, adapted to sense breathing of a subject during a clinical episode, and to generate a signal responsively thereto; and
a control unit, adapted to:
receive the signal,
determine at least one breathing pattern of the subject responsive to
20 the signal,
compare the breathing pattern with a baseline breathing pattern, and
detect an abnormal breathing pattern associated with a condition of
the subject, at least in part responsively to the comparison, the condition
selected from the list consisting of: chronic obstructive pulmonary disease
25 (COPD), cystic fibrosis (CF), diabetes, and epilepsy.

149. The apparatus according to claim 148, wherein the breathing pattern includes a breathing rate pattern of the subject, wherein the baseline breathing pattern includes a baseline breathing rate pattern, and wherein the control unit is adapted to compare the breathing rate pattern with the baseline breathing rate pattern.

150. The apparatus according to any one of claims 148-149, wherein the breathing sensor comprises a motion sensor, adapted to sense breathing-related body motion of the subject indicative of the breathing.
- 5 151. The apparatus according to claim 150, wherein the breathing sensor is adapted to acquire the body motion data without contacting the subject or clothes the subject is wearing.
152. The apparatus according to claim 151, wherein the breathing sensor comprises a pressure gauge.
- 10 153. The apparatus according to claim 152, wherein the pressure gauge is configured to measure a pressure at a mattress upon which the subject lies.
154. The apparatus according to claim 152, comprising a mattress, adapted to support the subject lying thereon, wherein the pressure gauge is configured to measure a pressure at the mattress.
- 15 155. The apparatus according to claim 152, wherein the pressure gauge is configured to measure a pressure under a mattress upon which the subject lies.
156. The apparatus according to claim 152, comprising a mattress, adapted to support the subject lying thereon, wherein the pressure gauge is configured to measure a pressure under the mattress.
- 20 157. The apparatus according to claim 152, wherein the pressure gauge is configured to measure a pressure under a mattress covering upon which the subject lies.
158. The apparatus according to claim 152, comprising a mattress covering, adapted for the subject to lie thereon, wherein the pressure gauge is configured to measure a pressure under the mattress covering.

FIG. 1




FIG. 2




FIG. 3





FIG. 4



专利名称(译)	用于预测和监测呼吸表现的临床事件的技术		
公开(公告)号	EP1786315A4	公开(公告)日	2010-03-03
申请号	EP2005703156	申请日	2005-01-31
[标]申请(专利权)人(译)	EARLYSENS		
申请(专利权)人(译)	EARLYSENS有限公司		
当前申请(专利权)人(译)	EARLYSENS有限公司		
[标]发明人	LANGE DANIEL H GROSS YOSEF HALPERIN AVNER		
发明人	LANGE, DANIEL, H. GROSS, YOSEF HALPERIN, AVNER		
IPC分类号	A61B5/00 A61B5/08 A61B5/113 A61B10/00 G06F17/00		
CPC分类号	A61B5/0823 A61B5/113 A61B5/4094 A61B5/7264 A61B5/7275 G16H50/20		
优先权	60/541779 2004-02-05 US		
其他公开文献	EP1786315A2		
外部链接	Espacenet		

摘要(译)

提供了一种用于预测临床发作的开始的方法，该方法包括感测对象的呼吸，响应于感测到的呼吸确定对象的至少一个呼吸模式，将呼吸模式与基线呼吸模式进行比较，以及预测该发作的开始至少部分地响应于比较。还描述了其他实施例。