



US 20100100105A1

(19) **United States**(12) **Patent Application Publication**
Bates et al.(10) **Pub. No.: US 2010/0100105 A1**(43) **Pub. Date: Apr. 22, 2010**(54) **APPARATUS AND METHOD FOR TREATING
OCCLUDED INFECTION COLLECTIONS OF
THE DIGESTIVE TRACT**(76) Inventors: **Douglas Bates**, La Jolla, CA (US);
Steven F. Bierman, Del Mar, CA
(US)


Correspondence Address:

KNOBBE MARTENS OLSON & BEAR LLP
2040 MAIN STREET, FOURTEENTH FLOOR
IRVINE, CA 92614 (US)(21) Appl. No.: **12/603,469**(22) Filed: **Oct. 21, 2009****Related U.S. Application Data**(60) Provisional application No. 61/107,261, filed on Oct.
21, 2008.**Publication Classification**(51) **Int. Cl.****A61B 17/22** (2006.01)**A61F 2/04** (2006.01)**A61F 2/82** (2006.01)(52) **U.S. Cl. 606/114; 623/23.7**

(57)

ABSTRACT

A device and method for treating an occluded infection collection along the digestive tract with a stent delivered to an appendix, diverticulum or other anatomical structure of the digestive tract. A device and method to remove an obstruction from an occluded infection collection along the digestive tract with an obstruction removal device delivered to an appendix, diverticulum or other anatomical structure of the digestive tract.








FIG. 5




FIG. 6




FIG. 7




FIG. 8







FIG. 9






FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27

APPARATUS AND METHOD FOR TREATING OCCLUDED INFECTION COLLECTIONS OF THE DIGESTIVE TRACT

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 61/107,261, filed Oct. 21, 2008, which is hereby incorporated by reference in its entirety.

BACKGROUND

[0002] 1. Field of the Invention

[0003] The invention relates generally to the field of treatment of occluded infection collections, abscesses, inflamed and/or diseased projections in pouch-like portions of the digestive tract, such as an appendix or a diverticular abscess or diverticulum, and more specifically to devices and methods that provide an opening to drain inflamed anatomical structures in the digestive tract without resorting to open surgery or to removal of the anatomical structures.

[0004] 2. Description of the Related Art

[0005] The appendix is a small, tube-like structure attached to the first part of the large intestine, also called the colon. The appendix is typically located in the lower right portion of the abdomen. Removal of the appendix appears to cause no change in digestive function. Appendicitis (or epityphlitis) is an inflammation of the appendix. The cause of appendicitis relates to blockage of the inside of the appendix, known as the lumen. The blockage leads to increased pressure, impaired blood flow, and inflammation. If the blockage is not treated, gangrene and rupture (breaking or tearing) of the appendix can result. Most commonly, feces or calcification blocks the inside of the appendix. Also, bacterial or viral infections in the digestive tract can lead to swelling and inflammation, which squeeze the appendix and cause obstruction to the more distal appendix. This swelling of lymph nodes is known as lymphoid hyperplasia. Traumatic injury to the abdomen may lead to appendicitis in a small number of people. Genetics may be a factor in others. For example, appendicitis that runs in families may result from a genetic variant that predisposes a person to obstruction of the appendiceal lumen.

[0006] In the past, appendicitis has lacked an effective medical therapy, so appendicitis is often considered a medical emergency. While some try to treat it with antibiotics, drug-only therapy techniques have had somewhat limited success. Prior treatment of appendicitis required the removal of the appendix by open surgery, laparotomy or laparoscopy. An appendectomy (or appendectomy) is normally performed as an emergency procedure, when the patient is suffering from acute appendicitis. When treated promptly, most patients recover without difficulty. If treatment is delayed, the appendix can burst, causing infection and even death. Appendicitis is the most common acute surgical emergency of the abdomen.

[0007] X rays, ultrasound, and computed tomography (CT) scans can produce images of the abdomen. Plain x rays can show signs of obstruction, perforation (a hole), foreign bodies, and in rare cases, an appendicolith, which is a calcification in the appendix. Ultrasound may show appendiceal inflammation or fluid and can diagnose gall bladder disease and pregnancy. By far the most common test used, however, is the CT scan. This test provides a series of cross-sectional images of the body and can identify many abdominal condi-

tions and facilitate diagnosis when the clinical impression is in doubt. In selected cases, MRI may be used and particularly in women when the cause of the symptoms may be either the appendix or an inflamed ovary or fallopian tube, laparoscopy may be necessary. Laparoscopy avoids radiation, but requires general anesthesia.

[0008] The most serious complication of appendicitis is rupture. The appendiceal wall bursts or tears if appendicitis is not diagnosed quickly and goes untreated. Infants, young children, and older adults are at highest risk. A ruptured appendix can lead to peritonitis and abscess. Peritonitis is a dangerous infection that happens when bacteria and other contents of the torn appendix leak into the abdomen. In people with appendicitis, an abscess usually takes the form of a swollen mass filled with fluid and bacteria. In a few patients, complications of appendicitis can lead to organ failure and death.

SUMMARY

[0009] In view of the aforementioned shortcomings of traditional treatments and recent purported actions of the appendix, a need exists for apparatus, systems, and methods that can treat appendicitis or other inflammation of structures of the digestive tract less invasively than appendectomies or surgical removal and more effectively than drug treatments alone.

[0010] Accordingly, there is provided in accordance with one aspect of the present invention a method for treating an occluded digestive tract projection. The method involves accessing a segment of the digestive track (preferably in a minimally invasive manner) and advancing a catheter to a proximal end of the digestive track projection (e.g., the appendix). A stent is deployed from the catheter into a lumen of the digestive tract projection, preferably at the proximal end of the projection. The stent conducts therethrough body fluids, bacteria and inflammatory components from the digestive tract projection to the digestive track (e.g., to the large intestine). In this manner, the stent can be used to drain an inflamed digestive track projection in a minimally invasive manner.

[0011] In some embodiments, the stent, the catheter, or another implant delivered during the same procedure can also be used to deliver a therapeutic agent (e.g., a drug) to the area in or around the digestive track projection. For example, the stent can elute an antibiotic to the tissue forming the digestive track projection to treat infection or other agents to reduce tissue inflammation.

[0012] In some embodiments, parts or the whole of the stent can be bioabsorbable and dissolve over a period of time (e.g., over a few weeks or months). In this manner, the stent can maintain patency of the proximal opening into the digestive track projection when infected, but dissolve once the infection has been treated. The shape of the implant can facilitate anchoring the implant in place when the tissue forming the proximal opening into the digestive track projection is inflamed and then releasing when tissue inflammation subsides.

[0013] In some embodiments, the stent can be removed through a second procedure. For this purpose, the stent can include structure that facilitates removal of the stent from the proximal opening into the digestive track projection. For example, the stent can include a tab or leash that can be grasped to pull the stent from the proximal opening. Such a tab or leash also can cause the inward collapse or movement

of some or all of the stent structure to allow the stent's removal in a less traumatic procedure.

[0014] In some embodiments, the stent includes one or more valves operating in one or more lumens to control fluid flow through the stent. For example, the stent can include a one-way valve to allow fluid flow out of the digestive track projection, but inhibit reverse flow.

[0015] The implantation procedure in some embodiments also can include the step of removing or macerating obstructions within the digestive track projection. For example, a macerating element can be delivered along with or integrated into the catheter for insertion into a proximal opening of or a space within the digestive track projection. The macerating element can either remove the obstruction from the digestive track projection or macerate. In either event, the obstruction or pieces thereof can be left within the body for natural discharge or can be removed (e.g., by aspiration or basket capture) by the catheter.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] These and other features, embodiments, and advantages of the present invention will now be described in connection with preferred embodiments of the invention, in reference to the accompanying drawings. The illustrated embodiments, however, are merely examples and are not intended to limit the invention.

[0017] FIG. 1 illustrates a schematic view of a human digestive tract.

[0018] FIG. 2 illustrates a schematic partially exploded sectioned view of a portion of a digestive tract near an appendix.

[0019] FIG. 3 illustrates a schematic partially exploded sectioned view of a portion of a digestive tract with diverticulum.

[0020] FIG. 4 is a schematic cross-sectional side view of a digestive tract projection, such as an appendix or diverticulum or other similar structures.

[0021] FIG. 5 is a schematic cross-sectional side view of an inflamed, infected, irritated or occluded projection of the digestive tract.

[0022] FIG. 6 is a schematic side perspective view of a digestive tract stent according to one embodiment of the present invention.

[0023] FIG. 7 is a schematic cross sectional side view of a digestive tract stent according to one embodiment of the present invention.

[0024] FIG. 8 is a schematic cross sectional side view of a digestive tract stent with a valve in a decreased flow configuration according to one embodiment of the present invention.

[0025] FIG. 9 is a schematic cross sectional side view of the digestive tract stent with a valve of FIG. 8 with the valve in an increased flow configuration.

[0026] FIG. 10 is a schematic side view of a mesh digestive tract stent according to one embodiment of the present invention.

[0027] FIG. 11 is a schematic side view of a mesh digestive tract stent according to one embodiment of the present invention with constant lumen width.

[0028] FIG. 12 is a schematic side view of a mesh digestive tract stent according to one embodiment of the present invention with linearly varying lumen width.

[0029] FIG. 13 is a schematic side view of a mesh digestive tract stent according to one embodiment of the present invention with varying lumen width.

[0030] FIG. 14 is a schematic side view of a mesh digestive tract stent with varying lumen width according to one embodiment of the present invention in a compressed configuration and in an expanded configuration.

[0031] FIG. 15 is a schematic side view of a mesh digestive tract stent with constant lumen width according to one embodiment of the present invention in a compressed configuration and in an expanded configuration.

[0032] FIG. 16 is a schematic side view of a digestive tract stent with one or more anchors according to one embodiment of the present invention.

[0033] FIG. 17 is a schematic side view of an hourglass shaped digestive tract stent according to one embodiment of the present invention.

[0034] FIG. 18 is a schematic side view of a digestive tract stent with a tether according to one embodiment of the present invention.

[0035] FIGS. 19-22 are schematic side view partial cross sectional views of the delivery of a stent to a digestive tract projection lumen using a catheter and a guide wire according to one embodiment of the present invention.

[0036] FIG. 23 is a schematic partially cross-sectional side view of an obstruction in a digestive tract projection lumen.

[0037] FIG. 24 is a schematic partially cross-sectional side view of an obstruction removal device according to one embodiment of the present invention.

[0038] FIG. 25 is a schematic side view of an obstruction removal device with two fingers for manipulating, grabbing, macerating, crushing, and/or compressing the obstruction according to one embodiment of the present invention.

[0039] FIG. 26 is a schematic perspective view of an obstruction removal device with a plurality of linked fingers for manipulating, grabbing, macerating, crushing, and/or compressing the obstruction according to one embodiment of the present invention.

[0040] FIG. 27 is a schematic side view of an obstruction removal device with a cutter according to one embodiment of the present invention.

[0041] Throughout the figures, the same reference numerals and characters, unless otherwise stated, are used to denote like features, elements, components or portions of the illustrated embodiments. In certain instances, similar names may be used to describe similar components with different reference numerals which have certain common or similar features. Moreover, while the subject invention will now be described in detail with reference to the figures, it is done so in connection with the illustrative embodiments. It is intended that changes and modifications can be made to the described embodiments without departing from the true scope and spirit of the subject invention as defined by the appended claims.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[0042] As should be understood in view of the following detailed description, this application is primarily directed to apparatuses, systems and methods for treating anatomical structures of the digestive tract. Embodiments of the present invention are directed to less-invasive technologies and methods of treating inflamed, infected and/or diseased portions of the digestive tract. The digestive tract can have one or more blind ended projections that are naturally formed or that develop with aging as a response to exposure to bacteria or other environmental factors. Certain projections from the digestive tract, such as the appendix or diverticular abscesses

(or diverticulum) can have worm-like or mushroom-like appearances in which recesses or pouches extend into or through the walls of the digestive tract. Various embodiments of the present invention are used to treat projections, or pouch-like extensions, of the digestive tract, such as an appendix or a diverticulum, with a device to reduce inflammation from an infection or other trauma without resorting to open surgery to remove portions of the digestive tract.

[0043] FIG. 1 illustrates a schematic representation of a human digestive tract. The digestive tract can be viewed as extending from the mouth, through the throat, down the esophagus 10 into the stomach 20 and to the small intestine 30, proceeding through the large intestine 40 (or colon) to the rectum 50 and terminating at the anus 60. Certain pouch-like structures or projections can extend from or through the walls of the digestive tract, such as at the large intestine 40. For example, some of these projections include the appendix 70 and diverticular abscesses (or diverticulum 80). Additional background information on the appendix 70 and diverticulum 80 can be found in libraries or on the Internet at sites such as Wikipedia or public documents from the National Institutes of Health.

[0044] The appendix 70 is a blind ended tube or lumen in fluid connection with the digestive tract. FIG. 2 illustrates a portion of a digestive tract near an appendix 70. Other names for the appendix 70 include the vermiform appendix and cecal appendix. In most people, the appendix 70 develops embryologically and can be found near the junction of the small intestine and the large intestine. The appendix 70 averages 10 cm in length, but can range from 2 to 26 cm. The diameter of the appendix 70 is usually between 7 and 8 mm. The appendix 70 is generally located in the lower right quadrant of the abdomen, or more specifically, the right iliac fossa. Its position within the abdomen corresponds to a point on the surface known as McBurney's point. While the base of the appendix 70 is at a fairly constant location, roughly 2 cm below the ileocaecal valve, the location of the tip of the appendix 70 can vary from being retrocaecal (in roughly three quarters of the population) to being in the pelvis to being extraperitoneal. In rare individuals with situs inversus, the appendix 70 may be located in the lower left side.

[0045] The understanding of the function of the appendix 70 is undergoing revisions in the scientific community. A common view was that the appendix 70 was a vestigial, or useless feature with no identified purpose. Doctors, professors and scientists have speculated that the appendix 70 was used in distant ancestors for digesting food, but could not determine how the appendix 70 assisted in the digestive process. More recently, scientists have demonstrated that the appendix 70 may currently serve a purpose for the immune system and/or for maintaining proper beneficial bacteria levels in the digestive tract. For example, studies have shown endocrine cell found in the appendix 70 of 11 week old fetuses that may help biological control and homeostatic mechanism. Some propose that the appendix 70 functions as a lymphatic organ, and studies have shown that the appendix 70 has a relatively high number infection-fighting lymphoid cells, suggesting that it might play a role in the immune system. Some suggest the appendix 70 plays a role in both manufacturing hormones in fetal development as well as functioning to 'train' the immune system, exposing the body to antigens in order that it can produce antibodies. Some doctors have stopped removing the appendix 70 during other surgical procedures as a routine precaution, because it can be

successfully transplanted into the urinary tract to rebuild a sphincter muscle and reconstruct a functional bladder. The benefits of keeping the appendix 70 intact in the body are increasing.

[0046] Although it was long accepted that the immune tissue, called gut associated lymphoid tissue, surrounding the appendix 70 and elsewhere in the gut carries out a number of important functions, explanations were lacking for the distinctive shape of the appendix 70 and its apparent lack of importance as judged by an absence of side-effects following appendectomy. Some scientists have proposed that the appendix 70 serves as a safe haven for useful bacteria when illness flushes those bacteria from the rest of the intestines. This proposal is based on a new understanding of how the immune system supports the growth of beneficial intestinal bacteria, in combination with many well-known features of the appendix 70, including its architecture and its association with copious amounts of immune tissue. Such a function is expected to be useful in a culture lacking modern sanitation and healthcare practice, where diarrhea may be prevalent. Current epidemiological data show that diarrhea is one of the leading causes of death in developing countries, indicating that a role of the appendix 70 as an aid in recovering beneficial bacteria following diarrhea may be extremely important in the absence of modern health and sanitation practices.

[0047] Other anatomical structures in the digestive tract may benefit from a device, system, and/or method of treating the appendix 70. Many people have small pouches in their colons that bulge outward through weak spots, like an inner tube that pokes through weak places in a tire. Each pouch is called a diverticulum 80. Pouches (plural) are called diverticula 80. FIG. 3 shows diverticula 80 in the colon. The condition of having diverticula 80 is called diverticulosis. About 10 percent of Americans over the age of 40 have diverticulosis. The condition becomes more common as people age. About half of all people over the age of 60 have diverticulosis.

[0048] When the pouches become infected or inflamed, the condition is called diverticulitis. This happens in 10 to 25 percent of people with diverticulosis. Diverticulosis and diverticulitis are also called diverticular disease. Most people with diverticulosis do not have any discomfort or symptoms. However, symptoms may include mild cramps, bloating, and constipation. The most common sign is tenderness around the left side of the lower abdomen. If infection is the cause, fever, nausea, vomiting, chills, cramping, and constipation may occur as well. The severity of symptoms depends on the extent of the infection and complications. Diverticulitis can lead to bleeding, infections, perforations or tears, or blockages. These complications always require treatment to prevent them from progressing and causing serious illness.

[0049] The infection causing diverticulitis can form an abscess in the colon. An abscess is an infected area with pus that may cause swelling and destroy tissue. Sometimes the infected diverticula 80 may develop small holes, called perforations. These perforations allow pus to leak out of the colon into the abdominal area. If the abscess does not clear up with antibiotics, the doctor may need to drain it. To drain the abscess, the doctor uses a needle and a small tube called a catheter. The doctor inserts the needle through the skin and drains the fluid through the catheter. This procedure is called percutaneous catheter drainage. Sometimes surgery is needed to clean the abscess and, if necessary, remove part of the colon. A large abscess can become a serious problem if the

infection leaks out and contaminates areas outside the colon. Infection that spreads into the abdominal cavity is called peritonitis. Peritonitis requires immediate surgery to clean the abdominal cavity and remove the damaged part of the colon. Without surgery, peritonitis can be fatal.

[0050] A fistula is an abnormal connection of tissue between two organs or between an organ and the skin. When damaged tissues come into contact with each other during infection, they sometimes stick together. If they heal that way, a fistula forms. When diverticulitis-related infection spreads outside the colon, the colon's tissue may stick to nearby tissues. The organs usually involved are the bladder, small intestine, and skin. The most common type of fistula occurs between the bladder and the colon. It affects men more than women. This type of fistula can result in a severe, long-lasting infection of the urinary tract. The problem can be corrected with surgery to remove the fistula and the affected part of the colon.

[0051] The scarring caused by infection may cause partial or total blockage of the large intestine. When this happens, the colon is unable to move bowel contents normally. When the obstruction totally blocks the intestine, emergency surgery is necessary. Partial blockage is not an emergency, so the surgery to correct it can be planned.

[0052] Treatment for diverticulitis focuses on clearing up the infection and inflammation, resting the colon, and preventing or minimizing complications. An attack of diverticulitis without complications may respond to antibiotics within a few days if treated early enough. An acute attack with severe pain or severe infection may require a hospital stay. Most acute cases of diverticulitis are treated with antibiotics and a liquid diet. If attacks are severe or frequent, the doctor may advise surgery. The surgeon removes the affected part of the colon and joins the remaining sections. This type of surgery, called colon resection, aims to keep attacks from coming back and to prevent complications. The doctor may also recommend surgery for complications of a fistula or intestinal obstruction. If antibiotics do not correct an attack, emergency surgery may be required. Other reasons for emergency surgery include a large abscess, perforation, peritonitis, or continued bleeding.

[0053] Emergency surgery usually involves two operations. The first surgery will clear the infected abdominal cavity and remove part of the colon. Because of infection and sometimes obstruction, it is not safe to rejoin the colon during the first operation. Instead, the surgeon creates a temporary hole, or stoma, in the abdomen. The end of the colon is connected to the hole, a procedure called a colostomy, to allow normal eating and bowel movements. The stool goes into a bag attached to the opening in the abdomen. In the second operation, the surgeon rejoins the ends of the colon.

[0054] FIG. 4 is a schematic cross-sectional side view of a general digestive tract projection 90, such as an appendix 70 or diverticulum 80 or other similar structures. The digestive tract projection 90 extends from digestive tract lumen 91 from a digestive tract wall 92 with a projection lumen 93 extending from a proximal section 94 to a distal section 95 surrounded by a digestive tract projection wall 96. When the digestive tract projection 90 is healthy, or not inflamed or infected it can perform its function. However, as illustrated in FIG. 5, the digestive tract projection 90 can become inflamed, infected, irritated or occluded, closing a portion of the lumen 93 at the proximal section 94, distal section 95, or both. The digestive

tract projection wall 96 can swell and become very sensitive, and potentially perforate or rupture if untreated.

[0055] In one embodiment a digestive tract stent 100 has a first, or proximal end 110, a second, or distal end 120, a lumen 130 and a side wall 140. In various embodiments, the stent 100 can have a circular, elliptical, oval, square, rectangular, parallelogram, pentagonal, hexagonal, octagonal or other shape cross-section transverse its longitudinal axis 150. In FIG. 6 the stent 100 has a substantially circular cross-section. FIG. 7 illustrates a sectional view of the stent 100 of FIG. 6 with the side wall 140 having an external surface 141 and an internal surface 142 along the lumen 130. In one embodiment the stent 100 has an extrudable cross-sectional shape with constant width, depth, and/or thickness. In various embodiments the width, height, wall thickness, diameter, circumference, or other geometric characteristics of the stent 100 can be varied. In one embodiment the stent 100 can have an hour-glass shape.

[0056] In various embodiments treatment of an infection, inflamed, diseased or swollen digestive tract projection 90 with a stent is advantageous over existing surgical removal of the digestive tract projection 90 include, but are not limited to, an easier set up, no requirement for an operating room, wider availability, less cost, procedures can be performed under conscious sedation instead of general anesthesia, procedures are less invasive, reduced risk of complications, reduced risk of bleeding, less expertise required for treatment, a gastrointestinal doctor can perform the procedures, quicker recovery time, less lost time for the patient. Longer term benefits include, but are not limited to, keeping the digestive tract projection 90. Those who have an appendectomy before age 10 are relatively protected from Crohn's disease, while those who have one later are at increased risk of Crohn's. One long term benefit of stent treatment of a digestive tract projection is that surgical complications are reduced or are eliminated. There is a reduced risk of adhesions that may cause small or large bowel obstruction.

[0057] In various embodiments the stent 100 is in the range of 0.25 cm-26 cm in length. In one embodiment the stent 100 is roughly 1 cm in length. In one embodiment the stent 100 is roughly 5 cm in length. In one embodiment the stent 100 is roughly 10 cm in length. In various embodiments the stent 100 length is in the range of 3 to 8 cm long. In one embodiment the stent 100 has a width or diameter the same or larger than that of the digestive tract projection lumen 93 width or diameter. Diameter can be measured in French. In one embodiment the stent 100 is delivered by a duodenoscope, which generally currently able to accept stents (in a collapsed or full expansion configuration) with a diameter less than or equal to 12 Fr. In various embodiments the digestive tract projection lumen 93 width or diameter is in the range of 6-9 mm. In one embodiment the stent 100 width or diameter is roughly 7 mm. In one embodiment the stent 100 width or diameter is roughly 8 mm. In one embodiment the stent is formed of a laser cut material. In one embodiment, the stent is formed of interwoven wires.

[0058] In one embodiment the stent 100 is made of plastic. In various embodiments the stent 100 can be made of Teflon, polyethylene, polyurethane, polyflex, or other materials. In one embodiment the stent 100 is made of metal. In various embodiments the stent 100 can be made of stainless steel, nickel-titanium alloy, cobalt-tungsten alloy, cobalt-chromium alloy, tantalum alloy, gold, titanium, or other materials. In one embodiment the stent 100 is repositionable during

deployment. In one embodiment the stent **100** is repositionable after deployment. In one embodiment the stent **100** is removable after deployment. In one embodiment the stent **100** is repositionable after deployment in a separate procedure. In one embodiment the stent **100** is made of a bio-absorbable material. In one embodiment the stent **100** is made of Vicryl (polyglactin 910). In various embodiments the stent **100** is made of polyester, polyorthoester, polyanhydride, collagen, or other biodegradable material. In one embodiment the stent **100** can be absorbed in a number of days to a number of weeks. In one embodiment the stent **100** dissolves after a period of time. In one embodiment the stent **100** dissolves after a few days. In one embodiment the stent **100** dissolves after a few weeks.

[0059] In one embodiment the stent **100** is made of a rigid material, such as a plastic, polymer, or metal. In one embodiment the stent **100** is made of an elastic material, such as certain plastics, polymers, or metals with flexible characteristics. In one embodiment the stent **100** is made of a shape memory material, such as Nitinol. In one embodiment, an elastic or deformable stent **100** can have a first configuration and a second configuration. In one embodiment the first configuration can have a small cross sectional profile to facilitate insertion over a guidewire or in a catheter into a digestive tract projection lumen **93**. In one embodiment the second configuration can have a large cross sectional profile to expand the height, width or depth of a digestive tract projection lumen **93**. In one embodiment a stent **100** can be expanded or reduced by gas, liquid, and/or fluid pressure. In one embodiment a stent **100** can be expanded or reduced by a balloon. In one embodiment a stent **100** is self expanding.

[0060] In one embodiment the stent **100** is at least partially coated with a low-friction substance (e.g., PTFE) to facilitate insertion of the stent **100** into a digestive tract projection lumen **93**. In one embodiment the stent **100** is at least partially coated with a low-friction substance on the distal end **120** of the outer wall **141** of the stent **100** to facilitate insertion of the stent **100** into a digestive tract projection lumen **93**. In one embodiment the stent **100** is at least partially coated with a high-friction substance to help anchor the stent **100** into a digestive tract projection lumen **93**. In one embodiment the stent **100** is at least partially coated with a high-friction substance on the proximal end **110** of the outer wall **141** of the stent **100** to help anchor the stent **100** into a digestive tract projection lumen **93**. In one embodiment the stent **100** is at least partially coated with a drug coating to release or elute therapeutic agents, such as an antibiotic, an antimicrobial agent, a growth factors, a growth inhibitors, an antiseptic agent, an anti-inflammatory agent, an antirestinosus agent, or an antioxidant. In one embodiment, the stent **100** is at least partially coated with a hydrophilic polymer coat. In one embodiment the stent **100** is at least partially coated with a silver coat. In one embodiment the stent **100** is at least partially coated with a bacterial delivery coat.

[0061] In one embodiment the stent **100** is sterile. In one embodiment the stent **100** comprises media eluting material. In one embodiment the stent **100** comprises pores on the surface or other drug storage systems for drug loading. In one embodiment the media is a drug or therapeutic agent. In one embodiment the media is one or more antibiotics. In one embodiment the media is a growth factor. In one embodiment the media is growth inhibitor. In one embodiment the media is bacteria to provide for beneficial flora to the digestive tract. In one embodiment the media is an antimicrobial agent. In one

embodiment the media is delivered over a time-delayed or extended time-release means. In one embodiment an antibiotic is initially released, followed by a time-delayed release of beneficial flora after the antibiotics have dispersed.

[0062] In one embodiment, a media or therapeutic agent is delivered to the digestive tract projection through one or more injection ports on a catheter. In one embodiment, the media or therapeutic agent is delivered to the digestive tract projection through one or more injection ports on an expandable balloon. In one embodiment, a media or therapeutic agent is delivered to the digestive tract projection through one or more injection ports on a stent. In one embodiment, the media or therapeutic agent is delivered simultaneous with the delivery of the stent **100**. In one embodiment, the media or therapeutic agent is delivered before or after the delivery of the stent **100**. Suitable media or therapeutic agents include, but are not limited to, an antibiotic agent, beneficial flora, an antimicrobial agent, a growth factors, a growth inhibitors, an antiseptic agent, an anti-inflammatory agent, an antirestinosus agent, or an antioxidant.

[0063] In one embodiment the stent **100** can have a valve **160** disposed in the lumen **130**. In one embodiment the valve **160** is disposed between the proximal end **110** and the distal end **120** of the stent **100**. In one embodiment the valve **100** is located at or near the proximal end **110**. In one embodiment the valve **100** is located at or near the distal end **120**. In one embodiment the valve **100** is a one-way valve that allows fluid or particle flow in one direction and reduces or eliminates flow in the opposite direction. In one embodiment the valve **100** is a one-way valve that promotes drainage of the digestive tract projection lumen **93**. In one embodiment the valve **100** is a one-way valve that prevents reflux of digestive tract lumen **91** contents into the digestive tract projection lumen **93**.

[0064] In one embodiment valve **100** is duck-bill valve. In one embodiment valve **100** has two or more opposed surfaces that can touch each other to occlude or restrict the lumen **130**, and that can be actuated by pressure differential between the proximal end **110** and the distal end **120** of the stent **100**. In one embodiment deformation of the stent **100** can open or close the valve **160**. In one embodiment valve **100** is a flap valve. In one embodiment valve **100** has one or more biased or spring loaded or deflectable surfaces. As shown in FIG. 8 a valve **100** can be configured to restrict or fully occlude flow within the lumen **130** between a proximal lumen section **131** and a distal lumen section **132**, and as shown in FIG. 9 the valve **100** can be configured to allow flow from the distal lumen section **132** to the proximal lumen section **131**.

[0065] In one embodiment a mesh digestive tract stent **200** includes an expandable mesh structure. In one embodiment a mesh stent **200** is capable of being delivered in a small profile and expanding into a large profile to facilitate flow in at least one direction in a digestive tract projection lumen **93**. In one embodiment a mesh stent **200** is self expanding. In one embodiment a mesh stent **200** expanded or reduced by gas, liquid, and/or fluid pressure. In one embodiment a mesh stent **200** can be expanded or reduced by a balloon (e.g. by an expandable balloon placed inside the stent). In one embodiment a mesh stent **200** has a nonabrasive or atraumatic distal and/or proximal end. Mesh stent **200** can have many of the characteristics described in relation to stent **100**. In any of the stent embodiments described herein, features of a stent in one embodiment can be combined with features of a stent in another embodiment. In one embodiment a mesh digestive tract stent **200** has a first, or proximal end **210**, a second, or

distal end **220**, a lumen **230** and a mesh wall **240**. In various embodiments, the mesh stent **200** can have a circular, elliptical, oval, square, rectangular, parallelogram, pentagonal, hexagonal, octagonal or other shape cross-section transverse its longitudinal axis **250**. In FIG. **10** the mesh stent **200** has a substantially circular cross-section. In one embodiment a mesh stent **200a** can have a constant lumen width, as shown in FIG. **11**. In some embodiment a width reflects a diameter. In various embodiments, either the proximal end **210** width or the distal end **220** width can be larger than the other. In various embodiments, a width of the stent **200** between the proximal end **210** and the distal end **220** can be larger or smaller than either or both of the proximal end **210** width or the distal end **220** width. In one embodiment a mesh stent **200b** can have a linearly varying lumen width, as shown in FIG. **12**. In one embodiment a mesh stent **200c** can have a varying lumen width, as shown in FIG. **13**. In one embodiment a mesh stent **200** can have an expanded configuration **200d** and a compressed configuration **200e**, as shown in FIGS. **14** and **15**.

[**0066**] FIG. **16** illustrates one embodiment a digestive tract stent **300** with a proximal end **310** and a distal end **320** has one or more anchors. In one embodiment, the stent **300** has one or more proximal anchors **343**. In one embodiment, the stent **300** has one or more distal anchors **344**. In one embodiment the anchors are use to prevent migration in one or more directions. Any of the anchors can be a rough surface, coating, barb, or surface feature that helps retain the stent **300** in the digestive tract projection lumen **93**. Any of the anchors can be disposed at an end or anywhere along the length of the stent **300**. The angle of a barb or orientation of an anchor can be in any direction.

[**0067**] In one embodiment a digestive tract stent **400** with a proximal end **410** and a distal end **420** has an hourglass configuration, as illustrated in FIG. **17**. In one embodiment the hourglass configuration can be configured to help retain the stent **400** in a position in the digestive tract projection lumen **93** by holding a portion of the tissue of the digestive tract projection lumen **93** between the proximal end **410** and a distal end **420**. In one embodiment the hourglass configuration can have one or more anchors to help retain the stent **400** in a position in the digestive tract projection lumen **93**. In any embodiment a stent can have a marker **430**. In one embodiment the marker **430** is radiopaque to facilitate visualization under fluoroscopy or other scanning or visualization techniques. In various embodiments the marker **430** is a dot, a ring, a shape for visualization and/or identification of the stent and/or stent orientation. In various embodiments, the visualization is facilitated by using stent wire with radiopaque cores or radiopaque coatings or platings.

[**0068**] In one embodiment a digestive tract stent **500** with a proximal end **510** and a distal end **520** has a tether **530**, as illustrated in FIG. **18**. In various embodiments the tether can be attached to any portion of the stent **500** to facilitate repositioning or removal of the stent **500**. In one embodiment the tether **530** has a snare feature **535**, such as a loop, that can be used to grab the tether **530**. In one embodiment the tether **530** is connected to the distal end **520** at a tether interface **532**. The tether **530** can be moved in a proximal direction to collapse or reduce the cross sectional size or profile of the distal end **502** of the stent **500** in order to facilitate insertion, repositioning, and/or removal of the stent **500**. In one embodiment, the stent **500** can be delivered with the tether **530** being pulled or moved proximally so the distal end **520** is small to facilitate insertion into a lumen. In one embodiment the tether **530** can

be pulled or moved proximally to reduce the size of the distal end **520** to facilitate removal of the stent **500** from a lumen. In one embodiment the tether **530** can be pulled or moved proximally to reduce the size of the distal end **520** to disengage a distal anchor (such as distal anchor **344**) to facilitate removal of the stent **500** from a lumen. The tether interface **532** can be located anywhere on the stent **500**.

[**0069**] In one embodiment of a digestive tract stent and method for delivery and/or removal of a stent, a spiral shaped device and methods associated with the deployment of a device for deployment which can be configured in a first delivery configuration to and/or from a second deployed spiral configuration is described in U.S. Pat. No. 5,536,274 issued Jul. 16, 1996 to Neuss can be used. U.S. Pat. No. 5,536,274 is incorporated by references as a whole, including FIGS. 1-6 and FIGS. 11a-11e, col. 2, l. 13 to col. 3, line 30, col. 3 line 31-col. 4 line 49, col. 4, ll. 63-65, col. 7, line 59 to col. 8, line 28 and col. 8, line 41 to col. 9, line 47.

[**0070**] FIGS. **19-22** illustrate various potential steps in one embodiment of the delivery of a stent **100** to a digestive tract projection lumen **93** using a catheter **600** and a guide wire **610** according to one embodiment of the present invention. In various embodiments a catheter **600** with a stent **100** can be directed to the digestive tract lumen **91** near a digestive tract projection lumen **93** opening from an endoscope, from an oral endoscopic path, from an anal endoscopic path, from the mouth, from the anus, from laparoscopic insertion, from piercing the digestive tract superior to or inferior to the digestive tract projection lumen **93** opening and inserting the catheter **600**. In one embodiment the laparoscopic insertion is made in the stomach. In one embodiment the laparoscopic insertion is made in the small intestine. In one embodiment the laparoscopic insertion is made in the large intestine superior or inferior to the digestive tract projection lumen. In one embodiment access to the digestive tract projection lumen **93** is provided by piercing the projection wall **96**. In one embodiment, piercing the projection wall **96** is accomplished with a cannulated needle and/or a guidewire. In one embodiment the catheter **600** has a guide wire **610** which can be inserted into the digestive tract projection lumen **93** to create a path for the stent **100** to be delivered. In one embodiment aspects of a procedure can be conducted under endoscopy. In one embodiment the procedure is conducted using a radiographically direct procedure, for example, under fluoroscopy. Fluoroscopy and other radiographically direct procedures each have an added benefit of being able to visualize bleeding or perforations. This approach would eliminate the need to use a large amount of injected air or gas via an endoscopic tube, and would reduce the risk of pushing fecal material through a weakness in the bowel wall or into the appendix. In one embodiment aspects of a procedure can be conducted under computed tomography (CT) guidance. In one embodiment aspects of a procedure can be conducted under ultrasound imagery. In one embodiment aspects of a procedure can be conducted under MRI. In one embodiment aspects of a procedure can be conducted in a catheter lab. In all such embodiments where access is gained through a naturally occurring orifice within the body, the treatment method eliminates the need for an operating room to treat the occluded digestive tract projection (e.g., an appendicitis). Such procedures only require a procedure room, but do not need a sterile environment. On the other hand, prior courses to treat an appendicitis

involved invasive surgery, which required an operating room with the accompanying sterile environment, procedure and expense.

[0071] In a further embodiment, the placement of the stent **100** can occur after an incision has been made partially through a wall of the connection portion between the digestive tract lumen **91** and the appendix **70** (or the diverticulum **80**) in a manner similar to a papillotomy. Such an incision will increase the size of the lumen **93**, once stented open, between the digestive tract lumen **91** and the appendix **70** (or the diverticulum **80**). A larger lumen size can enhance the release of matter from the appendix **70** and reduce the likelihood of later inflammation of the appendix **70** (or the diverticulum **80**).

[0072] FIG. 23 illustrates an obstruction **97**, which can be a stone, calcification, abscess, pus, or other material which can reside in or block the digestive tract projection lumen **93**. In one embodiment an obstruction removal device **700** can be directed to the digestive tract projection lumen **93** to remove or reduce the size of the obstruction **97**. In various embodiments a obstruction removal device **700** can be directed to the digestive tract lumen **91** near a digestive tract projection lumen **93** opening from an endoscope, from an oral endoscopic path, from an anal endoscopic path, from the mouth, from the anus, from laparoscopic insertion, from piercing the digestive tract superior to or inferior to the digestive tract projection lumen **93** opening and inserting the obstruction removal device **700**. In one embodiment the obstruction removal device **700** has a guide wire (not illustrated here) which can be inserted into the digestive tract projection lumen **93** to create a path for the obstruction removal device **700** to get near the obstruction **97**, as shown in FIG. 24. In one embodiment the obstruction removal device **700** is a suction device which can aspirate the obstruction **97** from the digestive tract projection lumen **93** with pressure differential. In one embodiment the obstruction removal device **700** has a sharp point or edge for piercing or cutting the obstruction **97**. In one embodiment the obstruction removal device **700** has a gripping mechanism for grabbing the obstruction **97**. In one embodiment the obstruction removal device **700** has two or more fingers **710** for manipulating, grabbing, macerating, crushing, and/or compressing the obstruction **97**. See FIG. 25. In various embodiments the obstruction removal device **700** can have two, three, four or more fingers **710**. In one embodiment the obstruction removal device **700** has a hinged linkage with two, three, or more links **720** for manipulating, grabbing, macerating, crushing, and/or compressing the obstruction **97**. See FIG. 26. In one embodiment the obstruction removal device **700** includes one or more nets. In one embodiment the obstruction removal device **700** includes one or more snares or baskets. Suitable baskets are disclosed in U.S. Pat. No. 5,330,482, which is hereby incorporated by reference. In one embodiment the obstruction removal device **700** includes one or fibers for ensnaring the obstruction **97**. In one embodiment the obstruction removal device **700** includes a medium or media for dissolving the obstruction **97**. In one embodiment the obstruction removal device **700** includes a cutter **730**. See FIG. 27. In one embodiment the cutter **730** is a linear cutter. In one embodiment the cutter **730** is a rotational cutter. In one embodiment the cutter **730** is a reciprocating cutter. In one embodiment the cutter **730** comprises one or more blades. In one embodiment the cutter **730** comprises one or more gear teeth. In one embodiment the obstruction removal device **700** includes an atraumatic distal tip to

facilitate insertion into a swollen lumen. Any of the obstruction removal device **700** embodiments can be combined.

[0073] It will be understood that the foregoing is only illustrative of the principles of the invention, and that various modifications, alterations, and combinations can be made by those skilled in the art without departing from the scope and spirit of the invention. Accordingly, it is not intended that the invention be limited, except as by the appended claims.

What is claimed is:

1. A method of treating an occluded digestive tract projection, comprising:

advancing a catheter to the proximal end of a digestive tract projection; and
deploying a stent from the catheter to the lumen of the digestive tract projection.

2. The method of claim 1, wherein the digestive tract projection is an appendix.

3. The method of claim 1, wherein the digestive tract projection is a diverticulum.

4. The method of claim 1, further comprising the delivery of a therapeutic agent.

5. The method of claim 1, wherein the catheter is advanced using an endoscope.

6. The method of claim 1, wherein the catheter is advanced through a laparoscopic insertion.

7. A stent for treating an occluded digestive tract projection, comprising a proximal end portion, a distal end portion, a lumen extending between the proximal and distal end portions; and a side wall sized and configured for opening a lumen of a digestive tract projection.

8. The stent of claim 7, wherein the digestive tract projection is an appendix.

9. The stent of claim 7, wherein the digestive tract projection is a diverticulum.

10. The stent of claim 7, wherein at least the proximal and distal end portions are formed of a biocompatible material.

11. The stent of claim 7, wherein at least the proximal and distal end portions are formed of a bioabsorbable material.

12. The stent of claim 7 comprising a therapeutic agent that elutes from the stent when implanted.

13. The stent of claim 7 comprising a delivery mechanism wherein a beneficial flora is released before or after a therapeutic agent is released.

14. The stent of claim 7, further comprising one or more coating selected from the group consisting of a biocompatible coating, a low-friction coating, a high-friction coating, a hydrophilic polymer coating, a silver coating, and a therapeutic agent coating.

15. The stent of claim 7, further comprising a one-way valve.

16. The stent of claim 7, further comprising a securing feature.

17. The stent of claim 7, further comprising a tether.

18. The stent of claim 7, wherein the stent has regions of varying width.

19. A method of removing an obstruction of an occluded digestive tract projection, comprising:

advancing a catheter to the proximal end of a digestive tract projection;

deploying a obstruction removal device in to the lumen of the digestive tract projection;

reducing the size of the obstruction; and

removing at least part of the obstruction from the lumen of the digestive tract projection.

20. The method of claim 19, wherein the digestive tract projection is an appendix.

21. The method of claim 19, wherein the digestive tract projection is a diverticulum.

22. The method of claim 19, further comprising the delivery of a therapeutic agent.

23. The method of claim 19, wherein the obstruction removal device is advanced using an endoscope.

24. The method of claim 19, wherein the obstruction removal device is advanced through a laparoscopic insertion.

25. The method of claim 19, further comprising positioning a stent to the lumen of the digestive tract projection after removing at least part of the obstruction.

26. An obstruction removal device for removing an obstruction of an occluded digestive tract projection, comprising:

a catheter;

a macerating element sized and configured for insertion in to an opening a lumen of a digestive tract projection.

27. The obstruction removal device of claim 26, wherein the digestive tract projection is an appendix.

28. The obstruction removal device of claim 26, wherein the digestive tract projection is a diverticulum.

29. The obstruction removal device of claim 26, further comprising stent and a stent deployment device.

30. The obstruction removal device of claim 26, further comprising a therapeutic agent delivery mechanism.

* * * * *

专利名称(译)	用于治疗消化道的闭塞感染集合的装置和方法		
公开(公告)号	US20100100105A1	公开(公告)日	2010-04-22
申请号	US12/603469	申请日	2009-10-21
[标]申请(专利权)人(译)	BATES DOUGLAS 比尔曼史蒂芬°F		
申请(专利权)人(译)	BATES DOUGLAS 比尔曼史蒂芬°F		
当前申请(专利权)人(译)	BATES DOUGLAS 比尔曼史蒂芬°F		
[标]发明人	BATES DOUGLAS BIERMAN STEVEN F		
发明人	BATES, DOUGLAS BIERMAN, STEVEN F.		
IPC分类号	A61B17/22 A61F2/04 A61F2/82		
CPC分类号	A61F2/04 A61M27/002 A61F2002/045		
优先权	61/107261 2008-10-21 US		
其他公开文献	US8343088		
外部链接	Espacenet USPTO		

摘要(译)

一种用于治疗沿消化道的闭塞感染集合的装置和方法，其中支架被输送到消化道的阑尾，憩室或其他解剖结构。一种装置和方法，用于沿着消化道从闭塞的感染集合中移除阻塞物，其中阻塞物移除装置被输送到消化道的阑尾，憩室或其他解剖结构。

