



(12) **EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 158(3) EPC

(43) Date of publication:  
**21.06.2006 Bulletin 2006/25**

(21) Application number: **04788274.1**

(22) Date of filing: **29.09.2004**

(51) Int Cl.:  
**C12N 15/09** <sup>(1990.01)</sup> **A61K 45/00** <sup>(1985.01)</sup>  
**A61P 1/16** <sup>(2000.01)</sup> **A61P 11/06** <sup>(2000.01)</sup>  
**A61P 31/12** <sup>(2000.01)</sup> **A61P 31/14** <sup>(2000.01)</sup>  
**A61P 35/00** <sup>(2000.01)</sup> **A61P 37/04** <sup>(2000.01)</sup>  
**A61P 37/06** <sup>(2000.01)</sup> **A61P 37/08** <sup>(2000.01)</sup>  
**C07K 14/705** <sup>(1995.01)</sup> **C07K 16/28** <sup>(1995.01)</sup>  
**C12N 1/15** <sup>(1990.01)</sup> **C12N 1/19** <sup>(1990.01)</sup>  
**C12N 1/21** <sup>(1990.01)</sup> **C12N 5/00** <sup>(1980.01)</sup>  
**C12P 21/02** <sup>(1980.01)</sup> **C12Q 1/02** <sup>(1980.01)</sup>  
**G01N 33/15** <sup>(1980.01)</sup> **G01N 33/50** <sup>(1980.01)</sup>

(86) International application number:  
**PCT/JP2004/014207**

(87) International publication number:  
**WO 2005/030955 (07.04.2005 Gazette 2005/14)**

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
HU IE IT LI LU MC NL PL PT RO SE SI SK TR**

(30) Priority: **29.09.2003 JP 2003338331**

(71) Applicants:  
• **CHUGAI SEIYAKU KABUSHIKI KAISHA**  
Tokyo, 115-8543 (JP)  
• **MATSUSHIMA, Kouji**  
Matsudo-shi,  
Chiba 271-0092 (JP)  
• **Hashimoto, Shinichi**  
Tokyo 1100001 (JP)

(72) Inventors:  
• **MATSUSHIMA, Kouji**  
Matsudo-shi,  
Chiba 2710092 (JP)  
• **HASHIMOTO, Shinichi**  
Tokyo 110-0001 (JP)

- **TSUCHIYA, Masayuki,**  
c/o Chugai Seiyaku K. K.  
Gotenba-shi,  
Shizuoka 4120038 (JP)
- **HIRATA, Yuichi,**  
c/o Chugai Seiyaku K. K.  
Niihari-gunk,  
Ibaraki 3004101 (JP)
- **YOSHIDA, Kenji,**  
c/o Chugai Seiyaku K. K.  
Niihari-gun,  
Ibaraki 3004101 (JP)
- **OJIMA, Kazuyuki,**  
c/o Chugai Seiyaku K. K.  
Niihari-gun,  
Ibaraki 3004101 (JP)

(74) Representative: **Vossius & Partner**  
**Siebertstrasse 4**  
**81675 München (DE)**

Remarks:  
Applicant has requested a correction according to rule 88 EPC.

(54) **PROTEIN EXPRESSED IN NK CELL**

(57) Purified immunocytes were analyzed for expression frequencies, and the NKIR gene expressed specifically in natural killer (NK) cells was successfully identified. The NKIR gene encodes a receptor. Agonists and

antagonists for the receptor can be identified by using the receptor.

**Description**Technical Field

5 **[0001]** The present invention relates to novel proteins expressed in NK cells, DNAs encoding the proteins, methods for producing the molecules, and uses thereof.

Background Art

10 **[0002]** Cytotoxic T cells recognize complexes between major histocompatibility complex (MHC) class I molecules and specific antigen peptides (foreign substances), and are thereby activated to exert cytotoxic activity against target cells. In contrast, natural killer (NK) cells typically damage target cells expressing no MHC class I molecules. The expression of such MHC class I molecules on normal cell surface is suppressed by viral infection or canceration. NK cells exert cytotoxic activity against such abnormal cells with decreased MHC class I molecule expression levels. It is therefore  
15 thought that NK cells play a central role in the innate immunity mechanism by eliminating cancerated cells or cells infected with viruses. NK cells were in fact identified as cells having the activity of spontaneously damaging certain cancer cells (see Non-Patent Document 1). Furthermore, such *in vivo* roles of NK cells are supported by the disease called Chediak-Higashi syndrome, which is caused by a deficiency in NK cell activity (Chediak-Higashi syndrome is due to a chromosome abnormality and enters the advanced stage when the patient is over 10 years old, making it impossible to control viral  
20 infection, and resulting in malignant lymphoma-like pathological changes and death by pancytopenia after 2 to 3 months).

**[0003]** Recently, transgenic mice lacking NK1.1+CD3-, a marker of NK cells, were generated. Analyses of their characteristics revealed that: NK cells

(i) play an important role in suppressing metastasis and growth of cancer cells; and

25 (ii) are major producers of interferon (IFN)  $\gamma$  in response to bacterial endotoxins (see Non-Patent Document 2).

**[0004]** Meanwhile, it has recently been found that a fourth lymphocyte, NKT cell, that has a NK cell receptor as well as an identical T cell receptor and characteristics of both innate and acquired immunities, are distributed in a wide variety of organs, including liver and bone marrow. In addition, these cells have been found to be involved in immunotolerance,  
30 and in many diseases such as autoimmune diseases, hepatitis, and infections. Based on studies using mice that develop multiple sclerosis early in life, it has been found that there is a certain relationship between NKT cells, which are immune cells included in lymphocytes, and the onset of multiple sclerosis (see Non-Patent Document 3). In addition, it has been suggested that asthma may be preventable by inactivating NKT cells, because no allergic airway hyperreactivity, which is a major asthma symptom, is detected in NKT-deficient mice (see Non-Patent Document 4). The revealed activation  
35 mechanism for NKT cells shows that unlike NK cells, NKT cells are activated through stimulation of the T cell receptor or through  $\alpha$ -galactosylceramide ( $\alpha$ -GalCer), which is a glucolipid presented to the CD1d receptor on dendritic cells (see Non-Patent Document 3). Meanwhile, like NK cells, NKT cells exert cytotoxic activity against cells with decreased MHC class I molecule expression levels, and is therefore thought to regulate cytotoxic activity by almost the same mechanism as the killer cell inhibitory receptor (hereinafter abbreviated as "KIR"), which has been identified as an inhibitory regulator  
40 molecule involved in the cytotoxic activity of lymphocyte populations. However, NKT subsets that do not express known KIRs are also known to exist. In addition, much was unclear about signal cascades and inhibitory receptors that act negatively in the activation, necessitating further studies.

**[0005]** While ligand substances that are effective for NKT cell activation have been found, low-molecular-weight ligands that specifically activate NK cells are yet to be identified.

45 Non-Patent Document 1: Trinchieri G., Adv Immunol. (1989), 47, 187-376

Non-Patent Document 2: Sungjin Kim, Koho Iizuka, Hector L. Aguila, Irving L. Weissman, and Wayne M. Yokoyama, Proc. Natl.Acad.Sci. (2000), 97, 2731-2736

50 Non-Patent Document 3: Michishige Harada, Masaru Taniguchi, Protein, Nucleic acid, and Enzyme (Tanpakushitsu Kakusan Kouso) (2002), 47, 2109-2116

Non-Patent Document 4: Akbari O, Stock P, Meyer E, Kronenberg M, Sidobre S, Nakayama T, Taniguchi M, Grusby MJ, DeKruyff RH, Umetsu DT, Nat. Med. (2003), 9, 582-588

Disclosure of the InventionProblems to be Solved by the Invention

55 **[0006]** The present invention was achieved in view of the above circumstances. An objective of the present invention

is to isolate receptor molecules that specifically activate or inactivate NK cells. More specifically, an objective is to provide novel receptor proteins expressed in NK cells, DNAs encoding the proteins, and uses of the molecules.

#### Means to Solve the Problems

5

**[0007]** The NK cell activation mechanism has been suggested to take place as a result of the regulation between the positive signal cascade, which follows phosphorylation by protein tyrosine kinase, and the negative signal cascade, which results from the dephosphorylation of the phosphorylated signaling molecule. In mouse, the inhibitory receptors on NK cells which recognize molecules driving the negative cascade are the group of Ly49 molecules (a type-C lectin family having the extracellular type-C lectin domain). In humans, they are the KIR molecule family having the extracellular immunoglobulin (Ig) domain. Details of KIR molecules and of MHC allotypes recognized by them are gradually being revealed. As apparent from the fact that NK cells exert cytotoxic activity against cells with decreased MHC class I molecule expression, most KIR molecules recognize the polymorphic classic MHC class I molecules expressed in almost all types of cells, to transmit negative signals into cells. Meanwhile, some KIR molecular species have non-classic MHC class I molecules as ligands. Accordingly, it is thought that the whole KIR molecule group plays a central role in the surveillance mechanism against autologous but abnormal cells in the natural immune system.

10

15

**[0008]** Like Ly49, KIR molecules also have in the cytoplasmic domain a functional sequence comprising the V/I XYXX L/V (V, valine; I, isoleucine; Y, tyrosine; L, leucine; and X, arbitrary amino acid) motif called "ITIM". When the tyrosine residue in the ITIM motif is phosphorylated, the protein tyrosine dephosphorylation enzyme SHP-1, which comprises a SH2 domain, binds to this site. The activation of NK cells is suppressed by the SHP-1-mediated dephosphorylation of tyrosine-phosphorylated protein (SLP-76 molecule is believed to be a promising candidate) that is essential for the activation of the positive signal cascade in NK cells.

20

**[0009]** The present inventors conducted dedicated studies to solve the above-described objectives. Referring to the analysis data of Serial Analysis of Gene Expression (SAGE) (Hashimoto S, Blood (2003), 101, 3509-3513) for purified immune cells (monocytes/macrophages, T cells, dendritic cells, natural killer cells, neutrophils, etc.) disclosed at <http://www.prevent.m.u-tokyo.ac.jp>, many tags presumed to derive from unknown genes, some with characteristic expression profiles, were identified in addition to sequence tags for known gene sequences.

25

**[0010]** Of these, the present inventors succeeded in identifying the NKIR gene expressed specifically in natural killer (NK) cells. Some characteristics of its deduced amino acid sequence suggested that the gene was a homologue of killer cell inhibitory receptor (hereinafter abbreviated as "KIR") that had been identified as a molecule that expresses in NK cells and a subset of T cells, and functions as an inhibitory regulator molecule for the cytotoxic activity of lymphocyte populations. The NKIR gene was mapped on chromosome 1, while most members belonging to the KIR molecule group are clustered on chromosome 19. The present inventors also analyzed characteristics of the gene to explore the possibility of applying the gene to drug discovery. Some sequences considered to be of splicing variants of the gene of the present invention (WO 01/49728) have been reported. The NCBI Annotation Project has also assigned a splicing variant (LOC343413) through predictions based on genomic sequences. However, the functions of these splicing variants still remain unknown. In addition, to date there is no known sequence perfectly identical to the gene of the present invention.

30

35

**[0011]** As described above, NK cells have anti-tumor and antiviral activities. Therefore, the anti-tumor and antiviral effects based on suppression of KIR function using an antagonistic antibody for KIR are two potential clinical applications of the present invention. Alternatively, when NKT cells are targeted, the present invention is expected to have potential uses against viral diseases such as hepatitis C or cancers by using antagonistic antibodies, or against allergic diseases, asthma, and autoimmune diseases by using agonistic antibodies.

40

**[0012]** The present invention relates to novel proteins expressed in NK cells, DNAs encoding the proteins, methods for producing the molecules, and uses thereof. Specifically, the present invention relates to:

45

[1] a DNA of any one of:

- (a) a DNA encoding a protein comprising the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6;
- (b) a DNA comprising a coding region of the nucleotide sequence of any one of SEQ ID NOs: 1, 3, and 5;
- (c) a DNA encoding a protein which comprises an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6, and which is functionally equivalent to a protein comprising the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6; and
- (d) a DNA which hybridizes under stringent conditions to a DNA comprising the nucleotide sequence of any one of SEQ ID NOs: 1, 3, and 5;

50

55

[2] a DNA encoding a fragment of a protein comprising the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6;

[3] a protein encoded by the DNA of [1] or [2];

## EP 1 672 065 A1

[4] a vector comprising the DNA of [1] or [2];

[5] a host cell comprising the DNA of [1] or [2], or the vector of [4];

[6] a method for producing the protein of [3], wherein the method comprises the steps of:

- 5           culturing the host cell of [5], and  
          collecting the protein from the host cell or a culture supernatant thereof;

[7] an antibody which binds to the protein of [3];

10           [8] a method for identifying a ligand for the protein of [3], wherein the method comprises the steps of:

- (a) contacting a candidate compound with the protein of [3] or a cell expressing the protein of [3]; and  
          (b) determining whether the candidate compound binds to the protein of [3] or the cell expressing the protein of [3];

15           [9] a method for identifying an agonist for the protein of [3], wherein the method comprises the steps of:

- (a) contacting a candidate compound with a cell expressing the protein of [3]; and  
          (b) determining whether the candidate compound generates a signal that is an indicator of activation of the protein of [3];

20           [10] a method for identifying an antagonist for the protein of [3], wherein the method comprises the steps of:

- (a) contacting a candidate compound with a cell expressing the protein of [3]; and  
          (b) determining whether a signal as an indicator of activation of the protein of [3] is reduced as compared with a detection result obtained in absence of the candidate compound;

25           [11] a ligand for the protein of [3], which can be identified by the method of [8];  
          [12] an agonist for the protein of [3], which can be identified by the method of [9];  
          [13] an antagonist for the protein of [3], which can be identified by the method of [10];  
          [14] a kit to be used in the method of any one of [8] to [10], wherein the kit comprises at least one of:

- 30           (a) the protein of [3]; and  
          (b) the host cell of [5];

35           [15] an immunosuppressant, which comprises as an active ingredient an agonist for the protein of [3] (or the agonist of [12]);

          [16] a therapeutic agent for an allergic disease or autoimmune disease, wherein the method comprises as an active ingredient an agonist for the protein of [3] (or the agonist of [12]);

          [17] an immunopotentiator, which comprises as an active ingredient an antagonist for the protein of [3] (or the antagonist of [13]);

40           [18] an anti-tumor or antiviral agent, which comprises as an active ingredient an antagonist for the protein of [3] (or the antagonist of [13]);

          [19] a DNA, which comprises at least 15 nucleotides and which is complementary to the DNA of [1] or [2], or to a complementary strand thereof; and

45           [20] a diagnostic reagent for a disease associated with an abnormality in the expression or activity of a gene encoding the protein of [3], wherein the diagnostic reagent comprises the DNA of [19] or the antibody of [7].

**[0013]** In an alternative embodiment, the present invention provides:

- 50           (a) use of the proteins of the present invention in screening for an agonist or antagonist for the proteins;  
          (b) use of an agonist for the proteins of the present invention in the production of immunosuppressants;  
          (c) use of an agonist for the proteins of the present invention in the production of therapeutic agents for allergic diseases or autoimmune diseases;  
          (d) use of an antagonist for the proteins of the present invention in the production of immunopotentiators; and  
55           (e) use of an antagonist for the proteins of the present invention in the production of anti-tumor or antiviral agents.

**[0014]** The present invention also relates to the following methods:

- (a) a method for suppressing immunological function, which comprises the step of administering an agonist for the

proteins of the present invention to subjects (patients or the like);  
 (b) a method for treating (or preventing) allergic or autoimmune diseases, which comprises the step of administering an agonist for the proteins of the present invention to subjects (patients or the like);  
 (c) a method for enhancing immunological function, which comprises the step of administering an antagonist for the proteins of the present invention to subjects (patients or the like); and  
 (d) a method for treating (or preventing) cancers (tumors) or viral diseases, which comprises the step of administering an antagonist for the protein of the present invention to subjects (patients or the like).

Brief Description of the Drawings

**[0015]**

Fig. 1 is a photograph showing an SDS-PAGE electrophoresis analysis result for NKIR fusion protein purified using a His trap column.  
 Fig. 2 is a photograph showing the detection of NKIR protein using an anti-NKIR polyclonal antibody.  
 Fig. 3 is a photograph showing an RT-PCR analysis result for tissue expression. Fig. 3A shows MTC panels I and II; and Fig. 3B shows Immune System and Blood Fraction.

Each lane is as follows:

**[0016]**

Fig. 3A:  
 Lane 1, heart; lane 2, brain; lane 3, placenta; lane 4, lung; lane 5, liver; lane 6, skeletal muscle; lane 7, kidney; lane 8, pancreas; lane 9, spleen; lane 10, thymus; lane 11, prostate; lane 12, testis; lane 13, ovary; lane 14, small intestine; lane 15, large intestine; and lane 16, peripheral blood leukocyte.  
 Fig. 3B:  
 Lane 1, spleen; lane 2, lymph nodes; lane 3, thymus; lane 4, peripheral blood leukocytes; lane 5, tonsilla; lane 6, fetal liver; lane 7, bone marrow; lane 8, mononuclear cells; lane 9, resting CD8+ cells; lane 10, resting CD4+ cells; lane 11, resting CD14+ cells; lane 12, resting CD19+ cells; lane 13, activated CD19+ cells; lane 14, activated mononuclear cells; lane 15, activated CD4+ cells; and lane 16, activated CD8+ cells.  
 Fig. 4 is a photograph showing the detection of natural NKIR protein expressed in NK-92 cell line using an anti-NKIR polyclonal antibody.  
 Fig. 5 is a graph showing flow cytometric analysis of NK-92 cell line using an anti-NKIR polyclonal antibody.  
 Fig. 6 shows a flow cytometric analysis for the transient expression of the NKIR gene in the COS-7 cell line.  
 Fig. 7 shows a result of a BLAST search using human NKIR as a query, which is aligned with a matching mouse chromosomal sequence.  
 Fig. 8 shows the alignment of human and mouse NKIR sequences.  
 Fig. 9 is a graph showing a flow cytometric analysis of each transformant cell line.  
 Fig. 10 is a histogram showing an assay result for ITIM activity determined using luciferase activity as an indicator.  
 Fig. 11-1 is a diagram showing CD8 chimeric structures of three clones selected in Example 16.  
 Fig. 11-2 is a graph showing a FACS analysis result for clones as described below using anti-CD8 antibody, LT8 (Serotec), and FITC-conjugated goat anti-mouse IgG antibody (Coulter) used in Example 17. Fig. 11-2A, F11 clone in the absence of LT8 (primary antibody); Fig. 11-2B, F11 clone in the presence of LT8 (primary antibody); Fig. 11-2C, CD8 chimeric clone in the absence of LT8 (primary antibody); and Fig. 11-2D, CD8 chimeric clone in the presence of (primary antibody).  
 Fig. 12 is a histogram showing a result of a determination of ITIM function activity using the reporter assay described in Example 17. LT8, anti-CD8 antibody (LT8); and CL, cross-linker (rat anti-mouse IgG1 antibody).

Best Mode for Carrying Out the Invention

**[0017]** The present inventors cloned genes specifically expressed in NK cells from a human spleen cDNA library using PCR. The yielded clones included multiple clones that were thought to be splicing variants. Hence, the inventors carried out 5' RACE to obtain the native sequence, and identified NKIR gene that was thought to belong to the KIR family. The nucleotide sequence of the gene is shown in SEQ ID NO: 1, and the amino acid sequence of the protein encoded by the gene is shown in SEQ ID NO: 2. Furthermore, the present inventors recloned the full-length NKIR gene by 5'- and 3'-RACE methods using total RNAs prepared from NK-92 cell line, and obtained a clone comprising a sequence that had a 36-nucleotide signal-like sequence at the 5' end and approximately 500-nucleotide sequence extension at the 3'

end. The nucleotide sequence of the clone is shown in SEQ ID NO: 3, and the amino acid sequence of the protein encoded by the nucleotide sequence is shown in SEQ ID NO: 4. Furthermore, the present inventors carried out cloning of mouse NKIR sequence. The nucleotide sequence thus obtained is shown in SEQ ID NO: 5, and the amino acid sequence of the protein encoded by the nucleotide sequence is shown in SEQ ID NO: 6.

**[0018]** The present invention provides novel proteins expressed in NK cells, and DNAs encoding the proteins. In a preferred embodiment, DNAs of the present invention are as follows:

- (a) a DNA encoding a protein comprising the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6;
- (b) a DNA comprising a coding region of the nucleotide sequence of any one of SEQ ID NOs: 1, 3, and 5;
- (c) a DNA encoding a protein which comprises an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid of any one of SEQ ID NOs: 2, 4, and 6, and which is functionally equivalent to a protein comprising the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6; and
- (d) a DNA which hybridizes under stringent conditions to a DNA comprising the nucleotide sequence of any one of SEQ ID NOs: 1, 3, and 5

(more preferably, a DNA which hybridizes under stringent conditions to a DNA comprising the nucleotide sequence of any one of SEQ ID NOs: 1, 3, and 5, and which encodes a protein functionally equivalent to a protein comprising the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6).

**[0019]** The present invention includes proteins that are functionally equivalent to the proteins expressed in NK cells (proteins comprising the amino acid sequence of SEQ ID NO: 2, 4, or 6; hereinafter sometimes referred to as "NKIR protein"), which were identified by the present inventors. Such proteins include, for example, mutants of the present proteins and homologues thereof derived from species other than human and mouse. Herein, the phrase "functionally equivalent" means that a protein of interest has a biological or biochemical activity identical to that of NKIR protein. Such activities include, for example: activities that KIR has; the activity of suppressing the cytotoxic activity of NK cells; and the activity of a KIR molecule to suppress activated signaling in immunocytes that have similar intracellular signaling for activation, such as T cell and mast cell hosts, into which the KIR molecule is introduced. Therefore, whether a protein of interest has a biological or biochemical activity equivalent to that of a protein identified by present inventors can be evaluated by, for example, a method that detects the inhibitory activity against the activation signaling of immunocytes to which the molecule (protein of interest) has been introduced by transduction and such, by measuring the cytoplasmic calcium concentration (Bruhns P., Marchetti P., Fridman WH., Vivier E., Daeron M., *J Immunol.* (1999), 162, 3168-3175), or by assaying the expression level of a reporter gene comprising an NF-AT cis sequence under the regulation of the calcineurin cascade (Fry AM., Lanier LL., Weiss A., *J Exp Med.* (1996), 184, 295-300). In addition, when mast cells are used as host cells, a method that assays <sup>3</sup>H-labeled serotonin to detect release of serotonin downstream of Ca cascade (Blery M., Delon J., Trautmann A., Cambiaggi A., Olcese L., Biassoni R., Moretta L., Chavrier P., Moretta A., Daeron M., Vivier E., *J Biol Chem.* (1997), 272, 8989-8996) can be used.

**[0020]** To prepare a protein functionally equivalent to another protein, for example, methods for introducing mutations into amino acids in proteins are well known to those skilled in the art. Specifically, those skilled in the art can prepare a protein functionally equivalent to proteins comprising the amino acid sequence of SEQ ID NO: 2, 4, or 6 by introducing an appropriate mutation into an amino acid(s) of the sequences using site-directed mutagenesis (Hashimoto-Gotoh T. *et al.* *Gene* 152: 271-275(1995); Zoller M.J. and Smith M. *Methods Enzymol.* 100: 468-500(1983); Kramer W. *et al.* *Nucleic Acids Res.* 12: 9441-9456 (1984); Kramer W. and Fritz H.J. *Methods Enzymol.* 154: 350-367(1987); Kunkel T.A. *Proc. Natl. Acad. Sci. USA* 82: 488-492 (1985); Kunkel T.A. *Methods Enzymol.* 85: 2763-2766 (1988)). Amino acid mutations may also occur in nature. Thus, both artificially synthesized and naturally occurring proteins comprising an amino acid sequence in which one or more amino acids in the sequence of the NKIR protein identified by the present inventors are mutated, and which is functionally equivalent to the MC-PIR1 or MC-PIR2 protein, are also included in the present invention. In such a mutant protein, the number of mutated amino acids is usually 50 or less, preferably 30 or less, and more preferably 10 or less (for example, five amino acids or less).

**[0021]** Sites to be mutagenized are not particularly limited, but preferably are those other than motifs and domains as described below.

**[0022]** In mutating an amino acid, it is preferable to change it into another amino acid that allows the properties of the amino acid side chain to be conserved. For example, amino acid side chain characteristics are: side chains having hydrophobic amino acid residues (A, I, L, M, F, P, W, Y, V), hydrophilic residues (R, D, N, C, E, Q, G, H, K, S, T), residues with an aliphatic side chain (G, A, V, L, I, P), residues with a side chain containing a hydroxyl group (S, T, Y), residues with a side chain containing sulfur (C, M), residues with a side chain containing a carboxylic acid and amide group (D, N, E, Q), basic residues (R, K, H), and aromatic residues (H, F, Y, W) (amino acids are shown using the one letter code in the parentheses).

**[0023]** It is already known that a protein having a modified amino acid sequence, in which one or more amino acids are deleted, added, and/or substituted with another amino acid, can maintain the original biological activity (Mark D.F.

*et al.* Proc. Natl. Acad. Sci. USA 81: 5662-5666 (1984); Zoller M.J. and Smith M. Nucleic Acids Res. 10: 6487-6500 (1982); Wang A. *et al.* Science 224: 1431-1433; Dalbadie-McFarland G. *et al.* Proc. Natl. Acad. Sci. USA 79: 6409-6413 (1982)).

5 [0024] A protein comprising an amino acid sequence in which multiple amino acid residues are added to the sequence of NKIR protein includes fusion proteins comprising the protein. Fusion proteins such as those between the proteins of this invention and other peptides or proteins are included in the present invention. To produce a fusion protein, a DNA encoding the NKIR protein (comprising the amino acid sequence according to SEQ ID NO: 2, 4, or 6) and a DNA encoding another peptide or protein are ligated so that their frames match, and introduced into an expression vector to express in a host. Any method commonly known to those skilled in the art can be used. Any peptide or protein may be used for making fusion proteins with proteins of this invention.

10 [0025] Known peptides that can be used as peptides that are fused to the proteins of the present invention include, for example, FLAG (Hopp, T. P. *et al.*, Biotechnology (1988) 6, 1204-1210), 6xHis containing six histidine (HIS) residues, 10xHis, HA (Influenza agglutinin), human c-myc fragment, VSP-GP fragment, p18HIV fragment, T7-tag, HSV-tag, E-tag, SV40T antigen fragment, Ick tag,  $\alpha$ -tubulin fragment, B-tag, Protein C fragment, and the like. Examples of proteins that may be fused to proteins of the present invention include GST (glutathione-S-transferase), HA (Influenza agglutinin), immunoglobulin constant region,  $\beta$ -galactosidase, MBP (maltose-binding protein), and such. Fusion proteins can be prepared by fusing commercially available DNA encoding the fusion peptides or proteins discussed above, with the DNA encoding the proteins of the present invention, and expressing the prepared fused DNA.

15 [0026] An alternative method known in the art to isolate functionally equivalent proteins is, for example, the method using hybridization (Sambrook, J. *et al.*, Molecular Cloning 2nd ed. 9.47-9.58, Cold Spring Harbor Lab. Press, 1989). One skilled in the art can readily isolate a DNA having high homology with an entire or partial DNA sequence (SEQ ID NO: 1, 3, or 5) that encodes the NKIR protein from homogenous or heterogenous organism-derived DNA samples, and isolate proteins functionally equivalent to the NKIR protein using the isolated DNA.

20 [0027] The present invention includes proteins encoded by DNA that hybridize with DNA encoding the NKIR protein, and which are functionally equivalent to the NKIR protein. Such proteins include, for example, homologues in humans, mice, or other mammals (for example, a protein encoded by a rat, rabbit, bovine, or simian homologous gene).

25 [0028] The conditions for hybridization used for isolating a DNA encoding a protein functionally equivalent to the NKIR protein can be appropriately selected by those skilled in the art. For example, low stringent conditions may be used for hybridization. Low stringent conditions are post-hybridization washing in 0.1x SSC, 0.1% SDS at 42°C, for example, and preferably in 0.1x SSC, 0.1% SDS at 50°C. Highly stringent conditions are more preferable, which are washing in 5x SSC, 0.1% SDS at 65°C, for example. Under these conditions, a DNA having a higher homology can be efficiently obtained by increasing the temperature. Multiple factors including the temperature, salt concentration, and such are considered to affect the stringency of hybridization; one skilled in the art can achieve similar stringencies by appropriately selecting these factors.

30 [0029] Furthermore, by using a gene amplification technique (PCR)(Current protocols in Molecular Biology edit. Ausubel *et al.* (1987) Publish. John Wiley & Sons Section 6.1-6.4) in place of hybridization, DNA fragments that are highly homologous to a DNA sequence (SEQ ID NO: 1, 3, or 5) encoding a protein identified by the present inventors can be isolated using primers designed based on portions of the DNA sequence encoding the protein identified by the present inventors, to obtain a protein that is functionally equivalent to the protein identified by the present inventors based on the DNA fragment.

35 [0030] Proteins of the present invention may be "mature" proteins or portions of larger proteins, such as fusion proteins. The proteins of the present invention may comprise a leader sequence, a pro-sequence, a sequence useful for purification, such as multiple histidine residues, or a sequence attached to ensure stability during recombinant production.

40 [0031] Normally, such a protein encoded by the DNA isolated using the above hybridization techniques or gene amplification, and which is functionally equivalent to the NKIR protein, has a high homology with the protein (comprising the amino acid sequence of SEQ ID NO: 2, 4, or 6) at the amino acid level. The proteins of this invention include proteins functionally equivalent to the NKIR protein, and having a high homology with the protein at the amino acid level. High homology normally means an identity of at least 50% or more at the amino acid level, preferably 75% or more, more preferably 85% or more, and most preferably 95% or more. Homology between proteins can be determined according to the algorithm described in literature (Wilbur W. J. and Lipman D. J. Proc. Natl. Acad. Sci. USA 80: 726-730 (1983)).

45 [0032] The identity of amino acid sequences can be determined, for example, using the BLAST algorithm of Karlin and Altschul (Proc. Natl. Acad. Sci. USA 87:2264-2268, 1990; and Proc. Natl. Acad. Sci. USA 90:5873-5877, 1993). Based on this algorithm, a program called BLASTX has been developed (Altschul *et al.* J. Mol. Biol. 215: 403-410, 1990). When amino acid sequences are analyzed using BLASTX, parameters are set, for example, as follows: score = 50 and wordlength = 3. When BLAST and Gapped BLAST programs are used, default parameters of each program may be used. The specific procedures of these analytic methods are known (<http://www.ncbi.nlm.nih.gov>).

50 [0033] The proteins of the present invention may have variations in the amino acid sequence, molecular weight, isoelectric point, or presence or composition of sugar chains, depending on the cell or host used for producing it, or the

method of purification, as described later on. Nevertheless, such proteins are included in the present invention as long as they are functionally equivalent to the proteins of the present invention.

5 [0034] The proteins of the present invention can be prepared as recombinant proteins or natural proteins, by methods well known to those skilled in the art. A recombinant protein can be prepared, for example, by: inserting a DNA that encodes a protein of the present invention (for example, the DNA comprising the nucleotide sequence of SEQ ID NO: 1, 3, or 5), into an appropriate expression vector; introducing the vector into an appropriate host cell; collecting the thus obtained recombinants; obtaining an extract thereof; and purifying the protein by subjecting the extract to a chromatography. Examples of chromatographies are ion exchange chromatography, reverse phase chromatography, gel filtration, or affinity chromatography utilizing a column to which antibodies against proteins of the present invention are immobilized, or combinations of more than one of the aforementioned columns.

10 [0035] When the proteins of the present invention are expressed within host cells (for example, animal cells or *E. coli*) as fusion proteins with the glutathione-S-transferase protein, or as a recombinant protein supplemented with multiple histidines, the expressed recombinant protein can be purified using a glutathione column or nickel column. After purifying the fusion protein, it is also possible to exclude regions other than the objective protein by cutting with thrombin or factor-Xa as required.

15 [0036] A natural protein may be isolated by a method known to those skilled in the art, for example, through purification by applying tissues or cell extracts expressing proteins of this invention onto an affinity column in which an antibody (described below) capable of binding to the proteins has been immobilized. Both monoclonal and polyclonal antibodies can be used.

20 [0037] The present invention also provides fragments (partial peptides) of the proteins of the present invention. Such fragments are proteins that comprise an amino acid sequence identical to a partial but not full-length amino acid sequence of the above-described proteins of the present invention. The fragments of the proteins of the present invention are typically protein fragments consisting of a sequence of 8 or more amino acid residues, preferably 12 or more amino acid residues (for example, 15 or more amino acid residues). The preferred fragment includes, for example, truncated polypeptides that comprise an amino acid sequence with deletion of either or both of stretches of amino or carboxy terminal residues. Fragments having structural or functional features are also preferred, such as those comprising an  $\alpha$  helix and an  $\alpha$  helix-forming region, a  $\beta$  sheet and a  $\beta$  sheet-forming region, a turn and a turn-forming region, a coil and a coil-forming region, a hydrophilic region, a hydrophobic region, an  $\alpha$  amphipathic region, a  $\beta$  amphipathic region, a variable region, a surface-forming region, a substrate-binding region, or a highly antigenic region. Biologically active fragments are also preferred. Such biologically active fragments are those having the activity of the proteins of the present invention, which include fragments having similar activity, fragments with enhanced activity, and fragments in which unfavorable activities have been decreased. Such fragments include, for example, those having the activity of initiating intracellular signaling in response to ligand binding. The fragments also include those exhibiting antigenicity or immunogenicity in animals, in particular humans. Preferably, such a protein fragment has a biological and biochemical activity, including antigenic activity, of a protein of the present invention. Mutants of the identified sequences and fragments are also included in the present invention. The preferred mutants are those that are different from the original sequence due to substitution between amino acids belonging to the same type, i.e., those in which a certain residue has been substituted by an alternative residue having similar properties. Such substitutions are typically those between Ala, Val, Leu, and Ile; Ser and Thr; the acidic residues Asp and Glu; Asn and Gln; the basic residues Lys and Arg; and the aromatic residues Phe and Tyr.

35 [0038] The above-described protein fragments of the present invention are not limited to any particular fragments. However, the fragments preferably comprise at least a motif or domain as described below.

40 [0039] The above-described fragments may be useful, for example, for preparing antibodies against the proteins of this invention, in screenings for compounds that can be ligands that bind to the proteins, and in screenings for agonists or antagonists of the proteins.

45 [0040] The protein fragments (partial peptides) of this invention can be produced using genetic engineering, by commonly known peptide synthesis methods, or digesting proteins of this invention with appropriate peptidases. Synthesis of protein fragments (partial peptides) may be performed, for example, by either solid phase synthesis or liquid phase synthesis.

50 [0041] Herein, the term "ligands" refers to molecules that bind to the proteins of the present invention. The ligands include natural and synthetic ligands. The term "agonists" refers to molecules that bind to and activate the proteins of the present invention; and "antagonists" refers to molecules that inhibit the activation of the proteins of the present invention.

55 [0042] A DNA encoding a protein of the present invention would be useful not only for producing the protein *in vivo* or *in vitro* as described above, but also for applications in gene therapy of a disease caused by an abnormal function of the gene encoding the protein or a disease that can be treated with the protein, DNA diagnostics, etc. DNAs of this invention can take any form as long as they encode proteins of this invention. cDNAs synthesized from mRNAs, genomic DNAs, and chemically-synthesized DNAs can be used. In addition, DNAs comprising any nucleotide sequence based

on the degeneracy of genetic code are included as long as they encode proteins of this invention.

**[0043]** DNAs of this invention can be prepared by methods commonly known to those skilled in the art. For example, they may be prepared by making a cDNA library from cells expressing proteins of this invention, and performing hybridization using a partial nucleotide sequence of the DNA (for example, SEQ ID NO: 1, 3, or 5) as a probe. The cDNA library may be prepared, for example, according to the method described in literature (Sambrook J. *et al.* Molecular Cloning, Cold Spring Harbor Laboratory Press (1989)), or obtained from a commercial source. Alternatively, the DNAs of this invention may be prepared as follows: RNA is prepared from cells expressing proteins of this invention, from which cDNA is synthesized using reverse transcriptase. Then, oligo DNA is synthesized based on the sequence of the DNA (for example, SEQ ID NO: 1, 3, or 5), and used as a primer in a PCR reaction to amplify a cDNA encoding a protein of this invention.

**[0044]** Furthermore, the coding region of the cDNA can be determined by determining the nucleotide sequence of the obtained cDNA, and the amino acid sequences of proteins of this invention can be thus obtained. In addition, the obtained cDNA may be used as a probe for screening a genomic DNA library to isolate a genomic DNA.

**[0045]** Specific procedures are as follows: First, mRNA is isolated from a cell, tissue, or organ expressing a protein of this invention (for example, NK cells or tissues in which expression was detected in the Example below). mRNA may be isolated by preparing total RNA using a commonly known method such as guanidine ultracentrifugation (Chirgwin J. M. *et al.* Biochemistry 18: 5294-5299 (1979)), or AGPC method (Chomczynski P. and Sacchi N. Anal. Biochem. 162: 156-159 (1987)), and then purifying mRNA from total RNA using an mRNA Purification Kit (Pharmacia), etc. Alternatively, mRNA may be directly prepared using the QuickPrep mRNA Purification Kit (Pharmacia).

**[0046]** The obtained mRNA is used to synthesize cDNA using reverse transcriptase. cDNA may be synthesized by using a commercially available kit such as the AMV Reverse Transcriptase First-strand cDNA Synthesis Kit (Seikagaku Kogyo). Alternatively, using a primer described herein, or such, cDNA may be synthesized and amplified following the 5'-RACE method (Frohman M. A. *et al.* Proc. Natl. Acad. Sci. U.S.A. 85:8998-9002 (1988); Belyavsky A. *et al.* Nucleic Acids Res. 17:2919-2932 (1989)) using the 5'-Ampli FINDER RACE Kit (Clontech) and the polymerase chain reaction (PCR). A desired DNA fragment is prepared from PCR products and ligated with a vector DNA to produce a recombinant vector construct. This construct is used to transform *E. coli* or such, and desired recombinant vectors are prepared from a selected colony/colonies. The nucleotide sequence of the desired DNA can be verified by conventional methods such as dideoxynucleotide chain termination.

**[0047]** The nucleotide sequences of DNAs of the present invention may be designed to be expressed more efficiently by taking into account the frequency of codon usage in the host used for the expression (Grantham R. *et al.* Nucleic Acids Res. 9: r43-74 (1981)). The DNAs of the present invention may be altered by a commercially available kit or a conventional method. For instance, the DNAs may be altered by digestion with restriction enzymes, insertion of a synthetic oligonucleotide or an appropriate DNA fragment, addition of a linker, or insertion of an initiation codon (ATG) and/or a stop codon (TAA, TGA, or TAG).

**[0048]** In addition, the DNAs of this invention include a DNA that hybridizes with a DNA comprising the nucleotide sequence shown as SEQ ID NO: 1, 3, or 5 and encodes a protein functionally equivalent to an above-described protein of this invention. Hybridization conditions may be appropriately chosen by one skilled in the art. Specifically, the above-described specific conditions may be used. Under these conditions, the higher the temperature, the higher the homology of the obtained DNA would be. The above hybridizing DNA is preferably a naturally occurring DNA, for example, a cDNA or genomic DNA.

**[0049]** The present invention also provides vectors into which a DNA of the present invention has been inserted. Vectors of the present invention are useful to maintain DNAs of the present invention in a host cell, or to express proteins of the present invention.

**[0050]** When the host cell is *E. coli* and a vector of the present invention is amplified and produced in a large amount in *E. coli* (e.g., JM109, DH5 $\alpha$ , HB101, or XL1Blue), the vector should have "ori" to be amplified in *E. coli* and a marker gene for selecting transformed *E. coli* (e.g., a drug-resistance gene selected by a drug such as ampicillin, tetracycline, kanamycin, chloramphenicol, or the like). For example, M13-series vectors, pUC-series vectors, pBR322, pBluescript, pCR-Script, etc. can be used. In addition to the vectors described above, pGEM-T, pDIRECT, and pT7 can also be used for subcloning and extracting cDNA. When a vector is used to produce a protein of the present invention, an expression vector is especially useful. For example, an expression vector to be expressed in *E. coli* should have the above characteristics to be amplified in *E. coli*. When *E. coli* such as JM109, DH5 $\alpha$ , HB101, or XL1 Blue are used as a host cell, the vector should have a promoter, for example, the lacZ promoter (Ward *et al.*, Nature (1989) 341, 544-546; FASEB J (1992) 6, 2422-2427), araB promoter (Better *et al.*, Science (1988) 240, 1041-1043), or T7 promoter or the like, that can efficiently express the desired gene in *E. coli*. In this respect, pGEX-5X-1 (Pharmacia), "QIAexpress system" (Qiagen), pEGFP and pET (in this case, the host is preferably BL21 which expresses T7 RNA polymerase), for example, can be used in addition to the above vectors.

**[0051]** Additionally, the vector may also contain a signal sequence for polypeptide secretion. An example of a signal sequence that directs the protein to be secreted to the periplasm of the *E. coli* is the pelB signal sequence (Lei, S. P. *et*

*al.*, J. Bacteriol. (1987) 169, 4379). Means for introducing the vectors into target host cells include, for example, the calcium chloride method and the electroporation method.

**[0052]** In addition to *E. coli*, for example, expression vectors derived from mammals (for example, pcDNA3 (Invitrogen) and pEGF-BOS (Nucleic Acids. Res. 1990, 18 (17), p5322), pEF, pCDM8), expression vectors derived from insect cells (for example, "Bac-to-BAC baculovirus expression system" (GIBCO BRL), pBacPAK8), from plants (for example pMH1, pMH2), from animal viruses (for example, pHSV, pMV, pAdexLcw), from retroviruses (for example, pZIPneo), from yeast (for example, "Pichia Expression Kit" (Invitrogen), pNV11, SP-Q01), and from *Bacillus subtilis* (for example, pPL608, pKTH50) can be used as vectors for producing proteins of the present invention.

**[0053]** For the purpose of expression in animal cells such as CHO, COS, or NIH3T3 cells, the vector should have a promoter necessary for expression in such cells, for example, the SV40 promoter (Mulligan *et al.*, Nature (1979) 277, 108), the MMLV-LTR promoter, the EF1 $\alpha$  promoter (Mizushima *et al.*, Nucleic Acids Res. (1990) 18, 5322), the CMV promoter, and the like, and preferably a marker gene for selecting transformants (for example, a drug resistance gene selected by a drug (e.g., neomycin, G418)). Examples of known vectors with these characteristics include, for example, pMAM, pDR2, pBK-RSV, pBK-CMV, pOPRSV, and pOP 13.

**[0054]** In addition, when the aim is to stably express a gene and at the same time increase the copy number of the gene in cells, one can use the method for introducing, into CHO cells in which the nucleic acid synthesizing pathway is deleted, a vector comprising the complementary DHFR gene (for example pCHO I) and then amplifying this by methotrexate (MTX). Furthermore, when the aim is to transiently express a gene, one can use the method for transfecting a vector comprising a replication origin of SV40 (pcD, etc.) into COS cells comprising the SV40 T antigen expressing gene on the chromosome. The replication origin may also be derived from the polyoma virus, adenovirus, bovine papilloma virus (BPV), and the like. Furthermore, the expression vector may carry, as a selection marker, the aminoglycoside transferase (APH) gene, thymidine kinase (TK) gene, *E. coli* xanthine-guanine-phosphoribosyl transferase (Ecogpt) gene, dihydrofolate reductase (dhfr) gene, and such, for increasing the copy number in the host cell system.

**[0055]** DNAs of the present invention can further be expressed *in vivo* in animals, for example, by inserting the DNAs into an appropriate vector and introducing it into living bodies by methods such as the retrovirus method, the liposome method, the cationic liposome method, and the adenovirus method. Gene therapy against diseases attributed to mutation of genes encoding proteins of the present invention can be thus accomplished. An adenovirus vector (for example pAdexlcw) or retrovirus vector (for example, pZIPneo) can be given as an example of a vector, but the vector is not restricted thereto. General gene manipulations, such as insertion of DNAs of the present invention to a vector, can be performed according to conventional methods (Molecular Cloning, 5. 61-5. 63). Administration into a living body can be either an *ex vivo* method, or *in vivo* method.

**[0056]** The present invention further provides host cells into which vectors of the present invention have been transfected. The host cells into which vectors of the present invention are transfected are not particularly limited. For example, *E. coli*, various animal cells and such can be used. The host cells of the present invention can be used, for example, as a production system for producing or expressing proteins of the present invention. The present invention provides methods for producing proteins of the present invention both *in vitro* and *in vivo*. For *in vitro* production, eukaryotic cells or prokaryotic cells can be used as host cells.

**[0057]** Useful eukaryotic cells may be animal, plant, or fungi cells. Animal cells include, for example, mammalian cells such as CHO (J. Exp. Med. 108:945 (1995)), COS, 3T3, myeloma, baby hamster kidney (BHK), HeLa, or Vero cells; amphibian cells such as *Xenopus* oocytes (Valle *et al.* Nature 291:340-358 (1981)); or insect cells such as Sf9, Sf21, or Tn5 cells. CHO cells lacking the DHFR gene (dhfr-CHO) (Proc. Natl. Acad. Sci. U.S.A. 77:4216-4220 (1980)) or CHO K-1 (Proc. Natl. Acad. Sci. U.S.A. 60:1275 (1968)) may also be used. Of animal cells, CHO cells are particularly preferable for large scale expression. A vector can be transfected into host cells by, for example, the calcium phosphate method, the DEAE-dextran method, the cationic liposome DOTAP (Boehringer Mannheim), the electroporation method, the lipofection method, and such.

**[0058]** As plant cells, plant cells derived from *Nicotiana tabacum* are known as protein-production systems, and may be used as callus cultures. As fungi cells, yeast cells such as *Saccharomyces*, including *Saccharomyces cerevisiae*, or filamentous fungi such as *Aspergillus*, including *Aspergillus niger*, are known and may be used herein.

**[0059]** Useful prokaryotic cells include bacterial cells such as *E. coli*, for example, JM109, DH5 $\alpha$ , and HB101. Other bacterial systems include *Bacillus subtilis*.

**[0060]** These cells are transformed by a desired DNA, and the resulting transformants are cultured *in vitro* to obtain the protein. Transformants can be cultured using known methods. Culture medium for animal cells include, for example, DMEM, MEM, RPMI 1640, and IMDM. These may be used with or without a serum supplement such as the fetal calf serum (FCS). The pH of the culture medium is preferably between about 6 and 8. Such cells are typically cultured at about 30 to 40°C for about 15 to 200 hr, and the culture medium may be replaced, aerated, or stirred if necessary.

**[0061]** Animal or plant hosts may be used for the *in vivo* production. For example, a desired DNA can be transfected into an animal or plant host. Encoded proteins are produced *in vivo*, and then recovered. These animal and plant hosts are included in the host cells of the present invention.

**[0062]** Animals used for the production system described above include, but are not limited to, mammals and insects. Mammals such as goats, pigs, sheep, mice, and cows may be used (Vicki Glaser, SPECTRUM Biotechnology Applications (1993)). Alternatively, the mammals may be transgenic animals.

5 **[0063]** For instance, a desired DNA may be prepared as a fusion gene, by fusing it with a gene such as the goat  $\beta$  casein gene which encodes a protein specifically produced into milk. DNA fragments comprising the fusion gene are injected into goat embryos, which are then implanted in female goats. Proteins are recovered from milk produced by the transgenic goats (i.e., those born from the goats that had received the embryos) or from their offspring. To increase the amount of milk containing the proteins produced by the transgenic goats, appropriate hormones may be administered to them (Ebert K. M. *et al.* Bio/Technology 12:699-702 (1994)).

10 **[0064]** Alternatively, insects, such as the silkworm, may be used. A DNA encoding a desired protein inserted into baculovirus can be used to transfect silkworms, and the desired protein may be recovered from their body fluid (Susumu M. *et al.* Nature 315: 592-594 (1985)).

15 **[0065]** As plants, for example, tobacco can be used. When using tobacco, a DNA encoding a desired protein may be inserted into a plant expression vector such as pMON530, which is introduced into bacteria such as *Agrobacterium tumefaciens*. Then, the bacteria are used to transfect a tobacco plant such as *Nicotiana tabacum*, and a desired polypeptide is recovered from the leaves (Julian K.-C. Ma *et al.*, Eur. J. Immunol. 24: 131-138 (1994)).

**[0066]** Proteins of the present invention obtained as above may be isolated from the inside or outside (such as culture medium) of host cells, and purified as a substantially pure homogeneous protein.

20 **[0067]** The present invention provides methods for producing proteins of the present invention, which comprise the steps of: culturing host cells of the present invention; and collecting thus produced proteins from the host cells or culture supernatant thereof.

25 **[0068]** The method for protein isolation and purification is not limited to any specific method, and any standard method may be used. For instance, column chromatography, filter, ultrafiltration, salt precipitation, solvent precipitation, solvent extraction, distillation, immunoprecipitation, SDS-polyacrylamide gel electrophoresis, isoelectric point electrophoresis, dialysis, and recrystallization may be appropriately selected and combined to isolate and purify the protein.

30 **[0069]** Examples of chromatographies include, for example, affinity chromatography, ion-exchange chromatography, hydrophobic chromatography, gel filtration, reverse phase chromatography, adsorption chromatography, and such (Strategies for Protein Purification and Characterization: A Laboratory Course Manual. Ed. Daniel R. Marshak *et al.*, Cold Spring Harbor Laboratory Press (1996)). These chromatographies may be performed by a liquid chromatography such as HPLC and FPLC. Thus, the present invention provides highly purified proteins prepared by the above methods.

35 **[0070]** Proteins of the present invention may be optionally modified or partially deleted by treating them with an appropriate protein modification enzyme before or after purification. Useful protein modification enzymes include, but are not limited to, trypsin, chymotrypsin, lysylendopeptidase, protein kinase, glucosidase and such.

**[0071]** The present invention provides antibodies that bind to the proteins of the present invention. The antibodies can take any form such as monoclonal or polyclonal antibodies, and includes antiserum obtained by immunizing animals such as a rabbit with proteins of the present invention, all classes of polyclonal and monoclonal antibodies, human antibodies, and humanized antibodies produced by genetic recombination.

40 **[0072]** Proteins of the present invention used as antigens to obtain an antibody may be derived from any animal species, but are preferably derived from a mammal such as a human, mouse, or rat, more preferably a human. A human-derived protein may be obtained from the nucleotide or amino acid sequences disclosed herein.

**[0073]** According to the present invention, the proteins to be used as immunizing antigens may be a complete protein or a partial peptide of the proteins. A partial peptide may comprise, for example, the amino (N)-terminal or carboxy (C)-terminal fragment of proteins of the present invention. Herein, an antibody is defined as a protein that reacts with either the whole proteins of the present invention, or a fragment of the proteins.

45 **[0074]** Genes encoding proteins of the present invention or their fragment may be inserted into a known expression vector, which is then used to transform a host cell as described herein. The desired protein or its fragment may be recovered from the outside or inside of host cells by any standard method, and may subsequently be used as an antigen. Alternatively, cells expressing the protein or their lysates, or a chemically synthesized protein may be used as antigen. In the case of a short peptide, it is preferably bound to an appropriate carrier protein such as keyhole limpet hemocyanin, bovine serum albumin, and ovalbumin before using as antigen.

50 **[0075]** Any mammalian animal may be immunized with the antigen, but preferably, the compatibility with parental cells used for cell fusion is taken into account. In general, animals of Rodentia, Lagomorpha, or Primates are used.

55 **[0076]** Animals of Rodentia include, for example, mice, rats, and hamsters. Animals of Lagomorpha include, for example, rabbits. Animals of Primates include, for example, monkeys of Catarrhini (old world monkeys) such as *Macaca fascicularis*, rhesus monkeys, sacred baboons, and chimpanzees.

**[0077]** Methods for immunizing animals with antigens are known in the art. Intraperitoneal injection or subcutaneous injection of antigens is a standard method of immunization for mammals. More specifically, antigens may be diluted and suspended in an appropriate amount of phosphate buffered saline (PBS), saline, etc. If desired, the antigen suspension

may be mixed with an appropriate amount of a standard adjuvant such as Freund's complete adjuvant, made into an emulsion, and then administered to mammalian animals. Preferably, this is followed by several administrations of antigen mixed with an appropriately amount of Freund's incomplete adjuvant every 4 to 21 days. An appropriate carrier may also be used for immunization. After immunizing as above, serum is examined by a standard method for an increase in the amount of desired antibodies.

**[0078]** Polyclonal antibodies against the proteins of the present invention may be prepared by collecting blood from the immunized mammal after verifying an increase of desired antibodies in the serum, and by separating serum from the blood by any conventional method. Polyclonal antibodies include serum containing polyclonal antibodies, as well as fractions containing the polyclonal antibodies isolated from the serum. Immunoglobulin G or M can be prepared from a fraction which recognizes only the proteins of the present invention using, for example, an affinity column coupled with proteins of the present invention, and further purifying this fraction using a protein A or protein G column.

**[0079]** To prepare monoclonal antibodies, immunocytes are collected from the mammal immunized with the antigen and checked for an increase in the level of desired antibodies in the serum as described above, and are subjected to cell fusion. The immune cells used for cell fusion are preferably obtained from the spleen. Other preferred parental cells to be fused with the above immunocytes include, for example, mammalian myeloma cells, and more preferably myeloma cells having an acquired property for the selection of fused cells by drugs.

**[0080]** The above immunocytes and myeloma cells can be fused according to known methods, for example, the method of Milstein *et al.* (Galfre, G. and Milstein, C., *Methods Enzymol.* (1981) 73, 3-46).

**[0081]** Resulting hybridomas obtained by the cell fusion may be selected by cultivating them in a standard selection medium such as the HAT medium (hypoxanthine, aminopterin, and thymidine containing medium). The cell culture is typically continued in the HAT medium for several days to several weeks, which is sufficient to allow all the other cells, with the exception of the desired hybridoma (non-fused cells), to die. Then, standard limiting dilution is performed to screen and clone a hybridoma producing the desired antibody.

**[0082]** In addition to the above method in which a non-human animal is immunized with an antigen for preparing a hybridoma, a hybridoma producing a desired human antibody that is able to bind to a protein can be obtained by the following method. First, human lymphocytes such as those infected by the EB virus may be immunized with a protein, protein expressing cells, or their lysates *in vitro*. Then, the immunized lymphocytes are fused with human-derived myeloma cells that are capable of indefinite division, such as U266, to yield the desired hybridoma (Unexamined Published Japanese Patent Application No. (JP-A) Sho 63-17688).

**[0083]** The obtained hybridomas are subsequently transplanted into the abdominal cavity of a mouse and the ascites are harvested. The obtained monoclonal antibodies can be purified by, for example, ammonium sulfate precipitation, a protein A or protein G column, DEAE ion exchange chromatography, or an affinity column to which proteins of the present invention are coupled. The antibodies of the present invention can be used not only for purification and detection of the proteins of the present invention, but also as candidates for agonists and antagonists of the proteins. In addition, these antibodies can be applied to the antibody treatment for diseases related to the proteins of the present invention. When an obtained antibody is to be administered to the human body (antibody treatment), a human antibody or a humanized antibody is preferable to reduce immunogenicity.

**[0084]** For example, transgenic animals having a repertoire of human antibody genes may be immunized with an antigen selected from a protein, cells expressing the protein or their lysates. Antibody producing cells are then collected from the animals and fused with myeloma cells to obtain hybridomas, from which human antibodies against the protein can be prepared (see WO92-03918, WO93-2227, WO94-02602, WO94-25585, WO96-33735, and W096-34096).

**[0085]** Alternatively, an immunocyte that produces antibodies, such as an immunized lymphocyte, may be immortalized by an oncogene and used for preparing monoclonal antibodies.

**[0086]** Monoclonal antibodies thus obtained can also be recombinantly prepared using genetic engineering techniques (see, for example, Borrebaeck C. A. K. and Larrick J. W. *Therapeutic Monoclonal Antibodies*, published in the United Kingdom by MacMillan Publishers LTD (1990)). A DNA encoding an antibody may be cloned from an immunocyte, such as a hybridoma or an immunized lymphocyte producing the antibody, inserted into an appropriate vector, and introduced into host cells to prepare a recombinant antibody. The present invention also provides recombinant antibodies prepared as described above.

**[0087]** Furthermore, antibodies of the present invention may be fragments of antibodies or modified antibodies, so long as they bind to one or more of the proteins of the invention. For instance, the antibody fragment may be Fab, F(ab')<sub>2</sub>, Fv, or single chain Fv (scFv), in which Fv fragments from H and L chains are ligated by an appropriate linker (Huston J. S. *et al.* *Proc. Natl. Acad. Sci. U.S.A.* 85:5879-5883 (1988)). More specifically, an antibody fragment may be generated by treating an antibody with an enzyme such as papain or pepsin. Alternatively, a gene encoding the antibody fragment may be constructed, inserted into an expression vector, and expressed in an appropriate host cell (see, for example, Co M. S. *et al.* *J. Immunol.* 152:2968-2976 (1994); Better M. and Horwitz A. H. *Methods Enzymol.* 178:476-496 (1989); Pluckthun A. and Skerra A. *Methods Enzymol.* 178:497-515 (1989); Lamoyi E. *Methods Enzymol.* 121:652-663 (1986); Rousseaoux J. *et al.* *Methods Enzymol.* 121:663-669 (1986); Bird R. E. and Walker B. W. *Trends Biotechnol.* 9:

132-137 (1991)).

**[0088]** An antibody may be modified by conjugation with a variety of molecules, such as polyethylene glycol (PEG). The term "antibodies" of the present invention also comprises such modified antibodies. The modified antibody can be obtained by chemically modifying an antibody. These modification methods are conventional in the field.

**[0089]** Alternatively, antibodies of the present invention may be obtained as chimeric antibodies between a variable region derived from a nonhuman antibody and a constant region derived from a human antibody. It can also be obtained as a humanized antibody comprising a complementarity-determining region (CDR) derived from a nonhuman antibody, a frame work region (FR) and a constant region derived from a human antibody. Such antibodies can be prepared using a known technology.

**[0090]** Antibodies obtained as above may be purified to homogeneity. For example, the separation and purification of the antibody can be performed according to separation and purification methods used for general proteins. For example, the antibody may be separated and isolated by appropriately selecting and combining column chromatographies such as affinity chromatography, filtration, ultrafiltration, salting-out, dialysis, SDS polyacrylamide gel electrophoresis, isoelectric focusing, and others (Antibodies: A Laboratory Manual. Ed Harlow and David Lane, Cold Spring Harbor Laboratory, 1988), but the chromatographies are not limited thereto. The concentration of the thus obtained antibodies can be determined by measuring the absorbance, by an enzyme-linked immunosorbent assay (ELISA), and such.

**[0091]** A protein A column or protein G column can be used as the affinity column. Examples of protein A columns include, for example, Hyper D, POROS, and Sepharose F.F. (Pharmacia).

**[0092]** Examples of chromatographies other than affinity chromatography includes, for example, ion-exchange chromatography, hydrophobic chromatography, gel filtration, reverse-phase chromatography, adsorption chromatography, and the like (Strategies for Protein Purification and Characterization: A Laboratory Course Manual. Ed Daniel R. Marshak *et al.*, Cold Spring Harbor Laboratory Press, 1996). The chromatographies can be carried out by a liquid-phase chromatography, such as HPLC, FPLC.

**[0093]** For example, measurement of absorbance, enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), radioimmunoassay (RIA), and/or immunofluorescence may be used to measure the antigen binding activity of antibodies of the invention. In ELISA, antibodies of the present invention are immobilized on a plate, proteins of the present invention are applied to the plate, and then samples containing a desired antibody, such as culture supernatants of antibody producing cells or purified antibodies, are applied. Then, a secondary antibody that recognizes the primary antibody and is labeled with an enzyme such as alkaline phosphatase is applied, and the plate is incubated. Next, after washing, an enzyme substrate such as *p*-nitrophenyl phosphate is added to the plate, and the absorbance is measured to evaluate the antigen binding activity of the sample. A fragment of the protein, such as a C-terminal fragment may be used as the protein. BIAcore (Pharmacia) may be used to evaluate the activity of antibodies according to the present invention.

**[0094]** These methods allow the detection or measurement of proteins of the present invention by exposing antibodies of the present invention to samples assumed to contain the proteins of the present invention, and detecting or measuring the immune complexes formed by the antibodies and the proteins. Because the methods for detecting or measuring the proteins according to the present invention can specifically detect or measure proteins, the methods may be useful in a variety of experiments in which the proteins of the present invention are used.

**[0095]** Furthermore, the present invention provides DNAs comprising at least 15 nucleotides, which is complementary to DNAs (comprising, for example, the nucleotide sequence according to SEQ ID NO: 1, 3, or 5) of the present invention or a complementary strand thereof.

**[0096]** Herein, "complementary strand" means a strand that is opposite relative to the other strand in a double-stranded nucleic acid composed of A:T (U in the case of RNA) and G:C base pairs. In addition, being "complementary" is not limited to having complete complementarity in a continuous region of at least 15 nucleotides, but it can also mean having a homology of at least 70%, preferably at least 80%, more preferably 90%, and most preferably 95% or higher at the nucleotide level. Homology can be determined using the algorithm described herein.

**[0097]** Such nucleic acids include: probes or primers used for detecting or amplifying DNAs encoding proteins of this invention; probes or primers used for detecting DNA expression; or nucleotides or nucleotide derivatives (for example, antisense oligonucleotides or ribozymes, or DNA encoding them) used for regulating the expression of proteins of this invention. Such nucleic acids may also be useful for preparing DNA chips.

**[0098]** When using as a primer, the 3'-region can be made complementary and a recognition site for a restriction enzyme, or a tag can be attached to the 5'-region.

**[0099]** The present invention also provides diagnostic reagents for diseases associated with abnormalities in the expression of the genes encoding the proteins of the present invention or abnormalities in the activity (function) of the proteins of the present invention.

**[0100]** In one embodiment, the diagnostic reagents comprise a DNA that comprises at least 15 nucleotides and hybridizes to a DNA encoding a protein of the present invention as described above or to a DNA comprising the expression regulatory region. The DNA can be used as: a probe for detecting a gene encoding a protein of the present invention

or a DNA comprising its expression regulatory region; or a primer for amplifying a gene encoding a protein of the present invention or its expression regulatory region in the diagnostic methods of the present invention described above. DNAs of the present invention can be prepared, for example, using commercially available DNA synthesizers. The probe can also be prepared as a double-stranded DNA fragment obtained by restriction enzyme treatment or such. When the DNAs of the present invention are used as probes, they are preferably used after being appropriately labeled. The labeling methods include, for example, those that comprise phosphorylating the 5' end of DNAs with <sup>32</sup>P using T4 polynucleotide kinase and methods (random priming method and such) that comprise the step of incorporating substrate nucleotides labeled with isotopes such as <sup>32</sup>P, fluorescent dye, biotin, or such using DNA polymerase such as Klenow enzyme, and as a primer random hexamer oligonucleotide or such.

**[0101]** In an alternative embodiment, the diagnostic reagents of the present invention comprise the antibodies described below that bind to the proteins of the present invention. The antibodies are used to detect the proteins of the present invention in the above-described diagnostic methods of the present invention. There is no limitation on the type of the antibodies, as long as they can detect the proteins of the present invention. The diagnostic antibodies include polyclonal and monoclonal antibodies. The antibodies may be labeled if required.

**[0102]** The diagnostic reagents described above may comprise, in addition to a DNA or an antibody which is an active ingredient, for example, sterilized water, saline, vegetable oil, detergent, lipid, solubilizing agent, buffering agent, protein stabilizing agent (such as BSA and gelatin), and preservative, if required.

**[0103]** The receptor proteins of the present invention are also useful in identifying their ligands, agonists, and antagonists. The present invention provides (screening) methods for identifying ligands, agonists, and antagonists for the proteins of the present invention using the proteins.

**[0104]** In a preferred embodiment of a method of the present invention for identifying ligands, first, candidate compounds (test samples) are contacted with a protein of the present invention or cells expressing the protein; and second, whether the candidate compounds bind to the protein of the present invention or cells expressing the protein is determined (the binding activity is evaluated). Then, compounds having binding activity are selected as ligands.

**[0105]** Proteins of the present invention used for the identification (screening) methods of the present invention may be recombinant proteins or naturally occurring proteins. They may also be partial peptides. They can be expressed on the cell surface, or contained in the membrane fraction. A candidate compound is not limited to any particular sample; it can be, for example, a cell extract, a cell culture supernatant, a product of a fermentation microorganism, a marine organism extract, a plant extract, a purified or crude protein, a peptide, a non-peptide compound, a synthetic low molecular weight compound, or a natural compound. The proteins of this invention can be contacted with the candidate compound (test sample) as a purified protein, soluble protein, in a form bound to a carrier, as a fusion protein with another protein, in a form expressed on the cell surface, or as a form contained in the membrane fraction.

**[0106]** As methods of screening for proteins that, for example, bind to proteins of the present invention using proteins of the present invention, many methods well known to those skilled in the art can be used. Such a screening can be conducted by, for example, the immunoprecipitation method, specifically, in the following manner. Genes encoding proteins of the present invention are expressed in animal cells or such, by inserting the gene into a foreign gene expression vector, such as pSV2neo, pcDNA 1, and pCD8. The promoters to be used for the expression are not limited as long as they are promoters generally used and include, for example, the SV40 early promoter (Rigby in Williamson (ed.), Genetic Engineering, vol. 3. Academic Press, London, p. 83-141 (1982)), the EF-1 $\alpha$  promoter (Kim *et al.*, Gene 91, p217-223 (1990)), the CAG promoter (Niwa *et al.* Gene 108, p. 193-200 (1991)), the RSV LTR promoter (Cullen Methods in Enzymology 152, p. 684-704 (1987)) the SR $\alpha$  promoter (Takebe *et al.*, Mol. Cell. Biol. 8, p. 466 (1988), the CMV immediate early promoter (Seed and Aruffo Proc. Natl. Acad. Sci. USA 84, p. 3365-3369 (1987)), the SV40 late promoter (Gheysen and Fiers J. Mol. Appl. Genet. 1, p. 385-394 (1982)), the Adenovirus late promoter (Kaufman *et al.*, Mol. Cell. Biol. 9, p. 946 (1989)), the HSV TK promoter, and such.

**[0107]** The introduction of the gene into animal cells to express a foreign gene can be performed according to any method, for example, the electroporation method (Chu G. *et al.* Nucl. Acids Res. 15, 1311-1326 (1987)), the calcium phosphate method (Chen, C and Okayama, H. Mol. Cell. Biol. 7, 2745-2752 (1987)), the DEAE dextran method (Lopata, M. A. *et al.* Nucl. Acids Res. 12, 5707-5717 (1984)), Sussman, D. J. and Milman, G. Mol. Cell. Biol. 4, 1642-1643 (1985)), the Lipofectin method (Derijard, B. Cell 7, 1025-1037 (1994); Lamb, B. T. *et al.* Nature Genetics 5, 22-30 (1993): Rabindran, S. K. *et al.* Science 259, 230-234 (1993)), and such.

**[0108]** Proteins of the present invention can be expressed as fusion proteins comprising a recognition site (epitope) of a monoclonal antibody by introducing, to the N- or C- terminus of the protein, an epitope of a monoclonal antibody whose specificity has been revealed. A commercially available epitope-antibody system can be used (Experimental Medicine 13, 85-90 (1995)). Vectors that can express a fusion protein with, for example,  $\beta$ -galactosidase, maltose-binding protein, glutathione S-transferase, green fluorescence protein (GFP) and such through multiple cloning sites are commercially available.

**[0109]** Method for preparing a fusion protein by introducing only a small epitope portion consisting of several to a dozen amino acids so as to not change, as much as possible, the property of the proteins of the present invention by

the fusion, have also been reported. Epitopes, such as polyhistidine (His-tag), influenza aggregate HA, human c-myc, FLAG, Vesicular stomatitis virus glycoprotein (VSV-GP), T7 gene 10 protein (T7-tag), human herpes simplex virus glycoprotein (HSV-tag), E-tag (an epitope on monoclonal phage), and such, and monoclonal antibodies recognizing them can be used as epitope-antibody systems for screening proteins binding to the proteins of the present invention (Experimental Medicine 13, 85-90 (1995)).

**[0110]** In immunoprecipitation, an immune complex is formed by adding these antibodies to a cell lysate prepared by using an appropriate detergent. The immune complex consists of a protein of the present invention, a protein that can bind to the protein, and an antibody. Immunoprecipitation can also be conducted by using antibodies against proteins of the present invention, besides using antibodies against the above epitopes. Antibodies against proteins of the present invention can be prepared, for example, by introducing a gene encoding the proteins into an appropriate *E. coli* expression vector, expressing the gene in *E. coli*, purifying the expressed protein, and immunizing rabbits, mice, rats, goats, chicken and such with the protein. The antibodies can also be prepared by immunizing animals of above with synthesized partial peptides of the proteins of the present invention.

**[0111]** An immune complex can be precipitated, for example by Protein A Sepharose or Protein G sepharose when the antibody is a mouse IgG antibody. If the proteins of the present invention are prepared as fusion proteins with an epitope such as GST, an immune complex can be formed in the same manner as when using the antibody against a protein of the present invention, by using a substance that specifically binds to the epitope, such as glutathione-Sepharose 4B.

**[0112]** Immunoprecipitation can be performed by following or according to, for example, methods in literature (Harlow, E. and Lane, D.: Antibodies pp. S11-552, Cold Spring Harbor Laboratory publications, New York (1988)).

**[0113]** SDS-PAGE is commonly used for analyzing immunoprecipitated proteins. The bound protein can be analyzed by the molecular weight of the protein using a gel with an appropriate concentration. Since the protein bound to proteins of the present invention is difficult to detect by a common staining method such as Coomassie staining or silver staining, the detection sensitivity of the protein can be improved by culturing cells in a culture medium containing the radioactive isotope <sup>35</sup>S-methionine or <sup>35</sup>S-cysteine, labeling proteins within the cells, and detecting the proteins. Once the molecular weight of the protein has been revealed, the target protein can be purified directly from the SDS-polyacrylamide gel and its sequence can be determined.

**[0114]** In addition, as a method for isolating a protein capable of binding to the protein using a protein of this invention, western blotting may be used (Skolnik E.Y. *et al.* Cell 65: 83-90 (1991)). Specifically, a cDNA library using a phage vector ( $\lambda$ gt11, ZAP, and the like) can be prepared using a cell, tissue, or organ expected to express a protein capable of binding to a protein of this invention (for example, NK cells). The cDNA library can be expressed on LB-agarose, and then the expressed protein can be immobilized onto a filter. Then, a purified and labeled protein of this invention can be incubated with the above filter. Finally, a plaque expressing a protein capable of binding to the protein of this invention can be detected by the label. For labeling a protein of this invention, methods that make use of: the binding between biotin and avidin; an antibody specifically binding the protein of this invention, or a peptide or polypeptide fused to the protein (for example, GST); radioisotopes; or fluorescence, or the like, can be used.

**[0115]** In another embodiment of the above-described identification (screening) methods of this invention, the two-hybrid system using cells may be used (Fields S. and Sternglanz R. Trends Genet. 10: 286-292 (1994); Dalton S. and Treisman R. Characterization of SAP-1, a protein recruited by serum response factor to the c-fos serum response element. Cell 68: 597-612 (1992); "MATCHMAKER Two-Hybrid System"; "Mammalian MATCHMAKER Two-Hybrid Assay Kit"; "MATCHMAKER One-Hybrid System" (all from Clontech); and "HybriZAP Two-Hybrid Vector System" (Stratagene)). In the two-hybrid system, proteins of this invention or a partial peptide may be expressed in yeast cells as a fusion protein with the SRF DNA binding domain, or GAL4 DNA binding domain. A cDNA library in which the protein is expressed as a fusion between the VP16 or GAL4 transcription activation domain is prepared from cells in which a protein capable of binding to proteins of this invention is expected to be present. The library is transfected into yeast cells, and cDNA derived from the library is isolated from a positive clone detected (when a protein capable of binding to the proteins of this invention is expressed in yeast cells, binding of the two proteins activates a reporter gene, which is used to detect a positive clone). Isolated cDNA may be introduced and expressed in *E. coli* to obtain a protein encoded by the cDNA. The reporter gene used in the two-hybrid system may be, for example, a gene such as HIS3, Ade2, LacZ, CAT, luciferase, and PAI-1 (plasminogen activator inhibitor type I), but is not limited thereto. Such a screening using the two-hybrid system may be performed using a mammalian cell other than yeast.

**[0116]** Candidate compounds binding to proteins of the present invention can be screened using affinity chromatography. For example, the proteins of the present invention may be immobilized on a carrier of an affinity column, and a candidate compound presumed to express a protein capable of binding to the proteins of the invention, is applied to the column. Herein, candidate compounds may be, for example, cell extracts, cell lysates, etc. After loading the candidate compound, the column is washed, and proteins bound to the proteins of the present invention can be prepared.

**[0117]** The amino acid sequence of the obtained protein is analyzed, an oligo DNA is synthesized based on the sequence, and cDNA libraries are screened using the oligo DNA as a probe to obtain a DNA encoding the protein.

5 [0118] A biosensor using the surface plasmon resonance phenomenon may be used as a means for detecting or quantifying the bound candidate compound in the present invention. When such a biosensor is used, the interaction between the proteins of the present invention and a candidate compound can be observed real-time as a surface plasmon resonance signal, using only a minute amount of protein and without labeling (for example, BIAcore, Pharmacia). Therefore, it is possible to evaluate the binding between the proteins of the present invention and a candidate compound using a biosensor such as BIAcore.

[0119] Ligands that can be identified by the above-described methods of the present invention are also included in the present invention.

10 [0120] In a preferred embodiment, a method for identifying agonists for a protein of the present invention comprises the steps of: first, contacting candidate compounds with cells expressing the protein of the present invention; and determining whether the candidate compounds generate a signal that is an indicator of activation of the protein of the present invention. Specifically, the method comprise the steps of: incubating candidate compounds with a receptor protein of the present invention; and determining whether the candidate compounds are agonists using as an indicator the presence or absence of a signal generated by the protein of the present invention in response to agonist stimulation.

15 [0121] The present inventors found that immunosuppression was induced by the signal generated by the receptor proteins of the present invention. Thus, the presence or absence of the signal in the present invention can be detected, for example, by determining whether the immunosuppression is induced. An example study is, inducing experimental autoimmune encephalomyelitis (EAE) using a humanized model mouse where the human immune system had been reconstituted\_ (note: mouse and human immune systems are not usually compatible) by transplanting human hematopoietic stem cells into immunodeficient mice (Hiramatsu H, Nishikomori R, Heike T, Ito M, Kobayashi K, Katamura K, and Nakahata T, Blood (2003), 102, 873-880), and then administering candidate agonist substances to such mice to evaluate them for EAE (quantitation of hind leg paralysis, urinary incontinence, and weight loss). When the above-described signal is detected by a method described above, the candidate compound is evaluated to be an agonist. Such agonists that can be identified by the above-described methods are also included in the present invention.

25 [0122] The present invention also provides immunosuppressants comprising an agonist of a protein of the present invention, as an active ingredient. The immunosuppressants of the present invention are expected to be effective in the treatment of, for example, allergic diseases (for example, allergic asthma) and autoimmune diseases. Thus, the present invention provides therapeutic agents for allergic or autoimmune diseases, which comprise an agonist for the proteins of the present invention, as an active ingredient.

30 [0123] In a preferred embodiment, the methods for identifying antagonists for the proteins of the present invention comprise the steps of: first contacting candidate compound(s) with cells expressing a protein of the present invention; and second determining whether the signal which is an indicator of activation of the protein of the present invention is reduced compared to when the candidate compound(s) is absent.

35 [0124] It is thought that, for example, an immunopotential effect (such as antiviral or anti-tumor activity) is induced as a result of a reduction (suppression) of the signal generated by a receptor protein of the present invention. Thus, whether the above-described signal of the present invention is reduced can be determined, for example, by detecting the enhanced immunopotential effect. Examples include: an *in-vitro* study where the cytotoxicity against cells (for example, NK sensitive cell lines such as K-562 cell line) targeted by NK cells is determined and compared in the presence or absence of antagonist candidate substances; and an *in-vivo* study where human cells infected with viruses or human cancer tissues are transplanted into the above-described humanized model mice comprising the reconstructed human immune system, antagonist candidate substances are administered to the mice, and then cytotoxicity against the human cells infected with viruses or cancer tissues transplanted into the mice is evaluated.

40 [0125] When the above-described signal is found to be decreased in the methods as described above, the candidate compound is determined to be an antagonist. Such antagonists that can be identified by the above-described methods are also included in the present invention.

45 [0126] The present invention also provides immunopotentiators comprising an agonist for a protein of the present invention, as an active ingredient. The immunopotentiators of the present invention are expected to be, for example, effective anti-tumor or antiviral agents. Thus, the present invention provides anti-tumor agents and antiviral agents that comprise an agonist for a protein of the present invention, as an active ingredient.

50 [0127] The methods of screening for molecules that bind when an immobilized protein of the present invention is exposed to synthetic chemical compounds, or natural substance banks, or a random phage peptide display library, or the methods of screening using high-throughput based on combinatorial chemistry techniques (Wrighton Nc, Farrel FX, Chang R, Kashyap AK, Barbone FP, Mulcahy LS, Johnson DL, Barret RW, Jolliffe LK, Dower WJ; Small peptides as potent mimetics of the protein hormone erythropoietin, Science (UNITED STATES) Jul 26 1996, 273 p458-64, Verdine GL., The combinatorial chemistry of nature. Nature (ENGLAND) Nov 7 1996, 384, p11-13, Hogan JC Jr., Directed combinatorial chemistry. Nature (ENGLAND) Nov 7 1996, 384 p17-9) to isolate proteins such as agonists and antagonists that bind to proteins of the present invention are well known to those skilled in the art.

55 [0128] The present invention also provides kits to be used in the identification (screening) methods as described

above. The kits comprise a protein of the present invention or cells expressing a protein of the present invention. The kits may comprise candidate compounds for a ligand, agonist, or antagonist for NKIR protein.

**[0129]** When administering a protein of this invention, a compounds isolated by a identification (screening) method of the present invention, or a pharmaceutical agent as a pharmaceutical for humans and other mammals such as mice, rats, guinea-pigs, rabbits, chicken, cats, dogs, sheep, pigs, cattle, monkeys, baboons and chimpanzees, the protein or the isolated compound can be directly administered or formulated into a dosage form using a known pharmaceutical preparation method. For example, according to the need, the pharmaceutical can be taken orally, as a sugar-coated tablet, capsule, elixir or microcapsule, or non-orally, in the form of an injection of a sterile solution or suspension with water or any other pharmaceutically acceptable liquid. For example, the compounds can be mixed with pharmacologically acceptable carriers or medium, specifically, sterilized water, saline, plant oils, emulsifiers, suspending agents, surfactants, stabilizers, flavoring agents, excipients, vehicles, preservatives, binders and such, in a unit dose form required for generally accepted drug implementation. The amount of active ingredient in these preparations facilitates the acquisition of a suitable dosage within the indicated range.

**[0130]** Examples of additives that can be mixed to tablets and capsules are, binders such as gelatin, corn starch, tragacanth gum and arabic gum; excipients such as crystalline cellulose; swelling agents such as corn starch, gelatin and alginic acid; lubricants such as magnesium stearate; sweeteners such as sucrose, lactose or saccharin; flavoring agents such as peppermint, Gaultheria adenothrix oil and cherry. When the unit dosage form is a capsule, a liquid carrier such as oil can also be included in the above ingredients. Sterile composites for injections can be formulated following normal drug implementations using vehicles such as distilled water used for injections.

**[0131]** Saline, glucose, and other isotonic liquids including adjuvants such as D-sorbitol, D-mannnose, D-mannitol, and sodium chloride, can be used as aqueous solutions for injections. These can be used in conjunction with suitable solubilizers such as alcohol, specifically ethanol, polyalcohols such as propylene glycol and polyethylene glycol, and non-ionic surfactants such as Polysorbate 80 (TM) and HCO-50.

**[0132]** Sesame oil or Soy-bean oil can be used as a oleaginous liquid and may be used in conjunction with benzyl benzoate or benzyl alcohol as a solubilizer and may be formulated with a buffer such as phosphate buffer and sodium acetate buffer; a pain-killer, such as procaine hydrochloride; a stabilizer, such as benzyl alcohol, phenol; and an anti-oxidant. The prepared injection may be filled into a suitable ampule.

**[0133]** Methods well known to those skilled in the art may be used to administer the inventive pharmaceutical to patients, for example as intraarterial, intravenous, percutaneous injections and also as intranasal, transbronchial, intramuscular, percutaneous, or oral administrations. The dosage and method of administration vary according to the body weight and age of the patient and the administration method; however, these can be routinely selected by one skilled in the art. If said compound is encodable by a DNA, the DNA can be inserted into a vector for gene therapy and the vector administered to perform the therapy. The dosage and method of administration vary according to the body weight, age, and symptoms of the patient, but one skilled in the art can select them suitably.

**[0134]** The dose per time of a protein of this invention may vary depending on the type of recipient, target organ, disease condition, and administration method. For example, when injecting into a normal adult (body weight: 60 kg), it may be administered at about 100  $\mu$ g to 20 mg per day.

**[0135]** The dose of a compound that binds to a protein of this invention, or that of a compound that regulates the activity of a protein of this invention varies depending on the type of disease. For example, the compound may be administered orally into a normal adult (body weight: 60 kg) at about 0.1 to 100 mg per day, preferably at about 1.0 to 50 mg per day, and more preferably at about 1.0 to 20 mg per day.

**[0136]** When administered parenterally, the dose per time may vary depending on the recipient, target organ, disease condition, and administration method. For example, an appropriate dose can be, as an intravenous injection into a normal adult (body weight: 60 kg), usually about 0.01 to 30 mg per day, preferably about 0.1 to 20 mg per day, and more preferably about 0.1 to 10 mg per day. For other animals, an amount converted to dose per 60 kg body weight, or dose per body surface area may be applied.

**[0137]** All prior-art documents cited herein are incorporated herein by reference.

### Examples

**[0138]** Hereinbelow, the present invention is specifically described using Examples, but it is not to be construed as being limited thereto.

[Example 1] Cloning of full-length sequence

**[0139]** The present inventors found sequence tags for unidentified genes specific to NK cells (Table 1) among serial analysis of gene-expression (SAGE) data for immune cells disclosed at <http://www.prevent.m.u-tokyo.ac.jp>, and also discovered that the tags were present in the sequence AX191619 in the International publication (WO 0149728). Table

## EP 1 672 065 A1

1 shows selective expression of the sequence tag (TGCCGCATAA) in an NK cell-derived library.

Table 1

	TGCCGCATAA	Total numbers of analyzed tags
Premature dendritic cells	0	50795
GM-CSF-induced macrophages	0	50041
LPS-stimulated monocytes	0	30885
Mature dendritic cells	0	27602

M-CSF-induced macrophages	0	46833
Monocytes	0	51228
Langerhans-like cells	0	44873
CD4 T cells (naive)	0	41789
CD4 T cells (memory, CCR4 negative)	0	27733
CD4 T cells (memory, CCR4 positive)	0	25415
Granulocytes	0	23608
Activated T cells (TH1)	0	26498
Activated T cells (TH2)	0	25371
NK cells	6	29878

**[0140]** Primers SA1 (5'-TTGAATTCACACACCCACAGGACCTGCAGCTGAA-3' / SEQ ID NO: 7), and SA2 (5'-TTGGATCCACTGAAGGACCCACAGAAAGAGTTGA-3' / SEQ ID NO: 8) were designed based on the data described above to amplify the entire coding region, and the gene was cloned by PCR from a cDNA library derived from human spleen. The nucleotide sequence of the yielded clone is shown in SEQ ID NO: 1, and the amino acid sequence of the protein encoded by the nucleotide sequence is shown in SEQ ID NO: 2. The procedures are specifically described below.

### (1) Construction of pGEMTE\_NK1

**[0141]** PCR was carried out under the following reaction conditions.

Template: marathon-ready cDNA human spleen (CLONTECH)

Primers: SA1 <=> SA2

Reaction condition:

96°C for 1 minute;

5 cycles of 96°C for 30 seconds and 72°C for 4 minutes;

5 cycles of 96°C for 30 seconds and 70°C for 4 minutes; and

25 cycles of 96°C for 20 seconds and 68°C for 4 minutes.

**[0142]** PCR was carried out using TaKaRa Ex Taq (buffer and dNTP mixture were supplied with the kit). A band of about 1.5 kb band was excised after agarose gel electrophoresis. After purification with QIAquick Gel Extraction Kit, the PCR product was inserted into pGEM T-Easy Vector to construct pGEMTE\_NK1. The nucleotide sequence was confirmed using the primers SA1 to 7, T7, and SP6.

SA3: 5'-ACCCTGAGATGTCAGACAAAG-3'/SEQ ID NO:9SA4: 5'-GCCACCTCACACCAGTAAAG-34/SEQ ID NO: 11

SA6: 5'-TGGAGCTGTGGGTGGTATCTG-3'/SEQ ID NO: 12

SA7: 5'-AGAACCTCAAAGAGGAGTGAA-3'/SEQ ID NO: 13

T7 promoter primer: 5'-ATTATGCTGAGTGATATCCC-3'/SEQ ID NO: 14

SP6 promoter primer: 5'-ATTTAGGTGACACTATAGAA-3'/SEQ ID NO: 15

### (2) Construction of pCOS\_NK1

**[0143]** pGEMTE-NK1 was digested with *EcoRI* and *BamHI* and subjected to agarose gel electrophoresis. The resulting

## EP 1 672 065 A1

band of about 1.5 kb was then excised. After purification with QIAquick Gel Extraction Kit, the DNA fragment was inserted between *EcoRI* and *BamHI* sites of pCOS1 to yield pCOS\_NK1.

### (3) Construction of pCHO2-NK1-FLAG

**[0144]** PCR was carried out under the following reaction conditions.  
Template: pGEMTE-NK1

**Primers: SAS1 (5'-GGGAATTCATGTTGCCATCTTTAGTTCC-3' / SEQ ID NO: 16) <=>**  
**SAS2 (5'-AAGGATCCACTCCTCTCTCTGGAGAC-3' / SEQ ID NO: 17)**

Reaction conditions:

94°C for 2 minutes; and  
25 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 68°C for 1 minute.

PCR was carried out using KOD plus (TOYOBO; buffer, dNTPs, and MgSO<sub>4</sub> were attached thereto). After purification with QIAquick PCR Purification Kit (QIAGEN), the PCR product was digested with *EcoRI* and *BamHI* and subjected to agarose gel electrophoresis, and then a band of about 1 kb was excised from the gel. After purification with QIAquick Gel Extraction Kit, the DNA fragment was inserted between *EcoRI* and *BamHI* sites of pCHO2-FLAG to yield pCHO2-NK1-FLAG. The nucleotide sequence was confirmed using the primers SAS1, SAS2, S3, S4, S5, EF1 $\alpha$ , and polyA.

EF1 $\alpha$ : 5'-GCCTCAGACAGTGGTTCAAA-3' / SEQ ID NO: 18

IgG1 polyA: 5'-AGAACCATCACAGTCTCGCA-3' / SEQ ID NO: 19

### (4) Construction of pCHO2-SGNK1-FLAG

**[0145]** PCR was carried out under the following reaction conditions.

A:

Template: pCOS2-SGhMPL-FLAG

**Primers: Hmpl-sig1 (5'-AAGAATTCCACCATGGCTGGACCTGCCAC-3' / SEQ ID NO: 20)**

**<=> NK1-sig2 (5'-ACAGGGTTTGGCCAGGCTTGGGCTTCCTGCACTGTCCAGAG-3' /**  
**SEQ ID NO: 21)**

B:

Template: pGEMTE-NK1

**Primers: NK1-sig1 (5'-GCAGGAAGCCCAAGCCTGGCCAAACCCTGT-3' / SEQ ID NO:**  
**22) <=> SAS2**

C:

Template: Mixture of products of reaction series A and B

Primers: Hmp1-sig1 <=> SAS2

Reaction condition: 94°C for 2 minutes; and 25 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 68°C for 1 minute.

**[0146]** PCR was carried out using KOD plus (TOYOBO; buffer, dNTPs, and MgSO<sub>4</sub> were attached thereto). After the PCR product of reaction series C was purified with QIAquick PCR Purification Kit (QIAGEN), it was digested with *EcoRI* and *BamHI*, and subjected to agarose gel electrophoresis. A band of about 0.9 kb was excised from the gel. After purification with QIAquick Gel Extraction Kit, the DNA fragment was inserted between *EcoRI* and *BamHI* sites of pCHO2-FLAG to yield pCHO2-SGNK1-FLAG. The nucleotide sequence was confirmed using the primers, NK1-sig1, NK1-sig2, SAS2, S3, S4, S5, EF1 $\alpha$ , and polyA.

**[0147]** The nucleotide sequences of the clones obtained through steps (1) to (4) as described above were presumed to encode a putative transmembrane protein consisting of 429 amino acids. This sequence was compared with the

## EP 1 672 065 A1

sequence (WO 01/49728) determined by the Sagami Chemical Research Center. The result showed that the central portion of the sequence is identical to the Sagami sequence, but the N-terminal sequences are different. Sequences variations are also found between the proteins encoded. All clones obtained above share the C-terminal sequence, but some clones were deduced to be splicing variants based on the 5'-end sequence. Thus, to determine the original

sequence, 5' RACE was carried out by the following procedure.

**[0148]** RNA (0.4 µg/µl, 20 µl in DEPC-DDW) was extracted from NK cells using RNA-Bee\_RNA ISOLATION REAGENT (Tel-Test). cDNA (5'-RACE-Ready cDNA, 100 µl in Tricine-EDTA Buffer) was synthesized from 1 µg of the RNA using SMART RACE cDNA Amplification Kit (CLONTECH). 5'-RACE PCR was carried out under the following reaction conditions:

1st round:

Template: 2.5 µl of 5'-RACE-Ready cDNA

Primers: 10x Universal Primer A Mix (attached; used at 1x) <=> SAS2

10x Universal Primer A Mix:

(long, 0.4 µM:

5'-CTAATACGACTCACTATAGGGCAAGCAGTGGTATCAACGCAGAGT-3' / SEQ ID NO: 23)

(short, 2 µM: 5'-CTAATACGACTCACTATAGGGC-3'/SEQ ID NO: 24)

2nd round:

Template: 5 µl of 1st round PCR product

Primers: Nested Universal Primer A (NUP) (5'-AAGCAGTGGTATCAACGCAGAGT-3' /SEQ ID NO: 25) <=> SA4

Reaction conditions:

94°C for 30 seconds;

5 cycles of 94°C for 5 seconds and 72°C for 3 minutes;

5 cycles of 94°C for 5 seconds, 70°C for 10 seconds, and 72°C for 3 minutes;

30 cycles of 94°C for 5 seconds, 68°C for 10 seconds, and 72°C for 3 minutes; and 72°C for 7 minutes.

**[0149]** The 2nd round PCR product was electrophoresed in an agarose gel, and a band of about 650bp was excised. After purification with QIAquick Gel Extraction Kit, the resulting fragment was inserted into pGEM T-Easy Vector. The nucleotide sequence was determined using the primers SA3, 4, T7, SP6, and NUP.

**[0150]** The result showed that all coding sequences of the 5 clones whose sequences were verified were identical to the nucleotide sequence described above (of these, one clone was found to be shorter at the 5' end). Thus, it is safe to conclude that the yielded clone described above is identical to the nucleotide sequence truly expressed.

**[0151]** Motif searches using Prosite, Pfam, Psort, and the like, revealed the following sequence features:

N-glycosylation sites: (N<sup>[^P]</sup>[ST]<sup>[^P]</sup>) 60, NQTL; 245, NHSA; and 268, NYSC

ITIM motifs (Yxx[VL]): 351, YANV; and 366, YSVV

Ig-like domains: 27-80; 120-177; and 216-273

Transmembrane domain: 309-325

**[0152]** Based on the above-described sequence information, the isolated gene as described above was inferred to be a member of suppressive receptors (KIR) that recognize, as a ligand, classic MHC class I belonging to the FcR superfamily. Since this gene is expressed specifically in NK cells, hereinafter it is called the NKIR gene.

[Example 2] Detection of NKIR protein using rabbit polyclonal antibody

**[0153]** An *E. coli* expression system was constructed, and a fusion protein comprising the NKIR extracellular domain was expressed and purified (Fig. 1). The procedure is described below in detail.

(1) Construction of *E. coli* expression plasmid pET32a-NK-sol for NKIR fusion protein

**[0154]** PCR was carried out under the following reaction conditions.

Template: pGEMTE-NK1

Primers: NKfusion (5'-CTCGGATCCTTGCCATCTTTAGTTCCTGTGTT-3' / SEQ ID NO: 26) <=> NKr2 (5'-GCTGTCGACTTAGTTGCTGGCGGGAGTGAACAAGAC-3' / SEQ ID NO: 27)

Reaction conditions:

94°C for 2 minutes; and

25 cycles of 94°C for 20 seconds, 60°C for 30 seconds, and 68°C for 2 minutes.

**[0155]** PCR was carried out using KOD plus (TOYOBO; buffer, dNTPs, and MgSO<sub>4</sub> were also supplied with the kit). After purification with MicroSpin S-300 HR column (Amersham Biosciences), the PCR product was digested with *Bam*HI and *Sal*I, and subjected to agarose gel electrophoresis. A band of about 0.9 kb was excised from the gel. After purification with a MicroSpin S-300 HR column, the fragment was inserted between *Bam*HI and *Sal*I sites of pET-32(a) (Novagen) to yield pET32a-NK-sol.

(2) Construction of animal cell expression plasmid pCOS2-NK-FLAG for NKIR protein

**[0156]** PCR was carried out under the following reaction conditions.

Template: pGEMTE-NK1

Primers: NKflag

(5'-GCGAATTCCACCATGGACTACAAAGACGATGACGACAAGTTGCCATCTTTAGTTCCTGTGTT-3' / SEQ ID NO: 28) <=> NKr1

(5'-CGTGTCGACTCACTAGCAGAGAACCTCCTCACAGTC-3'/SEQ ID NO: 29)

Reaction conditions:

94°C for 2 minutes; and

25 cycles of 94°C for 20 seconds, 60°C for 30 seconds, and 68°C for 2 minutes.

**[0157]** PCR was carried out using KOD plus (TOYOBO; buffer, dNTPs, and MgSO<sub>4</sub> were attached thereto). After purification with MicroSpin S-300 HR column (Amersham Biosciences), the PCR product was digested with *Eco*RI and *Sal*I and subjected to agarose gel electrophoresis. A band of about 1.3 kb was excised from the gel. After purification with a MicroSpin S-300 HR column, the fragment was inserted between *Eco*RI and *Sal*I sites of pCOS2 to yield pCOS2-NK-FLAG.

(3) Construction of NKIR fusion protein and rabbit polyclonal NKIR antibody

**[0158]** *E. coli* BL21 (DE3) was transformed with the expression plasmid pET32a-NK-sol for thioredoxin fusion protein. The *E. coli* was cultured overnight in LB medium containing 50 µg/ml ampicillin, and the suspension was diluted to the final concentration of 1% with LB medium containing 50 µg/ml ampicillin for large scale culture. Cells were cultured until the absorbance at 600 nm reached 0.4. Then, IPTG was added at the final concentration of 1 mM to induce the expression of the protein. After 4.5 hours of culture, bacterial cells were precipitated and collected by centrifugation. The cells were suspended in PBS. The bacterial cells were sonicated in a sonicator, and then centrifuged. The resulting supernatant was discarded, and the precipitated fraction was collected. The cells were lysed with phosphate buffer (PBS) containing 7 M urea, and the lysate was filtered with a 0.45-µm filter. The NKIR fusion protein was isolated from the filtrate using HisTrap kit (Amersham Biosciences). The eluate was dialyzed against 50 mM Tris (pH8.3), and thus a sample of solubilized fusion protein was obtained. SDS-PAGE, followed by Coomassie staining, confirmed that the fusion protein obtained had the expected molecular weight.

**[0159]** Rabbits were immunized using the NKIR fusion protein as an immunogen to prepare polyclonal antibodies. The antigen protein solution was adjusted to 0.2 mg/0.5 ml, and 0.5 ml of Freund's complete adjuvant (Becton Dickinson) was added thereto. 1 ml of the mixture was inoculated subcutaneously (day 1, 4, and 11). On day 19, 26, and 33, 0.05 mg of the antigen protein was intravenously injected. After blood sampling, increase in antibody titers was confirmed. Then, the blood was collected and the antiserum was loaded onto protein A affinity column to purify the antibody protein. Western blotting as described below was used to confirm that the obtained polyclonal antibody binds to NKIR expressed transiently in the animal cell COS-7.

**[0160]** pCOS2-NK-FLAG was introduced into COS-7 cells using FuGENE6 (Roche Diagnostics) according to the

## EP 1 672 065 A1

manufacturer's instructions. After two days of culture, a lysis solution (10% glycerol, 50 mM Tris (pH7.6), 150 mM NaCl, 5 mM NP-40, and protease inhibitors (complete)) was added to the cells. Then, the resulting suspension was centrifuged to obtain a supernatant comprising soluble membrane protein components. Anti-FLAG M2 antibody resins were added to the soluble fraction to immunoprecipitate FLAG-NKIR. The obtained sample was fractionated by SDS-PAGE, and then transferred onto a PVDF membrane. Detection was carried out using anti-NKIR polyclonal antibody. As the primary antibody, the rabbit-derived polyclonal antibody (4.1 mg/ml IgG) was diluted 1000 times. As the secondary antibody, HRP-conjugated anti-rabbit antibody (Amersham Biosciences) was diluted 3000 times. The ECL Western Blotting Detection System (Amersham Biosciences) was used in the detection. In a control experiment, the presence of FLAG-NKIR protein was confirmed using 1000-times diluted anti-FLAG M2 antibody (Sigma) and 3000-times diluted HRP-conjugated anti-mouse antibody (Amersham Biosciences) as the secondary antibody (Fig. 2).

[Example 3] Tissue expression analysis and expression analysis ofNK cell lines

**[0161]** Tissue expression profiles were analyzed by PCR using the commercially available cDNA panel (Multiple Tissue cDNA (MTC) panel; Clontech Laboratories, Inc.) as a template.

**[0162]** PCR was carried out under the following reaction conditions. Template: MTC panel I, II, Human Immune System and Human Blood Fraction (Clontech Laboratories, Inc.)

**Primers: NKIR07 (5'-AGGTCAGAGTGCAGGCTCCTGTATC-3' / SEQ ID NO: 30) <=>**  
**NKIR08 (5'-TAGAACTGTCCTTTCTCCCCACGGT-3' / SEQ ID NO: 31)**

Reaction conditions:

94°C for 30 seconds; and

35 cycles of 94°C for 30 seconds and 65°C for 2 minutes.

**[0163]** PCR was carried out using TaKaRa Ex Taq (buffer and dNTP mixture were provided). The reaction volume was 50  $\mu$ l. 5  $\mu$ l of the reaction solution was electrophoresed in a 1% agarose gel. A 0.6 kb band was observed. G3PDH primers attached to the MTC panel were used in a control reaction.

The result showed that the gene was specifically expressed in the immune system such as in the spleen and leukocytes (Fig. 3). When the number of subjects in the analysis was increased using subsets of "Blood Fraction" and "Immune System" of the above-described MTC panel, the gene was found to be specifically expressed in monocytes and in resting CD8+ for the "Blood Fraction" and in lymph nodes in addition to spleen and leukocytes for the "Immune System" (Fig. 3).

Western blotting analysis was carried out using a cell lysate prepared from NK-92 cell line, an established natural killer (NK) cell line, purchased from ATCC (catalog No: CRL-2407).

NK-92 cell line was cultured in NK-92 passaging medium ( $\alpha$ -MEM medium (Invitrogen) comprising 12.5% fetal calf serum (Invitrogen), 12.5% horse serum (Invitrogen), 2 mM glutamine (Invitrogen), 0.1 mg/l penicillin (Invitrogen), 0.1 mg/l streptomycin (Invitrogen), 1 mM sodium pyruvate (Invitrogen), 100  $\mu$ M 2-mercaptoethanol (Invitrogen), 2 mM folic acid (SIGMA-ALDRICH), and 20 mM myo-inositol (SIGMA-ALDRICH)) in the presence of 10 ng/ml interleukin-2 (SIGMA-ALDRICH). The resulting  $1 \times 10^7$  cells were suspended in 500  $\mu$ l of NP40 lysis buffer (1% NP40 / 150 mM NaCl / 50 mM Tris-HCl (pH8.0)), and allowed to stand on ice for 30 minutes. The suspension was then fractionated through micro centrifugation at 15,000 rpm to obtain a supernatant. The supernatant was used as a cell lysate in Western blotting analysis.

The protein concentration in the cell lysate was determined by using the Dc Protein Assay Kit (BIO-RAD Laboratories) and, as a control, bovine serum albumin (Fraction V) from PIERCE.

PAG-Mini (gel with a gradient of 4% to 20%; Daiichi Pure Chemicals Co. Ltd.) and 10  $\mu$ g or 20  $\mu$ g of lysate of NK-92 cell line were used in SDS-PAGE for the Western blotting analysis. The same immunogen as that used to immunize rabbits in the polyclonal antibody preparation was diluted 200 times with NP40 lysis buffer, and then 1  $\mu$ l of the resulting sample was simultaneously used as a positive control. The electrophoresis was carried out at 20 mA. After electrophoresis, the proteins were transferred onto PVDF membrane (Hybond-P, Amersham Biosciences) from the gel using SEMI-DRY TRANSFER CELL (BIO-RAD Laboratories) under the condition of 20 volts for 45 minutes. Western blotting was carried out using ECL plus Western Blotting Detection System (Amersham Biosciences) according to the method described in the manual. However, ECL-Advance blocking agent (Amersham Biosciences) was used as a blocking reagent at the concentration of 2%. 1,000-time diluted polyclonal antibody (4.1 mg/ml IgG) derived from rabbit described above was used as the primary antibody, while 3,000-time diluted anti-rabbit Ig derived from donkey (horseradish peroxidase linked whole antibody; Amersham Biosciences) was used as the secondary antibody. It was confirmed that there are molecular species that cross-reacted with the anti-NKIR polyclonal antibody.

**[0169]** The result showed that NK-92 cell line expresses an about 60-kDa protein cross-reactive to the polyclonal antibody obtained from the rabbit immunized with NKIR protein expressed in *E. coli* (Fig. 4).

**[0170]** Furthermore, to confirm that the NKIR molecule is expressed on the surface of NK cells, flow cytometric analysis of NK-92 cell line was carried out using the anti-NKIR polyclonal antibody.

**[0171]** After  $5 \times 10^5$  cells were suspended in 100  $\mu$ l of FACS buffer (phosphate buffered saline comprising 2.5% fetal calf serum and 0.02%  $\text{NaN}_3$ ), the anti-NKIR polyclonal antibody was added thereto at the final concentration of 82  $\mu$ g/ml. As a negative control, cells to which a purified rabbit IgG had been added at the same concentration were prepared. The cell samples were allowed to stand on ice for one hour, and then subjected to centrifugation at a low speed (at 300 x g for 5 minutes) to collect cells. The cells were washed with FACS buffer, and centrifuged again at low speed to collect the cells. Then, the collected cells were suspended in FACS buffer comprising an FITC-conjugated goat anti-rabbit IgG antibody (Beckman Coulter) at the final concentration of 14  $\mu$ g/ml, and allowed to stand on ice for 30 minutes. The cells were collected by centrifugation at low speed, and washed with FACS buffer. The cells were suspended in 500  $\mu$ l of FACS buffer, and analyzed by flow cytometry (Beckman Coulter; EPICS). The result showed that the NKIR molecule was expressed on the surface of the cells (Fig. 5).

[Example 4] Cloning of the full-length NKIR gene by RACE using NK-92 cell line-derived cDNA

**[0172]** The full-length NKIR gene was cloned again by 5'- and 3'-RACE using total RNAs prepared from NK-92 cell line.

**[0173]** Using RNeasy (QIAGEN) kit, 377.7  $\mu$ g of total RNAs was prepared from  $5.4 \times 10^7$  cells of NK-92 cell line cultured by the method in Example 3. 5'- and 3'-RACE were carried out using SMART RACE cDNA Amplification Kit (Clontech) and as a template 1  $\mu$ g of RNA prepared as described above. Gene-specific primers used in 5'- and 3'-RACE were NKIR08 and NKIR07, respectively. In each reaction, a major amplification product of about 1.3 kb was obtained in the 1st round PCR. The amplified fragment was excised from the agarose gel, purified with QIAquick (QIAGEN) kit, and then cloned into pCR2.1-TOPO (Invitrogen). The yielded eight transformants derived from TOP10F' were cultured in 2 ml of LB medium comprising 0.1 mg/ml ampicillin. Plasmids were prepared using QIAprep kit (QIAGEN) from the cultured *E. coli*, and then sequenced. As a result, 5'-RACE yielded a cDNA sequence comprising an insert that is 36 nucleotides longer than the previously identified sequence at the 5' end. The cDNA was used as pTOPONKIR626 in subsequent analyses. 3'-RACE yielded the cDNA clone pTOPONKIR620 comprising a downstream extension of about 500 nucleotides. The nucleotide sequence is shown in SEQ ID NO: 3 and the amino acid sequence of the protein encoded by the nucleotide sequence is shown in SEQ ID NO: 4 in the Sequence Listing. This sequence was confirmed to be located in the NKIR region of chromosome 1 in the human genome.

**[0174]** Then, focusing on the insertion sequence of 36 nucleotides at the 5' end, expression vectors respectively comprising the sequence identified in Example 1 and newly isolated sequence were constructed using pCOS 1 as the vector backbone to use in the transient expression in COS-7 cells.

**[0175]** PCR was carried out using pGEM-TE NK1 as a template and the following primers (0.2  $\mu$ M each): NKIR09 (5'-GAATTCACACACCCACAGGACCTGCA-3' / SEQ ID NO: 32) and NKIR10 (5'-GGATCCACTGAAGGACCCACA-GAAAG-3' / SEQ ID NO: 33). The HF Polymerase kit (Clontech) was used as PCR kit. The reaction conditions were: denaturation at 94°C for 30 seconds; 25 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 1 minute; followed by extension at 72°C for 5 minutes. The reaction solution was subjected to agarose gel electrophoresis, and the yielded 1.5 kb fragment was purified using QIAquick Gel Extraction kit (QIAGEN). The fragment was then cloned into pCR2.1-TOPO (Invitrogen) and introduced into *E. coli* strain TOP10F'. Plasmid was prepared from the resulting ampicillin-resistant strain using QIAprep Miniprep kit (QIAGEN), and then sequenced. The PCR error-free clone pTOPONKIR219 was selected for subsequent analyses.

**[0176]** Then, 5  $\mu$ g of pTOPONKIR219 and 1.15  $\mu$ g of pBluescript II SK+ (Stratagene) were digested with 20 units each of *Eco*RI and *Bam*HI at 37°C for one hour. After agarose gel electrophoresis, the resulting 1.5 kb and 3.0 kb fragments were purified using QIAquick Gel Extraction kit (QIAGEN). Then, ligation was carried out using a 1  $\mu$ l aliquot of each eluate (40  $\mu$ l) and LigaFAST Ligation kit (Promega), and introduced into the *E. coli* TOP10F' strain. Plasmids were prepared from the resulting ampicillin-resistant strains using QIAprep Miniprep kit (QIAGEN). The clones were tested by digestion with the restriction enzyme *Pst*I. The clone pBSNKIR224 gave a 1.3 kb fragment by the digestion, and therefore used in the subsequent analyses. The 4 kb and 0.4 kb fragments obtained respectively from pBSNKIR224 and pTOPONKIR626 by double digestion with *Bst*P1 and *Bgl*II were then isolated by agarose gel electrophoresis and purified with QIAquick Gel Extraction kit (QIAGEN). The fragments were ligated using LigaFAST Ligation kit (Promega), and introduced into *E. coli* DH5 $\alpha$ . Plasmids were prepared from the resulting ampicillin-resistant strains using QIAprep Miniprep kit (QIAGEN), and then sequenced. The clone pBSNKIRfu11605 comprising a 36-bp insert at the 5' end was used in the subsequent analyses.

**[0177]** Finally, 1.4-kb *Eco*RI-*Not*I fragments derived from pBSNKIR224 and pBSNKIRfu11605 were inserted between *Eco*RI and *Not*I sites of pCOS1 to construct the expression vectors pCOSNKIR610 and pCOSNKIRfu11610, respectively. The two plasmids were transfected into COS-7 cells as donor DNAs by lipofection method using MIRUS TransIT-LT1

(PanVera) according to the manufacturer's instructions, and the cells were cultured for 2 days. Then, the cells were trypsinized using a trypsin/EDTA solution (Invitrogen), and washed twice with DMEM (Invitrogen) comprising 10% fetal calf serum (Invitrogen). Flow cytometric analyses were then carried out by the same method as described in Example 3.

5 **[0178]** A cell fraction was found to be shifted in the FITC detection only when COS-7 cells were transfected with the expression plasmid for NKIR isolated from NK-92, which comprises the insertion sequence of 36 nucleotides at the 5' end. Accordingly, the result showed that the clone comprising the 36-nucleotide insertion functions as a secretory form (Fig. 6).

10 [Example 5] Cloning of mouse NKIR sequence

**[0179]** The mouse genomic sequence was searched by BLAST (tblastn) using the amino acid sequence of human NKIR as a query. A region on chromosome 1 that gave a hit over the entire sequence was identified (Fig. 7). With respect to the chromosomal structure, this mouse chromosomal region matches the human chromosomal region on which human NKIR has been mapped.

15 **[0180]** Primers (mNKIRf1 (5'-CTCAGTAAAGGCAGAGTGGAGTACC-3' / SEQ ID NO: 34) <=> mNKIRr1 (5'-ATACATTAGAACCACAGCCGCAATG-3' / SEQ ID NO: 35)) were designed based on the putative translated region. The gene was cloned from a mouse spleen cDNA library by PCR amplification and sequenced. The presence of spliced transcripts was verified.

20 **[0181]** 5'- and 3'-RACE PCR were carried out using mouse spleen Marathon-Ready cDNA (Clontech) as a template under the following reaction conditions: 1 st round PCR:  
 Template: 2.5 µl of Marathon Ready cDNA  
 Primers:

25 **API (5'-CCATCCTAATACGACTCACTATAGGGC-3' / SEQ ID NO: 36) <=> mNKIRf1 for 3' RACE**

30 API <=> mNKIRr1 for 5' RACE

2nd round PCR:

Template: 2.5 µl of 30-time diluted 1st round PCR product

Primers:

AP2(5'ACTCACTATAGGGCG-3'/SEQ ID NO: 37 <=>mNKIRf3

35 **(5'-CTCAAGAAGTTCCTTGGTTGTCTC-3' / SEQ ID NO: 38) for 3' RACE**

**AP2 <=> mNKIRr3 (5'-GCCAGATAGTTAGCATGTTGCTCTTG-3' / SEQ ID NO: 39) for 5' RACE**

40 1 st round PCR:

Reaction conditions:

94°C for 1 minute; and

45 35 cycles of 94°C for 10 seconds and 68°C for 3 minutes.

2nd round PCR:

Reaction conditions:

94°C for 1 minute; and

20 cycles of 94°C for 10 seconds and 68°C for 3 minutes.

50 **[0182]** A reaction solution was prepared and PCR was carried out using TaKaRa LA Taq (TAKARA; buffer, dNTPs, and MgCl<sub>2</sub> were attached thereto) according to the manufacturer's instructions. The products of 2nd round PCR were electrophoresed in an agarose gel, and the band resulting from the PCR amplification was excised. After purification with QIAquick Gel Extraction Kit, the product was inserted into pGEM T-Easy Vector. DH5α was transformed, and plasmids were prepared from the resulting clones and sequenced.

55 **[0183]** As a result, the clone was deduced to be a membrane protein comprising 268 amino acids. The nucleotide sequence of the obtained clone is shown in SEQ ID NO: 5 and the amino acid sequence of the protein encoded by the nucleotide sequence is shown in SEQ ID NO: 6 in the Sequence Listing. When the human and mouse sequences are compared (Fig. 8), the mouse sequence has a structure that is shorter than that of the human sequence at both N and

## EP 1 672 065 A1

C termini. The mouse sequence comprises a single N-glycosylation site, a single ITIM motif, a single transmembrane domain, and two Ig-like domains in the equivalent region.

**[0184]** Motif searches by Prosite, Pfam, Psort, and the like revealed the following sequence features:

N-glycosylation sites: (N[<sup>^</sup>P][ST][<sup>^</sup>P]) 180, NYSC; 188, NISR

ITIM motif: (Yxx[VL]) 259, YANV

Ig-like domains: 33-89 and 128-185

Transmembrane domain: 221-237

[Example 6] Cloning of 2KIR3DL

**[0185]** An assay system used for detecting ITIM activity comprises a modified method using T cells as recipient cells which comprises the step of determining luciferase activity under the control of the NFAT cascade (Fry AM, Lanier LL, Weiss A., J Exp Med. (1996), 184, 295-300).

**[0186]** 2KIR3DL, a known KIR gene, was cloned using a human spleen cDNA library. PCR was carried out under the following reaction conditions:

Template: Human Spleen Marathon-Ready cDNA (Clontech)

**Primers: p58KIR01 (5'-GAATTCATGTCGCTCATGGTCGTCAG-3' / SEQ ID NO: 40) <=>**

**p58KIR02 (5'-GGATCCTCAGGGCTCAGCATTGGAA-3' / SEQ ID NO: 41)**

Reaction conditions:

94°C for 30 seconds;

30 cycles of 94°C for 15 seconds, 52°C for 30 seconds, and 72°C for 1 minute; and

72°C for 5 minutes.

**[0187]** PCR was carried out using HF polymerase (Clontech). A 1-kb fragment was separated by agarose gel electrophoresis, and then purified with QIAquick (QIAGEN). The purified product was cloned into pCR2.1-TOPO, and introduced into *E. coli* TOP10F'. The plasmid was prepared from the transformant using QIAprep (QIAGEN) kit, and a clone comprising a PCR error-free sequence (pTOPO58KIR303) was selected to use in the subsequent analyses.

[Example 7] Construction of an in-frame fusion

**[0188]** An in-frame fusion between the cytoplasmic ITIM motif of NKIR and the extracellular domain of 2KIRDL3 obtained in Example 6 was constructed by the following procedure.

**[0189]** First, fusion PCR was carried out under the following reaction conditions. 1 st round PCR A

Template: pTOP058KIR303

Primers: p58KIR01 <=> p58NKIR04

**(5'-AGGGGCCAGCTTTTCTCCAGCGATGAAGGAGAAAGA-3' / SEQ ID NO: 42)**

**1st round PCR B**

Template: pBSNKIR224

**Primers: p58KIR03 (5'-TCTTCTTTCTCCTTCATCGCTGGAGAAAAGCTGGGCCCT-3' /**

**SEQ ID NO: 43) <=> T3+ (5'-GCAATTAACCCTCACTAAAGGGAAC-3' / SEQ ID NO: 44)**

**[0190]** The condition used for both reactions is as follows:

94°C for 30 seconds,

30 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 45 seconds; and

72°C for 2 minutes.

**[0191]** PCR was carried out using HF polymerase (Clontech). 0.8- and 0.4-kb fragments derived from PCR A and B, respectively, were separated by agarose gel electrophoresis, and then purified using QIAquick (QIAGEN). A 10- $\mu$ l aliquot of each purified product (50  $\mu$ l) was used as a template in the second round of PCR described below.

## EP 1 672 065 A1

Template: 10  $\mu$ l of each reaction product described above

**[0192]** The reaction conditions were as follows:

94°C for 30 seconds; and

15 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 90 seconds.

**[0193]** After the reaction, the template described below was added to 50  $\mu$ l of the reaction mixture at the final concentration of 1  $\mu$ M.

Primers: p58KIR01  $\leftrightarrow$  T3+

Reaction conditions:

94°C for 30 seconds;

35 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 2 minutes; and

72°C for 4 minutes.

**[0194]** A 1.2-kb fragment was separated by agarose gel electrophoresis, and then purified using QIAquick (QIAGEN). The fragment was cloned into pCR2.1-TOPO, and introduced into *E. coli* TOP10F'. Plasmid was prepared from the transformant using QIAprep (QIAGEN) kit, and a clone comprising a PCR error-free sequence (pBSKIR58NKIR314) was selected and used in the subsequent analyses.

**[0195]** Next, 1  $\mu$ g each of pCXND3 and pBSKIR58NKIR314 were digested with 20 units of *Eco*RI and *Not*I at 37°C for one hour, and then electrophoresed in an agarose gel. The resulting 7.8- and 1.2-kb fragments were purified using QIAquick Gel Extraction kit (QIAGEN). Then, ligation was carried out using a 1- $\mu$ l aliquot of each eluate (40  $\mu$ l) and LigaFAST Ligation kit (Promega), and the resulting plasmid was introduced into *E. coli* strain DH5 $\alpha$ . Ampicillin-resistant strains were saved and colony PCR was carried out using p58KIR01 and p58NKIR10 primers (0.5  $\mu$ M each). This treatment was carried out using Premix ExTaq (TaKaRa) as PCR polymerase, under conditions of: denaturation at 94°C for 5 minutes; followed by 35 cycles of 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 90 seconds. One fifth of the resulting reaction product was subjected to agarose gel electrophoresis, and the clone pCXND3KIR58NKIR313 that gave a 1.3-kb PCR fragment was used in the subsequent analyses.

[Example 8] Preparation of stable transformants

**[0196]** ChimeraA10, a candidate strain for a stable transformant, was prepared from a T cell line Jurkat using as a donor DNA pCXND3KIR58NKIR313 obtained in Example 7 described above. The procedure used to obtain the strain is described below.

**[0197]** Transduction was achieved by electroporation using pCXND3KIR58NKIR313 as donor DNA and Jurkat strain as recipient cell. 20  $\mu$ g of the donor DNA was pre-digested with 20 units of *Pvu*II (TaKaRa) at 37°C for one hour. After chloroform/phenol treatment followed by ethanol precipitation, the DNA was dissolved in 20  $\mu$ l of sterilized water. As recipient cells, Jurkat cells were passaged in RPMI1640 medium comprising 10% FBS, washed with potassium-based phosphate buffered saline (K-PBS), and then suspended in K-PBS at the cell density of 10<sup>7</sup> cells/ml to prepare 0.8 ml of a cell suspension. Electroporation was carried out using Gene Pulsar II (Bio-Rad) under the pulse condition of 0.3 kV and 950  $\mu$ FD. After pulse application, the cells were suspended in 48 ml of passaging medium and cultured overnight under a condition without selective pressure. Then, the candidate ChimeraA10 cell line for the stable transformant was obtained under a selective pressure using geneticin (Invitrogen) at the final concentration of 400  $\mu$ g/ml. Flow cytometric analysis was carried out by the same procedure as described in Example 3 except for using GL183 monoclonal antibody (Beckman Coulter) as the primary antibody and FITC-conjugated anti-rabbit IgG antibody (Beckman Coulter) as the secondary antibody.

**[0198]** As a result, an FITC-stained cell fraction was detected, indicating that the above-described fusion protein was expressed in ChimeraA10 cell line (Fig. 9).

[Example 9] Preparation of dual transformant

**[0199]** The luciferase reporter plasmid pNFATlucZEOR324 was transformed into ChimeraA10 strain obtained in Example 8 as recipient cell to obtain a dual transformant. pNFATlucZEOR324 was constructed by the procedure as described below.

**[0200]** 2.5  $\mu$ g of the luciferase reporter gene pNFAT-TA-Luc (included in Mercury Pathway Profiling Luciferase system 2 (Clontech)) and 5  $\mu$ g of pCOSIIZEO (prepared from SCS-110 strain (Stratagene), a *dam*- strain) were digested with 20 units each of *Acc*I (TaKaRa) and *Cla*I (TaKaRa), respectively, at 37°C for one hour, and then subjected to agarose gel electrophoresis. 5- and 2.2-kb fragments obtained respectively by the digestion were purified using Gel Extraction Kit (QIAGEN). The fragment derived from pNFAT-TA-Luc was treated with calf intestine alkaline phosphatase at 37°C for 30 minutes, and then purified using QIAquick Nucleotide Removal kit (QIAGEN). The two fragments were ligated using LigaFAST Ligation kit (Promega), and introduced into *E. coli* strain DH5 $\alpha$ . Plasmid was prepared from the resulting ampicillin-resistant strain using QIAprep Miniprep kit (QIAGEN). The plasmid was tested by double-digestion with the

restriction enzymes *EcoRI* and *SaI*. A clone that gave 4.15- and 3.15-kb fragments by the digestion was named the pNFAT1ucZEO324 clone comprising the insert in a forward orientation; and a clone that gave 5.15- and 2.15-kb fragments by the digestion was pNFAT1ucZEO324 clone comprising the insert in an reverse orientation. Both plasmids were used in the subsequent analyses.

5 **[0201]** Transduction was conducted by electroporation using pNFAT1ucZEO324 as donor DNA and chimeraA10 cell line as recipient cell. 20 µg of the donor DNA was pre-digested with 20 units of *PvuII* (TaKaRa) at 37°C for 1 hour. After chloroform/phenol treatment followed by ethanol precipitation, the DNA was dissolved in 20 µl of sterilized water. ChimeraA10 cells (recipient cells) were passaged in RPMI1640 medium comprising 10% FBS and 400 µg/ml geneticin, washed with potassium-based phosphate buffered saline (K-PBS), and suspended in K-PBS at the cell density of 10<sup>7</sup> cells/ml to prepare 0.8 ml of a cell suspension. Electroporation was carried out using Gene Pulsar II (Bio-Rad) under the pulse condition of 0.3 kV and 950 µFD. After pulse application, the cells were suspended in 48 ml of passaging medium comprising 400 µg/ml geneticin and cultured overnight under a condition without selective pressure. Then, the candidate cell line for the stable transformant was obtained under a selective pressure using zeocin (Invitrogen) at the final concentration of 100 µg/ml. Genomic DNA was prepared from the resulting zeocin-resistant cell line using DNeasy Tissue Kit (QIAGEN). PCR was carried out using a 5-µl aliquot of the final eluate (0.4 ml). Premix ExTaq was used as the PCR polymerase. Luc01 (5'-TTCATACAGAAGGCGTGGAG-3' / SEQ ID NO: 45) and Luc02 (5'-CGTTCGCGGGCG-CAACTGCA-3' / SEQ ID NO: 46) were used as primers. The reaction was carried out under conditions of: denaturation at 94°C for 5 minutes; and 25 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 45 seconds; followed by extension at 72°C for 5 minutes. Clones which gave a 0.5-kb PCR fragment in agarose gel electrophoresis were selected, and then the final screening was carried out by detecting luciferase activity.

10 **[0202]** The candidate cell line for the stable transformant with pNFAT1ucZEO324, which was obtained from chimeraA10 cell line, was suspended at the concentration of 6.7 x 10<sup>4</sup> cells/ml in a passaging medium containing 400 µg/ml geneticin and 100 µg/ml zeocin. A 75-µl aliquot of the cell suspension was plated into each well of anti-human CD3-coated microtiter plates (Beckton Dickinson) and non-coated immunomicroplates to carry out luciferase assay. After 20 hours of cultivation at 37°C, an equal volume (75 µl) of Dual-Glo Luciferase Buffer (Promega) was added to each well. The samples were allowed to stand for 10 minutes, and then the chemiluminescence was measured using the luminometer MicroLumat LB96P (EG&G Berthold) for 5 seconds for each well.

15 **[0203]** The chimeraA10ZEOR12 cell line that exhibited intense chemiluminescence only when cultured in anti-human CD3-coated microtiterplates was selected as a stable transformant. The cell line was analyzed by flow cytometry by the same procedure as described above.

20 **[0204]** The result showed that the cell line also expresses the fusion protein that is expressed in ChimeraA10 cell line (Fig. 9).

[Example 11] Assay for the activity of ITIM derived from NKIR

25 **[0205]** A functional evaluation of the NKIR-derived ITIM motif was done using chimeraA10ZEOR12 cell line. The method used is described below.

30 **[0206]** The chimeraA10ZEOR12 cell line was suspended at a concentration of 5 x 10<sup>5</sup> cells/ml in a passaging medium containing 400 µg/ml geneticin and 100 µg/ml zeocin. A 100-µl aliquot of the suspension was added to each well of anti-human CD3-coated microtiterplates (Beckton Dickinson). After the cells were cultured at 37°C for 6 hours, GL183 antibody (Beckman Coulter) was added thereto at the final concentration of 1 µg/ml. The cells were cultured at 37°C overnight. Then, the cells were washed twice with passaging medium containing 400 µg/ml geneticin and 100 µg/ml zeocin. After the washed cells were suspended in 100 µl of the medium containing rat anti-mouse IgG antibody at the concentration of 0, 2, or 10 µg/ml, luciferase activity was assayed according to the same procedure as described in Example 10.

35 **[0207]** The result showed that crosslinking with the rat anti-mouse IgG antibody inhibited the luciferase activity by 33.3% in average for three cases (Fig. 10), indicating that the ITIM motif within the cytoplasmic domain of NKIR molecule had ITIM activity.

40 [Example 12] Cloning of CD8α chain

**[0208]** The functional assay system for ITIM activity was the same as that described in Example 6, which comprises a modified method using T cells as recipient cells and features determination of luciferase activity under the control of NFAT cascade (Fry AM, Lanier LL, Weiss A., J Exp Med. (1996), 184, 295-300).

45 **[0209]** The known CD8α chain gene was cloned using resting CD8+ Marathon cDNA library (Clontech) as a template. PCR was carried out under the following reaction conditions.

Primers: CD01 (5'-GAATTCATGGCCTTACCAGTGACCGC-3' / SEQ ID NO: 47) <=>  
 CD02 (5'-GGATCCTTAGACGTATCTCGCCGAAA-3' / SEQ ID NO: 48)

5

Reaction conditions:  
 94°C for 30 seconds;  
 30 cycles of 94°C for 15 seconds, 50°C for 30 seconds, and 72°C for 30 seconds; and  
 72°C for 4 minutes.

10 **[0210]** PCR was carried out using GC polymerase (Takara Shuzo). A 0.7-kb fragment was separated by agarose gel electrophoresis, and then purified using QIAquick (QIAGEN). The purified product was cloned into pCR2.1-TOPO, and introduced into *E. coli* TOP10F'. Plasmid was prepared from the transformant using QIAprep (QIAGEN) kit, and a clone comprising a PCR error-free sequence (pCD8full0113) was selected and used in the subsequent analyses.

15 [Example 13] Construction of expression vector for CD8-NKIR fusion protein

**[0211]** An expression vector for fusion protein of the cytoplasmic ITIM motif of NKIR and the extracellular domain of CD8 $\alpha$  chain obtained in Example 12 was constructed by the following procedure.

20 **[0212]** First, fusion PCR was carried out under the following reaction conditions. 1 st round PCR A  
 Template: pCD8fu110 113

Primers: CD03 (5'-GAATTCCACCATGGCCTTACCAGTGACCGC-3' / SEQ ID NO: 49)  
 <=> CDNKIR12 (5'-ACCAGCCAGTTGCTGGCGGGGTCCAGCCCCCTCGTGTGCA-3' /  
 SEQ ID NO: 50)

25

1 st round PCR B  
 Template: pBSNKIR224  
 Primers: CDNKIR11

30

(5'-TGCACACGAGGGGGCTGGACCCCGCCAGCAACTGGCTGGT-3') / SEQ ID NO:  
 51) <=> T3+ (SEQ ID NO: 44)

35

**[0213]** The conditions used for both reactions are as follows:  
 94°C for 30 seconds,  
 30 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 1 minute; and  
 72°C for 4 minutes.

40

**[0214]** PCR was carried out using GC polymerase (Takara Shuzo). 0.5- and 0.6-kb fragments derived from PCR A and B, respectively, were separated by agarose gel electrophoresis, and then purified using QIAquick (QIAGEN). A 10- $\mu$ l aliquot of each purified product (50  $\mu$ l) was used as a template in the second round PCR described below. Template: 10  $\mu$ l each of the products of PCR A and B described above

45

The reaction conditions used are as follows:  
 94°C for 30 seconds; and  
 15 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 90 seconds.

**[0215]** After reaction, the primers described below were added to 50  $\mu$ l of the reaction mixture at the final concentration of 1  $\mu$ M.

50

Primers: CD03 <=> T3+  
 Reaction conditions:  
 94°C for 30 seconds;  
 35 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 30 seconds; and  
 72°C for 4 minutes.

55

**[0216]** A 1.1-kb fragment was separated by agarose gel electrophoresis, and then purified using QIAquick (QIAGEN). The fragment was cloned into pCR2.1-TOPO, and introduced into *E. coli* TOP10F'. Plasmid was prepared from the transformant using QIAprep (QIAGEN) kit, and a clone comprising a PCR error-free sequence (pTOPOCD8NKIRfull) was selected and used in the subsequent analyses.

## EP 1 672 065 A1

5 [0217] Next, 1 µg each of pCXND3 and pTOPOCD8NKIRfull were digested with 20 units each of *EcoRI* and *NotI* at 37°C for one hour, and then electrophoresed in an agarose gel. The resulting 7.8- and 1.1-kb fragments were purified using QIAquick Gel Extraction kit (QIAGEN). Then, ligation was carried out using a 1-µl aliquot of each eluate (40 µl) and LigaFAST Ligation kit (Promega) and introduced into *E. coli* strain DH5α. Ampicillin-resistant strains were saved, and the clone pCXND3CD8NKIRfull of interest that was found to carry a 1.1-kb insert was used in the subsequent analyses.

[Example 14] Construction of expression vector for CD8-KIR fusion protein

10 [0218] An expression vector for the fusion protein between the cytoplasmic ITIM motif of KIR and the extracellular domain of CD8α chain obtained in Example 12 was constructed by the following procedure. The construct is used as a positive control in the ITIM functional assay.

[0219] First, fusion PCR was carried out under the following reaction conditions. 1 st round PCR A

Template: pCD8full0113

15 Primers: CD03 (SEQ ID NO: 49) <=> CDKIR12

(5'-ATCAGAACATGCAGCAGGTGTCTTCCAGCCCCCTCGTGTGCA-3')/SEQ ID NO:52°

1 st round PCR B

Template: pBSKIR306

Primers: CDKIR11

20 (5'-TGCACACGAGGGGGCTGGACAGACACCTGCATGTTCTGAT-3') / SEQ ID NO: 53)

<=> T3+ (SEQ ID NO: 44)

25 [0220] The conditions used for both reactions are as follows:

94°C for 30 seconds;

30 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 1 minute; and

72°C for 4 minutes.

30 [0221] PCR was carried out using GC polymerase (Takara Shuzo). 0.5- and 0.4-kb fragments derived from PCR A and B, respectively, were separated by agarose gel electrophoresis, and then purified using QIAquick (QIAGEN). A 10-µl aliquot of each purified product (50 µl) was used as a template in the second round PCR described below. Template: 10 µl each of the products of PCR A and B described above

The reaction conditions used are as follows:

35 94°C for 30 minutes; and

15 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 90 seconds.

[0222] After reaction, the primers described below were added to 50 µl of the reaction mixture at the final concentration of 1 µM.

Primers: CD03 <=> T3+

40 Reaction conditions:

94°C for 30 seconds;

35 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 30 seconds; and

72°C for 4 minutes.

45 [0223] A 0.9-kb fragment was separated by agarose gel electrophoresis, and purified using QIAquick (QIAGEN). The fragment was cloned into pCR2.1-TOPO, and introduced into *E. coli* TOP10F'. Plasmid was prepared from the transformant using QIAprep (QIAGEN) kit. The 0.9-kb *EcoRI-NotI* insertion fragment from a clone comprising a PCR error-free sequence was inserted between *EcoRI* and *NotI* sites of pCXND3, and the resulting pCXND3CD8KIRfull was used in the subsequent analyses.

50 [Example 15] Establishment of a cell line stably expressing NFAT-luciferase reporter

[0224] Transformation was carried out by electroporation using the luciferase reporter plasmid pNFATlucZEOF324 obtained in Example 9 as donor DNA and Jurkat cell line as recipient cells. 20 µg of the donor DNA was pre-digested with 20 units of *PvuII* (TaKaRa) at 37°C for 1 hour. After chloroform/phenol treatment followed by ethanol precipitation, the DNA was dissolved in 20 µl of sterilized water. As recipient cells, Jurkat cells were passaged in RPMI1640 medium containing 10% FCS, washed with potassium-based phosphate buffered saline (K-PBS), and suspended in K-PBS at the cell density of 10<sup>7</sup> cells/ml to prepare 0.8 ml of a cell suspension. Electroporation was carried out using Gene Pulsar II (Bio-Rad) under the condition of 0.3 kV and 950 µFD. After pulse application, the cells were suspended in 48 ml of

passaging medium and cultured overnight under a condition without selective pressure. Then, a candidate cell line for the stable transformant was obtained under a selective pressure using zeocin (Invitrogen) at a final concentration of 100  $\mu\text{g/ml}$ . Genomic DNA was prepared from the zeocin-resistant strain using DNeasy Tissue Kit (QIAGEN). PCR was carried out using a 5- $\mu\text{l}$  aliquot of the final eluate (0.4 ml).

Premix ExTaq was used as the PCR polymerase. Luc01 (5'-TTCATACAGAAGGCGTGGAG-3' / SEQ ID NO: 45) and Luc02 (5'-CGTTCGCGGGCGCAACTGCA-3' / SEQ ID NO: 46) were used as primers. The reaction was carried out under conditions of: denaturation at 94°C for 5 minutes; 25 cycles of 94°C for 15 seconds, 55°C for 30 seconds, and 72°C for 45 seconds; followed by extension at 72°C for 5 minutes. Clones which gave a 0.5-kb PCR fragment in agarose gel electrophoresis were selected to use in the final screening by detecting luciferase activity.

**[0225]** A candidate cell line for the stable transformant with pNFATlucZEO324, which was obtained from the Jurkat cell line, was suspended at the concentration of  $6.7 \times 10^4$  cells/ml in a passaging medium containing 100  $\mu\text{g/ml}$  zeocin. A 75- $\mu\text{l}$  aliquot of the cells was plated into each well of anti-human CD3-coated microtiterplates (Beckton Dickinson) and non-coated immunomicroplates to carry out luciferase assay. After 20 hours of cultivation at 37°C, an equal volume (75  $\mu\text{l}$ ) of Dual-Glo Luciferase Reagent (Promega) was added to each well. The samples were allowed to stand for 10 minutes, and then the chemiluminescence was measured using the luminometer MicroLumat LB96P (EG&G Berthold) for 5 seconds for each well.

**[0226]** The F11 strain that exhibited intense chemiluminescence only when cultured in anti-human CD3-coated microtiterplates was selected as a stable transformant.

[Example 16] Preparation of dual transformant

**[0227]** F11 strain obtained in Example 15 as a recipient cell was transformed with pCXND3CD8NKIRfull for CD8-NKIR fusion obtained in Example 13 and pCXND3CD8KIRfull for CD8-KIR fusion obtained in Example 14 to yield a dual transformant.

**[0228]** 20  $\mu\text{g}$  each of pCXND3CD8NKIRfu11 and pCXND3CD8KIRfu11 were digested with 20 units of *PvuI* (Takara Shuzo) at 37°C for 2 hours. The digests were purified by treatment with an equal volume of phenol/chloroform (50% (v/v); Nacalai Tesque). The DNAs were precipitated using 1/10 volume of 3 M sodium acetate (Nacalai Tesque) and two volumes of ethanol (Nacalai Tesque), and then dried and dissolved in 20  $\mu\text{l}$  of sterilized water. Electroporation and subsequent establishment were achieved under the same condition as that described in Example 9, except for using as a passaging medium RPMI 1640 medium containing 100  $\mu\text{g/ml}$  zeocin, 10% (v/v) inactivated calf serum, and 1% (v/v) penicillin and streptomycin solutions and as a selection agent 700  $\mu\text{g/ml}$  geneticin. The drug-resistant clones obtained from single colonies were analyzed by FACS using anti-CD8-FITC antibody (Becton Dickinson), and 6 to 8 expression clones were selected for each.

**[0229]** Three CD8 chimeric clones (NKIR#16 and NKIR#19 as transformants with pCXND3CD8NKIRfull; and KIR#24 as a transformant with pCXND3CD8KIRfull) exhibiting reporter activity were finally selected from the transformant cell lines by the reporter activity assay described in Example 15. The structures of the three CD8 chimeric clones are shown in Fig. 11-1. Fig. 11-2 shows a result of FACS analysis of these clones using anti-CD8 antibody, LT8 (Serotec), and FITC-conjugated goat anti-mouse IgG antibody (Coulter) to be used in Example 17. All three CD8 chimeric clones were found to be stained specifically with LT8.

[Example 17] Assay of NKIR-derived ITIM activity

**[0230]** A luciferase reporter assay for NKIR-derived ITIM activity was carried out using the dual transformants NKIR#16, NKIR#19, and KIR#24 obtained in Example 16 and the host F11 strain.

**[0231]** The host F11 strain was grown in a passaging medium (RPMI 1640 medium containing 100  $\mu\text{g/ml}$  zeocin, 10% (v/v) inactivated calf serum, and 1% (v/v) penicillin and streptomycin solutions). The CD8 chimeric clones were grown in the above-described culture medium containing 700  $\mu\text{g/ml}$  geneticin. The cells were suspended in each of the growth media at the cell density of  $5.33 \times 10^5$  cells/ml. The suspensions were aliquoted into flat-bottomed 96-well plates (37.5  $\mu\text{l/well}$ ), and cultured at 37°C for 16 hours. 12.5- $\mu\text{l}$  aliquots (final concentration of 1.5  $\mu\text{g/ml}$ ) of anti-CD8 antibody (LT8, Serotec, MCA1226XZ) diluted to 6  $\mu\text{g/ml}$  with each of the growth media were added as the primary antibody to the wells. Each growth medium containing no antibody was added to the control group. After 1 hour of culture at 37°C, 12.5  $\mu\text{l}$  aliquots (final concentration of 1.2  $\mu\text{g/ml}$ ) of rabbit anti-mouse IgG1 antibody (H143.225.8, Southern Biotech, 1145-01) diluted to 6  $\mu\text{g/ml}$  with each medium were added as a cross-linker to the wells. Each growth medium containing no antibody was added to the control group. After 1 hour of cultivation at 37°C, 12.5  $\mu\text{l}$  aliquots (final concentration of 40  $\mu\text{g/ml}$ ) of ConA (SIGMA) diluted to 240  $\mu\text{g/ml}$  with each growth medium were added thereto. After 8 or 10 hours of cultivation at 37°C, an equal volume (75  $\mu\text{l}$ ) of luciferase reagent (Promega) was added to each well, and the resulting mixtures were allowed to stand at room temperature for 10 minutes. Then, the luminescence was measured by the same procedure as described in Example 15.

5 [0232] The result of the reporter assay is shown in Fig. 12. All assays were carried out in triplicats and the averages are shown in histograms. The SD values are indicated by error bars. When the results of 8 and 10 hours of ConA stimulation were compared, the effect after 10 hours was found to be higher. Nonetheless, since the two results had roughly the same tendency, the 10-hour Con A stimulation is specifically described below. The host F11 exhibited almost constant activity regardless of the presence of LT8 antibody or cross-linker. In contrast, the activity of CD8-KIRfull transformant CD8 chimeric clone KIR#24 (the positive control clone) was found to be reduced in the presence of LT8, and as was expected, the reduction in the activity was not influenced by the presence of the cross-linker. Meanwhile, both activities of the two CD8-NKIRfull-transfected CD8 chimeric clones were found to be reduced in the presence of LT8, and the reduction in the activity was not influenced by the cross-linker. While the degree of reduction for the CD-KIR transformant #24 positive control was 16.6%, that for the CD-NKIR transformants #16 and #19 was 21.2% and 30.2%, respectively. Thus, the sequence comprising the cytoplasmic ITIM motif of NKIR molecule was suggested to have ITIM activity identical to or greater than that of the known cytoplasmic ITIM of KIR2DL3 molecule.

15 Industrial Applicability

20 [0233] The present invention provides novel proteins expressed in NK cells, DNAs encoding the proteins, vectors comprising the DNAs, host cells containing the vectors, and methods for producing the proteins. The present invention also provides methods for identifying compounds which bind to the proteins or regulate the activity of the proteins. The proteins of the present invention and DNAs, and compounds that bind to the proteins of the present invention or regulate the activity of the proteins are expected to be applicable in the development of new preventive and therapeutic agents for diseases associated with the proteins of the present invention.

EP 1 672 065 A1

SEQUENCE LISTING

5 <110> CHUGAI SEIYAKU KABUSHIKI KAISHA  
 <120> PROTEIN EXPRESSED IN NK CELL  
 <130> C1-A0308P  
 10 <150> JP 2003-338331  
 <151> 2003-09-29  
 <160> 53  
 15 <170> PatentIn version 3.1  
 <210> 1  
 <211> 1473  
 <212> DNA  
 20 <213> Homo sapiens  
 <220>  
 <221> CDS  
 <222> (87)..(1376)  
 <223>  
 25 <400> 1  
 acacacccac aggacctgca gctgaacgaa gttgaagaca actcaggaga tctgttgaa 60  
 agagaacgat agaggaaaat atatga atg ttg cca tct tta gtt ccc tgt gtt 113  
 Met Leu Pro Ser Leu Val Pro Cys Val  
 30 1 5  
 ggg aaa act gtc tgg ctg tac ctc caa gcc tgg cca aac cct gtg ttt 161  
 Gly Lys Thr Val Trp Leu Tyr Leu Gln Ala Trp Pro Asn Pro Val Phe  
 10 15 20 25  
 35 gaa gga gat gcc ctg act ctg cga tgt cag gga tgg aag aat aca cca 209  
 Glu Gly Asp Ala Leu Thr Leu Arg Cys Gln Gly Trp Lys Asn Thr Pro  
 30 35 40  
 40 ctg tct cag gtg aag ttc tac aga gat gga aaa ttc ctt cat ttc tct 257  
 Leu Ser Gln Val Lys Phe Tyr Arg Asp Gly Lys Phe Leu His Phe Ser  
 45 50 55  
 aag gaa aac cag act ctg tcc atg gga gca gca aca gtg cag agc cgt 305  
 Lys Glu Asn Gln Thr Leu Ser Met Gly Ala Ala Thr Val Gln Ser Arg  
 45 60 65 70  
 ggc cag tac agc tgc tct ggg cag gtg atg tat att cca cag aca ttc 353  
 Gly Gln Tyr Ser Cys Ser Gly Gln Val Met Tyr Ile Pro Gln Thr Phe  
 75 80 85  
 50 aca caa act tca gag act gcc atg gtt caa gtc caa gag ctg ttt cca 401  
 Thr Gln Thr Ser Glu Thr Ala Met Val Gln Val Gln Glu Leu Phe Pro  
 90 95 100 105  
 55 cct cct gtg ctg agt gcc atc ccc tct cct gag ccc cga gag ggt agc 449

EP 1 672 065 A1

	Pro	Pro	Val	Leu	Ser	Ala	Ile	Pro	Ser	Pro	Glu	Pro	Arg	Glu	Gly	Ser	
					110					115					120		
5	ctg	gtg	acc	ctg	aga	tgt	cag	aca	aag	ctg	cac	ccc	ctg	agg	tca	gcc	497
	Leu	Val	Thr	Leu	Arg	Cys	Gln	Thr	Lys	Leu	His	Pro	Leu	Arg	Ser	Ala	
				125					130					135			
10	ttg	agg	ctc	ctt	ttc	tcc	ttc	cac	aag	gac	ggc	cac	acc	ttg	cag	gac	545
	Leu	Arg	Leu	Leu	Phe	Ser	Phe	His	Lys	Asp	Gly	His	Thr	Leu	Gln	Asp	
			140					145					150				
15	agg	ggc	cct	cac	cca	gaa	ctc	tgc	atc	ccg	gga	gcc	aag	gag	gga	gac	593
	Arg	Gly	Pro	His	Pro	Glu	Leu	Cys	Ile	Pro	Gly	Ala	Lys	Glu	Gly	Asp	
		155				160						165					
20	tct	ggg	ctt	tac	tgg	tgt	gag	gtg	gcc	cct	gag	ggt	ggc	cag	gtc	cag	641
	Ser	Gly	Leu	Tyr	Trp	Cys	Glu	Val	Ala	Pro	Glu	Gly	Gly	Gln	Val	Gln	
	170					175					180					185	
25	aag	cag	agc	ccc	cag	ctg	gag	gtc	aga	gtg	cag	gct	cct	gta	tcc	cgt	689
	Lys	Gln	Ser	Pro	Gln	Leu	Glu	Val	Arg	Val	Gln	Ala	Pro	Val	Ser	Arg	
					190					195					200		
30	cct	gtg	ctc	act	ctg	cac	cac	ggg	cct	gct	gac	cct	gct	gtg	ggg	gac	737
	Pro	Val	Leu	Thr	Leu	His	His	Gly	Pro	Ala	Asp	Pro	Ala	Val	Gly	Asp	
				205					210					215			
35	atg	gtg	cag	ctc	ctc	tgt	gag	gca	cag	agg	ggc	tcc	cct	ccg	atc	ctg	785
	Met	Val	Gln	Leu	Leu	Cys	Glu	Ala	Gln	Arg	Gly	Ser	Pro	Pro	Ile	Leu	
			220					225					230				
40	tat	tcc	ttc	tac	ctt	gat	gag	aag	att	gtg	ggg	aac	cac	tca	gct	ccc	833
	Tyr	Ser	Phe	Tyr	Leu	Asp	Glu	Lys	Ile	Val	Gly	Asn	His	Ser	Ala	Pro	
		235					240					245					
45	tgt	ggt	gga	acc	acc	tcc	ctc	ctc	ttc	cca	gtg	aag	tca	gaa	cag	gat	881
	Cys	Gly	Gly	Thr	Thr	Ser	Leu	Leu	Phe	Pro	Val	Lys	Ser	Glu	Gln	Asp	
	250					255					260				265		
50	gct	ggg	aac	tac	tcc	tgc	gag	gct	gag	aac	agt	gtc	tcc	aga	gag	agg	929
	Ala	Gly	Asn	Tyr	Ser	Cys	Glu	Ala	Glu	Asn	Ser	Val	Ser	Arg	Glu	Arg	
				270					275					280			
55	agt	gag	ccc	aag	aag	ctg	tct	ctg	aag	ggt	tct	caa	gtc	ttg	ttc	act	977
	Ser	Glu	Pro	Lys	Lys	Leu	Ser	Leu	Lys	Gly	Ser	Gln	Val	Leu	Phe	Thr	
				285					290					295			
60	ccc	gcc	agc	aac	tgg	ctg	ggt	cct	tgg	ctt	cct	gcg	agc	ctg	ctt	ggc	1025
	Pro	Ala	Ser	Asn	Trp	Leu	Val	Pro	Trp	Leu	Pro	Ala	Ser	Leu	Leu	Gly	
			300					305					310				
65	ctg	atg	ggt	att	gct	gct	gca	ctt	ctg	ggt	tat	gtg	aga	tcc	tgg	aga	1073
	Leu	Met	Val	Ile	Ala	Ala	Ala	Leu	Leu	Val	Tyr	Val	Arg	Ser	Trp	Arg	
		315					320					325					
70	aaa	gct	ggg	ccc	ctt	cca	tcc	cag	ata	cca	ccc	aca	gct	cca	ggt	gga	1121
	Lys	Ala	Gly	Pro	Leu	Pro	Ser	Gln	Ile	Pro	Pro	Thr	Ala	Pro	Gly	Gly	
	330					335					340				345		
75	gag	cag	tgc	cca	cta	tat	gcc	aac	gtg	cat	cac	cag	aaa	ggg	aaa	gat	1169
	Glu	Gln	Cys	Pro	Leu	Tyr	Ala	Asn	Val	His	His	Gln	Lys	Gly	Lys	Asp	
					350					355					360		

EP 1 672 065 A1

gaa ggt gtt gtc tac tct gtg gtg cat aga acc tca aag agg agt gaa 1217  
 Glu Gly Val Val Tyr Ser Val Val His Arg Thr Ser Lys Arg Ser Glu  
 365 370 375

5 gcc agg tct gct gag ttc acc gtg ggg aga aag gac agt tct atc atc 1265  
 Ala Arg Ser Ala Glu Phe Thr Val Gly Arg Lys Asp Ser Ser Ile Ile  
 380 385 390

10 tgt gcg gag gtg aga tgc ctg cag ccc agt gag gtt tca tcc acg gag 1313  
 Cys Ala Glu Val Arg Cys Leu Gln Pro Ser Glu Val Ser Ser Thr Glu  
 395 400 405

15 gtg aat atg aga agc agg act ctc caa gaa ccc ctt agc gac tgt gag 1361  
 Val Asn Met Arg Ser Arg Thr Leu Gln Glu Pro Leu Ser Asp Cys Glu  
 410 415 420 425

20 gag gtt ctc tgc tag tgatggtgtt ctctatcaa cacacgcca cccccagtct 1416  
 Glu Val Leu Cys

ccagtgctcc tcaggaagac agtgggggtcc tcaactcttt ctgtgggtcc ttcagtg 1473

<210> 2  
 <211> 429  
 <212> PRT  
 <213> Homo sapiens

<400> 2  
 Met Leu Pro Ser Leu Val Pro Cys Val Gly Lys Thr Val Trp Leu Tyr  
 1 5 10 15

30 Leu Gln Ala Trp Pro Asn Pro Val Phe Glu Gly Asp Ala Leu Thr Leu  
 20 25 30

Arg Cys Gln Gly Trp Lys Asn Thr Pro Leu Ser Gln Val Lys Phe Tyr  
 35 40 45

40 Arg Asp Gly Lys Phe Leu His Phe Ser Lys Glu Asn Gln Thr Leu Ser  
 50 55 60

Met Gly Ala Ala Thr Val Gln Ser Arg Gly Gln Tyr Ser Cys Ser Gly  
 65 70 75 80

45 Gln Val Met Tyr Ile Pro Gln Thr Phe Thr Gln Thr Ser Glu Thr Ala  
 85 90 95

Met Val Gln Val Gln Glu Leu Phe Pro Pro Pro Val Leu Ser Ala Ile  
 100 105 110

50 Pro Ser Pro Glu Pro Arg Glu Gly Ser Leu Val Thr Leu Arg Cys Gln  
 115 120 125

Thr Lys Leu His Pro Leu Arg Ser Ala Leu Arg Leu Leu Phe Ser Phe  
 130 135 140

55 His Lys Asp Gly His Thr Leu Gln Asp Arg Gly Pro His Pro Glu Leu  
 145 150 155 160

Cys Ile Pro Gly Ala Lys Glu Gly Asp Ser Gly Leu Tyr Trp Cys Glu  
 165 170 175

EP 1 672 065 A1

```

Val Ala Pro Glu Gly Gly Gln Val Gln Lys Gln Ser Pro Gln Leu Glu
180                               185                               190
5  Val Arg Val Gln Ala Pro Val Ser Arg Pro Val Leu Thr Leu His His
195                               200                               205

Gly Pro Ala Asp Pro Ala Val Gly Asp Met Val Gln Leu Leu Cys Glu
210                               215                               220

10 Ala Gln Arg Gly Ser Pro Pro Ile Leu Tyr Ser Phe Tyr Leu Asp Glu
225                               230                               235

Lys Ile Val Gly Asn His Ser Ala Pro Cys Gly Gly Thr Thr Ser Leu
245                               250                               255

15 Leu Phe Pro Val Lys Ser Glu Gln Asp Ala Gly Asn Tyr Ser Cys Glu
260                               265                               270

Ala Glu Asn Ser Val Ser Arg Glu Arg Ser Glu Pro Lys Lys Leu Ser
275                               280                               285

20 Leu Lys Gly Ser Gln Val Leu Phe Thr Pro Ala Ser Asn Trp Leu Val
290                               295                               300

Pro Trp Leu Pro Ala Ser Leu Leu Gly Leu Met Val Ile Ala Ala Ala
305                               310                               315                               320

25 Leu Leu Val Tyr Val Arg Ser Trp Arg Lys Ala Gly Pro Leu Pro Ser
325                               330                               335

Gln Ile Pro Pro Thr Ala Pro Gly Gly Glu Gln Cys Pro Leu Tyr Ala
340                               345                               350

30 Asn Val His His Gln Lys Gly Lys Asp Glu Gly Val Val Tyr Ser Val
355                               360                               365

Val His Arg Thr Ser Lys Arg Ser Glu Ala Arg Ser Ala Glu Phe Thr
370                               375                               380

35 Val Gly Arg Lys Asp Ser Ser Ile Ile Cys Ala Glu Val Arg Cys Leu
385                               390                               395                               400

Gln Pro Ser Glu Val Ser Ser Thr Glu Val Asn Met Arg Ser Arg Thr
405                               410                               415

40 Leu Gln Glu Pro Leu Ser Asp Cys Glu Glu Val Leu Cys
420                               425

<210> 3
<211> 2005
45 <212> DNA
<213> Homo sapiens

<220>
<221> CDS
50 <222> (88)..(1410)
<223>

<400> 3
cacacacca caggacctgc agctgaacga agttgaagac aactcaggag atctgttggg      60
55 aagagaacga .tagaggaaaa tatatga atg ttg cca tct tta ggc ccc atg ctg      114

```

EP 1 672 065 A1

Met Leu Pro Ser Leu Gly Pro Met Leu  
1 5

5	ctc tgg acg gct gtg ctg ctc ttt gtt ccc tgt gtt ggg aaa act gtc Leu Trp Thr Ala Val Leu Leu Phe Val Pro Cys Val Gly Lys Thr Val 10 15 20 25	162
10	tgg ctg tac ctc caa gcc tgg cca aac cct gtg ttt gaa gga gat gcc Trp Leu Tyr Leu Gln Ala Trp Pro Asn Pro Val Phe Glu Gly Asp Ala 30 35 40	210
15	ctg act ctg cga tgt cag gga tgg aag aat aca cca ctg tct cag gtg Leu Thr Leu Arg Cys Gln Gly Trp Lys Asn Thr Pro Leu Ser Gln Val 45 50 55	258
20	aag ttc tac aga gat gga aaa ttc ctt cat ttc tct aag gaa aac cag Lys Phe Tyr Arg Asp Gly Lys Phe Leu His Phe Ser Lys Glu Asn Gln 60 65 70	306
25	act ctg tcc atg gga gca gca aca gtg cag agc cgt ggc cag tac agc Thr Leu Ser Met Gly Ala Ala Thr Val Gln Ser Arg Gly Gln Tyr Ser 75 80 85	354
30	tgc tct ggg cag gtg atg tat att cca cag aca ttc aca caa act tca Cys Ser Gly Gln Val Met Tyr Ile Pro Gln Thr Phe Thr Gln Thr Ser 90 95 100 105	402
35	gag act gcc atg gtt caa gtc caa gag ctg ttt cca cct cct gtg ctg Glu Thr Ala Met Val Gln Val Gln Glu Leu Phe Pro Pro Pro Val Leu 110 115 120	450
40	agt gcc atc ccc tct cct gag ccc cga gag ggt agc ctg gtg acc ctg Ser Ala Ile Pro Ser Pro Glu Pro Arg Glu Gly Ser Leu Val Thr Leu 125 130 135	498
45	aga tgt cag aca aag ctg cac ccc ctg agg tca gcc ttg agg ctc ctt Arg Cys Gln Thr Lys Leu His Pro Leu Arg Ser Ala Leu Arg Leu Leu 140 145 150	546
50	ttc tcc ttc cac aag gac ggc cac acc ttg cag gac agg ggc cct cac Phe Ser Phe His Lys Asp Gly His Thr Leu Gln Asp Arg Gly Pro His 155 160 165	594
55	cca gaa ctc tgc atc ccg gga gcc aag gag gga gac tct ggg ctt tac Pro Glu Leu Cys Ile Pro Gly Ala Lys Glu Gly Asp Ser Gly Leu Tyr 170 175 180 185	642
60	tgg tgt gag gtg gcc cct gag ggt ggc cag gtc cag aag cag agc ccc Trp Cys Glu Val Ala Pro Glu Gly Gly Gln Val Gln Lys Gln Ser Pro 190 195 200	690
65	cag ctg gag gtc aga gtg cag gct cct gta tcc cgt cct gtg ctc act Gln Leu Glu Val Arg Val Gln Ala Pro Val Ser Arg Pro Val Leu Thr 205 210 215	738
70	ctg cac cac ggg cct gct gac ccc gct gtg ggg gac atg gtg cag ctc Leu His His Gly Pro Ala Asp Pro Ala Val Gly Asp Met Val Gln Leu 220 225 230	786
75	ctc tgt gag gca cag agg ggc tcc cct ccg atc ctg tat tcc ttc tac Leu Cys Glu Ala Gln Arg Gly Ser Pro Pro Ile Leu Tyr Ser Phe Tyr 235 240 245	834

EP 1 672 065 A1

ctt gat gag aag att gtg ggg aac cac tca gct ccc tgt ggt gga acc 882  
 Leu Asp Glu Lys Ile Val Gly Asn His Ser Ala Pro Cys Gly Gly Thr  
 250 255 260 265  
 5 acc tcc ctc ctc ttc cca gtg aag tca gaa cag gat gct ggg aac tac 930  
 Thr Ser Leu Leu Phe Pro Val Lys Ser Glu Gln Asp Ala Gly Asn Tyr  
 270 275 280  
 10 tcc tgc gag gct gag aac agt gtc tcc aga gag agg agt gag ccc aag 978  
 Ser Cys Glu Ala Glu Asn Ser Val Ser Arg Glu Arg Ser Glu Pro Lys  
 285 290 295  
 aag ctg tct ctg aag ggt tct caa gtc ttg ttc act ccc gcc agc aac 1026  
 Lys Leu Ser Leu Lys Gly Ser Gln Val Leu Phe Thr Pro Ala Ser Asn  
 300 305 310  
 15 tgg ctg gtt cct tgg ctt cct gcg agc ctg ctt ggc ctg atg gtt att 1074  
 Trp Leu Val Pro Trp Leu Pro Ala Ser Leu Leu Gly Leu Met Val Ile  
 315 320 325  
 20 gct gct gca ctt ctg gtt tat gtg aga tcc tgg aga aaa gct ggg ccc 1122  
 Ala Ala Ala Leu Leu Val Tyr Val Arg Ser Trp Arg Lys Ala Gly Pro  
 330 335 340 345  
 ctt cca tcc cag ata cca ccc aca gct cca ggt gga gag cag tgc cca 1170  
 Leu Pro Ser Gln Ile Pro Pro Thr Ala Pro Gly Gly Glu Gln Cys Pro  
 350 355 360  
 25 cta tat gcc aac gtg cat cac cag aaa ggg aaa gat gaa ggt gtt gtc 1218  
 Leu Tyr Ala Asn Val His His Gln Lys Gly Lys Asp Glu Gly Val Val  
 365 370 375  
 30 tac tct gtg gtg cat aga acc tca aag agg agt gaa gcc agg tct gct 1266  
 Tyr Ser Val Val His Arg Thr Ser Lys Arg Ser Glu Ala Arg Ser Ala  
 380 385 390  
 gag ttc acc gtg ggg aga aag gac agt tct atc atc tgt gcg gag gtg 1314  
 Glu Phe Thr Val Gly Arg Lys Asp Ser Ser Ile Ile Cys Ala Glu Val  
 395 400 405  
 35 aga tgc ctg cag ccc agt gag gtt tca tcc acg gag gtg aat atg aga 1362  
 Arg Cys Leu Gln Pro Ser Glu Val Ser Ser Thr Glu Val Asn Met Arg  
 410 415 420 425  
 40 agc agg act ctc caa gaa ccc ctt agc gac tgt gag gag gtt ctc tgc 1410  
 Ser Arg Thr Leu Gln Glu Pro Leu Ser Asp Cys Glu Glu Val Leu Cys  
 430 435 440  
 tagtgatggt gttctcctat caacacacgc ccacccccag tctccagtgc tcctcaggaa 1470  
 45 gacagtgggg tcctcaactc tttctgtggg tccttcagtg tccaagccc agcatcacag 1530  
 agccccctga gcccttgctc tggtcaggag cacctgaacc ctgggttctt ttcttagcag 1590  
 aagaccaacc aatggaatgg gaagggagat gctcccacca acacacacac ttaggttcaa 1650  
 50 tcagtgcacac tggacacata agccacagat gtcttctttc catacaagca tgtagttcg 1710  
 ccccaatata catatatata tgaatatagtc atgtgcccga taacaacatt tcagtccagt 1770  
 atagactgca tacacaacag tggtoccata agactgtaat ggagtttaaa aattcctact 1830  
 55 gcctagtgat atcatagttg ccttaacatc ataacacaac acatttctca cgcgtttgtg 1890

EP 1 672 065 A1

gtgatgctgg tacaacaag ctacagcgcc gctagtcata tacaatatata gcacatacaa 1950  
 5 ttatgtacag tacactatac ttgataatga taataaacia ctatgttact ggttt 2005

<210> 4  
 <211> 441  
 <212> PRT  
 <213> Homo sapiens

<400> 4  
 Met Leu Pro Ser Leu Gly Pro Met Leu Leu Trp Thr Ala Val Leu Leu  
 1 5 10 15  
 Phe Val Pro Cys Val Gly Lys Thr Val Trp Leu Tyr Leu Gln Ala Trp  
 20 25 30  
 Pro Asn Pro Val Phe Glu Gly Asp Ala Leu Thr Leu Arg Cys Gln Gly  
 35 40 45  
 20 Trp Lys Asn Thr Pro Leu Ser Gln Val Lys Phe Tyr Arg Asp Gly Lys  
 50 55 60  
 Phe Leu His Phe Ser Lys Glu Asn Gln Thr Leu Ser Met Gly Ala Ala  
 65 70 75 80  
 25 Thr Val Gln Ser Arg Gly Gln Tyr Ser Cys Ser Gly Gln Val Met Tyr  
 85 90 95  
 Ile Pro Gln Thr Phe Thr Gln Thr Ser Glu Thr Ala Met Val Gln Val  
 100 105 110  
 30 Gln Glu Leu Phe Pro Pro Pro Val Leu Ser Ala Ile Pro Ser Pro Glu  
 115 120 125  
 Pro Arg Glu Gly Ser Leu Val Thr Leu Arg Cys Gln Thr Lys Leu His  
 130 135 140  
 35 Pro Leu Arg Ser Ala Leu Arg Leu Leu Phe Ser Phe His Lys Asp Gly  
 145 150 155 160  
 His Thr Leu Gln Asp Arg Gly Pro His Pro Glu Leu Cys Ile Pro Gly  
 165 170 175  
 40 Ala Lys Glu Gly Asp Ser Gly Leu Tyr Trp Cys Glu Val Ala Pro Glu  
 180 185 190  
 Gly Gly Gln Val Gln Lys Gln Ser Pro Gln Leu Glu Val Arg Val Gln  
 195 200 205  
 45 Ala Pro Val Ser Arg Pro Val Leu Thr Leu His His Gly Pro Ala Asp  
 210 215 220  
 50 Pro Ala Val Gly Asp Met Val Gln Leu Leu Cys Glu Ala Gln Arg Gly  
 225 230 235 240  
 Ser Pro Pro Ile Leu Tyr Ser Phe Tyr Leu Asp Glu Lys Ile Val Gly  
 245 250 255  
 55 Asn His Ser Ala Pro Cys Gly Gly Thr Thr Ser Leu Leu Phe Pro Val



EP 1 672 065 A1

	35		40		45															
5	act	tta	cag	ctc	ttt	tac	act	ttc	tac	aag	gac	aac	cat	gtc	att	caa				309
	Thr	Leu	Gln	Leu	Phe	Tyr	Thr	Phe	Tyr	Lys	Asp	Asn	His	Val	Ile	Gln				
	50				55						60				65					
	gac	agg	agt	ccc	cac	tca	gta	ttt	tct	gca	gaa	gcc	aag	gag	gaa	aac				357
	Asp	Arg	Ser	Pro	His	Ser	Val	Phe	Ser	Ala	Glu	Ala	Lys	Glu	Glu	Asn				
				70						75					80					
10	tct	ggg	ctc	tac	cag	tgt	atg	gtg	gac	act	gag	gat	ggc	tta	att	cag				405
	Ser	Gly	Leu	Tyr	Gln	Cys	Met	Val	Asp	Thr	Glu	Asp	Gly	Leu	Ile	Gln				
			85						90					95						
15	aaa	aaa	agt	ggc	tat	ctg	gat	atc	cag	ttc	tgg	act	cct	gta	tcc	cat				453
	Lys	Lys	Ser	Gly	Tyr	Leu	Asp	Ile	Gln	Phe	Trp	Thr	Pro	Val	Ser	His				
			100					105					110							
	cct	gtg	ctc	act	ctg	caa	cat	gaa	gcc	act	aac	ctt	gct	gta	gga	gac				501
	Pro	Val	Leu	Thr	Leu	Gln	His	Glu	Ala	Thr	Asn	Leu	Ala	Val	Gly	Asp				
			115				120					125								
20	aag	gtg	gag	ttc	ctc	tgt	gag	gcc	cac	cag	ggc	tcc	ctt	cca	atc	ttt				549
	Lys	Val	Glu	Phe	Leu	Cys	Glu	Ala	His	Gln	Gly	Ser	Leu	Pro	Ile	Phe				
					135						140				145					
25	tac	tca	ttc	tac	att	aat	gga	gaa	atc	cta	ggg	aaa	ccc	ctg	gct	ccc				597
	Tyr	Ser	Phe	Tyr	Ile	Asn	Gly	Glu	Ile	Leu	Gly	Lys	Pro	Leu	Ala	Pro				
				150					155					160						
	tct	ggc	aga	gct	gcc	tcc	ctc	cta	gcc	tca	gta	aag	gca	gag	tgg	agt				645
	Ser	Gly	Arg	Ala	Ala	Ser	Leu	Leu	Ala	Ser	Val	Lys	Ala	Glu	Trp	Ser				
			165					170						175						
30	acc	aag	aac	tat	tcc	tgt	gaa	gct	aaa	aac	aac	atc	tcc	aga	gaa	ata				693
	Thr	Lys	Asn	Tyr	Ser	Cys	Glu	Ala	Lys	Asn	Asn	Ile	Ser	Arg	Glu	Ile				
			180					185					190							
35	agt	gag	ctc	aag	aag	ttc	ccc	ttg	ggt	gtc	tca	ggt	act	gcc	tgg	atc				741
	Ser	Glu	Leu	Lys	Lys	Phe	Pro	Leu	Val	Val	Ser	Gly	Thr	Ala	Trp	Ile				
			195			200						205								
	aag	agc	aac	atg	cta	act	atc	tgg	cta	cct	gca	agc	ctg	ctt	gga	ggg				789
	Lys	Ser	Asn	Met	Leu	Thr	Ile	Trp	Leu	Pro	Ala	Ser	Leu	Leu	Gly	Gly				
			210			215					220				225					
40	atg	gtc	att	gcg	gct	gtg	ggt	cta	atg	tat	ttc	ttc	aaa	ccc	tgt	aaa				837
	Met	Val	Ile	Ala	Ala	Val	Val	Leu	Met	Tyr	Phe	Phe	Lys	Pro	Cys	Lys				
				230						235					240					
45	aag	cat	gcc	aga	cct	gag	atg	ccc	acc	cta	aaa	gag	cca	gac	agt	ttt				885
	Lys	His	Ala	Arg	Pro	Glu	Met	Pro	Thr	Leu	Lys	Glu	Pro	Asp	Ser	Phe				
			245					250						255						
	cta	tat	gta	tcg	ggt	gat	aat	cga	aga	tat	aaa	tga	gattcccacc							931
	Leu	Tyr	Val	Ser	Val	Asp	Asn	Arg	Arg	Tyr	Lys									
50			260					265												
	aatgatttgg attcaaaaac caggacctgc caagatcccc ttggtcttta ggatcatgct																			991
	ctgtgtagt gcaatgtctt cctccagcat atactcaact ccagctccca gcctccacc																			1051
55	tccagcactc agcagtggtc ccaagttctc cctgcaggtc acccagttcc tagcccagca																			1111

EP 1 672 065 A1

gtgaggaagc ccatatgctc tattcctggc cagggctcct gaactgtggg ttctcttctg 1171  
 5 agcgggaaac caaacaatgg tgtgggaatg aacaatttcc accttgatac atacatatac 1231  
 acatgcacac acacaaacaa acacacatac acacacactt ccagatgtaa cattgtacac 1291  
 agagccacag ttatcttctt taagtacaaa aggaaaaggg ttttcacctc cagatagaca 1351  
 10 gataatagat acacagacac acaagacaga tagatgatag ataacatata gattagatag 1411  
 ataatagata gatggtagat aggtagatgg atgatagata gatagataga ttggatagat 1471  
 agatagatag atagatagat agatagatag ataataacat gacagataag atgatagaaa 1531  
 15 taagatacga ta 1543

<210> 6  
 <211> 268  
 <212> PRT  
 20 <213> Mus musculus

<400> 6  
 Met Leu Leu Trp Met Val Leu Leu Leu Cys Asp Ser Met Val Glu Ala  
 1 5 10 15  
 25 Gln Glu Leu Phe Pro Asn Pro Glu Leu Thr Glu Phe Thr Asn Ser Glu  
 20 25 30  
 Thr Met Asp Val Ile Leu Lys Cys Thr Ile Lys Val Asp Pro Lys Asn  
 35 40 45  
 30 Pro Thr Leu Gln Leu Phe Tyr Thr Phe Tyr Lys Asp Asn His Val Ile  
 50 55 60  
 Gln Asp Arg Ser Pro His Ser Val Phe Ser Ala Glu Ala Lys Glu Glu  
 65 70 75 80  
 35 Asn Ser Gly Leu Tyr Gln Cys Met Val Asp Thr Glu Asp Gly Leu Ile  
 85 90 95  
 Gln Lys Lys Ser Gly Tyr Leu Asp Ile Gln Phe Trp Thr Pro Val Ser  
 100 105 110  
 40 His Pro Val Leu Thr Leu Gln His Glu Ala Thr Asn Leu Ala Val Gly  
 115 120 125  
 Asp Lys Val Glu Phe Leu Cys Glu Ala His Gln Gly Ser Leu Pro Ile  
 130 135 140  
 45 Phe Tyr Ser Phe Tyr Ile Asn Gly Glu Ile Leu Gly Lys Pro Leu Ala  
 145 150 155 160  
 Pro Ser Gly Arg Ala Ala Ser Leu Leu Ala Ser Val Lys Ala Glu Trp  
 165 170 175  
 50 Ser Thr Lys Asn Tyr Ser Cys Glu Ala Lys Asn Asn Ile Ser Arg Glu  
 180 185 190  
 Ile Ser Glu Leu Lys Lys Phe Pro Leu Val Val Ser Gly Thr Ala Trp  
 195 200 205  
 55

EP 1 672 065 A1

Ile Lys Ser Asn Met Leu Thr Ile Trp Leu Pro Ala Ser Leu Leu Gly  
 210 215 220

5 Gly Met Val Ile Ala Ala Val Val Leu Met Tyr Phe Phe Lys Pro Cys  
 225 230 235 240

Lys Lys His Ala Arg Pro Glu Met Pro Thr Leu Lys Glu Pro Asp Ser  
 245 250 255

10 Phe Leu Tyr Val Ser Val Asp Asn Arg Arg Tyr Lys  
 260 265

15 <210> 7  
 <211> 34  
 <212> DNA  
 <213> Artificial

<220>  
 <223> an artificially synthesized primer sequence

20 <400> 7  
 ttgaattcac acaccacag gacctgcagc tga . 34

25 <210> 8  
 <211> 34  
 <212> DNA  
 <213> Artificial

<220>  
 <223> an artificially synthesized primer sequence

30 <400> 8  
 ttggatccac tgaagaccc acagaaagag ttga 34

35 <210> 9  
 <211> 21  
 <212> DNA  
 <213> Artificial

<220>  
 <223> an artificially synthesized primer sequence

40 <400> 9  
 accctgagat gtcagacaaa g 21

45 <210> 10  
 <211> 20  
 <212> DNA  
 <213> Artificial

<220>  
 <223> an artificially synthesized primer sequence

50 <400> 10  
 gccacctcac accagtaaag 20

55 <210> 11

EP 1 672 065 A1

<211> 21  
<212> DNA  
<213> Artificial

5  
<220>  
<223> an artificially synthesized primer sequence

<400> 11  
cctccgatcc tgtattcctt c 21

10  
<210> 12  
<211> 21  
<212> DNA  
<213> Artificial

15  
<220>  
<223> an artificially synthesized primer sequence

<400> 12  
tggagctgtg ggtggtatct g 21

20  
<210> 13  
<211> 21  
<212> DNA  
<213> Artificial

25  
<220>  
<223> an artificially synthesized primer sequence

<400> 13  
agaacctcaa agaggagtga a 21

30  
<210> 14  
<211> 20  
<212> DNA  
<213> Artificial

35  
<220>  
<223> an artificially synthesized primer sequence

<400> 14  
attatgctga gtgatatccc 20

40  
<210> 15  
<211> 20  
<212> DNA  
<213> Artificial

45  
<220>  
<223> an artificially synthesized primer sequence

<400> 15  
atttagtga cactatagaa 20

50  
<210> 16  
<211> 28  
<212> DNA  
<213> Artificial

55

EP 1 672 065 A1

<220>  
<223> an artificially synthesized primer sequence

5 <400> 16  
gggaattcat gttgccatct ttagttcc 28

<210> 17  
<211> 26  
10 <212> DNA  
<213> Artificial

<220>  
<223> an artificially synthesized primer sequence

15 <400> 17  
aaggatccac tcctctctct ggagac 26

<210> 18  
20 <211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> an artificially synthesized primer sequence

25 <400> 18  
gcctcagaca gtggttcaaa 20

<210> 19  
30 <211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> an artificially synthesized primer sequence

35 <400> 19  
agaaccatca cagtctcgca 20

<210> 20  
40 <211> 29  
<212> DNA  
<213> Artificial

<220>  
<223> an artificially synthesized primer sequence

45 <400> 20  
aagaattcca ccatggctgg acctgccac 29

<210> 21  
50 <211> 41  
<212> DNA  
<213> Artificial

<220>  
55 <223> an artificially synthesized primer sequence

EP 1 672 065 A1

5 <400> 21  
acagggtttg gccaggettg ggcttctgc actgtccaga g 41

<210> 22  
<211> 30  
<212> DNA  
<213> Artificial

10 <220>  
<223> an artificially synthesized primer sequence

<400> 22  
gcaggaagcc caagcctggc caaacctgt 30

15 <210> 23  
<211> 45  
<212> DNA  
<213> Artificial

20 <220>  
<223> an artificially synthesized primer sequence

<400> 23  
ctaatacgac tcactatagg gcaagcagtg gtatcaacgc agagt 45

25 <210> 24  
<211> 22  
<212> DNA  
<213> Artificial

30 <220>  
<223> an artificially synthesized primer sequence

<400> 24  
ctaatacgac tcactatagg gc 22

35 <210> 25  
<211> 23  
<212> DNA  
<213> Artificial

40 <220>  
<223> an artificially synthesized primer sequence

<400> 25  
aagcagtggg atcaacgcag agt 23

45 <210> 26  
<211> 33  
<212> DNA  
<213> Artificial

50 <220>  
<223> an artificially synthesized primer sequence

<400> 26  
ctcggatcct tgccatcttt agttccctgt gtt 33

55

EP 1 672 065 A1

5 <210> 27  
 <211> 36  
 <212> DNA  
 <213> Artificial  
  
 <220>  
 <223> an artificially synthesized primer sequence  
  
 10 <400> 27  
 gctgtcgact tagttgctgg cgggagtgaa caagac 36  
  
 15 <210> 28  
 <211> 63  
 <212> DNA  
 <213> Artificial  
  
 <220>  
 <223> an artificially synthesized primer sequence  
  
 20 <400> 28  
 gcgaattcca ccatggacta caaagacgat gacgacaagt tgccatcttt agttccctgt 60  
  
 gtt 63  
  
 25 <210> 29  
 <211> 36  
 <212> DNA  
 <213> Artificial  
  
 30 <220>  
 <223> an artificially synthesized primer sequence  
  
 <400> 29  
 cgtgtcgact cactagcaga gaacctcctc acagtc 36  
  
 35 <210> 30  
 <211> 25  
 <212> DNA  
 <213> Artificial  
  
 40 <220>  
 <223> an artificially synthesized primer sequence  
  
 <400> 30  
 aggtcagagt gcaggctcct gtatc 25  
  
 45 <210> 31  
 <211> 25  
 <212> DNA  
 <213> Artificial  
  
 50 <220>  
 <223> an artificially synthesized primer sequence  
  
 <400> 31  
 tagaactgtc ctttctcccc acggt 25  
  
 55

EP 1 672 065 A1

5 <210> 32  
<211> 26  
<212> DNA  
<213> Artificial  
  
<220>  
<223> an artificially synthesized primer sequence  
  
<400> 32  
10 gaattcacac acccacagga cctgca 26  
  
<210> 33  
<211> 26  
<212> DNA  
15 <213> Artificial  
  
<220>  
<223> an artificially synthesized primer sequence  
  
<400> 33  
20 ggatccactg aaggaccac agaaag 26  
  
<210> 34  
<211> 25  
<212> DNA  
25 <213> Artificial  
  
<220>  
<223> an artificially synthesized primer sequence  
  
<400> 34  
30 ctcagtaaag gcagagtgga gtacc 25  
  
<210> 35  
<211> 25  
<212> DNA  
35 <213> Artificial  
  
<220>  
<223> an artificially synthesized primer sequence  
  
<400> 35  
40 atacattaga accacagccg caatg 25  
  
<210> 36  
<211> 27  
<212> DNA  
45 <213> Artificial  
  
<220>  
<223> an artificially synthesized primer sequence  
  
<400> 36  
50 ccatcctaatac gactcact atagggc 27  
  
<210> 37  
55 <211> 23

EP 1 672 065 A1

<212> DNA  
<213> Artificial

5 <220>  
<223> an artificially synthesized primer sequence

<400> 37  
actcactata gggctcgagc ggc 23

10 <210> 38  
<211> 26  
<212> DNA  
<213> Artificial

15 <220>  
<223> an artificially synthesized primer sequence

<400> 38  
ctcaagaagt tccccttggt tgtctc 26

20 <210> 39  
<211> 26  
<212> DNA  
<213> Artificial

25 <220>  
<223> an artificially synthesized primer sequence

<400> 39  
gccagatagt tagcatgttg ctcttg 26

30 <210> 40  
<211> 26  
<212> DNA  
<213> Artificial

35 <220>  
<223> an artificially synthesized primer sequence

<400> 40  
gaattcatgt cgctcatggt cgtcag 26

40 <210> 41  
<211> 26  
<212> DNA  
<213> Artificial

45 <220>  
<223> an artificially synthesized primer sequence

<400> 41  
ggatcctcag ggctcagcat ttggaa 26

50 <210> 42  
<211> 40  
<212> DNA  
<213> Artificial

55

EP 1 672 065 A1

<220>  
<223> an artificially synthesized primer sequence

5 <400> 42  
aggggcccag cttttctcca gcgatgaagg agaaagaaga 40

<210> 43  
<211> 40  
10 <212> DNA  
<213> Artificial

<220>  
<223> an artificially synthesized primer sequence

15 <400> 43  
tcttctttct cttcatcgc tggagaaaag ctgggccct 40

<210> 44  
<211> 25  
20 <212> DNA  
<213> Artificial

<220>  
<223> an artificially synthesized primer sequence

25 <400> 44  
gcaattaacc ctactaaag ggaac 25

<210> 45  
<211> 20  
30 <212> DNA  
<213> Artificial

<220>  
<223> an artificially synthesized primer sequence

35 <400> 45  
ttcatacaga aggcgtggag 20

<210> 46  
<211> 20  
40 <212> DNA  
<213> Artificial

<220>  
<223> an artificially synthesized primer sequence

45 <400> 46  
cgttcgcggg cgcaactgca 20

<210> 47  
<211> 26  
50 <212> DNA  
<213> Artificial

<220>  
<223> an artificially synthesized primer sequence

55

EP 1 672 065 A1

<400> 47  
gaattcatgg ccttaccagt gaccgc 26

5  
<210> 48  
<211> 26  
<212> DNA  
<213> Artificial

10  
<220>  
<223> an artificially synthesized primer sequence

<400> 48  
ggatccttag acgtatctcg ccgaaa 26

15  
<210> 49  
<211> 30  
<212> DNA  
<213> Artificial

20  
<220>  
<223> an artificially synthesized primer sequence

<400> 49  
gaattccacc atggccttac cagtgaccgc 30

25  
<210> 50  
<211> 40  
<212> DNA  
<213> Artificial

30  
<220>  
<223> an artificially synthesized primer sequence

<400> 50  
accagccagt tgctggcggg gtccagcccc ctcggtgtgca 40

35  
<210> 51  
<211> 40  
<212> DNA  
<213> Artificial

40  
<220>  
<223> an artificially synthesized primer sequence

<400> 51  
tgcacacgag ggggctggac cccgccagca actggctggt 40

45  
<210> 52  
<211> 39  
<212> DNA  
<213> Artificial

50  
<220>  
<223> an artificially synthesized primer sequence

<400> 52  
atcagaacat gcaggtgtct tccagcccc tcggtgtgca 39

55

5  
 <210> 53  
 <211> 40  
 <212> DNA  
 <213> Artificial

<220>  
 <223> an artificially synthesized primer sequence

10  
 <400> 53  
 tgcacacgag ggggctggac agacacctgc atgttctgat 40

15

**Claims**

- 20  
 1. A DNA of any one of:
- 25  
 (a) a DNA encoding a protein comprising the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6;  
 (b) a DNA comprising a coding region of the nucleotide sequence of any one of SEQ ID NOs: 1, 3, and 5;  
 (c) a DNA encoding a protein which comprises an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6, and which is functionally equivalent to a protein comprising the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6; and  
 (d) a DNA which hybridizes under stringent conditions to a DNA comprising the nucleotide sequence of any one of SEQ ID NOs: 1, 3, and 5.
- 30  
 2. A DNA encoding a fragment of a protein comprising the amino acid sequence of any one of SEQ ID NOs: 2, 4, and 6.
3. A protein encoded by the DNA of claim 1 or 2.
- 35  
 4. A vector comprising the DNA of claim 1 or 2.
5. A host cell comprising the DNA of claim 1 or 2, or the vector of claim 4.
6. A method for producing the protein of claim 3, wherein the method comprises the steps of:
- 40  
 culturing the host cell of claim 5, and  
 collecting the protein from the host cell or a culture supernatant thereof.
7. An antibody which binds to the protein of claim 3.
- 45  
 8. A method for identifying a ligand for the protein of claim 3, wherein the method comprises the steps of:
- (a) contacting a candidate compound with the protein of claim 3 or a cell expressing the protein of claim 3; and  
 (b) determining whether the candidate compound binds to the protein of claim 3 or the cell expressing the protein of claim 3.
- 50  
 9. A method for identifying an agonist for the protein of claim 3, wherein the method comprises the steps of:
- (a) contacting a candidate compound with a cell expressing the protein of claim 3; and  
 (b) determining whether the candidate compound generates a signal that is an indicator of activation of the protein of claim 3.
- 55  
 10. A method for identifying an antagonist for the protein of claim 3, wherein the method comprises the steps of:

**EP 1 672 065 A1**

- (a) contacting a candidate compound with a cell expressing the protein of claim 3; and
- (b) determining whether a signal as an indicator of activation of the protein of claim 3 is reduced as compared with a detection result obtained in absence of the candidate compound.

- 5   **11.** A ligand for the protein of claim 3, which can be identified by the method of claim 8.
- 12.** An agonist for the protein of claim 3, which can be identified by the method of claim 9.
- 13.** An antagonist for the protein of claim 3, which can be identified by the method of claim 10.
- 10   **14.** A kit to be used in the method of any one of claims 8 to 10, wherein the kit comprises at least one of:
- (a) the protein of claim 3; and
  - (b) the host cell of claim 5.
- 15   **15.** An immunosuppressant, which comprises as an active ingredient an agonist for the protein of claim 3.
- 16.** A therapeutic agent for an allergic disease or autoimmune disease, wherein the method comprises as an active ingredient an agonist for the protein of claim 3.
- 20   **17.** An immunopotentiator, which comprises as an active ingredient an antagonist for the protein of claim 3.
- 18.** An anti-tumor or antiviral agent, which comprises as an active ingredient an antagonist for the protein of claim 3.
- 25   **19.** A DNA, which comprises at least 15 nucleotides and which is complementary to the DNA of claim 1 or 2, or to a complementary strand thereof.
- 20.** A diagnostic reagent for a disease associated with an abnormality in the expression or activity of a gene encoding the protein of claim 3, wherein the diagnostic reagent comprises the DNA of claim 19 or the antibody of claim 7.
- 30

35

40

45

50

55

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2004/014207

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> Int.Cl <sup>7</sup> C12N15/09, A61K45/00, A61P1/16, A61P11/06, A61P31/12, A61P31/14, A61P35/00, A61P37/04, A61P37/06, A61P37/08, C07K14/705, C07K16/28, C12N1/15, C12N1/19, C12N1/21, C12N5/00, According to International Patent Classification (IPC) or to both national classification and IPC										
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) Int.Cl <sup>7</sup> C12N15/09, A61K45/00, C07K14/705, C07K16/28, C12N1/15, C12N1/19, C12N1/21, C12N5/00, C12P21/02, C12Q1/02, G01N33/15, G01N33/50  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched										
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CA/MEDLINE/BIOSIS/EMBASE/BIOTECHABS/WPIDS (STN), GenBank/EMBL/DDBJ/GeneSeq, SwissProt/PIR/GeneSeq										
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>										
<table border="1"> <thead> <tr> <th>Category*</th> <th>Citation of document, with indication, where appropriate, of the relevant passages</th> <th>Relevant to claim No.</th> </tr> </thead> <tbody> <tr> <td>X</td> <td>EP 1201681 A (Millennium Pharmaceuticals, Inc.), 02 May, 2002 (02.05.02), Full text; Seq. ID. Nos. 6, 34 &amp; US 2003175890 A</td> <td>1-10, 14, 19, 20</td> </tr> <tr> <td>P, X</td> <td>WO 03/89624 A (UAB RESEARCH FOUNDATION), 30 October, 2003 (30.10.03), Full text; Seq. ID. Nos. 27, 28 (Family: none)</td> <td>1-7, 19, 20</td> </tr> </tbody> </table>	Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	X	EP 1201681 A (Millennium Pharmaceuticals, Inc.), 02 May, 2002 (02.05.02), Full text; Seq. ID. Nos. 6, 34 & US 2003175890 A	1-10, 14, 19, 20	P, X	WO 03/89624 A (UAB RESEARCH FOUNDATION), 30 October, 2003 (30.10.03), Full text; Seq. ID. Nos. 27, 28 (Family: none)	1-7, 19, 20	
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.								
X	EP 1201681 A (Millennium Pharmaceuticals, Inc.), 02 May, 2002 (02.05.02), Full text; Seq. ID. Nos. 6, 34 & US 2003175890 A	1-10, 14, 19, 20								
P, X	WO 03/89624 A (UAB RESEARCH FOUNDATION), 30 October, 2003 (30.10.03), Full text; Seq. ID. Nos. 27, 28 (Family: none)	1-7, 19, 20								
<input type="checkbox"/> Further documents are listed in the continuation of Box C.										
<input type="checkbox"/> See patent family annex.										
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family										
Date of the actual completion of the international search 15 October, 2004 (15.10.04)	Date of mailing of the international search report 02 November, 2004 (02.11.04)									
Name and mailing address of the ISA/ Japanese Patent Office	Authorized officer									
Facsimile No.	Telephone No.									

Form PCT/ISA/210 (second sheet) (January 2004)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2004/014207

Continuation of A. CLASSIFICATION OF SUBJECT MATTER  
(International Patent Classification (IPC))

Int.Cl<sup>7</sup> C21P21/02, C12Q1/02, G01N33/15, G01N33/50

(According to International Patent Classification (IPC) or to both national classification and IPC)

Continuation of Box No.II-2 of continuation of first sheet(2)

the meaning within Patent Corporation Treaty Article 5.

Continuation of Box No.III of continuation of first sheet(2)

is a technical relationship between the inventions of the parts relating to a protein as set forth in claim 1 comprising an amino acid sequence represented by SEQ ID NO:2 or 4 and the inventions of the parts relating to the protein as set forth in claim 1 comprising the amino acid sequence represented by SEQ ID NO:6, among the inventions as set forth in claims 1 to 20, involving one or more of the same or corresponding special technical features. Thus, these groups of inventions are not considered as being so linked as to form a single general inventive concept.

Such being the case, the present international application does not comply with the requirement of unity of invention.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2004/014207

**Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.b of the first sheet)**

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:
- a. type of material
- a sequence listing
- table(s) related to the sequence listing
- b. format of material
- in written format
- in computer readable form
- c. time of filing/furnishing
- contained in the international application as filed
- filed together with the international application in computer readable form
- furnished subsequently to this Authority for the purposes of search
2.  In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:



专利名称(译)	在NK细胞中表达的蛋白质		
公开(公告)号	<a href="#">EP1672065A1</a>	公开(公告)日	2006-06-21
申请号	EP2004788274	申请日	2004-09-29
[标]申请(专利权)人(译)	松岛浩二 桥本SHINICHI		
申请(专利权)人(译)	中外SEIYAKU株式会社 松岛，口技 桥本真一		
当前申请(专利权)人(译)	中外SEIYAKU株式会社 松岛，口技 桥本真一		
[标]发明人	MATSUSHIMA KOUJI HASHIMOTO SHINICHI TSUCHIYA MASAYUKI C O CHUGAI SEIYAKU K K		
发明人	MATSUSHIMA, KOUJI HASHIMOTO, SHINICHI TSUCHIYA, MASAYUKI, C/O CHUGAI SEIYAKU K. K. HIRATA, YUICHI, C/O CHUGAI SEIYAKU K. K. YOSHIDA, KENJI, C/O CHUGAI SEIYAKU K. K. OJIMA, KAZUYUKI, C/O CHUGAI SEIYAKU K. K.		
IPC分类号	C12N15/09 A61K45/00 A61P1/16 A61P11/06 A61P31/12 A61P31/14 A61P35/00 A61P37/04 A61P37/06 A61P37/08 C07K14/705 C07K16/28 C12N1/15 C12N1/19 C12N1/21 C12N5/00 C12P21/02 C12Q1/02 G01N33/15 G01N33/50 A61K31/00 C12Q1/68 G01N33/53 G01N33/564 G01N33/566		
CPC分类号	A61K31/00 A61P1/16 A61P11/06 A61P25/18 C07K14/705 G01N33/564 G01N33/566 G01N2500/00		
代理机构(译)	法思博事务所		
优先权	2003338331 2003-09-29 JP		
其他公开文献	EP1672065A4 EP1672065B1		
外部链接	<a href="#">Espacenet</a>		

摘要(译)

分析纯化的免疫细胞的表达频率，并成功鉴定在天然杀伤（NK）细胞中特异性表达的NKIR基因。NKIR基因编码受体。可以通过使用受体来鉴定受体的激动剂和拮抗剂。

Primer: SAS1 (5'-GGGAATCAATGTTGCCATCTTAGTTC-3' / SEQ ID NO: 16) <=>

SAS2 (5'-AAGGATCCATCCCTCTCTGGAGAG-3' / SEQ ID NO: 17)

