

(19)



(11)

EP 1 147 421 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
26.09.2007 Bulletin 2007/39

(51) Int Cl.:
G01N 33/574^(2006.01) G01N 33/68^(2006.01)

(21) Application number: **00901009.1**

(86) International application number:
PCT/CA2000/000060

(22) Date of filing: **24.01.2000**

(87) International publication number:
WO 2000/043789 (27.07.2000 Gazette 2000/30)

(54) METHOD FOR DIAGNOSING ENDOMETRIOSIS

VERFAHREN ZUR DIAGNOSE VON ENDOMETRIOSE

PROCEDE POUR LE DIAGNOSTIC DE L'ENDOMETRIOSE

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
 MC NL PT SE**

(30) Priority: **25.01.1999 US 117031 P**

(43) Date of publication of application:
24.10.2001 Bulletin 2001/43

(73) Proprietor: **Siemens Medical Solutions
 Diagnostics
 Tarrytown, NY 10591 (US)**

(72) Inventors:
 • **GAGNE, Danièle
 Montréal, Québec H1Y 2K7 (CA)**
 • **GOSSELIN, Diane
 Pointe Calumet, Québec J0N 1G2 (CA)**
 • **HUGO, Patrice
 Sainte Dorothée/Laval, Québec H7X 3T3 (CA)**
 • **MIRON, Pierre
 Laval, Québec H7L 4W7 (CA)**

(74) Representative: **Goulard, Sophie
 Cabinet ORES
 36, rue de St Pétersbourg
 75008 Paris Cedex (FR)**

(56) References cited:
EP-A- 0 387 027 WO-A-96/20404
WO-A-98/10291

- **CRUSE, J.M. AND LEWIS, R.E.:** "Illustrated dictionary of immunology", CRC PRESS , 1995 XP002136669 page 53 -page 62
- **OOSTERLYNCK D J ET AL:** "FLOW CYTOMETRY ANALYSIS OF LYMPHOCYTE SUBPOPULATIONS IN PERIONEAL FLUID OF WOMEN WITH ENDOMETRIOSIS" AMERICAN JOURNAL OF REPRODUCTIVE IMMUNOLOGY, DK,MUNKSGAARD INTERNATIONAL PUBLISHERS, COPENHAGEN, vol. 31, no. 1, 1 January 1994 (1994-01-01), pages 25-31, XP000568653 ISSN: 1046-7408 cited in the application
- **OOSTERLYNCK D J ET AL:** "IMMUNOHISTOCHEMICAL CHARACTERIZATION OF LEUCOCYTE SUBPOPULATIONS INENDOMETRIOTIC LESIONS" ARCHIVES OF GYNECOLOGY AND OBSTETRICS,DE,SPRINGER VERLAG, BERLIN, vol. 253, no. 4, 1 January 1993 (1993-01-01), pages 197-206, XP000568666 ISSN: 0932-0067
- **MING-YIH WU ET AL:** 'The suppression of peritoneal cellular immunity in women with endometriosis could be restored after gonadotropin releasing hormone agonist treatment' AMERICAN JOURNAL OF REPRODUCTIVE IMMUNOLOGY vol. 35, no. 6, 1996, pages 510 - 515
- **OTA H. ET AL:** 'Expression of gamma-delta T cells and adhesion molecules in endometriotic tissue in patients with endometriosis and adenomyosis' AMERICAN JOURNAL OF REPRODUCTIVE IMMUNOLOGY vol. 35, no. 5, 1996, pages 477 - 482

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 147 421 B1

Description**BACKGROUND OF THE INVENTION**5 (a) Field of the Invention

[0001] The invention relates to a method for the diagnosis of endometriosis using blood and endometrial leukocyte markers.

10 (b) Description of Prior Art

[0002] Endometriosis is one of the most common gynecological disorders, affecting up to 15% of women within reproductive age. It is closely associated with severe pelvic pain, dysmenorrhea, dyspareunia, infertility and several other symptoms such as intraperitoneal bleeding, back pain, constipation and/or diarrhea. It is a major threat to physical, psychological and social integrity of the patients.

[0003] Endometriosis is characterized by the implantation and growth of endometrial cells (which normally constitute the lining of the uterus) in extra-uterine sites such as the peritoneal cavity. Although the etiology and pathogenesis of endometriosis remain mainly unclear, the theory of retrograde menstruation is the most widely accepted to explain the presence of ectopic endometrial cells in the peritoneal cavity. However, this phenomenon occurs in most women and, thus, several other factors must be invoked to explain the implantation of endometrial cells and the subsequent development of endometriotic lesions. It is generally believed that initiation of endometriosis implies a complex cascade of events requiring several essential features. Retrogradely seeded endometrial cells must remain viable, be capable of adhering to the mesothelium and of proliferating. Local degradation of the extracellular matrix, as well as extensive vascularization, are also believed to play an essential role in promoting the invasion of the peritoneal cavity by endometrial cells. Furthermore, once implanted, ectopic endometrial cells must have the capacity to counteract the cytolytic action of the immune system. Indeed, this is supported by the observation of several immunological abnormalities in patients with endometriosis.

[0004] At present, direct visualization of the endometriotic lesions under surgical procedures (laparoscopy or laparotomy) is the golden standard and the only reliable method available to diagnose endometriosis. However, this method is highly invasive (i.e. surgery under general anesthesia), costly (i.e. direct cost and indirect cost due to convalescence) and requires a well-trained surgeon who has the ability to identify endometriotic lesions with a variety of appearances. The type of lesions, their size and their localization will determine the stage of the disease (stage I minimal, stage II mild, stage III moderate, stage IV severe). However, there is still no clear consensus on how these parameters correlate with the stage of the disease and the prognostic of endometriosis. In addition, early or minimal endometriosis (which can involve microlesions) can be hardly diagnosed by surgical methods, as they are unlikely to be detected by direct visualization. Indeed, several studies have reported microscopic endometriotic lesions that were not detected laparoscopically. Because the diagnosis of endometriosis by surgical procedures is difficult, costly and invasive, in some cases, several physicians and patients tend to avoid it or at least seriously delay it. Hence, the length of time between the onset of symptoms and the diagnosis can be as long as 8 to 12 years. The possibility to diagnose endometriosis at an early stage would certainly improve the efficacy of the treatments, and reduce dramatically the number of years during which patients endure acute or chronic pain.

[0005] Imaging methods such as transvaginal ultrasound and magnetic resonance imaging have been designed for the diagnosis of endometriosis. However, these techniques can only be reliable for the detection of large (> 1 cm diameter) endometriomas lesions detected among a very small proportion of patients with endometriosis. Moreover, the high cost of these techniques has limited their use for the diagnosis of endometriosis.

[0006] Serum proteins such as CA-125 and placental protein-14 have been proposed as diagnostic markers for endometriosis. Elevated levels of CA-125 have been observed in serum, menstrual effluent and peritoneal fluid of patients with endometriosis. However, these markers, when used alone, are of very limited value for a diagnosis test. Indeed, these markers are not suitable for screening or diagnostic purposes because they provide poor sensitivity. Furthermore, levels of CA-125 and placental protein-14 vary according to several factors such as the assay, the stage of the disease and the menstrual cycle. Finally these markers are known to be modulated by conditions other than endometriosis.

[0007] High concentrations of antibodies to endometrial antigens were found in the serum of patients with endometriosis, and thus were proposed as markers for a diagnostic test (International patent application publications WO 94/28021 and WO 92/18535). However, the levels of specificity and sensibility with these tests remain very low. In most cases, the antigens recognized by these antibodies are still poorly characterized or yet totally unknown.

[0008] In U.S. patent No. 5,478,725, low levels of $\alpha\beta3$ integrin expression in endometrial samples during the secretory phase of the menstrual cycle is described as a predictor of endometriosis in infertile but not in fertile patients with endometriosis. This observation was associated with milder form of endometriosis (stages I and II) only and, thus, is not

useful to detect advanced stages of the disease. Moreover, this method yielded a specificity of 91% but a very low sensitivity (38%).

[0009] Taking into account that a number immunological abnormalities have been reported in patients with endometriosis, it is conceivable that the proportion of leukocyte populations and/or their activation status may be modulated during the course of the disease and, thus, may provide some diagnostic value. Previous flow cytometric studies have shown that some T lymphocyte subpopulations (CD8+, CD45+/HLADR+, CD45+/CD3+/HLADR+ or CD3+/CD25+) can be slightly modulated in the peritoneal fluid of subjects with endometriosis relative to normal controls (Oosterlynck D.J., et al., Am J reprod. Immunol., 31: 25-31, 1994; Becker J.L., et al., Am J Reprod. Immunol., 34: 179-187, 1995; Wu M.Y., et al., Am. j. Reprod. Immunol. 35: 510-516, 1996). However, these observations have limited value for the diagnosis of endometriosis because peritoneal fluid collection is an invasive, non-conventional procedure. Proportions of leukocyte populations have also been studied in peripheral blood and endometrium of patients with endometriosis. Wu et al., (supra) have reported a modest but significant decrease in the proportion of CD3+ T lymphocytes expressing either CD69 or CD25 activation marker in the blood of patients with advanced endometriosis but not in patients with mild stage of endometriosis or normal controls. This difference was observed in advanced cases of endometriosis only and it was too modest to be used as a diagnostic marker. In contrast, Oosterlynck et al., (Oosterlynck D.J., et al., Am J reprod. Immunol., 31: 25-31, 1994) and Ho et al. (Ho H.N., et al., Hum Reprod., 97: 2528-2533, 1997) reported no significant difference in term of T lymphocyte subpopulations when comparing endometriosis subjects with normal controls. These inconsistent results may be explained by the very low number of samples tested in these studies.

[0010] Several studies have investigated whether leukocytes are also modulated in eutopic endometrium from patients with endometriosis. Results arising from these studies are contradictory, probably due to the fact that in most cases the methods used were only semi-quantitative and the number of samples tested were very low. For instance, by means of immunohistochemistry, Ota et al. (Ota H., et al., Am J Reprod. Immunol., 35: 477-482, 1996) have reported that the number of CD3+, CD4+, or CD8+ T lymphocytes, cells bearing adhesion molecules (i.e. ICAM-1, LFA-1, CD2) or CD68+ cells were upregulated in the endometrium of patients with endometriosis compared with infertile controls. In contrast, several other studies using similar techniques have reported no difference in the proportion of T lymphocyte subsets (Klentzeris L.D., et al., Eur. J Obstet gynecol Reprod Biol., 63:41-47, 1995; Jones R.K., et al., Fertil Steril, 66:81-89, 1996). In addition, a decrease in CD3 positive T cells has been shown by flow cytometry analysis but no difference in the proportion of CD4+, CD8+ stromal leukocytes in the endometrium of patients with endometriosis compared with fertile controls. When these observations are tentatively used in a diagnostic test, they give only low levels of sensibility and specificity because of a significant overlap between the groups.

[0011] Therefore, the diagnostic methods presented in the literature so far do not solve the problems encountered with the diagnosis of endometriosis by surgical procedures. It thus remains imperative to be provided with a less invasive, cheaper and reliable method that could allow detection of females suffering from endometriosis as early as possible.

SUMMARY OF THE INVENTION

[0012] One aim of the present invention is to provide a less invasive, cheaper and reliable method that could allow detection of females suffering from endometriosis as early as possible.

[0013] In accordance with the present invention there is provided a method for determining likelihood of endometriosis in a sample of eutopic uterine endometrial tissues or a sample of blood of a female subject, characterized in that it comprises the steps of:

- a) measuring in said sample a quantitative level of at least two different selected surface antigens from blood leukocytes and/or eutopic endometrial leukocytes, as defined in claim 1 ;
- b) establishing a cutoff value for each leukocyte marker in the combination;
- c) comparing the proportion obtained in step b) for each leukocyte marker to a predetermined cutoff value, wherein a positive result gives a score of 1 whereas a negative result gives a score of 0;
- d) obtaining a diagnostic value by adding the scores of all the markers of the combination and converting it in percentage; and
- e) comparing the final diagnostic value to an established threshold value wherein a positive diagnosis of endometriosis is given when the final diagnostic value exceeds the threshold value established for the combination of leukocyte markers whereas a negative diagnosis of endometriosis is given when the final diagnostic value is lower than the threshold value established for the combination of leukocyte markers.

[0014] Further, in accordance with the present invention, there is also provided a method for determining endometriosis in a sample of eutopic uterine endometrial tissues or a sample of blood of a female subject, comprising:

- a) measuring in said sample a quantitative level of at least two different selected surface antigens from blood

leukocytes and/or eutopic endometrial leukocytes, as defined in claim 3;
 b) establishing a predictive model for endometriosis by including each marker of said combination in a logistic regression equation, as defined in claim 3;
 c) comparing the probability of having endometriosis to a threshold value wherein a positive diagnosis of endometriosis is given when the probability of having endometriosis value exceeds the threshold value established for said combination of markers whereas a negative diagnosis of endometriosis is given when the probability of having endometriosis value is lower than the threshold value established for said combination of markers.

[0015] For the purpose of the present invention, the following symbol "/" is intended to mean a ratio between an expression in front of the symbol and another expression after the symbol.

BRIEF DESCRIPTION OF THE DRAWING

[0016]

Fig. 1 illustrates a predictive algorithm for the diagnosis of endometriosis.

DETAILED DESCRIPTION OF THE INVENTION

[0017] In accordance with the present invention, there is provided reliable diagnostic test for endometriosis that is less invasive and less costly than the actual surgical procedure accepted as the golden standard. An extensive study was undertaken by means of flow cytometric analysis, in which the proportion of several blood and endometrial leukocyte subsets was compared in patients with endometriosis and normal controls.

[0018] The present invention identifies a series of leukocyte subsets that can be used as markers in a diagnostic test for endometriosis. These leukocyte subsets are defined according to the expression of cell surface antigens. Several cell surface antigens may define the same population of cells, and thus they are included in the present invention.

[0019] Any other antibodies or molecules recognizing the same antigen or a different epitope, isoform, subunit, chain, glycosylation or phosphorylation form or an allelic variant of the same antigen, a member of the same complex, or an antigen with the same cell distribution is also included in the present invention.

[0020] Further in accordance with the present invention, there is provided examples showing how at least two different surface antigens from blood and/or endometrial leukocytes can be used in combinations in a diagnostic test for endometriosis (Tables 1 and 2).

TABLE 1
Levels of sensibility and specificity provided by several examples of endometrial and/or blood marker combinations used as a diagnostic method for endometriosis

<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
Endometrial leukocyte markers			
1. CD4+ (<17%) ² CD8+CD69- (<21%) CD13+CD45RO- (<17.5%)	>67%	90%	67%
2. CD4+ (<15.5%) CD8+CD69- (<21%) CD56+CD122- (>19%) CD3+CD45RA- (<35%) CD13+CD45RO- (<17.5%)	>60%	89%	65%
3. CD4+ (<17%)	>67%	88%	65%

EP 1 147 421 B1

(continued)

Levels of sensibility and specificity provided by several examples of endometrial and/or blood marker combinations used as a diagnostic method for endometriosis

	<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
5	CD8+CD69- (<21%) CD13-CD122+ (>28%) CD13+CD45RO- (<17.5%)			
10	4. CD4+ (<17%) CD8+CD69- (<21%) CD14+CD13-CD16b- (>14.5%)	>67%	89%	63%
15	5. CD3+CD16- (<40%) CD13+CD45RO- (<13.5%) CD3+ (<40%) CD8+ (<20%) CD3+CD69+ (<15%)	>55%	84%	62%
20	6. CD3+ (<40%) CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD3-CD20- (>56%)	>65%	84%	63%
25	7. CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD3+CD5+ (<37%) CD3+CD122- (<42.5%) CD3-CD20- (>56%) CD3+CD45RO- (<30%)	>65%	81 %	65%
30	8. CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD3+CD5+ (<37%) CD3+CD122- (<42.5%)	>60%	82%	64%
35	9. CD3+CD20-CD5- (>7.7%) CD4+CD13- (<20.5%) CD56-CD122- (<47%)	>60%	81%	66%
40				
45				
50				
55				

EP 1 147 421 B1

(continued)

Levels of sensibility and specificity provided by several examples of endometrial and/or blood marker combinations used as a diagnostic method for endometriosis

	<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
5	10. CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%)	>60%	80%	65%
10	CD4+CD45RA- (<16%) CD3+CD45RO- (<30%)			
15	11. CD3+ (<40%) CD8+CD69- (<18%) CD3-CD4-CD45RO+ (>56%) Ratio CD13+/CD3+ (>0.675%) CD13+CD45RO- (<21 %)	>35%	79%	67%
20				
25	12. CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD3-CD5- (>54%) CD20-CD5+ (<44%)	>70%	81%	61%
30				
35	13. CD8+ (<20%) ² CD5+ (<37%) CD3-CD20- (>58%) CD3-HLADR- (>54.5%)	>51%	81%	60%
40				
45	14. CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD5+ (<40%)	>60%	81%	60%
50				
55	15. CD4+ (<17%) CD13-CD122+ (>28%) CD8+CD69- (<19.5%) CD3+CD45RA- (<37%)	>50%	76%	71 %
	16. CD4+ (<15.5%) CD8+CD69- (<21%) CD13-CD122+ (>28%) CD3+CD45RA- (<35%) CD13+CD45RO- (<17.5%)	>35%	71%	78%

EP 1 147 421 B1

(continued)

Levels of sensibility and specificity provided by several examples of endometrial and/or blood marker combinations used as a diagnostic method for endometriosis

	<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
5	17.			
	CD4+ (<15.5%)	>40%	70%	78%
10	CD8+CD69- (<21 %)			
	CD56-CD122- (<47%)			
	CD3+CD45RA- (<35%)			
	CD13+CD45RO- (<17.5%)			
15	18.			
	CD3+(<40%)	>35%	72%	76%
	CD4+ (<17%)			
	CD3+CD8+ (<16%)			
20	CD13+CD45RO- (<21%)			
	CD3+CD5+ (<37%)			
25	19.			
	CD3+(<40%)	>40%	74%	74%
	CD4+ (<17%)			
	CD3+CD8+ (<16%)			
	CD13+CD45RO- (<21%)			
30	20.			
	CD3+(<40%)	>40%	75%	73%
	CD3+CD8+ (<16%)			
	CD13+CD45RO- (<21%)			
35	21.			
	CD3+ (<40%)	>25%	71%	69%
	CD5+ (<40%)			
	CD3+CD5+ (<37%)			
40	CD69+ (<33%)			
	CD4-CD69+ (<35%)			
45	22.			
	CD3+ (<40%)	>30%	68%	83%
	CD3+CD8+ (<13.5%)			
	CD13+CD45RO- (<17.5%)			
50	23.			
	CD3+ (<40%)	>30%	61%	86%
	CD3+CD8+ (<16%)			
	CD13+CD45RO- (<17.5%)			
55	24.			
	CD3+ (<40%)	>22%	62%	80%

EP 1 147 421 B1

(continued)

Levels of sensibility and specificity provided by several examples of endometrial and/or blood marker combinations used as a diagnostic method for endometriosis

	<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
5	CD3+CD8+ (<16%)			
	CD13+CD45RO- (<17.5%)			
10	CD3-CD20- (>58%)			
	CD56-CD16- (<46%)			
	25.			
	CD3+CD16- (<47.5%)	>45%	66%	79%
15	CD3-CD4-CD45RO+ (>31.5%)			
	CD3+ (<40%)			
	CD8+ (<20%)			
	CD3+CD69+ (<15%)			
20	CD13+CD45RO- (<17.5%)			
	26.			
	CD3+ (<40%)	>40%	66%	75%
25	CD3-CD45RO+ (>15%)			
	Ratio CD13+/CD3+ (>0.675%)			
	CD3+CD8+ (<16%)			
30	CD8+CD69- (<21 %)			
	27.			
	CD3+CD20-CD5- (>7.7%)	>45%	61%	80%
35	CD4+CD13- (<20.5%)			
	CD56-CD122- (<47%)			
	CD4+CD45RO- (<16%)			
	28.			
40	CD3+ (<40%)	>20%	61%	86%
	CD3+CD8+ (<16%)			
	CD13+CD45RO- (<17.5%)			
45	CD3-CD20- (>58%)			
	29.			
	CD13+CD45RO- (<17.5%)	>70%	90%	54%
50	CD4+CD45RA- (<16%)			
	CD3+CD122- (42.5%)			
	CD8+CD69- (<21%)			
	30.			
55	CD3+CD8+ (<16%)	>70%	84%	60%

EP 1 147 421 B1

(continued)

Levels of sensibility and specificity provided by several examples of endometrial and/or blood marker combinations used as a diagnostic method for endometriosis

	<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
5	CD13+CD45RO- (<17.5%)			
	CD4+CD45RA- (<16%)			
10	CD3+CD45RO- (<30%)			
	CD3-CD5- (>54%)			
	Blood leukocyte markers			
15	31.			
	CD3-CD5+ (>14.5%)	>55%	66%	60%
	CD3-CD45RA- (>14.5%)			
	CD3-CD44+ (>13%)			
20	CD13+ (>17.5%)			
	CD3-CD57-CD44- (<41.3%)			
	32.			
25	CD3-CD45RA- (>17%)	>22%	61%	64%
	CD20-CD44+ (>17%)			
	CD20-HLADR+ (>20%)			
30	CD3-CD4-CD44+ (>40.5%)			
	CD36-HLADR+ (<5.6%)			
	33.			
35	CD3-CD45RA- (>14.5%)	>40%	62%	64%
	CD3-CD45RO+ (>19%)			
	CD20-HLADR+ (>14.5%)			
	Blood (in Italics) and endometrial leukocyte markers			
40	34.			
	CD57+ (>10%)	>50%	76%	72%
	<i>CD14+</i> (>10%)			
45	CD3-CD69+ (>17.5%)			
	CD3+ (<40%)			
	CD4+ (<15.5%)			
50	CD3+CD8+HLADR- (<35%)			
	35.			
	<i>CD3-CD69+</i> (>17.5%)	>33%	70%	79%
	CD3+ (<40%)			
55	CD4+ (<15.5%)			
	CD3+CD8+HLADR- (<35%)			

EP 1 147 421 B1

(continued)

Levels of sensibility and specificity provided by several examples of endometrial and/or blood marker combinations used as a diagnostic method for endometriosis

5	<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
	36.			
	CD4-CD36+ (>14.4%)	>43%	70%	74%
10	CD3-CD69+ (>17.5%)- CD8+ (<20%) CD13+ (>29%) CD3+ (<40%) CD16+ (>27%)			
15	CD69+(<33%) CD5+ (<40%)			
	37.			
20	CD3-CD45RA- (>14.5%) CD3-CD45RO+ (>19%) CD20-HLADR+ (>14.5%) CD14+CD44+ (>15%) CD8+ (<20%) CD5+ (<37%)	>50%	73%	71%
25	CD3-CD20- (>58%) CD3-HLADR- (>54.5%)			

¹ Value above which a diagnosis of endometriosis is given.

² Cutoff point established for each individual marker.

TABLE 2

Examples of logistic regression models provided by endometrial or blood leukocyte markers for the identification of patients with endometriosis

35	<u>Marker combination</u>	<u>B value</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>	<u>Number of sample tested</u>
	Endometrial leukocyte markers					
40	Combination no. 1					
	1. CD3+ (<40%) ²	-7.9747	>.55	83%	79%	41
45	2. CD3-CD5- (>60%)	7.2921				
	3. CD13+CD45RO- (<17.5%)	-0.1410				
50	4. CD3-CD20- (>58.%)	-1.6259				
	Interaction of 1 to 4					
55	Constant = 2.0516					
	Combination no. 2					

EP 1 147 421 B1

(continued)

Examples of logistic regression models provided by endometrial or blood leukocyte markers for the identification of patients with endometriosis

5	Marker combination	B value	Threshold value ¹	Specificity	Sensibility	Number of sample tested
	1. CD3+ (<40%)	-6.7753	>.55	74%	73%	67
	2. CD3-CD5- (>60%)	5.8240				
10	3. CD13+CD45RO- (<17.5%)	-1.9298				
	Interaction 1 to 4	-0.0262				
15	Constant =	2.8385				
	2.7910					
	Combination no. 3					
20	1. CD3+CD8+ (<16%)	-0.1308	>.50	84%	72%	51
	2. CD13+CD45RO- (<17.5%)	-2.6688				
25	3. CD3+CD5+ (<37%)	-1.1778				
	Constant =					
	3.1417					
30	Combination no. 4					
	1. CD3+ (<40%)	-1.6965	>.50	78%	75%	81
35	2. Length of menstruation (>7days)	-1.8160				
	3. CD13+CD20- (<21%)	-1.9656				
40	4. Pelvic pain ³	10.3064				
	Constant =					
	3.1984					
45	Blood leukocyte markers					
	Combination no. 1					
50	1. CD14+CD44+ (>15%)	0.9298	>0.55	80%	70%	140
	2. CD57+ (>10%)	0.7423				
	3. CD3-CD45RA- (>12%)	-0.8147				
55	4. CD14+ (>10%)	0.8629				
	Combination no. 2					

EP 1 147 421 B1

(continued)

Examples of logistic regression models provided by endometrial or blood leukocyte markers for the identification of patients with endometriosis

Marker combination	B value	Threshold value ¹	Specificity	Sensibility	Number of sample tested
1. CD14+ (>10%)	10.5891	>.50	65%	71%	125
2. CD57+ (>10%)	0.7326				
3. CD3+CD69+ (>17.5%)	0.6899				
4. CD3+HLADR+ (<4%)	1.2004				
5. CD3-CD45RA- (>12%)	-0.1137				
Constant = -1.2062					
Combination no.3					
1. CD14+ (>10%)	1.1994	>.55	76%	75%	142
2. CD57+ (>10%)	0.8080				
3. CD3+HLADR- (<4%)	1.3593				
4. CD3-CD45RA- (>12%)	-0.63				
5. Pelvic pain	2.1506				
6. Length of menstruation (>7d)	.7489				
Constant = -1.771					
Combination no.4					
1. CD14+ (>10%)	.9727	>.50	71%	78%	141
2. CD57+ (>10%)	.4489				
3. CD3+CD69+ (>17.5%)	.8129				
4. CD3+HLADR- (<4%)	1.3368				
5. CD3-CD45RA- (>12%)	-0.8805				
6. Pelvic pain	2.1574				
7. Age (>40)	1.5164				
Constant = -1.7686					

¹ Value above which a diagnosis of endometriosis is given.

² Cutoff point established for each individual marker.

³ presence of pain at any time other than menstruation and intercourse

[0021] The predictive models for endometriosis were established according to the following equation:

$$P(r) = \frac{e^{c + B1*(marker1) + B2*(marker2) + \dots + Bn*(marker n)}}{1 + e^{c + B1*(marker1) + B2*(marker2) + \dots + Bn*(marker n)}}$$

Where:

P(r) = probability of having endometriosis;
 c = constant established for a particular combination;
 B = coefficient of regression; and
 n = total number of markers in the combination.

5
 [0022] In the present invention, a series of endometrial and peripheral blood leukocyte subpopulations for which proportions were modulated in patients with endometriosis (stage I-IV; I-II or III-IV) compared with those of normal controls, have been identified. The novelty of the present invention is to use these leukocyte subpopulations in combination, as markers for the diagnosis of endometriosis. Moreover, risk factors for endometriosis identified amongst personal information and menstrual characteristics were shown to be of significant value when use in combination with blood or endometrial leukocyte subsets in a predictive test for endometriosis.

10 [0023] Two methods were used for the combination of markers.

15 **Method 1**

[0024] A cutoff point is established for the proportion of each combination of leukocyte markers in order to obtain the best discrimination between patients with endometriosis and controls. The proportion obtained for each marker is compared to the cutoff point. A positive test result gives a score of 1, whereas a negative test result gives a score of 0. The diagnostic value is obtained by adding the scores of all the markers of a particular combination and converting it in percentage. The final diagnostic value is then compared to a threshold value that was established to provide the best levels of sensibility and specificity. A positive diagnosis of endometriosis is given when the final diagnostic value exceeds the threshold value established for a particular combination of markers. On the opposite, a negative diagnosis of endometriosis is given when the final diagnostic value is lower than the threshold value (see Fig. 1).

20 [0025] Therefore, according to a first embodiment of the invention, there is provided a method for determining endometriosis in a sample of eutopic uterine endometrial tissues or a sample of blood of a female subject, comprising:
 25 a) measuring in said sample a quantitative level of at least two different surface antigens from blood leukocytes and/or eutopic endometrial leukocytes, said surface antigens from blood leukocytes and/or eutopic endometrial leukocytes being selected from the leukocyte marker combinations defined in the following Table:

30

Leukocyte Marker Combinations	
Endometrial leukocyte markers	
35	CD4+, CD8+CD69-, and CD13+CD45RO-
	CD4+, CD8+CD69-, CD56+CD122-, CD3+CD45RA-, and CD13+CD45RO-
	CD4+, CD8+CD69-, CD13-CD122+, and CD13+CD45RO-
	CD4+ CD8+CD69-, and D14+CD13-CD16b-
	CD3+CD16-, CD13+CD45RO-, CD3+, CD8+, and CD3+CD69+
	CD3+, CD3+CD8+, CD13+CD45RO-, and CD3-CD20-
40	CD3+CD8+, CD13+CD45RO-, CD3+CD5+, CD3+CD122-, CD3-CD20-, and CD3+CD45RO-
	CD3+CD8+, CD13+CD45RO-, CD3+CD5+, and CD3+CD122-
	CD3+CD20-CD5-, CD4+CD13-, and CD56-CD122-
	CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, and CD3+CD45RO-
45	CD3+, CD8+CD69-, CD3-CD4-CD45RO+, ratio CD13+/CD3+, and CD13+CD45RO-
	CD3+CD8+, CD13+CD45RO-, CD3-CD5-, and CD20-CD5+
	CD8+, CD5+, CD3-CD20-, and CD3-HLADR-
	CD3+CD8+, CD13+CD45RO-, and CD5+
	CD4+, CD13-CD122+, CD8+CD69-, and CD3+CD45RA-
50	CD4+, CD8+CD69-, CD13-CD122+, CD3+CD45RA-, and CD13+CD45RO-
	CD4+, CD8+CD69-, CD56-CD122-, CD3+CD45RA-, CD13+CD45RO-
	CD3+, CD4+, CD3+CD8+, CD13+CD45RO-, and CD3+CD5+
	CD3+, CD4+, CD3+CD8+, and CD13+CD45RO-
	CD3+, CD3+CD8+, and CD13+CD45RO-
55	CD3+, CD5+, CD3+CD5+, CD69+, and CD4-CD69+
	CD3+, CD3+CD8+, and CD13+CD45RO-
	CD3+, CD3+CD8+, and CD13+CD45RO-

(continued)

<p>5 10 15</p>	<p>Endometrial leukocyte markers CD3+, CD3+CD8+, CD13+CD45RO-, CD3-CD20-, and CD56-CD16- CD3+CD16-, CD3-CD4-CD45RO+, CD3+, CD8+, CD3+CD69+, and CD13+CD45RO- CD3+, CD3-CD45RO+, ratio CD13+/CD3+, CD3+CD8+, and CD8+CD69- CD3+CD20-CD5-, CD4+CD13-, CD56-CD122-, and CD4+CD45RO- CD3+, CD3+CD8+, CD13+CD45RO-, and CD3-CD20- CD13+CD45RO-, CD4+CD45RA-, CD3+CD122-, and CD8+CD69- CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, CD3+CD45RO-, and CD3-CD5- CD3+, CD3-CD5-, CD13+CD45RO-, and CD3-CD20- CD3+, CD3-CD5-, and CD13+CD45RO- CD3+CD8+, CD13+CD45RO-, and CD3+CD5+ CD3+ and CD13+CD20.</p>
<p>20 25</p>	<p>Blood Leukocyte markers CD3-CD5+, CD3-CD45RA-, CD3-CD44+, CD13+, and CD3-CD57-CD44- CD3-CD45RA-, CD20-CD44+, CD20-HLADR+, CD3-CD4-CD44+, and CD36-HLADR+ CD3-CD45RA-, CD3-CD45RO+, and CD20-HLADR+ CD14+CD44+, CD57+, CD3-CD45RA, and CD14+ CD14+, CD57+, CD3+CD69+, CD3+HLADR+, and CD3-CD45RA- CD14+, CD57+, CD3+HLADR- and CD3-CD45RA- CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-</p>
<p>30</p>	<p>Blood (in italics) and endometrial leukocyte markers <i>CD57+</i>, <i>CD14+</i>, <i>CD3-CD69+</i>, CD3+, CD4+, and CD3+CD8+HLADR- <i>CD3-CD69+</i>, CD3+, CD4+, and CD3+CD8+HLADR- <i>CD4-CD36+</i>, <i>CD3-CD69+</i>, CD8+, CD13+, CD3+, CD16+, CD69+, and CD5+ <i>CD3-CD45RA-</i>, <i>CD3-CD45RO+</i>, CD20-HLADR+, <i>CD14+CD44+</i>, CD8+, CD5+, CD3-CD20-, and CD3-HLADR-</p>

the measurement of the quantitative level of the combination of endometrial leukocyte markers CD3+, CD13+CD20 being associated with length of menstruation (> 7 days) and pelvic pain at any time other than menstruation and intercourse ; the measurement of the quantitative level of the combination of blood leukocyte markers CD14+, CD57+, CD3+HLADR-, CD3-CD45RA-being associated with pelvic pain at any time other than menstruation and intercourse, and length of menstruation (>7 days); and the measurement of the quantitative level of the combination of blood leukocyte markers CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA- being associated with pelvic pain at any time other than menstruation and intercourse, and age > 40,

b) establishing a cutoff value for the proportion of each leukocyte marker in the combination,

c) comparing the proportion obtained in step (b) for each leukocyte marker to a predetermined cutoff value wherein a positive result gives a score of 1 whereas a negative result gives a score of 0,

d) obtaining a diagnostic value by adding the scores of all the markers of the combination and converting it in percentage,

e) comparing the final diagnostic value to an established threshold value wherein a positive diagnosis of endometriosis is given when the final diagnostic value exceeds the threshold value established for the combination of leukocyte markers whereas a negative diagnosis of endometriosis is given when the final diagnostic value is lower than the threshold value established for the combination of leukocyte markers.

Method 2

[0026] A predictive model for endometriosis is established by including each marker of a particular combination in the following logistic regression equation:

$$P(r) = \frac{e^{c + B1*(marker1) + B2 (marker2) + \dots Bn (marker n)}}{1 + e^{c + B1*(marker1) + B2 (marker2) + \dots Bn (marker n)}}$$

(continued)

Blood Leukocyte markers

CD3-CD5+, CD3-CD45RA-, CD3-CD44+, CD13+, and CD3-CD57-CD44-
 CD3-CD45RA-, CD20-CD44+, CD20-HLADR+, CD3-CD4-CD44+, and CD36-HLADR+
 CD3-CD45RA-, CD3-CD45RO+, and CD20-HLADR+
 CD14+CD44+, CD57+, CD3-CD45RA, and CD14+
 CD14+, CD57+, CD3+CD69+, CD3+HLADR+, and CD3-CD45RA-
 CD14+, CD57+, CD3+HLADR-, CD3-CD45RA-
 CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-

Blood (in italics) and endometrial leukocyte markers

CD57+, *CD14+*, *CD3-CD69+*, CD3+, CD4+, and CD3+CD8+HLADR-
CD3-CD69+, CD3+, CD4+, and CD3+CD8+HLADR-
CD4-CD36+, *CD3-CD69+*, CD8+, CD13+, CD3+, CD16+, CD69+, and CD5+
CD3-CD45RA-, *CD3-CD45RO+*, *CD20-HLADR+*, *CD14+CD44+*, CD8+, CD5+, CD3-CD20-, and CD3-HLADR-

the measurement of the quantitative level of the combination of endometrial leukocyte markers CD3+, CD13+CD20 being associated with length of menstruation (> 7 days) and pelvic pain at any time other than menstruation and intercourse ; the measurement of the quantitative level of the combination of blood leukocyte markers CD14+, CD57+, CD3+HLADR-, CD3-CD45RA-being associated with pelvic pain at any time other than menstruation and intercourse, and length of menstruation (>7 days); and the measurement of the quantitative level of the combination of blood leukocyte markers CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA- being associated with pelvic pain at any time other than menstruation and intercourse, and age > 40,

b) establishing a predictive model for endometriosis by including each marker of said combination in the following logistic regression equation :

$$P(r) = \frac{e^{c+B1*(marker1)+B2*(marker2)+...Bn*(marker n)}}{1 + e^{c+B1*(marker1)+B2*(marker2)+...Bn*(marker n)}}$$

Where:

- P (r) = probability of having endometriosis;
- c = constant established for a particular combination;
- B = coefficient of regression;
- n = total number of markers in the combination,

c) comparing the probability of having endometriosis (P (r)) to a threshold value wherein a positive diagnosis of endometriosis is given when the P (r) value exceeds the threshold value established for said combination of markers whereas a negative diagnosis of endometriosis is given when the P (r) value is lower than the threshold value established for said combination of markers.

[0029] In the present invention, there is reported a series of 101 endometrial CD45+ leukocyte populations and 96 blood mononuclear CD45+ leukocyte populations which were shown by flow cytometric analysis to be modulated in patients with endometriosis (stage I, II, III or IV) compared with normal controls and, thus are good candidate markers for the diagnosis of endometriosis (Tables 3, 4, 5, and 6). The innovative feature of the present invention is to use these markers in combination to increase their level of sensibility and specificity in the diagnostic test.

TABLE 3
Endometrial leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P1	area under ROC curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo stage I-IV	Control	Endo								
CD3+	47.7±12.3	38.7±12.6	58	88	3.37×10 ⁻⁵	0.703	3.3×10 ⁻⁵	<40	84	55	6.5	(2.9-14.9)
CD4+	18.3±5.6	15.7±5.9	55	88	0.008	0.632	0.008	<17	63	63	3.0	(1.5-6.2)
CD5+	45.1±11.6	36.3±12.3	46	74	1.6×10 ⁻⁴	0.702	0.0002	<37	80	53	4.2	(1.8-9.7)
CD8+	24.3±8.5	18.5±8.6	54	87	1.4×10 ⁻⁴	0.688	0.00019	<40	73	58	3.9	(1.7-8.8)
CD3+CD4+	17.2±5.5	14.4±5.7	55	88	0.004	0.641	0.004	<20	74	62	4.8	(2.3-10.2)
CD3+CD4-	29.9±9.6	23.6±9.1	55	88	1.2×10 ⁻⁴	0.687	0.00017	<15	67	57	2.7	(1.3-5.5)
CD3-CD4-	51.1±14.2	60.8±12.6	55	88	4.1×10 ⁻⁵	0.698	6.9×10 ⁻⁵	<24	80	52	4.4	(2.0-9.6)
CD3+CD8+	18.9±7.5	13.7±7.7	54	84	1.1×10 ⁻⁴	0.714	2.3×10 ⁻⁵	<61	83	50	5.1	(2.2-11.7)
CD3+CD8-	26.1±7.8	23.1±7.1	54	84	0.022	0.609		<16	70	70	5.6	(2.6-11.8)
CD3-CD8-	49.6±12.0	58.2±13.0	54	84	1.3×10 ⁻⁴	0.688	1.9×10 ⁻⁴	<13.5	81	54	5.3	(2.3-11.9)
CD3+CD69+	20.4±9.6	15.5±8.0	44	76	0.003	0.642	0.010	<23.5	68	51	2.2	(1.1-4.7)
CD3+CD122-	41.4±10.0	34.4±12.4	29	53	0.011	0.669	0.012	>53.5	70	63	4.0	(1.9-8.5)
CD3+HLADR-	38.1±10.3	30.6±12.3	51	80	2.8×10 ⁻⁴	0.681	0.0005	<42.5	64	76	2.9	(1.1-7.5)
CD3-HLADR-	46.4±13.0	55.6±13.5	51	80	1.9×10 ⁻⁴	0.693	0.0002	<35	72	63	4.0	(1.8-8.5)
CD3+CD45RA+	7.4±4.7	5.7±3.0	56	85	0.018	0.608	0.030	>54.5	80	51	4.1	(1.8-9.3)
CD3+CD45RA-	40.3±11.2	32.7±12.0	56	85	2.5×10 ⁻⁴	0.684	0.0002	<4.9	77	40	2.2	(1.0-4.7)
			2	0				<37	69	66	4.7	(2.2-9.7)
								<35	73	60	4.1	(1.9-8.5)

(continued)
 Endometrial leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P1	area under ROC curve ²	P3	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo stage I-IV	Control	Endo								
CD3-CD45RA-	31.4±12.7	39.6±13.4	56	85	4.2x10 ⁻⁴	0.667	8.2 x10 ⁻⁴	>32	51	69	2.3	(1.1-4.6)
CD3+CD45RO-	31.0±11.1	25.0±9.8	50	73	0.002	0.661	0.002	<28	65	62	3.1	(1.4-6.6)
CD3+CD16-	45.5±12.1	37.2±13.1	57	83	2.1x10 ⁻⁴	0.680	0.0003	<30	55	70	2.7	(1.3-5.7)
CD3+CD56-	46.5±12.2	38.5±12.8	56	83	3.2x10 ⁻⁴	0.674	0.00053	<40	75	58	4.2	(2.0-8.9)
CD3+CD5+	41.9±11.4	33.3±12.3	45	74	2.1x10 ⁻⁴	0.695	0.00036	<47.5	40	80	2.6	(1.2-5.6)
CD3-CD5-	50.6±12.0	59.0±12.9	45	74	0.001	0.690	0.00052	>60	82	51	4.6	(1.9-11.2)
CD4+CD69-	16.4±4.8	13.8±5.1	37	72	0.012	0.648	0.012	>54	66	66	4.3	(1.9-9.6)
CD4+CD45RA-	16.7±5.3	14.2±5.7	54	85	0.010	0.632	0.009	<14	78	53	4.1	(1.6-10.1)
CD8+CD69-	24.0±7.9	18.9±8.3	30	59	0.007	0.687	0.004	<16	62	66	3.3	(1.6-6.7)
CD8+HLADR-	23.3±7.7	18.1±8.5	49	79	0.001	0.673	0.001	<18	83	53	5.9	(1.9-17.6)
CD8-HLADR-	61.6±9.5	68.1±9.5	49	79	2.2x10 ⁻⁴	0.675	0.0009	<19.5	76	59	5.1	(1.9-13.9)
CD13-CD122+	27.0±9.8	33.6±18.7	32	58	0.031	0.605		<21	65	68	4.2	(1.6-10.7)
CD13-CD122-	47.1±14.4	40.4±15.0	32	58	0.043	0.635	0.035	<18	77	54	4.1	(1.8-9.2)
CD20-CD5+	44.6±12.0	36.4±12.6	41	66	0.001	0.681	0.002	>61.5	52	77	3.8	(1.7-8.3)
								>28	64	59	2.5	(1.0-6.2)
								<46	58	64	2.6	(1.1-6.2)
								<41	60	62	2.6	(1.2-5.7)

(continued)
 Endometrial leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P1	area under ROC curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo stage I-IV	Control	Endo								
CD20-CD5-	52.0 ± 12.8	60.9 ± 12.9	41	66	0.001	0.692	0.0009	<44	57	71	3.5	(1.5-7.9)
CDS6-CD16+	22.0 ± 2	27.0 ± 16.6	56	84	0.044	0.571		>60	77	50	3.6	(1.5-8.6)
CD56-CD16-	51.6 ± 2	42.8 ± 5	56	84	1.2 × 10 ⁻⁴	0.687	1.8 × 10 ⁻⁴	<46	71	57	3.3	(1.6-6.7)
ratio CD3/CD45RO	1.5 ± 1.0	1.2 ± 0.7	51	80	0.020	0.626	0.015					
CD14+CD13-	1.4 ± 0.9	2.3 ± 1.8	21	36	0.041	0.639						
CD3+CD20-	44.4 ± 11.2	36.9 ± 13.5	24	45	0.024	0.667	0.023	<40	78	58	4.1	(1.4-12.3)
CD3-CD20-	52.3 ± 5	61.0 ± 9	24	45	0.016	0.669	0.022	>58	83	53	6.2	(1.8-21.2)
CD3-CD4-CD45RA+	40.1 ± 7	34.9 ± 15	51	79	0.046	0.618	0.023	>56	70	60	4.5	(1.4-13.5)
CD3-CD4-CD45RA-	57.9 ± 14.1	63.3 ± 15	51	79	0.042	0.620	0.021					
CD3+CD8+CD69-	40.4 ± 10.2	35.1 ± 11.7	29	56	0.039	0.635	0.042	<34.5	75	52	3.4	(1.2-9.2)
CD3+CD8+HLADR-	39.5 ± 9.2	33.6 ± 11.5	48	74	0.003	0.665	0.002	<35	72	55	3.5	(1.6-7.7)
CD3+CD8-HLADR-	43.1 ± 7.5	46.3 ± 10	48	74		0.603		>47	69	50	2.2	(1.0-4.7)
CD3-CD8-HLADR-	76.7 ± 8.9	80.3 ± 7.9	48	74	0.021	0.597						
CD14+CD13-CD16b-	19.8 ± 4	33.4 ± 22	19	36	0.026	0.703	0.014	>23	68	69	4.9	(1.4-16.3)
CD4+CD14-	20.7 ± 7.8	14.8 ± 5.8	14	23	0.014	0.738	0.017	>14.5	53	83	4.6	(1.4-15.6)
								<16.6	79	65	6.9	(1.5-32.0)

(continued)
 Endometrial leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P1	area under ROC curve ²	P3	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo stage I-IV	Control	Endo								
CD4-CD14-	75.7 ± 8.1	81.0 ± 5.7	14	23	0.025	0.711	0.033	>76	57	83	6.3	(1.4-28.7)
CD4+HLADR-	16.0 ± 5.7	12.3 ± 4.1	14	30	0.018	0.715	0.023	<17	57	87	8.7	(1.9-38.6)
CD13-CD69+	54.2 ± 14.3	42.3 ± 18.2	14	30	0.039	0.705	0.030	<51	71	70	5.8	(1.4-23.6)
CD13+CD45RO-	22.1 ± 8.8	15.5 ± 10.7	25	54	0.009	0.746	4.6 x 10 ⁻⁴	<17.5	76	70	7.5	(2.5-22.3)
CD56-CD122-	49.0 ± 12.2	42.8 ± 15.8	29	51		0.631		<21	52	80	4.2	(1.5-11.8)
CD3+CD69-	26.4 ± 7.0	23.3 ± 8.4	44	76	0.036	0.612	0.041	<27	57	69	3.6	(1.4-9.3)
CD4+CD45RO-	14.9 ± 5.3	13.0 ± 6.2	47	74		0.615	0.034	<16	51	72	2.6	(1.2-5.6)
CD56+CD122+	3.5 ± 2.2	2.6 ± 1.7	29	51		0.651	0.025	<3.0	55	73	3.3	(1.3-8.5)
CD3-CD56+CD16+	8.2 ± 3.8	6.8 ± 4.0	53	78	0.049	0.615	0.026	<6.5	72	55	3.0	(1.4-6.2)
CD3-CD56+CD122+	3.6 ± 2.0	2.6 ± 1.8	27	49	0.033	0.638	0.048	<2.7	63	65	3.2	(1.2-8.5)
CD14+CD13+	3.8 ± 1.9	3.0 ± 2.6	21	36		0.672	0.032	<2.3	81	53	4.8	(1.3-16.9)
CD3+CD20-CD5+	93.0 ± 2.7	88.0 ± 13.8	24	40		0.659	0.034	<91.5	79	55	4.6	(1.4-14.9)
CD4-CD13+CD16+	31.0 ± 14.0	22.4 ± 18.9	16	36		0.699	0.023					
CD69+	41.8 ± 12.9	38.8 ± 17.8	43	78		0.557		<33	81	41	3.0	(1.2-7.4)
ratio CD13/CD3	0.56 ± 0.54	0.78 * 0.71	46	78		0.596		>0.68	80	40	2.6	(1.1-6.1)
CD3-CD20-CD5-	88.0 ± 6.9	90.7 ± 5.2	24	40		0.598		>84	37	90	5.4	(1.4-20.3)
CD3+CD20-CD5-	5.1 ± 2.3	9.5 ± 13.6	24	40		0.683	0.015	>7.7	87	50	9.9	(2.1-48.1)

(continued)
 Endometrial leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P ¹	area under ROC curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo stage I-IV	Control	Endo								
CD4+CD13-	17.5 ± 6.9	15.8 ± 5.4	36	63		0.594		<20.5	42	86	4.3	(1.6-11.3)
CD3+CD44-	41.7 ± 12.0	38.3 ± 13.6	31	56		0.596		<37.8	74	50	2.9	(1.1-7.5)
CD56+	26.2 ± 12.5	30.2 ± 17.2	57	87		0.562		>32	81	41	2.9	(1.3-6.5)
CD13-CD45RO+	21.4 ± 8.7	26.3 ± 11.3	25	54		0.625		>28	80	45	3.2	(1.0-9.8)
CD56+CD69-	19.6 ± 12.9	24.3 ± 14.3	33	53		0.610		>26	85	40	3.4	(1.1-10.2)
CD13-CD16+	8.0 ± 7.5	6.7 ± 3.3	39	71		0.562		<6	72	51	2.6	(1.1-6.1)
CD56+CD122-	21.8 ± 11.8	28.7 ± 17.3	29	51		0.621		>19	59	71	3.1	(1.2-7.9)
CD3+CD4-CD69+	37.2 ± 10.0	34.0 ± 13.8	34	66		0.572		>18	55	72	3.3	(1.3-8.5)
CD4-CD13-CD16+	8.5 ± 3.6	8.7 ± 10.8	16	36		0.642		<7.1	75	58	4.2	(1.1-15.6)
CD4-CD13-CD16-	54.5 ± 13.0	62.2 ± 21.4	16	36		0.655		>65	81	56	5.4	(1.3-22.3)
CD14+CD13+CD16b+	11.0 ± 12.5	6.5 ± 6.2	19	36		0.616		<16	32	94	7.8	(1.4-43.9)
CD56-CD122-	49.0 ± 12.2	42.8 ± 15.8	29	51		0.631		<47	65	68	3.6	(1.4-9.3)
CD4-CD69-	40.7 ± 12.9	46.4 ± 18.6	37	72		0.593		>47	76	47	2.8	(1.2-6.7)
CD3-CD45RO+	23.0 ± 12.5	27.1 ± 16.3	50	73		0.556		>15	34	80	2.0	(0.9-4.5)

5
10
15
20
25
30
35
40
45
50
55

(continued)
Endometrial leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P ¹	area under ROC curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo stage I-IV	Control	Endo								
CD4-CD69+	39.3 ± 11.8	37.7 ± 18.0	37	72		0.462		<35	68	46	1.6	(0.7-3.5)
CD3-CD4-CD45RO+	41.3 ± 18.1	43.4 ± 20.9	43	72		0.530		>31.5	33	71	1.2	(0.5-2.7)
								>56	81	28	1.6	(0.6-4.0)

¹ P value (when ≤ 0.05) obtained in a student "t" test when mean proportion found in patients with endometriosis stage I-IV was compared to normal controls.
² Discriminative value of each marker established by area under ROC curve.
³ P value (when ≤ 0.05), significance of area under ROC curve.
⁴ Confidence interval for odds ratio.

TABLE 4
Peripheral blood leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (%± leukocyte subsets)		s.d.) of		Number of samples tested		P ¹	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Control	Endo. Stage I-IV	Control	Endo. Stage I-IV	Control	Endo. Stage I-IV							
CD3+	66.6 ± 8.5	64.5 ± 8.7	0.032	132	172	0.570	0.037						
CD8+	17.3 ± 5.2	16.4 ± 4.8	0.039	129	172	0.549							
CD13+	16.0 ± 6.0	17.6 ± 6.5	0.020	122	155	0.575	0.032	>17.5	63	51	1.8	(1.1-2.9)	
CD14+	11.8 ± 4.9	13.4 ± 6.0	0.006	124	167	0.575	0.029	>10	45	71	2.0	(1.3-3.3)	
CD20+	5.7 ± 3.1	4.8 ± 2.3	0.006	124	162	0.582	0.017	<6	39	74	1.8	(1.1-3.0)	
CD36+	15.7 ± 6.8	17.2 ± 7.3	0.009	112	140	0.560		>19	77	37	2.1	(1.2-3.6)	
CD44+	17.1 ± 5.6	19.1 ± 6.5	0.023	113	148	0.585	0.018	>18.5	61	51	1.7	(1.0-2.7)	
CD57+	80 ± 3.9	9.2 ± 4.9	0.023	114	148	0.569		>10	75	39	1.8	(1.1-3.1)	
CD69+	19.4 ± 8.2	21.0 ± 7.1	0.017	109	144	0.590	0.014	>21.5	71	45	2.0	(1.2-3.4)	
CD122+	29.2 ± 8.4	31.2 ± 11.7	0.044	122	166	0.567		>34	74	42	2.1	(1.2-3.4)	
CD3+CD5+	66.6 ± 8.5	63.7 ± 10.4	0.017	115	146	0.586	0.017	<69	44	70	1.8	(1.1-2.9)	
CD3+CD45RA-	39.3 ± 9.5	37.2 ± 8.3	0.044	124	168	0.583	0.015	<42	40	72	1.7	(1.1-2.8)	
CD3+CD56-	65.3 ± 8.8	63.2 ± 8.6	0.035	126	169	0.571	0.037	<68	42	71	1.8	(1.1-2.9)	
CD3+CD57-	63.7 ± 8.3	60.7 ± 9.7	0.009	113	146	0.592	0.011	<67	40	77	2.3	(1.3-3.9)	

(continued)
Peripheral blood leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (%± leukocyte subsets)		s.d.) of Number of samples tested		P ¹	Control	Endo-Stage I-IV	area under curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Control	Endo-Stage I-IV	P ¹	Endo-Stage I-IV										
CD3+CD69-	60.1 ± 9.6	57.9 ± 9.2	0.021	141	0.023	107	141	0.584	0.023	<58	69	42	1.6	(1.0-2.7)
CD3+CD122-	62.2 ± 8.4	59.6 ± 9.8	0.006	121	0.004	121	164	0.578	0.024	<4	40	77	2.2	(1.3-3.7)
CD3+HLADR+	3.9 ± 1.4	3.5 ± 1.2	0.003	115	0.001	115	154	0.601	0.001	>14.5	50	72	2.5	(1.5-4.2)
CD3-CD5+	15.5 ± 5.3	18.3 ± 8.4	0.036	123	0.037	123	146	0.619	0.037	>11.5	44	73	2.2	(1.4-3.7)
CD3-CD16+	23.3 ± 7.9	25.4 ± 8.7	0.001	111	0.002	111	143	0.615	0.002	>17.5	75	41	2.1	(1.2-3.6)
CD3-CD44+	13.1 ± 5.3	15.6 ± 6.3	0.023	113	0.044	113	146	0.573	0.044	>19	77	44	2.6	(1.5-4.5)
CD3-CD57+	4.2 ± 2.5	5.0 ± 3.1	0.006	107	0.006	107	141	0.602	0.006	>14.5	61	60	2.3	(1.4-3.7)
CD3-CD69+	14.2 ± 5.6	16.4 ± 6.5	0.006	117	0.008	117	148	0.595	0.008	>25	73	39	1.7	(1.0-2.8)
CD3-CD45RO+	16.0 ± 5.6	18.1 ± 6.8	0.043	132	0.050	132	171	0.566	0.050	>16.5	69	51	2.4	(1.4-4.1)
CD3-CD4-	31.1 ± 9.5	33.3 ± 9.0		122		122	166	0.568	0.050					
CD3-CD8-	33.4 ± 9.2	35.5 ± 10.0		124		124	168	0.595	0.006					
CD3-CD45RA-	14.2 ± 5.2	16.2 ± 6.2	0.004	126	0.006	126	169	0.562	0.006					
CD3-CD56-	21.6 ± 6.3	23.1 ± 7.0	0.030	108	0.030	108	131	0.586	0.023					
CD4-CD13+	14.5 ± 6.0	16.3 ± 6.8												

(continued)
Peripheral blood leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (%± leukocyte subsets)		s.d.) of		Number of samples tested		P ¹	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Control	Endo-Stage I-IV	Control	Endo-Stage I-IV	Control	Endo-Stage I-IV							
CD4-CD36+	13.8 ± 7.4	19.0 ± 6.2	13	21	0.037	0.009	0.771	>19	92	62	19.5	(2.1-179.9)	
CD4-CD69+	16.0 ± 6.1	18.1 ± 6.5	94	120	0.021	0.009	0.603	>19	75	45	2.4	(1.3-4.3)	
CD4-CD45RO+	23.4 ± 6.7	27.0 ± 7.7	27	50	0.043	0.620							
CD4-CD45RA-	22.7 ± 6.5	24.4 ± 8.3	125	168		0.562							
CD8-CD44+	16.8 ± 5.5	18.9 ± 6.7	88	119	0.017	0.588							
CD8-CD44-	66.4 ± 7.0	64.4 ± 6.6	88	119	0.039	0.584		<68	48	70	2.1	(1.2-3.7)	
CD13+CD44+	13.3 5.3	15.4 ± 5.9	96	121	0.006	0.605		>11	38	76	1.9	(1.1-3.4)	
CD13+HLADR+	13.0 ± 5.3	14.7 ± 6.1	108	135	0.024	0.581		>15.5	71	42	1.8	(1.1-3.1)	
CD13+CD16-	1.4 ± 0.9	3.7 ± 4.6	21	37	0.005	0.721		>2	76	57	4.2	(1.3-13.9)	
CD13-HLADR+	8.2 ± 3.2	7.0 ± 3.0	108	135	0.005	0.619		<8	50	70	2.4	(1.4-4.0)	
CD13-CD44-	79.9 ± 6.2	76.9 ± 8.6	96	121	0.004	0.606		<82	39	76	2.0	(1.1-3.6)	
CD14+HLADR+	11.0 ± 4.6	12.7 ± 5.9	110	147	0.009	0.587		>9.5	48	70	2.2	(1.3-3.6)	
CD14+CD44+	10.7 ± 4.9	12.9 ± 5.7	85	123	0.003	0.612		>15	85	33	2.7	(1.3-5.4)	
CD14+CD45RO+	12.2 ± 4.4	14.3 ± 6.1	102	118	0.004	0.586							

(continued)
Peripheral blood leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (%± leukocyte subsets)		s.d.) of		Number of samples tested		P ¹	Control	Endo-Stage I-IV	area under curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Control	Endo-Stage I-IV	Control	Endo-Stage I-IV	Control	Endo-Stage I-IV										
CD14+CD16	0.7 ± 0.8	2.7 ± 5.4	0.023	0.023	23	41	0.603									
CD14+CD122	0.4 ± 0.3	2.4 ± 5.2	0.029	0.001	14	34	0.648	>0.7	86	47	5.3	(1.0-27.5)				
CD14-HLADR+	10.3 ± 3.3	9.1 ± 2.7	0.001	0.023	109	147	0.611	<8.5	69	47	2.0	(1.2-3.3)				
CD14-CD44-	83.5 ± 5.3	81.6 ± 6.3	0.023	0.005	85	123	0.588	<80	79	37	2.2	(1.2-4.2)				
CD20+HLADR+	5.2 ± 2.9	4.3 ± 2.1	0.005	0.033	106	141	0.596	<5.5	39	75	1.9	(1.1-3.3)				
CD20+CD44-	5.0 ± 2.8	4.2 ± 2.2	0.033	0.0004	95	126	0.571	<4	62	55	2.0	(1.2-3.4)				
CD20-CD44+	14.4 ± 5.1	17.3 ± 6.2	0.0004	0.016	95	126	0.636	>17	74	47	2.5	(1.4-4.4)				
CD20-CD69+	16.4 ± 4.3	21.0 ± 6.2	0.016	0.004	14	29	0.719	>16	57	76	4.2	(1.1-16.3)				
CD20-HLADR+	15.4 ± 4.9	17.6 ± 6.9	0.004	0.008	106	141	0.591	>14.5	49	70	2.2	(1.3-3.7)				
CD20-CD44-	79.5 ± 5.1	77.5 ± 6.1	0.008	0.065	95	126	0.590	<75.6	71	52	11.6	(2.3-57.0)				
CD20-CD69-	79.4 ± 6.7	75.3 ± 6.5	0.065	0.0005	14	29	0.711	<5.6	77	40	2.2	(1.2-4.0)				
CD36-HLADR+	7.8 ± 3.2	6.4 ± 2.0	0.0005	0.083	95	121	0.629	>20.5	85	52	6.0	(1.1-33.3)				
CD56-CD69+	18.1 ± 4.2	21.8 ± 6.7	0.083	0.028	13	23	0.667	>23	74	40	2.0	(1.2-3.3)				
CD56-CD122+	19.5 ± 5.7	21.4 ± 8.6	0.028		113	154	0.578									

(continued)
Peripheral blood leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (%± leukocyte subsets)		s.d.) of		Number of samples tested		P ¹	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Control	Endo-Stage I-IV	Control	Endo-Stage I-IV	Control	Endo-Stage I-IV							
CD56-CD69-	67.9 ± 5.3	62.7 ± 9.4	13	23	0.075	0.093	0.671	0.093	<64.5	67	51	2.1	(1.3-3.5)
CD56-CD122-	67.7 ± 8.3	64.7 ± 11.5	113	154	0.014	0.017	0.586	0.017	<64.5	67	51	2.1	(1.3-3.5)
CD57-CD44-	74.3 ± 6.9	71.7 ± 7.7	84	117	0.015	0.014	0.602	0.014	<76	44	71	1.9	(1.1-3.5)
CD3+CD57- HLADR+	4.3 ± 1.5	4.0 ± 1.4	96	118	0.032	0.034	0.584	0.034	<3.7	72	43	1.9	(1.1-3.5)
CD3-CD4-CD44+	34.4 ± 11.0	38.1 ± 12.6	88	112	0.032	0.037	0.577	0.037	>40.5	69	45	1.8	(1.0-3.3)
CD3-CD4-CD44-	58.3 ± 12.2	54.9 ± 12.7	88	112			0.572						
CD3-CD56+CD16-	1.2 ± 0.6	1.4 ± 1.0	121	163	0.023	0.024	0.553	0.024					
CD3-CD56-CD122-	23.4 ± 11.1	21.2 ± 11.4	113	148			0.581	0.024					
CD3-CD57-CD44-	48.7 ± 10.1	44.5 ± 10.3	84	114	0.004	0.008	0.610	0.008	<41.3	80	41	2.8	(1.4-5.3)
CD14+CD20+CD44-	0.2 ± 0.2	0.1 ± 0.1	65	75	0.036	0.037	0.613	0.037					
CD14+CD20-CD44+	91.8 ± 4.6	93.5 ± 3.5	65	75	0.016	0.021	0.614	0.021	>95	75	43	2.3	(1.1-4.7)
CD14+CD20-CD44-	5.9 ± 4.5	4.6 ± 3.3	65	75	0.037	0.035	0.604	0.035	<3	75	43	2.3	(1.1-4.7)
CD14-CD13- HLADR+	8.7 ± 3.5	7.6 ± 2.6	77	82	0.021	0.029	0.600	0.029	<7	71	45	2.1	(1.1-4.0)
Ratio CD13/CD3	0.25 ± 0.12	0.29 ± 0.14	121	153	0.025	0.018	0.583	0.018	>0.30	71	43	1.8	(1.1-3.0)

5
10
15
20
25
30
35
40
45
50
55

(continued)
Peripheral blood leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (%± leukocyte subsets)		s.d.) of		Number of samples tested		P ¹	Control	Endo. Stage I-IV	area under curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Control	Endo. Stage I-IV	Control	Endo. Stage I-IV	Control	Endo. Stage I-IV										
Ratio CD13/CD8	1.04 ± 0.55	1.20 ± 0.69	0.039	0.039	114	153	0.039	0.574	0.039	0.574	0.039	>0.14	40	74	1.9	(1-1-3.1)
Ratio CD14/CD3	0.19 ± 0.10	0.22 ± 0.12	0.011	0.011	123	165	0.028	0.575	0.028	0.575	0.028	>0.14	40	74	1.9	(1-1-3.1)
Ratio CD14/CD8	0.78 ± 0.47	0.91 ± 0.55	0.040	0.040	116	164	0.036	0.574	0.036	0.574	0.036	>0.14	40	74	1.9	(1-1-3.1)

¹ P value (when ≤ 0.05) obtained in a student "t" test when mean proportion of leukocyte subsets was compared between patients with endometriosis (stage I-IV) and normal controls.

² Discriminative value of each marker established by area under ROC curve.

³ P value (when ≤ 0.05), significance of area under ROC curve.

⁴ Confidence interval for odds ratio.

TABLE 5
Endometrial leukocyte populations used as markers to discriminate between patients with endometriosis stage I-II or stage III-IV and normal subjects

Leukocyte Subsets	Controls	Mean proportion (% ± s.d.) of leukocyte subsets		P^1	Endo stage III-IV	P^2	Number of samples tested		Endo area under ROC curve	P	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
		Endo stage I-II	Endo I-II				Control	Endo III-IV							
CD3-CD44-	50.0 ± 12.9	50.5 ± 11.9	61.8 ± 16.6	0.015	0.015	31	43	13	0.766	0.007	>48	52	92	12.9	(1.4-113.8)
CD3+HLADR+	8.3 ± 5.7	6.1 ± 2.6	0.013	7.6 ± 3.3	0.013	51	57	23	0.587		<5.5	67	48.3		
CD3+CD45RO+	14.9 ± 8.4	12.0 ± 5.6	0.043	13.2 ± 9.5	0.043	50	50	23	0.565		>15	37.3	71		
CD4+CD13+	2.3 ± 1.9	2.2 ± 1.7	1.4 ± 1.1	0.025	0.025	36	44	19	0.658		<1.7	50	79	3.8	(1.0-13.8)
CD13+HLADR+	5.7 ± 2.7	5.3 ± 2.3	3.6 ± 1.3	0.017	0.017	27	37	12	0.740	0.021	<5	54	92	13	(1.4-117.2)
CD14+HLADR+	14.7 ± 7.5	10.2 ± 3.9	0.049	10.8 ± 5.1	0.049	14	24	4	0.679		<13	57	75		
CD56+HLADR+	2.8 ± 1.7	1.5 ± 0.7	0.011	4.6 ± 3.3	0.011	16	17	5	0.741	0.018	<1.7	75	71	7.2	(1.5-33.6)
CD56-CD44-	63.5 ± 13.0	63.5 ± 13.4	51.1 ± 20.4	0.023	0.023	30	41	12	0.708	0.041	<65	58	83	6.8	(1.2-37.5)
CD3-CD4+CD45RA-	1.5 ± 0.9	1.5 ± 0.9	1.0 ± 0.7	0.022	0.022	51	54	25	0.668	0.021	<0.7	84	40	3.4	(1.1-10.7)
CD3+CD8+HLADR+	15.9 ± 8.1	15.8 ± 8.6	23.6 ± 12.1	0.011	0.011	48	52	22	0.701	0.009	>14.8	55	73	3.3	(1.1-10.1)
CD3-CD8+HLADR-	9.5 ± 6.5	8.6 ± 6.1	6.1 ± 5.2	0.036	0.036	48	52	22	0.691	0.013	<5	78	59	5.0	(1.6-15.4)
CD3+CD56+CD16+	3.0 ± 3.4	2.9 ± 3.2	6.0 ± 5.7	0.027	0.027	53	56	22	0.670	0.024	>1.8	49	77	3.5	(1.0-10.3)
CD3+CD56-CD16+	6.6 ± 7.5	7.0 ± 6.2	11.0 ± 9.1	0.350	0.350	53	56	22	0.698	0.009	>9	84	50	5.4	(1.7-17.3)
CD3+CD56-CD16-	87.0 ± 8.1	87.2 ± 7.1	79.5 ± 12.9	0.016	0.016	53	56	22	0.695	0.010	>83	81	50	4.1	(1.4-12.5)

5
10
15
20
25
30
35
40
45
50
55

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		p ¹	Endo stage III-IV	p ²	Number of samples tested		area under ROC curve	P	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo stage I-II				Control	Endo I-II							
CD3+CD56-CD44+	7.0 ± 6.1	6.1 ± 4.8		13.2 ± 10.5	0.025	28	38	0.721	0.034	<5	42	92		
CD3+CD56-CD44-	87.1 ± 6.1	88.5 ± 6.2		78.0 ± 13.9	0.049	28	38	0.668		>81	83	58	7.0	(1.5-33.7)
CD3-CD56-HLADR-	29.7 ± 15.6	42.7 ± 19.6	0.050	29.3 ± 21.3		15	17	0.706	0.047	<36	73	59		
CD3+CD56-CD122+	12.1 ± 7.5	11.9 ± 6.3		21.7 ± 14.0	0.005	27	33	0.715	0.024	>20	87	50	6.7	(1.4-31.7)
CD3+CD56-CD122-	81.9 ± 8.2	83.0 ± 6.8		68.0 ± 21.7	0.024	27	33	0.715	0.024	>76.5	71	63	4.0	(1.1-15.5)
CD4-CD16-	51.5 ± 12.9	59.1 ± 12.7	0.033	49.3 ± 21.4		24	31	0.659	0.045	<58	76	47		
CD14+CD13+CD16b-	62.7 ± 20.7	56.2 ± 25.8		38.2 ± 26.4	0.011	19	26	0.753	0.035	>61	69	80	8.8	(1.3-57.4)
CD16+	27.4 ± 12.4	30.0 ± 16.2		35.4 ± 19.5		58	62	0.603		>39	85	39	3.5	(1.2-10.5)
CD45RA+	28.3 ± 9.1	27.6 ± 9.0		27.9 ± 16.0		56	60	0.562		>27	50	51		
CD45RO+	38.4 ± 13.6	38.0 ± 15.5		44.6 ± 18.1		50	54	0.620		<23.5	75	48	2.8	(1.0-7.5)
CD13+	24.8 ± 12.7	25.5 ± 14.7		31.3 ± 20.3		47	57	0.620		>52	88	43	4.4	(1.3-14.4)
						47	57	0.620		>29	68	34		

¹ P value obtained in a student "t" test when mean % of leukocyte subsets found in patients with endometriosis stage I-II was compared to normal controls.

² P value obtained in a student "t" test when mean % of leukocyte subsets found in patients with endometriosis stage III-IV was compared to normal controls.

TABLE 6
Peripheral blood leukocyte populations used as markers to discriminate between patients with endometriosis stage I-II or stage III-IV and normal subjects

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		P^1	Endo stage III-IV	P^2	Number of samples tested		Endo area under ROC curve	P	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI)*
	Controls	Endo stage I-II				Control	Endo I-II							
CD3-CD57+CD44-	12.5 ± 6.2	14.9 ± 7.5	0.030	12.2 ± 6.9	0.037	84	81	0.586	0.008	>15.6	74	43	2.1	(1.1-4.1)
CD14-CD13-HLADR-	85.6 ± 5.0	87.6 ± 3.9	0.016	85.9 ± 3.3	0.012	77	54	0.637	0.008	>88.3	74	47	2.6	(1.1-5.3)
CD14-CD20+CD44-	5.7 ± 2.9	4.7 ± 2.6	0.043	5.6 ± 2.7	0.037	65	50	0.607	0.049	<4.5	66	53	2.2	(1.0-4.7)
HLADR+	21.6 ± 6.0	21.7 ± 6.6		23.8 ± 6.7	0.037	120	102	0.586		>24.5	73	45	2.2	(1.1-4.5)
CD3+CD8-	49.9 ± 8.5	49.0 ± 9.2		46.3 ± 8.8	0.012	122	113	0.654	0.002	<51.5	48	85	5.2	(2.2-12.1)
CD3+CD44-	60.2 ± 8.6	59.2 ± 8.8		57.2 ± 8.0	0.047	111	98	0.605	0.047	<61.2	51	73	2.8	(1.3-6.1)
CD3+HLADR-	63.3 ± 8.7	63.0 ± 9.1		59.6 ± 7.9	0.009	121	102	0.628	0.010	<63.5	55	73	3.2	(1.5-6.6)
CD3-HLADR+	18.1 ± 5.8	18.7 ± 6.5		20.8 ± 6.9	0.010	121	102	0.607	0.030	>21.5	73	45	2.3	(1.1-4.6)
CD3+CD16+	9.3 ± 13.3	8.4 ± 12.1		6.0 ± 4.3	0.016	123	113	0.528		>3.7	31	81		
CD3-CD57-	28.3 ± 6.9	29.0 ± 7.2		32.3 ± 9.5	0.004	113	101	0.644	0.007	>28	53	71	2.7	(1.2-5.7)
CD4-HLADR+	19.8 ± 6.0	19.8 ± 6.2		22.0 ± 6.6	0.036	112	96	0.588		>19.3	48	67		
CD4+CD45RA-	31.4 ± 8.1	29.7 ± 7.2		27.7 ± 8.5	0.006	125	114	0.625	0.010	<29.5	60	60	2.3	(1.2-4.5)
CD4+CD45RO+	19.5 ± 5.6	20.5 ± 6.4		16.2 ± 4.3	0.029	27	31	0.645		<19.8	50	90	8.5	(1.6-44.5)
CD13-CD16+	18.9 ± 8.2	16.8 ± 8.7		14.3 ± 3.1	0.035	21	28	0.698		<13	86	56	7.5	(1.2-45.1)

5
10
15
20
25
30
35
40
45
50
55

Leukocyte Subsets	Controls	Mean proportion (% ± s.d.) of leukocyte subsets		P ¹	Endo stage III-IV	P ²	Number of samples tested		area under ROC curve	P	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
		Endo stage I-II	Endo stage I-II				Control	Endo I-II							
CD14-CD69-	81.0 ± 6.1	77.4 ± 9.1	76.0 ± 6.7	0.031	20	32	14	0.729	0.025	<82	50	86	6.0	(1.1-34.0)	
CD20-HLADR-	79.3 ± 5.2	78.5 ± 7.0	77.1 ± 6.6	0.025	106	97	44	0.573		<80.3	41	72			
CD57-CD44+	17.2 ± 5.5	18.3 ± 6.4	19.7 ± 7.7	0.046	84	82	35	0.573		>21.6	80	40	2.7	(1.1-6.7)	
CD3-CD4+CD45RA-	6.9 ± 4.4	6.4 ± 4.0	5.4 ± 3.5	0.043	122	109	50	0.587		<5.5	59	61	2.3	(1.1-4.5)	
CD3-CD8-CD44+	37.2 ± 11.9	37.2 ± 12.7	43.2 ± 12.6	0.019	87	80	32	0.588		>40.5	54	63			
CD3-CD57-CD44+	38.5 ± 12.1	40.3 ± 12.5	44.0 ± 12.7	0.029	84	81	33	0.575		>44	63	58			
CD14+CD20+CD44+	2.1 ± 1.3	2.1 ± 1.5	1.4 ± 0.7	0.001	65	52	23	0.647		<2.1	39	91	6.4	(1.3-30.9)	

¹ P value obtained in a student "t" test when mean proportion found in patients with endometriosis stage I-II was compared to normal controls.

² P value obtained in a student t test when the % leukocyte subsets found in patients with endometriosis stage III-IV was compared to normal controls.

[0030] Cutoff points established for each individual marker are presented in Table 3, 4, 5, 6 and threshold value established for a particular marker combination are presented in Table 1. Any other cutoff points or threshold values providing a valuable diagnostic test for endometriosis are meant to be included in the present application.

5 [0031] In accordance with the present invention, there is provided a series of 34 different combinations of endometrial leukocyte markers (Tables 1 and 2), 7 combinations of blood leukocyte markers (Table 1) and 4 combination of endometrial and blood leukocyte markers providing a diagnostic test with levels of sensibility and specificity up to 89 and 90%, respectively. The different marker combinations of the present invention may serve several important clinical needs. Hence in the general population, these markers could be used to evaluate the risk factor to develop endometriosis or to identify women with high likelihood of suffering of the disease. Furthermore in patients with endometriosis, these markers could serve to monitor the disease or to give a prognosis.

10

Study subjects and samples

[0032] Uterine endometrial tissues were obtained from 146 subjects undergoing laparoscopy or laparotomy. The experimental group was formed of 88 subjects with endometriosis stage I-IV confirmed by laparoscopy or laparotomy and the control group consist of 58 healthy subjects who underwent surgery for tubal ligation (or reanastomosis) and had no clinical evidence, nor family history of endometriosis. Table 7 gives details concerning the age, menstrual cycle and indication of laparoscopy or laparotomy for the subjects included in experimental and control groups.

15

20

25

30

35

40

45

50

55

TABLE 7
Description of the experimental groups used in the analysis of endometrial leukocyte populations

Experimental groups	Number of subjects	Mean age ±s.d.	Menstrual cycle		Percentage of patients*			Other**
			ES ¹	LS ² *	ligation or reanastomosis	Hysterectomy and/or ovariectomy	Diagnostic laparoscopy	
Controls	58	34.2 ± 5.3	54.5%	45.5%	100%			
Endometriosis I-IV	88	34.4 ± 6.8	47.0%	53.0%	21.6%	22.7%	52.3%	3.4%
Stage I-II	63	34.4 ± 7.3	50.9%	49.1%	28.6%	22.2%	47.7%	1.5%
Stage III-IV	25	34.4 ± 5.4	36.4%	63.6%	4.0%	24.0%	64.0%	8.0%

¹Early secretory (days 14-21)
² Late secretory (days 22-28)
* % patients among control or endometriosis groups

EP 1 147 421 B1

[0033] Endometrial biopsies were taken with a Pipet Curette (Millex) (approximately 0.5g of tissue). All samples were harvested in the secretory phase (day 14-28) of the menstrual cycle as confirmed by histological evaluation. The samples were collected into sterile RPMI-1640 medium (Gibco) supplemented with 2% heat-inactivated fetal calf serum (Bio-Media) and 1% penicillin-streptomycin and kept at 4°C until cell isolation.

5 **[0034]** Blood samples were obtained from 172 subjects with endometriosis (stage I-IV) confirmed by laparoscopy or laparotomy and from 132 healthy subjects with no evidence of endometriosis at surgery, and no family history of endometriosis. Blood samples (30 ml) were collected in heparin-tubes (Vacutainer™, Becton Dickinson) and kept at 20°C until mononuclear cell separation. The age, menstrual dating and indication for laparoscopy of the subjects included in the study are given in Table 8.

10

15

20

25

30

35

40

45

50

55

TABLE 8
Description of the subjects included in the study of peripheral blood mononuclear leukocytes

Experimental group	Number of subjects	Mean age ± s.d.	Menstrual cycle		ligation or reanastomosis	Percentage of patients**		
			Proliferative **	Secretory**		Hysterectomy and/or ovariectomy	Diagnostic laparoscopy	Other
Control	132	34.30 ± 5.5	43.8 %	56.2%	100%			
Endometriosis I-IV	172	36.40*	42.8%	57.2%	22.1%	33.7%	38.9%	5.3%
Stage I-II	116	35.96 ± 6.39	41.0%	59.0%	31.1%	30.2%	37.1%	1.6%
Stage III-IV	56	34.30 ± 5.5	46.2%	53.8%	3.6%	41.1%	42.9%	12.4%

* P versus normal = 0.002

** % of patients amongst control or endometriosis groups

Stromal cell preparation from endometrial samples

[0035] Endometrial tissue samples were mechanically disrupted with a Pyrex™ glass Broeck tissue grinder (Fisher) to obtain a single cell suspension. Stromal cell fraction was isolated by filtration through a 250 µm stainless steel sieve (Millipore) to retain the glandular fraction and was washed twice with 10 ml phosphate buffered saline (PBS) (Sigma) containing 1% BSA (Boehringer Mannheim), 0.1% sodium azide (Fisher) (thereafter called PBS washing buffer).

Isolation of mononuclear cells from peripheral blood

[0036] Blood samples were diluted 1:1 with Hank's Balanced Salt Solution (HBSS) (Gibco), layered on an equal volume of Ficoll-Paque™ (Pharmacia Biotech) and centrifuged at 1500 rpm for 40 minutes at room temperature. Leukocytes were isolated at the interface of Ficoll and HBSS and they were washed in 50 ml of HBSS. Contaminating red blood cells were lysed with 6 ml of ammonium chloride solution (0.15M) (6 minutes at room temperature). The peripheral blood mononuclear cells were then washed twice in 10 ml PBS and resuspended in PBS washing buffer.

Endometrial and peripheral blood leukocyte surface antigen staining

[0037] Endometrial stromal cells or peripheral blood mononuclear cells were distributed in 5 ml tubes (1 to 1.5 x 10⁶ cells/tube) or in 96 well plates (5 x 10⁵ cells/well), respectively and incubated in the presence of 0.1 µg of human γ-globulin for 5 minutes at room temperature. The cells were then incubated 30 minutes in the dark (at room temperature for endometrial cells and at 4°C for peripheral blood mononuclear cells) with a panel of 4 different mouse monoclonal antibodies (MAbs) in a total volume of 100 µl. The cell samples were stained with mouse anti-human CD45 MAbs conjugated to peridinin chlorophyl protein (PerCP) and with several sets of three different mouse MAbs labeled with distinct fluorochromes (fluorescein isothiocyanate - FITC-, phycoerythrin -PE or with phycoerythrin-texas red -ECD-) directed toward cell surface markers for specific cell populations such as T lymphocytes, B lymphocytes, NK cells, macrophages and/or activation markers (Table 9).

TABLE 9
Description of mouse monoclonal antibodies used for immunophenotyping

Specificity	Clone	Isotype	Supplier	Fluorochrome
CD3	HIT3A	mouse IgG2a	Beckman/Coulter	ECD
CD4	SK3	mouse IgG1	Becton Dickinson	PE
CD5	BL1A	mouse IgG2a	Beckman/Coulter	FITC
CD8	SK1	mouse IgG1	Becton Dickinson	PE
CD13	SJ1D1	mouse IgG1	Beekman/Coulter	RPE
CD14	RM052	mouse IgG2a	Beckman/Coulter	PE
CD16	NKP15	mouse IgG1	Becton Dickinson	FITC
CD16B	1d3	mouse IgM	Beckman/Coulter	FITC
CD20	H299	mouse IgG2a	Beckman/Coulter	RD1
CD36	SMf	mouse IgM	Sigma	RPE
CD44	L178	mouse IgG1	Becton Dickinson	FITC
CD45	2DI	mouse IgG1	Becton Dickinson	PerCP
CD45-RA	ALB11	mouse IgG1	Beckman/Coulter	FITC
CD45-RO	UCHLI	mouse IgG2a	Beckman/Coulter	FITC
CD56	N901(NKH-1)	mouse IgG1	Beckman/Coulter	PE
CD57	VC1.1	mouse IgM	Sigma	RPE
CD69	L78	mouse IgG1	Becton Dickinson	FITC
CD122	2RB	mouse IgG1	Beckman/Coulter	FITC
HLA-DR	L243	mouse IgG2a	Becton Dickinson	FITC

[0038] Table 10 below lists the distribution of the antigens listed in Table 9.

TABLE 10
Main distribution of antigens

Antigen	Main Cell Distribution
5 CD3	Expressed on all mature T cells associated with TCR complex (α/b , γ/δ)
CD4	Expressed on T helper lymphocytes. It can be also expressed on cells of the monocyte/macrophage lineage
CD5	Found on all mature T lymphocytes and a subset of B lymphocytes
10 CD8	Found on a subset of T lymphocytes called suppressor/cytotoxic T cells.
CD13	Detected on most cells of myeloid origin polymorphonuclear cells or cells of the monocyte/macrophage lineage. Member of metalloproteinase family
15 CD14	Expressed strongly on the surface of monocytes Found on most tissue macrophages Weakly expressed on the surface of granulocytes and B lymphocytes Receptor for lipopolysacharride-(LPS) and LPS binding protein
CD16	Expressed mainly on NK cells, monocytes macrophages and polymorphonuclear leukocytes Low affinity receptor for IgG
20 CD 16b	Found on granulocytes including polymorphonuclear cells (PMN)
CD20	Present on all B lymphocytes
CD36	Expressed on platelets, monocytes or macrophages, microvascular endothelial cells, mammary endothelial cells, during stages of erythroid cell development
25 CD44	Widely expressed on the surface of most cell types. Including most leukocytes and epithelial cells. Family of core/link peptidoglycan
30 CD45	Present on the surface of all leukocytes
CD45RA	Isoforms of CD45 Found on naive/resting T cells Also expressed on B lymphocytes and monocytes
CD45RO	Isoforms of CD45 expressed on memory/activated T cells also expressed on monocytes
35 CD56	Marker for NK cells Can also be found on a population of T lymphocytes
CD57	Found on a subset of cells with natural killer activity
CD69	Expressed on activated leukocytes including T cells, B cells, NK cells, neutrophils, eosinophils and cells of the monocyte/macrophage lineage. Activation marker detected early after cell activation
40 CD122	Expressed on NK cells B, T lymphocytes or monocytes/macrophages Component of the IL-15 receptor
HLADR+	HLA class II molecule Found on antigen presenting cells or on other cells upon activation such as T cells.

45 **[0039]** Blood cells were washed twice with 0.15 ml of PBS washing buffer. Endometrial cell samples were incubated with a red blood cell lysing solution, (FACS™ Lysing Solution, Becton Dickinson) for 10 minutes at room temperature in the dark and washed with 3 ml of PBS washing buffer. Endometrial and blood cells were fixed in 1% paraformaldehyde (diluted in PBS) at a concentration of 1×10^6 cells/ml and kept at 4°C in the dark until the immunofluorescence reactivity was determined by flow cytometry.

Flow cytometry analysis

55 **[0040]** The immunofluorescence reactivity was carried out on a Coulter EPICS XL™ flow cytometer (Coulter Corporation, Hialeah, FL) equipped with an argon laser operating at 488 nm, 15 mW and detectors at 525, 575, 610, and 675 nm. Calibration of the flow cytometer parameters for forward scatter, side scatter and fluorescence were the same for all the samples. Cells expressing CD45 pan leukocyte antigen were gated using the Coulter system II software. The

percentage of cells bearing markers for T, B lymphocytes, macrophages or NK cells and/or activation markers was evaluated within the CD45 positive populations only. A minimum of 6000 CD45+ cells were analyzed for each sample.

Use of leukocyte markers in a diagnostic test for endometriosis

[0041] A cutoff point was established for the proportion of the endometrial or blood leukocyte subpopulations identified as diagnostic markers. The value obtained for each marker is compared to the cutoff point (Fig. 1). A positive result was given when the proportion of a particular leukocyte subset fulfills the condition established by the cutoff point (for example < 40% for CD3+ cells). When these markers are used in combination, a positive result for each marker gives a score of 1, whereas a negative result gives a score of 0. A diagnosis of endometriosis is given, when the final diagnostic score obtained from adding the results of all the markers of a particular combination is higher than a predetermined threshold value. The levels of sensibility and/or specificity measured for the marker combination represents the number of positive test results obtained among the patients already confirmed with endometriosis and the number of negative test results among the subjects within the control group, respectively.

[0042] Endometrial and blood leukocyte markers can be used in combinations in logistic regression model:

$$P(r) = \frac{e^{c + B1*(marker1) + B2 (marker2) + \dots Bn (marker n)}}{1 + e^{c + B1*(marker1) + B2 (marker2) + \dots Bn (marker n)}}$$

Where:

P(r) = probability of having endometriosis

c = constant established for a particular combination

B = coefficient of regression

n = total number of markers in the combination

[0043] The probability of having endometriosis (P(r)) is then compared to a threshold value that provides the best discriminative value. A positive diagnosis of endometriosis is given when the P(r) value exceeds the threshold value established for a particular combination of markers. Alternatively, a negative diagnosis of endometriosis is given when the P(r) value is lower than the threshold value.

Results

[0044] Endometrial and blood leukocyte subsets defined as good potential markers for the diagnosis of endometriosis are presented in Tables 3 and 4 respectively. Selection of these markers was done on the basis of a significative difference in the mean proportion of leukocyte subsets between patients with endometriosis (stage I-IV) and control groups. In addition, several endometrial and blood markers were also selected according to the area under the ROC curve, an indication of the discriminative value of the markers. The ROC curve allowed the determination of one or more cutoff proportion that best discriminate between patients with endometriosis (stage I-IV) and normal controls. In an attempt to use these differences for identifying patients with endometriosis, a positive test result was given when the proportion measured for a particular leukocyte subset fulfills the condition established by the cutoff point (for example < 40% for CD3+ cells). The levels of specificity and sensibility were calculated for each marker used alone to diagnose endometriosis and are presented in Table 3 (for endometrial leukocyte markers) and Table 4 (for blood leukocyte markers). Moreover, a significant odds ratio calculated with a particular cutoff point gave an additional indication that the leukocyte markers selected in Tables 3 and 4 are associated with an increased risk to develop endometriosis and can, thus, be used for identifying women with high likelihood of suffering of endometriosis.

[0045] The mean proportion of some endometrial (Table 5) and blood (Table 6) leukocyte subsets was found to be significantly modulated only in patients with stage I-II endometriosis or with stage III-IV endometriosis when compared to normal controls. Nevertheless, these markers remains good candidates for a diagnostic test for endometriosis, but their use may be limited to a specific stage of the disease.

[0046] Several of the endometrial and blood leukocyte markers were found to be even more reliable as diagnostic markers when they are analyzed in combination with other markers. Table 1 gives a series of 33 combinations in which endometrial or blood leukocyte markers are used in a diagnostic test for endometriosis. For each marker, a positive test result (as described above) gives a score of 1, whereas a negative test result gives a score of 0. The final diagnostic value obtained from adding the scores of all the markers of a particular combination is then compared to a threshold value, which is indicated in Table 1.

[0047] A diagnosis of endometriosis is given, when the diagnostic value exceeds the threshold value established for each set of combination markers- The use of leukocyte marker subsets in combination in this new method clearly improves the levels of sensibility and/or specificity for diagnosing endometriosis. Table 1 also provides 4 examples showing that blood leukocyte markers, when used in combination with endometrial markers, can also increase the predictive value of the diagnostic test.

[0048] The present invention also demonstrates that logistic regression models can also be used to combine endometrial as well as blood leukocyte markers for the development of a predictive model of endometriosis (Table 2). In some cases, these models need to be adjusted with risk factors associated with endometriosis such as the length of the menstrual cycle, the duration of menstruation, pain during intercourse and age. In some instances, these factors were shown to increase the predictive value of the model.

[0049] The present invention identifies several examples of marker combinations, which give rise to diagnostic methods yielding improved levels of sensibility and specificity. Indeed, the different marker combinations of the present invention may serve different clinical applications including screening, diagnosis, monitoring and prognosis of endometriosis.

Claims

1. A method for determining endometriosis in a sample of eutopic uterine endometrial tissues or a sample of blood of a female subject, comprising:

a) measuring in said sample a quantitative level of at least two different surface antigens from blood leukocytes and/or eutopic endometrial leukocytes, said surface antigens from blood leukocytes and/or eutopic endometrial leukocytes being selected from the leukocyte marker combinations defined in the following Table:

Leukocyte Marker Combinations	
Endometrial leukocyte markers	
	CD4+, CD8+CD69-, and CD13+CD45RO-
	CD4+, CD8+CD69-, CD56+CD122-, CD3+CD45RA-, and CD13+CD45RO-,
	CD4+, CD8+CD69-, CD13-CD122+, and CD13+CD45RO-
	CD4+ CD8+CD69-, and D14+CD13-CD16b-
	CD3+CD16-, CD13+CD45RO-, CD3+, CD8+, and CD3+CD69+
	CD3+, CD3+CD8+, CD13+CD45RO-, and CD3-CD20-
	CD3+CD8+, CD13+CD45RO-, CD3+CD5+, CD3+CD122-, CD3-CD20-, and CD3+CD45RO-
	CD3+CD8+, CD13+CD45RO-, CD3+CD5+, and CD3+CD122-
	CD3+CD20-CD5-, CD4+CD13-, and CD56-CD122-
	CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, and CD3+CD45RO-
	CD3+, CD8+CD69-, CD3-CD4-CD45RO+, ratio CD13+/CD3+, and CD13+CD45RO-
	CD3+CD8+, CD13+CD45RO-, CD3-CD5-, and CD20-CD5+
	CD8+, CD5+, CD3-CD20-, and CD3-HLADR-
	CD3+CD8+, CD13+CD45RO-, and CD5+
	CD4+, CD13-CD122+, CD8+CD69-, and CD3+CD45RA-
	CD4+, CD8+CD69-, CD13-CD122+, CD3+CD45RA-, and CD13+CD45RO-
	CD4+, CD8+CD69-, CD56-CD122-, CD3+CD45RA-, CD13+CD45RO-
	CD3+, CD4+, CD3+CD8+, CD13+CD45RO-, and CD3+CD5+
	CD3+, CD4+, CD3+CD8+, and CD13+CD45RO-
	CD3+, CD3+CD8+, and CD13+CD45RO-
	CD3+, CD5+, CD3+CD5+, CD69+, and CD4-CD69+
	CD3+, CD3+CD8+, and CD13+CD45RO-
	CD3+, CD3+CD8+, and CD13+CD45RO-
	CD3+, CD3+CD8+, CD13+CD45RO-, CD3-CD20-, and CD56-CD16-
	CD3+CD16-, CD3-CD4-CD45RO+, CD3+, CD8+, CD3+CD69+, and CD13+CD45RO-
	CD3+, CD3-CD45RO+, ratio CD13+/CD3+, CD3+CD8+, and CD8+CD69-
	CD3+CD20-CD5-, CD4+CD13-, CD56-CD122-, and CD4+CD45RO-
	CD3+, CD3+CD8+, CD13+CD45RO-, and CD3-CD20-

(continued)

<p>Endometrial leukocyte markers CD13+CD45RO-, CD4+CD45RA-, CD3+CD122-, and CD8+CD69- CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, CD3+CD45RO-, and CD3-CD5- CD3+, CD3-CD5-, CD13+CD45RO-, and CD3-CD20- CD3+, CD3-CD5-, and CD13+CD45RO- CD3+CD8+, CD13+CD45RO-, and CD3+CD5+ CD3+ and CD13+CD20.</p>
<p>Blood Leukocyte markers CD3-CD5+, CD3-CD45RA-, CD3-CD44+, CD13+, and CD3-CD57-CD44- CD3-CD45RA-, CD20-CD44+, CD20-HLADR+, CD3-CD4-CD44+, and CD36-HLADR+ CD3-CD45RA-, CD3-CD45RO+, and CD20-HLADR+ CD14+CD44+, CD57+, CD3-CD45RA, and CD14+ CD14+, CD57+, CD3+CD69+, CD3+HLADR+, and CD3-CD45RA- CD14+, CD57+, CD3+HLADR- and CD3-CD45RA- CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-</p>
<p>Blood (in italics) and endometrial leukocyte markers <i>CD57+, CD14+, CD3-CD69+</i>, CD3+, CD4+, and CD3+CD8+HLADR- <i>CD3-CD69+</i>, CD3+, CD4+, and CD3+CD8+HLADR- <i>CD4-CD36+</i>, <i>CD3-CD69+</i>, CD8+, CD13+, CD3+, CD16+, CD69+, and CD5+ <i>CD3-CD45RA-</i>, <i>CD3-CD45RO+</i>, <i>CD20-HLADR+</i>, <i>CD14+CD44+</i>, CD8+, CD5+, CD3-CD20-, and CD3-HLADR-</p>

the measurement of the quantitative level of the combination of endometrial leukocyte markers CD3+, CD13+CD20 being associated with length of menstruation (> 7 days) and pelvic pain at any time other than menstruation and intercourse ; the measurement of the quantitative level of the combination of blood leukocyte markers CD14+, CD57+, CD3+HLADR-, CD3-CD45RA- being associated with pelvic pain at any time other than menstruation and intercourse, and length of menstruation (>7 days); and the measurement of the quantitative level of the combination of blood leukocyte markers CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA- being associated with pelvic pain at any time other than menstruation and intercourse, and age > 40,

b) establishing a cutoff value for the proportion of each leukocyte marker in the combination,

c) comparing the proportion obtained in step (b) for each leukocyte marker to a predetermined cutoff value wherein a positive result gives a score of 1 whereas a negative result gives a score of 0,

d) obtaining a diagnostic value by adding the scores of all the markers of the combination and converting it in percentage,

e) comparing the final diagnostic value to an established threshold value wherein a positive diagnosis of endometriosis is given when the final diagnostic value exceeds the threshold value established for the combination of leukocyte markers whereas a negative diagnosis of endometriosis is given when the final diagnostic value is lower than the threshold value established for the combination of leukocyte markers.

2. The method of claim 1, wherein said proportion of leukocytes marker is determined by using at least one antibody specific for said at least one specific population of leukocytes.

3. A method for determining endometriosis in a sample of eutopic uterine endometrial tissues or a sample of blood of a female subject, comprising:

a) measuring in said sample a quantitative level of at least two different surface antigens from blood leukocytes and/or eutopic endometrial leukocytes, said surface antigens from blood leukocytes and/or eutopic endometrial leukocytes being selected from the leukocyte marker combinations defined in the following Table:

Leukocyte Marker Combinations
<p>Endometrial leukocyte markers CD4+, CD8+CD69-, and CD13+CD45RO- CD4+, CD8+CD69-, CD56+CD122-, CD3+CD45RA-, and CD13+CD45RO-,</p>

(continued)

<p>5 10 15 20 25 30 35</p>	<p>Endometrial leukocyte markers CD4+, CD8+CD69-, CD13-CD122+, and CD13+CD45RO- CD4+ CD8+CD69-, and D14+CD13-CD16b- CD3+CD16-, CD13+CD45RO-, CD3+, CD8+, and CD3+CD69+ CD3+, CD3+CD8+, CD13+CD45RO-, and CD3-CD20- CD3+CD8+, CD13+CD45RO-, CD3+CD5+, CD3+CD122-, CD3-CD20-, and CD3+CD45RO- CD3+CD8+, CD13+CD45RO-, CD3+CD5+, and CD3+CD122- CD3+CD20-CD5-, CD4+CD13-, and CD56-CD122- CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, and CD3+CD45RO- CD3+, CD8+CD69-, CD3-CD4-CD45RO+, ratio CD13+/CD3+, and CD13+CD45RO- CD3+CD8+, CD13+CD45RO-, CD3-CD5-, and CD20-CD5+ CD8+, CD5+, CD3-CD20-, and CD3-HLADR- CD3+CDB+, CD13+CD45RO-, and CD5+ CD4+, CD13-CD122+, CD8+CD69-, and CD3+CD45RA- CD4+, CD8+CD69-, CD13-CD122+, CD3+CD45RA-, and CD13+CD45RO- CD4+, CD8+CD69-, CD56-CD122-, CD3+CD45RA-, CD13+CD45RO- CD3+, CD4+, CD3+CD8+, CD13+CD45RO-, and CD3+CD5+ CD3+, CD4+, CD3+CD8+, and CD13+CD45RO- CD3+, CD3+CD8+, and CD13+CD45RO- CD3+, CD5+, CD3+CD5+, CD69+, and CD4-CD69+ CD3+, CD3+CD8+, and CD13+CD45RO- CD3+, CD3+CD8+, and CD13+CD45RO- CD3+, CD3+CD8+, CD13+CD45RO-, CD3-CD20-, and CD56-CD16- CD3+CD16-, CD3-CD4-CD45RO+, CD3+, CD8+, CD3+CD69+, and CD13+CD45RO- CD3+, CD3-CD45RO+, ratio CD13+/CD3+, CD3+CD8+, and CD8+CD69- CD3+CD20-CD5-, CD4+CD13-, CD56-CD122-, and CD4+CD45RO- CD3+, CD3+CD8+, CD13+CD45RO-, and CD3-CD20- CD13+CD45RO-, CD4+CD45RA-, CD3+CD122-, and CD8+CD69- CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, CD3+CD45RO-, and CD3-CD5- CD3+, CD3-CD5-, CD13+CD45RO-, and CD3-CD20- CD3+, CD3-CD5-, and CD13+CD45RO- CD3+CD8+, CD13+CD45RO-, and CD3+CD5+ CD3+ and CD13+CD20-</p>
<p>40 45</p>	<p>Blood Leukocyte markers CD3-CD5+, CD3-CD45RA-, CD3-CD44+, CD13+, and CD3-CD57-CD44- CD3-CD45RA-, CD20-CD44+, CD20-HLADR+, CD3-CD4-CD44+, and CD36-HLADR+ CD3-CD45RA-, CD3-CD45RO+, and CD20-HLADR+ CD14+CD44+, CD57+, CD3-CD45RA, and CD14+ CD14+, CD57+, CD3+CD69+, CD3+HLADR+, and CD3-CD45RA- CD14+, CD57+, CD3+HLADR-, CD3-CD45RA- CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-</p>
<p>50 55</p>	<p>Blood (in italics) and endometrial leukocyte markers <i>CD57+, CD14+, CD3-CD69+, CD3+, CD4+, and CD3+CD8+HLADR-</i> <i>CD3-CD69+, CD3+, CD4+, and CD3+CD8+HLADR-</i> <i>CD4-CD36+, CD3-CD69+, CD8+, CD13+, CD3+, CD16+, CD69+, and CD5+</i> <i>CD3-CD45RA-, CD3-CD45RO+, CD20-HLADR+, CD14+CD44+, CD8+, CD5+, CD3-CD20-, and CD3-HLADR-</i></p> <p>the measurement of the quantitative level of the combination of endometrial leukocyte markers CD3+, CD13+CD20 being associated with length of menstruation (> 7 days) and pelvic pain at any time other than menstruation and intercourse ; the measurement of the quantitative level of the combination of blood leukocyte</p>

EP 1 147 421 B1

markers CD14+, CD57+, CD3+HLADR-, CD3-CD45RA-being associated with pelvic pain at any time other than menstruation and intercourse, and length of menstruation (>7 days); and the measurement of the quantitative level of the combination of blood leukocyte markers CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA- being associated with pelvic pain at any time other than menstruation and intercourse, and age > 40, establishing a predictive model for endometriosis by including each marker of said combination in the following logistic regression equation :

$$P(r) = \frac{e^{c+B1*(marker1)+B2*(marker2)+...Bn*(marker n)}}{1 + e^{c+B1*(marker1)+B2*(marker2)+...Bn*(marker n)}}$$

Where:

- P (r) = probability of having endometriosis;
- c = constant established for a particular combination;
- B = coefficient of regression;
- n = total number of markers in the combination,

comparing the probability of having endometriosis (P (r)) to a threshold value wherein a positive diagnosis of endometriosis is given when the P (r) value exceeds the threshold value established for said combination of markers whereas a negative diagnosis of endometriosis is given when the P (r) value is lower than the threshold value established for said combination of markers.

4. The method of claim 3, wherein said quantitative level is measured by using at least one antibody directed against said specific populations of leukocytes.

Patentansprüche

1. Verfahren zur Feststellung von Endometriose in einer Probe von eutopischen uterinen endometrialen Geweben oder einer Blutprobe eines weiblichen Subjekts, umfassend:

a) Messen eines quantitativen Levels von wenigstens zwei verschiedenen Oberflächenantigenen aus Blutleukozyten und/oder eutopischen endometrialen Leukozyten in der Probe, wobei die Oberflächenantigene aus Blutleukozyten und/oder eutopischen endometrialen Leukozyten aus den in der nachstehenden Tabelle definierten Leukozytenmarkerkombinationen ausgewählt werden:

Leukozytenmarkerkombinationen
<p>Endometriale leukozytenmarker</p> <p>CD4+, CD8+CD69-, und CD13+CD45RO-</p> <p>CD4+, CD8+CD69-, CD56+CD122-, CD3+CD45RA-, und CD13+CD45RO-,</p> <p>CD4+, CD8+CD69-, CD13-CD122+, und CD13+CD45RO-</p> <p>CD4+ CD8+CD69-, und D14+CD13-CD16b-</p> <p>CD3+CD16-, CD13+CD45RO-, CD3+, CD8+, und CD3+CD69+</p> <p>CD3+, CD3+CD8+, CD13+CD45RO-, und CD3-CD20-</p> <p>CD3+CD8+, CD13+CD45RO-, CD3+CD5+, CD3+CD122-, CD3-CD20-, und CD3+CD45RO-</p> <p>CD3+CD8+, CD13+CD45RO-, CD3+CD5+, und CD3+CD122-</p> <p>CD3+CD20-CD5-, CD4+CD13-, und CD56-CD122-</p> <p>CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, und CD3+CD45RO-</p> <p>CD3+, CD8+CD69-, CD3-CD4-CD45RO+, Verhältnis CD13+/CD3+, und CD13+CD45RO-</p> <p>CD3+CD8+, CD13+CD45RO-, CD3-CD5-, und CD20-CD5+</p> <p>CD8+, CD5+, CD3-CD20-, und CD3-HLADR-</p> <p>CD3+CD8+, CD13+CD45RO-, and CD5+</p>

(fortgesetzt)

Endometriale Leukozytenmarker

CD4+, CD13-CD122+, CD8+CD69-, und CD3+CD45RA-
 CD4+, CD8+CD69-, CD13-CD122+, CD3+CD45RA-, und CD13+CD45RO-
 CD4+, CD8+CD69-, CD56-CD122-, CD3+CD45RA-, CD13+CD45RO-
 CD3+, CD4+, CD3+CD8+, CD13+CD45RO-, und CD3+CD5+
 CD3+, CD4+, CD3+CD8+, und CD13+CD45RO-
 CD3+, CD3+CD8+, und CD13+CD45RO-
 CD3+, CD5+, CD3+CD5+, CD69+, und CD4-CD69+
 CD3+, CD3+CD8+, und CD13+CD45RO-
 CD3+, CD3+CD8+, und CD13+CD45RO-
 CD3+, CD3+CD8+, CD13+CD45RO-, CD3-CD20-, und CD56-CD16-
 CD3+CD16-, CD3-CD4-CD45RO+, CD3+, CD8+, CD3+CD69+, und CD13+CD45RO-
 CD3+, CD3-CD45RO+, Verhältnis CD13+/CD3+, CD3+CD8+, und CD8+CD69-
 CD3+CD20-CD5-, CD4+CD13-, CD56-CD122-, und CD4+CD45RO-
 CD3+, CD3+CD8+, CD13+CD45RO-, und CD3-CD20-
 CD13+CD45RO-, CD4+CD45RA-, CD3+CD122-, und CD8+CD69-
 CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, CD3+CD45RO-, und CD3-CD5-
 CD3+, CD3-CD5-, CD13+CD45RO-, und CD3-CD20-
 CD3+, CD3-CD5-, und CD13+CD45RO-
 CD3+CD8+, CD13+CD45RO-, und CD3+CD5+
 CD3+ und CD13+CD20.

Blutleukozytenmarker

CD3-CD5+, CD3-CD45RA-, CD3-CD44+, CD13+, und CD3-CD57-CD44-
 CD3-CD45RA-, CD20-CD44+, CD20-HLADR+, CD3-CD4-CD44+, und CD36-HLADR+
 CD3-CD45RA-, CD3-CD45RO+, und CD20-HLADR+
 CD14+CD44+, CD57+, CD3-CD45RA, und CD14+
 CD14+, CD57+, CD3+CD69+, CD3+HLADR+, and CD3-CD45RA-
 CD14+, CD57+, CD3+HLADR- und CD3-CD45RA-
 CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-

Blut- (in Kursivschrift) und endometriale Leukozytenmarker

CD57+, CD14+, CD3-CD69+, CD3+, CD4+, und CD3+CD8+HLADR-
CD3-CD69+, CD3+, CD4+, und CD3+CD8+HLADR-
CD4-CD36+, *CD3-CD69+*, CD8+, CD13+, CD3+, CD16+, CD69+, und CD5+
CD3-CD45RA-, *CD3-CD45RO+*, *CD20-HLADR+*, *CD14+CD44+*, CD8+, CD5+, CD3-CD20-, und CD3-HLADR-

Wobei die Messung des quantitativen Levels der Kombination der endometriale Leukozytenmarker CD3+, CD13+CD20 mit der Länge der Menstruation (> 7 Tage) und Beckenschmerzen zu jeder beliebigen Zeit, außer der Menstruation und dem Geschlechtsverkehr, assoziiert ist; die Messung des quantitativen Levels der Kombination der Blutleukozytenmarker CD14+, CD57+, CD3+HLADR-, CD3-CD45RA- mit Beckenschmerzen zu jeder beliebigen Zeit, außer der Menstruation und dem Geschlechtsverkehr, und der Länge der Menstruation (> 7 Tage) assoziiert ist; und die Messung des quantitativen Levels der Kombination der Blutleukozytenmarker CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA- mit Beckenschmerzen zu jeder beliebigen Zeit, außer der Menstruation und dem Geschlechtsverkehr, und Alter > 40 assoziiert ist,

b) Etablieren eines Grenzwertes für den Anteil jedes Leukozytenmarkers in der Kombination,
 c) Vergleichen des in Stufe (b) für jeden Leukozytenmarker erhaltenen Anteils mit einem vorher festgelegten Grenzwert, wobei ein positives Ergebnis eine Bewertung von 1 ergibt, während ein negatives Ergebnis eine Bewertung von 0 ergibt,
 d) Erhalten eines diagnostischen Werts durch Addieren der Bewertungen aller Marker der Kombination und Umwandeln in eine Prozentzahl,
 e) Vergleichen des diagnostischen Endwertes mit einem etablierten Schwellenwert, wobei eine positive Diagnose der Endometriose vorliegt, wenn der diagnostische Endwert den für die Kombination von Leukozytenmarkern etablierten Schwellenwert überschreitet, während eine negative Diagnose der Endometriose vorliegt,

EP 1 147 421 B1

wenn der diagnostische Endwert niedriger als der für die Kombination von Leukozytenmarkern etablierte Schwellenwert ist.

2. Verfahren nach Anspruch 1, wobei der Anteil an Leukozytenmarker bestimmt wird, indem wenigstens ein Antikörper, der für die wenigstens eine spezifische Population von Leukozyten spezifisch ist, verwendet wird.

3. Verfahren zur Feststellung von Endometriose in einer Probe von eutopischen uterinen endometrialen Geweben oder einer Blutprobe eines weiblichen Subjekts, umfassend:

a) Messen eines quantitativen Levels von wenigstens zwei verschiedenen Oberflächenantigenen aus Blutleukozyten und/oder eutopischen endometrialen Leukozyten in der Probe, wobei die Oberflächenantigene aus Blutleukozyten und/oder eutopischen endometrialen Leukozyten aus den in der nachstehenden Tabelle definierten Leukozytenmarkerkombinationen ausgewählt werden:

Leukozytenmarkerkombinationen	
	Endometriale leukozytenmarker
	CD4+, CD8+CD69-, und CD13+CD45RO-
20	CD4+, CD8+CD69-, CD56+CD122-, CD3+CD45RA-, und CD13+CD45RO-, CD4+, CD8+CD69-, CD13-CD122+, und CD13+CD45RO-
	CD4+ CD8+CD69-, und D14+CD13-CD16b-
	CD3+CD16-, CD13+CD45RO-, CD3+, CD8+, und CD3+CD69+
25	CD3+, CD3+CD8+, CD13+CD45RO-, und CD3-CD20-
	CD3+CD8+, CD13+CD45RO-, CD3+CD5+, CD3+CD122-, CD3-CD20-, und CD3+CD45RO-
	CD3+CD8+, CD13+CD45RO-, CD3+CD5+, und CD3+CD122-
	CD3+CD20-CD5-, CD4+CD13-, und CD56-CD122-
	CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, und CD3+CD45RO-
30	CD3+, CD8+CD69-, CD3-CD4-CD45RO+, Verhältnis CD13+/CD3+, und CD13+CD45RO-
	CD3+CD8+, CD13+CD45RO-, CD3-CD5-, und CD20-CD5+
	CD8+, CD5+, CD3-CD20-, und CD3-HLADR-
	CD3+CD8+, CD13+CD45RO-, und CD5+
35	CD4+, CD13-CD122+, CD8+CD69-, und CD3+CD45RA-
	CD4+, CD8+CD69-, CD13-CD122+, CD3+CD45RA-, und CD13+CD45RO-
	CD4+, CD8+CD69-, CD56-CD122-, CD3+CD45RA-, CD13+CD45RO-
	CD3+, CD4+, CD3+CD8+, CD13+CD45RO-, und CD3+CD5+
	CD3+, CD4+, CD3+CD8+, und CD13+CD45RO-
40	CD3+, CD3+CD8+, und CD13+CD45RO-
	CD3+, CD5+, CD3+CD5+, CD69+, und CD4-CD69+
	CD3+, CD3+CD8+, und CD13+CD45RO-
	CD3+, CD3+CD8+, und CD13+CD45RO-
45	CD3+, CD3+CD8+, CD13+CD45RO-, CD3-CD20-, und CD56-CD16-
	CD3+CD16-, CD3-CD4-CD45RO+, CD3+, CD8+, CD3+CD69+, und CD13+CD45RO-
	CD3+, CD3-CD45RO+, Verhältnis CD13+/CD3+, CD3+CD8+, und CD8+CD69-
	CD3+CD20-CD5-, CD4+CD13-, CD56-CD122-, und CD4+CD45RO-
	CD3+, CD3+CD8+, CD13+CD45RO-, und CD3-CD20-
50	CD13+CD45RO-, CD4+CD45RA-, CD3+CD122-, und CD8+CD69-
	CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, CD3+CD45RO-, und CD3-CD5-
	CD3+, CD3-CD5-, CD13+CD45RO-, und CD3-CD20-
	CD3+, CD3-CD5-, und CD13+CD45RO-
55	CD3+CD8+, CD13+CD45RO-, und CD3+CD5+
	CD3+ und CD13+CD20
Blutleukozytenmarker	
CD3-CD5+, CD3-CD45RA-, CD3-CD44+, CD13+, und CD3-CD57-CD44-	

(fortgesetzt)

Blutleukozytenmarker

CD3-CD45RA-, CD20-CD44+, CD20-HLADR+, CD3-CD4-CD44+, und CD36-HLADR+
 CD3-CD45RA-, CD3-CD45RO+, und CD20-HLADR+
 CD14+CD44+, CD57+, CD3-CD45RA, und CD14+
 CD14+, CD57+, CD3+CD69+, CD3+HLADR+, und CD3-CD45RA-
 CD14+, CD57+, CD3+HLADR-, CD3-CD45RA-
 CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-

Blut- (in Kursivschrift) und endometriale Leukozytenmarker

CD57+, *CD14+*, *CD3-CD69+*, CD3+, CD4+, und CD3+CD8+HLADR-
CD3-CD69+, *CD3+*, CD4+, und CD3+CD8+HLADR-
 CD4-CD36+, *CD3-CD69+*, CD8+, CD13+, CD3+, CD16+, CD69+, und CD5+
CD3-CD45RA-, *CD3-CD45RO+*, *CD20-HLADR+*, *CD14+CD44+*, CD8+, CD5+, CD3-CD20-, und CD3-HLADR-

wobei die Messung des quantitativen Levels der Kombination der endometrialen Leukozytenmarker CD3+, CD13+CD20 mit der Länge der Menstruation (> 7 Tage) und Beckenschmerzen zu jeder beliebigen Zeit, außer der Menstruation und dem Geschlechtsverkehr, assoziiert ist; die Messung des quantitativen Levels der Kombination der Blutleukozytenmarker CD14+, CD57+, CD3+HLADR-, CD3-CD45RA- mit Beckenschmerzen zu jeder beliebigen Zeit, außer der Menstruation und dem Geschlechtsverkehr, und der Länge der Menstruation (> 7 Tage) assoziiert ist; und die Messung des quantitativen Levels der Kombination der Blutleukozytenmarker CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA- mit Beckenschmerzen zu jeder beliebigen Zeit, außer der Menstruation und dem Geschlechtsverkehr, und Alter > 40 assoziiert ist, Etablieren eines Vorhersagemodells für Endometriose durch Einschließen jedes Markers der Kombination in die nachstehende logistische Regressionsgleichung:

$$P(r) = \frac{e^{c+B1*(\text{Marker } 1)+B2*(\text{Marker } 2)+\dots+Bn*(\text{Marker } n)}}{1 + e^{c+B1*(\text{Marker } 1)+B2*(\text{Marker } 2)+\dots+Bn*(\text{Marker } n)}}$$

worin:

- P (r) = Wahrscheinlichkeit dafür, Endometriose zu haben;
- c = Konstante, etabliert für eine bestimmte Kombination;
- B = Regressionskoeffizient;
- n = Gesamtzahl der Marker in der Kombination,

Vergleichen der Wahrscheinlichkeit dafür, Endometriose zu haben (P (r)) mit einem Schwellenwert, wobei eine positive Diagnose der Endometriose vorliegt, wenn der P (r)-Wert den für die Kombination der Marker etablierten Schwellenwert überschreitet, während eine negative Diagnose der Endometriose vorliegt, wenn der P (r)-Wert niedriger ist, als der für die Kombination der Marker etablierte Schwellenwert.

4. Verfahren nach Anspruch 3, wobei der quantitative Level gemessen wird, indem wenigstens ein Antikörper, der gegen die spezifischen Populationen von Leukozyten gerichtet ist, verwendet wird.

Revendications

1. Procédé permettant de déterminer l'endométriose dans un échantillon de tissus de l'endomètre utérin eutopique ou dans un échantillon de sang d'un sujet féminin, comprenant les étapes consistant à :

a) mesurer dans ledit échantillon un niveau quantitatif d'au moins deux différents antigènes de surface provenant de leucocytes du sang et/ou de leucocytes de l'endomètre eutopique, lesdits antigènes de surface provenant des leucocytes du sang et/ou des leucocytes de l'endomètre eutopique étant choisis à partir des combinaisons

EP 1 147 421 B1

de marqueurs leucocytaires définies dans le tableau suivant :

<u>Combinaisons de marqueurs leucocytaires</u>	
5	<p>Marqueurs leucocytaires de l'endomètre</p> <p>CD4+, CD8+CD69-, et CD13+CD45RO-</p>
10	<p>CD4+, CD8+CD69-, CD56+CD122-, CD3+CD45RA-, et CD13+CD45RO-, CD4+, CD8+CD69-, CD13-CD122+, et CD13+CD45RO- CD4+ CD8+CD69-, et D14+CD13-CD16b-</p>
15	<p>CD3+CD16-, CD13+CD45RO-, CD3+, CD8+, et CD3+CD69+ CD3+, CD3+CD8+, CD13+CD45RO-, et CD3-CD20- CD3+CD8+, CD13+CD45RO-, CD3+CD5+, CD3+CD122-, CD3-CD20-, et CD3+CD45RO- CD3+CD8+, CD13+CD45RO-, CD3+CD5+, et CD3+CD122- CD3+CD20-CD5-, CD4+CD13-, et CD56-CD122-</p>
20	<p>CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, et CD3+CD45RO- CD3+, CD8+CD69-, CD3-CD4-CD45RO+, rapport CD13+/CD3+, et CD13+CD45RO- CD3+CD8+, CD13+CD45RO-, CD3-CD5-, et CD20-CD5+ CD8+, CD5+, CD3-CD20-, et CD3-HLADR-</p>
25	<p>CD3+CD8+, CD13+CD45RO-, et CD5+ CD4+, CD13-CD122+, CD8+CD69-, et CD3+CD45RA- CD4+, CD8+CD69-, CD13-CD122+, CD3+CD45RA-, et CD13+CD45RO- CD4+, CD8+CD69-, CD56-CD122-, CD3+CD45RA-, CD13+CD45RO- CD3+, CD4+, CD3+CD8+, CD13+CD45RO-, et CD3+CD5+</p>
30	<p>CD3+, CD4+, CD3+CD8+, et CD13+CD45RO- CD3+, CD3+CD8+, et CD13+CD45RO- CD3+, CD5+, CD3+CD5+, CD69+, et CD4-CD69+ CD3+, CD3+CD8+, et CD13+CD45RO- CD3+, CD3+CD8+, et CD13+CD45RO-</p>
35	<p>CD3+, CD3+CD8+, CD13+CD45RO-, CD3-CD20-, et CD56-CD16- CD3+CD16-, CD3-CD4-CD45RO+, CD3+, CD8+, CD3+CD69+, et CD13+CD45RO- CD3+, CD3-CD45RO+, rapport CD13+/CD3+, CD3+CD8+, et CD8+CD69- CD3+CD20-CD5-, CD4+CD13-, CD56-CD122-, et CD4+CD45RO- CD3+, CD3+CD8+, CD13+CD45RO-, et CD3-CD20- CD13+CD45RO-, CD4+CD45RA-, CD3+CD122-, et CD8+CD69- CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, CD3+CD45RO-, et CD3-CD5- CD3+, CD3-CD5-, CD13+CD45RO-, et CD3-CD20-</p>
40	<p>CD3+, CD3-CD5-, et CD13+CD45RO- CD3+CD8+, CD13+CD45RO-, et CD3+CD5+ CD3+ et CD13+CD20.</p>
45	<p>Marqueurs leucocytaires sanguins</p> <p>CD3-CD5+, CD3-CD45RA-, CD3-CD44+, CD13+, et CD3-CD57-CD44- CD3-CD45RA-, CD20-CD44+, CD20-HLADR+, CD3-CD4-CD44+, et CD36-HLADR+ CD3-CD45RA-, CD3-CD45RO+, et CD20-HLADR+ CD14+CD44+, CD57+, CD3-CD45RA, et CD14+ CD14+, CD57+, CD3+CD69+, CD3+HLADR+, et CD3-CD45RA-</p>
50	<p>CD14+, CD57+, CD3+HLADR- et CD3-CD45RA- CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-</p>
55	<p>Marqueurs leucocytaires du sang (en italiques) et de l'endomètre</p> <p><i>CD57+</i>, <i>CD14+</i>, <i>CD3-CD69+</i>, CD3+, CD4+, et CD3+CD8+HLADR- <i>CD3-CD69+</i>, CD3+, CD4+, et CD3+CD8+HLADR- <i>CD4-CD36+</i>, <i>CD3-CD69+</i>, CD8+, CD13+, CD3+, CD16+, CD69+, et CD5+ <i>CD3-CD45RA-</i>, <i>CD3-CD45RO+</i>, <i>CD20-HLADR+</i>, <i>CD14+CD44+</i>, CD8+, CD5+, CD3-CD20-, et CD3-HLADR-</p>

EP 1 147 421 B1

la mesure du niveau quantitatif de la combinaison de marqueurs leucocytaires de l'endomètre CD3+, CD13+CD20 étant associée à la durée de la menstruation (> 7 jours) et à une douleur pelvienne à tout moment autre que pendant la menstruation et le rapport sexuel ; la mesure du niveau quantitatif de la combinaison de marqueurs leucocytaires du sang CD14+, CD57+, CD3+HLADR-, CD3-CD45RA-, étant associée à une douleur pelvienne à tout moment autre que pendant la menstruation et le rapport sexuel, et à la durée de la menstruation (> 7 jours) ; et la mesure du niveau quantitatif de la combinaison de marqueurs leucocytaires du sang CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-, étant associée à une douleur pelvienne à tout moment autre que pendant la menstruation et le rapport sexuel, et à un âge > 40 ans,

b) établir une valeur limite pour la proportion de chaque marqueur leucocytaire dans la combinaison,

c) comparer la proportion obtenue à l'étape (b) pour chaque marqueur leucocytaire à une valeur limite prédéterminée, dans lequel un résultat positif donne une note de 1 tandis qu'un résultat négatif donne une note de 0, d) obtenir une valeur diagnostique par addition des notes de tous les marqueurs de la combinaison et conversion de celle-ci en pourcentage,

e) comparer la valeur diagnostique finale à une valeur seuil établie, dans lequel un diagnostic positif d'endométriose est établi lorsque la valeur diagnostique finale dépasse la valeur seuil établie pour la combinaison de marqueurs leucocytaires tandis qu'un diagnostic négatif d'endométriose est établi lorsque la valeur diagnostique finale est inférieure à la valeur seuil établie pour la combinaison de marqueurs leucocytaires.

2. Procédé selon la revendication 1, dans lequel ladite proportion de marqueurs leucocytaires est déterminée en utilisant au moins un anticorps spécifique de ladite au moins une population spécifique de leucocytes.

3. Procédé permettant de déterminer l'endométriose dans un échantillon de tissus de l'endomètre utérin eutopique ou dans un échantillon de sang d'un sujet féminin, comprenant les étapes consistant à :

a) mesurer dans ledit échantillon un niveau quantitatif d'au moins deux différents antigènes de surface provenant de leucocytes du sang et/ou de leucocytes de l'endomètre eutopique, lesdits antigènes de surface provenant des leucocytes du sang et/ou des leucocytes de l'endomètre eutopique étant choisis à partir des combinaisons de marqueurs leucocytaires définies dans le tableau suivant :

Combinaisons de marqueurs leucocytaires	
Marqueurs leucocytaires de l'endomètre	
	CD4+, CD8+CD69-, et CD13+CD45RO-
	CD4+, CD8+CD69-, CD56+CD122-, CD3+CD45RA-, et CD13+CD45RO-,
	CD4+, CD8+CD69-, CD13-CD122+, et CD13+CD45RO-
	CD4+ CD8+CD69-, et D14+CD13-CD16b-
	CD3+CD16-, CD13+CD45RO-, CD3+, CD8+, et CD3+CD69+
	CD3+, CD3+CD8+, CD13+CD45RO-, et CD3-CD20-
	CD3+CD8+, CD13+CD45RO-, CD3+CD5+, CD3+CD122-, CD3-CD20-, et CD3+CD45RO-
	CD3+CD8+, CD13+CD45RO-, CD3+CD5+, et CD3+CD122-
	CD3+CD20-CD5-, CD4+CD13-, et CD56-CD122-
	CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, et CD3+CD45RO-
	CD3+, CD8+CD69-, CD3-CD4-CD45RO+, rapport CD13+/CD3+, et CD13+CD45RO-
	CD3+CD8+, CD13+CD45RO-, CD3-CD5-, et CD20-CD5+
	CD8+, CD5+, CD3-CD20-, et CD3-HLADR-
	CD3+CD8+, CD13+CD45RO-, et CD5+
	CD4+, CD13-CD122+, CD8+CD69-, et CD3+CD45RA-
	CD4+, CD8+CD69-, CD13-CD122+, CD3+CD45RA-, et CD13+CD45RO-
	CD4+, CD8+CD69-, CD56-CD122-, CD3+CD45RA-, CD13+CD45RO-
	CD3+, CD4+, CD3+CD8+, CD13+CD45RO-, et CD3+CD5+
	CD3+, CD4+, CD3+CD8+, et CD13+CD45RO-
	CD3+, CD3+CD8+, et CD13+CD45RO-
	CD3+, CD5+, CD3+CD5+, CD69+, et CD4-CD69+
	CD3+, CD3+CD8+, et CD13+CD45RO-
	CD3+, CD3+CD8+, et CD13+CD45RO-

(suite)

Marqueurs leucocytaires de l'endomètre

CD3+, CD3+CD8+, CD13+CD45RO-, CD3-CD20-, et CD56-CD16-
 CD3+CD16-, CD3-CD4-CD45RO+, CD3+, CD8+, CD3+CD69+, et CD13+CD45RO-
 CD3+, CD3-CD45RO+, rapport CD13+/CD3+, CD3+CD8+, et CD8+CD69-
 CD3+CD20-CD5-, CD4+CD13-, CD56-CD122-, et CD4+CD45RO-
 CD3+, CD3+CD8+, CD13+CD45RO-, et CD3-CD20-
 CD13+CD45RO-, CD4+CD45RA-, CD3+CD122-, et CD8+CD69-
 CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, CD3+CD45RO-, et CD3-CD5-
 CD3+, CD3-CD5-, CD13+CD45RO-, et CD3-CD20-
 CD3+, CD3-CD5-, et CD13+CD45RO-
 CD3+CD8+, CD13+CD45RO-, et CD3+CD5+
 CD3+ et CD13+CD20

Marqueurs leucocytaires sanguins

CD3-CD5+, CD3-CD45RA-, CD3-CD44+, CD13+, et CD3-CD57-CD44-
 CD3-CD45RA-, CD20-CD44+, CD20-HLADR+, CD3-CD4-CD44+, et CD36-HLADR+
 CD3-CD45RA-, CD3-CD45RO+, et CD20-HLADR+
 CD14+CD44+, CD57+, CD3-CD45RA, et CD14+
 CD14+, CD57+, CD3+CD69+, CD3+HLADR+, et CD3-CD45RA-
 CD14+, CD57+, CD3+HLADR-, CD3-CD45RA-
 CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-

Marqueurs leucocytaires du sang (en italiques) et de l'endomètre

CD57+, *CD14+*, *CD3-CD69+*, CD3+, CD4+, et CD3+CD8+HLADR-
CD3-CD69+, CD3+, CD4+, et CD3+CD8+HLADR-
 CD4-CD36+, *CD3-CD69+*, CD8+, CD13+, CD3+, CD16+, CD69+, et CD5+
 CD3-CD45RA-, *CD3-CD45RO+*, CD20-HLADR+, CD14+CD44+, CD8+, CD5+, CD3-CD20-, et CD3-HLADR-

la mesure du niveau quantitatif de la combinaison de marqueurs leucocytaires de l'endomètre CD3+, CD13+CD20 étant associée à la durée de la menstruation (> 7 jours) et à une douleur pelvienne à tout moment autre que pendant la menstruation et le rapport sexuel ; la mesure du niveau quantitatif de la combinaison de marqueurs leucocytaires du sang CD14+, CD57+, CD3+HLADR-, CD3-CD45RA-, étant associée à une douleur pelvienne à tout moment autre que pendant la menstruation et le rapport sexuel, et à la durée de la menstruation (> 7 jours); et la mesure du niveau quantitatif de la combinaison de marqueurs leucocytaires du sang CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-, étant associée à une douleur pelvienne à tout moment autre que pendant la menstruation et le rapport sexuel, et à un âge > 40 ans, établir un modèle de prévision pour l'endométriose en incluant chaque marqueur de ladite combinaison dans l'équation de régression logistique suivante :

$$P(r) = \frac{e^{c+B1*(marqueur\ 1)+B2*(marqueur\ 2)+...Bn*(marqueur\ n)}}{1 + e^{c+B1*(marqueur\ 1)+B2*(marqueur\ 2)+...Bn*(marqueur\ n)}}$$

où :

P (r)= la probabilité d'avoir l'endométriose ;
 c = la constante établie pour une combinaison particulière ;
 B = le coefficient de régression;
 n = le nombre total de marqueurs dans la combinaison,

comparer la probabilité d'avoir l'endométriose (P(r)) à une valeur seuil, dans lequel un diagnostic positif d'endométriose est établi lorsque la valeur P(r) dépasse la valeur seuil établie pour ladite combinaison de marqueurs, tandis

EP 1 147 421 B1

qu'un diagnostic négatif d'endométriose est établi lorsque la valeur $P(r)$ est inférieure à la valeur seuil établie pour ladite combinaison de marqueurs.

4. Procédé selon la revendication 3, dans lequel ledit niveau quantitatif est mesuré en utilisant au moins un anticorps dirigé contre lesdites populations spécifiques de leucocytes.

5

10

15

20

25

30

35


40

45

50

55

PREDICTIVE ALGORITHM FOR THE DIAGNOSIS OF ENDOMETRIOSIS



¹ Proportion of cells expressing a specific marker, or a given subset defined by markers within the leucocyte population (CD45+) in the peripheral blood or the stromal fraction of the endometrium.
² A positive test result is given when the proportion of a leucocyte subset fulfills the condition established by the cutoff point.

Fig. 1

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 9428021 A [0007]
- WO 9218535 A [0007]
- US 5478725 A [0008]

Non-patent literature cited in the description

- **OOSTERLYNCCK D.J. et al.** *Am J reprod. Immunol.*, 1994, vol. 31, 25-31 [0009] [0009]
- **BECKER J.L. et al.** *Am J Reprod. Immunol.*, 1995, vol. 34, 179-187 [0009]
- **WU M.Y. et al.** *Am. j. Reprod. Immunol.*, 1996, vol. 35, 510-516 [0009]
- **HO H.N. et al.** *Hum Reprod.*, 1997, vol. 97, 2528-2533 [0009]
- **OTA H. et al.** *Am J Reprod. Immunol.*, 1996, vol. 35, 477-482 [0010]
- **KLENTZERIS L.D. et al.** *Eur. J Obstet gynecol Reprod Biol.*, 1995, vol. 63, 41-47 [0010]
- **JONES R.K. et al.** *Fertil Steril*, 1996, vol. 66, 81-89 [0010]

专利名称(译)	诊断子宫内膜异位症的方法		
公开(公告)号	EP1147421B1	公开(公告)日	2007-09-26
申请号	EP2000901009	申请日	2000-01-24
[标]申请(专利权)人(译)	procrea能够		
申请(专利权)人(译)	PROCREA生物科学公司.		
当前申请(专利权)人(译)	西门子医疗系统集团诊断		
[标]发明人	GAGNE DANIELE GOSELIN DIANE HUGO PATRICE MIRON PIERRE		
发明人	GAGNE, DANIELE GOSELIN, DIANE HUGO, PATRICE MIRON, PIERRE		
IPC分类号	G01N33/574 G01N33/68 G01N33/53		
CPC分类号	G01N33/57442 G01N2333/705 G01N2800/364		
优先权	60/117031 1999-01-25 US		
其他公开文献	EP1147421A1		
外部链接	Espacenet		

摘要(译)

本发明涉及使用血液和子宫内膜白细胞标记物或其组合诊断子宫内膜异位症的方法和试剂盒。标记物是来自子宫内膜或血液白细胞的表面抗原。

$$P(r) = \frac{e^{c + B1^*(marker1) + B2^*(marker2) + \dots + Bn^*(marker n)}}{1 + e^{c + B1^*(marker1) + B2^*(marker2) + \dots + Bn^*(marker n)}}$$

$$1 + e^{c + B1^*(marker1) + B2^*(marker2) + \dots + Bn^*(marker n)}$$