



(11) **EP 1 745 287 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
22.09.2010 Bulletin 2010/38

(21) Application number: **05747945.3**

(22) Date of filing: **11.05.2005**

(51) Int Cl.:
G01N 33/53 (2006.01)

(86) International application number:
PCT/US2005/016436

(87) International publication number:
WO 2005/111612 (24.11.2005 Gazette 2005/47)

(54) **DIAGNOSING AND MONITORING INFLAMMATORY DISEASES BY MEASURING COMPLEMENT COMPONENTS ON WHITE BLOOD CELLS**

DIAGNOSE UND ÜBERWACHUNG VON ENTZÜNDLICHEN KRANKHEITEN MITTELS MESSUNG DER ZUSÄTZLICHEN KOMPONENTEN VON LEUKOZYTEN

DIAGNOSTIC ET SUIVI DE MALADIES INFLAMMATOIRES PAR LA MESURE DE COMPOSANTS DU COMPLEMENT SUR DES LEUCOCYTES

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU MC NL PL PT RO SE SI SK TR

(30) Priority: **11.05.2004 US 570406 P**

(43) Date of publication of application:
24.01.2007 Bulletin 2007/04

(60) Divisional application:
10005232.3 / 2 214 014

(73) Proprietor: **University of Pittsburgh - Of the Commonwealth System of Higher Education Pittsburgh, PA 15260 (US)**

(72) Inventors:
• **AHEARN, Joseph M. Sewickley, Pennsylvania 15143 (US)**
• **MANZI, Susan M. Wexford, Pennsylvania 15090 (US)**
• **LIU, Chau-Ching Pittsburgh, Pennsylvania 15217 (US)**

(74) Representative: **Baldock, Sharon Claire Boulton Tennant Verulam Gardens 70 Gray's Inn Road London WC1X 8BT (GB)**

(56) References cited:
WO-A-03/022223 US-A1- 2005 037 441 US-A1- 2005 042 602

- **MARQUART H V ET AL: "COMPLEMENT RECEPTOR EXPRESSION AND ACTIVATION OF THE COMPLEMENT CASCADE ON B LYMPHOCYTES FROM PATIENTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS (SLE)" CLINICAL AND EXPERIMENTAL IMMUNOLOGY, OXFORD, GB, vol. 101, no. 1, 1995, pages 60-65, XP009044164 ISSN: 0009-9104**
- **MANZI S ET AL: "New insights into complement: a mediator of injury and marker of disease activity in Systemic Lupus Erythematosus" LUPUS, BASINGSTOKE, GB, vol. 13, no. 5, 1 May 2004 (2004-05-01), pages 298-303, XP002983171 ISSN: 0961-2033**
- **KATYAL MONIKA ET AL: "Complement receptor 1 in autoimmune disorders" CURRENT SCIENCE (BANGALORE), vol. 81, no. 8, 25 October 2001 (2001-10-25), pages 907-914, XP002470821 ISSN: 0011-3891 & DATABASE INTERNET [Online] Sage Journals Online; 27 February 2008 (2008-02-27), retrieved from HTTP: //LUP.SAGEPUB.COM/CONTENT/VOL13/ISSUE 5/**
- **DATABASE MEDLINE [Online] MELICONI ET AL: 'Complement Activation Products in idiopathic pulmonary fibrosis: relevance of fragment Ba to disease severity.', XP002990742 Database accession no. (90367342) & CLINICAL IMMUNOLOGY AND IMMUNOPATHOLOGY. vol. 57, no. 1, October 1990, pages 64 - 73**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 745 287 B1

- DATABASE MEDLINE [Online] MCCARTHY ET AL: 'Intrauterine devices and pelvic inflammatory disease.', XP002990743 Database accession no. (85046338) & AUSTRALIAN AND NEW ZEALAND JOURNAL OF OBSTETRICS AND GYNAECOLOGY. vol. 24, no. 2, May 1984, pages 106 - 110
- CLEMENZA, LILIANA; ISENMAN, DAVID E.: "The C4A and C4B Isotypic Forms of Human Complement Fragment C4b Have the Same Intrinsic Affinity for Complement Receptor 1 (CR1/CD35)" JOURNAL OF IMMUNOLOGY, vol. 172, 1 February 2004 (2004-02-01), pages 1670-1680,
- VAN DEN ELSEN JEAN M H; MARTIN ALBERTO; WONG VERONICA; CLEMENZA LILIANA; ROSE DAVID R; ISENMAN DAVID E: "X-ray crystal structure of the C4d fragment of human complement component C4" JOURNAL OF MOLECULAR BIOLOGY, vol. 322, no. 5, 4 October 2002 (2002-10-04), pages 1103-1115, ISSN: 0022-2836
- RODGAARD A ET AL: "COMPLEMENT RECEPTOR TYPE 1 CR1 CD35 EXPRESSION ON PERIPHERAL T LYMPHOCYTES BOTH CD4 AND CD8-POSITIVE CELLS EXPRESS CR1" COMPLEMENT AND INFLAMMATION, vol. 8, no. 5-6, 1991, pages 303-309, ISSN: 1012-8204
- WILSON J G ET AL: "DECREASED EXPRESSION OF THE COMPLEMENT C-3B-C-4B RECEPTOR CR-1 AND THE C-3D RECEPTOR CR-2 ON B LYMPHOCYTES AND OF CR-1 ON NEUTROPHILS OF PATIENTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS" ARTHRITIS AND RHEUMATISM, vol. 29, no. 6, 1986, pages 739-747, ISSN: 0004-3591
- YASKANIN D D ET AL: "DISTRIBUTION AND QUANTITATIVE EXPRESSION OF THE COMPLEMENT RECEPTOR TYPE 1 CR1 ON HUMAN PERIPHERAL BLOOD T LYMPHOCYTES" CELLULAR IMMUNOLOGY, vol. 142, no. 1, 1992, pages 159-176, ISSN: 0008-8749

Description

FIELD OF THE INVENTION

5 **[0001]** The invention is related to methods of diagnosing inflammatory diseases or conditions by determining levels of components of the complement pathway on the surface of white blood cells.

BACKGROUND OF THE INVENTION

10 **[0002]** This invention relates to the diagnosis and/or monitoring of patients with immunologic inflammatory conditions and diseases, *e.g.*, systemic lupus erythematosus (SLE). Inflammation is a characteristic of virtually every immune system disease or condition and every infectious disease or condition. Many chronic inflammatory conditions and diseases, *i.e.*, immune system diseases or conditions and infectious diseases or conditions, cause damage to multiple organ systems and are difficult to diagnose. Because the symptoms of many immunologic inflammatory conditions and diseases overlap there is a great need for diagnostic methods for rapidly and reliably diagnosing and monitoring specific immune system disease and conditions. The present invention solves this and other needs.

15 **[0003]** The present invention provides a method for diagnosing or monitoring systemic lupus erythematosus (SLE) comprising:

- 20 (a) determining, in a blood sample from the individual the level of the complement pathway component C4d on the surface of a T lymphocyte or a B lymphocyte in the sample, and
 (b) comparing said determinations with, respectively, the level of the complement pathway component C4d on the surface of a control T lymphocyte or B lymphocyte, wherein an increase in the level of complement component C4d indicates that the individual has systemic lupus erythematosus (SLE).

25 **[0004]** Diagnosis of SLE can also be accomplished by determining the level of C4d on a T lymphocyte or a B lymphocyte in combination with at least one other complement pathway component. In one embodiment the levels of complement components C4d and C3d are determined to diagnose or monitor SLE. C4d levels can be determined as above. C3d levels can be determined using *e.g.*, antibodies specific for C3d. The antibodies can be labelled for detection and *e.g.*, polyclonal or monoclonal antibodies can be used.

30 **[0005]** In another embodiment, the T lymphocyte or B lymphocyte is isolated using an antibody specific for the lymphocyte, *e.g.*, anti-CD3, CD4, CD8 or CD19 antibodies.

[0006] In a further aspect the invention provides a computer program which, when executed by a computer carries out a method according to the invention, said computer program comprising:

- 35 (a) code for receiving data corresponding to a determination of a complement component C4d deposited on the surface of a T lymphocyte or a B lymphocyte;
 (b) code for retrieving a reference value for the level of the complement component C4d deposited on the surface of a T lymphocyte or a B lymphocyte of individuals;
 40 (c) code for comparing the data in (a) with the reference value in (b).

[0007] Also provided is a computer readable medium storing a computer program according to the above.

[0008] Marquart et al., Clin. Exp. Immunol. 101, 60-65 (1995) describes increased deposition of C3dg on B-lymphocytes in SLE.

45 **[0009]** Manzi et al., Lupus 13, 298-303 (2004) describes assays to detect C4d on erythrocytes for diagnosis and monitoring of disease activity in SLE.

[0010] Katyal et al., Current Science 81 (8), 907-914 (2001) discloses that complement receptor 1 is expressed on monocytes and B- and T-lymphocytes, amongst other cell types.

50 **[0011]** van den Elsen et al., J. Mol. Biol. 332, 1103-1115 (2002) shows the X-ray crystal structure of the C4d fragment of human C4.

BRIEF DESCRIPTION OF THE DRAWINGS

55 **[0012]** Figure 1 demonstrates that complement pathway component C4d is deposited specifically on peripheral blood T lymphocytes of patients with SLE. Deposition of C4d on peripheral blood T lymphocytes was determined by a 2-color flow cytometric assay. T lymphocytes were identified using a FITC-conjugated monoclonal antibody specific for CD3 (a surface marker for T lymphocytes), and C4d deposited on these cells was determined using a monoclonal anti-C4d antibody followed by a PE-conjugated secondary antibody. Data shown are C4d-specific mean fluorescence (SMF) of

peripheral blood T lymphocytes derived from patients with SLE (n=87, diamonds), patients with other autoimmune diseases (n=31, squares), or healthy controls (n=19, triangles). A cutpoint of specific mean fluorescence was empirically determined to distinguish individuals with C4d-positive T lymphocytes (SMF > 58) from those with C4d-negative T lymphocytes (SMF < 58). The frequencies of individuals with C4d-positive T lymphocytes among the groups was compared using the Chi-square test, and the p values for each pair compared are shown above the horizontal line. The mean value of C4d-specific fluorescence on T lymphocytes among the groups was compared using the Students' T test and the respective p values are shown below the horizontal line.

[0013] Figure 2 demonstrates that complement pathway component C3d is deposited specifically on peripheral blood T lymphocytes of patients with SLE. Deposition of C3d on peripheral blood T lymphocytes was determined by a 2-color flow cytometric assay. T lymphocytes were identified using a FITC-conjugated monoclonal antibody specific for CD3 (a surface marker for T lymphocytes), and C3d deposited on these cells was determined using a monoclonal anti-C4d antibody followed by a PE-conjugated secondary antibody. Data shown are C3d-specific mean fluorescence of peripheral blood T lymphocytes derived from patients with SLE (n=87, diamonds), patients with other autoimmune diseases (n=31, squares), or healthy controls (n=19, triangles). A cutpoint of specific mean fluorescence was empirically determined to distinguish individuals with C3d-positive T lymphocytes (SMFC > 3) from those with C3d-negative T lymphocytes (SMFC < 3). The frequencies of individuals with C4d-positive T lymphocytes among the groups was compared using the Chi-square test, and the p values for each pair compared are shown above the horizontal line. The mean value of C3d-specific fluorescence on T lymphocytes among the groups was compared using the Students' T test and the respective p values are shown below the horizontal line.

[0014] Figure 3 demonstrates that complement ligands C4d and C3d are deposited specifically on peripheral blood T lymphocytes of patients with SLE. Deposition of C3d and C4d on peripheral blood T lymphocytes was determined by a 2-color flow cytometric assay. T lymphocytes were identified using a FITC-conjugated monoclonal antibody specific for CD3 (a surface marker for T lymphocytes), and C3d/C4d deposited on these cells was determined using a monoclonal anti-C3d (or anti-C4d) antibody, followed by a PE-conjugated secondary antibody. Data shown are C3d- and C4d-specific median fluorescence of peripheral blood T lymphocytes derived from patients with SLE (n=87), patients with other autoimmune diseases (n=31), or healthy controls (n=19). Cutpoints of specific mean fluorescence were empirically determined to distinguish individuals with C3d-positive T lymphocytes (SMFC > 3) or individuals with C4d-positive T lymphocytes (SMFI > 58) from those with C3d-negative or C4d-negative T lymphocytes. The mean value of C3d-specific fluorescence on T lymphocytes among the groups was compared using the Students' T test and the respective p values are <0.0001 (not shown).

[0015] Figure 4 provides levels of C4d on the Surface of T Lymphocytes, B Lymphocytes, and Monocytes. C4d on different types of cells were determined by a 3-color flow cytometric assay using monoclonal antibodies specific for cell-specific surface markers and C4d or isotype control immunoglobulins. Levels of C4d were calculated as specific median fluorescence intensity (SMFI)= anti-C4d median fluorescence intensity - isotype Ig median fluorescence intensity. ^aT cells were identified by electronic gating of cells positively stained by a monoclonal anti-CD3 antibody. ^bB cells were identified by electronic gating of cells positively stained by a monoclonal anti-CD19 antibody. ^cMonocytes were identified by forward and side scattering and negative staining by anti- Cd3. ^dPatients with other inflammatory diseases such as inflammatory myopathies, Sjogren's syndrome, vasculitis, Raynaud's phenomenon, and cardiovascular disease. ^eStudent t test; patients with SLE vs. patients with other diseases ^fStudent t test; patients with SLE vs. healthy controls.

[0016] Figure 5 provides levels of C3d on the Surface of T Lymphocytes, B Lymphocytes, and Monocytes. C3d on different types of cells were determined by a 3-color flow cytometric assay using monoclonal antibodies specific for cell-specific surface markers and C3d or isotype control immunoglobulins. Levels of C4d were calculated as specific median fluorescence intensity (SMFI)= anti-C4d median fluorescence intensity - isotype Ig median fluorescence intensity. ^aT cells were identified by electronic gating of cells positively stained by a monoclonal anti-CD3 antibody. ^bB cells were identified by electronic gating of cells positively stained by a monoclonal anti-CD19 antibody. ^cMonocytes were identified by forward and side scattering and negative staining by anti-Cd3. ^dPatients with other inflammatory diseases such as inflammatory myopathies, Sjogren's syndrome, vasculitis, Raynaud's phenomenon, and cardiovascular disease. ^eStudent t test; patients with SLE vs. patients with other diseases ^fStudent t test; patients with SLE vs. healthy controls

[0017] Figure 6 Deposition of C4d on Peripheral Blood T cells, depicts the data of figure 1 using a logarithmic scale on the Y-axis providing a clearer picture of the differences between the healthy controls and the diseased states.

[0018] Figure 7 Deposition of C3d on Peripheral Blood T cells, depicts the data of figure 2 using a logarithmic scale on the Y-axis providing a clearer picture of the differences between the healthy controls and the diseased states.

DETAILED DESCRIPTION OF THE INVENTION

Introduction

5 **[0019]** This disclosure provides methods of diagnosing and monitoring inflammatory diseases or conditions by determining the level of at least one complement component on the surface of a white blood cell. Previously, complement component C4d and CR1 levels on erythrocytes were determined and used to diagnose systemic lupus erythematosus (SLE) in individuals. See, e.g., WO03/022223 published March 20, 2003.

10 This disclosure is the first to describe diagnosing and monitoring inflammatory diseases or conditions by determining the level of at least one complement pathway component on the surface of a white blood cell.

[0020] In diagnosing the occurrence, or previous occurrence, of an inflammatory disease or condition, the level of at least one complement pathway component deposited on surfaces of white blood cells in a sample is determined. This determination is then compared with the quantities of the same complement pathway component found on the surfaces of white blood cells of individuals not having the inflammatory disease or condition.

15 **[0021]** In monitoring disease activity of a patient with an inflammatory disease or condition, a determination of at least one complement pathway component is made in the patient's blood sample, and is then compared with determinations of the quantities of the same complement pathway component on surfaces of white blood cells in a sample obtained from the same patient in the past. Comparison can also be made to quantities of the same complement pathway component found on the surfaces of white blood cells of individuals not having the inflammatory disease or condition.

20 **[0022]** The methods of this disclosure can be used to diagnose and/or monitor SLE by determining the level of the complement pathway component C4d and, optionally complement component C3d on lymphocytes. Because SLE is a serious health problem, there is a need for relatively accurate and early diagnosis of this condition. Likewise, the ability to monitor the activity of this disease is of great importance. The methods of this disclosure can also be used to diagnose and/or monitor SLE by determining the level of C4d and the level of complement pathway component C3d on lymphocytes.

Definitions

25 **[0023]** As used herein, an "inflammatory disease or condition" refers to any immune disease or condition that causes increased inflammation in an individual. An inflammatory disease or condition also refers to any infectious disease or condition that causes increased inflammation in an individual. In some embodiments the inflammatory disease or condition is a "chronic inflammatory disease or condition." A chronic inflammatory disease or condition is an inflammatory condition that does not resolve after a period of weeks, months or longer. Chronic inflammatory conditions can follow an acute inflammatory condition, or for some diseases or conditions can occur in the absence of an acute inflammatory disease or condition. An inflammatory disease or condition includes the following: SLE, rheumatoid arthritis, vasculitis (and its specific forms such as Wegener's granulomatosis), scleroderma, myositis, serum sickness, transplant rejection, sickle cell anemia, gout, complications of pregnancy such as pre-eclampsia, multiple sclerosis, cardiovascular disease, infectious disease such as hepatitis C virus infection, etc. Each of these diseases or conditions can also be described as chronic inflammatory diseases or conditions.

30 **[0024]** As used herein a "white blood cell" refers to circulating blood cells that are not erythrocytes or reticulocytes, e.g., lymphocytes, e.g., T and B cells, NK cells, eosinophils, basophils, granulocytes, neutrophils, monocytes, macrophages, megakaryocytes, plasma cells, circulating endothelial cells, and stem cells.

35 **[0025]** As used herein a "control white blood cell" refers to a white blood cell as defined above that is isolated from an individual who does not have an inflammatory disease or condition. When an inflammatory disease or condition is being monitored in a patient, a control white blood cell can also refer to a white blood cell isolated from the same patient at an earlier time, e.g., weeks, months, or years earlier.

40 **[0026]** As used herein the "complement pathway or system" refers to a complex network of more than 30 functionally linked proteins that interact in a highly regulated manner to provide many of the effector functions of humoral immunity and inflammation, thereby serving as the major defense mechanism against bacterial and fungal infections. This system of proteins acts against invasion by foreign organisms via three distinct pathways: the classical pathway (in the presence of antibody) or the alternative pathway (in the absence of antibody) and the lectin pathway. Once activated, the proteins within each pathway form a cascade involving sequential self-assembly into multimolecular complexes that perform various functions intended to eradicate the foreign antigens that initiated the response. For a review of the complement pathway, see, e.g., Sim and Tsiftoglou, Biochem. Soc. Trans. 32:21-27 (2004).

45 **[0027]** The classical pathway is usually triggered by an antibody bound to a foreign particle. It consists of several components that are specific to the classical pathway and designated C1, C4, C2. Sequentially, binding of C1q to an antigen-antibody complex results in activation of C1r and C1s (both are serine proteases), and activated C1s cleaves C4 and C2 into, respectively, C4a and C4b and C2a and C2b. Fragments C4b and C2a assemble to form C4b2a, which cleaves protein C3 into C3a and C3b, which completes activation of the classical pathway. Fragments C4b and C3b are

subject to further degradation by Factor I. This factor cleaves C4b to generate C4d and also cleaves C3b, to generate iC3b followed by C3d. Thus, activation of the classical pathway of complement can lead to deposition of a number of fragments, such as C4d, iC3b, and C3d, on immune complexes or other target surfaces. Such targets include cells circulating in the blood, *e.g.*, lymphocytes and other white blood cells, erythrocytes and platelets.

[0028] As used herein a "component of the complement pathway" includes proteins C1, C4, C2, C3 and fragments thereof, *e.g.*, C1q, C1r, C1s, C4a, C4b, C2a, C2b, C4b2a, C3a, C3b, C4c, C4d, iC3b, C3d, C3i, C3dg. Also included are C5, C5b, C6, C7, C8, C9, C1inh, MASP2, CR1, DAF, MCP, CD59, C3aR, C5aR, C1qR, CR2, CR3, and CR4, as well as other complement pathway components, receptors and ligands not listed specifically herein.

[0029] As used herein, "systemic lupus erythematosus", "SLE", or "lupus" is the prototypic autoimmune disease resulting in multiorgan involvement. This anti-self response is characterized by autoantibodies directed against a variety of nuclear and cytoplasmic cellular components. These autoantibodies bind to their respective antigens, forming immune complexes which circulate and eventually deposit in tissues. This immune complex deposition and consequential activation of the complement system causes chronic inflammation and tissue damage.

[0030] Diagnosing and monitoring disease activity are problematic in patients with SLE. Diagnosis is problematic because of the broad spectrum of disease ranging from subtle or vague symptoms to life threatening multi-organ failure. Moreover, other inflammatory diseases with multi-system involvement can be mistaken for lupus, or vice versa. Criteria were developed for the purpose of disease classification in 1971 (Cohen, AS, et al., 1971. *Bull.Rheum.Dis.* 21:643-648) and revised in 1982 (Tan, EM, et al., 1982. *Arth.Rheum.* 25:1271-1277) and 1997 (Hochberg, MC. 1997. *Arth.Rheum.* 40:1725). These criteria are meant to ensure that patients from different geographic locations are comparable. Of the eleven criteria, the presence of four or more, either serially or simultaneously, is sufficient for classification of a patient as having SLE. Although these criteria serve as useful reminders of those features that distinguish SLE from other related autoimmune diseases, they are unavoidable fallible. Determining the presence or absence of the criteria often depends on physicians' interpretation and judgment. The range of clinical manifestations in SLE is much greater than that described by the eleven criteria and each manifestation can vary in the level of activity and severity from one patient to another. Furthermore, symptoms of SLE often evolve over the course of disease. There is no definitive test for SLE to date, and, thus, it is often misdiagnosed. This disclosure, however, provides efficient and accurate methods for diagnosis of SLE and other inflammatory diseases and conditions.

[0031] SLE progresses in a series of flares, or periods of acute illness, followed by remissions. The symptoms of a flare, which vary considerably between patients and even within the same patient, include malaise, fever, symmetric joint pain, and photosensitivity (development of rashes after brief sun exposure). Other symptoms of SLE include hair loss, ulcers of mucous membranes and inflammation of the lining of the heart and lungs which leads to chest pain. Red blood cells, platelets and white blood cells can be targeted in lupus, resulting in anemia and bleeding problems. More seriously, immune complex deposition and chronic inflammation in the blood vessels can lead to kidney involvement and occasionally failure requiring dialysis or kidney transplantation. Since the blood vessel is a major target of the autoimmune response in SLE, premature strokes and heart disease are not uncommon. Over time, however, these flares can lead to irreversible organ damage.

[0032] "Antibody" refers to a polypeptide comprising a framework region from an immunoglobulin gene or fragments thereof that specifically binds and recognizes an antigen. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon, and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively. Typically, the antigen-binding region of an antibody will be most critical in specificity and affinity of binding.

[0033] An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kD) and one "heavy" chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (V_L) and variable heavy chain (V_H) refer to these light and heavy chains respectively.

[0034] Antibodies exist, *e.g.*, as intact immunoglobulins or as a number of well-characterized fragments produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce $F(ab)'_2$, a dimer of Fab which itself is a light chain joined to V_H-C_H1 by a disulfide bond. The $F(ab)'_2$ may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the $F(ab)'_2$ dimer into an Fab' monomer. The Fab' monomer is essentially Fab with part of the hinge region (*see* Fundamental Immunology (Paul ed., 3d ed. 1993). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized *de novo* either chemically or by using recombinant DNA methodology. Thus, the term antibody, as used herein, also includes antibody fragments either produced by the modification of whole antibodies, or those synthesized *de novo* using recombinant DNA methodologies (*e.g.*, single chain Fv) or those identified using phage display libraries (*see, e.g.*, McCafferty et al., *Nature* 348:552-554 (1990))

[0035] For preparation of antibodies, *e.g.*, recombinant, monoclonal, or polyclonal antibodies, many techniques known in the art can be used (*see, e.g.*, Kohler & Milstein, *Nature* 256:495-497 (1975); Kozbor et al., *Immunology Today* 4: 72 (1983); Cole et al., pp. 77-96 in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985); Coligan, *Current Protocols in Immunology* (1991); Harlow & Lane, *Antibodies, A Laboratory Manual* (1988); and Goding, *Monoclonal Antibodies: Principles and Practice* (2d ed. 1986)). The genes encoding the heavy and light chains of an antibody of interest can be cloned from a cell, *e.g.*, the genes encoding a monoclonal antibody can be cloned from a hybridoma and used to produce a recombinant monoclonal antibody. Gene libraries encoding heavy and light chains of monoclonal antibodies can also be made from hybridoma or plasma cells. Random combinations of the heavy and light chain gene products generate a large pool of antibodies with different antigenic specificity (*see, e.g.*, Kuby, *Immunology* (3rd ed. 1997)). Techniques for the production of single chain antibodies or recombinant antibodies (U.S. Patent 4,946,778, U.S. Patent No. 4,816,567) can be adapted to produce antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized or human antibodies (*see, e.g.*, U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, Marks et al., *Bio/Technology* 10:779-783 (1992); Lonberg et al., *Nature* 368:856-859 (1994); Morrison, *Nature* 368:812-13 (1994); Fishwild et al., *Nature Biotechnology* 14:845-51 (1996); Neuberger, *Nature Biotechnology* 14:826 (1996); and Lonberg & Huszar, *Intern. Rev. Immunol.* 13:65-93 (1995)). Alternatively, phage display technology can be used to identify antibodies and heteromeric Fab fragments that specifically bind to selected antigens (*see, e.g.*, McCafferty et al., *Nature* 348:552-554 (1990); Marks et al., *Biotechnology* 10:779-783 (1992)). Antibodies can also be made bispecific, *i.e.*, able to recognize two different antigens (*see, e.g.*, WO 93/08829, Traunecker et al., *EMBO J.* 10:3655-3659 (1991); and Suresh et al., *Methods in Enzymology* 121:210 (1986)). Antibodies can also be heteroconjugates, *e.g.*, two covalently joined antibodies, or immunotoxins (*see, e.g.*, U.S. Patent No. 4,676,980, WO 91/00360; WO 92/200373; and EP 03089).

[0036] In one embodiment, the antibody is conjugated to an "effector" moiety. The effector moiety can be any number of molecules, including labeling moieties such as radioactive labels or fluorescent labels for use in diagnostic assays.

[0037] The phrase "specifically (or selectively) binds" to an antibody or "specifically (or selectively) immunoreactive with," when referring to a protein or peptide, refers to a binding reaction that is determinative of the presence of the protein, often in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and more typically more than 10 to 100 times background. Specific binding to an antibody under such conditions requires an antibody that is selected for its specificity for a particular protein. For example, polyclonal antibodies raised to a component of the complement pathway or to a marker of a white blood cell, polymorphic variants, alleles, orthologs, and conservatively modified variants, or splice variants, or portions thereof, can be selected to obtain only those polyclonal antibodies that are specifically immunoreactive with the component of the complement pathway or the marker of a white blood cell and not with other proteins. This selection may be achieved by subtracting out antibodies that cross-react with other molecules. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select antibodies specifically immunoreactive with a protein (*see, e.g.*, Harlow & Lane, *Antibodies, A Laboratory Manual* (1988) for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity).

[0038] An "antigen" is a molecule that is recognized and bound by an antibody, *e.g.*, peptides, carbohydrates, organic molecules, or more complex molecules such as glycolipids and glycoproteins. The part of the antigen that is the target of antibody binding is an antigenic determinant and a small functional group that corresponds to a single antigenic determinant is called a hapten.

[0039] A "label" is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include ³²P, ¹²⁵I, fluorescent dyes, electron-dense reagents, enzymes (*e.g.*, as commonly used in an ELISA), biotin, digoxigenin, or haptens and proteins for which antisera or monoclonal antibodies are available (*e.g.*, antibody specific for a component of the complement pathway or a marker of a white blood cell can be made detectable, *e.g.*, by incorporating a radiolabel or fluorescent label into the antibody, and used to detect component of the complement pathway or the marker of a white blood cell specifically reactive with the labeled antibody). A labeled secondary antibody can also be used to detect an antibody specific for a component of the complement pathway or a marker of a white blood cell.

[0040] The term "contact" or "contacting" is used herein interchangeably with the following: combined with, added to, mixed with, passed over, incubated with, flowed over, *etc.*

[0041] The term "immunoassay" is an assay that uses an antibody to specifically bind an antigen. The immunoassay is characterized by the use of specific binding properties of a particular antibody to isolate, target, and/or quantify the antigen.

[0042] In both instances, when speaking of "determination or determining" and "quantity," we mean to include both an amount or quantity of material. When more than one complement pathway component is measured, *e.g.*, C4d and C3d "determination or determining" and "quantity," mean in addition, or alternatively, a ratio of a first complement pathway component to a second complement pathway component, *e.g.*, a ratio of C4d to C3d.

Determination of the level of a component of the complement pathway on a white blood cell.

[0043] The invention involves conducting assays on white blood cells obtained from patients to determine levels of complement pathway components. The levels of the complement components are then used to diagnose or monitor an inflammatory disease or condition in an individual.

[0044] Samples of blood are obtained from the patient and are treated with EDTA (ethylenediaminetetraacetate) to inhibit complement activation. The samples are maintained at room temperature or under cold conditions. Assays are run preferably within 24 hours.

[0045] In some embodiments the white blood cells are isolated from other components of the blood sample. For example, white blood cells (the buffy coat) can be isolated from plasma and from red blood cells by centrifugation. Each type of white blood cell can be isolated through the use of antibodies against known cell surface markers that are specific for that cell type, *e.g.*, a lymphocyte. Antibodies against cell surface markers of white blood cells are known to those of skill and are commercially available, *e.g.* from BD Immunocytometry Systems. For example, cell surface markers CD3, CD4, CD8, and CD19 are specific for lymphocytes and monoclonal antibodies specific for CD3, CD4, CD8, and CD19 are available from BD Immunocytometry Systems, San Jose, CA.

[0046] Isolation of white blood cells can be done by attaching an antibody specific to a cell's surface marker to a solid support, then contacting a sample containing the white blood cells with the linked antibody. Contaminating cells are washed away allowing the isolated white blood cells to be collected.

[0047] In some embodiments, FACS is used to isolate a white blood cell, *e.g.*, a lymphocyte. The term "FACS" refers to fluorescence activated cell sorting, a technique used to separate cells according to their content of particular molecules of interest. The molecule of interest can be specific for a type of cell or for particular cell state. The molecule of interest can be fluorescently labeled directly by binding to a fluorescent dye, or by binding to a second molecule, which has been fluorescently labeled, *e.g.*, an antibody, lectin or aptamer that has been fluorescently labeled and that specifically binds to the molecule of interest. Thus, white blood cell specific markers can be used to isolate specific white blood cells from other cells in a blood sample.

[0048] Isolation of white blood cells also refers to gating techniques used to assay a particular cell population during flow cytometric analysis. A labeled marker specific for a white blood cell population of interest is used to analyze those cells from a population. A second labeled marker is then used to determine the level of a component of the complement pathway on the surface of the white blood cell.

[0049] The determination of the level of a component of the complement pathway may be done by a number of methods including flow cytometry, ELISA using white blood cell lysates, and radioimmunoassay. In one embodiment of this invention, the determination of the levels of a component of the complement pathway is made using flow cytometric methods, with measurements taken by direct or indirect immunofluorescence using polyclonal or monoclonal antibodies specific for the component of the complement pathway. The mean fluorescence channel (MFC) for the white blood cell component of the complement pathway is determined. Determination of complement components, *e.g.*, C4d, CR1, and, on the surface of red blood cells or platelets is described in WO03/022223, published March 20, 2003 and in USSN 60/463,447, filed April 16, 2003.

[0050] In one embodiment, levels of the complement pathway component C4d or complement component C3d are determined on the surface of lymphocytes to diagnose or monitor the progression of SLE in individuals. The lymphocytes are isolated or detected using lymphocyte specific antibodies *e.g.*, anti-CD3, CD4, CD8, or CD19 antibodies. In some embodiments, complement pathway component C4d and complement pathway component C3d levels are determined. Determination of C4d and C3d levels can be done by a number of methods including flow cytometry, ELISA using lymphocyte lysates, and radioimmunoassay. In one embodiment of this invention, the determination of the levels of C4d and C3d is made using flow cytometric methods, with measurements taken by direct or indirect immunofluorescence using polyclonal or monoclonal antibodies specific for C4d or C3d. The mean fluorescence channel (MFC) for lymphocyte C4d or C3d is determined. The same type of assay may be used for diagnosis or for monitoring disease activity in patients known to have SLE.

Kits

[0051] Kits for conducting the assays for both the diagnosing of inflammatory disease and monitoring of inflammatory disease activity are described but not claimed. Said kits will use any of the various reagents needed to perform the methods described herein. For example using the immunofluorescence assays, the kits will generally comprise a conjugate of a monoclonal antibody specific for complement pathway component (*e.g.*, anti-C4d or C3d antibodies) with a fluorescent moiety, and preferably also a conjugate of a monoclonal antibody specific for a white blood cell of interest (*e.g.*, lymphocytes using, *e.g.*, anti-CD3, CD4, CD8, and CD19 antibodies) with a different fluorescent moiety. Additionally, the kits can comprise instructional material for the user and such other material as may be needed in carrying out assays of this type, for example, buffers, radiolabelled antibodies, colorimeter reagents etc.

[0052] The antibodies for use in these methods and kits are known. For example, monoclonal antibodies specific for CD3, CD4, CD8, and CD19 are available from Becton Dickinson Immunocytometry Systems, San Jose, CA. Anti-C4d and anti-C3d antibodies are available from Quidel Corp. in San Diego, California and are generally described in Rogers, J., N. Cooper, et al. PNAS 89:10016-10020, (1992); Schwab, C. et al. Brain. Res. 707(2):196 (1996); Gemmell, C. J. Biomed Mater. Res. 37:474-480, (1997); and, Stoltzner, S.E., et al. Am. J. Path. 156:489-499, (2000).

Diagnostic methods

[0053] Diagnosis of a patient with SLE is carried out by comparing the determination of C4d (and optionally C3d) with a base value or range of values for the quantities of C4d (and optionally C3d) typically present on the surfaces of T lymphocytes or B lymphocytes in normal individuals.

[0054] For example, diagnosis of a patient with SLE is carried out by comparing the determination of C4d and, optionally C3d with a base value or range of values for the quantities of C4d and C3d typically present on the surfaces of lymphocytes in normal individuals. In normal individuals, low levels of C4d may occasionally be detected but C3d is not present. When using flow cytometric measurement with indirect immunofluorescence, the MFC of C4d and C3d on lymphocytes of healthy individuals ranged from 1.25 to 58.63 (mean 17.02) and -3.42 to 2.67 (mean 0.52), respectively. (Tables 1 and 3). The MFC of lymphocytes C4d and C3d in patients having SLE was higher than that of healthy individuals and ranged from -2.62 to 1057.19 (mean 201.06) and -3.95 to 318.18 (mean 62.42), respectively (Tables 2 and 3).

Monitoring of patients

[0055] A particular feature of the methods of this invention is to indicate or reflect inflammatory activity that has occurred in the patient during the preceding several weeks or even several months. It is possible, using this procedure, to identify the occurrence of a flare-up of an inflammatory disease or condition, such as SLE, during the previous few weeks or possibly even the previous several months due to persistence of components of the complement pathway deposited on the surface of white blood cells, e.g., C4d and C3d deposited on the surface of lymphocytes.

Automation and computer software

[0056] The determinations of complement pathway components e.g., C4d and, optionally C3d, and the diagnostic and disease activity monitoring methods described above can be carried out manually, but often are conveniently carried out using an automated system and/or equipment, in which the blood sample is analyzed automatically to make the necessary determination or determinations, and the comparison with the base or reference value is carried out automatically, using computer software appropriate to that purpose.

[0057] Also described is a method for diagnosing or monitoring SLE in an individual comprising (a) automatically determining, in a blood sample from the individual containing lymphocytes, complement components C4d and, optionally C3d deposited on surfaces of lymphocytes in the sample, and (b) automatically comparing said determinations with reference values for components C4d and C3d on surfaces of lymphocytes.

(1): a computer readable medium, comprising:

code for receiving data corresponding to a determination of complement components C4d and, optionally C3d deposited on surfaces of lymphocytes; code for retrieving a reference value for complement components C4d and, optionally C3d deposited on surfaces of lymphocytes of individuals; and code for comparing the data in (a) with the reference value of (b).

[0058] Computer software, or computer-readable media for use in the methods of this invention include:

(1): a computer readable medium, comprising:

code for receiving data corresponding to a determination of complement components C4d and, optionally C3d deposited on surfaces of lymphocytes; code for retrieving a reference value for complement components C4d and, optionally C3d deposited on surfaces of lymphocytes of individuals; and code for comparing the data in (a) with the reference value of (b).

[0059] In embodiments of the invention, one or more reference values may be stored in a memory associated with a digital computer. After data corresponding to determinations of complement pathway components is obtained (e.g., from an appropriate analytical instrument), the digital computer may compare the complement pathway component data with

one or more appropriate reference values. After this comparison takes place, the digital computer can automatically determine if the data corresponding to the determination of t complement pathway component is associated with an inflammatory disease or condition of interest.

5 [0060] In some embodiments of the invention, one or more reference values may be stored in a memory associated with a digital computer. After data corresponding to determinations of complement C4d and C3d is obtained (e.g., from an appropriate analytical instrument), the digital computer may compare the C4d and C3d data with one or more appropriate reference values. After this comparison takes place, the digital computer can automatically determine if the data corresponding to the determination of complement C4d and C3d is associated with SLE.

10 [0061] Accordingly, some embodiments of the invention may be embodied by computer code that is executed by a digital computer. The digital computer may be a micro, mini or large frame computer using any standard or specialized operating system such as a Windows™ based operating system. The code may be stored on any suitable computer readable media. Examples of computer readable media include magnetic, electronic, or optical disks, tapes, sticks, chips, etc. The code may also be written by those of ordinary skill in the art and in any suitable computer programming language including, C, C++, etc.

15 EXAMPLES AND EXPERIMENTAL DATA

[0062] The following examples are provided by way of illustration only and not by way of limitation. Those of skill will readily recognize a variety of noncritical parameters that could be changed or modified to yield essentially similar results.

20 Example 1: Assays of Lymphocyte C4d and C3d in Healthy Controls: Negative

25 [0063] Nineteen healthy individuals were studied. As shown in Table 1, C3d was not detected or barely detectable on lymphocytes of each of the nineteen healthy individuals. Samples of 3 ml of EDTA-anticoagulated peripheral blood were taken from each individual and used as a source of lymphocytes and other white blood cells. The lymphocytes were washed and resuspended in FACS buffer. Levels of C4d, C3d, and CD3 were measured by two color indirect immunofluorescence using monoclonal antibodies specific for C4d, C3d, and CD3, respectively. Levels of C4d and C3d are quantitated by flow cytometry using a FACSCalibur cytometer (Becton Dickinson). The lymphocytes were identified by forward and side scatter and CD3-fluorescence, and the mean fluorescence channel (MFC) was determined for C4d and C3d.

30 [0064] More particularly, blood drawn into a Vacutainer™ containing EDTA (Becton Dickinson, Franklin Lakes, NJ) was centrifuged at 200 xg. The buffy coat containing white blood cells (WBC) were carefully collected, transferred into a fresh tube, and washed with phosphate buffered saline (PBS). After removal of contaminating erythrocytes by hypotonic lysis, the remaining leukocytes were washed with PBS and immunofluorescently labeled using different combinations of specific antibodies for a two-color flow cytometric analysis. Antibodies used in the initial study included: 1) antibodies specific for cell lineage markers, e.g., anti-CD3, anti-CD4, and anti-CD8 for T lymphocytes, anti-CD19 for B lymphocytes, and anti-CD14 for monocytes (BD PharMingen, San Diego, CA), and 2) antibodies reactive to complement C4d or C3d (Quidel, San Diego, CA), or the isotype control mouse IgG MOPC21. The stained cells were then analyzed using a FACSCalibur™ flow cytometer and the CellQuest™ software (Becton Dickinson Immunocytometry Systems, San Jose, CA). Lymphocytes were electronically gated by forward scatter properties and expression of CD3 (or CD4, CD8, CD19), specific markers for lymphocyte subsets. Nonspecific binding of immunoglobulins to lymphocytes was determined by performing identical assays in parallel using the isotype control antibody MOPC21 (obtained from ATCC). Specific binding of anti-C4d and anti-C3d were determined by subtracting the MFC obtained with MOPC21 from the MFC obtained with anti-C4d and anti-C3d respectively.

45 Example 2: Assays of Lymphocyte C4d/C3d to Distinguish Patients with SLE from Healthy Controls

[0065] This example describes conducting assays on patients to diagnose SLE, and to establish reference values or ranges of values for complement components C4d and C3d.

50 [0066] For this purpose, we recruited 87 patients with SLE from our outpatient office. A single determination of lymphocyte C4d/C3d was made in 87 individuals who met ACR criteria for the diagnosis of SLE (Table 2) and in 19 healthy controls (Table 1). The mean values of C4d and C3d for patients with SLE and healthy controls are shown in Table 3. Whereas the mean value for C4d and C3d in healthy individuals was 17.02 and 0.52, respectively, the mean value for C4d and C3d among patients with SLE was 201.06 and 62.42, respectively (both p= 0.0001).

EP 1 745 287 B1

Table 1. Levels of C4d and C3d on the Surface of Lymphocytes From Healthy Controls

Healthy Control	C4d Level (SMFC) ^a	C3d Level (SMFC)
2005	58.23	-2.18
2007	2.36	-1.19
2008	7.9	0.35
2009	1.25	1.81
2022	10.45	1.56
2003	12.95	2.67
2034	2.95	0.45
2021	20.12	-1.37
2028	5.66	-1.21
2006	3.34	ND ^b
2013	7.12	1.62
2011	24.00	-3.42
2037	4.91	-1.97
2010	22.00	3.11
2025	37.41	1.57
2026	34.55	2.57
2036	22.86	1.55
2029	20.91	1.31
2035	24.36	2.22

a: specific mean fluorescence channel
b: not done

Table 2. Levels of C4d and C3d on the Surface of Lymphocytes from SLE Patients

SLE Patient	C4d Level (SMFC)	C3d Level (SMFC)
1039	768.02	68.07
107	646.51	253.23
1078	12.71	1.53
1089	0.26	-3.8
1072	12.47	-2.61
1012	31.48	15.65
1019	30.84	-9.50
1037	0.04	-3.86
1047	355.46	222.75
1003	894.69	144.30
1006	ND	133.26
1079	14.80	16.17
1052	548.01	216.67
1038	300.77	348.40
1063	20.80	15.09
1095	3.63	33.13
1097	34.73	25.87
1092	46.44	20.25
1016	352.77	318.28
1093	-2.62	1.06
1094	15.49	6.52
1034	17.92	8.10
1066	540.76	74.75
1009	1183.30	65.33

EP 1 745 287 B1

(continued)

	SLE Patient	C4d Level (SMFC)	C3d Level (SMFC)
	1014	141.62	54.80
5	1031	82.67	64.40
	1086	5.56	-1.29
	1098	16.01	0.29
	1099	8.79	7.15
	1015	1709.54	258.91
10	1100	19.99	18.49
	1101	118.94	19.61
	1102	191.47	29.13
	1053	107.76	42.44
15	1059	108.89	106.60
	1084	61.48	29.58
	1103	43.90	-1.74
	1104	119.49	4.61
	1105	910.93	202.49
20	1107	44.64	0.00
	1109	40.17	17.67
	1110	244.78	88.44
	1111	18.05	5.77
25	1085	289.48	39.73
	1043	6.45	5.46
	1056	-1.45	-1.75
	1106	66.56	12.77
	1114	241.22	25.51
30	1017	166.47	43.17
	1021	201.31	111.77
	1032	371.08	77.65
	1045	4.75	5.23
35	1115	57.70	20.08
	1116	76.77	23.16
	1117	55.45	9.87
	1118	48.58	-1.94
	1030	50.43	15.72
40	1061	361.15	165.87
	1119	95.70	29.16
	1121	1057.19	263.21
	1122	600.58	262.34
45	1124	23.24	24.69
	1125	186.13	65.40
	1036	763.59	185.30
	1044	37.00	12.39
	1055	19.95	ND
50	1126	235.13	23.29
	1132	745.91	244.91
	1133	15.36	-1.62
	1136	31.47	12.88
55	1137	314.99	258.50
	1138	11.33	0.01
	1140	11.98	-0.62
	1013	115.8	100.20

EP 1 745 287 B1

(continued)

SLE Patient	C4d Level (SMFC)	C3d Level (SMFC)
1082	339.3	49.31
1048	5.07	-2.48
1060	5.96	3.59
1141	5.85	-2.56
1142	17.23	-3.95
1143	1.84	0.60
1080	49.98	32.36
1144	-1.76	-0.05
1145	-0.63	-0.67
1146	82.63	15.68
1147	497.35	227.98
1037	1.71	-1.42
1150	202.73	101.35

Example 3: Assay of Lymphocyte C4d/C3d for Distinguishing Patients with SLE from Patients with Other Diseases

[0067] These studies of patients with SLE vs. healthy controls were followed by studies to compare patients with SLE with patients diagnosed with diseases other than SLE (n=31; patients with rheumatoid arthritis, scleroderma, or inflammatory myositis). A single determination of lymphocyte C4d/C3d was made, using the same assay (Table 4). The mean values of C4d and C3d for patients with SLE, as compared with patients with other diseases are shown in Table 3. Whereas the mean value for C4d and C3d in patients with other diseases were 29.60 and 12.73, respectively, the mean value for C4d and C3d among patients with SLE was 201.06 and 62.42, respectively (both p< 0.0001).

Table 3. Analysis of Lymphocyte C4d and C3d Levels

	C4d Level (SMFC) ^a mean		C3d Level (SMFC) ^b mean	
	+/- SD	range	+/- SD	range
SLE	201.06 +/- 313.56	-2.62 - 1057.19	62.42 +/- 89.25	-3.95 - 318.18
Other Diseases	29.60 +/- 55.13	0 - 263.95	12.73 +/- 29.07	-4.95 - 141.82
Healthy Control	17.02 +/- 14.86	1.25 - 58.23	0.52 +/- 1.94	-3.42 - 2.67

*Leukocytes stained with FITC-anti-CD3 and PE-anti-C4d or PE-anti-C3d were subjected to 2-color flow cytometric analysis. CD3+ T lymphocytes were electronically gated and analyzed for the levels of C4d and C3d deposited on the surface.

a: Specific mean fluorescence intensity (SMFC) of the C4d levels detected on CD3+ T lymphocytes

b: SMFC of the C3d levels detected on CD3+ T lymphocytes

Table 4. Levels of C4d and C3d on the Surface of Lymphocytes from Patients with Other Diseases

Other Diseases	C4d Level (SMFC)	C3d Level (SMFC)
17001	2.88	0.24
13010	71.03	10.62
3042	10.11	4.91
4001	8.17	-0.92
5001	15.83	-0.12
6013	20.37	19.91
6008	10.88	-4.95
6014	19.17	4.38
3022	12.92	-4.60
4025	26.00	7.2
13032	25.41	56.25

EP 1 745 287 B1

(continued)

	Other Diseases	C4d Level (SMFC)	C3d Level (SMFC)
	6017	43.70	26.23
5	4033	11.97	3.44
	4028	173.06	7.98
	6011	9.11	2.20
	18002	15.33	8.38
10	4002	85.99	16.51
	8021	4.98	11.22
	15005	1.41	-1.71
	3029	10.33	0.00
	3030	4.41	-3.04
15	3031	263.95	141.82
	3032	0.00	-3.12
	3034	7.37	70.23
	3035	14.01	9.63
20	4026	0.42	-0.24
	6002	2.07	-0.52
	6008	10.93	2.76
	6015	3.98	0.70
	4030	21.01	3.12
25	15003	10.88	0.76

Example 4: Assay of Lymphocyte C4d for Distinguishing Patients with SLE

30 **[0068]** The complement pathway component C4d is deposited specifically on peripheral blood T lymphocytes of patients with SLE. Deposition of C4d on peripheral blood T lymphocytes was determined by a 2-color flow cytometric assay. T lymphocytes were identified using a FITC-conjugated monoclonal antibody specific for CD3 (a surface marker for T lymphocytes), and C4d deposited on these cells was determined using a monoclonal anti-C4d antibody followed by a PE-conjugated secondary antibody. Results are shown in Figure 1. Data shown are C4d-specific mean fluorescence (SMF) of peripheral blood T lymphocytes derived from patients with SLE (n=87), patients with other autoimmune diseases (n=31), or healthy controls (n=19). A cutpoint of specific mean fluorescence was empirically determined to distinguish individuals with C4d-positive T lymphocytes (SMF > 58) from those with C4d-negative T lymphocytes (SMF < 58). While none of the healthy controls had C4d levels above the cutpoint (0/19), almost half of the SLE patients had C4d levels above the cutpoint (42/87), and some of the patients with other autoimmune diseases had C4d levels above the cutpoint (4/31). The frequencies of individuals with C4d-positive T lymphocytes among the groups was compared using the Chi-square test, and the p values for each pair compared are shown above the horizontal line. The mean value of C4d-specific fluorescence on T lymphocytes among the groups was compared using the Students' T test and the respective p values are shown below the horizontal line. Figure 6 presents this data with a logarithmic scale on the X axis.

Example 5: Assay of Lymphocyte C3d for Distinguishing Patients with SLE and other inflammatory diseases.

45 **[0069]** The complement pathway component C3d is deposited specifically on peripheral blood T lymphocytes of patients with SLE and on patients with other inflammatory diseases. Deposition of C3d on peripheral blood T lymphocytes was determined by a 2-color flow cytometric assay. T lymphocytes were identified using a FITC-conjugated monoclonal antibody specific for CD3 (a surface marker for T lymphocytes), and C3d deposited on these cells was determined using a monoclonal anti-C4d antibody followed by a PE-conjugated secondary antibody. Results are shown in Figure 2. Data shown are C3d-specific mean fluorescence of peripheral blood T lymphocytes derived from patients with SLE (n=87), patients with other autoimmune diseases (n=31), or healthy controls (n=19). A cutpoint of specific mean fluorescence was empirically determined to distinguish individuals with C3d-positive T lymphocytes (SMFC > 3) from those with C3d-negative T lymphocytes (SMFC < 3). While none of the healthy controls had C3d levels above the cutpoint (0/19), almost 75% of the SLE patients had C3d levels above the cutpoint (65/87), and about half of the patients with other autoimmune diseases had C3d levels above the cutpoint (15/31). The frequencies of individuals with C4d-positive T lymphocytes among the groups was compared using the Chi-square test, and the p values for each pair compared are shown above the horizontal line. The mean value of C3d-specific fluorescence on T lymphocytes among the groups was compared

using the Students' T test and the respective p values are shown below the horizontal line. Figure 7 presents this data with a logarithmic scale on the X axis.

Example 6: Summary of assays using T-lymphocytes and comparison to assays using B lymphocytes or monocytes.

[0070] Complement ligands C4d and C3d are deposited specifically on peripheral blood T lymphocytes of patients with SLE. Figure 3 shows deposition of C3d and C4d on peripheral blood T lymphocytes that was determined by a 2-color flow cytometric assay. T lymphocytes were identified using a FITC-conjugated monoclonal antibody specific for CD3 (a surface marker for T lymphocytes), and C3d/C4d deposited on these cells was determined using a monoclonal anti-C3d (or anti-C4d) antibody, followed by a PE-conjugated secondary antibody. Data shown are C3d- and C4d-specific median fluorescence of peripheral blood T lymphocytes derived from patients with SLE (n=87), patients with other autoimmune diseases (n=31), or healthy controls (n=19). Cutpoints of specific mean fluorescence were empirically determined to distinguish individuals with C3d-positive T lymphocytes (SMFC > 3) or individuals with C4d-positive T lymphocytes (SMFI >58) from those with C3d-negative or C4d-negative T lymphocytes. The mean value of C3d specific fluorescence on T lymphocytes among the groups was compared using the Students' T test and the respective p values are <0.0001 (not shown). C4d levels were significantly higher on T lymphocytes from patients with SLE, as compared to patients with other diseases or as compared to healthy controls. C3d levels were also significantly higher on T lymphocytes from patients with SLE, as compared to patients with other diseases or as compared to healthy controls.

[0071] C4d levels were determined on the surface of T lymphocytes, B lymphocytes, and monocytes of patients with SLE, patients with other diseases, and healthy controls. C4d on different types of cells were determined by a 3-color flow cytometric assay using monoclonal antibodies specific for cell-specific surface markers and C4d or isotype control immunoglobulins. Levels of C4d were calculated as specific median fluorescence intensity (SMFI)= anti-C4d median fluorescence intensity - isotype Ig median fluorescence intensity. T cells were identified by electronic gating of cells positively stained by a monoclonal anti-CD3 antibody. B cells were identified by electronic gating of cells positively stained by a monoclonal anti-CD 19 antibody. Monocytes were identified by forward and side scattering and negative staining by anti- Cd3. Patients with other inflammatory diseases such as inflammatory myopathies, Sjogren's syndrome, vasculitis, Raynaud's phenomenon, and cardiovascular disease. eStudent t test; patients with SLE vs. patients with other diseases. [C4d levels were significantly higher on T lymphocytes, B lymphocytes, and monocytes from patients with SLE, as compared to patients with other diseases or as compared to healthy controls.

[0072] C3d levels were determined on the surface of T lymphocytes, B lymphocytes, and monocytes of patients with SLE, patients with other diseases, and healthy controls. C3d on different types of cells were determined by a 3-color flow cytometric assay using monoclonal antibodies specific for cell-specific surface markers and C3d or isotype control immunoglobulins. Levels of C4d were calculated as specific median fluorescence intensity (SMFI)= anti-C4d median fluorescence intensity - isotype Ig median fluorescence intensity. T cells were identified by electronic gating of cells positively stained by a monoclonal anti-CD3 antibody. B cells were identified by electronic gating of cells positively stained by a monoclonal anti-CD19 antibody. Monocytes were identified by forward and side scattering and negative staining by anti-Cd3. Patients with other inflammatory diseases such as inflammatory myopathies, Sjogren's syndrome, vasculitis, Raynaud's phenomenon, and cardiovascular disease. C3d levels were also significantly higher on T lymphocytes, B lymphocytes, and monocytes from patients with SLE, as compared to patients with other diseases or as compared to healthy controls.

[0073] From the figures and data, it can be seen that complement activation participates in a broad range of normal and abnormal inflammatory and immune processes. Therefore, abnormal patterns of complement activation products and complement receptors on white blood cells are useful in the diagnosis and/or monitoring of inflammatory and immune diseases other than systemic lupus erythematosus. The data in Figures 1-4 support this. Although the highest levels of C3d and C4d on peripheral blood T cells occurs in patients with SLE, abnormal levels of C3d are detected on T cells obtained from 15/31 (48.4%) of patients with other diseases. In addition, mean levels of C3d on T cells obtained from patients with other disease are significantly higher than mean levels of C3d on T cells obtained from healthy controls (p=0.027).

Claims

1. A method for diagnosing or monitoring systemic lupus erythematosus (SLE) comprising:

- (a) determining, in a blood sample from the individual the level of the complement pathway component C4d on the surface of a T lymphocyte or a B lymphocyte in the sample, and
- (b) comparing said determinations with, respectively, the level of the complement pathway component C4d on the surface of a control T lymphocyte or B lymphocyte, wherein an increase in the level of complement component

EP 1 745 287 B1

C4d indicates that the individual has systemic lupus erythematosus (SLE).

2. The method of claim 1, wherein the white blood cells are isolated using an anti-CD3 antibody.
- 5 3. The method of claim 1 or 2, wherein the level of complement component C4d is determined using an antibody specific for complement component C4d.
4. The method of claim 3, wherein the C4d antibody is labelled.
- 10 5. The method of claim 3, wherein the C4d antibody is a monoclonal antibody.
6. The method of any one of the preceding claims, wherein the level of at least one other complement component is determined.
- 15 7. The method of claim 6, wherein the level of complement component C3d is determined.
8. The method of claim 7, wherein the level of complement component C3d is determined using an antibody specific for complement component C3d.
- 20 9. The method of claim 8, wherein the C3d antibody is labelled.
10. The method of claim 8, wherein the C3d antibody is a monoclonal antibody.
- 25 11. The method of claim 1 comprising:
 - (a) determining, in a blood sample from the individual the level of the complement pathway component C4d on the surface of a T lymphocyte in the sample, and
 - (b) comparing said determinations with, respectively, the level of the complement pathway component C4d on the surface of a control T lymphocyte, wherein an increase in the level of complement component C4d indicates that the individual has systemic lupus erythematosus (SLE).
- 30 12. The method of claim 1 comprising:
 - (a) determining, in a blood sample from the individual the level of the complement pathway component C4d on the surface of a B lymphocyte in the sample, and
 - (b) comparing said determinations with, respectively, the level of the complement pathway component C4d on the surface of a control B lymphocyte, wherein an increase in the level of complement component C4d indicates that the individual has systemic lupus erythematosus (SLE).
- 35 13. A method according to any one of claims 1 to 10 wherein said determining and comparing is carried out by a computer program, said computer program comprising:
 - (a) code for receiving data corresponding to a determination of a complement component C4d deposited on the surface of a T lymphocyte or a B lymphocyte;
 - 45 (b) code for retrieving a reference value for the level of the complement component C4d deposited on the surface of a T lymphocyte or a B lymphocyte of individuals;
 - (c) code for comparing the data in (a) with the reference value in (b).
- 50 14. A computer program which, when executed by a computer carries out a method according to any one of claims 1 to 10, said computer program comprising:
 - (a) code for receiving data corresponding to a determination of a complement component C4d deposited on the surface of a T lymphocyte or a B lymphocyte;
 - (b) code for retrieving a reference value for the level of the complement component C4d deposited on the surface of a T lymphocyte or a B lymphocyte of individuals;
 - 55 (c) code for comparing the data in (a) with the reference value in (b).
15. A computer readable medium storing a computer program according to claim 14.

Patentansprüche

1. Methode zum Diagnostizieren oder Überwachen von systemischem Lupus erythematoses (SLE), umfassend:

5 (a) Bestimmen in einer Blutprobe von einem Individuum des Niveaus der C4d-Komponente des Komplementaktivierungswegs auf der Oberfläche einer T-Lymphozyte oder einer B-Lymphozyte in der Probe, und
(b) Vergleichen dieser Bestimmungen mit dem Niveau der C4d-Komponente des Komplementaktivierungswegs auf der Oberfläche einer Kontroll-T-Lymphozyte oder -B-Lymphozyte, wobei eine Erhöhung des Niveaus von Komplementkomponente C4d darauf hindeutet, dass das Individuum unter systemischem Lupus erythematoses
10 (SLE) leidet.

2. Methode nach Anspruch 1, wobei die weißen Blutkörperchen mit einem Anti-CD3-Antikörper isoliert werden.

3. Methode nach Anspruch 1 oder 2, wobei das Niveau von Komplementkomponente C4d mit einem für die Komplementkomponente C4d spezifischen Antikörper bestimmt wird.
15

4. Methode nach Anspruch 3, wobei der C4d-Antikörper markiert ist.

5. Methode nach Anspruch 3, wobei der C4d-Antikörper ein monoklonaler Antikörper ist.
20

6. Methode nach einem der vorstehenden Ansprüche, wobei das Niveau mindestens einer anderen Komplementkomponente bestimmt wird.

7. Methode nach Anspruch 6, wobei das Niveau der Komplementkomponente C3d bestimmt wird.
25

8. Methode nach Anspruch 7, wobei das Niveau der Komplementkomponente C3d mit einem C3d-spezifischen Antikörper bestimmt wird.

9. Methode nach Anspruch 8, wobei der C3d-Antikörper markiert ist.
30

10. Methode nach Anspruch 8, wobei der C3d-Antikörper ein monoklonaler Antikörper ist.

11. Methode nach Anspruch 1, umfassend:

35 (a) Bestimmen in einer Blutprobe von einem Individuum des Niveaus der C4d-Komponente des Komplementaktivierungswegs auf der Oberfläche einer T-Lymphozyte in der Probe, und
(b) Vergleichen dieser Bestimmungen mit dem Niveau der C4d-Komponente des Komplementaktivierungswegs auf der Oberfläche einer Kontroll-T-Lymphozyte, wobei eine Erhöhung des Niveaus der Komplementkomponente C4d, darauf hindeutet, dass das Individuum unter systemischem Lupus erythematoses (SLE) leidet.
40

12. Methode nach Anspruch 1, umfassend:

(a) Bestimmen in einer Blutprobe von einem Individuum des Niveaus der C4d-Komponente des Komplementaktivierungswegs auf der Oberfläche einer B-Lymphozyte in der Probe, und
45 (b) Vergleichen dieser Bestimmungen mit dem Niveau der C4d-Komponente des Komplementaktivierungswegs auf der Oberfläche einer Kontroll-B-Lymphozyte, wobei eine Erhöhung des Niveaus der Komplementkomponente C4d darauf hindeutet, dass das Individuum unter systemischem Lupus erythematoses (SLE) leidet.

13. Methode nach einem der Ansprüche 1 bis 10, wobei das Bestimmen und Vergleichen von einem Computerprogramm ausgeführt wird, wobei das Computerprogramm umfasst:
50

(a) Code zum Empfangen von Daten, die einer Bestimmung einer auf der Oberfläche einer T-Lymphozyte oder B-Lymphozyte aufgebracht C4d-Komplementkomponente entsprechen;
(b) Code zum Gewinnen eines Referenzwerts des Niveaus von auf der Oberfläche einer T-Lymphozyte oder
55 B-Lymphozyte von Individuen aufgebracht C4d-Komplementkomponente;
(c) Code zum Vergleichen der unter (a) erhaltenen Daten mit dem Referenzwert unter (b).

14. Computerprogramm, das, wenn von einem Computer ausgeführt, eine Methode nach einem der Ansprüche 1 bis

10 ausführt, wobei das Computerprogramm umfasst:

- 5 (a) Code zum Empfangen von Daten, die einer Bestimmung einer auf der Oberfläche einer T-Lymphozyte oder B-Lymphozyte aufgetragenen C4d-Komplementkomponente entsprechen;
(b) Code zum Gewinnen eines Referenzwerts des Niveaus von auf der Oberfläche einer T-Lymphozyte oder B-Lymphozyte von Individuen aufgetragenen C4d-Komplementkomponente;
(c) Code zum Vergleichen der unter (a) erhaltenen Daten mit dem Referenzwert unter (b).

10 15. Computerlesbares Medium, das ein Computerprogramm nach Anspruch 14 speichert.

Revendications

15 1. Procédé pour diagnostiquer ou surveiller un lupus érythémateux disséminé (LED) comprenant :

- 20 (a) la détermination, dans un échantillon de sang d'un individu, du taux du composant C4d des voies du complément à la surface d'un lymphocyte T ou d'un lymphocyte B dans l'échantillon, et
(b) la comparaison desdites déterminations avec, respectivement, le taux de composant C4d des voies du complément à la surface d'un lymphocyte T ou d'un lymphocyte B témoin, une augmentation du taux du composant C4d du complément indiquant que l'individu a un lupus érythémateux disséminé (LED).

2. Procédé de la revendication 1, dans lequel les globules blancs sont isolés au moyen d'un anticorps anti-CD3.

25 3. Procédé de la revendication 1 ou 2, dans lequel le taux du composant C4d du complément est déterminé au moyen d'un anticorps spécifique pour le composant C4d du complément.

4. Procédé de la revendication 3, dans lequel l'anticorps C4d est marqué.

30 5. Procédé de la revendication 3, dans lequel l'anticorps C4d est un anticorps monoclonal.

6. Procédé d'une quelconque des revendications précédentes, dans lequel le taux d'au moins un autre composant du complément est déterminé.

35 7. Procédé de la revendication 6, dans lequel le taux du composant C3d du complément est déterminé.

8. Procédé de la revendication 7, dans lequel le taux du composant C3d du complément est déterminé au moyen d'un anticorps spécifique pour le composant C3d du complément.

40 9. Procédé de la revendication 8, dans lequel l'anticorps C3d est marqué.

10. Procédé de la revendication 8, dans lequel l'anticorps C3d est un anticorps monoclonal.

45 11. Procédé de la revendication 1 qui comprend :

- (a) la détermination, dans un échantillon de sang d'un individu, du taux de composant C4d des voies du complément à la surface d'un lymphocyte T dans l'échantillon, et
(b) la comparaison desdites déterminations avec, respectivement, le taux de composant C4d des voies du complément à la surface d'un lymphocyte T témoin, **caractérisé en ce qu'**une augmentation du taux du composant C4d du complément indique que l'individu a un lupus érythémateux disséminé (LED).

50 12. Procédé de la revendication 1 qui comprend :

- 55 (a) la détermination, dans un échantillon de sang d'un individu, du taux de composant C4d des voies du complément à la surface d'un lymphocyte B dans l'échantillon, et
(b) la comparaison desdites déterminations avec, respectivement, le taux de composant C4d des voies du complément à la surface d'un lymphocyte B témoin, **caractérisé en ce qu'**une augmentation du taux du composant C4d du complément indique que l'individu a un lupus érythémateux disséminé (LED).

EP 1 745 287 B1

13. Procédé d'après une quelconque des revendications 1 à 10, dans lequel ladite détermination et ladite comparaison sont réalisées par un programme informatique, ledit programme informatique comprenant :

- 5
- (a) un code pour recevoir des données qui correspondent à une détermination d'un composant C4d de complément déposé à la surface d'un lymphocyte T ou d'un lymphocyte B ;
 - (b) un code pour extraire une valeur de référence pour le taux d'un composant C4d de complément déposé à la surface d'un lymphocyte T ou d'un lymphocyte B d'individus ;
 - (c) un code pour comparer les données de (a) avec la valeur de référence de (b).

10 14. Programme informatique qui, lorsqu'il est exécuté par un ordinateur applique un procédé d'après une quelconque des revendications 1 à 10, ledit programme informatique comprenant :

- 15
- (a) un code pour recevoir des données qui correspondent à une détermination d'un composant C4d de complément déposé à la surface d'un lymphocyte T ou d'un lymphocyte B ;
 - (b) un code pour extraire une valeur de référence pour le taux d'un composant C4d de complément déposé à la surface d'un lymphocyte T ou d'un lymphocyte B d'individus ;
 - (c) un code pour comparer les données de (a) avec la valeur de référence de (b).

20 15. Médium lisible par ordinateur qui stocke un programme informatique d'après la revendication 14.

25

30

35

40

45

50

55

Deposition of C4d on Peripheral Blood T Lymphocytes




FIG. 1




FIG. 2




FIG. 3

	T cell ^a -C4d Mean +/- SD (range)	p value	B cell ^b -C4d Mean +/- SD (range)	p value	Monocyte ^c - C4d Mean +/- SD (range)	p value
SLE (n=125)	13.89 +/- 26.50 (0 - 154.0)		62.55 +/-0 98.12 (0 - 66.30)		11.1 +/- 13.08 (0.58 - 60.31)	
Other Diseases ^d (n=45)	1.44 +/- 1.53 (0 - 7.62)	<0.0001 ^e	11.27 +/- 9.98 (0 - 44.15)	<0.0001 ^e	3.83 +/- 2.81 (0 - 13.59)	<0.0001 ^e
Healthy Controls (n=20)	1.40 +/- 0.91 (0.06 - 4.12)	<0.0001 ^f	7.67 +/- 5.10 (2.39 - 25.32)	<0.0001 ^f	3.86 +/- 2.40 (1.26 - 10.75)	<0.0001 ^f

FIG. 4

	T cell ^a -C4d Mean +/- SD (range)	p value	B cell ^b -C4d Mean +/- SD (range)	p value	Monocyte ^c - C4d Mean +/- SD (range)	P value
SLE (n=125)	3.26 +/- 5.87 (0 - 52.70)		18.16 +/- 12.37 (0 - 70.27)		2.88 +/- 2.69 (0 - 27.33)	
Other Diseases ^d (n=45)	0.94 +/- 1.07 (0 - 5.71)	<0.00003 ^e	13.47 +/- 12.01 (0 - 56.38)	0.36 ^e	1.96 +/- 1.34 (0 - 6.03)	<0.004 ^e
Healthy Controls (n=20)	0.78 +/- 0.45 (0 - 1.67)	<0.0001 ^f	7.22 +/- 2.90 (3.91 - 15.35)	<0.0001 ^f	2.4 +/- 1.39 (0.17 - 6.13)	<0.038 ^f

FIG. 5




FIG. 6




FIG. 7

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 03022223 A [0019] [0049]
- US 4946778 A [0035]
- US 4816567 A [0035]
- US 5545807 A [0035]
- US 5545806 A [0035]
- US 5569825 A [0035]
- US 5625126 A [0035]
- US 5633425 A [0035]
- US 5661016 A [0035]
- WO 9308829 A [0035]
- US 4676980 A [0035]
- WO 9100360 A [0035]
- WO 92200373 A [0035]
- EP 03089 A [0035]
- US SN60463447 A [0049]

Non-patent literature cited in the description

- **Marquart et al.** *Clin. Exp. Immunol.*, 1995, vol. 101, 60-65 [0008]
- **Manzi et al.** *Lupus*, 2004, vol. 13, 298-303 [0009]
- **Katyal et al.** *Current Science*, 2001, vol. 81 (8), 907-914 [0010]
- **van den Elsen et al.** *J. Mol. Biol.*, 2002, vol. 332, 1103-1115 [0011]
- **Sim ; Tsiftoglou.** *Biochem. Soc. Trans.*, 2004, vol. 32, 21-27 [0026]
- **Cohen, AS et al.** *Bull.Rheum.Dis.*, 1971, vol. 21, 643-648 [0030]
- **Tan, EM et al.** *Arth.Rheum.*, 1982, vol. 25, 1271-1277 [0030]
- **Hochberg, MC.** *Arth.Rheum.*, 1997, vol. 40, 1725 [0030]
- *Fundamental Immunology*. 1993 [0034]
- **McCafferty et al.** *Nature*, 1990, vol. 348, 552-554 [0034] [0035]
- **Kohler ; Milstein.** *Nature*, 1975, vol. 256, 495-497 [0035]
- **Kozbor et al.** *Immunology Today*, 1983, vol. 4, 72 [0035]
- **Cole et al.** *Monoclonal Antibodies and Cancer Therapy*. Alan R. Liss, Inc, 1985, 77-96 [0035]
- **Coligan.** *Current Protocols in Immunology*. 1991 [0035]
- **Harlow ; Lane.** *Antibodies, A Laboratory Manual*. 1988 [0035] [0037]
- **Goding.** *Monoclonal Antibodies: Principles and Practice*. 1986 [0035]
- **Kuby.** *Immunology*. 1997 [0035]
- **Marks et al.** *Bio/Technology*, 1992, vol. 10, 779-783 [0035]
- **Lonberg et al.** *Nature*, 1994, vol. 368, 856-859 [0035]
- **Morrison.** *Nature*, 1994, vol. 368, 812-13 [0035]
- **Fishwild et al.** *Nature Biotechnology*, 1996, vol. 14, 845-51 [0035]
- **Neuberger.** *Nature Biotechnology*, 1996, vol. 14, 826 [0035]
- **Lonberg ; Huszar.** *Intern. Rev. Immunol.*, 1995, vol. 13, 65-93 [0035]
- **Marks et al.** *Biotechnology*, 1992, vol. 10, 779-783 [0035]
- **Traunecker et al.** *EMBO J.*, 1991, vol. 10, 3655-3659 [0035]
- **Suresh et al.** *Methods in Enzymology*, 1986, vol. 121, 210 [0035]
- **Rogers, J., N. Cooper et al.** *PNAS*, 1992, vol. 89, 10016-10020 [0052]
- **Schwab, C. et al.** *Brain. Res.*, 1996, vol. 707 (2), 196 [0052]
- **Gemmell, C. J.** *Biomed Mater. Res.*, 1997, vol. 37, 474-480 [0052]
- **Stoltzner, S.E. et al.** *Am. J. Path.*, 2000, vol. 156, 489-499 [0052]

专利名称(译)	通过测量白细胞上的补体成分来诊断和监测炎症性疾病		
公开(公告)号	EP1745287B1	公开(公告)日	2010-09-22
申请号	EP2005747945	申请日	2005-05-11
[标]申请(专利权)人(译)	匹兹堡大学		
申请(专利权)人(译)	匹兹堡大学		
当前申请(专利权)人(译)	匹兹堡大学 - 英联邦系统的高等教育的		
[标]发明人	AHEARN JOSEPH M MANZI SUSAN M LIU CHAU CHING		
发明人	AHEARN, JOSEPH M. MANZI, SUSAN M. LIU, CHAU-CHING		
IPC分类号	G01N33/53 G01N33/569		
CPC分类号	G01N33/56972 G01N33/564 G01N2333/4716 Y10S435/967 Y10S435/973 Y10S436/811 Y10S436/821 Y10T436/101666		
优先权	60/570406 2004-05-11 US		
其他公开文献	EP1745287A4 EP1745287A1		
外部链接	Espacenet		

摘要(译)

本发明涉及通过确定白细胞表面上补体途径的组分水平来监测免疫学，血液学和炎症性疾病或状况的方法。

