

(19)



(11)

EP 1 964 931 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
29.12.2010 Bulletin 2010/52

(51) Int Cl.:
C12Q 1/68^(2006.01) G01N 33/53^(2006.01)
G01N 33/00^(2006.01)

(21) Application number: **08008370.2**

(22) Date of filing: **08.06.2006**

(54) **Method for predicting immune response to neoplastic disease based on mRNA expression profile in neoplastic cells and stimulated leukocytes**

Verfahren zur Vorhersage der Immunreaktion auf neoplastische Krankheiten auf der Grundlage des mRNA-Ausdrucksprofil in neoplastischen Zellen und stimulierten Leukozyten

Procédé de prédiction de réponse immune pour maladie néoplastique basée sur un profil d'expression ARNm dans des cellules néoplastiques et leucocytes stimulés

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR

(30) Priority: **08.06.2005 US 688744 P**
11.11.2005 US 735508 P

(43) Date of publication of application:
03.09.2008 Bulletin 2008/36

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
06772657.0 / 1 904 657

(73) Proprietors:
• **Hitachi Chemical Research Center, Inc.**
Irvine CA 92617 (US)
• **Hitachi Chemical Company, Ltd.**
Tokyo 163-0449 (JP)

(72) Inventor: **Mitsubishi, Masato**
Irvine, CA 92614 (US)

(74) Representative: **Elsy, David et al**
Withers & Rogers LLP
Goldings House
2 Hays Lane
London SE1 2HW (GB)

(56) References cited:
WO-A-03/099312 WO-A-2005/044792
WO-A-2006/110091

- **SCHIMANSKI CARL C ET AL: "Effect of chemokine receptors CXCR4 and CCR7 on the metastatic behavior of human colorectal cancer." CLINICAL CANCER RESEARCH : AN OFFICIAL JOURNAL OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH 1 MAR 2005, vol. 11, no. 5, 1 March 2005 (2005-03-01), pages 1743-1750, XP002504646 ISSN: 1078-0432**
- **LEUNG SUET YI ET AL: "Expression profiling identifies chemokine (C-C motif) ligand 18 as an independent prognostic indicator in gastric cancer." GASTROENTEROLOGY AUG 2004, vol. 127, no. 2, August 2004 (2004-08), pages 457-469, XP002504647 ISSN: 0016-5085**
- **MORAN CHRISTOPHER J ET AL: "RANTES expression is a predictor of survival in stage I lung adenocarcinoma." CLINICAL CANCER RESEARCH : AN OFFICIAL JOURNAL OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH DEC 2002, vol. 8, no. 12, December 2002 (2002-12), pages 3803-3812, XP002504648 ISSN: 1078-0432**
- **LE YINGYING ET AL: "Chemokines and chemokine receptors: their manifold roles in homeostasis and disease." CELLULAR & MOLECULAR IMMUNOLOGY APR 2004 LNKD-PUBMED:16212895, vol. 1, no. 2, April 2004 (2004-04), pages 95-104, ISSN: 1672-7681**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 964 931 B1

Description

BACKGROUND OF THE INVENTION

5 Field of the Invention

[0001] The present invention relates to a method for predicting the immune response in a mammal to neoplastic disease based on the expression profile of tumor necrosis factor receptor (TNF-R) superfamily mRNA in neoplastic tissue, and of tumor necrosis factor (TNF) superfamily mRNA in circulating leukocytes. In the method, samples of
10 neoplastic tissue are obtained and the TNF-R superfamily subtype mRNAs expressed in those tissues are assessed. Furthermore, whole blood of the mammal is subjected to a stimulus that activates T-cells in the blood and the TNF superfamily subtype mRNAs that exhibit a significant change in expression level in response to the stimulus are identified. Individuals that exhibit a change in expression level in TNF superfamily subtypes that correlate with the TNF-R superfamily subtypes expressed in their tumor tissue are determined to have a likely lesser severity of prognosis for their disease.

15 Description of the Related Art

[0002] Different modalities of cancer therapies may be compared to varying types of uses of force in society. Chemotherapy is like the widespread use of military force in a city; the overwhelming power of the weapons involved may
20 cause collateral damage to civilians. In contrast, leukocytes in circulating peripheral blood, which are the primary killers of cancer cells in the human body, are like police officers patrolling city streets who deal with street crime. Likewise, when leukocytes encounter cancer cells in the body, these cells are the initial responders to the cancer targets. Leukocytes are classified into many classes and subclasses based on morphological analysis and characterization of cell surface markers using flow cytometry or immunohistochemical staining techniques. These classes and subclasses are like
25 identifying police officers by their uniforms and identification badges. The number of leukocytes per mm³ of peripheral blood corresponds to the number of officers in the city.

[0003] When a police officer encounters street criminals, he must deal with them with the weapons he has at hand. But police officers are not always carrying appropriate weaponry. Similarly, cytotoxic T-cells are not always provided with the proper anti-tumor factors to combat specific cancer cells. Cytotoxic T-cells recognize cancer cells via IgG Fc
30 receptors (FcR γ), when the cancer cells are coated with IgG. This process is termed antibody-dependent cell-mediated cytotoxicity (ADCC). In fact, IgG is frequently recognized around cancer margins by staining with anti-human IgG (see, for example, Richman AV, Immunofluorescence studies of benign and malignant human mammary tissue, J. Natl. Cancer Inst. 1976;57:263-7, and Koneval T, et al., Demonstration of immunoglobulin in tumor and marginal tissues of squamous cell carcinomas of the head and neck, J. Natl. Cancer Inst. 1977;59:1089-97). Infiltration of mononuclear leukocytes
35 into cancer lesions is also found in many cases. The FcR γ on the surface of cytotoxic T-cells are like a bag containing an assortment of weapons that a police officer carries. Different types of FcR γ , such as CD 16, CD32, and CD64 correspond to different types of bags. Cudgels are an initial weapon, and are held ready to attack at any time. In cytotoxic T-cells, the cudgel corresponds to perforin (see Nakanishi et al., Perforin expression in lymphocytes infiltrated to human colorectal cancer, Br. J. Cancer 1991;64:239-42), which is pre-synthesized and stored in the cytosol of cytotoxic T-cells,
40 and immediately released upon FcR γ activation. Other presynthesized "cudgels" ready for use include granzymes, proteases related to the digestive enzymes trypsin and chymotrypsin, which may act to trigger apoptosis in the target cell.

[0004] Officers also carry more powerful guns, which may correspond to tumor necrosis factors (TNF) in cytotoxic T-cells. Usually, the "bullets" in the cellular context (TNF subtypes) are not loaded in the gun, and are synthesized and released from cytotoxic T-cells only upon Fc receptor activation. TNF is capable of inducing apoptosis by interacting
45 with specific TNF receptors present on the surface of target cells. In order to maintain killing activities against a broad spectrum of target cells, different types of TNF ligands exist (as part of a TNF superfamily, abbreviated to TNFSF). According to GenBank and UniGene information (<http://www.ncbi.nlm.nih.gov>), the human TNF superfamily encompasses up to TNF superfamily 18, with some missing numbers (16 and 17) and multiple sequences within the same number (13A and 13B), for a total of 17 human members. For the corresponding TNF receptors (the TNF-R superfamily:
50 abbreviated to TNFRSF), the human TNF-R superfamily encompasses up to TNF-R superfamily subtype 21. Although TNF/TNF-R superfamily subtype interactions are not strictly specific, each ligand generally reacts with a specific receptor, as shown in Table 1, and over 300 different TNF superfamily subtype TNF-R superfamily combinations exist.

55 Table I. List of GenBank UniGene entry of TNFRSF and TNFSF mRNA.

TNFRSF (UniGene #)	Corresponding TNFSF (UniGene #)
1A (Hs.279594)	2 (Hs.241570)

EP 1 964 931 B1

(continued)

	TNFRSF (UniGene #)	Corresponding TNFSF (UniGene #)
	1B (Hs.256278)	2 (Hs.241570)
5	3 (Hs.11116)	1 (Hs.36)
		3 (Hs.376208)
	4 (Hs.129780)	4 (Hs.181097)
	5 (Hs.472860)	5 (Hs.652)
10	6 (Hs.244139)	6 (Hs.2007)
	7 (Hs.355307)	7 (Hs.501497)
	8 (Hs.1314)	8 (Hs.494901)
	9 (Hs.193418)	9 (Hs.1524)
	10A (Hs.401745)	10 (Hs.478275)
15	10B (Hs.521456)	10 (Hs.478275)
	10C (Hs.119684)	10 (Hs.478275)
	10C (Hs.119684)	10 (Hs.478275)
	10D (Hs.213467)	10 (Hs.478275)
20	11A (Hs.204044)	11 (Hs.333791)
	11B (Hs.81791)	11 (Hs.333791)
	12A (Hs.355899)	12 (NM003809)*
	14 (Hs.512898)	14 (Hs.129708)
	17 (Hs.2556)	13 (Hs.54673)
25		13B (Hs.525157)
	18 (Hs.212680)	18 (Hs.248197)
	25 (Hs.462529)	15 (Hs.241382)

*:GeneBank accession number (no UniGene entry was found).

30 **[0005]** The complete eradication of cancers may happen when the appropriate TNF superfamily subtype is released from infiltrating cytotoxic T-cells. The rare cases of miracle survivors of cancers may be those in which the TNF superfamily subtype TNF-R superfamily subtype combination is perfect.

35 **[0006]** Antibody-dependent cell-mediated cytotoxicity (ADCC) is another mechanism involved in the cytolytic function of leukocytes in conjunction with antibodies (see Perussia et al., Assays for antibody-dependent cell-mediated cytotoxicity (ADCC) and reverse ADCC (redirected cytotoxicity) in human natural killer cells, Methods Mol Biol. 2000;121:179-92). Once the Fab portion of IgG binds to the target cells, the Fc portion of IgG activates Fc receptors on the leukocytes, followed by activation of these leukocytes to attack target cells. Although ADCC has been studied for many years, the conditions in *in vitro* experiments involving ADCC is dependent on the ratio between effector cytotoxic T-cells and target cells in a pure cell system suspended in an artificial solution. Moreover, because cytolysis was generally quantitated by the release of isotope (chromium-51) from the target cells, it was not known whether the observed cytolysis happens via perforin or TNF. It is also not known which TNF superfamily subtypes are involved in ADCC.

40 **[0007]** Finally, a further important mechanism by which leukocytes combat cancer is the release of chemokines to recruit more leukocytes to the disease site. Many such cytokines have been discovered. Although correlations have not been established between specific chemokines and specific populations of recruited leukocytes, the phenomenon of chemokine release and leukocyte recruitment is important in combating cancer. An analogy might be made here to policemen confronted with a gang of criminals, who radio back to headquarters for more reinforcements.

45 **[0008]** In sum, although complicated cellular mechanisms are required to educate appropriate subsets of leukocytes to be able to recognize specific cancer markers, and appropriate chemotactic factors should be released to attract these mature leukocytes to the lesion, actual cancer killing happens in two distinct ways. Immediate cancer killing takes place by the release of perforin from these leukocytes, followed by slow and sustained induction of apoptosis via the synthesis and release of TNF (see Vujanovic, Role of TNF family ligands in antitumor activity of natural killer cells, Int. Rev. Immunol. 2001 Jun;20(3-4):415-37). TNF induces apoptosis via binding to the specific receptors on the target cell surface. This means that leukocytes must release specific TNF ligand(s) corresponding to the receptors present on the cancer cells. Although the usefulness of TNF/TNF receptor ratios as predictors of disease outcome has been considered (see McDermott, TNF and TNFR biology in health and disease, Cell. Mol. Biol. (Noisy-le-grand). 2001 Jun;47(4):619-35), little is known about the profile of TNF receptors in cancer cells and induction of ligands in leukocytes, nor is it known how appropriate populations of leukocytes are recruited.

SUMMARY OF THE INVENTION

[0009] Schimanski C et al (Clin. Cancer Res. (2005) 11, 1743-1750) discloses the effect of CXCR4 and CCR7 on human colorectal cancer.

[0010] WO 03/099312 discloses methods for improving therapeutic treatments of cancer by controlling chemokines.

[0011] Leung S et al (Gastroenterology (2004) 127 457-469) discloses the identification of chemokine (C-C motif) ligand 18 as a prognostic indicator of cancer.

[0012] Moran C et al (Clin. Cancer Res. (2002) 8, 3803-3812) discloses RANTES as a predictor of survival of stage I lung carcinoma.

[0013] WO 2005/044792 discloses the role of CCX-CKR2 in cancer.

[0014] WO 2006/110091 discloses the use of cytokine-like IL6 in detection of cancer.

[0015] If a pathological specimen shows IgG deposition around cancer margins, this means that the patient is at least capable of raising cancer-recognizing IgG. If the infiltration of mononuclear leukocytes exists at the margin of the cancer mass, this means that the patient's leukocytes are somehow attracted toward the cancer lesion, probably due to the release of chemotactic factors (C3a, C5a, C567) (see Becker, The relationship of the chemotactic behavior of the complement-derived factors, C3a, C5a, and C567, and a bacterial chemotactic factor to their ability to activate the proesterase 1 of rabbit polymorphonuclear leukocytes, J. Exp. Med. 1972;135:376-87) during complement activation of cancer-bound IgG. The type of TNF-R superfamily subtypes expressed on the surface of cancer cells can be identified by, for example, immunostaining (if antibodies are available), in-situ hybridization, or in-situ PCR (sequences are available from GenBank). Identification of the TNF superfamily subtypes expressed in infiltrating mononuclear cells is difficult to interpret, because FcR γ may not be activated yet. Since leukocytes in peripheral blood are like reserve police officers, a supply of appropriately and fully equipped cytotoxic T-cells in peripheral blood is a critical first step for the battle against cancer. Thus, this method makes it possible to predict anti-cancer immunity in each patient by quantitating the inducibility of the appropriate TNF superfamily subtypes in peripheral blood leukocytes.

[0016] The invention provides a method of determining the likely severity of prognosis for a neoplastic disease in a mammal as defined in the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017]

Figure 1A shows the results of an analysis of TNF superfamily mRNAs induced in peripheral blood leukocytes after exposure of whole blood to heat-aggregated IgG (HAG).

Figure 1B shows the results of a repetition of the analysis of Figure 1A.

Figure 2 shows the results of an analysis of TNF-R superfamily mRNAs expressed in tissue samples taken from various adenocarcinomas and squamous cell carcinomas.

Figure 3 shows the results of an analysis of TNF superfamily mRNAs induced in peripheral blood leukocytes after exposure of whole blood to anti-human T-cell receptor (TCR) monoclonal antibody.

Figure 4A shows the results of an analysis of the dose response and kinetics when peripheral blood leukocytes are exposed to anti-TCR antibody and TNF superfamily subtype 2 mRNA expression is quantified.

Figure 4B shows the results of an analysis of the kinetics when peripheral blood leukocytes are exposed to anti-TCR antibody and TNF superfamily subtype 2 mRNA expression is quantified.

Figure 5A shows the results of an analysis of the dose response and kinetics when peripheral blood leukocytes are exposed to HAG and TNF superfamily subtype 15 mRNA expression is quantified.

Figure 5B shows the results of an analysis of the kinetics when peripheral blood leukocytes are exposed to HAG and TNF superfamily subtype 15 mRNA expression is quantified.

Figure 6 shows the results of a repetition of the analysis of Figure 5 with respect to several TNF superfamily subtype mRNAs and control mRNA.

Figure 7 shows the results of an analysis of TNF superfamily mRNAs induced in peripheral blood leukocytes after exposure of whole blood to HAG.

Figure 8 shows the chemokine primer sequences employed in Embodiment 4.

Figure 9 shows the results of an analysis of chemokine mRNAs induced in peripheral blood leukocytes after exposure of whole blood to HAG.

Figure 10 shows the results of an analysis of TNF superfamily subtype 3 mRNA induced in peripheral blood leukocytes after exposure to HAG of whole blood aliquots previously stimulated with either a dietary supplement or a control stimulus.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

Quantitation of mRNA

5 **[0018]** The present inventors have developed a unique technology known as the Hem(A)⁺ system that is capable of quantitating minute changes of gene expression in whole blood leukocytes. This technology is described in detail in US 20040265864 and International Patent Application WO 2005042784 (see also Mitsuhashi, Absolute quantitation of mRNA in human blood leukocytes as a model for phenotypic gene expression-based diagnostics, Clin Chem, 52:4 (2006)).

10 **[0019]** In the quantitation technique, whole blood is applied to 96-well filterplates to trap leukocytes. Lysis buffer containing both synthetic external control RNA and specific reverse primers is added to the filterplates, and cell lysates are transferred to oligo(dT)-immobilized microplates for hybridization. The cDNA is then synthesized in the oligo(dT)-immobilized microplates from these primer sites. The solution, which contains specific reverse primer-primed cDNA, is used for real-time PCR. The plate containing oligo(dT)-primed immobilized cDNA may be stored as a cDNA bank.

15 **[0020]** In more detail, the assay procedure consists of 3 major steps: (a) leukocyte isolation and lysis on filterplates; (b) mRNA isolation, reverse primer hybridization, and cDNA synthesis in oligo(dT)-immobilized microplates; and (c) real-time quantitative PCR. Custom 96-well filterplates may be manufactured, for example by Whatman or Pall, by assembly with leukocyte reduction membranes (Leukosorb; Pall). These filterplates are placed over collection plates, and 150 μ L of 5 mmol/L Tris (pH 7.4) is applied to wet the filter membranes. After centrifugation at 120g for 1 min at 4 °C to remove the Tris solution from the membranes, 50 μ L of well-mixed whole blood sample is applied to each well and immediately centrifuged at 120g for 2 min at 4 °C. The wells are then washed once with 300 μ L of phosphate-buffered saline. After centrifugation at 2000g for 5 min at 4 °C to remove the saline solution, 60 μ L of stock lysis buffer [5 g/L N-lauroylsarcosine, 4X standard saline citrate, 10 mmol/L Tris-HCl (pH 7.4), 1 mmol/L EDTA, 1 mL/L IGEPAL CA-630 (substitute of NP-40), 1.79 mol/L guanidine thiocyanate (all from Sigma)], supplemented with 1 mL/L 2-mercaptoethanol (Bio-Rad), 0.5 g/L proteinase K (Pierce), 0.1 g/L salmon sperm DNA (5 Prime Eppendorf/Brinkman), 0.1 g/L Escherichia coli tRNA (Sigma), 5 nmol/L each of the specific reverse primers, and 10⁹ molecules/L of synthetic RNA34 (as external control), are added to each well of the filterplates. The plates are then incubated at 37 °C for 10 min, placed over oligo(dT)-immobilized microplates (GenePlate; RNAture), and centrifuged at 2000g for 5 min at 4 °C. After overnight storage at 4 °C, the microplates are washed 3 times with 100 μ L of plain lysis buffer and then 3 times with 150 μ L of wash buffer [0.5 mol/L NaCl, 10 mmol/L Tris (pH 7.4) 1 mmol/L EDTA] at 4 °C.

20 **[0021]** cDNA can be synthesized directly in each well by addition of 30 μ L of buffer containing 1 x reverse transcription buffer [50 mM KC1, 10 mM Tris-HCl (pH 8.3), 5.5 mM MgCl₂, 1 mL/L Tween 20], 1.25 mM of each deoxynucleoside triphosphate, 4 units of rRNasin, and 80 U of MMLV reverse transcriptase (Promega; without primers) and incubation at 37 °C for 2 h. From each 30- μ L reaction, 4 μ L of cDNA is transferred directly to 384-well PCR plates, and 5 μ L of TaqMan universal master mixture (Applied Biosystems) and 1 μ L of oligonucleotide cocktail (5 μ M each of the forward and reverse primers, and 1-2 μ M TaqMan probe) are added. PCR may be carried out in a PRISM 7900HT (Applied Biosystems), with 1 cycle of 95 °C for 10 min followed by 45 cycles of 95 °C for 30 s, 55 °C for 30 s, and 60 °C for 1 min. Each gene is amplified in separate wells. The cycle threshold (Ct), i.e., the cycle at which certain amounts of PCR products (based on fluorescence) are generated, is determined with analytical software (SDS; Applied Biosystems).

25 **[0022]** To construct calibration curves for quantification, long synthetic DNA oligonucleotides containing sequences of the forward and reverse primers and TaqMan probes are synthesized for each target. The TaqMan probe for control RNA34, for example, is CCAAGGCCAGCCCTCACACA (SEQ ID No. 1). Alternatively, SYBRGreen PCR may be employed. Each oligonucleotide may be purified by HPLC to a purity of >95%. Each PCR contains 10-10⁶ molecules/well of these template oligonucleotides. The calibration curve for control RNA34 generated by DNA oligonucleotides may be used to convert the Ct values of the samples to molecules/PCR well (4 μ L of cDNA), which are then multiplied by 7.5 (30 divided by 4) to obtain molecules/sample (30 μ L of cDNA). The percentage recovery of RNA34 is obtained by dividing these values by the amounts of RNA34 in the original 60 μ L of lysis buffer (6 x 10⁵ = 10⁷/mL x 60 μ L). For native mRNA, the molecules/PCR well are determined as described above with the respective calibration curves, and then these values are converted to molecules per microliter of blood by multiplying by 7.5 (30 divided by 4), dividing by the percentage recovery of RNA34 in each sample, and dividing by the volume of blood added to each well of the filterplates (usually 50 μ L). The mean (SD) is calculated from triplicate aliquots of whole blood and the Student t-test may be used for statistical analyses.

30 **[0023]** In embodiments, the expression profile of TNF receptor superfamily (TNFRSF) mRNA was characterized among various cancer specimens by using the mRNA quantitation system (Hem(A)⁺) described above. Then the inducibility of TNF superfamily (TNFSF) mRNA was quantitated by stimulating heparinized whole blood with mouse anti-human T cell receptor monoclonal antibody or heat-aggregated IgG as *ex vivo* models of cytotoxic T-cell mediated reaction and antibody dependent cell cytotoxicity (ADCC), respectively. Significant induction or reduction in the expression levels was calculated using statistical methods well known in the art; a p value of 0.05 or less was employed. The

substantial individual-to-individual variation shown indicates that embodiments of the present invention may be employed in laying the groundwork for individualized cancer immunotherapy.

Example 1

5 [0024] The mRNA quantitation method described above was employed to quantitate all types of TNF superfamily mRNA, both with and without *ex vivo* stimulation using heat-aggregated IgG (HAG), which was employed as a model of immune complex stimulation (see Ostreiko et al., Production and characterization of heat-aggregated IgG complexes with predetermined molecular masses: light-scattering study, Immunol Lett. 1987;15:311-6). The immune complex is
10 formed when IgG molecules bound to a target cell bind with high avidity to Fc receptors on a leukocyte, causing cross-linking of the Fc receptors and generation of a leukocyte activation signal.

Triplicate aliquots of 50 μ L each of heparinized whole blood was incubated with 200 μ g/mL HAG at 37°C for 2 hours. Then blood samples were applied to 96-well filterplates to trap leukocytes, and cell lysates were transferred to 96-well oligo(dT)-immobilized microplates for isolation of poly(A)⁺ mRNA as described above. cDNA was synthesized on the
15 microplates and was used for real-time polymerase chain reaction (PCR) in 384-well plates by monitoring SYBR Green fluorescence (see Morrison et al., Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification, Biotechniques 1998;24:954-8, 960, 96). In brief, SYBR Green PCR was conducted by diluting cDNA 3-4 fold in water, and transferring 4 μ L cDNA solution directly to 384-well PCR plates, to which 5 μ L of a master mix (BioRad, Hercules, CA) and 1 μ L of oligonucleotide cocktail (15 μ M each of forward and reverse primer) were applied, and
20 conducting PCR in PRISM 7900HT (ABI), with one cycle of 95°C for 10 min followed by 45 cycles of 95°C for 30 sec and 60°C for 1 min. Each gene was amplified in separate wells.

[0025] The Ct was determined by analytical software (SDS, ABI). The PCR conditions were carefully optimized, so that a single appropriate peak was detected on melting curve analysis without any primer dimers. To validate the assay conditions, synthetic RNA34 was spiked into the lysis buffer, and this RNA34 was also quantitated by real time PCR.
25 The sequence specificity of primers was confirmed by blast analysis (<http://www.ncbi.nlm.nih.gov/BLAST>). The Ct was set equal to the PCR cycle where certain amounts of PCR products were generated, and delta Ct (Δ Ct) was also calculated by subtracting the Ct values of the un-stimulated samples from those of the stimulated samples. Since Ct is a log scale, 1 Δ Ct means double or one half in quantity, and a negative Δ Ct indicates an increase in expression.

[0026] The results are shown in Figures 1A and 1B. In the Figures, TNFSF mRNAs (1-18) and spiked control RNA34
30 were quantitated by SYBR Green real time PCR. Δ Ct values of 3, 2, 1, 0, -1, -2, and -3 indicate 1/8, 1/4, 1/2, 0, 2, 4, 8 fold increases, respectively. In Figure 1A, each symbol is the mean value from each individual. The \circ symbol indicates a significant increase ($p < 0.05$), while the \triangle symbol indicates a significant decrease ($p < 0.05$), and X indicates no change. Figure 1B shows the results of repeating the analysis twice in 2 consecutive days using the same individuals. The symbols are the mean \pm standard deviation from each individual. \bullet : TNFSF-2 (TNF α), \blacktriangle : TNFSF-8, \blacklozenge : TNFSF-15, respectively. The solid line is the line where two data sets are identical. The dotted lines indicate $\pm 0.5 \Delta$ Ct.
35

[0027] This technique was capable of detecting basal levels of all members of TNF superfamily mRNA in blood leukocytes. As shown in Figure 1A, significant induction of TNF superfamily mRNA was identified when whole blood was exposed to HAG. Although large individual-to-individual variation exists, the dominant TNF superfamily mRNAs were TNF superfamily subtypes 2 (TNF α), 8, and 15. These mRNAs exhibited a HAG-induced response in, respectively,
40 80% (12/15), 73% (11/15), and 60% (9/15) of the tested individuals. TNF superfamily subtypes 1, 3, 4, 6 (i.e., FasL), 7, 9, 10 (i.e., TRAIL), 14, and 18 mRNAs exhibited a HAG-induced response in at least one tested individual. TNF superfamily subtypes 5, 11, 12, 13, and 13B mRNAs exhibited no HAG-induced response in any of the tested individuals. At least one of TNF superfamily subtypes 2, 8 and 15 was induced by HAG exposure in every tested individual. For example, three individuals that exhibited no HAG-induced TNF superfamily subtype 2 response all exhibited a subtype 15 response,
45 whereas six individuals that exhibited no HAG-induced subtype 15 response exhibited a subtype 2 response. Four individuals in which TNF superfamily subtype 8 exhibited no HAG-induced response did exhibit a response in either subtype 2 or subtype 15. The Δ Ct of the control, spiked RNA34, was all within ± 0.5 , as shown in Figure 1A. In order to further validate these experiments, blood samples were obtained from the same individuals twice in 2 consecutive days. As shown in Figure 1B, the values for TNF superfamily subtypes 2, 8, 15 and RNA34 were reproducible within \pm
50 0.5 standard deviation. The variation within triplicate whole blood aliquots in each sample was also less than 0.5 Δ Ct (Figure 1B, X-Y bar in each symbol).

[0028] This method may be applied in the following manner in an embodiment. Specimens of cancers removed from mammalian patients undergoing surgical removal or biopsy are analyzed for IgG deposition, infiltration of mononuclear leukocytes, and the type of TNF-R superfamily subtypes expressed on cancer cells. Blood samples are then obtained
55 from the same individuals, and subjected to analysis to determine which types of TNF superfamily mRNA are induced by HAG stimulation and to what degree. Individuals are classified into a group in which the induced TNF superfamily and the expressed TNF-R match and a group in which they do not match. The results of a comparison of clinical outcomes in the 2 groups will reveal that the matching group exhibits a lesser severity of prognosis. If the cancer mass is small,

EP 1 964 931 B1

such as with early invisible metastatic lesions, appropriate TNF superfamily-mediated immune system attack may be sufficient to prevent cancer relapse. The clinical outcomes compared should thus include the length of remission, and frequency of relapse (metastasis).

5 Example 2

10 **[0029]** DNA sequences of TNF-R superfamily and TNF superfamily mRNA were downloaded from GenBank, as shown in Table 1, and the primer sequences shown in Table 2 below were designed by Primer Express (Applied Biosystems) and HYBsimulator (RNAture). These sequences were designed at common regions among multiple transcripts if such variants exist. No amplification happened without cDNA (no primer dimer), and a single peak was detected at melting curve analysis.

Table 2: Primer sequences used

Target mRNA	Forward	Reverse
15 TNFRSF-1A	CCTGCCAGGAGAAACAGAACA (SEQ ID No. 2)	GGAGACACACTCGTTTTCTCTTAGAA (SEQ ID No. 3)
TNFRSF-1B	CAAGCCAGCTCCACAATGG (SEQ ID No. 4)	TGACCGAAAGGCACATTCT (SEQ ID No. 5)
20 TNFRSF-3	CCTCCCGGGCTCTCTACAC (SEQ ID No. 6)	TCATGGGTGATAAATTGGTTCCT (SEQ ID No. 7)
TNFRSF-4	ACGACGTGGTCAGCTCCAA (SEQ ID No. 8)	GCGGCAGACTGTGTCCTGTGT (SEQ ID No. 9)
25 TNFRSF-5	GGCCAAGAAGCCAACCAATA (SEQ ID No. 10)	GAAGATCGTCGGGAAAATTGAT (SEQ ID No. 11)
TNFRSF-6	TGGCATCAACTTCATGGAAAGA (SEQ ID No. 12)	GCAAGAGTACAAAGATTGGCTTTTT (SEQ ID No. 13)
TNFRSF-7	CTGCAGAGCCTTGTCGTTACAG (SEQ ID No. 14)	GCTCCGGTTTTCGGTAATCC (SEQ ID No. 15)
30 TNFRSF-8	GGTTGAGGCAGCAAACAGATG (SEQ ID No. 16)	GCCTGGTGGTTAAGGTCTGATG (SEQ ID No. 17)
TNFRSF-9	CGTCGACTGCGTTGCTCTT (SEQ ID No. 18)	TTCTGCCCCGTTTAAACAACAG (SEQ ID No. 19)
35 TNFRSF-10A	TGAGGACAATGCTCACAACGA (SEQ ID No. 20)	TTGCTGCTCAGAGACGAAAGTG (SEQ ID No. 21)
TNFRSF-10B	CTGAGACAGTGCTTCGATGACTTT (SEQ ID No. 22)	CCATGAGGCCCAACTTCT (SEQ ID No. 23)
TNFRSF-10C	GGAAGTGTAGCAGGTGCCCTAGT (SEQ ID No. 24)	ACCAAATTCTTCAACACACTGGATAT (SEQ ID No. 25)
40 TNFRSF-10D	ATGGACTTACGAGGGTTCGACTTAG (SEQ ID No. 26)	GGAAAAGAGATGTACAGCCTACAGTAGTAAGC (SEQ ID No. 27)
TNFRSF-11A	GGAAACAGTAACTCCACGTTTCATCT (SEQ ID No. 28)	GCGAGGTCTGGCTGACGTA (SEQ ID No. 29)
45 TNFRSF-11B	TGCAAACCCAGTGACCAGATC (SEQ ID No. 30)	AAGGTGTCTTGGTCGCCATT (SEQ ID No. 31)
TNFRSF-12A	CGCTGATCCAGTGACAATGTG (SEQ ID No. 32)	GCGTCTGGGAGGCAGAGA (SEQ ID No. 33)
50 TNFRSF-14	CAGGGAGCCTCGTCATCGT (SEQ ID No. 34)	CACCCCTTGGCTTTCTTCTTT (SEQ ID No. 35)
TNFSF-17	GGAGGAAGGCACAACCAT (SEQ ID No. 36)	GCAGCTGGCAGGCTCTTG (SEQ ID No. 37)
TNFRSF-18	AGTTTTGGCTTCCAGTGTATCGA (SEQ ID No. 38)	GCAGTCTGTCCAAGGTTTGCA (SEQ ID No. 39)
55 TNFRSF-25	CCTGCTCGCCCCTATCG (SEQ ID No. 40)	TTCACCCCTCTCGACATTC (SEQ ID No. 41)
TNFSF-1	CAGCTATCCACCCACACAGATG (SEQ ID No. 42)	CGAAGGCTCCAAAGAAGACAGT (SEQ ID No. 43)

EP 1 964 931 B1

(continued)

Target mRNA	Forward	Reverse
5 TNFSF-2	CGAAGGCTCCAAAGAAGACAGT (SEQ ID No. 44)	CAGGGCAATGATCCCAAAGT (SEQ ID No. 45)
TNFSF-3	AGGGTGTACGTCAACATCAGTCA (SEQ ID No. 46)	CACGGCCCCAAAGAAGGT (SEQ ID No. 47)
TNFSF-4	GCCCCTCTTCCAACCTGAAGAA (SEQ ID No. 48)	GGTATTGTCAGTGGTCACATTCAAG (SEQ ID No. 49)
10 TNFSF-5	CCACAGTTCGCCAAACCT (SEQ ID No. 50)	CACCTGGTTGCAATTCAAATACTC (SEQ ID No. 51)
TNFSF-6	TGGCAGCATCTTCACTTCTA AATG (SEQ ID No. 52)	GAAATGAGTCCCCAAAACAT CTCT (SEQ ID No. 53)
15 TNFSF-7	CACACTCTGCACCAACCTCACT (SEQ ID No. 54)	TGCACTCCAAAGAAGGTCTCATC (SEQ ID No. 55)
TNFSF-8	ACCACCATATCAGTCAATGTGGAT (SEQ ID No. 56)	GAAGATGGACAACACATTCTCAAGA (SEQ ID No. 57)
TNFSF-9	AGCTACAAAGAGGACACGAAGGA (SEQ ID No. 58)	CGCAGCTCTAGTTGAAAGAAGACA (SEQ ID No. 59)
20 TNFSF-10	GGGAATATTTGAGCTTAAGGAAAATG (SEQ ID No. 60)	AAAAGGCCCCGAAAAAACTG (SEQ ID No. 61)
TNFSF-12	TACTGTCAGGTGCACTTTGATGAG (SEQ ID No. 62)	CGCAGTGGCTGAGAATTCCT (SEQ ID No. 63)
25 TNFSF-13	ATATGGTGTCCGAATCCAGGAT (SEQ ID No. 64)	CCTGACCCATGGTGAAAGTCA (SEQ ID No. 65)
TNFSF-13B	ATGCCTGAAACACTACCCAATAATT (SEQ ID No. 66)	GCAAGTTGGAGTTCATCTCCTTCT (SEQ ID No. 67)
TNFSF-14	CGTCCGTGTGCTGGATGA (SEQ ID No. 68)	CATGAAAGCCCCGAAAGTAAGAC (SEQ ID No. 69)
30 TNFSF-15	TGCGAAGTAGGTAGCAACTGGTT (SEQ ID No. 70)	CCATTAGCTTGTCCCCTTCTTG (SEQ ID No. 71)
TNFSF-18	CGGCTGTATAAAAAACAAAGACATGAT (SEQ ID No. 72)	TCCCCAACATGCAATTCATAAG (SEQ ID No. 73)
35 RNA34	AGCCCCCTCACTCCCAA (SEQ ID No. 74)	GGGTGCTGTGCTTCTGTGAAC (SEQ ID No. 75)

[0030] The quantity of each mRNA was determined by the method described above as modified below. In brief, approximately 125 mg of frozen cancer specimens were mixed with 1 mL lysis buffer supplemented with 1% 2-mecaptoethanol (Bio Rad), 0.5 mg/mL proteinase K (Pierce,) 0.1 m/mL salmon sperm DNA (5 Prime Eppendorf/Brinkmann), 0.1 mg/mL *E. coli* tRNA (Sigma), 10 nM each of specific reserve TNF-R superfamily subtype primers, and 10⁷ molecules/mL of synthetic RNA34 as an external control, and homogenized by Polytron (Brinkmann). Seventy μ L of these lysates were applied to oligo(dT)-immobilized microplate (GnePlate, RNature) for mRNA purification. Following overnight incubation at 4°C, the microplates were washed with 100 μ L plain lysis buffer 3 times, and then with 150 μ L wash buffer (0.5 mol/L NaCl, 10mmol/L Tris, pH 7.4, 1 mmol/L EDTA) 3 times at 4°C. The cDNA was directly synthesized in each well by adding 30 μ L buffer containing 1x RT-buffer (50mmol/L KCl, 10 mmol/L Tris-HCl, pH 8.3, 5.5 mmol/L MgCl₂, no dithiothreitol), 1.25 mmol/L each of dNTP, 4 units rRNasin, and 80 units pf MMLV reverse transcriptase (Promega) (without primers), and incubation at 37°C for 2 hours. 100 μ L of water was added to these 30 μ L of cDNA, and 4 μ L cDNA were transferred to 384-well PCR plates, to which 5 μ L iTaq SYBR master mix (BioRad) and 1 μ L primer cocktail (10 μ mol/L each of forward and reverse primer) were applied, and PCR was conducted in a PRISM 7900HT (Applied Biosystems), with one cycle of 95°C for 10 min followed by 45 cycles of 95°C for 30 sec, and 60°C for 1 min. Each gene was amplified in separate wells. The cycle threshold (Ct), which was the cycle of PCR to generate certain amounts of PCR products (fluorescence), was determined using analytical software (SDS, Applied Biosystems). No amplification was confirmed without cDNA (no primer dimers), and a single peak was detected at melting curve analysis. In order to compare multiple samples among 20 different mRNAs, each Ct was subtracted by the Ct of TNFRSF1A from the same sample (Δ Ct, Y-axis). The positive Ct means lower expression than TNFRSF1A, and 1 Δ Ct means one half of expression. Each symbol represents the mean value from single specimen. The left-hand side ($\circ\blacklozenge$) of each column in Figure 2

shows the results from samples of adenocarcinoma including colon, liver, stomach, uterus, and 3 cases of breast cancer (◆), and melanoma. The right-hand side (Δ) shows the results from squamous cell carcinoma, including lung, pharynx, tongue and cervical cancer samples.

[0031] As for the analysis of the induction of TNF superfamily mRNA in heparinized whole blood by anti-human T cell receptor (TCR) monoclonal antibody, the results of which are shown in Figure 3, sixty μL of whole blood was stimulated at 37°C for only 2 hours in triplicate with 5 $\mu\text{g}/\text{mL}$ anti-TCR α/β or control IgG. Fifty μL of blood samples were applied to filterplate to trap leukocytes followed by lysis in the membranes by adding lysis buffer, as described above. The purification of poly(A)⁺ mRNA, cDNA synthesis, and real time PCR were conducted as described above. The ΔCt was calculated by subtracting the Ct values of control IgG from the Ct of anti-TCR α/β . Each symbol represents the mean ΔCt from triplicate blood aliquots of single individual. Synthetic RNA34 spiked into the lysis buffer was also quantitated to validate each assay as an external control.

[0032] The cycle threshold (Ct), which was the cycle of PCR to generate certain amounts of PCR products (fluorescence), was determined using analytical software (SDS, Applied Biosystems). Although DNA chip technology (see Schena et al, Quantitative monitoring of gene expression patterns with a complementary DNA microarray, Science 1995; 270:467-70) is capable of analyzing many genes simultaneously, it lacks the capability of exact quantitation, and generally requires more than 3 folds increases as meaningful results. In contrast, the DNA quantitation technology described above provides more accurate quantitation by considering mRNA recovery and cDNA synthesis efficiency in each sample, and as small as 150% (1.5-fold) changes can be detected with statistical significance. Thus, each mRNA was analyzed by SYBR Green real time PCR (see Morrison et al., Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification, Biotechniques 1998 Jun;24(6):954-8, 960, 962) individually from each cDNA prepared by this method, as described above.

[0033] Figure 2 shows the results of TNF-R superfamily mRNA expression in various cancers (colon, breast, liver, stomach, uterus, melanoma, lung, pharynx, tongue, and cervix). In order to compare multiple samples among 20 different mRNAs, the Ct of TNF-R superfamily subtype 1A from the same sample was subtracted from each observed Ct (ΔCt , Y-axis). A positive Ct value means a lower level of expression than TNFR superfamily subtype 1A, and 1 ΔCt means one-half the level of expression. As shown in Fig. 2, almost all cancer tissues expressed high levels of TNF-R superfamily subtypes 1A, 3, 12A, and 14. This indicates that if cancer-infiltrating leukocytes are capable of producing TNF superfamily subtype 1 (i.e., lymphotoxin) (corresponding to TNF-R superfamily subtype 3), TNF superfamily subtype 2 (i.e., TNF α) (corresponding to TNF-R superfamily subtypes 1A and 1B), TNF superfamily subtype 12 (corresponding to TNF-R superfamily subtype 12A), TNF superfamily subtype 14 (corresponding to TNF-R superfamily subtype 14) or a combination of these ligands at infiltrating sites, any of these types of cancers may be eradicated. Although the expression levels of TNF-R superfamily subtypes 4, 5, 8, 9, 11A and 11B, and 17 were very low in the assessed samples (Fig. 2), the possibility that other samples may express these receptor subtypes cannot be ruled out. Accordingly, the inducibility of TNF superfamily subtype 4 (corresponding to TNF-R superfamily subtype 4), TNF superfamily subtype 5 (i.e., CD40 ligand) (corresponding to TNF-R superfamily subtype 5), TNF superfamily subtype 8 (i.e., CD30 ligand) (for TNF-R superfamily subtype 8), and TNF superfamily subtypes 13 and 13B (corresponding to TNF-R superfamily subtype 17) in leukocytes may also be important for cancer immunotherapy. Although huge individual-to-individual variation exists, the inducibility in leukocytes of TNF superfamily subtype 6 (i.e., Fas ligand) (for TNF-R superfamily subtype 6), TNF superfamily subtype 7 (i.e., CD27 ligand) (for TNF-R superfamily subtype 7), TNF superfamily subtype 11 (for TNF-R superfamily subtypes 11A and 11B), TNF superfamily subtype 18 (for TNF-R superfamily subtype 18), and TNF superfamily subtype 15 (i.e., ligand of DR3) (for TNF-R superfamily subtype 25) may be important for cancers in which expression levels of the corresponding receptors are high. Although TNF-R superfamily subtype 10A mRNA exhibited high expression in cancers, the mRNA expression of decoy receptor TNF-R superfamily subtype 10B (see Sheridan et al., Control of TRAIL-induced apoptosis by a family of signaling and decoy receptors, Science 1997 Aug 8;277(5327): 818-21 and Pan et al., An antagonist decoy receptor and a death domain-containing receptor for TRAIL, Science 1997 Aug 8;277(5327):815-8) was higher than TNF-R superfamily subtype 10A. Nevertheless, TNF-R superfamily subtype 10 (i.e., TRAIL) may still be a useful target for therapy of some cancers. There is no major difference between adenocarcinoma/melanoma (shown on the left-hand side of each column in Figure 2) and squamous cell carcinoma (shown on the right-hand side of each column in the Figure).

[0034] The T-cell receptor (TCR) is the cell surface molecule of cytotoxic T cells that recognizes specific foreign non-self molecules. The TCR's role in the initial steps of antigen presentation, as well as cancer killing, is well known (see, e.g., Mami-Chouaib, Antitumor cytotoxic T-lymphocyte response in human lung carcinoma: identification of a tumor-associated antigen, Immunol Rev. 2002 Oct;188:114-21). Although typical experiments in cellular immunology employ pure systems using isolated cells suspended in an artificial solution, the mRNA assay employed in the present embodiments permits the identification of changes in various mRNA levels after *ex vivo* stimulation of whole blood. Because the latter is more physiological than the former pure system, the present embodiments employ a simulation of TCR-target molecule interactions in whole blood, where cell-to-cell and cell-to-plasma interactions exist.

[0035] Since whole blood contains not only cytotoxic T cells, but also other types of leukocytes, the TCR was stimulated

specifically by anti-human α/β TCR mouse monoclonal IgG1 k ("anti-TCR," obtainable from BioLegend). As a control, the same concentrations of purified mouse IgG1 k (also obtainable from BioLegend) were also used. 60 μ L of whole blood was stimulated at 37°C for only 2 hours in triplicate with 5 μ g/mL anti-TCR or control

[0036] IgG. The dose and incubation time were determined by preliminary analysis of the effect of anti-TCR antibody on TNFSF mRNA expression in peripheral blood leukocytes, the results of which are shown in Figure 4A and 4B. With respect to dose response and kinetics, as shown in Figure 4A, triplicate aliquots of 60 μ L each of heparinized whole blood was mixed with PBS (○), 10 (■) or 1 (◆) μ g/mL mouse anti-human α/β TCR IgG1k, or 10 () or 1 (0) μ g/mL purified mouse IgG1k, and incubated at 37°C for 0-7 hours. Then TNFSF-2 mRNA was quantitated as described above. With respect to kinetics, as shown in Figure 4B, the Δ Ct of TNFSF-2 (•), TNFSF-5 (○), TNFSF-6 (◇), TNFSF-9 (), and TNFSF-14 (▲) were calculated by subtracting the Ct values of control IgG therefrom. Each data was the mean \pm standard deviation from triplicate aliquots of whole blood. The Δ Ct was calculated by subtracting the Ct values obtained using control IgG from the Ct values obtained using anti-TCR.

[0037] As shown in Fig. 3, anti-TCR stimulation specifically induced TNF superfamily subtypes 2, 5, 6, 9, 10, and 14. The results of TNF superfamily subtypes 2 and 14 are particularly interesting, because the mRNA expression levels of these receptors were high in all types of cancers (see Fig. 2). Simultaneous activation of multiple TNF/TNF-R superfamily subtype-related cascades may prevent resistance mechanisms of cancer cells, similarly to combination chemotherapy. The results obtained with control synthetic RNA34 spiked into the lysis buffer showed very small changes, less than ± 0.3 Δ Ct (RNA34 in Fig. 3), suggesting that the assay system is reliable. More importantly, the TCR responses exhibited substantial individual-to-individual variation, and 6 out of 9 (66%) and 5 out of 9 (56%) individuals failed to show anti-TCR-mediated induction of TNF superfamily subtypes 2 and 14, respectively. Furthermore, the individuals who showed negative TNF superfamily subtype 2 responses also showed negative TNF superfamily subtype 14 responses. However, these individuals did show other positive TNF superfamily ligands.

Example 3

[0038] In this embodiment, as in Example 1, heat-aggregated human IgG (HAG), which is used widely as a model of the immune complex as found in ADCC, was used to stimulate the Fc receptors of leukocytes in whole blood. In brief, human IgG (Sigma) was suspended in PBS at 20 mg/mL and heated at 63°C for 20 min. Sixty μ L of whole blood was stimulated at 37°C for only 2-4 hours in triplicate with 200 μ g/mL HAG or control phosphate-buffered saline (PBS). The dose of HAG and incubation time were determined by preliminary analysis, as shown in Figures 5 and 6.

[0039] Figure 5A shows the results of an analysis of the dose response. Triplicate aliquots of 60 μ L each of heparinized whole blood were mixed with various concentrations of human IgG (○), or HAG (●) and incubated at 37°C for 2 hours. Figure 5B shows the results of an analysis of kinetics. Triplicate aliquots of 60 μ L each of heparinized whole blood was mixed with PBS (○), or 200 mg/mL HAG (●) and incubated at 37°C for 0-12 hours. TNFSF-15 mRNA was then quantitated. Each data was the mean \pm standard deviation from triplicate aliquots of whole blood.

[0040] Figure 6 shows the results of an analysis of reproducibility. Blood was drawn from the same individuals within 1-3 days, and HAG-induced TNFSF mRNA and control RNA34 was quantitated. The symbols indicate ○: RNA34, ●: TNFSF-2, ◆: TNFSF-8, and ▲: TNFSF-15, respectively. Each data was the mean \pm standard deviation from triplicate aliquots of whole blood. The solid line is the line where two data are identical. The dotted lines indicate ± 0.5 Δ Ct.

[0041] As shown in Fig. 7, HAG mainly induced expression of TNF superfamily subtypes 2, 8, 14, 15, and 18. The induction of TNF superfamily subtype 15 was higher than that of the TNF superfamily subtypes obtained with anti-TCR stimulation (Fig. 3). Moreover, HAG responses exhibited substantial individual-to-individual variation. Some individuals expressed TNF superfamily subtype 15 alone, and others expressed TNF superfamily subtypes 2 and 8 without TNF superfamily subtype 15 responses. Interestingly, all individuals showed at least 1 TNF superfamily response. The results of control RNA34 spiked into the lysis buffer showed very small changes of less than ± 0.6 Δ Ct (RNA34 in Fig. 3), suggesting that the assay system is reliable. In order to test the reproducibility of the assay, blood was drawn from the same individuals twice within 1-3 days. However, the results were very similar, with Δ Ct differences of less than 0.5 (Fig. 6). Among nine individuals whose blood was tested independently using HAG and anti-TCR stimulation, the TNF superfamily subtype 15 expression response induced by HAG stimulation did not correlate with the TNF superfamily subtypes 2 and 14 expression response induced by anti-TCR stimulation.

Example 4

[0042] In this embodiment, whole blood from several individuals was subjected to HAG stimulation, and the changes in the mRNA levels of a group of chemokines were assessed in the same manner as described in Example 1. RNA34 and CD4 mRNA were employed as controls. The forward and reverse primer sequences for the assayed chemokine mRNAs are shown in Figure 8.

[0043] The results of the analysis are shown in Figure 9. In the Figure, the circles indicate individuals in which the

particular chemokine mRNA level was significantly induced by HAG stimulation. Triangles indicate a significant reduction in the mRNA level after stimulus, while X symbols indicate no significant change. As can be seen from Figure 9, the responses were varied among individuals, although there were points of commonality: for example, all of the tested individuals showed significant induction of CCL-3 and -20; CXCL-1, -2, and -3; and IL-8 and -1B.

[0044] These results indicate that it will be possible to identify the likelihood that a cancer patient will be able to recruit more leukocytes to the diseased area, and if a connection is established between specific chemokines and specific leukocyte populations, data such as that shown in Figure 9 could be used to assess the likelihood of recruiting leukocyte populations capable of effectively combating the tumor, based on TNF/TNF-R matching analysis between the leukocyte population recruited and the tumor cells.

Example 5

[0045] In this embodiment, the effects of various dietary components such as supplements on the HAG-induced changes in TNF superfamily subtype 3 mRNA levels were assessed. "Dietary components" refers to any compound or substance that may be ingested by a mammal, while "dietary supplements" indicates those beneficial dietary components, such as vitamins and natural extracts, that are used to supplement the diet of mammals, as well as other chemical compounds such as pharmaceuticals. As such, "dietary component" is broader and subsumes "dietary supplements" within it. The dietary components employed were: vitamin A (10 nM, final concentration), genistein (soy) (100 nM), curcumin (spice turmeric) (100 nM), and quercetin (plant pigments flavonoids) (100 nM). All of these dietary components have reported effects on the immune system or cancer or both. Vitamin A is known to stimulate the immune system. Genistein has been found in some studies to have anticarcinogenic activity; possible mechanisms of action include upregulation of apoptosis, inhibition of angiogenesis, inhibition of DNA topoisomerase II and inhibition of protein tyrosine kinases. Curcumin has proapoptotic effects in neoplastic cells and interferes with the activity of the transcription factor NF- κ B, which is often highly overexpressed in such cells. Quercetin has been shown to boost natural killer cell activity in rats and to inhibit degranulation of mast cells, basophils and neutrophils. In this embodiment, phosphate-buffered saline was employed as a control.

[0046] Heparinized whole blood was pre-incubated with various dietary supplements at 37 °C for 1 hour (at the blood concentrations described above), then stimulated with 1.2 μ L of heat-aggregated IgG for four hours. The blood was then incubated at 37 °C for 2 hours. The level of TNF superfamily subtype 3 mRNA was then assessed using the method described in Example 1. The primer sequences are given in Table 2 above. TNF superfamily subtype 3 is also known as lymphotoxin-alpha (LT alpha), tumor necrosis factor-beta (TNF-beta), and lymphotoxin-beta (LT beta). Secreted LT alpha assembles as a soluble homotrimer, LT alpha 3. Secreted LT alpha also complexes with the membrane-associated LT beta to generate two types of heterotrimers, LTalpha 1/beta and LT alpha 2/beta 1. Subtype 3 is expressed by activated naive CD4 cells, unpolarized IL-2-secreting effectors, and Th1 effectors, and subtype 3-specific receptors are expressed by some tumor cells.

[0047] The results are shown in Figure 10. In Figure 10, the open circles indicate TNF superfamily subtype 3 values obtained using a PBS control, while closed circles indicate subtype 3 values obtained when using the HAG stimulus. Basal levels of subtype 3 mRNA were unchanged by pretreatment with dietary supplements (open circles). However, the results indicate that HAG caused a significant ($p=0.03$) induction of TNF superfamily subtype 3 expression in blood exposed to quercetin. If this individual also had tumor cells expressing the appropriate TNF-R subtype (subtype 3), an increase in the expression of this gene may increase apoptosis of the tumor cells, so that quercetin would be a good candidate for inclusion in a diet having anti-cancer properties for this individual. The Y-axis values in Figure 10 show the cycle threshold (Ct). Each symbol is the mean \pm S.D. from triplicate aliquots of 50 μ L heparinized whole blood.

[0048] One of skill in the art will appreciate that it would also be possible to screen pharmaceuticals and other chemical compounds for similar effects on the expression of particular TNF superfamily subtypes or chemokines, so that the system of the present embodiment may be broadly employed to screen for substances useful in mounting an immune response to cancer in a particular individual.

[0049] Cancer infiltrating leukocytes are often observed in surgically removed tissue specimens by pathological examinations. However, these findings are not always an unambiguous good sign for patients: cancer progressions sometimes occur even when leukocyte infiltration exists. Pathologists are often puzzled when this occurs. The present embodiments provide a further refinement of this technique: matching TNF-R superfamily subtypes in cancer cells and TNF superfamily subtypes in infiltrating leukocytes may be a good prognostic sign, particularly when chemokine induction is also demonstrated. Pathological analyses of infiltrating leukocytes are difficult to interpret, because we do not know whether each leukocyte encounters cancer cells and whether the TNF superfamily cascade is activated or not. Quantitated base line TNF superfamily mRNA levels are also difficult to interpret. Instead, the embodiments of the present invention employ an analysis of functional changes in the expression of TNF superfamily mRNA subtypes after anti-TCR or HAG stimulation. Since infiltrating leukocytes come from the blood stream, the *ex vivo* functional assays of the present embodiments will be useful tools for the analysis of immune functions not only in cancer patients, but also in patients

suffering from other diseases such as autoimmune diseases, inflammation, transplantation, etc., in terms of diagnostics, therapeutic monitoring, prognostic markers, and identification of effective immunomodulatory drugs, monoclonal antibody-based medicines, gene therapies and vaccines. The assay is simple and physiological (2-4 hours incubation of whole blood without isolation of leukocytes), accurate with meaningful statistical analysis (triplicate aliquots of whole blood as starting materials), and sensitive (60 μ L whole blood is enough for all 17 TNF superfamily subtype mRNA quantitations). Due to an extremely small variation in the mRNA data among triplicate aliquots of whole blood, the assay permits the identification of significant changes in gene expression as small as 0.5-1.0 Δ Ct in many cases, which is far better than DNA chip-based methodologies or conventional real time PCR. The number of NK cells or cytotoxic T-cells can be quantitated by flow cytometric analysis or immunohistochemical staining, where each cell population is identified by specific cell surface markers. However, it is unclear whether these marker-positive cells really have the expected function at the time of the analysis. Although the method of the present embodiments does not identify a specific cell population with the stimulation responses of each TNF superfamily subtype mRNA, it shows the function as a whole, which is useful for the understanding of individual people. The wide individual variations obtained may lead to the discovery of the specific genetic makeup of individuals. Although the TNF superfamily response showed good reproducibility within 1-3 days (Fig. 6), it can also change over a long period of time as well as by administration of dietary supplements and exercise, suggesting the involvement of non-genetic factors. Thus, the methods of the present embodiments represent an advance toward future individualized cancer immunotherapies.

[0050] Such individualized cancer immunotherapies will be based on increasing the expression of the appropriate TNF superfamily subtypes in blood leukocytes, as well as recruiting subtype-specific leukocytes via appropriate chemokine signaling. The system and method of the embodiments described above will be useful in drug screening, validation, and clinical trials. Moreover, the same system can be applicable to autoimmune diseases, where inhibition of appropriate TNF superfamily will be the drug target.

[0051] An abundant supply of appropriately equipped police officers in the city is only meaningful when they are dispatched to the crime scene. If an officer encounters a group of criminals, his or her primary job is to call help to bring more colleagues to the scene rather than to try to arrest them alone. For cytotoxic T-cells, this action may correspond to the release of chemotactic factors, such as CCL and CXCL chemokines, and interleukins (ILs) (sequences are available from GenBank and UniGene). Thus, monitoring HAG inducibility of CCLs, CXCLs, and in peripheral blood leukocyte is also important. Like a police dispatcher, an oncologist may be able to orchestrate each patient's own immune cells by monitoring their supply in peripheral blood.

SEQUENCE LISTING

[0052]

<110> Hitachi Chemical Research Center, Inc.
Hitachi chemical Co. Ltd.
<120> Method for predicting immune response to neoplastic disease based on MRNA, expression profile in neoplastic cells and stimulated leukocytes
<130> P718364EP-PCT-DIV1
<150> US60/688,744
<151> 2005-06-08
<150> EP06772657.0
<151> 2006-06-08
<150> US60/735,508
<151> 2005-11-11
<160> 167
<170> PatentIn version 3.3
<210> 1
<211> 21
<212> DNA
<213> Artificial
<220>
<223> Primer
<400> 1
ccaaggccca gccctcacac a 21
<210> 2
<211> 21
<212> DNA

EP 1 964 931 B1

<213> Artificial
 <220>
 <223> Oligonucleotide
 <400> 2
 5 cctgccagga gaaacagaac a 21
 <210> 3
 <211> 26
 <212> DNA
 <213> Artificial
 10 <220>
 <223> Oligonucleotide
 <400> 3
 ggagacacac tcgtttctc ttagaa 26
 <210> 4
 15 <211> 19
 <212> DNA
 <213> Artificial
 <220>
 <223> Oligonucleotide
 20 <400> 4
 caagccagct ccacaatgg 19
 <210> 5
 <211> 20
 <212> DNA
 25 <213> Artificial
 <220>
 <223> Oligonucleotide
 <400> 5
 30 tgaccgaaag gcacattct 20
 <210> 6
 <211> 19
 <212> DNA
 <213> Artificial
 <220>
 35 <223> Oligonucleotide
 <400> 6
 cctcccgggc tctctacac 19
 <210> 7
 <211> 23
 40 <212> DNA
 <213> Artificial
 <220>
 <223> Oligonucleotide
 <400> 7
 45 tcatgggtga taaattggt cct 23
 <210> 8
 <211> 19
 <212> DNA
 <213> Artificial
 50 <220>
 <223> Oligonucleotide
 <400> 8
 55 acgacgtggt cagctccaa 19
 <210> 9
 <211> 21

EP 1 964 931 B1

<212> DNA
<213> Artificial

<220>
5 <223> Oligonucleotide

<400> 9
gcggcagact gtgtcctgtg t 21

10 <210> 10
<211> 20
<212> DNA
<213> Artificial

15 <220>
<223> Oligonucleotide

<400> 10
20 ggccaagaag ccaaccaata 20

<210> 11
<211> 22
<212> DNA
<213> Artificial

25 <220>
<223> Oligonucleotide

<400> 11
30 gaagatcgtc gggaaaattg at 22

<210> 12
<211> 22
<212> DNA
35 <213> Artificial

<220>
<223> Oligonucleotide

40 <400> 12
tgcatcaac tcatggaaa ga 22

<210> 13
<211> 25
45 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

50 <400> 13
gcaagagtac aaagattggc tttt 25

<210> 14
55 <211> 22
<212> DNA
<213> Artificial

EP 1 964 931 B1

<220>
<223> Oligonucleotide

<400> 14
5 ctgcagagcc ttgctgttac ag 22

<210> 15
<211> 20
<212> DNA
10 <213> Artificial

<220>
<223> Oligonucleotide

<400> 15
15 gctccggttt tcggaatcc 20

<210> 16
<211> 21
20 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 16
25 ggtaggca gcaaacagat g 21

<210> 17
30 <211> 22
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 17
35 gcctggtggt taaggctga tg 22

<210> 18
40 <211> 19
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 18
45 cgtcgactgc gttgctctt 19

<210> 19
50 <211> 21
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

55

EP 1 964 931 B1

<400> 19
ttctgccccg ttaacaaca g 21

5 <210> 20
<211> 21
<212> DNA
<213> Artificial

10 <220>
<223> Oligonucleotide

<400> 20
tgaggacaat gtcacaacg a 21

15 <210> 21
<211> 22
<212> DNA
<213> Artificial

20 <220>
<223> Oligonucleotide

<400> 21
ttgctgctca gagacgaaag tg 22

25 <210> 22
<211> 24
<212> DNA
<213> Artificial

30 <220>
<223> Oligonucleotide

<400> 22
ctgagacagt gcttcgatga cttt 24

35 <210> 23
<211> 19
<212> DNA
<213> Artificial

40 <220>
<223> Oligonucleotide

45 <400> 23
ccatgaggcc caacttct 19

50 <210> 24
<211> 23
<212> DNA
<213> Artificial

55 <220>
<223> Oligonucleotide

<400> 24
ggaagtgtag caggtgccct agt 23

EP 1 964 931 B1

<210> 25
<211> 26
<212> DNA
<213> Artificial
5

<220>
<223> Oligonucleotide

<400> 25
10 accaaattct tcaacacact ggatat 26

<210> 26
<211> 25
<212> DNA
15 <213> Artificial

<220>
<223> Oligonucleotide

20 <400> 26
atggacttac gaggttcga ctag 25

<210> 27
<211> 32
25 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

30 <400> 27
ggaaaagaga tgtacagcct acagtagtaa gc 32

<210> 28
35 <211> 25
<212> DNA
<213> Artificial

<220>
40 <223> Oligonucleotide

<400> 28
ggaaacagta actccagtt catct 25

45 <210> 29
<211> 19
<212> DNA
<213> Artificial

50 <220>
<223> Oligonucleotide

<400> 29
55 gcgaggctg gctgacgta 19

<210> 30
<211> 21
<212> DNA

EP 1 964 931 B1

<213> Artificial

<220>
<223> Oligonucleotide

5

<400> 30
tgcaaaccca gtgaccagat c 21

<210> 31
<211> 20
<212> DNA
<213> Artificial

10

<220>
<223> Oligonucleotide

15

<400> 31
aaggtgtctt ggtcgccatt 20

20

<210> 32
<211> 21
<212> DNA
<213> Artificial

25

<220>
<223> Oligonucleotide

<400> 32
cgctgatcca gtgacaatgt g 21

30

<210> 33
<211> 18
<212> DNA
<213> Artificial

35

<220>
<223> Oligonucleotide

<400> 33
gcgtctggga ggcagaga 18

40

<210> 34
<211> 19
<212> DNA
<213> Artificial

45

<220>
<223> Oligonucleotide

<400> 34
cagggagcct cgtcacgt 19

50

<210> 35
<211> 21
<212> DNA
<213> Artificial

55

<220>

EP 1 964 931 B1

<223> Oligonucleotide

<400> 35
cacccttgg ctttctt t 21

5

<210> 36
<211> 18
<212> DNA
<213> Artificial

10

<220>
<223> Oligonucleotide

<400> 36
ggaggaaggc gcaacat 18

15

<210> 37
<211> 18
<212> DNA
<213> Artificial

20

<220>
<223> Oligonucleotide

<400> 37
gcagctggca ggctcttg 18

25

<210> 38
<211> 23
<212> DNA
<213> Artificial

30

<220>
<223> Oligonucleotide

<400> 38
agtttggct tccagtgtat cga 23

35

<210> 39
<211> 21
<212> DNA
<213> Artificial

40

<220>
<223> Oligonucleotide

<400> 39
gcagtctgtc caagtttgc a 21

45

<210> 40
<211> 17
<212> DNA
<213> Artificial

50

<220>
<223> Oligonucleotide

<400> 40

55

EP 1 964 931 B1

cctgctcgcc cctatcg 17

<210> 41
<211> 20
5 <212> DNA
<213> Artificial

<220>
10 <223> Oligonucleotide

<400> 41
ttcaccctcct ctcgacattc 20

<210> 42
15 <211> 22
<212> DNA
<213> Artificial

<220>
20 <223> Oligonucleotide

<400> 42
cagctatcca cccacacaga tg 22

25 <210> 43
<211> 22
<212> DNA
<213> Artificial

30 <220>
<223> Oligonucleotide

<400> 43
35 cgaaggctcc aaagaagaca gt 22

<210> 44
<211> 22
<212> DNA
40 <213> Artificial

<220>
<223> Oligonucleotide

<400> 44
45 cgaaggctcc aaagaagaca gt 22

<210> 45
<211> 20
<212> DNA
50 <213> Artificial

<220>
<223> Oligonucleotide

55 <400> 45
cagggcaatg atcccaaagt 20

<210> 46

<211> 23
<212> DNA
<213> Artificial

5 <220>
<223> Oligonucleotide

<400> 46
agggtgtacg tcaacatcag tca 23

10 <210> 47
<211> 18
<212> DNA
<213> Artificial

15 <220>
<223> Oligonucleotide

20 <400> 47
cacggcccca aagaaggt 18

<210> 48
<211> 21
<212> DNA
25 <213> Artificial

<220>
<223> Oligonucleotide
30 <400> 48
gcccctcttc caactgaaga a 21

<210> 49
<211> 25
35 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

40 <400> 49
ggtattgtca gtggtcacat tcaag 25

<210> 50
45 <211> 19
<212> DNA
<213> Artificial

<220>
50 <223> Oligonucleotide

<400> 50
ccacagtcc gccaaacct 19

55 <210> 51
<211> 24
<212> DNA

EP 1 964 931 B1

<213> Artificial

<220>
<223> Oligonucleotide

5

<400> 51
cacctggtg caattcaat actc 24

<210> 52
<211> 24
<212> DNA
<213> Artificial

10

<220>
<223> Oligonucleotide

15

<400> 52
tggcagcatc ttcactcta aatg 24

20

<210> 53
<211> 24
<212> DNA
<213> Artificial

25

<220>
<223> Oligonucleotide

30

<400> 53
gaaatgagtc cccaaaacat ctct 24

35

<210> 54
<211> 22
<212> DNA
<213> Artificial

40

<220>
<223> Oligonucleotide

45

<400> 54
cacactctgc accaacctca ct 22

<210> 55
<211> 23
<212> DNA
<213> Artificial

50

<220>
<223> Oligonucleotide

55

<400> 55
tgactccaa agaaggtctc atc 23

<210> 56
<211> 24
<212> DNA
<213> Artificial

<220>

<223> Oligonucleotide
 <400> 56
 5 accaccatat cagtcaatgt ggat 24
 <210> 57
 <211> 25
 <212> DNA
 <213> Artificial
 10
 <220>
 <223> Oligonucleotide
 <400> 57
 15 gaagatggac aacacattct caaga 25
 <210> 58
 <211> 23
 <212> DNA
 20 <213> Artificial
 <220>
 <223> Oligonucleotide
 25 <400> 58
 agctacaaag aggacacgaa gga 23
 <210> 59
 <211> 24
 30 <212> DNA
 <213> Artificial
 <220>
 <223> Oligonucleotide
 35 <400> 59
 cgcagctcta gttgaaagaa gaca 24
 <210> 60
 40 <211> 26
 <212> DNA
 <213> Artificial
 <220>
 45 <223> Oligonucleotide
 <400> 60
 gggaaatatt gagcttaagg aaaatg 26
 <210> 61
 <211> 20
 <212> DNA
 <213> Artificial
 50
 <220>
 55 <223> Oligonucleotide
 <400> 61

EP 1 964 931 B1

aaaaggcccc gaaaaaactg 20

<210> 62
<211> 24
5 <212> DNA
<213> Artificial

<220>
10 <223> Oligonucleotide

<400> 62
tactgtcagg tgcacttga tgag 24

<210> 63
15 <211> 20
<212> DNA
<213> Artificial

<220>
20 <223> Oligonucleotide

<400> 63
cgcagtggct gagaattcct 20

<210> 64
25 <211> 22
<212> DNA
<213> Artificial

<220>
30 <223> Oligonucleotide

<400> 64
35 atatggtgtc cgaatccagg at 22

<210> 65
<211> 21
<212> DNA
40 <213> Artificial

<220>
<223> Oligonucleotide

<400> 65
45 cctgacccat ggtgaaagtc a 21

<210> 66
<211> 25
<212> DNA
50 <213> Artificial

<220>
<223> Oligonucleotide

<400> 66
55 atgcctgaaa cactacccaa taatt 25

<210> 67

<211> 24
<212> DNA
<213> Artificial

5 <220>
<223> Oligonucleotide

<400> 67
gcaagtgga gttcatctcc ttct 24

10 <210> 68
<211> 18
<212> DNA
<213> Artificial

15 <220>
<223> Oligonucleotide

20 <400> 68
cgcccgtgtg ctggatga 18

<210> 69
<211> 22
<212> DNA
25 <213> Artificial

<220>
<223> Oligonucleotide

30 <400> 69
catgaaagcc ccgaagtaag ac 22

<210> 70
<211> 23
35 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

40 <400> 70
tgccaagtag gtagcaactg gtt 23

<210> 71
45 <211> 22
<212> DNA
<213> Artificial

50 <220>
<223> Oligonucleotide

<400> 71
ccattagctt gtccccttct tg 22

55 <210> 72
<211> 26
<212> DNA
<213> Artificial

EP 1 964 931 B1

<220>
<223> Oligonucleotide

<400> 72
5 cggctgtata aaaacaaaga catgat 26

<210> 73
<211> 22
<212> DNA
10 <213> Artificial

<220>
<223> Oligonucleotide

<400> 73
15 tccccaacat gcaattcata ag 22

<210> 74
<211> 18
20 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 74
25 agccccctca ctccaaa 18

<210> 75
30 <211> 21
<212> DNA
<213> Artificial

<220>
35 <223> Oligonucleotide

<400> 75
gggtgctgtg ctctgtgaa c 21

<210> 76
40 <211> 15
<212> DNA
<213> Artificial

<220>
45 <223> Oligonucleotide

<400> 76
50 ggctgcgcc ttgga 15

<210> 77
<211> 19
<212> DNA
55 <213> Artificial

<220>
<223> Oligonucleotide

EP 1 964 931 B1

<400> 77
gggcagtgcc tcagcatt 19

5
<210> 78
<211> 20
<212> DNA
<213> Artificial

10
<220>
<223> Oligonucleotide

<400> 78
ccattgtggc caaggagatc 20

15
<210> 79
<211> 19
<212> DNA
<213> Artificial

20
<220>
<223> Oligonucleotide

<400> 79
tgtccaggtg gtccatgga 19

25
<210> 80
<211> 28
<212> DNA
<213> Artificial

30
<220>
<223> Oligonucleotide

<400> 80
cacagaattt catagctgac tactttga 28

35
<210> 81
<211> 22
<212> DNA
<213> Artificial

40
<220>
<223> Oligonucleotide

45
<400> 81
tcgcttggtt aggaagatga ca 22

50
<210> 82
<211> 23
<212> DNA
<213> Artificial

55
<220>
<223> Oligonucleotide

<400> 82
ggtattccaa accaaaagaa gca 23

EP 1 964 931 B1

<210> 83
<211> 26
<212> DNA
<213> Artificial
5
<220>
<223> Oligonucleotide
<400> 83
10 gttcagttcc aggtcataca cgtact 26
<210> 84
<211> 21
<212> DNA
15 <213> Artificial
<220>
<223> Oligonucleotide
20 <400> 84
agtcgtcttt gtcacccgaa a 21
<210> 85
<211> 26
25 <212> DNA
<213> Artificial
<220>
<223> Oligonucleotide
30 <400> 85
agctcatctc caaagagttg atgtac 26
<210> 86
35 <211> 19
<212> DNA
<213> Artificial
<220>
40 <223> Oligonucleotide
<400> 86
tgtgctgacc ccacacaga 19
45 <210> 87
<211> 23
<212> DNA
<213> Artificial
50 <220>
<223> Oligonucleotide
<400> 87
55 gctttggagt ttgggtttc ttg 23
<210> 88
<211> 26
<212> DNA

EP 1 964 931 B1

<213> Artificial

<220>
<223> Oligonucleotide

5

<400> 88
agagctacac aagaatcacc aacatc 26

<210> 89
<211> 19
<212> DNA
<213> Artificial

10

<220>
<223> Oligonucleotide

15

<400> 89
agacctcctt gccccgttt 19

20

<210> 90
<211> 22
<212> DNA
<213> Artificial

25

<220>
<223> Oligonucleotide

<400> 90
cccagaaagc tgtgatcttc aa 22

30

<210> 91
<211> 21
<212> DNA
<213> Artificial

35

<220>
<223> Oligonucleotide

<400> 91
tctgcaccc acttctctt g 21

40

<210> 92
<211> 21
<212> DNA
<213> Artificial

45

<220>
<223> Oligonucleotide

<400> 92
ccaaactggg caaggagatc t 21

50

<210> 93
<211> 24
<212> DNA
<213> Artificial

55

<220>

<223> Oligonucleotide
 <400> 93
 5 ggcccagggtg tttcatataa ttct 24
 <210> 94
 <211> 27
 <212> DNA
 <213> Artificial
 10
 <220>
 <223> Oligonucleotide
 <400> 94
 15 tgcttcacct acactaccta caagatc 27
 <210> 95
 <211> 18
 <212> DNA
 <213> Artificial
 20
 <220>
 <223> Oligonucleotide
 <400> 95
 25 gacaattccg ggcttga 18
 <210> 96
 <211> 22
 <212> DNA
 <213> Artificial
 30
 <220>
 <223> Oligonucleotide
 <400> 96
 35 ccaagccagg tgtcatatc ct 22
 <210> 97
 <211> 23
 <212> DNA
 <213> Artificial
 40
 <220>
 <223> Oligonucleotide
 <400> 97
 45 tgagtagggc ttcagcttt tca 23
 <210> 98
 <211> 22
 <212> DNA
 <213> Artificial
 50
 <220>
 <223> Oligonucleotide
 <400> 98
 55

EP 1 964 931 B1

caaggatccc aacctacctt tg 22

<210> 99
<211> 27
5 <212> DNA
<213> Artificial

<220>
10 <223> Oligonucleotide

<400> 99
ggttgacat tcttgctgt aataatt 27

<210> 100
15 <211> 18
<212> DNA
<213> Artificial

<220>
20 <223> Oligonucleotide

<400> 100
gggcagggcc atctgttc 18

25 <210> 101
<211> 24
<212> DNA
<213> Artificial

30 <220>
<223> Oligonucleotide

<400> 101
35 tctcaaggct tgcaggat ttaa 24

<210> 102
<211> 27
<212> DNA
<213> Artificial

40 <220>
<223> Oligonucleotide

<400> 102
45 cagattccac aaaagttcat agttgac 27

<210> 103
<211> 19
<212> DNA
50 <213> Artificial

<220>
<223> Oligonucleotide

55 <400> 103
ccggcctctc ttggttagg 19

<210> 104

<211> 24
 <212> DNA
 <213> Artificial

5 <220>
 <223> Oligonucleotide

<400> 104
 ctgctgtagt gtcaccaca ctga 24

10 <210> 105
 <211> 22
 <212> DNA
 <213> Artificial

15 <220>
 <223> Oligonucleotide

20 <400> 105
 cagtctctgg atgatgcggt ct 22

25 <210> 106
 <211> 28
 <212> DNA
 <213> Artificial

<220>
 <223> Oligonucleotide

30 <400> 106
 gatacacaga ccgtattctt catcctaa 28

35 <210> 107
 <211> 26
 <212> DNA
 <213> Artificial

40 <220>
 <223> Oligonucleotide

<400> 107
 tgaagatga tagcattgat gtcaca 26

45 <210> 108
 <211> 21
 <212> DNA
 <213> Artificial

50 <220>
 <223> Oligonucleotide

<400> 108
 cgctctcagg cagagctatg t 21

55 <210> 109
 <211> 21
 <212> DNA
 <213> Artificial

EP 1 964 931 B1

<220>
<223> Oligonucleotide

5 <400> 109
ctgtccaga tgctcatca g 21

<210> 110
<211> 19
<212> DNA
10 <213> Artificial

<220>
<223> Oligonucleotide

15 <400> 110
gcgcgtggtg aaacactc 19

<210> 111
<211> 22
20 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

25 <400> 111
atcggcacag atctcctat cc 22

<210> 112
30 <211> 20
<212> DNA
<213> Artificial

<220>
35 <223> Oligonucleotide

<400> 112
cgaagcatcc cgtgtcact 20

40 <210> 113
<211> 18
<212> DNA
<213> Artificial

45 <220>
<223> Oligonucleotide

<400> 113
50 gatgacacc ggcttga 18

<210> 114
<211> 22
<212> DNA
55 <213> Artificial

<220> <223>
Oligonucleotide

EP 1 964 931 B1

<400> 114
caggagtgat cttcaccacc aa 22

5 <210> 115
<211> 20
<212> DNA
<213> Artificial

10 <220>
<223> Oligonucleotide

<400> 115
ggcgtccagg ttctcatgt 20

15 <210> 116
<211> 17
<212> DNA
<213> Artificial

20 <220>
<223> Oligonucleotide

<400> 116
gcgcgctgg acttacc 17

25 <210> 117
<211> 23
<212> DNA
<213> Artificial

30 <220>
<223> Oligonucleotide

<400> 117
gtagaatatt gcagcaggca gat 23

35 <210> 118
<211> 22
<212> DNA
<213> Artificial

40 <220>
<223> Oligonucleotide

<400> 118
ctgcttcaa tacagccaca ag 22

45 <210> 119
<211> 24
<212> DNA
<213> Artificial

50 <220>
<223> Oligonucleotide

55 <400> 119
gagcagctgt tactggtgaa ttca 24

EP 1 964 931 B1

<210> 120
<211> 19
<212> DNA
<213> Artificial
5
<220>
<223> Oligonucleotide

<400> 120
10 cgtgctcac ctggctcaa 19

<210> 121
<211> 21
<212> DNA
15 <213> Artificial

<220>
<223> Oligonucleotide

20 <400> 121
ggtgctcaaa ccactgtgac a 21

<210> 122
<211> 22
25 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

30 <400> 122
ggaaatgttt gccacaggaa ga 22

<210> 123
35 <211> 21
<212> DNA
<213> Artificial

<220>
40 <223> Oligonucleotide

<400> 123
tgttcgtgt tcccctgat g 21

45 <210> 124
<211> 16
<212> DNA
<213> Artificial

50 <220>
<223> Oligonucleotide

<400> 124
55 ccactgcgcc caaacc 16

<210> 125
<211> 20
<212> DNA

EP 1 964 931 B1

<213> Artificial

<220>
<223> Oligonucleotide

5

<400> 125
gcaggattga ggcaagcttt 20

<210> 126
<211> 18
<212> DNA
<213> Artificial

10

<220>
<223> Oligonucleotide

15

<400> 126
cccttgcca ctgaactg 18

20

<210> 127
<211> 23
<212> DNA
<213> Artificial

25

<220>
<223> Oligonucleotide

<400> 127
tggatgtct tgagtgat tcc 23

30

<210> 128
<211> 23
<212> DNA
<213> Artificial

35

<220>
<223> oligonucleotide

<400> 128
ggaattcacc tcaagaacat cca 23

40

<210> 129
<211> 21
<212> DNA
<213> Artificial

45

<220>
<223> Oligonucleotide

50

<400> 129
gtggctatga cttcggttg g 21

<210> 130
<211> 18
<212> DNA
<213> Artificial

55

<220>

EP 1 964 931 B1

<223> Oligonucleotide

<400> 130
ccgtcccagg cacatcac 18

5

<210> 131
<211> 23
<212> DNA
<213> Artificial

10

<220>
<223> Oligonucleotide

<400> 131
gcaaatttc ctccattct tca 23

15

<210> 132
<211> 19
<212> DNA
<213> Artificial

20

<220>
<223> oligonucleotide

<400> 132
agagctgcgt tgcgttgt 19

25

<210> 133
<211> 22
<212> DNA
<213> Artificial

30

<220>
<223> Oligonucleotide

<400> 133
tggcgaacac ttgcagatta ct 22

35

<210> 134
<211> 20
<212> DNA
<213> Artificial

40

<220>
<223> oligonucleotide

<400> 134
cagagctgcg ttgcactgt 20

45

<210> 135
<211> 23
<212> DNA
<213> Artificial

50

<220>
<223> oligonucleotide

<400> 135

55

EP 1 964 931 B1

acacctgcag ttaccaatc gtt 23

<210> 136
<211> 22
5 <212> DNA
<213> Artificial

<220>
10 <223> Oligonucleotide

<400> 136
tctggaattc atcccaaaaa ca 22

<210> 137
15 <211> 21
<212> DNA
<213> Artificial

<220>
20 <223> Oligonucleotide

<400> 137
acttcgactt gttgcaatg g 21

25 <210> 138
<211> 25
<212> DNA
<213> Artificial

30 <220>
<223> Oligonucleotide

<400> 138
35 ccacctacaa tcctgaaag acctt 25

<210> 139
<211> 27
<212> DNA
40 <213> Artificial

<220>
<223> Oligonucleotide

<400> 139
45 cagtgtagca atgattcaa tttctc 27

<210> 140
<211> 21
<212> DNA
50 <213> Artificial

<220>
<223> Oligonucleotide
<400> 140
55 tccacgtggt gagatcattg c 21

<210> 141

EP 1 964 931 B1

<211> 21
<212> DNA
<213> Artificial

5 <220>
<223> Oligonucleotide

<400> 141
tcttgatggc cttcgattct g 21

10 <210> 142
<211> 21
<212> DNA
<213> Artificial

15 <220>
<223> Oligonucleotide

20 <400> 142
aggacgctgt cttgcatag g 21

<210> 143
<211> 25
<212> DNA
25 <213> Artificial

<220>
<223> oligonucleotide

30 <400> 143
gcatcgttgt cctttathtt ctttc 25

<210> 144
<211> 19
35 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

40 <400> 144
tgccagagcc aacgtcaag 19

<210> 145
45 <211> 19
<212> DNA
<213> Artificial

<220>
50 <223> Oligonucleotide

<400> 145
cagccgggct acaatctga 19

55 <210> 146
<211> 22
<212> DNA
<213> Artificial

EP 1 964 931 B1

<220>
<223> Oligonucleotide

<400> 146
5 gacgctcat tgatcgaatt ca 22

<210> 147
<211> 28
<212> DNA
10 <213> Artificial

<220>
<223> oligonucleotide

<400> 147
15 ttctccaga ctatgattc tttcttg 28

<210> 148
<211> 22
20 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 148
25 aagctggaaa tgaagccaaa gt 22

<210> 149
30 <211> 20
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 149
35 acacgctctt ggtggtgatg 20

<210> 150
40 <211> 19
<212> DNA
<213> Artificial

<220>
<223> oligonucleotide

<400> 150
45 cccacagcca ggacatcag 19

<210> 151
50 <211> 22
<212> DNA
<213> Artificial

<220>
55 <223> Oligonucleotide

EP 1 964 931 B1

<400> 151
cttgcacagc acataggaaa gg 22

5 <210> 152
<211> 27
<212> DNA
<213> Artificial

10 <220>
<223> oligonucleotide

<400> 152
tgctaaagaa cttagatgtc agtgcacat 27

15 <210> 153
<211> 23
<212> DNA
<213> Artificial

20 <220>
<223> Oligonucleotide

<400> 153
tggtcactc tcaatcactc tca 23

25 <210> 154
<211> 20
<212> DNA
<213> Artificial

30 <220>
<223> Oligonucleotide

<400> 154
gctcttgag ctgcctacgt 20

35 <210> 155
<211> 24
<212> DNA
<213> Artificial

40 <220>
<223> Oligonucleotide

45 <400> 155
aaggtctctt tcaccaatgc actt 24

50 <210> 156
<211> 24
<212> DNA
<213> Artificial

55 <220>
<223> Oligonucleotide

<400> 156
tcatcactgg tcttttgag tttg 24

EP 1 964 931 B1

<210> 157
<211> 20
<212> DNA
<213> Artificial
5
<220>
<223> Oligonucleotide
<400> 157
10 tctgcacagc tctggcttgt 20
<210> 158
<211> 20
<212> DNA
15 <213> Artificial
<220>
<223> Oligonucleotide
20 <400> 158
gcaggccctg aattcaaca 20
<210> 159
<211> 29
25 <212> DNA
<213> Artificial
<220>
<223> Oligonucleotide
30 <400> 159
gaagtatgca gagcttgatt ttagttta 29
<210> 160
35 <211> 17
<212> DNA
<213> Artificial
<220>
40 <223> Oligonucleotide
<400> 160
cggatcatctg ccgcaaa 17
45 <210> 161
<211> 20
<212> DNA
<213> Artificial
50 <220>
<223> Oligonucleotide
<400> 161
55 cccattcgct ccaagatgag 20
<210> 162
<211> 25
<212> DNA

EP 1 964 931 B1

<213> Artificial

<220>
<223> Oligonucleotide

5

<400> 162
tgaagtgctt tctcttgag ttaca 25

<210> 163
<211> 26
<212> DNA
<213> Artificial

10

<220>
<223> Oligonucleotide

15

<400> 163
cattccatt agaagacaaa ctgttg 26

20

<210> 164
<211> 22
<212> DNA
<213> Artificial

25

<220>
<223> Oligonucleotide

<400> 164
aaaacctctt gggaagcatg ag 22

30

<210> 165
<211> 20
<212> DNA
<213> Artificial

35

<220>
<223> Oligonucleotide

<400> 165
gggaccccga ggacagtact 20

40

<210> 166
<211> 24
<212> DNA
<213> Artificial

45

<220>
<223> Oligonucleotide

<400> 166
gaagatggaa aagcgattg tctt 24

50

<210> 167
<211> 21
<212> DNA
<213> Artificial

55

<220>

<223> Oligonucleotide

<400> 167

gggcatgtt tctgcttgag a 21

5

SEQUENCE LISTING

[0053]

10 <110> Hitachi Chemical Research Center, Inc.
 Hitachi chemical Co. Ltd.
 <120> Method for predicting immune response to neoplastic disease based on mRNA expression profile in neoplastic
 cells and stimulated leukocytes
 <130> P718364EP-PCT-DIV1
 15 <150> US60/688,744
 <151> 2005-06-08
 <150> EP06772657.0
 <151> 2006-06-08
 <150> US60/735,508
 20 <151> 2005-11-11
 <160> 167
 <170> PatentIn version 3.3
 <210> 1
 <211> 21
 25 <212> DNA
 <213> Artificial
 <220>
 <223> Primer
 <400> 1
 30 ccaaggccca gccctcacac a 21
 <210> 2
 <211> 21
 <212> DNA
 <213> Artificial
 35 <220>
 <223> Oligonucleotide
 <400> 2
 cctgccagga gaaacagaac a 21
 <210> 3
 40 <211> 26
 <212> DNA
 <213> Artificial
 <220>
 <223> Oligonucleotide
 45 <400> 3
 ggagacacac tcgtttctc ttagaa 26
 <210> 4
 <211> 19
 <212> DNA
 50 <213> Artificial
 <220>
 <223> Oligonucleotide
 <400> 4
 caagccagct ccacaatgg 19
 55 <210> 5
 <211> 20
 <212> DNA
 <213> Artificial

EP 1 964 931 B1

<220>
<223> Oligonucleotide
<400> 5
5 tgaccgaaag gcacattcct 20
<210> 6
<211> 19
<212> DNA
<213> Artificial
<220>
10 <223> Oligonucleotide
<400> 6
cctcccgggc tctctacac 19
<210> 7
<211> 23
15 <212> DNA
<213> Artificial
<220>
<223> Oligonucleotide
<400> 7
20 tcatgggtga taaattggt cct 23
<210> 8
<211> 19
<212> DNA
<213> Artificial
25 <220>
<223> Oligonucleotide
<400> 8
acgacgtggt cagctccaa 19
30 <210> 9
<211> 21
<212> DNA
<213> Artificial
35 <220>
<223> Oligonucleotide
<400> 9
40 gcggcagact ggtcctgtg t 21
<210> 10
<211> 20
<212> DNA
<213> Artificial
45 <220>
<223> Oligonucleotide
<400> 10
50 ggccaagaag ccaaccaata 20
<210> 11
<211> 22
<212> DNA
55 <213> Artificial
<220>
<223> Oligonucleotide

EP 1 964 931 B1

<400> 11
gaagatcgtc gggaaaattg at 22

5 <210> 12
<211> 22
<212> DNA
<213> Artificial

10 <220>
<223> Oligonucleotide

<400> 12
tgcatcaac ttcattgaaa ga 22

15 <210> 13
<211> 25
<212> DNA
<213> Artificial

20 <220>
<223> oligonucleotide

<400> 13
gcaagagtac aaagattggc tttt 25

25 <210> 14
<211> 22
<212> DNA
<213> Artificial

30 <220>
<223> Oligonucleotide

<400> 14
ctgcagagcc ttgtcgttac ag 22

35 <210> 15
<211> 20
<212> DNA
<213> Artificial

40 <220>
<223> oligonucleotide

<400> 15
gctccggtt tcgtaatcc 20

45 <210> 16
<211> 21
<212> DNA
<213> Artificial

50 <220>
<223> Oligonucleotide

55 <400> 16
ggttgaggca gcaaacagat g 21

EP 1 964 931 B1

<210> 17
<211> 22
<212> DNA
<213> Artificial
5
<220>
<223> oligonucleotide
<400> 17
10 gcctggtggt taaggtctga tg 22
<210> 18
<211> 19
<212> DNA
15 <213> Artificial
<220>
<223> oligonucleotide
20 <400> 18
cgtcgactgc gttgctctt 19
<210> 19
<211> 21
25 <212> DNA
<213> Artificial
<220>
<223> Oligonucleotide
30 <400> 19
ttctgccccg ttaacaaca g 21
<210> 20
35 <211> 21
<212> DNA
<213> Artificial
<220>
40 <223> Oligonucleotide
<400> 20
tgaggacaat gtcacaacg a 21
45 <210> 21
<211> 22
<212> DNA
<213> Artificial
50 <220>
<223> Oligonucleotide
<400> 21
55 ttgctgctca gagacgaaag tg 22
<210> 22
<211> 24
<212> DNA

EP 1 964 931 B1

<213> Artificial

<220>
<223> Oligonucleotide

5

<400> 22
ctgagacagt gcttcgatga ctt 24

<210> 23
<211> 19
<212> DNA
<213> Artificial

10

<220>
<223> Oligonucleotide

15

<400> 23
ccatgaggcc caacttct 19

20

<210> 24
<211> 23
<212> DNA
<213> Artificial

25

<220>
<223> Oligonucleotide

30

<400> 24
ggaagtgtag caggtgccct agt 23

35

<210> 25
<211> 26
<212> DNA
<213> Artificial

40

<220>
<223> Oligonucleotide

45

<400> 25
accaaattct tcaacacact ggatat 26

<210> 26
<211> 25
<212> DNA
<213> Artificial

50

<220>
<223> Oligonucleotide

55

<400> 26
atggacttac gaggttcga cttag 25

<210> 27
<211> 32
<212> DNA
<213> Artificial

<220>

EP 1 964 931 B1

<223> Oligonucleotide

<400> 27
5 ggaaaagaga tgtacagcct acagtagtaa gc 32

<210> 28
<211> 25
<212> DNA
<213> Artificial

10 <220>
<223> Oligonucleotide

<400> 28
15 ggaaacagta actccacgtt catct 25

<210> 29
<211> 19
<212> DNA
20 <213> Artificial

<220>
<223> Oligonucleotide

25 <400> 29
gcgaggctg gctgacgta 19

<210> 30
<211> 21
<212> DNA
30 <213> Artificial

<220>
<223> Oligonucleotide

35 <400> 30
tgcaaacca gtgaccagat c 21

<210> 31
40 <211> 20
<212> DNA
<213> Artificial

<220>
45 <223> Oligonucleotide

<400> 31
aaggtgtctt ggtcgccatt 20

50 <210> 32
<211> 21
<212> DNA
<213> Artificial

55 <220>
<223> oligonucleotide

<400> 32

EP 1 964 931 B1

cgctgatcca gtgacaatgt g 21

<210> 33
<211> 18
5 <212> DNA
<213> Artificial

<220>
10 <223> oligonucleotide

<400> 33
gcgtctggga ggcagaga 18

<210> 34
15 <211> 19
<212> DNA
<213> Artificial

<220>
20 <223> Oligonucleotide

<400> 34
cagggagcct cgtcatcgt 19

25 <210> 35
<211> 21
<212> DNA
<213> Artificial

30 <220>
<223> oligonucleotide

<400> 35
35 cacccttgg ctttctct t 21

<210> 36
<211> 18
<212> DNA
40 <213> Artificial

<220>
<223> Oligonucleotide

45 <400> 36
ggaggaaggc gcaacat 18

<210> 37
<211> 18
<212> DNA
50 <213> Artificial

<220>
<223> Oligonucleotide

55 <400> 37
gcagctggca ggctcttg 18

<210> 38

<211> 23
<212> DNA
<213> Artificial

5 <220>
<223> Oligonucleotide

<400> 38
agtttggct tccagtgtat cga 23

10 <210> 39
<211> 21
<212> DNA
<213> Artificial

15 <220>
<223> oligonucleotide

20 <400> 39
gcagtctgtc caaggttgc a 21

25 <210> 40
<211> 17
<212> DNA
<213> Artificial

<220>
<223> oligonucleotide

30 <400> 40
cctgctcgcc cctatcg 17

35 <210> 41
<211> 20
<212> DNA
<213> Artificial

40 <220>
<223> oligonucleotide

<400> 41
ttcaccctcct ctcgacattc 20

45 <210> 42
<211> 22
<212> DNA
<213> Artificial

50 <220>
<223> Oligonucleotide

<400> 42
cagctatcca cccacacaga tg 22

55 <210> 43
<211> 22
<212> DNA
<213> Artificial

EP 1 964 931 B1

<220>
<223> Oligonucleotide

<400> 43
5 cgaaggctcc aaagaagaca gt 22

<210> 44
<211> 22
<212> DNA
10 <213> Artificial

<220>
<223> oligonucleotide

<400> 44
15 cgaaggctcc aaagaagaca gt 22

<210> 45
<211> 20
20 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 45
25 cagggcaatg atcccaaagt 20

<210> 46
30 <211> 23
<212> DNA
<213> Artificial

<220>
35 <223> Oligonucleotide

<400> 46
agggtgtacg tcaacatcag tca 23

<210> 47
40 <211> 18
<212> DNA
<213> Artificial

<220>
45 <223> Oligonucleotide

<400> 47
50 cacggcccca aagaaggt 18

<210> 48
<211> 21
<212> DNA
55 <213> Artificial

<220>
<223> Oligonucleotide

EP 1 964 931 B1

<400> 48
gccctcttc caactgaaga a 21

5
<210> 49
<211> 25
<212> DNA
<213> Artificial

10
<220>
<223> Oligonucleotide

<400> 49
ggtattgtca ggtgcacat tcaag 25

15
<210> 50
<211> 19
<212> DNA
<213> Artificial

20
<220>
<223> Oligonucleotide

<400> 50
ccacagtcc gccaaacct 19

25
<210> 51
<211> 24
<212> DNA
<213> Artificial

30
<220>
<223> Oligonucleotide

<400> 51
cacctggtg caattcaat actc 24

35
<210> 52
<211> 24
<212> DNA
<213> Artificial

40
<220>
<223> Oligonucleotide

45
<400> 52
tggcagcatc ttcactcta aatg 24

50
<210> 53
<211> 24
<212> DNA
<213> Artificial

55
<220>
<223> Oligonucleotide

<400> 53
gaaatgagtc cccaaaacat ctct 24

<210> 54
<211> 22
<212> DNA
<213> Artificial
5
<220>
<223> Oligonucleotide
<400> 54
10 cacactctgc accaacctca ct 22
<210> 55
<211> 23
<212> DNA
15 <213> Artificial
<220>
<223> Oligonucleotide
20 <400> 55
tgcactccaa agaaggtctc atc 23
<210> 56
<211> 24
25 <212> DNA
<213> Artificial
<220>
<223> Oligonucleotide
30 <400> 56
accacatat cagtcaatgt ggat 24
<210> 57
35 <211> 25
<212> DNA
<213> Artificial
<220>
40 <223> oligonucleotide
<400> 57
gaagatggac aacacattct caaga 25
45 <210> 58
<211> 23
<212> DNA
<213> Artificial
50 <220>
<223> Oligonucleotide
<400> 58
55 agctacaaag aggacacgaa gga 23
<210> 59
<211> 24
<212> DNA

EP 1 964 931 B1

<213> Artificial

<220>
<223> Oligonucleotide

5

<400> 59
cgcagctcta gttgaaagaa gaca 24

<210> 60
<211> 26
<212> DNA
<213> Artificial

10

<220>
<223> oligonucleotide

15

<400> 60
gggaatatt gagctaagg aaaatg 26

20

<210> 61
<211> 20
<212> DNA
<213> Artificial

25

<220>
<223> Oligonucleotide

30

<400> 61
aaaaggcccc gaaaaaactg 20

35

<210> 62
<211> 24
<212> DNA
<213> Artificial

40

<220>
<223> Oligonucleotide

45

<400> 62
tactgtcagg tgcacttga tgag 24

<210> 63
<211> 20
<212> DNA
<213> Artificial

50

<220>
<223> oligonucleotide

55

<400> 63
cgcagtggtc gagaattcct 20

<210> 64
<211> 22
<212> DNA
<213> Artificial

<220>

<223> Oligonucleotide
 <400> 64
 5 atatggtgtc cgaatccagg at 22
 <210> 65
 <211> 21
 <212> DNA
 <213> Artificial
 10
 <220>
 <223> oligonucleotide
 <400> 65
 15 cctgacccat ggtgaaagtc a 21
 <210> 66
 <211> 25
 <212> DNA
 20 <213> Artificial
 <220>
 <223> Oligonucleotide
 25 <400> 66
 atgcctgaaa cactaccaa taatt 25
 <210> 67
 <211> 24
 30 <212> DNA
 <213> Artificial
 <220>
 <223> Oligonucleotide
 35 <400> 67
 gcaagtgga gttcatctcc ttct 24
 <210> 68
 <211> 18
 40 <212> DNA
 <213> Artificial
 <220>
 45 <223> Oligonucleotide
 <400> 68
 cgccgtgtg ctggatga 18
 <210> 69
 <211> 22
 <212> DNA
 <213> Artificial
 50
 <220>
 <223> Oligonucleotide
 55
 <400> 69

EP 1 964 931 B1

catgaaagcc ccgaagtaag ac 22

<210> 70
<211> 23
5 <212> DNA
<213> Artificial

<220>
10 <223> Oligonucleotide

<400> 70
tgccaagtag gtagcaactg gtt 23

<210> 71
15 <211> 22
<212> DNA
<213> Artificial

<220>
20 <223> oligonucleotide

<400> 71
ccattagctt gtccccttct tg 22

<210> 72
25 <211> 26
<212> DNA
<213> Artificial

<220>
30 <223> Oligonucleotide

<400> 72
35 cggctgtata aaaacaaaga catgat 26

<210> 73
<211> 22
<212> DNA
<213> Artificial

<220>
40 <223> Oligonucleotide

<400> 73
45 tcccaacat gcaattcata ag 22

<210> 74
<211> 18
<212> DNA
50 <213> Artificial

<220>
<223> Oligonucleotide

<400> 74
55 agccccctca ctccaaa 18

<210> 75

EP 1 964 931 B1

<211> 21
<212> DNA
<213> Artificial

5 <220>
<223> Oligonucleotide

<400> 75
gggtgctgtg cttctgtgaa c 21

10 <210> 76
<211> 15
<212> DNA
<213> Artificial

15 <220>
<223> Oligonucleotide

20 <400> 76
ggcctgcgcc ttgga 15

<210> 77
<211> 19
<212> DNA
25 <213> Artificial

<220>
<223> oligonucleotide

30 <400> 77
gggcagtgcc tcagcattt 19

<210> 78
<211> 20
35 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

40 <400> 78
ccattgtggc caaggagatc 20

<210> 79
45 <211> 19
<212> DNA
<213> Artificial

50 <220>
<223> oligonucleotide

<400> 79
tgtccaggtg gtccatgga 19

55 <210> 80
<211> 28
<212> DNA
<213> Artificial

EP 1 964 931 B1

<220>
<223> Oligonucleotide

<400> 80
5 cacagaattt catagctgac tactttga 28

<210> 81
<211> 22
<212> DNA
10 <213> Artificial

<220>
<223> Oligonucleotide

<400> 81
15 tcgcttggtt aggaagatga ca 22

<210> 82
<211> 23
20 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 82
25 ggtattccaa accaaaagaa gca 23

<210> 83
<211> 26
30 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 83
35 gttcagttcc aggtcataca cgtact 26

<210> 84
<211> 21
40 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 84
45 agtcgtcttt gtcaccgaa a 21

<210> 85
<211> 26
50 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

55

EP 1 964 931 B1

<400> 85
agctcatctc caaagagttg atgtac 26

5
<210> 86
<211> 19
<212> DNA
<213> Artificial

10
<220>
<223> oligonucleotide

15
<400> 86
tgtgctgacc ccacacaga 19

20
<210> 87
<211> 23
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

25
<400> 87
gcttggagt ttgggtttc ttg 23

30
<210> 88
<211> 26
<212> DNA
<213> Artificial

<220>
<223> oligonucleotide

35
<400> 88
agagctacac aagaatcacc aacatc 26

40
<210> 89
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

45
<400> 89
agacctcctt gccccgttt 19

50
<210> 90
<211> 22
<212> DNA
<213> Artificial

55
<220>
<223> Oligonucleotide

<400> 90

EP 1 964 931 B1

cccagaaagc tgtgatcttc aa 22

<210> 91
<211> 21
5 <212> DNA
<213> Artificial

<220>
10 <223> oligonucleotide

<400> 91
tctgcaccc acttctctt g 21

<210> 92
15 <211> 21
<212> DNA
<213> Artificial

<220>
20 <223> oligonucleotide

<400> 92
ccaaactggg caaggagatc t 21

25 <210> 93
<211> 24
<212> DNA
<213> Artificial

30 <220>
<223> Oligonucleotide

<400> 93
35 ggcccagggtg ttcatataa ttct 24

<210> 94
<211> 27
<212> DNA
<213> Artificial

40 <220>
<223> Oligonucleotide

<400> 94
45 tgcttcacct acactaccta caagatc 27

<210> 95
<211> 18
<212> DNA
50 <213> Artificial

<220>
<223> oligonucleotide

55 <400> 95
gacaattccg ggcttgga 18

<210> 96

EP 1 964 931 B1

<211> 22
<212> DNA
<213> Artificial

5 <220>
<223> Oligonucleotide

<400> 96
ccaagccagg tgtcatattc ct 22

10 <210> 97
<211> 23
<212> DNA
<213> Artificial

15 <220>
<223> oligonucleotide

20 <400> 97
tgagtagggc ttcagctttt tca 23

25 <210> 98
<211> 22
<212> DNA
<213> Artificial

<220>
<223> oligonucleotide

30 <400> 98
caaggatccc aacctacctt tg 22

35 <210> 99
<211> 27
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

40 <400> 99
ggttgaccat tcttgctgt aataatt 27

45 <210> 100
<211> 18
<212> DNA
<213> Artificial

50 <220>
<223> Oligonucleotide

<400> 100
gggcagggcc atctgttc 18

55 <210> 101
<211> 24
<212> DNA
<213> Artificial

EP 1 964 931 B1

<220>
<223> oligonucleotide

<400> 101
5 tctcaaggct ttgcaggat ttaa 24

<210> 102
<211> 27
<212> DNA
10 <213> Artificial

<220>
<223> Oligonucleotide

<400> 102
15 cagattccac aaaagttcat agttgac 27

<210> 103
<211> 19
20 <212> DNA
<213> Artificial

<220>
<223> oligonucleotide

<400> 103
25 ccggcctctc ttggttagg 19

<210> 104
30 <211> 24
<212> DNA
<213> Artificial

<220>
35 <223> oligonucleotide

<400> 104
ctgctgtagt gttcaccaca ctga 24

<210> 105
40 <211> 22
<212> DNA
<213> Artificial

<220>
45 <223> Oligonucleotide

<400> 105
50 cagtctctgg atgatgcgtt ct 22

<210> 106
<211> 28
<212> DNA
55 <213> Artificial

<220>
<223> Oligonucleotide

EP 1 964 931 B1

<400> 106
gatacacaga ccgtattctt catcctaa 28

5 <210> 107
<211> 26
<212> DNA
<213> Artificial

10 <220>
<223> Oligonucleotide

<400> 107
tgaaagatga tagcattgat gtcaca 26

15 <210> 108
<211> 21
<212> DNA
<213> Artificial

20 <220>
<223> oligonucleotide

25 <400> 108
cgctctcagg cagagctatg t 21

30 <210> 109
<211> 21
<212> DNA
<213> Artificial

35 <220>
<223> Oligonucleotide

<400> 109
cttgtccaga tgctgcatca g 21

40 <210> 110
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

45 <400> 110
gcgctggtg aaacacttc 19

50 <210> 111
<211> 22
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

55 <400> 111
atcggcacag atctccttat cc 22

EP 1 964 931 B1

<210> 112
<211> 20
<212> DNA
<213> Artificial
5
<220>
<223> Oligonucleotide
<400> 112
10 cgaagcatcc cgtgtcact 20
<210> 113
<211> 18
<212> DNA
15 <213> Artificial
<220>
<223> Oligonucleotide
20 <400> 113
gatgacaccc ggcttga 18
<210> 114
<211> 22
25 <212> DNA
<213> Artificial
<220>
<223> oligonucleotide
30 <400> 114
caggagtgat cttcaccacc aa 22
<210> 115
35 <211> 20
<212> DNA
<213> Artificial
<220>
40 <223> Oligonucleotide
<400> 115
ggcgtccagg ttctcatgt 20
45 <210> 116
<211> 17
<212> DNA
<213> Artificial
50 <220>
<223> Oligonucleotide
<400> 116
55 gcgcgctgg acttacc 17
<210> 117
<211> 23
<212> DNA

EP 1 964 931 B1

<213> Artificial

<220>
<223> Oligonucleotide

5

<400> 117
gtagaatattc gcagcaggca gat 23

<210> 118
<211> 22
<212> DNA
<213> Artificial

10

<220>
<223> Oligonucleotide

15

<400> 118
ctgcttccaa tacagccaca ag 22

20

<210> 119
<211> 24
<212> DNA
<213> Artificial

25

<220>
<223> Oligonucleotide

30

<400> 119
gagcagctgt tactggtgaa ttca 24

35

<210> 120
<211> 19
<212> DNA
<213> Artificial

40

<220>
<223> Oligonucleotide

45

<400> 120
cgtgcttcac ctggctcaa 19

<210> 121
<211> 21
<212> DNA
<213> Artificial

50

<220>
<223> Oligonucleotide

55

<400> 121
ggtgctcaaa ccactgtgac a 21

<210> 122
<211> 22
<212> DNA
<213> Artificial

<220>

EP 1 964 931 B1

<223> Oligonucleotide

<400> 122
5 ggaaatggtt gccacaggaa ga 22

<210> 123
<211> 21
<212> DNA
<213> Artificial

10 <220>
<223> oligonucleotide

<400> 123
15 tgttctgtg tccccctgat g 21

<210> 124
<211> 16
<212> DNA
20 <213> Artificial

<220>
<223> Oligonucleotide

25 <400> 124
ccactgcgcc caaacc 16

<210> 125
<211> 20
<212> DNA
30 <213> Artificial

<220>
<223> Oligonucleotide

35 <400> 125
gcaggattga ggcaagcttt 20

<210> 126
<211> 18
<212> DNA
40 <213> Artificial

<220>
45 <223> Oligonucleotide

<400> 126
cccctggcca ctgaactg 18

<210> 127
<211> 23
<212> DNA
50 <213> Artificial

55 <220>
<223> Oligonucleotide

<400> 127

EP 1 964 931 B1

5
10
15
20
25
30
35
40
45
50
55

tggatgtct tgaggtgaat tcc 23

<210> 128
<211> 23
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 128
ggaattcacc tcaagaacat cca 23

<210> 129
<211> 21
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 129
gtggctatga cttcggttg g 21

<210> 130
<211> 18
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 130
ccgtcccagg cacatcac 18

<210> 131
<211> 23
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 131
gcaaatttc cttccattct tca 23

<210> 132
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 132
agagctgcgt tgcgtttgt 19

<210> 133

EP 1 964 931 B1

<211> 22
<212> DNA
<213> Artificial

5 <220>
<223> Oligonucleotide

<400> 133
tggcgaacac ttgcagatta ct 22

10 <210> 134
<211> 20
<212> DNA
<213> Artificial

15 <220>
<223> Oligonucleotide

20 <400> 134
cagagctgcg ttgcactgt 20

<210> 135
<211> 23
<212> DNA
25 <213> Artificial

<220>
<223> Oligonucleotide

30 <400> 135 23
acacctgcag ttaccaatc gtt 23

<210> 136
<211> 22
35 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

40 <400> 136
tctggaattc atcccaaaaa ca 22

<210> 137
45 <211> 21
<212> DNA
<213> Artificial

<220>
50 <223> Oligonucleotide

<400> 137 21
acttcgactt gttgcaatg g 21

55 <210> 138
<211> 25
<212> DNA
<213> Artificial

EP 1 964 931 B1

<220>
<223> Oligonucleotide

<400> 138
5 ccacctaca tcctgaaag acct 25

<210> 139
<211> 27
<212> DNA
10 <213> Artificial

<220>
<223> oligonucleotide

<400> 139
15 cagtgtagca atgattcaa tttctc 27

<210> 140
<211> 21
20 <212> DNA
<213> Artificial

<220>
<223> Oligonucleotide

<400> 140
25 tccacgtgt gagatcattg c 21

<210> 141
30 <211> 21
<212> DNA
<213> Artificial

<220>
35 <223> oligonucleotide

<400> 141
tcttgatggc cttcgattct g 21

<210> 142
40 <211> 21
<212> DNA
<213> Artificial

<220>
45 <223> oligonucleotide

<400> 142
50 aggacgctgt cttgcatag g 21

<210> 143
<211> 25
<212> DNA
55 <213> Artificial

<220>
<223> oligonucleotide

EP 1 964 931 B1

<400> 143
gcatcgtgt cctttatctt ctttc 25

5 <210> 144
<211> 19
<212> DNA
<213> Artificial

10 <220>
<223> Oligonucleotide

<400> 144
tgccagagcc aacgtcaag 19

15 <210> 145
<211> 19
<212> DNA
<213> Artificial

20 <220>
<223> Oligonucleotide

<400> 145
cagccgggct acaatctga 19

25 <210> 146
<211> 22
<212> DNA
<213> Artificial

30 <220>
<223> Oligonucleotide

<400> 146
gacgcttcat tgatcgaatt ca 22

35 <210> 147
<211> 28
<212> DNA
<213> Artificial

40 <220>
<223> oligonucleotide

<400> 147
ttctccaga ctatgattc tttcttg 28

45 <210> 148
<211> 22
<212> DNA
<213> Artificial

50 <220>

55 <223> Oligonucleotide

<400> 148
aagctggaaa tgaagccaaa gt 22

EP 1 964 931 B1

<210> 149
<211> 20
<212> DNA
<213> Artificial
5
<220>
<223> Oligonucleotide
<400> 149
10 acacgctctt ggtgggatg 20
<210> 150
<211> 19
<212> DNA
15 <213> Artificial
<220>
<223> Oligonucleotide
20 <400> 150
cccacagcca ggacatcag 19
<210> 151
<211> 22
25 <212> DNA
<213> Artificial
<220>
<223> Oligonucleotide
30 <400> 151
cttgcacagc acataggaaa gg 22
<210> 152
35 <211> 27
<212> DNA
<213> Artificial
<220>
40 <223> oligonucleotide
<400> 152
tgctaaagaa cctagatgtc agtgcacat 27
45 <210> 153
<211> 23
<212> DNA
<213> Artificial
50 <220>
<223> Oligonucleotide
<400> 153
55 tggccactc tcaatcactc tca 23
<210> 154
<211> 20
<212> DNA

EP 1 964 931 B1

<213> Artificial

<220>
<223> Oligonucleotide

5

<400> 154
gctcttgag ctgcctacgt 20

<210> 155
<211> 24
<212> DNA
<213> Artificial

10

<220>
<223> oligonucleotide

15

<400> 155
aaggtctctt tcaccaatgc actt 24

20

<210> 156
<211> 24
<212> DNA
<213> Artificial

25

<220>
<223> Oligonucleotide

<400> 156
tcactcactgg tcttttgag ttg 24

30

<210> 157
<211> 20
<212> DNA
<213> Artificial

35

<220>
<223> Oligonucleotide

<400> 157
tctgcacagc tctggctgt 20

40

<210> 158
<211> 20
<212> DNA
<213> Artificial

45

<220>
<223> Oligonucleotide

<400> 158
gcaggccctg aattcaaca 20

50

<210> 159
<211> 29
<212> DNA
<213> Artificial

55

<220>

<223> oligonucleotide
 <400> 159
 gaagtatgca gagcttgatt ttagttta 29
 5
 <210> 160
 <211> 17
 <212> DNA
 <213> Artificial
 10
 <220>
 <223> Oligonucleotide
 <400> 160
 cggtcacatcg ccgcaaa 17
 15
 <210> 161
 <211> 20
 <212> DNA
 <213> Artificial
 20
 <220>
 <223> Oligonucleotide
 <400> 161
 cccattcgct ccaagatgag 20
 25
 <210> 162
 <211> 25
 <212> DNA
 <213> Artificial
 30
 <220>
 <223> Oligonucleotide
 <400> 162
 tgaagtgctt tctctggag ttaca 25
 35
 <210> 163
 <211> 26
 <212> DNA
 <213> Artificial
 40
 <220>
 <223> Oligonucleotide
 <400> 163
 cattccatt agaagacaaa ctgttg 26
 45
 <210> 164
 <211> 22
 <212> DNA
 <213> Artificial
 50
 <220>
 <223> oligonucleotide
 <400> 164

aaaacctctt gggaagcatg ag 22

<210> 165
 <211> 20
 <212> DNA
 <213> Artificial

<220>
 <223> oligonucleotide

<400> 165
 gggaccccga ggacagtact 20

<210> 166
 <211> 24
 <212> DNA
 <213> Artificial

<220>
 <223> Oligonucleotide

<400> 166
 gaagatggaa aagcgatttg tctt 24

<210> 167
 <211> 21
 <212> DNA
 <213> Artificial

<220>
 <223> oligonucleotide

<400> 167
 gggcatgttt tctgcttgag a 21

Claims

1. A method of determining the likely severity of prognosis for a neoplastic disease in a mammal, comprising:

determining a level of mRNA expression of a plurality of chemokines selected from the group consisting of CCL3, CCL20, CXCL1, CXCL2, CXCL3 in a first sample of whole blood from the mammal;
 exposing a second sample of whole blood of the mammal to a stimulus of heat-aggregated human IgG;
 determining a level of mRNA expression of the plurality of chemokines in the second sample;
 determining a likely lesser severity of prognosis when chemokines that exhibit a significant induction in mRNA expression level in said whole blood between said first and second samples are identified.

2. The method of claim 1, wherein exposing the second sample of whole blood includes addition of heparin.

Patentansprüche

1. Ein Verfahren zur Bestimmung der wahrscheinlichen Schwere bei der Vorhersage für eine neoplastische Erkrankung eines Säugers, umfassend:

Bestimmen eines Niveaus einer mRNA-Expression einer Vielzahl von Chemokinen, die aus der Gruppe ausgewählt sind, die aus CCL3, CCL20, CXCL1, CXCL2 und CXCL3 besteht, in einer ersten Probe aus Vollblut von dem Säuger;

EP 1 964 931 B1

Aussetzen einer zweiten Probe aus Vollblut des Säugers einem Reiz aus wärme-aggregiertem humanem IgG;
Bestimmen eines Niveaus einer mRNA-Expression der Vielzahl von Chemokinen in der zweiten Probe;
Bestimmen einer wahrscheinlich geringeren Schwere bei der Vorhersage, wenn Chemokine, die eine signifi-
5 kante Induktion bei dem Niveau der mRNA-Expression in dem Vollblut zwischen den ersten und zweiten Proben
zeigen, identifiziert werden.

2. Das Verfahren nach Anspruch 1, worin das Aussetzen der zweiten Probe aus Vollblut einen Zusatz von Heparin
einschließt.

10

Revendications

1. Procédé pour déterminer la gravité probable du pronostic d'une affection néoplastique chez le mammifère compre-
nant les étapes consistant à :

15

- déterminer un niveau d'expression d'ARNm d'un ensemble de chémokines choisies dans le groupe formé par
CCL3, CCL20, CXCL1, CXCL2, CXCL3 dans un premier échantillon de sang total du mammifère ;
- exposer un second échantillon du sang total du mammifère à un stimulus d'IgG humaine agglomérée par la
chaleur ;
- 20 - déterminer le niveau d'expression de l'ARNm de l'ensemble des chémokines dans le second échantillon ;
- déterminer une gravité inférieure probable du pronostic lorsque des chémokines qui présentent une induction
significative du niveau d'expression de l'ARNm dans le sang total entre le premier et le second échantillon sont
identifiées.

20

25

2. Procédé conforme à la revendication 1, selon lequel l'étape d'exposition du second échantillon de sang total comporte
l'addition d'héparine.

30

35

40

45

50

55




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7

EP 1 964 931 B1

CCL-1	MM282:	GGCCTGCGCCTTGG A	(SEQ ID No. 76)
	MM283:	GGGCAGTGCCTCAGCATT	(SEQ ID No. 77)
CCL-2	MM284:	CCATTGTGGCCAAGGAGATC	(SEQ ID No. 78)
	MM285:	TGTCCAGGTGGTCCATGGA	(SEQ ID No. 79)
CCL-3	MM286:	ACAGAATTTTCATAGCTGACTACTTTGA	(SEQ ID No. 80)
	MM287:	TCGCTTGGTTAGGAAGATGACA	(SEQ ID No. 81)
CCL-4	MM288:	GGTATTCCAAACCAAAGAAGCA	(SEQ ID No. 82)
	MM289:	G TTCAGTTC CAGTTCATACACGTACT	(SEQ ID No. 83)
CCL-5	MM290:	AGTCGTCTTTGT CACCCGAAA	(SEQ ID No. 84)
	MM291:	AGTCATCTCCAAAGAGTTGATGTAC	(SEQ ID No. 85)
CCL-7	MM292:	TGTGCTGACCCACACAGA	(SEQ ID No. 86)
	MM293:	GCTTTGGAGTTTGGGTTTCTTG	(SEQ ID No. 87)
CCL-8	MM294:	AGAGCTACACAAGAATCACCAACATC	(SEQ ID No. 88)
	MM295:	AGACCTCCTTGCCCCGTTT	(SEQ ID No. 89)
CCL-11	MM296:	CCCAGAAAGCTGTGATCTTCAA	(SEQ ID No. 90)
	MM297:	TCCTGCACCCACTTCTTCTTG	(SEQ ID No. 91)
CCL-13	MM298:	CCAAACTGGGCAAGGAGATCT	(SEQ ID No. 92)
	MM299:	GGCCCAGGTGTTTCATATAATTCT	(SEQ ID No. 93)
CCL-14	MM300:	TGCTTCACCTACACTACCTACAAGATC	(SEQ ID No. 94)
	MM301:	GACAATTC CGGGCTTGG A	(SEQ ID No. 95)
CCL-15	MM302:	CCAAGCCAGGTGTCATATTCCT	(SEQ ID No. 96)
	MM303:	TGAGTAGGGCTTCAGCTTTTCA	(SEQ ID No. 97)
CCL-16	MM304:	CAAGGATCCCAACCTACCTTTG	(SEQ ID No. 98)
	MM305:	GGTTGACCAATCTTTGCTGTAATAATT	(SEQ ID No. 99)
CCL-17	MM306:	GGGCAGGGCCATCTGTTC	(SEQ ID No. 100)
	MM307:	TCTCAAGGCTTTGCAGGTATTTAA	(SEQ ID No. 101)
CCL-18	MM308:	CAGATTCACAAAAGTTCATAGTTGAC	(SEQ ID No. 102)
	MM309:	CCGGCCTCTCTTGGTTAGG	(SEQ ID No. 103)
CCL-19	MM310:	CTGCTGTAGTGTTCACCACACTGA	(SEQ ID No. 104)
	MM311:	CAGTCTCTGGATGATGCGTTCT	(SEQ ID No. 105)
CCL-20	MM312:	ATACACAGACCGTATTCTTCATCCTAA	(SEQ ID No. 106)
	MM313:	TGAAAGATGATAGCATTGATGTCACA	(SEQ ID No. 107)
CCL-21	MM314:	CGCTTCAGGCAGAGCTATGT	(SEQ ID No. 108)
	MM315:	CTTGTCCAGATGCTGCATCAG	(SEQ ID No. 109)
CCL-22	MM316:	GCGCGTGGTGAAACACTTC	(SEQ ID No. 110)
	MM317:	ATCGGCACAGATCTCCTTATCC	(SEQ ID No. 111)
CCL-23	MM318:	CGAAGCATCCCGTGTTCACT	(SEQ ID No. 112)
	MM319:	GATGACACCCGGCTTGG A	(SEQ ID No. 113)
CCL-24	MM320:	CAGGAGTGATCTTCACCACCAA	(SEQ ID No. 114)
	MM321:	GGCGTCCAGGTTCTTCATGT	(SEQ ID No. 115)
CCL-25	MM322:	GCGCGCTGGACTTACC	(SEQ ID No. 116)
	MM323:	GTAGAATATCGCAGCAGGCAGAT	(SEQ ID No. 117)
CCL-26	MM324:	CTGCTTCCAATACAGCCACAAG	(SEQ ID No. 118)
	MM325:	GAGCAGCTGTACTGGTGAATTCA	(SEQ ID No. 119)
CCL-27	MM326:	CGTGCTTCACCTGGCTCAA	(SEQ ID No. 120)
	MM327:	GGTGCTCAAACCACTGTGACA	(SEQ ID No. 121)
CCL-28	MM328:	GGAAATGTTTGCCACAGGAAGA	(SEQ ID No. 122)
	MM329:	TGTTTCGTGTTTCCCCTGATG	(SEQ ID No. 123)

FIG 8A

CXCL-1	MM330:	CCACTGCGCCCAAACC	(SEQ ID No. 124)
	MM331:	GCAGGATTGAGGCAAGCTTT	(SEQ ID No. 125)
CXCL-2	MM332:	CCCCTGGCCACTGAACTG	(SEQ ID No. 126)
	MM333:	TGGATGTTCTTGAGGTGAATTCC	(SEQ ID No. 127)
CXCL-3	MM334:	GGAATTCACCTCAAGAACATCCA	(SEQ ID No. 128)
	MM335:	GTGGCTATGACTTCGGTTTGG	(SEQ ID No. 129)
CXCL-4	MM336:	CCGTCCCAGGCACATCAC	(SEQ ID No. 130)
	MM337:	GCAAATTTTCTTCCATTCTTCA	(SEQ ID No. 131)
CXCL-5	MM338:	AGAGCTGCGTTGCGTTTGT	(SEQ ID No. 132)
	MM339:	TGGCGAACACTTGCAGATTACT	(SEQ ID No. 133)
CXCL-6	MM340:	CAGAGCTGCGTTGCACCTTGT	(SEQ ID No. 134)
	MM341:	ACACCTGCAGTTTACCAATCGTT	(SEQ ID No. 135)
CXCL-7	MM342:	TCTGGAATTCATCCCAAAAACA	(SEQ ID No. 136)
	MM343:	ACTTCGACTTGGTTGCAATGG	(SEQ ID No. 137)
CXCL-9	MM344:	CCACCTACAATCCTTGAAGACCTT	(SEQ ID No. 138)
	MM345:	CAGTGTAGCAATGATTTCAATTTTCTC	(SEQ ID No. 139)
CXCL-10	MM346:	TCCACGTGTGAGATCATTGC	(SEQ ID No. 140)
	MM347:	TCTTGATGGCCTTCGATTCTG	(SEQ ID No. 141)
CXCL-11	MM348:	AGGACGCTGTCTTTGCATAGG	(SEQ ID No. 142)
	MM349:	GCATCGTTGTCTTTATTTTCTTTC	(SEQ ID No. 143)
CXCL-12	MM350:	TGCCAGAGCCAACGTCAAG	(SEQ ID No. 144)
	MM351:	CAGCCGGGCTACAATCTGA	(SEQ ID No. 145)
CXCL-13	MM352:	GACGCTTCATTGATCGAATTCA	(SEQ ID No. 146)
	MM353:	TTCTTCCAGACTATGATTTCTTTTCTTG	(SEQ ID No. 147)
CXCL-14	MM354:	AAGCTGGAATGAAGCCAAAGT	(SEQ ID No. 148)
	MM355:	ACACGCTCTTGGTGGTGATG	(SEQ ID No. 149)
CXCL-16	MM356:	CCCACAGCCAGGACATCAG	(SEQ ID No. 150)
	MM357:	CTTGACACAGCACATAGGAAAGG	(SEQ ID No. 151)
IL-8	MM358:	TGCTAAAGAACTTAGATGTCAGTGCAT	(SEQ ID No. 152)
	MM359:	TGGTCCACTCTCAATCACTCTCA	(SEQ ID No. 153)
IL-5	MM360:	GCTCTTGGAGCTGCCTACGT	(SEQ ID No. 154)
	MM361:	AAGGTCTCTTTCACCAATGCACCTT	(SEQ ID No. 155)
IL-6	MM362:	TCATCACTGGTCTTTTGGAGTTTG	(SEQ ID No. 156)
	MM363:	TCTGCACAGCTCTGGCTTGT	(SEQ ID No. 157)
IL-12A	MM364:	GCAGGCCCTGAATTTCAACA	(SEQ ID No. 158)
	MM365:	GAAGTATGCAGAGCTTGATTTTAGTTTTA	(SEQ ID No. 159)
IL-12B	MM366:	CGGTCACTCTGCCGCAA	(SEQ ID No. 160)
	MM367:	CCCATTTCGCT CCAAGATGAG	(SEQ ID No. 161)
IL-15	MM368:	TGAAGTGCTTTCTCTTGGAGTTACA	(SEQ ID No. 162)
	MM369:	CATTCCCATTAGAAGACAAACTGTTG	(SEQ ID No. 163)
IL-16c	MM370:	AAAACCTCTTGGGAAGCATGAG	(SEQ ID No. 164)
	MM371:	GGGACCCCGAGGACAGTACT	(SEQ ID No. 165)
IL-1B	MM372:	GAAGATGGAAAAGCGATTTGTCTT	(SEQ ID No. 166)
	MM373:	GGGCATGTTTTCTGCTTGAGA	(SEQ ID No. 167)

FIG 8B




FIGURE 9




FIGURE 10

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 03099312 A [0010]
- WO 2005044792 A [0013]
- WO 2006110091 A [0014]
- US 20040265864 A [0018]
- WO 2005042784 A [0018]
- US 60688744 B [0052] [0053]
- EP 06772657 A [0052] [0053]
- US 60735508 B [0052] [0053]

Non-patent literature cited in the description

- **Richman AV.** Immunofluorescence studies of benign and malignant human mammary tissue. *J. Natl. Cancer Inst.*, 1976, vol. 57, 263-7 [0003]
- **Koneval T et al.** Demonstration of immunoglobulin in tumor and marginal tissues of squamous cell carcinomas of the head and neck. *J. Natl. Cancer Inst.*, 1977, vol. 59, 1089-97 [0003]
- **Nakanishi et al.** Perforin expression in lymphocytes infiltrated to human colorectal cancer. *Br. J. Cancer*, 1991, vol. 64, 239-42 [0003]
- **Perussia et al.** Assays for antibody-dependent cell-mediated cytotoxicity (ADCC) and reverse ADCC (redirected cytotoxicity) in human natural killer cells. *Methods Mol Biol.*, 2000, vol. 121, 179-92 [0006]
- **Vujanovic.** Role of TNF family ligands in antitumor activity of natural killer cells. *Int. Rev. Immunol.*, June 2001, vol. 20 (3-4), 415-37 [0008]
- **McDermott.** TNF and TNFR biology in health and disease. *Cell. Mol. Biol. (Noisy-le-grand)*, June 2001, vol. 47 (4), 619-35 [0008]
- **Schimanski C et al.** *Clin. Cancer Res.*, 2005, vol. 11, 1743-1750 [0009]
- **Leung S et al.** *Gastroenterology*, 2004, vol. 127, 457-469 [0011]
- **Moran C et al.** *Clin. Cancer Res.*, 2002, vol. 8, 3803-3812 [0012]
- **Becker.** The relationship of the chemotactic behavior of the complement-derived factors, C3a, C5a, and C567, and a bacterial chemotactic factor to their ability to activate the proesterase 1 of rabbit polymorphonuclear leukocytes. *J. Exp. Med.*, 1972, vol. 135, 376-87 [0015]
- **Mitsuhashi.** Absolute quantitation of mRNA in human blood leukocytes as a model for phenotypic gene expression-based diagnostics. *Clin Chem*, 2006, vol. 52, 4 [0018]
- **Ostreiko et al.** Production and characterization of heat-aggregated IgG complexes with predetermined molecular masses: light-scattering study. *Immunol Lett.*, 1987, vol. 15, 311-6 [0024]
- **Morrison et al.** Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification. *Biotechniques*, 1998, vol. 24, 954-8960, 96 [0024]
- **Schena et al.** Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science*, 1995, vol. 270, 467-70 [0032]
- **Morrison et al.** Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification. *Biotechniques*, June 1998, vol. 24 (6), 954-8960, 962 [0032]
- **Sheridan et al.** Control of TRAIL-induced apoptosis by a family of signaling and decoy receptors. *Science*, 08 August 1997, vol. 277 (5327), 818-21 [0033]
- **Pan et al.** An antagonist decoy receptor and a death domain-containing receptor for TRAIL. *Science*, 08 August 1997, vol. 277 (5327), 815-8 [0033]
- **Mami-Chouaib.** Antitumor cytotoxic T-lymphocyte response in human lung carcinoma: identification of a tumor-associated antigen. *Immunol Rev.*, October 2002, vol. 188, 114-21 [0034]

专利名称(译)	基于肿瘤细胞和刺激的白细胞中的mRNA表达谱预测对肿瘤疾病的免疫应答的方法		
公开(公告)号	EP1964931B1	公开(公告)日	2010-12-29
申请号	EP2008008370	申请日	2006-06-08
[标]申请(专利权)人(译)	日立化成研究中心公司 日立化成工业株式会社		
申请(专利权)人(译)	日立化成研究中心, INC. 日立化成工业有限公司.		
当前申请(专利权)人(译)	日立化成研究中心, INC. 日立化成工业有限公司.		
[标]发明人	MITSUHASHI MASATO		
发明人	MITSUHASHI, MASATO		
IPC分类号	C12Q1/68 G01N33/53 G01N33/00		
CPC分类号	G01N33/505 C12Q1/6806 C12Q1/6886 C12Q2600/118 C12Q2600/158 G01N33/574		
代理机构(译)	ELSY, DAVID		
优先权	60/735508 2005-11-11 US 60/688744 2005-06-08 US		
其他公开文献	EP1964931A3 EP1964931A2		
外部链接	Espacenet		

摘要(译)

肿瘤坏死因子 (TNF) 能够通过与其细胞表面上的特异性TNF受体相互作用来诱导细胞凋亡。因为TNF配体和受体的多个成员存在于每个超家族中, 所以存在超过300种不同的配体 - 受体组合。活化的血液白细胞产生TNF作为对癌症的免疫应答的一部分, 以及产生趋化因子以吸引其他白细胞到达该位点。公开了一种方法, 当全血暴露于热聚集的IgG或抗T细胞受体抗体作为免疫系统相互作用的模型时, 检测血液白细胞中各种TNF超家族亚型和趋化因子mRNA的显著诱导。在TNF亚型和诱导的趋化因子中观察到显著的个体间变异。由于外周血白细胞是抗癌免疫细胞的供应, 因此血液中适当的TNF配体和趋化因子的离体诱导能力的定量将在个体化的癌症免疫疗法中 useful。如果肿瘤块很小, 例如早期隐形转移灶, 则适当的TNF攻击可能足以预防复发。




FIGURE 1