



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


<p>(51) International Patent Classification ⁷ : G01N 33/574</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/43789 (43) International Publication Date: 27 July 2000 (27.07.00)</p>
<p>(21) International Application Number: PCT/CA00/00060 (22) International Filing Date: 24 January 2000 (24.01.00) (30) Priority Data: 60/117,031 25 January 1999 (25.01.99) US (71) Applicant (for all designated States except US): PROCREA BIOSCIENCES INC. [CA/CA]; 1100 Beaumont, Suite 305, Ville Mont-Royal, Québec H3P 3H5 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): GAGNÉ, Danièle [CA/CA]; 6280 9ème avenue, Montréal, Québec H1Y 2K7 (CA). GOSSELIN, Diane [CA/CA]; 100, rue de la Plage, Pointe Calumet, Québec J0N 1G2 (CA). HUGO, Patrice [CA/CA]; 820, rue Pierre, Sainte Dorothée/Laval, Québec H7X 3T3 (CA). MIRON, Pierre [CA/CA]; 2474, boulevard des oiseaux, Laval, Québec H7L 4W7 (CA). (74) Agents: MURPHY, Kevin, P. et al.; Swabey Ogilvy Renault, Suite 1600, 1981 McGill College Avenue, Montreal, Québec H3A 2Y3 (CA).</p>	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	

(54) Title: METHOD AND DIAGNOSTIC KIT FOR DIAGNOSIS OF ENDOMETRIOSIS

(57) Abstract

The invention relates to a method and a kit for the diagnosis of endometriosis using blood and endometrial leukocyte markers or a combination thereof. The marker is a surface antigen from endometrial or blood leukocytes.

PREDICTIVE ALGORITHM FOR THE DIAGNOSIS OF ENDOMETRIOSIS



¹ Proportion of cells expressing a specific marker, or a given subset defined by markers within the leukocyte population (CD45+) in the peripheral blood or the stromal fraction of the endometrium.

² A positive test result is given when the proportion of a leukocyte subset fulfills the condition established by the cutoff point.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHOD AND DIAGNOSTIC KIT FOR
DIAGNOSIS OF ENDOMETRIOSIS

BACKGROUND OF THE INVENTION

5 (a) Field of the Invention

The invention relates to a method and a kit for the diagnosis of endometriosis using blood and endometrial leukocyte markers.

(b) Description of Prior Art

10 Endometriosis is one of the most common gynecological disorders, affecting up to 15% of women within reproductive age. It is closely associated with severe pelvic pain, dysmenorrhea, dyspareunia, infertility and several other symptoms such as
15 intraperitoneal bleeding, back pain, constipation and/or diarrhea. It is a major threat to physical, psychological and social integrity of the patients.

Endometriosis is characterized by the implantation and growth of endometrial cells (which
20 normally constitute the lining of the uterus) in extra-uterine sites such as the peritoneal cavity. Although the etiology and pathogenesis of endometriosis remain mainly unclear, the theory of retrograde menstruation is the most widely accepted to
25 explain the presence of ectopic endometrial cells in the peritoneal cavity. However, this phenomenon occurs in most women and, thus, several other factors must be invoked to explain the implantation of endometrial cells and the subsequent development of endometriotic
30 lesions. It is generally believed that initiation of endometriosis implies a complex cascade of events requiring several essential features. Retrogradely seeded endometrial cells must remain viable, be capable of adhering to the mesothelium and of
35 proliferating. Local degradation of the extracellular matrix, as well as extensive vascularization, are also

believed to play an essential role in promoting the invasion of the peritoneal cavity by endometrial cells. Furthermore, once implanted, ectopic endometrial cells must have the capacity to counteract
5 the cytolytic action of the immune system. Indeed, this is supported by the observation of several immunological abnormalities in patients with endometriosis.

At present, direct visualization of the
10 endometriotic lesions under surgical procedures (laparoscopy or laparotomy) is the golden standard and the only reliable method available to diagnose endometriosis. However, this method is highly invasive (i.e. surgery under general anesthesia), costly (i.e.
15 direct cost and indirect cost due to convalescence) and requires a well-trained surgeon who has the ability to identify endometriotic lesions with a variety of appearances. The type of lesions, their size and their localization will determine the stage
20 of the disease (stage I minimal, stage II mild, stage III moderate, stage IV severe). However, there is still no clear consensus on how these parameters correlate with the stage of the disease and the prognostic of endometriosis. In addition, early or
25 minimal endometriosis (which can involve microlesions) can be hardly diagnosed by surgical methods, as they are unlikely to be detected by direct visualization. Indeed, several studies have reported microscopic endometriotic lesions that were not detected
30 laparoscopically. Because the diagnosis of endometriosis by surgical procedures is difficult, costly and invasive, in some cases, several physicians and patients tend to avoid it or at least seriously delay it. Hence, the length of time between the onset
35 of symptoms and the diagnosis can be as long as 8 to

12 years. The possibility to diagnose endometriosis at an early stage would certainly improve the efficacy of the treatments, and reduce dramatically the number of years during which patients endure acute or chronic
5 pain.

Imaging methods such as transvaginal ultrasound and magnetic resonance imaging have been designed for the diagnosis of endometriosis. However, these techniques can only be reliable for the detection of
10 large (> 1 cm diameter) endometriomas lesions detected among a very small proportion of patients with endometriosis. Moreover, the high cost of these techniques has limited their use for the diagnosis of endometriosis.

15 Serum proteins such as CA-125 and placental protein-14 have been proposed as diagnostic markers for endometriosis. Elevated levels of CA-125 have been observed in serum, menstrual effluent and peritoneal fluid of patients with endometriosis. However, these
20 markers, when used alone, are of very limited value for a diagnosis test. Indeed, these markers are not suitable for screening or diagnostic purposes because they provide poor sensitivity. Furthermore, levels of CA-125 and placental protein-14 vary according to
25 several factors such as the assay, the stage of the disease and the menstrual cycle. Finally these markers are known to be modulated by conditions other than endometriosis.

High concentrations of antibodies to
30 endometrial antigens were found in the serum of patients with endometriosis, and thus were proposed as markers for a diagnostic test (International patent application publications WO 94/28021 and WO 92/18535). However, the levels of specificity and sensibility
35 with these tests remain very low. In most cases, the

antigens recognized by these antibodies are still poorly characterized or yet totally unknown.

In U.S. patent No. 5,478,725, low levels of $\alpha v \beta 3$ integrin expression in endometrial samples during the secretory phase of the menstrual cycle is described as a predictor of endometriosis in infertile but not in fertile patients with endometriosis. This observation was associated with milder form of endometriosis (stages I and II) only and, thus, is not useful to detect advanced stages of the disease. Moreover, this method yielded a specificity of 91% but a very low sensitivity (38%).

Taking into account that a number immunological abnormalities have been reported in patients with endometriosis, it is conceivable that the proportion of leukocyte populations and/or their activation status may be modulated during the course of the disease and, thus, may provide some diagnostic value. Previous flow cytometric studies have shown that some T lymphocyte subpopulations (CD8+, CD45+/HLADR+, CD45+/CD3+/HLADR+ or CD3+/CD25+) can be slightly modulated in the peritoneal fluid of subjects with endometriosis relative to normal controls (Oosterlynck D.J., et al., *Am J reprod. Immunol.*, **31**: 25-31, 1994; Becker J.L., et al., *Am J Reprod. Immunol.*, **34**: 179-187, 1995; Wu M.Y., et al., *Am. j. Reprod. Immunol.* **35**: 510-516, 1996). However, these observations have limited value for the diagnosis of endometriosis because peritoneal fluid collection is an invasive, non-conventional procedure. Proportions of leukocyte populations have also been studied in peripheral blood and endometrium of patients with endometriosis. Wu et al., (supra) have reported a modest but significant decrease in the proportion of CD3+ T lymphocytes expressing either CD69 or CD25

activation marker in the blood of patients with advanced endometriosis but not in patients with mild stage of endometriosis or normal controls. This difference was observed in advanced cases of
5 endometriosis only and it was too modest to be used as a diagnostic marker. In contrast, Oosterlynck et al., (Oosterlynck D.J., et al., *Am J reprod. Immunol.*, **31**: 25-31, 1994) and Ho et al. (Ho H.N., et al., *Hum Reprod.*, **97**: 2528-2533, 1997) reported no significant
10 difference in term of T lymphocyte subpopulations when comparing endometriosis subjects with normal controls. These inconsistent results may be explained by the very low number of samples tested in these studies.

Several studies have investigated whether
15 leukocytes are also modulated in eutopic endometrium from patients with endometriosis. Results arising from these studies are contradictory, probably due to the fact that in most cases the methods used were only semi-quantitative and the number of samples tested
20 were very low. For instance, by means of immunohistochemistry, Ota et al. (Ota H., et al., *Am J Reprod. Immunol.*, **35**: 477-482, 1996) have reported that the number of CD3+, CD4+, or CD8+ T lymphocytes, cells bearing adhesion molecules (i.e. ICAM-1, LFA-1,
25 CD2) or CD68+ cells were upregulated in the endometrium of patients with endometriosis compared with infertile controls. In contrast, several other studies using similar techniques have reported no difference in the proportion of T lymphocyte subsets
30 (Klentzeris L.D., et al., *Eur. J Obstet gynecol Reprod Biol.*, **63**:41-47, 1995; Jones R.K., et al., *Fertil Steril*, **66**:81-89, 1996). In addition, a decrease in CD3 positive T cells has been shown by flow cytometry analysis but no difference in the proportion of CD4+,
35 CD8+ stromal leukocytes in the endometrium of patients

with endometriosis compared with fertile controls. When these observations are tentatively used in a diagnostic test, they give only low levels of sensibility and specificity because of a significant
5 overlap between the groups.

Therefore, the diagnostic methods presented in the literature so far do not solve the problems encountered with the diagnosis of endometriosis by surgical procedures. It thus remains imperative to be
10 provided with a less invasive, cheaper and reliable method that could allow detection of females suffering from endometriosis as early as possible.

SUMMARY OF THE INVENTION

15 One aim of the present invention is to provide a less invasive, cheaper and reliable method that could allow detection of females suffering from endometriosis as early as possible.

In accordance with the present invention there
20 is provided a specific blood and/or endometrial leukocyte marker for endometriosis selected from the group consisting of CD3+, CD4+, CD5+, CD8+, CD13+, CD14+, CD20+, CD36+, CD44+, CD56+, CD57+, CD69+, CD122+, HLADR+, CD16+, CD45RA+, CD45RO+, CD56-CD122+,
25 CD3+CD4-CD69+, CD3-CD8+HLADR-, CD3+CD4+, CD3+CD4-, CD3-CD4-, CD3+CD5+, CD3-CD5+, CD3-CD5-, CD3+CD8+, CD3+CD8-, CD3-CD8-, CD3+CD16+, CD3-CD16+, CD3+CD16-, CD3+CD20-, CD3-CD20-, CD3+CD44-, CD3-CD44+, CD3-CD44-, CD3+CD45RA-, CD3-CD45RA-, CD3+CD45RA+, CD3+CD45RO+,
30 CD3-CD45RO+, CD3+CD45RO-, CD3+CD56-, CD3-CD56-, CD3+CD57-, CD3-CD57+, CD3-CD57-, CD3+CD69-, CD3+CD69+, CD3-CD69+, CD3+CD122-, CD3+HLADR+, CD3+HLADR-, CD3-HLADR+, CD3-HLADR-, CD4+CD13+, CD4+CD13-, CD4-CD13+, CD4+CD14-, CD4-CD14-, CD4-CD16-, CD4-CD36+,
35 CD4+CD45RA-, CD4-CD45RA-, CD4+CD45RO+, CD4+CD45RO-,

CD4-CD45RO+, CD4+CD69-, CD4-CD69+, CD4-CD69-,
 CD4+HLADR-, CD4-HLADR+, CD8-CD44+, CD8-CD44-,
 CD8+CD69-, CD8+HLADR-, CD8-HLADR-, CD13+CD16-, CD13-
 CD16+, CD13+CD44+, CD13-CD44-, CD13+CD45RO-, CD13-
 5 CD45RO+, CD13-CD69+, CD13-CD122+, CD13-CD122-,
 CD13+HLADR+, CD13-HLADR+, CD14+CD13+, CD14+CD13-,
 CD14+CD16-, CD14+CD44+, CD14-CD44-, CD14+CD45RO+,
 CD14-CD69-, CD14+CD122-, CD14+HLADR+, CD14-HLADR+,
 CD20-CD5+, CD20-CD5-, CD20+CD44-, CD20-CD44+, CD20-
 10 CD44-, CD20-CD69+, CD20-CD69-, CD20+HLADR+, CD20-
 HLADR+, CD20-HLADR-, CD36-HLADR+, CD56-CD16+, CD56-
 CD16-, CD56-CD44-, CD56+CD69-, CD56-CD69+, CD56-CD69-,
 CD56+CD122+, CD56+CD122-, CD56-CD122-, CD56+HLADR+,
 CD57-CD44+, CD57-CD44-, CD3-CD4-CD44+, CD3-
 15 CD4+CD45RA-, CD3-CD4-CD45RA+, CD3-CD4-CD45RA-, CD3-
 CD4-CD45RO+, CD3-CD8-CD44+, CD3+CD8+CD69-,
 CD3+CD8+HLADR-, CD3+CD8-HLADR+, CD3+CD8-HLADR-, CD3-
 CD8-HLADR-, CD3+CD20-CD5+, CD3+CD20-CD5-, CD3-CD20-
 CD5-, CD3+CD56+CD16+, CD3+CD56-CD16+, CD3+CD56-CD16-,
 20 CD3-CD56+CD16+, CD3-CD56+CD16-, CD3+CD56-CD44+,
 CD3+CD56-CD44-, CD3+CD56-CD122+, CD3+CD56-CD122-, CD3-
 CD56+CD122+, CD3-CD56-CD122-, CD3-CD56-HLADR-, CD3-
 CD57+CD44-, CD3-CD57-CD44+, CD3-CD57-CD44-, CD3+CD57-
 HLADR+, CD4-CD13+CD16+, CD4-CD13-CD16+, CD4-CD13-CD16-
 25 , CD14+CD13+CD16b+, CD14+CD13+CD16b-, CD14+CD13-CD16b-
 , CD14-CD13-HLADR+, CD14-CD13-HLADR-, CD14+CD20+CD44+,
 CD14+CD20+CD44-, CD14+CD20-CD44+, CD14+CD20-CD44-,
 CD14-CD20+CD44-, ratio CD3/CD45RO, Ratio CD13/CD3,
 Ratio CD13/CD8, Ratio CD14/CD3, and Ratio CD14/CD8.

30 Also in accordance with the present invention,
 there is provided a diagnostic method for the
 detection of endometriosis in a patient sample. The
 method comprises the step of detecting at least one
 specific marker as described above, whereby detection

of this specific marker is indicative of endometriosis.

Further in accordance with the present invention, there is provided a diagnostic method for
5 the detection of endometriosis in a patient sample. The method comprises the step of detecting at least two different surface antigens from blood or endometrial leukocytes, whereby detection of at least two different surface antigens is indicative of
10 endometriosis.

In accordance with a further embodiment of the invention, there is provided a diagnostic method for the detection of endometriosis in a patient sample. The method comprises the step of detecting a specific
15 marker combination for endometriosis as defined above, whereby detection of this combination is indicative of endometriosis.

Further in accordance with the present invention, there is provided a diagnostic kit for the
20 detection of endometriosis. The kit comprises an antibody specific for the specific maker as described above. Preferably, the kit comprises at least two different antibodies, each specific for different surface antigens as defined in the specific marker
25 combination defined above. Most preferably, the specific marker combination of the diagnostic kit is selected from the combination described below in Tables 1 and 2.

For the purpose of the present invention, the
30 following symbol "/" is intended to mean a ratio between an expression in front of the symbol and another expression after the symbol.

BRIEF DESCRIPTION OF THE DRAWING

Fig. 1 illustrates a predictive algorithm for the diagnosis of endometriosis.

5 DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, there is provided reliable diagnostic test for endometriosis that is less invasive and less costly than the actual surgical procedure accepted as the golden standard.

10 An extensive study was undertaken by means of flow cytometric analysis, in which the proportion of several blood and endometrial leukocyte subsets was compared in patients with endometriosis and normal controls.

15 The present invention identifies a series of leukocyte subsets that can be used as markers in a diagnostic test for endometriosis. These leukocyte subsets are defined according to the expression of cell surface antigens. Several cell surface antigens
20 may define the same population of cells, and thus they are included in the present invention.

Any other antibodies or molecules recognizing the same antigen or a different epitope, isoform, subunit, chain, glycosylation or phosphorylation form
25 or an allelic variant of the same antigen, a member of the same complex, or an antigen with the same cell distribution is also included in the present invention.

Further in accordance with the present
30 invention, there is provided examples showing how at least two different surface antigens from blood and/or endometrial leukocytes can be used in combinations in a diagnostic test for endometriosis (Tables 1 and 2).

TABLE 1

Levels of sensibility and specificity provided by several examples of endometrial and/or blood marker combinations used as a diagnostic method for endometriosis

5

<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
Endometrial leukocyte markers			
1.			
CD4+ (<17%) ²	>67%	90%	67%
CD8+CD69- (<21%)			
CD13+CD45RO- (<17.5%)			
2.			
CD4+ (<15.5%)	>60%	89%	65%
CD8+CD69- (<21%)			
CD56+CD122- (>19%)			
CD3+CD45RA- (<35%)			
CD13+CD45RO- (<17.5%)			
3.			
CD4+ (<17%)	>67%	88%	65%
CD8+CD69- (<21%)			
CD13-CD122+ (>28%)			
CD13+CD45RO- (<17.5%)			
4.			
CD4+ (<17%)	>67%	89%	63%
CD8+CD69- (<21%)			
CD14+CD13-CD16b- (>14.5%)			
5.			
CD3+CD16- (<40%)	>55%	84%	62%
CD13+CD45RO- (<13.5%)			
CD3+ (<40%)			
CD8+ (<20%)			
CD3+CD69+ (<15%)			
6.			
CD3+ (<40%)	>65%	84%	63%
CD3+CD8+ (<16%)			
CD13+CD45RO- (<17.5%)			
CD3-CD20- (>56%)			
7.			
CD3+CD8+ (<16%)	>65%	81%	65%
CD13+CD45RO- (<17.5%)			
CD3+CD5+ (<37%)			
CD3+CD122- (<42.5%)			
CD3-CD20- (>56%)			
CD3+CD45RO- (<30%)			

<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
8. CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD3+CD5+ (<37%) CD3+CD122- (<42.5%)	>60%	82%	64%
9. CD3+CD20-CD5- (>7.7%) CD4+CD13- (<20.5%) CD56-CD122- (<47%)	>60%	81%	66%
10. CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD4+CD45RA- (<16%) CD3+CD45RO- (<30%)	>60%	80%	65%
11. CD3+ (<40%) CD8+CD69- (<18%) CD3-CD4-CD45RO+ (>56%) Ratio CD13+/CD3+ (>0.675%) CD13+CD45RO- (<21%)	>35%	79%	67%
12. CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD3-CD5- (>54%) CD20-CD5+ (<44%)	>70%	81%	61%
13. CD8+ (<20%) ² CD5+ (<37%) CD3-CD20- (>58%) CD3-HLADR- (>54.5%)	>51%	81%	60%
14. CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD5+ (<40%)	>60%	81%	60%
15. CD4+ (<17%) CD13-CD122+ (>28%) CD8+CD69- (<19.5%) CD3+CD45RA- (<37%)	>50%	76%	71%

<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
16. CD4+ (<15.5%) CD8+CD69- (<21%) CD13-CD122+ (>28%) CD3+CD45RA- (<35%) CD13+CD45RO- (<17.5%)	>35%	71%	78%
17. CD4+ (<15.5%) CD8+CD69- (<21%) CD56-CD122- (<47%) CD3+CD45RA- (<35%) CD13+CD45RO- (<17.5%)	>40%	70%	78%
18. CD3+ (<40%) CD4+ (<17%) CD3+CD8+ (<16%) CD13+CD45RO- (<21%) CD3+CD5+ (<37%)	>35%	72%	76%
19. CD3+ (<40%) CD4+ (<17%) CD3+CD8+ (<16%) CD13+CD45RO- (<21%)	>40%	74%	74%
20. CD3+ (<40%) CD3+CD8+ (<16%) CD13+CD45RO- (<21%)	>40%	75%	73%
21. CD3+ (<40%) CD5+ (<40%) CD3+CD5+ (<37%) CD69+ (<33%) CD4-CD69+ (<35%)	>25%	71%	69%
22. CD3+ (<40%) CD3+CD8+ (<13.5%) CD13+CD45RO- (<17.5%)	>30%	68%	83%
23. CD3+ (<40%) CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%)	>30%	61%	86%

<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
24. CD3+ (<40%) CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD3-CD20- (>58%) CD56-CD16- (<46%)	>22%	62%	80%
25. CD3+CD16- (<47.5%) CD3-CD4-CD45RO+ (>31.5%) CD3+ (<40%) CD8+ (<20%) CD3+CD69+ (<15%) CD13+CD45RO- (<17.5%)	>45%	66%	79%
26. CD3+ (<40%) CD3-CD45RO+ (>15%) Ratio CD13+/CD3+ (>0.675%) CD3+CD8+ (<16%) CD8+CD69- (<21%)	>40%	66%	75%
27. CD3+CD20-CD5- (>7.7%) CD4+CD13- (<20.5%) CD56-CD122- (<47%) CD4+CD45RO- (<16%)	>45%	61%	80%
28. CD3+ (<40%) CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD3-CD20- (>58%)	>20%	61%	86%
29. CD13+CD45RO- (<17.5%) CD4+CD45RA- (<16%) CD3+CD122- (42.5%) CD8+CD69- (<21%)	>70%	90%	54%
30. CD3+CD8+ (<16%) CD13+CD45RO- (<17.5%) CD4+CD45RA- (<16%) CD3+CD45RO- (<30%) CD3-CD5- (>54%)	>70%	84%	60%

<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
Blood leukocyte markers			
31.			
CD3-CD5+ (>14.5%)	>55%	66%	60%
CD3-CD45RA- (>14.5%)			
CD3-CD44+ (>13%)			
CD13+ (>17.5%)			
CD3-CD57-CD44- (<41.3%)			
32.			
CD3-CD45RA- (>17%)	>22%	61%	64%
CD20-CD44+ (>17%)			
CD20-HLADR+ (>20%)			
CD3-CD4-CD44+ (>40.5%)			
CD36-HLADR+ (<5.6%)			
33.			
CD3-CD45RA- (>14.5%)	>40%	62%	64%
CD3-CD45RO+ (>19%)			
CD20-HLADR+ (>14.5%)			
Blood (in italics) and endometrial leukocyte markers			
34.			
<i>CD57+</i> (>10%)	>50%	76%	72%
<i>CD14+</i> (>10%)			
<i>CD3-CD69+</i> (>17.5%)			
CD3+ (<40%)			
CD4+ (<15.5%)			
CD3+CD8+HLADR- (<35%)			
35.			
<i>CD3-CD69+</i> (>17.5%)	>33%	70%	79%
CD3+ (<40%)			
CD4+ (<15.5%)			
CD3+CD8+HLADR- (<35%)			

<u>Marker combination</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>
36. CD4-CD36+ (>14.4%) CD3-CD69+ (>17.5%) CD8+ (<20%) CD13+ (>29%) CD3+ (<40%) CD16+ (>27%) CD69+(<33%) CD5+ (<40%)	>43%	70%	74%
37. CD3-CD45RA- (>14.5%) CD3-CD45RO+ (>19%) CD20-HLADR+ (>14.5%) CD14+CD44+ (>15%) CD8+ (<20%) CD5+ (<37%) CD3-CD20- (>58%) CD3-HLADR- (>54.5%)	>50%	73%	71%

¹ Value above which a diagnosis of endometriosis is given.

² Cutoff point established for each individual marker.

5

TABLE 2

Examples of logistic regression models provided by endometrial or blood leukocyte markers for the identification of patients with endometriosis

<u>Marker combination</u>	<u>B value</u>	<u>Threshold value¹</u>	<u>Specificity</u>	<u>Sensibility</u>	<u>Number of sample tested</u>
<u>Endometrial leukocyte markers</u>					
<u>Combination no. 1</u>					
1. CD3+ (<40%) ²	-7.9747	>.55	83%	79%	41
2. CD3-CD5- (>60%)	7.2921				
3. CD13+CD45RO- (<17.5%)	-0.1410 -1.6259				
4. CD3-CD20- (>58%)	9.5142				
Interaction of 1 to 4 Constant = 2.0516					

Marker combination	B value	Threshold value ¹	Specificity	Sensibility	Number of sample tested
<u>Combination no. 2</u>					
1. CD3+ (<40%)	-6.7753	>.55	74%	73%	67
2. CD3-CD5- (>60%)	5.8240				
3. CD13+CD45RO- (<17.5%)	-1.9298 -0.0262				
Interaction 1 to 4	2.8385				
Constant = 2.7910					
<u>Combination no. 3</u>					
1. CD3+CD8+ (<16%)	-0.1308	>.50	84%	72%	51
2. CD13+CD45RO- (<17.5%)	-2.6688 -1.1778				
3. CD3+CD5+ (<37%)					
Constant = 3.1417					
<u>Combination no. 4</u>					
1. CD3+ (<40%)	-1.6965	>.50	78%	75%	81
2. Length of menstruation (>7days)	-1.8160 -1.9656				
3. CD13+CD20- (<21%)	10.3064				
4. Pelvic pain ³					
Constant = 3.1984					
<u>Blood leukocyte markers</u>					
<u>Combination no. 1</u>					
1. CD14+CD44+ (>15%)	0.9298	>0.55	80%	70%	140
2. CD57+ (>10%)	0.7423				
3. CD3-CD45RA- (>12%)	-0.8147				
4. CD14+ (>10%)	0.8629				
<u>Combination no. 2</u>					
1. CD14+ (>10%)	10.5891	>.50	65%	71%	125
2. CD57+ (>10%)	0.7326				
3. CD3+CD69+ (>17.5%)	0.6899				
4. CD3+HLADR+ (<4%)	1.2004				
5. CD3-CD45RA- (>12%)	-0.1137				
Constant = -1.2062					

Marker combination	B value	Threshold value ¹	Specificity	Sensibility	Number of sample tested
<u>Combination no.3</u>					
1. CD14+ (>10%)	1.1994	>.55	76%	75%	142
2. CD57+ (>10%)	0.8080				
3. CD3+HLADR- (<4%)	1.3593				
4. CD3-CD45RA- (>12%)	-0.63				
5. Pelvic pain	2.1506				
6. Length of menstruation (>7d)	.7489				
Constant = -1.771					
<u>Combination no.4</u>					
1. CD14+ (>10%)	.9727	>.50	71%	78%	141
2. CD57+ (>10%)	.4489				
3. CD3+CD69+ (>17.5%)	.8129 1.3368				
4. CD3+HLADR- (<4%)	-0.8805				
5. CD3-CD45RA- (>12%)	2.1574				
6. Pelvic pain	1.5164				
7. Age (>40)					
Constant = -1.7686					

¹ Value above which a diagnosis of endometriosis is given.

² Cutoff point established for each individual marker.

³ presence of pain at any time other than menstruation and intercourse

The predictive models for endometriosis were established according to the following equation:

$$P(r) = \frac{e^{c + B1*(marker1) + B2*(marker2) + \dots Bn*(marker n)}}{1 + e^{c + B1*(marker1) + B2*(marker2) + \dots Bn*(marker n)}}$$

Where: P(r) = probability of having endometriosis;

c = constant established for a particular combination;

B = coefficient of regression; and

n = total number of markers in the combination.

In the present invention, a series of endometrial and peripheral blood leukocyte subpopulations for which proportions were modulated in

patients with endometriosis (stage I-IV; I-II or III-IV) compared with those of normal controls, have been identified. The novelty of the present invention is to use these leukocyte subpopulations, either alone or in combination, as markers for the diagnosis of endometriosis. Moreover, risk factors for endometriosis identified amongst personal information and menstrual characteristics were shown to be of significant value when used in combination with blood or endometrial leukocyte subsets in a predictive test for endometriosis.

Two methods were used for the combination of markers.

Method 1

A cutoff point is established for the proportion of each leukocyte markers in order to obtain the best discrimination between patients with endometriosis and controls. The proportion obtained for each marker is compared to the cutoff point. A positive test result gives a score of 1, whereas a negative test result gives a score of 0. The diagnostic value is obtained by adding the scores of all the markers of a particular combination and converting it in percentage. The final diagnostic value is then compared to a threshold value that was established to provide the best levels of sensibility and specificity. A positive diagnosis of endometriosis is given when the final diagnostic value exceeds the threshold value established for a particular combination of markers. On the opposite, a negative diagnosis of endometriosis is given when the final diagnostic value is lower than the threshold value (see Fig. 1).

Method 2

A predictive model for endometriosis is established by including each marker of a particular combination in the following logistic regression
 5 equation:

$$P(r) = \frac{e^{c + B1*(marker1) + B2*(marker2) + \dots + Bn*(marker n)}}{1 + e^{c + B1*(marker1) + B2*(marker2) + \dots + Bn*(marker n)}}$$

10 Where:

P(r) = probability of having endometriosis;
 c = constant established for a particular combination;
 B = coefficient of regression; and
 n = total number of markers in the combination.

15

The probability of having endometriosis (P(r)) is then compared to a threshold value that provides the best discriminative value. A positive diagnosis of endometriosis is given when the P(r) value exceeds the
 20 threshold value established for a particular combination of markers. Alternatively, a negative diagnosis of endometriosis is given when the P(r) value is lower than the threshold value.

In the present invention, there is reported a
 25 series of 102 endometrial CD45+ leukocyte populations and 93 blood mononuclear CD45+ leukocyte populations which were shown by flow cytometric analysis to be modulated in patients with endometriosis compared with normal controls and, thus are candidate markers for
 30 the diagnosis of endometriosis (Tables 3, 4, 5, and 6). An innovative feature of the present invention is to use these markers in combination to increase their level of sensibility and specificity in the diagnostic test.

TABLE 3
Endometrial leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Endo stage I-IV	P ¹	Number of samples tested		area under ROC curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo			Control	Endo							
CD3+	47.7 ± 12.3	38.7 ± 12.6	3.37x10 ⁻⁵	58	88	0.703	3.3X10 ⁻⁵	<40	84	55	6.5	(2.9-14.9)	
CD4+	18.3 ± 5.6	15.7 ± 5.9	0.008	55	88	0.632	0.008	<17 <15.5	63 72	63 52	3.0 2.9	(1.5-6.2) (1.4-6.0)	
CD5+	45.1 ± 11.6	36.3 ± 12.3	1.6x10 ⁻⁴	46	74	0.702	0.0002	<37 <40	80 73	53 58	4.2 3.9	(1.8-9.7) (1.7-8.8)	
CD8+	24.3 ± 8.5	18.5 ± 8.6	1.4x10 ⁻⁴	54	87	0.688	0.00019	<20	74	62	4.8	(2.3-10.2)	
CD3+CD4+	17.2 ± 5.5	14.4 ± 5.7	0.004	55	88	0.641	0.004	<15	67	57	2.7	(1.3-5.5)	
CD3+CD4-	29.9 ± 9.6	23.6 ± 9.1	1.2x10 ⁻⁴	55	88	0.687	0.00017	<24	80	52	4.4	(2.0-9.6)	
CD3-CD4-	51.1 ± 14.2	60.8 ± 12.6	4.1x10 ⁻⁵	55	88	0.698	6.9x10 ⁻⁵	>61	83	50	5.1	(2.2-11.7)	
CD3+CD8+	18.9 ± 7.5	13.7 ± 7.7	1.1x10 ⁻⁴	54	84	0.714	2.3 x10 ⁻⁵	<16 <13.5	70 81	70 54	5.6 5.3	(2.6-11.8) (2.3-11.9)	
CD3+CD8-	26.1 ± 7.8	23.1 ± 7.1	0.022	54	84	0.609		<23.5	68	51	2.2	(1.1-4.7)	
CD3-CD8-	49.6 ± 12.0	58.2 ± 13.0	1.3x10 ⁻⁴	54	84	0.688	1.9 x10 ⁻⁴	>53.5	70	63	4.0	(1.9-8.5)	
CD3+CD69+	20.4 ± 9.6	15.5 ± 8.0	0.003	44	76	0.642	0.010	<15	67	53	2.4	(1.1-5.2)	
CD3+CD122-	41.4 ± 10.0	34.4 ± 12.4	0.011	29	53	0.669	0.012	<42.5	64	76	2.9	(1.1-7.5)	
CD3+HLADR-	38.1 ± 10.3	30.6 ± 12.3	2.8x10 ⁻⁴	51	80	0.681	0.0005	<35	72	63	4.0	(1.8-8.5)	
CD3-III-ADR-	46.4 ± 13.0	55.6 ± 13.5	1.9x10 ⁻⁴	51	80	0.693	0.0002	>54.5	80	51	4.1	(1.8-9.3)	

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Endo stage I-IV	P ¹	Number of samples tested		area under ROC curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo			Control	Endo							
CD3+CD45RA+	7.4 ± 4.7	5.7 ± 3.0	0.018		56	85	0.608	0.030	<4.9	77	40	2.2	(1.0-4.7)
CD3+CD45RA-	40.3 ± 11.2	32.7 ± 12.0	2.5x10 ⁻⁴		56	85	0.684	0.0002	<37	69	66	4.7	(2.2-9.7)
CD3-CD45RA-	31.4 ± 12.7	39.6 ± 13.4	4.2x10 ⁻⁴		56	85	0.667	8.2 x 10 ⁻⁴	>32	73	60	4.1	(1.9-8.5)
CD3+CD45RO-	31.0 ± 11.1	25.0 ± 9.8	0.002		50	73	0.661	0.002	<28	65	62	3.1	(1.4-6.6)
CD3+CD16-	45.5 ± 12.1	37.2 ± 13.1	2.1x10 ⁻⁴		57	83	0.680	0.0003	<30	55	70	2.7	(1.3-5.7)
CD3+CD56-	46.5 ± 12.2	38.5 ± 12.8	3.2x10 ⁻⁴		56	83	0.674	0.00053	<38	80	49	4.5	(2.0-9.9)
CD3+CD5+	41.9 ± 11.4	33.3 ± 12.3	2.1x10 ⁻⁴		45	74	0.695	0.00036	<40	77	58	4.2	(2.0-8.9)
CD3-CD5-	50.6 ± 12.0	59.0 ± 12.9	0.001		45	74	0.690	0.00052	<47.5	82	66	4.3	(1.9-9.6)
CD4+CD69-	16.4 ± 4.8	13.8 ± 5.1	0.012		37	72	0.648	0.012	<14	66	66	2.6	(1.2-5.6)
CD4+CD45RA-	16.7 ± 5.3	14.2 ± 5.7	0.010		54	85	0.632	0.009	<16	78	53	4.1	(1.6-10.1)
CD8+CD69-	24.0 ± 7.9	18.9 ± 8.3	0.007		30	59	0.687	0.004	>60	62	66	3.3	(1.6-6.7)
CD8+HLADR-	23.3 ± 7.7	18.1 ± 8.5	0.001		49	79	0.673	0.001	>54	83	53	5.9	(1.9-17.6)
CD8-HLADR-	61.6 ± 9.5	68.1 ± 9.5	2.2x10 ⁻⁴		49	79	0.675	0.0009	<18	76	59	5.1	(1.9-13.9)
CD13-CD122+	27.0 ± 9.8	33.6 ± 18.7	0.031		32	58	0.605		<21	65	68	4.2	(1.6-10.7)
									<18	77	54	4.1	(1.8-9.2)
									>61.5	52	77	3.8	(1.7-8.3)
									>28	64	59	2.5	(1.0-6.2)

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Endo stage I-IV	P ¹	Number of samples tested		area under ROC curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo			Control	Endo							
CD13-CD122-	47.1 ± 14.4	40.4 ± 15.0	0.043	32	58	0.635	0.035	<46	58	64	2.6	(1.1-6.2)	
CD20-CD5+	44.6 ± 12.0	36.4 ± 12.6	0.001	41	66	0.681	0.002	<41	60	62	2.6	(1.2-5.7)	
CD20-CD5-	52.0 ± 12.8	60.9 ± 12.9	0.001	41	66	0.692	0.0009	>60	57	71	3.5	(1.5-7.9)	
CD56-CD16+	22.0 ± 12.2	27.0 ± 16.6	0.044	56	84	0.571			77	50	3.6	(1.5-8.6)	
CD56-CD16-	51.6 ± 12.2	42.8 ± 13.5	1.2x10 ⁻⁴	56	84	0.687	1.8 x 10 ⁻⁴	<46	71	57	3.3	(1.6-6.7)	
ratio CD3+/CD45RO+	1.5 ± 1.0	1.2 ± 0.7	0.020	51	80	0.626	0.015						
CD14+CD13-	1.4 ± 0.9	2.3 ± 1.8	0.041	21	36	0.639							
CD3+CD20-	44.4 ± 11.2	36.9 ± 13.5	0.024	24	45	0.667	0.023	<40	78	58	4.1	(1.4-12.3)	
CD3-CD20-	52.3 ± 11.5	61.0 ± 14.9	0.016	24	45	0.669	0.022	>58	83	53	6.2	(1.8-21.2)	
CD3-CD4-CD45RA+	40.1 ± 13.7	34.9 ± 15.1	0.046	51	79	0.618	0.023	>56	70	60	4.5	(1.4-13.5)	
CD3-CD4-CD45RA-	57.9 ± 14.1	63.3 ± 15.2	0.042	51	79	0.620	0.021						
CD3+CD8+CD69-	40.4 ± 10.2	35.1 ± 11.7	0.039	29	56	0.635	0.042	<34.5	75	52	3.4	(1.2-9.2)	
CD3+CD8+HLADR-	39.5 ± 9.2	33.6 ± 11.5	0.003	48	74	0.665	0.002	<35	72	55	3.5	(1.6-7.7)	
CD3+CD8-HLADR-	43.1 ± 7.5	46.3 ± 10.9		48	74	0.603		>47	69	50	2.2	(1.0-4.7)	
CD3-CD8-HLADR-	76.7 ± 8.9	80.3 ± 7.9	0.021	48	74	0.597							
CD14+CD13-CD16b-	19.8 ± 16.4	33.4 ± 22.9	0.026	19	36	0.703	0.014	>23	68	69	4.9	(1.4-16.3)	
								>14.5	53	83	4.6	(1.4-15.6)	

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P ¹	area under ROC curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo stage I-IV	Control	Endo								
CD4+CD14-	20.7 ± 7.8	14.8 ± 5.8	14	23	0.014	0.738	0.017	<16.6	79	65	6.9	(1.5-32.0)
CD4-CD14-	75.7 ± 8.1	81.0 ± 5.7	14	23	0.025	0.711	0.033	>76	57	83	6.3	(1.4-28.7)
CD4+HLADR-	16.0 ± 5.7	12.3 ± 4.1	14	30	0.018	0.715	0.023	<17	57	87	8.7	(1.9-38.6)
CD13-CD69+	54.2 ± 14.3	42.3 ± 18.2	14	30	0.039	0.705	0.030	<51	71	70	5.8	(1.4-23.6)
CD13+CD45RO-	22.1 ± 8.8	15.5 ± 10.7	25	54	0.009	0.746	4.6 x 10 ⁻⁴	<17.5	76	70	7.5	(2.5-22.3)
CD56-CD122-	49.0 ± 12.2	42.8 ± 15.8	29	51	0.014	0.631		<21	52	80	4.2	(1.5-11.8)
CD3+CD69-	26.4 ± 7.0	23.3 ± 8.4	44	76	0.036	0.612	0.041	<27	57	67	2.2	(1.0-4.8)
CD4+CD45RO-	14.9 ± 5.3	13.0 ± 6.2	47	74		0.615	0.034	<16	51	72	2.6	(1.2-5.6)
CD56+CD122+	3.5 ± 2.2	2.6 ± 1.7	29	51	0.049	0.651	0.025	<3.0	55	73	3.3	(1.3-8.5)
CD3-CD56+CD16+	8.2 ± 3.8	6.8 ± 4.0	53	78	0.049	0.615	0.026	<6.5	72	55	3.0	(1.4-6.2)
CD3-CD56+CD122+	3.6 ± 2.0	2.6 ± 1.8	27	49	0.033	0.638	0.048	<2.7	63	65	3.2	(1.2-8.5)
CD14+CD13+	3.8 ± 1.9	3.0 ± 2.6	21	36		0.672	0.032	<2.3	81	53	4.8	(1.3-16.9)
CD3+CD20-CD5+	93.0 ± 2.7	88.0 ± 13.8	24	40		0.659	0.034	<91.5	79	55	4.6	(1.4-14.9)
CD4-CD13+CD16+	31.0 ± 14.0	22.4 ± 18.9	16	36		0.699	0.023					
CD69+	41.8 ± 12.9	38.8 ± 17.8	43	78		0.557		<33	81	41	3.0	(1.2-7.4)
ratio CD13+/CD3+	0.56 ± 0.54	0.78 ± 0.71	46	78		0.596		>0.68	80	40	2.6	(1.1-6.1)

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P ³	area under ROC curve ²	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo stage I-IV	Control	Endo							
CD3-CD20-CD5-	88.0 ± 6.9	90.7 ± 5.2	24	40		0.598	>84	37	90	5.4	(1.4-20.3)
CD3+CD20-CD5-	5.1 ± 2.3	9.5 ± 13.6	24	40	0.015	0.683	>7.7	87	50	9.9	(2.1-48.1)
CD4+CD13-	17.5 ± 6.9	15.8 ± 5.4	36	63		0.594	<20.5	42	86	4.3	(1.6-11.3)
CD3+CD44-	41.7 ± 12.0	38.3 ± 13.6	31	56		0.596	<37.8	74	50	2.9	(1.1-7.5)
CD56+	26.2 ± 12.5	30.2 ± 17.2	57	87		0.562	>32	81	41	2.9	(1.3-6.5)
CD13-CD45RO+	21.4 ± 8.7	26.3 ± 11.3	25	54		0.625	>28	80	45	3.2	(1.0-9.8)
CD56+CD69-	19.6 ± 12.9	24.3 ± 14.3	33	53		0.610	>26	85	40	3.4	(1.1-10.2)
CD13-CD16+	8.0 ± 7.5	6.7 ± 3.3	39	71		0.562	<6	72	51	2.6	(1.1-6.1)
CD56+CD122-	21.8 ± 11.8	28.7 ± 17.3	29	51		0.621	>19	59	71	3.1	(1.2-7.9)
CD3+CD4-CD69+	37.2 ± 10.0	34.0 ± 13.8	34	66		0.572	<33.5	73	49	2.6	(1.1-6.4)
CD4-CD13-CD16+	8.5 ± 3.6	8.7 ± 10.8	16	36		0.642	<7.1	75	58	4.2	(1.1-15.6)
CD4-CD13-CD16-	54.5 ± 13.0	62.2 ± 21.4	16	36		0.655	>65	81	56	5.4	(1.3-22.3)

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		p ¹	area under ROC curve ²	p ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Controls	Endo stage I-IV	Control	Endo								
CD14+CD13+CD16b+	11.0 ± 12.5	6.5 ± 6.2	19	36		0.616		<16	32	94	7.8	(1.4-43.9)
CD4-CD69-	40.7 ± 12.9	46.4 ± 18.6	37	72		0.593		>47	76	47	2.8	(1.2-6.7)
CD3-CD45RO+	23.0 ± 12.5	27.1 ± 16.3	50	73		0.556		>15	34	80		
CD4-CD69+	39.3 ± 11.8	37.7 ± 18.0	37	72		0.462		<35	68	46		
CD3-CD4-CD45RO+	41.3 ± 18.1	43.4 ± 20.9	43	72		0.530		>31.5 >56	33 81	71 28		

¹ P value (when ≤ 0.05) obtained in a student "t" test when mean proportion found in patients with endometriosis stage I-IV was compared to normal controls.
² Discriminative value of each marker established by area under ROC curve.
³ P value (when ≤ 0.05), significance of area under ROC curve.
⁴ Confidence interval for odds ratio.

TABLE 4
Peripheral blood leukocyte populations proposed as good predictive markers for the identification of patients with endometriosis

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P ¹	area under curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Control	Endo. Stage I-IV	Control	Endo. Stage I-IV								
CD3+	66.6 ± 8.5	64.5 ± 8.7	132	172	0.032	0.570	0.037					
CD8+	17.3 ± 5.2	16.4 ± 4.8	129	172		0.549		<18.9	33	77	1.7	(1.0-2.8)
CD13+	16.0 ± 6.0	17.6 ± 6.5	122	155	0.039	0.575	0.032	>17.5	63	51	1.8	(1.1-2.9)
CD14+	11.8 ± 4.9	13.4 ± 6.0	124	167	0.020	0.575	0.029	>10	45	71	2.0	(1.3-3.3)
CD20+	5.7 ± 3.1	4.8 ± 2.3	124	162	0.006	0.582	0.017	<6	39	74	1.8	(1.1-3.0)
CD36+	15.7 ± 6.8	17.2 ± 7.3	112	140		0.560		>19	77	37	2.1	(1.2-3.6)
CD44+	17.1 ± 5.6	19.1 ± 6.5	113	148	0.009	0.585	0.018	>18.5	61	51	1.7	(1.0-2.7)
CD57+	8.0 ± 3.9	9.2 ± 4.9	114	148	0.023	0.569		>10	75	39	1.8	(1.1-3.1)
CD69+	19.4 ± 8.2	21.0 ± 7.1	109	144		0.590	0.014	>21.5	71	45	2.0	(1.2-3.4)
CD122+	29.2 ± 8.4	31.2 ± 11.7	122	166		0.567		>34	74	42	2.1	(1.2-3.4)
CD3+CD5+	66.6 ± 8.5	63.7 ± 10.4	115	146	0.017	0.586	0.017	<69	44	70	1.8	(1.1-2.9)
CD3+CD45RA-	39.3 ± 9.5	37.2 ± 8.3	124	168	0.044	0.583	0.015	<42	40	72	1.7	(1.1-2.8)
CD3+CD56-	65.3 ± 8.8	63.2 ± 8.6	126	169	0.035	0.571	0.037	<68	42	71	1.8	(1.1-2.9)
CD3+CD57-	63.7 ± 8.3	60.7 ± 9.7	113	146	0.009	0.592	0.011	<67	40	77	2.3	(1.3-3.9)
CD3+CD69-	60.1 ± 9.6	57.9 ± 9.2	107	141		0.584	0.023					
CD3+CD122-	62.2 ± 8.4	59.6 ± 9.8	121	164	0.021	0.578	0.024	<58	69	42	1.6	(1.0-2.7)
CD3+HLADR+	3.9 ± 1.4	3.5 ± 1.2	121	154	0.006	0.601	0.004	<4	40	77	2.2	(1.3-3.7)
CD3-CD5+	15.5 ± 5.3	18.3 ± 8.4	115	146	0.003	0.619	0.001	>14.5	50	72	2.5	(1.5-4.2)
CD3-CD16+	23.3 ± 7.9	25.4 ± 8.7	123	166	0.036	0.572	0.037					
CD3-CD44+	13.1 ± 5.3	15.6 ± 6.3	111	143	0.001	0.615	0.002	>11.5	44	73	2.2	(1.4-3.7)
CD3-CD57+	4.2 ± 2.5	5.0 ± 3.1	113	146	0.023	0.573	0.044					
CD3-CD69+	14.2 ± 5.6	16.4 ± 6.5	107	141	0.006	0.602	0.006	>17.5	75	41	2.1	(1.2-3.6)
CD3-CD45RO+	16.0 ± 5.6	18.1 ± 6.8	117	148	0.006	0.595	0.008	>19	77	44	2.6	(1.5-4.5)
CD3-CD4-	31.1 ± 9.5	33.3 ± 9.0	132	171	0.043	0.566	0.050					
CD3-CD8-	33.4 ± 9.2	35.5 ± 10.0	122	166		0.568	0.050					
CD3-CD45RA-	14.2 ± 5.2	16.2 ± 6.2	124	168	0.004	0.595	0.006	>14.5	61	60	2.3	(1.4-3.7)
CD3-CD56-	21.6 ± 6.3	23.1 ± 7.0	126	169		0.562		>25	73	39	1.7	(1.0-2.8)
CD4-CD13+	14.5 ± 6.0	16.3 ± 6.8	108	131	0.030	0.586	0.023	>16.5	69	51	2.4	(1.4-4.1)
CD4-CD36+	13.8 ± 7.4	19.0 ± 6.2	13	21	0.037	0.771	0.009	>19	92	62	19.5	(2.1-179.9)

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P ¹	area under curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
	Control	Endo. Stage I-IV	Control	Endo. Stage I-IV								
CD4-CD69+	16.0 ± 6.1	18.1 ± 6.5	94	120	0.021	0.603	0.009	>19	75	45	2.4	(1.3-4.3)
CD4-CD45RO+	23.4 ± 6.7	27.0 ± 7.7	27	50	0.043	0.620						
CD4-CD45RA-	22.7 ± 6.5	24.4 ± 8.3	125	168		0.562						
CD8-CD44+	16.8 ± 5.5	18.9 ± 6.7	88	119	0.017	0.588	0.030	<68	48	70	2.1	(1.2-3.7)
CD8-CD44-	66.4 ± 7.0	64.4 ± 6.6	88	119	0.039	0.584	0.040	>11	38	76	1.9	(1.1-3.4)
CD13+CD44+	13.3 ± 5.3	15.4 ± 5.9	96	121	0.006	0.605	0.008	>15.5	71	42	1.8	(1.1-3.1)
CD13+HLADR+	13.0 ± 5.3	14.7 ± 6.1	108	135	0.024	0.581	0.031	>2	76	57	4.2	(1.3-13.9)
CD13+CD16-	1.4 ± 0.9	3.7 ± 4.6	21	37	0.005	0.727	0.004	<8	50	70	2.4	(1.4-4.0)
CD13+HLADR+	8.2 ± 3.2	7.0 ± 3.0	108	135	0.005	0.619	0.001	<82	39	76	2.0	(1.1-3.6)
CD13-CD44-	79.9 ± 6.2	76.9 ± 8.6	96	121	0.004	0.606	0.007	>9.5	48	70	2.2	(1.3-3.6)
CD14+HLADR+	11.0 ± 4.6	12.7 ± 5.9	109	147	0.009	0.587	0.017	>15	85	33	2.7	(1.3-5.4)
CD14+CD44+	10.7 ± 4.9	12.9 ± 5.7	85	123	0.003	0.612	0.006					
CD14+CD45RO+	12.2 ± 4.4	14.3 ± 6.1	102	118	0.004	0.586	0.028					
CD14+CD16-	0.7 ± 0.8	2.7 ± 5.4	23	41	0.023	0.603		>0.7	86	47	5.3	(1.0-27.5)
CD14+CD122-	0.4 ± 0.3	2.4 ± 5.2	14	34	0.029	0.648		<8.5	69	47	2.0	(1.2-3.3)
CD14+HLADR+	10.3 ± 3.3	9.1 ± 2.7	109	147	0.001	0.611	0.002	<80	79	37	2.2	(1.2-4.2)
CD14-CD44-	83.5 ± 5.3	81.6 ± 6.3	85	123	0.023	0.588	0.032	<5.5	39	75	1.9	(1.1-3.3)
CD20+HLADR+	5.2 ± 2.9	4.3 ± 2.1	106	141	0.005	0.596	0.010	<4	62	55	2.0	(1.2-3.4)
CD20+CD44-	5.0 ± 2.8	4.2 ± 2.2	95	126	0.033	0.571		>17	74	47	2.5	(1.4-4.4)
CD20-CD44+	14.4 ± 5.1	17.3 ± 6.2	95	126	0.0004	0.636	0.001	>16	57	76	4.2	(1.1-16.3)
CD20-CD69+	16.4 ± 4.3	21.0 ± 6.2	14	29	0.016	0.719	0.021	>14.5	49	70	2.2	(1.3-3.7)
CD20-HLADR+	15.4 ± 4.9	17.6 ± 6.9	106	141	0.004	0.591	0.014					
CD20-CD44-	79.5 ± 5.1	77.5 ± 6.1	95	126	0.008	0.590	0.022	<75.6	71	52	11.6	(2.3-57.0)
CD20-CD69-	79.4 ± 6.7	75.3 ± 6.5	14	29	0.065	0.711	0.027	<5.6	77	40	2.2	(1.2-4.0)
CD36-HLADR+	7.8 ± 3.2	6.4 ± 2.0	95	121	0.0005	0.629	0.001	>20.5	85	52	6.0	(1.1-33.3)
CD56-CD69+	18.1 ± 4.2	21.8 ± 6.7	13	23	0.083	0.667		>23	74	40	2.0	(1.2-3.3)
CD56-CD122+	19.5 ± 5.7	21.4 ± 8.6	113	154	0.028	0.578	0.030					
CD56-CD69-	67.9 ± 5.3	62.7 ± 9.4	13	23	0.075	0.671	0.093	<64.5	67	51	2.1	(1.3-3.5)
CD56-CD122-	67.7 ± 8.3	64.7 ± 11.5	113	154	0.014	0.586	0.017	<76	44	71	1.9	(1.1-3.5)
CD57-CD44-	74.3 ± 6.9	71.7 ± 7.7	84	117	0.015	0.602	0.014	<3.7	72	43	1.9	(1.1-3.5)
CD3+CD57-HLADR+	4.3 ± 1.5	4.0 ± 1.4	96	118	0.032	0.584	0.034	>40.5	69	45	1.8	(1.0-3.3)
CD3-CD4-CD44+	34.4 ± 11.0	38.1 ± 12.6	88	112	0.023	0.577						
CD3-CD56+CD16-	1.2 ± 0.6	1.4 ± 1.0	121	163	0.023	0.553						
CD3-CD56-CD122-	23.4 ± 11.1	21.2 ± 11.4	113	148	0.004	0.581	0.024					
CD3-CD57-CD44-	48.7 ± 10.1	44.5 ± 10.3	84	114	0.004	0.610	0.008	<41.3	80	41	2.8	(1.4-5.3)

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested		P ¹	area under curve ²	P ³	Cutoff point	Specificity	Sensitivity	Odds ratio	CI ⁴
	Control	Endo. Stage I-IV	Control	Endo. Stage I-IV								
CD14+CD20+CD44-	0.2 ± 0.2	0.1 ± 0.1	65	75	0.036	0.613	0.037	>95	75	43	2.3	(1.1-4.7)
CD14+CD20-CD44+	91.8 ± 4.6	93.5 ± 3.5	65	75	0.016	0.614	0.021	>95	75	43	2.3	(1.1-4.7)
CD14+CD20-CD44-	5.9 ± 4.5	4.6 ± 3.3	65	75	0.037	0.604	0.035	<3	75	43	2.1	(1.1-4.0)
CD14-CD13-HLADR+	8.7 ± 3.5	7.6 ± 2.6	77	82	0.021	0.600	0.029	<7	71	45	1.8	(1.1-3.0)
Ratio CD13+/CD3+	0.25 ± 0.12	0.29 ± 0.14	121	153	0.025	0.583	0.018	>0.30	71	43	1.9	(1.1-3.1)
Ratio CD13+/CD8+	1.04 ± 0.55	1.20 ± 0.69	114	153	0.039	0.574	0.039	>0.14	40	74	1.9	(1.1-3.1)
Ratio CD14+/CD3+	0.19 ± 0.10	0.22 ± 0.12	123	165	0.011	0.575	0.028	>0.14	40	74	1.9	(1.1-3.1)
Ratio CD14+/CD8+	0.78 ± 0.47	0.91 ± 0.55	116	164	0.040	0.574	0.036	>0.14	40	74	1.9	(1.1-3.1)

¹ P value (when ≤ 0.05) obtained in a student "t" test when mean proportion of leukocyte subsets was compared between patients with endometriosis (stage I-IV) and normal controls.

² Discriminative value of each marker established by area under ROC curve.

³ P value (when ≤ 0.05), significance of area under ROC curve.

⁴ Confidence interval for odds ratio.

TABLE 5
Endometrial leukocyte populations used as markers to discriminate between patients with endometriosis stage I-II or stage III-IV and normal subjects

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets		Number of samples tested				Cutoff point	Specificity	Sensitivity	Odds ratio (CI) ⁴				
	Controls	Endo stage I-II	P ¹	Endo stage III-IV	P ²	Control					Endo I-II	Endo III-IV	area under ROC curve	P
CD3-CD44-	50.0 ± 12.9	50.5 ± 11.9	61.8 ± 16.6	0.015	31	43	13	0.766	0.007	>48	52	92	12.9	(1.4-113.8)
CD3+HLADR+	8.3 ± 5.7	6.1 ± 2.6	0.013	7.6 ± 3.3	51	57	23	0.587		<5.5	67	48.3		
CD3+CD45RO+	14.9 ± 8.4	12.0 ± 5.6	0.043	13.2 ± 9.5	50	50	23	0.565		>15	37.3	71		
CD4+CD13+	2.3 ± 1.9	2.2 ± 1.7	0.025	1.4 ± 1.1	36	44	19	0.658		<1.7	50	79	3.8	(1.0-13.8)
CD13+HLADR+	5.7 ± 2.7	5.3 ± 2.3	0.017	3.6 ± 1.3	27	37	12	0.740	0.021	<5	54	92	13	(1.4-117.2)
CD14-HLADR+	14.7 ± 7.5	10.2 ± 3.9	0.049	10.8 ± 5.1	14	24	4	0.679		<13	57	75		
CD56+HLADR+	2.8 ± 1.7	1.5 ± 0.7	0.011	4.6 ± 3.3	16	17	5	0.741	0.018	<1.7	75	71	7.2	(1.5-33.6)
CD56-CD44-	63.5 ± 13.0	63.5 ± 13.4	0.023	51.1 ± 20.4	30	41	12	0.708	0.041	<65	58	83	6.8	(1.2-37.5)
CD3-CD4+CD45RA-	1.5 ± 0.9	1.5 ± 0.9	0.022	1.0 ± 0.7	51	54	25	0.668	0.021	<0.7	84	40	3.4	(1.1-10.7)
CD3+CD8-HLADR+	15.9 ± 8.1	15.8 ± 8.6	0.011	23.6 ± 12.1	48	52	22	0.701	0.009	>14.8	55	73	3.3	(1.1-10.1)
CD3-CD8+HLADR-	9.5 ± 6.5	8.6 ± 6.1	0.036	6.1 ± 5.2	48	52	22	0.691	0.013	<5	78	59	5.0	(1.6-15.4)
CD3+CD56+CD16+	3.0 ± 3.4	2.9 ± 3.2	0.027	6.0 ± 5.7	53	56	22	0.670	0.024	>1.8	49	77	3.5	(1.0-10.3)
CD3+CD56-CD16+	6.6 ± 7.5	7.0 ± 6.2	0.350	11.0 ± 9.1	53	56	22	0.698	0.009	>9	84	50	5.4	(1.7-17.3)
CD3+CD56-CD16-	87.0 ± 8.1	87.2 ± 7.1	0.016	79.5 ± 12.9	53	56	22	0.695	0.010	>83	81	50	4.1	(1.4-12.5)

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets				Number of samples tested				Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴	
	Controls	Endo stage I-II	P ¹	Endo stage III-IV	P ²	Control	Endo I-II	Endo III-IV						area under ROC curve
CD3+CD56-CD44+	7.0 ± 6.1	6.1 ± 4.8		13.2 ± 10.5	0.025	28	38	12	0.721	0.034	<5	42	92	
CD3+CD56-CD44-	87.1 ± 6.1	88.5 ± 6.2		78.0 ± 13.9	0.049	28	38	12	0.668		>81	83	58	7.0 (1.5-33.7)
CD3-CD56-HLADR-	29.7 ± 15.6	42.7 ± 19.6	0.050	29.3 ± 21.3		15	17	6	0.706	0.047	<36	73	59	
CD3+CD56-CD122+	12.1 ± 7.5	11.9 ± 6.3		21.7 ± 14.0	0.005	27	33	16	0.715	0.024	>20	87	50	6.7 (1.4-31.7)
CD3+CD56-CD122-	81.9 ± 8.2	83.0 ± 6.8		68.0 ± 21.7	0.024	27	33	16	0.715	0.024	>76.5	71	63	4.0 (1.1-15.5)
CD4-CD16-	51.5 ± 12.9	59.1 ± 12.7	0.033	49.3 ± 21.4		24	31	11	0.659	0.045	<58	76	47	
CD14+CD13+CD16b-	62.7 ± 20.7	56.2 ± 25.8		38.2 ± 26.4	0.011	19	26	10	0.753	0.035	>61	69	80	8.8 (1.3-57.4)
CD16+	27.4 ± 12.4	30.0 ± 16.2		35.4 ± 19.5		58	62	23	0.603		>39	85	39	3.5 (1.2-10.5)
CD45RA+	28.3 ± 9.1	27.6 ± 9.0		27.9 ± 16.0		56	60	25	0.562		<23.5	75	48	2.8 (1.0-7.5)
CD45RO+	38.4 ± 13.6	38.0 ± 15.5		44.6 ± 18.1		50	54	24	0.620		>52	88	43	4.4 (1.3-14.4)
CD13+	24.8 ± 12.7	25.5 ± 14.7		31.3 ± 20.3		47	57	22			>29	68	34	

¹ P value obtained in a student "t" test when mean % of leukocyte subsets found in patients with endometriosis stage I-II was compared to normal controls.
² P value obtained in a student "t" test when mean % of leukocyte subsets found in patients with endometriosis stage III-IV was compared to normal controls.

TABLE 6
Peripheral blood leukocyte populations used as markers to discriminate between patients with endometriosis stage I-II or stage III-IV and normal subjects

Leukocyte Subsets	Mean proportion (% ± s.d.) of leukocyte subsets										Number of samples tested				Odds ratio	Sensitivity	Specificity	Cutoff point	P	area under ROC curve	(CI) ⁴
	Controls	Endo stage I-II	P ¹	Endo stage III-IV	P ²	Control	Endo I-II	Endo III-IV	area under ROC curve	P	Cutoff point	Specificity	Sensitivity	Odds ratio							
CD3-CD57+CD44-	12.5 ± 6.2	14.9 ± 7.5	0.030	12.2 ± 6.9	0.037	84	81	33	0.586	0.008	>15.6	74	43	2.1	(1.1-4.1)						
CD14-CD13-HLADR-	85.6 ± 5.0	87.6 ± 3.9	0.016	85.9 ± 3.3	0.012	77	54	28	0.637	0.008	>88.3	74	47	2.6	(1.1-5.3)						
CD14-CD20+CD44-	5.7 ± 2.9	4.7 ± 2.6	0.043	5.6 ± 2.7	0.037	65	50	23	0.607	0.049	<4.5	66	53	2.2	(1.0-4.7)						
HLADR+	21.6 ± 6.0	21.7 ± 6.6		23.8 ± 6.7	0.037	120	102	52	0.586		>24.5	73	45	2.2	(1.1-4.5)						
CD3+CD8-	49.9 ± 8.5	49.0 ± 9.2		46.3 ± 8.8	0.012	122	113	53	0.654	0.002	<51.5	48	85	5.2	(2.2-12.1)						
CD3+CD44-	60.2 ± 8.6	59.2 ± 8.8		57.2 ± 8.0	0.047	111	98	45	0.605	0.047	<61.2	51	73	2.8	(1.3-6.1)						
CD3+HLADR-	63.3 ± 8.7	63.0 ± 9.1		59.6 ± 7.9	0.009	121	102	52	0.628	0.010	<63.5	55	73	3.2	(1.5-6.6)						
CD3-HLADR+	18.1 ± 5.8	18.7 ± 6.5		20.8 ± 6.9	0.010	121	102	52	0.607	0.030	>21.5	73	45	2.3	(1.1-4.6)						
CD3+CD16+	9.3 ± 13.3	8.4 ± 12.1		6.0 ± 4.3	0.016	123	113	53	0.528		>3.7	31	81								
CD3-CD57-	28.3 ± 6.9	29.0 ± 7.2		32.3 ± 9.5	0.004	113	101	45	0.644	0.007	>28	53	71	2.7	(1.2-5.7)						
CD4-HLADR+	19.8 ± 6.0	19.8 ± 6.2		22.0 ± 6.6	0.036	112	96	49	0.588		>19.3	48	67								
CD4+CD45RA-	31.4 ± 8.1	29.7 ± 7.2		27.7 ± 8.5	0.006	125	114	54	0.625	0.010	<29.5	60	60	2.3	(1.2-4.5)						
CD4+CD45RO+	19.5 ± 5.6	20.5 ± 6.4		16.2 ± 4.3	0.029	27	31	19	0.645		<19.8	50	90	8.5	(1.6-44.5)						
CD13-CD16+	18.9 ± 8.2	16.8 ± 8.7		14.3 ± 3.1	0.035	21	28	9	0.698		<13	86	56	7.5	(1.2-45.1)						

Leukocyte Subsets	Controls	Mean proportion (% ± s.d.) of leukocyte subsets		P ²	Number of samples tested				P	area under ROC curve	Cutoff point	Specificity	Sensitivity	Odds ratio	(CI) ⁴
		Endo stage I-II	Endo stage III-IV		Control	Endo I-II	Endo III-IV								
CD14-CD69-	81.0 ± 6.1	77.4 ± 9.1	76.0 ± 6.7	0.031	20	32	14	0.729	0.025	<82	50	86	6.0	(1.1-34.0)	
CD20-HLADR-	79.3 ± 5.2	78.5 ± 7.0	77.1 ± 6.6	0.025	106	97	44	0.573		<80.3	41	72			
CD57-CD44+	17.2 ± 5.5	18.3 ± 6.4	19.7 ± 7.7	0.046	84	82	35	0.573		>21.6	80	40	2.7	(1.1-6.7)	
CD3-CD4+CD45RA-	6.9 ± 4.4	6.4 ± 4.0	5.4 ± 3.5	0.043	122	109	50	0.587		<5.5	59	61	2.3	(1.1-4.5)	
CD3-CD8-CD44+	37.2 ± 11.9	37.2 ± 12.7	43.2 ± 12.6	0.019	87	80	32	0.588		>40.5	54	63			
CD3-CD57-CD44+	38.5 ± 12.1	40.3 ± 12.5	44.0 ± 12.7	0.029	84	81	33	0.575		>44	63	58			
CD14+CD20+CD44+	2.1 ± 1.3	2.1 ± 1.5	1.4 ± 0.7	0.001	65	52	23	0.647		<2.1	39	91	6.4	(1.3-30.9)	

¹ P value obtained in a student "t" test when mean proportion found in patients with endometriosis stage I-II was compared to normal controls.

² P value obtained in a student t test when the % leukocyte subsets found in patients with endometriosis stage III-IV was compared to normal controls.

Cutoff points established for each individual marker are presented in Table 3, 4, 5, 6 and threshold value established for a particular marker combination are presented in Table 1. Any other cutoff points or
5 threshold values providing a valuable diagnostic test for endometriosis are meant to be included in the present invention

In accordance with a preferred embodiment of the present invention, there is provided a series of
10 34 different combinations of endometrial leukocyte markers (Tables 1 and 2), 7 combinations of blood leukocyte markers (Tables 1 and 2) and 4 combinations of endometrial and blood leukocyte markers providing a diagnostic test with levels of sensibility and
15 specificity up to 89 and 90%, respectively. The different marker combinations of the present invention may serve several important clinical needs. Hence in the general population, these markers could be used to evaluate the risk factor to develop endometriosis or
20 to identify women with high likelihood of suffering from the disease. Furthermore in patients with endometriosis, these markers could serve to monitor the disease or to give a prognosis.

Study subjects and samples

25 Uterine endometrial tissues were obtained from 146 subjects undergoing laparoscopy or laparotomy. The experimental group was formed of up to 88 subjects with endometriosis stage I-IV confirmed by laparoscopy or laparotomy and the control group consists of up to
30 58 healthy subjects who underwent surgery for tubal ligation (or reanastomosis) and had no clinical evidence, nor family history of endometriosis. Table 7 gives details concerning the age, menstrual cycle and indication of laparoscopy or laparotomy for the
35 subjects included in experimental and control groups.

TABLE 7
Description of the experimental groups used in the
analysis of endometrial leukocyte populations

Experimental groups	Number of subjects	Mean age \pm s.d.	Menstrual cycle		Percentage of patients* Indication of laparoscopy			
			ES ^{1*}	LS ^{2*}	ligation or reanastomosis	Hysterectomy and/or ovariectomy	Diagnostic laparoscopy	Other**
Controls	58	34.2 \pm 5.3	54.5 %	45.5%	100%			
Endometriosis								
Stage I-IV	88	34.4 \pm 6.8	47.0%	53.0%	21.6%	22.7%	52.3%	3.4%
Stage I-II	63	34.4 \pm 7.3	50.9%	49.1%	28.6%	22.2%	47.7%	1.5%
Stage III-IV	25	34.4 \pm 5.4	36.4%	63.6%	4.0%	24.0%	64.0%	8.0%

5 1 Early secretory (days 14-21)

2 Late secretory (days 22-28)

* % patients among control or endometriosis groups

Endometrial biopsies were taken with a Pipet
 10 Curette (Milex) (approximately 0.5g of tissue). All
 samples were harvested in the secretory phase (day 14-
 28) of the menstrual cycle as confirmed by
 histological evaluation. The samples were collected
 into sterile RPMI-1640 medium (Gibco) supplemented
 15 with 2% heat-inactivated fetal calf serum (Bio-Media)
 and 1% penicillin-streptomycin and kept at 4°C until
 cell isolation.

Blood samples were obtained from up to 172
 subjects with endometriosis (stage I-IV) confirmed by
 20 laparoscopy or laparotomy and from up to 132 healthy
 subjects with no evidence of endometriosis at surgery,
 and no family history of endometriosis. Blood samples
 (30 ml) were collected in heparin-tubes (Vacutainer™,
 Becton Dickinson) and kept at 20°C until mononuclear
 25 cell separation. The age, menstrual dating and
 indication for laparoscopy of the subjects included in
 the study are given in Table 8.

TABLE 8
Description of the subjects included in the study of
peripheral blood mononuclear leukocytes

Experimental groups	Number of subjects	Mean age ± s.d.	Menstrual cycle		Percentage of patients*			Other
			Proliferative*	Secretory*	Indication of laparoscopy ligation or reanastomosis	Hysterectomy and/or ovariectomy	Diagnostic laparoscopy	
Control	132	34.30 ± 5.5	43.8 %	56.2%	100%			
Endometriosis								
Stage I-IV	172	36.40*	42.8%	57.2%	22.1%	33.7%	38.9%	5.3%
Stage I-II	116	35.96 ± 6.39	41.0%	59.0%	31.1%	30.2%	37.1%	1.6%
Stage III-IV	56	34.30 ± 5.5	46.2%	53.8%	3.6%	41.1%	42.9%	12.4%

5 * % of patients amongst control or endometriosis groups

Stromal cell preparation from endometrial samples

Endometrial tissue samples were mechanically disrupted with a Pyrex™ glass Broeck tissue grinder (Fisher) to obtain a single cell suspension. Stromal cell fraction was isolated by filtration through a 250 µm stainless steel sieve (Millipore) to retain the glandular fraction and was washed twice with 10 ml phosphate buffered saline (PBS) (Sigma) containing 1% BSA (Boehringer Mannheim), 0.1% sodium azide (Fisher) (thereafter called PBS washing buffer).

Isolation of mononuclear cells from peripheral blood

Blood samples were diluted 1:1 with Hank's Balanced Salt Solution (HBSS) (Gibco), layered on an equal volume of Ficoll-Paque™ (Pharmacia Biotech) and centrifuged at 1500 rpm for 40 minutes at room temperature. Leukocytes were isolated at the interface of Ficoll and HBSS and they were washed in 50 ml of HBSS. Contaminating red blood cells were lysed with 6 ml of ammonium chloride solution (0.15M) (6 minutes at room temperature). The peripheral blood mononuclear cells were then washed twice in 10 ml PBS and resuspended in PBS washing buffer.

Endometrial and peripheral blood leukocyte surface antigen staining

Endometrial stromal cells or peripheral blood mononuclear cells were distributed in 5 ml tubes (1 to 5
1.5 x 10⁶ cells/tube) or in 96 well plates (5 x 10⁵ cells/well), respectively and incubated in the presence of 0.1 µg of human γ-globulin for 5 minutes at room temperature. The cells were then incubated 30 minutes in the dark (at room temperature for
10 endometrial cells and at 4°C for peripheral blood mononuclear cells) with a panel of 4 different mouse monoclonal antibodies (MAbs) in a total volume of 100 µl. The cell samples were stained with mouse anti-human CD45 MAbs conjugated to peridinin chlorophyll
15 protein (PerCP) and with several sets of three different mouse MAbs labeled with distinct fluorochromes (fluorescein isothiocyanate -FITC-, phycoerythrin -PE or with phycoerythrin-texas red -ECD-) directed toward cell surface markers for
20 specific cell populations such as T lymphocytes, B lymphocytes, NK cells, macrophages and/or activation markers (Table 9).

TABLE 9
Description of mouse monoclonal antibodies used for immunophenotyping

Specificity	Clone	Isotype	Supplier	Fluoro-chrome
CD3	HIT3A	mouse IgG2a	Beckman/Coulter	ECD
CD4	SK3	mouse IgG1	Becton Dickinson	PE
CD5	BL1A	mouse IgG2a	Beckman/Coulter	FITC
CD8	SK1	mouse IgG1	Becton Dickinson	PE
CD13	SJ1D1	mouse IgG1	Beckman/Coulter	RPE
CD14	RM052	mouse IgG2a	Beckman/Coulter	PE
CD16	NKP15	mouse IgG1	Becton Dickinson	FITC
CD16B	1D3	mouse igM	Beckman/Coulter	FITC
CD20	H299	mouse IgG2a	Beckman/Coulter	RD1
CD36	SMf	mouse IgM	Sigma	RPE
CD44	L178	mouse IgG1	Becton Dickinson	FITC
CD45	2D1	mouse IgG1	Becton Dickinson	PerCP
CD45-RA	ALB11	mouse IgG1	Beckman/Coulter	FITC
CD45-RO	UCHL1	mouse IgG2a	Beckman/Coulter	FITC
CD56	N901(NKH-1)	mouse IgG1	Beckman/Coulter	PE
CD57	VC1.1	mouse IgM	Sigma	RPE
CD69	L78	mouse IgG1	Becton Dickinson	FITC
CD122	2RB	mouse IgG1	Beckman/Coulter	FITC
HLA-DR	L243	mouse IgG2a	Becton Dickinson	FITC

5 Table 10 below lists the distribution of the antigens listed in Table 9.

TABLE 10
Main distribution of antigens

Antigen	Main Cell Distribution
CD3	Expressed on all mature T cells associated with TCR complex (α/β , γ/δ)
CD4	Expressed on T helper lymphocytes. It can be also expressed on cells of the monocyte/macrophage lineage
CD5	Found on all mature T lymphocytes and a subset of B lymphocytes
CD8	Found on a subset of T lymphocytes called suppressor/cytotoxic T cells.

Antigen	Main Cell Distribution
CD13	Detected on most cells of myeloid origin polymorphonuclear cells or cells of the monocyte/macrophage lineage. Member of metalloproteinase family
CD14	Expressed strongly on the surface of monocytes Found on most tissue macrophages Weakly expressed on the surface of granulocytes and B lymphocytes Receptor for lipopolysaccharide (LPS) and LPS binding protein
CD16	Expressed mainly on NK cells, monocytes macrophages and polymorphonuclear leukocytes Low affinity receptor for IgG
CD16b	Found on granulocytes including polymorphonuclear cells (PMN)
CD20	Present on all B lymphocytes
CD36	Expressed on platelets, monocytes or macrophages, microvascular endothelial cells, mammary endothelial cells, during stages of erythroid cell development
CD44	Widely expressed on the surface of most cell types. Including most leukocytes and epithelial cells. Family of core/link peptidoglycan
CD45	Present on the surface of all leukocytes
CD45RA	Isoforms of CD45 Found on naive/resting T cells Also expressed on B lymphocytes and monocytes
CD45RO	Isoforms of CD45 expressed on memory/activated T cells also expressed on monocytes
CD56	Marker for NK cells Can also be found on a population of T lymphocytes
CD57	Found on a subset of cells with natural killer activity
CD69	Expressed on activated leukocytes including T cells, B cells, NK cells, neutrophils, eosinophils and cells of the monocyte/macrophage lineage. Activation marker detected early after cell activation
CD122	Expressed on NK cells B, T lymphocytes or monocytes/macrophages Component of the IL-15 receptor
HLADR+	HLA class II molecule Found on antigen presenting cells or on other cells upon activation such as T cells.

Blood cells were washed twice with 0.15 ml of PBS washing buffer. Endometrial cell samples were

incubated with a red blood cell lysing solution, (FACS™ Lysing Solution, Becton Dickinson) for 10 minutes at room temperature in the dark and washed with 3 ml of PBS washing buffer. Endometrial and blood
5 cells were fixed in 1% paraformaldehyde (diluted in PBS) at a concentration of 1×10^6 cells/ml and kept at 4°C in the dark until the immunofluorescence reactivity was determined by flow cytometry.

Flow cytometry analysis

10 The immunofluorescence reactivity was carried out on a Coulter EPICS XL™ flow cytometer (Coulter Corporation, Hialeah, FL) equipped with an argon laser operating at 488 nm, 15 mW and detectors at 525, 575, 610, and 675 nm. Calibration of the flow cytometer
15 parameters for forward scatter, side scatter and fluorescence were the same for all the samples. Cells expressing CD45 pan leukocyte antigen were gated using the Coulter system II software. The percentage of cells bearing markers for T, B lymphocytes,
20 macrophages or NK cells and/or activation markers was evaluated within the CD45 positive populations only. A minimum of 6000 CD45+ cells were analyzed for each sample.

Use of leukocyte markers in a diagnostic test for endometriosis

25 A cutoff point was established for the proportion of the endometrial or blood leukocyte subpopulations identified as diagnostic markers. The value obtained for each marker is compared to the
30 cutoff point (Fig. 1). A positive result was given when the proportion of a particular leukocyte subset fulfills the condition established by the cutoff point (for example < 40% for CD3+ cells). When these markers are used in combination, a positive result for each
35 marker gives a score of 1, whereas a negative result

gives a score of 0. A diagnosis of endometriosis is given, when the final diagnostic score obtained from adding the results of all the markers of a particular combination is higher than a predetermined threshold value. The levels of sensibility and/or specificity measured for the marker combination represents the number of positive test results obtained among the patients already confirmed with endometriosis and the number of negative test results among the subjects within the control group, respectively.

Endometrial and blood leukocyte markers can be used in combinations in logistic regression model:

$$P(r) = \frac{e^{c + B_1(\text{marker1}) + B_2(\text{marker2}) + \dots + B_n(\text{marker n})}}{1 + e^{c + B_1(\text{marker1}) + B_2(\text{marker2}) + \dots + B_n(\text{marker n})}}$$

Where:

$P(r)$ = probability of having endometriosis

c = constant established for a particular combination

B = coefficient of regression

n = total number of markers in the combination

The probability of having endometriosis ($P(r)$) is then compared to a threshold value that provides the best discriminative value. A positive diagnosis of endometriosis is given when the $P(r)$ value exceeds the threshold value established for a particular combination of markers. Alternatively, a negative diagnosis of endometriosis is given when the $P(r)$ value is lower than the threshold value.

Results

Endometrial and blood leukocyte subsets defined as good potential markers for the diagnosis of endometriosis are presented in Tables 3 and 4 respectively. Selection of these markers was done on

the basis of 1) a significant difference in the mean proportion of leukocyte subsets between patients with endometriosis (stage I-IV) and control groups; 2) several endometrial and blood leukocyte markers were also selected according to the area under the ROC curve (>0.500), an indication of the discriminative value of the markers. The ROC curve allowed the determination of one or more cutoffs (proportion % of leukocyte subpopulations) that best discriminate between patients with endometriosis (stage I-IV) and normal controls. In an attempt to use these differences for identifying patients with endometriosis, a positive test result was given when the proportion measured for a particular leukocyte subset fulfills the condition established by the cutoff point (for example $< 40\%$ for CD3+ cells). The levels of specificity and sensibility were calculated for each marker used alone to diagnose endometriosis and are presented in Table 3 (for endometrial leukocyte markers) and Table 4 (for blood leukocyte markers). Moreover, 3) a significant odds ratio calculated with a particular cutoff point gave an additional indication that the leukocyte markers selected in Tables 3 and 4 are associated with an increased risk to develop endometriosis and can, thus, be used for identifying women with high likelihood of suffering from endometriosis.

The mean proportion of some endometrial (Table 5) and blood (Table 6) leukocyte subsets was found to be significantly modulated only in patients with stage I-II endometriosis or with stage III-IV endometriosis when compared to normal controls. These markers remain good candidates for a diagnostic test for endometriosis, but their use may be limited to a specific stage of the disease.

Several of endometrial and blood leukocyte markers were found to be even more reliable as diagnostic markers when they are analyzed in combination with other markers. Table 1 gives a series of 33 combinations in which endometrial or blood leukocyte markers are used in a diagnostic test for endometriosis. For each marker, a positive test result (as described above; see also Fig. 1) gives a score of 1, whereas a negative test result gives a score of 0. The final diagnostic value obtained from adding the scores of all the markers of a particular combination is then compared to a threshold value, which is indicated in Table 1. A diagnosis of endometriosis is given, when the diagnostic value exceeds the threshold value established for each set of combination markers. The use of leukocyte marker subsets in combination in this new method clearly improves the levels of sensibility and/or specificity for diagnosing endometriosis. Table 1 also provides 4 examples showing that blood leukocyte markers, when used in combination with endometrial markers, can also increase the predictive value of the diagnostic test.

The present invention also demonstrate that logistic regression models can also be used to combine endometrial as well as blood leukocyte markers for the development of a predictive model of endometriosis (Table 2). In some cases, these models need to be adjusted with risk factors associated with endometriosis such as the length of the menstrual cycle, the duration of menstruation, pain (during intercourse, menstruation or in other circumstances) and age. In some instances, these factors were shown to increase the predictive value of the model.

The present invention identifies several examples of marker combinations, which give rise to diagnostic methods yielding improved levels of sensibility and specificity. Indeed, the different
5 marker combinations of the present invention may serve different clinical applications including screening, diagnosis, monitoring and prognosis of endometriosis.

While the invention has been described in connection with specific embodiments thereof, it will be
10 understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure
15 as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth, and as follows in the scope of the appended claims. For example, other blood or endometrial
20 markers, alone or in combination, might also be suitable to practice the method of the present invention, and are thus intended to be included in the present invention.

WHAT IS CLAIMED IS:

1. An endometrial or blood leukocyte marker for endometriosis characterized in that said marker is a surface antigen from endometrial or blood leukocytes.

2. The marker of claim 1, wherein the surface antigen from blood leukocytes is selected from the group consisting of CD3+, CD8+, CD13+, CD14+, CD20+, CD36+, CD44+, CD57+, CD69+, CD122+, CD3+CD5+, CD3+CD45RA-, CD3+CD56-, CD3+CD57-, CD3+CD69-, CD3+CD122-, CD3+HLADR+, CD3-CD5+, CD3-CD16+, CD3-CD44+, CD3-CD57+, CD3-CD69+, CD3-CD45RO+, CD3-CD4-, CD3-CD8-, CD3-CD45RA-, CD3-CD56-, CD4-CD13+, CD4-CD36+, CD4-CD69+, CD4-CD45RO+, CD4-CD45RA-, CD8-CD44+, CD8-CD44-, CD13+CD44+, CD13+HLADR+, CD13+CD16-, CD13-HLADR+, CD13-CD44-, CD14+HLADR+, CD14+CD44+, CD14+CD45RO+, CD14+CD16-, CD14+CD122-, CD14-HLADR+, CD14-CD44-, CD20+HLADR+, CD20+CD44-, CD20-CD44+, CD20-CD69+, CD20-HLADR+, CD20-CD44-, CD20-CD69-, CD36-HLADR+, CD56-CD69+, CD56-CD122+, CD56-CD69-, CD56-CD122-, CD57-CD44-, CD3+CD57-HLADR+, CD3-CD4-CD44+, CD3-CD56+CD16-, CD3-CD56-CD122-, CD3-CD57-CD44-, CD14+CD20+CD44-, CD14+CD20-CD44+, CD14+CD20-CD44-, CD14-CD13-HLADR+, Ratio CD13+/CD3+, Ratio CD13+/CD8+, Ratio CD14+/CD3+, Ratio CD14+/CD8+, CD3-CD57+CD44-, CD14-CD13-HLADR-, CD14-CD20+CD44-, HLADR+, CD3+CD8-, CD3+CD44-, CD3+HLADR-, CD3-HLADR+, CD3+CD16+, CD3-CD57-, CD4-HLADR+, CD4+CD45RA-, CD4+CD45RO+, CD13-CD16+, CD14-CD69-, CD20-HLADR-, CD57-CD44+, CD3-CD4+CD45RA-, CD3-CD8-CD44+, CD3-CD57-CD44+, and CD14+CD20+CD44+.

3. The marker of claim 2, wherein the surface antigen is chosen from CD3-CD57+CD44-, CD14-CD13-

HLADR-, CD14-CD20+CD44-, HLADR+, CD3+CD8-, CD3+CD44-, CD3+HLADR-, CD3-HLADR+, CD3+CD16+, CD3-CD57-, CD4-HLADR+, CD4+CD45RA-, CD4+CD45RO+, CD13-CD16+, CD14-CD69-, CD20-HLADR-, CD57-CD44+, CD3-CD4+CD45RA-, CD3-CD8-CD44+, CD3-CD57-CD44+, and CD14+CD20+CD44+.

4. The marker of claim 1, wherein the surface antigen from endometrial leukocytes is selected from the group consisting of CD3+, CD4+, CD5+, CD8+, CD3+CD4+, CD3+CD4-, CD3-CD4-, CD3+CD8+, CD3+CD8-, CD3-CD8-, CD3+CD69+, CD3+CD122-, CD3+HLADR-, CD3-HLADR-, CD3+CD45RA+, CD3+CD45RA-, CD3-CD45RA-, CD3+CD45RO-, CD3+CD16-, CD3+CD56-, CD3+CD5+, CD3-CD5-, CD4+CD69-, CD4+CD45RA-, CD8+CD69-, CD8+HLADR-, CD8-HLADR-, CD13-CD122+, CD13-CD122-, CD20-CD5+, CD20-CD5-, CD56-CD16+, CD56-CD16-, ratio CD3+/CD45RO+, CD14+CD13-, CD3+CD20-, CD3-CD20-, CD3-CD4-CD45RA+, CD3-CD4-CD45RA-, CD3+CD8+CD69-, CD3+CD8+HLADR-, CD3+CD8-HLADR-, CD3-CD8-HLADR-, CD14+CD13-CD16b-, CD4+CD14-, CD4-CD14-, CD4+HLADR-, CD13-CD69+, CD13+CD45RO-, CD56-CD122-, CD3+CD69-, CD4+CD45RO-, CD56+CD122+, CD3-CD56+CD16+, CD3-CD56+CD122+, CD14+CD13+, CD3+CD20-CD5+, CD4-CD13+CD16+, CD69+, ratio CD13+/CD3+, CD3-CD20-CD5-, CD3+CD20-CD5-, CD4+CD13-, CD3+CD44-, CD56+, CD13-CD45RO+, CD56+CD69-, CD13-CD16+, CD56+CD122-, CD3+CD4-CD69+, CD4-CD13-CD16+, CD4-CD13-CD16-, CD14+CD13+CD16b+, CD4-CD69-, CD3-CD45RO+, CD4-CD69+, CD3-CD4-CD45RO+, CD3+HLADR+, CD3+CD45RO+, CD4+CD13+, CD13+HLADR+, CD14-HLADR+, CD56+HLADR+, CD56-CD44-, CD3-CD4+CD45RA-, CD3+CD8-HLADR+, CD3-CD8+HLADR-, CD3+CD56+CD16+, CD3+CD56-CD16+, CD3+CD56-CD16-, CD3+CD56-CD44+, CD3+CD56-CD44-, CD3-CD56-HLADR-, CD3+CD56-CD122+, CD3+CD56-CD122-, CD4-CD16-, CD14+CD13+CD16b-, CD3-CD44-, CD16+, CD45RA+, CD45RO+, and CD13+.

5. The marker of claim 4, wherein the surface antigen is chosen from CD3+HLADR+, CD3+CD45RO+, CD4+CD13+, CD13+HLADR+, CD14-HLADR+, CD56+HLADR+, CD56-CD44-, CD3-CD4+CD45RA-, CD3+CD8-HLADR+, CD3-CD8+HLADR-, CD3+CD56+CD16+, CD3+CD56-CD16+, CD3+CD56-CD16-, CD3+CD56-CD44+, CD3+CD56-CD44-, CD3-CD56-HLADR-, CD3+CD56-CD122+, CD3+CD56-CD122-, CD4-CD16-, CD14+CD13+CD16b-, CD3-CD44-, CD16+, CD45RA+, CD45RO+, and CD13+.

6. A combination of markers for endometriosis comprising at least two different surface antigens from blood or endometrial leukocytes.

7. The combination of claim 6, wherein the surface antigens are selected from the group consisting of CD3+, CD4+, CD5+, CD8+, CD13+, CD14+, CD20+, CD36+, CD44+, CD56+, CD57+, CD69+, CD122+, HLADR+, CD16+, CD45RA+, CD45RO+, CD56-CD122+, CD3-CD8+HLADR-, CD3+CD4-CD69+, CD3+CD4+, CD3+CD4-, CD3-CD4-, CD3+CD5+, CD3-CD5+, CD3-CD5-, CD3+CD8+, CD3+CD8-, CD3-CD8-, CD3+CD16+, CD3-CD16+, CD3+CD16-, CD3+CD20-, CD3-CD20-, CD3+CD44-, CD3-CD44+, CD3-CD44-, CD3+CD45RA-, CD3-CD45RA-, CD3+CD45RA+, CD3+CD45RO+, CD3-CD45RO+, CD3+CD45RO-, CD3+CD56-, CD3-CD56-, CD3+CD57-, CD3-CD57+, CD3-CD57-, CD3+CD69-, CD3+CD69+, CD3-CD69+, CD3+CD122-, CD3+HLADR+, CD3+HLADR-, CD3-HLADR+, CD3-HLADR-, CD4+CD13+, CD4+CD13-, CD4-CD13+, CD4+CD14-, CD4-CD14-, CD4-CD16-, CD4-CD36+, CD4+CD45RA-, CD4-CD45RA-, CD4+CD45RO+, CD4+CD45RO-, CD4-CD45RO+, CD4+CD69-, CD4-CD69+, CD4-CD69-, CD4+HLADR-, CD4-HLADR+, CD8-CD44+, CD8-CD44-, CD8+CD69-, CD8+HLADR-, CD8-HLADR-, CD13+CD16-, CD13-CD16+, CD13+CD44+, CD13-CD44-, CD13+CD45RO-, CD13-

CD45RO+, CD13-CD69+, CD13-CD122+, CD13-CD122-,
 CD13+HLADR+, CD13-HLADR+, CD14+CD13+, CD14+CD13-,
 CD14+CD16-, CD14+CD44+, CD14-CD44-, CD14+CD45RO+,
 CD14-CD69-, CD14+CD122-, CD14+HLADR+, CD14-HLADR+,
 CD20-CD5+, CD20-CD5-, CD20+CD44-, CD20-CD44+, CD20-
 CD44-, CD20-CD69+, CD20-CD69-, CD20+HLADR+, CD20-
 HLADR+, CD20-HLADR-, CD36-HLADR+, CD56-CD16+, CD56-
 CD16-, CD56-CD44-, CD56+CD69-, CD56-CD69+, CD56-CD69-,
 CD56+CD122+, CD56+CD122-, CD56-CD122-, CD56+HLADR+,
 CD57-CD44+, CD57-CD44-, CD3-CD4-CD44+, CD3-CD4+CD45RA-
 , CD3-CD4-CD45RA+, CD3-CD4-CD45RA-, CD3-CD4-CD45RO+,
 CD3-CD8-CD44+, CD3+CD8+CD69-, CD3+CD8+HLADR-, CD3+CD8-
 HLADR+, CD3+CD8-HLADR-, CD3-CD8-HLADR-, CD3+CD20-CD5+,
 CD3+CD20-CD5-, CD3-CD20-CD5-, CD3+CD56+CD16+,
 CD3+CD56-CD16+, CD3+CD56-CD16-, CD3-CD56+CD16+, CD3-
 CD56+CD16-, CD3+CD56-CD44+, CD3+CD56-CD44-, CD3+CD56-
 CD122+, CD3+CD56-CD122-, CD3-CD56+CD122+, CD3-CD56-
 CD122-, CD3-CD56-HLADR-, CD3-CD57+CD44-, CD3-CD57-
 CD44+, CD3-CD57-CD44-, CD3+CD57-HLADR+, CD4-
 CD13+CD16+, CD4-CD13-CD16+, CD4-CD13-CD16-,
 CD14+CD13+CD16b+, CD14+CD13+CD16b-, CD14+CD13-CD16b-,
 CD14-CD13-HLADR+, CD14-CD13-HLADR-, CD14+CD20+CD44+,
 CD14+CD20+CD44-, CD14+CD20-CD44+, CD14+CD20-CD44-,
 CD14-CD20+CD44-, ratio CD3+/CD45RO+, Ratio CD13+/CD3+,
 Ratio CD13+/CD8+, Ratio CD14+/CD3+, and Ratio
 CD14+/CD8+.

8. A combination of markers for endometriosis selected from the group consisting of:

- a) CD4+, CD8+CD69-, and CD13+CD45RO-;
- b) CD4+, CD8+CD69-, CD56+CD122-, CD3+CD45RA-,
and CD13+CD45RO-;
- c) CD4+, CD8+CD69-, CD13-CD122+, and
CD13+CD45RO-;
- d) CD4+, CD8+CD69-, and CD14+CD13-CD16b-;

- e) CD3+CD16-, CD13+CD45RO-, CD3+, CD8+, and CD3+CD69+;
- f) CD3+, CD3+CD8+, CD13+CD45RO-, and CD3-CD20-;
- g) CD3+CD8+, CD13+CD45RO-, CD3+CD5+, CD3+CD122-, CD3-CD20-, and CD3+CD45RO-;
- h) CD3+CD8+, CD13+CD45RO-, CD3+CD5+, and CD3+CD122-;
- i) CD3+CD20-CD5-, CD4+CD13-, and CD56-CD122-;
- j) CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, and CD3+CD45RO-;
- k) CD3+, CD8+CD69-, CD3-CD4-CD45RO+, Ratio CD13+/CD3+, and CD13+CD45RO-;
- l) CD3+CD8+, CD13+CD45RO-, CD3-CD5-, and CD20-CD5+;
- m) CD8+, CD5+, CD3-CD20-, and CD3-HLADR-;
- n) CD3+CD8+, CD13+CD45RO-, and CD5+;
- o) CD4+, CD13-CD122+, CD8+CD69-, and CD3+CD45RA-;
- p) CD4+, CD8+CD69-, CD13-CD122+, CD3+CD45RA-, and CD13+CD45RO-;
- q) CD4+, CD8+CD69-, CD56-CD122-, CD3+CD45RA-, and CD13+CD45RO-;
- r) CD3+, CD4+, CD3+CD8+, CD13+CD45RO-, and CD3+CD5+;
- s) CD3+, CD4+, CD3+CD8+, and CD13+CD45RO-;
- t) CD3+, CD3+CD8+, and CD13+CD45RO-;
- u) CD3+, CD5+, CD3+CD5+, CD69+, and CD4-CD69+;
- v) CD3+, CD3+CD8+, and CD13+CD45RO-;
- w) CD3+, CD3+CD8+, and CD13+CD45RO-;
- x) CD3+, CD3+CD8+, CD13+CD45RO-, CD3-CD20-, and CD56-CD16-;
- y) CD3+CD16-, CD3-CD4-CD45RO+, CD3+, CD8+, CD3+CD69+, and CD13+CD45RO-;
- z) CD3+, CD3-CD45RO+, Ratio CD13+/CD3+, CD3+CD8+, and CD8+CD69-;

- aa) CD3+CD20-CD5-, CD4+CD13-, CD56-CD122-, and CD4+CD45RO-;
- bb) CD3+, CD3+CD8+, CD13+CD45RO-, and CD3-CD20-;
- cc) CD13+CD45RO-, CD4+CD45RA-, CD3+CD122-, and CD8+CD69-;
- dd) CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, CD3+CD45RO-, and CD3-CD5-;
- ee) CD3-CD5+, CD3-CD45RA-, CD3-CD44+, CD13+, and CD3-CD57-CD44-;
- ff) CD3-CD45RA-, CD20-CD44+, CD20-HLADR+, CD3-CD4-CD44+, and CD36-HLADR+;
- gg) CD3-CD45RA-, CD3-CD45RO+, and CD20-HLADR+;
- hh) CD57+, CD14+, CD3-CD69+, CD3+, CD4+, and CD3+CD8+HLADR-;
- ii) CD3-CD69+, CD3+, CD4+, and CD3+CD8+HLADR-;
- jj) CD4-CD36+, CD3-CD69+, CD8+, CD13+, CD3+, CD16+, CD69+, and CD5+;
- kk) CD3-CD45RA-, CD3-CD45RO+, CD20-HLADR+, CD14+CD44+, CD8+, CD5+, CD3-CD20-, and CD3-HLADR-;
- ll) CD3+, CD3-CD5-, CD13+CD45RO-, and CD3-CD20-;
- mm) CD3+, CD3-CD5-, and CD13+CD45RO-;
- nn) CD3+CD8+, CD13+CD45RO-, and CD3+CD5+;
- oo) CD3+, Length of menstruation, CD13+CD20-, and Pelvic pain;
- pp) CD14+CD44+, CD57+, CD3-CD45RA-, and CD14+;
- qq) CD14+, CD57+, CD3+CD69+, CD3+HLADR+, and CD3-CD45RA-;
- rr) CD14+, CD57+, CD3+HLADR-, CD3-CD45RA-, Pelvic pain, and Length of menstruation; and
- ss) CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-CD45RA-, Pelvic pain, and Age.

9. A diagnostic method for the detection of endometriosis in a patient sample, said method

comprising the step of detecting at least one marker of any one of claims 1, 2, 3, 4, and 5, whereby detection of said specific marker is indicative of endometriosis.

10. A diagnostic method for the detection of endometriosis in a patient sample, said method comprising the step of detecting at least two different surface antigens from blood or endometrial leukocytes, whereby detection of said at least two different surface antigens is indicative of endometriosis.

11. A diagnostic method for the detection of endometriosis in a patient sample, said method comprising the step of detecting a combination for endometriosis as defined in any one of claims 6, 7, and 8, whereby detection of said combination is indicative of endometriosis.

12. A diagnostic kit for the detection of endometriosis, said kit comprising an antibody specific for a marker as defined in any one of claims 1, 2, 3, 4, and 5.

13. A diagnostic kit for the detection of endometriosis, said kit comprising at least two different antibodies, each specific for different surface antigens as defined in the combination of claim 6, 7 or 8.

14. The diagnostic kit of claim 13, wherein said specific marker combination is selected from the group consisting of:

- a) CD4+, CD8+CD69-, and CD13+CD45RO-;

- b) CD4+, CD8+CD69-, CD56+CD122-, CD3+CD45RA-, and CD13+CD45RO-;
- c) CD4+, CD8+CD69-, CD13-CD122+, and CD13+CD45RO-;
- d) CD4+, CD8+CD69-, and CD14+CD13-CD16b-;
- e) CD3+CD16-, CD13+CD45RO-, CD3+, CD8+, and CD3+CD69+;
- f) CD3+, CD3+CD8+, CD13+CD45RO-, and CD3-CD20-;
- g) CD3+CD8+, CD13+CD45RO-, CD3+CD5+, CD3+CD122-, CD3-CD20-, and CD3+CD45RO-;
- h) CD3+CD8+, CD13+CD45RO-, CD3+CD5+, and CD3+CD122-;
- i) CD3+CD20-CD5-, CD4+CD13-, and CD56-CD122-;
- j) CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, and CD3+CD45RO-;
- k) CD3+, CD8+CD69-, CD3-CD4-CD45RO+, Ratio CD13+/CD3+, and CD13+CD45RO-;
- l) CD3+CD8+, CD13+CD45RO-, CD3-CD5-, and CD20-CD5+;
- m) CD8+, CD5+, CD3-CD20-, and CD3-HLADR-;
- n) CD3+CD8+, CD13+CD45RO-, and CD5+;
- o) CD4+, CD13-CD122+, CD8+CD69-, and CD3+CD45RA-;
- p) CD4+, CD8+CD69-, CD13-CD122+, CD3+CD45RA-, and CD13+CD45RO-;
- q) CD4+, CD8+CD69-, CD56-CD122-, CD3+CD45RA-, and CD13+CD45RO-;
- r) CD3+, CD4+, CD3+CD8+, CD13+CD45RO-, and CD3+CD5+;
- s) CD3+, CD4+, CD3+CD8+, and CD13+CD45RO-;
- t) CD3+, CD3+CD8+, and CD13+CD45RO-;
- u) CD3+, CD5+, CD3+CD5+, CD69+, and CD4-CD69+;
- v) CD3+, CD3+CD8+, and CD13+CD45RO-;
- w) CD3+, CD3+CD8+, and CD13+CD45RO-;

- x) CD3+, CD3+CD8+, CD13+CD45RO-, CD3-CD20-, and CD56-CD16-;
- y) CD3+CD16-, CD3-CD4-CD45RO+, CD3+, CD8+, CD3+CD69+, and CD13+CD45RO-;
- z) CD3+, CD3-CD45RO+, Ratio CD13+/CD3+, CD3+CD8+, and CD8+CD69-;
- aa) CD3+CD20-CD5-, CD4+CD13-, CD56-CD122-, and CD4+CD45RO-;
- bb) CD3+, CD3+CD8+, CD13+CD45RO-, and CD3-CD20-;
- cc) CD13+CD45RO-, CD4+CD45RA-, CD3+CD122-, and CD8+CD69-;
- dd) CD3+CD8+, CD13+CD45RO-, CD4+CD45RA-, CD3+CD45RO-, and CD3-CD5-;
- ee) CD3-CD5+, CD3-CD45RA-, CD3-CD44+, CD13+, and CD3-CD57-CD44-;
- ff) CD3-CD45RA-, CD20-CD44+, CD20-HLADR+, CD3-CD4-CD44+, and CD36-HLADR+;
- gg) CD3-CD45RA-, CD3-CD45RO+, and CD20-HLADR+;
- hh) CD57+, CD14+, CD3-CD69+, CD3+, CD4+, and CD3+CD8+HLADR-;
- ii) CD3-CD69+, CD3+, CD4+, and CD3+CD8+HLADR-;
- jj) CD4-CD36+, CD3-CD69+, CD8+, CD13+, CD3+, CD16+, CD69+, and CD5+;
- kk) CD3-CD45RA-, CD3-CD45RO+, CD20-HLADR+, CD14+CD44+, CD8+, CD5+, CD3-CD20-, and CD3-HLADR-;
- ll) CD3+, CD3-CD5-, CD13+CD45RO-, and CD3-CD20-;
- mm) CD3+, CD3-CD5-, and CD13+CD45RO-;
- nn) CD3+CD8+, CD13+CD45RO-, and CD3+CD5+;
- oo) CD3+, Length of menstruation, CD13+CD20-, and Pelvic pain;
- pp) CD14+CD44+, CD57+, CD3-CD45RA-, and CD14+;
- qq) CD14+, CD57+, CD3+CD69+, CD3+HLADR+, and CD3-CD45RA-;

- rr) CD14+, CD57+, CD3+HLADR-, CD3-CD45RA-,
Pelvic pain, and Length of menstruation; and
- ss) CD14+, CD57+, CD3+CD69+, CD3+HLADR-, CD3-
CD45RA-, Pelvic pain, and Age.


15. Use of a marker as defined in any one of claims 1, 2, 3, 4, and 5 for identifying a woman suffering or likely to suffer from endometriosis.

16. Use of a combination as defined in any one of claims 6, 7, and 8 for identifying a woman suffering or likely to suffer from endometriosis.

17. Use of a diagnostic kit as defined in any one of claims 12, 13, and 14 for identifying a woman suffering or likely to suffer from endometriosis.

PREDICTIVE ALGORITHM FOR THE DIAGNOSIS OF ENDOMETRIOSIS

Example



¹ Proportion of cells expressing a specific marker, or a given subset defined by markers within the leucocyte population (CD45+) in the peripheral blood or the stromal fraction of the endometrium.

² A positive test result is given when the proportion of a leucocyte subset fulfills the condition established by the cutoff point.

Fig. 1

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 00/00060

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 G01N33/574

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CRUSE, J.M. AND LEWIS, R.E.: "Illustrated dictionary of immunology", CRC PRESS, 1995 XP002136669 page 53 -page 62	1
X	OOSTERLYNCK D J ET AL: "FLOW CYTOMETRY ANALYSIS OF LYMPHOCYTE SUBPOPULATIONS IN PERIONEAL FLUID OF WOMEN WITH ENDOMETRIOSIS" AMERICAN JOURNAL OF REPRODUCTIVE IMMUNOLOGY, DK, MUNKSGAARD INTERNATIONAL PUBLISHERS, COPENHAGEN, vol. 31, no. 1, 1 January 1994 (1994-01-01), pages 25-31, XP000568653 ISSN: 1046-7408 cited in the application the whole document	1,9-17

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

28 April 2000

Date of mailing of the international search report

13/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hoekstra, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 00/00060

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	OOSTERLYNCK D J ET AL: "IMMUNOHISTOCHEMICAL CHARACTERIZATION OF LEUCOCYTE SUBPOPULATIONS IN ENDOMETRIOTIC LESIONS" ARCHIVES OF GYNECOLOGY AND OBSTETRICS, DE, SPRINGER VERLAG, BERLIN, vol. 253, no. 4, 1 January 1993 (1993-01-01), pages 197-206, XP000568666 ISSN: 0932-0067 the whole document	1,9-17
X	EP 0 387 027 A (ADEZA BIOMEDICAL CORP) 12 September 1990 (1990-09-12) claim 1	1,9,12, 15
X	WO 96 20404 A (INST MEDECINE DE LA REPRODUCTI ; MIRON PIERRE (CA); LACHAPELLE MARI) 4 July 1996 (1996-07-04) claim 1	1,9,12, 15
A	WO 98 10291 A (LARSEN PETER MOSE ; BYRJALSEN INGER (DK); CENTER FOR CLINICAL & BAS) 12 March 1998 (1998-03-12) the whole document	1,9-17

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA 00/00060

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.: 2-8
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see further information sheet

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 2-8

The subject-matter of claims 2-8 relate to a physical entity (markers or combination of markers defined in claim 1 as surface antigens) which is contradicted by reference to absence of markers ("-") and even "pelvic pain" and "age" which can clearly not be considered to characterise a physical entity or a surface antigen. It appears that the only support in the description for matter for which protection can be sought are methods to identify the presence or absence of markers (Articles 5 and 6 PCT).

The search has been limited to surface antigens per se (claim 1) and to the methods and kits of claims 9-17.

The ISA thus considers that the claims and description fail to comply with the prescribed requirements (Article 5 and 6 PCT) to such an extent that a meaningful search for claims 2-8 cannot be carried out.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/CA 00/00060

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0387027 A	12-09-1990	AU 5068090 A CA 2011704 A JP 3202769 A NZ 232801 A	13-09-1990 07-09-1990 04-09-1991 29-01-1992
WO 9620404 A	04-07-1996	US 5618680 A AU 4295696 A CA 2206420 A	08-04-1997 19-07-1996 04-07-1996
WO 9810291 A	12-03-1998	AU 4127597 A EP 0931262 A	26-03-1998 28-07-1999

专利名称(译)	用于诊断子宫内膜异位症的方法和诊断试剂盒		
公开(公告)号	EP1147421A1	公开(公告)日	2001-10-24
申请号	EP2000901009	申请日	2000-01-24
[标]申请(专利权)人(译)	procrea能够		
申请(专利权)人(译)	PROCREA生物科学公司.		
当前申请(专利权)人(译)	西门子医疗系统集团诊断		
[标]发明人	GAGNE DANIELE GOSELIN DIANE HUGO PATRICE MIRON PIERRE		
发明人	GAGNE, DANIELE GOSELIN, DIANE HUGO, PATRICE MIRON, PIERRE		
IPC分类号	G01N33/53 G01N33/574 G01N33/68		
CPC分类号	G01N33/57442 G01N2333/705 G01N2800/364		
优先权	60/117031 1999-01-25 US		
其他公开文献	EP1147421B1		
外部链接	Espacenet		

摘要(译)

本发明涉及使用血液和子宫内膜白细胞标记物或其组合诊断子宫内膜异位症的方法和试剂盒。标记物是来自子宫内膜或血液白细胞的表面抗原。