



(11)

EP 1 892 297 A2

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
27.02.2008 Bulletin 2008/09

(21) Application number: **07109568.1**

(22) Date of filing: **06.06.2000**

(51) Int Cl.:
C12N 15/12 ^(2006.01) **C07K 14/435** ^(2006.01)
A61K 38/17 ^(2006.01) **G01N 33/50** ^(2006.01)
C12Q 1/68 ^(2006.01) **C07K 16/18** ^(2006.01)
C12N 5/20 ^(2006.01)

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

(30) Priority: **09.06.1999 GB 9913425**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
00934824.4 / 1 187 916

(71) Applicant: **Henogen S.A.**
6041 Gosselies (BE)

(72) Inventors:
• **Godfroid, Edmond**
1020 Bruxelles (BE)

• **Bollen, Alex**
1701 Itterbeek (BE)
• **Leboulle, Gérard**
1190 Bruxelles (BE)

(74) Representative: **pronovem**
Office Van Malderen
Avenue Josse Goffin 158
1082 Bruxelles (BE)

Remarks:

This application was filed on 04 - 06 - 2007 as a divisional application to the application mentioned under INID code 62.

(54) **Identification and molecular characterisation of proteins, expressed in the tick salivary glands**

(57) The invention relates to the molecular characterisation of DNA sequences which encode proteins expressed in the salivary glands of ticks, more particularly the *Ixodes ricinus* arthropod tick. Genes which are induced in the salivary glands of *Ixodes ricinus* during the slow-feeding phase of the blood meal were characterized. The cloning of these genes was carried out by setting up a subtractive cDNA library by subtracting uninduced cDNAs (synthesised from mRNAs expressed in

the salivary glands of unfed ticks) from induced cDNAs (synthesised from mRNAs expressed in the salivary glands at the end of the slow-feeding phase). A full-length cDNA library was set up starting from incomplete sequences deemed of interest and identified in the subtractive cDNA library. Also included in the invention are the various uses of the newly identified polynucleotides and derived polypeptides.

EP 1 892 297 A2

Description

FIELD OF INVENTION.

5 **[0001]** The invention relates to the molecular characterisation of DNA sequences which encode proteins expressed in the salivary glands of ticks, more particularly the *Ixodes ricinus* arthropod tick. These proteins are involved in the complex mechanism of interaction between this arthropod and its mammalian host. The invention relates to newly identified polynucleotides, polypeptides encoded by them and the use of such polynucleotides and polypeptides, and to their production.

BACKGROUND OF THE INVENTION.

10 **[0002]** Ticks are ectoparasites which infest a large number of animals such as mammals, birds, reptiles and amphibians. They are present in almost every area of the world. The tick feeding process itself is often harmful to the host. In addition, many of these ticks are vectors of viruses, bacteria and protozoa that cause host morbidity and, in some cases, mortality, particularly in humans and livestock animals. There are three families of ticks : the *Ixodidae* or hard ticks, the *Argasidae* or soft ticks, and the *Nuttalliellidae*. The life cycle of all ticks involves four stages (egg - larva - nymph - adult). In the majority of species, as *Ixodes ricinus*, the ticks drop off the host animal after each blood meal. The larvae hatch from the eggs and climb the vegetation where they come into easy reach of passing animals. Once on the host, they attach themselves and feed on blood. Nymphs and adults feed on other hosts and employ the same methods of host seeking. Mating of adults often takes place on the host while attached and feeding. Egg laying occurs after detachment.

15 **[0003]** The tick salivary gland play an important role in the accomplishment of the blood meal and in the transmission of pathogens. During the blood meal, *Ixodidae* ticks concentrate the blood by using special mechanisms which eliminate the excess of water and ions through salivary glands. A striking modification of the morphology and the physiology of the salivary glands occurs during this blood meal. The cytoplasm and the nucleus of several salivary gland cells enhance in volume leading to an important increase in the size and the weight of the salivary glands. The messenger RNA (mRNA) synthesis is also induced, resulting in the expression of new proteins. At the end of the meal, the degeneration of the salivary glands is probably caused by the 20-hydroxyecdysone, also called "salivary degenerating factor".

20 **[0004]** The salivary gland is rich in bio-active factors : cement, enzymes, enzyme inhibitors, histamine agonist and antagonist, anticoagulant factors, modulating factors of the host immune response, prostaglandin. Some of these interactive factors are already present in the salivary glands of unfed ticks; but others, mainly proteins, are induced during the feeding phase of the blood meal. These induced-proteinic factors seem to play an important role in the modulation of the host immune response. One of these factors, a 65 kDa protein, was isolated by Brossard and co-workers (Ganapamo *et al.*, 1997). This protein induces, *in vitro*, a specific CD4⁺ T cell proliferation of lymph node cells from mice infested with *I. ricinus* ticks (Ganapamo *et al.*, 1997). These cells produce high levels of interleukin-4 (IL-4) and low levels of interferon- γ (IFN- γ) when they are stimulated with concanavalin-A (Con A). This suggest a T_H2 polarisation of the cytokine pattern (Ganapamo *et al.*, 1995). The production of IL-5 and IL-10 confirms this phenomenon (Ganapamo *et al.*, 1996). This polarised response could constitute favourable conditions for the transmission of some pathogens.

25 **[0005]** The inhibition of the alternative pathway of complement activation, the decrease of the synthesis of antibodies induced by thymo-dependent antigens in infested animals, and the decrease of the proliferative activity of T-lymphocytes stimulated with mitogens, contribute to the setting up of these processes (Wikel *et al.*, 1996; Brossard and Wikel, 1997). Furthermore, prostaglandins (PGE₂) and salivary proteins are involved in the suppression of the immune response. In addition, it is known that some proteinic factors expressed by the salivary glands stimulate the growth of *Borrelia burgdorferi*, the causal agent of Lyme disease, which is the main human pathogen transmitted by the *Ixodidae* ticks (De Silva *et al.*, 1995)

SUMMARY OF THE INVENTION.

30 **[0006]** The *Ixodes ricinus* tick is the known vector of *Borrelia burgdorferi* sensu lato, the causal agent of Lyme disease. Tick salivary glands play an important role in the blood meal and in pathogen transmission. In the salivary glands, several genes are induced during the feeding process leading to the expression of new proteins which can play an important role in the modulation of the host immune and haemostatic responses.

35 **[0007]** It is accordingly important to identify and characterize those genes which are induced during the feeding process as well as those proteins which are expressed under those circumstances and to take benefit of the interesting properties of the substances thus identified.

40 **[0008]** Accordingly, in a first aspect, the present invention relates to a method for producing a library of cDNAs which are induced in the salivary glands of a tick, preferably an *Ixodes ricinus* tick, during the tick feeding phase which comprises :

- a) selectively cloning mRNAs induced during the tick feeding phase to obtain a corresponding cDNA library ;
- b) cloning full-length cDNAs corresponding to some incomplete cDNA sequences identified in the library obtained in step a).

5 **[0009]** More particularly, the genes induced are those induced during the slow-feeding phase of the blood meal.
[0010] In a preferred aspect, the method comprises:

- a) synthesising uninduced cDNAs starting from mRNAs expressed in the salivary gland of unfed ticks;
- 10 b) synthesising induced cDNAs starting from mRNAs expressed in the salivary gland of fed ticks ;
- c) subtracting said uninduced cDNAs from said induced cDNAs;
- d) isolating and cloning specifically induced cDNAs, thus obtaining a subtractive library;
- 15 e) obtaining corresponding full-length induced cDNA ;
- f) sequencing and comparing said full-length induced DNA molecules with known-polypeptide and polynucleotide sequences.

20 **[0011]** In a further preferred aspect, a full-length cDNA library is set up and screened by means of incomplete cDNAs isolated from the subtractive library.

[0012] More particularly, the method comprises:

- 25 a) randomly sequencing a number of clones of said subtractive library ;
- b) comparing their DNA and amino acid translated sequences with DNA and protein databases ;
- c) identifying distinct family sequences ;
- 30 d) characterising their corresponding full-length mRNA sequence.

[0013] Other features of said method are described hereafter.

35 **[0014]** According to another aspect, the invention relates to libraries obtained from fed tick salivary glands and particularly whenever obtained by a method as described above or equivalent thereto and to the use of such libraries in identifying genes induced during tick feeding.

[0015] The sequences identified by the above described methods can be divided into three classes : i) the first one refers to putative anticoagulant and anti-complement sequences; ii) the second one refers to putative immunomodulatory sequences; iii) the last class comprises sequences presenting low or no homologies to known polynucleotide and polypeptide sequences

40 **[0016]** According to other aspects, the present invention relates to polynucleotides isolated or isolatable from the tick salivary glands, polypeptides encoded by said polynucleotides as well as to the use of said polynucleotides and said polypeptides.

45 **[0017]** According to still another aspect, the invention comprises vectors including said nucleic acids or said polynucleotides.

[0018] A still further aspect of the invention includes cell lines transfected with said vectors.

[0019] Still another aspect of the invention includes methods of producing polypeptides by culturing above cell lines, possibly including additional steps of purifying polypeptide.

[0020] In another aspect, the inventions pertains to antibodies to said polypeptides.

50 **[0021]** In a further aspect, the invention includes hybridoma cell lines expressing antibodies.

[0022] In a still further aspect, the invention relates to compositions including said nucleic acids, polynucleotides, polypeptides or antibodies and pharmaceutically acceptable carrier.

[0023] Another aspect of the invention includes above compositions for use in a method of medical treatment.

55 **[0024]** Said polynucleotides, said polypeptides, said uses and other aspects of the invention including those stated hereabove are futher defined and described hereafter.

BRIEF DESCRIPTION OF THE DRAWINGS.

[0025]

5 Figure 1.
 RACE assay (Frohman et al., 1995) specific to Seq 16 and 24. The reverse transcription step was carried out using
 10 ng of mRNAs extracted from salivary gland of engorged ticks. The brightest bands represent the cDNA fragments
 corresponding to the 3' end of the targeted mRNA. The amplified products were subjected to agarose gel electro-
 phoresis followed by staining the DNA fragments by ethidium bromide. Molecular weight markers (M) were the
 10 Smart ladder (Life technologies, Rockville, Maryland, USA). Arrows indicate the position of the expected amplified
 products.

Figure 2.
 15 Differential expression analysis of the 5 full-length selected cDNAs and 9 cDNA fragment isolated in the subtractive
 library. PCR assays were carried out using as DNA template cDNAs obtained from a reverse transcription procedure
 on mRNAs extracted from salivary glands either of engorged (E) or of unfed (UF) ticks. These RNA messengers
 were also used as template in reverse transcription assays. Ten microliter of both PCR and RT-PCR mixture were
 subjected to agarose gel electrophoresis and ethidium bromide staining for the detection of amplified DNA products.
 20 [++] strongly positive; [+] positive; [-] negative.

DESCRIPTION OF THE INVENTION.

[0026] We have characterised genes which are induced in the salivary glands of *Ixodes ricinus* during the slow-feeding
 phase of the blood meal. The cloning of these genes was carried out by setting up two complementary DNA (cDNA)
 25 libraries. The first one is a subtractive library based on the methodology described by Lisitsyn et al. (Science 259,
 946-951, 1993) and improved by Diatchenko et al. (Proc. Natl. Acad. Sci. USA 93, 6025-6030, 1996). This library cloned
 selectively induced mRNA during the tick feeding phase. The second library is a full-length cDNA library which has been
 constructed by using the basic property of mRNAs (presence of a polyA tail in its 3' end and the cap structure in its 5'
 end). This cDNA library permitted the cloning of full-length cDNAs, corresponding to some incomplete cDNA sequences
 30 deemed of interest, and identified in the subtractive cDNA library.

[0027] The subtractive library was set up by subtracting uninduced-cDNAs (synthesised from mRNAs expressed in
 the salivary glands of unfed ticks) from induced-cDNAs (synthesised from mRNAs expressed in the salivary gland at
 the end of the slow-feeding phase). The induced-cDNAs was digested by a restriction enzyme, divided into two aliquots,
 and distinctively modified by the addition of specific adapters. As for the induced-cDNAs, the uninduced cDNAs was
 35 also digested by the same restriction enzyme and then mixed in excess to each aliquot of modified induced-cDNA. Each
 mixture of uninduced-/induced-cDNAs was subjected to a denaturation step, immediately followed by an hybridisation
 step, leading to a capture of homologous induced-cDNAs by the uninduced-cDNA. Each mixture was then mixed together
 and subjected again to a new denaturation/hybridisation cycle. Among the hybridised cDNA molecules, this final mixture
 comprises induced-cDNAs with different adapters at their 5' and 3' end. These relevant cDNAs were amplified by
 40 polymerase chain reaction (PCR), using primers specific to each adapter located at each end of the cDNA molecules.
 The PCR products were then ligated into the pCRII™ vector by A-T cloning and cloned in an TOP-10 E. coli strain. The
 heterogeneity of this subtractive library was evaluated by sequencing the recombinant clones. The "induced" property
 of these cDNA sequences was checked by reverse transcription-PCR (RT-PCR) on mRNA extracted from salivary glands
 of engorged and unfed ticks. Finally, the full-length induced-cDNA was obtained by screening the expression library
 45 using, as a probe, some incomplete induced-cDNAs isolated from the subtractive library. These full-length induced DNA
 molecules were sequenced and compared to known polypeptide and polynucleotide sequences.

[0028] The full-length cDNA library was set up by using the strategy developed in the "CapFinder PCR cDNA Library
 Construction Kit" (Clontech). This library construction kit utilises the unique CapSwitch™ oligonucleotide (patent pending)
 in the first-strand synthesis, followed by a long-distance PCR amplification to generate high yields of full-length, double-
 50 stranded cDNAs. All commonly used cDNA synthesis methods rely on the ability of reverse transcriptase to transcribe
 mRNA into single stranded DNA in the first-strand reaction. However, because the reverse transcriptase cannot always
 transcribe the entire mRNA sequence, the 5' ends of genes tend to be under-represented in cDNA population. This is
 particularly true for long mRNAs, especially if the first-strand synthesis is primed with oligo(dT) primers only, or if the
 mRNA has a persistent secondary structure. Furthermore, the use of T4 DNA polymerase to generate blunt cDNA ends
 55 after second-strand synthesis commonly results in heterogeneous 5' ends that are 5-30 nucleotides shorter than the
 original mRNA (D'Alessio, 1988). In the CapFinder cDNA synthesis method, a modified oligo(dT) primer is used to prime
 the first-strand reaction, and the CapSwitch oligonucleotide acts as a short, extended template at the 5' end for the
 reverse transcriptase. When the reverse transcriptase reaches the 5' end of the mRNA, the enzyme switches templates

and continues replicating to the end of the CapSwitch oligonucleotide. This switching in most cases occurs at the 7-methylguanosine cap structure, which is present at the 5' end of all eukaryotic mRNAs (Furuichi & Miura, 1975). The resulting full-length single stranded cDNA contains the complete 5' end of the mRNA as well as the sequence complementary to the CapSwitch oligonucleotide, which then serves as a universal PCR priming site (CapSwitch anchor) in the subsequent amplification. The CapSwitch-anchored single stranded cDNA is used directly (without an intervening purification step) for PCR. Only those oligo(dT)-primed single stranded cDNAs having a CapSwitch anchor sequence at the 5' end can serve as templates and be exponentially amplified using the 3' and 5' PCR primers. In most cases, incomplete cDNAs and cDNA transcribed from polyA⁻ RNA will not be recognised by the CapSwitch anchor and therefore will not be amplified.

[0029] At the end of these reactions, the full-length cDNA PCR products was ligated into the pCRII cloning vector (Invitrogen) and used for the transformation of XL2 *E. coli* strain. The full-length cDNA library was then screened by using, as a probe, the incomplete induced-cDNAs isolated from the subtractive library.

[0030] Eighty-nine clones of subtractive library were randomly sequenced, and their DNA and amino acid translated sequences were compared to DNA and protein databases. Among these, 27 distinct family sequences were identified, and 3 of them were selected for further characterisation of their corresponding full-length mRNA sequence. These 3 sequences matched the sequence of i) the human tissue factor pathway inhibitor (TFPI), ii) the human thrombin inhibitor gene, and iii) a snake venom zinc dependant metallopeptidase protein. These genes encode proteins which could be involved in the inhibition of the blood coagulation. The other 24 family sequences presented low or no homologies with polynucleotide and polypeptide sequences existing in databases. Screening of the full-length cDNA library using oligonucleotide probes specific to the 3 previously selected subtractive clones led to the recovery of the corresponding full-length cDNAs. Random screening of this library led to the selection of 2 other clones. One is closely homologous to an interferon-like protein, whereas the other shows homologies to the *Rattus norvegicus* leucocyte common antigen related protein.

Definitions.

[0031] "Putative anticoagulant, anti-complement and immunomodulatory" polypeptides refer to polypeptides having the amino acid sequence encoded by the genes defined in the table. These present homologies with anticoagulant, anti-complement and immunomodulatory polypeptides already existing in databases. These polypeptides belong to the Class I and Class II sequences (see table)

[0032] "Putative anticoagulant, anti-complement and immunomodulatory" cDNAs refer to polynucleotides having the nucleotide sequence defined in the table, or allele variants thereof and/or their complements. These present homologies with anticoagulant, anti-complement and immunomodulatory polynucleotides already existing in databases. These cDNAs belong to the Class I and Class II sequences (see table)

[0033] Some polypeptide or polynucleotide sequences present low or no homologies with already existing polypeptides or polynucleotides in databases. These belong to the Class III (see table).

[0034] « Polypeptide » refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslational natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a hem moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-linkings, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenylation, sulfation, transfer-RNA mediated addition of amino of amino acids to proteins such as arginylation, and ubiquitination. See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Comany, New York, 1993 and Wolt, F., Posttranslational Protein Modifications : Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", Meth

Enzymol (1990) 182 : 626-646 and Rattan et al, "Protein Synthesis : Posttranslational Modifications and Aging", Ann NY Acad Sci (1992) 663 : 48-62.

[0035] "Polynucleotide" generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double- stranded RNA, and RNA that is a mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "Polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term Polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "Polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

[0036] "Variant" as the term is used herein, is a Polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions (preferably conservative), additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis. Variants should retain one or more of the biological activities of the reference polypeptide. For instance, they should have similar antigenic or immunogenic activities as the reference polypeptide. Antigenicity can be tested using standard immunoblot experiments, preferably using polyclonal sera against the reference polypeptide. The immunogenicity can be tested by measuring antibody responses (using polyclonal sera generated against the variant polypeptide) against purified reference polypeptide in a standard ELISA test. Preferably, a variant would retain all of the above biological activities.

[0037] "Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identify" *per se* has an art-recognized meaning and can be calculated using published techniques. See, e.g. : (COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING : INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds, Humana Press, New Jersey, 1994; SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heijne, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds, M Stockton Press, New York, 1991). While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans (Carillo, H., and Lipton, D., SIAM JApplied Math (1998) 48 : 1073). Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48 : 1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCG program package (Devereux, J., et al., J Molec Biol (1990) 215 : 403). Most preferably, the program used to determine identity levels was the GAP program, as was used in the Examples below.

[0038] As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include an average up to five point mutations per each 100 nucleotides of the reference nucleotide sequence. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

Polypeptides of the invention.

[0039] The present invention relates to proteins (or polypeptides) secreted by *I. ricinus* salivary glands. These polypeptides include the polypeptides encoded by the cDNAs defined in the table; as well polypeptides comprising the amino acid sequence encoded by the cDNAs defined in the table; and polypeptides comprising the amino acid sequence which have at least 75 % identity to that encoded by the cDNAs defined in the table over their entire length, and preferably at least 80 % identity, and more preferably at least 90 % identity. Those with 95-99 % are highly preferred.

[0040] The *I. ricinus* salivary gland polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It may be advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which help in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

[0041] Fragments of the *I. ricinus* salivary gland polypeptides are also included in the present invention. A fragment is a polypeptide having an amino acid sequence that is the same as part, but not all, of the amino acid sequence of the aforementioned *I. ricinus* salivary gland polypeptides. As with *I. ricinus* salivary gland polypeptides, fragment may be "free-standing" or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of the polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

[0042] Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of the *I. ricinus* salivary gland polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus and / or transmembrane region or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus. Also preferred are fragments characterised by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate *I. ricinus* salivary gland protein activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that antigenic or immunogenic in an animal or in a human.

[0043] Preferably, all of these polypeptide fragments retain parts of the biological activity (for instance antigenic or immunogenic) of the *I. ricinus* salivary gland polypeptides, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions - i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination. Most preferred variants are naturally occurring allelic variants of the *I. ricinus* salivary gland polypeptide present in *I. ricinus* salivary glands.

[0044] The *I. ricinus* salivary gland polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

Polynucleotides of the invention.

[0045] Another aspect of the invention relates to *I. ricinus* salivary gland cDNAs (polynucleotides). These include isolated polynucleotides which encode *I. ricinus* salivary gland polypeptides and fragments respectively, and polynucleotides closely related thereto. More specifically, *I. ricinus* salivary gland cDNAs of the invention include a polynucleotide comprising the nucleotide sequence of cDNAs defined in the table, encoding a *I. ricinus* salivary gland polypeptide. The *I. ricinus* salivary gland cDNAs further include a polynucleotide sequence that has at least 75% identity over its entire length to a nucleotide sequence encoding the *I. ricinus* salivary gland polypeptide encoded by the cDNAs defined in the table, and a polynucleotide comprising a nucleotide sequence that is at least 75% identical to that of the cDNAs defined in the table. In this regard, polynucleotides at least 80% identical are particularly preferred, and those with at least 90% are especially preferred. Furthermore, those with at least 95% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred, with at least 99% being the most preferred. Also included under *I. ricinus* salivary gland cDNAs is a nucleotide sequence which has sufficient identity to a nucleotide sequence of a cDNA defined in the table to hybridize under conditions useable for amplification or for use as a probe

or marker. The invention also provides polynucleotides which are complementary to such *I. ricinus* salivary gland cDNAs.

[0046] The nucleotide sequence encoding *I. ricinus* salivary gland polypeptide encoded by the cDNAs defined in the table may be identical to the polypeptide encoding sequence contained in the genes defined in the table, or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide encoded by the genes defined in the table respectively.

[0047] When the polynucleotides of the invention are used for the recombinant production of an *I. ricinus* salivary gland polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or preproprotein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz et al, Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag, or is glutathione-s-transferase. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polydenylation signals, ribosome binding sites and sequences that stabilize mRNA.

[0048] Further preferred embodiments are polynucleotides encoding *I. ricinus* salivary gland protein variants comprising the amino acid sequence of the *I. ricinus* salivary gland polypeptide encoded by the cDNAs defined by the table respectively in which several, 10-25, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination. Most preferred variant polynucleotides are those naturally occurring *I. ricinus* sequences that encode allelic variants of the *I. ricinus* salivary gland proteins in *I. ricinus*.

[0049] The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridisation will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

[0050] Polynucleotides of the invention which are identical or sufficiently identical to a nucleotide sequence of any gene defined in the table or a fragment thereof, may be used as hybridisation probes for cDNA clones encoding *I. ricinus* salivary gland polypeptides respectively and to isolate cDNA clones of other genes (including cDNAs encoding homologs and orthologs from species other than *I. ricinus*) that have a high sequence similarity to the *I. ricinus* salivary gland cDNAs. Such hybridisation techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides, preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides. In one embodiment, to obtain a polynucleotide encoding *I. ricinus* salivary gland polypeptide, including homologs and orthologs from species other than *I. ricinus*, comprises the steps of screening an appropriate library under stringent hybridisation conditions with a labelled probe having a nucleotide sequence contained in one of the gene sequences defined by the table, or a fragment thereof; and isolating full-length cDNA clones containing said polynucleotide sequence. Thus in another aspect, *I. ricinus* salivary gland polynucleotides of the present invention further include a nucleotide sequence comprising a nucleotide sequence that hybridise under stringent condition to a nucleotide sequence having a nucleotide sequence contained in the cDNAs defined in the table, or a fragment thereof. Also included with *I. ricinus* salivary gland polypeptides are polypeptides comprising amino acid sequences encoded by nucleotide sequences obtained by the above hybridisation conditions. Such hybridisation techniques are well known to those of skill in the art. Stringent hybridisation conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising : 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1xSSC at about 65°C.

[0051] The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

Diagnostic Assays

[0052] This invention also relates to the use of *I. ricinus* salivary gland polypeptides, or *I. ricinus* salivary gland polynucleotides, for use as diagnostic reagents.

[0053] Materials for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy.

[0054] Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease which comprises :

- (a) an *I. ricinus* salivary gland polynucleotide, preferably the nucleotide sequence of one of the gene sequences defined by the table, or a fragment thereof;

- (b) a nucleotide sequence complementary to that of(a);
 (c) an *I. ricinus* salivary gland polypeptide, preferably the polypeptide encoded by one of the gene sequences defined in the table, or a fragment thereof;
 (d) an antibody to an *I. ricinus* salivary gland polypeptide, preferably to the polypeptide encoded by one of the gene sequences defined in the table; or
 (e) a phage displaying an antibody to an *I. ricinus* salivary gland polypeptide, preferably to the polypeptide encoded by one of the cDNAs sequences defined in the table.

[0055] It will be appreciated that in any such kit, (a), (b), (c), (d) or (e) may comprise a substantial component.

[0056] The anti-*I. ricinus* salivary gland polypeptide antibodies may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the *I. ricinus* salivary gland polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL TDM adjuvant. The immunization protocol may be selected by one skilled in the art without undue experimentation.

[0057] The anti-*I. ricinus* salivary gland polypeptide antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature. 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro.

[0058] The immunizing agent will typically include the *I. ricinus* salivary gland polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, Monoclonal Antibodies : Principles and Practice Academic Press, (1986) pp. 59-103). Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

[0059] Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Rockville, Maryland. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, J Immunol, 133:3001 (1984) ; Brodeur et al., Monoclonal Antibody Production Techniques and Applications Marcel Dekker, Inc., New York, (1987) pp 51-63).

[0060] The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against *I. ricinus* salivary gland polypeptide. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal Biochem., 107:220 (1980).

[0061] After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, supra). Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively the hybridoma cells may be grown in vivo as ascites in a mammal.

[0062] The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

[0063] The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U. S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred

source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567 ; Morrison et al., *supra*) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

[0064] The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

[0065] In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

[0066] As an alternative or supplement to immunising a mammal with a peptide, an antibody specific for a protein may be obtained from a recombinantly produced library of expressed immunoglobulin variable domains, e.g. using lambda bacteriophage or filamentous bacteriophage which display functional immunoglobulin bindings domains on their surfaces ; for instance see WO92/01047. The library may be native, that is constructed from sequences obtained from an organism which has not been immunised with any of the proteins (or fragments), or may be one constructed using sequences obtained from an organism which has been exposed to the antigen of interest.

Vaccines.

[0067] Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with *I. ricinus* salivary gland polypeptide or epitope-bearing fragments, analogs, outer-membrane vesicles or cells (attenuated or otherwise), adequate to produce antibody and/or T cell immune response to protect said animal from bacteria and viruses which could be transmitted during the blood meal of *I. ricinus* and related species. In particular the invention relates to the use of *I. ricinus* salivary gland polypeptides encoded by the cDNAs defined in the table. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which comprises, delivering *I. ricinus* salivary gland polypeptide via a vector directing expression of *I. ricinus* salivary gland polynucleotide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases (Lyme disease, tick encephalitis virus disease, ...).

[0068] A further aspect of the invention relates to an immunological composition or vaccine formulation which, when introduced into a mammalian host, induces an immunological response in that mammal to a *I. ricinus* salivary gland polypeptide wherein the composition comprises a *I. ricinus* salivary gland cDNA, or *I. ricinus* salivary gland polypeptide or epitope-bearing fragments, analogs, outer-membrane vesicles or cells (attenuated or otherwise) the vaccine formulation may further comprise a suitable carrier. The *I. ricinus* salivary gland polypeptide vaccine composition is preferably administered orally or parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example; sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity to the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

[0069] Yet another aspect relates to an immunological/vaccine formulation which comprises the polynucleotide of the invention. Such techniques are known in the art, see for example Wolff et al , Sciences, (1990) 247 : 1465-8. **Therapeutics.**

[0070] Another aspect of the invention related to the use of these *I. ricinus* salivary gland polypeptides as therapeutics agent. In considering the particular potential therapeutic areas for the likely products, the hospital disciplines cover by these products are : haematology (particularly coagulation clinics), transplantation (for immunosuppression control), rheumatology (for anti-inflammatories), and general treatment (for specific or improved anesthetics).

EP 1 892 297 A2

Table 1 : Sequences identified in the subtractive and the cDNA full-length libraries.

Gene	Motifs	Similar sequences in databases	Score	Class
5		No significant identity		III
		No significant identity		III
		No significant identity		III
		No significant identity		III
10	Prokariotic mbrane lipoprotein lipid attachment site	No significant identity		III
		R. melioli Nitrogen fixation (fixF)	0.00089	III
15		Human Apolipoprotein B-100	0.0045	III
		Hu. mRNA for cAMP response element (CRE-BP1) binding prot	0.057	III
	Kunitz family of serine protease inhibitor	Human BAC clone GS345D13 H. sap Tissue factor Pathway Inhibitor PI-2	4.7 ⁻¹³ 4 ¹²	I
20	Prokariotic mbrane lipoprotein lipid attachment site	No significant identity		III
		Pea mRNA for GTP binding prot.	0.48	III
		No significant identity		III
25		IL-11 R-Béta gene	0.18	II
		No significant identity		III
		C. gloeosporioides cutinase gene	0.082	III
30		No significant identity		III
		Mouse mRNA for secretory prot cont. thranspondine motifs	0.014	III
35	Zinc dependent metallopeptidase family	B. jararaca mRNA for jararhagin Agkistrodon contortrix metalloproteinase precursor	1.1 ⁻⁵ 3.9 ⁻⁵	II
		O. aries gene for ovine INF-alpha	0.7	II
40		Interferon-omega 45	0.88	II
		Interferon-omega20	0.89	II
		RCPT PGE2	0.85	III
		PGE Rcpt EP2	0.85	III
45		No significant identity		III
		IgG1 L chain directed against human IL2 rcpt Tac p	0.19	II
		rot Var region of light chain of MAK447/179	0.2	II
50		No significant identity		III
		No significant identity		III
		Mus Musculus neuroactin	o.42	III
55		No significant identity		III

EP 1 892 297 A2

(continued)

Gene	Motifs	Similar sequences in databases	Score	Class
Seq. 24		H. sapiens thrombin inhibitor	2,1 ⁻¹²	I
		Cytoplasmic antiproteinase 38 kDa intracellular serine prot.	2.3 ⁻¹²	I
Seq. 25		No significant identity		III
Seq. 26		No significant identity		III
Seq.27		Mus musculus transcription factor ELF3 (fasta)	0.053	III
Seq. 28		Homo sapiens putative interferon-related protein (SM15) mRNA	1.70E-22	II
Seq. 29		R.norvegicus mRNA for leucocyte common antigen-related protein	4.80E-09	II

Class I : putative anticoagulant homologs ; Class II : putative immunomodulatory homologs ; Class III : low or no homologies found in the databases.

TABLE 2. Biological characteristics of the selected clones

The full sequences of 5 selected clones were compared with EMBL/GenBank data bases using the tFasta and Blastp algorithms. These sequences were analysed for the presence of a Kozak consensus sequence indicated in the column entitled "Nucleotide in position -3". By applying the algorithm of the GCG*pnognam the deduced amino sequences were analysed for the presence of specific protein units on motifs (algorithm motif) and for the presence of a signal peptide sequence (von Heijne and McGeoch analysis).

Clone	Full-length sequences similarly to databases to databases	IFasta/ Blastp Scores ^a	QRF (aa)	Motifs motifs	Signal peptide scores	Sp length/ Prob.	Nucleotide in position -3 ^c
Seq28	<i>Homo sapiens</i> putative interferon related gene (SKMc15) [U09585]	1.8 10 ³⁸ /1.10 ⁻⁷¹	426		5.4 d Fe	48 aa / 84 10 ⁻¹	G
Seq29	<i>R. norvegicus</i> leukocyte common antigen (LAR) mRNA [X83546]	7.8.10 ⁻¹¹ / N 7,8.10 ⁻¹¹ / N	274		10.2/S	18 aa / 7.4.10 ⁻⁷	A
Seq 16	Mouse mRNA for secretory protein containing thrombospondin motives [D67076]	0.002/6.10 ⁻⁷	489	Metallopeptidase	7.9/S	19 aa/ 7.4.10 ⁻⁴	G
Seq24	Pig leukocyte elastase inhibitor mRNA. [P80229]	0/7.10 ⁻⁶⁷ 0/7.10 ⁻⁶⁷	378	Serpin	8.5/S	51 aa / 3.28. 10 ⁻³ 51 aa/ 3,28.10 ⁻³	A A

(continued)

5	Clone	Full-length sequences similarly to databases to databases	IFasta/ Blastp Scores ^a	QRF (aa)	Motifs motifs	Signal peptide scores	Sp length/ Prob.	Nucleotide in position -3 ^c
10	Seq7	Human Tissue Factor Pathway Inhibitor [P48307]	4.8.10 ⁻¹² / 2.10 ⁻⁵	87	Kunitz	6.5/S	19 aa/ 1.8.10 ⁻¹	G

a No score (N)

b Succeeded (S) and Failed (F)

c Guanine (G) and Adenine (A)

d von Heijne analysis

McGeoch analysis

* GCG = Genetic Computer Group, Madison Wisconsin, 10.0 version in Unix, January 1999.

20 EXAMPLES

[0071] Biological materials used in this study.

[0072] The salivary glands of 5 day engorged or unfed free of pathogen Ixodes ricinus female adult ticks were used in this work. When removed, these glands were immediately frozen in liquid nitrogen and stored at -80°C. To extract RNA messengers (mRNA), the salivary glands were crushed in liquid nitrogen using a mortar and a pestle. The mRNAs were purified by using an oligo-dT cellulose (Fast Track 2.0 kit, Invitrogen, Groningen, The Netherlands). Two micrograms of mRNAs were extracted from 200 salivary glands of fed ticks; and 1.5 µg of mRNAs were also extracted from 1,000 salivary glands of unfed ticks.

30 Example 1 : Construction of a Representational

[0073] Difference Analysis (RDA) subtractive library.

[0074] All procedures were performed as described by Hubank and Schatz (1994). Double-stranded cDNAs were synthesised using the Superscript Choice System (Life Technologies, Rockville, Maryland, USA). The cDNAs were digested with DpnII restriction enzyme, ligated to R-linkers, amplified with R-24 primers (Hubank and Schatz, 1994), and finally digested again with the same enzyme to generate a "tester" pool consisting of cDNAs from salivary glands of fed ticks and a "driver" pool consisting of cDNAs from salivary glands of unfed ticks. The first round of the subtractive hybridisation process used a tester/driver ratio of 1:100. The second and third rounds utilised a ratio of 1:400 and 1:200,000, respectively. After three cycles of subtraction and amplification, the DpnII-digested differential products were subdivided according to size into 4 different fractions on a 1.7% electrophoresis agarose gel, and subcloned into the BamHI site of the pTZ19r cloning vector. The ligated product was used to transform TOP-10 E. coli competent cells (Invitrogen, Groningen, The Netherlands). Nine thousand six hundred clones of this subtractive library were randomly selected, and individually put in 100 microplates and stored at -80°C. This subtractive library was analysed by sequencing 89 randomly chosen clones, using M13 forward and reverse primers specific to a region located in the pT19r cloning vector. The DNA sequences of these 89 clones were compared, and 27 distinct family sequences were identified. Homology of these sequences to sequences existing in databases is presented in Table 1. The subtractive sequences 1 to 27 are presented in the sequence-listing file (except for sequences 16 and 24 whose complete mRNA sequences are presented; see also Example 2). Three sequences (Seq 7, 16 and 24) were selected for further characterisation of their corresponding full-length mRNA sequence. These 3 sequences matched the sequence of i) the human tissue factor pathway inhibitor (TFPI), ii) a snake venom zinc dependant metalloproteinase protein, and iii) the human thrombin inhibitor protein, corresponding to Seq 7, 16 and 24, respectively. These genes encode proteins which could be involved in the inhibition of the blood coagulation.

55 Example 2 : Construction of the full length cDNA library and recovery of full length cDNAs sequence.s by screening of this full length cDNA library.

[0075] This library was set up using mRNAs extracted from salivary glands of engorged ticks. The mRNAs (80 ng) were subjected to reverse transcription using a degenerated oligo-dT primer (5'-A(T)30VN-3'), the SmartTM oligonucle-

otide (Clontech, Palo Alto, USA), and the Superscript II reverse transcriptase (Life Technologies, Rockville, Maryland, USA). The single strand cDNA mixture was used as template in a hot start PCR assay including the LA Taq polymerase (Takara, Shiga, Japan), the modified oligo-dT primer and a 3'-'Smart' primer specific to a region located at the 5' end of the Smart™ oligonucleotide. The PCR protocol applied was : 1 min at 95°C, followed by 25 sec at 95°C / 5 min at 68°C, 25 times, and 10 min at 72°C. The amplified double stranded cDNA mixture was purified with a Centricon 30 concentrator (Millipore, Bedford, USA). The eDNAs were divided into 4 fractions ranging from 0.3 to 0.6 kb, 0.6 to 1 kb, 1 kb to 2 kb and 2 kb to 4 kb on a 0,8% high grade agarose electrophoresis gel and recovered separately by using the Qiaex II extraction kit (Qiagen, Hilden, Germany). The 4 fractions were ligated individually into the pCRII cloning vector included in the TOPO cloning kit (Invitrogen, Groningen, The Netherlands). The ligated fractions were then used to transform XL2-Blue ultracompetent E. coli cells (Stratagene, Heidelberg, Germany). The resulted recombinant clones were stored individually in microplates at -80°C. Ten clones were randomly chosen for partial or complete sequencing. As a result of this procedure, 2 cDNA sequences (Seq 28 and Seq 29, see Table 1) were selected for their homology to sequence databases. One is closely homologous to an interferon-like protein (Seq 28), whereas the other shows homologies to the *Rattus norvegicus* leucocyte common antigen-related protein (Seq 29).

[0076] The 4 different fractions of the full-length cDNA library were screened with radio-labelled oligonucleotide probes specific to selected clones identified in the subtractive cDNA library. The labelling of these oligo probes was performed as described in Current Protocols in Molecular Biology (Ausubel et al, 1995, J. Wiley and sons, Eds). These 4 different fractions were then plated on nitrocellulose membranes and grown overnight at 37°C. These membranes were denatured in NaOH 0.2M / NaCl 1.5M, neutralised in Tris 0.5M pH 7.5-NaCl 1.5M and fixed in 2X SSC (NaCl 0.3 M/ Citric Acid Trisodium di-hydrated 0.03 M). The membranes were heated for 90 min. at 80°C, incubated in a pre-hybridisation solution (SSC 6X, Denhardt's 10X, SDS 0,1%) at 55°C for 90 min., and finally put overnight in a preheated hybridisation solution containing a specific radio-labelled oligonucleotide probe at 55 °C. The hybridised membranes were washed 3 times in a SSC 6X solution at 55 °C for 10 min, dried and exposed on Kodak X-OMAT film overnight at -80°C. The full length cDNA library was also analysed by sequencing a set of clones. The resulted DNA sequences were compared to ENBL/ Gene Bank databases and were used to set up oligonucleotide probes to recover other corresponding clones. In this way, the complete consensus mRNA sequence of the Seq 28 and 29 was confirmed by the recovery of two other clones corresponding to these sequences. Only one full-length cDNA clone corresponding to the subtractive clone 16 was isolated. Therefore, to identify the complete sequence of the Seq 16 and 24, the Rapid Amplification of cDNA Ends (RACE) method was applied.

[0077] The RACE methodology was performed as described by Frohman et al. (1995). The reverse transcription step was carried out using 10 ng of mRNAs extracted from salivary gland of engorged ticks and the Thermoscript Reverse transcriptase (Life technologies, Rockville, Maryland, USA). All gene specific primers (GSP) had an 18 base length with a 61% G/C ratio. The amplified products were subjected to an agarose gel electrophoresis and recovered by using an isotachophorese procedure. The cDNAs were cloned into the pCRII-TOPO cloning vector (Invitrogen, Groningen, The Netherlands). To identify the consensus cDNA sequence, different clones were sequenced., and their sequence was compared to their known corresponding sequence. Therefore, the complete cDNA sequences of the clones 16 and 24 isolated in the subtractive library were obtained by this RACE procedure (figure 1).

Example 3 : Analysis of the full sequences of 5 selected clones.

[0078] The sequences of selected clones (Seq. 7, 16, 24, 28 and 29) allowed identification of the open reading frames, from which the amino sequence were deduced. These potential translation products have a size between 87 and 489 amino acids (see table 2). In order to evaluate, in silico, their respective properties, the amino acid sequences and the nucleotide sequences of said 5 open frames were compared with the databases using the tFasta and Blastp algorithms. These comparisons show that Seq. 7 is highly homologous to the human Tissue Factor Pathway Inhibitor (TFPI). TFPI is an inhibitor of serine proteases having 3 domains of the Kunitz Protease Inhibitor (KPI) type. Each of these units or motifs has a particular affinity for different types of proteases. The first and second KPI domains are responsible for the respective inhibition of VIIa and Xa coagulation factors. The third KPI domain apparently has no inhibitory activity. It should be noted that the sequence coded by the Seq. 7 clone is homologous to the regio of the first KPI domain of TFPI and that the KPI is perfectly kept therein. This similarity suggests that the Seq. 7 protein is a potential factor VIIa inhibitor.

[0079] The amino sequence deduced from the Seq. 28 clone has a great homology with 3 database sequences, namely : mouse TIS7 protein, rat PC4 protein and human SKMcl5 protein. These 3 proteins are described as putative interferon type factors. They possess very well conserved regions of the B2 interferon protein. Therefore, it is proposed that the Seq. 28 protein has immunomodulatory activity.

[0080] Sequences 16 and 24 were compared with databases thus showing their homology respectively with *Gloydius halys* (sub-order of ophidians) M12b metallopeptidase and porcine elastase inhibitor belonging to the super-family of the serine protease inhibitors (Serpin). The amino sequences of these 2 clones also have specific units of said families. It is proposed that said proteins have anticoagulant and immuno-modulatory properties.

[0081] Finally, the Seq. 29 clone has a weak homology with the *R. norvegicus* leucocyte common antigen (LAR). The latter is an adhesion molecule. It is thus possible that the Seq. 29 protein has immunomodulatory properties related to those expressed by the LAR protein.

[0082] Due to their potential properties, most of the proteins examined are expected to be secreted in the tick saliva during the blood meal. Accordingly, tests were made for finding the presence of a signal peptide at the beginning of the deduced amino sequences. All of the results by the Von Heijne analysis method were positive. By the McGeoch method, signal peptide sequences were detected for the Seq. 7, 16, 24 and 29 deduced amino sequences. It seems that said proteins are secreted in the tick salivary gland. Furthermore, the presence of a Kozak consensus sequence was observed upstream of the coding sequences of all examined clones. This indicates that their mRNAs potentially could be translated to proteins.

Example 4 : Evaluation of the differential expression of the cDNA clones isolated in the subtractive and full-length cDNA libraries.

[0083] The differential expression of the mRNAs corresponding to the 5 full-length selected clones (Seq 7, 16, 24, 28 and 29) and of 9 subtractive clones was assessed using a PCR and a RT-PCR assays (figure 2).

[0084] The PCR assays were carried out using as DNA template cDNAs obtained from a reverse transcription procedure on mRNAs extracted from salivary glands either of engorged or of unfed ticks. Each PCR assay included pair of primers specific to each target subtractive or cDNAs full-length sequence. PCR assays were performed in a final volume of 50 μ l containing 1 μ M primers, 0.2 mM deoxynucleotide (dATP, dCTP, dGTP and dTTP; Boehringer Mannheim GmbH, Mannheim, Germany), PCR buffer (10 mM Tris-HCl, 50 mM KCl, 2.5 mM MgCl₂, pH 8.3) and 2.5 U of Taq DNA polymerase (Boehringer mannheim GmbH, Mannheim, Germany). DNA samples were amplified for 35 cycles under the following conditions : 94°C for 1 min., 72 °C for 1 min. and 64 °C for 1 min, followed by a final elongation step of 72 °C for 7 min.

[0085] The RT-PCR assay was carried out on the 5 selected full-length cDNA clones and on 5 cDNA subtractive clones. The mRNAs used as template in the reverse transcription assay was extracted from salivary glands of engorged and unfed *I. ricinus* ticks. The reverse transcription assays were performed using a specific primer (that target one the selected sequences) and the "Thermoscript Reverse transcriptase" (Life technologies, Rockville, Maryland, USA) at 60°C for 50 min. Each PCR assay utilised the reverse transcription specific primer and an another specific primer. The PCR assays were performed in a final volume of 50 μ l containing 1 μ M primers, 0.2 mM deoxynucleotide (dATP, dCTP, dGTP and dTTP; Boehringer Mannheim GmbH, Mannheim, Germany), PCR buffer (10 mM Tris-HCl, 50 mM KCl, 2.5 mM MgCl₂, pH 8.3) and 2.5 U of Expand High Fidelity polymerase (Roche, Bruxelles, Belgium). Single stranded DNA samples were amplified for 30 cycles under the following conditions : 95°C for 1 min., 72 °C for 30 sec. and 60°C for 1 min, followed by a final elongation step of 72 °C for 7 min. The figure 2 shows that the expression of the selected sequences is induced in salivary glands of 5 day engorged ticks, except for the sequence 28 that is expressed at a similar level in salivary glands of engorged and unfed ticks. The expression of the other mRNAs could be either induced specifically or increased during the blood meal.

Example 5 : Expression of recombinant proteins in mammal cells.

[0086] The study of the properties of isolated sequences involves the expression thereof in expression systems allowing large amounts of proteins encoded by these sequences to be produced and purified. These experiments are described hereafter as well as in example 6.

5.1. Sub-cloning of the sequences in the p CDNA3. 1-His/V5 (Invitrogen) vector.

[0087] The DNA sequences of the 5 selected clones (Seq. 7, 16, 24, 28 and 29) were transferred into the p CDNA3.1 His/V5 expression vector. Said vector allows the expression of heterologous proteins fused to a tail of 6 histidines as well as to the V5 epitope in eucaryotic cells. The different DNAs were produced by RT-PCR by using primers specific to the corresponding clones. These primers were constructed so as to remove the stop codon of each open reading frame or phase in order to allow the protein to be fused to the 6xHIS/Epitope V5 tail. In addition, the primers contained restriction sites adapted to the cloning in the expression vector. Care was taken to use, when amplifying, a high fidelity (Promega) Pfu polymerase.

[0088] The transient expression of the Seq. 16 and 24 recombinant proteins was measured after transfection of the Seq. 16 and Seq 24 - pCDNA3.1-His/V5 constructions in COS1 cells, using Fugen 6 (Boehringer). The protein extracts of the culture media corresponding to times 24, 48 and 72 hours after transfection were analysed on acrylamide gel by staining with Coomassie blue or by Western blot using on the one hand an anti-6x histidine antibody or on the other hand Nickel chelate beads coupled to alkaline phosphatase. These analyses showed the expression of said proteins in

the cell culture media.

Example 6 : Expression of proteins in *E. coli*

5 6.1. Insertion of coding sequences into the pMAL-C2E expression vector.

[0089] It also was decided to express various proteins in bacteria by using the pMAL-C2E (NEB) vector. Said vector expresses the various proteins fused with the Maltose-Binding-Protein (MBP). The protein of interest then could be separated from the MBP thanks to a site separating the MBP from the protein, said site being specific to protease enterokinase.

10 [0090] In order to express optimally the 5 sequences examined, using the p MAL-C2E vector, PCR primer pairs complementary to 20 bases located upstream of the STOP codon and to 20 bases located downstream of the ATG of the open reading frame or phase were constructed. In that way, the amplified CDNA fragments only comprised the coding sequence of the target mRNA provided with its stop codon. The protein of interest was fused to MBP by its N-terminal end. On the other hand, since these primers contained specific restriction sites specific to the expression vector, it was possible to effect direct cloning of the cDNAs. The use of Pfu DNA polymerase (Promega) made it possible to amplify the cDNAs without having to fear for errors introduced into the amplified sequences.

15 [0091] The coding sequences of clones Seq 7, 16, 24 and 29 were reconstructed in that way. Competent TGI cells of *E. coli* were transferred using these constructions. Enzymatic digestions of these mini-preparations of plasmidic DNA made it possible to check that the majority of clones Seq. 7, 16, 24 and 29-p-MALC2-E effectively were recombinant.

20 6.2. Expression of recombinant proteins.

[0092] Starting from various constructions cloned in *E. coli* TG1, the study of the expression of recombinant proteins fused with MBP was initiated for all sequences of interest (i.e. Seq. 7, 16, 24 and 29) except for Seq. 28. The culture of representative clones of Seq. 7, 16, 24 and 29 as well as negative controls (non recombinant plasmids) were undertaken to induce the expression of recombinant proteins therein. These cultures were centrifuged and the sediments (culots) were separated from the media for being suspended in 15mM pH7.5 Tris and passed to the French press. The analysis of these samples by (10%) acrylamide gel coloured with Coomassie blue or by Western Blot using rabbit anti-MBP antibodies, showed the expression of recombinant proteins Seq. 7 (~50kDa), Seq. 16 (~92kDa), Seq 24 (~80kDa) and Seq 29 (~67kDa).

30 Example 7 : Production of antibodies.

[0093] The Seq. 7, 16 and 24 protein were injected into groups of 4 mice with the purpose of producing antibodies directed against said proteins. The first antigen injected was made with the complete Freund adjuvant. Two weeks later, a recall injection was made with incomplete Freund adjuvant. The sera of mice injected with Seq. 16 provided positive tests for anti-MBP antibodies.

40

45

50

55

SEQUENCE LISTING

5 (1) INFORMATION FOR SEQ ID N°: 1

(i) SEQUENCE CHARACTERISTICS :

- 10 (a) LENGTH : 194 base pairs
 (b) TYPE : nucleic acid
 (c) STRANDEDNESS : single
 (d) TOPOLOGY : linear

15 (ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°1 :

20 1 ATACCTTCCA CTTGTAGCCC TTCCTCATCC GATATGGTGA CGGATGCCAT
 51 TGCATCCTCG TCGTGGAAGA GGTCTCTTTC TAAATAAGAC CCATCCATAT
 25 101 ATGTGTGTTT GCGAATGCCG TCGACGTAGC TCCTGACTAG AAACCTCGTCG
 151 GCTAGGACAG AACTTTTCTT CAGGTTTAGC GTAATGTCCT CGTT

30 (2) INFORMATION FOR SEQ ID N°: 2

(i) SEQUENCE CHARACTERISTICS :

- 35 (a) LENGTH : 607 base pairs
 (b) TYPE : nucleic acid
 (c) STRANDEDNESS : single
 (d) TOPOLOGY : linear

40 (ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°2 :

45 1 TACCNGGGAA TCCAAAACCA ATTTTATTG GAACTCCAC GTCTTCTTCA
 51 AGGCGGTGGC ACCTCTGCAT TTATGAAGTT CGTCTTGGCA TTTATTTTT
 50 101 TGCTTCTTTC ATTGCRGAAC TCGCAAATGC ACTTCCCCTG CTTGTCGCAT
 151 TTCGCCCAA AAGCGCATGG CATTCCTTCC GGCAGATTAA CTTTTTCAA
 55 201 TTCACGGTTC TGAACCAATA ATAGATCGTG GCAATGTTTG TGCTGTTTGC

EP 1 892 297 A2

5
10
15
20

251 GATTTGCAAA CCAGCTGTAG CCACCATTGG ACTCAAAGGT GCGCACAACA
301 TGGCGCCGAA CTGTGAAAAA CAAATTAAGG CTNCTTTGTA ATAACGCTAG
351 TCTTGGTACG CCGTTAGAGG TCGATGTCGC GCCTCGCGAT TGCAAAGTGA
401 CTTGCACTTA TCAAGCTCCT GGAGAAAAAT GGGTGCAACG GGGGGATCAG
451 CGTTTGTACT TGCAAACATT TGTGGAGACG GTAAACCWGT ATTTGCGGGA
501 ACTCAGATGC TCCAGCGTGA AGCTCGTCTT AATAAAAGTT GTAAATTGGA
551 GTATNGATGA AGAACTGAAA TTCGAGGCAT TTAGAAACAC CACGAGAAGC
601 AGCGGAA

(3) INFORMATION FOR SEQ ID N°: 3

25
30

(i) SEQUENCE CHARACTERISTICS :
(a) LENGTH : 259 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

35
(iii) SEQUENCE DESCRIPTION : SEQ ID N°3 :

40
45
50
55

1 GATCCTACGC CTGAAAATGA GTGTCCATCG TCTTCACATA GTGCCACATT
51 GTAATTGGTA CAAGCTCCAT TTCGTCAGC GCTGTTTGTG ATGCTGCCGC
101 CTACTTTTCC TTCGGCACTC CATAAGTTAA ACCCTGTCAT TATAAGTGTG
151 ATTGCCGTAT CTCGGCTGAA TGGGTTCAT TTTTCTCTTA AATAATCAGC
201 TGTCCATATT CCATGTATTG TGTTCATGAG TATGTGATTC TCATCGTATA
251 TCTTCGCCT

(4) INFORMATION FOR SEQ ID N°: 4

5 (i) SEQUENCE CHARACTERISTICS :
(a) LENGTH : 170 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
10 (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

15 (iii) SEQUENCE DESCRIPTION : SEQ ID N°4 :

1 CCACTCGAAA ATGGAGGCTT TGAAACATTT CAGTAGCCCT GTGAACTCTG
20 51 GCTTTGCAAT GTAACAGCAA AAACACTTAC AGTTGAAGGG TGCAGTGTCA
101 GACGCTATGG AAGTTGCATC CACGAGCACR ACCCTGATTA CTACTGGCCA
25 151 CGTTGCTRTC CGGGTCTGCC

(5) INFORMATION FOR SEQ ID N°: 5

30 (i) SEQUENCE CHARACTERISTICS :
(a) LENGTH : 168 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
35 (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

40 (iii) SEQUENCE DESCRIPTION : SEQ ID N°5 :

1 GTATGTTACC ATGTCCAACC CGGTTATTAA ATACACCAAG TCGTAGGATT
45 51 TGTAGGCAGC TGCATTGCCC TTGACGTACT CTCTCAACGT TGCCAAGGAC
101 TCAGGCCCAT AAATGTAGTG GGGTTGACCT TGAACCTCTC GTAAAAAGCG
50 151 TTCTTTCTCC GTCGTGAG

55

(6) INFORMATION FOR SEQ ID N°: 6

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 247 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°6 :

1 CCGAAMATAA AACTTAGTCT CACCAATATA CGTTTGCCTA ACGCGAAGGA
51 ACAGGCACAA ATATACTACG AGCAGGACAT TCTCAAGAAC ACGGTTACAG
101 GAGTGTGGAC GAGAATTCAC TCAAAATATC CGTTCCTGA AGATGAGGGA
151 ATTAACTGT
201 AGCGGCACC ATAACAAACA GCGCTGAGAA AAGCGGAGCT TGTACGA

(7) INFORMATION FOR SEQ ID N°: 7

5 (i) SEQUENCE CHARACTERISTICS :

(a) LENGTH : 261 base pairs

(b) TYPE : nucleic acid

10 (c) STRANDEDNESS : single

(d) TOPOLOGY : linear

15 (ii) MOLECULE TYPE : cDNA

(iii) FEATURE :

(a) NAME/KEY : CDS

20 (b) LOCATION : 1 .. 261

(iv) SEQUENCE DESCRIPTION : SEQ ID N°7

25

1 ATG CCT TTT ATT TTC GTG GTG AGC TTA GTC ATT GTG GCC TGC ATC GTG
 GTA GAC ACA

30 Met Pro Phe Ile Phe Val Val Ser Leu Val Ile Val Ala Cys Ile Val
 Val Asp Thr

35 58 GCC AAC CAC AAA GGT AGA GGG CGG CCT GCG AAG TGT AAA CTT CCT CCG
 GAC GAC GGA

Ala Asn His Lys Gly Arg Gly Arg Pro Ala Lys Cys Lys Leu Pro Pro
 Asp Asp Gly

40

115 CCA TGC AGA GCA CGA ATT CCG AGT TAC TAC TTT GAT AGA AAA ACC AAA
 ACG TGC AAG

45 Pro Cys Arg Ala Arg Ile Pro Ser Tyr Tyr Phe Asp Arg Lys Thr Lys
 Thr Cys Lys

50 172 GAG TTT ATG TAT GGC GGA TGC GAA GGA AAC GAA AAC AAT TTT GAA AAC
 ATA ACT ACG

Glu Phe Met Tyr Gly Gly Cys Glu Gly Asn Glu Asn Asn Phe Glu Asn
 55 Ile Thr Thr

EP 1 892 297 A2

229 TGC CAA GAG GAA TGC AGA GCA AAA AAA GTC TAG
Cys Gln Glu Glu Cys Arg Ala Lys Lys Val End

5

(8) INFORMATION FOR SEQ ID N°: 8

10

(i) SEQUENCE CHARACTERISTICS :

(a) LENGTH : 292 base pairs

(b) TYPE : nucleic acid

15

(c) STRANDEDNESS : single

(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

20

(iii) SEQUENCE DESCRIPTION : SEQ ID N°8 :

1 CATCGMAGCC ATAGTATATT TTGCACTTGT CTTCCGTTTC GTCGTAGTAG

25

51 GACCGATTCC ACATTGTAGT ACACCAGTCA CTTATATCCT GCGGGCGGTG

101 CTTGCATTTG TCCTGAACAA ATCTTCCAGA GCGCTTGTGG CAGGCCTCCT

30

151 GGGAAATAGAA GCGGTTCTCT CCTCCGCATC TCCATTTGGA ATCATAGAAA

201 CATCTTTCAG TTTGAATATT GTAGCGATAA TAATCGGTAT CAGTTTCTTT

35

251 GCATGGTCTT GGGAGGGGTT TGGCGCAGGG GCCGTATTCA GG

(9) INFORMATION FOR SEQ ID N°: 9

40

(i) SEQUENCE CHARACTERISTICS :

(a) LENGTH : 270 base pairs

(b) TYPE : nucleic acid

45

(c) STRANDEDNESS : single

(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

50

(iii) SEQUENCE DESCRIPTION : SEQ ID N°9 :

1 GGTAATAGTT GTCAAATTC ATTAATGTAT CCTGAAATGT GACCATATCT

55

51 TTGTTTCCC TGTAATAATCT CATAAAAGGC TGTGTGTTTT CCTTAAGAAG

EP 1 892 297 A2

101 TGTAACAGCC ACGATGGTCA ATCTCACGGA TGGATGTGTG ACACTTTTAT
5 151 ATCTCAGGTT TGCCGACATT GCCATTACAG ATAAATAGTT GATAATTTCT
201 TTCTTGTTAT AGTTGTAAGC AGGGCATGTT GTTGCATCAA GCACCCACATG
10 251 CACTTCAGGC AATATGGTTT

(10) INFORMATION FOR SEQ ID N°: 10

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 316 base pairs
(b) TYPE : nucleic acid
20 (c) STRANDEDNESS : single
(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

25 (iii) SEQUENCE DESCRIPTION : SEQ ID N°10 :

1 AGAAAGCAGT CATATTGGCC ATCCACAGGT CACAATGGTT CTCTCCTTGA
30 51 CCTGGCATCG GGATTGGAAG TATGGTGCAG TTCACGTAGT TGGAATACAA
101 CACGAAATGT GTTCGTTGGT ACGCCAATAG GGGTTCTCGC AAAGAACATA
35 151 TCATTTGGAG GAAGGCGTAG TCCGTCGAGA TATCCCAAAA CTAGGGTTTC
201 ATTGCGTGCG AACCAACTGC CCCCACTTCT GTATGTGTAC TGTAAGGAGT
40 251 RGTGGAACGG YGTCCTCTTT GCCATAACCT TGAAGTTTTC ACACTGCAGA
301 GGATTACCTC TCAAAA

45 (11) INFORMATION FOR SEQ ID N°: 11

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 241 base pairs
50 (b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

55 (ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°11 :

5 1 AAGGTAGCAA GGGTGGTAGG CTTTCCTCAC AAAGAGTCTG GCTTCCGTGA
 51 TAACCATATC CATTGCTGAC CGTATACCCG TCATCCAACG TCAATTGTGT
 10 101 TACAAGGCAG ATAATGTCAA AATGGCTCTG GTCCTATAA TAGTCGGATA
 151 ATGTAGAAAT CGCTCCATGT GGCCAAATAG ATGTTCTCTT TTCATACTGT
 15 201 TTTAACTTTA ATTGTAGGTC CGCCTCGTTC TCGAGGTATG T

(12) INFORMATION FOR SEQ ID N°: 12

20 (i) SEQUENCE CHARACTERISTICS :
 (a) LENGTH : 636 base pairs
 (b) TYPE : nucleic acid
 25 (c) STRANDEDNESS : single
 (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

30 (iii) SEQUENCE DESCRIPTION : SEQ ID N°12 :

1 TTCCCNAAT TGGCCTTGGC ANNCTTGCAA GTCGACNCTA GAGGCTCCGA
 35 51 AGATGGACAG ATTGCGCATG AAATATTTGA AATCGAGCAG AATGGTGATT
 101 TTAGGAGCGA TTATATTGTG CCACCCAGTT TGAAAGTGCA- AGAACGCACA
 40 151 GTGGTTTACC GTAACAAGTA CACCAGAGTT CCTGTAAATT TTACCGTCGA
 201 AGTTGCCATG CTGATTGATA AGTATTTATA CWAGGAGTTC AAGAACGAGA
 45 251 GCCACATCGT ACCGTACCTG GCTATGATAC TGACTTTGAT AAATCTGAGG
 301 TATGCCGACA CACATGACCC GTACATCCAG TTTCTTCTCA CACAAGTGTT
 50 351 CGTGGGGA AW WCTGGCGATC ATATGGGCCA CATGCCCTTC CGACGAGCGT
 401 TCTTGTTT CAG GCGCCGGCAT TATGCGCAGT TTAGGCCCAA TMACACCTTC
 55 451 CACTTGTAAT TCTCCGTTGT TGGATAGTGT AAGTGAGGCC ATTGCATCAG

EP 1 892 297 A2

501 CATCCTGGAA GARGCCTTCC TCCAAGTAGG AACCGCCCAT TTAGGTTTGC

551 TTTCCCAATC CGCCAATTTA ANTTTTAAAA AAAATTCCCC CCCCAAAAAT

601 TAATTTTTTT TAAAGGTGGA TTGTGATTTC TCCGTT

(13) INFORMATION FOR SEQ ID N°: 13

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 432 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°13 :

1 GATCCCAAAA GTGCCCTGG ARGACGGTT ACATCATGAG CTACGTCATA
51 AACTTCAAAA ACCACTTCAA ATTTTCTCCC TGCTGTGTAG AATCAATTGC
101 ATTCGTGCGA CGAGAGCGGG ACTGCCTCTA CAAAGTCAAT GCCAAGGATG
151 CTGTAAAAAG CCTAATATCT CTGCCCGGAT TTAGGATATC GCCAACGAGT
201 TTCTGTCAAT TTATGCATCC GCTTTACCGC GGTGTCCATA GCGATAAGAA
251 AGCAGGTCTG TCCGATTGCG TACAGACGTG TAGAACGGCC AAAAATCGAC
301 GAGGAGGCTA CCATTTCATG ATTCACGCGG CACTTGACGG GGTTCCTTGC
351 GACAAGAGAA ACCCCAAGAA GGCCTGCATA AACGGGAAAT GCACCCTCCT
401 TAAGAGCATG CCCACAGAA CGTACCGGGA AT

(14) INFORMATION FOR SEQ ID N°: 14

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 466 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°14 :

1 AGGGCGTTCT TTGCTTYACA GGAACRGCA TATGGGCCAC GTGACCTTCC
 51 AATGACCGCT CCAAATCTGG CATAGGTTGA AYTGGCAAGT CGTGGCGCAG
 101 CAGGCCTYCC ACATTCACTC CATCCTCGTC TTTTAGGATG ACTGCCGCCA
 151 TTTGTTTTGT ATCGTGGTAC AGGTGTTTGT TATGGTCCGA GCCGTGCACA
 201 TAAGTATTGA CCAACGATCG GCCGAATGAT TACGGCTCAC CAAACACATC
 251 AAATACCCCC GTCAAGTCAA GAGCTGGAAG CACAAAGCAT AGTATGTACA
 301 AGATACCCTT GGAAATCTTT CCCGAAGTTC ACCTTGTGGT GGACAGCACA
 351 TTTGCCAAAG CTTTTAAATT TGACGTGTAC AAAGTAACGC GTTACTTGGC
 401 AGTGCTTACA AATGCGGCTA ATCTTAGGTA TGCCAGCTTC GTATTTCCAA
 451 AAGTACAGCT CAGGAT

(15) INFORMATION FOR SEQ ID N°: 15

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 377 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°15 :

1 CTCGTCCACA CATTCTCCTA AAATGCAAGC CTTTTTTTTT CCACAAGGTG

EP 1 892 297 A2

51 TACCGTGGAC TACTACTGAGT CTCCAATAAA TATGTTTTCC GGTGCAATTT
101 ACCTTGCAAGT CTTTGACGCC GTATGTAGGG TCAGCGTGCA TGCCTTCGTC
5
151 GTACATATAC ACCCTCTGAC AGTAGTTGCT CAGTGTGTC ATCCTACCAG
201 GAAGCTTAGA CGAACGTTTT ATTGTTTTTG TCGTGTATCG TTCTCTAAGG
10
251 CATTGAATT CCGGACGGTT GTAGAGGTTT CTGACTTCTC GCTGGCAGCA
301 ATAAGAGAAC TGATACTGGC GCTCGTCTTG CATCTTGIAA CTCATGAGGT
15
351 ATCCGTCATC CCATGGGCAG TCCGCAG

20 (16) INFORMATION FOR SEQ ID N°: 16

(i) SEQUENCE CHARACTERISTICS :

- 25 (a) LENGTH : 1670 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

30 (ii) MOLECULE TYPE : cDNA

(iii) FEATURE :

- (a) NAME/KEY : CDS
(b) LOCATION : 54 .. 1520

35 (iv) SEQUENCE DESCRIPTION : SEQ ID N°16 :

1 AAGGAAGAAG TTAGGCGTAG GCTTTGGGAA ACCGGTCATC CTCGAAACCA GAG
40
54 ATG TCG GGA CTC AGC CTG AAA TTG TGG ATT GTA GCG TTC TTT TCT
Met Ser Gly Leu Ser Leu Lys Leu Trp Ile Val Ala Phe Phe Ser
45
99 TTC TGC TTG GCC GAG AAA GAG CAT GGG ATC GTG TAC CCC AGG ATG CTT
Phe Cys Leu Ala Glu Lys Glu His Gly Ile Val Tyr Pro Arg Met Leu
147 GAA AGC AGA GCA GCA ACT GGA GAG AGA ATG CTT AAA ATC AAC GAT GAC
50
Glu Ser Arg Ala Ala Thr Gly Glu Arg Met Leu Lys Ile Asn Asp Asp
195 CTG ACG TTG ACG CTG CAG AAG AGT AAG GTC TTC GCT GAC GAC TTT CTC
Leu Thr Leu Thr Leu Gln Lys Ser Lys Val Phe Ala Asp Asp Phe Leu
55
243 TTC AGC ACG ACC GAC GGA ATT GAA CCT ATT GAT TAC TAC ATC AAA GCC

EP 1 892 297 A2

Phe Ser Thr Thr Asp Gly Ile Glu Pro Ile Asp Tyr Tyr Ile Lys Ala
 5 291 GAA GAC GCT GAA CGT GAC ATC TAC CAC GAC GCA ACT CAC ATG GCA TCA
 Glu Asp Ala Glu Arg Asp Ile Tyr His Asp Ala Thr His Met Ala Ser
 339 GTA AGG GTA ACG GAC GAT GAT GGC GTG GAA GTG GAA GGA ATT CTT GGA
 10 Val Arg Val Thr Asp Asp Asp Gly Val Glu Val Glu Gly Ile Leu Gly
 387 GAG AGG CTT CGT GTT AAA CCT TTG CCG GCA ATG GCC CGC AGC AGC GAT
 Glu Arg Leu Arg Val Lys Pro Leu Pro Ala Met Ala Arg Ser Ser Asp
 15 435 GGC CTC AGA CCG CAT ATG TTG TAC GAA GTC GAC GCA CAC GAA AAC GGC
 Gly Leu Arg Pro His Met Leu Tyr Glu Val Asp Ala His Glu Asn Gly
 483 CGG CCA CAT GAT TAT GGT TCA CCG AAC ACA ACA AAT ACC CCC GTA GAG
 20 Arg Pro His Asp Tyr Gly Ser Pro Asn Thr Thr Asn Thr Pro Val Glu
 531 AGA AGA GCT GGA GGC ACA GAA CCC CAG ATG TAC AAG ATA CCA GCG GAA
 Arg Arg Ala Gly Gly Thr Glu Pro Gln Met Tyr Lys Ile Pro Ala Glu
 25 579 ATC TAT CCC GAA GTT TAC CTT GTG GCG GAT AGT GCC TTT GCC AAA GAA
 Ile Tyr Pro Glu Val Tyr Leu Val Ala Asp Ser Ala Phe Ala Lys Glu
 627 TTT AAC TTT GAT GTG AAC GCC GTT ACG CGT TAC TTC GCA GTG CTT ACA
 30 Phe Asn Phe Asp Val Asn Ala Val Thr Arg Tyr Phe Ala Val Leu Thr
 675 AAT GCG GCT AAT CTT AGG TAT GAA AGC TTC AAA TCT CCA AAG GTA CAG
 35 Asn Ala Ala Asn Leu Arg Tyr Glu Ser Phe Lys Ser Pro Lys Val Gln
 723 CTC AGG ATC GTT GGC ATA ACG ATG AAC AAA AAC CCA GCA GAC GAG CCA
 Leu Arg Ile Val Gly Ile Thr Met Asn Lys Asn Pro Ala Asp Glu Pro
 40 771 TAC ATT CAC AAT ATA CGG GGA TAT GAG CAG TAC GGG AAT ATT TTG TTT
 Tyr Ile His Asn Ile Arg Gly Tyr Glu Gln Tyr Arg Asn Ile Leu Phe
 819 AAG GAA ACA CTG GAG GAT TTC AAC ACT CAG ATG AAG TCA AAA CAT TTT
 45 Lys Glu Thr Leu Glu Asp Phe Asn Thr Gln Met Lys Ser Lys His Phe
 867 TAT CGT ACT GCC GAT ATC GTG TTT CTC GTG ACA GCA AAA AAT ATG TCC
 Tyr Arg Thr Ala Asp Ile Val Phe Leu Val Thr Ala Lys Asn Met Ser
 50 915 GAA TGG GTT GGT AGC ACA CTA CAA TCA TGG ACT GGC GGG TAC GCT TAC
 Glu Trp Val Gly Ser Thr Leu Gln Ser Trp Thr Gly Gly Tyr Ala Tyr
 963 GTA GGA ACA GCG TGT TCC GAA TGG AAA GTA GGA ATG TGT GAA GAC CGA
 55 Val Gly Thr Ala Cys Ser Glu Trp Lys Val Gly Met Cys Glu Asp Arg

EP 1 892 297 A2

5
1011 CCG ACA AGC TAT TAC GGA GCT TAC GTT TTC GCC CAT GAG CTG GCG CAT
Pro Thr Ser Tyr Tyr Gly Ala Tyr Val Phe Ala His Glu Leu Ala His

1059 AAT TTG GGT TGT CAA CAC GAT GGA GAT GGT GCC AAT AGC TGG GTG AAA
Asn Leu Gly Cys Gln His Asp Gly Asp Gly Ala Asn Ser Trp Val Lys

10
1107 GGG CAC ATC GGA TCT GCG GAC TGC CCA TGG GAT GAC GGA TAC CTT ATG
Gly His Ile Gly Ser Ala Asp Cys Pro Trp Asp Asp Gly Tyr Leu Met

15
1155 AGC TAC AAG ATG GAA GAC GAG CGC CAG TAT AAG TTT TCT CCC TAC TGC
Ser Tyr Lys Met Glu Asp Glu Arg Gln Tyr Lys Phe Ser Pro Tyr Cys

1203 CAG AGA GAA GTC AGG AAC CTC TAC AGG CGT CCG GAA TTC AAA TGC CTC
Gln Arg Glu Val Arg Asn Leu Tyr Arg Arg Pro Glu Phe Lys Cys Leu

20
1251 ACT GAA CGA AAA GCG AAA AAA ACA ATC CGC TCG TCT AAG CTA CCT GGT
Thr Glu Arg Lys Ala Lys Lys Thr Ile Arg Ser Ser Lys Leu Pro Gly

25
1299 GTG ATG ACA TCA TCG AGC AAC TAT TGC CGG AGG GTG TAC ATG TAC GAA
Val Met Thr Ser Ser Ser Asn Tyr Cys Arg Arg Val Tyr Met Tyr Glu

30
1347 AAA GGC ATG CAC GCC GAC GAG GCA TaT GGC GTC AAG GAC TGC AGG GTA
Lys Gly Met His Ala Asp Glu Ala Tyr Gly Val Lys Asp Cys Arg Val

35
1395 AAA TGC ACC ACC ACA TCA AGA ATG TAT TGG CTA CTC GGT GTA GTC GAC
Lys Cys Thr Thr Thr Ser Arg Met Tyr Trp Leu Leu Gly Val Val Asp

1443 GGT ACA CCT TGC GGA AAT GGA AAG GCT TGC ATT CTT GGG AAA TGC AGG
Gly Thr Pro Cys Gly Asn Gly Lys Ala Cys Ile Leu Gly Lys Cys Arg

40
1491 AAC AAA ATC AAA ATA AGC AAG AAG GAC TGA GAGGTTGATA ATATCAAATT
Asn Lys Ile Lys Ile Ser Lys Lys Asp End

1541 AATCATGATA TTTCAACCAC ATGACTTCGT GCTCAACTGG TAGCCCCAAA TAAATTTTAA

45
1601 AAAAAATCCC AATATGCGTG GTAGAAAAAG CAGCAAACAA TAAAACTTCT AAAAAATGTCT

50
1661 TGCAAAAAATG

55 (17) INFORMATION FOR SEQ ID N°: 17

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 158 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°17 :

1 CACCAGTGAT GCTTATTGTT GCACTGCACT TGTTGATAAT ATCCGGTCGT
51 CGAATTGCAC TTCGGA ACTT CCACTCCAAC TTGGCGAGCC GTGGATTTTG
101 ACTTCTCGTG ATGCTCCACC AGACAGTTGC AGGACTTCAG CTGCCTAGAT
151 GGAGCCTT

(18) INFORMATION FOR SEQ ID N°: 18

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 146 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°18 :

1 CTGTTGTTGA ACTGAAATAA ATAACAAAAA AATCATAAAG NTGGAGGAAA
51 GATGATCGAN TCCCCGCCCC TTGACAATCG TCCGATAAAA ACCAACTATA
101 TTCNGTCCTT TTTACAAACA ATTCCAANTG TCTGACCGAA CCGCGA

(19) INFORMATION FOR SEQ ID N°: 19

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 140 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°19 :

1 CTNGGACGAN GTCCTATGAC TTGCGCTTAN GTTTCTTAGT CTCCTTCGGT
51 TTCTTCTTTT TTTGCTTCGG TTTTTCGGTG GGCGCAGGTG TATAGTCATC
101 AGTGTCGGTG GGCCCATCCG AATGAGTTGT CAAATGACAT

(20) INFORMATION FOR SEQ ID N°: 20

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 143 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°20 :

1 TGCCGAAAAA TAAGGATGAT TTGACGTTGA CTCTGCAGAA GAGTAAGGTT
51 TTCACCGACA GTTTTCTGTT TAGCAGGACG AAGGATAACG AGCCTATCGA
101 TTACTACGTG AGAGCCGAAG ATGCCGAACG AGACATATAT CAC

(21) INFORMATION FOR SEQ ID N°: 21

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 140 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°21 :

5
1 TGTTCCTACA GACTCGACGT TTCGAGCTTG CTCGCCATTT MAAGACAACG
51 CACTCACAGA ATATTTAAGT GCGTTCGTGA WAGCTGTGGG CTTACGATTG
10
101 CAGGCGCTTC ANTCACCAGC TGTGATATTA MAGTTCCTAG

15
(22) INFORMATION FOR SEQ ID N°: 22

(i) SEQUENCE CHARACTERISTICS :

- 20
(a) LENGTH : 144 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

25
(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°22 :

30
1 TCACGATAGT TGAAACGTTG AAACCTGAAA TACTGCCACA GTCGTTGGAT
51 GCTTCAGAAC TGCTAAGAAC TTCACACTTT GCAAGAAGTW CCAAAATGAA
35
101 AGCCGCGATG ACCGATGATT TAGCTTCCAT CTTCTATCAC TTGA

(23) INFORMATION FOR SEQ ID N°: 23

40
(i) SEQUENCE CHARACTERISTICS :

- 45
(a) LENGTH : 95 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

50
(iii) SEQUENCE DESCRIPTION : SEQ ID N°23 :

55
1 GACCACCCCG TCCGAACTTG CTAAAKCAAG CAATGGAGTG AGGTGTTCTA
51 TCGGGGTTGA TTACACCAAT GGCCTGCGT GGTGCGTGGT GATT

EP 1 892 297 A2

509 TTT GCA GGA AAT GCC GAC AGG GTC CGT CTG GAG GTC AAT GCC TGG GTT
Phe Ala Gly Asn Ala Asp Arg Val Arg Leu Glu Val Asn Ala Trp Val

5
557 GAG GAA GTC ACC AGG TCA AAG ATC AGG GAC CTG CTC GCA CCT GGA ACT
Glu Glu Val Thr Arg Ser Lys Ile Arg Asp Leu Leu Ala Pro Gly Thr

10
605 GTT GAT TCA TCG ACA TCA CTT ATA TTA GTG AAT GCC ATT TAC TTC AAA
Val Asp Ser Ser Thr Ser Leu Ile Leu Val Asn Ala Ile Tyr Phe Lys

15
653 GGT CTG TGG GAT TCT CAG TTC AAG CCT AGT GCT ACG AAG CCG GGA GAT
Gly Leu Trp Asp Ser Gln Phe Lys Pro Ser Ala Thr Lys Pro Gly Asp

20
701 TTT CAC TTG ACA CCA CAG ACC TCA AAG AAA GTG GAC ATG ATG CAC CAG
Phe His Leu Thr Pro Gln Thr Ser Lys Lys Val Asp Met Met His Gln

25
749 GAA GGG GAC TTC AAG ATG GGT CAC TGC AGC GAC CTC AAG GTC ACT GCG
Glu Gly Asp Phe Lys Met Gly His Cys Ser Asp Leu Lys Val Thr Ala

30
797 CTT GAG ATA CCC TAC AAA GGC AAC AAG ACG TCG ATG GTC ATT CTC CTG
Leu Glu Ile Pro Tyr Lys Gly Asn Lys Thr Ser Met Val Ile Leu Leu

35
845 CCC GAA GAT GTA GAG GGA CTC TCA GTC CTG GAG GAA CAC TTG ACC GCT
Pro Glu Asp Val Glu Gly Leu Ser Val Leu Glu Glu His Leu Thr Ala

40
893 CCG AAA CTG TCG GCT CTG CTC GGC GGC ATG TAT GCG ACG TCC GAT GTC
Pro Lys Leu Ser Ala Leu Leu Gly Gly Met Tyr Ala Thr Ser Asp Val

45
941 AAC TTG CGC TTG CCG AAG TTC AAA CTA GAG CAG TCC ATA GGT TTG AAG
Asn Leu Arg Leu Pro Lys Phe Lys Leu Glu Gln Ser Ile Gly Leu Lys

50
989 GAT GTA CTG ATG GCG ATG GGA GTC AAG GAT TTC TTC ACG TCC CTT GCA
Asp Val Leu Met Ala Met Gly Val Lys Asp Phe Phe Thr Ser Leu Ala

55
1037 GAT CTT TCT GGC ATC AGC GCT GCG GGG AAT CTG TGC GCT TCG GAT GTC
Asp Leu Ser Gly Ile Ser Ala Ala Gly Asn Leu Cys Ala Ser Asp Val

1085 ATC CAC AAG GCT TTT GTG GAA GTT AAT GAG GAG GGC ACA GAG GCT GCA
Ile His Lys Ala Phe Val Glu Val Asn Glu Glu Gly Thr Glu Ala Ala

1133 GCT GCC ACT GCC ATA CCC ATT ATG TTG ATG TGT GCG AGA TTT CCA CAG
Ala Ala Thr Ala Ile Pro Ile Met Leu Met Cys Ala Arg Phe Pro Gln

1181 GTG GTG AAC TTT TTC GTT GAC CGC CCA TTC ATG TTC TTG ATC CAC AGC
Val Val Asn Phe Phe Val Asp Arg Pro Phe Met Phe Leu Ile His Ser

1229 CAT GAT CCA GAT GTT GTT CTC TTC ATG GGA TCC ATC CGT GAG CTC TAA

EP 1 892 297 A2

His Asp Pro Asp Val Val Leu Phe Met Gly Ser Ile Arg Glu Leu End

1277 AAAGCATATT CTTAACGGCG GCCAATCAGT CTGTGGAGTT ATCTCTTAGT CACTAATGTG

1337 TAACAATTCT GCAATATTCA GCTTGTGTAT TTCAGTAACT TGCTAGATCT TTGTGTTGTT

1397 GATGTTAGGC TTCTTGCG

(25) INFORMATION FOR SEQ ID N°: 25

(i) SEQUENCE CHARACTERISTICS :

(a) LENGTH : 200 base pairs

(b) TYPE : nucleic acid

(c) STRANDEDNESS : single

(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°25 :

1 ACCGTAACCA AAATTGTTTC TTTCCAGAAG AATGGTTCAA ACTTTTCAAA

51 CAGATTTGGG AAACTCTTCT TGCAC TT TA AAATCCAATC TACAATCTTT

101 CCTCGCACTT CTGAATTGCA TTCCAGTTA GCTTCCAAGC AAACCTCTTT

151 TGGCAACTCC AGCCGTACTC CATTTCGGCA TACCACAGTG CATGCAC TTG

(26) INFORMATION FOR SEQ ID N°: 26

(i) SEQUENCE CHARACTERISTICS :

(a) LENGTH : 241 base pairs

(b) TYPE : nucleic acid

(c) STRANDEDNESS : single

(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°26 :

1 CGTATTCTTT GAAGATTTGT ATACGAAACA TAAATTCGTC ATGCATACTT

51 TTGATGGTTA CACGACATGC GAAGCTGCCG ACAAAGAAGA CTGGGAAGAT

EP 1 892 297 A2

101 AAGAAGCACC TAGTTACGGT AGTGCGTGGA CCGGATAAAC GAAAGTACAC
5
151 GTTTCTACGC AACATTCTCA CCTTACAACG GAGAGTGAGA GTTAGCAAAA
201 CAATGATTGA GCTCGTACGG AACATGTCCT GTAGGACATT T

10
(27) INFORMATION FOR SEQ ID N°: 27

15 (i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 313 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
20 (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

25 (iii) SEQUENCE DESCRIPTION : SEQ ID N°27 :

1 AAGCANCCGG ACTACCTGCT TGAAAACGTT GTACGGGCAA ACTTGGACGG
51 AAAACTCCCA GATGCTACTC CAGTTCCTCC CGGAAGCTAC ACGTACGCTG
30 101 AGAATGATAA CTTACCTGC TATTCCAGAA GTACACCGTT TCCGGATGGG
151 GTGAATGTTG TATAACGGCT GCTGGGTGCG GAAGACTATG ATGGATTACG
35 201 CAAAAAAGTT CTAAACGAGT TGTTTCCCAT CCCGGAAAGT CTGCTGTATG
251 CTGACATGAT GCGACTTGTG GCTAAGAAAAG ACAGAGTTGA TCACACTAGT
40 301 GGATGACCTG GGA

45 (28) INFORMATION FOR SEQ ID N°: 28

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 2417 base pairs
50 (b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

55 (ii) MOLECULE TYPE : cDNA

EP 1 892 297 A2

(iii) FEATURE :

- (a) NAME/KEY : CDS
- (b) LOCATION : 218 .. 1495

(iv) SEQUENCE DESCRIPTION : SEQ ID N°28 :

5

10

15

20

25

30

35

40

45

50

55

```

1 GTCGTAGTCG TAGTCGTAGT CAGTTGCGCA TCGCGGGGCG TTCCTGTCT TTCTTGCCCTT
61 TCTGCAGTCG TTCACCAACA TGTGGATACA GCTCCGGAGA TTTGTAAACA AATACTGCAC
121 TTTTAAGCAA GACTTGATAT TTAGATCGAT ATCCTCCTGT TGTCCGTCTT GATTAATCGG
181 CTCTTTAGGG TTTTLAGAAT AGGCTTTTCG GTACGAG ATG CCC AAA GGA AAG AGG
Met Pro Lys Gly Lys Arg
236 GGA CCC AAA GCA GGT GGC GCC GCG CGC GGT GGC CGG TGC GAG GCC AGC
Gly Pro Lys Ala Gly Gly Ala Ala Arg Gly Gly Arg Cys Glu Ala Ser
284 CTG GCT CCG TCg TCC AGC GAC GAG GAG TCC AAC GCA GAC ACG GCG AGC
Leu Ala Pro Ser Ser Ser Asp Glu Glu Ser Asn Ala Asp Thr Ala Ser
332 GTG CTG AGC TGC GCC TCG GAG TCT CGC TGT GGC AGT GAC GGC ACC GTT
Val Leu Ser Cys Ala Ser Glu Ser Arg Cys Gly Ser Asp Gly Thr Val
380 GGA GAC CCA GAA GCG GAG GAG GCT GTG CTG CAT GAC GAC TTT GAA GAC
Gly Asp Pro Glu Ala Glu Glu Ala Val Leu His Asp Asp Phe Glu Asp
428 AAA CTC AAG GAG GCC ATC GAC GGA GCT TCG CAG AAG AGT GCC AAA GGA
Lys Leu Lys Glu Ala Ile Asp Gly Ala Ser Gln Lys Ser Ala Lys Gly
476 CGG CTG TCG TGC CTG GAG GCG ATT CGC AAG GCC TTT TcC ACC AAA TAC
Arg Leu Ser Cys Leu Glu Ala Ile Arg Lys Ala Phe Ser Thr Lys Tyr
524 CTG TAC GAC TTC CTC ATG GAC AGA CCG AGC ACG GTG TGC GAC CTG GTG
Leu Tyr Asp Phe Leu Met Asp Arg Pro Ser Thr Val Cys Asp Leu Val
572 GAG CGT GGG GTG CGC AAG GGC CGA GGG GAG GAG GCG GCC CTG TGC GCC
Glu Arg Gly Val Arg Lys Gly Arg Gly Glu Glu Ala Ala Leu Cys Ala
620 ACT CTC GGG GCC CTG GCC TGC GTC CAG CTC GGG GTC GGG GCC GAG GCG
Thr Leu Gly Ala Leu Ala Cys Val Gln Leu Gly Val Gly Ala Glu Ala
668 GAC GCC CTG TTC GAC GCC CTG CGC CAG CCG CTC TGC ACT TTG CTG CTT
Asp Ala Leu Phe Asp Ala Leu Arg Gln Pro Leu Cys Thr Leu Leu Leu
716 GAC GGG GCC CAG GGG CCC TCC CCC AGG GCC AGG TGT GCC ACT GCC CTC

```

EP 1 892 297 A2

Asp Gly Ala Gln Gly Pro Ser Pro Arg Ala Arg Cys Ala Thr Ala Leu

5 764 GGC CTC TGC TGc TTC GTG GTG GAC TCG GAC AAC CAG CTG GTG CTG CAG
Gly Leu Cys Cys Phe Val Val Asp Ser Asp Asn Gln Leu Val Leu Gln

10 812 CCG TGC ATG GAG GTG CTC TGG CAG GTG GTG GGT GCC AAG GCG GGC CCC
Pro Cys Met Glu Val Leu Trp Gln Val Val Gly Ala Lys Ala Gly Pro

15 860 GGC TCT CCG GTG CTC CAG GCA GCG GCC CTG CTC GCC TGG GGC CTC CTG
Gly Ser Pro Val Leu Gln Ala Ala Ala Leu Leu Ala Trp Gly Leu Leu

20 908 CTC AGC GTG GCT CCC GTC GAC CGc CTG CTG GCG CTc ACG CGC ACG CAC
Leu Ser Val Ala Pro Val Asp Arg Leu Leu Ala Leu Thr Arg Thr His

25 956 CTG CCC CGG CTG CAG GAG CTG CTG GAG AGC CCC GAC CTG GAC CTG CGC
Leu Pro Arg Leu Gln Glu Leu Leu Glu Ser Pro Asp Leu Asp Leu Arg

30 1004 ATT GCG GCC GGG GAG GTG ATC GCC GTC ATG TAC GAG GGG GCC AGG GAC
Ile Ala Ala Gly Glu Val Ile Ala Val Met Tyr Glu Gly Ala Arg Asp

35 1052 TAC GAC GAG GAC TTT GAG GAG CCC TCG GAG TCC CTG TGT GCC CAG CTG
Tyr Asp Glu Asp Phe Glu Glu Pro Ser Glu Ser Leu Cys Ala Gln Leu

40 1100 CGC CAG CTG GCC ACG GAC AGC CAG AAG TTT CGG GCC AAG AAG GAG CGG
Arg Gln Leu Ala Thr Asp Ser Gln Lys Phe Arg Ala Lys Lys Glu Arg

45 1148 CGC CAG CAG CGC TCC ACC TTC AGG GAC GTC TAC CGG GCC GTC AGG GAG
Arg Gln Gln Arg Ser Thr Phe Arg Asp Val Tyr Arg Ala Val Arg Glu

50 1196 GGG GCC TCT CCC GAC GTG AGC GTC AAG TTT GGC CGG GAA GTC CTG GAA
Gly Ala Ser Pro Asp Val Ser Val Lys Phe Gly Arg Glu Val Leu Glu

55 1244 CTG GAC ACC TGG AGT CGC AAG CTG CAG TAC GAC GCT TTC TGC CAG CTG
Leu Asp Thr Trp Ser Arg Lys Leu Gln Tyr Asp Ala Phe Cys Gln Leu

1292 CTG GGC TCC GGC ATG AAC CTG CAC CTG GCC GTG AAC GAG CTG CTG AGG
Leu Gly Ser Gly Met Asn Leu His Leu Ala Val Asn Glu Leu Leu Arg

1340 GAC ATC TTT GAA CTG GGG CAG GTG CTG GCA ACC GAG GAC CAC ATT ATC
Asp Ile Phe Glu Leu Gly Gln Val Leu Ala Thr Glu Asp His Ile Ile

1388 TCC AAG ATC ACC AAG TTC GAA AGG CAC ATG GTG AAC ATG GCC AGC TGC
Ser Lys Ile Thr Lys Phe Glu Arg His Met Val Asn Met Ala Ser Cys

1436 CGG GCC CGC ACC AAG ACA CGC AAC CGG CTG AGG GAC AAG CGC GCC GAC
Arg Ala Arg Thr Lys Thr Arg Asn Arg Leu Arg Asp Lys Arg Ala Asp

EP 1 892 297 A2

1484 GTG GTC GCC TGA ACCTGCGGAG GGATGCTTAG CTATGCACTC GCCGGCCTAC
Val Val Ala End

5

1536 CCTGGCGGGA CTCGATGCCA CTCACGAGTC GGCGCTCGCA AATTCGCCGC CCATCGTTAC

1596 GCAATGGGAG ACAAAGCTGC TTTTGGCATT ACCGTTTGAG GTCGGCTCCA ACCCATAGAT

10

1656 GAATTTCTTT TTTGTGGCCG TTTCTGGGTT ACATGTTTTG GGGGAAGGGA GTGGAAGTGT

1716 CCGGTTCTTT GGCACACGTC AGGTTGCTCT TGATGCGCGA CGTGCTTGTA TTTGGGtACT

15

1776 GCCGACACCA AGCGTTTCGG CGATTCCTGG AAAAGAGTGC CTCTCGCTCG ACGTTTGGTT

1836 GTTTTCTGCG TGGTCCGTCG TCGACCTTCG TTCGTCCAAA GACGCCGTCC GGTTCAtAC

20

1896 TCCCCCGCGC ACACATATCG AGGCCAATTA AATTGCTAAG GGTGCCGTTG TCGTGATCT

1956 GGCAGGCTCA GAAGTGGCTT ATTTGTCTTT TAATTTTGCC GATGCACGCA AAAATTGTCA

25

2016 TTTCTTGAAA GTTTCTCTTT TATTGCGTAC ACAATTCAAC TTTTATGTAA TTTCTGATGG

2076 TCTGTTTTAC GTGTCCGTGT GTAAAACGTA ACTTTGGAAG AATTTTTATG CACACTGAAC

30

2136 AAACGCTCGG TCCTGGGGTT GAAAGTGCTC GGTGTGTGCA TGAGCTAAAG TGCAACTGCT

2196 TTGTTCCGAA GTTTTTCTAG TCGCCGAAAT GTACCATTGT GGACCTTGTT GCGAGAGACC

35

2256 TTGGTCTTCT GGGGGAGCTG CTGTAGCGTG GCAAGCCACT ATTTTGGGAG CGACATTGCA

2316 GAGAAAATCG GCTTTTAGAA AGGCACCTGC GCGGCGAGTG GACGTTTTTT CGTATATACT

40

2376 GCGAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AA

(29) INFORMATION FOR SEQ ID N°: 29

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 933 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) FEATURE :

EP 1 892 297 A2

(a) NAME/KEY : CDS
 (b) LOCATION : 32 .. 853

5 (iv) SEQUENCE DESCRIPTION : SEQ ID N°29 :

1 GATTGGGAAC CTCCTATTCC TCACTTGAAA C ATG GCT GGA CTC CGC TCC
 Met Ala Gly Leu Arg Ser

10
 50 TGC ATC CTC CTG GCT CTT GCC ACT AGT GCC TTC GCC GGC TAC CTT CAC
 Cys Ile Leu Leu Ala Leu Ala Thr Ser Ala Phe Ala Gly Tyr Leu His

15
 98 GGT GGC CTT ACC CAC GGC GCT GGG TAC GGT TAC GGT GTC GGC TAC GGT
 Gly Gly Leu Thr His Gly Ala Gly Tyr Gly Tyr Gly Val Gly Tyr Gly

20
 146 TCC GGC CTT GGC TAT GGC CTT GGC TAC GGT TCC GGC CTT GGC TAT GGA
 Ser Gly Leu Gly Tyr Gly Leu Gly Tyr Gly Ser Gly Leu Gly Tyr Gly

25
 194 CAT GCT GTT GGC CTT GGA CAC GGC TTT GGC TAT TCT GGT CTG ACC GGC
 His Ala Val Gly Leu Gly His Gly Phe Gly Tyr Ser Gly Leu Thr Gly

30
 242 TAC AGT GTG GCT GCC CCA GCT AGC TAC GCC GTT GCT GCT CCA GCC GTC
 Tyr Ser Val Ala Ala Pro Ala Ser Tyr Ala Val Ala Ala Pro Ala Val

35
 290 AGC CGC ACC GTT TCC ACT TAC CAC GCT GCT CCA GCT GTG GCC ACC TAC
 Ser Arg Thr Val Ser Thr Tyr His Ala Ala Pro Ala Val Ala Thr Tyr

40
 338 GCC GCT GCT CCT GTC GCC ACC TAT GGT GTT GCT CCA GCT GTC ACT AGG
 Ala Ala Ala Pro Val Ala Thr Tyr Ala Val Ala Pro Ala Val Thr Arg

45
 386 GTT TCC CCC GTT CGC GCC GCC CCA GCT GTG GCC ACG TAC GCC GCC GCT
 Val Ser Pro Val Arg Ala Ala Pro Ala Val Ala Thr Tyr Ala Ala Ala

50
 434 CCA GTG GCC ACC TAC GCC GCT GCT CCA GCT GTG ACC AGG GTG TCC ACC
 Pro Val Ala Thr Tyr Ala Ala Ala Pro Ala Val Thr Arg Val Ser Thr

55
 482 ATT CAC GCT GCC CCG GCT GTG GCC AAT TAC GCC GTC GCT CCA GTC GCC
 Ile His Ala Ala Pro Ala Val Ala Asn Tyr Ala Val Ala Pro Val Ala

530 ACC TAT GCC GCT GCT CCA GCT GTG ACC AGG GTG TCC ACC ATC CAC GCC
 Thr Tyr Ala Ala Ala Pro Ala Val Thr Arg Val Ser Thr Ile His Ala

578 GCT CCA GCC GTG GCT AGC TAC CAG ACC TAC CAC GCT CCA GCT GTC GCC
 Ala Pro Ala Val Ala Ser Tyr Gln Thr Tyr His Ala Pro Ala Val Ala

626 ACT GTG GCT CAT GCT CCA GCT GTG GCC AGC TAC CAG ACC TAC CAC GCT
 Thr Val Ala His Ala Pro Ala Val Ala Ser Tyr Gln Thr Tyr His Ala

EP 1 892 297 A2

5
674 GCC CCA GCC GTG GCT ACC TAC GCC CAT GCC GCT CCC GTC TAC GGC TAT
Ala Pro Ala Val Ala Thr Tyr Ala His Ala Ala Pro Val Tyr Gly Tyr

10
722 GGT GTC GGT ACC CTC GGA TAT GGT GTC GGC CAC TAC GGC TAC GGA CAC
Gly Val Gly Thr Leu Gly Tyr Gly Val Gly His Tyr Gly Tyr Gly His

15
770 GGT CTT GGC AGC TAC GGC CTG AAC TAC GGT TAC GGC CTC GGC ACC TAC
Gly Leu Gly Ser Tyr Gly Leu Asn Tyr Gly Tyr Gly Leu Gly Thr Tyr

818 GGT GAC TAC ACC ACC CTT CTC CGC AAG AAG AAG TAA ATGGCA CATCTCAAGA
15 Gly Asp Tyr Thr Thr Leu Leu Arg Lys Lys Lys End

870 GAGCCCAT TG GACTGCCATC GACATTCTTC TTCAATAAAA GAGCCCGAAG ATGGCATTAT

20
930 TTTT

Bibliography.

- 25
[0094]
- 30 Ganapamo et al, 1997 : Identification of an Ixodes ricinus salivary gland fraction through its ability to stimulate CD4 T cells present in BALB/c mice lymph nodes draining the tick fixation site. Ganapamo-F; Rutti-B; Brossard-M. Parasitology; 1997 Jul; 115 (Pt 1): 91-6
- 35 Ganapamo et al, 1995 : In vitro production of interleukin-4 and interferon-gamma by lymph node cells from BALB/c mice infested with nymphal Ixodes ricinus ticks. Ganapamo-F; Rutti-B; Brossard-M. Immunology; 1995 May; 85(1): 120-4
- 40 Ganapamo et al, 1996 : Immunosuppression and cytokine production in mice infested with Ixodes ricinus ticks: a possible role of laminin and interleukin-10 on the in vitro responsiveness of lymphocytes to mitogens. Ganapamo-F; Rutti-B; Brossard-M. Immunology; 1996 Feb; 87(2): 259-63
- 45 Wikel et al, 1996 : Host immunity to ticks. Wikel-SK. Annu-Rev-Entomol.; 1996; 41: 1-22
- Wikel and Brossard, 1997 : Immunology of interactions between ticks and hosts. Brossard-M; Wikel-SK. Med-Vet-Entomol.; 1997 Jul; 11(3): 270-6
- 50 De Silva et al, 1995 : Growth and migration of Borrelia burgdorferi in Ixodes ticks during blood feeding. Aravinda M De Silva, Erol Fikrig. Am. J. Trop. Med. Hyg.; 53(4), 1995 pp 397-404
- Hubank and Schatz, 1994 : Identifying differences in mRNA expression by representational difference analysis of cDNA. Hubank-M; Schatz-DG. Nucleic-Acids-Res.; 1994 Dec 25; 22(25): 5640-8
- 55 Frohman, 1995 : Rapid amplification of cDNA Ends. In PCR Primer. A laboratory manual (Dieffenbach, C. W. and Dveksler, G. S., eds), pp.381-409, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

SEQUENCE LISTING

5

(1) INFORMATION FOR SEQ ID N°: 1

(i) SEQUENCE CHARACTERISTICS :

10

(a) LENGTH : 194 base pairs

(b) TYPE : nucleic acid

(c) STRANDEDNESS : single

(d) TOPOLOGY : linear

15

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°1 :

20

1 ATACCTTCCA CTTGTAGCCC TTCCTCATCC GATATGGTGA CCGATGCCAT

51 TGCATCCTCG TCGTGGAAGA GGTCTCTTC TAAATAAGAC CCATCCATAT

25

101 ATGTGTGTTT GCGAATGCCG TCGACGTAGC TCCTGACTAG AAACTCGTCC

151 GCTAGGACAG AACTTTTCTT CAGGTTTAGC GTAATGTCCT CGTT

30

(2) INFORMATION FOR SEQ ID N°: 2

(i) SEQUENCE CHARACTERISTICS :

35

(a) LENGTH : 607 base pairs

(b) TYPE : nucleic acid

(c) STRANDEDNESS : single

(d) TOPOLOGY : linear

40

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°2 :

45

1 TACCNGGGAA TCCAAAACCA ATTTTATTG GAACTTCCAC GTCTTCTTCA

51 AGGCGGTGGC ACCTCTGCAT TTATGAAGTT CGTCTTGGCA TTTTATTTTT

50

101 TGCTTCTTTC ATTGCRGAAC TCGCAAATGC ACTTCCCGTG CTTGTGCGCAT

151 TTCGCCCAA AAGCGCATGG CATTCTTCC GGCAGATTAA CTTTTCAAAA

55

201 TTCACGGTTC TGAACCAATA ATAGATCGTG GCAATGTTT TGCTGTTTGC

EP 1 892 297 A2

5
10
15
20

251 GATTTGCAAA CCAGCTGTAG CCACCATTGG ACTCAAAGGT GCGCACAACA
301 TGGCGCCGAA CTGTGAAAAA CAAATTAAGG CTNCTTTGTA ATAACGCTAG
351 TCTTGGTACG CCGTTAGAGG TCGATGTCCG GCCTCGCGAT TGCAAAGTCA
401 CTTGCACTTA TCAAGCTCCT GGAGAAAAAT GGGTGCAACG GGGGGATCAG
451 CGTTTGTACT TGCAAACATT TGTGGAGACG GTAAACCWGT ATTTCCGCGA
501 ACTCAGATGC TCCAGCGTGA AGCTCGTCTT AATAAAAAGTT GTAAATTCTGA
551 GTATNGATGA AGAACTGAAA TTCGAGGCAT TTAGAAACAC CACGAGAAGC
601 AGCGGAA

25 (3) INFORMATION FOR SEQ ID N°: 3

25

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 259 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

30

(ii) MOLECULE TYPE : cDNA

35

(iii) SEQUENCE DESCRIPTION : SEQ ID N°3 :

40

1 GATCCTACGC CTGAAAATGA GTGTCCATCG TCTTCACATA GTGCCACATT
51 GTAATTGGTA CAAGCTCCAT TTTCGTCAGC GCTGTTTGTGTT ATGCTGCCGC
101 CTACTTTTCC TTCGGCACTC CATAAGTTAA ACCCTGTCAT TATAAGTGTG
151 ATTGCCGTAT CTCGGCTGAA TGGGTTCCAT TTTTCTCTTA AATAATCAGC
201 TGTCCATATT CCATGTATTG TGTTTCATGAG TATGTGATTC TCATCGTATA
50 251 TCTTCGCCT

55

(4) INFORMATION FOR SEQ ID N°: 4

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 170 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°4 :

1 CCACTCGAAA ATGGAGGCTT TGAACATTT CAGTACCCCT GTGAACTCTG
51 GCTTTGCAAT GTAACAGCAA AAACACTTAC AGTTGAAGGG TGCAGTGTCA
101 GACGCTATGG AAGTTGCATC CACGAGCACR ACCCTGATTA CTACTGGCCA
151 CGTTGCTRTC CGGGTGCTCC

(5) INFORMATION FOR SEQ ID N°: 5

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 168 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°5 :

1 GTATGTTACC ATGTCCAACC CGGTTATTAA ATACACCAAG TCGTAGGATT
51 TGTAGGCAGC TGCATTGCCC TTGACGTACT CTCTCAACGT TGCCAAGGAC
101 TCAGGCCCAT AAATGTAGTG GGGTTGACCT TGA ACTCTTC GTAAAAAGCG
151 TTCTTCTCC GTCGTGAG

(6) INFORMATION FOR SEQ ID N°: 6

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 247 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°6 :

1 CCGAAMATAA AACTTAGTCT CACCAATATA CGTTTGCCTA ACGCGAAGGA
51 ACAGGCACAA ATATACTACG AGCACGACAT TCTCAAGAAC ACGGTTACAG
101 GAGTGTGGAC GAGAATTCAC TCAAAATATC CGTTCCTGA AGATGAGGGA
151 ATTACACTGA TAATGACAGG GTTTGATTTA TGGAGTGCCG ATTTAACTGT
201 AGGCGGCACC ATAACAAACA GCGCTGAGAA AAGCGGAGCT TGTACGA

(7) INFORMATION FOR SEQ ID N°: 7

5 (i) SEQUENCE CHARACTERISTICS :
 (a) LENGTH : 261 base pairs
 (b) TYPE : nucleic acid
 10 (c) STRANDEDNESS : single
 (d) TOPOLOGY : linear

15 (ii) MOLECULE TYPE : cDNA

(iii) FEATURE :
 (a) NAME/KEY : CDS
 20 (b) LOCATION : 1 .. 261

(iv) SEQUENCE DESCRIPTION : SEQ ID N°7

25

1 ATG CCT TTT ATT TTC GTG GTG AGC TTA GTC ATT GTG GCC TGC ATC GTG
 GTA GAC ACA
 30 Met Pro Phe Ile Phe Val Val Ser Leu Val Ile Val Ala Cys Ile Val
 Val Asp Thr

35 58 GCC AAC CAC AAA GGT AGA GGG CGG CCT GCG AAG TGT AAA CTT CCT CCG
 GAC GAC GGA
 Ala Asn His Lys Gly Arg Gly Arg Pro Ala Lys Cys Lys Leu Pro Pro
 40 Asp Asp Gly

45 115 CCA TGC AGA GCA CGA ATT CCG AGT TAC TAC TTT GAT AGA AAA ACC AAA
 ACG TGC AAG
 Pro Cys Arg Ala Arg Ile Pro Ser Tyr Tyr Phe Asp Arg Lys Thr Lys
 Thr Cys Lys

50 172 GAG TTT ATG TAT GGC GGA TGC GAA GGA AAC GAA AAC AAT TTT GAA AAC
 ATA ACT ACG
 Glu Phe Met Tyr Gly Gly Cys Glu Gly Asn Glu Asn Asn Phe Glu Asn
 55 Ile Thr Thr

EP 1 892 297 A2

229 TGC CAA GAG GAA TGC AGA GCA AAA AAA GTC TAG

Cys Gln Glu Glu Cys Arg Ala Lys Lys Val End

5

(8) INFORMATION FOR SEQ ID N°: 8

10

(i) SEQUENCE CHARACTERISTICS :

(a) LENGTH : 292 base pairs

(b) TYPE : nucleic acid

(c) STRANDEDNESS : single

(d) TOPOLOGY : linear

15

(ii) MOLECULE TYPE : cDNA

20

(iii) SEQUENCE DESCRIPTION : SEQ ID N°8 :

1 CATCGMAGCC ATAGTATATT TTGCACTTGT CTTCGGTTTC GTCGTAGTAG

25

51 GACCGATTCC ACATTGTAGT ACACCAGTCA CTTATATCCT GCGGGCGGGTG

101 CTTCGATTTG TCCTGAACAA ATCTTCCACA GCGCTTGTCG CACGCCTCCT

30

151 GGGAAATAGAA CGCGTTCTCT CCTCCGCATC TCCATTTGGA ATCATAGAAA

201 CATCTTTCAG TTTGAATATT GTAGCGATAA TAATCGGTAT CAGTTTCTTT

35

251 GCATGGTCCT GGGAGGGGTT TGGCGCAGGG GCCGTATTCA GG

(9) INFORMATION FOR SEQ ID N°: 9

40

(i) SEQUENCE CHARACTERISTICS :

(a) LENGTH : 270 base pairs

(b) TYPE : nucleic acid

(c) STRANDEDNESS : single

(d) TOPOLOGY : linear

45

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°9 :

50

1 GGTAATAGTT GTCAAATTCC ATTAATGTAT CCTGAAATGT GACCATATCT

55

51 TTGTTTCCCC TGTAATAATCT CATAAAAGGC TGTGTGTTTT CCTTAAGAAG

EP 1 892 297 A2

101 TGTAACAGCC ACGATGGTCA ATCTCACGGA TGGATGTGTG ACACTTTTAT
5 151 ATCTCAGGTT TGCCGACATT GCCATTACAG ATAAATAGTT GATAATTTCT
201 TTCTTGTTAT AGTTGTAAGC AGCCGATGTT GTTGCATCAA GCACCACATG
10 251 CACTTCAGGC AATATGGTTT

(10) INFORMATION FOR SEQ ID N°: 10

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 316 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°10 :

1 AGAAAGCAGT CATATTGGCC ATCCACAGGT CACAATGGTT CTCTCCTTGA
30 51 CCTGGCATCG GGATTCGAAG TATGGTGCAG TTCACGTAGT TGAATACAA
101 CACGAAATGT GTTCGTTGGT ACGCCAATAG GGGTTCTCGC AAAGAACATA
35 151 TCATTTGGAG GAAGGCGTAG TCCGTCGAGA TATCCAAAA CTAGGGTTTC
201 ATTGCGTGGC AACCAACTGC CCCCACTTCT GTATGTGTAC TGTAAGGAGT
40 251 RGTGAAACGG YGTCCTCTTT CCCATAACCT TGAAGTTTTC ACACTGCAGA
301 GGATTACCTC TCAAAA

(11) INFORMATION FOR SEQ ID N°: 11

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 241 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°11 :

5
 1 AAGGTAGCAA GGGTGGTAGG CTTTCCTCAC AAAGAGTCTG GCTTCCGTGA
 51 TAACCATATC CATTTCCTCAC CGTATACCGG TCATCCAACG TCAATTGTGT
 10
 101 TACAAGGCAG ATAATGTCAA AATGGCTCTG GTCCTATAA TAGTCGGATA
 151 ATGTAGAAAT CGCTCCATGT GGCCAAATAG ATGTTCTCTT TTCATACTGT
 15
 201 TTTAACTTTA ATTGTAGGTC CGCCTCGTTC TCGAGGTATG T

(12) INFORMATION FOR SEQ ID N°: 12

20
 (i) SEQUENCE CHARACTERISTICS :
 (a) LENGTH : 636 base pairs
 (b) TYPE : nucleic acid
 (c) STRANDEDNESS : single
 25
 (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

30
 (iii) SEQUENCE DESCRIPTION : SEQ ID N°12 :

1 TTCCCCNAAT TGGCCTTGCG ANNCTTGCAA GTCGACNCTA GAGGCTCCGA
 35
 51 AGATGGACAG ATTGCGCATG AAATATTTGA AATCGAGCAG AATGGTGATT
 101 TTAGGAGCGA TTATATTGTG CCACCCAGTT TGAAAGTGCA AGAACGCACA
 151 GTGGTTTACC GTAACAAGTA CACCAGAGTT OCTGTAAATT TTACCGTCGA
 40
 201 AGTTGCCATG CTGATTGATA AGTATTTATA CWAGGAGTTC AAGAACGAGA
 251 GCCACATCGT ACCGTACCTG GCTATGATAC TGACTTTGAT AAATCTGAGG
 45
 301 TATGCCGACA CACATGACCC GTACATCCAG TTTCTTCTCA CACAAGTGTT
 351 CGTGGGGA AW WCTGGCGATC ATATGGGCCA CATGCCCTTC CGACGAGCGT
 50
 401 TCTTGTTT CAG GCGCCGGCAT TATGCGCAGT TTAGGCCCAA TMACACCTTC
 451 CACTTGTAAT TCTCCGTTGT TGGATAGTGT AAGTGAGGCC ATTGCATCAG

55

EP 1 892 297 A2

501 CATCGTGGAA GARGCCTTCC TCCAAGTAGG AACCGCCCAT TTAGGTTTGC
551 TTTCCCAATC CGCCAATTTA ANTTTTAAAA AAAATTCCCC CCCGAAAAAT
601 TAATTTTTTT TAAAGGTGGA TTGTGATTC TCCGTT

(13) INFORMATION FOR SEQ ID N°: 13

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 432 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°13 :

1 GATCCCAAAA GTGCCCTGG ARCGACGGTT ACATCATGAG CTACGTCATA
51 AACTTCAAAA ACCACTTCAA ATTTTCTCCC TGCTGTGTAG AATCAATTCC
101 ATTCTGTGCA CGAGAGCGGG ACTGCCCTCTA CAAAGTCAAT GCCAAGGATG
151 CTGTAAAAAG CCTAATATCT CTGCCGGGAT TTAGGATATC GCCAACGAGT
201 TTCTGTCAAT TTATGCATCC GCTTTACCGC GGTGTCCATA GCGATAAGAA
251 AGCAGGTCTG TCCGATTGCG TACAGACGTG TAGAACGGCC AAAAATCGAC
301 GAGGAGGCTA CCATTCATGG ATTCACGCGG CACTTGACGG GGTTCCTTGC
351 GACAAGAGAA AGCCCAAGAA GGCCTGCATA AACGGGAAAT GCACCCTCCT
401 TAAGAGCATG CCCACAGAA CGTACCGGGA AT

(14) INFORMATION FOR SEQ ID N°: 14

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 466 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°14 :

1 AGGGCGTCT TTGCTTYACA GGGAACRGCA TATGGGCCAC GTGACCTTCC
51 AATGACCGCT CCAAATCTGG CATAGGTGA AYTCCGAAGT CGTGGCGCAG
101 CAGGCCTYCC ACATTCACTC CATCCTCGTC TTTTAGGATG ACTGCCGCCA
151 TTTGTTTTGT ATCGTGGTAC AGGTGTTTGT TATGGTCCGA GCCGTGACA
201 TAAGTATTGA CCAACGATCG GCCGAATGAT TACGGCTCAC CAAACACATC
251 AAATACCCCC GTCAAGTCAA GAGCTGGAAG CACAAAGCAT AGTATGTACA
301 AGATACCCCTT GGAAATCTTT CCGGAAGTTC ACCTTGTGGT GGACAGCACA
351 TTTGCCAAAG CTTTTAAATT TGACGTGTAC AAAGTAACGC GTTACTTCGC
401 AGTGCTTACA AATGCGGCTA ATCTTAGGTA TGCCAGCTTC GTATTTCCAA
451 AAGTACAGCT CAGGAT

(15) INFORMATION FOR SEQ.ID N°: 15

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 377 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°15 :

1 CTCGTCCACA CATTCTCCTA AAATGCAAGC CTTTTTTTTTC CCACAAGGTG

EP 1 892 297 A2

51 TACCGTCCGAC TACTACTGAGT CTCCAATAAA TATGTTTTCC GGTGCAATTT
101 ACCTTGCAGT CTTTGACGCC GTATGTAGGG TCAGCGTGCA TGCCTTCGTC
5
151 GTACATATAC ACCCTCTGAC AGTACTTGCT CAGTGTGTC ATCCTACCAG
201 GAAGCTTAGA CGAACGTTTT ATTGTTTTTG TCGTGATCG TTCTCTAAGG
10
251 CATTGGAATT CCGGACGGTT GTAGAGGTTT CTGACTTCTC GCTGGCAGCA
301 ATAAGAGAAC TGATACTGGC GCTCGTCTTG CATCTTGTAAT CTCATGAGGT
15
351 ATCCGTCATC CCATGGGCAG TCCGCAG

(16) INFORMATION FOR SEQ ID N°: 16

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 1670 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) FEATURE :

- (a) NAME/KEY : CDS
(b) LOCATION : 54 .. 1520

(iv) SEQUENCE DESCRIPTION : SEQ ID N°16 :

1 AAGGAAGAAG TTAGGCGTAG GCTTTGGGAA ACCGGTCATC CTCGAAACCA GAG
54 ATG TCG GGA CTC AGC CTG AAA TTG TGG ATT GTA GCG TTC TTT TCT
Met Ser Gly Leu Ser Leu Lys Leu Trp Ile Val Ala Phe Phe Ser
99 TTC TGC TTG GCC GAG AAA GAG CAT GGG ATC GTG TAC CCC AGG ATG CTT
Phe Cys Leu Ala Glu Lys Glu His Gly Ile Val Tyr Pro Arg Met Leu
147 GAA AGC AGA GCA GCA ACT GGA GAG AGA ATG CTT AAA ATC AAC GAT GAC
Glu Ser Arg Ala Ala Thr Gly Glu Arg Met Leu Lys Ile Asn Asp Asp
195 CTG ACG TTG ACG CTG CAG AAG AGT AAG GTC TTC GCT GAC GAC TTT CTC
Leu Thr Leu Thr Leu Gln Lys Ser Lys Val Phe Ala Asp Asp Phe Leu
243 TTC AGC ACG ACC GAC GGA ATT GAA CCT ATT GAT TAC TAC ATC AAA GCC

EP 1 892 297 A2

Phe Ser Thr Thr Asp Gly Ile Glu Pro Ile Asp Tyr Tyr Ile Lys Ala

5 291 GAA GAC GCT GAA CGT GAC ATC TAC CAC GAC GCA ACT CAC ATG GCA TCA
Glu Asp Ala Glu Arg Asp Ile Tyr His Asp Ala Thr His Met Ala Ser

10 339 GTA AGG GTA ACG GAC GAT GAT GGC GTG GAA GTG GAA GGA ATT CTT GGA
Val Arg Val Thr Asp Asp Asp Gly Val Glu Val Glu Gly Ile Leu Gly

15 387 GAG AGG CTT CGT GTT AAA CCT TTG CCG GCA ATG GCC CGC AGC AGC GAT
Glu Arg Leu Arg Val Lys Pro Leu Pro Ala Met Ala Arg Ser Ser Asp

20 435 GGC CTC AGA CCG CAT ATG TTG TAC GAA GTC GAC GCA CAC GAA AAC GGC
Gly Leu Arg Pro His Met Leu Tyr Glu Val Asp Ala His Glu Asn Gly

25 483 CGG CCA CAT GAT TAT GGT TCA CCG AAC ACA ACA AAT ACC CCC GTA GAG
Arg Pro His Asp Tyr Gly Ser Pro Asn Thr Thr Asn Thr Pro Val Glu

30 531 AGA AGA GCT GGA GGC ACA GAA CCC CAG ATG TAC AAG ATA CCA GCG GAA
Arg Arg Ala Gly Gly Thr Glu Pro Gln Met Tyr Lys Ile Pro Ala Glu

35 579 ATC TAT CCC GAA GTT TAC CTT GTG GCG GAT AGT GCC TTT GCC AAA GAA
Ile Tyr Pro Glu Val Tyr Leu Val Ala Asp Ser Ala Phe Ala Lys Glu

40 627 TTT AAC TTT GAT GTG AAC GCC GTT ACG CGT TAC TTC GCA GTG CTT ACA
Phe Asn Phe Asp Val Asn Ala Val Thr Arg Tyr Phe Ala Val Leu Thr

45 675 AAT GCG GCT AAT CTT AGG TAT GAA AGC TTC AAA TCT CCA AAG GTA CAG
Asn Ala Ala Asn Leu Arg Tyr Glu Ser Phe Lys Ser Pro Lys Val Gln

50 723 CTC AGG ATC GTT GGC ATA ACG ATG AAC AAA AAC CCA GCA GAC GAG CCA
Leu Arg Ile Val Gly Ile Thr Met Asn Lys Asn Pro Ala Asp Glu Pro

55 771 TAC ATT CAC AAT ATA CGG GGA TAT GAG CAG TAC CGG AAT ATT TTG TTT
Tyr Ile His Asn Ile Arg Gly Tyr Glu Gln Tyr Arg Asn Ile Leu Phe

819 AAG GAA ACA CTG GAG GAT TTC AAC ACT CAG ATG AAG TCA AAA CAT TTT
Lys Glu Thr Leu Glu Asp Phe Asn Thr Gln Met Lys Ser Lys His Phe

867 TAT CGT ACT GCC GAT ATC GTG TTT CTC GTG ACA GCA AAA AAT ATG TCC
Tyr Arg Thr Ala Asp Ile Val Phe Leu Val Thr Ala Lys Asn Met Ser

915 GAA TGG GTT GGT AGC ACA CTA CAA TCA TGG ACT GGC GGG TAC GCT TAC
Glu Trp Val Gly Ser Thr Leu Gln Ser Trp Thr Gly Gly Tyr Ala Tyr

963 GTA GGA ACA GCG TGT TCC GAA TGG AAA GTA GGA ATG TGT GAA GAC CGA
Val Gly Thr Ala Cys Ser Glu Trp Lys Val Gly Met Cys Glu Asp Arg

EP 1 892 297 A2

5
1011 CCG ACA AGC TAT TAC GGA GCT TAC GTT TTC GCC CAT GAG CTG GCG CAT
Pro Thr Ser Tyr Tyr Gly Ala Tyr Val Phe Ala His Glu Leu Ala His

10
1059 AAT TTG GGT TGT CAA CAC GAT GGA GAT GGT GCC AAT AGC TGG GTG AAA
Asn Leu Gly Cys Gln His Asp Gly Asp Gly Ala Asn Ser Trp Val Lys

15
1107 GGG CAC ATC GGA TCT GCG GAC TGC CCA TGG GAT GAC GGA TAC CTT ATG
Gly His Ile Gly Ser Ala Asp Cys Pro Trp Asp Asp Gly Tyr Leu Met

20
1155 AGC TAC AAG ATG GAA GAC GAG CGC CAG TAT AAG TTT TCT CCC TAC TGC
Ser Tyr Lys Met Glu Asp Glu Arg Gln Tyr Lys Phe Ser Pro Tyr Cys

25
1203 CAG AGA GAA GTC AGG AAC CTC TAC AGG CGT CCG GAA TTC AAA TGC CTC
Gln Arg Glu Val Arg Asn Leu Tyr Arg Arg Pro Glu Phe Lys Cys Leu

30
1251 ACT GAA CGA AAA GCG AAA AAA ACA ATC CGC TCG TCT AAG CTA CCT GGT
Thr Glu Arg Lys Ala Lys Lys Thr Ile Arg Ser Ser Lys Leu Pro Gly

35
1299 GTG ATG ACA TCA TCG AGC AAC TAT TGC CGG AGG GTG TAC ATG TAC GAA
Val Met Thr Ser Ser Ser Asn Tyr Cys Arg Arg Val Tyr Met Tyr Glu

40
1347 AAA GGC ATG CAC GCC GAC GAG GCA TaT GGC GTC AAG GAC TGC AGG GTA
Lys Gly Met His Ala Asp Glu Ala Tyr Gly Val Lys Asp Cys Arg Val

45
1395 AAA TGC ACC ACC ACA TCA AGA ATG TAT TGG CTA CTC GGT GTA GTC GAC
Lys Cys Thr Thr Thr Ser Arg Met Tyr Trp Leu Leu Gly Val Val Asp

50
1443 GGT ACA CCT TGC GGA AAT GGA AAG GCT TGC ATT CTT GGG AAA TGC AGG
Gly Thr Pro Cys Gly Asn Gly Lys Ala Cys Ile Leu Gly Lys Cys Arg

55
1491 AAC AAA ATC AAA ATA AGC AAG AAG GAC TGA GAGGTTGATA ATATCAAATT
Asn Lys Ile Lys Ile Ser Lys Lys Asp End

60
1541 AATCATGATA TTTCAACCAC ATGACTTCGT GCTCAACTGG TAGCCCCAAA TAAATTTTAA

65
1601 AAAAAATCCC AATATGCGTG GTAGAAAAAG CAGCAAACAA TAAACTTCT AAAAAATGTCT

70
1661 TGCAAAAATG

(17) INFORMATION FOR SEQ ID N°: 17

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 158 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°17 :

1 CACCAGTGAT GCTTATTGTT GCACTGCACT TGTTGATAAT ATCGGGTCGT
51 CGAATTGCAC TTCGGAACTT CCACTCCAAC TTGGCGAGCC GTGGATTTTG
101 ACTTCTCGTG ATGCTCCACC AGACAGTTGC AGGACTTCAG CTGCCTAGAT
151 GGAGCCTT

(18) INFORMATION FOR SEQ ID N°: 18

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 146 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°18 :

1 CTGTTGTTGA ACTGAAATAA ATAACAAAAA AATCATAAAG NTGGAGGAAA
51 GATGATCGAN TCCCCGCCCC TTGACAATCG TCCGATAAAA ACCAACTATA
101 TTCNGTCCTT TTTACAAACA ATTCCAANTG TCTGACCGAA CCGCGA

(19) INFORMATION FOR SEQ ID N°: 19

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 140 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°19 :

1 CTNGGAGGAN GTCCTATGAC TTGGGCTTAN GTTTCTTAGT CTTCTTCGGT
51 TTCTTCTTTT TTTGCTTCGG TTTTTCGGTG GCGGCAGGTG TATAGTCATC
101 AGTGTCCGGT GGGCCATCCG AATGAGTTGT CAAATGACAT

(20) INFORMATION FOR SEQ ID N°: 20

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 143 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°20 :

1 TGCCGAAAAA TAACGATGAT TTGACGTTGA CTCTGCAGAA GAGTAAGGTT
51 TTCACCGACA GTTTTCTGTT TAGCAGGACG AAGGATAACG AGCCTATCGA
101 TTACTACGTG AGAGCCGAAG ATGCCGAACG AGACATATAT CAC

(21) INFORMATION FOR SEQ ID N°: 21

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 140 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°21 :

5
1 TGTTGCTACA GACTCGACGT TTGGAGCTTG CTGGCCATTT MAAGACAACG
51 CACTCACAGA ATATTTAAGT GCGTTCGTGA WAGCTGTGGG CTTACGATTG
10
101 CAGGCGCTTC ANTCACCAGC TGTGATATTA MAGTTCCTAG

15
(22) INFORMATION FOR SEQ ID N°: 22

(i) SEQUENCE CHARACTERISTICS :

- 20
(a) LENGTH : 144 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

25
(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°22 :

30
1 TCAGGATAGT TGAAAGGTTG AAACCTGAAA TACTCCCACA GTCGTTGGAT
51 GCTTCAGAAC TGCTAAGAAC TTCACACTTT GCAAGAAGTW CCAAAATGAA
35
101 AGCCGCGATG ACCGATGATT TAGCTTCCAT CTTCTATCAC TTGA

(23) INFORMATION FOR SEQ ID N°: 23

40
(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 95 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

45
(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°23 :

50
1 GACCACCCCG TCCGAACTTG CTAAAKCAAG CAATGGAGTG AGGTGTTCTA
55
51 TGCGGGTTGA TTACACCAAT GGCGCTGCGT GGTGCGTGGT GATTT

EP 1 892 297 A2

509 TTT GCA GGA AAT GCC GAC AGG GTC CGT CTG GAG GTC AAT GCC TGG GTT
Phe Ala Gly Asn Ala Asp Arg Val Arg Leu Glu Val Asn Ala Trp Val

5

557 GAG GAA GTC ACC AGG TCA AAG ATC AGG GAC CTG CTC GCA CCT GGA ACT
Glu Glu Val Thr Arg Ser Lys Ile Arg Asp Leu Leu Ala Pro Gly Thr

10

605 GTT GAT TCA TCG ACA TCA CTT ATA TTA GTG AAT GCC ATT TAC TTC AAA
Val Asp Ser Ser Thr Ser Leu Ile Leu Val Asn Ala Ile Tyr Phe Lys

15

653 GGT CTG TGG GAT TCT CAG TTC AAG CCT AGT GCT ACG AAG CCG GGA GAT
Gly Leu Trp Asp Ser Gln Phe Lys Pro Ser Ala Thr Lys Pro Gly Asp

20

701 TTT CAC TTG ACA CCA CAG ACC TCA AAG AAA GTG GAC ATG ATG CAC CAG
Phe His Leu Thr Pro Gln Thr Ser Lys Lys Val Asp Met Met His Gln

25

749 GAA GGG GAC TTC AAG ATG GGT CAC TGC AGC GAC CTC AAG GTC ACT GCG
Glu Gly Asp Phe Lys Met Gly His Cys Ser Asp Leu Lys Val Thr Ala

30

797 CTT GAG ATA CCC TAC AAA GGC AAC AAG ACG TCG ATG GTC ATT CTC CTG
Leu Glu Ile Pro Tyr Lys Gly Asn Lys Thr Ser Met Val Ile Leu Leu

35

845 CCC GAA GAT GTA GAG GGA CTC TCA GTC CTG GAG GAA CAC TTG ACC GCT
Pro Glu Asp Val Glu Gly Leu Ser Val Leu Glu Glu His Leu Thr Ala

40

893 CCG AAA CTG TCG GCT CTG CTC GGC GGC ATG TAT GCG ACG TCC GAT GTC
Pro Lys Leu Ser Ala Leu Leu Gly Gly Met Tyr Ala Thr Ser Asp Val

45

941 AAC TTG CGC TTG CCG AAG TTC AAA CTA GAG CAG TCC ATA GGT TTG AAG
Asn Leu Arg Leu Pro Lys Phe Lys Leu Glu Gln Ser Ile Gly Leu Lys

50

989 GAT GTA CTG ATG GCG ATG GGA GTC AAG GAT TTC TTC ACG TCC CTT GCA
Asp Val Leu Met Ala Met Gly Val Lys Asp Phe Phe Thr Ser Leu Ala

55

1037 GAT CTT TCT GGC ATC AGC GCT GCG GGG AAT CTG TGC GCT TCG GAT GTC
Asp Leu Ser Gly Ile Ser Ala Ala Gly Asn Leu Cys Ala Ser Asp Val

1085 ATC CAC AAG GCT TTT GTG GAA GTT AAT GAG GAG GGC ACA GAG GCT GCA
Ile His Lys Ala Phe Val Glu Val Asn Glu Glu Gly Thr Glu Ala Ala

1133 GCT GCC ACT GCC ATA CCC ATT ATG TTG ATG TGT GCG AGA TTT CCA CAG
Ala Ala Thr Ala Ile Pro Ile Met Leu Met Cys Ala Arg Phe Pro Gln

1181 GTG GTG AAC TTT TTC GTT GAC CGC CCA TTC ATG TTC TTG ATC CAC AGC
Val Val Asn Phe Phe Val Asp Arg Pro Phe Met Phe Leu Ile His Ser

1229 CAT GAT CCA GAT GTT GTT CTC TTC ATG GGA TCG ATC CGT GAG CTC TAA

EP 1 892 297 A2

His Asp Pro Asp Val Val Leu Phe Met Gly Ser Ile Arg Glu Leu End

5
1277 AAAGCATATT CTTAACGGCG GCCAATCAGT CTGTGGAGTT ATCTCTTAGT CACTAATGTG
1337 TAACAATTCT GCAATATTCA GCTTGTGTAT TTCAGTAACT TGCTAGATCT TTGTGTTGTT
10
1397 GATGTTAGGC TTCTTGCG

(25) INFORMATION FOR SEQ ID N°: 25

15
(i) SEQUENCE CHARACTERISTICS :
(a) LENGTH : 200 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
20
(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

25
(iii) SEQUENCE DESCRIPTION : SEQ ID N°25 :

1 ACCGTAACCA AAATTGTTTC TTTCCAGAAG AATGGTTCAA ACTTTTCAAA
51 CAGATTTCCG AAACTCTTCT TGCACTTTAA AAATCCAATC TACAATCTTT
30
101 CCTGGCACTT CTGAATTGCA TTCCAGTTTA CCTTCCAAGC AAACCTCTTT
151 TGGCAACTCC AGCCGTACTC CATTTCGGCA TACCACAGTG CATGCACTTG
35

(26) INFORMATION FOR SEQ ID N°: 26

40
(i) SEQUENCE CHARACTERISTICS :
(a) LENGTH : 241 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
45
(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

50
(iii) SEQUENCE DESCRIPTION : SEQ ID N°26 :

1 CGTATTCTTT GAAGATTTGT ATACGAAACA TAAATTCGTC ATGCATACTT
51 TTGATGGTTA CACGACATGC GAAGCTGCCG ACAAAGAAGA CTGGGAAGAT
55

EP 1 892 297 A2

101 AAGAAGCACC TAGTTACGGT AGTGCCTGGA CCGGATAAAC GAAAGTACAC
5
151 GTTTCTACGC AACATTCTCA CCTTACAACG GAGAGTGAGA GTTAGCAAAA
201 CAATGATTGA GCTCGTACGG AACATGTCCT GTAGGACATT T
10

(27) INFORMATION FOR SEQ ID N°: 27

(i) SEQUENCE CHARACTERISTICS :

- 15
20
(a) LENGTH : 313 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) SEQUENCE DESCRIPTION : SEQ ID N°27 :

25
1 AAGCANCCGG ACTACCTGCT TGAAAACGTT GTACGGGCAA ACTTGGACGG
51 AAAACTGCCA GATGCTACTC CAGTTCCTCC CGGAAGCTAC ACGTACGCTG
30
101 AGAATGATAA CTTCACCTGC TATTCCAGAA GTACACCGTT TCCGGATGGG
151 GTGAATGTTG TATAACGGCT GCTGGGTGCG GAAGACTATG ATGGATTACG
35
201 CAAAAAAGTT CTAAACGAGT TGTTCCCAT CCCGGAAAAG CTGCTGTATG
251 CTGACATGAT GCGACTTGTG GCTAAGAAAAG ACAGAGTTGA TCACACTAGT
40
301 GGATGACCTG GGA

(28) INFORMATION FOR SEQ ID N°: 28

(i) SEQUENCE CHARACTERISTICS :

- 45
50
(a) LENGTH : 2417 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

55

EP 1 892 297 A2

(iii) FEATURE :

(a) NAME/KEY : CDS

(b) LOCATION : 218 .. 1495

(iv) SEQUENCE DESCRIPTION : SEQ ID N°28 :

5

10

15

20

25

30

35

40

45

50

55

1 GTCG TAGTCG TAGTCGTAGT CAGTTGCGCA TCGCGGGGGC TTCCTGTCT TTCTTGCCTT
61 TCTGCAGTCG TTCACCAACA TGTGGATACA GCTCCGGAGA TTTGTAAACA AATACTGCAC
121 TTTTAAGCAA GACTTGATAT TTAGATCGAT ATCCTCCTGT TGTCCGTCTT GATTAATCGG
181 CICTTTAGGG TTTT TAGAAT AGGCTTTTCG GTACGAG ATG CCC AAA GGA AAG AGG
Met Pro Lys Gly Lys Arg
236 GGA CCC AAA GCA GGT GGC GCC GCG CGG GGT GGC CGG TGC GAG GCC AGC
Gly Pro Lys Ala Gly Gly Ala Ala Arg Gly Gly Arg Cys Glu Ala Ser
284 CTG GCT CCG TCg TCC AGC GAC GAG GAG TCC AAC GCA GAC ACG GCG AGC
Leu Ala Pro Ser Ser Ser Asp Glu Glu Ser Asn Ala Asp Thr Ala Ser
332 GTG CTG AGC TGC GCC TCG GAG TCT CGC TGT GGC AGT GAC GGC ACC GTT
Val Leu Ser Cys Ala Ser Glu Ser Arg Cys Gly Ser Asp Gly Thr Val
380 GGA GAC CCA GAA GCG GAG GAG GCT GTG CTG CAT GAC GAC TTT GAA GAC
Gly Asp Pro Glu Ala Glu Glu Ala Val Leu His Asp Asp Phe Glu Asp
428 AAA CTC AAG GAG GCC ATC GAC GGA GCT TCG CAG AAG AGT GCC AAA GGA
Lys Leu Lys Glu Ala Ile Asp Gly Ala Ser Gln Lys Ser Ala Lys Gly
476 CGG CTG TCG TGC CTG GAG GCG ATT CGC AAG GCC TTT TcC ACC AAA TAC
Arg Leu Ser Cys Leu Glu Ala Ile Arg Lys Ala Phe Ser Thr Lys Tyr
524 CTG TAC GAC TTC CTC ATG GAC AGA CCG AGC ACG GTG TGC GAC CTG GTG
Leu Tyr Asp Phe Leu Met Asp Arg Pro Ser Thr Val Cys Asp Leu Val
572 GAG CGT GGG GTG CGC AAG GGC CGA GGG GAG GAG GCG GCC CTG TGC GCC
Glu Arg Gly Val Arg Lys Gly Arg Gly Glu Glu Ala Ala Leu Cys Ala
620 ACT CTC GGG GCC CTG GCC TGC GTC CAG CTC GGG GTC GGG GCC GAG GCG
Thr Leu Gly Ala Leu Ala Cys Val Gln Leu Gly Val Gly Ala Glu Ala
668 GAC GCC CTG TTC GAC GCC CTG CGC CAG CCG CTC TGC ACT TTG CTG CTT
Asp Ala Leu Phe Asp Ala Leu Arg Gln Pro Leu Cys Thr Leu Leu Leu
716 GAC GGG GCC CAG GGG CCC TCC CCC AGG GCC AGG TGT GCC ACT GCC CTC

EP 1 892 297 A2

Asp Gly Ala Gln Gly Pro Ser Pro Arg Ala Arg Cys Ala Thr Ala Leu

5 764 GGC CTC TGC TGC TTC GTG GTG GAC TCG GAC AAC CAG CTG GTG CTG CAG
Gly Leu Cys Cys Phe Val Val Asp Ser Asp Asn Gln Leu Val Leu Gln

812 CCG TGC ATG GAG GTG CTC TGG CAG GTG GTG GGT GCC AAG GCG GGC CCC
10 Pro Cys Met Glu Val Leu Trp Gln Val Val Gly Ala Lys Ala Gly Pro

860 GGC TCT CCG GTG CTC CAG GCA GCG GCC CTG CTC GCC TGG GGC CTC CTG
Gly Ser Pro Val Leu Gln Ala Ala Ala Leu Leu Ala Trp Gly Leu Leu

15 908 CTC AGC GTG GCT CCC GTC GAC CGC CTG CTG GCG CTC ACG CGC ACG CAC
Leu Ser Val Ala Pro Val Asp Arg Leu Leu Ala Leu Thr Arg Thr His

956 CTG CCC CGG CTG CAG GAG CTG CTG GAG AGC CCC GAC CTG GAC CTG CGC
20 Leu Pro Arg Leu Gln Glu Leu Leu Glu Ser Pro Asp Leu Asp Leu Arg

1004 ATT GCG GCC GGG GAG GTG ATC GCC GTC ATG TAC GAG GGG GCC AGG GAC
Ile Ala Ala Gly Glu Val Ile Ala Val Met Tyr Glu Gly Ala Arg Asp

25 1052 TAC GAC GAG GAC TTT GAG GAG CCC TCG GAG TCC CTG TGT GCC CAG CTG
Tyr Asp Glu Asp Phe Glu Glu Pro Ser Glu Ser Leu Cys Ala Gln Leu

1100 CGC CAG CTG GCC ACG GAC AGC CAG AAG TTT CGG GCC AAG AAG GAG CGG
30 Arg Gln Leu Ala Thr Asp Ser Gln Lys Phe Arg Ala Lys Lys Glu Arg

1148 CGC CAG CAG CGC TCC ACC TTC AGG GAC GTC TAC CGG GCC GTC AGG GAG
Arg Gln Gln Arg Ser Thr Phe Arg Asp Val Tyr Arg Ala Val Arg Glu

35 1196 GGG GCC TCT CCC GAC GTG AGC GTC AAG TTT GGC CGG GAA GTC CTG GAA
Gly Ala Ser Pro Asp Val Ser Val Lys Phe Gly Arg Glu Val Leu Glu

1244 CTG GAC ACC TGG AGT CGC AAG CTG CAG TAC GAC GCT TTC TGC CAG CTG
40 Leu Asp Thr Trp Ser Arg Lys Leu Gln Tyr Asp Ala Phe Cys Gln Leu

1292 CTG GGC TCC GGC ATG AAC CTG CAC CTG GCC GTG AAC GAG CTG CTG AGG
Leu Gly Ser Gly Met Asn Leu His Leu Ala Val Asn Glu Leu Leu Arg

45 1340 GAC ATC TTT GAA CTG GGG CAG GTG CTG GCA ACC GAG GAC CAC ATT ATC
Asp Ile Phe Glu Leu Gly Gln Val Leu Ala Thr Glu Asp His Ile Ile

1388 TCC AAG ATC ACC AAG TTC GAA AGG CAC ATG GTG AAC ATG GCC AGC TGC
50 Ser Lys Ile Thr Lys Phe Glu Arg His Met Val Asn Met Ala Ser Cys

1436 CGG GCC CGC ACC AAG ACA CGC AAC CGG CTG AGG GAC AAG CGC GCC GAC
Arg Ala Arg Thr Lys Thr Arg Asn Arg Leu Arg Asp Lys Arg Ala Asp

55

EP 1 892 297 A2

1484 GTG GTC GCC TGA ACCTGCGGAG GGATGCTTAG CTATGCACTC GCCGGCCTAC
Val Val Ala End

1536 CCTGGCGGGA CTCGATGCCA CTCACGAGTC GGCGCTCGCA AATTCGCCGC CGATCGTTAC

1596 GCAATGGGAG ACAAAGCTGC TTTTGGCATT ACCGTTTGAG GTCGGCTCCA ACCCATAGAT

1656 GAATTTCTTT TTTGTGGCCG TTTCTGGGTT ACATGTTTTG GGGGAAGGGA GTGGAACGTG

1716 CCGGTTCTTT GGCACACGTC AGGTGCTCT TGATGCGCGA CGTGCTTGTA TTTGGGtACT

1776 GCCGACACCA AGCGTTTCGG CGATTCCCTGG AAAAGAGTGC CTCTCGCTCG ACGTTTGgTT

1836 GTTTTCTGCG TGGTCCGTCG TCGACCTTCG TTCGTCCAAA GACGCCGTCC GTTTTCAtAC

1896 TCCCCCCC GC ACACATATCG AGGCCAATTA AATTGCTAAG GGTGCCGTTG TCGTGCATCT

1956 GGCAGGCTCA GAAGTGGCTT ATTTGTCTTT TAATTTTGCC GATGCACGCA AAAATTGTCA

2016 TTTCTTGAAA GTTTCTCTTT TATTGCGTAC ACAATTCAAC TTTTATGTAA TTTCTGATGG

2076 TCTGTTTTAC GTGTGCGTGT GTAAAACGTA ACTTTGGAAG AATTTTTATG CACACTGAAC

2136 AAACGCTCGG TCCTGGGGTT GAAAGTGCTC GGIGTGTCGCA TGAGCTAAAG TGCAACTGCT

2196 TTGTTCCGAA GGTTTTCTAG TCGCCGAAAT GTACCATTGT GGACCTTGTG GCGAGAGACC

2256 TTGGTCTTCT GGGGGAGCTG CTGTAGCGTG GCAAGCCACT ATTTTGGGAG CGACATTGCA

2316 GAGAAAATCG GCTTTTAGAA AGGCACCTGC GCGGCGAGTG GACGTTTTTT CGIATATACT

2376 GCGAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AA

(29) INFORMATION FOR SEQ ID N°: 29

(i) SEQUENCE CHARACTERISTICS :

- (a) LENGTH : 933 base pairs
- (b) TYPE : nucleic acid
- (c) STRANDEDNESS : single
- (d) TOPOLOGY : linear

(ii) MOLECULE TYPE : cDNA

(iii) FEATURE :

EP 1 892 297 A2

(a) NAME/KEY : CDS
 (b) LOCATION : 32 .. 853

5 (iv) SEQUENCE DESCRIPTION : SEQ ID N°29 :

1 GATTGGGAAC CTCCTATTCC TCACTTGAAA C ATG GCT GGA CTC CGC TCC
 Met Ala Gly Leu Arg Ser

10 50 TGC ATC CTC CTG GCT CTT GCC ACT AGT GCC TTC GCC GGC TAC CTT CAC
 Cys Ile Leu Leu Ala Leu Ala Thr Ser Ala Phe Ala Gly Tyr Leu His

15 98 GGT GGC CTT ACC CAC GGC GCT GGG TAC GGT TAC GGT GTC GGC TAC GGT
 Gly Gly Leu Thr His Gly Ala Gly Tyr Gly Tyr Gly Val Gly Tyr Gly

20 146 TCC GGC CTT GGC TAT GGC CTT GGC TAC GGT TCC GGC CTT GGC TAT GGA
 Ser Gly Leu Gly Tyr Gly Leu Gly Tyr Gly Ser Gly Leu Gly Tyr Gly

25 194 CAT GCT GTT GGC CTT GGA CAC GGC TTT GGC TAT TCT GGT CTG ACC GGC
 His Ala Val Gly Leu Gly His Gly Phe Gly Tyr Ser Gly Leu Thr Gly

30 242 TAC AGT GTG GCT GCC CCA GCT AGC TAC GCC GTT GCT GCT CCA GCC GTC
 Tyr Ser Val Ala Ala Pro Ala Ser Tyr Ala Val Ala Ala Pro Ala Val

35 290 AGC CGC ACC GTT TCC ACT TAC CAC GCT GCT CCA GCT GTG GCC ACC TAC
 Ser Arg Thr Val Ser Thr Tyr His Ala Ala Pro Ala Val Ala Thr Tyr

40 338 GCC GCT GCT CCT GTC GCC ACC TAT GCT GTT GCT CCA GCT GTC ACT AGG
 Ala Ala Ala Pro Val Ala Thr Tyr Ala Val Ala Pro Ala Val Thr Arg

45 386 GTT TCC CCC GTT CGC GCC GCC CCA GCT GTG GCC ACG TAC GCC GCC GCT
 Val Ser Pro Val Arg Ala Ala Pro Ala Val Ala Thr Tyr Ala Ala Ala

50 434 CCA GTC GCC ACC TAC GCC GCT GCT CCA GCT GTG ACC AGG GTG TCC ACC
 Pro Val Ala Thr Tyr Ala Ala Ala Pro Ala Val Thr Arg Val Ser Thr

55 482 ATT CAC GCT GCC CCG GCT GTG GCC AAT TAC GCC GTC GCT CCA GTC GCC
 Ile His Ala Ala Pro Ala Val Ala Asn Tyr Ala Val Ala Pro Val Ala

530 ACC TAT GCC GCT GCT CCA GCT GTG ACC AGG GTG TCC ACC ATC CAC GCC
 Thr Tyr Ala Ala Ala Pro Ala Val Thr Arg Val Ser Thr Ile His Ala

578 GCT CCA GCC GTG GCT AGC TAC CAG ACC TAC CAC GCT CCA GCT GTC GCC
 Ala Pro Ala Val Ala Ser Tyr Gln Thr Tyr His Ala Pro Ala Val Ala

626 ACT GTG GCT CAT GCT CCA GCT GTG GCC AGC TAC CAG ACC TAC CAG GCT
 Thr Val Ala His Ala Pro Ala Val Ala Ser Tyr Gln Thr Tyr His Ala

5
 674 GCC CCA GCC GTG GCT ACC TAC GCC CAT GCC GCT CCC GTC TAC GGC TAT
 Ala Pro Ala Val Ala Thr Tyr Ala His Ala Ala Pro Val Tyr Gly Tyr

722 GGT GTC GGT ACC CTC GGA TAT GGT GTC GGC CAC TAC GGC TAC GGA CAC
 Gly Val Gly Thr Leu Gly Tyr Gly Val Gly His Tyr Gly Tyr Gly His

10
 770 GGT CTT GGC AGC TAC GGC CTG AAC TAC GGT TAC GGC CTC GGC ACC TAC
 Gly Leu Gly Ser Tyr Gly Leu Asn Tyr Gly Tyr Gly Leu Gly Thr Tyr

15
 818 GGT GAC TAC ACC ACC CTT CTC CGC AAG AAG AAG TAA ATGGCA CATCTCAAGA
 Gly Asp Tyr Thr Thr Leu Leu Arg Lys Lys Lys End

870 GAGCCCATTG GACTGCCATC GACATTCTTC TTCAATAAAA GAGCCCGAAG ATGGCATTAT

20
 930 TTTT

25 **Claims**

1. A polynucleotide obtained from tick salivary gland and presenting more than 75% identity with the nucleotide sequence SEQ. ID. NO.7 or a sequence complementary thereto.
- 30 2. The polynucleotide according to claim 1, which presents at least 80% identity with SEQ.ID.NO.7 nucleotide sequence or a sequence complementary thereto.
3. The polynucleotide according to claim 1 or 2, which is at least 90% identical with SEQ.ID.NO.7 nucleotide sequence or a sequence complementary thereto.
- 35 4. The polynucleotide according to any one of the preceding claims 1 to 3, which is at least 95% identical with SEQ.ID.NO. 7 nucleotide sequence or a sequence complementary thereto.
- 40 5. The polynucleotide according to any one of the preceding claims 1 to 4, which is at least 98-99% identical with SEQ.ID.NO.7 nucleotide sequence or a sequence complementary thereto.
- 45 6. The polynucleotide according to any one of the preceding claims 1 to 5, which is at least 99% identical with SEQ.ID.NO. 7 nucleotide sequence or a sequence complementary thereto.
7. A polynucleotide having the nucleotide sequence SEQ.ID.NO.7 or a sequence complementary thereto.
8. A tick salivary gland polypeptide encoded by the polynucleotide according to any one of the preceding claims 1 to 7.
9. A polypeptide according to claim 8, modified by or linked to at least one substitution group, preferably selected from the group consisting of amide, acetyl, phosphoryl, and/or glycosyl groups.
- 50 10. The polypeptide according to claim 8 or 9 in the form of a "mature" protein.
11. The polypeptide according to claim 8 or 9 as part of a larger protein.
- 55 12. The polypeptide according to claim 8 or 9 as part of a fusion protein.
13. The polypeptide according to any one of the preceding claims 8 to 12 further including at least one additional amino

acid sequence which contains secretor or leader sequences, pro-sequences, sequences which help in purification such as multiple histidine residues, or additional sequences for stability during recombination protection.

- 5
14. A vector comprising at least one element selected from the group consisting of the polynucleotide according to any one of the preceding claims 1 to 7 or the polypeptide according to any one of the preceding claims 8 to 13.
15. A cell line transfected by the vector according to claim 14.
- 10
16. An antibody raised against *Ixodes ricinus* polypeptide according to any of the preceding claims 8 to 13.
17. A hybridoma cell line expressing the antibody according to the claim 16.
18. A pharmaceutical composition comprising an adequate pharmaceutical carrier and an element selected from the group consisting of the polynucleotide according to any of the preceding claims 1 to 7 or the polypeptide according to any of the preceding claims 8 to 13.
- 15
19. The pharmaceutical composition according to claim 18 which presents anti coagulant properties.
20. An immunological composition or vaccine for inducing an immunological response in a mammalian host to a tick salivary gland polypeptide which comprises at least one element of the group consisting of :
- 20
- a) a tick salivary gland cDNA according to any of the preceding claims 1 to 7,
 - b) a tick salivary gland polypeptide according to any of the preceding claims 8 to 13,
 - c) epitope-bearing fragments, outer-membrane vesicles or cells (attenuated or otherwise) of components a) or b) ; and
 - d) possibly a carrier.
- 25
21. Use of the immunological composition or vaccine according to claim 20 or the vector according to the claim 14 for the manufacture of a medicament to produce antibody and/or T-cell immune response to protect a mammal from bacteria and viruses transmitted during the blood meal of *Ixodes ricinus* and related species and in the treatment or the prevention of lyme diseases or tick encephalitis virus diseases.
- 30
22. A diagnostic kit for detecting a disease or susceptibility to a disease induced or transmitted by tick, especially *Ixodes ricinus*, which comprises:
- 35
- a) an *Ixodes ricinus* salivary gland polynucleotide according to any of the preceding claims 1 to 7,
 - b) a nucleotide sequence complementary to that of a),
 - c) an *Ixodes ricinus* salivary gland polypeptide according to any one of the preceding claims 8 to 13,
 - d) the antibody according to claim 16,
 - e) a phage displaying the antibody according to claim 16.
- 40
23. A polynucleotide which is identical to the nucleotide sequence according to any one of the preceding claims 1 to 7 for use as hybridisation probe for cDNA clones encoding ticks, more particularly *Ixodes ricinus* salivary gland polypeptide, or for isolating clones or other genes similar to tick salivary gland cDNAs.
- 45
- 50
- 55




Figure 1.




FIGURE 2.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 4816567 A [0063] [0063]
- WO 9201047 A [0066]

Non-patent literature cited in the description

- **LISITSYN et al.** *Science*, 1993, vol. 259, 946-951 [0026]
- **DIATCHENKO et al.** *Proc. Natl. Acad. Sci. USA*, 1996, vol. 93, 6025-6030 [0026]
- **PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES.** W. H. Freeman and Comany, 1993 [0034]
- **Posttranslational Protein Modifications : Perspectives and Prospects. POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS.** Academic Press, 1983, 1-12 [0034]
- **SEIFTER et al.** Analysis for protein modifications and nonprotein cofactors. *Meth Enzymol*, 1990, vol. 182, 626-646 [0034]
- **RATTAN et al.** Protein Synthesis : Posttranslational Modifications and Aging. *Ann NY Acad Sci*, 1992, vol. 663, 48-62 [0034]
- **COMPUTATIONAL MOLECULAR BIOLOGY.** Oxford University Press, 1988 [0037]
- **BIOCOMPUTING : INFORMATICS AND GENOME PROJECTS.** Academic Press, 1993 [0037]
- **COMPUTER ANALYSIS OF SEQUENCE DATA, PART I.** Humana Press, 1994 [0037]
- **SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY.** Academic Press, 1987 [0037]
- **SEQUENCE ANALYSIS PRIMER.** M Stockton Press, 1991 [0037]
- **CARILLO, H. ; LIPTON, D.** *SIAM J Applied Math*, 1998, vol. 48, 1073 [0037]
- **Guide to Huge Computers.** Academic Press, 1994 [0037]
- **CARILLO, H. ; LIPTON, D.** *SIAM J Applied Math*, 1988, vol. 48, 1073 [0037]
- **DEVEREUX, J. et al.** *J Molec Biol*, 1990, vol. 215, 403 [0037]
- **GENTZ.** *Proc Natl Acad Sci USA*, 1989, vol. 86, 821-824 [0047]
- **KOHLER ; MILSTEIN.** *Nature*, 1975, vol. 256, 495 [0057]
- **GODING.** *Monoclonal Antibodies : Principles and Practice.* Academic Press, 1986, 59-103 [0058]
- **KOZBOR.** *J Immunol*, 1984, vol. 133, 3001 [0059]
- **BRODEUR et al.** *Monoclonal Antibody Production Techniques and Applications.* Marcel Dekker, Inc, 1987, 51-63 [0059]
- **MUNSON ; POLLARD.** *Anal Biochem.*, 1980, vol. 107, 220 [0060]
- **WOLFF.** *Sciences*, 1990, vol. 247, 1465-8 [0069]
- **Current Protocols in Molecular Biology.** J. Wiley and sons, 1995 [0076]
- **GANAPAMO ; GANAPAMO-F ; RUTTI-B ; BROS-SARD-M et al.** Identification of an Ixodes ricinus salivary gland fraction through its ability to stimulate CD4 T cells present in BALB/c mice lymph nodes draining the tick fixation site. *Parasitology*, July 1997, vol. 115 (1), 91-6 [0094]
- **GANAPAMO ; GANAPAMO-F ; RUTTI-B ; BROS-SARD-M et al.** In vitro production of interleukin-4 and interferon-gamma by lymph node cells from BALB/c mice infested with nymphal Ixodes ricinus ticks. *Immunology*, May 1995, vol. 85 (1), 120-4 [0094]
- **GANAPAMO ; GANAPAMO-F ; RUTTI-B ; BROS-SARD-M et al.** Immunosuppression and cytokine production in mice infested with Ixodes ricinus ticks: a possible role of laminin and interleukin-10 on the in vitro responsiveness of lymphocytes to mitogens. *Immunology*, February 1996, vol. 87 (2), 259-63 [0094]
- **WIKEL ; WIKEL-SK et al.** Host immunity to ticks. *Annu-Rev-Entomol.*, 1996, vol. 41, 1-22 [0094]
- **WIKEL ; BROSSARD ; BROSSARD-M ; WIKEL-SK.** Immunology of interactions between ticks and hosts. *Med-Vet-Entomol*, July 1997, vol. 11 (3), 270-6 [0094]
- **DE SILVA ; ARAVINDA M DE SILVA ; EROL FIKRIG et al.** Growth and migration of Borrelia burgdorferi in Ixodes ticks during blood feeding. *Am. J. Trop. Med. Hyg.*, 1995, vol. 53 (4), 397-404 [0094]
- **HUBANK ; SCHATZ ; HUBANK-M ; SCHATZ-DG.** Identifying differences in mRNA expression by representational difference analysis of cDNA. *Nucleic-Acids-Res.*, 25 December 1994, vol. 22 (25), 5640-8 [0094]
- **Rapid amplification of cDNA Ends. FROHMAN.** In PCR Primer. A laboratory manual. Cold Spring Harbor Laboratory Press, 1995, 381-409 [0094]

专利名称(译)	蛋白质的鉴定和分子表征，在蜱唾液腺中表达		
公开(公告)号	EP1892297A2	公开(公告)日	2008-02-27
申请号	EP2007109568	申请日	2000-06-06
[标]申请(专利权)人(译)	HENOGEN		
申请(专利权)人(译)	HENOGEN S.A.		
当前申请(专利权)人(译)	HENOGEN S.A.		
[标]发明人	GODFROID EDMOND BOLLEN ALEX LEBOULLE GERARD		
发明人	GODFROID, EDMOND BOLLEN, ALEX LEBOULLE, GÉRARD		
IPC分类号	C12N15/12 C07K14/435 A61K38/17 G01N33/50 C12Q1/68 C07K16/18 C12N5/20 G01N33/53 A61K38/00 A61K39/00 A61K48/00 A61P7/02 A61P23/00 A61P29/00 A61P31/04 A61P31/12 A61P37/02 C12N1/15 C12N1/19 C12N1/21 C12N5/10 C12N15/09 G01N33/566		
CPC分类号	A61K38/00 A61K2121/00 A61P7/02 A61P23/00 A61P29/00 A61P31/04 A61P31/12 A61P37/02 C07K14/43527 Y02A50/401 Y10S435/975		
优先权	1999013425 1999-06-09 GB		
其他公开文献	EP1892297A3 EP1892297B1		
外部链接	Espacenet		

摘要(译)

本发明涉及DNA序列的分子表征，所述DNA序列编码在蜱的唾液腺中表达的蛋白质，更特别是蓖麻节肢动物蜱（Ixodes ricinus节肢动物蜱）。在血液的缓慢喂养阶段期间，在蓖麻毒素的唾液腺中诱导的基因被表征。这些基因的克隆是通过从诱导的cDNA中减去未诱导的cDNA（从未诱导的唾液腺中表达的mRNA合成）建立一个减去的cDNA文库（从慢速结束时在唾液腺中表达的mRNA合成）- 喂养阶段）。从感兴趣的不完整序列开始建立全长cDNA文库，并在消减cDNA文库中鉴定。本发明还包括新鉴定的多核苷酸和衍生多肽的各种用途。

SEQUENCE LISTING

```
(1) INFORMATION FOR SEQ ID N°: 1
(i) SEQUENCE CHARACTERISTICS :
(a) LENGTH : 194 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear
(ii) MOLECULE TYPE : cDNA
(iii) SEQUENCE DESCRIPTION : SEQ ID N°1 :
1 ATACCTTCCA CTTGTAGCCC TTCTCATCC GATATGGTGA CCGATGCCAT
31 TGCATCCTCC TCGTGAAGA GGTCTCTTC TAAATAAGAC CCGATCCATAT
101 ATGTGTGTTT GCGAATGCCG TCGAGTAGC TCCTGACTAG AAAGTCGTCC
151 CCTAGGACAG AACTTTTCCT CAGCTTAGC STAATGTCTT CATT

(2) INFORMATION FOR SEQ ID N°: 2
(i) SEQUENCE CHARACTERISTICS :
(a) LENGTH : 607 base pairs
(b) TYPE : nucleic acid
(c) STRANDEDNESS : single
(d) TOPOLOGY : linear
(ii) MOLECULE TYPE : cDNA
(iii) SEQUENCE DESCRIPTION : SEQ ID N°2 :
1 TACGNGGAA TCGAAAACGA ATTTTATTG GAACTGCCAC GTCTTCTCA
51 AGCGGGTGGC ACCTCTGCAT TTATGAAATG CCTCTTGCCA TTTTATTTT
101 TGCTTCTTTC ATTGCRGAAC TCGCAAAATCG ACTTCCCGTG CTTGTGGCAT
151 TTGCGCCCAA AAGCGCATGG CATTCTCTCC GGCAGATTAA CTTTTTGAAA
201 TTCAGGCTC TGAACCAATA ATAGATCGTG GCAATGTTTG TCTGTGTTTC
```