



(11) **EP 3 088 896 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
06.11.2019 Bulletin 2019/45

(51) Int Cl.:
G01N 33/68 (2006.01) G01N 33/53 (2006.01)
G01N 33/574 (2006.01)

(21) Application number: **13900340.4**

(86) International application number:
PCT/JP2013/085115

(22) Date of filing: **27.12.2013**

(87) International publication number:
WO 2015/097862 (02.07.2015 Gazette 2015/26)

(54) **METHOD FOR ANALYZING INFORMATION FOR DIAGNOSIS AND KIT THEREFOR**

VERFAHREN ZUR AUSWERTUNG VON INFORMATIONEN ZUR DIAGNOSE UND KIT DAFÜR
PROCÉDÉ D'ANALYSE D'INFORMATIONS POUR DIAGNOSTIC ET SA TROUSSE

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(43) Date of publication of application:
02.11.2016 Bulletin 2016/44

(73) Proprietor: **Konica Minolta, Inc.**
Tokyo 100-7015 (JP)

(72) Inventors:
• **YAMASHITA, Katsuko**
Yokohama-shi
Kanagawa 245-0053 (JP)
• **KAYA, Takatoshi**
Tokyo 100-7015 (JP)
• **IDE, Youichi**
Tokyo 100-7015 (JP)
• **KANEKO, Tomonori**
Tokyo 100-7015 (JP)

(74) Representative: **Gille Hrabal**
Brucknerstrasse 20
40593 Düsseldorf (DE)

(56) References cited:
WO-A1-2010/100862 WO-A1-2011/052244
WO-A1-2012/173228 WO-A1-2012/173228
WO-A1-2013/054281 WO-A2-2007/129114
JP-B2- 5 231 247 US-A1- 2011 294 141
US-A1- 2013 045 543

- **ATSUSHI MATSUDA ET AL: "Wisteria floribunda agglutinin-positive mucin 1 is a sensitive biliary marker for human cholangiocarcinoma", HEPATOLOGY, vol. 52, no. 1, 1 July 2010 (2010-07-01), pages 174-182, XP55030660, ISSN: 0270-9139, DOI: 10.1002/hep.23654**
- **Patricia A Burke ET AL: "Characterization of MUC1 glycoprotein on prostate cancer for selection of targeting molecules", INTERNATIONAL JOURNAL OF ONCOLOGY, 1 July 2006 (2006-07-01), pages 49-55, XP055367367, Retrieved from the Internet: URL:https://www.spandidos-publications.com/ijco/29/1/49/download [retrieved on 2017-04-25]**
- **DEVINE P L ET AL: "Prostate-specific antigen (PSA) and cancer-associated serum antigen (CASA) in distinguishing benign and malignant prostate disease", INTERNATIONAL JOURNAL OF BIOLOGICAL MARKERS, WICHTIG EDITORE, MILAN, IT, vol. 10, no. 4, 1 January 1995 (1995-01-01), pages 221-225, XP008181066, ISSN: 0393-6155**
- **KUNO ATSUSHI ET AL.: 'Wisteria floribunda agglutinin-positive mucin 1 is a sensitive biliary marker for human cholangiocarcinoma' HEPATOLOGY vol. 52, no. 1, 2010, pages 174 - 182, XP055030660**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 3 088 896 B1

Description

TECHNICAL FIELD

[0001] The present invention relates to a method of analyzing diagnostic information pertaining to a benign prostate disease and prostate cancer. The present invention also relates to a method of analyzing diagnostic information based on the blood concentration of mucin-1 having a specific sugar residue.

BACKGROUND ART

[0002] In order to allow proteins bearing a principal role in the biological life functions to orderly and properly exhibit their functions *in vivo*, post-translation modifications including glycosylation play an extremely important role. With regard to the post-translation modifications, the following findings were gradually made in recent years. The majority of the proteins in a living body are modified with sugar chains, and those sugar chains attached to the proteins play important roles in various aspects of biological phenomena, including protein stability, binding with hormones, binding with toxins, viral infection, mycoplasma infection, bacterial infection, protozoan infestation, fertilization, development and differentiation, cancer cell metastasis, apoptosis and the like. Even when proteins have the same amino acid sequence and the same name, such proteins are modified with a wide variety of sugar chains and, depending on the condition of the protein-producing cells, the structures of the sugar chains are variable and the proteins thus have different roles *in vivo*.

[0003] The relationships between such changes in sugar chains and diseases have also been gradually elucidated. For example, Patent Document 1 describes as follows with regard to prostate-specific antigens (hereinafter, referred to as "PSAs") which indicate that a subject has a prostate disease. That is, it is described that, as compared to blood samples originated from prostatic hyperplasia patients, those blood samples originated from prostate cancer patients contain a greater amount of a prostate-specific antigen having a specific sugar residue, i.e., an N-acetyl-D-galactosamine β 1-4 N-acetylglucosamine (hereinafter, referred to as "LacdiNAc") residue (this prostate-specific antigen is hereinafter referred to as "LacdiNAc-PSA") and/or a fucose- α (1,2)-galactose β 1 \rightarrow 4 N-acetylglucosamine residue, in its sugar chain. This means that the onset of prostate cancer changes the sugar chains of PSAs and this leads to an increase in the amount of PSAs having the above-described specific sugar residue(s), as a result of which a high concentration of PSAs having the specific sugar residue (s) is observed in blood samples of prostate cancer patients. On the other hand, since the onset of prostatic hyperplasia does not cause such a change in the sugar chain, prostatic hyperplasia patients are not observed with a change in the concentration of PSAs having the specific

sugar residue(s). On the basis of this, Patent Document 1 discloses a method of distinguishing prostate cancer by fraction measurement of sugar chains, which method is capable of distinguishing a prostate cancer patient from a prostatic hyperplasia patient by measuring, in their blood samples, the concentration of PSAs having the above-described specific sugar residue(s). In addition, for example, a method of identifying liver cancer by fraction measurement of α -fetoprotein (AFP) sugar chains and a method of identifying adenocarcinoma by fraction measurement of carcinoembryonic antigen sugar chains have also been proposed.

[0004] Further, mucin-1 (hereinafter, referred to as "MUC1"), which is one of mucins that are high-molecular-weight glycoproteins, is also known as a tumor-associated antigen. MUC1 is widely expressed in the epithelial cells of normal glands, and it is known that the expression of MUC1 is drastically increased when the cells are malignant, for example, in breast cancer, ovarian cancer, lung cancer, pancreas cancer and bladder cancer; and that the glycosylation pattern changes in breast cancer and the like. Patent Documents 2 and 3 disclose an aptamer ligand and an anti-MUC1 antibody, both of which can be used for the quantification of such MUC1, respectively. Moreover, Patent Document 4 describes that MUC1 and the like can be used as biomarkers of prostate cancer and colorectal cancer.

[0005] For specific detection of a glycoprotein containing a specific sugar residue in its sugar chain, proteins called lectin that are capable of specifically recognizing and binding to such sugar residue are widely utilized. This is because it is very difficult to prepare an antibody whose epitope is a sugar chain, particularly an antibody whose epitope is a specific sugar residue, and such an antibody is thus hardly available. Lectins not only are inexpensive and available in a large amount but also have excellent stability and can thus be stored over a long time.

[0006] For example, *Wisteria floribunda* lectin (*Wisteria floribunda* agglutinin; hereinafter, referred to as "WFA") is known to have N-acetylgalactosamine as its primary binding sugar residue. Patent Document 1 discloses a method in which WFA having such a property is bound to a carrier and loaded to a column and a PSA having a LacdiNAc residue in a side chain of an asparagine-linked sugar chain is subsequently fractionated and quantified by ELISA or the like. In addition, Patent Document 5 discloses a method in which a solid-phase anti-PSA antibody and a fluorescently labeled WFA are allowed to form a sandwich complex with a PSA having a LacdiNAc residue in a side chain of its sugar chain and this PSA having the specific sugar residue is then quantified by SPFS (Surface Plasmon-field enhanced Fluorescence Spectroscopy).

PRIOR ART REFERENCES

PATENT DOCUMENTS

[0007]

[Patent Document 1] WO 2010/090264
 [Patent Document 2] JP 2009-535051 A
 [Patent Document 3] WO 2011/135869
 [Patent Document 4] JP 2013-526852 A
 [Patent Document 5] JP 2013-076666 A

[0008] Moreover, WO 2011/052244 A discloses a method for distinguishing prostate cancer from prostatic hypertrophy, wherein the amount of PSA having N-acetylgalactosamine residues is determined using WFA lectin.

[0009] WO 2013/054281 A discloses a method to diagnose the presence of prostate cancer, whereby in the method the PSA-IgM occurring in a sample of a subject with prostate cancer is quantified.

[0010] "Wisteria floribunda agglutinin-positive mucin 1 is a sensitive biliary marker for human cholangiocarcinoma", Atsushi Matsuda et al., Hepatology, Vol. 52, No. 1, pages 174-182 discloses a sandwich ELISA assay combining the use of an anti-mucin-1 antibody and WFA.

SUMMARY OF THE INVENTION

PROBLEMS TO BE SOLVED BY THE INVENTION

[0011] Prostate cancer occurs primarily in men of 60 years and older. In Western countries, prostate cancer is the second leading cause of cancer-related death in men after lung cancer, and early detection of prostate cancer is thus desired. As a method of diagnosing prostate cancer, conventionally, a method in which all PSAs contained in a blood sample or the like are quantified as a total PSA value and this value is used as an index for determination is widely employed. Further, there is a method in which PSAs contained in a blood sample or the like that do not bind to α 1-antichymotrypsin are quantified as free PSAs and the ratio of the amount of the free PSAs with respect to the total PSA amount (free PSA/total PSA ratio) is used as an index for determination.

[0012] For example, (1) a method of determining a high probability of prostate cancer based on a total PSA value of 10 ng/mL or greater and (2) a method of determining a high probability of prostate cancer based on a total PSA value of 4 to 10 ng/mL (gray zone) and a free PSA/total PSA ratio of 25% or lower have been employed. For a patient determined with a high probability of prostate cancer, a prostate gland biopsy is performed to obtain a definitive diagnosis. However, in the above-described diagnostic methods, there are relatively many cases where a patient of a benign disease such as prostatic hyperplasia, not prostate cancer, is also determined with a high probability of prostate cancer. Since a prostate gland bi-

opsy is also performed for such patients of a benign disease and undue burden is thus placed on the patients, a diagnostic method capable of easily and highly accurately distinguishing prostate cancer from a benign disease has been desired.

[0013] In this respect, as described above, Patent Documents 1 and 5 disclose methods of distinguishing prostate cancer from prostatic hyperplasia, in which a lectin having an affinity for N-acetylgalactosamine residue, such as WFA, is brought into contact with a sample possibly containing a PSA; a PSA having an affinity for the lectin, namely a PSA having an N-acetylgalactosamine residue in its sugar chain, is quantified; and then prostate cancer is distinguished from prostatic hyperplasia based on the absolute amount of the specific PSA, that is, the concentration of the specific PSA in the blood sample, or the ratio of the amount of the specific PSA with respect to the amount of total PSA or free PSA. As a result of performing these methods, a prostatic hyperplasia patient who is potentially determined to have prostate cancer based only on the amount of total PSA or free PSA is correctly diagnosed with prostatic hyperplasia, not prostate cancer.

[0014] Although diagnostic methods with an improved accuracy of discriminating between prostate cancer and a benign prostate disease have been proposed as described above, an analysis method by which novel diagnostic information that enables to further improve the discrimination accuracy is continuously desired.

[0015] An object of the present invention is to provide an analysis method by which novel diagnostic information that enables to accurately distinguish prostate cancer from a benign prostate disease can be obtained.

TECHNICAL SOLUTION

[0016] The present inventors discovered that, as compared to a blood sample originated from a prostate cancer patient or healthy individual, a blood sample originated from a benign prostate disease patient contains a greater amount of WFA-reactive MUC1, that is, MUC1 presumed to have an N-acetyl-D-galactosamine \rightarrow serine (threonine) (GalNAc-Ser (Thr)) residue in its sugar chain (hereinafter, abbreviated as "Tn-MUC1"). Based on this discovery, the present inventors also discovered that a benign prostate disease and prostate cancer can be distinguished and diagnosed with good accuracy by quantifying the WFA-reactive MUC1 in a blood sample and utilizing this information, thereby completing the present invention. That is, the present invention enables to prevent a benign prostate disease patient from being misdiagnosed with prostate cancer and receiving unnecessary medical interventions.

[0017] In one aspect, the present invention provides methods of analyzing diagnostic information according to the following [1] to [7].

[1] A method of analyzing diagnostic information, the

method comprising: measuring the concentration of a mucin-1 (Tn-MUC1), which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine (threonine) residue, in a blood sample originated from a subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that a disease affecting the subject to be diagnosed is not prostate cancer but a benign prostate disease when the measured value of the concentration of Tn-MUC1 is greater than the threshold value.

[2] A method of analyzing diagnostic information, the method using a combination of: a method of analyzing diagnostic information that comprises: measuring the concentration of a mucin-1 (Tn-MUC1), which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine (threonine) residue, in a blood sample originated from a subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that a disease affecting the subject to be diagnosed is not prostate cancer but a benign prostate disease when the measured value of the concentration of Tn-MUC1 is greater than the threshold value; and a method of analyzing diagnostic information that comprises: measuring the concentration of all mucin-1s (total MUC1) in the blood sample originated from the subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that the disease affecting the subject to be diagnosed is not prostate cancer but the benign prostate disease when the measured value of the concentration of total MUC1 is greater than the threshold value.

[3] A method of analyzing diagnostic information for estimating which of a benign prostate disease and prostate cancer is affecting a subject to be diagnosed, the method using a combination of:

a method of analyzing diagnostic information that comprises: measuring the concentration of a mucin-1 (Tn-MUC1), which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine (threonine) residue, in a blood sample originated from a subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that a disease affecting the subject to be diagnosed is not prostate cancer but the benign prostate disease when the measured value of the concentration of Tn-MUC1 is greater than the threshold value; and diagnostic information relating to a prostate-specific antigen which indicates that the subject to be diagnosed has a prostate disease.

[4] A method of analyzing diagnostic information for estimating which of a benign prostate disease and prostate cancer is affecting a subject to be diag-

nosed, the method using a combination of:

a method of analyzing diagnostic information that comprises: measuring the concentration of a mucin-1 (Tn-MUC1), which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine (threonine) residue, in a blood sample originated from the subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that a disease affecting the subject to be diagnosed is not prostate cancer but the benign prostate disease when the measured value of the concentration of Tn-MUC1 is greater than the threshold value; and

a method of analyzing diagnostic information that comprises: measuring the concentration of a prostate-specific antigen (LacdiNAc-PSA) which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine β 1-4 *N*-acetylglucosamine residue in the blood sample originated from the subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that the disease affecting the subject to be diagnosed is not the benign prostate disease but prostate cancer when the measured value of the concentration of LacdiNAc-PSA is greater than the threshold value.

[5] The method of analyzing diagnostic information according to any one of [1] to [4], wherein the lectin that recognizes and binds to the *N*-acetyl-D-galactosamine → serine (threonine) residue is *Wisteria floribunda* lectin.

[6] The method of analyzing diagnostic information according to any one of [1] to [5], wherein the benign prostate disease is prostatic hyperplasia.

[7] The method of analyzing diagnostic information according to any one of [1] to [6], wherein the Tn-MUC1 or total MUC1 is reactive with an anti-KL-6 monoclonal antibody.

In another aspect, the present invention provides the use of a diagnostic kits for performing the method of analyzing diagnostic information as mentioned above under [1] or diagnostic kits according to the following [8] to [13].

[8] The use of a diagnostic kit for performing the method of analyzing diagnostic information according to [1], the kit comprising: an anti-mucin-1 antibody and a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine (threonine) residue for quantifying Tn-MUC1.

[9] A diagnostic kit for performing the method of analyzing diagnostic information according to [2], the kit comprising: an anti-mucin-1 antibody and a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine (threonine) residue for quantifying-

ing Tn-MUC1; and an anti-mucin-1 antibody for quantifying total MUC1.

[10] A diagnostic kit for performing the method of analyzing diagnostic information according to [3], the kit comprising: an anti-mucin-1 antibody and a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine (threonine) residue for quantifying Tn-MUC1; and an anti-prostate-specific antigen antibody for quantifying a prostate-specific antigen.

[11] A diagnostic kit for performing the method of analyzing diagnostic information according to [4], the kit comprising: an anti-mucin-1 antibody and a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine (threonine) residue for quantifying Tn-MUC1; and an anti-prostate-specific antigen antibody and a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine β1-4 *N*-acetylglucosamine residue for quantifying LacdiNAc-PSA.

[12] The use of the diagnostic kit of [8] or the diagnostic kit according to any one of [9] to [11], wherein the lectin that recognizes and binds to the *N*-acetyl-D-galactosamine → serine (threonine) residue is *Wisteria floribunda* lectin.

[13] The use of the diagnostic kit of [8] or the diagnostic kit according to any one of [9] to [12], wherein the anti-mucin-1 antibody is an anti-KL-6 monoclonal antibody.

[14] The use of the diagnostic kit of [8] or the diagnostic kit according to any one of [8] to [13], which further comprises an instruction manual that describes information required for performing the method of analyzing diagnostic information.

ADVANTAGEOUS EFFECTS OF INVENTION

[0018] The use of the diagnostic information obtained by the analysis method of the present invention, by itself, enables to determine a benign prostate disease with high accuracy; however, by using such diagnostic information in combination with conventional prostate cancer-related diagnostic information that leads to an incorrect estimation of a benign prostate disease patient as a prostate cancer patient, the possibility of such a misdiagnosis given by a conventional diagnostic method can be drastically reduced.

BRIEF DESCRIPTION OF DRAWINGS

[0019]

Fig. 1 is a scatter diagram that shows the results of the quantification step (1) of Example: Quantification of Tn-MUC1. It is noted here that "PC" represents the samples collected from prostate cancer patients and "BPH" represents the samples collected from prostatic hyperplasia patients.

Fig. 2 is a scatter diagram that shows, in an integrated manner, the results of the quantification step (1)

of Example: Quantification of Tn-MUC1 (abscissa) and the results of the quantification step (2): Quantification of Total MUC1(KL6) (ordinate). It is noted here that "PC" represents the samples collected from prostate cancer patients and "BPH" represents the samples collected from prostatic hyperplasia patients.

Fig. 3 is a scatter diagram that shows, in an integrated manner, the results of the quantification step (1) of Example: Quantification of Tn-MUC1 (abscissa) and the results of the quantification step (3): Quantification of Total PSA (ordinate). It is noted here that "PC" represents the samples collected from prostate cancer patients and "BPH" represents the samples collected from prostatic hyperplasia patients.

Fig. 4 is a scatter diagram that shows, in an integrated manner, the results of the quantification step (1) of Example: Quantification of Tn-MUC1 (abscissa) and the results of the quantification step (4): Quantification of LacdiNAc-PSA (ordinate). It is noted here that "PC" represents the samples collected from prostate cancer patients and "BPH" represents the samples collected from prostatic hyperplasia patients.

MODE FOR CARRYING OUT THE INVENTION

[0020] In the present specification, *α*-*N*-acetyl-D-galactosamine → serine (threonine), *N*-acetyl-D-galactosamine β1-4 *N*-acetylglucosamine, mucin-1 and *Wisteria floribunda* lectin are abbreviated as "Tn", "LacdiNAc", "MUC1" and "WFA", respectively. The notation "*α*-*N*-acetyl-D-galactosamine → serine (threonine)" may indicate "*α*-*N*-acetyl-D-galactosamine → serine" or "*α*-*N*-acetyl-D-galactosamine → threonine".

[0021] Further, MUC1 which is reactive with a lectin that recognizes a Tn residue as represented by WFA is referred to as "Tn-MUC1", and all of MUC1s including Tn-MUC1 and other MUC1s are referred to as "total MUC1". A PSA which is reactive with a lectin that recognizes a LacdiNAc residue and used in the present invention as represented by WFA is referred to as "LacdiNAc-PSA", and all of PSAs including LacdiNAc-PSA and other PSAs are referred to as "total PSA".

-Analysis Method-

[0022] The first embodiment of the method of analyzing diagnostic information according to the present invention is a method of analyzing diagnostic information which comprises: measuring the concentration of a mucin-1 which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine (threonine) residue, namely Tn-MUC1, in a blood sample originated from a subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that a disease affecting the subject to be diagnosed is not prostate cancer but a benign prostate disease when

the measured value of the concentration of Tn-MUC1 is greater than the threshold value.

[0023] The second embodiment of the method of analyzing diagnostic information according to the present invention is a method of analyzing diagnostic information which uses a combination of: a method of analyzing diagnostic information obtained by the first embodiment; and a method of analyzing diagnostic information that comprises: measuring the concentration of all of MUC1s, namely total MUC1, in a blood sample; comparing the thus measured value with a threshold value; and estimating that the disease affecting the subject to be diagnosed is not prostate cancer but the benign prostate disease when the measured value of the concentration of total MUC1 is greater than the threshold value.

[0024] In order to more clearly determine which of a benign prostate disease and prostate cancer is affecting the subject to be diagnosed, the diagnostic information that is obtained by the first embodiment of the analysis method of the present invention based on the Tn-MUC1 concentration in a blood sample can be utilized in combination with prostate cancer-related diagnostic information provided by a prior art that cannot clearly distinguish a benign prostate disease from prostate cancer.

[0025] All of prior arts have a possibility of incorrectly estimating a benign prostate disease as prostate cancer, and the prostate cancer-related diagnostic information provided by a prior art is not particularly restricted and can be obtained by any known method of analyzing diagnostic information. For example, diagnostic information based on the concentration of all of PSAs in a blood sample, which is conventionally and commonly used, or diagnostic information based on the concentration of a specific sugar residue-containing PSA in a blood sample, which is described in WO 2010/090264 (Patent Document 1) and JP 2013-76666 A (Patent Document 5), can be utilized.

[0026] Accordingly, the third embodiment of the method of analyzing diagnostic information according to the present invention can be, for example, a method of analyzing diagnostic information for estimating which of a benign prostate disease and prostate cancer is affecting a subject to be diagnosed, the method using a combination of: a method of analyzing diagnostic information obtained by the first embodiment; and a method of analyzing diagnostic information that comprises: measuring the concentration of all of PSAs, namely total PSA, in a blood sample originated from the subject to be diagnosed; comparing the thus measured value with a threshold value and; and estimating that the disease affecting the subject to be diagnosed is not the benign prostate disease but prostate cancer when the measured value of the concentration of total PSA is greater than the threshold value.

[0027] In the above-described third embodiment, as a method of analyzing diagnostic information that is used in combination with a method of analyzing diagnostic information obtained by the first embodiment, a method of using the value of total PSA was exemplified; however,

as described above, a method of analyzing diagnostic information that uses the ratio of the amount of free PSAs with respect to the total PSA amount (free PSA/total PSA ratio) as well as other known method of analyzing diagnostic information may be used.

[0028] The fourth embodiment of the method of analyzing diagnostic information according to the present invention is a method of analyzing diagnostic information for estimating which of a benign prostate disease and prostate cancer is affecting a subject to be diagnosed, the method using a combination of: a method of analyzing diagnostic information obtained by the first embodiment; and a method of analyzing diagnostic information that comprises: measuring the concentration of a PSA which is reactive with a lectin that recognizes and binds to a LacdiNAc residue, namely LacdiNAc-PSA, in a blood sample originated from the subject to be diagnosed; comparing the thus measured value with a threshold value and; and estimating that the disease affecting the subject to be diagnosed is not the benign prostate disease but prostate cancer when the measured value of the concentration of LacdiNAc-PSA is greater than the threshold value.

[0029] In the present invention, it is preferred that "Tn-MUC1" and "total MUC1" be reactive with an "anti-KL-6 monoclonal antibody". That is, in the measurement of the concentration of Tn-MUC1 and that of total MUC1, an anti-MUC1 antibody is usually used as an antibody for immobilization or labeling, and it is preferred to use an anti-KL-6 monoclonal antibody as the anti-MUC1 antibody.

[0030] There are several MUC1 variants; however, anti-MUC1 antibodies include monoclonal antibodies that recognize and bind to a moiety common to all of or the majority of the variants as an epitope, as well as monoclonal antibodies that bind to some of the variants, such as an anti-KL-6 monoclonal antibody that recognizes a sialylated sugar chain antigen KL-6 (an antigen composed of an amino acid sequence PDTRPAP and a sialylated sugar chain bound to the threonine (T) thereof, which is Neu5Ac α 2-3Gal β 1-3GalNAc α sugar chain) as an epitope and an anti-CA15-3 monoclonal antibody. In the present invention, a variety of anti-MUC1 antibodies can be used for the analysis of diagnostic information and, as one of such embodiments, a highly reliable analysis can be performed using an anti-KL-6 monoclonal antibody. In the following descriptions of the present specification, as a preferred embodiment, the term "anti-MUC1 antibody" is interchangeable with the term "anti-KL-6 monoclonal antibody".

[0031] Specifically, in the above-described second embodiment, when an "anti-KL-6 monoclonal antibody" is used as an "anti-MUC1 antibody" for measuring the concentration of "total MUC1" in the mode of a sandwich assay such as SPFS or ELISA, the "total MUC1" for which the concentration is thus measured means all of MUC1s that are reactive with the anti-KL-6 monoclonal antibody (hereinafter, referred to as "total MUC1 (KL6)"). Further,

in the above-described first and second embodiments and the below-described third and fourth embodiments, when an "anti-KL-6 monoclonal antibody" is used as an "anti-MUC1 antibody" for measuring the concentration of "Tn-MUC1" in the mode of a sandwich assay such as SPFS or ELISA, the "Tn-MUC1" for which the concentration is thus measured means an MUC1 which is reactive not only with the anti-KL-6 monoclonal antibody but also with a lectin that recognizes and binds to a Tn residue. This is common not only to those embodiments where, as in the below-described first method of quantifying Tn-MUC1, fractionation using a lectin that recognizes and binds to a Tn residue is not performed in advance and a sandwich complex of Tn-MUC1 is formed by an anti-MUC1 antibody for immobilization and the above-described prescribed lectin for detection, but also those embodiments where fractionation is performed in advance using a lectin that recognizes and binds to a Tn residue and then a sandwich complex of Tn-MUC1 is formed using an anti-MUC1 antibody for immobilization and an anti-MUC1 antibody for detection. Even when the "anti-MUC1 antibody" is other antibody, the "Tn-MUC1" for which the concentration is measured in the mode of a sandwich assay is substantially, among those MUC1 variants capable of binding with the antibody, one which is reactive with a lectin that recognizes and binds to a Tn residue.

[0032] The matters relating to the first to fourth embodiments will now be described sequentially.

• Blood Sample

[0033] In the analysis method of the present invention, as a sample for the measurement of the concentration of a specific glycoprotein, a blood sample originated from a subject to be diagnosed is used.

[0034] The subject to be diagnosed is typically a human; however, it may also be a non-human mammal, examples of which include human disease model animals such as mice, rats, guinea pigs, rabbits, goats, cats, dogs, pigs and monkeys.

[0035] An example of a preferred human subject to be diagnosed is a patient for whom it is necessary to improve the diagnostic accuracy and to clarify whether or not the patient has prostate cancer because, although other diagnostic method strongly indicates that the patient has prostate cancer, the possibility of the affecting disease being a benign prostate disease cannot be eliminated.

[0036] The benign prostate disease is typically prostatic hyperplasia (benign prostatic hyperplasia: BPH); however, examples thereof also include other prostate gland-related benign diseases, such as prostatitis.

[0037] The blood sample may be whole blood, or serum or plasma prepared from whole blood. Further, the blood sample may contain an anticoagulant that is added at the time of blood collection, as well as a diluent, a reagent and the like that are added as required after the collection. The blood sample can be collected and pre-

pared in accordance with a known method.

• Lectin

[0038] In the analysis method of the present invention, a lectin that recognizes and binds to a Tn residue is used for the quantification of Tn-MUC1. A representative example of such a lectin is WFA; however, lectins that can be used in the present invention are not restricted thereto.

That is, any lectin can be used in the present invention in the same manner as WFA, as long as the lectin is capable of recognizing and binding to a Tn residue contained in a sugar chain of Tn-MUC1 and can be utilized for the measurement of the concentration of Tn-MUC1 in a blood sample.

[0039] WFA is a lectin that is obtained from *Wisteria floribunda*, particularly seeds thereof, through separation, extraction and purification, and WFA is commercially available as well. Lectins other than WFA that may be used in the present invention can also be obtained from natural products by performing separation, extraction and purification as required, or can be purchased as commercially available products.

[0040] Further, in the fourth embodiment of the analysis method of the present invention, a lectin that recognizes and binds to a LacdiNAc residue is used for the quantification of LacdiNAc-PSA. A representative example of such a lectin is WFA; however, lectins that can be used in the present invention are not restricted thereto.

That is, any lectin can be used in the present invention in the same manner as WFA, as long as the lectin is capable of recognizing and binding to a LacdiNAc residue contained in a sugar chain of LacdiNAc-PSA and can be utilized for the measurement of the concentration of LacdiNAc-PSA in a blood sample.

[0041] Examples of the lectin used in the fourth embodiment of the analysis method of the present invention also include *Trichosanthes japonica* lectin-II (TJA-II) which recognizes and binds to both a β -GalNAc residue and a fucose- α (1,2)-galactose residue.

[0042] As described above, WFA is a lectin that is capable of recognizing and binding to the *N*-acetyl-D-galactosamine residue of both Tn and LacdiNAc and is thus suitable for achieving the object of the present invention because it can be commonly used for both the quantification of MUC1 (Tn-MUC1) having a Tn residue in its sugar chain and the quantification of PSA (LacdiNAc-PSA) having a LacdiNAc residue in its sugar chain. That is, a fluorescent-labeling lectin prepared from WFA can be commonly used for fluorescent labeling of Tn-MUC1 and fluorescent labeling of LacdiNAc-PSA. An embodiment that uses such a fluorescent-labeling lectin is preferred because reagents required for the analysis can be reduced and the measurement process can be simplified.

The use of a fluorescent-labeling lectin enables to simultaneously quantify plural measurement targets using a single SPFS measuring member as described below and to thereby simplify the measurement process.

[0043] WFA also recognizes and binds to galactose in addition to a Tn residue and a LacdiNAc residue; however, such binding does not cause a problem in the execution of the present invention because the strength of the binding is extremely weak and the binding constant of galactose is lower than that of WFA by several orders.

[0044] As for a lectin other than WFA, it is preferred that the lectin be capable of recognizing and binding to only a Tn residue and a LacdiNAc residue among sugar residues; however, any lectin can be used in the present invention as long as the ability of the lectin to recognize and bind to a sugar residue other than Tn residue and LacdiNAc residue is sufficiently weak in a range that causes substantially no problem in the execution of the present invention.

• Quantification Method

[0045] When performing the analysis method of the present invention, it is necessary to measure and quantify an item(s) that is/are required for each embodiment and selected from four items of measurement targets contained in a blood sample, which are Tn-MUC1, total MUC1, LacdiNAc-PSA and total PSA, or all of these four items of measurement targets. The method of quantifying these measurement targets is not particularly restricted as long as it is capable of yielding measured values with an accuracy sufficient for performing the analysis in each embodiment, and such a method can be appropriately selected from known quantification methods.

[0046] Examples of a preferred method of quantifying the measurement targets include SPFS (Surface Plasmon-field enhanced Fluorescence Spectroscopy) that is capable of quantifying a measurement target with high sensitivity and high accuracy. SPFS is a method which utilizes a phenomenon that, when an excitation light is irradiated to a metal thin film formed on a dielectric member at an angle that causes attenuated total reflection (ATR), an evanescent wave transmitting through the metal thin film is enhanced by several ten times to several hundred times due to resonance with surface plasmon, so as to efficiently excite a fluorescent substance labeling a measurement target captured in the vicinity of the metal thin film. By measuring the intensity of the fluorescence emitted from the fluorescent substance, the measurement target can be quantified based on the measured value. As required, this measured value is compared with a value obtained by measuring the fluorescence intensity using a standard sample having a known concentration, and the measured value can thereby be converted into the concentration of the measurement target in a blood sample. Such SPFS is extremely sensitive as compared to common fluorescent labeling methods and the like; therefore, it is capable of determining the concentration of a measurement target even when the measurement target is contained in a sample at an extremely low concentration.

[0047] Among the four items which are the Tn-MUC1

concentration, the total MUC1 concentration, the LacdiNAc-PSA concentration and the total PSA concentration, for a plurality of measurement items that are required depending on each embodiment, the data required for final analysis may be obtained by repeatedly quantifying a single item or a plurality of items which do not meet the requirement for the final analysis by themselves; however, from the standpoints of an improvement in the efficiency of the measurement process, the reliability of the measurement results and the like, it is preferred to quantify a plurality of measurement items that are required depending on each embodiment altogether at once.

[0048] For example, by forming a plurality of assay areas corresponding to each measurement item on a single SPFS measuring member (hereinafter, referred to as "sensor chip"), a plurality of measurement items can be quantified simultaneously. Such an embodiment is preferred because, as compared to an embodiment where a plurality of sensor chips are used to quantify the respective measurement targets, not only the cost and time required for the quantification can be cut down by reduction in the required members and reagents as well as the number of steps, but also variations in the measured values that are attributed to the differences among sensor chips can be reduced by the use of a single sensor chip.

[0049] Further, in the present invention, diagnostic information is analyzed based on the concentrations of Tn-MUC1, total MUC1, LacdiNAc-PSA and total PSA in a blood sample; however, it is also possible to, for the analysis, directly use "measured values" obtained by a quantification method such as SPFS in accordance with a prescribed measurement protocol, without converting each "measured value" into "concentration" expressed in a prescribed unit such as ng/mL. For example, in SPFS, a measured value of fluorescence intensity is usually expressed in an arbitrary unit (a.u.), and this measured value that reflects the concentration of a measurement target in a blood sample may be used to set a threshold value and perform the analysis of diagnostic information.

• Embodiments Based on SPFS

[0050] The fundamental constitutions for performing SPFS, namely embodiments of a sensor chip, a measuring apparatus, a system, measurement procedure and the like, are known, and they can be applied to appropriately carry out the present invention. In SPFS, a measurement target is generally quantified in the mode of a sandwich assay where a ternary complex composed of an immobilization probe (antibody), the measurement target and a fluorescent-labeling probe (lectin or antibody) is formed.

[0051] Among the Tn-MUC1, total MUC1, LacdiNAc-PSA and total PSA, for the quantification of Tn-MUC1 and LacdiNAc-PSA that are glycoproteins having a specific sugar residue, there are two representative quantification methods.

[0052] The first quantification method is a method in

which, without fractionating glycoproteins using a lectin that recognizes and binds to a specific sugar residue, the entire blood sample (glycoproteins) is subjected to SPFS and, among those glycoproteins that are captured by an immobilization probe (antibody), ones having a specific sugar residue are quantified by fluorescently labeling them with a fluorescent-labeling probe (lectin) that binds to the specific sugar residue. Although glycoproteins that do not have the specific sugar residue also bind to the immobilization probe, since the fluorescent-labeling probe (lectin) recognizing the specific sugar residue does not bind thereto, such glycoproteins are not quantified. For example, JP 2013-76666 A (Patent Document 5) discloses a method of quantifying a specific glycoprotein (PSA) by SPFS using a specific lectin (e.g., WFA) in the same manner as in the above-described first quantification method, and reference can be made to this method.

[0053] The second quantification method is a method in which, after fractionating a glycoprotein having a specific sugar chain in advance using a lectin that recognizes and binds to the specific sugar residue, only the thus obtained fractions of the glycoprotein are subjected to SPFS and the glycoprotein having the specific sugar chain that is captured by an immobilization probe (antibody) is quantified by fluorescently labeling the glycoprotein with a fluorescent-labeling probe (lectin) that binds thereto. Both of these methods can quantify Tn-MUC1 or LacdiNAc-PSA, which shows reactivity between a lectin recognizing a specific sugar residue and an anti-MUC1 antibody or an anti-PSA antibody. For example, WO2010/090264 (Patent Document 1) discloses a method in which a fraction containing a specific glycoprotein (PSA) is purified using a column (lectin affinity column) loaded with a carrier bound with a specific lectin (e.g., WFA) and the thus purified glycoprotein is quantified by ELISA, and reference can be made to this method.

[0054] A fractionation method using a lectin affinity column is summarized as follows. First, in accordance with a conventional method, a lectin affinity column is prepared by allowing a carrier for the column to react with a lectin that recognizes and binds to a specific sugar residue corresponding to a measurement target. Next, in accordance with a conventional method, a blood sample is applied to the thus prepared lectin affinity column so as to allow the measurement target having the specific sugar residue in the blood sample to bind to the lectin. Fractions containing substances that do not bind to the lectin (unbound fractions) are recovered as effluent by adding a washing buffer to the column. Then, fractions containing the measurement target bound with the lectin (bound fractions) are recovered as eluent by adding an elution buffer to the column. In this elution buffer, a hapten sugar capable of dissociating the measurement target bound with the lectin is dissolved. The hapten sugar may be selected in accordance with the lectin and, for example, when WFA is used as the lectin, GalNAc can be used as the hapten sugar.

(1) Quantification of Tn-MUC1

[0055] Examples of the first method of quantifying Tn-MUC1 include a method in which, the entire blood sample is subjected to SPFS without fractionation of Tn-MUC1 and then a ternary sandwich immunocomplex composed of an anti-MUC1 antibody immobilized on the surface of a sensor chip (immobilized anti-MUC1 antibody), Tn-MUC1 and a fluorescent substance-bound lectin for Tn-MUC1 (fluorescently labeled lectin for Tn-MUC1) is formed. Although MUC1s other than Tn-MUC1 also bind to the immobilized anti-MUC1 antibody, since the fluorescently labeled lectin for Tn-MUC1 does not bind thereto, the above-described complex is not formed and such MUC1s are thus not quantified.

[0056] The anti-MUC1 antibody can be prepared by a known method, or it can be purchased as a commercially available product. From the standpoint of the measurement stability, a monoclonal antibody is more preferably used than a polyclonal antibody. Examples of a commercially available anti-MUC1 monoclonal antibody include VU-3D1, VU-12E1, VU-11E2, VU-12E1, VU-3C6, VU-13F11, VU-5F12, VU-4C2, VU-2G7, VU-4H5, VU-11D1, M4021209, M3A106, M9E7, Ma552, VU-3C6, SM3, DF3, No. 1, No. 3, M2C5, 3H2740, 175C5, 126E7, C595, 139H2, 9A102, 8L820, M2F1, M4H2, 1B7, M10G4, EPR1023, EP1024Y, X325, 115D8, OC125 and X75. All of these anti-MUC1 monoclonal antibodies correspond to anti-KL-6 monoclonal antibodies.

[0057] The fluorescently labeled lectin for Tn-MUC1 can be prepared by binding a desired fluorescent substance to a lectin that recognizes a Tn residue as represented by WFA in accordance with a known method and, in this process, a commercially available fluorescent substance labeling kit or the like can also be used. The fluorescent substance is not particularly restricted and, for example, a known fluorescent dye that is capable of emitting an appropriate fluorescence in SPFS can be used.

[0058] Examples of the second method of quantifying Tn-MUC1 include a method in which, after a Tn-MUC1 fraction is prepared from a blood sample using a lectin for Tn-MUC1 in advance, the Tn-MUC1 fraction is subjected to SPFS and a ternary sandwich immunocomplex composed of an immobilized anti-MUC1 antibody, Tn-MUC1 and an anti-MUC1 antibody bound with a fluorescent substance (fluorescently labeled anti-MUC1 antibody) is formed.

[0059] The immobilized anti-MUC1 antibody used in this second method of quantifying Tn-MUC1 may be of the same type as the one used in the above-described first method of quantifying Tn-MUC1.

[0060] The fluorescently labeled anti-MUC1 antibody can be prepared by binding a desired fluorescent substance to an anti-MUC1 antibody in accordance with a known method and, in this process, a commercially available fluorescent substance labeling kit or the like can also be used. The fluorescent substance is not particularly restricted and, for example, a known fluorescent dye

that is capable of emitting an appropriate fluorescence in SPFS can be used. It is noted here that the type of the anti-MUC1 antibody of the fluorescently labeled anti-MUC1 antibody may be the same as or different from that of the immobilized anti-MUC1 antibody. Since MUC1s are extremely large molecules and contain a common repeating structure, they can be immobilized and fluorescently labeled with the same type of anti-MUC1 antibody.

(2) Quantification of Total MUC1

[0061] Examples of a method of quantifying the total MUC1 include a method in which a ternary sandwich immunocomplex composed of an anti-MUC1 antibody immobilized on the surface of a sensor chip (immobilized anti-MUC1 antibody), MUC1 and an anti-MUC1 antibody bound with a fluorescent substance (fluorescently labeled anti-MUC1 antibody) is formed.

[0062] The immobilized anti-MUC1 antibody used for the quantification of the total MUC1 may be of the same type as the one used for the quantification of Tn-MUC1. Further, the fluorescently labeled anti-MUC1 antibody (fluorescent substance and/or anti-MUC1 antibody) used for the quantification of the total MUC1 may be of the same type as the fluorescently labeled anti-MUC1 antibody (fluorescent substance and/or anti-MUC1 antibody) used in the second method of quantifying Tn-MUC1.

(3) Quantification of LacdiNAc-PSA

[0063] Examples of the first method of quantifying LacdiNAc-PSA include a method in which the entire blood sample (PSA) is subjected to SPFS without fractionation of LacdiNAc-PSA and then a ternary sandwich immunocomplex composed of an anti-PSA antibody immobilized on the surface of a sensor chip (immobilized anti-PSA antibody), LacdiNAc-PSA and a phosphor-bound lectin for LacdiNAc-PSA (fluorescently labeled lectin for LacdiNAc-PSA) is formed.

[0064] The anti-PSA antibody can be prepared by a commonly used known method, or it can be purchased as a commercially available product. From the standpoint of the measurement stability, a monoclonal antibody is more preferably used than a polyclonal antibody. Examples of a commercially available anti-PSA monoclonal antibody include PS2, PS3, PS4, PS5, PS6, PS15, 2H9, 3B4, 5A6, 5G6, 8G4, 9A8, 9G2, PS1, 8A6, 2H9, 1H12 and No. 79. Further, in order not to prevent the fluorescently labeled lectin for LacdiNAc-PSA from recognizing and binding to a specific sugar residue in a sugar chain, it is preferred to use an antibody whose epitope is the protein moiety, not the sugar chain, of PSA.

[0065] The fluorescently labeled lectin for LacdiNAc-PSA can be prepared by binding a desired fluorescent substance to a lectin that recognizes a LacdiNAc residue as represented by WFA in accordance with a commonly used known method and, in this process, a commercially

available fluorescent substance labeling kit or the like can also be used. The fluorescent substance is not particularly restricted and, for example, a known fluorescent dye that is capable of emitting an appropriate fluorescence in SPFS can be used.

[0066] Examples of the second method of quantifying LacdiNAc-PSA include a method in which, after a LacdiNAc-PSA fraction is prepared from a blood sample using a lectin for LacdiNAc-PSA in advance, the LacdiNAc-PSA fraction is subjected to SPFS and then a ternary sandwich immunocomplex composed of an immobilized anti-PSA antibody, LacdiNAc-PSA and an anti-PSA antibody bound with a fluorescent substance (fluorescently labeled anti-PSA antibody) is formed.

[0067] The immobilized anti-PSA antibody used in this second method of quantifying LacdiNAc-PSA may be of the same type as the one used in the above-described first method of quantifying LacdiNAc-PSA.

[0068] The fluorescently labeled anti-PSA antibody can be prepared by binding a desired fluorescent substance to an anti-PSA antibody in accordance with a known method and, in this process, a commercially available fluorescent substance labeling kit or the like can also be used. The fluorescent substance is not particularly restricted and, for example, a known fluorescent dye that is capable of emitting an appropriate fluorescence in SPFS can be used.

[0069] In the present invention, the fluorescently labeled lectin for LacdiNAc-PSA (fluorescent substance and/or lectin) that is used for the quantification of LacdiNAc-PSA may be of the same type as the fluorescently labeled lectin for Tn-MUC1 (fluorescent substance and/or lectin) that is used for the quantification of Tn-MUC1 and, for example, WFA can be commonly used as these lectins.

[0070] However, depending on the embodiment of the present invention, for acquisition of prostate cancer-related diagnostic information, not only PSA having a LacdiNAc residue but also both of PSA having a LacdiNAc residue and PSA having a fucose- α (1,2)-galactose residue can be the subject of quantification (see Patent Documents 1 and 5). In that case, the fluorescently labeled lectin for LacdiNAc-PSA can also be prepared using a lectin that recognizes both of the LacdiNAc residue and the fucose- α (1,2)-galactose residue, such as *Trichosanthes japonica* lectin-II.

(4) Quantification of Total PSA

[0071] Examples of a method of quantifying the total PSA include a method in which a ternary sandwich immunocomplex composed of an anti-PSA antibody immobilized on the surface of a sensor chip (immobilized anti-PSA antibody), PSA and an anti-PSA antibody bound with a phosphor (fluorescently labeled anti-PSA antibody) is formed.

[0072] The immobilized anti-PSA antibody used for the quantification of the total PSA may be of the same type

as the one used for the quantification of LacdiNAc-PSA. Further, the fluorescently labeled anti-PSA antibody (fluorescent substance and/or anti-PSA antibody) used for the quantification of the total PSA may be of the same type as the fluorescently labeled anti-PSA antibody (fluorescent substance and/or anti-MUC1 antibody) used in the second method of quantifying LacdiNAc-PSA.

• Embodiments Based on Other Quantification Method

[0073] Examples of a method used for the quantification of a measurement target other than SPFS include ELISA that is widely used as a method of quantifying a biologically relevant substance. In ELISA, a measurement target is generally quantified in the mode of a sandwich assay where a ternary complex composed of an immobilization probe (antibody), the measurement target and an enzyme-labeling probe (antibody or lectin) is formed, and the measurement target is quantified based on the amount of a dye (signal intensity) generated by reaction between the enzyme of the enzyme-labeling probe and a prescribed substrate.

[0074] In cases where Tn-MUC1 or LacdiNAc-PSA is quantified by ELISA, as in the respective second methods of quantifying Tn-MUC1 or LacdiNAc-PSA, it is required to separate in advance those fractions containing Tn-MUC1 or LacdiNAc-PSA from all of MUC1s or PSAs contained in a blood sample by utilizing a lectin that recognizes and binds to a specific sugar residue, for example, a lectin affinity column on which WFA is supported.

[0075] Basic methods for performing ELISA and fractionation are known, and the present invention can be appropriately carried out by applying these methods. For example, reference can be made to WO 2010/090264 (Patent Document 1) which describes, for example, a method of quantifying a specific glycoprotein, PSA, by utilizing ELISA and a lectin affinity column on which a specific lectin is supported.

• Threshold Value

[0076] The threshold value that is used in the first embodiment as well as in the second, third and fourth embodiments utilizing the first embodiment for the purpose of determining whether or not a measured value of the Tn-MUC1 concentration in a blood sample indicates the presence of a certain disease can be set by a commonly used method. For example, the Tn-MUC1 concentration is measured for blood samples originated from a plurality of patients who have been definitively diagnosed with prostate cancer and blood samples originated from a plurality of patients who have been definitively diagnosed with a benign prostate disease such as prostatic hyperplasia, and the concentration at which the measured data of the latter are included at a certain reference ratio or higher is determined and defined as the threshold value for estimating that a subject has a benign prostate disease. This reference ratio of the measured data corre-

sponds to the reliability relating to the estimation of a benign prostate disease, that is, the probability of correct estimation. A more reliable threshold value can be set by increasing the number of samples that represent the population of the measured data. When the Tn-MUC1 concentration measured for a particular blood sample is compared against a threshold value set in this manner and the measured value of the Tn-MUC1 concentration is greater than the threshold value, it can be estimated with a certain probability that this blood sample came from a subject to be diagnosed who has a benign prostate disease. It would be ideal if, from the measured data, a threshold value could be set such that a determination of 100% positive can be given when a blood sample originated from a subject to be diagnosed who has a benign prostate disease is analyzed and a determination of 100% negative can be given when a blood sample originated from a subject to be diagnosed who has prostate cancer is analyzed; however, it is realistically difficult to set such a threshold value. Taking into consideration the desired accuracy of the analysis, such a threshold value with which a positive determination can be obtained with a desirably high probability when a blood sample originated from a subject to be diagnosed who has a benign prostate disease is analyzed or, in other words, such a threshold value with which a probability of getting a false-positive determination can be minimized for the analysis of a blood sample originated from a subject to be diagnosed who has prostate cancer, can be set.

[0077] The threshold value used in the second embodiment against which the measured value of the total MUC1 concentration in a blood sample is compared, the threshold value used in the third embodiment against which the measured value of the total PSA concentration in a blood sample is compared, and the threshold value used in the fourth embodiment against which the measured value of the LacdiNAc-PSA concentration in a blood sample is compared can also be set in the same manner as described above.

-Kit-

[0078] In order to efficiently performing the method of analyzing diagnostic information according to the present invention, a kit can be configured by putting together necessary reagents and the like. The analysis of diagnostic information is typically performed using a quantification method based on a sandwich assay such as SPFS or ELISA and, as required, a fractionation method using a lectin affinity column in combination. Accordingly, the kit typically contains, as main constituents, reagents for the formation of a sandwich immunocomplex suitable for these methods, namely an antibody, which is a biologically relevant substance for immobilization, and a lectin or antibody, which is a biologically relevant substance for detection, and the kit may also contain, as required, a lectin for the preparation of a lectin affinity column as a constituent.

[0079] A kit for performing the first embodiment of the method of analyzing diagnostic information (first embodiment of the kit) can be configured to contain at least an anti-MUC1 antibody and a lectin that recognizes and binds to a Tn residue as reagents for the quantification of Tn-MUC1.

[0080] In a case where Tn-MUC1 is quantified in accordance with the above-described first method, the anti-MUC1 antibody contained in the kit is used for immobilization and binding on the surface of an SPFS sensor chip or ELISA substrate. Meanwhile, the lectin that recognizes and binds to a Tn residue is bound with a fluorescent substance to prepare a fluorescently labeled lectin for Tn-MUC1 when the quantification method is SPFS, or it is bound with an enzyme that reacts with a prescribed substrate to prepare an enzyme-labeled lectin for Tn-MUC1 and used for detection when the quantification method is ELISA.

[0081] In a case where Tn-MUC1 is quantified in accordance with the above-described second method, the anti-MUC1 antibody contained in the kit is used for immobilization and binding on the surface of an SPFS sensor chip or ELISA substrate, as well as for the preparation of a fluorescently labeled anti-MUC1 antibody where the anti-MUC1 antibody is bound with a fluorescent substance when the quantification method is SPFS, or for the preparation of an enzyme-labeled anti-MUC1 antibody where the anti-MUC1 antibody is bound with an enzyme that reacts with a prescribed substrate and for detection when the quantification method is ELISA. Accordingly, in this case, the anti-MUC1 antibody comprises an anti-MUC1 antibody for immobilization and an anti-MUC1 antibody for detection; however, these anti-MUC1 antibodies may be of the same type. Meanwhile, the lectin that recognizes and binds to a Tn residue is used for the production of a lectin affinity column used for the preparation of Tn-MUC1 fractions.

[0082] A kit for performing the second embodiment of the method of analyzing diagnostic information (second embodiment of the kit) can be configured to contain at least the following reagents: an anti-MUC1 antibody and a lectin that recognizes and binds to a Tn residue as reagents for the quantification of Tn-MUC1; and an anti-MUC1 antibody as a reagent for the quantification of the total MUC1.

[0083] With regard to the reagents for the quantification of Tn-MUC1, the same descriptions as provided above for the first embodiment of the kit are applied.

[0084] The anti-MUC1 antibody contained as a reagent for the quantification of the total MUC1 is used for immobilization and binding on the surface of an SPFS sensor chip or ELISA substrate, as well as for the preparation of a fluorescently labeled anti-MUC1 antibody where the anti-MUC1 antibody is bound with a fluorescent substance when the quantification method is SPFS, or for the preparation of an enzyme-labeled anti-MUC1 antibody where the anti-MUC1 antibody is bound with an enzyme that reacts with a prescribed substrate and for

detection when the quantification method is ELISA. The anti-MUC1 antibody for the quantification of the total MUC1 may be of the same type as the anti-MUC1 antibody for the quantification of Tn-MUC1.

[0085] A kit for performing the third embodiment of the method of analyzing diagnostic information (third embodiment of the kit) can be configured to contain at least the following reagents: an anti-MUC1 antibody and a lectin that recognizes and binds to a Tn residue as reagents for the quantification of Tn-MUC1; and an anti-PSA antibody as a reagent for the quantification of PSA.

[0086] With regard to the reagents for the quantification of Tn-MUC1, the same descriptions as provided above for the first embodiment of the kit are applied.

[0087] For example, in the quantification of the total PSA, the anti-PSA antibody contained as a reagent for the quantification of PSA is used for immobilization and binding on the surface of an SPFS sensor chip or ELISA substrate, as well as for the preparation of a fluorescently labeled anti-PSA antibody where the anti-PSA antibody is bound with a fluorescent substance when the quantification method is SPFS, or for the preparation of an enzyme-labeled anti-PSA antibody where the anti-PSA antibody is bound with an enzyme that reacts with a prescribed substrate and for detection when the quantification method is ELISA. Accordingly, in this case, the anti-PSA antibody comprises an anti-PSA antibody for immobilization and an anti-PSA antibody for detection.

[0088] A kit for performing the fourth embodiment of the method of analyzing diagnostic information (fourth embodiment of the kit) can be configured to contain at least the following reagents: an anti-MUC1 antibody and a lectin that recognizes and binds to a Tn residue as reagents for the quantification of Tn-MUC1; and an anti-PSA antibody and a lectin that recognizes and binds to a LacdiNAc residue as reagents for the quantification of LacdiNAc-PSA.

[0089] With regard to the reagents for the quantification of Tn-MUC1, the same descriptions as provided above for the first embodiment of the kit are applied.

[0090] In a case where LacdiNAc-PSA is quantified in accordance with the above-described first method, the anti-PSA antibody contained in the kit is used for immobilization and binding on the surface of an SPFS sensor chip or ELISA substrate. Meanwhile, the lectin that recognizes and binds to a LacdiNAc residue is bound with a fluorescent substance to prepare a fluorescently labeled lectin for LacdiNAc-PSA when the quantification method is SPFS, or it is bound with an enzyme that reacts with a prescribed substrate to prepare an enzyme-labeled lectin for LacdiNAc-PSA and used for detection when the quantification method is ELISA.

[0091] In a case where LacdiNAc-PSA is quantified in accordance with the above-described second method, the anti-PSA antibody contained in the kit is used for immobilization and binding on the surface of an SPFS sensor chip or ELISA substrate, as well as for the preparation of a fluorescently labeled anti-PSA antibody where the

anti-PSA antibody is bound with a fluorescent substance when the quantification method is SPFS, or for the preparation of an enzyme-labeled anti-PSA antibody where the anti-PSA antibody is bound with an enzyme that reacts with a prescribed substrate and for detection when the quantification method is ELISA. Accordingly, in this case, the anti-PSA antibody comprises an anti-PSA antibody for immobilization and an anti-PSA antibody for detection. Meanwhile, the lectin that recognizes and binds to a LactiNac residue is used for the production of a lectin affinity column used for the preparation of LactiNac-PSA fractions.

[0092] As the lectin that recognizes and binds to a Tn residue, which is contained in the first to fourth embodiments of the kit, WFA is preferred. The lectin that recognizes and binds to a Tn residue and the lectin that recognizes and binds to a LactiNac residue, which are contained in the fourth embodiments of the kit, may be of the same type and, for example, WFA can be commonly used as these lectins.

[0093] As the anti-MUC1 antibody contained in the first to fourth embodiments of the kit, an anti-KL-6 monoclonal antibody is preferred. Particularly, in the embodiment where the kit contains both an anti-MUC1 antibody for immobilization and an anti-MUC1 antibody for detection, it is preferred that both of these anti-MUC1 antibodies be anti-KL-6 monoclonal antibodies.

[0094] The kits for performing the method of analyzing diagnostic information according to the respective embodiments may also contain, as required, a reagent other than the above-described ones, an instrument, an instruction manual and/or the like. Examples of such a reagent and instrument include: a reagent for immobilizing a prescribed antibody on the surface of an SPFS sensor chip or ELISA substrate; a reagent for binding a fluorescent substance to a prescribed lectin or antibody; a blocking liquid for inhibition of non-specific adsorption to the surface of an SPFS sensor chip or ELISA substrate; a washing liquid for liquid transfer after application of a blood sample and a reagent (s) in each step of SPFS; a reagent for binding a prescribed lectin to a carrier of a lectin affinity column; a reagent for washing a lectin affinity column or for preparing an elution buffer; a treatment liquid for a blood sample; an instrument for reaction of these reagents; a sensor chip for SPFS; a substrate for ELISA; and a carrier of a lectin affinity column. Further, in the instruction manual, information required for performing the method of analyzing diagnostic information according to each embodiment of the present invention, for example, methods (protocols) for using the above-described reagents and instruments and threshold values relating to the analysis of diagnostic information, can be described.

EXAMPLES

[Production/Preparation Steps]

5 (1) Production of Plasmon Excitation Sensor

[0095] After plasma-cleaning a 1 mm-thick glass transparent planar substrate ("S-LAL 10" manufactured by Ohara Inc., refractive index = 1.72), a chromium thin film was formed on one side of this substrate by sputtering, and a gold thin film was further formed on the surface of this chromium thin film by sputtering. The chromium thin film had a thickness of 1 to 3 nm and the gold thin film had a thickness of 44 to 52 nm. The substrate on which a gold thin film was formed in this manner was immersed in an ethanol solution containing 10-carboxy-1-decanethiol (concentration = 25 mg/mL) for at least 24 hours to form a SAM film on the surface of the gold thin film. The substrate was then removed from the ethanol solution and washed with ethanol and isopropanol, followed by drying using an air gun.

(2) Production of Flow Channel-type Measuring Member for SPFS

[0096] On the surface of the plasmon excitation sensor (gold thin film) produced in the above-described step, a 0.5 mm-thick sheet-form silicone rubber spacer, which had a hole of 2 mm in width and 14 mm in length that served as a flow channel, was placed. A 2 mm-thick polymethyl methacrylate plate having a through-hole on both ends was further placed and press-adhered thereon, and the spacer and the plate were screw-fixed on the plasmon excitation sensor, thereby producing a flow channel-type measuring member for SPFS comprising a flow channel through which samples, reagents and the like could be transferred via the through-holes.

(3) Production of Fluorescently Labeled WFA

[0097] WFA (L-1350, manufactured by Vector Laboratories, Inc.) was fluorescently labeled using Alexa Fluor (trademark) 647 Protein Labeling Kit (manufactured by Invitrogen Corp.). The procedure of this process was in accordance with the protocol attached to the kit. In order to remove unreacted lectin and fluorescent dye, the reaction product was purified using an ultrafiltration membrane (manufactured by Nihon Millipore K.K.), and an Alexa Fluor 647-labeled WFA solution was thereby obtained. After determining the concentration of Alexa Fluor 647-labeled WFA in the thus obtained solution by measuring the absorbance, the solution was stored at 4°C. This fluorescently labeled WFA was used as a fluorescently labeled lectin for LactiNac-PSA.

55

(4) Production of Fluorescently Labeled Anti-MUC1 Antibody

[0098] An Alexa Fluor 647-labeled anti-MUC1 antibody was produced in the same manner as in the above step (3), except that a commercially available anti-MUC1 monoclonal antibody (anti-KL-6 monoclonal antibody) was used in place of WFA. It is noted here that, this anti-MUC1 monoclonal antibody was also used for the formation of an anti-MUC1 antibody-immobilized region as described below.

(5) Production of Fluorescently Labeled Anti-PSA Antibody

[0099] An Alexa Fluor 647-labeled anti-PSA antibody was produced in the same manner as in the above step (3), except that a commercially available anti-PSA monoclonal antibody (Clone No.79, 2.5 mg/mL; manufactured by Mikuri Immunolaboratory, Ltd.) was used in place of WFA.

(6) Preparation of Blood Samples

[0100] Serum samples originated from patients who had been definitively diagnosed with prostate cancer and serum samples originated from patients who had been definitively diagnosed with prostatic hyperplasia were each 2- to 10-fold diluted with phosphate buffer to prepare serum samples.

(7) Production of WFA Affinity Column

[0101] A WFA affinity column was produced by coupling WFA (L-1350, manufactured by Vector Laboratories, Inc.) to a Sepharose column (CNBr-Sepharose, manufactured by GE Healthcare) in accordance with the protocol attached to this product.

(8) Preparation of WFA-bound Fractions

[0102] The WFA affinity column (1-mL capacity) produced in (7) was equilibrated with a 0.1% bovine serum albumin (BSA)-containing phosphate buffer at 4°C. Each serum sample prepared in (6) was applied to this column and maintained for 30 minutes. Then, the column was washed with 5-fold volume of a washing buffer (0.1% BSA-containing phosphate buffer) and fractionated into 1-mL fractions, thereby obtaining WFA-unbound fractions. The column was brought back to room temperature, subsequently subjected to elution with 5-fold volume of an elution buffer (phosphate buffer containing 10 mM of GalNAc and 0.1% of BSA) and then fractionated into 1-mL fractions, thereby obtaining WFA-bound fractions.

[Quantification Steps]

[0103] Using the blood samples prepared in the above

production/preparation step (6) or the WFA-bound fractions prepared in the above step (8), the concentration was measured for each of Tn-MUC1, total MUC1, Lac-diNAc-PSA and total PSA by the below-described procedures.

(1) Quantification of Tn-MUC1

(1-1) Formation of Anti-MUC1 Antibody-Immobilized Region

[0104] In the flow channel of the flow channel-type measuring member for SPFS that was produced in the above production/preparation step (2), using a peristaltic pump, ultrapure water and then phosphate-buffered physiological saline were circulated for 10 minutes and 20 minutes, respectively, at room temperature (25°C) and a flow rate of 500 μ L/min, thereby equilibrating the surface of the plasmon excitation sensor. Subsequently, 5 mL of a phosphate-buffered physiological saline containing 50 mM of *N*-hydroxysuccinimide and 100 mM of ethyl(dimethylaminopropyl)carbodiimide was circulated for 20 minutes. Thereafter, 2.5 mL of a commercially available anti-MUC1 monoclonal antibody (anti-KL-6 monoclonal antibody) solution was circulated for 30 minutes to immobilize the antibody on the SAM film, thereby forming an anti-MUC1 antibody-immobilized region. Further, a phosphate-buffered physiological saline containing 1% by weight of bovine serum albumin was circulated for 30 minutes to perform a non-specific adsorption-inhibiting treatment in the flow channel.

(1-2) Measurement by SPFS

[0105] Sample Application Step: In the flow channel in which the anti-MUC1 antibody-immobilized region was formed, the WFA-bound fractions prepared in the above production/preparation step (8) were circulated for 25 minutes.

[0106] First Washing Step: A Tris-buffered physiological saline containing 0.05% by weight of Tween (registered trademark) 20 was circulated for 10 minutes.

[0107] Blank Measurement Step: With the flow channel being filled with the above-described Tris-buffered physiological saline, the metal thin film was irradiated with a laser light having a wavelength of 635 nm, whose photon amount was adjusted by an optical filter, from the back side of the plasmon excitation sensor through a prism. Then, using a CCD image sensor (manufactured by Texas Instruments Inc.) that was equipped with a $\times 20$ objective lens and arranged above the assay area, the intensity of the light passing through a filter arranged for cutting wavelengths of non-fluorescent components was measured. Since the thus measured value is a noise value that does not include the fluorescence emitted by the phosphor labeling Tn-MUC1, the measured value is defined as "blank value".

[0108] Labeling Step: Then, 5 mL of a phosphate-buff-

ered physiological saline containing the Alexa Fluor 647-labeled anti-MUC1 antibody produced in the above production/preparation step (4) (1 ng/mL) was circulated for 20 minutes.

[0109] Second Washing Step: A Tris-buffered physiological saline containing 0.05% by weight of Tween (registered trademark) 20 was circulated for 20 minutes.

[0110] Signal Measurement Step: With the flow channel being filled with the above-described Tris-buffered physiological saline, the light intensity was measured in the same manner as in the above-described blank measurement step. The thus measured value corresponds to a signal value that includes the fluorescence emitted by the phosphor labeling Tn-MUC1.

(2) Quantification of Total MUC1

(2-1) Formation of Anti-MUC1 Antibody-Immobilized Region

[0111] An anti-MUC1 antibody-immobilized region was formed in the same manner as in the above-described step (1-1) for the quantification of Tn-MUC1.

(2-2) Measurement by SPFS

[0112] The total MUC1 was quantified in the same manner as in the above-described step (1-2) for the quantification of Tn-MUC1, except that the blood samples prepared in the production/preparation step (6) were used in the sample application step in place of the WFA-bound fractions.

(3) Quantification of Total PSA

(3-1) Formation of Anti-PSA Antibody-Immobilized Region

[0113] An anti-PSA antibody-immobilized region was formed in the same manner as in the above-described step (1-1) for the quantification of Tn-MUC1, except that a commercially available anti-PSA monoclonal antibody was used in place of the anti-MUC1 monoclonal antibody.

(3-2) Measurement by SPFS

[0114] The total PSA was quantified in the same manner as in the above-described step (1-2) for the quantification of Tn-MUC1, except that the blood samples prepared in the production/preparation step (6) were used in the sample application step in place of the WFA-bound fractions and that the Alexa Fluor 647-labeled anti-PSA antibody produced in the production/preparation step (5) was used in the labeling step in place of the Alexa Fluor 647-labeled anti-MUC1 antibody.

(4) Quantification of LacdiNac-PSA

(4-1) Formation of Anti-PSA Antibody-Immobilized Region

[0115] An anti-PSA antibody-immobilized region was formed in the same manner as in the above-described step (1-1) for the quantification of Tn-MUC1, except that a commercially available anti-PSA monoclonal antibody was used in place of the anti-MUC1 monoclonal antibody.

(4-2) Measurement by SPFS

[0116] LacdiNac-PSA was quantified in the same manner as in the above-described step (1-2) for the quantification of Tn-MUC1, except that the blood samples prepared in the production/preparation step (6) were used in the sample application step in place of the WFA-bound fractions and that the Alexa Fluor 647-labeled WFA (fluorescently labeled lectin for LacdiNac-PSA) produced in the production/preparation step (3) was used in the labeling step in place of the Alexa Fluor 647-labeled anti-MUC1 antibody.

[Analysis Steps]

[0117] Fig. 1 shows the results of the quantification step (1): Quantification of Tn-MUC1. It is shown that the concentration of WFA-reactive Tn-MUC1 was higher in the blood samples of prostatic hyperplasia patients than in those of prostate cancer patients. Therefore, it is seen that, based on these results, the concentration corresponding to a value in a range of, for example, 1 to 1.5 units can be set as the threshold value used in the first embodiment of the analysis method of the present invention.

[0118] Fig. 2 shows, in an integrated manner, the results of the quantification step (1): Quantification of Tn-MUC1 (ordinate) and the results of the quantification step (2): Quantification of Total MUC1 (KL6) (abscissa). It is shown that the concentration of WFA-reactive Tn-MUC1 was higher in the blood samples of prostatic hyperplasia patients than in those of prostate cancer patients. That is, it is seen that, based on these results, the first threshold value in the analysis method of the present invention can be set in a range of, for example, 1 to 1.5 units. Further, it is also shown that the total MUC1 (KL6) concentration was higher in the blood samples of prostatic hyperplasia patients than in those of prostate cancer patients. That is, it is seen that, based on these results, the second threshold value in the analysis method of the present invention can be set in a range of, for example, 4.5 to 5 units. Therefore, it is understood that, in the second embodiment of the analysis method according to the present invention, the possibility of misdiagnosing a prostatic hyperplasia patient as a prostate cancer patient can be further reduced by using a combination of the first and second threshold values.

[0119] Fig. 3 shows, in an integrated manner, the results of the quantification step (1): Quantification of Tn-MUC1 (abscissa) and the results of the quantification step (3): Quantification of Total PSA (ordinate). It is shown that the concentration of WFA-reactive Tn-MUC1 was higher in the blood samples of prostatic hyperplasia patients than in those of prostate cancer patients. That is, it is seen that, based on these results, the threshold value used in the analysis method of the present invention can be set in a range of, for example, 1 to 1.5 units. Therefore, it is understood that, in the third embodiment of the analysis method according to the present invention, even when the threshold value relating to the total PSA, which have been used in conventional diagnosis, includes a gray zone and this yields false-positive results in which a certain number of prostatic hyperplasia patients are diagnosed with prostate cancer, these false-positive patients can be correctly diagnosed with prostatic hyperplasia by using the Tn-MUC1-related threshold value in combination.

[0120] Fig. 4 shows, in an integrated manner, the results of the quantification step (1): Quantification of Tn-MUC1 (abscissa) and the results of the quantification step (4): Quantification of LacdiNAC-PSA (ordinate). It is shown that the concentration of WFA-reactive Tn-MUC1 was higher in the blood samples of prostatic hyperplasia patients than in those of prostate cancer patients. That is, it is seen that, based on these results, the threshold value used in the analysis method of the present invention can be set in a range of, for example, 1 to 1.5 units. Therefore, it is understood that, in the fourth embodiment of the analysis method according to the present invention, even when the threshold value relating to LacdiNAC-PSA, which is disclosed in Patent Document 1, includes a gray zone and this yields false-positive results in which a certain number of prostatic hyperplasia patients are diagnosed with prostate cancer, these false-positive patients can be correctly diagnosed with prostatic hyperplasia by using the Tn-MUC1-related threshold value in combination.

Claims

1. A method of analyzing diagnostic information, said method comprising:

measuring the concentration of the mucin-1, Tn-MUC1, which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine/threonine residue, in a blood sample originated from a subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that a disease affecting said subject to be diagnosed is not prostate cancer but a benign prostate disease when said measured value of said concentration of Tn-MUC1 is greater

than said threshold value.

2. A method of analyzing diagnostic information, said method using a combination of:

a method of analyzing diagnostic information that comprises: measuring the concentration of the mucin-1, Tn-MUC1, which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine/threonine residue, in a blood sample originated from a subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that a disease affecting said subject to be diagnosed is not prostate cancer but a benign prostate disease when said measured value of said concentration of Tn-MUC1 is greater than said threshold value; and

a method of analyzing diagnostic information that comprises: measuring the concentration of total MUC1 in said blood sample originated from said subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that said disease affecting said subject to be diagnosed is not prostate cancer but said benign prostate disease when said measured value of said concentration of total MUC1 is greater than said threshold value.

3. A method of analyzing diagnostic information for estimating which of a benign prostate disease and prostate cancer is affecting a subject to be diagnosed, said method using a combination of:

a method of analyzing diagnostic information that comprises: measuring the concentration of the mucin-1, Tn-MUC1, which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine/ threonine residue, in a blood sample originated from said subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that a disease affecting said subject to be diagnosed is not prostate cancer but said benign prostate disease when said measured value of said concentration of Tn-MUC1 is greater than said threshold value; and

diagnostic information relating to a prostate-specific antigen which estimates that said subject to be diagnosed has a prostate cancer.

4. A method of analyzing diagnostic information for estimating which of a benign prostate disease and prostate cancer is affecting a subject to be diagnosed, said method using a combination of:

a method of analyzing diagnostic information that comprises: measuring the concentration of

- the mucin-1, Tn-MUC1, which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine/ threonine residue, in a blood sample originated from said subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that a disease affecting said subject to be diagnosed is not prostate cancer but said benign prostate disease when said measured value of said concentration of Tn-MUC1 is greater than said threshold value; and
- a method of analyzing diagnostic information that comprises: measuring the concentration of a prostate-specific antigen (LacdiNAc-PSA) which is reactive with a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine β 1-4 *N*-acetylglucosamine residue in said blood sample originated from said subject to be diagnosed; comparing the thus measured value with a threshold value; and estimating that said disease affecting said subject to be diagnosed is not said benign prostate disease but prostate cancer when said measured value of said concentration of LacdiNAc-PSA is greater than said threshold value.
5. The method of analyzing diagnostic information according to any one of claims 1 to 4, wherein said lectin that recognizes and binds to said *N*-acetyl-D-galactosamine → serine/threonine residue is *Wisteria floribunda* lectin.
6. The method of analyzing diagnostic information according to any one of claims 1 to 5, wherein said benign prostate disease is prostatic hyperplasia.
7. The method of analyzing diagnostic information according to any one of claims 1 to 6, wherein said Tn-MUC1 or total MUC1 is reactive with an anti-KL-6 monoclonal antibody.
8. Use of a diagnostic kit for performing the method of analyzing diagnostic information according to claim 1, said kit comprising:
- an anti-mucin-1 antibody and
a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine/threonine residue for quantifying Tn-MUC1.
9. A diagnostic kit for performing the method of analyzing diagnostic information according to claim 2, said kit comprising:
- an anti-mucin-1 antibody and a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine/threonine residue for quantifying Tn-MUC1;
- and an anti-mucin-1 antibody for quantifying total MUC1.
10. A diagnostic kit for performing the method of analyzing diagnostic information according to claim 3, said kit comprising:
- an anti-mucin-1 antibody and a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine/threonine residue for quantifying Tn-MUC1;
- and
an anti-prostate-specific antigen antibody for quantifying a prostate-specific antigen.
11. A diagnostic kit for performing the method of analyzing diagnostic information according to claim 4, said kit comprising:
- an anti-mucin-1 antibody and a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine → serine/threonine residue for quantifying Tn-MUC1;
- and
an anti-prostate-specific antigen antibody and a lectin that recognizes and binds to an *N*-acetyl-D-galactosamine β 1-4 *N*-acetylglucosamine residue for quantifying LacdiNAc-PSA.
12. The use of the diagnostic of claim 8 or the diagnostic kit according to any one of claims 9 to 11, wherein said lectin that recognizes and binds to said *N*-acetyl-D-galactosamine → serine/threonine residue is *Wisteria floribunda* lectin.
13. The use of the diagnostic of claim 8 or the diagnostic kit according to any one of claims 9 to 12, wherein said anti-mucin-1 antibody is an anti-KL-6 monoclonal antibody.
14. The use of the diagnostic of claim 8 or the diagnostic kit according to any one of claims 9 to 13, which further comprises an instruction manual that describes information required for performing said method of analyzing diagnostic information.

Patentansprüche

1. Verfahren zur Analyse diagnostischer Informationen, wobei das Verfahren aufweist:
- Messen der Konzentration des Mucin-1, Tn-MUC1, das mit einem Lektin reagiert, welches einen *N*-Acetyl-D-Galactosamin → Serin/Threonin-Rest in einer von einem zu diagnostizierenden Patienten stammenden Blutprobe erkennt und sich daran bindet;

- Vergleichen des so gemessenen Wertes mit einem Schwellenwert; und
Einschätzen, dass eine den zu diagnostizierenden Patienten betreffende Krankheit kein Prostatakrebs, sondern eine gutartige Prostataerkrankung ist, wenn der gemessene Wert der Konzentration an Tn-MUC1 größer ist als der Schwellenwert.
2. Verfahren zur Analyse diagnostischer Informationen, wobei das Verfahren eine Kombination aus verwendet:
- einem Verfahren zur Analyse diagnostischer Informationen, welches aufweist: Messen der Konzentration des Mucin-1, Tn-MUC1, das mit einem Lektin reagiert, welches einen *N*-Acetyl-D-Galactosamin → Serin/Threonin-Rest in einer von einem zu diagnostizierenden Patienten stammenden Blutprobe erkennt und sich daran bindet; Vergleichen des so gemessenen Wertes mit einem Schwellenwert; und Einschätzen, dass eine den zu diagnostizierenden Patienten betreffende Krankheit kein Prostatakrebs, sondern eine gutartige Prostataerkrankung ist, wenn der gemessene Wert der Konzentration an Tn-MUC1 größer ist als der Schwellenwert; und
einem Verfahren zur Analyse diagnostischer Informationen, welches aufweist: Messen der Konzentration des gesamten MUC1 in der von dem zu diagnostizierenden Patienten stammenden Blutprobe; Vergleichen des so gemessenen Wertes mit einem Schwellenwert; und Einschätzen, dass die den zu diagnostizierenden Patienten betreffende Krankheit kein Prostatakrebs, sondern eine gutartige Prostataerkrankung ist, wenn der gemessene Wert der Konzentration an gesamtem MUC1 größer ist als der Schwellenwert.
3. Verfahren zur Analyse diagnostischer Informationen zur Einschätzung, ob eine gutartige Prostataerkrankung oder Prostatakrebs einen zu diagnostizierenden Patienten befallen hat, wobei das Verfahren eine Kombination aus verwendet:
- einem Verfahren zur Analyse diagnostischer Informationen, welches aufweist: Messen der Konzentration des Mucin-1, Tn-MUC1, das mit einem Lektin reagiert, welches einen *N*-Acetyl-D-Galactosamin → Serin/Threonin-Rest in einer von einem zu diagnostizierenden Patienten stammenden Blutprobe erkennt und sich daran bindet; Vergleichen des so gemessenen Wertes mit einem Schwellenwert; und Einschätzen, dass eine den zu diagnostizierenden Patienten betreffende Krankheit kein Prostatakrebs, sondern eine gutartige Prostataerkrankung ist, wenn der gemessene Wert der Konzentration an Tn-MUC1 größer ist als der Schwellenwert; und
einem Verfahren zur Analyse diagnostischer Informationen, welches aufweist: Messen der Konzentration eines Prostata-spezifischen Antigens (LacdiNAc-PSA), das mit einem Lektin reagiert, welches einen *N*-Acetyl-D-Galactosamin-β1-4-*N*-Acetylglucosamin-Rest in einer von einem zu diagnostizierenden Patienten stammenden Blutprobe erkennt und sich daran bindet; Vergleichen des so gemessenen Wertes mit einem Schwellenwert; und Einschätzen, dass eine den zu diagnostizierenden Patienten betreffende Krankheit nicht die gutartige Prostataerkrankung, sondern Prostatakrebs ist, wenn der gemessene Wert der Konzentration an LacdiNAc-PSA größer ist als der Schwellenwert.
5. Verfahren zur Analyse diagnostischer Informationen gemäß irgendeinem der Ansprüche 1 bis 4, wobei das Lektin, das den *N*-Acetyl-D-Galactosamin → Serin/Threonin-Rest erkennt und sich daran bindet, *Wisteria floribunda* Lektin ist.
6. Verfahren zur Analyse diagnostischer Informationen gemäß irgendeinem der Ansprüche 1 bis 5, wobei die gutartige Prostataerkrankung Prostatahyperplasie ist.
4. Verfahren zur Analyse diagnostischer Informationen zur Einschätzung, ob eine gutartige Prostataerkrankung oder Prostatakrebs einen zu diagnostizierenden Patienten befallen hat, wobei das Verfahren eine Kombination aus verwendet:
- einem Verfahren zur Analyse diagnostischer Informationen, welches aufweist: Messen der Konzentration des Mucin-1, Tn-MUC1, das mit einem Lektin reagiert, welches einen *N*-Acetyl-D-Galactosamin → Serin/Threonin-Rest in einer von einem zu diagnostizierenden Patienten stammenden Blutprobe erkennt und sich daran bindet; Vergleichen des so gemessenen Wertes mit einem Schwellenwert; und Einschätzen, dass eine den zu diagnostizierenden Patienten betreffende Krankheit kein Prostatakrebs, sondern eine gutartige Prostataerkrankung ist, wenn der gemessene Wert der Konzentration an Tn-MUC1 größer ist als der Schwellenwert; und
einem Verfahren zur Analyse diagnostischer Informationen, welches aufweist: Messen der Konzentration eines Prostata-spezifischen Antigens (LacdiNAc-PSA), das mit einem Lektin reagiert, welches einen *N*-Acetyl-D-Galactosamin-β1-4-*N*-Acetylglucosamin-Rest in einer von einem zu diagnostizierenden Patienten stammenden Blutprobe erkennt und sich daran bindet; Vergleichen des so gemessenen Wertes mit einem Schwellenwert; und Einschätzen, dass eine den zu diagnostizierenden Patienten betreffende Krankheit nicht die gutartige Prostataerkrankung, sondern Prostatakrebs ist, wenn der gemessene Wert der Konzentration an LacdiNAc-PSA größer ist als der Schwellenwert.

7. Verfahren zur Analyse diagnostischer Informationen gemäß irgendeinem der Ansprüche 1 bis 6, wobei das Tn-MUC1 oder das gesamte MUC1 reaktiv mit einem Anti-KL-6 monoklonalen Antikörper ist.

8. Verwendung eines diagnostischen Kits zur Durchführung des Verfahrens zur Analyse diagnostischer Informationen gemäß Anspruch 1, wobei das Kit aufweist:

einen Anti-Mucin-1-Antikörper und ein Lektin, das einen *N*-Acetyl-D-Galactosamin → Serin/Threonin-Rest erkennt und sich daran bindet, um Tn-MUC1 zu quantifizieren.

9. Diagnostisches Kit zur Durchführung des Verfahrens zur Analyse diagnostischer Informationen gemäß Anspruch 2, wobei das Kit aufweist:

einen Anti-Mucin-1-Antikörper und ein Lektin, das einen *N*-Acetyl-D-Galactosamin → Serin/Threonin-Rest erkennt und sich daran bindet, um Tn-MUC1 zu quantifizieren; und einen Anti-Mucin-1-Antikörper, um Gesamt-MUC1 zu quantifizieren.

10. Diagnostisches Kit zur Durchführung des Verfahrens zur Analyse diagnostischer Informationen gemäß Anspruch 3, wobei das Kit aufweist:

einen Anti-Mucin-1-Antikörper und ein Lektin, das einen *N*-Acetyl-D-Galactosamin → Serin/Threonin-Rest erkennt und sich daran bindet, um Tn-MUC1 zu quantifizieren; und einen Antikörper gegen ein Prostata-spezifisches Antigen, um ein Prostata-spezifisches Antigen zu quantifizieren.

11. Diagnostisches Kit zur Durchführung des Verfahrens zur Analyse diagnostischer Informationen gemäß Anspruch 4, wobei das Kit aufweist:

einen Anti-Mucin-1-Antikörper und ein Lektin, das einen *N*-Acetyl-D-Galactosamin → Serin/Threonin-Rest erkennt und sich daran bindet, um Tn-MUC1 zu quantifizieren; und einen Antikörper gegen ein Anti-Prostata-spezifisches Antigen und ein Lektin, welches einen *N*-Acetyl-D-Galactosamin-β1-4-*N*-Acetylglucosamin-Rest erkennt und sich daran bindet, um LacDiNAc-PSA zu quantifizieren.

12. Verwendung der Diagnostik gemäß Anspruch 8 oder des Diagnosekits gemäß irgendeinem der Ansprüche 9 bis 11, wobei das Lektin, das einen *N*-Acetyl-D-Galactosamin → Serin/Threonin-Rest erkennt

und sich daran bindet, ein *Wisteria floribunda* Lektin ist.

13. Verwendung der Diagnostik gemäß Anspruch 8 oder des Diagnosekits gemäß irgendeinem der Ansprüche 9 bis 12, wobei der Anti-Mucin-1-Antikörper ein Anti-KL-6 monoklonaler Antikörper ist.

14. Verwendung der Diagnostik gemäß Anspruch 8 oder des Diagnosekits gemäß irgendeinem der Ansprüche 9 bis 13, welches ferner eine Gebrauchsanweisung aufweist, die für die Durchführung des Verfahrens zur Analyse diagnostischer Informationen erforderliche Informationen beschreibt.

Revendications

1. Procédé d'analyse d'informations diagnostiques, ledit procédé comprenant les étapes de :

mesurer la concentration de mucine-1, Tn-MUC1, qui est réactive avec une lectine, qui reconnaît et se relie à un reste de *N*-acétyl-D-galactosamine → sérine/thréonine dans un échantillon de sang provenant d'un patient à diagnostiquer ;

comparer la valeur ainsi mesurée avec une valeur de seuil ; et

estimer que la maladie affectant ledit patient à diagnostiquer n'est pas du cancer de la prostate, mais une maladie bénigne de la prostate, si ladite valeur mesurée de ladite concentration de Tn-MUC1 est supérieure à ladite valeur de seuil.

2. Procédé d'analyse d'informations diagnostiques, ledit procédé utilisant une combinaison de :

un procédé d'analyse d'informations diagnostiques, qui comprend les étapes de : mesurer la concentration de mucine-1, Tn-MUC1, qui est réactive avec une lectine, qui reconnaît et se relie à un reste de *N*-acétyl-D-galactosamine → sérine/thréonine dans un échantillon de sang provenant d'un patient à diagnostiquer ; comparer la valeur ainsi mesurée avec une valeur de seuil ; et estimer que la maladie affectant ledit patient à diagnostiquer n'est pas du cancer de la prostate, mais une maladie bénigne de la prostate, si ladite valeur mesurée de ladite concentration de Tn-MUC1 est supérieure à ladite valeur de seuil ; et

un procédé d'analyse d'informations diagnostiques, qui comprend les étapes de : mesurer la concentration de MUC1 totale dans un échantillon de sang provenant d'un patient à diagnostiquer ; comparer la valeur ainsi mesurée avec une valeur de seuil ; et estimer que

ladite maladie affectant ledit patient à diagnostiquer n'est pas du cancer de la prostate, mais une maladie bénigne de la prostate, si ladite valeur mesurée de ladite concentration de MUC1 totale est supérieure à ladite valeur de seuil.

3. Procédé d'analyse d'informations diagnostiques pour estimer laquelle maladie parmi une maladie bénigne de la prostate et un cancer de la prostate affecte un patient à diagnostiquer, ledit procédé utilisant une combinaison de :

un procédé d'analyse d'informations diagnostiques, qui comprend les étapes de : mesurer la concentration de mucine-1, Tn-MUC1, qui est réactive avec une lectine, qui reconnaît et se relie à un reste de N-acétyl-D-galactosamine → sérine/thréonine dans un échantillon de sang provenant dudit patient à diagnostiquer ; comparer la valeur ainsi mesurée avec une valeur de seuil ; et estimer que la maladie affectant ledit patient à diagnostiquer n'est pas du cancer de la prostate, mais une maladie bénigne de la prostate, si ladite valeur mesurée de ladite concentration de Tn-MUC1 est supérieure à ladite valeur de seuil ; et

des informations diagnostiques concernant un antigène spécifique pour la prostate, qui estiment que ledit patient à diagnostiquer a un cancer de la prostate.

4. Procédé d'analyse d'informations diagnostiques pour estimer laquelle maladie parmi une maladie bénigne de la prostate et un cancer de la prostate affecte un patient à diagnostiquer, ledit procédé utilisant une combinaison de :

un procédé d'analyse d'informations diagnostiques, qui comprend les étapes de : mesurer la concentration de mucine-1, Tn-MUC1, qui est réactive avec une lectine, qui reconnaît et se relie à un reste de N-acétyl-D-galactosamine → sérine/thréonine dans un échantillon de sang provenant dudit patient à diagnostiquer ; comparer la valeur ainsi mesurée avec une valeur de seuil ; et estimer que la maladie affectant ledit patient à diagnostiquer n'est pas du cancer de la prostate, mais ladite maladie bénigne de la prostate, si ladite valeur mesurée de ladite concentration de Tn-MUC1 est supérieure à ladite valeur de seuil ; et

un procédé d'analyse d'informations diagnostiques, qui comprend les étapes de : mesurer la concentration d'un antigène spécifique pour la prostate (LacdiNAc-PSA), qui est réactif avec une lectine, qui reconnaît et se relie à un reste de N-acétyl-D-galactosamine β1-4 N-acétylglucosamine dans ledit échantillon de sang pro-

venant dudit patient à diagnostiquer ; comparer la valeur ainsi mesurée avec une valeur de seuil ; et estimer que ladite maladie affectant ledit patient à diagnostiquer n'est pas ladite maladie bénigne de la prostate, mais du cancer de la prostate, si ladite valeur mesurée de ladite concentration de LacdiNAc-PSA est supérieure à ladite valeur de seuil.

5. Procédé d'analyse d'informations diagnostiques selon l'une quelconque des revendications 1 à 4, dans lequel ladite lectine, qui reconnaît et se relie audit reste de N-acétyl-D-galactosamine → sérine/thréonine est de la lectine *Wisteria floribunda*.

6. Procédé d'analyse d'informations diagnostiques selon l'une quelconque des revendications 1 à 5, dans lequel ladite maladie bénigne de la prostate est de l'hyperplasie prostatique.

7. Procédé d'analyse d'informations diagnostiques selon l'une quelconque des revendications 1 à 6, dans lequel ladite Tn-MUC1 ou ladite MUC1 totale est réactive avec un anticorps monoclonal anti-KL-6.

8. Utilisation d'un kit de diagnostic pour effectuer le procédé d'analyse d'informations diagnostiques selon la revendication 1, ledit kit comprenant :

un anticorps anti-mucine-1 et une lectine, qui reconnaît et se relie à un reste de N-acétyl-D-galactosamine → sérine/thréonine pour quantifier Tn-MUC-1.

9. Utilisation d'un kit de diagnostic pour effectuer le procédé d'analyse d'informations diagnostiques selon la revendication 2, ledit kit comprenant :

un anticorps anti-mucine-1 et une lectine, qui reconnaît et se relie à un reste de N-acétyl-D-galactosamine → sérine/thréonine pour quantifier Tn-MUC-1 ; et un anticorps anti-mucine-1 pour quantifier MUC-1 totale.

10. Kit de diagnostic pour effectuer le procédé d'analyse d'informations diagnostiques selon la revendication 3, ledit kit comprenant :

un anticorps anti-mucine-1 et une lectine, qui reconnaît et se relie à un reste de N-acétyl-D-galactosamine → sérine/thréonine pour quantifier Tn-MUC-1 ; et un anticorps d'antigène spécifique pour l'antiprostata pour quantifier un antigène spécifique pour la prostate.

11. Kit de diagnostic pour effectuer le procédé d'analyse d'informations diagnostiques selon la revendication 4, ledit kit comprenant :
- un anticorps anti-mucine-1 et une lectine, qui reconnaît et se relie à un reste de N-acétyle-D-galactosamine → sérine/thréonine pour quantifier Tn-MUC-1 ; 5
 - et
 - un anticorps d'antigène spécifique pour l'anti-prostate et une lectine, qui reconnaît et se relie à un reste de N-acétyle-D-galactosamine β 1-4 N-acétylglucosamine pour quantifier LacdiNAc-PSA. 10
12. Utilisation du diagnostic selon la revendication 8 ou du kit de diagnostic selon l'une quelconque des revendications 9 à 11, dans lequel ladite lectine, qui reconnaît et se relie audit reste de N-acétyle-D-galactosamine → sérine/thréonine est de la lectine *Wisteria floribunda*. 15 20
13. Utilisation du diagnostic selon la revendication 8 ou du kit de diagnostic selon l'une quelconque des revendications 9 à 12, dans lequel ledit anticorps anti-mucine-1 est un anticorps monoclonal anti-KL-6. 25
14. Utilisation du diagnostic selon la revendication 8 ou du kit de diagnostic selon l'une quelconque des revendications 9 à 13, qui comprend en outre un manuel d'utilisation, qui décrit des informations nécessaires pour effectuer ledit procédé d'analyse d'informations diagnostiques. 30

35


40

45


50

55


[Fig. 1]




[Fig. 2]



[Fig. 3]



[Fig. 4]



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2010090264 A [0007] [0025] [0053] [0075]
- JP 2009535051 A [0007]
- WO 2011135869 A [0007]
- JP 2013526852 A [0007]
- JP 2013076666 A [0007] [0025] [0052]
- WO 2011052244 A [0008]
- WO 2013054281 A [0009]

Non-patent literature cited in the description


- **ATSUSHI MATSUDA et al.** Wisteria floribunda agglutinin-positive mucin 1 is a sensitive biliary marker for human cholangiocarcinoma. *Hepatology*, vol. 52 (1), 174-182 [0010]

专利名称(译)	诊断信息评估方法和套件		
公开(公告)号	EP3088896B1	公开(公告)日	2019-11-06
申请号	EP2013900340	申请日	2013-12-27
[标]申请(专利权)人(译)	柯尼卡株式会社		
申请(专利权)人(译)	柯尼卡美能达, INC.		
当前申请(专利权)人(译)	柯尼卡美能达, INC.		
[标]发明人	YAMASHITA KATSUKO KAYA TAKATOSHI IDE YOUICHI KANEKO TOMONORI		
发明人	YAMASHITA, KATSUKO KAYA, TAKATOSHI IDE, YOUICHI KANEKO, TOMONORI		
IPC分类号	G01N33/68 G01N33/53 G01N33/574		
CPC分类号	G01N33/57434 G01N33/6893 G01N2333/4725 G01N2800/342 G01N2333/42		
其他公开文献	EP3088896A1 EP3088896A4		
外部链接	Espacenet		


摘要(译)

本发明的一个目的是提供：一种新颖的分析诊断信息的方法，通过该方法可以高精度地区分前列腺癌和良性前列腺疾病。以及用于获得诊断信息的诊断套件。根据本发明的分析诊断信息的方法的一个实施方式是分析诊断信息的方法，该方法包括：测量与识别并结合到其上的凝集素反应的粘蛋白-1 (Tn-MUC1) 的浓度。来自待诊断对象的血液样本中的N-乙酰基-D-半乳糖胺丝氨酸 (苏氨酸) 残基；将如此测量的值与阈值进行比较；本发明的诊断试剂盒的一个实施方式是，当mucin-1的浓度的测量值大于阈值时，估计影响被诊断对象的疾病不是前列腺癌而是良性前列腺疾病。本发明是一种诊断试剂盒，其包括：用于定量Tn-MUC1的抗粘蛋白-1抗体；和用于检测的凝集素，该凝集素识别并结合N-乙酰基-D-半乳糖胺+'丝氨酸 (苏氨酸) 残基。

[Fig. 1]



[Fig. 2]



[Fig. 3]

