



EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

06.05.2020 Bulletin 2020/19

(21) Application number: **14884789.0**

(22) Date of filing: **04.12.2014**

(51) Int Cl.:

C07K 16/28 ^(2006.01) **A61K 39/395** ^(2006.01)
A61K 49/00 ^(2006.01) **C07K 16/46** ^(2006.01)
C07K 17/00 ^(2006.01) **C12N 15/13** ^(2006.01)
G01N 33/53 ^(2006.01)

(86) International application number:

PCT/CA2014/000861

(87) International publication number:

WO 2015/131257 (11.09.2015 Gazette 2015/36)

(54) **INSULIN-LIKE GROWTH FACTOR 1 RECEPTOR -SPECIFIC ANTIBODIES AND USES THEREOF**

FÜR INSULINÄHNLICHEN WACHSTUMSFAKTOR-1-REZEPTOR SPEZIFISCHE ANTIKÖRPER
UND VERWENDUNGEN DAVON

ANTICORPS SPÉCIFIQUES DU RÉCEPTEUR DU FACTEUR DE CROISSANCE 1 ANALOGUES À
L'INSULINE ET LEURS UTILISATIONS

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **06.03.2014 US 201461948818 P**

(43) Date of publication of application:

11.01.2017 Bulletin 2017/02

(73) Proprietor: **National Research Council of Canada
Ottawa, Ontario K1A 0R6 (CA)**

(72) Inventors:

- **STANIMIROVIC, Danica**
Ottawa, Ontario K4P 1R7 (CA)
- **KEMMERICH, Kristin**
Ottawa, Ontario K1Y 0A5 (CA)
- **HAQQANI, Arsalan S.**
Kanata, Ontario K2T 0A8 (CA)
- **SULEA, Traian**
Kirkland, Quebec H9J 3A2 (CA)
- **ARBABI-GHAHROUDI, Mehdi**
Ottawa, Ontario K1W 1K1 (CA)
- **MASSIE, Bernard**
Laval, Quebec H7H 1R2 (CA)
- **GILBERT, Rénald**
Montreal, Quebec H4V 2B4 (CA)

(74) Representative: **Fleuchaus, Andrea et al**

Fleuchaus & Gallo Partnerschaft mbB
Patentanwälte
Steinerstraße 15/Haus A
81369 München (DE)

(56) References cited:

WO-A1-2011/044542 WO-A1-2011/150061
US-A1- 2005 085 419

- **BURTRUM D ET AL:** "A fully human monoclonal antibody to the insulin-like growth factor I receptor blocks ligand-dependent signaling and inhibits human tumor growth in vivo", **CANCER RESEARCH, AMERICAN ASSOCIATION FOR CANCER RESEARCH, US**, vol. 63, no. 24, 15 December 2003 (2003-12-15), pages 8912-8921, XP002316542, ISSN: 0008-5472
- **GABATHULER R:** 'Approaches to transport therapeutic drugs across the blood-brain barrier to treat brain diseases.' **NEUROBIOL DIS.** vol. 37, no. 1, January 2010, ISSN 0969-9961 pages 48 - 57, XP026765785
- **WEBSTER CARL I ET AL:** "Brain penetration, target engagement, and disposition of the blood-brain barrier-crossing bispecific antibody antagonist of metabotropic glutamate receptor type 1", **FASEB JOURNAL**, vol. 30, no. 5, May 2016 (2016-05), pages 1927-1940,

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

DescriptionFIELD OF THE INVENTION

[0001] The present invention relates to Insulin-Like Growth Factor 1 Receptor-specific antibodies, fragments thereof, and uses thereof. More specifically, the present invention relates to Insulin-Like Growth Factor 1 Receptor-specific antibodies and fragments thereof that transmigrate the blood-brain barrier, and uses thereof.

BACKGROUND OF THE INVENTION

[0002] Neurodegenerative diseases, such as Alzheimer's and Parkinson's disease, are an increasing burden on our ageing society because there are currently no effective treatments for these disabling conditions. Treatment as well as early diagnosis of these and other diseases that originate in the brain remain challenging because the majority of suitable therapeutic molecules and diagnostics cannot penetrate the tight and highly restrictive blood-brain barrier (BBB) (Abbott, 2013). The BBB constitutes a physical barricade that is formed by brain endothelial cells (BECs) that line the blood vessels and connect with each other through tight junctions (Abbott, 2013). The tight junctions formed between the BECs are essential for the integrity of the BBB and prevent the paracellular transport of molecules larger than 500 daltons (Da). Because brain endothelial cells exhibit very low pinocytosis rates (Abbott, 2013), transcellular transport of larger molecules is limited to the highly specific receptor mediated transcytosis (RMT) pathway, and the passive, charge-based adsorption mediated transcytosis (Abbott, 2013; Pardridge, 2002). Additionally, the high density of efflux pumps, such as P-glycoprotein or the multi-drug resistance protein -1 (MDR-1), contribute to the removal of unwanted substances from the brain (Abbott, 2013).

[0003] While all these characteristics protect the brain from pathogens and toxins, they equally prevent the entry of most therapeutics. In fact, less than 5% of small molecule therapeutics and virtually none of the larger therapeutics can cross the BBB in pharmacologically relevant concentrations (i.e., sufficient to engage a central nervous system (CNS) target and elicit pharmacologic/therapeutic response) unless they are specifically 'ferried', that is, coupled to a transporter molecule.

[0004] Due to the lack of effective 'carriers' to transport molecules across the BBB, numerous drugs against neurodegenerative diseases have been 'shelved' or eliminated from further development as they cannot be delivered to the brain in sufficient amount.

[0005] Different approaches to deliver larger molecules into the brain have been explored. For example, the integrity of the BBB may be disrupted, resulting in a leaky BBB, which in turn allows for unrestricted, paracellular entry of larger molecules into the brain. Tight junctions can be successfully loosened or disrupted by various approaches. For example, injection of substances that induce osmotic shock (for example, mannitol, hypertonic solutions) into the blood stream causes cell shrinkage and results in the disruption of tight junctions, therefore severely compromising the BBB (Guillaume, 2010). Other modulators of tight junctions include alkylglycerols, bradykinin and several analogues thereof, as well as viruses that modulate expression of proteins involved in maintaining the tight junctions (Erdlenbruch et al., 2003; Preston et al., 2008; Gan et al., 2013). A more localized disruption of the BBB is possible through application of ultrasound (Nhan et al., 2013).

[0006] Recent scientific studies refer e.g. also to potential cellular receptors, specific antibodies thereto and their capacity to interfere with the BBB. For example, Burtum et al., 2003, Cancer Research 63, 8812-8921, US 2005/0085419 A1 and WO 2011/044542 A1 refer to antibody-receptor mediated endocytosis and potential treatments, further WO 2011/150061 A1 refers to a peptide-receptor mediated endocytosis and potential treatments.

[0007] Additionally, Gabathuler R. (2010, Neurobiology of Diseases 37, p48-57) reviews further potential approaches - iter alia - the use of single domain antibodies targeting the CDC50A for a non-invasive BBB transmigration.

[0008] However, the periods during which the BBB is disrupted are sufficient to alter brain homeostasis and allow harmful chemicals, toxins and pathogens to enter the brain; this can result in serious side-effects, e.g., seizures and brain swelling, infection and possibly permanent neuropathological changes. As would be evident to those of skill in the art, repeated treatments with these techniques for chronic and diffuse brain diseases affecting multiple brain regions are not practical. Most of these treatments are costly, necessitate hospitalisation, and some approaches require anesthesia.

[0009] Another approach for circumventing the BBB is direct injection of therapeutic molecules into the cerebrospinal fluid (CSF), the parenchymal space, or other parts of the brain. Several delivery methods have been developed, including: intracerebral (intra-parenchymal), intraventricular, and intrathecal delivery via infusion or convection-enhanced diffusion (CED) pumps. However, any type of direct injection into the brain or intracerebral implant is an invasive and costly procedure, as it requires hospitalization, anesthesia, and often surgery. Moreover, the poor diffusion rates of the therapeutics, particularly large biologics, within brain parenchyma limit the penetration of therapeutics to only small areas surrounding the site of injection/implantation. The correct placement of injections, catheters, and implants is challenging

yet crucial to achieve diffusion of the drug to the targeted region of the brain. Additionally, catheters and implants provide a site for infection and/or immune response against the foreign material.

[0010] In another attempt to increase delivery across the BBB, CNS drugs have been modified to increase their brain uptake. Such modifications can include a change of their surface charge, a reduction in molecule size, and change to the lipophilicity of the drugs. However, any modifications to increase brain penetration are also likely to alter the overall pharmacology of the drug, including its desired activity and/or specificity. In addition, lipophilic molecules are more prone to being exported from the brain through the P-glycoprotein efflux pump.

[0011] Finally, endogenous transport mechanisms across the BBB have been exploited. Physiological mechanisms that allow transport of large molecules across the BBB can be divided into the highly specific receptor mediated transcytosis (RMT) and the non-specific charge based adsorptive mediated endocytosis pathways. Endocytosis is triggered upon binding of the specific ligand to its receptor, or upon electrostatic interaction between the cationic ligand or drug and the anionic functional groups on the brain endothelial cell surface (luminal side), respectively. Subsequently, the newly formed endosome is transcytosed across the cell to the abluminal side, to release its cargo.

[0012] Because adsorptive mediated transcytosis is non-specific, charge-mediated interaction, it occurs in all vascular beds and organs, limiting the availability of drug for brain delivery. Therefore, exploiting the RMT pathway remains the only physiological, non-invasive yet highly receptor-specific brain delivery method.

[0013] Only a few receptors are presently known to undergo RMT at the BBB and 'ferry' across their natural ligands: the well-studied transferrin receptor (TfR), the insulin receptor (IR), low-density lipoprotein receptor related proteins 1 and 2 (LRP-1 and -2), diphtheria toxin receptor, and TMEM30A. Peptides, natural ligands, and antibodies or antibody fragments have been developed that bind to these receptors (Pardridge et al., 1991; Yu et al., 2011; Muruganandam et al., 2001; Abulrob et al., 2005; Demeule, 2008; Sumbria et al., 2013), functioning as drug-to-brain transporters that utilize endogenous RMT pathways. However, to date only a single peptide (Angiopep ANG1005, targeting LRP-1) has been analyzed in phase I clinical studies, while other candidates are being studied in laboratory settings. The RMT pathway appears to be the most promising pathway for drug transport to the brain, but current approaches have limitations, including: non-selective expression of the target receptor at the BBB, competition between the carrier and the natural ligands to the receptor, ineffective transcytosis of a receptor as well as lysosomal degradation of endocytosed carriers (Xiao and Gun, 2013).

[0014] The lack of high-capacity and high-selectivity BBB carriers delays the development of new therapeutics and diagnostics for diseases originating in the brain, including brain tumors and neurodegenerative diseases. There is clearly a need for a non-invasive method to deliver small and large therapeutic and diagnostic molecules in pharmacologically efficacious doses into the brain without disrupting the physiology and homeostasis of the BBB.

SUMMARY OF THE INVENTION

[0015] The present invention relates to Insulin-Like Growth Factor 1 Receptor (IGF1R) -specific antibodies and uses thereof. More specifically, the present invention relates to Insulin-Like Growth Factor 1 Receptor-specific antibodies and fragments thereof that transmigrate the blood-brain barrier, and uses thereof.

[0016] The present invention further provides isolated or purified antibodies, comprising

a complementarity determining region (CDR) 1 sequence of GGTVSPTA (SEQ ID NO:1);

a CDR2 sequence of ITWSRGTT (SEQ ID NO:2); and

a CDR3 sequence of AASTFLRILPEESAYTY (SEQ ID NO:3),

wherein the antibody specifically binds to the Insulin-Like Growth Factor 1 Receptor (IGF1R) and transmigrates the blood-brain barrier.

[0017] For example, and without wishing to be limiting in any manner, the isolated or purified antibody specific for IGF1R may be $X_1VX_2LX_3ESGGGLVQX_4GGSRLRLSCX_5X_6SGGTVSPTAMGWX_7RQAPGKX_8X_9EX_{10}VX_{11}HITWSRGTTX_{12}ASSVKX_{13}RFTISRDX_{14}X_{15}KNTX_{16}YLMNSLX_{17}X_{18}EDTAVYYCAAS\ TFLRILPEESAYTYWGQGT$ $X_{19}VTVSS$ (SEQ ID NO:4), where X_1 is E or Q; X_2 is K or Q; X_3 is V or E; X_4 is A or P; X_5 is A or E; X_6 is V or A; X_7 is V or F; X_8 is G or E; X_9 is L or R; X_{10} is F or W; X_{11} is G or S; X_{12} is V or Y; X_{13} is D or G; X_{14} is N or S; X_{15} is A or S; X_{16} is L or V; X_{17} is K or R; X_{18} is A or S; and X_{19} is L or Q. In more specific, non-limiting examples, the isolated or purified antibody may comprise a sequence selected from the group consisting of:

QVKLEESGGGLVQAGGSLRLSCEVSGGTVSPTAMGWFRQAPGKEREFVGH ITWSR
GTRVASSVKDRFTISRDSAKNTVYLQMNSLKSEDTAVYYCAASTFLRILPEESAYTY WGQGTQVTVSS (SEQ ID
NO:5), referred to herein as IGF1R-4;

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWVRQAPGKGLEWVGHITWSR
 GTTRYASSVKGRFTISRDN SKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTY WGQGT LVT VSS (SEQ ID
 NO:6), referred to herein as IGF1R-4_H2;

5 QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKGLEFVGHITWSR
 GTTRYASSVKGRFTISRDN SKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTY WGQGT LVT VSS (SEQ ID
 NO:7), referred to herein as IGF1R-4_H3;

10 QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKGLEFVGHITWSR
 GTTRYASSVKGRFTISRDN SKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTY WGQGT LVT VSS (SEQ ID
 NO:8), referred to herein as IGF1R-4_H4;

15 QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKEREFGHITWSR
 GTTRYASSVKGRFTISRDN SKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTY WGQGT LVT VSS (SEQ ID
 NO:9), referred to herein as IGF1R-4_H5; and

or a sequence having 90% identity thereto.

[0018] The isolated or purified antibody as described above is a single-domain antibody (sdAb) also named immunoglobulin single variable domain antibody; the sdAb may be of camelid origin.

20 **[0019]** The isolated or purified antibody of the present invention may be presented in a multivalent display format. In a multivalent display format, the antibody may be linked to a Fc fragment; the Fc fragment is the mouse Fc2b or human Fc1. For example, and without wishing to be limiting in any manner, the isolated or purified antibody in multivalent display may comprise the sequence of SEQ ID NO:10 (referred to herein as IGF1R-4 consensus-Fc fusion), SEQ ID NO:39 (referred to herein as Fc-IGF1R-4 consensus fusion), or 11 (referred to herein as IGF1R-4-Fc fusion).

25 **[0020]** The present invention also provides a nucleic acid molecule encoding the isolated or purified antibody as described herein. A vector comprising the nucleic acid molecule as just described is also provided.

[0021] The isolated or purified antibody as described herein may be immobilized onto a surface.

30 **[0022]** The present invention further provides the isolated or purified antibody as described herein linked to a cargo molecule; the cargo molecule may have a molecular weight in the range of about 1kD to about 200 kDa. The cargo molecule linked to the antibody may be a detectable agent, a therapeutic, a drug, a peptide, a growth factor, a cytokine, a receptor trap, a chemical compound, a carbohydrate moiety, an enzyme, an antibody or fragments thereof, a DNA-based molecule, a viral vector, or a cytotoxic agent; one or more liposomes or nanocarriers loaded with a detectable agent, a therapeutic, a drug, a peptide, an enzyme, and antibody or fragments thereof, a DNA-based molecule, a viral vector, or a cytotoxic agent; or one or more nanoparticle, nanowire, nanotube, or quantum dots.

35 **[0023]** Additionally, the present invention provides a composition comprising one or more than one isolated or purified antibody as described herein and a pharmaceutically-acceptable carrier, diluent, or excipient.

[0024] An *in vitro* method of detecting IGF1R is also provided, the method comprising

40 a) contacting a tissue sample with one or more than one isolated or purified antibody of as described herein linked to a detectable agent; and

b) detecting the detectable agent detecting the detectable agent linked to the antibody bound to IGF1R in the tissue sample.

45 **[0025]** In the method described above, the sample may be a serum sample, a vascular tissue sample, tumour tissue sample, or a brain tissue sample from a human or animal subject. In the method as described, the step of detecting (step b)) may be performed using optical imaging, immunohistochemistry, molecular diagnostic imaging, ELISA, imaging mass spectrometry, or other suitable method.

[0026] Further provided is an *in vivo* method of detecting IGF1R expression in a subject, the method comprising:

50 a) administering one or more than one isolated or purified antibody as described herein linked to a detectable agent to the subject; and

55 b) detecting the detectable agent linked to the antibody bound to IGF1R.

[0027] In the method described above, the step of detecting (step b)) may be performed using PET (positron emission tomography), SPECT (single-photon emission computed tomography), fluorescence imaging, or any other suitable method.

[0028] Presently provided is a method of transporting a molecule of interest across the blood-brain barrier (BBB), the method comprising:

- a) administering one or more than one isolated or purified antibody as described herein linked to the molecule of interest to a subject, where the antibody transmigrates the blood-brain barrier,

wherein the one or more than one antibody ferries the molecule of interest across the BBB. In the method as just described, the molecule of interest may have a molecular weight in the range of about 1 kD to about 200 kDa; the molecule of interest may be a detectable agent, a therapeutic, a drug, a peptide, a growth factor, a cytokine, a receptor trap, a chemical compound, a carbohydrate moiety, an enzyme, an antibody or fragment thereof, a DNA-based molecule, a viral vector, or a cytotoxic agent; one or more liposomes or nanocarriers loaded with a detectable agent, a therapeutic, a drug, a peptide, an enzyme, an antibody or fragment thereof, or a cytotoxic agent; or one or more nanoparticle, nanowire, nanotube, or quantum dots. In the method as described, the administration may be intravenous (iv), subcutaneous (sc), or intramuscular (im).

[0029] The disclosure also encompasses a method of quantifying an amount of a cargo molecule delivered across the BBB of a subject, wherein the cargo molecule is linked to one or more than one isolated or purified antibody as described herein, the method comprising

- a) collecting cerebrospinal fluid (CSF) from the subject; and

- b) using targeted proteomics methods to quantify the amount of the cargo molecule linked to one or more than one isolated or purified antibody or fragment in the CSF.

[0030] The cargo molecule may be any desired molecule, including the cargo molecules previously described; the antibody transmigrates the BBB; and the molecule may be "linked" to the antibody or fragment thereof, as previously described. In the above method, the CSF is collected from a subject using any suitable method known in the art. The amount of CSF required for targeted proteomics method in step b) may be between about 1 to 10 μ l. The targeted proteomics methods used to quantify the amount of the one or more than antibody linked to the cargo molecule may be any suitable method known in the art. For example and without wishing to be limiting, the targeted proteomics method may be a mass spectrometry method, such as multiple reaction monitoring - isotope labeled internal standards (MRM - ILIS).

[0031] The poor delivery of diagnostics or drugs across the tight and highly selective BBB compromises the development of treatments for brain diseases, such as, but not limiting to, brain tumors and neurodegenerative diseases. The lack of carriers to transport molecules across the BBB delays the development of new therapeutics and diagnostics for such diseases. As described herein, an IGF1 R-binding V_HH has been produced that provides an effective transport platform for delivery of drugs conjugated to the antibody across the BBB to their targets in the brain. The presently described antibody exploits the natural RMT pathway of the IGF1R from the luminal to abluminal side of the BBB-forming brain endothelial cells. Following binding of the antibody to IGF1R, RMT is initiated and the antibody, together with a conjugated molecule (cargo), is transcytosed through the cell to the abluminal side where they are both released into the brain microenvironment. The anti-IGF1R V_HH was confirmed to bind to IGF1R (Fig. 3C), internalize into BBB cells (Fig. 4), and cross to abluminal side of the *in vitro* BBB model (Fig. 6B). Drug-to-brain delivery studies *in vivo* also showed that the IGF1R V_HH 'carried' a conjugated peptide (Galanin; about 3kDa) as well as a large protein fusion (about 80kDa) across the BBB (Fig. 9A and B; Fig. 9C).

[0032] The results also show that the anti-IGF1R V_HH can be expressed in fusion with Fc (fragment crystallisable) fragment to prolong circulation half-life by about 75 fold (about 25h compared to about 20 min for V_HH alone). This high molecular weight fusion construct (about 80kDa) is also efficiently transported across the BBB. The long plasma half-life increases CSF exposure of the IGF1R V_HH -mFc (mFc = mouse Fc) conjugate significantly compared to the V_HH alone and is useful as a BBB delivery carrier for the treatment of chronic diseases with targets in the CNS. The conjugate is readily detected in brain parenchyma using immunofluorescence detection. The results demonstrate that the IGF1R V_HH carrier can "ferry" large molecules (similar in size to: antibodies, enzymes, growth factors, peptides, cytokines, receptor traps) across the BBB.

[0033] Thus, the antibody-delivery may not only be useful for short term treatment (e.g. epileptic seizure), but may also be useful for medium-term (e.g. cancer) and long-term (e.g. Alzheimer's or Parkinson's disease) treatments.

BRIEF DESCRIPTION OF THE DRAWINGS

[0034] These and other features of the invention will now be described by way of example, with reference to the appended drawings, wherein:

FIGURE 1 is a schematic diagram of the insulin-like growth factor 1 receptor (IGF1R). The IGF1R is found on the cell surface, and comprises two subunits: the alpha subunit and the beta subunit. The alpha subunit (comprising an extracellular part with the insulin-like growth factor 1 binding site) is connected by a disulphide bond to the beta subunit (comprising a small extracellular domain, a transmembrane region, and an intracellular portion). The IGF1 receptor can form a dimer. A 933 amino acid long fragment comprising the alpha subunit and the extracellular portion of the beta subunit, as indicated within the grey box, (M1 - F933, SwissProt Accession No. P08069; see FIGURE 2) was recombinantly produced and used for immunization of a llama.

FIGURE 2 shows the sequence of IGF1R (SwissProt Accession No. P08069). The 933 amino acid long protein fragment used for immunization and panning is shown in bold; the full ectodomain is 2 amino acids longer. The furin cleavage site, separating alpha and beta subunits, is shown in italicized lowercase letters. The signal peptide is shown in bold italics.

FIGURE 3A shows a size exclusion chromatogram of the IGF1R-binding V_HH IGF1R-4 and its humanized variants (H1, H2, H3, H4, H5, H6) run through a Superdex 75 column. The profile suggests that all the V_HHs except IGF1R H1 are monomeric and non-aggregating. FIGURE 3B shows the melting temperature (T_m) as determined by circular dichroism (CD) for IGF1R-4 V_HH and its humanized variants (H2, H3, H4, H5, H6). The proteins were heated to above 90°C and measurements were taken in the CD instrument to determine the melting curve (top curve) and the T_m. Subsequently, the proteins were cooled to room temperature, heated once more and analysed by CD (bottom curve for IGF1 R-4 VHH; line or dots around 0% refolded for humanized variants). This allowed the determination of the fraction of refolded protein, which was zero for humanized versions and 80% for IGF1R-4. FIGURE 3C shows a surface plasmon resonance (SPR) sensogram overlay for binding of 0.1 - 5 nM IGF1R-4 V_HH and its humanized variants (H2, H3, H4, H5, H6) to the recombinant extracellular portion of the human IGF1R fragment. The data fit well to a 1:1 model. FIGURE 3D shows SPR sensograms of IGF1R-4 V_HH binding to the recombinant extracellular portion of the human IGF1R fragment in presence of IGF1. IGF1 in 100-fold excess did not affect IGF1R-4 V_HH binding, showing that both bind to different epitopes on the receptor. FIGURE 3E shows that IGF1R-4 V_HH did not bind to the recombinant extracellular portion of the human Insulin Receptor (IR) immobilized on SPR surface, while it did bind to the control surface of human IGF1R.

FIGURE 4 shows imaging results of cell uptake of Cy-5.5-labeled IGF1 R-4 V_HH and control V_HH. Cy5.5-labeled IGF1R-4 V_HH (or FC5 V_HH as a positive control; Muruganandam et al., 2002; Haqqani et al., 2012) were incubated with SV40 immortalized rat brain endothelial cells (svARBEC) at 4°C (top panels) or at 37°C (bottom panels) to assess whether IGF1R-4 is internalized passively (4°C) or through active mechanisms (37°C) such as receptor mediated endocytosis. Co-staining with wheat germ agglutinin and DAPI was carried out to visualize the cell surface and the nucleus, respectively. Top panel: When incubated at 4°C, IGF1R-4 and FC5 V_HH were found outside the cells (arrowheads), bound to cell membrane (co-localized with the wheat germ agglutinin). Bottom panel: At 37°C both FC5 and IGF1R-4 V_HH accumulated inside the cells in vesicle-like structures (arrowheads), likely endosomes, suggesting internalization through an active transport mechanism.

FIGURE 5A shows the sequence for the C-terminal fusion of IGF1R-4 V_HH with a murine Fc fragment (IGF1R-4-mFc). A schematic representation of the assembled fusion protein is shown in FIGURE 5B. The IGF1R-4 V_HH is shown in black and the murine Fc (CH2 and CH3) is shown in grey in FIGURE 5A and B.

FIGURE 6A is a flowchart summarizing the use of the *in vitro* BBB model to assess the ability of various V_HHs to cross the BBB. Equimolar amounts (5.6 µM or 1.25 µM) of positive (FC5) and negative control (A20.1, a *Clostridium difficile* toxin A-binding V_HH; and EG2, an EGFR binding V_HH) V_HH and IGF1R-4 were tested simultaneously for their ability to cross a rat *in vitro* BBB model. SV40-immortalized brain endothelial cells from adult rat (svARBECs) are grown in a monolayer on the membrane of an insert in the presence of rat astrocyte-conditioned medium in the bottom chamber and standard medium in the top chamber. Following co-addition of equimolar amounts of the various V_HH to the luminal side of the BBB model, samples were taken from the bottom chamber after 15, 30, and 60 min. The concentrations of each V_HH were then quantified by mass spectrometry (multiple reaction monitoring - isotope labeled internal standards; MRM-ILIS) in these samples. The P_{app} value [Q_r/dt = cumulative amount in the receiver compartment versus time; A = area of the cell monolayer; C₀ = initial concentration of the dosing solution] is commonly used to determine the ability of a molecule to cross the BBB. FIGURE 6B shows the P_{app} values of the four co-administered V_HHs. IGF1R-4 V_HH has a significantly higher P_{app} value than FC5, while both negative controls have low P_{app} values, indicating facilitated transport of FC5 and IGF1R-4 V_HH compared to low non-specific transport or paracellular transport of control V_HHs. FIGURE 6C shows the P_{app} values of the humanized IGF1R-4 single-domain antibodies (H2, H3, H4, H5, H6) (black bars), compared to A20.1 V_HH (grey bars), which was tested as a control in

the same well (the average A20.1 value is indicated by a grey dotted line). The result clearly indicates that simply binding of IGF1R (as seen for the humanized 6 variant (H6)) is not sufficient to trigger transmigration of the BBB. FIGURE 6D shows the P_{app} values of the C-terminal fusion of IGF1R-4 V_{HH} to mouse Fc (black), compared to the A20.1-mouse Fc (A20.1mFc) (grey) which was tested in the same well and has a very low P_{app} value. IGF1R-4-mFc has a significantly higher P_{app} value than FC5-mFc (typically around 180 cm/min, as indicated by the grey dotted line). Another IGF1 R-binding V_{HH} , IGF1R-1 in fusion with Fc has only marginally higher P_{app} value than the internal negative control A20.1 V_{HH} . This result as well demonstrates that not all IGF1R-binding antibodies trans-migrate the BBB. The data further shows that IGF1R-4-mFc has a similar high rate of transport across the BBB compared to the monovalent IGF1R-4 V_{HH} . The results are average P_{app} values obtained in 3-6 independent experiments. Average P_{app} values of FC5 (V_{HH}) and A20.1 (V_{HH}) are also indicated by dotted lines for comparison.

FIGURE 7 (upper panels) shows *in vivo* whole body images of CD-1 mice injected intravenously with PBS (naive) or 2.5 mg/kg of either Cy5.5-IGF1R-4 V_{HH} or Cy5.5-A20.1 V_{HH} (schematic of injected molecules is shown on top) and scanned in eXplore Optix time-domain optical imager 30 min after injection. Bottom panels show ex-vivo brain images of the same animals obtained 1 h after V_{HH} injection and after transcardial perfusion to remove antibodies from the circulation. Arrows indicate high optical signal (light gray) of the fluorescent tracer in the head and in ex-vivo brain in the animal injected with Cy5.5-IGF1R-4, whereas it cannot be detected in the animal injected with the Cy5.5-A20.1 control (black).

FIGURE 8A shows the scheme for chemical synthesis of IGF1R-4 V_{HH} -Galanin conjugate. IGF1R-4 was first conjugated to the NHS group of a Sulfo-SMCC cross-linker (1); then maleimide-activated IGF1 R-4-sulfo-SMCC was conjugated to the reduced cysteine of Galanin (2). FIGURE 8B shows SDS-PAGE gel of IGF1R-4 (lane 2), IGF1R-4-SMCC (lane 3), and IGF1 R-4-Galanin conjugate (lane 4). The 'banding' pattern indicates the attachment of 1-2 galanin molecules per IGF1R-4. The same conjugation methods were used to link Galanin with IGF1 R-mFc.

FIGURE 9 shows IGF1 R-4-mediated brain delivery of the chemically conjugated peptide Galanin. FIGURE 9A is a graph showing the ability of IGF1R-4 to deliver pharmacologically efficacious doses of the analgesic peptide Galanin (3.2 kD) into the brain using Hargreaves pain model. In this model, localised chronic pain is induced in male Wistar rats (4-6 weeks age), by injecting 100 μ l of complete Freund's adjuvant (CFA) into the left plantar surface, causing a local inflammation within a few hours. Following tail vein injection of the BBB carrier V_{HH} -drug conjugate or Galanin alone, the rats are placed into Plexiglas enclosures set on a glass surface. A thermal stimulus is focused on the inflamed or contralateral paw via an angled mirror. The latency between stimulus application and paw withdrawal (lick or flick of paw) is interpreted as a measure of the analgesic effect (inhibition of thermal hyperalgesia).. The peptide Galanin alone cannot penetrate the BBB, as demonstrated by the lack of analgesic effect after systemic (tail-vein) injection of 1 mg/kg Galanin (solid black triangles). Systemic injection of IGF1 R-4-Galanin conjugate (3 mg/kg) inhibits thermal hyperalgesia, reaching maximal reversal 30 min after administration (solid black circles); the effect was more pronounced than that observed with the 6mg/kg of FC5-Galanin conjugate (white open squares). FIGURE 9B shows these results as area under the curve (AUC) of the response compared to the maximal possible effect (MPE; control paw). FIGURE 9C shows dose-dependent (2 mg/kg and 5 mg/kg) inhibition of thermal hyperalgesia in Hargreaves model of inflammatory pain with IGF1 R-4-mFc-Galanin conjugate administered iv. In contrast, A20.1-mFc-Galanin (5 mg/kg) did not induce any reversal of thermal hyperalgesia. FIGURE 9D shows percent reversal of thermal hyperalgesia (E_{max}) at the peak of response achieved with various doses of A20.1-mFc-Galanin and IGF1 R-4-mFc-Galanin.

FIGURE 10 shows plasma and CSF levels of IGF1R-4, and 'positive' control V_{HH} , FC5 and 'negative' control V_{HH} A20.1 30 min after systemic (tail vein) co-administration of 6 mg/kg of each antibody. The CSF was collected from cisterna magna. Plasma and CSF levels of IGF1R-4, FC5 and A20.1 were determined using the MRM-ILIS method that 'tracks' and quantifies specific protein peptide signatures. Albumin levels in the CSF were concurrently determined by MRM. All CSF samples having a plasma/CSF ratio lower than 1500 were excluded as potentially blood-contaminated. The CSF/serum ratio for IGF1R-4 was 1.2% compared to 0.9% for FC5 and 0.017% for A20.1.

FIGURE 11 shows immuno-detection of IGF1R-4mFc in brain sections 24 h after tail-vein administration of a 6 mg/kg dose. Sacrifice perfusion with PBS was carried out on the rats and brain sections (12 μ m) were obtained using the vibratome. IGF1R-4mFc was immuno-detected using an anti-mouse Fc antibody (red; red channel only shown in inserts). Blood vessels in the brain section (dentate gyrus, A; frontal cortex, B) were detected using lectin RCA1 (green). IGF1R-4mFc could be detected in both vessels and outside the vessels (i.e. in the brain parenchyma, transmigrated across the BBB) as indicated by arrowheads.

FIGURE 12 shows that IGF1R-4 does not interfere with insulin or IGF-1 signaling through either the insulin receptor or IGF1R. FIGURE 12A is a representative Western blot showing that neither IGF1 R-4, nor any of the other anti-IGF1 R V_HH tested (IGF1 R-1, -3, -5, or -6) induce downstream Akt phosphorylation alone at a concentration of 100nM. Neither does the presence of 100nM of IGF1 R-4, or any of the other anti-IGF1 R V_HHs inhibit Akt phosphorylation as induced by 10 μ g/ml of insulin. The quantitation of Western blot band densities from 3 independent experiments is shown in the bar graph (average $\pm$ SD) below the gel image. FIGURE 12B is a representative Western blot showing that neither IGF1 R-4, nor any of the other anti-IGF1 R V_HH tested (IGF1 R, -3, -5, or -6) at 100nM induce phosphorylation of Akt on their own and neither inhibit IGF-1 induced Akt phosphorylation (i.e. signaling) induced upon stimulation with 200ng/ml of IGF-1. The quantitation of Western blot band densities from 3 independent experiments is shown in the bar graph (average $\pm$ SD) below the gel image. FIGURE 12C shows Western blots probed for phosphorylated IGF1R. Cells were incubated with either 100nM or 500nM IGF1 R-4-mFc, or any of the other anti-IGF1R V_HH-mFc fusions (IGF1R-1, -3, or -4 -mFc) alone or stimulated with 200ng/ml IGF-1 in presence of the respective IGF1R-V_HH-mFc fusion proteins. The Western blots indicate that none of the fusion constructs inhibited IGF-1 induced phosphorylation of IGF1R and neither induced receptor phosphorylation on their own.

DETAILED DESCRIPTION OF THE INVENTION

[0035] The present invention relates to Insulin-Like Growth Factor 1 Receptor-specific antibodies, fragments thereof, and uses thereof. More specifically, the present invention relates to Insulin-Like Growth Factor 1 Receptor-specific antibodies or fragments thereof that transigrate the blood-brain barrier, and uses thereof.

[0036] The present invention provides an isolated or purified antibody, comprising

a complementarity determining region (CDR) 1 sequence of GGTVSPTA (SEQ ID NO:1);

a CDR2 sequence of ITWSRGTT (SEQ ID NO:2); and

a CDR3 sequence of AASTFLRLPEESAYTY (SEQ ID NO:3),

wherein the antibody specifically binds to the Insulin-Like Growth Factor 1 Receptor (IGF1R).

[0037] The term "antibody", also referred to in the art as "immunoglobulin" (Ig), as used herein refers to a protein constructed from paired heavy and light polypeptide chains; various Ig isotypes exist, including IgA, IgD, IgE, IgG, and IgM. When an antibody is correctly folded, each chain folds into a number of distinct globular domains joined by more linear polypeptide sequences. For example, the immunoglobulin light chain folds into a variable (V_L) and a constant (C_L) domain, while the heavy chain folds into a variable (V_H) and three constant (C_{H1}, C_{H2}, C_{H3}) domains. Interaction of the heavy and light chain variable domains (V_H and V_L) results in the formation of an antigen binding region (Fv). Each domain has a well-established structure familiar to those of skill in the art.

[0038] The light and heavy chain variable regions are responsible for binding the target antigen and can therefore show significant sequence diversity between antibodies. The constant regions show less sequence diversity, and are responsible for binding a number of natural proteins to elicit important biochemical events. The variable region of an antibody contains the antigen-binding determinants of the molecule, and thus determines the specificity of an antibody for its target antigen. The majority of sequence variability occurs in six hypervariable regions, three each per variable heavy (V_H) and light (V_L) chain; the hypervariable regions combine to form the antigen-binding site, and contribute to binding and recognition of an antigenic determinant. The specificity and affinity of an antibody for its antigen is determined by the structure of the hypervariable regions, as well as their size, shape, and chemistry of the surface they present to the antigen. Various schemes exist for identification of the regions of hypervariability, the two most common being those of Kabat and of Chothia and Lesk. Kabat et al (1991) define the "complementarity-determining regions" (CDR) based on sequence variability at the antigen-binding regions of the V_H and V_L domains. Chothia and Lesk (1987) define the "hypervariable loops" (H or L) based on the location of the structural loop regions in the V_H and V_L domains. As these individual schemes define CDR and hypervariable loop regions that are adjacent or overlapping, those of skill in the antibody art often utilize the terms "CDR" and "hypervariable loop" interchangeably, and they may be so used herein. The CDR/loops are referred to herein according to the more recent IMGT numbering system (Lefranc, M.-P. et al., 2003), which was developed to facilitate comparison of variable domains. In this system, conserved amino acids (such as Cys23, Trp41, Cys104, Phe/Trp118, and a hydrophobic residue at position 89) always have the same position. Additionally, a standardized delimitation of the framework regions (FR1: positions 1 to 26; FR2: 39 to 55; FR3: 66 to 104; and FR4: 118 to 129) and of the CDR (CDR1: 27 to 38, CDR2: 56 to 65; and CDR3: 105 to 117) is provided.

[0039] An "antibody fragment" as referred to herein may include any suitable antigen-binding antibody fragment known in the art. The antibody fragment may be a naturally-occurring antibody fragment, or may be obtained by manipulation

of a naturally-occurring antibody or by using recombinant methods. For example, an antibody fragment may include, but is not limited to a Fv, single-chain Fv (scFv; a molecule consisting of V_L and V_H connected with a peptide linker), Fab, $F(ab')_2$, single domain antibody (sdAb; a fragment composed of a single V_L or V_H), and multivalent presentations of any of these. Antibody fragments such as those just described may require linker sequences, disulfide bonds, or other type of covalent bond to link different portions of the fragments; those of skill in the art will be familiar with the requirements of the different types of fragments and various approaches and various approaches for their construction.

[0040] In a non-limiting example, the antibody fragment may be an sdAb derived from naturally-occurring sources. Heavy chain antibodies of camelid origin (Hamers-Casterman et al, 1993) lack light chains and thus their antigen binding sites consist of one domain, termed V_{HH} . sdAb have also been observed in shark and are termed V_{NAR} (Nuttall et al, 2003). Other sdAb may be engineered based on human Ig heavy and light chain sequences (Jespers et al, 2004; To et al, 2005). As used herein, the term "sdAb" includes those sdAb directly isolated from V_H , V_{HH} , V_L , or V_{NAR} reservoir of any origin through phage display or other technologies, sdAb derived from the aforementioned sdAb, recombinantly produced sdAb, as well as those sdAb generated through further modification of such sdAb by humanization, affinity maturation, stabilization, solubilization, camelization, or other methods of antibody engineering. Also encompassed by the present invention are homologues, derivatives, or fragments that retain the antigen-binding function and specificity of the sdAb.

[0041] SdAb possess desirable properties for antibody molecules, such as high thermostability, high detergent resistance, relatively high resistance to proteases (Dumoulin et al, 2002 and high production yield (Arbabi-Ghahroudi et al, 1997); they can also be engineered to have very high affinity by isolation from an immune library (Li et al, 2009) or by *in vitro* affinity maturation (Davies & Riechmann, 1996). Further modifications to increase stability, such as the introduction of non-canonical disulfide bonds (Hussack et al, 2011; Kim et al, 2012), may also be brought to the sdAb.

[0042] A person of skill in the art would be well-acquainted with the structure of a single-domain antibody (see, for example, 3DWT, 2P42 in Protein Data Bank). An sdAb comprises a single immunoglobulin domain that retains the immunoglobulin fold; most notably, only three CDR/hypervariable loops form the antigen-binding site. However, and as would be understood by those of skill in the art, not all CDR may be required for binding the antigen. For example, and without wishing to be limiting, one, two, or three of the CDR may contribute to binding and recognition of the antigen by the sdAb of the present invention. The CDR of the sdAb or variable domain are referred to herein as CDR1, CDR2, and CDR3, and numbered as defined by Lefranc, M.-P. et al. (2003).

[0043] The antibody of the present invention is specific for Insulin-Like Growth Factor 1 Receptor (IGF1R), a receptor found on cell surfaces. The IGF1R comprises an alpha subunit, which comprises an extracellular part having the insulin-like growth factor 1 binding site, connected by a disulphide bond to the beta subunit, which comprises a small extracellular domain, a transmembrane region, and an intracellular portion. The IGF1 receptor assembles into a homo-dimer or may form a heterodimer with the insulin receptor. The sequence of IGF1R may be, but is not limited to that shown in Figure 2 (SwissProt Accession No. P08069; SEQ ID NO:12), or a sequence substantially identical thereto.

[0044] The antibody as described herein should not interfere with signaling through the Insulin Receptor (IR) or IGF1R. Specifically, the antibodies or fragments thereof as described herein should not inhibit AKT phosphorylation induced by insulin, nor should they induce phosphorylation of the IR on their own or inhibit insulin-induced signaling; additionally, the antibodies or fragments thereof described herein should not inhibit IGF-1-induced phosphorylation of IGF1R. Moreover, they should not bind to the Insulin Receptor.

[0045] As previously stated, the antibody may be an sdAb. The sdAb may be of camelid origin or derived from a camelid V_{HH} , and thus may be based on camelid framework regions; alternatively, the CDR described above may be grafted onto V_{NAR} , V_{HH} , V_H or V_L framework regions. In yet another alternative, the hypervariable loops described above may be grafted onto the framework regions of other types of antibody fragments (Fv, scFv, Fab) of any source (for example, mouse) or proteins of similar size and nature onto which CDR can be grafted (for example, see Nicaise et al, 2004).

[0046] The present invention further encompasses an antibody that is "humanized" using any suitable method known in the art, for example, but not limited to CDR grafting and veneering. Humanization of an antibody or antibody fragment comprises replacing an amino acid in the sequence with its human counterpart, as found in the human consensus sequence, without loss of antigen-binding ability or specificity; this approach reduces immunogenicity of the antibody when introduced into human subjects. In the process of CDR grafting, one or more than one of the CDR defined herein may be fused or grafted to a human variable region (V_H , or V_L), to other human antibody (IgA, IgD, IgE, IgG, and IgM), to other human antibody fragment framework regions (Fv, scFv, Fab), or to other proteins of similar size and nature onto which CDR can be grafted (Nicaise et al, 2004). In such a case, the conformation of said one or more than one hypervariable loop is likely preserved, and the affinity and specificity of the sdAb for its target (i.e., IGF1R) is likely minimally affected. CDR grafting is known in the art and is described in at least the following: US Patent No. 6180370, US Patent No. 5693761, US Patent No. 6054297, US Patent No. 5859205, and European Patent No. 626390. Veneering, also referred to in the art as "variable region resurfacing", involves humanizing solvent-exposed positions of the antibody or fragment; thus, buried non-humanized residues, which may be important for CDR conformation, are preserved while

the potential for immunological reaction against solvent-exposed regions is minimized. Veneering is known in the art and is described in at least the following: US Patent No. 5869619, US Patent No. 5766886, US Patent No. 5821123, and European Patent No. 519596. Persons of skill in the art would also be amply familiar with methods of preparing such humanized antibody fragments and humanizing amino acid positions.

[0047] For example, and without wishing to be limiting in any manner, the isolated or purified antibody specific for IGF1R may be $X_1VX_2LX_3ESGGGLVQX_4GGSLRLSCX_5X_6SGGTVSPTAMGWX_7RQAPGKX_8X_9EX_{10}VX_{11}HITWSRGTTX_{12}ASSVKX_{13}RFTISRDX_{14}X_{15}KNTX_{16}YMQMNSLX_{17}X_{18}EDTAVYYCAAS\ TFLRILPEESAYTYWGQGT$ $X_{19}VTVSS$ (SEQ ID NO:4), where X_1 is E or Q; X_2 is K or Q; X_3 is V or E; X_4 is A or P; X_5 is A or E; X_6 is V or A; X_7 is V or F; X_8 is G or E; X_9 is L or R; X_{10} is F or W; X_{11} is G or S; X_{12} is V or Y; X_{13} is D or G; X_{14} is N or S; X_{15} is A or S; X_{16} is L or V; X_{17} is K or R; X_{18} is A or S; and X_{19} is L or Q. Alternatively, the isolated or purified antibody may comprise a sequence selected from the group consisting of:

QVKLEESGGGLVQAGGSLRLSCEVSGGTVSPTAMGWFRQAPGKEREFGVGH ITWSR
GTTTRVASSVKDRFTISRDSKNTVYLQMNSLKSEDTAVYYCAASTFLRILPEESAYTY WGQGTQVTVSS (SEQ ID
NO:5), referred to herein as IGF1R-4;

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWVRQAPGKGLEWVGHITWSR
GTRYASSVKGRFTISRDNKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTY WGQGTTLTVSS (SEQ ID
NO:6), referred to herein as IGF1 R-4_H2;

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKGLEFVGHITWSR
GTRYASSVKGRFTISRDNKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTY WGQGTTLTVSS (SEQ ID
NO:7), referred to herein as IGF1 R-4_H3;

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKGLEFVGHITWSR
GTRYASSVKGRFTISRDSKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTY WGQGTTLTVSS (SEQ ID
NO:8), referred to herein as IGF1R-4_H4;

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKEREFGVGHITWSR
GTRYASSVKGRFTISRDSKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTY WGQGTTLTVSS (SEQ ID
NO:9), referred to herein as IGF1 R-4_H5; and

or a sequence having 90% identity thereto.

[0048] A substantially identical sequence may comprise one or more conservative amino acid mutations. It is known in the art that one or more conservative amino acid mutations to a reference sequence may yield a mutant peptide with no substantial change in physiological, chemical, physico-chemical or functional properties compared to the reference sequence; in such a case, the reference and mutant sequences would be considered "substantially identical" polypeptides. A conservative amino acid substitution is defined herein as the substitution of an amino acid residue for another amino acid residue with similar chemical properties (e.g. size, charge, or polarity). These conservative amino acid mutations may be made to the framework regions of the sdAb while maintaining the CDR sequences listed above and the overall structure of the CDR of the antibody or fragment; thus the specificity and binding of the antibody are maintained.

[0049] In a non-limiting example, a conservative mutation may be an amino acid substitution. Such a conservative amino acid substitution may substitute a basic, neutral, hydrophobic, or acidic amino acid for another of the same group. By the term "basic amino acid" it is meant hydrophilic amino acids having a side chain pK value of greater than 7, which are typically positively charged at physiological pH. Basic amino acids include histidine (His or H), arginine (Arg or R), and lysine (Lys or K). By the term "neutral amino acid" (also "polar amino acid"), it is meant hydrophilic amino acids having a side chain that is uncharged at physiological pH, but which has at least one bond in which the pair of electrons shared in common by two atoms is held more closely by one of the atoms. Polar amino acids include serine (Ser or S), threonine (Thr or T), cysteine (Cys or C), tyrosine (Tyr or Y), asparagine (Asn or N), and glutamine (Gln or Q). The term "hydrophobic amino acid" (also "non-polar amino acid") is meant to include amino acids exhibiting a hydrophobicity of greater than zero according to the normalized consensus hydrophobicity scale of Eisenberg (1984). Hydrophobic amino acids include proline (Pro or P), isoleucine (Ile or I), phenylalanine (Phe or F), valine (Val or V), leucine (Leu or L), tryptophan (Trp or W), methionine (Met or M), alanine (Ala or A), and glycine (Gly or G). "Acidic amino acid" refers to hydrophilic amino acids having a side chain pK value of less than 7, which are typically negatively charged at physiological pH. Acidic amino acids include glutamate (Glu or E), and aspartate (Asp or D).

[0050] Sequence identity is used to evaluate the similarity of two sequences; it is determined by calculating the percent of residues that are the same when the two sequences are aligned for maximum correspondence between residue positions. Any known method may be used to calculate sequence identity; for example, computer software is available

to calculate sequence identity. Without wishing to be limiting, sequence identity can be calculated by software such as NCBI BLAST2 service maintained by the Swiss Institute of Bioinformatics (and as found at ca.expasy.org/tools/blast/), BLAST-P, Blast-N, or FASTA-N, or any other appropriate software that is known in the art.

[0051] The substantially identical sequences of the present invention may be at least 90% identical; in another example, the substantially identical sequences may be at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% identical, or any percentage therebetween, at the amino acid level to sequences described herein. Importantly, the substantially identical sequences retain the activity and specificity of the reference sequence. In a non-limiting embodiment, the difference in sequence identity may be due to conservative amino acid mutation(s). In a non-limiting example, the present invention may be directed to an antibody comprising a sequence at least 95%, 98%, or 99% identical to that of the antibodies described herein.

[0052] The antibody of the present invention may also comprise additional sequences to aid in expression, detection or purification of a recombinant antibody or fragment thereof. Any such sequences or tags known to those of skill in the art may be used. For example, and without wishing to be limiting, the antibody may comprise a targeting or signal sequence (for example, but not limited to ompA), a detection/purification tag (for example, but not limited to c-Myc, His₆, or His₆), or a combination thereof. In another example, the additional sequence may be a biotin recognition site such as that described by Cronan et al in WO 95/04069 or Voges et al in WO/2004/076670. As is also known to those of skill in the art, linker sequences may be used in conjunction with the additional sequences or tags, or may serve as a detection/purification tag.

[0053] The antibody of the present invention may also be in a multivalent display format, also referred to herein as multivalent presentation. Multimerization may be achieved by any suitable method of known in the art. For example, and without wishing to be limiting in any manner, multimerization may be achieved using self-assembly molecules such as those described in Zhang et al (2004a; 2004b) and WO2003/046560, where pentabodies are produced by expressing a fusion protein comprising the antibody of the present invention and the pentamerization domain of the B-subunit of an AB₅ toxin family (Merritt & Hol, 1995). A multimer may also be formed using the multimerization domains described by Zhu et al. (2010); this form, referred to herein as a "combody" form, is a fusion of the antibody or fragment of the present invention with a coiled-coil peptide resulting in a multimeric molecule (Zhu et al., 2010). Other forms of multivalent display are also encompassed by the present invention. For example, and without wishing to be limiting, the antibody may be presented as a dimer, a trimer, or any other suitable oligomer. This may be achieved by methods known in the art, for example direct linking connection (Nielson et al, 2000), c-jun/Fos interaction (de Kruif & Logtenberg, 1996), "Knob into holes" interaction (Ridgway et al, 1996).

[0054] Another method known in the art for multimerization is to dimerize the antibody using an Fc domain, for example, but not limited to human Fc domains. The Fc domains may be selected from various classes including, but not limited to, IgG, IgM, or various subclasses including, but not limited to IgG1, IgG2, etc. In this approach, the Fc gene is inserted into a vector along with the sdAb gene to generate a sdAb-Fc fusion protein (Bell et al, 2010; Iqbal et al, 2010); the fusion protein is recombinantly expressed then purified. For example, and without wishing to be limiting in any manner, multivalent display formats may encompass chimeric formats of anti-IGF1R-4 V_HH linked to an Fc domain, or bi- or tri-specific antibody fusions with two or three anti-IGF1R-4 V_HH recognizing unique epitopes. Such antibodies are easy to engineer and to produce, can greatly extend the serum half-life of sdAb, and may be excellent tumor imaging reagents (Bell et al., 2010).

[0055] The Fc domain in the multimeric complex as just described may be any suitable Fc fragment known in the art. The Fc fragment may be from any suitable source; for example, the Fc may be of mouse or human origin. In a specific, non-limiting example, the Fc may be the mouse Fc2b fragment or human Fc1 fragment (Bell et al, 2010; Iqbal et al, 2010). The Fc fragment may be fused to the N-terminal or C-terminal end of the anti-IGF1R-3 V_HH or humanized versions of the present invention. In a specific, non-limiting example, the multimerized isolated or purified antibody or fragment as just described may comprise the sequence of SEQ ID NO:10, 38, or 11.

[0056] Each subunit of the multimers described above may comprise the same or different antibodies or fragments thereof of the present invention, which may have the same or different specificity. Additionally, the multimerization domains may be linked to the antibody or antibody fragment using a linker, as required; such a linker should be of sufficient length and appropriate composition to provide flexible attachment of the two molecules, but should not hamper the antigen-binding properties of the antibody.

[0057] The antibody as described herein may transmigrate across the blood brain barrier. The brain is separated from the rest of the body by a specialized endothelial tissue known as the blood-brain barrier (BBB). The endothelial cells of the BBB are connected by tight junctions and efficiently prevent many therapeutic compounds from entering the brain. In addition to low rates of vesicular transport, one specific feature of the BBB is the existence of enzymatic barrier(s) and high level(s) of expression of ATP-dependent transporters on the abluminal (brain) side of the BBB, including P-glycoprotein (Gottesman et al., 1993; Watanabe, 1995), which actively transport various molecules from the brain into the blood stream (Samuels, 1993). Only small (<500 Daltons) and hydrophobic (Pardridge, 1995) molecules can more readily cross the BBB. Thus, the ability of the antibody as described above to specifically bind the surface receptor,

internalize into brain endothelial cells, and undergo transcytosis across the BBB by evading lysosomal degradation is useful in the neurological field.

[0058] The present invention also encompasses nucleic acid sequences encoding the molecules as described herein. Given the degeneracy of the genetic code, a number of nucleotide sequences would have the effect of encoding the polypeptide, as would be readily understood by a skilled artisan. The nucleic acid sequence may be codon-optimized for expression in various microorganisms. The present invention also encompasses vectors comprising the nucleic acids as just described. Furthermore, the invention encompasses cells comprising the nucleic acid and/or vector as described.

[0059] The present invention further encompasses the isolated or purified antibody or fragments thereof immobilized onto a surface using various methodologies; for example, and without wishing to be limiting, the antibody or fragment may be linked or coupled to the surface via His-tag coupling, biotin binding, covalent binding, adsorption, and the like. Immobilization of the antibody of the present invention may be useful in various applications for capturing, purifying or isolating proteins. The solid surface may be any suitable surface, for example, but not limited to the well surface of a microtiter plate, channels of surface plasmon resonance (SPR) sensor chips, membranes, beads (such as magnetic-based or sepharose-based beads or other chromatography resin), glass, plastic, stainless steel, a film, or any other useful surface such as nanoparticles, nanowires and cantilever surfaces.

[0060] The invention also encompasses the antibody as described above linked to a cargo molecule. The cargo molecule may be any suitable molecule, which is delivered across the BBB by the antibody or fragment thereof. The cargo molecule may have a molecular weight in the range of about 1kD to about 200 kDa; for example, the cargo molecule may have a molecular weight of about 1, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, or 200 kDa, or any weight therebetween, or any range of weights defined by any two aforementioned weights. In specific, non-limiting examples, the cargo molecule may have a molecular weight of 1 kDa (for example, but not limited to a small molecule such as Cy5.5), 1-10 kDa (for example, but not limited to a peptide such as galanin, 3kDa), about 80 kDa (for example, but not limited to a Fc fragment, enzyme, protein, antibody etc), or about 200kDa (for example, but not limited to a monoclonal antibody).

[0061] For example, and without wishing to be limiting in any manner, the cargo molecule may be a detectable agent, a therapeutic agent, a drug, a peptide, an enzyme, a growth factor, a cytokine, a receptor trap, an antibody (e.g., IgG, scFv, Fab, $V_{H}H$, V_{H} , V_{L} , etc) a chemical compound, a carbohydrate moiety, DNA-based molecules (anti-sense oligonucleotide, microRNA, siRNA, plasmid), a cytotoxic agent, viral vector (adeno-, lenti-, retro), one or more liposomes or nanocarriers loaded with any of the previously recited types of cargo molecules, or one or more nanoparticle, nanowire, nanotube, or quantum dots. The cargo molecule as described above may be a detectable agent. For example, the IGF1R-specific antibody may be linked to a radioisotope, a paramagnetic label, a fluorophore, a fluorescent agent, Near Infra-Red (NIR; for example Cy5.5) fluorochrome or dye, an echogenic microbubble, an affinity label, a detectable protein-based molecule, nucleotide, quantum dot, nanoparticle, nanowire, or nanotube or any other suitable agent that may be detected by imaging methods. The antibody may be linked to the cargo molecule using any method known in the art (recombinant technology, chemical conjugation, etc.).

[0062] The cargo molecule as described herein may be linked, also referred to herein as "conjugated", to the antibody by any suitable method known in the art. For example, and without wishing to be limiting, the cargo molecule may be linked to the peptide by a covalent bond or ionic interaction. The linkage may be achieved through a chemical cross-linking reaction, or through fusion using recombinant DNA methodology combined with any peptide expression system, such as bacteria, yeast or mammalian cell-based systems. When conjugating the cargo molecule to the antibody or fragment thereof, a suitable linker may be used. Methods for linking an antibody or fragment thereof to a cargo molecule such as a therapeutic or detectable agent would be well-known to a person of skill in the art.

[0063] In one non-limiting example, the cargo molecule may be a detectable label, a radioisotope, a paramagnetic label such as gadolinium or iron oxide, a fluorophore, Near Infra-Red (NIR) fluorochrome or dye, an echogenic microbubble, an affinity label (for example biotin, avidin, etc), enzymes, or any other suitable agent that may be detected by diagnostic imaging methods. In a specific, non-limiting example, the anti-IGF1R-4 or fragment thereof may be linked to a near infrared fluorescence (NIRF) imaging dye, for example and not wishing to be limiting Cy5.5, Alexa680, Dylight680, or Dylight800.

[0064] Thus, the present invention further provides an *in vitro* method of detecting IGF1R, comprising contacting a tissue sample with one or more than one isolated or purified antibody or fragment thereof of the present invention linked to a detectable agent. The IGF1 R-antibody complex can then be detected using detection and/or imaging technologies known in the art. The tissue sample in the method as just described may be any suitable tissue sample, for example but not limited to a serum sample, a vascular tissue sample, a tumour tissue sample, or a brain tissue sample; the tissue sample may be from a human or animal subject. The step of contacting is done under suitable conditions, known to those skilled in the art, for formation of a complex between the antibody and IGF1R. The step of detecting may be accomplished by any suitable method known in the art, for example, but not limited to optical imaging, immunohistochemistry, molecular diagnostic imaging, ELISA, imaging mass spectrometry, or other suitable method. For example,

and without wishing to be limiting in any manner, the isolated or purified antibody linked to a detectable agent may be used in immunoassays (IA) including, but not limited to enzyme IA (EIA), ELISA, "rapid antigen capture", "rapid chromatographic IA", and "rapid EIA". (For example, see Planche et al, 2008; Sloan et al, 2008; Rüssmann et al, 2007; Musher et al, 2007; Turgeon et al, 2003; Fenner et al, 2008).

[0065] The present invention also provides an *in vivo* method of detecting IGF1R expression in a subject. The method comprises administering one or more than one isolated or purified antibody as described herein linked to a detectable agent to the subject, then detecting the labelled antibody bound to IGF1R. The step of detecting may comprise any suitable method known in the art, for example, but not limited to PET, SPECT, or fluorescence imaging, or any other suitable method. The method as just described may be useful in detecting the expression of IGF1R in blood vessels or tissues, for example but not limited to tumor tissues.

[0066] The *in vivo* detection step in the methods described above may be whole body imaging for diagnostic purposes or local imaging at specific sites, such as but not limited to brain vessels or brain tumor vessels, in a quantitative manner to assess the progression of disease or host response to a treatment regimen. The detection step in the methods as described above may be immunohistochemistry, or a non-invasive (molecular) diagnostic imaging technology including, but not limited to:

- Optical imaging;
- Positron emission tomography (PET), wherein the detectable agent is an isotopes such as ^{11}C , ^{13}N , ^{15}O , ^{18}F , ^{64}Cu , ^{62}Cu , ^{124}I , ^{76}Br , ^{82}Rb and ^{68}Ga , with ^{18}F being the most clinically utilized;
- Single photon emission computed tomography (SPECT), wherein the detectable agent is a radiotracer such as $^{99\text{m}}\text{Tc}$, ^{111}In , ^{123}I , ^{201}Tl , ^{133}Xe , depending on the specific application;
- Magnetic resonance imaging (MRI), wherein the detectable agent may be, for example and not limited to gadolinium, iron oxide nanoparticles and carbon-coated iron-cobalt nanoparticles thereby increasing the sensitivity of MRI for the detection of plaques.
- Contrast-Enhanced Ultrasonography (CEUS) or ultrasound, wherein the detectable agent is at least one acoustically active and gas-filled microbubble. Ultrasound is a widespread technology for the screening and early detection of human diseases. It is less expensive than MRI or scintigraphy and safer than molecular imaging modalities such as radionuclide imaging because it does not involve radiation.

[0067] The present invention further provides a method of transporting a molecule of interest across the blood-brain barrier. The method comprises administering the molecule linked to an antibody as described herein to a subject; the antibody transmigrates the blood-brain barrier. The molecule may be any desired molecule, including the cargo molecules as previously described; the molecule may be "linked" to the antibody using any suitable method, including, but not limited to conjugation or expression in a fusion protein. The administration may be by any suitable method, for example parenteral administration, including but not limited to intravenous (iv), subcutaneous (sc), and intramuscular (im) administration. In this method, the antibody of the present invention 'ferries' the molecule of interest across the BBB to its brain target.

[0068] The present invention also encompasses a composition comprising one or more than one isolated or purified antibody as described herein. The composition may comprise a single antibody or fragment as described above, or may be a mixture of antibodies or fragments. Furthermore, in a composition comprising a mixture of antibodies or fragments of the present invention, the antibodies may have the same specificity, or may differ in their specificities; for example, and without wishing to be limiting in any manner, the composition may comprise antibodies or fragments thereof specific to IGF1R (same or different epitope).

[0069] The composition may also comprise a pharmaceutically acceptable diluent, excipient, or carrier. The diluent, excipient, or carrier may be any suitable diluent, excipient, or carrier known in the art, and must be compatible with other ingredients in the composition, with the method of delivery of the composition, and is not deleterious to the recipient of the composition. The composition may be in any suitable form; for example, the composition may be provided in suspension form, powder form (for example, but limited to lyophilised or encapsulated), capsule or tablet form. For example, and without wishing to be limiting, when the composition is provided in suspension form, the carrier may comprise water, saline, a suitable buffer, or additives to improve solubility and/or stability; reconstitution to produce the suspension is effected in a buffer at a suitable pH to ensure the viability of the antibody or fragment thereof. Dry powders may also include additives to improve stability and/or carriers to increase bulk/volume; for example, and without wishing to be limiting, the dry powder composition may comprise sucrose or trehalose. In a specific, non-limiting example, the composition may be so formulated as to deliver the antibody to the gastrointestinal tract of the subject. Thus, the composition

may comprise encapsulation, time-release, or other suitable technologies for delivery of the antibody or fragment thereof. It would be within the competency of a person of skill in the art to prepare suitable compositions comprising the present compounds.

[0070] The invention also encompasses a method of quantifying an amount of a cargo molecule delivered across the BBB of a subject, wherein the cargo molecule is linked to one or more than one isolated or purified antibody as described herein, the method comprising

a) collecting cerebrospinal fluid (CSF) from the subject; and

b) using targeted proteomics methods to quantify the amount of the cargo molecule linked to one or more than one antibody in the CSF.

[0071] The cargo molecule may be any desired molecule, including the cargo molecules, as previously described; the isolated or purified antibody transmigrates the blood-brain barrier; and the molecule may be "linked" to the antibody using any suitable method, including, but not limited to conjugation or expression in a fusion protein, as previously described. In the above method, the CSF is collected from a subject using any suitable method known in the art. The amount of CSF required for targeted proteomics method in step b) may be between about 1 to 10 μ l; for example, the amount of CSF required may be about 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, 9.0, 9.5, or 10 μ l, or any amount therebetween, or any range defined by the amount just described. The antibody or fragment linked to the cargo molecule may have been administered to the subject prior to collection of the CSF. A suitable delay between administration and delivery of the antibody or fragment linked to the cargo molecule across the BBB may be required. The delay may be at least 30 minutes following administration of the antibody or fragment linked to the cargo molecule; for example and without wishing to be limiting in any manner, the delay may be at least 30 minutes, 1 hour, 1.5 hour, 2 hours, 2.5 hours, 3 hours, 3.5 hours, 4 hours, 4.5 hours, or 5 hours. The targeted proteomics methods used to quantify the amount of the one or more than antibody linked to the cargo molecule may be any suitable method known in the art. For example and without wishing to be limiting, the targeted proteomics method may be a mass spectrometry method, such as but not limited to multiple reaction monitoring using an isotopically labeled internal standard (MRM-ILIS; see for example Haqqani et al., 2013). MRM is advantageous in that it allows for rapid, sensitive, and specific quantification of unlabelled targeted analytes (for example, an antibody as described herein) in a biological sample. The multiplexing capability of the assay may allow for quantification of both the antibody and the cargo molecule.

[0072] The present invention will be further illustrated in the following examples. However, it is to be understood that these examples are for illustrative purposes only and should not be used to limit the scope of the present invention in any manner.

Example 1: Purification of IGF1R recombinant fragment

[0073] A 933 amino acid long recombinant fragment of the extracellular domain of IGF1R (shown by the grey box in Figure 1; see also amino acids 1-933 of SEQ ID NO:12) was prepared. The fragment comprised an N-terminal 30 amino acid signal peptide, the full alpha subunit, a furin cleavage site (RKRR, SEQ ID NO:13; separating alpha and beta subunits), as well as the majority of the extracellular portion of the beta subunit (Figures 1 and 2).

[0074] *Cloning.* The sequence of the IGF1R ectodomain of interest was amplified by PCR using the following primers:

5'-CGGGATCCGCCACCATGAAGTCTGGCTCCGGAG-3' (forward; SEQ ID NO:14)

5'-GCTCTAGATCAGAAGTTTTCATATCCTGTTTTGG-3' (reverse; SEQ ID NO:15)

and subcloned into the SmaI site of Puc19. The IGF1R⁹³³ sequence was then sub-cloned into pCDN4/myc-His (Invitrogen) to generate pIGF1R⁹³³-His, which allows for expression of a His-tagged ectodomain as described previously (Samani et al. 2004).

[0075] *Transient transfection.* Lentivirus particles expressing IGF1R⁹³³-His were generated in the packaging cell line 293SF-PacLV, as detailed previously (Broussau et al., 2008). Briefly, the cells were transfected with the vector using polyethylenimine. Fresh medium (LC-SFM), containing 1 μ g/ml doxycycline and 10 μ g/ml cumate, was added to the cells 5 hours post-transfection and the supernatants containing LV particles were collected after 48-72 hours, concentrated by ultracentrifugation at 100,000xg for 2h at 4°C on a 20% sucrose cushion (Gaillet B et al. 2007), re-suspended in LC-SFM medium supplemented with 1% FBS and stored at -70°C until used.

[0076] *Stable expression.* A stable cell line, 293SF-cum2-CR5-IGF1R-His, was generated by transduction of 293SF-Cum2 cell lines with the respective lentivirus particles using a protocol described previously (Gaillet B. et al. 2010). Briefly, 0.5-1.0 X 10⁵ 293SF-Cum2 cells were seeded in 24 wells plates into 200 μ l of LC-SFM medium without dextran

sulfate. The LV suspension was prepared by mixing 200-500 μ L of LV with 8 μ g/mL of polybrene and incubating it for 30min at 37°C. The freshly made LV suspension was added to the cells 4 hours after seeding. After 24h, 500 μ L medium supplemented with dextran sulfate was added to the cells. To increase the level of expression the cells were re-transduced up to 6 times using the same protocol after 3-4 days of cell recovery. Finally the cells were expanded in 6-well plates and shaker flasks.

[00777] *Large scale protein production and purification.* The clone identified as the highest producer was expanded in shaker or spinner flasks. Protein production was initiated by the addition of 1 μ g/ml cumate in fresh medium, followed by a 24 h incubation at 37°C and a 4-8 day incubation at 30°C. Cells were removed by centrifugation and the supernatants filtered and concentrated (10x) using the Tangential Flow Filtration Systems (Pellicon ultrafiltration cassettes, EMD Millipore).

[0078] The IGF1R⁹³³-His was purified using a HisPrep column (GE Healthcare) according to the manufacturer's instructions. Briefly, the concentrated sample was applied to a His-prep FF (16/10) column (GE Healthcare), equilibrated and washed with 50 mM sodium phosphate, 300mM NaCl, 5mM imidazole pH 7.5 and eluted with the 50 mM sodium phosphate, 300mM NaCl, 500mM imidazole pH 7.5. A step elution with 0.1 M sodium citrate pH 4.5 to pH 2.5 was used to elute the protein and peak fractions were pooled. Buffer exchange was performed by ultrafiltration using a 50 kDa cut-off membrane or a desalting column with a buffer containing 50 mM sodium phosphate, 150 mM NaCl and 0.01% Tween-80, pH 7.2. Purity of both proteins was verified by SDS-PAGE and they were stored at -80°C until used (see subsequent examples).

Example 2: Llama immunization and serum response

[0079] To isolate V_HHs that targets the extracellular domain of IGF1R, a llama was immunized with the recombinant IGF1R⁹³³-His fragment obtained in Example 1.

[0080] One male llama (*Lama glama*) was immunized by sub-cutaneous, lower-back injection of IGF1R⁹³³-His recombinant antigen (Example 1). On day 1, 200 μ g of antigen diluted in PBS to 1 ml was injected together with 1 ml of Freund's Complete Adjuvant (Sigma, St. Louis, MO). Three further injections of 100 μ g of IGF1R⁹³³-His antigen plus Freund's Incomplete Adjuvant (Sigma) were performed on days 22, 36, and 50. A final injection of 100 μ g of antigen with no adjuvant was performed on day 77. Pre-immune blood was drawn before the first injection on day 1 and served as a negative control. Blood (10-15 ml) was collected on days 29, 43, 57 and 84. The blood from day 84 was processed immediately to isolate peripheral blood mononuclear cells (PBMC). The blood was diluted 1:1 with phosphate buffered saline (PBS) and PBMCs were purified from the blood using the Lymphoprep Tube (Axis Shield). The cells were counted and stored as aliquots of about 1×10^7 cells at -80°C for future use.

[0081] Pre-immune and post-immune total serum was analyzed for a specific response to IGF1 R⁹³³-His antigen by ELISA on day 57. Llama sera from day 84 were fractionated as previously described (Doyle et al, 2008). The resulting fractions, A1 (HCAb), A2 (HCAb), G1 (HCAb) and G2 (cIgG) were analyzed for specific binding to the IGF1R⁹³³-His antigen by ELISA. Briefly, 5 μ g of IGF1R⁹³³-His recombinant antigen diluted in PBS was incubated overnight (100 μ L/well, 18 h, 4°C) in 96 well Maxisorp plates (Nalgene, Nunc) to coat the plates. Plates were blocked with bovine serum albumin (BSA), washed with PBS-T (PBS + 0.05% (v/v) Tween-20), and serial dilutions of pre-immune total serum, post-immune total serum (day 57), and fractionated serum (day 84) were applied. After incubation at room temperature for 1.5 h the plates were washed with PBS-T, before goat anti-llama IgG (1:1,000 in PBS) was added and plates were incubated for 1 h at 37°C. After washing with PBS-T, pig anti-goat IgG-HRP conjugate (1:3,000 in PBS) was added and plates were incubated for 1 h at 37°C. A final PBS-T wash was carried out prior to the addition of 100 μ L/well TMB substrate (KPL, Gaithersburg, MD); the substrate was incubated for 10 min. The reaction was stopped with 100 μ L/well 1 M H₃PO₄. Absorbance was read at 450 nm.

Example 3: Library construction and selection of IGF1 R-binding V_HHs

[0082] A hyperimmunized llama V_HH library was constructed based on RNA isolated from the PBMCs in Example 2.

[0083] Library construction and panning was performed essentially as previously described (Arbabi-Ghahroudi et al, 2009a, 2009b; Tanha et al, 2003). Total RNA was isolated from approximately 10^7 PBMCs collected on day 84 post-immunization (Example 2) using the QIAamp RNA Blood Mini Kit (Qiagen). About 5 μ g of total RNA was used as template for first strand cDNA synthesis with oligo dT primers using the First-Strand cDNA Synthesis Kit (GE Healthcare). The cDNA was amplified by an equimolar mix of three variable region-specific sense primers:

MJ1: 5'-GCCCAGCCGGCCATGGCCSMKGTGCAGCTGGTGGAKTCTGGGGGA-3' (SEQ ID NO:16)

MJ2: 5'-GCCCAGCCGGCCATGGCCCAGGTAAAGCTGGAGGAGTCTGGGGGA-3' (SEQ ID NO:17)

MJ3: 5'-GCCCAGCCGGCCATGGCCCAGGCTCAGGTACAGCTGGTGGAGTCT-3' (SEQ ID NO:18),

and two antisense CH₂-specific primers:

CH₂: 5'-CGCCATCAAGGTACCAGTTGA-3' (SEQ ID NO:19)

CH₂b₃: 5'-GGGGTACCTGTCATCCACGGACCAGCTGA-3' (SEQ ID NO:20).

[0084] Briefly, the PCR reaction mixture was set up in a total volume of 50 μ l with the following components: 1-3 μ l cDNA, 5 pmol of MJ1-3 primer mixture, 5 pmol of CH₂ or CH₂b₃ primers, 5 μ l of 10x reaction buffer, 1 μ l of 10 mM dNTP, 2.5 unit of Taq DNA polymerase (Hoffmann-La Roche). The PCR protocol consisted of an (i) initial step at 94°C for 3 min, (ii) followed by 30 cycles of 94°C for 1 min, 55°C for 30 s, 72°C for 30 s and (iii) a final extension step at 72°C for 7 min. The amplified PCR products were run on a 2% agarose gel and two major bands were observed: a band of about 850 bp, corresponding to conventional IgG, and a second band of around 600 bp, corresponding to the V_HH-CH₂ region of camelid heavy chain antibodies. The smaller bands were cut and purified using the QIAquick Gel Extraction Kit (Qiagen) and re-amplified in a second PCR in a total volume of 50 μ l using 1 μ l (30ng) of DNA template, 5 pmol of each of MJ7 primer (5'-CATGTGTAGACTCGCG GCCCAGCCGGCCATGGCC-3' SEQ ID NO:21) and MJ8 primer (5'-CATGTGTAGATTCTGCGCCGGCCTGCGCTGAGGAGACGGTGACCTGG-3' SEQ ID NO:22), 5 μ l of 10x reaction buffer, 1 μ l of 10 mM dNTP, 2.5 unit of Taq DNA polymerase. The PCR protocol consisted of (i) an initial step at 94°C for 3 min, (ii) followed by 30 cycles of 94°C for 30 s, 57°C for 30 s and 72°C for 1 min and (iii) a final extension step at 72°C for 7 min. The amplified PCR products, ranging between 340 bp and 420 bp and corresponding to V_HH fragments of heavy chain antibodies, were purified using the QIAquick PCR Purification Kit (Qiagen), digested with *Sfi*I restriction enzyme (New England BioLabs) and re-purified using the same kit.

[0085] 80 μ g of pMED1 phagemid vector (Arbabi-Ghahroudi et al, 2009b) were digested with *Sfi*I overnight at 50°C. To minimize self-ligation, 20 units of *Xho*I and *Pst*I restriction enzymes were added to cut the excised fragment and the digestion reaction was incubated for an additional 2 h at 37°C. 60 μ g of digested phagemid DNA was ligated with 6 μ g of digested (*Sfi*I for 5h at 50°C) V_HH fragments (molar ratio 1:1) for 3 h at room temperature using LigaFast Rapid DNA Ligation System (Promega) according to the manufacturer's instructions. The ligated plasmids were purified using the QIAquick PCR Purification Kit (Qiagen), eluted in a final volume of 100 μ l, and transformed into electrocompetent TG1 *E. coli* (Stratagene) using 5 μ l of ligated DNA aliquot per transformation reaction, as described (Arbabi-Ghahroudi et al, 2009b). The size of the library was determined to be 5 x 10⁷ as described in (Arbabi-Ghahroudi et al, 2009b). 20 clones were sequenced and contained all unique V_HH sequences. The *E. coli*, containing the library was grown for 2-3 h at 37°C, 250 rpm in the presence of 2% (w/v) glucose. The bacteria were then pelleted, resuspended in 2xYT/Amp/Glu (2xYT medium with 100 μ g/ml ampicillin and 2% (w/v) glucose) with 35% (v/v) glycerol and stored at -80°C in small aliquots.

[0086] Panning experiments were essentially performed as described in (Arbabi et al, 1997). Two milliliters of the library (2.0 x 10¹⁰ bacteria) were thawed on ice and grown in 2xYT/Amp/Glu for about 2 h at 37°C (A₆₀₀ = 0.4 - 0.5). The *E. coli* were subsequently infected with 20x excess M13K07 helper phage (New England Biolabs) for 1 h at 37°C. The culture was then centrifuged at 4°C and infected bacterial pellets were re-suspended in 200 ml of 2xYT/Amp with 50 μ g/ml kanamycin and incubated at 37°C and 250 rpm. The phage particles in the culture supernatant were incubated with 1/5 volume of 20% PEG 8000/2.5M NaCl on ice for 1 h and centrifuged at 10,000 rpm for 15 min. The phage pellets were re-suspended in 1.5 ml of sterile PBS, titrated and used as input phage for panning. For panning round 1, 96-well Maxisorp™ plates were coated with 10 μ g of recombinant IGF1 R⁹³³-His per well in 100 μ l PBS overnight at 4°C. The wells were rinsed with PBS and blocked with PBS plus 1% (w/v) casein for 2 h at 37°C. Approximately 10¹² phages were added to the blocked wells and incubated for 2 h at 37°C. After 10x washing with PBS/0.1% (v/v) Tween 20, the bound phages were eluted with 0.1 M triethylamine, neutralized (50 μ l of 1M Tris-HCl, pH 7.4) and mixed with exponentially growing TG1 *E. coli*. Titration of eluted phage was performed and infected bacteria were superinfected with M13K07 and grown overnight at 37°C. The purified phage from the overnight culture was used as the input for the next round of panning. The panning was continued for three further rounds. The same protocol as described above was used, except that the amount of recombinant antigen used to coat the plates was reduced to 7 μ g, 5 μ g and 5 μ g for the second, third and fourth rounds of panning, respectively.

[0087] Individual TG1 colonies obtained after round four of panning were subjected to phage ELISA screening, essentially as described elsewhere (Doyle et al, 2008), with the exception that 5 μ g/ml of IGF1 R⁹³³-His recombinant antigen were used to coat the microtiter plates. All positive clones were sent for DNA sequencing. Unique clones that gave high phage ELISA signals were selected for large-scale expression and purification using known methods (see Example 4). A clone dubbed IGF1R-4 was identified for further study; its sequence is shown below.

QVKLEESGGGLVQAGGSLRLSCEVSGGTVSPTAMGWFRQAPGKEREFEVGHITWSR
 GTTRVASSVKDRFTISRDSAKNTVYLQMNSLKSEDTAVYYCAASTFLRILPEESAYTY
 WGQGTQVTVSS (SEQ ID NO:5)

Example 4: Humanization of IGF1R-4

[0088] To avoid potential immunogenicity of llama-derived IGF1R-4 when applied as BBB carrier for therapeutics, to the camelid-derived sdAb was "humanized" by mutation of "camelid" residues in the V_HH. It should be noted that, for the purpose of humanization, Kabat numbering (Kabat et al, 1991) was used for identification of CDR residues.

[0089] *3D-structure modeling of camelid V_HHS.* Template structures similar to IGF1R-4 V_HH were identified using BLAST searches against the Protein Data Bank (PDB). The 3D structure of the IGF1R-4 was approximated using homology modeling based on 4KRPJB (PDB code | Chain ID) as the main template, with additional information from 4FHBID. The IGF1R-4 V_HH structure was then built by mutating the main template structure to the IGF1R-4 sequence; this included 35 mutations at various positions. The IGF1R-4 V_HH model was then refined by energy minimization with the AMBER force-field and a stepwise release of constraints, ranging from the CDR loops, which were relaxed first, to the backbone heavy atoms of the framework region, which were fully relaxed only in the last stage. The CDR-H3 loop of the V_HH model was then refined by Monte-Carlo-minimization (MCM) conformational sampling, in which dihedral angles in the CDR-H3 region were sampled followed by energy minimization.

[0090] *Selection of the human heavy-chain framework for the camelid CDR.* Human heavy-chain framework was selected by standard sequence homology comparison against the human germline databases (VBASE), against other sequence databases (Genbank and SwissProt), and against the human framework consensus sequences. BLAST searches were conducted to retrieve sequence matches with highest homology in the framework region only (i.e., excluding CDR) while matching the length of the CDR. The closest human frameworks identified for IGF1R-4 V_HH corresponded to the human VH-3 subgroup. Several human germline VH-3 framework sequences that were most similar to IGF1R-4 V_HH were also retained in addition to the human VH-3 consensus sequence. The IGF1R-4 V_HH framework sequences required 18 mutations in order to arrive at the consensus human VH-3 sequence for 100% framework humanization.

[0091] *Identification of framework residues for back-mutations.* The IGF1R-4 V_HH model and its fully-humanized counterpart were characterized to estimate the humanness index, antigen contact propensity index, to delineate the CDR, canonical residues, unusual framework residues, potential glycosylation sites, buried residues, Vernier zone residues, and proximity to CDR. The analysis of these data suggested the design of several humanized variants for the anti-IGF1R V_HH, each variant having varying numbers of back-mutations to the parent camelid residues at various positions. 6 humanized variants were designed for IGF1R-4 V_HH (IGF1R-4_H1 to IGF1 R-4_H6), where variants contained up to 10 back-mutations. Some of these camelid back-mutations residues were buried inside the V_HH domain core and hence are not expected to induce an immune response.

Example 5: Expression and purification of selected V_HH constructs

[0092] IGF1R-4 identified in Example 3 and the humanized versions constructed in Example 4 (collectively referred to herein as "V_HH constructs") were sub-cloned into expression plasmids for protein expression and purification.

[0093] A phagemid vector containing the DNA of the V_HH IGF1R-4 clone was purified using the MiniPrep Kit (Qiagen). The IGF1 R-binding V_HH IGF1R-4 was PCR amplified from the pMED1 phagemid vector, adding an N-terminal BbsI cleavage site and a BamHI cleavage site at the C-terminus, using primers:

5'-TATGAAGACACCAGGCCAGGTAAAGCTGGAGGAGTCT-3' (forward; SEQ ID NO:23)

5'-TTGTTTCGGATCCTGAGGAGACGGTGACCTG-3' (reverse; SEQ ID NO:24)

[0094] The PCR fragment and the pSJF2H expression vector was digested with BbsI and BamHI restriction enzymes (NEB) according to the manufacturer's instructions. Following digestion, each digested IGF1R-4 V_HH fragment was ligated into the digested pSJF2H expression vector, using methods similar to those described in Arbabi-Ghahroudi et al. (2009b); the ligation products were then transformed into electro-competent TG1 E. coli. Clones were selected on LB agar plates + 100 µg/ml ampicillin and verified by DNA sequencing.

[0095] The humanized clones were synthesized and directly cloned into pSJF2H, similarly as described above and subsequently transformed into E. coli TG1 and selected as described above.

[0096] *Protein Expression.* All IGF1R-4 V_HH constructs were expressed in TG1 E. coli. An overnight culture in

LB/amp/glu medium (LB medium supplemented with 100 µg/ml ampicillin and 1% glucose) was subcultured at 1:100 dilution in 1L LB/amp/glu. Protein expression was induced at an OD₆₀₀ of 0.8 - 0.9 by the addition of IPTG to a final concentration of 0.2 mM. The culture was grown at 220 rpm overnight at 37°C. The bacteria were pelleted by centrifuging at 6000 rpm for 12 min; the pellet was re-suspended in 35 ml of cold TES buffer (0.2M Tris-Cl pH 8.0, 20% sucrose, 0.5mM EDTA). The suspension was incubated on ice and vortexed every 10 min for 1 hour. Then 45 ml of cold TES (1/8 volume of total volume) was added and immediately vortexed for 1 minute and for 15 seconds every 10 min thereafter for 1 hour to extract the protein from the periplasm. The resulting supernatant containing V_HH was filtered through a 0.22 µm membrane and dialysed overnight into immobilized metal-affinity chromatography (IMAC) buffer A (10 mM HEPES pH 7.0, 500 mM NaCl). The protein was purified using HiTrap Chelating HP columns (GE Healthcare) as described previously (Arbabi-Ghahroudi 2009b). Eluted protein fractions were analyzed by SDS-PAGE and Western blotting before being dialysed against PBS as described previously (Arbabi-Ghahroudi 2009b). The purified protein fractions were pooled and dialyzed against PBS + 3 mM EDTA, and the protein concentration was determined.

Example 6: Biophysical characterization of anti-IGF1R V_HH IGF1R-4

[0097] The anti-IGF1R V_HH IGF1R-4 constructs expressed and purified in Example 4 were characterized using size exclusion chromatography, melting temperature analysis, and surface plasmon resonance.

[0098] *Size exclusion chromatography:* Size exclusion chromatography employing Superdex™ 75 (GE Healthcare) was used to eliminate any possible aggregates prior to Surface Plasmon Resonance (SPR) analysis. The running buffer used was 10mM HEPES, pH7.4 containing 150mM NaCl, 3mM EDTA and 0.005% P20. Concentrations of fractions used for SPR analysis were determined by measuring absorbance at 280nm wavelength. The SEC analysis suggested that all IGF1R-4 V_HH constructs, except H1 were monomeric, based on the elution volume compared to standards (Figure 3A). IGF1R-4 H1 was therefore excluded from further evaluation.

[0099] *Melting temperature:* The thermal stability of the IGF1R-4 V_HH and humanized constructs was evaluated using melting temperature (T_m) measurement by CD spectroscopy. A Jasco J-815 spectropolarimeter equipped with a Peltier thermoelectric type temperature control system (Jasco, Easton, MD, USA) was used to carry out experiments. A CD cuvette with a path length of 1 mm was used. The spectra were recorded over a wavelength range of 180 - 260 nm with scanning speed of 50 nm/min, digital integration time (DIT) of 4 s, a band width of 1 nm, data pitch of 1 nm, and an integration time of 1 s. To measure melting temperature or T_m (Greenfield, 2006a; 2006b), CD spectra were recorded over a temperature range of 30°C to 96°C. All CD spectra were subtracted from the blank corresponding to buffer spectra. Measurements were performed with concentrations of 50 µg/mL V_HH in 100 mM sodium phosphate buffer, pH 7.4. Heat-induced protein denaturation was monitored at 210 nm for all variants. The fraction folded (ff) was obtained by a formula as described (Greenfield, 2006a; 2006b):

$$ff = ([\theta]_T - [\theta]_U) / ([\theta]_F - [\theta]_U) \quad \text{formula I}$$

where $[\theta]_T$ is the molar ellipticity at any temperature, $[\theta]_F$ is the molar ellipticity of the fully folded protein at 30°C and $[\theta]_U$ is the molar ellipticity of the unfolded protein at 90°C. Melting temperature (T_m) was obtained as a midpoint of the unfolding curve (fraction folded (ff) versus temperature) by a nonlinear regression curve fit (Boltzman sigmoidal equation) using the graphing software GraphPad Prism (version 4.02 for Windows). The melting temperatures (T_m) of V_HH were determined based on ellipticity data assuming a two-state system, which is in agreement with the observed denaturation curves corresponding to a sharp transition into denaturation. The T_m values were taken at midpoint of the sigmoidal denaturation curves of fraction folded (ff) versus temperature.

[0100] Results are shown in Figure 3B. The melting temperatures of humanized V_HH H3, H4, and H5 were improved (higher) in comparison to the IGF1R-4 V_HH, suggesting improved biophysical properties. Humanized constructs H2 and H6 had poorer melting temperatures compared to IGF1R-4 V_HH. In contrast to IGF1R-4 V_HH, none of the humanized V_HHs refolded after heat exposure.

[0101] *Surface Plasmon Resonance (SPR):* The binding of monomeric IGF1R-4 V_HH constructs to immobilized recombinant human IGF1R (Example 1) was determined by SPR using BIACORE 3000 (GE Healthcare). Approximately 3000 Resonance Units (RU) of recombinant human IGF1R were immobilized on a Sensor chip CM5. Immobilization was carried out at a concentration of 10 µg/ml in 10mM acetate at pH4.0 using the amine coupling kit supplied by the manufacturer. The remaining binding sites were blocked with 1 M ethanolamine pH 8.5. An ethanolamine blocked surface was used as a reference surface. For the binding studies, analyses were carried out at 25°C in 10mM HEPES, pH7.4 containing 150mM NaCl, 3mM EDTA and 0.005% surfactant P20 (Polyoxyethylenesorbitan; GE Healthcare). Various concentrations of the IGF1R-4 V_HH were injected over the immobilized human IGF1R and reference surfaces at a flow rate of 20 µl/min. Surfaces were regenerated with 10mM glycine pH 2.0 with a contact time of 24 seconds. Data were analyzed with BIAevaluation 4.1 software (GE Healthcare). The sensograms in Figure 3C show that the data fit well to

a 1:1 model, giving K_D and 'off-rates' shown Table 1. This indicates that the IGF1R-4 and the humanized variants are high-affinity single-domain antibodies binding to the extracellular domain of human IGF1R.

Table 1. Affinity of IGF1R-4 constructs for human IGF1R as determined by surface plasmon resonance.

	K_D (nM)	k_d (S ⁻¹)
IGF1R-4	0.32	2.9×10^{-4}
IGF1R-4 H2	1.07	6.1×10^{-4}
IGF1R-4 H3	0.39	3.4×10^{-4}
IGF1R-4 H4	0.45	3.9×10^{-4}
IGF1R-4 H5	0.52	4.6×10^{-4}
IGF1R-4 H6	0.52	3.8×10^{-4}

[0102] The SPR analyses were further used to demonstrate that IGF1R-4 V_HH does not bind to the same epitope on the receptor as the natural ligand IGF-1 (Figure 3D). The experiment was set up, carried out and analysed as described above. Binding to freshly immobilized human IGF1 R surface was studied by injection of human IGF1 ligand at a concentration of $25 \times K_D$ followed by co-injection of IGF1R-4 both at concentrations $25 \times K_D$, with flow rates of 20 μ L/min and injection times of 5 minutes. Surfaces were regenerated by washing with running buffer. The natural ligand IGF-1 bound the receptor with saturation reached at 70RU; the IGF1R-4 V_HH bound to the IGF1R - IGF-1 complex with the expected ~265RU (relative units; binding saturation). The simultaneous binding of both IGF1R-4 V_HH and IGF-1 to the receptor demonstrates that both bind to different epitopes.

[0103] The cross-reactivity of IGF1R-4 binding to human Insulin Receptor was also evaluated using SPR. The experiment was set up, carried out and analysed as described above. Briefly, besides human IGF1R, approximately 4000 Resonance Units (RU) of recombinant human Insulin Receptor (R&D systems) was immobilized on a separate cell on a Sensor chip CM5. Subsequently, different concentrations of IGF1R-4 were flowed over the surfaces (0.5-50nM). While binding to IGF1R could be observed at already 0.5nM IGF1R-4, no binding to the Insulin Receptor surface was observed even at 50nM (FIGURE 3E), suggesting that IGF1R-5 cannot bind the Insulin receptor.

Example 7: Internalization of the IGF1R-4 by brain endothelial cells

[0104] To determine whether IGF1R-4 is internalized into cells, svARBEC cells were incubated with Cy5-5-labelled IGF1 R-4.

[0105] IGF1R-4 V_HH was labeled with NHS-Cy5.5. The labeling was done through a stable amine bond between primary amines (at the N-terminus and/or on lysine side chains of the protein) and the NHS ester. Typically, 10% v/v of 1M carbonate/bicarbonate buffer (759mM bicarbonate, 258mM carbonate) pH 9.3 was added to 4mg of V_HH prepared in PBS (1.06mM KH₂PO₄, 154mM NaCl, 5.6mM Na₂HPO₄) pH7.4 and adjusted to a final concentration of 4mg/mL. The NHS-Cy5.5, dissolved in DMSO at 10mg/mL, was added at a 2X molar ratio of dye to protein. The mixture was incubated at room temperature for 2 h with several inversions in a 1.5mL microcentrifuge tube. Following the incubation, unbound dye and reaction bi-products were filtered using Zeba Spin Desalting Columns, 7K MWCO (Pierce) and measured using a Beckman DU530 spectrophotometer (Beckman Coulter). Cy5.5-labeled IGF1R-4 or FC5 as a positive control (1mg/ml) were incubated with SV40 immortalized rat brain endothelial cells (svARBEC) at 4°C (Figure 4, top panels), thus allowing only for passive, non-specific transport mechanism to occur, or at 37°C (Figure 4, bottom panels) to allow for active transport such as receptor mediated endocytosis to take place. Co-staining with wheat germ agglutinin and DAPI was carried out to visualize the cell surface and the nucleus, respectively. Cells were observed under fluorescent microscope and images were captured.

[0106] If incubated at 4°C, IGF1R-4 was found outside the cells co-localizing with the cell membrane stained with wheat germ agglutinin. In contrast, when incubated at 37°C, IGF1R-4 accumulated in vesicles inside the cells, likely endosomes, suggesting that the antibody internalized into cells through an active transport mechanism. Similar behaviour was observed for FC5, previously shown to enter the cells by energy-dependent endocytosis via clathrin-coated vesicles (Abulrob et al.2005)

Example 8: Production of a IGF1R-4-mFc construct

[0107] Constructs comprising IGF1R-4 V_HH fused to a murine antibody fragment crystallisable (Fc; mFc2b) was prepared, expressed, and isolated. The sequence of the IGF1R-4-mFc construct is shown in Figure 5A, with a schematic

of the molecule shown in Figure 5B. The fusion protein also comprised a N-terminal signal peptide (MEFGLSWVFL-VAILKGVQC; SEQ ID NO:37) that is not shown in the sequence of Figure 5A.

[0108] The IGF1R-4 cDNA was cloned into mammalian expression vector pTT5 (Durocher, 2002) containing the mouse Fc2b fragment. Polyplexes of the resulting vector were pre-formed by mixing 25 ml of plasmid DNA solution containing 187.5 μ g pTT5-IR5mFc2b, 56.25 μ g pTT-AKTdd (activated mutant of Protein Kinase B), 18.75 μ g pTTo-GFP (to monitor transfection efficiency), and 112.5 μ g of salmon testis DNA (Sigma-Aldrich); and 25 ml of PEI solution containing 1.125 mg of PEIproTM (PolyPlus Transfection), both made in F17 medium. The mixture was incubated for 10 minutes prior to addition to the cell culture. A 450 ml culture of CHO cells stably expressing a truncated EBNA1 protein (CHO-3E7) and grown in F17 medium (Invitrogen) was transfected with 50 ml of polyplexes. Twenty four hours post-transfection, the culture was fed with 12.5 ml of 40% (w/v) tryptone N1 (Organotechnie) solution and 1.25 ml of 200 mM valproic acid solution. The culture was harvested 8 days post-transfection and clarified by centrifugation. Clarified medium was filtered through a 0.22 μ m membrane prior to its application on a column packed with 5 ml of protein-A MabSelect SuRe resin (GE Healthcare). After loading, the column was washed with 5 volumes of phosphate-buffered saline pH 7.1 (PBS) and the antibody was eluted with 100 mM sodium citrate buffer pH 3.0. Fractions containing the eluted antibody were pooled and a buffer exchange was performed by loading on a desalting Econo-Pac column (BioRad) equilibrated in PBS. Desalted antibody was then sterile-filtered by passing through a Millex GP (Millipore) filter unit (0.22 μ m) and aliquoted.

Example 9: Transport of the IGF1R-4 across in vitro blood brain barrier model

[0109] To evaluate whether the IGF1R-4 V_{HH} , its humanized versions (H2 to H6), and IGF1R-4-mFc (Example 8) transmigrate the blood-brain barrier, an *in vitro* assay was used as described below. A flow chart summarizing the experiment is shown at Figure 6A.

[0110] SV40-immortalized Adult Rat Brain Endothelial Cells (Sv-ARBE) were used to generate an *in vitro* blood-brain barrier (BBB) model as described (Garberg et al., 2005; Haqqani et al., 2012). Sv-ARBE (80,000 cells/membrane) were seeded on a 0.1mg/mL rat tail collagen type I-coated tissue culture inserts (pore size-1 μ m; surface area 0.9 cm², Falcon) in 1 ml of growth medium. The bottom chamber of the insert assembly contained 2 ml of growth medium supplemented with the immortalized neonatal rat astrocytes-conditioned medium in a 1:1 (v/v) ratio. Equimolar amounts (5.6 μ M) of positive (FC5) or negative controls (A20.1, a *Clostridium difficile* toxin A binding V_{HH} ; and EG2, an EGFR binding V_{HH}), IGF1R-4 V_{HH} , humanized versions (H2-H6) or IGF1 R-4mFc were tested for their ability to cross this rat *in vitro* BBB model. Following exposure of equimolar amounts of the sdAbs to the luminal side of the BBB, samples were taken after 15, 30, and 60 min from the abluminal side. The sdAb content of each sample was then quantified by mass spectrometry (multiple reaction monitoring - isotope labeled internal standards; MRM - ILIS) as described by Haqqani et al. (2012) (see method description below).

[0111] *Determination of the apparent permeability coefficient:* Quantified values can be directly plotted or the P_{app} (apparent permeability coefficient) values can be determined with the given formula (Fig 6A) and plotted. The P_{app} value is commonly used to determine the ability of a molecule to cross the BBB. [Q_r/dt = cumulative amount in the receiver compartment versus time; A = area of the cell monolayer; C₀ = initial concentration of the dosing solution]. P_{app} values are a measure of the specific permeability of the compound across brain endothelial monolayer.

[0112] Results are shown in Figures 6B-D. The results given are average P_{app} values obtained from several independent experiments. Both negative controls have a very low P_{app} value, indicating that non-specific transport or paracellular transport of these V_{HH} s across the BBB model is minimal. IGF1R-4 V_{HH} has a high P_{app} value, indicating high rate of transport across the *in vitro* BBB model. The P_{app} for IGF1R-4 V_{HH} is 4-fold higher than that of a positive control - BBB-permeable V_{HH} FC5 (WO 02/057445). The results provide strong indication that IGF1R-4 undergoes a facilitated trans-cellular transport across brain endothelial cells *in vitro* and could have similar properties *in vivo*. Humanized IGF1R-4 V_{HH} variants H3, H4 and H5 showed similar (H3) or slightly lower P_{app} values compared to IGF1R-4 V_{HH} ; H2 variant showed a significantly attenuated ability to cross *in vitro* BBB model, whereas H6 variants completely lost this ability, showing the same P_{app} value as internal control V_{HH} A20.1 (Figure 6C) and was excluded from further study. The P_{app} value for IGF1R-4mFc (Figure 6D) was similar to that of a 6-fold smaller IGF1R-4 V_{HH} (Figure 6B), indicating that IGF1R-4 can 'carry' large (antibody-size) molecules across the BBB. P_{app} value for the positive control, FC5mFc was 3-fold lower than that for IGF1R-4mFc. It is worth noting that constructs comprising IGF1R-5 or a humanized version linked to a cargo molecule (MW ~110kDa or 180kDa) have also been shown to be ferried across the BBB (data not shown).

[0113] Another IGF1R-binding V_{HH} , IGF1R-1 was also tested as mFc fusion in this assay. The IGF1R-1mFc had only marginally higher P_{app} value than control A20.1 V_{HH} (Figure 6D), indicating that only a subset of IGF1 R-binding V_{HH} s isolated from immune llama library has BBB-transmigrating properties.

[0114] *Absolute quantitation of V_{HH} s using MRM-ILIS method.* The methods are all as described in Haqqani et al. (2012). Briefly, to develop the SRM (selected reaction monitoring also known as multiple reaction monitoring (MRM) assay for V_{HH} , each V_{HH} was first analyzed by nanoLC-MS/MS using data-dependent acquisition to identify all ionizable

peptides. For each peptide, the 3 to 5 most intense fragment ions were chosen. An initial SRM assay was developed to monitor these fragments at attomole amounts of the digest (about 100-300 amol). Fragments that showed reproducible intensity ratios at low amounts (i.e., had Pearson $r^2 \geq 0.95$ compared to higher amounts) were considered stable and were chosen for the final SRM assay. To further optimize the assay, elution times for each peptide were also included, with care taken to not choose peptides that have close m/z (mass-to-charge ratio) and elution times.

[0115] A typical multiplexed SRM analysis of V_HH in cell media or body fluids (serum or cerebrospinal fluid (CSF)) involved spiking known amount of ILIS (0.1-10 nM) followed by injecting 100-400 ng of CSF or cultured media proteins (0.3-1 μ L) or about 50-100 ng of serum proteins (1-3 nanoliters) into the nanoLC-MS system. The precursor m/z of each target peptide ion was selected in the ion trap (and the remaining unrelated ions were discarded) at the specified elution time for the target, followed by collision induced dissociation (CID) fragmentation, and selection of only the desired fragment ions in the ion trap for monitoring by the detector. For quantification analysis, raw files generated by the LTQ (ThermoFisher) were converted to the standard mass spectrometry data format mzXML and intensities were extracted using an in-house software called Q-MRM (Quantitative-MRM; see Haqqani et al. 2012), which is a modified version of MatchRx software. For each V_HH , extracted-ion chromatograms were generated for each of its fragment ion that consisted of combined intensities within 0.25 Da of the fragment m/z over the entire elution time. To obtain a final intensity value for each fragment, all intensities within 0.5 min of the expected retention times were summed. A V_HH was defined as detectable in a sample if the fragments of at least one of its peptides showed the expected intensity ratios, i.e., the final intensity values showed a strong Pearson correlation $r \geq 0.95$ and $p < 0.05$ compared with the final intensities values of its corresponding pure V_HH .

[0116] Samples containing mixtures of V_HH (media, serum, CSF) were reduced, alkylated and trypsin-digested as previously described (Haqqani et al., 2012; Gergov et al., 2003). The digests (tryptic peptides) were acidified with acetic acid (5% final concentration) and analyzed on a reversed-phase nanoAcquity UPLC (Waters, Milford, MA) coupled to LTQ XL ETD or LTQ Orbitrap ETD mass spectrometer (ThermoFisher, Waltham, MA). The desired aliquot of the sample was injected and loaded onto a 300 μ m I.D. x 0.5 mm 3 μ m PepMaps C18 trap (ThermoFisher) then eluted onto a 100 μ m I.D. x 10 cm 1.7 μ m BEH130C18 nanoLC column (Waters) using a gradient from 0% - 20% acetonitrile (in 0.1% formic) in 1 minute, 20% - 46% in 16 min, and 46% - 95% in 1 min at a flow rate of 400 nL/min. The eluted peptides were ionized into the mass spectrometer by electrospray ionization (ESI) for MS/MS and SRM analysis using CID for fragmentation of the peptide ions. The CID was performed with helium as collision gas at normalized collision energy of 35% and 30 ms of activation time. Ion injection times into linear ion trap were adjusted by the instrument using an automatic gain control (AGC) target value of 6×10^3 and a maximum accumulation time of 200 ms

[0117] The V_HH -specific peptides used for detection and quantification of each V_HH in multiplexed assay are shown in Table 2.

Table 2. Peptides used in nanoLC-SRM detection of FC5, FC5-ILIS, EG2, A20.1, IGF-1R-5 and albumin. (a) In various studies described, assays were multiplexed in different combinations for simultaneous monitoring in the same sample; (b) Heavy-labeled peptide; (c) Limits of detection and quantification of the SRM assay for each peptide ranged from 1.5-2.5 ng/ml. 1 ng/mL corresponds to about 60-70 pM of V_HH . A20-1 as described in Hussack et al, 2011b; EG2 as described in Iqbal et al, 2010.

Protein	Signatures	SEQ ID NO:	Unique
IGF1R-4	EFVGHITWSR	25	Yes
	LSCEVSGGTVSPTAMGWFR	26	Yes
FC5	ITWGGDNTFYSNSVK	27	Yes
FC5-ILIS	ITWGGDNTFYSNSVK ^(b)	27	Yes
A20.1	TTYADSVK	28	Yes
	EFVAAGSSTGR	29	Yes
	TFSMDPMAWFR	30	Yes
	DEYAYWGQGTQVTVSSGQAGQGSEK	31	Yes
EG2	DFSDYVMGWFR	32	Yes
	LEESGGGLVQAGDSLRL	33	Yes
	NMYYLQMNSLKPEDTAVYYCAVNSAGTYVSPR	34	Yes
Albumin	APQVSTPTLVEAAR	35	Yes

Example 10: In vivo and Ex-vivo Optical Imaging

[0118] To determine whether IGF1R-4 V_HH accumulates in the brain after intravenous (iv) injection, IGF1R-4 V_HH and negative control A20.1 V_HH were conjugated to the near-infrared fluorescent tracer, Cy5.5 and purified as described in Example 7. 2.5 mg/kg of either IGF1 R-Cy5.5 or A20.1 Cy5.5 (in 150 μ l volume) were injected via tail vein into CD-1 nude mice, and they were imaged live in prone position in eXplore Optix preclinical imager MX2 (Advanced Research Technologies, QC) 30 min after injection. One hour after injections, animals were anesthetized with 2% isoflurane (Baxter Canada, Mississauga, ON, Canada), cardiac puncture was made and 10 mL of saline (Baxter Canada, Mississauga, ON, Canada) supplemented with 1 EU/mL heparin (Organon, Toronto, ON, Canada) was perfused from the same needle at 2 mL/min to remove the blood (ascertained by the appearance of blood-free fluid at the site of right ventricle incision). The perfused brains were removed and imaged *ex vivo* using in eXplore Optix. The imaging protocols were as described in detail in Iqbal et al., 2011.

[0119] The results are shown in Figure 7. Mice injected with Cy5.5-IGF1R-4 showed enhanced optical signal in the head region *in vivo* (upper panels, arrow) and in *ex-vivo* brain after perfusion (bottom panels, arrows), whereas Cy5.5-A20.1 - injected mice showed only low background signal. The data suggest that IGF1R-4 V_HH can deliver imaging agent (Cy5.5, 1kD) across the blood-brain barrier *in vivo*.

Example 11: Conjugation of IGF1R-4 to Galanin

[0120] To determine whether IGF1R-4 V_HH can cross the blood-brain barrier (BBB) *in vivo* and 'ferry' across the BBB a molecule that cannot cross the BBB on its own, the neuropeptide Galanin was chemically conjugated to IGF1R-4 V_HH or to IGF1R-4-mFc and administered systemically. Galanin is a neuroactive peptide that produces analgesia by binding GalR1 and GalR2 expressed in brain tissue. When given peripherally, Galanin has no analgesic effects because it cannot cross the BBB on its own (Robertson et al., 2011).

[0121] The IGF1R-4 V_HH was conjugated to a rat Galanin (Gal) fragment with cysteamide modified C-terminus (Bio-matic) (GWTLSAGYLLGPHAIDNHRFS SDKHGLT-cysteamide, SEQ ID NO:36). The scheme for conjugation is shown in Figure 8A.

[0122] Briefly, 5mg of IGF1R-4 V_HH (Example 4) in 0.5X PBS, 2.5mM EDTA at [2mg/ml] were mixed with 436.4 μ l of 2.5mg/ml sulfosuccinimidyl-4-(N-maleimidomethyl)cyclohexane-1-carboxylate (Sulfo-SMCC) (7.5x excess molar ratio). The mixture was then flushed with nitrogen gas and incubated for 30 minutes at room temperature (RT) to allow the NHS ester arm of the Sulfo-SMCC to react with amines on the V_HH. Subsequently the unreacted Sulfo-SMCC was removed from the maleimide-activated IGF1R-5 V_HH using a 10ml 7K Zeba column (Pierce). Prior to sample loading, the column was washed 3 times with 5ml PBS and spun at 1000xg for 2 min. After sample loading, the column was topped-up with 200 μ l of PBS and was spun for 2 min at 1000xg. As for the IGF1R-Fc constructs, 5 mg were reacted with ~68 μ l of Sulfo-SMCC (6.5x excess molar ratio) in the same way as described above.

[0123] Separately and concurrently, cysteamide modified C-TERM galanin (Gal-cya) was prepared by dissolving 10 mg of lyophilized powder in 10 ml of endotoxin free water to make a 1mg/ml stock (the galanin-cya powder has a small amount of DTT to prevent disulfide bridge formation during purification). Finally, 100 μ l of 0.5M EDTA was added (5mM final concentration).

[0124] The purified maleimide-activated IGF1R-4 V_HH and IGF1R-Fc constructs (2.6 ml) were diluted to 5ml with 0.5X PBS, 2.5mM EDTA and then 5ml or 1ml Gal-cya respectively was added while vortexing. The samples were flushed with nitrogen, sealed and incubated overnight at 4°C. The next day, the unreacted Gal-cya was removed using Amicon-15 10K and 30K column (Millipore) respectively. The sample were added to the column and spun at 4000xg for 7 minutes until the volume was reduced to 5ml. 5ml of 0.5X PBS, 2.5mM EDTA was added to the remaining 5ml sample in the column's insert and was spun again until the sample was reduced to 4ml. The conjugated samples were then added to a 10ml 7K Zeba column (Pierce), prepared as described above, and then spun for 2 min at 1000xg.

[0125] Conjugated IGF1R-4-Gal and IGF1R-4 Fc-Gal samples were then ran on a 16% or 10% SDS-PAGE non-reducing gel and silver stained to confirm shift in molecular weight size after conjugation. The reaction was titrated to achieve ~1 to 2 galanin molecules per V_HH or IGF1R-Fc constructs. Results confirm the galanin load on the IGF1R-4 V_HH (see Figure 8B).

[0126] *Endotoxin removal and determination of endotoxin levels:* Endotoxins were removed using Amicon Ultra - molecular weight cut-off (MWCO) cellulose membrane spin columns (Millipore). First, 15ml of the V_HH preparation was passed through an Amicon-15-50K MWCO column by centrifugation at 4000xg for 10minutes; the elution was collected. This elution was then added to an Amicon-15-10K MWCO column and spun at 4000xg for 7-10minutes resulting in a reduction of the supernatant volume from 15 ml to 7.5ml. The supernatant volume in the column was adjusted back to 15 ml by adding PBS. The column was spun again as described above. The supernatant was collected and the endotoxin levels were measured with EndoSafe-PTS system using cartridges with a sensitivity range of 10-0.1EU/ml (Charles River Laboratories International). 25 μ l of sample was loaded in each of the 4 wells on the cartridge and diluted if necessary.

Only samples with EU<1 per 1 mg were used for animal studies.

Example 12: Transport of the IGF1R-4-Gal using the Hargreaves model

[0127] To evaluate whether the IGF1R-4-Gal or IGF1R-4-mFc-Gal (Example 11) transmigrate the blood-brain barrier, an *in vivo* assay, was utilized, previously described in International Patent Publication No. WO 2011/127580.

[0128] A rat model of inflammatory hyperalgesia, similar to that described by Hargreaves et al. (1988), was used. Animals were housed in groups of three (Hargreaves model) per polypropylene cage, and were allowed free access to food and water. Experiments were done in a 12 h light/dark cycle at a temperature of 24°C and a relative humidity of 50 ± 5%. All animal procedures were approved by the NRC's Animal Care Committee and were in compliance with the Canadian Council of Animal Care guidelines.

[0129] In this model, male Wistar rats, 6-8 weeks (weight range 230-250 g) old were injected with low volume (100 µl with a 30-gauge needle) of complete Freund's adjuvant (CFA; heat-killed *M. tuberculosis* (Sigma) suspended in oil:saline 1:1 emulsion) into the right hind paw under brief isoflurane anesthesia (3%). CFA induces the release of pro-inflammatory substances that activate nociceptors and create a chronic pain state and hyperalgesia (a heightened sensitivity to noxious heat). The paw withdrawal latency was measured by the application of a radiant stimulus in the plantar surface in both hind paws (inflamed and non-inflamed control paw) using the plantar Analgesia Meter equipment for paw stimulation (IITC Model # 336TG Life Science, Inc.). The time taken by the animal to respond by licking or flicking its paw was interpreted as positive response (paw withdrawal latency). The light-intensity lamp was adjusted to elicit baseline paw withdrawal latencies between 17 to 20 s in both hind paws before CFA administration. If a withdrawal response does not occur within 20 s, the light beam was automatically turned off to avoid tissue damage and the paw was assigned the maximum score.

[0130] Two days post-CFA injection and prior to the administration of the compounds, the animals were manipulated and acclimatized in the analgesia meter equipment for at least 60 min with the aim to reduce stress and prevent false positive responses. The baseline was measured in both paws to verify the developed pain (thermal hyperalgesia); the non-inflamed paw was used as a control against the injected paw. Animals with paw withdrawal latency of more than 6 s for the "inflamed paw" and less than 17s for the "normal paw" were excluded from the experiment.

[0131] To determine whether IGF1R-4-Gal is delivered across the blood brain barrier and can engage target receptors (GalR1 and 2) in brain parenchyma, the rats received one tail vein injection of IGF1 R-4-Galanin (3mg/kg) or IGF1 R-4mFc-Galanin (2 mg/kg or 5 mg/kg) or control compounds three days post-CFA injection. The paw withdrawal latency (PWL) was tested for each hind paw (inflamed and non-inflamed) every 15 min for 3 hours. An increased latency of paw withdrawal indicates suppression of thermal hyperalgesia, which can only be obtained through successful delivery of Galanin into the brain parenchyma by IGF1R-4. Galanin can only induce analgesia when present in the brain parenchyma and on its own cannot cross an intact BBB (Robertson et al., 2011).

[0132] Results were analyzed as temporal courses of Paw Withdrawal Latencies (PWL, sec) versus time (min or hrs) (Figures 9A and C). Figure 9B shows the results as area under the curve (AUC) and compares it to the % of Maximum Possible Effect (%MPE). Figure 9D shows the results as percent reversal (of thermal hyperalgesia) at the peak response (Emax).

[0133] The results show that intravenously administered galanin does not reduce pain compared to PBS. In contrast, a single injection of FC5-Gal or IGF1 R4-Gal produce measurable analgesic effect, suggesting that this V_HH 'ferries' Galanin across the BBB to produce an analgesic effect by binding to GalR1 and/or 2 in the brain parenchyma. The effect of IGF1R-4-Galanin is significantly more pronounced than that induced by FC5-Gal, suggesting that the IGF1R receptor has a higher rate of BBB transport than the putative FC5 receptor. The results demonstrate that IGF1R-4 V_HH can 'ferry' molecules of at least 3000 Da across the BBB using receptor-mediated transcytosis pathway (the combined MW of antibody-peptide conjugate is about 18 kDa). Active, receptor mediated transport is required since the BBB is known to prevent the passage of all hydrophilic molecules greater than 0.5 kDa. Similarly, IGF1R4-mFc-Galanin produces a dose-dependent suppression of thermal hyperalgesia whereas A201.mFc-Galanin is ineffective. This suggests that IGF1R-4 V_HH can ferry molecules similar in size to antibodies (~80kD) across the BBB. The IGF1 R-4mFc-Galanin was about 5-fold more potent in suppressing thermal hyperalgesia than IGF1 R-4-Galanin, suggesting that bi-valent display and extended plasma half-life contribute to improved efficacy of brain delivery.

Example 13: IGF1R-4 levels in CSF and plasma

[0134] An *in vivo* assay was carried out to determine whether IGF1R-4 V_HH is able to cross into the brain, and specifically into the cerebrospinal fluid (CSF), as well as to quantify its presence in CSF and serum.

[0135] Animals were housed singly in polypropylene cages, and were allowed free access to food and water. Experiments were done in a 12 h light/dark cycle at a temperature of 24°C and a relative humidity of 50 ± 5%. All animal procedures were approved by the NRC's Animal Care Committee and were in compliance with the Canadian Council

of Animal Care guidelines.

[0136] Male Wistar rats, 8-10 weeks of age (weight range, 230-250 g) were used. To sample CSF, the fur on the neck and head region of animals was shaved and they were then placed in a Plexiglas chamber and moderately anesthetized with 3% isoflurane; the CSF was collected essentially as described by Nirogi et al (2009). The anesthetized rat was placed in a metal frame instrument (generously provided by Dr. Vinicio Granados-Soto; CINVESTAV, Mexico) and immobilised using earbars. The position of the animal's head was maintained downward at approximately 45°. A depressible surface with the appearance of a rhomb between the occipital protuberance and the spine of the atlas was made visible by rubbing the cotton embedded in ethanol (75%) over this surface. 27G needle covered with PE-10 tubing (Becton Dickinson, Mississauga, ON, Canada) 10 cm in length and connected to a 100 cc insulin syringe was inserted horizontally and centrally into the cisterna magna for CSF collection without making incisions. Two resistance points (clicks) along the needle path can be easily felt, due to the tearing of the skin and the ripping of atlanto-occipital membrane. When the needle passed the second resistance point, CSF was collected (40-100 µL) through the needle by applying a gentle suction of the insulin syringe. After the CSF sampling, corresponding blood was collected by cardiac puncture after thoracotomy and placed in vacutainer tubes (Becton Dickinson, Mississauga, ON, Canada) with clot activator and gel, and then centrifuged at 3,000xg for 15 min. Serum was removed using a micropipette and rapidly frozen at - 80°C until further analyses.

[0137] Serum and CSF samples were analyzed by mass spectrometry and nanoLC-SRM based quantification as described in Example 9.

[0138] CSF collection is a delicate procedure during which CSF can be easily contaminated with blood. Since the amounts of V_H s are expected to be much smaller in the CSF (<0.1%) than blood, even a slight contamination with blood could seriously compromise the value of an individual CSF sample. It was therefore necessary to develop stringent exclusion criteria for blood-contaminated CSF samples. To evaluate blood-CSF albumin ratio, a nanoLC-SRM method was developed for quantifying albumin levels in plasma and CSF. An albumin peptide APQVSTPTLVEAAR (SEQ ID NO:37) was selected based on its unique retention time and m/z value (Mol Pharm) in order to have minimum interference with other peptide peaks in the multiplex assay. The intensity of the peptide was quantified in both CSF and plasma samples using SRM as described above. The albumin ratio was calculated as follows for each rat:

$$\text{Albumin Ratio} = \text{Intensity per nL of plasma analyzed} / \text{Intensity per nL of CSF analyzed}$$

A ratio of 1500 and below was considered as blood contaminated.

[0139] Results are shown in Figure 10. Animals were co-injected with 6 mg/kg each of IGF1R-4, FC5 or A20.1, and CSF and serum were collected 30 min after injection. The figure shows that CSF/serum ratio for IGF1R-4 is 1.2%, compared to 0.98% for FC5 and 0.017% for A20.1, suggesting that both IGF1R-4 and FC5 have higher CSF penetration and brain exposure from co-injected control antibody, A20.1.

Example 14: Immunodetection of IGF1R-4-mFc

[0140] To ascertain that high levels of IGF1 R-4-mFc detected in the CSF after peripheral administration originate at least in part from parenchymal extracellular space, in other words, that the intact construct had crossed the BBB, immunodetection of IGF1R-4-mFc in rat brains was performed.

[0141] Briefly, brains of animals were harvested immediately following animal perfusion with PBS 24h after 6mg/kg tail-vein injection of either IGF1 R-4-mFc or A20.1 mFc. The brains were frozen and sectioned on cryotome into 12 µm sections. Sections were fixed for 10 min RT in 100% Methanol, washed 3x in PBS and incubated for 1 h in 10% normal goat serum (NGS) containing 0.3% TritonX-100 in 1xPBS. Goat anti-m-IgG Fcy-cy3 (Cat#115-165-071, Jackson Immuno Research, lot#106360) 1:200 in 5%NGS containing 0.3% TritonX-100 in 1XPBS was applied overnight at 4°C Sections were washed three times in 1x Vasculature-staining lectin RCAI (Cat# FL-1081, Vector) 1:500 in 1xPBS was then added for 10 min. After washing three times with 1xPBS, sections were covered with a cover slip in Dako fluorescent mounting medium (Cat#S3023, Dako) and spiked with 2µg/mL Hoechst (Cat#H3570, Invitrogen) to stain nuclei. Images were captured with Olympus 1X81 Fluorescent Microscope using 10X and 60X Objectives and channels as shown in Table 3.

Table 3. Objectives and channels used for fluorescent microscopy.

	Fluorescent molecule	Excitation (nm)	Emmision (nm)
RCAI-vessels	FITC	495	518
Hoechst33342- nuclei	Hoechst	350	461
IGF1R-5-m-Fc	Cy3	531	593

[0142] Results are shown in Figure 11. Immunodetection of mouse Fc showed strong staining of brain vessels throughout different brain regions, as well as staining of perivascular brain parenchyma, indicating that the IGF1 R-4-mFc is accumulated in brain vessels and also transmigrated the BBB into surrounding brain parenchyma. In contrast, no mFc-specific staining could be detected in A20.1mFc-injected animals. The results support the assertion that increased CSF levels of IGF1R-4-mFc are indicative of the construct transmigration across the BBB. This is further strongly supported by the observation that galanin linked to IGF1R-4 induced pharmacological response (analgesia) on parenchymal GalR1 and GalR2 receptors. Collectively, *in vitro* BBB transmigration results, *in vivo* pharmacokinetic (serum/CSF levels) and pharmacodynamic (Hargreaves model) results demonstrate that IGF1R-4 V_HH transmigrates the intact BBB at significantly higher rates than other V_HHs via active receptor-mediated transcytosis triggered by its binding to IGF1R epitopes and that it can 'ferry' a range (1-80kD) of otherwise non-permeable molecules across the blood-brain barrier.

Example 15: IGF1R-4 effect on 'physiological' function of IGF1R

[0143] From a safety perspective, it is important to show that the antibody of the invention does not interfere with the physiological function of the receptor- i.e., signaling induced by its natural ligand, IGF-1 - when engaging its receptor for drug delivery via receptor-mediated transcytosis. In view of this, it is important to demonstrate that IGF1R-4 V_HH or IGF1 R-4-mFc do not interfere with physiological signaling through IGF1R or the related insulin receptor (IR) induced by their natural ligands.

[0144] To determine whether IGF1R-4 induces signaling through IGF1R or IR alone, or interferes with signalling as stimulated by the receptor's natural ligands, IGF-1 or insulin, their effect on phosphorylation of the receptors themselves or receptor-stimulated downstream kinase, Akt, was determined in SV-ARBEK cells.

[0145] SV-ARBEK were grown to confluence in M199 base medium supplemented with peptone, D-glucose, BME amino acids, BME vitamins, antibiotic/antimycotic solution and fetal bovine serum, according to art-known methods. The cells were switched into serum free medium 18 h prior to treatment. IGF1R-4 V_HH or IGF1R-4-Fc fusion (100nM or 500nM) was added to the cells 1 h prior to the addition of either 200ng/ml IGF-1, 10 μ g/ml of insulin or vehicle. The cells were incubated with ligands or vehicle for 20 minutes and then washed twice in Hank's balanced salt solution. Cells were subsequently lysed using 1x RIPA buffer (Cell Signaling Technology) supplemented with 1% Triton-x 100 and protease inhibitor cocktail (Sigma). The cells were given 2 x 20 second bursts in a water bath sonicator and lysates were clarified by centrifugation at 14,000rpm for 10 minutes. Protein concentration was determined using DC protein assay system (BIO-RAD laboratories). Equal μ g of protein samples were resolved on a 4-20% gradient SDS polyacrylamide gel at 125V and transferred to PVDF membrane. Phospho-Akt (Ser 473) was detected by overnight incubation in 1:1000 dilution of the primary antibody against this target (Cell Signaling Technology) followed by a 1 h incubation with goat anti-rabbit IgG-HRP secondary antibody then reacted with ECL Plus reagent and visualized on autoradiography film. Densitometry values were determined using Un-Scan-It software (Silk Scientific Inc.).

[0146] The results are shown in Figure 12. Western blot analyses of Akt phosphorylation showed that IGF1R-4 did not inhibit Akt phosphorylation induced by 10 μ g/ml of insulin or by 200ng/ml of IGF-1 when co-applied with at 100 nM of IGF1R-4 or IGF1R-4-mFc or 500nM IGF1R-4-mFc. Neither did either of the V_HH or Fc fusions induce Akt signalling on its own (Fig 12A, 12B and 12C, labelled "-4"). The results demonstrate that even in bivalent display in the Fc fusion format IGF1R-4 does not trigger receptor dimerization and down-stream signaling, and therefore does not interfere with the receptor function in the presence of the natural ligand. This feature of IGF1R-4 ('silent binder') is important for its application as a BBB carrier for therapeutics, since it confers a favourable safety profile.

REFERENCES

[0147]

Abbott NJ (2013) Blood-brain barrier structure and function and the challenges for CNS drug delivery. J Inherit Metab Dis. 36(3):437-49.

Abulrob A, Sprong H, Van Bergen en Henegouwen P, Stanimirovic D (2005) The blood-brain barrier transmigrating single domain antibody: mechanisms of transport and antigenic epitopes in human brain endothelial cells. J Neurochem. 2005 Nov;95(4):1201-14.

Arbabi-Ghahroudi, M. Desmyter A, Wyns L, Hamers R., and Muyldermans S (1997) Selection and identification of single domain antibody fragments from camel heavy-chain antibodies, FEBS Lett 414, 521-526

Arbabi-Ghahroudi, M., To, R., Gaudette, N., Hiram, T., Ding, W., MacKenzie, R., and Tanha, J. (2009a) Protein Eng. Des. Sel. 22, 59-66.

Arbabi-Ghahroudi, M., MacKenzie, R., and Tanha, J. (2009b) Methods Mol. Biol. 525, 187-216.

Bell, A., Wang, Z.J., Arbabi-Ghahroudi, M., Chang, T.A., Durocher, Y., Trojahn, U., Baardsnes, J., Jaramillo, M.L., Li, S., Baral, T.N., O'Connor-McCourt, M., Mackenzie, R., and Zhang, J. (2010) Cancer Lett. 289, 81-90.

Broussau, s., Jabbour, N., Lachapelle, G., Durocher, Y., Tom, R., Transfiguracion, J., Gilbert, R. and Massie, B. (2008) Mol Ther 16, 500-507.

Chothia, C., and Lesk, A.M. (1987) J. Mol. Biol. 196, 901-917.

Davies J., and L. Riechmann, Affinity improvement of single antibody VH domains: residues in all three hypervariable regions affect antigen binding. Immunotechnology 2 (1996) 169-179

De Kruif, J., and Logtenberg, T. (1996) J. Biol. Chem. 271, 7630-7634.

Demeule, M.; Currie, J. C.; Bertrand, Y.; Che, C.; Nguyen, T.; Regina, A.; Gabathuler, R.; Castaigne, J. P.; Beliveau, R. Involvement of the low-density lipoprotein receptor-related protein in the transcytosis of the brain delivery vector angiopep-2, J. Neurochem. 2008, 106, 1534-1544.

Dumoulin, M., Conrath, K., Van Meirhaighe, A., Meersman, F., Heremans, K., Frenken, L.G., Muyldermans, S., Wyns, L., and Matagne, A. (2002) Protein Sci 11, 500-515.

Durocher, Y., S. Perret, et al. (2002). "High-level and high-throughput recombinant protein production by transient transfection of suspension-growing human 293-EBNA1 cells." Nucleic Acids Res 30(2): E9.

Doyle, P.J., Arbabi-Ghahroudi, M., Gaudette, N., Furzer, G., Savard, M.E., Gleddie, S., McLean, M.D., MacKenzie, C.R., and Hall, J.C. (2008) Mol. Immunol. 45, 3703-3713.

Eisenberg, D., Schwarz, E., Komaromy, M., and Wall, R. (1984) J. Mol. Biol. 179, 125-142

Erdlenbruch B, Alipour M, Fricker G, Miller DS, Kugler W, Eibl H, Lakomek M (2003) Alkylglycerol opening of the blood-brain barrier to small and large fluorescence markers in normal and C6 glioma-bearing rats and isolated rat brain capillaries. Br J Pharmacol. 140(7):1201-10.

Fenner, L., Widmer, A.F., Goy, G., Rudin, S., and Frei, R. (2008) J. Clin. Microbiol. 46, 328-330.

Gaillet, B., Gilbert, R., Broussau, S., Pilotte, A., Malenfant, F., Mullick, A., Garnier, A., and Massie, B. (2010) Biotechnol Bioeng 106, 203-215.

Gan Y, Jing Z, Stetler RA, Cao G (2013) Gene delivery with viral vectors for cerebrovascular diseases. Front Biosci (Elite Ed). 5:188-203. Review.

Garberg, P.; Ball, M.; Borg, N.; Cecchelli, R.; Fenart, L.; Hurst, R. D.; Lindmark, T.; Mabondzo, A.; Nilsson, J. E.; Raub, T. J.; Stanimirovic, D.; Terasaki, T.; Oberg, J. O.; Osterberg, T. In vitro models for the blood-brain barrier, Toxicol. In Vitro 2005, 19, 299-334.

Gergov, M.; Ojanpera, I.; Vuori, E. Simultaneous screening for 238 drugs in blood by liquid chromatography-ion spray tandem mass spectrometry with multiple-reaction monitoring, J. Chromatogr. B Analyt. Technol. Biomed. Life Sci. 2003, 795, 41-53.

Gaillet B, Gilbert R, Amziani R, Guilbault C, Gadoury C, Caron AW, Mullick A, Garnier A, Massie B (2007) High-level recombinant protein production in CHO cells using an adenoviral vector and the cumate gene-switch. Biotechnol

Prog. Jan-23(1):200-9

Gottesman et al., Ann. Rev. Biochem., 62, 385-427 (1993)

5 Hamers-Casterman, C., Atarhouch, T., Muyldermans, S., Robinson, G., Hamers, C., Songa, E.B., Bendahman, N., and Hamers, R. (1993) Nature 363, 446-448.

10 Haqqani AS, Caram-Salas N, Ding W, Brunette E, Delaney CE, Baumann E, Boileau E, Stanimirovic D (2012) Multiplexed evaluation of serum and CSF pharmacokinetics of brain-targeting single-domain antibodies using a NanoLC-SRM-ILIS method. Mol Pharm. 2013 May 6;10(5):1542-56.

Hargreaves KM, Troullos ES, Dionne RA, Schmidt EA, Schafer SC, Joris JL (1988) Bradykinin is increased during acute and chronic inflammation: therapeutic implications. Clin Pharmacol Ther. 44(6):613-21.

15 Huang YL, Säljö A, Suneson A, Hansson HA (1995) A new approach for multiple sampling of cisternal cerebrospinal fluid in rodents with minimal trauma and inflammation. J Neurosci Methods. 63(1-2):13-22.

Hussack, G., Hiram, T., Ding, W., MacKenzie, R., and Tanha, J. (2011) PLoS ONE 6, e28218.

20 Hussack G, Arbabi-Ghahroudi M, van Faassen H, Songer JG, Ng KK, MacKenzie R, Tanha J (2011b) Neutralization of Clostridium difficile toxin A with single-domain antibodies targeting the cell receptor binding domain. J Biol Chem. 286(11): 8961-76.

25 Iqbal U. Abulrob A. Stanimirovic D B (2011) Integrated platform for brain imaging and drug delivery across the blood-brain barrier. Methods Mol. Biol. 686, 465-481.

Iqbal, U., Trojahn, U., Albaghdadi, H., Zhang, J., O'Connor, M., Stanimirovic, D., Tomanek, B., Sutherland, G., and Abulrob, A. (2010) Br. J. Pharmacol. 160, 1016-1028.

30 Jespers, L., Schon, O., Famm, K., and Winter, G. (2004) Nat. Biotechnol. 22, 1161-1165.

Kabat EA, Wu TT. Identical V region amino acid sequences and segments of sequences in antibodies of different specificities. Relative contributions of VH and VL genes, minigenes, and complementarity-determining regions to binding of antibody-combining sites. J Immunol. 1991;147:1709-19.

35 Kim, D.Y., Kandalaf, H., Ding, W., Ryan, S., van Fassen, H., Hiram, T., Foote, S.J., MacKenzie, R., and Tanha, J. (2012) PEDS advance access August 30, 2012, 1-9.

40 Kornhuber ME, Kornhuber J, Cimniak U (1986) A method for repeated CSF sampling in the freely moving rat. J Neurosci Methods. 17(1):63-8.

Lefranc, M.-P. et al., (2003) Dev. Comp. Immunol., 27, 55-77.

45 Li S, Zheng W, Kuolee R, Hiram T, Henry M, Makvandi-Nejad S, Fjallman T, Chen W, Zhang J. Pentabody-mediated antigen delivery induces antigen-specific mucosal immune response. Mol Immunol 2009;46:1718-26.

Merritt, E.A., and Hol, W.G. (1995) Curr. Opin. Struct. Biol. 5, 165-171.

50 Muruganandam A, Tanha J, Narang S, Stanimirovic D (2001) Selection of phage-displayed llama single-domain antibodies that transigrate across human blood-brain barrier endothelium. FASEB J. 2002 Feb;16(2):240-2.

Musher, D.M., Manhas, A., Jain, P., Nuila, F., Waqar, A., Logan, N., Marino, B., Graviss, E.A. (2007) J. Clin. Microbiol. 45, 2737-2739.

55 Nhan T, Burgess A, Cho EE, Stefanovic B, Lilge L, Hynynen K. (2013) Drug delivery to the brain by focused ultrasound induced blood-brain barrier disruption: Quantitative evaluation of enhanced permeability of cerebral vasculature using two-photon microscopy. J Control Release. 172(1):274-280.

Nicaise M, Valeio-Lepiniec M, Minard P, Desmadril M. (2004) Affinity transfer by CDR grafting on a nonimmunoglobulin scaffold. *Protein Sci.* 13(7): 1882-1891.

Nielsen, U.B., Adams, G.P., Weiner, L.M., and Marks, J.D. (2000) *Cancer Res.* 60, 6434-6440.

Nirogi, R.; Kandikere, V.; Mudigonda, K.; Bhyrapuneni, G.; Muddana, N.; Saralaya, R.; Benade, V. (2009) A simple and rapid method to collect the cerebrospinal fluid of rats and its application for the assessment of drug penetration into the central nervous system, *J. Neurosci. Methods*, 178, 116-119.

Nuttall, S.D., Krishnan, U.V., Doughty, L., Pearson, K., Ryan, M.T., Hoogenraad, N.J., Hattarki, M., Carmichael, J.A., Irving, R.A., and Hudson, P.J. (2003) *Eur. J. Biochem.* 270, 3543-3554.

Pardridge, W. M.; Buciak, J. L.; Friden, P. M. Selective transport of an anti-transferrin receptor antibody through the blood-brain barrier in vivo, *J. Pharmacol. Exp. Ther.* 1991, 259, 66-70.

Pardridge, W.M., *Adv. Drug Delivery Reviews*, 15, 5-36 (1995)

Pardridge, W. M. Drug and gene delivery to the brain: the vascular route, *Neuron*. 2002, 36, 555-558.

Planche, T., Aghaizu, A., Holliman, R., Riley, P., Poloniecki, J., Breathnach, A., and Krishna, S. (2008) *Lancet Infect. Dis.* 8, 777-784.

Preston E, Slinn J, Vinokourov I, Stanimirovic D. (2008) Graded reversible opening of the rat blood-brain barrier by intracarotid infusion of sodium caprate. *J Neurosci Methods*. 168(2):443-9.

Ridgway, J.B., Presta, L.G., and Carter, P. (1996) *Protein Eng.* 9, 617-621.

Robertson CR, Flynn SP, White HS, Bulaj G (2011) Anticonvulsant neuropeptides as drug leads for neurological diseases. *Nat Prod Rep.* 28(4):741-62.

Russmann, H., Panthel, K., Bader, R.C., Schmitt, C., and Schaumann, R. (2007) *Eur. J. Clin. Microbiol. Infect. Dis.* 26, 115-119.

Samani, A.A., Chevet, E., Fallavollita, L., Galipeau, J., and Brodt, P. (2004) *Cancer Research* 64, 3380-3385.

Samuels B.L., *J. Clin. Pharmacol. Ther.*, 54, 421-429 (1993)

Sloan, L.M., Duresko, B.J., Gustafson, D.R., and Rosenblatt, J.E. (2008) *J. Clin. Microbiol.* 46, 1996-2001.

Sumbria RK, Zhou QH, Hui EK, Lu JZ, Boado RJ, Pardridge WM.(2013) Pharmacokinetics and brain uptake of an IgG-TNF decoy receptor fusion protein following intravenous, intraperitoneal, and subcutaneous administration in mice. *Mol Pharm.* 10(4):1425-31.

Tanha, J., Muruganandam, A., and Stanimirovic, D. (2003) *Methods Mol. Med.* 89, 435-449.

To, R., Hiram, T., Arbabi-Ghahroudi, M., MacKenzie, R., Wang, P., Xu, P., Ni, F., and Tanha, J. (2005) *J. Biol. Chem.* 280, 41395-41403.

Turgeon, D.K., Novicki, T.J., Quick, J., Carlson, L., Miller, P., Ulness, B., Cent, A., Ashley, R., Larson, A., Coyle, M., Limaye, A.P., Cookson, B.T., and Fritsche, T.R. (2003) *J. Clin. Microbiol.* 41, 667-670.

Watanabe, T., *Acta Oncol.*, 34, 235-241 (1995)

Xiao G, Gan LS.(2013) Receptor-mediated endocytosis and brain delivery of therapeutic biologics. *Int J Cell Biol.* doi: 10.1155/2013/703545. Epub 2013 Jun 11. Yaksh TL, Rudy TA (1976) Chronic catheterization of the spinal subarachnoid space. *Physiol Behav.* 17(6):1031-6.

Yu, Y. J.; Zhang, Y.; Kenrick, M.; Hoyte, K.; Luk, W.; Lu, Y.; Atwal, J.; Elliott, J. M.; Prabhu, S.; Watts, R. J.; Dennis,

EP 3 114 141 B1

M. S. Boosting brain uptake of a therapeutic antibody by reducing its affinity for a transcytosis target, Sci. Transl. Med. 2011, 3, 84ra44.

5 Zhang, J., Li, Q., Nguyen, T.-D., Tremblay, T.-L., Stone, E., To, R., Kelly, J., and MacKenzie, C.R. (2004a) J. Mol. Biol. 341, 161-169.

Zhang, J., Tanha, J., Hirama, T., Khiew, N.H., To, R., Tong-Sevinc, H., Stone, E., Brisson, J.R., and MacKenzie, C.R. (2004b) J. Mol. Biol. 335, 49-56.

10 Zhu et al., Immunology and Cell Biology (2010) 88:667-675.

European Patent No. 519596

European Patent No. 626390

15 US Patent No. 5693761

US Patent No. 5766886

20 US Patent No. 5821123

US Patent No. 5859205

US Patent No. 5869619

25 US Patent No. 6054297

US Patent No. 6180370

30 WO 02/057445

WO 2011/127580

WO 95/04069

35 WO/2004/076670

WO2003/046560

40 SEQUENCE LISTING

[0148]

<110> National Research Council of Canada

45 Stanimirovic, Danica

Kemmerich, Kristin

Sulea, Traian

Arbabi-Ghahroudi, Mehdi

50 Haqqani, Arsalan

Gilbert, Renald

Massie, Bernard

<120> Insulin-Like Growth Factor 1 Receptor-Specific Antibodies and Uses Thereof

55 <130> 12599-98

<150> 61/948, 818

<151> 2014-03-06

EP 3 114 141 B1

<160> 38

<170> PatentIn version 3.5

5 <210> 1
<211> 8
<212> PRT
<213> Artificial

10 <220>
<223> IGF1R-4 CDR1

<400> 1

15 Gly Gly Thr Val Ser Pro Thr Ala
1 5

20 <210> 2
<211> 8
<212> PRT
<213> Artificial

25 <220>
<223> IGF1R-4 CDR2

<400> 2

30 Ile Thr Trp Ser Arg Gly Thr Thr
1 5

35 <210> 3
<211> 17
<212> PRT
<213> Artificial

40 <220>
<223> IGF1R-4 CDR3

<400> 3

45 Ala Ala Ser Thr Phe Leu Arg Ile Leu Pro Glu Glu Ser Ala Tyr Thr
1 5 10 15

50 Tyr

55 <210> 4
<211> 124
<212> PRT
<213> Artificial

<220>
<223> IGF1R-4 Consensus

5 <220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> Xaa is Glu or Gln

10 <220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Xaa is Lys or Gln

15 <220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> Xaa is Val or Glu

20 <220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa is Ala or Pro

25 <220>
<221> MISC_FEATURE
<222> (23)..(23)
<223> Xaa is Ala or Glu

30 <220>
<221> MISC_FEATURE
<222> (24)..(24)
<223> Xaa is Val or Ala

35 <220>
<221> MISC_FEATURE
<222> (37)..(37)
<223> Xaa is Val or Phe

40 <220>
<221> MISC_FEATURE
<222> (44)..(44)
<223> Xaa is Gly or Glu

45 <220>
<221> MISC_FEATURE
<222> (45)..(45)
<223> Xaa is Leu or Arg

50 <220>
<221> MISC_FEATURE
<222> (47)..(47)
<223> Xaa is Phe or Trp

55 <220>
<221> MISC_FEATURE
<222> (49)..(49)
<223> Xaa is Gly or Ser

EP 3 114 141 B1

<220>
 <221> MISC_FEATURE
 <222> (60)..(60)
 <223> Xaa is Val or Tyr
 5
 <220>
 <221> MISC_FEATURE
 <222> (66)..(66)
 <223> Xaa is Asp or Gly
 10
 <220>
 <221> MISC_FEATURE
 <222> (74)..(74)
 <223> Xaa is Asn or Ser
 15
 <220>
 <221> MISC_FEATURE
 <222> (75)..(75)
 <223> Xaa is Ala or Ser
 20
 <220>
 <221> MISC_FEATURE
 <222> (79)..(79)
 <223> Xaa is Leu or Val
 25
 <220>
 <221> MISC_FEATURE
 <222> (87)..(87)
 <223> Xaa is Lys or Arg
 30
 <220>
 <221> MISC_FEATURE
 <222> (88)..(88)
 <223> Xaa is Ala or Ser
 35
 <220>
 <221> MISC_FEATURE
 <222> (119)..(119)
 <223> Xaa is Leu or Gln
 40
 <400> 4

 Xaa Val Xaa Leu Xaa Glu Ser Gly Gly Gly Leu Val Gln Xaa Gly Gly
 45
 1 5 10 15

50

55

EP 3 114 141 B1

Ser Leu Arg Leu Ser Cys Xaa Xaa Ser Gly Gly Thr Val Ser Pro Thr

30

5

Ala Met Gly Trp Xaa Arg Gln Ala Pro Gly Lys Xaa Xaa Glu Xaa Val

45

10

15

Xaa His Ile Thr Trp Ser Arg Gly Thr Thr Arg Xaa Ala Ser Ser Val

60

20

Lys Xaa Arg Phe Thr Ile Ser Arg Asp Xaa Xaa Lys Asn Thr Xaa Tyr

80

25

30

Leu Gln Met Asn Ser Leu Xaa Xaa Glu Asp Thr Ala Val Tyr Tyr Cys

95

35

40

Ala Ala Ser Thr Phe Leu Arg Ile Leu Pro Glu Glu Ser Ala Tyr Thr

110

45

Tyr Trp Gly Gln Gly Thr Xaa Val Thr Val Ser Ser

120

50

<210> 5

$\langle 211 \rangle$ 124

<212> PRT

<213> Artificial

<220>

55

EP 3 114 141 B1

<223> IGF1R-4 VHH

<400> 5

5	Gln Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
	1 5 10 15
10	
	Ser Leu Arg Leu Ser Cys Glu Val Ser Gly Gly Thr Val Ser Pro Thr
15	
	20 25 30
20	
	Ala Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Phe Val
	35 40 45
25	
	Gly His Ile Thr Trp Ser Arg Gly Thr Thr Arg Val Ala Ser Ser Val
30	
	50 55 60
35	
	Lys Asp Arg Phe Thr Ile Ser Arg Asp Ser Ala Lys Asn Thr Val Tyr
40	
	65 70 75 80
45	
50	
55	

EP 3 114 141 B1

Leu Gln Met Asn Ser Leu Lys Ser Glu Asp Thr Ala Val Tyr Tyr Cys

85

90

95

5

Ala Ala Ser Thr Phe Leu Arg Ile Leu Pro Glu Glu Ser Ala Tyr Thr

10

100

105

110

15

Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser

115

120

20

<210> 6

<211> 124

<212> PRT

<213> Artificial

25

<220>

<223> IGF1R-4_H2

<400> 6

30

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

1

5

10

15

35

Ser Leu Arg Leu Ser Cys Ala Val Ser Gly Gly Thr Val Ser Pro Thr

40

20

25

30

45

50

55

EP 3 114 141 B1

Ala Met Gly Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35

40

45

5

Gly His Ile Thr Trp Ser Arg Gly Thr Thr Arg Tyr Ala Ser Ser Val

10

50

55

60

15

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Val Tyr

65

70

75

80

20

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

25

85

90

95

30

Ala Ala Ser Thr Phe Leu Arg Ile Leu Pro Glu Glu Ser Ala Tyr Thr

35

100

105

110

40

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser

115

120

45

<210> 7

<211> 124

<212> PRT

<213> Artificial

50

<220>

<223> IGF1R-4_H3

<400> 7

55

EP 3 114 141 B1

	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
	1				5					10					15	
5																
	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Val	Ser	Gly	Gly	Thr	Val	Ser	Pro	Thr
10																
				20					25					30		
15																
	Ala	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Phe	Val
20			35					40					45			
25	Gly	His	Ile	Thr	Trp	Ser	Arg	Gly	Thr	Thr	Arg	Tyr	Ala	Ser	Ser	Val
	50					55					60					
30																
	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Val	Tyr
35	65					70					75				80	
40																
	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
				85						90				95		
45																
	Ala	Ala	Ser	Thr	Phe	Leu	Arg	Ile	Leu	Pro	Glu	Glu	Ser	Ala	Tyr	Thr
				100					105					110		
50																
	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser				
55																
				115					120							

<210> 8
<211> 124

EP 3 114 141 B1

<212> PRT
<213> Artificial

<220>
<223> IGF1R-4_H4

<400> 8

5

10

15

20

25

30

35

40

45

50

55

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Val Ser Gly Gly Thr Val Ser Pro Thr

20 25 30

Ala Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Phe Val

35 40 45

EP 3 114 141 B1

Gly His Ile Thr Trp Ser Arg Gly Thr Thr Arg Tyr Ala Ser Ser Val

50

55

60

5

Lys Gly Arg Phe Thr Ile Ser Arg Asp Ser Ser Lys Asn Thr Val Tyr

10

65

70

75

80

15

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

20

85

90

95

25

Ala Ala Ser Thr Phe Leu Arg Ile Leu Pro Glu Glu Ser Ala Tyr Thr

100

105

110

30

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser

35

115

120

<210> 9

<211> 124

<212> PRT

40

<213> Artificial

<220>

<223> IGF1R-4_H5

45

<400> 9

50

55

EP 3 114 141 B1

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

5

Ser Leu Arg Leu Ser Cys Ala Val Ser Gly Gly Thr Val Ser Pro Thr
10 20 25 30

15

Ala Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Phe Val
35 40 45

20

Gly His Ile Thr Trp Ser Arg Gly Thr Thr Arg Tyr Ala Ser Ser Val
25 50 55 60

30

Lys Gly Arg Phe Thr Ile Ser Arg Asp Ser Ser Lys Asn Thr Val Tyr
65 70 75 80

35

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

45

Ala Ala Ser Thr Phe Leu Arg Ile Leu Pro Glu Glu Ser Ala Tyr Thr
100 105 110

50

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

55

<210> 10
 <211> 370
 <212> PRT
 <213> Artificial
 5
 <220>
 <223> IGF1R-4 consensus-Fc fusion
 <220>
 10 <221> MISC_FEATURE
 <222> (1)..(1)
 <223> Xaa is Glu or Gln
 <220>
 15 <221> MISC_FEATURE
 <222> (3)..(3)
 <223> Xaa is Lys or Gln
 <220>
 20 <221> MISC_FEATURE
 <222> (5)..(5)
 <223> Xaa is Val or Glu
 <220>
 25 <221> MISC_FEATURE
 <222> (14)..(14)
 <223> Xaa is Ala or Pro
 <220>
 30 <221> MISC_FEATURE
 <222> (23)..(23)
 <223> Xaa is Ala or Glu
 <220>
 35 <221> MISC_FEATURE
 <222> (24)..(24)
 <223> Xaa is Val or Ala
 <220>
 40 <221> MISC_FEATURE
 <222> (37)..(37)
 <223> Xaa is Val or Phe
 <220>
 45 <221> MISC_FEATURE
 <222> (44)..(44)
 <223> Xaa is Gly or Glu
 <220>
 50 <221> MISC_FEATURE
 <222> (45)..(45)
 <223> Xaa is Leu or Arg
 <220>
 55 <221> MISC_FEATURE
 <222> (47)..(47)
 <223> Xaa is Phe or Trp

5
 <220>
 <221> MISC_FEATURE
 <222> (49)..(49)
 <223> Xaa is Gly or Ser

 10
 <220>
 <221> MISC_FEATURE
 <222> (60)..(60)
 <223> Xaa is Val or Tyr

 15
 <220>
 <221> MISC_FEATURE
 <222> (66)..(66)
 <223> Xaa is Asp or Gly

 20
 <220>
 <221> MISC_FEATURE
 <222> (74)..(74)
 <223> Xaa is Asn or Ser

 25
 <220>
 <221> MISC_FEATURE
 <222> (75)..(75)
 <223> Xaa is Ala or Ser

 30
 <220>
 <221> MISC_FEATURE
 <222> (79)..(79)
 <223> Xaa is Leu or Val

 35
 <220>
 <221> MISC_FEATURE
 <222> (87)..(87)
 <223> Xaa is Lys or Arg

 40
 <220>
 <221> MISC_FEATURE
 <222> (88)..(88)
 <223> Xaa is Ala or Ser

 45
 <220>
 <221> MISC_FEATURE
 <222> (119)..(119)
 <223> Xaa is Leu or Gln

 50
 <400> 10

 55

EP 3 114 141 B1

Xaa Val Xaa Leu Xaa Glu Ser Gly Gly Gly Leu Val Gln Xaa Gly Gly

1 5 10 15

5

Ser Leu Arg Leu Ser Cys Xaa Xaa Ser Gly Gly Thr Val Ser Pro Thr

10

20 25 30

15

Ala Met Gly Trp Xaa Arg Gln Ala Pro Gly Lys Xaa Xaa Glu Xaa Val

20 35 40 45

25

30

35

40

45

50

55

EP 3 114 141 B1

Xaa His Ile Thr Trp Ser Arg Gly Thr Thr Arg Xaa Ala Ser Ser Val

50

55

60

5

Lys Xaa Arg Phe Thr Ile Ser Arg Asp Xaa Xaa Lys Asn Thr Xaa Tyr

10

65

70

75

80

15

Leu Gln Met Asn Ser Leu Xaa Xaa Glu Asp Thr Ala Val Tyr Tyr Cys

85

90

95

20

Ala Ala Ser Thr Phe Leu Arg Ile Leu Pro Glu Glu Ser Ala Tyr Thr

25

100

105

110

30

Tyr Trp Gly Gln Gly Thr Xaa Val Thr Val Ser Ser Met Thr Val Asp

115

120

125

35

Lys Lys Leu Glu Pro Ser Gly Pro Ile Ser Thr Ile Asn Pro Cys Pro

40

130

135

140

45

Pro Cys Lys Glu Cys His Lys Cys Pro Ala Pro Asn Leu Glu Gly Gly

145

150

155

160

50

Pro Ser Val Phe Ile Phe Pro Pro Asn Ile Lys Asp Val Leu Met Ile

55

165

170

175

EP 3 114 141 B1

	Ser	Leu	Thr	Pro	Lys	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	Glu	Asp
5					180				185						190	
10	Asp	Pro	Asp	Val	Gln	Ile	Ser	Trp	Phe	Val	Asn	Asn	Val	Glu	Val	His
				195				200						205		
15																
20	Thr	Ala	Gln	Thr	Gln	Thr	His	Arg	Glu	Asp	Tyr	Asn	Ser	Thr	Ile	Arg
		210					215							220		
25	Val	Val	Ser	Thr	Leu	Pro	Ile	Gln	His	Gln	Asp	Trp	Met	Ser	Gly	Lys
		225				230					235				240	
30																
35	Glu	Phe	Lys	Cys	Lys	Val	Asn	Asn	Lys	Asp	Leu	Pro	Ser	Pro	Ile	Glu
				245						250					255	
40																
45	Arg	Thr	Ile	Ser	Lys	Ile	Lys	Gly	Leu	Val	Arg	Ala	Pro	Gln	Val	Tyr
				260					265					270		
50																
	Ile	Leu	Pro	Pro	Pro	Ala	Glu	Gln	Leu	Ser	Arg	Lys	Asp	Val	Ser	Leu
				275					280					285		
55																

EP 3 114 141 B1

	Thr	Cys	Leu	Val	Val	Gly	Phe	Asn	Pro	Gly	Asp	Ile	Ser	Val	Glu	Trp
5																
10	Thr	Ser	Asn	Gly	His	Thr	Glu	Glu	Asn	Tyr	Lys	Asp	Thr	Ala	Pro	Val
15																
20	Leu	Asp	Ser	Asp	Gly	Ser	Tyr	Phe	Ile	Tyr	Ser	Lys	Leu	Asn	Met	Lys
25	Thr	Ser	Lys	Trp	Glu	Lys	Thr	Asp	Ser	Phe	Ser	Cys	Asn	Val	Arg	His
30																
35	Glu	Gly	Leu	Lys	Asn	Tyr	Tyr	Leu	Lys	Lys	Thr	Ile	Ser	Arg	Ser	Pro
40																
45																
50																
55																

<210> 11
 <211> 370
 <212> PRT
 <213> Artificial

 <220>
 <223> IGF1R-4-Fc fusion

 <400> 11

EP 3 114 141 B1

Gln Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly

1 5 10 15

5

Ser Leu Arg Leu Ser Cys Glu Val Ser Gly Gly Thr Val Ser Pro Thr

10

20 25 30

15

Ala Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Phe Val

20

35 40 45

25

Gly His Ile Thr Trp Ser Arg Gly Thr Thr Arg Val Ala Ser Ser Val

50 55 60

30

Lys Asp Arg Phe Thr Ile Ser Arg Asp Ser Ala Lys Asn Thr Val Tyr

35

65 70 75 80

40

Leu Gln Met Asn Ser Leu Lys Ser Glu Asp Thr Ala Val Tyr Tyr Cys

85 90 95

45

Ala Ala Ser Thr Phe Leu Arg Ile Leu Pro Glu Glu Ser Ala Tyr Thr

50

100 105 110

55

EP 3 114 141 B1

Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser Met Thr Val Asp
115 120 125

5

Lys Lys Leu Glu Pro Ser Gly Pro Ile Ser Thr Ile Asn Pro Cys Pro
130 135 140

10

15

Pro Cys Lys Glu Cys His Lys Cys Pro Ala Pro Asn Leu Glu Gly Gly
145 150 155 160

20

Pro Ser Val Phe Ile Phe Pro Pro Asn Ile Lys Asp Val Leu Met Ile
165 170 175

25

30

Ser Leu Thr Pro Lys Val Thr Cys Val Val Val Asp Val Ser Glu Asp
180 185 190

35

40

Asp Pro Asp Val Gln Ile Ser Trp Phe Val Asn Asn Val Glu Val His
195 200 205

45

Thr Ala Gln Thr Gln Thr His Arg Glu Asp Tyr Asn Ser Thr Ile Arg
210 215 220

50

55

EP 3 114 141 B1

Val Val Ser Thr Leu Pro Ile Gln His Gln Asp Trp Met Ser Gly Lys
225 230 235 240

5

Glu Phe Lys Cys Lys Val Asn Asn Lys Asp Leu Pro Ser Pro Ile Glu
245 250 255

10

Arg Thr Ile Ser Lys Ile Lys Gly Leu Val Arg Ala Pro Gln Val Tyr
260 265 270

15

Ile Leu Pro Pro Pro Ala Glu Gln Leu Ser Arg Lys Asp Val Ser Leu
275 280 285

25

Thr Cys Leu Val Val Gly Phe Asn Pro Gly Asp Ile Ser Val Glu Trp
290 295 300

30

Thr Ser Asn Gly His Thr Glu Glu Asn Tyr Lys Asp Thr Ala Pro Val
305 310 315 320

35

Leu Asp Ser Asp Gly Ser Tyr Phe Ile Tyr Ser Lys Leu Asn Met Lys
325 330 335

45

Thr Ser Lys Trp Glu Lys Thr Asp Ser Phe Ser Cys Asn Val Arg His
340 345 350

50

55

EP 3 114 141 B1

Glu Gly Leu Lys Asn Tyr Tyr Leu Lys Lys Thr Ile Ser Arg Ser Pro

355

360

365

5

Gly Lys

10

370

<210> 12

<211> 1367

15

<212> PRT

<213> Artificial

<220>

<223> IGF1R

20

<400> 12

Met Lys Ser Gly Ser Gly Gly Gly Ser Pro Thr Ser Leu Trp Gly Leu

25

1

5

10

15

30

Leu Phe Leu Ser Ala Ala Leu Ser Leu Trp Pro Thr Ser Gly Glu Ile

20

25

30

35

Cys Gly Pro Gly Ile Asp Ile Arg Asn Asp Tyr Gln Gln Leu Lys Arg

40

35

40

45

45

50

55

EP 3 114 141 B1

Leu Glu Asn Cys Thr Val Ile Glu Gly Tyr Leu His Ile Leu Leu Ile

50

55

60

5

Ser Lys Ala Glu Asp Tyr Arg Ser Tyr Arg Phe Pro Lys Leu Thr Val

10

65

70

75

80

15

Ile Thr Glu Tyr Leu Leu Leu Phe Arg Val Ala Gly Leu Glu Ser Leu

20

85

90

95

25

Gly Asp Leu Phe Pro Asn Leu Thr Val Ile Arg Gly Trp Lys Leu Phe

100

105

110

30

Tyr Asn Tyr Ala Leu Val Ile Phe Glu Met Thr Asn Leu Lys Asp Ile

35

115

120

125

40

Gly Leu Tyr Asn Leu Arg Asn Ile Thr Arg Gly Ala Ile Arg Ile Glu

130

135

140

45

Lys Asn Ala Asp Leu Cys Tyr Leu Ser Thr Val Asp Trp Ser Leu Ile

50

145

150

155

160

55

EP 3 114 141 B1

Leu Asp Ala Val Ser Asn Asn Tyr Ile Val Gly Asn Lys Pro Pro Lys

165

170

175

5

Glu Cys Gly Asp Leu Cys Pro Gly Thr Met Glu Glu Lys Pro Met Cys

10

180

185

190

15

Glu Lys Thr Thr Ile Asn Asn Glu Tyr Asn Tyr Arg Cys Trp Thr Thr

195

200

205

20

Asn Arg Cys Gln Lys Met Cys Pro Ser Thr Cys Gly Lys Arg Ala Cys

25

210

215

220

30

Thr Glu Asn Asn Glu Cys Cys His Pro Glu Cys Leu Gly Ser Cys Ser

225

230

235

240

35

Ala Pro Asp Asn Asp Thr Ala Cys Val Ala Cys Arg His Tyr Tyr Tyr

40

245

250

255

45

Ala Gly Val Cys Val Pro Ala Cys Pro Pro Asn Thr Tyr Arg Phe Glu

260

265

270

50

Gly Trp Arg Cys Val Asp Arg Asp Phe Cys Ala Asn Ile Leu Ser Ala

55

275

280

285

EP 3 114 141 B1

	Glu Ser Ser Asp Ser Glu Gly Phe Val Ile His Asp Gly Glu Cys Met	
	290	295 300
5		
	Gln Glu Cys Pro Ser Gly Phe Ile Arg Asn Gly Ser Gln Ser Met Tyr	
10	305	310 315 320
15		
	Cys Ile Pro Cys Glu Gly Pro Cys Pro Lys Val Cys Glu Glu Glu Lys	
	325	330 335
20		
	Lys Thr Lys Thr Ile Asp Ser Val Thr Ser Ala Gln Met Leu Gln Gly	
25	340	345 350
30		
	Cys Thr Ile Phe Lys Gly Asn Leu Leu Ile Asn Ile Arg Arg Gly Asn	
35	355	360 365
40		
	Asn Ile Ala Ser Glu Leu Glu Asn Phe Met Gly Leu Ile Glu Val Val	
	370	375 380
45		
	Thr Gly Tyr Val Lys Ile Arg His Ser His Ala Leu Val Ser Leu Ser	
50	385	390 395 400
55		

EP 3 114 141 B1

Phe Leu Lys Asn Leu Arg Leu Ile Leu Gly Glu Glu Gln Leu Glu Gly
405 410 415

5

Asn Tyr Ser Phe Tyr Val Leu Asp Asn Gln Asn Leu Gln Gln Leu Trp
420 425 430

10

Asp Trp Asp His Arg Asn Leu Thr Ile Lys Ala Gly Lys Met Tyr Phe
435 440 445

15

20

Ala Phe Asn Pro Lys Leu Cys Val Ser Glu Ile Tyr Arg Met Glu Glu
450 455 460

25

30

Val Thr Gly Thr Lys Gly Arg Gln Ser Lys Gly Asp Ile Asn Thr Arg
465 470 475 480

35

Asn Asn Gly Glu Arg Ala Ser Cys Glu Ser Asp Val Leu His Phe Thr
485 490 495

40

45

Ser Thr Thr Thr Ser Lys Asn Arg Ile Ile Ile Thr Trp His Arg Tyr
500 505 510

50

Arg Pro Pro Asp Tyr Arg Asp Leu Ile Ser Phe Thr Val Tyr Tyr Lys
515 520 525

55

EP 3 114 141 B1

	Glu	Ala	Pro	Phe	Lys	Asn	Val	Thr	Glu	Tyr	Asp	Gly	Gln	Asp	Ala	Cys
	530						535					540				
5																
	Gly	Ser	Asn	Ser	Trp	Asn	Met	Val	Asp	Val	Asp	Leu	Pro	Pro	Asn	Lys
10	545					550					555				560	
15																
	Asp	Val	Glu	Pro	Gly	Ile	Leu	Leu	His	Gly	Leu	Lys	Pro	Trp	Thr	Gln
20					565					570					575	
25																
	Tyr	Ala	Val	Tyr	Val	Lys	Ala	Val	Thr	Leu	Thr	Met	Val	Glu	Asn	Asp
30					580					585				590		
35																
	His	Ile	Arg	Gly	Ala	Lys	Ser	Glu	Ile	Leu	Tyr	Ile	Arg	Thr	Asn	Ala
40					595					600				605		
45																
	Ser	Val	Pro	Ser	Ile	Pro	Leu	Asp	Val	Leu	Ser	Ala	Ser	Asn	Ser	Ser
50					610					615				620		
55																
	Ser	Gln	Leu	Ile	Val	Lys	Trp	Asn	Pro	Pro	Ser	Leu	Pro	Asn	Gly	Asn
60	625					630					635				640	
65																

EP 3 114 141 B1

Leu Ser Tyr Tyr Ile Val Arg Trp Gln Arg Gln Pro Gln Asp Gly Tyr

645

650

655

5

Leu Tyr Arg His Asn Tyr Cys Ser Lys Asp Lys Ile Pro Ile Arg Lys

10

660

665

670

15

Tyr Ala Asp Gly Thr Ile Asp Ile Glu Glu Val Thr Glu Asn Pro Lys

675

680

685

20

Thr Glu Val Cys Gly Gly Glu Lys Gly Pro Cys Cys Ala Cys Pro Lys

25

690

695

700

30

Thr Glu Ala Glu Lys Gln Ala Glu Lys Glu Glu Ala Glu Tyr Arg Lys

705

710

715

720

35

Val Phe Glu Asn Phe Leu His Asn Ser Ile Phe Val Pro Arg Pro Glu

40

725

730

735

45

Arg Lys Arg Arg Asp Val Met Gln Val Ala Asn Thr Thr Met Ser Ser

740

745

750

50

Arg Ser Arg Asn Thr Thr Ala Ala Asp Thr Tyr Asn Ile Thr Asp Pro

55

755

760

765

EP 3 114 141 B1

Glu Glu Leu Glu Thr Glu Tyr Pro Phe Phe Glu Ser Arg Val Asp Asn

770

775

780

5

Lys Glu Arg Thr Val Ile Ser Asn Leu Arg Pro Phe Thr Leu Tyr Arg

10

785

790

795

800

15

Ile Asp Ile His Ser Cys Asn His Glu Ala Glu Lys Leu Gly Cys Ser

20

805

810

815

25

Ala Ser Asn Phe Val Phe Ala Arg Thr Met Pro Ala Glu Gly Ala Asp

820

825

830

30

Asp Ile Pro Gly Pro Val Thr Trp Glu Pro Arg Pro Glu Asn Ser Ile

35

835

840

845

40

Phe Leu Lys Trp Pro Glu Pro Glu Asn Pro Asn Gly Leu Ile Leu Met

850

855

860

45

Tyr Glu Ile Lys Tyr Gly Ser Gln Val Glu Asp Gln Arg Glu Cys Val

50

865

870

875

880

55

EP 3 114 141 B1

Ser Arg Gln Glu Tyr Arg Lys Tyr Gly Gly Ala Lys Leu Asn Arg Leu

885

890

895

5

Asn Pro Gly Asn Tyr Thr Ala Arg Ile Gln Ala Thr Ser Leu Ser Gly

10

900

905

910

15

Asn Gly Ser Trp Thr Asp Pro Val Phe Phe Tyr Val Gln Ala Lys Thr

915

920

925

20

Gly Tyr Glu Asn Phe Ile His Leu Ile Ile Ala Leu Pro Val Ala Val

25

930

935

940

30

Leu Leu Ile Val Gly Gly Leu Val Ile Met Leu Tyr Val Phe His Arg

945

950

955

960

35

Lys Arg Asn Asn Ser Arg Leu Gly Asn Gly Val Leu Tyr Ala Ser Val

40

965

970

975

45

Asn Pro Glu Tyr Phe Ser Ala Ala Asp Val Tyr Val Pro Asp Glu Trp

980

985

990

50

Glu Val Ala Arg Glu Lys Ile Thr Met Ser Arg Glu Leu Gly Gln Gly

55

995

1000

1005

EP 3 114 141 B1

Ser Phe Gly Met Val Tyr Glu Gly Val Ala Lys Gly Val Val Lys

1010

1015

1020

5

Asp Glu Pro Glu Thr Arg Val Ala Ile Lys Thr Val Asn Glu Ala

1025

1030

1035

10

15

Ala Ser Met Arg Glu Arg Ile Glu Phe Leu Asn Glu Ala Ser Val

1040

1045

1050

20

Met Lys Glu Phe Asn Cys His His Val Val Arg Leu Leu Gly Val

1055

1060

1065

25

30

Val Ser Gln Gly Gln Pro Thr Leu Val Ile Met Glu Leu Met Thr

1070

1075

1080

35

40

Arg Gly Asp Leu Lys Ser Tyr Leu Arg Ser Leu Arg Pro Glu Met

1085

1090

1095

45

Glu Asn Asn Pro Val Leu Ala Pro Pro Ser Leu Ser Lys Met Ile

1100

1105

1110

50

55

EP 3 114 141 B1

Gln Met Ala Gly Glu Ile Ala Asp Gly Met Ala Tyr Leu Asn Ala

1115

1120

1125

5

Asn Lys Phe Val His Arg Asp Leu Ala Ala Arg Asn Cys Met Val

10

1130

1135

1140

15

Ala Glu Asp Phe Thr Val Lys Ile Gly Asp Phe Gly Met Thr Arg

1145

1150

1155

20

Asp Ile Tyr Glu Thr Asp Tyr Tyr Arg Lys Gly Gly Lys Gly Leu

25

1160

1165

1170

30

Leu Pro Val Arg Trp Met Ser Pro Glu Ser Leu Lys Asp Gly Val

1175

1180

1185

35

Phe Thr Thr Tyr Ser Asp Val Trp Ser Phe Gly Val Val Leu Trp

40

1190

1195

1200

45

Glu Ile Ala Thr Leu Ala Glu Gln Pro Tyr Gln Gly Leu Ser Asn

1205

1210

1215

50

Glu Gln Val Leu Arg Phe Val Met Glu Gly Gly Leu Leu Asp Lys

55

1220

1225

1230

EP 3 114 141 B1

Pro Asp Asn Cys Pro Asp Met Leu Phe Glu Leu Met Arg Met Cys

1235

1240

1245

5

Trp Gln Tyr Asn Pro Lys Met Arg Pro Ser Phe Leu Glu Ile Ile

1250

1255

1260

10

15

Ser Ser Ile Lys Glu Glu Met Glu Pro Gly Phe Arg Glu Val Ser

1265

1270

1275

20

Phe Tyr Tyr Ser Glu Glu Asn Lys Leu Pro Glu Pro Glu Glu Leu

1280

1285

1290

25

30

Asp Leu Glu Pro Glu Asn Met Glu Ser Val Pro Leu Asp Pro Ser

1295

1300

1305

35

40

Ala Ser Ser Ser Ser Leu Pro Leu Pro Asp Arg His Ser Gly His

1310

1315

1320

45

Lys Ala Glu Asn Gly Pro Gly Pro Gly Val Leu Val Leu Arg Ala

1325

1330

1335

50

55

EP 3 114 141 B1

Ser Phe Asp Glu Arg Gln Pro Tyr Ala His Met Asn Gly Gly Arg

1340

1345

1350

5

Lys Asn Glu Arg Ala Leu Pro Leu Pro Gln Ser Ser Thr Cys

10

1355

1360

1365

<210> 13

<211> 4

15

<212> PRT

<213> Artificial

<220>

<223> Furin cleavage site

20

<400> 13

Arg Lys Arg Arg

25

1

<210> 14

<211> 33

<212> DNA

30

<213> Artificial

<220>

<223> Forward primer

35

<400> 14

cgggatccgc caccatgaag tctggctccg gag 33

<210> 15

<211> 34

40

<212> DNA

<213> Artificial

<220>

<223> Reverse primer

45

<400> 15

gctctagatc agaagtttc atatcctgtt ttgg 34

<210> 16

50

<211> 45

<212> DNA

<213> Artificial

<220>

55

<223> MJ1 primer

<400> 16

gcccagccgg ccatggccsm kgtgcagctg gtggaktctg gggga 45

EP 3 114 141 B1

<210> 17
 <211> 45
 <212> DNA
 <213> Artificial
 5
 <220>
 <223> MJ2 primer
 <400> 17
 10 gccagccgg ccatggcca ggtaaagctg gaggagtctg gggga 45
 <210> 18
 <211> 45
 <212> DNA
 15 <213> Artificial
 <220>
 <223> MJ3 primer
 20 <400> 18
 gccagccgg ccatggcca ggctcaggtg cagctggtgg agtct 45
 <210> 19
 <211> 21
 25 <212> DNA
 <213> Artificial
 <220>
 <223> CH2 primer
 30 <400> 19
 cgccatcaag gtaccagtg a 21
 <210> 20
 35 <211> 29
 <212> DNA
 <213> Artificial
 <220>
 40 <223> CH2b3 primer
 <400> 20
 ggggtacctg tcatccacgg accagctga 29
 45 <210> 21
 <211> 34
 <212> DNA
 <213> Artificial
 50 <220>
 <223> MJ7 primer
 <400> 21
 55 catgtgtaga ctgcggccc agccggccat ggcc 34
 <210> 22
 <211> 47
 <212> DNA

EP 3 114 141 B1

<213> Artificial
 <220>
 <223> MJ8 primer
 5
 <400> 22
 catgtgtaga ttcctggccg gcctggcctg aggagacggt gacctgg 47
 <210> 23
 10 <211> 38
 <212> DNA
 <213> Artificial
 <220>
 15 <223> Forward primer
 <400> 23
 tatgaagaca ccaggcccag gtaaagctgg aggagtct 38
 20 <210> 24
 <211> 30
 <212> DNA
 <213> Artificial
 25 <220>
 <223> Reverse primer
 <400> 24
 ttgttcggat cctgaggaga cggtgacctg 30
 30 <210> 25
 <211> 10
 <212> PRT
 <213> Artificial
 35 <220>
 <223> IGF1R-4 peptide
 <400> 25
 40
 Glu Phe Val Gly His Ile Thr Trp Ser Arg
 1 5 10
 45 <210> 26
 <211> 19
 <212> PRT
 <213> Artificial
 50 <220>
 <223> IGF1R-4 peptide
 <400> 26
 55

EP 3 114 141 B1

Leu Ser Cys Glu Val Ser Gly Gly Thr Val Ser Pro Thr Ala Met Gly

1 5 10 15

5

Trp Phe Arg

10

<210> 27
 <211> 15
 <212> PRT
 <213> Artificial

15

<220>
 <223> FC5 and FC5-ILIS peptide

<400> 27

20

Ile Thr Trp Gly Gly Asp Asn Thr Phe Tyr Ser Asn Ser Val Lys

1 5 10 15

25

<210> 28
 <211> 9
 <212> PRT
 <213> Artificial

30

<220>
 <223> A20.1 peptide

<400> 28

35

Thr Thr Tyr Tyr Ala Asp Ser Val Lys

1 5

40

<210> 29
 <211> 11
 <212> PRT
 <213> Artificial

45

<220>
 <223> A20.1 peptide

<400> 29

50

Glu Phe Val Ala Ala Gly Ser Ser Thr Gly Arg

1 5 10

55

<210> 30
 <211> 11
 <212> PRT
 <213> Artificial

EP 3 114 141 B1

<220>

<223> A20.1 peptide

<400> 30

5

Thr Phe Ser Met Asp Pro Met Ala Trp Phe Arg

1 5 10

10

<210> 31

<211> 26

<212> PRT

<213> Artificial

15

<220>

<223> A20.1 peptide

<400> 31

20

Asp Glu Tyr Ala Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser

1 5 10 15

25

Gly Gln Ala Gly Gln Gly Ser Glu Gln Lys

30

20 25

<210> 32

35

<211> 11

<212> PRT

<213> Artificial

<220>

40

<223> EG2 peptide

<400> 32

Asp Phe Ser Asp Tyr Val Met Gly Trp Phe Arg

45

1 5 10

<210> 33

50

<211> 16

<212> PRT

<213> Artificial

<220>

55

<223> EG2 peptide

<400> 33

EP 3 114 141 B1

Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Asp Ser Leu Arg

1 5 10 15

5

<210> 34
 <211> 32
 <212> PRT
 <213> Artificial

10

<220>
 <223> EG2 peptide

15

<400> 34

Asn Met Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala

1 5 10 15

20

Val Tyr Tyr Cys Ala Val Asn Ser Ala Gly Thr Tyr Val Ser Pro Arg

20 25 30

25

<210> 35
 <211> 14
 <212> PRT
 <213> Artificial

30

<220>
 <223> Albumin peptide

35

<400> 35

Ala Pro Gln Val Ser Thr Pro Thr Leu Val Glu Ala Ala Arg

1 5 10

40

<210> 36
 <211> 29
 <212> PRT
 <213> Artificial

45

<220>
 <223> Galanin-cysteamide

50

<400> 36

55

EP 3 114 141 B1

Gly Trp Thr Leu Asn Ser Ala Gly Tyr Leu Leu Gly Pro His Ala Ile

1 5 10 15

5

Asp Asn His Arg Ser Phe Ser Asp Lys His Gly Leu Thr

10

20 25

<210> 37

<211> 19

15

<212> PRT

<213> Artificial

<220>

<223> Signal peptide

20

<400> 37

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Leu Lys Gly

25

1 5 10 15

30

Val Gln Cys

35

<210> 38

<211> 371

<212> PRT

<213> Artificial

40

<220>

<223> Fc-IGF1R-4 consensus fusion

<220>

<221> MISC_FEATURE

45

<222> (248)..(248)

<223> Xaa is Glu or Gln

<220>

<221> MISC_FEATURE

50

<222> (250)..(250)

<223> Xaa is Lys or Gln

<220>

<221> MISC_FEATURE

55

<222> (252)..(252)

<223> Xaa is Val or Glu

<220>

	<221> MISC_FEATURE
	<222> (261)..(261)
	<223> Xaa is Ala or Pro
5	<220>
	<221> MISC_FEATURE
	<222> (270)..(270)
	<223> Xaa is Ala or Glu
10	<220>
	<221> MISC_FEATURE
	<222> (271)..(271)
	<223> Xaa is Val or Ala
15	<220>
	<221> MISC_FEATURE
	<222> (284)..(284)
	<223> Xaa is Val or Phe
20	<220>
	<221> MISC_FEATURE
	<222> (291)..(291)
	<223> Xaa is Gly or Glu
25	<220>
	<221> MISC_FEATURE
	<222> (292)..(292)
	<223> Xaa is Leu or Arg
30	<220>
	<221> MISC_FEATURE
	<222> (294)..(294)
	<223> Xaa is Phe or Trp
35	<220>
	<221> MISC_FEATURE
	<222> (296)..(296)
	<223> Xaa is Gly or Ser
40	<220>
	<221> MISC_FEATURE
	<222> (307)..(307)
	<223> Xaa is Val or Tyr
45	<220>
	<221> MISC_FEATURE
	<222> (313)..(313)
	<223> Xaa is Asp or Gly
50	<220>
	<221> MISC_FEATURE
	<222> (321)..(321)
	<223> Xaa is Asn or Ser
55	<220>
	<221> MISC_FEATURE
	<222> (322)..(322)
	<223> Xaa is Ala or Ser

EP 3 114 141 B1

<220>
 <221> MISC_FEATURE
 <222> (326)..(326)
 <223> Xaa is Leu or Val
 5
 <220>
 <221> MISC_FEATURE
 <222> (334)..(334)
 <223> Xaa is Lys or Arg
 10
 <220>
 <221> MISC_FEATURE
 <222> (335)..(335)
 <223> Xaa is Ala or Ser
 15
 <220>
 <221> MISC_FEATURE
 <222> (366)..(366)
 <223> Xaa is Leu or Gln
 20
 <400> 38

 Glu Pro Ser Gly Pro Ile Ser Thr Ile Asn Pro Cys Pro Pro Cys Lys
 25 1 5 10 15

 30
 Glu Cys His Lys Cys Pro Ala Pro Asn Leu Glu Gly Gly Pro Ser Val
 20 25 30
 35

 Phe Ile Phe Pro Pro Asn Ile Lys Asp Val Leu Met Ile Ser Leu Thr
 40 35 40 45

 45
 Pro Lys Val Thr Cys Val Val Val Asp Val Ser Glu Asp Asp Pro Asp
 50 55 60
 50

 Val Gln Ile Ser Trp Phe Val Asn Asn Val Glu Val His Thr Ala Gln
 65 70 75 80

EP 3 114 141 B1

Thr Gln Thr His Arg Glu Asp Tyr Asn Ser Thr Ile Arg Val Val Ser

85

90

95

5

Thr Leu Pro Ile Gln His Gln Asp Trp Met Ser Gly Lys Glu Phe Lys

100

105

110

10

15

Cys Lys Val Asn Asn Lys Asp Leu Pro Ser Pro Ile Glu Arg Thr Ile

115

120

125

20

Ser Lys Ile Lys Gly Leu Val Arg Ala Pro Gln Val Tyr Ile Leu Pro

130

135

140

25

30

Pro Pro Ala Glu Gln Leu Ser Arg Lys Asp Val Ser Leu Thr Cys Leu

145

150

155

160

35

40

Val Val Gly Phe Asn Pro Gly Asp Ile Ser Val Glu Trp Thr Ser Asn

165

170

175

45

Gly His Thr Glu Glu Asn Tyr Lys Asp Thr Ala Pro Val Leu Asp Ser

180

185

190

50

55

EP 3 114 141 B1

Asp Gly Ser Tyr Phe Ile Tyr Ser Lys Leu Asn Met Lys Thr Ser Lys
 195 200 205
 5
 Trp Glu Lys Thr Asp Ser Phe Ser Cys Asn Val Arg His Glu Gly Leu
 210 215 220
 10
 Lys Asn Tyr Tyr Leu Lys Lys Thr Ile Ser Arg Ser Pro Gly Gly Gly
 225 230 235 240
 15
 Gly Ser Gly Gly Gly Gly Ser Xaa Val Xaa Leu Xaa Glu Ser Gly Gly
 245 250 255
 20
 Gly Leu Val Gln Xaa Gly Gly Ser Leu Arg Leu Ser Cys Xaa Xaa Ser
 260 265 270
 25
 Gly Gly Thr Val Ser Pro Thr Ala Met Gly Trp Xaa Arg Gln Ala Pro
 275 280 285
 30
 Gly Lys Xaa Xaa Glu Xaa Val Xaa His Ile Thr Trp Ser Arg Gly Thr
 290 295 300
 35
 Thr Arg Xaa Ala Ser Ser Val Lys Xaa Arg Phe Thr Ile Ser Arg Asp
 305 310 315 320
 40
 45
 50
 55

EP 3 114 141 B1

Xaa Xaa Lys Asn Thr Xaa Tyr Leu Gln Met Asn Ser Leu Xaa Xaa Glu

325

330

335

5

Asp Thr Ala Val Tyr Tyr Cys Ala Ala Ser Thr Phe Leu Arg Ile Leu

10

340

345

350

15

Pro Glu Glu Ser Ala Tyr Thr Tyr Trp Gly Gln Gly Thr Xaa Val Thr

20

355

360

365

25

Val Ser Ser

370

30

Claims

1. An isolated or purified immunoglobulin single variable domain antibody, comprising a complementarity determining region (CDR) 1 sequence of GGTVSPTA (SEQ ID NO:1); a CDR2 sequence of ITWSRGTT (SEQ ID NO:2); and
35 a CDR3 sequence of AASTFLRILPEESAYTY (SEQ ID NO:3), wherein the antibody is specific for Insulin-Like Growth Factor 1 Receptor (IGF1R) and wherein the antibody transmigrates the blood-brain barrier.

40

2. The isolated or purified antibody according to claim 1, comprising the sequence X₁VX₂LX₃ESGGGLVQX₄GGSLRLSCX₅X₆SGGTVSPTAMGWX₇RQAPGKX₈X₉EX₁₀VX₁₁HITWSRGTTX₁₂AS SVKX₁₃RFTISRDX₁₄X₁₅KNTX₁₆YLQMNSLX₁₇X₁₈EDTAVYYCAASTFLRILPEESAYTYWGQGT X₁₉VTVSS (SEQ ID NO:4), where X₁ is E or Q; X₂ is K or Q; X₃ is V or E; X₄ is A or P; X₅ is A or E; X₆ is V or A; X₇ is V or F; X₈ is G or E; X₉ is L or R; X₁₀ is F or W; X₁₁ is G or S; X₁₂ is V or Y; X₁₃ is D or G; X₁₄ is N or S; X₁₅ is A or S; X₁₆ is L or V; X₁₇ is K or R; X₁₈ is A or S; and X₁₉ is L or Q.

45

3. The isolated or purified antibody any one of claims 1 to 2, comprising a sequence selected from the group consisting of:

QVKLEESGGGLVQAGGSLRLSCEVSGGTVSPTAMGWFRQAPGKEREFVGHITWSRGTTRVASSVKDRFT

50

ISRDSAKNTVYLQMNSLKSEDTAVYYCAASTFLRILPEESAYTYWGQGTQVTVSS (SEQ ID NO:5);

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWVRQAPGKGLEWVGHITWSRGTTTRYASSVKGRF

55

TISRDNKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQGTQVTVSS (SEQ ID NO:6);

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKGLEFVGHITWSRGTTTRYASSVKGRFT
ISRDNSKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQGLTVTVSS (SEQ ID NO:7);

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKGLEFVGHITWSRGTTTRYASSVKGRFT
ISRDSSKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQGLTVTVSS (SEQ ID NO:8);

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKEREFVGHITWSRGTTTRYASSVKGRFT
ISRDSSKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQGLTVTVSS (SEQ ID NO:9);

and/
or a sequence having 90% identity thereto.

4. The isolated or purified antibody according to any one of claims 1 to 3, wherein the antibody is in a multivalent display format.
5. The isolated or purified antibody according to claim 4, wherein the antibody is linked to a Fc fragment.
6. A nucleic acid molecule encoding the isolated or purified antibody of any one of claims 1 to 5.
7. A vector comprising the nucleic acid molecule of claim 6.
8. The isolated or purified antibody according to any one of claims 1 to 5, wherein the antibody is linked to a cargo molecule.
9. The isolated or purified antibody according to claim 8, wherein the cargo molecule has a molecular weight in the range of about 1kD to about 200 kDa.
10. The isolated or purified antibody according to claim 9 wherein the cargo molecule is selected from the group consisting of a detectable agent, a therapeutic, a drug, a peptide, a growth factor, a cytokine, a receptor trap, a chemical compound, a carbohydrate moiety, one or more liposomes or nanocarriers loaded with a detectable agent, a therapeutic, a drug, a peptide, an enzyme, an antibody or fragment thereof, a DNA-based molecule, a viral vector, or a cytotoxic agent, or one or more nanoparticle, nanowire, nanotube, or quantum dots.
11. A composition comprising one or more than one isolated or purified antibody of any one of claims 1 to 5 and 9 to 10 and a pharmaceutically acceptable carrier, diluent, or excipient.
12. The isolated or purified antibody of any one of claims 1 to 5 and 9 to 10 or the composition according to claim 11 for use in medicine.
13. An *in vitro* method of detecting Insulin-Like Growth Factor 1 Receptor (IGF1R), comprising
 - a) contacting a tissue sample with one or more than one isolated or purified antibody of any one of claims 1 to 5 linked to a detectable agent; and
 - b) detecting the detectable agent linked to the antibody bound to IGF1R in the tissue sample.

Patentansprüche

1. Isolierter oder gereinigter Immunglobulin-Variable-Einzeldomäne-Antikörper, umfassend eine Komplementarität-bestimmende-Region- (CDR-) 1-Sequenz aus GGTVSPTA (SEQ ID NO:1); eine CDR2-Sequenz aus ITWSRGTT (SEQ ID NO:2) und eine CDR3-Sequenz aus AASTFLRILPEESAYTY (SEQ ID NO:3),

wobei der Antikörper für Insulinartiger-Wachstumsfaktor-1-Rezeptor (IGF1R) spezifisch ist und wobei der Antikörper die Blut-Hirn-Schranke transmigriert.

2. Isolierter oder gereinigter Antikörper gemäß Anspruch 1, umfassend die Sequenz
 X_1 VX₂ LX₃ E SGGGLVQX₄ GGSLRLSCX₅ X₆ SGGTVSPTAMGWX₇ RQAPGKX₈ X₉ EX₁₀ VX₁₁ HITWSRGTTTRX₁₂ ASSVKX₁₃ RFTISRDX₁₄ X₁₅ KNTX₁₆ YLQMNSLX₁₇ X₁₈ EDTAVYYCAASTFLRILPEESAYTYWGQGT X₁₉ VTVSS (SEQ ID NO:4), wobei X₁ E oder Q ist; X₂ K oder Q ist; X₃ V oder E ist; X₄ A oder P ist; X₅ A oder E ist; X₆ V oder A ist; X₇ V oder F ist; X₈ G oder E ist; X₉ L oder R ist; X₁₀ F oder W ist; X₁₁ G oder S ist; X₁₂ V oder Y ist; X₁₃ D oder G ist; X₁₄ N oder S ist; X₁₅ A oder S ist; X₁₆ L oder V ist; X₁₇ K oder R ist; X₁₈ A oder S ist und X₁₉ L oder Q ist.
3. Isolierter oder gereinigter Antikörper gemäß einem der Ansprüche 1 bis 2, umfassend eine Sequenz, die aus der Gruppe ausgewählt ist, welche besteht aus:

QVKLEESGGGLVQAGGSLRLSCEVSGGTVSPTAMGWFRQAPGKEREFGHITWSRGTTTRVASSVKD
 RFTISRDSAKNTVYLQMNSLKSEDTAVYYCAASTFLRILPEESAYTYWGQGTQVTVSS (SEQ ID
 NO:5);

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWVRQAPGKGLEWVGHITWSRGTTTRYASSVK
 GRFTISRDN SKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQGT LVTVSS (SEQ ID NO:
 6);

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKGLEFVGHITWSRGTTTRYASSVKG
 RFTISRDN SKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQGT LVTVSS (SEQ ID
 NO:7);

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKGLEFVGHITWSRGTTTRYASSVKG
 RFTISRDS SKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQGT LVTVSS (SEQ ID
 NO:8);

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKEREFGHITWSRGTTTRYASSVKG
 RFTISRDS SKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQGT LVTVSS (SEQ ID
 NO:9);

und/oder eine Sequenz, die damit zu 90% identisch ist.

4. Isolierter oder gereinigter Antikörper gemäß einem der Ansprüche 1 bis 3, wobei der Antikörper in einem multivalenten Displayformat ist.
5. Isolierter oder gereinigter Antikörper gemäß Anspruch 4, wobei der Antikörper mit einem Fc-Fragment verknüpft ist.
6. Nukleinsäuremolekül, das den isolierten oder gereinigten Antikörper nach einem der Ansprüche 1 bis 5 kodiert.
7. Vektor, der das Nukleinsäuremolekül nach Anspruch 6 umfasst.
8. Isolierter oder gereinigter Antikörper gemäß einem der Ansprüche 1 bis 5, wobei der Antikörper mit einem Frachtmolekül verknüpft ist.

9. Isolierter oder gereinigter Antikörper gemäß Anspruch 8, wobei das Frachtmolekül ein Molekulargewicht im Bereich von circa 1 kDa bis circa 200 kDa hat.

10. Isolierter oder gereinigter Antikörper gemäß Anspruch 9, wobei das Frachtmolekül aus der Gruppe ausgewählt ist, welche besteht aus einem detektierbaren Wirkstoff, einem Therapeutikum, einem Arzneistoff, einem Peptid, einem Wachstumsfaktor, einem Zytokin, einer Rezeptorfalle, einer chemischen Verbindung, einem Kohlenhydratanteil, einem oder mehreren Liposomen oder Nanoträgern, beladen mit einem detektierbaren Wirkstoff, einem Therapeutikum, einem Arzneistoff, einem Peptid, einem Enzym, einem Antikörper oder Fragment desselben, einem DNA-basierten Molekül, einem viralen Vektor, oder einem zytotoxischen Wirkstoff, oder einem oder mehreren Nanopartikeln, Nanodraht, Nanoröhrchen oder Quantenpunkten.

11. Zusammensetzung, die einen oder mehr als einen isolierten oder gereinigten Antikörper nach einem der Ansprüche 1 bis 5 und 9 bis 10 und einen pharmazeutisch akzeptablen Träger, Verdünnungsmittel oder Hilfsstoff umfasst.

12. Isolierter oder gereinigter Antikörper nach einem der Ansprüche 1 bis 5 und 9 bis 10 oder die Zusammensetzung gemäß Anspruch 11 zur Verwendung in der Medizin.

13. In-vitro-Verfahren zum Detektieren von Insulinartiger-Wachstumsfaktor-1-Rezeptor (IGF1R), umfassend

- a) Kontaktieren einer Gewebeprobe mit einem oder mehr als einem isolierten oder gereinigten Antikörper nach einem der Ansprüche 1 bis 5, der mit einem detektierbaren Wirkstoff verknüpft ist; und
- b) Detektieren des detektierbaren Wirkstoffs, der mit dem an IGF1R gebundenen Antikörper verknüpft ist, in der Gewebeprobe.

Revendications

1. Anticorps isolé ou purifié à domaine individuel variable d'immunoglobuline, comprenant une séquence de région déterminante de complémentarité (CDR) 1 de GGTVSPTA (N° ID SEQ : 1) ;
une séquence CDR2 de ITWSRGTT (N° ID SEQ : 2) ; et
une séquence CDR3 de AASTFLRILPEESAYTY (N° ID SEQ : 3),
dans lequel l'anticorps est spécifique pour le récepteur de facteur 1 de croissance analogue à l'insuline (IGF1R) et dans lequel l'anticorps transmigre la barrière hématoencéphalique.

2. Anticorps isolé ou purifié selon la revendication 1, comprenant la séquence
X₁VX₂LX₃ESGGGLVQX₄GGSLRLSCX₅X₆SGGTVSPTAMGWX₇RQAPGKX₈X₉EX₁₀VX₁₁HI
TWSRGTTX₁₂ASSVKX₁₃RFTISRDX₁₄X₁₅KNTX₁₆YLMNSLX₁₇X₁₈EDTAVWCAASTFLRIL
PEESAYTYWGQGTX₁₉VTVSS (ID. de SEQ. N° : 4), où X₁ est E ou Q ; X₂ est K ou Q ; X₃ est V ou E ; X₄ est A ou P ; X₅ est A ou E ; X₆ est V ou A ; X₇ est V ou F ; X₈ est G ou E ; X₉ est L ou R ; X₁₀ est F ou W ; X₁₁ est G ou S ; X₁₂ est V ou Y ; X₁₃ est D ou G ; X₁₄ est N ou S ; X₁₅ est A ou S ; X₁₆ est L ou V ; X₁₇ est K ou R ; X₁₈ est A ou S ; et X₁₉ est L ou Q.

3. Anticorps isolé ou purifié selon une quelconque des revendications 1 à 2, comprenant une séquence sélectionnée à partir du groupe se composant de :

QVKLEESGGGLVQAGGSLRLSCEVSGGTVSPTAMGWFRQAPGKEREFVGHITWSRGTT
RVASSVKDRFTISRDSAKNTVYLMNSLKS EDTAVYYCAASTFLRILPEESAYTYWGQGT
QVTVSS (ID. de SEQ. N° : 5) ;

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWVRQAPGKGLEWVGHITWSRGTT
TRYASSVKGRFTISRDN SKNTVYLMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQG
TLVTVSS (ID. de SEQ. N° : 6) ;

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKGLFVGHITWSRGT
TRYASSVKGRFTISRDNKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQG
TLVTVSS (ID. de SEQ. N° : 7) ;

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKGLFVGHITWSRGT
TRYASSVKGRFTISRDNSSKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQG
TLVTVSS (ID. de SEQ. N° : 8) ;

QVQLVESGGGLVQPGGSLRLSCAVSGGTVSPTAMGWFRQAPGKEREFVGHITWSRGT
TRYASSVKGRFTISRDNSSKNTVYLQMNSLRAEDTAVYYCAASTFLRILPEESAYTYWGQG
TLVTVSS (ID. de SEQ. N° : 9) ;

et/ou une séquence possédant une identité à 90 % de celui-ci.

4. Anticorps isolé ou purifié selon une quelconque des revendications 1 à 3, dans lequel l'anticorps est dans un format de manifestation polyvalente.
5. Anticorps isolé ou purifié selon la revendication 4, dans lequel l'anticorps est relié à un fragment Fc.
6. Molécule d'acide nucléique codant l'anticorps isolé ou purifié selon une quelconque des revendications 1 à 5.
7. Vecteur comprenant la molécule d'acide nucléique selon la revendication 6.
8. Anticorps isolé ou purifié selon une quelconque des revendications 1 à 5, dans lequel l'anticorps est relié à une molécule cargo.
9. Anticorps isolé ou purifié selon la revendication 8, dans lequel la molécule cargo possède un poids moléculaire dans la fourchette d'environ 1kD à environ 200 kDa.
10. Anticorps isolé ou purifié selon la revendication 9, dans lequel la molécule est sélectionnée à partir du groupe se composant d'un agent détectable, d'un thérapeutique, d'un médicament, d'un peptide, d'un facteur de croissance, d'une cytokine, d'un récepteur piège, d'un composé chimique, d'une moitié de carbohydrate, d'un ou plusieurs liposomes ou nano-porteurs chargés d'un agent détectable, d'un thérapeutique, d'un médicament, d'un peptide, d'une enzyme, d'un anticorps ou d'un fragment de celui-ci, d'une molécule à base d'ADN, d'un vecteur viral, ou d'un agent cytotoxique, ou d'une ou plusieurs nanoparticules, d'un nanofil, d'un nanotube, ou de points quantiques.
11. Composition comprenant un ou plus d'un anticorps isolé ou purifié selon une quelconque des revendications 1 à 5 et 9 à 10 et un support, un diluant ou un excipient pharmaceutiquement acceptables.
12. Anticorps isolé ou purifié selon une quelconque des revendications 1 à 5 et 9 à 10 ou composition selon la revendication 11 en vue de l'utilisation dans la médecine.
13. Méthode in vitro de détection du récepteur de facteur 1 de croissance analogue à l'insuline (IGF1R), comprenant
 - a) la mise en contact d'un échantillon de tissu avec un ou plus d'un anticorps isolé ou purifié selon une quelconque des revendications 1 à 5 relié à un agent détectable ; et
 - b) la détection de l'agent détectable relié à l'anticorps lié à IGF1R dans l'échantillon de tissu.




FIG. 1

MKSGSGGGSP	TSLWGLLFLS	AALSLWPTSG	EICGPGIDIR	NDYQQLKRLE	NCTVIEGYLH	60
ILLISKAEDY	RSYRFPKLTV	ITEYLLLFRV	AGLESLGDLF	PNLTVIRGWK	LFYNYALVIF	120
EMTNLKDIGL	YNLRNITRGA	IRIEKNADLC	YLSTVDWSLI	LDAVSNNYIV	GNKPPKECGD	180
LCPGTMEEKP	MCEKTTINNE	YNYRCWTTNR	CQKMCPSTCG	KRACTENNEC	CHPECLGSCS	240
APDNDTACVA	CRHYYYAGVC	VPACPPNTYR	FEGWRCVDRD	FCANILSAES	SDSEG FVIHD	300
GECMQECPSG	FIRNGSQSMY	CIPCEGPCPK	VCEEEKTKTK	IDSV TSAQML	QGCTIFKGNL	360
LINIRRGNNI	ASELENFMGL	IEVVTGYVKI	RHSHALVSL	FLKNLRLILG	EEQLEGNYSF	420
YVLDNQNLQQ	LWDWDHRNLT	IKAGKMYFAF	NPKLCVSEIY	RMEEVTGTKG	RQSKGDINTR	480
NNGERASCES	DVLHFTSTTT	SKNR IIITWH	RYP PPDYRDL	ISFTVYYKEA	PFKNVTEYDG	540
QDACGSNSWN	MVDVDLPPNK	DVEPGILLHG	LKPWTQYAVY	VKAVTLTMVE	NDHIRGAKSE	600
ILYIRTNASV	PSIPLDVLSA	SNSSSQLIVK	WNPPSLPNGN	LSYYIVRWQR	QPQDGYLYRH	660
NYCSKDKIPI	RKYADGTIDI	EEVTENPKTE	VCGGEKG PCC	ACP KTEAEKQ	AEKEEA EYRK	720
VFENFLHNSI	FVPRPERkrr	DVMQVANTTM	SSRSRNTTAA	DTYNITDPEE	LETEY PFFES	780
RVDNKERTVI	SNLRPFTLYR	IDIHSCNHEA	EKLGC SASNF	VFARTMPAEG	ADDIPGPVTW	840
EPRPENSIFL	KWPEPENPNG	LILMYEIKYG	SQVEDQRECV	SRQEYRK YGG	AKLNRLNPGN	900
YTARIQATSL	SGNGSWTDPV	FFYVQAKTGY	ENFIHLIIAL	PVAVLLIVGG	LVIMLYVFHR	960
KRNNSRLGNG	VLYASVNPEY	FSAADVYPD	EWEVAREKIT	MSRELQGGSF	GMVYEGVAKG	1020
VVKDEPETRV	AIKTVNEAAS	MRERIEFLNE	ASVMKEFNCH	HVVRL LGVVS	QGQPTLVIME	1080
LMTRGDLKSY	LRSLRP EMEN	NPVLAPPSLS	KMIQMAGEIA	DGMAYLNANK	FVHRDLAARN	1140
CMVAEDFTVK	IGDFGMTRDI	YETDY YRKGG	KGLLPVRWMS	PESLKDGVFT	TYSDVWSFGV	1200
VLWEIATLAE	QPYQGLSNEQ	VLR FVMEGGL	LDKPDNCPDM	LFELMRMCWQ	YNPKMRPSFL	1260
EIISSIKEEM	EPGFREV SFY	YSEENKLPEP	EELDLEPENM	ESVPLDPSAS	SSSLPLPDRH	1320
SGHKAENGP	PGVLVLRASF	DERQPYAHMN	GGRKNERALP	LPQSSTC		1367

FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C




FIG 3D




FIG 3E




FIG. 4

QVKLEESGGGLVQAGGSLRLSCEVSGGTVSPTAMGWFRQAPGKEREFVGHITWSRGTTTRVASS
 VKDRFTISRDSAKNTVYLQMNSLKSEDTAVYYCAASTFLRILPEESAYTYWGQGTQVTVSSMTVD
 KKLEPSGPISTINPCPPCKECHKCPAPNLEGGPSVFIFPPNIKDVLMLSLTPKVTCVVVDVSEDDPDV
 QISWVFVNNVEVHTAQQTQTHREDYNSTIRVVSTLPIQHQQDWMSGKEFKCKVNNKDLPSPIERTISK
 IKGLVRAPQVYILPPPAEQLSRKDVSLTCLVVGFNPGDISVEWTSNGHTEENYKDTAPVLDSGGSY
 FIYSKLNMKTSKWEKTDSEFSCNVRHEGLKNYYLKKTISRSPGK

FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D




FIG. 7




FIG. 8A




FIG. 8B




FIG. 9A




FIG. 9B




FIG 9C




FIG 9D




FIG 10.




FIG. 11




FIG. 12A




FIG. 12B




FIG. 12C

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 20050085419 A1 [0006]
- WO 2011044542 A1 [0006]
- WO 2011150061 A1 [0006]
- US 6180370 B [0046] [0147]
- US 5693761 A [0046] [0147]
- US 6054297 A [0046] [0147]
- US 5859205 A [0046] [0147]
- EP 626390 A [0046] [0147]
- US 5869619 A [0046] [0147]
- US 5766886 A [0046] [0147]
- US 5821123 A [0046] [0147]
- EP 519596 A [0046] [0147]
- WO 9504069 A, Cronan [0052] [0147]
- WO 2004076670 A, Voges [0052] [0147]
- WO 2003046560 A [0053] [0147]
- WO 02057445 A [0112] [0147]
- WO 2011127580 A [0127] [0147]
- WO 61948818 A [0148]

Non-patent literature cited in the description

- **BURTUM et al.** *Cancer Research*, 2003, vol. 63, 8812-8921 [0006]
- **GABATHULER R.** *Neurobiology of Diseases*, 2010, vol. 37, 48-57 [0007]
- **ABBOTT NJ.** Blood-brain barrier structure and function and the challenges for CNS drug delivery. *J Inherit Metab Dis*, 2013, vol. 36 (3), 437-49 [0147]
- **ABULROB A ; SPRONG H ; VAN BERGEN EN HENEGOUWEN P ; STANIMIROVIC D.** The blood-brain barrier transmigration single domain antibody: mechanisms of transport and antigenic epitopes in human brain endothelial cells. *J Neurochem.*, November 2005, vol. 95 (4), 1201-14 [0147]
- **ARBABI-GHAHROUDI ; M. DESMYTER A ; WYNS L ; HAMERS R. ; MUYLDERMANS S.** Selection and identification of single domain antibody fragments from camel heavy-chain antibodies. *FEBS Lett*, 1997, vol. 414, 521-526 [0147]
- **ARBABI-GHAHROUDI, M. ; TO, R. ; GAUDETTE, N. ; HIRAMA, T. ; DING, W. ; MACKENZIE, R. ; TANHA, J.** *Protein Eng. Des. Sel.*, 2009, vol. 22, 59-66 [0147]
- **ARBABI-GHAHROUDI, M. ; MACKENZIE, R. ; TANHA, J.** *Methods Mol. Biol.*, 2009, vol. 525, 187-216 [0147]
- **BELL, A. ; WANG, Z.J. ; ARBABI-GHAHROUDI, M. ; CHANG, T.A. ; DUROCHER, Y. ; TROJAHN, U. ; BAARDSNES, J. ; JARAMILLO, M.L. ; LI, S. ; BARAL, T.N.** *Cancer Lett.*, 2010, vol. 289, 81-90 [0147]
- **BROUSSAU, S. ; JABBOUR, N. ; LACHAPPELLE, G. ; DUROCHER, Y. ; TOM, R. ; TRANSGURATION, J. ; GILBERT, R. ; MASSIE, B.** *Mol Ther*, 2008, vol. 16, 500-507 [0147]
- **CHOTHIA, C. ; LESK, A.M.** *J. Mol. Biol.*, 1987, vol. 196, 901-917 [0147]
- **DAVIES J. ; L. RIECHMANN.** Affinity improvement of single antibody VH domains: residues in all three hypervariable regions affect antigen binding. *Immunotechnology*, 1996, vol. 2, 169-179 [0147]
- **DE KRUIF, J. ; LOGTENBERG, T.** *J. Biol. Chem.*, 1996, vol. 271, 7630-7634 [0147]
- **DEMEULE, M. ; CURRIE, J. C. ; BERTRAND, Y. ; CHE, C. ; NGUYEN, T. ; REGINA, A. ; GABATHULER, R. ; CASTAIGNE, J. P. ; BELIVEAU, R.** Involvement of the low-density lipoprotein receptor-related protein in the transcytosis of the brain delivery vector angiopep-2. *J. Neurochem.*, 2008, vol. 106, 1534-1544 [0147]
- **DUMOULIN, M. ; CONRATH, K. ; VAN MEIRHAIGHE, A. ; MEERSMAN, F. ; HEREMANS, K. ; FRENKEN, L.G. ; MUYLDERMANS, S. ; WYNS, L. ; MATAGNE, A.** *Protein Sci*, 2002, vol. 11, 500-515 [0147]
- **DUROCHER, Y. ; S. PERRET et al.** High-level and high-throughput recombinant protein production by transient transfection of suspension-growing human 293-EBNA1 cells. *Nucleic Acids Res*, 2002, vol. 30 (2), E9 [0147]
- **DOYLE, P.J. ; ARBABI-GHAHROUDI, M. ; GAUDETTE, N. ; FURZER, G. ; SAVARD, M.E. ; GLEDIE, S. ; MCLEAN, M.D. ; MACKENZIE, C.R. ; HALL, J.C.** *Mol. Immunol.*, 2008, vol. 45, 3703-3713 [0147]
- **EISENBERG, D. ; SCHWARZ, E. ; KOMAROMY, M. ; WALL, R.** *J. Mol. Biol.*, 1984, vol. 179, 125-142 [0147]

- **ERDLENBRUCH B ; ALIPOUR M ; FRICKER G ; MILLER DS ; KUGLER W ; EIBL H ; LAKOMEK M.** Alkylglycerol opening of the blood-brain barrier to small and large fluorescence markers in normal and C6 glioma-bearing rats and isolated rat brain capillaries. *Br J Pharmacol.*, 2003, vol. 140 (7), 1201-10 [0147]
- **FENNER, L. ; WIDMER, A.F. ; GOY, G. ; RUDIN, S. ; FREI, R.** *J. Clin. Microbiol.*, 2008, vol. 46, 328-330 [0147]
- **GAILLET, B. ; GILBERT, R. ; BROUSSAU, S. ; PILOTTE, A. ; MALENFANT, F. ; MULLICK, A. ; GARNIER, A. ; MASSIE, B.** *Biotechnol Bioeng*, 2010, vol. 106, 203-215 [0147]
- Gene delivery with viral vectors for cerebrovascular diseases. **GAN Y ; JING Z ; STETLER RA ; CAO G.** *Front Biosci.* 2013, vol. 5, 188-203 [0147]
- **GARBERG, P. ; BALL, M. ; BORG, N. ; CECHEL-LI, R. ; FENART, L. ; HURST, R. D. ; LINDMARK, T. ; MABONDZO, A. ; NILSSON, J. E. ; RAUB, T. J.** In vitro models for the blood-brain barrier, *Toxicol. In Vitro*, 2005, vol. 19, 299-334 [0147]
- **GERGOV, M. ; OJANPERA, I. ; VUORI, E.** Simultaneous screening for 238 drugs in blood by liquid chromatography-ion spray tandem mass spectrometry with multiple-reaction monitoring. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.*, 2003, vol. 795, 41-53 [0147]
- **GAILLET B ; GILBERT R ; AMZIANI R ; GUILBAULT C ; GADOURY C ; CARON AW ; MULLICK A ; GARNIERA ; MASSIE B.** High-level recombinant protein production in CHO cells using an adenoviral vector and the cumate gene-switch. *Biotechnol Prog.*, January 2007, vol. 23 (1), 200-9 [0147]
- **GOTTESMAN et al.** *Ann. Rev. Biochem.*, 1993, vol. 62, 385-427 [0147]
- **HAMERS-CASTERMAN, C. ; ATARHOUCHE, T. ; MUYLDERMANS, S. ; ROBINSON, G. ; HAMERS, C. ; SONGA, E.B. ; BENDAHMAN, N. ; HAMERS, R.** *Nature*, 1993, vol. 363, 446-448 [0147]
- **HAQQANIAS ; CARAM-SALAS N ; DING W ; BRUNETTE E ; DELANEY CE ; BAUMANN E ; BOILEAU E ; STANIMIROVIC D.** Multiplexed evaluation of serum and CSF pharmacokinetics of brain-targeting single-domain antibodies using a NanoLC-SRM-ILIS method. *Mol Pharm.*, 2012, vol. 10 (5), 1542-56 [0147]
- **HARGREAVES KM ; TROULLOS ES ; DIONNE RA ; SCHMIDT EA ; SCHAFER SC ; JORIS JL.** Bradykinin is increased during acute and chronic inflammation: therapeutic implications. *Clin Pharmacol Ther.*, 1988, vol. 44 (6), 613-21 [0147]
- **HUANG YL ; SÄLJÖ A ; SUNESON A ; HANSSON HA.** A new approach for multiple sampling of cisternal cerebrospinal fluid in rodents with minimal trauma and inflammation. *J Neurosci Methods*, 1995, vol. 63 (1-2), 13-22 [0147]
- **HUSSACK, G. ; HIRAMA, T. ; DING, W. ; MACKENZIE, R. ; TANHA, J.** *PLoS ONE*, 2011, vol. 6, e28218 [0147]
- **HUSSACK G ; ARBABI-GHAHROUDI M ; VAN FAASSEN H ; SONGER JG ; NG KK ; MACKENZIE R ; TANHA J.** Neutralization of Clostridium difficile toxin A with single-domain antibodies targeting the cell receptor binding domain. *J Biol Chem.*, 2011, vol. 286 (11), 8961-76 [0147]
- **IQBAL U ; ABULROB A ; STANIMIROVIC D B.** Integrated platform for brain imaging and drug delivery across the blood-brain barrier. *Methods Mol. Biol.*, 2011, vol. 686, 465-481 [0147]
- **IQBAL, U. ; TROJAHN, U. ; ALBAGHDADI, H. ; ZHANG, J. ; O'CONNOR, M. ; STANIMIROVIC, D. ; TOMANEK, B. ; SUTHERLAND, G. ; ABULROB, A.** *Br. J. Pharmacol.*, 2010, vol. 160, 1016-1028 [0147]
- **JESPEERS, L. ; SCHON, O. ; FAMM, K. ; WINTER, G.** *Nat. Biotechnol.*, 2004, vol. 22, 1161-1165 [0147]
- **KABAT EA ; WU TT.** Identical V region amino acid sequences and segments of sequences in antibodies of different specificities. Relative contributions of VH and VL genes, minigenes, and complementarity-determining regions to binding of antibody-combining sites. *J Immunol.*, 1991, vol. 147, 1709-19 [0147]
- **KIM, D.Y. ; KANDALAFT, H. ; DING, W. ; RYAN, S. ; VAN FASSEN, H. ; HIRAMA, T. ; FOOTE, S.J. ; MACKENZIE, R. ; TANHA, J.** *PEDS advance access*, 30 August 2012, 1-9 [0147]
- **KORNHUBER ME ; KORNHUBER J ; CIMNIAK U.** A method for repeated CSF sampling in the freely moving rat. *J Neurosci Methods*, 1986, vol. 17 (1), 63-8 [0147]
- **LEFRANC, M.-P. et al.** *Dev. Comp. Immunol.*, 2003, vol. 27, 55-77 [0147]
- **LI S ; ZHENG W ; KUOLEE R ; HIRAMA T ; HENRY M ; MAKVANDI-NEJAD S ; FJALLMAN T ; CHEN W ; ZHANG J.** Pentabody-mediated antigen delivery induces antigen-specific mucosal immune response. *Mol Immunol*, 2009, vol. 46, 1718-26 [0147]
- **MERRITT, E.A. ; HOL, W.G.** *Curr. Opin. Struct. Biol.*, 1995, vol. 5, 165-171 [0147]
- **MURUGANANDAM A ; TANHA J ; NARANG S ; STANIMIROVIC D.** Selection of phage-displayed llama single-domain antibodies that transmute across human blood-brain barrier endothelium. *FASEB J.*, 2001, vol. 16 (2), 240-2 [0147]
- **MUSHER, D.M. ; MANHAS, A. ; JAIN, P. ; NILA, F. ; WAQAR, A. ; LOGAN, N. ; MARINO, B. ; GRAVISS, E.A.** *J. Clin. Microbiol.*, 2007, vol. 45, 2737-2739 [0147]
- **NHAN T ; BURGESS A ; CHO EE ; STEFANOVIC B ; LILGE L ; HYNYNEN K.** Drug delivery to the brain by focused ultrasound induced blood-brain barrier disruption: Quantitative evaluation of enhanced permeability of cerebral vasculature using two-photon microscopy. *J Control Release*, 2013, vol. 172 (1), 274-280 [0147]

- **NICAISE M ; VALEIO-LEPINIEC M ; MINARD P ; DESMADRIL M.** Affinity transfer by CDR grafting on a nonimmunoglobulin scaffold. *Protein Sci.*, 2004, vol. 13 (7), 1882-1891 [0147]
- **NIELSEN, U.B. ; ADAMS, G.P. ; WEINER, L.M. ; MARKS, J.D.** *Cancer Res.*, 2000, vol. 60, 6434-6440 [0147]
- **NIROGI, R. ; KANDIKERE, V. ; MUDIGONDA, K. ; BHYRAPUNENI, G. ; MUDDANA, N. ; SARALAYA, R. ; BENADE, V.** A simple and rapid method to collect the cerebrospinal fluid of rats and its application for the assessment of drug penetration into the central nervous system. *J. Neurosci. Methods*, 2009, vol. 178, 116-119 [0147]
- **NUTTALL, S.D. ; KRISHNAN, U.V. ; DOUGHTY, L. ; PEARSON, K. ; RYAN, M.T. ; HOOGENRAAD, N.J. ; HATTARKI, M. ; CARMICHAEL, J.A. ; IRVING, R.A. ; HUDSON, P.J.** *Eur. J. Biochem.*, 2003, vol. 270, 3543-3554 [0147]
- **PARDRIDGE, W. M. ; BUCIAK, J. L. ; FRIDEN, P. M.** Selective transport of an anti-transferrin receptor antibody through the blood-brain barrier in vivo. *J. Pharmacol. Exp. Ther.*, 1991, vol. 259, 66-70 [0147]
- **PARDRIDGE, W.M.** *Adv. Drug Delivery Reviews*, 1995, vol. 15, 5-36 [0147]
- **PARDRIDGE, W. M.** Drug and gene delivery to the brain: the vascular route. *Neuron*, 2002, vol. 36, 555-558 [0147]
- **PLANCHE, T. ; AGHAIZU, A. ; HOLLIMAN, R. ; RILEY, P. ; POLONIECKI, J. ; BREATHNACH, A. ; KRISHNA, S.** *Lancet Infect. Dis.*, 2008, vol. 8, 777-784 [0147]
- **PRESTON E ; SLINN J ; VINOKOUROV I ; STANIMIROVIC D.** Graded reversible opening of the rat blood-brain barrier by intracarotid infusion of sodium caprate. *J Neurosci Methods.*, 2008, vol. 168 (2), 443-9 [0147]
- **RIDGWAY, J.B. ; PRESTA, L.G. ; CARTER, P.** *Protein Eng.*, 1996, vol. 9, 617-621 [0147]
- **ROBERTSON CR ; FLYNN SP ; WHITE HS ; BULAJ G.** Anticonvulsant neuropeptides as drug leads for neurological diseases. *Nat Prod Rep.*, 2011, vol. 28 (4), 741-62 [0147]
- **RUSSMANN, H. ; PANTHEL, K. ; BADER, R.C. ; SCHMITT, C. ; SCHAUMANN, R.** *Eur. J. Clin. Microbiol. Infect. Dis.*, 2007, vol. 26, 115-119 [0147]
- **SAMANI, A.A. ; CHEVET, E. ; FALLAVOLLITA, L. ; GALIPEAU, J. ; BRODT, P.** *Cancer Research*, 2004, vol. 64, 3380-3385 [0147]
- **SAMUELS B.L.** *J. Clin. Pharmacol. Ther.*, 1993, vol. 54, 421-429 [0147]
- **SLOAN, L.M. ; DURESKO, B.J. ; GUSTAFSON, D.R. ; ROSENBLATT, J.E.** *J. Clin. Microbiol.*, 2008, vol. 46, 1996-2001 [0147]
- **SUMBRIA RK ; ZHOU QH ; HUI EK ; LU JZ ; BOADO RJ ; PARDRIDGE WM.** Pharmacokinetics and brain uptake of an IgG-TNF decoy receptor fusion protein following intravenous, intraperitoneal, and subcutaneous administration in mice. *Mol Pharm.*, 2013, vol. 10 (4), 1425-31 [0147]
- **TANHA, J. ; MURUGANANDAM, A. ; STANIMIROVIC, D.** *Methods Mol. Med.*, 2003, vol. 89, 435-449 [0147]
- **TO, R. ; HIRAMA, T. ; ARBABI-GHAHROUDI, M. ; MACKENZIE, R. ; WANG, P. ; XU, P. ; NI, F. ; TANHA, J.** *J. Biol. Chem.*, 2005, vol. 280, 41395-41403 [0147]
- **TURGEON, D.K. ; NOVICKI, T.J. ; QUICK, J. ; CARLSON, L. ; MILLER, P. ; ULNESS, B. ; CENT, A. ; ASHLEY, R. ; LARSON, A. ; COYLE, M. ; LIMAYE, A.P.** *J. Clin. Microbiol.*, 2003, vol. 41, 667-670 [0147]
- **WATANABE, T.** *Acta Oncol.*, 1995, vol. 34, 235-241 [0147]
- **XIAO G ; GAN LS.** Receptor-mediated endocytosis and brain delivery of therapeutic biologics. *Int J Cell Biol.*, 2013 [0147]
- **YAKSH TL ; RUDY TA.** Chronic catheterization of the spinal subarachnoid space. *Physiol Behav.*, 1976, vol. 17 (6), 1031-6 [0147]
- **YU, Y. J. ; ZHANG, Y. ; KENRICK, M. ; HOYTE, K. ; LUK, W. ; LU, Y. ; ATWAL, J. ; ELLIOTT, J. M. ; PRABHU, S. ; WATTS, R. J.** Boosting brain uptake of a therapeutic antibody by reducing its affinity for a transcytosis target. *Sci. Transl. Med.*, 2011, vol. 3, 84ra44 [0147]
- **ZHANG, J. ; LI, Q. ; NGUYEN, T.-D. ; TREMBLAY, T.-L. ; STONE, E. ; TO, R. ; KELLY, J. ; MACKENZIE, C.R.** *J. Mol. Biol.*, 2004, vol. 341, 161-169 [0147]
- **ZHANG, J. ; TANHA, J. ; HIRAMA, T. ; KHIEW, N.H. ; TO, R. ; TONG-SEVINC, H. ; STONE, E. ; BRISSON, J.R. ; MACKENZIE, C.R.** *J. Mol. Biol.*, 2004, vol. 335, 49-56 [0147]
- **ZHU et al.** *Immunology and Cell Biology*, 2010, vol. 88, 667-675 [0147]

专利名称(译)	胰岛素样生长因子1受体特异性抗体及其用途		
公开(公告)号	EP3114141A1	公开(公告)日	2017-01-11
申请号	EP2014884789	申请日	2014-12-04
[标]申请(专利权)人(译)	加拿大国家研究委员会		
申请(专利权)人(译)	加拿大国家研究委员会的		
当前申请(专利权)人(译)	加拿大国家研究委员会的		
[标]发明人	STANIMIROVIC DANICA KEMMERICH KRISTIN HAQQANI ARSALAN S SULEA TRAIAN ARBABI GHAHROUDI MEHDI MASSIE BERNARD GILBERT RENALD		
发明人	STANIMIROVIC, DANICA KEMMERICH, KRISTIN HAQQANI, ARSALAN S. SULEA, TRAIAN ARBABI-GHAHROUDI, MEHDI MASSIE, BERNARD GILBERT, RÉNALD		
IPC分类号	C07K16/28 A61K39/395 A61K49/00 C07K16/46 C07K17/00 C12N15/13 G01N33/53		
CPC分类号	A61K47/6849 A61K49/0002 A61K49/0058 A61K2039/505 A61P25/00 A61P25/16 A61P25/28 C07K16/2863 C07K16/465 C07K2317/22 C07K2317/24 C07K2317/35 C07K2317/51 C07K2317/52 C07K2317/565 C07K2317/569 C07K2317/64 C07K2317/70 C07K2317/77 C07K2317/90 C07K2317/92 C07K2317/94 G01N33/57492 G01N33/68 G01N33/74 G01N2333/71 G01N2333/72 G01N2570/00 G01N2800/28		
优先权	61/948818 2014-03-06 US		
其他公开文献	EP3114141B1 EP3114141A4		
外部链接	Espacenet		

摘要(译)

血脑屏障 (BBB) 可以阻止大于500 Dal吨的分子从血液转运到大脑。受体介导的转胞吞作用 (RMT) 促进与形成BBB的脑内皮细胞上的受体结合的特定分子穿过BBB的转运。 识别出胰岛素样生长因子1受体 (IGF 1R) 结合抗体或其片段，其通过RMT迁移BBB。 抗体或片段用于跨BBB递送货物分子，其中该货物分子可以是治疗剂或可检测剂。 该抗体是骆驼VHH，通过用933个氨基酸的IGF 1R多肽免疫美洲驼而制备。 骆驼科动物VHH的人源化形式也被产生。