

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 May 2007 (18.05.2007)

PCT

(10) International Publication Number
WO 2007/056114 A2

(51) International Patent Classification:
G01N 33/569 (2006.01)

(74) Agent: **SLEATH, Janet**; 1201 Third Avenue, Suite 330,
Seattle, WA 98101 (US).

(21) International Application Number:
PCT/US2006/042907

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
2 November 2006 (02.11.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/733,119 3 November 2005 (03.11.2005) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): **INBIOS INTERNATIONAL, INC.** [US/US]; 562 1st Avenue South, Suite 600, Seattle, Washington 98104 (US).

(72) Inventors; and


(75) Inventors/Applicants (for US only): **HOUGHTON, Raymond L.** [US/US]; 562 1st Avenue South, Suite 600, Seattle, WA 98104 (US). **REED, Steven G.** [US/US]; 562 1st Avenue South, Suite 600, Seattle, WA 98104 (US). **RAY-CHAUDHURI, Syamal** [IN/US]; 562 1st Avenue South, Suite 600, Seattle, WA 98104 (US).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS AND METHODS FOR THE DETECTION OF *TRYPANOSOMA CRUZI* INFECTION



(57) Abstract: Compositions comprising fusion polypeptides of *T. cruzi* epitopes are provided, together with methods for the use of such compositions in the diagnosis of *T. cruzi* infection and in screening blood supplies. Diagnostic kits comprising such compositions are also provided.

WO 2007/056114 A2

**COMPOSITIONS AND METHODS FOR THE
DETECTION OF *TRYPANOSOMA CRUZI* INFECTION**

**5 STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR
DEVELOPMENT**

The U.S. Government has a paid-up license in this invention and the right in limited circumstances to require the patent owner to license others on reasonable terms as
10 provided for by the terms of Grant No. 2 R 44AIO52683-02 awarded by the National Institutes of Health.

FIELD OF THE INVENTION

The present invention relates generally to the diagnosis of *Trypanosoma cruzi* (*T.*
15 *cruzi*) infection. More specifically, the invention relates to the use of *T. cruzi* antigenic polypeptides and fusion polypeptides in methods for screening individuals and blood supplies for *T. cruzi* infection.

BACKGROUND OF THE INVENTION

20 Protozoan parasites are a serious health threat in many areas of the world. *Trypanosoma cruzi* (*T. cruzi*) is one such parasite that infects millions of individuals. Ten to thirty percent of individuals infected with *T. cruzi* develop chronic symptomatic Chagas' disease, which may in turn lead to heart disease and a variety of immune system disorders. *T. cruzi* infection has long been a public health problem in Central and South
25 America. It is estimated that 18 million people worldwide are chronically infected with *T. cruzi*, but available drug treatments lack efficacy and often cause serious side effects.

The most significant route of transmission in areas where the disease is endemic is through contact with an infected triatomid insect. However in other areas blood transfusions are the dominant means of transmission. Accordingly, in order to inhibit the
30 transmission of *T. cruzi*, it is necessary to develop accurate methods for both diagnosing *T. cruzi* infection in individuals and for screening blood supplies. Blood bank screening is particularly important in South America, where 0.1%-62% of blood samples may be infected and where the parasite is frequently transmitted by blood transfusion. Due to high flow of immigrants to the US from many Central and South American countries
35 where *T. cruzi* infection is endemic, the US blood supply is becoming at high risk for contamination from *T. cruzi* infected blood donors. While there are a few tests available

for diagnosing infection in individuals, there is currently no FDA approved test available in the US for blood donor screening for *T. cruzi* infection.

The diagnosis of *T. cruzi* infection has been problematic, since accurate methods for detecting the parasite that are suitable for routine use have been unavailable. During the acute phase of infection, which may last for decades, the infection may remain quiescent and the host may be asymptomatic. As a result, serological tests for *T. cruzi* infection are the most reliable and the most commonly used form of diagnosis. Such diagnoses are complicated, however, by the complex life cycle of the parasite and the diverse immune responses of the host. The parasite passes through an epimastigote stage in the insect vector and two main stages in the mammalian host. One host stage is present in blood (the trypomastigote stage), while a second stage is intracellular (the amastigote stage). The multiple stages result in a diversity of antigens being presented by the parasite during infection. In addition, immune responses to protozoan infection are complex, involving both humoral and cell-mediated responses to the array of parasite antigens.

While detection of antibodies against parasite antigens is the most common and reliable method of diagnosing clinical and subclinical infections, current tests for *T. cruzi* infection are generally insensitive, lack specificity, and are not suitable for screening of blood supplies. Most serological tests use whole or lysed *T. cruzi* and require positive results on two of three tests, including complement fixation, indirect immunofluorescence, passive agglutination or ELISA, to accurately detect *T. cruzi* infection. The cost and difficulty of such tests has prevented the screening of blood or sera in many endemic areas.

US Patents 5,876,734 and 6,228,601 disclose compositions useful for diagnosing Chagas' disease that comprise a non-repetitive region of the *T. cruzi* protein TCR27, and fusion polypeptides including such regions. US Patent 6,419,933 discloses a fusion polypeptide referred to as TcF that contains the four antigenic *T. cruzi* peptides PEP-2, TcD, TcE and TcLo1.2, together with methods for the use of the fusion polypeptide in the detection of *T. cruzi* infection. While TcF is highly reactive with *T. cruzi*-infected sera from South America, it exhibits low activity, and is occasionally negative with, Central American sera. US Patent 6,458,922 discloses an assay for *T. cruzi* infection that employs compositions comprising at least six antigenic *T. cruzi* peptides selected from the group consisting of: SAPA, CRA, FRA, TcD, Tc24, Ag39 and MAP. Published US Patent Application No. US-2004/0132077-A1 discloses recombinant polypeptides and

fusion polypeptides (referred to as FP3, FP4, FP5, FP6, FP7, FP8, FP9 and FP10) useful for diagnosing *T. cruzi* infection. The disclosed fusion polypeptides comprise modified versions of previously identified *T. cruzi* epitopes, including TCR27, TCR39, SAPA and MAP.

5

SUMMARY OF THE INVENTION

The present invention provides compositions and methods for detecting *T. cruzi* infection in individuals and in biological samples, including blood supplies. The inventive compositions may be employed to detect *T. cruzi* infection in all geographical areas where Chagas' disease is present and with improved sensitivity compared to assays currently in use.

In one embodiment, the inventive compositions comprise the *T. cruzi* fusion polypeptide TcF (SEQ ID NO: 1), or a variant thereof, and at least one peptide selected from the group consisting of: SAPA (SEQ ID NO: 2); Pep30 (SEQ ID NO: 3); Pep36 (SEQ ID NO: 4); KMP-11 (SEQ ID NO: 5); Peptide 1 (SEQ ID NO: 6; also referred to as FRA; Lafaille et al., *Mol. Biochem. Parasitol.* 35:127-136, 1989); a modified version of peptide 1 (SEQ ID NO: 7); and variants thereof. The fusion polypeptide TcF and the at least one peptide may be present as individual components within the composition or may be linked to form a fusion polypeptide. In certain embodiments, the inventive compositions comprise the fusion polypeptide TcF in combination with the peptides Pep30, Pep36 and SAPA. Such compositions may, for example, include the fusion polypeptide of SEQ ID NO: 8 (referred to herein as ITC-6). In alternative embodiments, the inventive compositions include a fusion polypeptide comprising ITC-6 in combination with at least one repeat of the peptide KMP-11, and/or the peptide 1 sequences of SEQ ID NO: 6 or 7. The amino acid sequence of a representative fusion polypeptide comprising ITC-6 plus one repeat of KMP-11 (referred to as ITC7.1) is provided in SEQ ID NO: 15, with the amino acid sequence of a representative fusion polypeptide comprising ITC-6 plus two repeats of KMP-11 (referred to as ITC7.2) being provided in SEQ ID NO: 17. SEQ ID NO: 19 is the amino acid sequence of a representative fusion polypeptide comprising ITC7.2 plus peptide 1 (referred to as ITC8.2), with the corresponding DNA sequence being provided in SEQ ID NO: 18. The amino acid sequence of a shortened version of ITC8.2, referred to as ITC8.1, is provided in SEQ ID NO: 20. SEQ ID NO: 21 and 22 are the DNA and amino acid sequences, respectively, for a shortened version of ITC8.1.

Polynucleotides encoding the inventive fusion polypeptides, expression vectors comprising such polynucleotides, and host cells transformed or transfected with such expression vectors are also provided by the present invention.

As described in detail below, the inventors have determined that the inventive compositions may be employed to effectively detect *T. cruzi* infection in a biological sample. Accordingly, in one aspect, the present invention provides methods for detecting *T. cruzi* infection in a biological sample, comprising: (a) contacting the biological sample with a composition of the present invention; and (b) detecting in the biological sample the presence of antibodies that bind to an epitope present within the inventive composition, thereby detecting *T. cruzi* infection in the biological sample.

In a further aspect, diagnostic kits for detecting *T. cruzi* infection in a biological sample are provided, such kits comprising: (a) a composition of the present invention; and (b) a detection reagent.

The inventive compositions may also comprise at least one component selected from the group consisting of: physiologically acceptable carriers and immunostimulants. Methods for inducing protective immunity against Chagas' disease in a patient by administering such compositions are also encompassed by the present invention.

The above-mentioned and additional features of the present invention and the manner of obtaining them will become apparent, and the invention will be best understood by reference to the following more detailed description. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows the reactivity of the fusion polypeptides TcF and ITC-6 with *T. cruzi*-infected sera and control sera from non-infected individuals, as determined by ELISA.

Figs. 2A-C show the reactivity of TcF and ITC-6 with various sera as determined by ELISA. The RIPA status of the individual sera are shown in Table 1.

Fig. 3 shows the reactivity of the fusion polypeptides TcF, ITC-6, ITC7.1 and ITC7.2 on a panel of sera as determined by ELISA.

Fig. 4 shows the reactivity of the fusion polypeptides ITC7.2 and the shortened version of ITC8.2 provided in SEQ ID NO: 22 on a panel of sera as determined by ELISA.

Fig. 5 shows the activity of non-absorbed and absorbed peptide 1-positive sera with ITC7.2, ITC8.1 and shortened ITC8.2 (SEQ ID NO: 22).

Figs. 6A and 6B show the isolated DNA sequence and corresponding amino acid sequence, respectively, for TC5. The insert is shown in bold font, and the flanking sequence in non-bold font.

Figs. 7A and 7B show the isolated DNA sequence and corresponding amino acid sequence, respectively, for TC48. The insert is shown in bold font, and the flanking sequence in non-bold font.

Figs. 8A and 8B show the isolated DNA sequence and corresponding amino acid sequence, respectively, for TC60. The insert is shown in bold font, and the flanking sequence in non-bold font.

Figs. 9A and 9B show the isolated DNA sequence and corresponding amino acid sequence, respectively, for TC70. The insert is shown in bold font, and the flanking sequence in non-bold font.

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for detecting *T. cruzi* infection in individuals and for screening blood supplies for *T. cruzi* infection. The compositions of this invention generally comprise the known *T. cruzi* fusion polypeptide TcF (US Patent 6,419,933; SEQ ID NO: 1), or a variant thereof, and at least one antigenic *T. cruzi* epitope or polypeptide selected from the group consisting of: SAPA (SEQ ID NO: 2); Pep30 (SEQ ID NO: 3); Pep36 (SEQ ID NO: 4); KMP-11 (SEQ ID NO: 5); peptide 1 (SEQ ID NO: 6); modified peptide 1 (SEQ ID NO: 7); and variants thereof. The use of one or more additional epitopes from *T. cruzi* polypeptides, either prior to or in combination with one or more of the specific *T. cruzi* polypeptides disclosed herein, in order to enhance the sensitivity and specificity of detection is also contemplated by and encompassed within the present invention. The fusion polypeptide TcF and the at least one peptide may be present as individual components within the inventive composition or may be linked to form a fusion polypeptide. Fusion polypeptides comprising more than one repeat of the *T. cruzi* antigenic epitopes or polypeptides are also contemplated and encompassed by the present invention, as are fusion polypeptides in which the peptides are linked in an order which differs from those shown in the specific fusion polypeptide sequences provided herein.

In one embodiment, the inventive compositions include a fusion polypeptide comprising TcF, Pep30, Pep36 and SAPA, herein referred to as ITC-6 (SEQ ID NO: 8). In alternative embodiments, the inventive compositions include a fusion protein selected from the group consisting of SEQ ID NO: 15 (referred to as ITC7.1), SEQ ID NO: 17 (referred to as ITC7.2), SEQ ID NO: 19 (referred to as ITC8.2), and SEQ ID NO: 20 (referred to as ITC8.1). The DNA sequences for ITC-6, ITC7.1, ITC7.2 and ITC8.2 are provided in SEQ ID NO: 11, 14, 16 and 18, respectively.

As described in US Patent 6,419,933, the disclosure of which is hereby incorporated by reference, the fusion polypeptide TcF includes four antigenic epitopes, or peptides, referred to as PEP2, TcD, TcE and TcLo1.2. In an alternative embodiment, the inventive compositions may include these four individual peptides in place of the fusion protein TcF.

As used herein, the term "polypeptide" encompasses amino acid chains of any length, including full-length proteins, wherein amino acid residues are linked by covalent peptide bonds. Polypeptides disclosed herein may be naturally purified products, or may be produced partially or wholly using recombinant techniques. Such polypeptides may be glycosylated with mammalian or other eukaryotic carbohydrates or may be non-glycosylated. A polypeptide comprising an epitope may consist entirely of the epitope or may contain additional sequences. The additional sequences may be derived from the native antigen or may be heterologous, and such sequences may (but need not) be antigenic.

As used herein, a "fusion polypeptide" is a polypeptide in which epitopes of different antigens, or variants thereof, are joined, for example through a peptide linkage, into a single amino acid chain. The amino acid chain thus formed may be either linear or branched. The epitopes may be joined directly (*i.e.*, with no intervening amino acids) or may be joined by way of a linker sequence that does not significantly alter the antigenic properties of the epitopes. The peptide epitopes may also be linked through non-peptide linkages, such as hetero- or homo-bifunctional agents that chemically or photochemically couple between specific functional groups on the peptide epitopes such as through amino, carboxyl, or sulfhydryl groups. Bifunctional agents which may be usefully employed in the combination polypeptides of the present invention are well known to those of skill in the art. Epitopes may also be linked by means of a complementary ligand/anti-ligand pair, such as avidin/biotin, with one or more epitopes being linked to a first member of the ligand/anti-ligand pair and then being bound to the complementary member of the

ligand/anti-ligand pair either in solution or in solid phase. A fusion polypeptide may contain epitopes of one or more other *T. cruzi* antigens, linked to an epitope described herein.

A polynucleotide encoding a fusion protein of the present invention is constructed using known recombinant DNA techniques to assemble separate polynucleotides encoding the first and second polypeptides into an appropriate expression vector. The 3' end of a polynucleotide encoding a first polypeptide is ligated, with or without a peptide linker, to the 5' end of a polynucleotide encoding a second polypeptide so that the reading frames of the sequences are in phase to permit mRNA translation of the two polynucleotides into a single fusion protein that retains the biological activity of both the first and the second polypeptides.

As noted above, a peptide linker sequence may be employed to separate the first and the second polypeptides by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea *et al.*, *Gene* 40:39-46, 1985; Murphy *et al.*, *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may be from 1 to about 50 amino acids in length. Peptide linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference. The ligated polynucleotides encoding the fusion proteins are cloned into suitable expression systems using techniques known to those of ordinary skill in the art.

The present invention further provides polynucleotides that encode a polypeptide or fusion polypeptide of the present invention. Polynucleotides that comprise complements of such polynucleotide sequences, reverse complements of such

polynucleotide sequences, or reverse sequences of such polynucleotide sequences, together with variants of such sequences, are also provided.

The definition of the terms “complement(s),” “reverse complement(s),” and “reverse sequence(s),” as used herein, is best illustrated by the following example. For the sequence 5' AGGACC 3', the complement, reverse complement, and reverse sequence are as follows:

complement	3' TCCTGG 5'
reverse complement	3' GGTCCT 5'
reverse sequence	5' CCAGGA 3'.

Preferably, sequences that are complements of a specifically recited polynucleotide sequence are complementary over the entire length of the specific polynucleotide sequence.

The term “polynucleotide(s),” as used herein, means a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases and includes DNA and corresponding RNA molecules, including HnRNA and mRNA molecules, both sense and anti-sense strands, and comprehends cDNA, genomic DNA and recombinant DNA, as well as wholly or partially synthesized polynucleotides. An HnRNA molecule contains introns and corresponds to a DNA molecule in a generally one-to-one manner. An mRNA molecule corresponds to an HnRNA and DNA molecule from which the introns have been excised. A polynucleotide may consist of an entire gene, or any portion thereof. Operable anti-sense polynucleotides may comprise a fragment of the corresponding polynucleotide, and the definition of “polynucleotide” therefore includes all such operable anti-sense fragments.

All of the polypeptides, fusion polypeptides and polynucleotides described herein are isolated and purified, as those terms are commonly used in the art. Preferably, the polypeptides, fusion polypeptides and polynucleotides are at least about 80% pure, more preferably at least about 90% pure, and most preferably at least about 99% pure.

The compositions and methods of the present invention also encompass variants of the above polypeptides, fusion polypeptides and polynucleotides.

As used herein, the term “variant” comprehends nucleotide or amino acid sequences different from the specifically identified sequences, wherein one or more nucleotides or amino acid residues is deleted, substituted, or added. Variants may be naturally occurring allelic variants, or non-naturally occurring variants. Variant sequences (polynucleotide or polypeptide) preferably exhibit at least 75%, more

preferably at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably yet at least 95%, and most preferably, at least 98% identity to a sequence of the present invention. The percentage identity is determined by aligning the two sequences to be compared as described below, determining the number of identical residues in the aligned portion, dividing that number by the total number of residues in the inventive (queried) sequence, and multiplying the result by 100. In addition to exhibiting the recited level of sequence identity, variant sequences of the present invention preferably exhibit a functionality that is substantially similar to the functionality of the specific sequences disclosed herein. Variant fusion polypeptide sequences thus preferably retain the antigenic and diagnostic properties of the fusion polypeptides disclosed herein. Preferably a variant polypeptide or fusion polypeptide sequence will generate at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably yet at least 95% and most preferably 100% of the response generated by the specifically identified polypeptide or fusion polypeptide sequence in an antibody binding assay, such as an ELISA assay. Such variants may generally be identified by modifying one of the polypeptide or fusion polypeptide sequences disclosed herein, and evaluating the antigenic and/or diagnostic properties of the modified polypeptide or fusion polypeptide using, for example, the representative procedures described herein. Suitable assays for evaluating reactivity with *T. cruzi*-infected sera, such as an enzyme linked immunosorbent assay (ELISA), are described in more detail below, and in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988.

Variant sequences generally differ from the specifically identified sequence only by conservative substitutions, deletions or modifications. As used herein, a "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the polypeptide to be substantially unchanged. In general, the following groups of amino acids represent conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. Variants may also, or alternatively, contain other modifications, including the deletion or addition of amino acids that have minimal influence on the antigenic properties, secondary structure and hydropathic nature of the polypeptide. For example, a polypeptide may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be

conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (*e.g.*, poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

5 Polypeptide and polynucleotide sequences may be aligned, and percentages of identical nucleotides in a specified region may be determined against another polynucleotide, using computer algorithms that are publicly available. Two exemplary algorithms for aligning and identifying the identity of polynucleotide sequences are the BLASTN and FASTA algorithms. The alignment and identity of polypeptide sequences
10 may be examined using the BLASTP and algorithm. BLASTX and FASTX algorithms compare nucleotide query sequences translated in all reading frames against polypeptide sequences. The FASTA and FASTX algorithms are described in Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444-2448, 1988; and in Pearson, *Methods in Enzymol.* 183:63-98, 1990. The FASTA software package is available from the University of
15 Virginia, Charlottesville, VA 22906-9025. The FASTA algorithm, set to the default parameters described in the documentation and distributed with the algorithm, may be used in the determination of polynucleotide variants. The readme files for FASTA and FASTX Version 2.0x that are distributed with the algorithms describe the use of the algorithms and describe the default parameters.

20 The BLASTN software is available on the NCBI anonymous FTP server and is available from the National Center for Biotechnology Information (NCBI), National Library of Medicine, Building 38A, Room 8N805, Bethesda, MD 20894. The BLASTN algorithm Version 2.0.6 [Sep-10-1998] and Version 2.0.11 [Jan-20-2000] set to the default parameters described in the documentation and distributed with the algorithm, is
25 preferred for use in the determination of variants according to the present invention. The use of the BLAST family of algorithms, including BLASTN, is described at NCBI's website and in the publication of Altschul, *et al.*, "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs," *Nucleic Acids Res.* 25:3389-3402, 1997.

30 The "hits" to one or more database sequences by a queried sequence produced by BLASTN, BLASTP, FASTA, or a similar algorithm, align and identify similar portions of sequences. The hits are arranged in order of the degree of similarity and the length of sequence overlap. Hits to a database sequence generally represent an overlap over only a fraction of the sequence length of the queried sequence.

The percentage identity of a polynucleotide or polypeptide sequence is determined by aligning polynucleotide and polypeptide sequences using appropriate algorithms, such as BLASTN or BLASTP, respectively, set to default parameters; identifying the number of identical nucleic or amino acids over the aligned portions; dividing the number of identical nucleic or amino acids by the total number of nucleic or amino acids of the polynucleotide or polypeptide of the present invention; and then multiplying by 100 to determine the percentage identity.

In general, *T. cruzi* polypeptides and fusion polypeptides, and polynucleotide sequences encoding such polypeptides and fusion polypeptides, may be prepared using any of a variety of procedures. For example, a *T. cruzi* cDNA or genomic DNA expression library may be screened with pools of sera from *T. cruzi*-infected individuals. Such screens may generally be performed using techniques well known to those of ordinary skill in the art, such as those described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, N.Y., 1989. Briefly, the bacteriophage library may be plated and transferred to filters. The filters may then be incubated with serum and a detection reagent. In the context of this invention, a "detection reagent" is any compound capable of binding to the antibody-antigen complex, which may then be detected by any of a variety of means known to those of ordinary skill in the art. Typical detection reagents for screening purposes contain a "binding agent," such as Protein A, Protein G, IgG or a lectin, coupled to a reporter group. Preferred reporter groups include, but are not limited to, enzymes, substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. More preferably, the reporter group is horseradish peroxidase (HRP), which may be detected by incubation with a substrate such as tetramethylbenzidine (TMB) or 2,2'-azino-di-3-ethylbenzthiazoline sulfonic acid. Plaques containing cDNAs that express a protein that binds to an antibody in the serum may be isolated and purified by techniques known to those of ordinary skill in the art. Appropriate methods may be found, for example, in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, N.Y., 1989.

Alternatively, polynucleotides encoding the fusion polypeptides disclosed herein may be amplified from *T. cruzi* genomic DNA or cDNA via polymerase chain reaction (PCR). For this approach, sequence-specific primers may be designed based on the polynucleotide sequence, and may be purchased or synthesized. An amplified portion of the DNA sequences may then be used to isolate the full length genomic or cDNA clones

using well known techniques, such as those described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY (1989).

Polypeptides having fewer than about 100 amino acids, and generally fewer than about 50 amino acids, can be synthesized using, for example, the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. See Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963. Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied Biosystems Division, Foster City, CA.

Polypeptides and fusion polypeptides may also be produced recombinantly by inserting a polynucleotide that encodes the fusion polypeptide into an expression vector and expressing the antigen in an appropriate host. Any of a variety of expression vectors known to those of ordinary skill in the art may be employed. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, mycobacteria, insect, yeast or a mammalian cell line such as COS or CHO. The polynucleotides expressed in this manner may encode naturally occurring antigens, portions of naturally occurring antigens, or other variants thereof.

Expressed polypeptides and fusion polypeptides are generally isolated in substantially pure form. Preferably, the polypeptides and fusion polypeptides are isolated to a purity of at least 80% by weight, more preferably, to a purity of at least 95% by weight, and most preferably to a purity of at least 99% by weight. In general, such purification may be achieved using, for example, the standard techniques of ammonium sulfate fractionation, SDS-PAGE electrophoresis, and affinity chromatography.

The present invention further provides methods for detecting *T. cruzi* infection in individuals and blood supplies. *T. cruzi* infection may be detected in any biological sample that contains antibodies. Preferably, the sample is blood, serum, plasma, saliva, cerebrospinal fluid or urine. More preferably, the sample is a blood or serum sample obtained from a patient or a blood supply. Briefly, *T. cruzi* infection may be detected using any one or more of the polypeptides or fusion polypeptides described above, or variants thereof, to determine the presence or absence of antibodies to the polypeptide or fusion polypeptide in the sample, relative to a predetermined cut-off value.

There are a variety of assay formats known to those of ordinary skill in the art for using purified antigen to detect antibodies in a sample. *See, e.g., Harlow and Lane, Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In a preferred embodiment, the assay involves the use of polypeptide or fusion polypeptide immobilized on a solid support to bind to and remove the antibody from the sample. The bound antibody may then be detected using a detection reagent that binds to the antibody/ fusion polypeptide complex and contains a detectable reporter group. Suitable detection reagents include antibodies that bind to the antibody/fusion polypeptide complex and free polypeptide labeled with a reporter group (*e.g., in a semi-competitive assay*). Alternatively, a competitive assay may be utilized, in which an antibody that binds to the fusion polypeptide is labeled with a reporter group and allowed to bind to the immobilized antigen after incubation of the antigen with the sample. The extent to which components of the sample inhibit the binding of the labeled antibody to the fusion polypeptide is indicative of the reactivity of the sample with the immobilized fusion polypeptide.

The solid support may be any solid material known to those of ordinary skill in the art to which the fusion polypeptide may be attached. For example, the solid support may be a test well in a microtiter plate, or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, formed of glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681.

The fusion polypeptide may be bound to the solid support using a variety of techniques known to those in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "bound" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the antigen and functional groups on the support or may be a linkage by way of a cross-linking agent). Binding by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the polypeptide, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of fusion polypeptide ranging from

about 10 ng to about 1 µg, and preferably about 100 ng, is sufficient to bind an adequate amount of antigen. Nitrocellulose will bind approximately 100 µg of protein per cm³.

Covalent attachment of the polypeptide or fusion polypeptide to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the fusion polypeptide. For example, the fusion polypeptide may be bound to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the fusion polypeptide (*see, e.g.*, Pierce Immunotechnology Catalog and Handbook (1991) at A12-A13).

In certain embodiments, the assay is an enzyme linked immunosorbent assay (ELISA). This assay may be performed by first contacting a polypeptide or fusion polypeptide that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that antibodies to the polypeptide or fusion polypeptide within the sample are allowed to bind to the immobilized polypeptide or fusion polypeptide. Unbound sample is then removed from the immobilized polypeptide or fusion polypeptide and a detection reagent capable of binding to the immobilized antibody-polypeptide complex is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the specific detection reagent.

Once the polypeptide or fusion polypeptide is immobilized on the support, the remaining protein binding sites on the support are typically blocked using any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20TM (Sigma Chemical Co., St. Louis, MO). The immobilized polypeptide or fusion polypeptide is then incubated with the sample, and antibody (if present in the sample) is allowed to bind to the antigen. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is that period of time that is sufficient to detect the presence of *T. cruzi* antibody within a *T. cruzi*-infected sample. Preferably, the contact time is sufficient to achieve a level of binding that is at least 95% of that achieved at equilibrium between bound and unbound antibody. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined

by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20TM. Detection reagent may then be added to the solid support. An appropriate detection reagent is any compound that binds to the immobilized antibody-polypeptide complex and that can be detected by any of a variety of means known to those in the art. Preferably, the detection reagent contains a binding agent (such as, for example, Protein A, Protein G, immunoglobulin, lectin or free antigen) conjugated to a reporter group. Preferred reporter groups include enzymes (such as horseradish peroxidase), substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. The conjugation of binding agent to reporter group may be achieved using standard methods known to those of ordinary skill in the art. Common binding agents may also be purchased conjugated to a variety of reporter groups from many sources (*e.g.*, Zymed Laboratories, San Francisco, CA and Pierce, Rockford, IL).

The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound antibody. An appropriate amount of time may generally be determined from the manufacturer's instructions or by assaying the level of binding that occurs over a period of time. Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of *T. cruzi* antibodies in the sample, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. This cut-off value is preferably the average mean signal obtained when the immobilized antigen is incubated with samples from an uninfected patient. In general, a sample generating a signal that is three standard deviations above the mean is considered positive

for *T. cruzi* antibodies and *T. cruzi* infection. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, p. 106-7 (Little Brown and Co., 1985). Briefly, in this embodiment, the cut-off value may be
5 determined from a plot of pairs of true positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method
10 may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for *T. cruzi* infection.

In a related embodiment, the assay is performed in a flow-through or strip test
15 format, wherein the polypeptide or fusion polypeptide is immobilized on a membrane such as nitrocellulose. In the flow-through test, antibodies within the sample bind to the immobilized polypeptide or fusion polypeptide as the sample passes through the membrane. A detection reagent (*e.g.*, protein A-colloidal gold) then binds to the antibody-polypeptide complex as the solution containing the detection reagent flows
20 through the membrane. The detection of bound detection reagent may then be performed as described above. In the strip test format, one end of the membrane to which polypeptide or fusion polypeptide is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing detection reagent and to the area of immobilized polypeptide or fusion polypeptide. Concentration
25 of detection reagent at the polypeptide or fusion polypeptide indicates the presence of *T. cruzi* antibodies in the sample. Such tests can typically be performed with a very small amount (*e.g.*, one drop) of patient serum or blood.

In yet another aspect of this invention, methods are provided for detecting *T. cruzi* in a biological sample using monospecific antibodies (which may be polyclonal or
30 monoclonal) to one or more *T. cruzi* polypeptides or fusion polypeptides. Antibodies to purified or synthesized polypeptides may be prepared by any of a variety of techniques known to those of ordinary skill in the art. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In one such technique, an immunogen comprising the antigenic polypeptide is initially injected into any of a wide

variety of mammals (e.g., mice, rats, rabbits, sheep and goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for the antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (i.e., reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction.

Monospecific antibodies to epitopes of one or more of the polypeptides or fusion polypeptides described herein may be used to detect *T. cruzi* infection in a biological

sample using any of a variety of immunoassays, which may be direct or competitive. Suitable biological samples for use in this aspect of the present invention are as described above. Briefly, in one direct assay format, a monospecific antibody may be immobilized on a solid support (as described above) and contacted with the sample to be tested. After
5 removal of the unbound sample, a second monospecific antibody, which has been labeled with a reporter group, may be added and used to detect bound antigen. In an exemplary competitive assay, the sample may be combined with the monoclonal or polyclonal antibody, which has been labeled with a suitable reporter group. The mixture of sample and antibody may then be combined with polypeptide antigen immobilized on a suitable
10 solid support. Antibody that has not bound to an antigen in the sample is allowed to bind to the immobilized antigen, and the remainder of the sample and antibody is removed. The level of antibody bound to the solid support is inversely related to the level of antigen in the sample. Thus, a lower level of antibody bound to the solid support indicates the presence of *T. cruzi* in the sample. To determine the presence or absence of *T. cruzi*
15 infection, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. Such cut-off values may generally be determined as described above. Any of the reporter groups discussed above in the context of ELISAs may be used to label the monospecific antibodies, and binding may be detected by any of a variety of techniques
20 appropriate for the reporter group employed. Other formats for using monospecific antibodies to detect *T. cruzi* in a sample will be apparent to those of ordinary skill in the art, and the above formats is provided solely for exemplary purposes.

In another aspect of this invention, compositions are provided for the prevention or treatment of *T. cruzi* infection, and complications thereof, in a mammal. Such
25 compositions generally comprise one or more fusion polypeptides disclosed herein, together with at least one component selected from the group consisting of: physiologically acceptable carriers and immunostimulants.

Routes and frequency of administration and fusion polypeptide doses will vary from individual to individual and may parallel those currently being used in immunization
30 against other protozoan infections. In general, the compositions may be administered by injection (e.g., intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (e.g., by aspiration), transdermally, orally or by transcutaneous patch as described, for example, in US Patents 5,910,306 and 5,980,898, the disclosures of which are hereby incorporated by reference. Between 1 and 4 doses may be administered for a 2-6 week

period. Preferably, two doses are administered, with the second dose 2-4 weeks later than the first. A suitable dose is an amount of fusion polypeptide that is effective to raise antibodies in a treated mammal that are sufficient to protect the mammal from *T. cruzi* infection for a period of time. In general, the amount of fusion polypeptide present in a dose ranges from about 1 pg to about 100 mg per kg of host, typically from about 10 pg to about 1 mg, and preferably from about 100 pg to about 1 µg. Suitable dose sizes will vary with the size of the animal, but will typically range from about 0.01 mL to about 5 mL for 10-60 kg animal.

While any suitable carrier known to those of ordinary skill in the art may be employed in the compositions of this invention, the type of carrier will vary depending on the mode of administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (e.g., polylactic galactide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

Any of a variety of immunostimulants may be employed in the compositions of this invention to nonspecifically enhance the immune response. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a nonspecific stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis*. Suitable adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Freund's Complete Adjuvant (Difco Laboratories) and Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ). Other suitable adjuvants include alum, biodegradable microspheres, monophosphoryl lipid A and quil A.

The following Examples are offered by way of illustration and not by way of limitation.

Example 1

DESIGN AND PREPARATION OF A MULTI-EPITOPE FUSION POLYPEPTIDE

Serological expression studies employing *T. cruzi*-infected sera from Mexico and Central America that show little or no reactivity with the *T. cruzi* fusion protein TcF (SEQ ID NO: 1) were carried out to identify known *T. cruzi* antigens that will

complement TcF with the goal of achieving high sensitivity and specificity. TcF (SEQ ID NO: 1), SAPA (SEQ ID NO: 2), Pep30 (SEQ ID NO: 3) and Pep36 (SEQ ID NO: 4) were found to have very good specificity and to complement each other in samples where TcF reactivity is low or zero.

5 Following these studies, a fusion polypeptide containing the TcF, Pep30, Pep36 and SAPA sequences was prepared in two steps.

 In step one, SAPA, Pep30 and Pep36 were fused together with EcoR1 and Xho1 restriction sites to provide the polypeptide sequence of SEQ ID NO: 9 (referred to as SAPA3036; DNA sequence for *E. coli* expression provided in SEQ ID NO: 10) and
10 cloned into a pUC related vector using standard techniques. In step two, SAPA3036 was digested with EcoR1 and Xho1, and subcloned into the 3' end of TcF (SEQ ID NO: 1). The amino acid sequence of the resulting fusion polypeptide (referred to as ITC-6) is provided in SEQ ID NO: 8, with the corresponding DNA sequence being provided in SEQ ID NO: 11. After sequence verification, expression was carried out in the RosettaTM
15 pLysS *E. coli* strain (Novagen, Madison, Wisconsin). The ITC-6 fusion polypeptide has sufficient flexibility at the DNA level for insertion of other antigenic epitopes, such as the KMP-11c *T. cruzi* specific sequence (SEQ ID NO: 5; Thomas et al., *Clin. Exp. Immunol.* 123:465-471, 2001), the Peptide 1 sequence (SEQ ID NO: 6) and the modified Peptide 1 sequence (SEQ ID NO: 7).

20 Oligonucleotides (SEQ ID NO: 12 and 13) were designed to PCR the SAPA3036 fusion protein for cloning directly into the pET28 vector in order to check expression and viability without the TcF component. The resulting PCR product was digested with Nde1 and Xho1 prior to cloning into pET28. There was no problem with expression of SAPA3036 without TcF.

25 It is envisioned that the order of the peptides in the recombinant fusion polypeptide ITC-6 could be altered without significantly changing the activity of the polypeptide. Also, the inclusion of a Gly-Cys-Gly linkage between the peptides may enhance solid phase binding without significantly affecting the activity of the polypeptide.

Example 2

DETECTION OF T. CRUZI INFECTION IN SERA USING ITC7.1 AND ITC 7.2

 The reactivity of sera from *T. cruzi*-infected individuals and control sera from non-infected individuals against the fusion polypeptides TcF and ITC-6 was determined

by ELISA using the procedure described above. The results of this study are shown in Fig. 1. The *T. cruzi*-infected sera RR39, 40218, 63221, PMT201-13, RR66, 40219, 74043 and PMT201-9 had all been shown to be positive for *T. cruzi* by radio immunoprecipitation assay (RIPA). The serum 63225 is consensus positive, with the RIPA status being equivocal. NHS86, NHS92 and PMT201-15 are all normal sera. The 197001, 312001, 38001 and 505001 sera are all from individuals infected with visceral leishmaniasis (VL). The sera were assayed at 1/100 final dilution. Goat anti-human IgG-HRP was used for detection followed by the TMB substrate. As can be seen from Fig. 1, ITC-6 is capable of recognizing sera that are negative or low with TcF.

In subsequent studies, the reactivities of 56 sera (American Red Cross, 42; BBI 14) from the US, Central and South America previously determined to be positive by RIPA, and four endemic control sera from Mexico and Central American countries were tested against the recombinants TcF and ITC-6 by ELISA. The RIPA status of the individual sera are shown in Table 1 below. The coating concentration was 100 ng per well. All sera were tested at 1/100 final dilution. Following incubation, assays were developed using anti-human IgG-HRP and TMB substrate.

The results of these studies are shown in Fig. 2A-C. As predicted, ITC-6 detected *T. cruzi*-infected sera much better than TcF. All the endemic control sera were negative with TcF and ITC-6. Because of very low background and cross-reactivity with TcF and ITC-6, these can be used even at 200 ng per well. While there was 100% concordance with RIPA, with some sera (RR mix and Teragenix mix), the OD was low. These are mixes of several low titer sera and therefore represent dilutions of individual sera that were subsequently used in serological expression screening for additional antigens. In these cases, it is likely that use of other peptides, such as KMP-11 and peptide 1, in combination with ITC-6 will produce higher OD.

Table 1: RIPA status of different sera from North, South and Central America

Serial Number	Serum ID	RIPA status
RR115	<i>T. cruzi</i> (ARC)	Pos
RR99	<i>T. cruzi</i> (ARC)	Pos
RR148	<i>T. cruzi</i> (ARC)	Pos
RR131	<i>T. cruzi</i> (ARC)	Pos
RR52	<i>T. cruzi</i> (ARC)	Pos
RR167	<i>T. cruzi</i> (ARC)	Pos
RR111	<i>T. cruzi</i> (ARC)	Pos
RR71	<i>T. cruzi</i> (ARC)	Pos
RR161	<i>T. cruzi</i> (ARC)	Pos

Serial Number	Serum ID	RIPA status
RR78	<i>T. cruzi</i> (ARC)	Pos
RR04	<i>T. cruzi</i> (ARC)	Pos
RR38	<i>T. cruzi</i> (ARC)	Pos
RR182	<i>T. cruzi</i> (ARC)	Pos
RR102	<i>T. cruzi</i> (ARC)	Pos
RR112	<i>T. cruzi</i> (ARC)	Pos
RR164	<i>T. cruzi</i> (ARC)	Pos
CRR94	<i>T. cruzi</i> (ARC)	Pos
RR119	<i>T. cruzi</i> (ARC)	Pos
RR86	<i>T. cruzi</i> (ARC)	Pos
RR85	<i>T. cruzi</i> (ARC)	Pos
RR34	<i>T. cruzi</i> (ARC)	Pos
RR109	<i>T. cruzi</i> (ARC)	Pos
RR73	<i>T. cruzi</i> (ARC)	Pos
RR190	<i>T. cruzi</i> (ARC)	Pos
CRR101	<i>T. cruzi</i> (ARC)	Pos
RR171	<i>T. cruzi</i> (ARC)	Pos
RR128	<i>T. cruzi</i> (ARC)	Pos
RR105	<i>T. cruzi</i> (ARC)	Pos
RR172	<i>T. cruzi</i> (ARC)	Pos
RR174	<i>T. cruzi</i> (ARC)	Pos
RR150	<i>T. cruzi</i> (ARC)	Pos
RR181	<i>T. cruzi</i> (ARC)	Pos
RR75	<i>T. cruzi</i> (ARC)	Pos
RR57	<i>T. cruzi</i> (ARC)	Pos
RR107	<i>T. cruzi</i> (ARC)	Pos
RR39	<i>T. cruzi</i> (ARC)	Pos
RR189	<i>T. cruzi</i> (ARC)	Pos
RR163	<i>T. cruzi</i> (ARC)	Pos
RR158	<i>T. cruzi</i> (ARC)	Pos
RR157	<i>T. cruzi</i> (ARC)	Pos
RR26	<i>T. cruzi</i> (ARC)	Pos
RR94	<i>T. cruzi</i> (ARC)	Pos
PMT201-1	<i>T. cruzi</i> (BBI)	Pos
PMT201-2	<i>T. cruzi</i> (BBI)	Pos
PMT201-3	<i>T. cruzi</i> (BBI)	Pos
PMT201-4	<i>T. cruzi</i> (BBI)	Pos
PMT201-5	<i>T. cruzi</i> (BBI)	Pos
PMT201-6	<i>T. cruzi</i> (BBI)	Pos
PMT201-7	<i>T. cruzi</i> (BBI)	Pos
PMT201-8	<i>T. cruzi</i> (BBI)	Pos
PMT201-9	<i>T. cruzi</i> (BBI)	Pos
PMT201-10	<i>T. cruzi</i> (BBI)	Pos
PMT201-11	<i>T. cruzi</i> (BBI)	Pos
PMT201-12	<i>T. cruzi</i> (BBI)	Pos
PMT201-13	<i>T. cruzi</i> (BBI)	Pos
PMT201-14	<i>T. cruzi</i> (BBI)	Pos
PMT201-15	Control serum (BBI)	Neg
NHS1	Control serum	neg
NHS2	Control serum	neg

Serial Number	Serum ID	RIPA status
NHS3	Control serum	neg
	Low <i>T. cruzi</i> positive control	RR used in expression cloning
	Low <i>T. cruzi</i> positive control	Teragenix Mix <i>T. cruzi</i> used in expression cloning

In Fig. 3, the reactivities of TcF and ITC-6 are compared with those of ITC-7.1 and 7.2 on a panel of sera. ITC-7.1 and 7.2 contain one and two repeats of the *T. cruzi* specific KMP-11 sequence. Assays were performed as described above. The reactivity of these sera with TcF, ITC-6, ITC-7.1 and 7.2 are compared in Fig. 3. The constructs ITC-7.1 and 7.2 showed comparable activity to ITC-6 with indications of possibly higher signal.

Example 3

DETECTION OF *T. CRUZI* INFECTION IN SERA USING ITC 7.2 AND ITC 8.2

Recombinant protein for ITC8.2 (SEQ ID NO: 19) was expressed as described in Example 6 below. This protein was then used as the solid phase antigen in an ELISA assay and compared with ITC7.2 (SEQ ID NO: 17) which differs from ITC8.2 by not incorporating peptide 1 (SEQ ID NO: 6). Included in the group was a serum sample (sample no. 7190014) which was known to react with peptide 1 as well as other components of the multiepitope recombinant proteins. The results of this study are shown in Fig. 4. As shown, comparable activity was seen with ITC8.2 and ITC7.2.

To ensure that peptide 1 is in fact functional in ITC8.1 (containing the short form of peptide 1 provided in SEQ ID NO: 7) and ITC8.2 (containing the full length peptide 1 provided in SEQ ID NO: 6), absorption studies were performed with a known peptide 1-reactive serum (sample no. 7190014, from the University of Chile) absorbed with beads coated with ITC7.2. Preparation of ITC7.2 coated beads treated with serum #7190014 was carried out as follows:

- 1) 100ul of ITC-7.2 coated beads was centrifuged at 2000 rpm for 1min.
- 2) The supernatant buffer was carefully removed and 1/10 diluted serum sample no. 7190014 in 10mM Tris, pH8.0 was added to the ITC7.2 coated beads.
- 3) The mixture was gently rocked for approximately 1 hr at RT.
- 4) The mixture tube was centrifuged again at 2000rpm for 1min.
- 5) The supernatant was further diluted to a final dilution of 1/20 and 1/100 with Sample Dilution Buffer (SDB) for IgG, 15mM EDTA. Non-treated

serum #7190014 was also directly diluted to 1/20 and 1/100 in serum dilution buffer.

All samples were then tested in an ELISA with 100µl per well of diluted sera being added to coated ITC7.2, shortened ITC8.1 (SEQ ID NO: 22) and ITC8.2 (SEQ ID NO: 19) microtiter strips (3.12 ng/well). *T. cruzi* ELISA was performed as outlined above in Example 1. Fig. 5 shows the activity of non-absorbed and absorbed peptide 1-positive sera. Residual peptide 1 reactivity was observed only in ITC8.2, indicating that the shortened form of peptide 1 in shortened ITC8.1 was either unreactive or less active than the larger peptide 1 sequence present in ITC8.2.

Example 4

IMMUNOBLOT WITH ITC RECOMBINANT PROTEINS

Colloidal gold immunoblots were performed to determine if ITC-6 and ITC7.2 recombinant proteins detected *T. cruzi* positive sera that were low reactive or negative by the parent recombinant protein TcF. Also tested were 12 visceral leishmaniasis (VL) serum samples and sera from four normal donors. The data is shown in Table 2 below.

Briefly, the recombinant proteins were sprayed onto nitrocellulose, blocked with nonfat dried milk, washed in PBS Tween and cut into strips. Strips were incubated in serum at a 1/50 dilution for 15 minutes at 37°C. Strips were then washed in PBS Tween and further incubated for 6 min at ambient temperature with PA-gold (OD5) with ECL 10%. Blots were washed with PBS Tween 20 and observed for the presence or absence of a colored line.

The data (expressed as line intensity) indicate that both ITC6 and ITC7.2 greatly improve the detectability of *T. cruzi* positive sera as compared to TcF without jeopardizing specificity as indicated by no change in reactivity profile with both VL and normal donor sera.

Table 2

Sample	Antigen		
	ITC 6	ITC 7.2	TcF
74042	++++	++++	±
40222	++	++	±
74044	++++	++++	±
40119	++++	++++	-
40224	++	++	-
74040	++++	++++	+
RR148	-	-	-
RR174	+	+	-
RR115	++	++	±
RR161	+++	+++	-
RR131	+++	+++	±
VL474	-	-	-
VL485	-	-	-
VL487	-	-	-
VL494	-	-	-
VL507	-	-	-
VL461	-	-	-
VL453	-	-	-
VL495	-	-	-
VL494	-	-	-
VL477	-	-	-
VL488	-	-	-
VL505	-	-	-
NHS108	-	-	-
NHS117	-	-	-
NHS126	-	-	-
NHS135	-	-	-

5

Example 5**RAPID TEST DATA WITH ITC8.2**

Rapid lateral flow immunoassays were performed using dipsticks prepared with ITC8.2 on the membrane and compared with radioimmunoprecipitation data, immunofluorescence, Lysate and ITC8.2 ELISA. The comparative data is shown in Table 3, below. Good correlation was seen between all assay formats.

10

Table 3

		Chile				
	ELISA ITC8.2	IFAT titer	ELISA lysate	Peptide 1 ELISA	RIPA	Dipstick ITC8.2
	3.13ng/well					
7290140	2.988	320	+	+	+	4+
7290191	2.636	40	+	+	+	3+
7290192	2.785	40	+	+	+	4+
7290083	2.613	40	+	-	+	2+
7290062	1.417	40	+	-	+	1+
7190033	2.713	80	+	+	+	3+
7190024	2.677	40	+	+	+	3+
7190016	2.755	80	+	+	+	3+
7190120(733)	0.055	-	-	-	-	-
7190026	1.966	40	+	+	+	2+
7290129	2.714	20	+	+	+	3+
7290189	1.856	20	+	+	+	1+
7290187	2.436	80	+	+	+	2+
7190012	2.732	160	+	-	+	4+
7190047	2.090	20	+	+	+	2+
7190052	0.060	-	-	-	-	-
7290183	0.574	40	+	-	+	2+
7290147	2.772	+	+	+	+	3+
7190045	2.576	20	+	+	+	4+
7190035	2.432	20	+	+	+	2+
7290144	2.412	+	+	+	+	4+
7190117(736)	0.053	-	-	-	-	+/-
7190114	1.524	80	+		-	1+
7190042	2.426	20	+	+	+	2+
7290146	2.872	160	+	+	+	4+
7190113(755)	0.064	-	-	-	-	-
7290145	2.063	20	+	+	+	2+
7190036	2.515	20	+	+	+	3+
7190119	0.044	80	+	-	-	-
7190116	0.087	320	+	-	-	-
7190115	0.110	20	E	-	-	-
7290190	2.472	160	+	+	+	4+
7190118	0.090	20	E	-	-	+/-
7190014	2.818	40	+	+	+	4+
823	0.051	-	-	-	-	-
838	0.051	-	-	-	-	-
Panel #5	1.576	ND	ND	ND	ND	ND
Panel #6	1.197	ND	ND	ND	ND	ND
Panel #7	0.535	ND	ND	ND	ND	ND

5

Example 6**EXPRESSION OF ITC8.2 IN SUMO EXPRESSION SYSTEM AND IMPROVED SPECIFICITY**

Two expression systems were compared to determine if improved expression of ITC8.2 was possible and also to improve upon specificity. *T. cruzi* positive and negative sera, and an antibody dilutional panel were run in the assay. Negative sera included problematic normal sera. The systems compared were the expression of a Hexahistidine tagged ITC8.2 in a pET17 vector versus an expression system using the SUMO

expression system (no His Tag; Life Sensors, Inc., Malvern, PA) wherein the SUMO fragment was excised with the appropriate protease. The data, which is shown in Table 4, demonstrates the potential for improvement in specificity achieved when using the SUMO-derived recombinant versus preparing the hexahistidine ITC8.2 in a pET vector system.

Table 4

		ITC8.2 (His Tag)	ITC8.2 (SUMO)
	ELISA	20min	20min
7290140	P	4+	4+
7290062	P	1+	2+
7190120(733)	N	-	-
7190052	N	-	-
7290183	P	2+	2+
7190117(736)	N	+/-	-
7190114	P	1+	2+
7190113(755)	N	-	-
7190119	N	-	-
7190116	N	-	-
7190115	N	-	-
7190118	N	+/-?	-
823	N	-	-
838	N	-	-
Chagas D#2	P	4+	4+
Chagas D#4	P	3+	3+
Chagas D#5	P	2.5+	1+
Chagas D#6	P	1.5+	+
BM123204		-	-
BM123207		+/-	-
BM123210		-	-
SeraCare#5		+	-
SeraCare#22		+	-
SeraCare#122		+/-	-
SeraCare#154		+/-	-

Example 7

SEQUENCES DERIVED FROM SEROLOGICAL EXPRESSION CLONING

***T. cruzi* library preparation:**

A genomic random shear expression library was constructed by sonicating genomic DNA from *Trypanosoma cruzi* CL strain. Sonication produced fragment sizes

of 0.5-2.0 kb. Fifteen micrograms of sonicated DNA was treated with T4 polymerase (NEB) for 15 minutes at 12°C followed by incubation for 20 minutes at 75°C to produce blunt ended fragments. EcoRI adaptors were then ligated to the fragments and adaptors were phosphorylated with *E. coli* polynucleotide kinase. Fragments were next
 5 fractionated with a Sephacryl S400 column and finally ligated to a Lambda ZAP Express (Stratagene) vector. Ligated vector was packaged with Gigapack III Gold packaging extract (Stratagene).

Screening:

The amplified library was plated on LB agarose plates at a concentration of 20,000 plaque forming units (PFU) per 35 plates. After incubation at 42°C for 4 hrs, nitrocellulose filters soaked in 10mM IPTG were added and the plates were incubated at 37°C overnight. Filters were removed and washed 3x with PBS containing 0.1% Tween 20 (PBST), blocked for 1 hr with 1% BSA in PBST, washed 3x with PBST, blocked
 15 another 1 hr with 1% Tween 20 in PBS, washed 3x with PBST and then incubated overnight at 4°C in serum, patient pool #1 (RR mix) and/or pool #2 (Teragenix mix). Both patient serum pools were obtained from RIPA-confirmed low reactive *T. cruzi* sera. The following day, after washing 3x with PBST, filters were incubated in an alkaline phosphatase secondary antibody goat anti human Ig (IgG, IgA, IgM) for 1hr at room
 20 temperature. Filters were finally washed 3x with PBST, 2x with AP buffer and developed with BCIP/NBT. Positive clones were purified using the same technique. Phagemid were excised, and resulting plasmid DNA was sequenced and searched against the *T. cruzi* databases.

Features and outcome of library screening:

Lambda vector:	Lambda Zap Express (Stratagene)
Plasmid vector:	pBK-CMV (kanamycin)
DNA:	Genomic <i>T. cruzi</i>
Library titer:	2.5×10^8 / ml (amplified) (total of 30 ml)
Insert size:	0.5-2.0 Kbp (average = 1.1)
Screened:	20,000 pfu per 35 plates
Serum:	Patient pool #1 and pool #2 (from normal donors) 1:200 dilution
Primary Picks:	31 (Human Ig) from patient pool#1 (from 15 plates)

47 (Human Ig) from patient pool#2 (from 20 plates)

Purified Secondary: 38 (weak – strong signal)

Submitted for Sequencing: 12

Table 5: Hits from *T. cruzi* genomic library screening with pooled serum from infected patients

Clone	Score	Blastn	Homology	Size (kDa)	GenBank
Tc-2	1816	Tc00.1047053 509181.9	dispersed gene family protein (DGF-1 pseudogene)		M90534 <i>T. cruzi</i> protein 1 of DGF-1
Tc-5	883	Tc00.1047053 507757.10	hypothetical protein	14.2	Novel
Tc-11	3025	Tc00.1047053 509181.9	dispersed gene family protein (DGF-1 pseudogene)		M90534 <i>T. cruzi</i> protein 1 of DGF-1
Tc-12	3784	Tc00.1047053 509181.9	dispersed gene family protein (DGF-1 pseudogene)		M90534 <i>T. cruzi</i> protein 1 of DGF-1
Tc-13	2765	Tc00.1047053 509181.9	dispersed gene family protein (DGF-1 pseudogene)		M90534 <i>T. cruzi</i> protein 1 of DGF-1
Tc-14	2330	Tc00.1047053 509659.20	dispersed gene family protein (DGF-1 pseudogene)		M90534 <i>T. cruzi</i> protein 1 of DGF-1
Tc-15	3277	Tc00.1407053 511211.170	Heat shock 70	73.2	X58715 <i>T. cruzi</i> hsp 70 mRNA for 70 kDa HSP
Tc-19					
Tc-25	2964	Tc00.1407053 507713.30	Heat shock 85	81	M15346 <i>T. cruzi</i> 85 kDa heat shock protein
Tc-26	3175	Tc00.1407053 510271.20	dispersed gene family protein (DGF-1 pseudogene)		
Tc-27	3362	Tc00.1407053 503739.20	trans-sialidase		
Tc-31	3276		microtubule-associated protein	125.5	

Additional sequences, referred to as Tc48, Tc60 and Tc70, were also identified. These sequences, plus Tc5, were of the most interest for further evaluation. The DNA sequences for Tc5, Tc48, Tc60 and Tc70 are provided in SEQ ID NO: 23, 25, 28 and 30, respectively, with the amino acid sequences for Tc5, Tc60 and Tc70 being provided in SEQ ID NO: 24, 29 and 31. The identified partial amino acid sequence of Tc48 is provided in SEQ ID NO: 26, with the corresponding full-length sequence obtained from the public database being provided in SEQ ID NO: 27.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purpose of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

5 SEQ ID NO: 1-31 are set out in the attached Sequence Listing. The codes for polynucleotide and polypeptide sequences used in the attached Sequence Listing confirm to WIPO Standard ST.25 (1988), Appendix 2.

10 All references disclosed herein, including patent references and non-patent references, are hereby incorporated by reference in their entirety as if each was incorporated individually.

We claim:

1. A fusion polypeptide comprising an amino acid sequence of SEQ ID NO: 1 and at least one amino acid sequence selected from the group consisting of: SEQ ID NO: 2-7, 24, 26, 27, 29 and 31.
2. The fusion polypeptide of claim 1, wherein the fusion polypeptide comprises the amino acid sequences of SEQ ID NO: 1, 2, 3 and 4.
3. The fusion polypeptide of claim 1, wherein the fusion polypeptide comprises the amino acid sequences of SEQ ID NO: 1, 2, 3, 4 and 5.
4. The fusion polypeptide of claim 1, wherein the fusion polypeptide comprises the amino acid sequences of SEQ ID NO: 1, 2, 3, 4, 5 and 6.
5. The fusion polypeptide of claim 1, wherein the fusion polypeptide comprises the amino acid sequences of SEQ ID NO: 1, 2, 3, 4, 5 and 7.
6. The fusion polypeptide of claim 1, wherein the fusion polypeptide comprises an amino acid sequence selected from the group consisting of: SEQ ID NO: 8, 15, 17, 19, and 20.
7. A fusion polypeptide comprising an amino acid sequence selected from the group consisting of:
 - (a) sequences having at least 85% identity to SEQ ID NO: 8, 15, 17, 19 or 20;
 - (b) sequences having at least 90% identity to SEQ ID NO: 8, 15, 17, 19 or 20; and
 - (c) sequences having at least 95% identity to SEQ ID NO: 8, 15, 17, 19 or 20,wherein the fusion polypeptide is capable of generating at least 80% of the response generated by a fusion polypeptide of SEQ ID NO: 8, 15, 17, 19 or 20 to *T. cruzi*-infected sera in an antibody binding assay.

8. An isolated polynucleotide sequence encoding a fusion polypeptide according to any one of claims 1-7.
9. A recombinant expression vector comprising a polynucleotide sequence according to claim 8.
10. A host cell transformed or transfected with an expression vector according to claim 9.
11. A method for detecting *T. cruzi* infection in a biological sample, comprising:
 - (a) contacting the biological sample with a fusion polypeptide of any one of claims 1-7; and
 - (b) detecting in the biological sample the presence of antibodies that bind to the fusion polypeptide, thereby detecting *T. cruzi* infection in the biological sample.
12. The method of claim 11, wherein the biological sample is selected from the group consisting of: blood, serum, plasma, saliva, cerebrospinal fluid and urine.
13. A diagnostic kit for detecting *T. cruzi* infection in a biological sample, comprising:
 - (a) a fusion polypeptide of any one of claims 1-7; and
 - (b) a detection reagent.
14. The kit of claim 13, wherein the detection reagent comprises a reporter group.
15. The kit of claim 14, wherein the reporter group is selected from the group consisting of: enzymes, substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups, and biotin.
16. A composition comprising a fusion protein according to any one of claims 1-7 and at least one component selected from the group consisting of: physiologically acceptable carriers and immunostimulants.

17. A method for the prevention or treatment of *T. cruzi* infection in a patient, comprising administering to the patient a composition of claim 16.

1/6








FIGURE 1

2/6



3/6



4/6

Figure 4





Figure 5



5/6

Figure 6A

ATGGGCAGCAGCCATCATCATCATCACAGCAGCGGCCTGGTGCCGCG
CGGCAGCCATATGGCCGCAGGCGGAGAGCTGGCGGCAACAGCAGCTG
CAGTGCGGGGACAGTTTGCATCGTGTGCCCCGTGGGGGAAGGACGAC
ATGGCGCAGGATGGGCTTCTGGAGGCTGAGCTTTTCACCCCGGTGGA
TGAGAGCCTCTAGTCCAGCAGACGTGCTGAGCATGGGGAGAAGGAGC
TGTGACCGCCCCGTATTTTGGCCTCAGAGCCAGAGTGCACCCTGTGC
AAGGCAAAAAGGGGCGAAGTCGACACAATGCGACTTGTCCACGAAGC
GCTTCTTCCTCCCATGCTCTGGCAGAGGCGTGCCGGGAGGATACTGC
GAACAGTTGCGCTTTTGTGTGGCTGTGGTGGCGGGGAGGGCGGAG
GGGCAGACTCAACTGAAAGCTT

T This is 'A' in XM803645
G This is 'A' in XM803645
G This is 'C' in XM803645

Figure 6B

MGSSHHHHHHSSGLVPRGSHMAAGGELAATAAAVRGQFASCARGGRTT
WRRMGFWRLSFSRWMRASSPADVLSMGRRSCDRPVFWPQSQSAPCRR
QKGAKSTQCDLSTKRFFLPSCGRGVPGGYCEQLRFCCWLWWRGGRRG
RLN*

Figure 7A

ATGGGCAGCAGCCATCATCATCATCACAGCAGCGGCCTGGTGCCGCG
CGGCAGCCATATGCCCAAAAAGACCGGTGGCAAAAAGAAGGGGCAAA
GTTCTCCGGATGGCTCTGAGCCGCGGAAACGAAAGAACAACAAAAG
GCGACAATGGAGCCGCGGGACGTGGATGAGATGCAGAAGCTGCAGG
AACTTTTAGGGGACGAGGAACAGCCGTTGGGTGTCTCCAAGAAATCG
CTAGAGGGCTTATTGTCCCTTCGGCAGCCGCAGGAGTTGGCGGTGAG
GCTTGCGCAATCTCTCTCCTCCCTGCGCGCGCGGCTTGCGGAGTTGG
AGTTGGAGAGGCTTAACCGTGGGAGCGAGGCGCCGGGGCTGTGCGAA
CATCGT...

Figure 7B

MGSSHHHHHHSSGLVPRGSHMPKKTGGKKKGQSSPDGSEPRKRKNNKKA
TMEPRDVEDEMQKLQELLGDEEQPLGVSKKSLEGLLSLRQPQELAVRLA
QSLSSLRARLAELELERLNRGSEAPGLSNI...

6/6

Figure 8A

CCATGGGCAGCAGCCATCATCATCATCATCACAGCAGCGGCCTGGTGCCG
CGCGGCAGCCATATGTGCATTGCTCTTGGCATCGTCGCGGAGGATGAT
GAGGCATGGCGAATTGCCAGCGAGGCTGTCGCTGCAGAAAAGGAGC
CTGTCTTCAGCGGGAACAACGGCCCTTTTGTAGATGTCTGGTTTGGC
GAACAGAACTCTTTGGCCTCGTTCAACGCGTTGCTCCAAACGACTTT
ATTCAGGTCGCCCAGGAGTGTGGCGAGAAGAGCGATGACGCCGCAG
CGACGTTGCGGATGCGTGTGACGCACAACGTCTCTTTTGTCTTCACC
TCTCGTCGGTGCCGCATGCGATGCTGCAGGCACGGGGAGCGCCCGA
GGACAAGTTTGTGAACTTCATGCAACTTGTCGTGGATTACGCTTCGCT
GCTGCGGCGCGGGATGAAGGATGAGTTTCTTGGCGTCGATCCCGAGT
CCGATGCGGAGTACATACGCTTCACGCCCCAGTGAAAGCTT

Figure 8B

MGSSHHHHHHSSGLVPRGSHMCIALGIVAEDDEAWRIASEAVAAEKEPVFS
GNNGPFVDVWFGEQKLFLVQRVAPNDFIQVAQECGEKSDDAAATLRM
RVTHNVSFVLHLSSVPHAMLQARGAPEDKFVNFMQLVVDYASLLRRGM
KDEFLGVDPESDAEYIRFTPQ*

Figure 9A

CCATGGGCAGCAGCCATCATCATCATCATCACAGCAGCGGCCTGGTGCCG
CGCGGCAGCCATATGTGTTCCCAATACGGACCCTTGAGGAACTCATCG
CAGCGTGGAAGACCGATCGCTGTGTCTGTTGCCAACTCGTGTCTTCG
CGTAGCGGCCACGCAGAAAGTTTCTAAGCCACCTTCTACAGTTCACC
CCCGGAATATTGGCCGGCAAGCGACTGAGGACTCGATGACCAATGAA
CTCAAAGGCCTTGCTGGAGTCTACCAGCACCAACGGAGCCCGATGGG
GTCTGCAGTGGAGCTGGCTTCCAACACCGCTCTTCCTGGGAAGGTTC
ACTTGGAATTAATCGTTTCTGTTATGCTCAAATTCGTTTATCAGGTGT
GCCAGCTGCATCGTCGTGGTACGCATACTACGGCACGCATACTGGTG
CGGACGAGGCGTCCACAGCCAAAGCAGTCTCCTCAATGCCATTTTCT
CAGCAACCTTACCCACGATGGAAGGACAGCTGATTGGTCGATATGAA
GCTT

Figure 9B

MGSSHHHHHHSSGLVPRGSHMCSQYGPLRNSSQRRGRPIAVSVANSCLRVA
ATQKVS KPSTVHPRNIGRQATEDSMTNELKGLAGVYQHQRSPMGSVE
LASNTALPGKVHLELIVSVMLKFVYQVCQLHRRGTHTTARILVRTRRPQ
PKQSPQCHFLSNLTHDGRTADWSI*

SEQUENCE LISTING

<110> Raychaudhuri, Syamal
Houghton, Raymond L.
Reed, Steven G.

<120> Compositions and Methods for the
Detection of *Trypanosoma Cruzi* Infection

<130> 13379.1001PCT

<150> 60/733,119

<151> 2005-11-03

<160> 31

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 86

<212> PRT

<213> T. Cruzi

<400> 1

Met	Gln	His	His	His	His	His	Gly	Asp	Lys	Pro	Ser	Pro	Phe	Gly
1			5					10					15	
Gln	Ala	Ala	Ala	Gly	Asp	Lys	Pro	Ser	Pro	Phe	Gly	Gln	Ala	Glu
		20					25					30		
Pro	Lys	Ser	Ala	Glu	Pro	Lys	Pro	Ala	Glu	Pro	Lys	Ser	Lys	Ala
		35				40					45			
Ile	Ala	Pro	Ala	Lys	Ala	Ala	Ala	Ala	Pro	Ala	Lys	Ala	Ala	Thr
	50				55				60					
Pro	Ala	Ser	Ser	Met	Pro	Ser	Gly	Thr	Ser	Glu	Glu	Gly	Ser	Arg
65					70				75					80
Gly	Ser	Ser	Met	Pro	Ala									
			85											

<210> 2

<211> 25

<212> PRT

<213> T. Cruzi

<400> 2

Tyr	Asp	Ser	Thr	Ala	His	Gly	Thr	Pro	Ser	Thr	Pro	Ala	Asp	Ser
1			5					10					15	
Ala	His	Ser	Thr	Pro	Ser	Thr	Pro	Ala						
		20					25							

<210> 3

<211> 29

<212> PRT

<213> T. Cruzi

<400> 3

Tyr	Glu	Lys	Gln	Lys	Ala	Ala	Glu	Ala	Thr	Lys	Val	Ala	Glu	Ala
1			5					10					15	
Lys	Gln	Arg	Ala	Ala	Glu	Ala	Thr	Lys	Val	Ala	Glu	Ala		
		20					25							

<210> 4
 <211> 38
 <212> PRT
 <213> T. Cruzei

<400> 4
 Ala Leu Pro Gln Glu Glu Gln Glu Asp Val Gly Pro Arg His Val Asp
 1 5 10 15
 Pro Asp His Phe Arg Ser Thr Thr Gln Asp Ala Tyr Arg Pro Val Asp
 20 25 30
 Pro Ser Ala Tyr Lys Arg
 35

<210> 5
 <211> 17
 <212> PRT
 <213> T. cruzi

<400> 5
 Lys Phe Ala Glu Leu Leu Glu Gln Gln Lys Asn Ala Gln Phe Pro Gly
 1 5 10 15
 Lys

<210> 6
 <211> 68
 <212> PRT
 <213> T. cruzi

<400> 6
 Met Glu Gln Glu Arg Arg Gln Leu Leu Glu Lys Asp Pro Arg Arg Asn
 1 5 10 15
 Ala Lys Glu Ile Ala Ala Leu Glu Glu Ser Met Asn Ala Arg Ala Gln
 20 25 30
 Glu Leu Ala Arg Glu Lys Lys Leu Ala Asp Arg Ala Phe Leu Asp Gln
 35 40 45
 Lys Pro Glu Arg Val Pro Leu Ala Asp Val Pro Leu Asp Asp Asp Ser
 50 55 60
 Asp Phe Val Ala
 65

<210> 7
 <211> 60
 <212> PRT
 <213> T. cruzi

<400> 7
 Arg Ala Gln Glu Leu Ala Arg Glu Lys Lys Leu Ala Asp Arg Ala Phe
 1 5 10 15
 Leu Asp Gln Lys Pro Glu Arg Val Pro Leu Ala Asp Val Pro Leu Asp
 20 25 30
 Asp Asp Ser Asp Phe Val Ala Met Glu Gln Glu Arg Arg Gln Leu Leu
 35 40 45
 Glu Lys Asp Pro Arg Arg Asn Ala Lys Glu Ile Ala
 50 55 60

<210> 8
 <211> 182
 <212> PRT
 <213> T. Cruzei

<400> 8

```

Met Gln His His His His His His Gly Asp Lys Pro Ser Pro Phe Gly
 1          5          10          15
Gln Ala Ala Ala Gly Asp Lys Pro Ser Pro Phe Gly Gln Ala Ala Glu
 20          25          30
Pro Lys Ser Ala Glu Pro Lys Pro Ala Glu Pro Lys Ser Lys Ala Ala
 35          40          45
Ile Ala Pro Ala Lys Ala Ala Ala Pro Ala Lys Ala Ala Thr Ala
 50          55          60
Pro Ala Ser Ser Met Pro Ser Gly Thr Ser Glu Glu Gly Ser Arg Gly
 65          70          75          80
Gly Ser Ser Met Pro Ala Glu Phe Tyr Asp Ser Thr Ala His Gly Thr
 85          90          95
Pro Ser Thr Pro Ala Asp Ser Ser Ala His Ser Thr Pro Ser Thr Pro
100          105          110
Ala Tyr Glu Lys Gln Lys Ala Ala Glu Ala Thr Lys Val Ala Glu Ala
115          120          125
Glu Lys Gln Arg Ala Ala Glu Ala Thr Lys Val Ala Glu Ala Ala Leu
130          135          140
Pro Gln Glu Glu Gln Glu Asp Val Gly Pro Arg His Val Asp Pro Asp
145          150          155          160
His Phe Arg Ser Thr Thr Gln Asp Ala Tyr Arg Pro Val Asp Pro Ser
165          170          175
Ala Tyr Lys Arg Lys Leu
180

```

<210> 9

<211> 92

<212> PRT

<213> T. Cruzei

<400> 9

```

Tyr Asp Ser Thr Ala His Gly Thr Pro Ser Thr Pro Ala Asp Ser Ser
 1          5          10          15
Ala His Ser Thr Pro Ser Thr Pro Ala Tyr Glu Lys Gln Lys Ala Ala
 20          25          30
Glu Ala Thr Lys Val Ala Glu Ala Glu Lys Gln Arg Ala Ala Glu Ala
 35          40          45
Thr Lys Val Ala Glu Ala Ala Leu Pro Gln Glu Glu Gln Glu Asp Val
 50          55          60
Gly Pro Arg His Val Asp Pro Asp His Phe Arg Ser Thr Thr Gln Asp
 65          70          75          80
Ala Tyr Arg Pro Val Asp Pro Ser Ala Tyr Lys Arg
 85          90

```

<210> 10

<211> 297

<212> DNA

<213> T. Cruzei

<400> 10

```

gaattctatg attccaccgc ccatggtacc cctagcaccc ccgccgattc ctccgcccat      60
tccaccccat ccaccccgag ctatgaaaaa caaaaagccg ccgaagccac caaagtagca      120
gaagccgaaa aacaacgtgc cgcagaagcc accaaagtag ccgaagccgc cctcccccaa      180
gaagaacaag aagatgtcgg tccccgccat gtagatccag accactttcg ctccaccaca      240
caagatgcct accgcccgat tgaccctctc gcctataaac gtaagctttg actcgag      297

```

<210> 11

<211> 549

<212> DNA

<213> T. cruzi

<400> 11

atgcagcatc	accaccatca	ccacggtgat	aagcctagcc	catttggtca	ggctgctgct	60
ggtgacaaac	cgtctccggt	cggtcaggct	gctgaaccga	aatctgctga	accgaaaccg	120
gctgaaccga	aaagcaaagc	tgctatcgct	ccggctaaag	ctgctgctgc	tccggctaaa	180
gctgctaccg	ctccggcctc	ttctatgccg	tctggtacct	ctgaagaagg	ttctcgtggt	240
ggttctttcta	tgccggccga	attctatgat	tccaccgccc	atggtacccc	tagcaccccc	300
gccgattcct	ccgcccattc	caccccatcc	accccgacct	atgaaaaaca	aaaagccgcc	360
gaagccacca	aagtagcaga	agccgaaaaa	caacgtgccg	cagaagccac	caaagtagcc	420
gaagccgccc	tcccccaaga	agaacaagaa	gatgtcggtc	cccgccatgt	agatccagac	480
cactttcgct	ccaccacaca	agatgcctac	cgccagttg	acccctccgc	ctataaacgt	540
aagctttga						549

<210> 12

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 12

caattacata	tgtatgattc	caccgccccat	ggt	33
------------	------------	-------------	-----	----

<210> 13

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 13

caattactcg	agtcaaagct	tacgtttata	ggc	33
------------	------------	------------	-----	----

<210> 14

<211> 621

<212> DNA

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 14

catatgcagc	atcaccacca	tcaccacggt	gataagccta	gcccathttgg	tcaggctgct	60
gctggtgaca	aaccgtctcc	gttcggtcag	gctgctgaac	cgaaatctgc	tgaaccgaaa	120
ccggctgaac	cgaaaagcaa	agctgctatc	gctccggcta	aagctgctgc	tgtccgggct	180
aaagctgcta	ccgctccggc	ctcttctatg	ccgtctggta	cctctgaaga	aggttctcgt	240
ggtggttctt	ctatgccggc	cgaattctat	gattccaccg	cccatggtag	ccctagcacc	300
cccgcggatt	cctccgcccc	ttccaccccc	tccaccccag	cctatgaaaa	acaaaaagcc	360
gccgaagcca	ccaaagtagc	agaagccgaa	aaacaacgtg	ccgcagaagc	caccaaagta	420
gccgaagccg	ccctccccca	agaagaacaa	gaagatgtcg	gtccccgcca	tgtagatcca	480
gaccactttc	gtccaccac	acaagatgcc	taccgcccag	ttgacccttc	cgccataaaa	540
cgtaagctta	aatttgcgga	actgctggaa	cagcagaaaa	acgcgcagtt	tccgggcaaa	600
gagctcacta	gttaactcga	g				621

<210> 15

<211> 203

<212> PRT

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 15

```

Met Gln His His His His His His Gly Asp Lys Pro Ser Pro Phe Gly
1      5      10      15
Gln Ala Ala Gly Asp Lys Pro Ser Pro Phe Gly Gln Ala Glu
20      25      30
Pro Lys Ser Ala Glu Pro Lys Pro Ala Glu Pro Lys Ser Lys Ala Ala
35      40      45
Ile Ala Pro Ala Lys Ala Ala Ala Ala Pro Ala Lys Ala Ala Thr Ala
50      55      60
Pro Ala Ser Ser Met Pro Ser Gly Thr Ser Glu Glu Gly Ser Arg Gly
65      70      75      80
Gly Ser Ser Met Pro Ala Glu Phe Tyr Asp Ser Thr Ala His Gly Thr
85      90      95
Pro Ser Thr Pro Ala Asp Ser Ser Ala His Ser Thr Pro Ser Thr Pro
100      105      110
Ala Tyr Glu Lys Gln Lys Ala Ala Glu Ala Thr Lys Val Ala Glu Ala
115      120      125
Glu Lys Gln Arg Ala Ala Glu Ala Thr Lys Val Ala Glu Ala Ala Leu
130      135      140
Pro Gln Glu Glu Gln Glu Asp Val Gly Pro Arg His Val Asp Pro Asp
145      150      155      160
His Phe Arg Ser Thr Gln Asp Ala Tyr Arg Pro Val Asp Pro Ser
165      170      175
Ala Tyr Lys Arg Lys Leu Lys Phe Ala Glu Leu Leu Glu Gln Gln Lys
180      185      190
Asn Ala Gln Phe Pro Gly Lys Glu Leu Thr Ser
195      200

```

<210> 16

<211> 672

<212> DNA

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 16

```

catatgcagc atcaccacca tcaccacggt gataagccta gcccatttgg tcaggctgct      60
gctggtgaca aaccgtctcc gttcgggtcag gctgctgaac cgaaatctgc tgaaccgaaa      120
ccggctgaac cgaaaagcaa agctgctatc gctccggcta aagctgctgc tgctccggct      180
aaagctgcta ccgctccggc ctcttctatg ccgtctggta cctctgaaga aggttctcgt      240
ggtggttctt ctatgccggc cgaattctat gattccaccg cccatgggtac ccctagcacc      300
cccgccgatt cctccgcccc ttccacccca tccacccag cctatgaaaa acaaaaagcc      360
gccgaagcca ccaaagtagc agaagccgaa aaacaacgtg ccgcagaagc caccaaagta      420
gccgaagccg ccctcccccagaagaacaa gaagatgtcg gtccccgcca tgtagatcca      480
gaccactttc gtcaccacac acaagatgcc taccgcccag ttgaccctc cgcctataaa      540
cgtaagctta aatttgcgga actgctggaa cagcagaaaa acgcgcagtt tccgggcaaa      600
gagctcaaat ttgcggaact gctggaacag cagaaaaaac cgcagtttcc gggcaaagag      660
ctcactagtt aa                                     672

```

<210> 17

<211> 219

<212> PRT

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 17

```

His Met Gln His His His His His His Gly Asp Lys Pro Ser Pro Phe
 1           5           10           15
Gly Gln Ala Ala Ala Gly Asp Lys Pro Ser Pro Phe Gly Gln Ala Ala
          20           25           30
Glu Pro Lys Ser Ala Glu Pro Lys Pro Ala Glu Pro Lys Ser Lys Ala
          35           40           45
Ala Ile Ala Pro Ala Lys Ala Ala Ala Pro Ala Lys Ala Ala Thr
          50           55           60
Ala Pro Ala Ser Ser Met Pro Ser Gly Thr Ser Glu Glu Gly Ser Arg
65           70           75           80
Gly Gly Ser Ser Met Pro Ala Glu Phe Tyr Asp Ser Thr Ala His Gly
          85           90           95
Thr Pro Ser Thr Pro Ala Asp Ser Ser Ala His Ser Thr Pro Ser Thr
          100          105          110
Pro Ala Tyr Glu Lys Gln Lys Ala Ala Glu Ala Thr Lys Val Ala Glu
          115          120          125
Ala Glu Lys Gln Arg Ala Ala Glu Ala Thr Lys Val Ala Glu Ala Ala
          130          135          140
Leu Pro Gln Glu Glu Gln Glu Asp Val Gly Pro Arg His Val Asp Pro
145          150          155          160
Asp His Phe Arg Ser Thr Thr Gln Asp Ala Tyr Arg Pro Val Asp Pro
          165          170          175
Ser Ala Tyr Lys Arg Lys Leu Lys Phe Ala Glu Leu Leu Glu Gln Gln
          180          185          190
Lys Asn Ala Gln Phe Pro Gly Lys Glu Leu Lys Phe Ala Glu Leu Leu
          195          200          205
Glu Gln Gln Lys Asn Ala Gln Phe Pro Gly Lys
          210          215

```

<210> 18

<211> 882

<212> DNA

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 18

```

catatgcagc atcaccacca tcaccacggt gataagccta gccatttgg tcaggctgct      60
gctggtgaca aaccgtctcc gttcggtcag gctgctgaac cgaaatctgc tgaaccgaaa      120
ccggctgaac cgaaaagcaa agctgctatc gctccggcta aagctgctgc tgctccggct      180
aaagctgcta ccgctccggc ctcttctatg ccgtctggta cctctgaaga aggttctcgt      240
ggtggttctt ctatgccggc cgaattctat gattccaccg cccatggtac ccctagcacc      300
cccgccgatt cctccgcca ttccacccca tccacccag cctatgaaaa acaaaaagcc      360
gccgaagcca ccaaagtagc agaagccgaa aaacaacgtg ccgcagaagc caccaaagta      420
gccgaagccg ccctccccc agaagaacaa gaagatgtcg gtccccgcca tgtagatcca      480
gaccactttc gtcaccacc acaagatgcc taccgcccag ttgaccctc cgcctataaa      540
cgtaagctta aatttgcgga actgctggaa cagcagaaaa acgcgcagtt tccgggcaaa      600
gagctcaaat ttgcggaact gctggaacag cagaaaaacg cgcagtttcc gggcaaagag      660
ctcactagta tggaacagga acgccgccag ctgctggaaa aagatccgcg ccgcaacgcg      720
aaagaaattg cggcgctgga agaaagcatg aacgcgcgcg cgcaggaact ggcgcgcgaa      780
aaaaaactgg cggatcgcgc gtttctggat cagaaaccgg aacgcgtccc gctggcggat      840
gtcccgcgtg atgatgatag cgattttgtc gcgtaactcg ag                        882

```

<210> 19

<211> 291

<212> PRT

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 19

His Met Gln His His His His His His Gly Asp Lys Pro Ser Pro Phe
 1 5 10 15
 Gly Gln Ala Ala Ala Gly Asp Lys Pro Ser Pro Phe Gly Gln Ala Ala
 20 25 30
 Glu Pro Lys Ser Ala Glu Pro Lys Pro Ala Glu Pro Lys Ser Lys Ala
 35 40 45
 Ala Ile Ala Pro Ala Lys Ala Ala Ala Pro Ala Lys Ala Ala Thr
 50 55 60
 Ala Pro Ala Ser Ser Met Pro Ser Gly Thr Ser Glu Glu Gly Ser Arg
 65 70 75 80
 Gly Gly Ser Ser Met Pro Ala Glu Phe Tyr Asp Ser Thr Ala His Gly
 85 90 95
 Thr Pro Ser Thr Pro Ala Asp Ser Ser Ala His Ser Thr Pro Ser Thr
 100 105 110
 Pro Ala Tyr Glu Lys Gln Lys Ala Ala Glu Ala Thr Lys Val Ala Glu
 115 120 125
 Ala Glu Lys Gln Arg Ala Ala Glu Ala Thr Lys Val Ala Glu Ala Ala
 130 135 140
 Leu Pro Gln Glu Glu Gln Glu Asp Val Gly Pro Arg His Val Asp Pro
 145 150 155 160
 Asp His Phe Arg Ser Thr Thr Gln Asp Ala Tyr Arg Pro Val Asp Pro
 165 170 175
 Ser Ala Tyr Lys Arg Lys Leu Lys Phe Ala Glu Leu Leu Glu Gln Gln
 180 185 190
 Lys Asn Ala Gln Phe Pro Gly Lys Glu Leu Lys Phe Ala Glu Leu Leu
 195 200 205
 Glu Gln Gln Lys Asn Ala Gln Phe Pro Gly Lys Glu Leu Thr Ser Met
 210 215 220
 Glu Gln Glu Arg Arg Gln Leu Leu Glu Lys Asp Pro Arg Arg Asn Ala
 225 230 235 240
 Lys Glu Ile Ala Ala Leu Glu Glu Ser Met Asn Ala Arg Ala Gln Glu
 245 250 255
 Leu Ala Arg Glu Lys Lys Leu Ala Asp Arg Ala Phe Leu Asp Gln Lys
 260 265 270
 Pro Glu Arg Val Pro Leu Ala Asp Val Pro Leu Asp Asp Asp Ser Asp
 275 280 285
 Phe Val Ala
 290

<210> 20

<211> 279

<212> PRT

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 20

His Met Gln His His His His His His Gly Asp Lys Pro Ser Pro Phe
 1 5 10 15
 Gly Gln Ala Ala Ala Gly Asp Lys Pro Ser Pro Phe Gly Gln Ala Ala
 20 25 30
 Glu Pro Lys Ser Ala Glu Pro Lys Pro Ala Glu Pro Lys Ser Lys Ala
 35 40 45
 Ala Ile Ala Pro Ala Lys Ala Ala Ala Ala Pro Ala Lys Ala Ala Thr
 50 55 60

Ala Pro Ala Ser Ser Met Pro Ser Gly Thr Ser Glu Glu Gly Ser Arg
65 70 75 80
Gly Gly Ser Ser Met Pro Ala Glu Phe Tyr Asp Ser Thr Ala His Gly
85 90 95
Thr Pro Ser Thr Pro Ala Asp Ser Ser Ala His Ser Thr Pro Ser Thr
100 105 110
Pro Ala Tyr Glu Lys Gln Lys Ala Ala Glu Ala Thr Lys Val Ala Glu
115 120 125
Ala Glu Lys Gln Arg Ala Ala Glu Ala Thr Lys Val Ala Glu Ala Ala
130 135 140
Leu Pro Gln Glu Glu Gln Glu Asp Val Gly Pro Arg His Val Asp Pro
145 150 155 160
Asp His Phe Arg Ser Thr Thr Gln Asp Ala Tyr Arg Pro Val Asp Pro
165 170 175
Ser Ala Tyr Lys Arg Lys Leu Lys Phe Ala Glu Leu Leu Glu Gln Gln
180 185 190
Lys Asn Ala Gln Phe Pro Gly Lys Glu Leu Lys Phe Ala Glu Leu Leu
195 200 205
Glu Gln Gln Lys Asn Ala Gln Phe Pro Gly Lys Arg Ala Gln Glu Leu
210 215 220
Ala Arg Glu Lys Lys Leu Ala Asp Arg Ala Phe Leu Asp Gln Lys Pro
225 230 235 240
Glu Arg Val Pro Leu Ala Asp Val Pro Leu Asp Asp Asp Ser Asp Phe
245 250 255
Val Ala Met Glu Gln Glu Arg Arg Gln Leu Leu Glu Lys Asp Pro Arg
260 265 270
Arg Asn Ala Lys Glu Ile Ala
275

<210> 21

<211> 786

<212> DNA

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 21

atgcagcatc	accaccatca	ccacggtgat	aagcctagcc	catttggtca	ggctgctgct	60
ggtgacaaac	cgtctccgtt	cggtcaggct	gctgaaccga	aatctgctga	accgaaaccg	120
gctgaaccga	aaagcaaagc	tgctatcgct	cgggctaaag	ctgctgctgc	tccggctaaa	180
gctgctaccg	ctccggcctc	ttctatgccg	tctggtacct	ctgaagaagg	ttctcgtggt	240
ggtttcttcta	tgccggccga	attctatgat	tccaccgccc	atggtacccc	tagcaccccc	300
gccgattcct	ccgcccattc	caccccatcc	accccagcct	atgaaaaaca	aaaagccgcc	360
gaagccacca	aagtagcaga	agccgaaaaa	caacgtgccg	cagaagccac	caaagtagcc	420
gaagccgccc	tcccccaaga	agaacaagaa	gatgtcggtc	cccgccatgt	agatccagac	480
cactttcgct	ccaccacaca	agatgcctac	cgcccagttg	acccctccgc	ctataaacgt	540
aagcttaaat	ttgcggaact	gctggaacag	cagaadaaacg	cgcagtttcc	gggcaaagag	600
ctcaaatttg	cggaactgct	ggaacagcag	aaaaacgcgc	agtttccggg	caaagagctc	660
actagtcgcg	cgcaggaact	ggcgcgcgaa	aaaaaactgg	cggatcgcgc	gtttctggat	720
cagaaaccgg	aacgcgtccc	gctggcggat	gtcccgcgtg	atgatgatag	cgattttgtc	780
gcgtaa						786

<210> 22

<211> 261

<212> PRT

<213> Artificial Sequence

<220>

<223> Made in a lab

<400> 22

```

Met Gln His His His His His His Gly Asp Lys Pro Ser Pro Phe Gly
 1          5          10          15
Gln Ala Ala Ala Gly Asp Lys Pro Ser Pro Phe Gly Gln Ala Ala Glu
          20          25          30
Pro Lys Ser Ala Glu Pro Lys Pro Ala Glu Pro Lys Ser Lys Ala Ala
          35          40          45
Ile Ala Pro Ala Lys Ala Ala Ala Ala Pro Ala Lys Ala Ala Thr Ala
          50          55          60
Pro Ala Ser Ser Met Pro Ser Gly Thr Ser Glu Glu Gly Ser Arg Gly
65          70          75          80
Gly Ser Ser Met Pro Ala Glu Phe Tyr Asp Ser Thr Ala His Gly Thr
          85          90          95
Pro Ser Thr Pro Ala Asp Ser Ser Ala His Ser Thr Pro Ser Thr Pro
          100          105          110
Ala Tyr Glu Lys Gln Lys Ala Ala Glu Ala Thr Lys Val Ala Glu Ala
          115          120          125
Glu Lys Gln Arg Ala Ala Glu Ala Thr Lys Val Ala Glu Ala Ala Leu
          130          135          140
Pro Gln Glu Glu Gln Glu Asp Val Gly Pro Arg His Val Asp Pro Asp
145          150          155          160
His Phe Arg Ser Thr Thr Gln Asp Ala Tyr Arg Pro Val Asp Pro Ser
          165          170          175
Ala Tyr Lys Arg Lys Leu Lys Phe Ala Glu Leu Leu Glu Gln Gln Lys
          180          185          190
Asn Ala Gln Phe Pro Gly Lys Glu Leu Lys Phe Ala Glu Leu Leu Glu
          195          200          205
Gln Gln Lys Asn Ala Gln Phe Pro Gly Lys Glu Leu Thr Ser Arg Ala
          210          215          220
Gln Glu Leu Ala Arg Glu Lys Lys Leu Ala Asp Arg Ala Phe Leu Asp
225          230          235          240
Gln Lys Pro Glu Arg Val Pro Leu Ala Asp Val Pro Leu Asp Asp Asp
          245          250          255
Ser Asp Phe Val Ala
          260

```

<210> 23

<211> 447

<212> DNA

<213> T. cruzi

<400> 23

```

atgggcagca gccatcatca tcatcatcac agcagcggcc tgggtgccgcg cggcagccat      60
atggccgcag gcggagagct ggcggcaaca gcagctgcag tgcggggaca gtttgcacgc      120
tgtgcccgtg ggggaaggac gacatggcgc aggatgggct tctggaggct gagcttttca      180
ccccggtgga tgagagcctc tagtccagca gacgtgctga gcatggggag aaggagctgt      240
gaccgccccg tattttggcc tcagagccag agtgcaccct gtcgaaggca aaaaggggag      300
aagtgcacac aatgcgactt gtccacgaag cgcttcttcc tcccatgctc tggcagaggg      360
gtgccgggag gatactgcga acagttgcgc ttttggtggt ggctgtggtg gcggggaggg      420
cggaggggca gactcaactg aaagctt                                447

```

<210> 24

<211> 146

<212> PRT

<213> T. cruzi

<400> 24

```

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
 1          5          10          15
Arg Gly Ser His Met Ala Ala Gly Gly Glu Leu Ala Ala Thr Ala Ala
          20          25          30

```

Ala Val Arg Gly Gln Phe Ala Ser Cys Ala Arg Gly Gly Arg Thr Thr
 35 40 45
 Trp Arg Arg Met Gly Phe Trp Arg Leu Ser Phe Ser Pro Arg Trp Met
 50 55 60
 Arg Ala Ser Ser Pro Ala Asp Val Leu Ser Met Gly Arg Arg Ser Cys
 65 70 75 80
 Asp Arg Pro Val Phe Trp Pro Gln Ser Gln Ser Ala Pro Cys Arg Arg
 85 90 95
 Gln Lys Gly Ala Lys Ser Thr Gln Cys Asp Leu Ser Thr Lys Arg Phe
 100 105 110
 Phe Leu Pro Cys Ser Gly Arg Gly Val Pro Gly Gly Tyr Cys Glu Gln
 115 120 125
 Leu Arg Phe Cys Cys Trp Leu Trp Trp Arg Gly Gly Arg Arg Gly Arg
 130 135 140
 Leu Asn
 145

<210> 25
 <211> 383
 <212> DNA
 <213> T. cruzi

<400> 25
 atgggcagcagccatcatca tcatcatcac agcagcggcc tggcgccgcg cggcagccat 60
 atgccccaaaa agaccggtgg caaaaagaag gggcaaagtt ctccgcatgg ctctgagccg 120
 cggaaacgaa agaacaacaa aaaggcgaca atggagccgc gggacgtgga tgagatgcag 180
 aagctgcagg aacttttagg ggacgaggaa cagccgttgg gtgtctccaa gaaatcgcta 240
 gagggttat tgtcccttcg gcagccgcag gagttggcgg tgaggcttgc gcaatctctc 300
 tcctccctgc gcgcgcggct tgcggagttg gagttggaga ggcttaaccg tgggagcgag 360
 gcgcgggggc tgtcgaacat cgt 383

<210> 26
 <211> 127
 <212> PRT
 <213> T. cruzi

<400> 26
 Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
 1 5 10 15
 Arg Gly Ser His Met Pro Lys Lys Thr Gly Gly Lys Lys Lys Gly Gln
 20 25 30
 Ser Ser Pro Asp Gly Ser Glu Pro Arg Lys Arg Lys Asn Asn Lys Lys
 35 40 45
 Ala Thr Met Glu Pro Arg Asp Val Asp Glu Met Gln Lys Leu Gln Glu
 50 55 60
 Leu Leu Gly Asp Glu Glu Gln Pro Leu Gly Val Ser Lys Lys Ser Leu
 65 70 75 80
 Glu Gly Leu Leu Ser Leu Arg Gln Pro Gln Glu Leu Ala Val Arg Leu
 85 90 95
 Ala Gln Ser Leu Ser Ser Leu Arg Ala Arg Leu Ala Glu Leu Glu Leu
 100 105 110
 Glu Arg Leu Asn Arg Gly Ser Glu Ala Pro Gly Leu Ser Asn Ile
 115 120 125

<210> 27
 <211> 400
 <212> PRT
 <213> T. cruzi

<400> 27
 Met Pro Lys Lys Thr Gly Gly Lys Lys Lys Gly Gln Ser Ser Pro Asp

1	5	10	15
Gly Ser Glu Pro Arg Lys Arg Lys Asn Asn Lys Lys Ala Thr Met Glu			
20	25	30	
Pro Arg Asp Val Asp Glu Met Gln Lys Leu Gln Glu Leu Leu Gly Asp			
35	40	45	
Glu Glu Gln Pro Leu Gly Val Ser Lys Lys Ser Leu Glu Gly Leu Leu			
50	55	60	
Ser Leu Arg Gln Pro Gln Glu Leu Ala Val Arg Leu Ala Gln Ser Leu			
65	70	75	80
Ser Ser Leu Arg Ala Arg Leu Ala Glu Leu Glu Leu Glu Arg Leu Asn			
85	90	95	
Arg Gly Ser Glu Ala Pro Gly Leu Ser Asn Ile Val Ala Arg Arg Ala			
100	105	110	
Gln Glu Lys Ala Glu Lys Leu Glu Leu Glu Ile Gln Lys Thr Glu Arg			
115	120	125	
Val Thr Arg Arg Leu Lys Ile Ile Ser Ser Leu Val Gly His Ile Ile			
130	135	140	
Arg Leu Arg Glu Lys Thr Leu Thr Glu Thr His Thr Ala Met Glu Ala			
145	150	155	160
Glu Val Gln Ser Leu Gln Glu Lys Ile Arg Val Asn Glu Glu Leu Ile			
165	170	175	
Arg Glu Arg Phe Val Ser Arg Val Asn Met Leu His Arg Tyr Trp Leu			
180	185	190	
Trp Arg Thr Leu Gln Glu Leu Gly Asp Gln Thr Val Gly Trp Thr Phe			
195	200	205	
Glu Glu Glu Leu Ala Arg Gly Pro Arg Tyr Arg Thr Leu Gly Val Gln			
210	215	220	
Asn Asn Ile Val Ser Glu Thr Leu Glu Gln Gln Leu Ser Trp Leu Leu			
225	230	235	240
Val Phe Ala Glu Lys Glu Lys Ile Phe Arg Glu His Val Arg Arg Leu			
245	250	255	
Glu Leu Leu Val Glu Glu Leu Thr Asp Ile Asn Asp Ala Leu Glu Glu			
260	265	270	
Ala Leu Thr Cys Arg Val Cys Gly Leu Leu Phe Glu Asp Pro Val Leu			
275	280	285	
Phe Trp Pro Cys Gly His Val Phe Cys Leu Val Cys Phe Asp Thr Leu			
290	295	300	
Ser Ile Ala Pro Ser Leu Phe Arg Cys Pro Thr Cys Gly Ser Met Gly			
305	310	315	320
Ser Glu Gly Tyr Val His Asn Leu Leu Ile Ala Glu Ser Val Ala Lys			
325	330	335	
Trp Met Phe Lys Asp Ala Gly Tyr Gly Asp Ile His Gly Ala Leu Ser			
340	345	350	
Leu Ile Arg Leu His Leu Ser Lys Phe Arg Lys Glu Val Ile Ser Ser			
355	360	365	
Arg Val Ala Gln Leu His Gln Gln Leu Thr Glu Ala Arg Gln Lys Glu			
370	375	380	
Thr Lys Val Glu Glu Leu Ser Gln Met Asp Ile Thr Tyr Arg Asp Phe			
385	390	395	400

<210> 28

<211> 515

<212> DNA

<213> T. cruzi

<400> 28

ccatggggcag	cagccatcat	catcatcatc	acagcagcgg	cctggtgccg	cgcggcagcc	60
atatgtgcat	tgctcttggc	atcgctcgcg	aggatgatga	ggcatggcga	attgccagcg	120
aggctgtcgc	tgcagaaaag	gagcctgtct	tcagcgggaa	caacggccct	ttttagatg	180
tctggtttgg	cgaacagaaa	ctctttggcc	tcgttcaacg	cgttgctcca	aacgacttta	240
ttcaggtcgc	ccaggagtgt	ggcgagaaga	gcgatgacgc	cgcagcgacg	ttgcggatgc	300

```

gtgtgacgca caacgtctct tttgtccttc acctctcgtc ggtgccgcat gcgatgctgc 360
aggcacgggg agcgcccgag gacaagtttg tgaacttcat gcaacttgtc gtggattacg 420
cttcgctgct gcggcgcggg atgaaggatg agtttcttg gctcgatccc gagtccgatg 480
cggagtacat acgcttcacg cccagtgaa agctt 515

```

<210> 29
 <211> 168
 <212> PRT
 <213> T. cruzi

<400> 29

Met	Gly	Ser	Ser	His	His	His	His	His	His	Ser	Ser	Gly	Leu	Val	Pro
1				5					10				15		
Arg	Gly	Ser	His	Met	Cys	Ile	Ala	Leu	Gly	Ile	Val	Ala	Glu	Asp	Asp
			20					25				30			
Glu	Ala	Trp	Arg	Ile	Ala	Ser	Glu	Ala	Val	Ala	Ala	Glu	Lys	Glu	Pro
			35				40				45				
Val	Phe	Ser	Gly	Asn	Asn	Gly	Pro	Phe	Val	Asp	Val	Trp	Phe	Gly	Glu
	50				55				60						
Gln	Lys	Leu	Phe	Gly	Leu	Val	Gln	Arg	Val	Ala	Pro	Asn	Asp	Phe	Ile
65				70					75					80	
Gln	Val	Ala	Gln	Glu	Cys	Gly	Glu	Lys	Ser	Asp	Asp	Ala	Ala	Ala	Thr
			85					90				95			
Leu	Arg	Met	Arg	Val	Thr	His	Asn	Val	Ser	Phe	Val	Leu	His	Leu	Ser
		100						105				110			
Ser	Val	Pro	His	Ala	Met	Leu	Gln	Ala	Arg	Gly	Ala	Pro	Glu	Asp	Lys
		115					120				125				
Phe	Val	Asn	Phe	Met	Gln	Leu	Val	Val	Asp	Tyr	Ala	Ser	Leu	Leu	Arg
	130				135					140					
Arg	Gly	Met	Lys	Asp	Glu	Phe	Leu	Gly	Val	Asp	Pro	Glu	Ser	Asp	Ala
145				150					155					160	
Glu	Tyr	Ile	Arg	Phe	Thr	Pro	Gln								
				165											

<210> 30
 <211> 526
 <212> DNA
 <213> T. cruzi

<400> 30

ccatgggcag	cagccatcat	catcatcatc	acagcagcgg	cctggtgccg	cgcggcagcc		60
atatgtgttc	ccaatacgga	cccttgagga	actcatcgca	gcgtggaaga	ccgatcgctg		120
tgtctgtgtg	caactcgtgt	cttcgcgtag	cggccacgca	gaaagtttct	aagccacctt		180
ctacagttca	cccccggaat	attggccggc	aagcgactga	ggactcgatg	accaatgaac		240
tcaaaggcct	tgctggagtc	taccagcacc	aacggagccc	gatgggggtc	gcagtggagc		300
tggtcttcaa	caccgctctt	cctgggaagg	ttcacttgga	attaatcggt	tctgttatgc		360
tcaaattcgt	ttatcaggtg	tgccagctgc	atcgctcggt	tacgcatact	acggcacgca		420
tactggtgcg	gacgagggcg	ccacagccaa	agcagtcctc	tcaatgccat	tttctcagca		480
accttaccca	cgatggaagg	acagctgatt	ggtcgatatg	aagctt			526

<210> 31
 <211> 172
 <212> PRT
 <213> T. cruzi

<400> 31

Met	Gly	Ser	Ser	His	His	His	His	His	Ser	Ser	Gly	Leu	Val	Pro
1				5				10				15		
Arg	Gly	Ser	His	Met	Cys	Ser	Gln	Tyr	Gly	Pro	Leu	Arg	Asn	Ser
			20				25				30			
Gln	Arg	Gly	Arg	Pro	Ile	Ala	Val	Ser	Val	Ala	Asn	Ser	Cys	Leu
														Arg

专利名称(译)	用于检测克氏锥虫感染的组合物和方法		
公开(公告)号	EP1945248A2	公开(公告)日	2008-07-23
申请号	EP2006836858	申请日	2006-11-02
[标]申请(专利权)人(译)	INBIOS INT		
申请(专利权)人(译)	inbios国际有限公司		
当前申请(专利权)人(译)	inbios国际有限公司		
[标]发明人	HOUGHTON RAYMOND L REED STEVEN G RAYCHAUDHURI SYAMAL		
发明人	HOUGHTON, RAYMOND L. REED, STEVEN G. RAYCHAUDHURI, SYAMAL		
IPC分类号	A61K39/00 A61K39/385 A61K39/002 C12P21/06 C12N15/00 C12N1/20 G01N33/53 C07H21/04 C07K14/44 A61K39/005 C12N15/62		
CPC分类号	A61K39/00 A61P31/12 A61P37/04 C07K14/44 C07K2319/40 G01N2333/44 Y02A50/414 Y10S435/975		
优先权	60/733119 2005-11-03 US		
其他公开文献	EP1945248A4		
外部链接	Espacenet		

摘要(译)

提供了包含克氏锥虫表位的融合多肽的组合物，以及这些组合物在诊断克氏锥虫感染和筛选血液供应中的用途。还提供了包含此类组合物的诊断试剂盒。