

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 February 2006 (23.02.2006)

PCT

(10) International Publication Number
WO 2006/020266 A2

(51) International Patent Classification:
A61K 39/395 (2006.01)

(74) Agent: FISH & NEAVE IP GROUP; ROPES & GRAY,
One International Place, Boston, MA 02110 (US).

(21) International Application Number:
PCT/US2005/025488

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 19 July 2005 (19.07.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/894,672 20 July 2004 (20.07.2004) US
10/996,316 23 November 2004 (23.11.2004) US
11/171,567 30 June 2005 (30.06.2005) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): ALEX-ION PHARMACEUTICALS, INC. [US/US]; 352 Knotter Drive, Cheshire, CT 06410 (US).

(72) Inventors; and


(75) Inventors/Applicants (for US only): BOWDISH, Katherine, S. [US/US]; 13773 Boquita Drive, Del Mar, CA 92014 (US). KRETZ-ROMMEL, Anke [CH/US]; 4535 Kensington Drive, San Diego, CA 92116 (US). MCWHIRTER, John [US/US]; 959 Wild Oak Road, Chula Vista, CA 91910 (US).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: POLYPEPTIDES AND ANTIBODIES DERIVED FROM CHRONIC LYMPHOCYTIC LEUKEMIA CELLS AND USES THEREOF



(57) Abstract: Small animal models for assessing immunomodulatory effects of compounds are provided.

WO 2006/020266 A2

**POLYPEPTIDES AND ANTIBODIES DERIVED FROM
CHRONIC LYMPHOCYTIC LEUKEMIA CELLS AND USES THEREOF**

5

RELATED APPLICATIONS

This application claims priority to U.S. Application Serial No. 11/171,567 filed June 30, 2005 which is a continuation in part of U.S. Application Serial No. 10/996,316 filed November 23, 2004 which is a continuation in part of U.S. Application Serial No. 10/894,672 filed July 20, 2004 which is a continuation in part of U.S. Application Serial No. 10/736,188 filed December 15, 2003 which is a continuation in part of U.S. Application Serial No. 10/379,151 filed on March 4, 2003 which, in turn, is a continuation in part of PCT/US01/47931 filed on December 10, 2001, which is an international application that claims priority to U. S. Provisional Application No. 60/254,113 filed December 8, 2000. The entire disclosures of the aforementioned U.S., international and provisional applications are incorporated herein by reference.

TECHNICAL FIELD

Cancer treatments using a therapy that provides a combination of two mechanisms are disclosed. More specifically, this disclosure relates to treating cancer using a therapy that: 1) interferes with the interaction between CD200 and its receptor to block immune suppression thereby promoting eradication of the cancer cells; and 2) directly kills the cancer cells either by a) complement-mediated or antibody-dependent cellular cytotoxicity or b) by targeting cells using a fusion molecule that includes a CD200-targeting portion.

25 BACKGROUND

Chronic Lymphocytic Leukemia (CLL) is a disease of the white blood cells and is the most common form of leukemia in the Western Hemisphere. CLL represents a diverse group of diseases relating to the growth of malignant lymphocytes that grow slowly but have an extended life span. CLL is classified in various categories that include, for example, B-cell chronic lymphocytic leukemia (B-CLL) of classical and mixed types, B-

cell and T-cell prolymphocytic leukemia, hairy cell leukemia, and large granular lymphocytic leukemia.

Of all the different types of CLL, B-CLL accounts for approximately 30 percent of all leukemias. Although it occurs more frequently in individuals over 50 years of age, it is increasingly seen in younger people. B-CLL is characterized by accumulation of B-lymphocytes that are morphologically normal but biologically immature, leading to a loss of function. Lymphocytes normally function to fight infection. In B-CLL, however, lymphocytes accumulate in the blood and bone marrow and cause swelling of the lymph nodes. The production of normal bone marrow and blood cells is reduced and patients often experience severe anemia as well as low platelet counts. This can pose the risk of life-threatening bleeding and the development of serious infections because of reduced numbers of white blood cells.

To further understand diseases such as leukemia it is important to have suitable cell lines that can be used as tools for research on their etiology, pathogenesis and biology. Examples of malignant human B-lymphoid cell lines include pre-B acute lymphoblastic leukemia (Reh), diffuse large cell lymphoma (WSU-DLCL2), and Waldenstrom's macroglobulinemia (WSU- WM). Unfortunately, many of the existing cell lines do not represent the clinically most common types of leukemia and lymphoma.

The use of Epstein Barr Virus (EBV) infection in vitro has resulted in some CLL derived cell lines, in particular B-CLL cells lines, that are representative of the malignant cells. The phenotype of these cell lines is different than that of the in vivo tumors and instead the features of B-CLL lines tend to be similar to those of lymphoblastoid cell lines. Attempts to immortalize B-CLL cells with the aid of EBV infection have had little success. The reasons for this are unclear but it is known that it is not due to a lack of EBV receptor expression, binding or uptake. Wells et al. found that B-CLL cells were arrested in the G1/S phase of the cell cycle and that transformation associated EBV DNA was not expressed. This suggests that the interaction of EBV with B-CLL cells is different from that with normal B cells. EBV-transformed CLL cell lines moreover appear to differentiate, possessing a morphology more similar to lymphoblastoid cell lines (LCL) immortalized by EBV.

An EBV-negative CLL cell line, WSU-CLL, has been established previously (Mohammad et al., (1996) *Leukemia* 10(1):130-7). However, no other such cell lines are known.

Various mechanisms play a role in the body's response to a disease state, including cancer and CLL. For example, CD4⁺ T helper cells play a crucial role in an effective immune response against various malignancies by providing stimulatory factors to effector cells. Cytotoxic T cells are believed to be the most effective cells to eliminate cancer cells, and T helper cells prime cytotoxic T cells by secreting Th1 cytokines such as IL-2 and IFN- γ . In various malignancies, T helper cells have been shown to have an altered phenotype compared to cells found in healthy individuals. One of the prominent altered features is decreased Th1 cytokine production and a shift to the production of Th2 cytokines. (See, e.g., Kiani, et al., *Haematologica* 88:754-761 (2003); Maggio, et al., *Ann Oncol* 13 Suppl 1:52-56 (2002); Ito, et al., *Cancer* 85:2359-2367 (1999); Podhorecka, et al., *Leuk Res* 26:657-660 (2002); Tatsumi, et al., *J Exp Med* 196:619-628 (2002); Agarwal, et al., *Immunol Invest* 32:17-30 (2003); Smyth, et al., *Ann Surg Oncol* 10:455-462 (2003); Contasta, et al., *Cancer Biother Radiopharm* 18:549-557 (2003); Lauerova, et al., *Neoplasma* 49:159-166(2002).) Reversing that cytokine shift to a Th1 profile has been demonstrated to augment anti-tumor effects of T cells. (See Winter, et al., *Immunology* 108:409-419 (2003); Inagawa, et al., *Anticancer Res* 18:3957-3964 (1998).)

Mechanisms underlying the capacity of tumor cells to drive the cytokine expression of T helper cells from Th1 to Th2 include the secretion of cytokines such as IL-10 or TGF- β as well as the expression of surface molecules interacting with cells of the immune system. OX-2/CD200, a molecule expressed on the surface of dendritic cells which possesses a high degree of homology to molecules of the immunoglobulin gene family, has been implicated in immune suppression (Gorczyński et al., *Transplantation* 65:1106-1114 (1998)) and evidence that OX-2/CD200-expressing cells can inhibit the stimulation of Th1 cytokine production has been provided. Gorczyński et al. demonstrated in a mouse model that infusion of OX-2/CD200 Fc suppresses the rejection of tumor cells in an animal model using leukaemic tumor cells (*Clin Exp Immunol* 126:220-229 (2001)).

Improved methods for treating individuals suffering from cancer or CLL are desirable, especially to the extent they can enhance the activity of T cells.

30

SUMMARY

In one embodiment a CLL cell line of malignant origin is provided that is not established by immortalisation with EBV. The cell line was derived from primary CLL cells and is deposited under ATCC accession no. PTA-3920. In a preferred embodiment,

the cell line is CLL-AAT. CLL-AAT is a B-CLL cell line, derived from a B-CLL primary cell.

In a further aspect, the CLL-AAT cell line is used to generate monoclonal antibodies useful in the diagnosis and/or treatment of CLL. Antibodies may be generated by using the cells as disclosed herein as immunogens, thus raising an immune response in animals from which monoclonal antibodies may be isolated. The sequence of such antibodies may be determined and the antibodies or variants thereof produced by recombinant techniques. In this aspect, "variants" includes chimeric, CDR-grafted, humanized and fully human antibodies based on the sequence of the monoclonal antibodies.

Moreover, antibodies derived from recombinant libraries ("phage antibodies") may be selected using the cells described herein, or polypeptides derived therefrom, as bait to isolate the antibodies on the basis of target specificity.

In a still further aspect, antibodies may be generated by panning antibody libraries using primary CLL cells, or antigens derived therefrom, and further screened and/or characterized using a CLL cell line, such as, for example, the CLL cell line described herein. Accordingly, a method for characterizing an antibody specific for CLL is provided, which includes assessing the binding of the antibody to a CLL cell line.

In a further aspect, there is provided a method for identifying proteins uniquely expressed in CLL cells employing the CLL-AAT cell line, by methods well known to those skilled in the art, such as by immunoprecipitation followed by mass spectroscopy analyses. Such proteins may be uniquely expressed in the CLL-AAT cell line, or in primary cells derived from CLL patients.

Small molecule libraries (many available commercially) may be screened using the CLL-AAT cell line in a cell-based assay to identify agents capable of modulating the growth characteristics of the cells. For example, the agents may be identified which modulate apoptosis in the CLL-AAT cell line, or which inhibit growth and/or proliferation thereof. Such agents are candidates for the development of therapeutic compounds.

Nucleic acids isolated from CLL-AAT cell lines may be used in subtractive hybridization experiments to identify CLL-specific genes or in micro array analyses (e.g., gene chip experiments). Genes whose transcription is modulated in CLL cells may be identified. Polypeptide or nucleic acid gene products identified in this manner are useful as leads for the development of antibody or small molecule therapies for CLL.

In a preferred aspect, the CLL-AAT cell line may be used to identify internalizing antibodies, which bind to cell surface components which are internalized by the cell. Such antibodies are candidates for therapeutic use. In particular, single-chain antibodies, which remain stable in the cytoplasm and which retain intracellular binding activity, may be screened in this manner.

In yet another aspect, a therapeutic treatment is described in which a patient is screened for the presence of a polypeptide that is upregulated by a malignant cancer cell and an antibody that interferes with the metabolic pathway of the upregulated polypeptide is administered to the patient.

The present disclosure further is directed to methods wherein a determination is made as to whether OX-2/CD200 is upregulated in a subject and, if so, administering to the subject a therapy that enhances immune response. Upregulation of OX2/CD200 can be determined by measuring OX2/CD200 levels directly, or by monitoring the level of any marker that correlates with OX2/CD200. Suitable immunomodulatory therapies include the administration of agents that block negative regulation of T cells or antigen presenting cells, administration of agents that enhance positive co-stimulation of T cells, cancer vaccines, general adjuvants stimulating the immune system or treatment with cytokines such as IL-2, GM-CSF and IFN-gamma. In particularly useful embodiments, the therapy that enhances immune response includes the administration of a polypeptide that binds to OX-2/CD200, optionally in combination with one or more other immunomodulatory therapies. In another embodiment, the polypeptide binds to an OX-2/CD200 receptor.

In another aspect, methods in accordance with this disclosure are used to treat a disease state in which OX-2/CD200 is upregulated in a subject by administering a polypeptide that binds to OX-2/CD200 or an OX-2/CD200 receptor to the subject afflicted with the disease state. In one embodiment, the disease state treated by these methods includes cancer, specifically, in other embodiments, CLL.

In a particularly useful embodiment, a cancer therapy in accordance with this disclosure includes i) administering an antibody that interferes with the interaction between CD200 and its receptor to block immune suppression, thereby promoting eradication of the cancer cells; and ii) administering a fusion molecule that includes a CD200-targeting portion to directly kill cancer cells. Alternatively, the antibody directly kills the cancer cells through complement-mediated or antibody-dependent cellular cytotoxicity.

In another embodiment in accordance with the present disclosure, methods are provided for monitoring the progress of a therapeutic treatment. The method involves administering an immunomodulatory therapy and determining OX-2/CD200 levels in a subject at least twice to determine the effectiveness of the therapy.

5 In another aspect, the present disclosure provides methods for assessing the immunomodulatory effect of molecules expressed by cancer cells. In these methods a molecule that is expressed or upregulated by a cancer cell is identified (e.g., experimentally or from a database). Cancer cells, lymphocytes and the molecule that is expressed or upregulated by a cancer cell are administered to a subject and the rate of
10 growth of the cancer cells is monitored. The number of lymphocytes administered is predetermined to be either a) sufficient to slow the growth of the cancer cells or b) insufficient to slow the growth of cancer cells. The molecule that is expressed or upregulated by a cancer cell can be administered as the molecule or the active portion of the molecule itself, or by administering cells that produce or express the molecule or
15 portions thereof naturally, or by administering cells that have been engineered to produce or express the molecule or portions thereof. If any change in the growth rate of the cancer cells is observed compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the molecule is deemed to have an immunomodulatory effect. For example, if the number of lymphocytes administered is
20 sufficient to slow the growth of the cancer cells and the rate of growth of the tumor cells observed is higher compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the molecule that is expressed or upregulated by a cancer cell is considered immunosuppressive. As another example, if the number of lymphocytes administered is insufficient to slow the growth of the cancer cells and the
25 rate of growth of the tumor cells observed is lowered compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the compound is considered immune enhancing. Once the immunomodulatory effect of the molecule is established, compounds that either enhance or inhibit the activity of the molecule can be identified in accordance with embodiments described herein. The enhancing or inhibiting
30 effect can be the result of direct interaction with the molecule expressed or upregulated by the cancer cell or may be the result of an interaction with other molecules in the metabolic pathway of the compound expressed or upregulated by the cancer cell.

In another aspect, the present disclosure provides methods for assessing the immunomodulatory effect of a compound. In these methods cancer cells, lymphocytes

and the compound to be assessed are administered to a subject and the rate of growth of the cancer cells is monitored. The number of lymphocytes administered is predetermined to be either a) sufficient to slow the growth of the cancer cells or b) insufficient to slow the growth of cancer cells. The compound to be assessed can be administered as the
5 compound itself, or by administering cells that produce the compound naturally, or by administering cells that have been engineered to produce the compound. If any change in the growth rate of the cancer cells is observed compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the compound is deemed to have an immunomodulatory effect. For example, if the number of lymphocytes
10 administered is sufficient to slow the growth of the cancer cells and the rate of growth of the tumor cells observed is higher compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the compound is considered immunosuppressive. As another example, if the number of lymphocytes administered is insufficient to slow the growth of the cancer cells and the rate of growth of the tumor
15 cells observed is lowered compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the compound is considered immune enhancing.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 schematically illustrates typical steps involved in cell surface panning of
20 antibody libraries by magnetically-activated cell sorting (MACS).

Fig. 2 is a graph showing the results of whole cell ELISA demonstrating binding of selected scFv clones to primary B-CLL cells and absence of binding to normal human PBMC. The designation 2°+3° in this and other figures refers to negative control wells stained with Mouse Anti-HA and detecting antimouse antibodies alone. The designation
25 RSC-S Library in this and other figures refers to soluble antibodies prepared from original rabbit scFv unpanned library. The designation R3/RSC-S Pool in this and other figures refers to soluble antibodies prepared from the entire pool of scFv antibodies from round 3 of panning. Anti-CD5 antibody was used as a positive control to verify that equal numbers of B-CLL and PBMC cells were plated in each well.

30 Figs. 3a and 3b show the results of whole cell ELISA comparing binding of selected scFv antibodies to primary B-CLL cells and normal primary human B cells. Anti-CD19 antibody was used as a positive control to verify that equal numbers of B-CLL and normal B cells were plated in each well. Other controls were as described in the legend to Fig. 2.

Figs. 4a and 4b show the results of whole cell ELISA used to determine if scFv clones bind to patient-specific (i.e. idiotype) or blood type-specific (i.e. HLA) antigens. Each clone was tested for binding to PBMC isolated from 3 different B-CLL patients. Clones that bound to only one patient sample were considered to be patient or blood type-specific.

Figs. 5a and 5b show the results of whole cell ELISA comparing binding of scFv clones to primary B-CLL cells and three human leukemic cell lines. Ramos is a mature B cell line derived from a Burkitt's lymphoma. RL is a mature B cell line derived from a non-Hodgkin's lymphoma. TF-1 is an erythroblastoid cell line derived from an erythroleukemia.

Figs. 6a, 6b and 6c show the results of whole cell ELISA comparing binding of scFv clones to primary B-CLL cells and CLL-AAT, a cell line derived from a B-CLL patient. TF-1 cells were included as a negative control.

Fig. 7 shows the binding specificity of scFv antibodies in accordance with this disclosure as analyzed by 3-color flow cytometry. In normal peripheral blood mononuclear cells, the antigen recognized by scFv-9 is moderately expressed on B lymphocytes and weakly expressed on a subpopulation of T lymphocytes. PBMC from a normal donor were analyzed by 3-color flow cytometry using anti-CD5-FITC, anti-CD19-PerCP, and scFv-9/Anti-HA-biotin/streptavidin-PE.

Figs. 8a, 8b and 8c show the expression levels of antigens recognized by scFv antibodies in accordance with this disclosure. The antigens recognized by scFv-3 and scFv-9 are overexpressed on the primary CLL tumor from which the CLL-AAT cell line was derived. Primary PBMC from the CLL patient used to establish the CLL-AAT cell line or PBMC from a normal donor were stained with scFv antibody and analyzed by flow cytometry. ScFv-3 and scFv-9 stain the CLL cells more brightly than the normal PBMC as measured by the mean fluorescent intensities.

Figs. 9A – 9C provide a summary of CDR sequences and binding specificities of selected scFv antibodies.

Fig. 10 is Table 2 which shows a summary of flow cytometry results comparing expression levels of scFv antigens on primary CLL cells vs. normal PBMC as described in Figs 8a-8c.

Fig. 11 is a Table showing a summary of flow cytometry results comparing expression levels of scFv-9 antigen with the percentage of CD38⁺ cells in peripheral blood mononuclear cells isolated from ten CLL patients.

Fig. 12 shows the identification of scFv antigens by immunoprecipitation and mass spectrometry. CLL-AAT cells were labeled with a solution of 0.5mg/ml sulfo-NHS-LC-biotin (Pierce) in PBS, pH8.0 for 30'. After extensive washing with PBS to remove unreacted biotin, the cells were disrupted by nitrogen cavitation and the
5 microsomal fraction was isolated by differential centrifugation. The microsomal fraction was resuspended in NP40 Lysis Buffer and extensively precleared with normal rabbit serum and protein A Sepharose. Antigens were immunoprecipitated with HA-tagged scFv antibodies coupled to Rat Anti-HA agarose beads (Roche). Following immunoprecipitation, antigens were separated by SDS-PAGE and detected by Western
10 blot using streptavidin-alkaline phosphatase (AP) or by Coomassie G-250 staining. ScFv-7, an antibody which doesn't bind to CLL-AAT cells, was used as a negative control. Antigen bands were excised from the Coomassie-stained gel and identified by mass spectrometry (MS). MALDI-MS was performed at the Proteomics Core Facility of The Scripps Research Institute (La Jolla, CA). μ LC/MS/MS was performed at the Harvard
15 Microchemistry Facility (Cambridge, MA).

Fig. 13 shows that three scFv antibodies bind specifically to 293-EBNA cells transiently transfected with a human OX-2/CD200 cDNA clone. An OX-2/CD200 cDNA was cloned from CLL cells by RT-PCR and inserted into the mammalian expression vector pCEP4 (Invitrogen). PCEP4-CD200 plasmid or the corresponding empty vector
20 pCEP4 was transfected into 293-EBNA cells using Polyfect reagent (QIAGEN). Two days after transfection, the cells were analyzed for binding to scFv antibodies by flow cytometry.

Fig. 14 shows that the presence of OX-2/CD200 transfected cells resulted in down-regulation of Th1 cytokines such as IL-2 and IFN- γ . Addition of the anti-OX-
25 2/CD200 antibody at 30 μ g/ml fully restored the Th1 response.

Fig. 15 shows that the presence of CLL cells in a mixed lymphocyte reaction resulted in down-regulation of the Th1 response for IL-2.

Fig. 16 shows that the presence of CLL cells in a mixed lymphocyte reaction resulted in down-regulation of the Th1 response for IFN- γ .

30 Figs. 17A and B show the mean +/-SD of tumor volumes for all groups of NOD/SCID mice were injected subcutaneously with 4×10^6 RAJI cells either in the presence or absence of human PBL cells.

Fig. 18 shows the results of statistical analyses performed using 2 parametric tests (Student's t-test and Welch's test) and one non-parametric test (the Wilcox test).

Fig. 19A shows ELISA results of representative IgG1 kappa clones after round 3 panning on CD200-Fc captured on goat anti-mouse IgG Fc antibody.

Fig. 19B shows ELISA results of representative IgG2a kappa clones after round 3 panning on CD200-Fc captured on goat anti-mouse IgG Fc antibody.

5 Fig. 19C shows ELISA results of representative IgG1 kappa clones after round 3 panning on CD200-Fc directly coated on microtiter wells.

Fig. 19D shows ELISA results of representative IgG2a kappa clones after round 3 panning on CD200-Fc directly coated on microtiter wells.

10 Fig. 20A shows flow cytometry results of representative IgG1 clones selected on CD200-Fc captured with goat anti-mouse IgG Fc.

Fig. 20B shows flow cytometry results of representative IgG2a clones selected on CD200-Fc captured with goat anti-mouse IgG Fc.

Fig. 20C shows flow cytometry results of representative IgG1 clones selected on directly coated CD200-Fc.

15 Fig. 20D shows flow cytometry results of representative IgG2a clones selected on directly coated CD200-Fc.

Fig. 21A shows deduced amino acid sequence of heavy chain complementarity regions of CD200-specific clones.

20 Fig. 21B shows deduced amino acid sequence of heavy chain complementarity regions of CD200-specific clones.

Fig. 22 shows ability of selected clones to block the interaction of CD200 with its receptor (CD200R) in a fluorescent bead assay.

Fig. 23 shows deduced amino acid sequences of selected CD200 Fabs for chimerization.

25 Fig. 24 shows ELISA results of chimeric IgG obtained from the culture supernatant of a small-scale transient transfection.

Fig. 25 shows bead inhibition assay results on purified IgG showing that all antibodies directed against CD200 blocked the receptor ligand interaction very well.

30 Figs. 26A and 26B show that the presence of CLL cells completely abrogated IFN-gamma and most of IL-2 production observed in the mixed lymphocyte reaction but that the presence of any of the antibodies allowed for production of these Th1 cytokines.

Fig. 26C shows that IL-10 production was downregulated in the presence of the antibodies.

Fig. 27 shows the ability to kill CD200 expressing tumor cells in an antibody-dependent cell-mediated cytotoxicity assay (ADCC). All of the mouse chimeric CD200 antibodies produced similar levels of lysis when cultured with CD200 positive cells.

Fig. 28 shows a representative example of 10 experiments using different PBL donors compared to a group that received tumor cells only, analyzed by a 2-tailed unpaired Student's t-test. Significant differences were observed in the groups that received 5 or 10 million PBLs, but not in the group that received 1 million PBLs from Day 32 on.

Fig. 29 shows a representative example of 10 experiments using different PBL donors compared to a group that received tumor cells only, analyzed by 2-tailed unpaired Student's t-test. Significant differences were observed in the groups that received 10 million PBLs for both donors, but not in the group that received 2 million PBLs from Day 8 on.

Fig. 30(a) shows RAJI cells infected with CD200 expressing tumor cells transduced from a lentivirus vector appeared to grow somewhat more slowly than its parental RAJI cells. The growth difference between the transduced and parental cells did not reach statistical significance.

Fig. 30(b) shows the presence of PBLs reduced tumor growth in the RAJI cells transduced with the reversed CD200 (non functional CD200) by up to 84% when 5 or 10×10^6 PBLs were injected indicating that this particular donor rejects RAJI tumor cells very strongly.

Fig. 30(c) shows results indicating that CD200 expression on tumor cells does indeed prevent the immune system from slowing tumor growth. Also this study demonstrates the usefulness of the RAJI/PBL model to assess immunosuppressive compounds or molecules.

Figs. 31(a)-(d) show whether the effects seen in the RAJI/PBL model can also be observed with other tumor cell models.

Fig. 31(a) shows Namalwa tumor cells resulted in rapid tumor growth with no significant difference between transduced and parental cells.

Fig. 31(b) shows the presence of PBLs slowed tumor growth by about 50%. The 2-tailed Student's t-test results showed that the differences between the PBL treated group versus groups that received only tumor cells were statistically significant.

Figs. 31(c) and (d) show tumor growth in the groups that received CD200 expressing Namalwa cells and PBLs was similar to the tumor growth in the group that

received Namalwa cells. These data confirm that CD200 expression on tumor cells prevents slowing of tumor growth by the human immune system.

Fig. 32 shows results demonstrating that the RAJI PBL model is an efficient way to assess efficacy of immune-enhancing compounds.

5 Figure 33 shows that CD200 expression on the tumor cells prevented the immune cells from reducing tumor growth.

Figure 34 shows that CD200 expression on the tumor cells prevented the immune cells from reducing tumor growth.

10 DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

In accordance with the present disclosure, methods are provided for determining whether OX-2/CD200 is upregulated in a subject and, if so, administering to the subject a therapy that enhances immune response. Illustrative examples of suitable immunomodulatory therapies include the administration of agents that block negative
15 regulation of T cells or antigen presenting cells (e.g., anti-CTLA4 antibodies, anti-PD-L1 antibodies, anti-PDL-2 antibodies, anti-PD-1 antibodies and the like) or the administration of agents that enhance positive co-stimulation of T cells (e.g., anti-CD40 antibodies or anti 4-1BB antibodies) or administration of agents that increase NK cell number or T-cell activity (e.g., anti-CD200 antibodies alone or in combination with
20 inhibitors such as IMiDs, thalidomide, or thalidomide analogs). Furthermore, immunomodulatory therapy could be cancer vaccines such as dendritic cells loaded with tumor cells, tumor RNA or tumor DNA, tumor protein or tumor peptides, patient derived heat-shocked proteins (hsp's) or general adjuvants stimulating the immune system at various levels such as CpG, Luivac, Biostim, Ribominy, Imudon, Bronchovaxom or any
25 other compound activating receptors of the innate immune system (e.g., toll like receptors). Also, immunomodulatory therapy could include treatment with cytokines such as IL-2, GM-CSF and IFN-gamma.

In particularly useful embodiments, the therapy that enhances immune response is the administration of a polypeptide that binds to OX-2/CD200, alone or in combination
30 with one of the previously mentioned immunomodulatory therapies. In general, the polypeptides utilized in the present disclosure can be constructed using different techniques which are known to those skilled in the art. In one embodiment, the polypeptides are obtained by chemical synthesis. In other embodiments, the polypeptides

are antibodies or constructed from a fragment or several fragments of one or more antibodies.

Preferably, the polypeptides utilized in the methods of the present disclosure are obtained from a CLL cell line. "CLL", as used herein, refers to chronic lymphocytic leukemia involving any lymphocyte including, but not limited to, various developmental stages of B cells and T cells including, but not limited to, B cell CLL ("B-CLL"). B-CLL, as used herein, refers to leukemia with a mature B cell phenotype which is CD5⁺, CD23⁺, CD20^{dim+}, sIg^{dim+} and arrested in G0/G1 of the cell cycle. In a further aspect, the CLL cell line is used to generate polypeptides, including antibodies, useful in the diagnosis and/or treatment of a disease state in which OX-2/CD200 is upregulated, including cancer and CLL.

As used herein, the term "antibodies" refers to complete antibodies or antibody fragments capable of binding to a selected target. Included are Fab, Fv, scFv, Fab' and F(ab')₂, monoclonal and polyclonal antibodies, engineered antibodies (including chimeric, CDR-grafted and humanized, fully human antibodies, and artificially selected antibodies), and synthetic or semi-synthetic antibodies produced using phage display or alternative techniques. Small fragments, such as Fv and scFv, possess advantageous properties for diagnostic and therapeutic applications on account of their small size and consequent superior tissue distribution.

Antibodies may be generated by using the cells as disclosed herein as immunogens, thus raising an immune response in animals from which monoclonal antibodies may be isolated. The sequence of such antibodies may be determined and the antibodies or variants thereof produced by recombinant techniques. In this aspect, "variants" includes chimeric, CDR-grafted, humanized and fully human antibodies based on the sequence of the monoclonal antibodies, as well as polypeptides capable of binding to OX-2/CD200.

Moreover, antibodies derived from recombinant libraries ("phage antibodies") may be selected using the cells described herein, or polypeptides derived therefrom, as bait to isolate the antibodies or polypeptides on the basis of target specificity.

In a still further aspect, antibodies or polypeptides may be generated by panning antibody libraries using primary CLL cells, or antigens derived therefrom, and further screened and/or characterized using a CLL cell line, such as, for example, the CLL cell line described herein. Accordingly, a method for characterizing an antibody or

polypeptide specific for CLL is provided, which includes assessing the binding of the antibody or polypeptide to a CLL cell line.

Preparation of Cell Lines

5 Cell lines may be produced according to established methodologies known to those skilled in the art. In general, cell lines are produced by culturing primary cells derived from a patient until immortalized cells are spontaneously generated in culture. These cells are then isolated and further cultured to produce clonal cell populations or cells exhibiting resistance to apoptosis.

10 For example, CLL cells may be isolated from peripheral blood drawn from a patient suffering from CLL. The cells may be washed, and optionally immunotyped in order to determine the type(s) of cells present. Subsequently, the cells may be cultured in a medium, such as a medium containing IL-4. Advantageously, all or part of the medium is replaced one or more times during the culture process. Cell lines may be isolated
15 thereby, and will be identified by increased growth in culture.

In one embodiment a CLL cell line of malignant origin is provided that is not established by immortalization with EBV. "Malignant origin" refers to the derivation of the cell line from malignant CLL primary cells, as opposed to non-proliferating cells which are transformed, for example, with EBV. Cell lines useful according to this
20 disclosure may be themselves malignant in phenotype, or not. A CLL cell having a "malignant" phenotype encompasses cell growth unattached from substrate media characterized by repeated cycles of cell growth and exhibits resistance to apoptosis. The cell line, which was derived from primary CLL cells, is deposited under ATCC accession no. PTA-3920. In a preferred embodiment, the cell line is CLL-AAT. CLL-AAT is a B-
25 CLL cell line, derived from a B-CLL primary cell.

In one embodiment, proteins uniquely expressed in CLL cells are identified employing the CLL-AAT cell line by methods well known to those skilled in the art, such as by immunoprecipitation followed by mass spectroscopy analyses. Such proteins may be uniquely expressed in the CLL-AAT cell line, or in primary cells derived from CLL
30 patients.

Small molecule libraries (many available commercially) may be screened using the CLL-AAT cell line in a cell-based assay to identify agents capable of modulating the growth characteristics of the cells. For example, the agents may be identified which modulate apoptosis in the CLL-AAT cell line, or which inhibit growth and/or

proliferation thereof. Such agents are candidates for the development of therapeutic compounds.

Nucleic acids isolated from CLL-AAT cell lines may be used in subtractive hybridization experiments to identify CLL-specific genes or in micro array analyses (e.g.,
5 gene chip experiments). Genes whose transcription is modulated in CLL cells may be identified. Polypeptide or nucleic acid gene products identified in this manner are useful as leads for the development of antibody or small molecule therapies for CLL.

In one embodiment, the CLL-AAT cell line may be used to identify internalizing antibodies, which bind to cell surface components and are then internalized by the cell.
10 Such antibodies are candidates for therapeutic use. In particular, single-chain antibodies, which remain stable in the cytoplasm and which retain intracellular binding activity, may be screened in this manner.

Preparation of Monoclonal Antibodies

15 Recombinant DNA technology may be used to improve the antibodies produced in accordance with this disclosure. Thus, chimeric antibodies may be constructed in order to decrease the immunogenicity thereof in diagnostic or therapeutic applications. Moreover, immunogenicity may be minimized by humanizing the antibodies by CDR grafting and, optionally, framework modification. See, U.S. Patent No. 5,225,539, the
20 contents of which are incorporated herein by reference.

Antibodies may be obtained from animal serum, or, in the case of monoclonal antibodies or fragments thereof produced in cell culture. Recombinant DNA technology may be used to produce the antibodies according to established procedure, in bacterial or preferably mammalian cell culture. The selected cell culture system preferably secretes
25 the antibody product.

In another embodiment, a process for the production of an antibody disclosed herein includes culturing a host, e.g. E. coli or a mammalian cell, which has been transformed with a hybrid vector. The vector includes one or more expression cassettes containing a promoter operably linked to a first DNA sequence encoding a signal peptide
30 linked in the proper reading frame to a second DNA sequence encoding the antibody protein. The antibody protein is then collected and isolated. Optionally, the expression cassette may include a promoter operably linked to polycistronic, for example bicistronic, DNA sequences encoding antibody proteins each individually operably linked to a signal peptide in the proper reading frame.

Multiplication of hybridoma cells or mammalian host cells in vitro is carried out in suitable culture media, which include the customary standard culture media (such as, for example Dulbecco's Modified Eagle Medium (DMEM) or RPMI 1640 medium), optionally replenished by a mammalian serum (e.g. fetal calf serum), or trace elements and growth sustaining supplements (e.g. feeder cells such as normal mouse peritoneal exudate cells, spleen cells, bone marrow macrophages, 2-aminoethanol, insulin, transferrin, low density lipoprotein, oleic acid, or the like). Multiplication of host cells which are bacterial cells or yeast cells is likewise carried out in suitable culture media known in the art. For example, for bacteria suitable culture media include medium LE, NZCYM, NZYM, NZM, Terrific Broth, SOB, SOC, 2 x YT, or M9 Minimal Medium. For yeast, suitable culture media include medium YPD, YEPD, Minimal Medium, or Complete Minimal Dropout Medium.

In vitro production provides relatively pure antibody preparations and allows scale-up to give large amounts of the desired antibodies. Techniques for bacterial cell, yeast, plant, or mammalian cell cultivation are known in the art and include homogeneous suspension culture (e.g. in an airlift reactor or in a continuous stirrer reactor), and immobilized or entrapped cell culture (e.g. in hollow fibres, microcapsules, on agarose microbeads or ceramic cartridges).

Large quantities of the desired antibodies can also be obtained by multiplying mammalian cells in vivo. For this purpose, hybridoma cells producing the desired antibodies are injected into histocompatible mammals to cause growth of antibody-producing tumors. Optionally, the animals are primed with a hydrocarbon, especially mineral oils such as pristane (tetramethyl-pentadecane), prior to the injection. After one to three weeks, the antibodies are isolated from the body fluids of those mammals. For example, hybridoma cells obtained by fusion of suitable myeloma cells with antibody-producing spleen cells from Balb/c mice, or transfected cells derived from hybridoma cell line Sp2/0 that produce the desired antibodies are injected intraperitoneally into Balb/c mice optionally pre-treated with pristane. After one to two weeks, ascitic fluid is taken from the animals.

The foregoing, and other, techniques are discussed in, for example, Kohler and Milstein, (1975) *Nature* 256:495-497; U.S. Patent No. 4,376,110; Harlow and Lane, *Antibodies: a Laboratory Manual*, (1988) Cold Spring Harbor, the disclosures of which are all incorporated herein by reference. Techniques for the preparation of recombinant antibody molecules are described in the above references and also in, for example

WO97/08320; U.S. Patent No. 5,427,908; U.S. Patent No. 5,508,717; Smith, 1985, Science, Vol. 225, pp 1315-1317; Parmley and Smith, 1988, Gene 73, pp 305-318; De La Cruz et al., 1988, Journal of Biological Chemistry, 263 pp 4318-4322; U.S. Patent No. 5,403,484; U.S. Patent No. 5223409; WO88/06630; WO92/15679; U.S. Patent No. 5780279; U.S. Patent No. 5571698; U.S. Patent No. 6040136; Davis et al., 1999, Cancer Metastasis Rev., 18(4):421-5; Taylor, et al., Nucleic Acids Research 20 (1992): 6287-6295; Tomizuka et al., Proc. Natl. Academy of Sciences USA 97(2) (2000): 722-727. The contents of all these references are incorporated herein by reference.

The cell culture supernatants are screened for the desired antibodies, preferentially by immunofluorescent staining of CLL cells, by immunoblotting, by an enzyme immunoassay, e.g. a sandwich assay or a dot-assay, or a radioimmunoassay.

For isolation of the antibodies, the immunoglobulins in the culture supernatants or in the ascitic fluid may be concentrated, e.g. by precipitation with ammonium sulfate, dialysis against hygroscopic material such as polyethylene glycol, filtration through selective membranes, or the like. If necessary and/or desired, the antibodies are purified by the customary chromatography methods, for example gel filtration, ion-exchange chromatography, chromatography over DEAE-cellulose and/or (immuno-) affinity chromatography, e.g. affinity chromatography with one or more surface polypeptides derived from a CLL cell line according to this disclosure, or with Protein-A or G.

Another embodiment provides a process for the preparation of a bacterial cell line secreting antibodies directed against the cell line characterized in that a suitable mammal, for example a rabbit, is immunized with pooled CLL patient samples. A phage display library produced from the immunized rabbit is constructed and panned for the desired antibodies in accordance with methods well known in the art (such as, for example, the methods disclosed in the various references incorporated herein by reference).

Hybridoma cells secreting the monoclonal antibodies are also contemplated. The preferred hybridoma cells are genetically stable, secrete monoclonal antibodies described herein of the desired specificity and can be activated from deep-frozen cultures by thawing and recloning.

In another embodiment, a process is provided for the preparation of a hybridoma cell line secreting monoclonal antibodies directed to the CLL cell line is described herein. In that process, a suitable mammal, for example a Balb/c mouse, is immunized with one or more polypeptides or antigenic fragments thereof derived from a cell described in this disclosure, the cell line itself, or an antigenic carrier containing a purified polypeptide as

described. Antibody-producing cells of the immunized mammal are grown briefly in culture or fused with cells of a suitable myeloma cell line. The hybrid cells obtained in the fusion are cloned, and cell clones secreting the desired antibodies are selected. For example, spleen cells of Balb/c mice immunized with the present cell line are fused with
5 cells of the myeloma cell line PAI or the myeloma cell line Sp2/0-Ag 14, the obtained hybrid cells are screened for secretion of the desired antibodies, and positive hybridoma cells are cloned.

Preferred is a process for the preparation of a hybridoma cell line, characterized in that Balb/c mice are immunized by injecting subcutaneously and/or intraperitoneally
10 between 10^6 and 10^7 cells of a cell line in accordance with this disclosure several times, e.g. four to six times, over several months, e.g. between two and four months. Spleen cells from the immunized mice are taken two to four days after the last injection and fused with cells of the myeloma cell line PAI in the presence of a fusion promoter, preferably polyethylene glycol. Preferably, the myeloma cells are fused with a three- to
15 twenty-fold excess of spleen cells from the immunized mice in a solution containing about 30% to about 50% polyethylene glycol of a molecular weight around 4000. After the fusion, the cells are expanded in suitable culture media as described hereinbefore, supplemented with a selection medium, for example HAT medium, at regular intervals in order to prevent normal myeloma cells from overgrowing the desired hybridoma cells.

20 In a further embodiment, recombinant DNA comprising an insert coding for a heavy chain variable domain and/or for a light chain variable domain of antibodies directed to the cell line described hereinbefore are produced. The term DNA includes coding single stranded DNAs, double stranded DNAs consisting of said coding DNAs and of complementary DNAs thereto, or these complementary (single stranded) DNAs
25 themselves.

Furthermore, DNA encoding a heavy chain variable domain and/or a light chain variable domain of antibodies directed to the cell line disclosed herein can be enzymatically or chemically synthesized DNA having the authentic DNA sequence coding for a heavy chain variable domain and/or for the light chain variable domain, or a
30 mutant thereof. A mutant of the authentic DNA is a DNA encoding a heavy chain variable domain and/or a light chain variable domain of the above-mentioned antibodies in which one or more amino acids are deleted or exchanged with one or more other amino acids. Preferably said modification(s) are outside the CDRs of the heavy chain variable domain and/or of the light chain variable domain of the antibody in humanization and

expression optimization applications. The term mutant DNA also embraces silent mutants wherein one or more nucleotides are replaced by other nucleotides with the new codons coding for the same amino acid(s). The term mutant sequence also includes a degenerate sequence. Degenerate sequences are degenerate within the meaning of the genetic code in that an unlimited number of nucleotides are replaced by other nucleotides without resulting in a change of the amino acid sequence originally encoded. Such degenerate sequences may be useful due to their different restriction sites and/or frequency of particular codons which are preferred by the specific host, particularly *E. coli*, to obtain an optimal expression of the heavy chain murine variable domain and/or a light chain murine variable domain.

The term mutant is intended to include a DNA mutant obtained by *in vitro* mutagenesis of the authentic DNA according to methods known in the art.

For the assembly of complete tetrameric immunoglobulin molecules and the expression of chimeric antibodies, the recombinant DNA inserts coding for heavy and light chain variable domains are fused with the corresponding DNAs coding for heavy and light chain constant domains, then transferred into appropriate host cells, for example after incorporation into hybrid vectors.

Recombinant DNAs including an insert coding for a heavy chain murine variable domain of an antibody directed to the cell line disclosed herein fused to a human constant domain μ , for example $\gamma 1$, $\gamma 2$, $\gamma 3$ or $\gamma 4$, preferably $\gamma 1$ or $\gamma 4$ are also provided.

Recombinant DNAs including an insert coding for a light chain murine variable domain of an antibody directed to the cell line disclosed herein fused to a human constant domain κ or λ , preferably κ are also provided

Another embodiment pertains to recombinant DNAs coding for a recombinant polypeptide wherein the heavy chain variable domain and the light chain variable domain are linked by way of a spacer group, optionally comprising a signal sequence facilitating the processing of the antibody in the host cell and/or a DNA coding for a peptide facilitating the purification of the antibody and/or a cleavage site and/or a peptide spacer and/or an effector molecule.

The DNA coding for an effector molecule is intended to be a DNA coding for the effector molecules useful in diagnostic or therapeutic applications. Thus, effector molecules which are toxins or enzymes, especially enzymes capable of catalyzing the activation of prodrugs, are particularly indicated. The DNA encoding such an effector

molecule has the sequence of a naturally occurring enzyme or toxin encoding DNA, or a mutant thereof, and can be prepared by methods well known in the art.

Uses of the Present Antibodies/Polypeptides

5 The polypeptides and/or antibodies utilized herein are especially indicated for diagnostic and therapeutic applications.

 The present antibodies can be administered as a therapeutic to cancer patients, especially, but not limited to CLL patients. In some embodiments, the antibodies are capable of interfering with the interaction of CD200 and its receptors. This interference
10 can block the immune suppressing effect of CD200. By improving the immune response in this manner, such antibodies can promote the eradication of cancer cells.

 The anti-CD200 antibody can also be administered in combination with other immunomodulatory compounds, vaccines or chemotherapy. For example, elimination of existing regulatory T cells with reagents such as anti-CD25 or cyclophosphamide is
15 achieved in one particularly useful embodiment before starting anti-CD200 treatment. Also, therapeutic efficacy of myeloablative therapies followed by bone marrow transplantation or adoptive transfer of T cells reactive with CLL cells is enhanced by anti-CD200 therapy. Furthermore, anti-CD200 treatment can substantially enhance efficacy of cancer vaccines such as dendritic cells loaded with CLL cells or proteins, peptides or
20 RNA derived from such cells, patient-derived heat-shocked proteins (hsp's), tumor peptides or protein. In other embodiments, an anti-CD200 antibody is used in combination with an immuno-stimulatory compound, such as CpG, toll-like receptor agonists or any other adjuvant, anti-CTLA-4 antibodies, and the like. In yet other embodiments, efficacy of anti-CD200 treatment is improved by blocking of
25 immunosuppressive mechanisms such as anti-PDL1 and/or 2 antibodies, anti-IL-10 antibodies, anti-IL-6 antibodies, and the like. In yet other embodiments, efficacy of anti-CD200 treatment is improved by administration of agents that increase NK cell number or T-cell such as the small molecule inhibitor IMiDs, thalidomide, or thalidomide analogs).

30 Anti-CD200 antibodies in accordance with the present disclosure can also be used as a diagnostic tool. For example, using blood obtained from patients with hematopoietic cancers, expression of CD200 can be evaluated on cancer cells by FACS analysis using anti-CD200 antibodies in combination with the appropriate cancer cell markers such as, e.g., CD38 and CD19 on CLL cells. Patients with CD200 levels at least 1.4-fold above

the levels found on normal B cells can be selected for treatment with anti-CD200 antibodies.

In another example of using the present anti-CD200 antibodies as a diagnostic tool, biopsies from patients with malignancies are obtained and expression of CD200 is
5 determined by FACS analysis using anti-CD200 antibodies. If tumor cells express CD200 at levels that are at least 1.4-fold higher compared to corresponding normal tissue, cancer patients are selected for immunomodulatory therapy. Immunomodulatory therapy can be anti-CD200 therapy, but can also be any other therapy affecting the patient's immune system. Examples of suitable immunomodulatory therapies include the
10 administration of agents that block negative regulation of T cells or antigen presenting cells (e.g., anti-CTLA4, anti-PD-L1, anti-PDL-2, anti-PD-1) or the administration of agents that enhance positive co-stimulation of T cells (e.g., anti-CD40 or anti 4-1BB). Furthermore, immunomodulatory therapy could be the administration of agents that increase NK cell number or T-cell activity (e.g., anti-CD200 antibodies alone or in
15 combination with inhibitors such as IMiDs, thalidomide, or thalidomide analogs) or the administration of agents that deplete regulatory T cells (e.g. anti-CD200 antibodies alone or in combination with ONTAK). Furthermore, immunomodulatory therapy could be cancer vaccines such as dendritic cells loaded with tumor cells, tumor RNA or tumor DNA, tumor protein or tumor peptides, patient derived heat-shocked proteins (hsp's) or
20 general adjuvants stimulating the immune system at various levels such as CpG, Luivac, Biostim, Ribominy, Imudon, Bronchovaxom or any other compound activating receptors of the innate immune system (e.g., toll like receptors). Also, therapy could include treatment with cytokines such as IL-2, GM-CSF and IFN-gamma.

In another embodiment in accordance with the present disclosure, methods are
25 provided for monitoring the progress and/or effectiveness of a therapeutic treatment. The method involves administering an immunomodulatory therapy and determining OX-2/CD200 levels in a subject at least twice to determine the effectiveness of the therapy. For example, pre-treatment levels of OX-2/CD200 can be ascertained and, after at least one administration of the therapy, levels of OX-2/CD200 can again be determined. A
30 decrease in OX-2/CD200 levels is indicative of an effective treatment. Measurement of OX-2/CD200 levels can be used by the practitioner as a guide for increasing dosage amount or frequency of the therapy. It should of course be understood that OX-2/CD200 levels can be directly monitored or, alternatively, any marker that correlates with OX-2/CD200 can be monitored.

The present antibodies also may be utilized to detect cancerous cells in vivo. This is achieved by labeling the antibody, administering the labeled antibody to a subject, and then imaging the subject. Examples of labels useful for diagnostic imaging in accordance with the present disclosure are radiolabels such as ^{131}I , ^{111}In , ^{123}I , $^{99\text{m}}\text{Tc}$, ^{32}P , ^{125}I , ^3H , ^{14}C ,
5 and ^{188}Rh , fluorescent labels such as fluorescein and rhodamine, nuclear magnetic resonance active labels, positron emitting isotopes detectable by a positron emission tomography ("PET") scanner, chemiluminescers such as luciferin, and enzymatic markers such as peroxidase or phosphatase. Short-range radiation emitters, such as isotopes detectable by short-range detector probes, such as a transrectal probe, can also be
10 employed. The antibody can be labeled with such reagents using techniques known in the art. For example, see Wensel and Meares, *Radioimmunoimaging and Radioimmunotherapy*, Elsevier, N.Y. (1983), which is hereby incorporated by reference, for techniques relating to the radiolabeling of antibodies. See also, D. Colcher et al., "Use of Monoclonal Antibodies as Radiopharmaceuticals for the Localization of Human
15 Carcinoma Xenografts in Athymic Mice", *Meth. Enzymol.* 121: 802-816 (1986), which is hereby incorporated by reference.

A radiolabeled antibody in accordance with this disclosure can be used for in vitro diagnostic tests. The specific activity of an antibody, binding portion thereof, probe, or ligand, depends upon the half-life, the isotopic purity of the radioactive label, and how the
20 label is incorporated into the biological agent. In immunoassay tests, the higher the specific activity, in general, the better the sensitivity. Procedures for labeling antibodies with the radioactive isotopes are generally known in the art.

The radiolabeled antibody can be administered to a patient where it is localized to cancer cells bearing the antigen with which the antibody reacts, and is detected or
25 "imaged" in vivo using known techniques such as radionuclear scanning using e.g., a gamma camera or emission tomography. See e.g., A. R. Bradwell et al., "Developments in Antibody Imaging", *Monoclonal Antibodies for Cancer Detection and Therapy*, R. W. Baldwin et al., (eds.), pp. 65-85 (Academic Press 1985), which is hereby incorporated by reference. Alternatively, a positron emission transaxial tomography scanner, such as
30 designated Pet VI located at Brookhaven National Laboratory, can be used where the radiolabel emits positrons (e.g., ^{11}C , ^{18}F , ^{15}O , and ^{13}N).

Fluorophore and chromophore labeled biological agents can be prepared from standard moieties known in the art. Since antibodies and other proteins absorb light having wavelengths up to about 310 nm, the fluorescent moieties should be selected to have substantial absorption at wavelengths above 310 nm and preferably above 400 nm.

5 A variety of suitable fluorescers and chromophores are described by Stryer, *Science*, 162:526 (1968) and Brand, L. et al., *Annual Review of Biochemistry*, 41:843-868 (1972), which are hereby incorporated by reference. The antibodies can be labeled with fluorescent chromophore groups by conventional procedures such as those disclosed in U.S. Patent Nos. 3,940,475, 4,289,747, and 4,376,110, which are hereby incorporated by
10 reference.

In other embodiments, bispecific antibodies are contemplated. Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the CD200 antigen on a cancer cell, the other one is for any other
15 antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are within the purview of those skilled in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, *Nature*, 305:537-539
20 (1983)). Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, including at least part of the hinge, CH2, and CH3 regions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into
25 separate expression vectors, and are co-transfected into a suitable host organism. For further details of illustrative currently known methods for generating bispecific antibodies see, for example, Suresh et al., *Methods in Enzymology*, 121:210 (1986); WO 96/27011; Brennan et al., *Science* 229:81 (1985); Shalaby et al., *J. Exp. Med.* 175:217-225 (1992); Kostelny et al., *J. Immunol.* 148(5):1547-1553 (1992); Hollinger et al., *Proc. Natl. Acad.*
30 *Sci. USA* 90:6444-6448 (1993); and Gruber et al., *J. Immunol.* 152:5368 (1994); and Tutt et al., *J. Immunol.* 147:60 (1991).

The present antibodies can also be utilized to directly kill or ablate cancerous cells in vivo. This involves administering the antibodies (which are optionally fused to a cytotoxic drug) to a subject requiring such treatment. Since the antibodies recognize CD200 on cancer cells, any such cells to which the antibodies bind are destroyed.

5 Where the antibodies are used alone to kill or ablate cancer cells, such killing or ablation can be effected by initiating endogenous host immune functions, such as complement-mediated or antibody-dependent cellular cytotoxicity. Assays for determining whether an antibody kills cells in this manner are within the purview of those skilled in the art.

10 The antibodies of the present disclosure may be used to deliver a variety of cytotoxic compounds. Any cytotoxic compound can be fused to the present antibodies. The fusion can be achieved chemically or genetically (e.g., via expression as a single, fused molecule). The cytotoxic compound can be a biological, such as a polypeptide, or a small molecule. As those skilled in the art will appreciate, for small molecules, chemical
15 fusion is used, while for biological compounds, either chemical or genetic fusion can be employed.

 Non-limiting examples of cytotoxic compounds include therapeutic drugs, a compound emitting radiation, molecules of plants, fungal, or bacterial origin, biological proteins, and mixtures thereof. The cytotoxic drugs can be intracellularly acting cytotoxic
20 drugs, such as short-range radiation emitters, including, for example, short-range, high-energy α -emitters. Enzymatically active toxins and fragments thereof are exemplified by diphtheria toxin A fragment, nonbinding active fragments of diphtheria toxin, exotoxin A (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, .alpha.-sacrin, certain *Aleurites fordii* proteins, certain *Dianthin* proteins, *Phytolacca americana*
25 proteins (PAP, PAPII and PAP-S), *Morodica charantia* inhibitor, curcin, crotin, *Saponaria officinalis* inhibitor, gelonin, mitogillin, restrictocin, phenomyacin, and enomyacin, for example. Procedures for preparing enzymatically active polypeptides of the immunotoxins are described in WO84/03508 and WO85/03508, which are hereby
30 incorporated by reference. Certain cytotoxic moieties are derived from adriamycin, chlorambucil, daunomycin, methotrexate, neocarzinostatin, and platinum, for example.

 Procedures for conjugating the antibodies with the cytotoxic agents have been previously described and are within the purview of one skilled in the art.

 Alternatively, the antibody can be coupled to high energy radiation emitters, for example, a radioisotope, such as ^{131}I , a γ -emitter, which, when localized at the tumor site,

results in a killing of several cell diameters. See, e.g., S. E. Order, "Analysis, Results, and Future Prospective of the Therapeutic Use of Radiolabeled Antibody in Cancer Therapy", *Monoclonal Antibodies for Cancer Detection and Therapy*, R. W. Baldwin et al. (eds.), pp 303-316 (Academic Press 1985), which is hereby incorporated by reference. Other
5 suitable radioisotopes include α -emitters, such as ^{212}Bi , ^{213}Bi , and ^{211}At , and β -emitters, such as ^{186}Re and ^{90}Y .

The route of antibody administration of the present antibodies (whether the pure antibody, a labeled antibody, an antibody fused to a toxin, etc.) is in accord with known methods, e.g., injection or infusion by intravenous, intraperitoneal, intracerebral,
10 intramuscular, subcutaneous, intraocular, intraarterial, intrathecal, inhalation or intralesional routes, or by sustained release systems. The antibody is preferably administered continuously by infusion or by bolus injection. One may administer the antibodies in a local or systemic manner.

The present antibodies may be prepared in a mixture with a pharmaceutically
15 acceptable carrier. Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition. This therapeutic composition can be administered intravenously or through the nose or lung, preferably as a liquid or powder aerosol (lyophilized). The composition may also be administered parenterally or
20 subcutaneously as desired. When administered systemically, the therapeutic composition should be sterile, pyrogen-free and in a parenterally acceptable solution having due regard for pH, isotonicity, and stability. These conditions are known to those skilled in the art.

Pharmaceutical compositions suitable for use include compositions wherein one or more of the present antibodies are contained in an amount effective to achieve their
25 intended purpose. More specifically, a therapeutically effective amount means an amount of antibody effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated. Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. Therapeutically effective dosages may be determined
30 by using in vitro and in vivo methods.

In some embodiments, present CD200 binding antibodies provide the benefit of blocking immune suppression in CLL by targeting the leukemic cells directly through CD200. Specifically, stimulating the immune system can allow the eradication of CLL cells from the spleen and lymph nodes. Applicants are unaware of any successful

eradication of CLL cells from these microenvironments having been achieved with agents that simply target B cells (such as alemtuzumab). In contrast, CLL reactive T cells can have better access to these organs than antibodies. In other embodiments, direct cell killing is achieved by tagging the CLL cells with anti-CD200 Abs.

5 In particularly useful embodiments, the combination of direct cell killing and driving the immune response towards a Th1 profile provides a particularly powerful approach to cancer treatment. Thus, in one embodiment, a cancer treatment is provided wherein an antibody or antibody fragment, which binds to CD200 and both a) blocks the interaction between CD200 and its receptor and b) directly kills the cancer cells
10 expressing CD200, is administered to a cancer patient. The mechanism by which the cancer cells are killed can include, but are not limited to antibody-dependent cellular cytotoxicity (ADCC) or complement dependent cytotoxicity (CDC); fusion with a toxin; fusion with a radiolabel; fusion with a biological agent involved in cell killing, such as granzyme B or perforin; fusion with a cytotoxic virus; fusion with a cytokine such as
15 TNF- α or IFN- α . In an alternative embodiment, a cancer treatment involves administering an antibody that both a) blocks the interaction between CD200 and its receptor and b) enhances cytotoxic T cell or NK cell activity against the tumor. Such enhancement of the cytotoxic T cell or NK cell activity may, for example, be combined by fusing the antibody with cytokines such as e.g. IL-2, IL-12, IL-18, IL-13, and IL-5. In
20 addition, such enhancement may be achieved by administration an anti-CD200 antibody in combination with inhibitors such as IMiDs, thalidomide, or thalidomide analogs.

 In yet another embodiment, the cancer treatment involves administering an antibody that both a) blocks the interaction between CD200 and its receptor and b) attracts T cells to the tumor cells. T cell attraction can be achieved by fusing the Ab with
25 chemokines such as MIG, IP-10, I-TAC, CCL21, CCL5 or LIGHT. The combined action of blocking immune suppression and killing directly through antibody targeting of the tumor cells is a unique approach that provides increased efficacy.

 While the above disclosure has been directed to antibodies, in some embodiments polypeptides derived from such antibodies can be utilized in accordance with the present
30 disclosure.

Uses of the CLL Cell Line

There are many advantages to the development of a CLL cell line, as it provides an important tool for the development of diagnostics and treatments for CLL, cancer, and other disease states characterized by upregulated levels of OX-2/CD200, e.g., melanoma.

5 A cell line according to this disclosure may be used for in vitro studies on the etiology, pathogenesis and biology of CLL and other disease states characterized by upregulated levels of OX-2/CD200. This assists in the identification of suitable agents that are useful in the therapy of these diseases.

10 The cell line may also be used to produce polypeptides and/or monoclonal antibodies for in vitro and in vivo diagnosis of CLL, cancer, and other disease states characterized by upregulated levels of OX-2/CD200 (e.g., melanoma), as referred to above, and for the screening and/or characterization of antibodies produced by other methods, such as by panning antibody libraries with primary cells and/or antigens derived from CLL patients.

15 The cell line may be used as such, or antigens may be derived therefrom. Advantageously, such antigens are cell-surface antigens specific for CLL. They may be isolated directly from cell lines according to this disclosure. Alternatively, a cDNA expression library made from a cell line described herein may be used to express CLL-specific antigens, useful for the selection and characterization of anti-CLL antibodies and
20 the identification of novel CLL-specific antigens.

Treatment of CLL using monoclonal antibody therapy has been proposed in the art. Recently, Hainsworth (*Oncologist* 5 (5) (2000) 376-384) has described the current therapies derived from monoclonal antibodies. Lymphocytic leukemia in particular is considered to be a good candidate for this therapeutic approach due to the presence of
25 multiple lymphocyte-specific antigens on lymphocyte tumors.

Existing antibody therapies (such as RituximabTM, directed against the CD20-antigen, which is expressed on the surface of B-lymphocytes) have been used successfully against certain lymphocytic disease. However, a lower density CD20 antigen is expressed on the surface of B-lymphocytes in CLL (Almasri et al., *Am. J. Hematol.*, 40
30 (4) (1992) 259-263).

The CLL cell line described herein thus permits the development of novel anti-CLL antibodies and polypeptides having specificity for one or more antigenic determinants of the present CLL cell line, and their use in the therapy and diagnosis of

CLL, cancer, and other disease states characterized by upregulated levels of OX-2/CD200.

The antibody or polypeptide may bind to a receptor with which OX-2/CD200 normally interacts, thereby preventing or inhibiting OX-2/CD200 from binding to the receptor. As yet another alternative, the antibody can bind to an antigen that modulates expression of OX-2/CD200, thereby preventing or inhibiting normal or increased expression of OX-2/CD200. Because the presence of OX-2/CD200 has been associated with reduced immune response, it would be desirable to interfere with the metabolic pathway of OX-2/CD200 so that the patient's immune system can defend against the disease state, such as cancer or CLL, more effectively.

In a particularly useful embodiment, the polypeptide binds to OX-2/CD200. In one embodiment, the polypeptide can be an antibody which binds to OX-2/CD200 and prevents or inhibits OX-2/CD200 from interacting with other molecules or receptors. As CLL cells and other cells overexpressing OX-2/CD200 greatly diminish the production of Th1 cytokines, the administration of anti-CD200 antibody or a polypeptide which binds to OX-2/CD200 to a subject having upregulated levels of OX-2/CD200 restores the Th1 cytokine profile. Thus, these polypeptides and/or antibodies can be useful therapeutic agents in the treatment of CLL and other cancers or diseases overexpressing OX-2/CD200.

Thus, in another embodiment, the method of the present disclosure includes the steps of screening a subject for the presence OX-2/CD200 and administering a polypeptide that binds to OX-2/CD200. It should of course be understood that the presence of OX-2/CD200 can be directly monitored or, alternatively, any marker that correlates with OX-2/CD200 can be detected. In a particularly useful embodiment, a CLL patient is screened for overexpression of OX-2/CD200 and an antibody that binds to OX-2/CD200 is administered to the patient. One such antibody is the commercially available anti-CD200 antibody from Serotec Inc. (3200 Atlantic Ave, Suite 105, Raleigh, NC 27604). As described in detail below, another such antibody is scFv-9 (see Fig. 9B) which binds to OX-2/CD200.

In another aspect, the present disclosure provides methods for assessing the immunomodulatory effect of molecules expressed by cancer cells. In these methods a molecule that is expressed or upregulated by a cancer cell is first identified. The molecule can be identified from a database or experimentally. Databases that identify molecules that are expressed or upregulated by cancer cells are known and include, for

example, the NCI60 cancer microarray project (<http://genome-www.stanford.edu/nci60/>) (Ross et al., Nature Genetics 24: 227-34, 2000), the Carcinoma classification (<http://www.gnf.org/cancer/epican/>) (Andrew I. Su et al., "Molecular Classification of Human Carcinomas by Use of Gene Expression Signatures." Cancer Research 61:7388-7393, 2001), and the Lymphoma/Leukemia molecular profiling project (<http://lmpp.nih.gov/lymphoma/>) (Alizadeh et al., Nature 403: 503-11, 2000).

Experimental methods useful for identifying molecules that are expressed or upregulated by cancer cells are also known and include, for example microarray experiments, quantitative PCR, FACS, and Northern analysis.

10 Cancer cells, lymphocytes and the previously identified molecule that is expressed by a cancer cell are administered to a subject and the rate of growth of the cancer cells is monitored. Any type of cancer cells can be employed in the present methods. In some embodiments, the cancer cells express an immunosuppressive compound. In particularly useful embodiments, the cancer cells express or even overexpress CD200. Suitable
15 cancer cells include, but are not limited to lymphoma cell lines such as the RAJI or Namalwa cell lines. The amount of cancer cells administered may range from about 1×10^6 to about 20×10^6 .

Any type of lymphocyte may be employed in the present process. Suitable lymphocytes include, for example, PBLs, T cells, cytotoxic T cells, dendritic cells or NK
20 cells. In particularly useful embodiments, the lymphocytes are human lymphocytes, specifically human PBLs. The number of lymphocytes administered is predetermined to be either a) sufficient to slow the growth of the cancer cells or b) insufficient to slow the growth of cancer cells. The amount of lymphocytes administered may be greater than or equal to the number of cancer cells administered when the number of lymphocytes
25 administered is intended to be sufficient to slow the growth of the cancer cells. In embodiments, the amount of lymphocytes administered may be from about 5×10^6 to about 10×10^6 when the number of lymphocytes administered is intended to be sufficient to slow the growth of the cancer cells. The amount of lymphocytes administered may be less than the number of cancer cells administered when the number of lymphocytes
30 administered is intended to be insufficient to slow the growth of the cancer cells. In embodiments, the amount of lymphocytes administered may range from about 1×10^6 to about 4×10^6 when the number of lymphocytes administered is intended to be insufficient to slow the growth of the cancer cells. The foregoing amounts are illustrative and other suitable amounts may be experimentally determined and used in the present methods.

The molecule that is expressed or upregulated by a cancer cell can be administered as the molecule or the active portion thereof itself (either isolated or recombinantly generated), or by administering cells that produce the molecule naturally, or by administering cells that have been engineered to produce the molecule or portions thereof. Methods for engineering cells to express a desired molecule are known to those skilled in the art. The amount of molecule administered can be any amount above the amount found in a healthy individual. For example, the amount of the molecule administered can be from about 1.4-fold above what is found in a healthy individual in the same type of cell to about 10,000-fold above what is found in a healthy individual in the same type of cell. It should be understood that the molecule being assessed may or may not be known to have some degree of immunomodulatory activity. Thus, the present methods may be used to confirm the immunomodulatory effect of a molecule as well as to determine such activity *ab initio*.

Any small animal may be chosen as the subject to which the cancer cells, lymphocytes and the molecule are administered. The subject may advantageously be immunocompromised. Suitable small animals include, for example, immunodeficient mice, irradiated rats, irradiated guinea pigs and the like.

The rate of cancer cell growth can be monitored using conventional techniques. For example, tumor growth can be monitored by measuring length and width with a caliper. Tumor volume can be calculated, for example, based on multiplying the length of the tumor by the width of the tumor and then multiplying by one-half the width of the tumor. The rate of cancer cell growth can be measured periodically, such as, for example, three times a week. The rate of growth of cancer cells when cancer cells and lymphocytes alone have been administered may be determined by administering cancer cells and lymphocytes alone to a control subject and periodically measuring tumor size. Alternatively, cancer cells and lymphocytes alone can be administered initially, and once a baseline growth rate is established, the molecule that is expressed or upregulated by a cancer cell can be subsequently administered to the subject and the rate of growth of the cancer cells after the second administration can be measured. Alternatively, with systemic models the rate of cancer cell growth can be monitored using FACS, survival or other conventional techniques.

If any change in the growth rate of the cancer cells is observed compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the molecule is deemed to have an immunomodulatory effect. If the number of lymphocytes

administered is sufficient to slow the growth of the cancer cells and the rate of growth of the tumor cells observed is higher compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the molecule that is expressed or upregulated by a cancer cell is considered immunosuppressive. If the number of lymphocytes administered is insufficient to slow the growth of the cancer cells and the rate of growth of the tumor cells observed is lowered compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the molecule is considered immune enhancing. Typically, the rate of cancer cell growth observed may be about 20 to about 1,000% higher than the rate of growth when the tumor cells and donor lymphocytes are administered alone for the immunosuppressing or immune enhancing effect to be statistically significant.

Once the immunomodulatory effect of the molecule is established, compounds that either enhance or inhibit the activity of the molecule can be identified in accordance with embodiments described herein. The compound that either enhances or inhibits the activity of the molecule previously found to have immunomodulatory effect can be any compound that alters the protein/protein interaction that provides the immunomodulatory effect. The enhancing or inhibiting effect can be the result of direct interaction with the compound expressed or upregulated by the cancer cell or may be the result of an interaction with other compounds in the metabolic pathway of the compound expressed or upregulated by the cancer cell.

For example, antibodies or functional antibody fragments can be identified that interact with the molecule previously found to have an immunomodulatory effect, the receptor with which the molecule interacts, or some other molecule in the metabolic pathway of the molecule responsible for the immunomodulatory effect. Techniques for making antibodies (including antibody libraries) and screening them for an inhibitory or enhancing effect will be apparent to those skilled in the art. As another example, small molecules may be screened for an inhibitory or enhancing effect. Techniques for screening small molecule libraries for an inhibitory or enhancing effect will be apparent to those skilled in the art.

In another aspect, methods for assessing the immunomodulatory effect of a compound are contemplated by the present disclosure. Demonstrating immunomodulatory properties of compounds or molecules acting on the human immune system is very challenging in small animal models. Often, the compounds do not act on the immune system of small animals, requiring reconstitution of the human immune

system in mice. Reconstitution can be accomplished by grafting of various fetal immune organs into mice or by injection of human lymphocytes, but none of the models described to date has proven to be useful for demonstrating immunomodulatory properties of compounds or molecules. The immune system is believed to play an important role in eradicating cancer cells. Cancer cells have found ways to evade the immune system by upregulation of immunosuppressive receptors.

The present methods of assessing the immunomodulatory effect of a compound is accomplished by a model that mimics the graft versus leukemia effect observed in patients with leukemia that are infused with donor lymphocytes (e.g., PBLs) resulting in remission in up to 80% of patients. The method involves administering cancer cells, lymphocytes and the compound to be assessed to a subject and the rate of growth of the cancer cells is monitored.

Any type of cancer cells can be employed in the present methods. In some embodiments, the cancer cells express an immunosuppressive compound. In particularly useful embodiments, the cancer cells express or even overexpress CD200. Suitable cancer cells include, but are not limited to lymphoma cell lines such as the RAJI or Namalwa cell lines. The amount of cancer cells administered may range from about 2×10^6 to about 20×10^6 .

Any type of lymphocyte may be employed in the present process. Suitable lymphocytes include, for example, PBLs, dendritic cells, T cells, cytotoxic T cells, NK cells. In particularly useful embodiments, the lymphocytes are human lymphocytes, specifically human PBLs. The number of lymphocytes administered is predetermined to be either a) sufficient to slow the growth of the cancer cells or b) insufficient to slow the growth of cancer cells. The amount of lymphocytes administered may be greater than or equal to the number of cancer cells administered when the number of lymphocytes administered is intended to be sufficient to slow the growth of the cancer cells. In embodiments, the amount of lymphocytes administered may be from about 5×10^6 to about 10×10^6 when the number of lymphocytes administered is intended to be sufficient to slow the growth of the cancer cells. The amount of lymphocytes administered may be less than the number of cancer cells administered when the number of lymphocytes administered is intended to be insufficient to slow the growth of the cancer cells. In embodiments, the amount of lymphocytes administered may range from about 1×10^6 to about 4×10^6 when the number of lymphocytes administered is intended to be insufficient

to slow the growth of the cancer cells. The foregoing amounts are illustrative and other suitable amounts may be experimentally determined and used in the present methods.

The compound being assessed can be any compound whose immunomodulatory effect is sought to be determined. Illustrative examples of compounds include the antibodies and peptides described herein above. The amount of the compound being assessed administered may range from about 1 mg/kg to about 200 mg/kg. The compound to be assessed can be administered as the compound itself, or by administering cells that produce the compound naturally, or by administering cells that have been engineered to produce the compound. It should be understood that the compound being assessed may or may not be known to have some degree of immunomodulatory activity. Thus, the present methods may be used to confirm the immunomodulatory effect of a compound as well as to determine such activity *ab initio*.

Any small animal may be chosen as the subject to which the cancer cells, lymphocytes and the compound are administered. The subject may advantageously be immunocompromised. Suitable small animals include, for example, immunodeficient mice, irradiated rats, irradiated guinea pigs and the like.

The rate of cancer cell growth can be monitored using conventional techniques. For example, tumor growth can be monitored by measuring length and width with a caliper. Tumor volume can be calculated, for example, based on multiplying the length of the tumor by the width of the tumor and then multiplying by one-half the width of the tumor. The rate of cancer cell growth can be measured periodically, such as, for example, three times a week. The rate of growth of cancer cells when cancer cells and lymphocytes alone have been administered may be determined by administering cancer cells and lymphocytes alone to a control subject and periodically measuring tumor size. Alternatively, cancer cells and lymphocytes alone can be administered initially, and once a baseline growth rate is established, the compound to be assessed can be subsequently administered to the subject and the rate of growth of the cancer cells after the second administration can be measured.

When injecting cancer cells such as the lymphoma cell lines RAJI or Namalwa into immune-deficient mice, administration of five to 10 million PBLs results in significantly slower tumor growth. In contrast, low PBL numbers (1-2 million depending on donor) do not slow tumor growth.

If any change in the growth rate of the cancer cells is observed compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the

molecule is deemed to have an immunomodulatory effect. If the number of lymphocytes administered is sufficient to slow the growth of the cancer cells and the rate of growth of the tumor cells observed is higher compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the compound is considered immunosuppressive. If the number of lymphocytes administered is insufficient to slow the growth of the cancer cells and the rate of growth of the tumor cells observed is lowered compared to the rate of growth when the tumor cells and donor lymphocytes are administered alone, the compound is considered immune enhancing.

In order that those skilled in the art may be better able to practice the compositions and methods described herein, the following examples are given for illustration purposes.

EXAMPLE 1

Isolation of Cell Line CLL-AAT

Establishment of the cell line

Peripheral blood from a patient diagnosed with CLL was obtained. The WBC count was 1.6×10^8 /ml. Mononuclear cells were isolated by Histopaque-1077 density gradient centrifugation (Sigma Diagnostics, St. Louis, MO). Cells were washed twice with Iscove's Modified Dulbecco's Medium (IMDM) supplemented with 10% heat-inactivated fetal bovine serum (FBS), and resuspended in 5 ml of ice-cold IMDM/10% FBS. Viable cells were counted by staining with trypan blue. Cells were mixed with an equal volume of 85% FBS/15% DMSO and frozen in 1 ml aliquots for storage in liquid nitrogen.

Immunophenotyping showed that >90% of the CD45⁺ lymphocyte population expressed IgD, kappa light chain, CD5, CD19, and CD23. This population also expressed low levels of IgM and CD20. Approximately 50% of the cells expressed high levels of CD38. The cells were negative for lambda light chain, CD10 and CD138

An aliquot of the cells was thawed, washed, and resuspended at a density of 10^7 /mL in IMDM supplemented with 20% heat-inactivated FBS, 2mM L-glutamine, 100 units/ml penicillin, 100 µg/ml streptomycin, 50 µM 2-mercaptoethanol, and 5 ng/ml recombinant human IL-4 (R & D Systems, Minneapolis, MN). The cells were cultured at 37°C in a humidified 5% CO₂ atmosphere. The medium was partially replaced every 4 days until steady growth was observed. After 5 weeks, the number of cells in the culture began to double approximately every 4 days. This cell line was designated CLL-AAT.

Characterization of the cell line

Immunophenotyping of the cell line by flow cytometry showed high expression of IgM, kappa light chain, CD23, CD38, and CD138, moderate expression of CD19 and CD20, and weak expression of IgD and CD5. The cell line was negative for lambda light chain, CD4, CD8, and CD10.

- 5 Immunophenotyping of the cell line was also done by whole cell ELISA using a panel of rabbit scFv antibodies that had been selected for specific binding to primary B-CLL cells. All of these CLL-specific scFv antibodies also recognized the CLL -AAT cell line. In contrast, the majority of the scFvs did not bind to two cell lines derived from B cell lymphomas: Ramos, a Burkitt's lymphoma cell line, and RL, a non-Hodgkin's
10 lymphoma cell line.

EXAMPLE 2

Selection of scFv Antibodies for B-CLL-specific Cell Surface Antigens using Antibody Phage Display and Cell Surface Panning

Immunizations and scFv antibody library construction

- Peripheral blood mononuclear cells (PBMC) were isolated from blood drawn from CLL patients at the Scripps Clinic (La Jolla, CA). Two rabbits were immunized with
15 2×10^7 PBMC pooled from 10 different donors with CLL. Three immunizations, two subcutaneous injections followed by one intravenous injection, were done at three week intervals. Serum titers were checked by measuring binding of serum IgG to primary CLL cells using flow cytometry. Five days after the final immunization, spleen, bone marrow, and PBMC were harvested from the animals. Total RNA was isolated from these tissues
20 using Tri-Reagent (Molecular Research Center, Inc). Single-chain Fv (scFv) antibody phage display libraries were constructed as previously described (Barbas et al., (2001) Phage Display: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York). For cell surface panning, phagemid particles from the reamplified library were precipitated with polyethylene glycol (PEG), resuspended in phosphate-
25 buffered saline (PBS) containing 1% bovine serum albumin (BSA), and dialysed overnight against PBS.

Antibody selection by cell surface panning

The libraries were enriched for CLL cell surface-specific antibodies by positive-negative selection with a magnetically-activated cell sorter (MACS) as described by Siegel et al. (1997, *J. Immunol. Methods* 206:73-85). Briefly, phagemid particles from the scFv antibody library were preincubated in MPBS (2% nonfat dry milk, 0.02% sodium azide in PBS, pH 7.4) for 1 hour at 25°C to block nonspecific binding sites. 5 Approximately 10^7 primary CLL cells were labeled with mouse anti-CD5 IgG and mouse anti-CD19 IgG conjugated to paramagnetic microbeads (Miltenyi Biotec, Sunnyvale, CA). Unbound microbeads were removed by washing. The labeled CLL cells (“target cells”) were mixed with an excess of “antigen-negative absorber cells”, pelleted, and 10 resuspended in 50 μ l (10^{10} - 10^{11} cfu) of phage particles. The absorber cells serve to soak up phage that stick non-specifically to cell surfaces as well as phage specific for “common” antigens present on both the target and absorber cells. The absorber cells used were either TF-1 cells (a human erythroleukemia cell line) or normal human B cells 15 isolated from peripheral blood by immunomagnetic negative selection (StemSep system, StemCell Technologies, Vancouver, Canada). The ratio of absorber cells to target cells was approximately 10 fold by volume. After a 30 minute incubation at 25°C, the cell/phage mixture was transferred to a MiniMACS MS+ separation column. The column was washed twice with 0.5 ml of MPBS, and once with 0.5 ml of PBS to remove the unbound phage and absorber cells. The target cells were eluted from the column in 1 ml 20 of PBS and pelleted in a microcentrifuge at maximum speed for 15 seconds. The captured phage particles were eluted by resuspending the target cells in 200 μ l of acid elution buffer (0.1 N HCl, pH adjusted to 2.2 with glycine, plus 1 mg/ml BSA). After a 10 minute incubation at 25°C, the buffer was neutralized with 12 μ L of 2M Tris base, pH10.5, and the eluted phage were amplified in *E. coli* for the next round of panning. For 25 each round of panning, the input and output phage titers were determined. The input titer is the number of reamplified phage particles added to the target cell/absorber cell mixture and the output titer is the number of captured phage eluted from the target cells. An enrichment factor (E) is calculated using the formula $E=(R_n \text{ output}/R_n \text{ input})/(R_1 \text{ output}/R_1 \text{ input})$, where R_1 = round 1 and R_n = round 2, 3, or 4. In most cases, an 30 enrichment factor of 10^2 - 10^3 fold should be attained by the third or fourth round.

Analysis of enriched antibody pools following panning

After 3-5 rounds of panning, the pools of captured phage were assayed for binding to CLL cells by flow cytometry and/or whole cell ELISA:

1. To produce an entire pool in the form of HA-tagged soluble antibodies, 2ml of a non-suppressor strain of *E. coli* (e.g. TOP10F') was infected with 1 μ l (10^9 - 10^{10} cfu) of phagemid particles. The original, unpanned library was used as a negative control. Carbenicillin was added to a final concentration of 10 μ M and the culture was incubated at 37°C with shaking at 250rpm for 1 hour. Eight ml of SB medium containing 50 μ g/ml carbenicillin was added and the culture was grown to an OD 600 of ~0.8. IPTG was added to a final concentration of 1mM to induce scFv expression from the Lac promoter and shaking at 37°C was continued for 4 hours. The culture was centrifuged at 3000xg for 15'. The supernatant containing the soluble antibodies was filtered and stored in 1 ml aliquots at -20°C.
2. Binding of the scFv antibody pools to target cells vs. absorber cells was determined by flow cytometry using high-affinity Rat Anti-HA (clone 3F10, Roche Molecular Biochemicals) as secondary antibody and PE-conjugated Donkey Anti-Rat as tertiary antibody.
3. Binding of the antibody pools to target cells vs. absorber cells was also determined by whole-cell ELISA as described below.

Screening individual scFv clones following panning

To screen individual scFv clones following panning, TOP10F' cells were infected with phage pools as described above, spread onto LB plates containing carbenicillin and tetracycline, and incubated overnight at 37°C. Individual colonies were inoculated into deep 96-well plates containing 0.6-1.0 ml of SB-carbenicillin medium per well. The cultures were grown for 6-8 hours in a HiGro shaking incubator (GeneMachines, San Carlos, CA) at 520 rpm and 37°C. At this point, a 90 μ l aliquot from each well was transferred to a deep 96-well plate containing 10 μ L of DMSO. This replica plate was stored at -80°C. IPTG was added to the original plate to a final concentration of 1 mM and shaking was continued for 3 hours. The plates were centrifuged at 3000xg for 15 minutes. The supernatants containing soluble scFv antibodies were transferred to another deep 96-well plate and stored at -20°C.

A sensitive whole-cell ELISA method for screening HA-tagged scFv antibodies was developed:

1. An ELISA plate is coated with concanavalin A (10mg/ml in 0.1 M NaHCO₃, pH8.6, 0.1 mM CaCl₂).
- 5 2. After washing the plate with PBS, 0.5-1x10⁵ target cells or absorber cells in 50µl of PBS are added to each well, and the plate is centrifuged at 250xg for 10 minutes.
3. 50µl of 0.02% glutaraldehyde in PBS are added and the cells are fixed overnight at 4°C.
4. After washing with PBS, non-specific binding sites are blocked with PBS containing
10 4% non-fat dry milk for 3 hours at room temperature.
5. The cells are incubated with 50µl of soluble, HA-tagged scFv or Fab antibody (TOP10F' supernatant) for 2 hours at room temperature, then washed six times with PBS.
6. Bound antibodies are detected using a Mouse Anti-HA secondary antibody (clone
15 12CA5) and an alkaline phosphatase (AP)-conjugated Anti-Mouse IgG tertiary antibody. An about 10-fold amplification of the signal is obtained by using AMDEX AP-conjugated Sheep Anti-Mouse IgG as the tertiary antibody (Amersham Pharmacia Biotech). The AMDEX antibody is conjugated to multiple AP molecules via a dextran backbone. Color is developed with the alkaline phosphatase substrate PNPP
20 and measured at 405nm using a microplate reader.

Primary screening of the scFv clones was done by ELISA on primary CLL cells versus normal human PBMC. Clones which were positive on CLL cells and negative on normal PBMC were rescreened by ELISA on normal human B cells, human B cell lines, TF-1 cells, and the CLL-AAT cell line. The clones were also rescreened by ELISA on
25 CLL cells isolated from three different patients to eliminate clones that recognized patient-specific or blood type-specific antigens. Results from representative ELISAs are shown in Figures 2-6 and summarized in Figs. 9A – 9C.

The number of unique scFv antibody clones obtained was determined by DNA fingerprinting and sequencing. The scFv DNA inserts were amplified from the plasmids
30 by PCR and digested with the restriction enzyme BstNI. The resulting fragments were separated on a 4% agarose gel and stained with ethidium bromide. Clones with different restriction fragment patterns must have different amino acid sequences. Clones with identical patterns probably have similar or identical sequences. Clones with unique BstNI fingerprints were further analyzed by DNA sequencing. Twenty-five different sequences

were found, which could be clustered into 16 groups of antibodies with closely related complementarity determining regions (Figs. 9A – 9C).

Characterization of scFv antibodies by flow cytometry

5 The binding specificities of several scFv antibodies were analyzed by 3-color flow cytometry (Fig. 7). PBMC isolated from normal donors were stained with FITC-conjugated anti-CD5 and PerCP-conjugated anti-CD19. Staining with scFv antibody was done using biotin-conjugated anti-HA as secondary antibody and PE-conjugated streptavidin. Three antibodies, scFv-2, scFv-3, and scFv-6, were found to specifically
10 recognize the CD19⁺ B lymphocyte population (data not shown). The fourth antibody, scFv-9, recognized two distinct cell populations: the CD19⁺ B lymphocytes and a subset of CD5⁺ T lymphocytes (Fig. 7). Further characterization of the T cell subset showed that it was a subpopulation of the CD4⁺CD8⁻ TH cells (data not shown).

To determine if the antigens recognized by the scFv antibodies were
15 overexpressed on primary CLL cells, PBMC from five CLL patients and five normal donors were stained with scFv and compared by flow cytometry (Fig. 8 and Table 2). By comparing the mean fluorescent intensities of the positive cell populations, the relative expression level of an antigen on CLL cells vs. normal cells could be determined. One antibody, scFv-2, consistently stained CLL cells less intensely than normal PBMC,
20 whereas scFv-3 and scFv-6 both consistently stained CLL cells more brightly than normal PBMC. The fourth antibody, scFv-9, stained two of the five CLL samples much more intensely than normal PBMC, but gave only moderately brighter staining for the other three CLL samples (Fig. 8 and Table 2). This indicates that the antigens for scFv-3 and scFv-6 are overexpressed approximately 1.4 fold on most if not all CLL tumors, whereas
25 scFv-9 is overexpressed 3 to 6-fold on a subset of CLL tumors.

CLL patients can be divided into two roughly equal groups: those with a poor prognosis (median survival time of 8 years) and those with a favorable prognosis (median survival time of 26 years). Several unfavorable prognostic indicators have been identified for CLL, most notably the presence of VH genes lacking somatic mutations and the
30 presence of a high percentage of CD38⁺ B cells. Since scFv-9 recognizes an antigen overexpressed in only a subset of CLL patients, it was sought to determine if scFv-9 antigen overexpression correlated with the percentage of CD38⁺ cells in blood samples from ten CLL patients (Fig. 11). The results indicate that scFv-9 antigen overexpression and percent CD38⁺ cells are completely independent of one another.

Identification of antigens recognized by scFv antibodies by immunoprecipitation (IP) and mass spectrometry (MS)

To identify the antigens for these antibodies, scFvs were used to immunoprecipitate the antigens from lysates prepared from the microsomal fraction of cell-surface biotinylated CLL-AAT cells (Fig. 12). The immunoprecipitated antigens were purified by SDS-PAGE and identified by matrix assisted laser desorption ionization mass spectrometry (MALDI-MS) or microcapillary reverse-phase HPLC nano-electrospray tandem mass spectrometry (μ LC/MS/MS) (data not shown). ScFv-2 immunoprecipitated a 110 kd antigen from both RL and CLL-AAT cells (Fig. 12). This antigen was identified by MALDI-MS as the B cell-specific marker CD19. ScFv-3 and scFv-6 both immunoprecipitated a 45 kd antigen from CLL-AAT cells (not shown). This antigen was identified by MALDI-MS as CD23, which is a known marker for CLL and activated B cells. ScFv-9 immunoprecipitated a 50 kd antigen from CLL-AAT cells (Fig. 12). This antigen was identified by μ LC/MS/MS as OX-2/CD200, a known marker for B cells, activated CD4⁺ T cells, and thymocytes. OX-2/CD200 is also expressed on some non-lymphoid cells such as neurons and endothelial cells.

EXAMPLE 3

The capability of cells overexpressing OX-2/CD200 to shift the cytokine response from a Th1 response (IL-2, IFN- γ) to a Th2 response (IL-4, IL-10) was assessed in a mixed lymphocyte reaction using monocyte-derived macrophages/dendritic cells from one donor and blood-derived T cells from a different donor. As a source of OX-2/CD200-expressing cells, either OX-2/CD200 transfected EBNA cells as described below or CLL patient samples were used.

25

Transfection of 293-EBNA cells

293-EBNA cells (Invitrogen) were seeded at 2.5×10^6 per 100mm dish. 24 hours later the cells were transiently transfected using Polyfect reagent (QIAGEN) according to the manufacturer's instructions. Cells were cotransfected with 7.2 μ g of OX-2/CD200 cDNA in vector pCEP4 (Invitrogen) and 0.8 μ g of pAdVantage vector (Promega). As a negative control, cells were cotransfected with empty pCEP4 vector plus pAdVantage. 48 hours after transfection, approximately 90% of the cells expressed OX-2/CD200 on their surface as determined by flow cytometry with the scFv-9 antibody.

30

Maturation of dendritic cells/macrophages from blood monocytes

Buffy coats were obtained from the San Diego Blood Bank and primary blood lymphocytes (PBL) were isolated using Ficoll. Cells were adhered for 1 hour in Eagles Minimal Essential Medium (EMEM) containing 2% human serum followed by vigorous washing with PBS. Cells were cultured for 5 days either in the presence of GM-CSF, IL-4 and IFN- γ or M-CSF with or without the addition of lipopolysaccharide (LPS) after 3 days. Matured cells were harvested and irradiated at 2000 RAD using a γ -irradiator (Shepherd Mark I Model 30 irradiator (Cs137)).

10 Mixed lymphocyte reaction

Mixed lymphocyte reactions were set up in 24 well plates using 500,000 dendritic cells/macrophages and 1×10^6 responder cells. Responder cells were T cell enriched lymphocytes purified from peripheral blood using Ficoll. T cells were enriched by incubating the cells for 1 hour in tissue culture flasks and taking the non-adherent cell fraction. 500,000 OX-2/CD200 transfected EBNA cells or CLL cells were added to the macrophages/dendritic cells in the presence or absence of 30 $\mu\text{g/ml}$ anti-CD200 antibody (scFv-9 converted to full IgG) 2-4 hours before the lymphocyte addition. Supernatants were collected after 48 and 68 hours and analyzed for the presence of cytokines.

20 Conversion of scFv-9 to full IgG

Light chain and heavy chain V genes of scFv-9 were amplified by overlap PCR with primers that connect the variable region of each gene with human lambda light chain constant region gene, and human IgG1 heavy chain constant region CH1 gene, respectively. Variable regions of light chain gene and heavy chain gene of scFv-9 were amplified with specific primers and the human lambda light chain constant region gene and the IgG1 heavy chain constant region CH1 gene were separately amplified with specific primers as follows:

30 R9VL-F1 QP: 5' GGC CTC TAG ACA GCC TGT GCT GAC TCA GTC
GCC CTC 3' (SEQ ID NO: 103);

R9VL/hCL2-rev: 5' CGA GGG GGC AGC CTT GGG CTG ACC TGT
GAC GGT CAG CTG GGT C 3' (SEQ ID NO: 104);

R9VL/hCL2-F: 5' GAC CCA GCT GAC CGT CAC AGG TCA GCC
CAA GGC TGC CCC CTC G 3' (SEQ ID NO: 105);

5 R9VH-F1: 5' TCT AAT CTC GAG CAG CAG CAG CTG ATG GAG
TCC G 3' (SEQ ID NO: 106);

R9VH/hCG-rev: 5' GAC CGA TGG GCC CTT GGT GGA GGC TGA
GGA GAC GGT GAC CAG GGT GC 3' (SEQ ID NO: 107);

10 R9VH/hCG-F: 5' GCA CCC TGG TCA CCG TCT CCT CAG CCT CCA
CCA AGG GCC CAT CGG TC 3' (SEQ ID NO: 108);

hCL2-rev : 5' CCA CTG TCA GAG CTC CCG GGT AGA AGT C 3'
(SEQ ID NO: 109);

15 hCG-rev : 5' GTC ACC GGT TCG GGG AAG TAG TC 3'
(SEQ ID NO: 110).

Amplified products were purified and overlap PCR was performed.

20 Final products were digested with Xba I/Sac I (light chain) and Xho I/Pin AI
(heavy chain) and cloned into a human Fab expression vector, PAX243hGL (see
published International Application WO 2004/078937, the disclosure of which is
incorporated herein by this reference). DNA clones were analyzed for PCR errors by
DNA sequencing. The hCMV IE promoter gene was inserted at Not I/ Xho I site (in front
25 of the heavy chain). The vector was digested with Xba I/Pin AI/EcoR I/Nhe I and a 3472
bp fragment containing the light chain plus the hCMV IE promoter and the heavy chain
gene was transferred to an IgG1 expression vector at the Xba I/Pin AI site.

Cytokine analysis

30 The effect of the scFv-9 converted to full IgG on the cytokine profile in the mixed
lymphocyte reaction was determined.

Cytokines such as IL-2, IFN- γ , IL-4, IL-10 and IL-6 found in the tissue culture
supernatant were quantified using ELISA. Matched capture and detection antibody pairs
for each cytokine were obtained from R+D Systems (Minneapolis, MN), and a standard

curve for each cytokine was produced using recombinant human cytokine. Anti-cytokine capture antibody was coated on the plate in PBS at the optimum concentration. After overnight incubation, the plates were washed and blocked for 1 hour with PBS containing 1 % BSA and 5% sucrose. After 3 washes with PBS containing 0.05% Tween, 5 supernatants were added at dilutions of two-fold or ten-fold in PBS containing 1% BSA. Captured cytokines were detected with the appropriate biotinylated anti-cytokine antibody followed by the addition of alkaline phosphatase conjugated streptavidin and SigmaS substrate. Color development was assessed with an ELISA plate reader (Molecular Devices).

10 As shown in Figure 14, the presence of OX-2/CD200 transfected but not untransfected cells resulted in down-regulation of Th1 cytokines such as IL-2 and IFN- γ . Addition of the anti-CD200 antibody at 30 μ g/ml fully restored the Th1 response, indicating that the antibody blocked interaction of OX-2/CD200 with its receptor.

As set forth in Figures 15 and 16, the presence of CLL cells in a mixed 15 lymphocyte reaction resulted in down-regulation of the Th1 response. (Figure 15 shows the results for IL-2; Figure 16 shows the results for IFN- γ). This was not only the case for cells over-expressing OX-2/CD200 (IB, EM, HS, BH), but also for CLL cells that did not overexpress OX-2/CD200 (JR, JG and GB) (the expression levels for these cells are set forth in Figure 11). However, the anti-CD200 antibody only restored the Th1 20 response in cells over-expressing OX-2/CD200, indicating that for patients overexpressing OX-2/CD200, abrogating OX-2/CD200 interaction with its receptor on macrophages was sufficient to restore a Th1 response. In patients that did not overexpress OX-2/CD200, other mechanisms appeared to be involved in down-regulating the Th1 response.

25

Animal Models To Test An Effect Of Anti-CD200 On Tumor Rejection

A model was established in which RAJI lymphoma tumor growth is prevented by the simultaneous injection of PBLs. NOD/SCID mice were injected subcutaneously with 4x10⁶ RAJI cells either in the presence or absence of human PBLs from different donors 30 at 1x10⁶, 5x10⁶ or 10x10⁶ cells. Tumor length and width as well as body weight was determined 3 times a week. Mean +/-SD of tumor volumes for all groups is shown in Figures 17 A and B. Statistical analysis was performed using 2 parametric tests (Student's t-test and Welch's test) and one non-parametric test (the Wilcox test). Results of the statistical analysis are found in Figure 18. RAJI cells form subcutaneous tumors

with acceptable variation. Rejection is dependent on the specific donor and the PBL cell number. 1×10^6 PBLs were insufficient to prevent tumor growth. Donor 2 at 5×10^6 PBLs from day 22-43 and donor 3 at 5×10^6 or 1×10^7 PBLs starting at day 36 significantly reduced tumor growth. Donor 4 is very close to being significant after day 48.

5 To test for an effect of anti-CD200, RAJI cells are stably transfected with CD200. Animals are injected as described in the previous paragraph. In the presence of CD200-transfected cells, tumors grow even in the presence of human PBLs. Anti-CD200 antibody is administered to evaluate tumor rejection in this model.

Also, a liquid tumor model is established. RAJI cells are injected intraperitoneally
10 into NOD/SCID mice. Cells disseminate to bone marrow, spleen, lymph node and other organs resulting in paralysis. Concurrent injection of human PBLs prevents or slows tumor growth. Tumor growth is monitored by assessing the mice for signs of movement impairment and paralysis. Once these signs are observed, mice are sacrificed and the number of tumor cells is assessed in various organs including bone marrow, spleen,
15 lymph nodes and blood by FACS analysis and PCR.

Similar to the subcutaneous model, CD200 transfected cells are injected intraperitoneally. They grow even in the presence of human PBLs. Treatment with anti-CD200 results in tumor rejection or slower tumor growth.

20

EXAMPLE 4

Library construction

A mouse was immunized alternately with baculovirus expressed recombinant CD200 extracellular domain fused to mouse IgG Fc (CD200-Fc) (Orbigen Inc., San Diego, CA) and 293-EBNA cells transiently transfected with a vector containing full
25 length CD200. Total RNA was prepared from mouse spleen using TRI reagent (Molecular Research Center, Inc., Cincinnati, OH) according to the manufacturer's protocol. Messenger RNA (mRNA) was purified using Oligotex (QIAGEN Inc., Valencia, CA) according to the manufacturer's manual. First strand cDNA was synthesized using SuperScript II RTase (Invitrogen Life Technologies, Carlsbad, CA)
30 according to the manufacturer's protocol. First strand cDNA was digested with restriction endonuclease and second strand cDNA was synthesized according to the method fully described in published PCT application WO03/025202A2, published March 27, 2003. Second strand cDNA was cleaned up with PCR purification kit (QIAGEN) and single primer amplification was performed according to the method described in published PCT

application WO03/025202A2, published March 27, 2003. Amplified products were pooled and purified with PCR purification kit. Kappa light chain was digested with Xba I and BspE I, and IgG1 and IgG2a heavy chains were digested with Xho I and Bln I. Digested fragments were purified from the agarose gel using Gel extraction kit
5 (QIAGEN) and cloned into PAX313m/hG vector as described in published PCT application WO/04078937A2 published September 16, 2004.

Library panning

The libraries (IgG1 kappa and IgG2a kappa) were panned on CD200-Fc either
10 directly coated on the microtiter wells (Costar Group, Bethesda, MD) or captured with goat anti-mouse IgG Fc specific antibody (Sigma-Aldrich Corp., St. Louis, MO). For the preparation of library phage, electrocompetent XL1-Blue cells (Stratagene, La Jolla, CA) were electroporated with library DNA and grown in SOC medium for 1 hour and in SB medium for 2 hours with carbenicillin. Phage production was induced with the addition of
15 VCS M13 helper phage (Amersham Biosciences Corp., Piscataway, NJ) and 1 mM IPTG at 30°C overnight. The culture was spun down and phage were precipitated with 4% polyethylene glycol and 3% NaCl. The phage were spun down and resuspended in 1% BSA/PBS containing unrelated antigen, FLJ32028 that is also baculovirus expressed extracellular domain fused to mouse IgG Fc (FLJ32028-Fc) (Orbigen, San Diego), as a
20 soluble competitor. For the panning on directly coated CD200-Fc, four wells were coated with 100 µl of CD200-Fc (5 µg/ml in 0.1 M NaHCO₃ pH8.6) at 4°C overnight. The wells were washed 5 times with phosphate buffered saline (PBS) pH7.0 and blocked with 1% bovine serum albumin (BSA)/PBS at 37°C for 1 hr. For the panning on CD200-Fc captured on goat anti-mouse IgG Fc, four microtiter wells were coated with 100 µl goat
25 anti-mouse IgG Fc (20 µg/ml in PBS) at 4°C overnight. The wells were washed 5 times with PBS and incubated with 100 µl CD200-Fc (20 µg/ml in PBS) for 1 hour at 37°C. The wells were washed 5 times with PBS and blocked with 1% BSA/PBS at 37°C for 1 hour. For both directly-coated and captured methods of panning, the blocker was replaced with the mixture of soluble Fabs obtained from the panning of another library (the library
30 described in Example 3 of PCT application serial No. PCT/US04/17118 filed June 2, 2004 (not yet published), the entire disclosure of which is incorporated herein by this reference) on FLJ32028 to mask epitopes on mouse IgG Fc and the wells were incubated for 30 min at 37°C. These masking Fabs were shown to also bind to CD200-Fc. Library

phage were added on top of the masking Fabs and the wells were incubated for approximately 1.5 hours at 37°C. The unbound phage were washed with PBS with increasing stringency (3 times in the first round, 5 times in the 2nd round and 10 times in the 3rd and the 4th rounds) with 5 minute incubation and pipetting up and down 5 times for each wash. The bound phage were eluted twice with 100 µl 0.1 M HCl with 1 mg/ml BSA, pH2.2 and neutralized with 2 M Tris Base pH 11.5. The freshly grown ER2738 cells were infected with eluted phage and titrated onto LB agarose plates containing carbenicillin and glucose. The remaining phage were propagated overnight at 30-37°C with the addition of VCS M13 helper phage and 1 mM IPTG for the next round of panning.

Library screening

Ninety five colonies from round 3 and 4 titration plates were grown in 1 ml SB containing 12.5 µg/ml tetracycline and 50 µg/ml carbenicillin for approximately 6 hours at 37°C. VCS M13 helper phage were added and the culture was incubated for 2 hours at 37°C. 1 mM IPTG and 70 µg/ml kanamycin were added and Fab-phage production was induced at 30°C overnight. Microtiter wells were coated with 50 µl of rabbit anti-mouse IgG F(ab')₂ (4 µg/ml in PBS), CD200-Fc (4 µg/ml in 0.1 M NaHCO₃ pH8.6), or FLJ32028-Fc (4 µg/ml in 0.1 M NaHCO₃ pH8.6) at 4°C overnight. The wells were washed 3 times with PBS and blocked with 100 µl 1% BSA/PBS for 1 hour at 37°C. The culture was spun down. The blocker was replaced with the culture supernatant containing Fab-phage and the wells were incubated for 1.5~2 hours at 37°C. The remaining Fab-phage was stored at -80°C for flow cytometry. The plates were washed 3 times with PBS and the binding was detected with 50 µl alkaline phosphatase (AP)-conjugated goat anti-mouse IgG F(ab')₂ antibody (Pierce)(1:500 in 1% BSA/PBS) for 1 hr at 37°C. The plates were washed 3 times with PBS and developed with AP substrate (Sigma-Aldrich) in pNPP buffer. Almost all of the clones from round 3 were already specifically positive to CD200 (Figs. 19A-D). Clones were also screened by high throughput flow cytometry analysis. One hundred microliters of 293 cells transiently transfected with CD200 (1 x 10⁵ cells) were aliquoted into a 96 well plate (Costar). Fifty microliters Fab-phage was added to the cells and mixed by pipetting and incubated on ice for 30 minutes. The cells were washed twice with 1% BSA/PBS containing 0.01% NaN₃. The cells were resuspended in 100 µl PE-conjugated goat anti-mouse IgG antibody (Sigma-Aldrich) in 1% BSA/PBS

containing 0.01% NaN₃ and incubated on ice for 30 minutes. The cells were washed twice with 1% BSA/PBS containing 0.01% NaN₃ and resuspended in 200 µl 1% paraformaldehyde in PBS. Representative clones showing positive binding to CD200 expressing cells are shown in Figs. 20A-D.

- 5 DNA sequences were analyzed and deduced amino acid sequences of the heavy chain were grouped according to the complementarity determining region 3 (CDR3) (Figs. 21A, B). They were divided into 17 groups.

Fluorescent bead assay

- 10 23 clones were selected for further analysis. They were cG2aR3B5, dG1R3A5, cG2aR3A2, dG2aR3B2, dG1R3A1, cG2aR3A1, cG2aR3B, dG1R3B, cG1R3A, cG1R3A, cG1R3A1, dG1R3B, dG1R3B, cG1R3C, dG2aR3C, dG2aR3A1, cG2aR3B, cG2aR3B, dG1R3B, cG2aR3B, cG2aR3C, dG1R3H, and dG2aR3A6. DNA of selected Fabs was digested with Spe I/Nhe I for gene III removal and soluble Fab expression and
- 15 purification. The purified Fabs were evaluated for their ability to block the interaction of CD200 with its receptor (CD200R) in a fluorescent bead assay. TransFluoSpheres carboxylate-modified microspheres (488/645) (Molecular Probes Invitrogen Detection Technologies, Eugene, OR) were coated with streptavidin followed by a biotin-labeled anti-human Fc antibody and baculovirus-produced CD200-Fc protein. 293 cells were
- 20 transiently transfected with CD200R. Cell surface expression was confirmed by FACS analysis. 1 million CD200-coated beads were pre-incubated with various amounts of anti-CD200 Fabs or chimeric IgG for 10 minutes before the addition of 50,000 CD200R transfected cells. After a 30 minute incubation at 37°C, the cells were washed in Tris buffer containing 1% BSA and analyzed using a FACS Calibur. Fabs c1A10, c2aB7, and
- 25 d1A5 showed the best blocking of CD200 and CD200R interaction at 6.7 µg/ml of Fab (Fig. 22). These clones are referred to as cG1R3A10, cG2aR3B7 and dG1R3A5, respectively in Figs. 21A and/or B.

Chimerization and IgG conversion

- 30 Six antibodies were selected for chimerization and IgG conversion. (See Fig. 23.) They were c1A10 (cG1R3A10), c2aA10 (cG2aR3A10), c2aB7 (cG2aR3B7), d1A5 (dG1R3A5), d1B5 (dG1R3B5), and d1B10 (dG1R3B10). For the chimerization, overlap PCR was performed to connect mouse kappa chain variable region and human kappa

chain constant region. Mouse heavy chain variable region was amplified with a 3' primer that contains a partial human IgG1 constant region and Apa I site for cloning. Amplified kappa chain fragments and heavy chain fragments were cloned into PAX243hGK vector (see published International Application WO 2004/078937) that contains human IgG1
5 constant region at Xba I/Not I for kappa light chain and Xho I/Apa I for heavy chain fragment. Binding of chimeric Fab to CD200 was confirmed by ELISA and flow cytometry. These chimeric Fabs were converted into IgG by insertion of human cytomegalovirus immediate early promoter (hCMV IE Pro) sequence for the heavy chain expression at Not I/Xho I, then the transfer of the light chain and heavy chain into
10 human IgG1 expression vector at Xba I/Pin AI sites. This vector has an additional hCMV IE Pro sequence upstream Xba I site for the light chain expression in mammalian cells. The DNA sequences were confirmed and maxi prep DNA was prepared using HiSpeed Maxi prep columns (QIAGEN) for mammalian cell transfection. Transient transfection was performed in 293-EBNA cells using Effectene (QIAGEN) according to the
15 manufacturer's protocol with the addition of pAdVantage vector (Promega US, Madison, WI). Stable cell line transfection was performed in NS0 cells using Effectene according to the manufacturer's protocol. After a small scale transient transfection, culture supernatant for each antibody was tested by ELISA (Fig. 24). After a large scale transient transfection, each IgG was purified from the culture supernatant by anti-human IgG
20 F(ab')₂ affinity column using FPLC (Amersham Biosciences).

The purified IgG were tested in a bead inhibition assay as described for the Fabs. All antibodies directed against CD200 blocked the receptor ligand interaction very well as shown below in Fig. 25.

25 Mixed lymphocyte reaction

Whether blocking of CD200 interaction with its receptor also prevents the cytokine shift from Th1 to Th2 observed in mixed lymphocytes reactions in the presence of CD200 was evaluated. Buffy coats were obtained from the San Diego Blood Bank and primary blood lymphocytes (PBL) were isolated using Histopaque (Sigma-Aldrich). Cells
30 were adhered for 1 h in EMEM containing 2% human serum followed by vigorous washing with PBS. Cells were cultured for 5 days in the presence of GM-CSF, IL-4 and IFN- γ . Matured cells were harvested and irradiated at 2000 RAD using a γ -irradiator (University of California San Diego). Mixed lymphocyte reactions were set up in 24 well plates using 500,000 dendritic cells and 1×10^6 responder cells. Responder cells were T

cell enriched lymphocytes purified from peripheral blood using Histopaque. T cells were enriched by incubating the cells for 1 hour in tissue culture flasks and taking the non-adherent cell fraction. Five hundred thousand CD200 expressing primary irradiated CLL cells were added to the dendritic cells in the presence or absence of various amounts of anti-CD200 antibodies 2-4 hours before the lymphocyte addition. Supernatants were collected after 48 and 68 hours and cytokines such as IL-2, IFN- γ , IL-4, IL-10 and IL-6 were quantified using ELISA. Matched capture and detection antibody pairs for each cytokine were obtained from R+D Systems (Minneapolis), and a standard curve for each cytokine was produced using recombinant human cytokine. Anti-cytokine capture antibody was coated on the plate in PBS at the optimum concentration. After overnight incubation, the plates were washed and blocked for 1h with PBS containing 1% BSA and 5% sucrose. After 3 washes with PBS containing 0.05% Tween, supernatants were added at the indicated dilutions in PBS containing 1% BSA. Captured cytokines were detected with the appropriate biotinylated anti-cytokine antibody followed by the addition of alkaline phosphatase conjugated streptavidin and SigmaS substrate. Color development was assessed with an ELISA plate reader (Molecular Devices Corp., Sunnyvale, CA). As shown in Figs. 26A and B, the presence of CLL cells completely abrogated IFN-gamma and most of IL-2 production observed in the mixed lymphocyte reaction. Presence of any of the antibodies allowed for production of these Th1 cytokines (Figs. 26A and B). In contrast, IL-10 production was downregulated in the presence of the antibodies. (See Fig. 26C.)

Antibody-dependent cell-mediated cytotoxicity assay

Furthermore, the six chimeric mouse anti-CD200 antibodies were evaluated for their ability to kill CD200 expressing tumor cells in an antibody-dependent cell-mediated cytotoxicity assay (ADCC). 293-EBNA cells transfected with CD200 were labeled with 100 μ Ci/million cells in 0.5ml medium for 1hr at 37°C. After 3 washes, cells were counted, resuspended in medium (RPMI supplemented with 10% human AB serum) at 0.2 million/ml and 50 μ l (10,000 cells/well) was dispensed in triplicate into a 96 well round bottom plate. 20 μ l of anti-CD200 antibodies were dispensed into each well so as to achieve a final concentration of 20 μ g/ml. Peripheral blood mononuclear cells (effector cells) were isolated on a Ficoll gradient, red blood cells were lysed with ammonium chloride, washed and resuspended in culture medium and 50 μ l of cells were dispensed

into each well. The assay plates were spun (1,500 rpm/5 minutes/low brake) and transferred to the cell culture incubator. After 4 hours, assay plates were spun as before. 36 μ l of the supernatants were transferred to pico plates and mixed with 250 μ l microscint-20 cocktail, and placed on the orbital shaker for 2 minutes and read on a Top
5 count. As illustrated in the Fig. 27, all of the mouse chimeric CD200 antibodies produced similar levels of lysis when cultured with CD200 positive cells. No lysis was observed with CD200 negative cells. In addition, the extent of lysis was statistically significant ($p < 0.05$) when compared to isotype control antibody, d2A6 (anti-FLJ32028 antibody).

10

EXAMPLE 5

Raji/PBL Model

NOD.CB17-Prkdc<scid> mice (Jackson Laboratory) were injected with 200 μ l RPMI containing 4×10^6 RAJI cells (ATCC) s.c. along with 0, 1, 5 or 10 million PBLs.
15 Nine or ten mice were included per group. PBLs were isolated from 250 ml whole ammonium chloride. Tumor growth was monitored three times a week by measuring length and width with a caliper. Tumor volume was calculated based on length x width x width/2.

Differences between the groups that were injected with PBLs compared to the
20 group that received tumor cells only were analyzed by 2-tailed unpaired Student's t-test. Significant differences were observed in the groups that received 5 or 10 million PBLs, but not in the group that received 1 million PBLs from Day 32 on. The data shown in Figure 28 are a representative example of 10 experiments using different PBL donors.

25 Namalwa PBL model

NOD.CB17-Prkdc<scid> mice (Jackson Laboratory, Bar Harbor, Maine) were injected with 200 μ l RPMI containing 4×10^6 Namalwa cells (ATCC) s.c. along with 0, 2 or 10 million PBLs. 9-10 mice were included per group. PBLs were isolated from 250 ml whole blood on a histopaque gradient followed by red blood cell lysis using 0.9%
30 ammonium chloride. Tumor growth was monitored three times a week by measuring length and width with a caliper. Tumor volume was calculated based on length x width x width/2.

Figure 29 shows differences between the groups that were injected with PBLs compared to the group that received tumor cells only analyzed by 2-tailed unpaired Student's t-test. Significant differences were observed in the groups that received 10 million PBLs for both donors, but not in the group that received 2 million PBLs from day 5 8 on.

Creation of stable CD200-expressing cell lines

Stable CD200-expressing Raji and Namalwa cell lines were generated using the Virapower Lentiviral Expression System (Invitrogen, Carlsbad, CA). A CD200 cDNA was isolated from primary CLL cells by RT-PCR using forward primer 10 5'-GACAAGCTTGCAAGGATGGAGAGGCTGGTGA-3' (SEQ ID NO: 212) and reverse primer 5'-GACGGATCCGCCCTTTTCCTCCTGCTTTTCTC-3' (SEQ ID NO: 213). The PCR product was cloned into the Gateway entry vector pCR8/GW/TOPO-TA and individual clones were sequenced. Clones with the correct sequence were recombined in both the sense and antisense orientations into the lentiviral vectors 15 pLenti6/V5/DEST and pLenti6/UbC/V5/DEST using Gateway technology (Invitrogen, Carlsbad, CA). The primary difference between these two vectors is the promoter used to drive CD200 expression: pLenti6/V5/DEST contains the human CMV immediate early promoter, whereas pLenti6/UbC/V5/DEST contains the human ubiquitin C promoter.

High-titer, VSV-G pseudotyped lentiviral stocks were produced by transient 20 cotransfection of 293-FT cells as recommended by the manufacturer. Raji or Namalwa cells were transduced by resuspending 10^6 cells in 1ml of growth medium containing 12 μ g/ml Polybrene and adding 1ml of lentiviral stock. After incubating the cells overnight at 37°C, the medium containing virus was removed and replaced with 4ml of fresh medium. Two days later, the infected cells were analyzed for CD200 expression by 25 flow cytometry. In all experiments, $\geq 70\%$ of the cells were CD200⁺, whereas CD200 was undetectable in the parental cell lines and in cells transduced with the negative control (antisense CD200) viruses.

To isolate clonal cell lines that overexpress CD200, the infected cells were selected with blasticidin for 13 days. The concentrations of blasticidin used were 6 μ g/ml 30 for Raji cells or 2 μ g/ml for Namalwa cells. Stable clones were then isolated by limiting dilution of the blasticidin-resistant cells into 96-well plates. Clones were screened in 96-

well format by flow cytometry using PE-conjugated Mouse Anti-Human CD200 (clone MRC OX104, Serotec) and a BD FACSCalibur equipped with a High Throughput Sampler. After screening a total of 2000 Raji and 2000 Namalwa clones, those clones with the highest CD200 expression were expanded for further characterization using
5 conventional techniques.

Immunosuppressive effect of CD200 in the RAJI/PBL model

In accordance with the methods disclosed, it has been demonstrated that CD200 is upregulated on CLL cells. Upregulation of this molecule might potentially be
10 immunosuppressive. To test whether cancer cells expressing CD200 prevent the immune system from eradicating the cancer cells, RAJI cells, that normally do not express CD200, were infected with a lentivirus vector system encoding for CD200 as described above. RAJI clones stably expressing CD200 were selected. As a control to ensure that there was no effect of vector infection, clones expressing a reversed, nonfunctional form of CD200
15 (CD200rev) were also selected. RAJI cells expressing CD200, CD200REV or the parental RAJI cells were injected subcutaneously into NOD.CB17-Prkdc<scid> mice. The following groups were included in the study:

group 1: 4×10^6 RAJI s.c.; 9 mice

20 This group was needed to ensure that the lentivirus transduced cells show similar growth as the parent cells.

group 2: 4×10^6 RAJICD200 s.c.; 9 mice

This group was needed to ensure that the CD200 transduced cells showed similar growth as the parent cells. Also, this group will give the maximum
25 tumor growth. Group 3 and group 4 were compared to this group.

group 3: 4×10^6 RAJICD200 + 5×10^6 PBL s.c.; 9 mice

This PBL number has been shown to reduce tumor growth in some mice in previous experiments. Rejection is not as strong as with 10 million cells, but in order to determine whether CD200 can affect only a certain number of cells,
30 and 5×10^6 is the minimum amount of PBLs which can be used to get rejection; rejection should be prevented by the presence of CD200.

group 4: 4×10^6 RAJICD200 + 10×10^6 PBL s.c.; 8 mice

This is the optimum number of PBLs to see rejection in the RAJI/PBL model.

The design was that CD200 expression would prevent this rejection.

group 5: 4×10^6 RAJICD200rev s.c.; 9 mice

This group was needed to ensure that the lentivirus transduced cells show similar growth as the parent cells.

5 group 6: 4×10^6 RAJICD200rev + 2×10^6 PBL s.c.; 9 mice

This number of PBLs should not result in strong rejection or reduction of tumor growth. This is the positive control for group 3 and group 4 (maximum expected tumor growth). If there is no rejection in this group, then the donor PBLs were hyperactivated to start with which could explain lack of an effect by CD200.

10 group 7: 4×10^6 RAJICD200rev + 5×10^6 PBL s.c.; 9 mice

Controls that any observed effects in the CD200 group 3 are really related to CD200 and not to lentivirus transduction.

group 8: 4×10^6 RAJICD200rev + 10×10^6 PBL s.c.; 8 mice

15 Controls that any observed effects in the CD200 group 4 are really related to CD200 and not to lentivirus transduction.

20 Animals were sacrificed at day 38 based on tumors reaching a size above acceptable limits. Tumors from 4 animals/group were removed. Two tumors/group were frozen in OCT, the other 2 were used to isolate cells and analyze by FACS for CD200 expression. Figures 30(a-c) demonstrate the results for this study.

Although RAJICD200 cells appeared to grow somewhat more slowly, the growth difference between transduced and parental cells did not reach statistical significance as shown in Figure 30(a).

25 The presence of PBLs slowed tumor growth by up to 84% when 5 or 10×10^6 PBLs were injected, although generally 10×10^6 PBL resulted in a stronger reduction over time compared to 5×10^6 PBL. The reduction in growth compared to the parent tumor cells was significant from day 20 on. 2×10^6 PBL resulted in a significant tumor growth reduction from d22-d29, but that reduction was overcome at later timepoints (Fig. 30(b)).

30 This study indicated that this particular donor rejects RAJI tumor cells very strongly.

Tumor growth in the groups that received CD200 expressing RAJI cells and PBLs was not significantly different from the tumor growth in the group that only received RAJI cells although mice that received 10×10^6 PBL showed a trend of reduced tumor growth, but the difference reached no statistical significance at any time point

after tumors reach 100 mm³. Every mouse in the group that received RAJI cells and 5x10⁶ PBL developed a second tumor, some mice as early as d7, while this was not observed in any other group. For analysis, the second tumor was added to the first tumor and the combined size is shown in Fig. 30(c).

5 These results indicate that CD200 expression on tumor cells does indeed prevent the immune system from slowing tumor growth. Also, this study demonstrates the usefulness of the RAJI/PBL model to assess immunosuppressive compounds or molecules.

10 Immunosuppressive effect of CD200 in the Namalwa/PBL model

To evaluate whether the effects seen in the RAJI/PBL model can also be observed in other tumor models, Namalwa tumor cells were also infected with the lentivirusCD200 system and stable clones selected. As a control to ensure that there is no effect of vector infection, clones expressing a reversed, nonfunctional form of CD200 (CD200rev) were also selected. NOD.CB17-Prkdc<scid> mice were injected according to the following scheme as shown in Figures 31(a)-(d):

group 1: 4 x 10⁶ Namalwa s.c.; 9 mice

20 This group was needed to ensure that the lentivirus transduced cells show similar growth as the parent cells.

group 2: 4 x 10⁶ Namalwa CD200 (1D12Ub) s.c.; 9 mice

This group was needed to ensure that the CD200 transduced cells showed similar growth as the parent cells. Also, this group will give the maximum tumor growth. Group 3 and group 4 were compared to this group.

25 group 3: 4 x 10⁶ Namalwa CD200 (1D12Ub)+ 5x10⁶ PBL s.c.; 9 mice

30 This PBL number has been shown to reduce tumor growth in some mice in previous experiments. Rejection is not as strong as with 10 million cells, but if CD200 can affect only a certain number of cells, and this is the minimum PBL we can use to get rejection; rejection will be prevented by the presence of CD200.

group 4: 4 x 10⁶ Namalwa CD200 (1D12Ub) + 10x10⁶ PBL s.c.; 8 mice

This is the optimum number of PBLs to see rejection in the Namalwa /PBL model. This was done to show that CD200 expression prevents this rejection.

group 5: 4×10^6 Namalwa CD200rev (C5UbreV) s.c.; 9 mice

This group was needed to ensure that the lentivirus transduced cells showed similar growth as the parent cells.

group 6: 4×10^6 Namalwa CD200rev + 2×10^6 PBL s.c.; 9 mice

5 This number of PBLs should not result in strong rejection or reduction of tumor growth. This is the positive control for group 3 and group 4 (maximum expected tumor growth). If there is no rejection in this group, then the donor PBLs were hyperactivated to start with which would explain lack of an effect by CD200.

10 group 7: 4×10^6 Namalwa CD200rev + 5×10^6 PBL s.c.; 9 mice

This group was needed as a control to detect any effects in the CD200 group 3 that were related to CD200 and not to lentivirus transduction.

group 8: 4×10^6 Namalwa CD200rev + 10×10^6 PBL s.c.; 8 mice

15 This group was needed as a control to detect any effects in the CD200 group 4 that were related to CD200 and not to lentivirus transduction.

Tumor length and width were assessed three times/week.

All tumor cells resulted in rapid tumor growth. There was no significant growth difference between transduced and parental cells. The tumor grows more aggressively
20 than previously observed as shown in Figure 31(a).

Figure 31(b) shows the presence of PBLs slows tumor growth by about 50%. This trend was observed from d12 on. The differences of the PBL treated group versus groups that received only tumor cells are statistically significant (as determined by 2-tailed Student's t-test) at d17 and d19 when 2 million or 10 million PBLs were injected.
25 Injection of 5 million PBLs resulted in tumor growth reduction, but did not reach significance.

Tumor growth in the groups that received CD200 expressing Namalwa cells and PBLs was similar to the tumor growth in the group that only received Namalwa cells (Figure 31(c)).

30 These data confirm that CD200 expression on tumor cells prevents slowing of tumor growth by the human immune system.

Blockage of the immunosuppressive effect of CD200 in the RAJI/PBL model by anti-CD200 antibodies

To evaluate whether anti-CD200 antibodies can block the immunosuppressive effect of CD200 expressed on tumor cells, RAJI cells transduced with CD200 were injected s.c. into NOD.CB17-Prkdc<scid> mice, and the ability of PBLs to reduce tumor growth in the presence or absence of chimeric anti-CD200 antibodies d1B5 and c2aB7 or a control antibody that does not bind tumor cells (alxn4100) was assessed. Antibodies were administered initially at 10 mg/kg or 2.5 mg/kg with the tumor cells, and then at concentrations indicated below twice/week i.v.. The following groups were set up:

10

1. 4×10^6 RAJICD200; 10 mice
2. 4×10^6 RAJICD200 + 5×10^6 PBL; 10 mice
3. 4×10^6 RAJICD200 + 5×10^6 PBL + 100 mg/kg d1B5; 12 mice
4. 4×10^6 RAJICD200 + 5×10^6 PBL + 20 mg/kg d1B5; 9 mice
- 15 5. 4×10^6 RAJICD200 + 5×10^6 PBL + 100 mg/kg c2aB7; 11 mice
6. 4×10^6 RAJICD200 + 5×10^6 PBL + 20 mg/kg c2aB7; 9 mice
7. 4×10^6 RAJICD200 + 5×10^6 PBL + 100 mg/kg alxn4100; 9 mice

Tumor length and width was measured 3 times a week, and the tumor volume was calculated by tumor length*width*width/2. Figure 33 shows that as expected, CD200 expression on the tumor cells prevented the immune cells from reducing tumor growth. However, addition of anti-CD200 antibodies reduced the tumor volume by 50-75 %. The reduction in growth by the antibodies was statistically significant as determined by Student's t-test and Mann Whitney test from day 18 on through the end of the study. In contrast, treatment with the control antibody did not reduce the tumor growth. These data demonstrate the usefulness of anti-CD200 in anti-cancer treatment. Also, this study demonstrates the usefulness of the RAJI/PBL model to assess immunomodulatory therapeutics.

30 Blockage of the immunosuppressive effect of CD200 in the Namalwa/PBL model by anti-CD200 antibodies

To evaluate whether the effect seen with the anti-CD200 antibodies in the RAJI/PBL model can also be observed in other tumor models, RAJI cells transduced with CD200 were injected s.c. into NOD.CB17-Prkdc<scid> mice, and the ability of PBLs to reduce tumor growth in the presence or absence of chimeric anti-CD200 antibodies d1B5 and c2aB7 or a control antibody that does not bind tumor cells (alxn4100) was assessed. Antibodies at concentrations indicated below were administered initially with the tumor

35

cells, and then twice/week s.c. within 0.5 cm of the tumor. The following groups were set up (10 mice/group unless indicated otherwise):

1. 4×10^6 NamalwaCD200
- 5 2. 4×10^6 NamalwaCD200 + 5×10^6 PBL; 12 mice
3. 4×10^6 NamalwaCD200 + 5×10^6 PBL + 10 mg/kg d1B5; 12 mice
4. 4×10^6 NamalwaCD200 + 5×10^6 PBL + 2.5 mg/kg d1B5
5. 4×10^6 NamalwaCD200 + 5×10^6 PBL + 10 mg/kg c2aB7; 12 mice
6. 4×10^6 NamalwaCD200 + 5×10^6 PBL + 2.5 mg/kg c2aB7
- 10 7. 4×10^6 NamalwaCD200 + 5×10^6 PBL + 10 mg/kg alxn4100

Tumor length and width was measured 3 times a week, and the tumor volume was calculated by tumor length*width*width/2. Figure 34 shows that as expected, CD200 expression on the tumor cells prevented the immune cells from reducing tumor growth. However, addition of anti-CD200 antibodies reduced the tumor volume by up to 97 %. The reduction in growth by the antibodies was statistically significant as determined by Student's t-test and Mann Whitney test from day 12 on through the end of the study. In contrast, treatment with the control antibody did not reduce the tumor growth. These data confirm the usefulness of anti-CD200 in anti-cancer treatment, and the use of the tumor/PBL models in assessing immunomodulatory therapeutics.

Detection of a potential immune-enhancing effect of compounds in the RAJI/PBL model

T cells isolated from PBLs using CD3 columns (Miltenyi) were incubated in vitro with an ascites preparation of AZND1 (anti-DC-SIGN antibody) or an ascites preparation of a control antibody (BB5). NOD.CB17-Prkdc<scid> mice were injected according to the following scheme:

- Group 6: 8 mice, 4×10^6 RAJI + 0.8×10^6 AZND1-potentiated T cells + 2×10^6 PBL s.c.
- Group 5: 10 mice, 4×10^6 RAJI s.c.
- 30 Group 4: 8 mice, 4×10^6 RAJI + 8×10^6 PBL s.c.
- Group 3: 8 mice, 4×10^6 RAJI + 0.8×10^6 fresh T cells s.c.
- Group 2: 8 mice, 4×10^6 RAJI + 0.8×10^6 AZND1-potentiated T cells s.c.
- Group 1: 8 mice, 4×10^6 RAJI + 0.8×10^6 BB5.1-potentiated T cells s.c.

Tumor length and width were measured 3 times/week.

While T cells incubated with the negative control BB5 did not reduce tumor growth, AZND1 treated T cells did reduce tumor growth significantly (See, Figure 32).

5 These results demonstrate that the RAJI PBL model can be used to assess efficacy of immune-enhancing compounds.

EXAMPLE 6

Determination of CD200 Upregulation in CLL Patients

10 Lymphocytes from 15 CLL patients were stained with FITC-conjugated anti-CD5 (e-bioscience), APC-conjugated anti-CD19 (e-bioscience) and PE-conjugated anti-CD200 (Serotec). Lymphocytes from healthy donors were stained accordingly. CD200 expression on CD5+CD19+ cells was determined. As shown in the table below, although the level of CD200 expression varied among CLL patient samples, all CLL samples showed elevated levels (1.6-4-fold range) higher CD200 expression compared to CD200
 15 expression on normal B cells. The CLL patients showing elevated levels of CD200 expression are selected for anti-CD200 treatment in accordance with the methods described herein.

FACS analysis of CD200 expression
 on B-CLL cells in comparison to normal B cells

20

CLL sample		Healthy donor	
Donor ID	B-CLL CD200 (GMFI)	Normal B CD200 (GMFI)	Ratio(CLL/normal B)
RC011731	93	58	1.6
RF020934	659	185	3.6
JA073031	334	64	5.2
GR011846	156	64	2.4
BB101735	420	95	4.4
DM6988172	290	97	2.9
MR8074020	403	97	4.2
CB8267677	300	97	3.1
GB1325248	178	77(7)	2.3
VN7029373	154	77(7)	2.0
DG8942820	146	77(7)	1.9
MM8451869	237	77(7)	3.1

JR4539931	215	77(7)	2.8
HS6787771	305	77(7)	4.0
VB040439	123	41	3.0
			MEAN= 3.1
			STDEV= 1.0

REFERENCES

The following references are incorporated herein by reference to more fully describe the state of the art to which the present invention pertains. Any inconsistency between these publications below or those incorporated by reference above and the present disclosure shall be resolved in favor of the present disclosure.

- 1) Agarwal, et al., (2003). Disregulated expression of the Th2 cytokine gene in patients with intraoral squamous cell carcinoma. *Immunol Invest* 32:17-30.
- 2) Almasri, NM et al. (1992). *Am J Hemato* 140 259-263.
- 10 3) Contasta, et al., (2003). Passage from normal mucosa to adenoma and colon cancer: alteration of normal sCD30 mechanisms regulating TH1/TH2 cell functions. *Cancer Biother Radiopharm* 18:549-557.
- 4) Gorczynski, et al., (1998). Increased expression of the novel molecule OX-2 is involved in prolongation of murine renal allograft survival. *Transplantation* 65:1106-15 1114.
- 5) Gorczynski, et al., (2001). Evidence of a role for CD200 in regulation of immune rejection of leukaemic tumour cells in C57BL/6 mice. *Clin Exp Immunol* 126:220-229.
- 6) Hainsworth, JD (2000). *Oncologist* 2000;5(5):376-84.
- 7) Inagawa, et al., (1998). Mechanisms by which chemotherapeutic agents augment the antitumor effects of tumor necrosis factor: involvement of the pattern shift of cytokines from Th2 to Th1 in tumor lesions. *Anticancer Res* 18:3957-3964.
- 20 8) Ito, et al., (1999). Lung carcinoma: analysis of T helper type 1 and 2 cells and T cytotoxic type 1 and 2 cells by intracellular cytokine detection with flow cytometry. *Cancer* 85:2359-2367.
- 25 9) Kiani, et al., (2003). Normal intrinsic Th1/Th2 balance in patients with chronic phase chronic myeloid leukemia not treated with interferon-alpha or imatinib. *Haematologica* 88:754-761.

- 10) Lauerova, et al., (2002). Malignant melanoma associates with Th1/Th2 imbalance that coincides with disease progression and immunotherapy response. *Neoplasma* 49:159-166.
- 11) Maggio, et al., (2002). Chemokines, cytokines and their receptors in Hodgkin's lymphoma cell lines and tissues. *Ann Oncol* 13 Suppl 1:52-56.
- 12) Nilsson, K (1992). *Burn Cell*. 5(1):25-41.
- 13) Podhorecka, et al., (2002). T type 1/type 2 subsets balance in B-cell chronic lymphocytic leukemia--the three-color flow cytometry analysis. *Leuk Res* 26:657-660.
- 14) Pu, QQ and Bezwoda, W (2000). *Anticancer Res*. 20(4):2569- 78.
- 10 15) Smyth, et al., (2003). Renal cell carcinoma induces prostaglandin E2 and T-helper type 2 cytokine production in peripheral blood mononuclear cells. *Ann Surg Oncol* 10:455-462.
- 16) Tatsumi, et al., (2002). Disease-associated bias in T helper type 1 (Th1)/Th2 CD4(+) T cell responses against MAGE-6 in HLA-DRB10401(+) patients with renal cell carcinoma or melanoma. *J Exp Med* 196:619-628.
- 15 17) Walls A Vet al. (1989). *Int. J Cancer* 44:846-853.
- 18) Winter, et al., (2003). Tumour-induced polarization of tumour vaccine-draining lymph node T cells to a type 1 cytokine profile predicts inherent strong immunogenicity of the tumour and correlates with therapeutic efficacy in adoptive transfer studies.
- 20 Immunology 108:409-419.

It will be understood that various modifications may be made to the embodiments disclosed herein. For example, as those skilled in the art will appreciate, the specific sequences described herein can be altered slightly without necessarily adversely affecting the functionality of the polypeptide, antibody or antibody fragment used in binding OX-2/CD200. For instance, substitutions of single or multiple amino acids in the antibody sequence can frequently be made without destroying the functionality of the antibody or fragment. Thus, it should be understood that polypeptides or antibodies having a degree of identity greater than 70% to the specific antibodies described herein are within the scope of this disclosure. In particularly useful embodiments, antibodies having an identity greater than about 80% to the specific antibodies described herein are contemplated. In other useful embodiments, antibodies having an identity greater than about 90% to the specific antibodies described herein are contemplated. Therefore, the above description should not be construed as limiting, but merely as exemplifications of

preferred embodiments. Those skilled in the art will envision other modifications within the scope and spirit of this disclosure.

Claims:

1. A method of treating cancer comprising:
administering to a subject afflicted with cancer a therapeutic composition that i) interferes with the interaction between CD200 and a CD200 receptor thereby
5 inhibiting the immune suppressing effect of CD200 and ii) kills cancer cells using a polypeptide fusion molecule that includes a portion that binds to OX-2/CD200 or an OX-2/CD200 receptor.
2. A method as in claim 1 wherein the step of administering a therapeutic
10 composition comprises administering to the subject an antibody that binds to OX-2/CD200.
3. A method as in claim 1 wherein the step of administering a therapeutic
composition comprises administering to the subject a monoclonal antibody that binds to
15 OX-2/CD200.
4. A method as in claim 1 wherein the step of administering a therapeutic
composition comprises administering to the subject an antibody that binds to an OX-
2/CD200 receptor.
20
5. A method as in claim 1 wherein the step of administering a therapeutic
composition comprises administering to the subject a monoclonal antibody that binds to
an OX-2/CD200 receptor.
- 25 6. A method as in claim 1 wherein the subject afflicted with cancer is a CLL
patient.
7. A method as in claim 1 wherein the fusion molecule comprises a toxin.
- 30 8. A method as in claim 1 wherein the fusion molecule comprises a high
energy radiation emitter.
9. A method as in claim 1 wherein the fusion molecule comprises a cytokine
or chemokine that enhances cytotoxic T cell or NK cell activity.

10. A method as in claim 1 wherein the fusion molecule comprises a chemokine that attracts T cells.

11. A method of treating cancer comprising administering an antibody that
5 binds to CD200 wherein a) the interaction between CD200 and its receptor is blocked and
b) cancer cells expressing CD200 are killed.

12. A method of treating cancer comprising administering an antibody that
10 binds CD200 wherein a) the interaction between CD200 and its receptor is blocked and b)
cytotoxic T cell or NK cell activity against the cancer is enhanced.

13. A method of treating cancer as in claim 12 wherein enhancement of the
cytotoxic T cell or NK cell activity is achieved by an antibody that binds CD200 fused
with a cytokine selected from the group consisting of IL-2, IL-12, IL-18, IL-13, and IL-5.

15

14. A method of treating cancer comprising and administering an antibody that
binds CD200 wherein a) the interaction between CD200 and its receptor is blocked and b)
T cells are attracted to the tumor cells.

15. A method of treating cancer as in claim 14 wherein T cell attraction is
20 achieved by an antibody that binds CD200 fused with a chemokine selected from the
group consisting of MIG, IP-10 and I-TAC.

16. A fusion molecule comprising:
25 a first portion that targets cells bearing the OX-2/CD200 antigen; and
a second portion that promotes the death of cells.

17. A fusion molecule as in claim 16 wherein the first portion comprises an
antibody that binds to OX-2/CD200.

30

18. A fusion molecule as in claim 16 wherein the first portion comprises a
monoclonal antibody that binds to OX-2/CD200.

19. A method of treating cancer comprising:
administering to a cancer patient an antibody that (i) interferes with the interaction
of CD200 and a CD200 receptor, thereby inhibiting the immune suppressing effect of
CD200 and (ii) kills cancer cells through complement-mediated cellular cytotoxicity or
5 antibody-dependent cellular cytotoxicity.

20. An antibody that (i) interferes with the interaction of CD200 and a CD200
receptor, thereby inhibiting the immune suppressing effect of CD200 and (ii) kills cancer
cells through complement-mediated cellular cytotoxicity or antibody-dependent cellular
10 cytotoxicity.

21. A composition comprising an antibody of claim 20 and a pharmaceutically
acceptable carrier.

15 22. A method as in claim 1 wherein the step of administering a therapeutic
composition comprises administering an antibody comprising a light chain CDR1 region
having a sequence selected from the group consisting of SEQ ID NOS: 5, 12 and 13.

20 23. A method as in claim 1 wherein the step of administering a therapeutic
composition comprises administering an antibody comprising a light chain CDR2 region
having a sequence selected from the group consisting of SEQ ID NOS: 21 and 23.

24. A method as in claim 1 wherein the step of administering a therapeutic
composition comprises administering an antibody comprising a light chain CDR3 region
25 having a sequence selected from the group consisting of SEQ ID NOS: 29, 37 and 38.

25. A method as in claim 1 wherein the step of administering a therapeutic
composition comprises administering an antibody comprising a heavy chain CDR1 region
having a sequence selected from the group consisting of SEQ ID NOS: 50, 55 and 56.

30 26. A method as in claim 1 wherein the step of administering a therapeutic
composition comprises administering an antibody comprising a heavy chain CDR2 region
having a sequence selected from the group consisting of SEQ ID NOS: 69, 74 and 75.

27. A method as in claim 1 wherein the step of administering a therapeutic composition comprises administering an antibody comprising a heavy chain CDR3 region having a sequence selected from the group consisting of SEQ ID NOS: 88, 93 and 94.

5 28. A fusion molecule as in claim 16 wherein the first portion comprises a polypeptide comprising one or more amino acid sequences selected from the group consisting of SEQ. ID. NOS: 5, 12,13, 21, 23, 29, 37, 38, 50, 55, 56, 69, 74, 75, 88, 93 and 94.

10 29. A method as in any of claims 2-5 wherein the therapeutic composition comprises a humanized antibody.

30. A method as in any of claims 2-5 wherein the therapeutic composition comprises an Fv, scFv, Fab' and F(ab')₂.

15

31. A method as in any of claims 11-15 wherein the antibody comprises a humanized antibody.

20 32. A method as in any of claims 11-15 wherein the antibody comprises an Fv, scFv, Fab' and F(ab')₂.

33. A fusion molecule as in any of claims 17-18 wherein the antibody comprises a humanized antibody.

25 34. A fusion molecule as in any of claims 17-18 wherein the antibody comprises an Fv, scFv, Fab' and F(ab')₂.

35. A method as in any of claims 22-27 wherein the therapeutic composition comprises a humanized antibody.

30

36. A method as in any of claims 22-27 wherein the therapeutic composition comprises an Fv, scFv, Fab' and F(ab')₂.

37. A composition as in claim 21 wherein the antibody comprises a humanized antibody.

38. A composition as in claim 21 wherein the antibody comprises an Fv, scFv,
5 Fab' and F(ab')₂.

39. A fusion molecule as in claim 28 wherein the polypeptide comprises a humanized antibody.

10 40. A fusion molecule as in claim 28 wherein the polypeptide comprises an Fv, scFv, Fab' and F(ab')₂.

41. An antibody that binds to a cell that expresses CD200 comprising a polypeptide sequence selected from the group consisting of SEQ ID NOS: 111 - 199.

15

42. An antibody as in claim 41 comprising a polypeptide sequence selected from the group consisting of SEQ ID NOS: 111, 114, 118, 133, 136, 140, 141, 149, 155, 162, 163, 165, 174, 181, 187, 188, 189, 195 and 199.

20 43. An antibody as in claim 41 comprising a polypeptide sequence selected from the group consisting of SEQ ID NOS: 200-211.

44. An antibody as in claim 41 wherein the antibody comprises a humanized antibody.

25

45. An antibody as in claim 41 wherein the antibody comprises an Fv, scFv, Fab' and F(ab')₂.

46. A method of treating cancer comprising:
30 administering to a subject afflicted with cancer a therapeutic composition that contains an antibody that binds to a cell that expresses CD200, the antibody being present in an amount sufficient to promote production of one or more Th1 cytokines.

47. A method as in claim 46 wherein the antibody comprises a polypeptide sequence selected from the group consisting of SEQ ID NOS: 111 - 199.

48. A method as in claim 46 wherein the antibody comprises a polypeptide
5 sequence selected from the group consisting of SEQ ID NOS: 111, 114, 118, 133, 136, 140, 141, 149, 155, 162, 163, 165, 174, 181, 187, 188, 189, 195 and 199.

49. A method as in claim 46 wherein the antibody comprises polypeptide sequence selected from the group consisting of SEQ ID NOS: 200-211.

10

50. A method as in claim 46 wherein the antibody comprises a humanized antibody.

51. A method as in claim 46 wherein the antibody comprises an Fv, scFv, Fab'
15 and F(ab')₂.

52. A method as in claim 46 wherein the cancer is CLL.

53. A method of killing tumor cells comprising:
20 administering to a subject afflicted with cancer a therapeutic composition that contains an antibody that binds to a cell that expresses CD200, the antibody being present in an amount sufficient to cause antibody-dependent cell-mediated cytotoxicity.

54. A method as in claim 53 wherein the antibody comprises a polypeptide
25 sequence selected from the group consisting of SEQ ID NOS: 111 - 199.

55. A method as in claim 53 wherein the antibody comprises a polypeptide sequence selected from the group consisting of SEQ ID NOS: 111, 114, 118, 133, 136,
30 140, 141, 149, 155, 162, 163, 165, 174, 181, 187, 188, 189, 195 and 199.

56. A method as in claim 53 wherein the antibody comprises a polypeptide sequence selected from the group consisting of SEQ ID NOS: 200-211.

57. A method as in claim 53 wherein the antibody comprises a humanized antibody.
58. A method as in claim 53 wherein the antibody comprises an Fv, scFv, Fab' and F(ab')₂.
59. A method as in claim 53 wherein the cancer is CLL.
60. A fusion molecule comprising:
a first portion that targets cells bearing the OX-2/CD200 antigen, the first portion comprising a polypeptide sequence selected from the group consisting of SEQ ID NOS: 111 - 199; and
a second portion that promotes the death of cells.
61. A fusion molecule as in claim 60 wherein the first portion comprises a polypeptide sequence selected from the group consisting of SEQ ID NOS: 111, 114, 118, 133, 136, 140, 141, 149, 155, 162, 163, 165, 174, 181, 187, 188, 189, 195 and 199.
62. A fusion molecule as in claim 61 wherein the first portion comprises a polypeptide sequence selected from the group consisting of SEQ ID NOS: 200-211.
63. A composition comprising an antibody comprising a polypeptide sequence selected from the group consisting of SEQ ID NOS: 111 - 199 and a pharmaceutically acceptable carrier.
64. A composition as in claim 63 wherein the antibody comprises a humanized antibody.
65. A composition as in claim 63 wherein the antibody comprises an Fv, scFv, Fab' and F(ab')₂.

66. A method of treating cancer comprising:
administering to a subject afflicted with cancer a therapeutic composition that contains an antibody that binds to a cell that expresses CD200, the antibody being present in an amount sufficient to promote production of one or more Th1 cytokines and to cause
5 complement-mediated or antibody-dependent cellular cytotoxicity.
67. A chimeric antibody that binds to a cell that expresses CD200 comprising a polypeptide sequence selected from the group consisting of SEQ ID NOS: 200-211.
- 10 68. A method comprising:
determining whether OX-2/CD200 is upregulated in a subject; and
administering to the subject of a therapy that enhances immune response.
- 15 69. A method as in claim 68 wherein the therapy comprises administering a polypeptide that binds to OX-2/CD200 or an OX-2/CD200 receptor, the polypeptide being administered in an amount effective to inhibit the immune-suppressing effect of OX-2/CD200.
- 20 70. A method as in claim 68 wherein the subject is a cancer patient.
71. A method as in claim 68 wherein the subject is a CLL patient.
72. A method as in claim 69 wherein the polypeptide comprises an antibody.
- 25 73. A method as in claim 69 wherein the polypeptide comprises a humanized antibody.
74. A method as in claim 69 wherein the polypeptide comprises an Fv, scFv, Fab' and F(ab')₂.

30

75. A method as in claim 68 wherein the step of determining whether OX-2/CD200 is upregulated in a subject comprises:
obtaining tissue from a cancer patient;
evaluating whether CD200 levels are at least 2-fold above the levels found on
5 corresponding normal tissue.

76. A method as in claim 75 wherein blood is the tissue obtained from a cancer patient.

10 77. A method as in claim 75 wherein tissue is obtained from a cancer patient by performing a biopsy.

78. A method as in claim 75 wherein CD200 levels are evaluated by conducting FACS analysis using anti-CD200 antibodies in combination with cancer cell
15 markers.

79. A method as in claim 75 wherein the cancer cell marker is CD38 or CD19.

80. A method as in claim 75 wherein the patient is afflicted with a
20 hematopoietic cancer.

81. A method as in claim 75 wherein the patient is afflicted with CLL.

82. A method as in claim 69 wherein the therapy comprises eliminating
25 existing regulatory T cells prior to administering a polypeptide that binds to OX-2/CD200 or an OX-2/CD200 receptor, the polypeptide being administered in an amount effective to inhibit the immune-suppressing effect of OX-2/CD200.

83. A method as in claim 82 wherein the therapy comprises administering a
30 regulatory T cell eliminating amount of a reagent selected from the group consisting of anti-CD25 antibodies and cyclophosphamide.

84. A method as in claim 69 wherein the therapy further comprises administering a myeloablative therapy.

85. A method as in claim 84 wherein the therapy further comprises a bone marrow transplant.

86. A method as in claim 84 wherein the therapy further comprises a transfer
5 of CLL reactive T cells.

87. A method as in claim 69 wherein the therapy further comprises administering a cancer vaccine.

10 88. A method as in claim 87 wherein the cancer vaccine is selected from the group consisting of dendritic cells loaded with CLL cells or proteins, peptides or RNA derived from dendritic cells loaded with CLL cells, patient-derived heat-shocked proteins, tumor peptides and tumor proteins.

15 89. A method as in claim 69 wherein the therapy further comprises administering an effective immune stimulating amount of an immuno-stimulatory compound.

20 90. A method as in claim 89 wherein the immuno-stimulatory compound is selected from the group consisting of CpG, toll-like receptor agonists and anti-CTLA-4 antibodies.

25 91. A method as in claim 69 wherein the therapy further comprises administering an effective immunosuppressive blocking amount of a compound selected from the group consisting of anti-PDL1 antibodies, anti-PDL1 antibodies, anti-IL-10 antibodies and anti-IL-6 antibodies.

30 92. A method as in claim 69 wherein the therapy comprises administering an effective amount of an agent capable of blocking negative regulation of T cells or antigen presenting cells.

93. A method as in claim 92 wherein the agent is selected from the group consisting of anti-CTLA4 antibodies, anti-PD-L1 antibodies, anti-PDL-2 antibodies and anti-PD-1 antibodies.

94. A method as in claim 69 wherein the therapy comprises administering an effective amount of an agent capable of enhancing positive co-stimulation of T cells.

5 95. A method as in claim 94 wherein the agent is selected from the group consisting of anti-CD40 antibodies and anti 4-1BB antibodies.

96. A method as in claim 69 wherein the therapy comprises administering a cancer vaccine.

10 97. A method as in claim 96 wherein the cancer vaccine is selected from the group consisting of dendritic cells loaded with CLL cells or proteins, peptides or RNA derived from dendritic cells loaded with CLL cells, patient-derived heat-shocked proteins, tumor peptides and tumor proteins.

15 98. A method as in claim 69 wherein the therapy comprises administering an effective amount of an agent capable of activating receptors of the innate immune system.

99. A method as in claim 98 wherein the agent is selected from the group consisting of CpG, Luivac, Biostim, Ribominyl, Imudon and Bronchovaxom.

20

100. A method as in claim 69 wherein the therapy comprises administering an effective immune-enhancing amount of a cytokine.

25 101. A method as in claim 100 wherein the cytokine is selected from the group consisting of IL-2, GM-CSF and IFN-gamma.

102. A method for monitoring the progress of a therapeutic treatment, the method comprising:

administering an immunomodulatory therapy to a subject; and

30 determining OX-2/CD200 levels in a subject at least twice to determine the effectiveness of the therapy.

103. A method as in claim 102 wherein OX-2/CD200 levels in a subject are determined before administration of the immunomodulatory therapy and then OX-2/CD200 levels in a subject are determined after at least one administration of the therapy.

5

104. A method as in claim 102 further comprising the step of adjusting the dosage amount or frequency of the immunomodulatory therapy.

105. A method as in claim 102 wherein OX-2/CD200 levels are determined by
10 detecting a marker that correlates with OX-2/CD200.

106. A method of assessing the immunomodulatory effect of a molecule expressed by cancer cells comprising:
administering a molecule naturally expressed by cancer cells, cancer cells and
15 lymphocytes to a subject; and
monitoring the rate of growth of the cancer cells,
wherein the molecule is deemed to have an immunomodulatory effect when any change in the growth rate of the cancer cells is observed compared to the rate of growth when the cancer cells and donor lymphocytes are administered alone.

20

107. A method as in claim 106 wherein the molecule is considered immune enhancing where a reduction in the rate of growth of the cancer cells is observed compared to the rate of growth when the cancer cells and donor lymphocytes are administered alone.

25

108. A method as in claim 106 wherein the molecule is considered immunosuppressive where a greater rate of growth of the cancer cells is observed compared to the rate of growth when the cancer cells and donor lymphocytes are administered alone.

30

109. A method as in claim 106 wherein the molecule naturally expressed by a cancer cell is administered by injecting cells that naturally express the compound.

110. A method as in claim 106 wherein the molecule naturally expressed by a cancer cell is administered by injecting cells that have been engineered to express the molecule.

5 111. A method as in claim 106 further comprising identifying the molecule expressed by cancer cells from a database.

112. A method as in claim 106 further comprising identifying the molecule expressed by cancer cells experimentally.

10

113. The method of claim 106, wherein the lymphocytes are human.

114. The method of claim 113 wherein the human lymphocytes are selected from the group consisting of peripheral blood lymphocytes (PBLs), T cells, B cells, and
15 dendritic cells.

115. The method of claim 106, wherein the lymphocytes are human peripheral blood lymphocytes (PBLs).

20 116. The method of claim 106 wherein the number of lymphocytes administered is insufficient to reduce the rate of growth of cancer cells.

117. The method of claim 106 wherein the number of lymphocytes administered is less than the number of cancer cells administered.

25

118. The method of claim 106 wherein the number of lymphocytes administered is from about 1 million cells to about 2 million cells.

119. The method of claim 106 wherein the number of lymphocytes
30 administered is sufficient to reduce the rate of growth of cancer cells.

120. The method of claim 106 wherein the number of lymphocytes administered is greater than or equal to the number of cancer cells administered.

121. The method of claim 106 wherein the number of lymphocytes administered is from about 5 million cells to about 10 million cells.

122. The method of claim 106, wherein administering of cancer cells comprises injecting at least one lymphoma cell line selected from the group consisting of RAJI and Namalwa into the subject.

123. The method of claim 106 wherein the subject is an immunodeficient mouse.

124. The method of claim 106 wherein the number of lymphocytes administered is sufficient to slow the growth of the cancer cells and the molecule expressed by cancer cells is considered immunosuppressive where the rate of growth of the cancer cells observed is higher compared to the rate of growth when the cancer cells and lymphocytes are administered alone.

125. The method of claim 106 wherein the number of lymphocytes administered is insufficient to slow the growth of the cancer cells and the molecule expressed by cancer cells is considered immune-enhancing where the rate of growth of the cancer cells observed is lowered compared to the rate of growth when the cancer cells and lymphocytes are administered alone.

126. A method as in claim 106 further comprising determining the rate of growth of the cancer cells by administering lymphocytes and cancer cells alone to a control subject.

127. A method as in claim 106 further comprising determining the rate of growth of the cancer cells by administering lymphocytes and cancer cells alone to the subject prior to administering the molecule that is expressed by cancer cells.

128. A method of assessing the immunomodulatory effect of a compound comprising the steps of:

administering a compound to be assessed, cancer cells and lymphocytes to a subject; and

monitoring the rate of growth of the cancer cells,

wherein the compound is deemed to have an immunomodulatory effect where any change in the growth rate of the cancer cells is observed compared to the rate of growth when the cancer cells and donor lymphocytes are administered alone.

5

129. The method of claim 128 wherein the compound is considered immune enhancing where a reduction in the rate of growth of the cancer cells is observed compared to the rate of growth when the cancer cells and donor lymphocytes are administered alone.

10

130. The method of claim 128 wherein the compound is considered immunosuppressive when the rate of growth of the cancer cells in the presence of the compound is greater than the rate of growth when the cancer cells and donor lymphocytes are administered alone.

15

131. The method of claim 128 wherein the compound to be assessed is an antibody.

132. The method of claim 128 wherein administering of cancer cells comprises injecting at least one lymphoma cell line selected from the group consisting of RAJI and Namalwa into the subject.

20

133. The method of claim 128 wherein the cancer cells administered express an immunosuppressive molecule.

25

134. The method of claim 128 wherein the cancer cells administered express CD200.

135. The method of claim 128 wherein the expression of CD 200 is upregulated by the cancer cells administered.

30

136. The method of claim 128 wherein the lymphocytes are human.

137. The method of claim 136 wherein the human lymphocytes are selected

from the group consisting of peripheral blood lymphocytes (PBLs), T cells, B cells, and dendritic cells.

138. The method of claim 128 wherein the lymphocytes are human peripheral
5 blood lymphocytes (PBLs).

139. The method of claim 138 wherein the number of human PBLs administered is less than the number of cancer cells administered.

10 140. The method of claim 138 wherein the number of human PBLs administered is from about 1 million cells to about 2 million cells.

141. The method of claim 138 wherein the number of human PBLs administered is greater than or equal to the number of cancer cells administered.

15

142. The method of claim 138 wherein the number of human PBLs administered is from about 5 million cells to about 10 million cells.

143. The method of claim 128 wherein the subject is an immunodeficient
20 mouse.

144. The method of claim 128 wherein the number of lymphocytes administered is sufficient to slow the rate of growth of the cancer cells when administered with the cancer cells alone, the compound is considered immunosuppressive when the
25 rate of growth of the cancer cells in the presence of the compound is greater than the rate of growth when the cancer cells and donor lymphocytes are administered alone.

145. The method of claim 128 wherein the number of lymphocytes administered is not sufficient to slow the rate of growth of cancer cells when administered with the cancer cells alone, and the compound is considered immune enhancing where a
30 reduction in the rate of growth of the cancer cells is observed compared to the rate of growth when the cancer cells and donor lymphocytes are administered alone.

146. A method of assessing the immunomodulatory effect of a compound comprising the steps of:

administering a compound to be assessed, cancer cells and less than about 2×10^6 lymphocytes to a subject; and

5 monitoring the rate of growth of the cancer cells,

wherein the compound is considered immune enhancing where a reduction in the rate of growth of the cancer cells is observed compared to the rate of growth when the cancer cells and donor lymphocytes are administered alone.

10 147. The method of claim 146 wherein the cancer cells administered express an immunosuppressive molecule.

148. The method of claim 146 wherein the cancer cells administered express CD200.

15

149. The method of claim 146 wherein the expression of CD200 is upregulated by the cancer cells administered.

150. A method of assessing the immunomodulatory effect of a compound comprising the steps of:

20

administering a compound to be assessed, cancer cells and more than about 5×10^6 lymphocytes to a subject; and

monitoring the rate of growth of the cancer cells,

25 wherein the compound is considered immunosuppressive when the rate of growth of the cancer cells in the presence of the compound is greater than the rate of growth when the cancer cells and donor lymphocytes are administered alone.

30

151. The method of claim 150 wherein the cancer cells administered express an immunosuppressive molecule.

30

152. The method of claim 150 wherein the cancer cells administered express CD200.

153. The method of claim 150 wherein the expression of CD200 is upregulated by the cancer cells administered.




FIG. 1






FIG. 2



Rabbit scFv Clone

FIG. 3a



rabbit scFv Clone

FIG. 3b





FIG. 4a

CELL ELISA 8/19/00



Rabbit scFv Clone

FIG. 5a






FIG. 5b

CELL ELISA 9/15/00




Rabbit scFv Clone

FIG. 6a



Rabbit scFv Clone



FIG. 6b



Rabbit scFv Clone



FIG. 6c

FL2: scFv-9/HA-biotin/SA-PE
 FL1: CD5-FITC
 FL3: CD19-PerCP



File: PBMC.015
 XParameter: FL1-H FL1-Height (Log)
 YParameter: FL2-H FL2-Height (Log)

Quad	Events	%Gated	%Total	XGeo Mean	YGeo Mean
UL	1881	9.40	5.84	6.45	118.74
UR	4368	21.84	13.56	266.89	45.49
LL	2831	14.16	8.79	6.65	7.40
LR	10920	54.60	33.90	282.52	5.72





File: PBMC.015
 XParameter: FL1-H FL1-Height (Log)
 YParameter: FL3-H FL3-Height (Log)

Quad	Events	%Gated	%Total	XGeo Mean	YGeo Mean
UL	1874	9.37	5.82	6.55	65.56
UR	409	2.04	1.27	50.57	55.81
LL	2838	14.19	8.81	6.57	4.19
LR	14879	74.39	46.19	291.30	2.17

File: PBMC.015
 XParameter: FL2-H FL2-Height (Log)
 YParameter: FL3-H FL3-Height (Log)

Quad	Events	%Gated	%Total	XGeo Mean	YGeo Mean
UL	171	0.85	0.53	10.16	54.88
UR	2112	10.56	6.56	137.20	64.47
LL	13744	68.72	42.67	6.08	2.52
LR	3973	19.86	12.33	41.31	2.06



Numerator Histogram

File: CLL.006

Denominator Histogram

File: CLL.004

Marker	Left	Right	Events	% Gated	% Total	Mean	Geo. Mean	CV	Median	Peak Ch
All	1	9910	10233	99.57	86.52	654.78	553.65	47.14	620.82	685
M1	28	9910	10028	97.58	84.79	667.84	596.59	44.59	626.43	685

Numerator Histogram

File: PBMC.008

Denominator Histogram

File: PBMC.006

Marker	Left	Right	Events	% Gated	% Total	Mean	Geo. Mean	CV	Median	Peak Ch
All	1	9910	20309	99.96	64.48	111.91	12.05	280.93	8.98	1
M1	28	9910	2403	11.83	7.63	881.67	708.77	45.81	897.69	1000





FIG. 8a

Key	Name	Parameter	Gate
---	CLL.004	FL2-H	G3
---	CLL.009	FL2-H	G3
---	PBMC.006	FL2-H	G1
---	PBMC.010	FL2-H	G1



Numerator Histogram
File: PBMC.010

Denominator Histogram
File: PBMC.006


Marker	Left	Right	Events	% Gated	% Total	Mean	Geo Mean	CV	Median	Peak Ch
All	1	9910	20389	100.12	64.95	34.40	8.94	351.33	8.35	1
M1	28	9910	1953	9.59	6.22	281.98	176.73	103.08	201.69	271

Numerator Histogram
File: CLL.009

Denominator Histogram
File: CLL.004

Marker	Left	Right	Events	% Gated	% Total	Mean	Geo Mean	CV	Median	Peak Ch
All	1	9910	10258	99.82	86.73	587.42	345.59	104.91	392.42	385
M1	28	9910	9788	95.24	82.76	614.87	403.58	100.46	414.18	385

FIG. 8b



Numerator Histogram
File: CLL.018

Denominator Histogram
File: CLL.004

Marker	Left	Right	Events	% Gated	% Total	Mean	Geo Mean	CV	Median	Peak Ch
All	1	9910	10272	99.95	86.85	568.19	481.42	48.32	528.03	528
M1	28	9910	10073	98.01	85.17	579.08	515.60	45.93	532.80	528
M2	103	9910	9924	96.57	83.91	587.05	534.83	44.25	537.61	528

Numerator Histogram
File: PBMC.014

Denominator Histogram
PBMC.006

Marker	Left	Right	Events	% Gated	% Total	Mean	Geo Mean	CV	Median	Peak Ch
All	1	9910	20356	100.19	64.63	34.36	15.09	173.49	12.30	1
M1	28	9910	5229	25.74	16.60	103.57	80.28	82.38	71.05	38
M2	103	9910	1893	9.32	6.01	190.40	179.03	45.79	177.83	168

FIG. 8c

Table 1. Summary of CLL scFv Clones

Pool	Clone	CLL	Primary B	CLL-AAT	RL (NHL)	Ramos (Burkitt's)	TF1	Patient-Specific	Expression Lost	Fingerprint
R3/RSC-S CLL-TF1	E1	++	++	++	-	-	-			1
	E2	+	-	+	-	-	-			2
	E4.1	++	+	++	±	±	±			3
	E11	++	+	++	-	-	-			4
	F1	+	-	+	-	-	-			5
	F2	++	+	++	-	-	-			6
	F12	++	+	++	-	-	-			7
	G7	+	+	+	+	+	+	+		8
	H6.1	++	+	++	+	+	-	-		9
	H12	++	+	++	-	-	-	-		10
	A2	++	++	++	+	+	-	-		11
	A9	++	+	++	-	-	-	-		12
	C6.1	-	-	nd	nd	nd	nd	+		13
	C7	-	-	nd	nd	nd	nd	+		14
	D7.1	+	+	+	+	+	+	+		15
	D8.1	-	+	nd	nd	nd	nd	+		16
	D11.1	++	+	++	-	-	-	-		17
	G10.1	+	+	+	-	-	-	-		18
	H6.2	++	++	++	-	-	-	-		19
	A3.1	++	+	++	-	-	-	-		20
	A5.2	++	+	++	-	-	-	-		21
	B3.1	+	+	+	-	-	-	-		22
	D2.1	+	+	+	-	-	-	-		23
	D5.1	+	+	+	±	±	±	±		24
E3	+	+	+	-	-	-	-		25	
E4.2	+	+	+	-	-	-	-		26	
G2.2	-	-	nd	nd	nd	nd	+		27	
H1	-	-	nd	nd	nd	nd	+		28	
H6.3	-	-	nd	nd	nd	nd	+		29	
A8	-	+	+	nd	nd	nd	+		30	
B12.1	+	+	++	+	+	+	±		31	
C12	++	+	++	+	+	+	±		32	
D1.1	+	+	+	nd	nd	nd	+		33	
D5.2	-	-	+	nd	nd	nd	+		34	
D8.2	++	+	++	+	+	+	±		(nd)	
F10	++	+	++	+	+	+	±			
A1.1	+	+	++	nd	nd	nd	nd			




FIG. 9a

Table 1.(cont'd) Summary of CLL scFv Clones

Pool	Clone	CLL	Primary B	CLL-AAT	RL (NHL)	Ramos (Burkitt's)	TF1	Patient-Specific	Expression Lost	Fingerprint
R5/RSC-L CLL-B	G9	±?	+	++	nd	nd	nd	±?		(nd)
	B1	+	++	++	nd	nd	nd			35
	B4.2	++	++	++	nd	nd	nd			36
	C10	++	+	++	nd	nd	nd			37
	D4.2	·	·	·	nd	nd	nd	±?	+	38
	D11.2	±?	·	+	nd	nd	nd	±?		39
	G1.2	++	+	++	nd	nd	nd			40
	D2.2	+	+	+	nd	nd	nd			41
	G12.1	+	+	+	+	+	·			42
	A1.2	±?	nd	++	nd	nd	nd	±?		43
R4/RSC-S CLL-B	A3.2	+	nd	++	+	+	+			44
	B3.2	·	nd	+	nd	nd	+	+		45
	B4.3	·	nd	++	nd	nd	nd			46
	B12.2	+	nd	++	+	+	+			47
	C4	++	nd	++	·	·	·			48
	E8.2	++	nd	++	·	·	·			46
	F7	+	nd	++	+	+	·			49
	D7.2	++	nd	++	+	+	±			50
	D12	++	nd	++	+	+	+			51
	E5	++	nd	++	+	+	·			52
	E6.2	+	nd	++	+	+	+			53
	E7.2	+	nd	++	+	+	+			54
	E5.2	+	nd	++	+	+	·			




FIG. 9a (Cont.)

Table 1. CDR Sequence of CLL Specific Rabbit scFv Antibodies

CLONE	LC-CDR1	(SEQ ID NO: 1)	LC-CDR2	(SEQ ID NO: 18)	LC-CDR3	(SEQ ID NO: 25)
A2c	TLSTGYSVGSYVIA	(SEQ ID NO: 2)	HSEAKHQGS	(SEQ ID NO: 19)	ATAHGSSSFHVH	(SEQ ID NO: 26)
G12.1c	QASESIRN---YLA	(SEQ ID NO: 3)	GASNL---ES	(SEQ ID NO: 20)	QSGDYSA---GLT	(SEQ ID NO: 27)
B4.2a	QASESIRN---YLA	(SEQ ID NO: 4)	GASNL---AS	(SEQ ID NO: 21)	QSGYISA---GLT	(SEQ ID NO: 28)
E1c	QASESISN---WLA	(SEQ ID NO: 5)	RASLTL---AS	(SEQ ID NO: 22)	QSGYISA---GVT	(SEQ ID NO: 29)
F2d	QASESISN---YLA	(SEQ ID NO: 6)	GASNL---ES	(SEQ ID NO: 23)	QSGYISA---GLT	(SEQ ID NO: 30)
F5e	QASQNIYS---NLA	(SEQ ID NO: 7)	LAFTL---AS	(SEQ ID NO: 24)	QGGDYSSSSSYGYG	(SEQ ID NO: 31)
H6.2b	QASQSVNN---LLA	(SEQ ID NO: 8)	GASNL---ES	(SEQ ID NO: 25)	QSGYISP---GVT	(SEQ ID NO: 32)
G10.1	QASESINN---YLA	(SEQ ID NO: 9)	GASNL---ES	(SEQ ID NO: 26)	QSGYISG---GAT	(SEQ ID NO: 33)
D11.1c	LASENVYS---AVA	(SEQ ID NO: 10)	GASDL---ES	(SEQ ID NO: 27)	Q-GYSSYPPT	(SEQ ID NO: 34)
A5.2c	LASENVYG---AVA	(SEQ ID NO: 11)	GASNL---ES	(SEQ ID NO: 28)	Q-GYSSYP-T	(SEQ ID NO: 35)
F1d	QASQSVNN---LLA	(SEQ ID NO: 12)	GASNL---ES	(SEQ ID NO: 29)	AGYKSSSTD-GIA	(SEQ ID NO: 36)
F1e	QASQSISN---LLA	(SEQ ID NO: 13)	GASNL---ES	(SEQ ID NO: 30)	QSGYISA---GHLT	(SEQ ID NO: 37)
E4.2	LASENVAS---TVS	(SEQ ID NO: 14)	GASNL---ES	(SEQ ID NO: 31)	LGGFGYSTT-GLT	(SEQ ID NO: 38)
E2c	TLSTGYSVGEYVW	(SEQ ID NO: 15)	HTDDIKHQGS	(SEQ ID NO: 32)	AIAHGTESSFHVV	(SEQ ID NO: 39)
A9c	TLSTGYSVGEYVW	(SEQ ID NO: 16)	HTDDIKHQGS	(SEQ ID NO: 33)	AIAHGTESSFHVV	(SEQ ID NO: 40)
E11e	TLRTGYSVGEYPLV	(SEQ ID NO: 17)	HTDDIKHQGS	(SEQ ID NO: 34)	ATGHGSSSAGVV	(SEQ ID NO: 41)
A1.1	LASEDIYS---GLS	(SEQ ID NO: 18)	GASNL---ES	(SEQ ID NO: 35)	LGGYPYSST-GTA	(SEQ ID NO: 42)
F5.2	QASQSVSN---LLA	(SEQ ID NO: 19)	GASNL---AS	(SEQ ID NO: 36)	QSGWYSA---GALT	(SEQ ID NO: 43)
F10b	QASQSVNN---LLA	(SEQ ID NO: 20)	RASTL---AS	(SEQ ID NO: 37)	QSGYIRA---GDLT	(SEQ ID NO: 44)
F7a	QASQSVSN---LLA	(SEQ ID NO: 21)	GASNL---ES	(SEQ ID NO: 38)	QSGYISA---GLT	(SEQ ID NO: 45)
F6b	QASQSVSN---LLA	(SEQ ID NO: 22)	GASNL---ES	(SEQ ID NO: 39)	QSGYISA---GLT	(SEQ ID NO: 46)
C12b	QASQSVSN---LLA	(SEQ ID NO: 23)	GASNL---ES	(SEQ ID NO: 40)	QSGYISA---GLT	(SEQ ID NO: 47)
D2.1b	QASQSVNN---LLA	(SEQ ID NO: 24)	GASNL---ES	(SEQ ID NO: 41)	QSGYISA---GLT	(SEQ ID NO: 48)
D1.1	QASEDIES---YLA	(SEQ ID NO: 25)	GASNL---ES	(SEQ ID NO: 42)	QSNAWSV---GMT	(SEQ ID NO: 49)
D2.2a	QSSQSIAGA---YLS	(SEQ ID NO: 26)	LASKL---AS	(SEQ ID NO: 43)	AAQYSGN---IYT	(SEQ ID NO: 50)
D2.2b	LASENVYG---AVA	(SEQ ID NO: 27)	GASNL---ES	(SEQ ID NO: 44)	Q-GYSSYP-T	(SEQ ID NO: 51)

FIG. 9B

CLONE	HC-CDR1	HC-CDR2	HC-CDR3
A2c	NYAMT (SEQ ID NO: 45)	IISSNGGA--DYASWAK (SEQ ID NO: 64)	DDEGYDDYGDYMGYFTL (SEQ ID NO: 84)
G12.1c	SYGLS (SEQ ID NO: 46)	YFDPVFGNI--YYATWVD (SEQ ID NO: 65)	DRIYVSSVG---YAFNL (SEQ ID NO: 85)
B4.2a	TYGVS (SEQ ID NO: 47)	YNDPIFGNT--YYATWVN (SEQ ID NO: 66)	DRAYASSG---YXXXX (SEQ ID NO: 86)
E1c	SNAMG (SEQ ID NO: 48)	IISSSGGT--YYASWAK (SEQ ID NO: 67)	DWIAAGKS----YGLDL (SEQ ID NO: 87)
F2d	TNAMG (SEQ ID NO: 49)	IISSSGST--YYASWAK (SEQ ID NO: 68)	DWIAAGKS----YGLDL (SEQ ID NO: 87)
E5e	SSDWIC (SEQ ID NO: 50)	CIYTGSSSSTWYASWAK (SEQ ID NO: 69)	RYTGDNG-----NL (SEQ ID NO: 88)
H6.2b	SDVIS (SEQ ID NO: 51)	YIYTGSGT--DYASWVN (SEQ ID NO: 70)	DAAYAGYGM----IFNL (SEQ ID NO: 89)
G10.1	SDVIS (SEQ ID NO: 51)	YIYTGSGT--DYASWVN (SEQ ID NO: 70)	DAAYAGYGM----IFNL (SEQ ID NO: 89)
D11.1c	TYAMG (SEQ ID NO: 52)	SIYASRSP--YYASWAK (SEQ ID NO: 71)	GDAGSIP-----YFKL (SEQ ID NO: 90)
A5.2c	TYAMG (SEQ ID NO: 52)	SIYASRSP--YYASWAK (SEQ ID NO: 71)	GDAGSIP-----YFKL (SEQ ID NO: 90)
F1d	SNAMT (SEQ ID NO: 53)	TIIYGDNT--YYASWAK (SEQ ID NO: 72)	GNV-----FSDL (SEQ ID NO: 91)
F1e	DFAMS (SEQ ID NO: 54)	VVYAGTRGDTYYANWAK (SEQ ID NO: 73)	GLT-----YYPL (SEQ ID NO: 92)
E4.2	DFAMS (SEQ ID NO: 54)	VVYAGTRGDTYYANWAK (SEQ ID NO: 73)	GLT-----YYPL (SEQ ID NO: 92)
E2c	SYGMN (SEQ ID NO: 55)	YIDPDYGST--YYASWVN (SEQ ID NO: 74)	GAYSGYPS-----YFNL (SEQ ID NO: 93)
A9c	SYGMN (SEQ ID NO: 55)	YIDPDYGST--YYASWVN (SEQ ID NO: 74)	GAYSGYPS-----YFNL (SEQ ID NO: 93)
E11e	SNAMS (SEQ ID NO: 56)	ITVPSGNV--YYASWAK (SEQ ID NO: 75)	G-----FFNL (SEQ ID NO: 94)
A1.1	TNAIS (SEQ ID NO: 57)	YSSYGNNA--HYTNWAK (SEQ ID NO: 76)	GNA-----YSDL (SEQ ID NO: 95)
F5.2	SNAMS (SEQ ID NO: 56)	IIIGSGT--YYANWAK (SEQ ID NO: 77)	DQPIIYGAYGDYGLATGTRLDL (SEQ ID NO: 96)
F10b	SYYMS (SEQ ID NO: 58)	IISSSGTS--YYATWAK (SEQ ID NO: 78)	DQPIIDAAYGDYGLATGTRLDL (SEQ ID NO: 97)
F7a	SYTMS (SEQ ID NO: 59)	IISSSGSA--YYATWAK (SEQ ID NO: 79)	DQPIITTDYGGYGIATGTRLDL (SEQ ID NO: 98)
F6b	SNALS (SEQ ID NO: 60)	IIVSGGTT--YYADWAK (SEQ ID NO: 80)	DQPIIYAGYGY---ATGTRLDL (SEQ ID NO: 99)
C12b	SNALS (SEQ ID NO: 60)	IIVSGGTT--YYADWAK (SEQ ID NO: 80)	DQPIIYAGYGY---ATGTRLDL (SEQ ID NO: 99)
D2.1b	TNAMS (SEQ ID NO: 61)	TIIYGTNA--YYASWAK (SEQ ID NO: 81)	GNT-----YSDL (SEQ ID NO: 100)
D1.1	SNAMS (SEQ ID NO: 56)	TIIYGTNA--YYASWAK (SEQ ID NO: 81)	GNT-----YSDL (SEQ ID NO: 100)
D2.2a	SSYWIC (SEQ ID NO: 62)	CIYTGSGSTYYASWAK (SEQ ID NO: 82)	AYIYGGYG-----FFDL (SEQ ID NO: 101)
D2.2b	NYGVN (SEQ ID NO: 63)	YIDPVFGST--YYASWVN (SEQ ID NO: 83)	EASFYY-----GMDL (SEQ ID NO: 102)

CLONE: designation of representative clone for sequence; LC: Ig light chain; HC: Ig heavy chain; CDR: complementarity determining region

FIG. 9B (Cont.)

Table 1(cont'd). Expression Pattern of CLL Specific Rabbit scFv Antibodies

<u>CLONE</u>	<u>CLL</u>	<u>B</u>	<u>RL</u>	<u>Ramos</u>	<u>TF-1</u>	<u>Ag</u>	<u>Linker</u>
A2c	+	+	++	+	-		S
G12.1c	+	+	+	+	-	CD19	L
B4.2a	+	nd	+	+	-		L
E1c	++	+	-	-	-	CD23	S
F2d	++	+	-	-	-		S
E5e	±	nd	-	-	-		S
H6.2b	++	++	-	-	-		S
G10.1	+	+	-	-	-		S
D11.1c	++	+	-	-	-	CD23	S
A5.2c	++	+	-	-	-		S
F1d	+	±	-	-	-		S
F1e	++	nd	-	-	-		S
E4.2	+	+	-	-	-		S
E2c	+	±	-	-	-		S
A9c	++	+	-	-	-		S
E11e	++	+	-	-	-		S
A1.1	+	+	nd	nd	nd		L
F5.2	+	nd	+	+	-		L
F10b	nd	nd	nd	nd	nd		L
F7a	nd	nd	nd	nd	nd		L
F6b	nd	nd	nd	nd	nd		L
C12b	nd	nd	nd	nd	nd		L
D2.1b	nd	nd	nd	nd	nd		S
D1.1	+	+	nd	nd	nd		L
D2.2a	nd	nd	nd	nd	nd		L
D2.2b	nd	nd	nd	nd	nd		S

CLONE: designation of representative clone for sequence

Expression pattern: binding of scFv antibodies to primary human cells and cell lines as determined by whole cell ELISA assay

CLL: chronic lymphocytic leukemias (primary tumors and CLL-AAT cell line)

B: normal, primary human B lymphocytes

RL: non-Hodgkin's lymphoma cell line

Ramos: Burkitt's lymphoma cell line

TF-1: human erythroleukemia cell line

Ag: antigen recognized by scFv antibody (determined by immunoprecipitation and mass spectrometry)

Linker: type of linker sequence between VL and VH regions. S, short linker; L, long linker

FIG. 9C

Table 2. Mean fluorescent intensities of B-CLL cells and normal PBMC labeled with scFv antibodies

Donor	scFv-2		scFv-3		scFv-6		scFv-9	
	ratio	ratio	ratio	ratio	ratio	ratio	ratio	ratio
CLL(ML)	590	0.83	398	2.2	284	2.1	511	6.4
PBMC-1	715		181		137		80	
CLL(JR)	311	0.85	207	2.4	nd	nd	117	1.7
PBMC-2	368		87		nd	nd	67	
CLL(HTS)	219	0.69	173	1.6	nd	nd	176	3.6
PBMC-3	317		106		nd	nd	49	
CLL(RE)	305	0.59	360	3	nd	nd	142	1.7
PBMC-4	513		121		nd	nd	81	
CLL(GB)	262	0.47	387	1.8	nd	nd	163	1.5
PBMC-5	563		212		nd	nd	106	

Primary PBMC from five patients diagnosed with CLL and five normal donors were analyzed by flow cytometry. The geometric mean fluorescent intensities were determined for cells stained with four different scFv antibodies. For scFvs that bind to antigens overexpressed on CLL cells, the CLL/PBMC ratio of fluorescent intensities is >1.0

FIG. 10

Comparison of scFv-9 antigen and CD38 expression on CLL cells.

Patient ID	%CD19+	%CD38+	%scFv-9+	ScFv-9 Level	CD38	ScFv-9
ML	80	40	98	266	Hi	Hi
IB	86	87	96	366	Hi	Hi
BH	76	56	86	284	Hi	Hi
JG	82	92	97	125	Hi	Lo
RE	87	97	100	125	Hi	Lo
EM	91	8	95	268	Lo	Hi
HS	76	11	94	268	Lo	Hi
MP	40	6	95	280	Lo	Hi
JR	81	12	92	124	Lo	Lo
GB	65	20	98	187	Lo	Lo

FIG. 11

Identification of scFv Antigens

- Cell-surface biotinylation (CLL-ATT cells)
- Membrane isolation (nitrogen cavitation, differential centrifugation)
- Immunoprecipitation with scFv-HA coupled to Anti-HA beads
- SDS-PAGE
- Detection by Coomassie-stain or AP-streptavidin Western blot
- MALDI-MS or LC-MS/MS to obtain peptide mass spectra/peptide sequences




FIG. 12

Key	Name	Parameter	Gate
293-pCEP4:	---	091102.002 FL2-H	G1
293-CD200:	---	091102.008 FL2-H	G1





FIG. 13




FIG. 14




FIG. 15




FIG. 16

- Group 1: 8 mice, 4×10^6 RAJI cells each
- Group 2: 9 mice, 4×10^6 RAJI + 1×10^6 PBL donor 1
- Group 3: 7 mice, 4×10^6 RAJI + 5×10^6 PBL donor 1
- Group 4: 6 mice, 4×10^6 RAJI + 1×10^6 PBL donor 2
- Group 5: 7 mice, 4×10^6 RAJI + 5×10^6 PBL donor 2




FIG. 17A

- Group 6: 10 mice, 4×10^6 RAJI cells each
- Group 7: 10 mice, 4×10^6 RAJI + 1×10^6 PBL donor 3
- Group 8: 10 mice, 4×10^6 RAJI + 5×10^6 PBL donor 3
- Group 9: 9 mice, 4×10^6 RAJI + 1×10^7 PBL donor 3
- Group 10: 10 mice, 4×10^6 RAJI + 1×10^6 PBL donor 4
- Group 11: 14 mice, 4×10^6 RAJI + 1×10^7 PBL donor 4





FIG. 17B

Statistics	p values are given											
	comparing group2-4 to 1	group1	group2	group3	group4	group5	group7	group8	group9	group10	group11	
d22	t-test	ns	ns	ns	ns	<0.0022	ns	<0.468	<0.865	<0.837	<0.055	<0.927
	Welch	ns	ns	ns	ns	<0.0031	ns	<0.416	<0.521	<0.83	<0.053	<0.588
d25	Wilcox	ns	ns	ns	ns	<0.0093	ns	<0.87	<0.564	<0.64	<0.0712	<0.936
	t-test	ns	ns	ns	ns	<0.0766	ns	<0.824	<0.123	<0.296	<0.31	<0.89
	Welch	ns	ns	ns	ns	<0.0788	ns	<0.45	<0.0727	<0.057	<0.296	<0.427
d27	Wilcox	ns	ns	ns	ns	<0.014	ns	<0.2	<0.059	<0.0546	<0.1775	<0.145
	t-test	ns	ns	ns	ns	<0.038	ns	<0.41	<0.1	<0.06	<0.274	<0.26
	Welch	ns	ns	ns	ns	<0.036	ns	<0.575	<0.0122	<0.0082	<0.357	<0.245
d29	Wilcox	ns	ns	ns	ns	<0.014	ns	<0.88	<0.0154	<0.0076	<0.327	<0.196
	t-test	ns	ns	ns	ns	<0.0121	ns	<0.653	<0.0288	<0.0095	<0.48	<0.059
	Welch	ns	ns	ns	ns	<0.017	ns	<0.4429	<0.0207	<0.0106	<0.3459	<0.0622
d32	Wilcox	ns	ns	ns	ns	<0.0093	ns	<0.445	<0.025	<0.004	<0.341	<0.05
	t-test	ns	ns	ns	ns	<0.0197	ns	<0.315	<0.035	<0.0021	<0.2775	<0.053
	Welch	ns	ns	ns	ns	<0.025	ns	<0.4	<0.0178	<0.0025	<0.0915	<0.0915
d34	Wilcox	ns	ns	ns	ns	<0.0059	ns		<0.035	<0.0019	<0.35	<0.082
	t-test	ns	ns	ns	ns	<0.03	ns		<0.0345	<0.007	<0.48	<0.059
	Welch	ns	ns	ns	ns	<0.035	ns		<0.0144	<0.008	<0.0566	<0.0566
d36	Wilcox	ns	ns	ns	ns	<0.0059	ns		<0.0144	<0.0012	<0.0622	<0.0622
	t-test	ns	ns	ns	ns	<0.0277	ns		<0.0178	<0.0069	<0.3459	<0.3459
	Welch	ns	ns	ns	ns	<0.0277	ns		<0.035	<0.0073	<0.341	<0.05
d39	Wilcox	ns	ns	ns	ns	<0.0328	ns		<0.0076	<0.0083	<0.2775	<0.053
	t-test	ns	ns	ns	ns	<0.0022	ns		<0.0076	<0.0056	<0.0915	<0.0915
	Welch	ns	ns	ns	ns	<0.0607	ns		<0.0099	<0.0025	<0.082	<0.082
d41	Wilcox	ns	ns	ns	ns	<0.064	ns		<0.0172	<0.0019	<0.35	<0.082
	t-test	ns	ns	ns	ns	<0.064	ns		<0.0025	<0.007	<0.48	<0.059
	Welch	ns	ns	ns	ns	<0.0289	ns		<0.0032	<0.008	<0.0566	<0.0566
d43	Wilcox	ns	ns	ns	ns	<0.0309	ns		<0.0041	<0.0012	<0.0622	<0.0622
	t-test	ns	ns	ns	ns	<0.0502	ns		<0.0055	<0.0069	<0.3459	<0.3459
	Welch	ns	ns	ns	ns	<0.0023	ns		<0.0076	<0.0073	<0.341	<0.05
d45	Wilcox	ns	ns	ns	ns	<0.0257	ns		<0.0133	<0.004	<0.2775	<0.053
	t-test	ns	ns	ns	ns	<0.0438	ns		<0.0103	<0.008	<0.0915	<0.0915
	Welch	ns	ns	ns	ns	<0.004	ns		<0.006	<0.0085	<0.082	<0.082
d48	Wilcox	ns	ns	ns	ns	<0.004	ns		<0.0106	<0.0142	<0.35	<0.082
	t-test	ns	ns	ns	ns	<0.004	ns		<0.0106	<0.0142	<0.35	<0.082
	Welch	ns	ns	ns	ns	<0.004	ns		<0.0106	<0.0142	<0.35	<0.082
d50	Wilcox	ns	ns	ns	ns	<0.004	ns		<0.0106	<0.0142	<0.35	<0.082
	t-test	ns	ns	ns	ns	<0.004	ns		<0.0106	<0.0142	<0.35	<0.082
	Welch	ns	ns	ns	ns	<0.004	ns		<0.0106	<0.0142	<0.35	<0.082


FIG. 18

ns=not significant
significant groups are shown in red




ELISA results of representative IgG1 kappa clones after round 3 panning on CD200-Fc captured on goat anti-mouse IgG Fc antibody

FIG. 19A




ELISA results of representative IgG2a kappa clones after round 3 panning on CD200-Fc captured on goat anti-mouse IgG Fc antibody

FIG. 19B



ELISA results of representative IgG1 kappa clones after round 3 panning on CD200-Fc directly coated on microtiter wells.

FIG. 19C



ELISA results of representative IgG2a kappa clones after round 3 panning on CD200-Fc directly coated on microtiter wells.

FIG. 19D

Flow cytometry results of representative IgG1 clones selected on CD200-Fc captured with goat anti-mouse IgG Fc




FIG. 20A

Flow cytometry results of representative IgG2a clones selected on CD200-Fc captured with goat anti-mouse IgG Fc




FIG. 20B

Flow cytometry results of representative IgG1 clones selected on directly coated CD200-Fc




FIG. 20C

Flow cytometry results of representative IgG2a clones selected on directly coated CD200-Fc




FIG. 20D

Deduced amino acid sequence of heavy chain complementarity regions of CD200-specific clones

Seq#	HCDR1	HCDR2	HCDR3	Fab	geo mean*
A1	GFNIKDYMH (SEQ ID NO:111)	WID?ENGDTKYAPKFKQG (SEQ ID NO:117)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3A2	291.8
B1	GFNIKDYMH (SEQ ID NO:111)	WID?ENGDTKYAPKFKQG (SEQ ID NO:117)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3A3	486.15
C1	GFNIKDYMH (SEQ ID NO:111)	WID?ENGDTKYAPKFKQG (SEQ ID NO:117)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3A4	340.61
B2	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3B3	131.84
E2	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3B6	576.09
G2	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3B9	407.58
A3	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3B12	363.52
C3	DFNIKDY?H (SEQ ID NO:112)	WIDPE?DDTKYAPKFKQG (SEQ ID NO:119)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3C2	271.55
D3	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3C3	452.53
F3	GLNIKDYMH (SEQ ID NO:113)	WIDPENGNTKYAPKFKQG (SEQ ID NO:120)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3C6	505.75
A5	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	cG2aR3A9	1684.92
A6	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	cG2aR3B5	1819.42
B7	GFNIKDYIH (SEQ ID NO:114)	WIDPDNGDTKYAPKFKRG (SEQ ID NO:121)	KNYVSDYNYFDV (SEQ ID NO:132)	dG1R3A2	1325
D7	GFNIKDYIH (SEQ ID NO:114)	WIDPDNGDTKYAPKFKRD (SEQ ID NO:122)	KNYVSDYNYFDV (SEQ ID NO:132)	dG1R3A4	1245
E7	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:133)	dG1R3A5	1969
G7	GFNIKDYIH (SEQ ID NO:115)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	dG1R3A7	1486
H7	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	dG1R3A8	1852
C8	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	dG1R3A12	981.26
D8	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	dG1R3B2	488.59
E8	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	dG1R3E3	790.06
D9	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	dG1R3B11	1328
E9	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	dG1R3B12	1333
F9	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	dG1R3C1	1782.8
F1	GFNIKDYMH (SEQ ID NO:111)	WIDPENGDTKYAPKFKQG (SEQ ID NO:118)	KNYVSNYNYFDV (SEQ ID NO:131)	cG1R3A9	505.72

* Geometric mean of flow cytometry signal.

FIG. 21a

Deduced amino acid sequence of heavy chain complementarity regions of CD200-specific clones

Seq#	HCDR1	HCDR2	HCDR3	Fab	geo mean*
A4	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDVTKNEKFRG (SEQ ID NO:123)	KRGGYDGAWFAY (SEQ ID NO:134)	cG2aR3A1	?
E4	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDVTKNEKFRG (SEQ ID NO:123)	KRGGYDGAWFAY (SEQ ID NO:134)	cG2aR3A5	2146.54
G4	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDVTKNEKFRG (SEQ ID NO:123)	KRGGYDGAWFAY (SEQ ID NO:134)	cG2aR3A7	586.74
C12	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDVTKNEKFRG (SEQ ID NO:123)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3B9	1563.74
D11	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDVTKNEKFRG (SEQ ID NO:123)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3B1	1942.44
D12	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDVTKNEKFRG (SEQ ID NO:123)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3B10	1668.3
B4	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDITVNEKFKG (SEQ ID NO:124)	KRGGYDGAWFAY (SEQ ID NO:134)	cG2aR3A2	2244.57
B12	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDITVNEKFKG (SEQ ID NO:124)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3B8	1281.05
E6	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDGSKYNEKFKG (SEQ ID NO:125)	KRGGYDGAWFAY (SEQ ID NO:134)	cG2aR3B9	1351.49
D4	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDGTYNEKFKG (SEQ ID NO:126)	KRGGYDGAWFAY (SEQ ID NO:134)	cG2aR3A4	2004.46
C11	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDGTYNEKFKG (SEQ ID NO:126)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3A1	1315.64
A10	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDGTRYNEKFKG (SEQ ID NO:127)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3A1	2043.86
B10	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDGTRYNEKFKG (SEQ ID NO:127)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3A2	2179.3
A12	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDGTRYNEKFKG (SEQ ID NO:127)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3B6	1625.73
F10	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDVTNYNEKFKG (SEQ ID NO:128)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3A7	1814.85
E11	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDGTNYNEKFKG (SEQ ID NO:129)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3B2	2272.54
H10	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDGTNYNEKFKG (SEQ ID NO:129)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3A9	1526.7
F11	GYTFTSYVMH (SEQ ID NO:116)	YINPYNDVTKYNEKFRG (SEQ ID NO:130)	KRGGYDGAWFAY (SEQ ID NO:134)	dG2aR3B3	1882.28

* Geometric mean of flow cytometry signal.

FIG. 21a (Cont.)

Deduced amino acid sequence of heavy chain complementarity regions of CD200-specific clones

Seq#	HCDR1	HCDR2	HCDR3	Fab	geo mean*
D2	GFNIKDHVMH (SEQ ID NO:135)	WIDPENGDTTEYAPKFKQG	(SEQ ID NO:154) FNGYYAMDY	(SEQ ID NO:178) cG1R3B5	299.95
A7	GFNIKDHVMH (SEQ ID NO:135)	WIDPENGDTTEYAPKFKQG	(SEQ ID NO:154) FNGYYAMDY	(SEQ ID NO:178) dG1R3A1	2075
C7	AFNIKDHVMH (SEQ ID NO:136)	WIDPENGDTTEYAPKFKQG	(SEQ ID NO:154) FNGYALDLY	(SEQ ID NO:179) dG1R3A3	624.84
A8	AFNIKDHVMH (SEQ ID NO:136)	WIDPESGDTTEYAPKFKQG	(SEQ ID NO:155) FNGYQALD?	(SEQ ID NO:180) dG1R3A9	1490
B5	AFNIKDHVMH (SEQ ID NO:136)	WIDPESGDTTEYAPKFKQG	(SEQ ID NO:155) FNGYQALDQ	(SEQ ID NO:181) cG2aR3A10	1386.23
H3	AFNIKDHVMH (SEQ ID NO:136)	WIDPESGDTTEYAPKFKQG	(SEQ ID NO:155) FNGYQALDQ	(SEQ ID NO:182) cG1R3C8	399.21
B8	AFNIKDHVMH (SEQ ID NO:136)	WIDPESGDTTEYAPKFKQG	(SEQ ID NO:155) FNGYQALDQ	(SEQ ID NO:181) dG1R3A11	929.93
B6	GFNLKDYVMH (SEQ ID NO:137)	WIDPENGNTTEYAPKFKQG	(SEQ ID NO:156) RNEYVTMDY	(SEQ ID NO:183) cG2aR3B6	1564.28
E5	GFNLKDYVMH (SEQ ID NO:111)	WIDPENGNTTEYAPKFKQ?	(SEQ ID NO:157) RNEYVTMDY	(SEQ ID NO:184) cG2aR3B1	1751.43
E12	GFNLKDYVMH (SEQ ID NO:111)	WIDPENGNTTEYAPKFKQG	(SEQ ID NO:156) RNEYVTMDY	(SEQ ID NO:183) dG2aR3B11	1462.43
A11	GFNLKDYVMH (SEQ ID NO:137)	WIDPENGNTTEYAPKFKQG	(SEQ ID NO:158) RNEYVTMDY	(SEQ ID NO:183) dG2aR2A10	1668.3
H8	AFNIKDHVMH (SEQ ID NO:136)	WIDPESGDTTEYAPKFKQG	(SEQ ID NO:159) FNGYQALDQ	(SEQ ID NO:181) dG1R3B6	1513
D1	GYTFDYWLH (SEQ ID NO:138)	TIDTSTGYTGYNQKFKG	(SEQ ID NO:160) GGDNYVWFAY	(SEQ ID NO:185) cG1R3A6	448.43
E1	GFTFSAAWMD (SEQ ID NO:139)	EIRSKANNAHTYYAASVKG	(SEQ ID NO:161) NGYDDGVFPDY	(SEQ ID NO:186) cG1R3A7	393.89
G1	GYTFEYTMH (SEQ ID NO:140)	GVNPNNGGALYNQKFKG	(SEQ ID NO:162) RSNVRYDDAMDY	(SEQ ID NO:187) cG1R3A10	274.99
A2	GFTFSGFAMS (SEQ ID NO:141)	SISGGTTYLLDSVKG	(SEQ ID NO:163) GNYVSGTSDY	(SEQ ID NO:188) cG1R3B2	476.04
G3	GFTFSGFAMS (SEQ ID NO:141)	SISGGTTYLLDSVKG	(SEQ ID NO:163) GNYVSGTSDY	(SEQ ID NO:188) cG1R3C7	638.59
F7	GFTFSGFAMS (SEQ ID NO:141)	SISGGTTYLLDSVKG	(SEQ ID NO:163) GNYVSGTSDY	(SEQ ID NO:188) dG1R3A6	1665
A9	GFTFGYAMS (SEQ ID NO:142)	SISGGSSAYPPDSVKG	(SEQ ID NO:164) GNYVSGTSDY	(SEQ ID NO:188) dG1R3B7	1994
B9	GFTFSGFAMS (SEQ ID NO:141)	SISGGTTYLLDSVKG	(SEQ ID NO:163) GNYVSGTSDY	(SEQ ID NO:188) dG1R3B9	1569
C9	GFTFSGFAMS (SEQ ID NO:141)	SISGGTTYLLDSVKG	(SEQ ID NO:163) GNYVSGTSDY	(SEQ ID NO:188) dG1R3B10	1839
G9	GFTFSGFAMS (SEQ ID NO:141)	SISGGTTYLLDSVKG	(SEQ ID NO:163) GNYVSGTSDY	(SEQ ID NO:188) dG1R3C2	1586
C2	GFNIKDYIYH (SEQ ID NO:114)	WIDPEIGATKVPKFKQG	(SEQ ID NO:165) LYGNVDRIYAMDY	(SEQ ID NO:189) cG1R3B4	385.24
G8	GFNIKDYIYH (SEQ ID NO:114)	WIDPEIGATKVPKFKQG	(SEQ ID NO:165) LYGNVDRIYAMDY	(SEQ ID NO:189) dG1R3B5	1451

* Geometric mean of flow cytometry signal.


FIG. 21b

Deduced amino acid sequence of heavy chain complementarity regions of CD200-specific clones

Seq#	HCDR1	HCDR2	HCDR3	Fab	geo mean*
F2	GFTFSHAMS (SEQ ID NO:143)	SISGGGTYYPNSVKGR (SEQ ID NO:166)	RGDYRYPYAMDY (SEQ ID NO:190)	cG1R3B7	415.38
H2	GFTFSHAMS (SEQ ID NO:143)	SISGGGTYYPNSVKGR (SEQ ID NO:166)	RGDYRYPYAMDY (SEQ ID NO:190)	cG1R3E10	109.14
E3	GFTFSHAMS (SEQ ID NO:143)	SISGGGTYYPNSVKGR (SEQ ID NO:166)	RGDYRYPYAMDY (SEQ ID NO:190)	cG1R3C4	589.64
H4	GYTFEFTMH (SEQ ID NO:144)	GINPENNGYSYNQKFKG (SEQ ID NO:167)	MITTGYHYAMDY (SEQ ID NO:191)	cG2aR3A8	682.7
H5	GYTFEFTMH (SEQ ID NO:144)	GINPENTGGYSYNQKFKG (SEQ ID NO:168)	MITTGYHYAMDY (SEQ ID NO:191)	cG2aR3B4	711.01
F6	GYTFEFTMH (SEQ ID NO:144)	GINPENTGGYSYNQKFKG (SEQ ID NO:168)	MITTGYHYAMDY (SEQ ID NO:191)	cG2aR3B10	737.93
G6	GYTFEFTMH (SEQ ID NO:144)	GINPENNGYSYNQKFKG (SEQ ID NO:167)	MITTGYHYAMDY (SEQ ID NO:191)	cG2aR3B12	278.53
C10	GYTFEFTMH (SEQ ID NO:144)	GINPENTGGYSYNQKFKG (SEQ ID NO:168)	MITTGYHYAMDY (SEQ ID NO:191)	dG2aR3A4	814.16
D10	GYTFEFTMH (SEQ ID NO:144)	GINPENT?GAYNQKFKG (SEQ ID NO:169)	MITTGYHYAMDY (SEQ ID NO:191)	dG2aR3A5	421.7
G12	GYTFEFTMH (SEQ ID NO:144)	GINPENTGGYSYNQKFKG (SEQ ID NO:168)	MITTGYHYAMDY (SEQ ID NO:191)	dG2aR3C1	1068.43
H12	GYTFEFTMH (SEQ ID NO:144)	GINPENTGGYSYNQKFKG (SEQ ID NO:168)	MITTGYHYAMDY (SEQ ID NO:191)	dG2aR3C	1106.7
C5	GYTFEFTMH (SEQ ID NO:145)	WISPGSLNTNNEKFKG (SEQ ID NO:169)	KARGDSGAWFAY (SEQ ID NO:192)	cG2aR3A11	1311.44
D5	GYTFEFTMH (SEQ ID NO:146)	WISPGSLNTNNEKFKG (SEQ ID NO:169)	KARGDSGAWFAY (SEQ ID NO:192)	cG2aR3A12	887.54
B11	GYTFEFTMH (SEQ ID NO:146)	WISPGSLNTNNEKFKG (SEQ ID NO:169)	KARGDSGAWFAY (SEQ ID NO:192)	dG2aR3A11	1588.4
F5	GYTFDYWMH (SEQ ID NO:147)	AIDTFDSNTKYNQKFKG (SEQ ID NO:170)	GVVDY (SEQ ID NO:193)	cG2aR3B2	1491.49
G10	GYTFDYWMH (SEQ ID NO:147)	AIDTFDSNTRYNQKFKG (SEQ ID NO:171)	GVVDY (SEQ ID NO:193)	dG2aR3A8	1424.53
H11	GYTFDYWMH (SEQ ID NO:147)	AIDTFDSNTRYNQKFKG (SEQ ID NO:172)	GVVDY (SEQ ID NO:193)	dG2aR3B5	1259.43
G5	GYTFDNIWH (SEQ ID NO:148)	TIDASDRYISYNQKFKG (SEQ ID NO:173)	LEGSGYGFAY (SEQ ID NO:194)	cG2aR3B3	1289.19
F8	GYTFDNIWH (SEQ ID NO:148)	TIDASDRYISYNQKFKG (SEQ ID NO:173)	LEGSGYGFAY (SEQ ID NO:194)	dG1R3B4	1340
C6	GYSTFDYLL (SEQ ID NO:149)	HIDPYYGSSNMLKFKG (SEQ ID NO:174)	SKRDYFDY (SEQ ID NO:195)	cG2aR3B7	1776.38
H6	GFNIKSYIH (SEQ ID NO:150)	WIDPENGTEYAPKFKG (SEQ ID NO:175)	CNFYGNPFYDY (SEQ ID NO:196)	cG2aR3C1	1833.02
F12	GFNIK?SYIH (SEQ ID NO:151)	WIDPENGTEYAPKFKG (SEQ ID NO:175)	CNFYANDPFYDY (SEQ ID NO:197)	dG2aR3B12	1350.25
H9	GYTFEFTMH (SEQ ID NO:152)	YINPSSGYTNNQKFKD (SEQ ID NO:176)	RPMTAGAWFAY (SEQ ID NO:198)	dG1R3H8	2470
E10	GYTFEFTMH (SEQ ID NO:153)	GINPNTGAYNNQKFKG (SEQ ID NO:177)	ITTVVGYYYAMDY (SEQ ID NO:199)	dG2aR3A6	796.81

* Geometric mean of flow cytometry signal.

FIG. 21b (Cont.)




R9.4 is a chimeric antibody derived from a rabbit scFv9. 9H is an anti-FLJ32028 antibody and was included as a negative control.

FIG. 22

Deduced amino acid sequences of selected CD200 Fabs for chimerization

Heavy chain								
Fab		FR1	CDR1	FR2	CDR2			
d1B10	LEVKLVEGGGLVKKPGGSLKLSCAAS	GFTFSGFAMS	WVRQTPFKRLEWVA	SISSGGTTYILDSVKG	(SEQ ID NO: 200)			
d1A5	LEVQLQOQSGAELVRSASVKLSCTAS	GFNKIDYYMH	WVKORPEQGLEWIG	WIDPENGDTKYAPKFGQ	(SEQ ID NO: 201)			
d1B5	LEVQLQOQSGAELVRSASVKLSCKAS	GFNKIDYYIH	WVKORPEQGLEWIG	WIDPEIGATKYVPKFGQ	(SEQ ID NO: 202)			
c2aB7	LEVQLQOQSGPELVKPGASLKMCKAS	GYSFTDYILL	WVKQNHGKSLEWIG	HIDPHYGSSNYNLKFKG	(SEQ ID NO: 203)			
c1A10	LEVQLQOQSGPELVKPGASVKISCKTS	GYTFTEYTMH	WVKQSHGKSLEWIG	GVNPNNGGALYNQKFKG	(SEQ ID NO: 204)			
c2aA10	LEVQLQOQSGAELVRSASVKLSCTAS	AFNIKDHMH	WVKORPEQGLEWIG	WIDPESGDTFYAPKFGQ	(SEQ ID NO: 205)			
Fab		FR3	CDR3	FR4				
d1B10	RFTISRDIARNILYLQMSLSRSEDAMVYCAR	GNYSGTSYDY	WGQGTLLTVSS	(SEQ ID NO: 200)	- cont'd)			
d1A5	KATMTADTSSNTAYLQLSLSLTSEDVAVYICNA	KNYYVSNYNFFDV	WGAGTIVTVSS	(SEQ ID NO: 201)	- cont'd)			
d1B5	KATMTTDTSSNTAYLQLSLSLTSEDVAVYICNA	LYGNIDRYAYAMDY	WGQGTSTVTVSS	(SEQ ID NO: 202)	- cont'd)			
c2aB7	KATLTVDKSSSTAYMQLNSLTSEDSAVYICGR	SKRDYFDY	WGQGTLLTVSS	(SEQ ID NO: 203)	- cont'd)			
c1A10	KATLTVDKSSSTAYMELRSLASEDSAVYICAR	RSNRYDDAMDY	WGQGTSTVTVSS	(SEQ ID NO: 204)	- cont'd)			
c2aA10	KATMTADISSNTAYLQLNSLTSEDVAVYICNA	FNGYQALDQ	WGQGTSTVTVSS	(SEQ ID NO: 205)	- cont'd)			
Light chains								
Fab		FR1	CDR1	FR2	CDR2			
d1B10	SRDIVLTQSPASLAVSLGQRATISC	RASESVDSYGNSEFMH	WYQOKPGQPPKLLIY	RASNLES	(SEQ ID NO: 206)			
d1A5	SREIVLTQSPATMSASPGKEKVTMTCTC	SASSVRYMY	WYQOKSSTSPKLIWY	DTSKLAS	(SEQ ID NO: 207)			
d1B5	SRDIVMTQSQKFMSTSVGDRVSIITC	KASQNVRTAVA	WYQOKPGQSPKALIIY	LASNRHT	(SEQ ID NO: 208)			
c2aB7	SRDIQMTQSPSSMYASLGERVTITC	KASQDINSYLS	WFQOKPGKSPKTLIIY	RANRLVD	(SEQ ID NO: 209)			
c1A10	SRDVVMTQTPLTFLSVTIGQPASISCTC	KSSQSLLDIDEKTYLN	WFLQRPQSPKRLIIY	LVSKLDS	(SEQ ID NO: 210)			
c2aA10	SREIVLTQSPAIMASALGERVTMTCTC	TASSVSSSYLH	WYQOKPGSSPKLIWY	STSNLAS	(SEQ ID NO: 211)			
Fab		FR3	CDR3	FR4				
d1B10	GIPARFSGSGSRRTDFTLLTINPVEADDAVATYYC	QQSNEDPRT	FGGGTKLEIKR	(SEQ ID NO: 206)	- cont'd)			
d1A5	GVPGRFSGSGSGNSYSLTISMEAEADVAATYYC	FQSGGYPLT	FGSGTKLEIKR	(SEQ ID NO: 207)	- cont'd)			
d1B5	GVPDRFTGSGSGTDFTLTISNVQSEDLADYFC	LQHWNYPLT	FGAGTKLELKR	(SEQ ID NO: 208)	- cont'd)			
c2aB7	GVPDRFSGSGGQDYSLTISLLEYEDMGIYYC	LQYDEFPPY	FGGGTKLEIKR	(SEQ ID NO: 209)	- cont'd)			
c1A10	GVPDRFTGSGSGTDFTLKISRVEAEDLGVYYC	WQGTFFPQT	FGGGTKLEIKR	(SEQ ID NO: 210)	- cont'd)			
c2aA10	GVPARFSGSGGTSYSLTISMEAEADAATYYC	RQYHRSPPIFT	FGSGTKLEIKR	(SEQ ID NO: 211)	- cont'd)			

FIG. 23



ELISA results of chimeric IgG obtained from the culture supernatant of a small-scale transient transfection.

FIG. 24




FIG. 26C




FIG. 25




FIG. 26A




FIG. 26B




FIG. 27

RAJI/PBL MODEL




FIG. 28

NAMALWA/ PBL MODEL




FIG. 29

IMMUNOSUPPRESSIVE EFFECT OF THE CD200 IN THE RAJI/PBL MODEL




FIG. 30a

IMMUNOSUPPRESSIVE EFFECT OF THE CD200 IN THE RAJI/PBL MODEL




FIG. 30b

IMMUNOSUPPRESSIVE EFFECT OF THE CD200 IN THE RAJI/PBL MODEL




FIG. 30c

IMMUNOSUPPRESSIVE EFFECT OF THE CD200 IN THE NAMALWA/PBL MODEL

Comparison of lentivirus transduced cells with parental cells




FIG. 31a

IMMUNOSUPPRESSIVE EFFECT OF THE CD200 IN THE NAMALWA/PBL MODEL




FIG. 31b

IMMUNOSUPPRESSIVE EFFECT OF THE CD200 IN THE NAMALWA/PBL MODEL




FIG. 31c

IMMUNOSUPPRESSIVE EFFECT OF THE CD200 IN THE NAMALWA/PBL MODEL




FIG. 31d




FIG. 32

Figure 33





Figure 34



专利名称(译)	衍生自慢性淋巴细胞白血病细胞的多肽和抗体及其用途		
公开(公告)号	EP1786464A2	公开(公告)日	2007-05-23
申请号	EP2005774630	申请日	2005-07-19
申请(专利权)人(译)	ALEXION制药公司.		
当前申请(专利权)人(译)	ALEXION制药公司.		
[标]发明人	BOWDISH KATHERINE S KRETZ ROMMEL ANKE MCWHIRTER JOHN MARUYAMA TOSHIAKI		
发明人	BOWDISH, KATHERINE, S. KRETZ-ROMMEL, ANKE MCWHIRTER, JOHN MARUYAMA, TOSHIAKI		
IPC分类号	A61K39/395 A61K39/00 A61K51/00 A61K51/10 C07K16/18 C07K16/28 C07K16/30 C07K16/46 G01N33/53 G01N33/574		
CPC分类号	C07K16/2803 A61K47/6849 A61K2039/505 C07K16/3061 C07K2317/24 C07K2317/622 C07K2317 /732 C07K2317/76 C07K2317/92 G01N33/57426 G01N2333/70596 G01N2800/52		
代理机构(译)	法思博事务所		
优先权	10/996316 2004-11-23 US 11/171567 2005-06-30 US 10/894672 2004-07-20 US		
其他公开文献	EP1786464A4		
外部链接	Espacenet		

摘要(译)

提供了用于评估化合物的免疫调节作用的小动物模型。