

(43) International Publication Date
18 September 2014 (18.09.2014)

WIP O | P C T

(10) International Publication Number
WO 2014/144821 A1

(51) International Patent Classification:

G01N 33/68 (2006.01) **C12Q 1/68** (2006.01)
G01N 33/53 (2006.01)

(21) International Application Number:

PCT/US2014/029392

(22) International Filing Date:

14 March 2014 (14.03.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/801,290	15 March 2013 (15.03.2013)	US
61/801,476	15 March 2013 (15.03.2013)	US
61/874,947	6 September 2013 (06.09.2013)	US


(71) Applicant: **INTERMUNE, INC.** [US/US]; 3280, Bayshore Boulevard, Brisbane, CA 94005-1021 (US).(72) Inventors: **KOSSEN, Karl**; 1533, La Mesa Dr., Burlingame, CA 94010 (US). **QIN, Xiaoli**; 426, Sonora Drive, San Mateo, CA 94402 (US). **LIM, Sharleen, R.**; 221, Cupertino Way, San Mateo, CA 94403 (US). **SEIWERT, Scott, D.**; 419, Seymour Street, Half Moon Bay, CA 94019 (US). **RUHRMUND, Donald**; 2178, 40th Avenue, San Francisco, CA 94116 (US).(74) Agents: **HETZER-EGGER, Claudia, H.** et al; Kilpatrick Townsend & Stockton LLP, Two Embarcadero Center, 8th Floor, San Francisco, CA 94111 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

[Continued on nextpage]

(54) Title: PROTEOMIC IPF MARKERS

FIG. 2



(57) Abstract: Provided herein are novel biomarkers for the identification, prediction or monitoring of fibrotic pulmonary diseases (e.g. idiopathic pulmonary fibrosis (IPF)). The biomarkers provided herein may further be used for therapeutic treatment of IPF.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

PROTEOMIC IPF MARKERS

CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] This application claims priority to US Provisional Application No. 61/801,290, filed March 15, 2013, US Provisional Application No. 61/801,476, filed March 15, 2013, and US Provisional Application No. 61/874,947, filed September 6, 2013, the disclosures of which are incorporated herein in their entireties.

BACKGROUND OF THE INVENTION

[0002] The idiopathic interstitial pneumonias (IIPs) represents a class of chronic pulmonary fibrotic disorders characterized by progressive scarring of the alveolar interstitium leading to severe dyspnea, hypoxemia, and death. Among interstitial lung disease idiopathic pulmonary fibrosis (IPF) is the most progressive disease with median survival of 2.5-3 years (ATS/ERS. Am J Respir Crit Care Med 2002; 165(2): 277-304). Patients diagnosed with IPF typically experience progressive pulmonary insufficiency, and most die of respiratory failure. The ratio of the estimated prevalence (90,000 individuals) and incidence (30,000 individuals) of IPF in the United States reflects this poor prognosis (Raghu G, Weycker D, Edelsberg J, Bradford WZ, Oster G. Incidence and prevalence of idiopathic pulmonary fibrosis. Am J Respir Crit Care Med 2006; 174(7): 810-816).. The clinical course of IPF is highly variable and may be characterized by periods of relative stability, chronic decline, or periods of rapid decline (known as acute exacerbations).

[0003] Clinical and physiological parameters (e.g., forced vital capacity, diffusing capacity of carbon monoxide DLco, six minute walk test distance) are currently used to monitor disease state and progression. These clinical assessments of IPF and risk prediction models that incorporate these components (or change therein) are relatively successful in staging patients and predicting risk of mortality. However, management approaches that are based on molecular information (such as that derived from protein markers) may enable better monitoring of disease progression and risk of progression in patients with IPF.

[0004] There is a need in the art for well-defined biomarkers to determine if a patient has or is at risk for developing a fibrotic pulmonary disease, to indicate the progressiveness of the disease, and/or to facilitate evaluation of responsiveness to therapy. The invention provided herein

addresses these and other needs in the art by providing, inter alia, novel protein biomarkers of fibrotic pulmonary diseases (e.g. IPF).

BRIEF SUMMARY OF THE INVENTION

- 5 **[0005]** In one aspect, a method of determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB in a subject that has or is at risk for developing a fibrotic pulmonary disease is provided. The method includes (i) obtaining a biological sample from said subject and (ii) determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB in the biological sample.
- 10 **[0006]** In another aspect, a method of determining whether a subject has or is at risk of developing a fibrotic pulmonary disease is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB in a subject; (ii) determining whether the expression level is increased or decreased relative to a standard control, wherein an elevated expression level of a fibrotic pulmonary
- 15 disease marker protein in Table 1A or a decreased expression level of a fibrotic pulmonary disease marker protein in Table IB relative to the standard control indicates that the subject has or is at risk of developing a fibrotic pulmonary disease; and (iii) based at least in part on the expression level in step (ii), determining whether the subject has or is at risk for developing a fibrotic pulmonary disease.
- 20 **[0007]** In another aspect, a method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB in a fibrotic pulmonary disease patient (ii) determining whether the expression level is modulated relative to a standard control, wherein a modulated
- 25 expression level of a fibrotic pulmonary disease marker protein in Table 1A or Table IB relative to the standard control indicates that the fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, and (iii) based at least in part on the expression level in step (ii), determining whether the fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease.
- 30 **[0008]** In another aspect, a method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease is provided. The method includes (i) determining a first expression level of a protein set forth in Table 1A or Table IB in

the patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A or Table IB in the patient at a second time point; and (iii) comparing the second expression level of a protein set forth in Table 1A or Table IB to the first expression level of a protein set forth in Table 1A or Table IB, wherein when the second expression level of a protein set forth in Table 1A is greater than the first level of a protein set forth in Table 1A; or wherein when the second expression level of a protein set forth in Table IB is smaller than the first level of a protein set forth in Table IB, the patient is at risk for progression of the fibrotic pulmonary disease.

[0009] In another aspect, a method of determining a fibrotic pulmonary disease activity in a patient is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB in a subject; (ii) determining whether the expression level is modulated relative to a standard control, thereby determining a fibrotic pulmonary disease activity in the patient; and (iii) based at least in part on the expression level in step (ii), determining the fibrotic pulmonary disease activity in the patient.

[0010] In another aspect, a method of determining a fibrotic pulmonary disease activity in a patient is provided. The method includes (i) determining a first expression level of a protein set forth in Table 1A or Table IB in the patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A or Table IB in the patient at a second time point; (iii) comparing the second expression level of a protein set forth in Table 1A or Table IB to the first expression level of a protein set forth in Table 1A or Table IB, thereby determining the fibrotic pulmonary disease activity in the patient.

[0011] In another aspect, a method of determining a change in fibrotic pulmonary disease activity in a patient is provided. The method includes (i) determining a first expression level of a protein set forth in Table 1A or Table IB in the patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A or Table IB in the patient at a second time point; and (iii) comparing the second expression level of a protein set forth in Table 1A or Table IB to the first expression level of a protein set forth in Table 1A or Table IB, thereby determining the a change in fibrotic pulmonary disease activity in the patient.

[0012] In another aspect, a method of determining a change in fibrotic pulmonary disease activity in a patient is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB in a subject;

(ii) determining whether the expression level is modulated relative to a standard control, thereby determining a change in fibrotic pulmonary disease activity in the patient; and (iii) based at least in part on the expression level in step (ii), determining the change in fibrotic pulmonary disease activity in the patient.

5 **[0013]** In another aspect, provided herein is a method of monitoring the effect of treatment for a fibrotic pulmonary disease in a patient undergoing fibrotic pulmonary disease therapy or a patient that has received fibrotic pulmonary disease therapy. The method includes (i) determining a first expression level of a protein set forth in Table 1A or Table IB in the patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A
10 or Table IB in the patient at a second time point; and (iii) comparing the second expression level of a protein set forth in Table 1A or Table IB to the first expression level of a protein set forth in Table 1A or Table IB, thereby determining the effect of treatment for a fibrotic pulmonary disease in the patient.

[0014] In another aspect, provided herein is a method of monitoring the effect of treatment for
15 a fibrotic pulmonary disease in a patient undergoing fibrotic pulmonary disease therapy or a patient that has received fibrotic pulmonary disease therapy. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB in a subject; (ii) determining whether the expression level is modulated relative to a standard control, thereby monitoring the effect of treatment for a fibrotic pulmonary disease
20 in the patient; and (iii) based at least in part on the expression level in step (ii), monitoring the effect of treatment for a fibrotic pulmonary disease in the patient.

[0015] In another aspect, a method of treating a fibrotic pulmonary disease in a subject in need thereof is provided. The method includes administering to the subject an effective amount of an modulator of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB,
25 thereby treating a fibrotic pulmonary disease in said subject.

[0016] In one aspect, a method of treating a fibrotic pulmonary disease in a subject in need thereof is provided. The method includes (i) determining whether a subject expresses an elevated level of a fibrotic pulmonary disease marker protein as set forth in Table 1A or a decreased level of a fibrotic pulmonary disease marker protein as set forth in Table IB relative to
30 a standard control, and (ii) when an elevated expression level of the fibrotic pulmonary disease marker protein of Table 1A or a decreased expression level of the fibrotic pulmonary disease marker protein of Table IB is found relative to the standard control, administering to the subject

a fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker protein set forth in Table 1A or an agonist of a fibrotic pulmonary disease marker protein set forth in Table 1B, thereby treating the subject.

[0017] In another aspect, a method of determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B in a subject that has or is at risk for developing a fibrotic pulmonary disease is provided. The method includes (i) obtaining a biological sample from the subject, and (ii) determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B in the biological sample.

[0018] In another aspect, a method of determining whether a subject has or is at risk of developing a fibrotic pulmonary disease is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject, (ii) determining whether said expression level is increased or decreased relative to a standard control, wherein an elevated expression level of a fibrotic pulmonary disease marker protein in Table 3A or a decreased expression level of a fibrotic pulmonary disease marker protein in Table 3B relative to the standard control indicates that the subject has or is at risk of developing a fibrotic pulmonary disease, and (iii) based at least in part on the expression level in step (ii), determining whether the subject has or is at risk for developing a fibrotic pulmonary disease.

[0019] In another aspect, a method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker protein set forth in Table 3A or 3B in a fibrotic pulmonary disease patient, (ii) determining whether the expression level is modulated relative to a standard control, wherein a modulated expression level of a fibrotic pulmonary disease marker protein in Table 3A or 3B relative to the standard control indicates that the fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, and (iii) based at least in part on the expression level in step (ii), determining whether the fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease.

[0020] In one aspect, a method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease is provided. The method includes (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in the patient at a first time point, (ii) determining a second expression level of a protein set forth in Table 3A

or Table 3B in the patient at a second time point, (iii) comparing the second expression level of a protein set forth in Table 3A or Table 3B to the first expression level of a protein set forth in Table 3A or Table 3B, wherein when the second expression level of a protein set forth in Table 3A is greater than the first level of a protein set forth in Table 3A, or wherein when the second expression level of a protein set forth in Table 3B is smaller than the first level of a protein set forth in Table 3B, the patient is at risk for progression of the fibrotic pulmonary disease.

[0021] In another aspect, a method of determining a fibrotic pulmonary disease activity in a patient is provided. The method includes, (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject; (ii) determining whether the expression level is modulated relative to a standard control, thereby determining a fibrotic pulmonary disease activity in the patient; and (iii) based at least in part on the expression level in step (ii), determining said fibrotic pulmonary disease activity in the patient.

[0022] In another aspect, a method of determining a fibrotic pulmonary disease activity in a patient is provided. The method includes (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in the patient at a first time point, (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in the patient at a second time point, (iii) comparing the second expression level of a protein set forth in Table 3A or Table 3B to the first expression level of a protein set forth in Table 3A or Table 3B, thereby determining the fibrotic pulmonary disease activity in the patient.

[0023] In another aspect, a method of determining a change in fibrotic pulmonary disease activity in a patient is provided. The method includes (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in the patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in the patient at a second time point; and (iii) comparing the second expression level of a protein set forth in Table 3A or Table 3B to the first expression level of a protein set forth in Table 3A or Table 3B, thereby determining a change in fibrotic pulmonary disease activity in a patient.

[0024] In another aspect, a method of determining a change in fibrotic pulmonary disease activity in a patient is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject; (ii) determining whether the expression level is modulated relative to a standard control, thereby determining a change in fibrotic pulmonary disease activity in the patient; and (iii) based at least

in part on the expression level in step (ii), determining the change in fibrotic pulmonary disease activity in the patient.

[0025] In another aspect, provided herein is a method of monitoring the effect of treatment for a fibrotic pulmonary disease in a patient undergoing fibrotic pulmonary disease therapy or a patient that has received fibrotic pulmonary disease therapy. The method includes (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in the patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in the patient at a second time point; and (iii) comparing the second expression level of a protein set forth in Table 3A or Table 3B to the first expression level of a protein set forth in Table 3A or Table 3B, thereby monitoring the effect of treatment for a fibrotic pulmonary disease in the patient.

[0026] In another aspect, provided herein is a method of monitoring the effect of treatment for a fibrotic pulmonary disease in a patient undergoing fibrotic pulmonary disease therapy or a patient that has received fibrotic pulmonary disease therapy. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject; (ii) determining whether the expression level is modulated relative to a standard control, thereby monitoring the effect of treatment for a fibrotic pulmonary disease in the patient; and (iii) based at least in part on the expression level in step (ii), monitoring the effect of treatment for a fibrotic pulmonary disease in the patient

[0027] In another aspect, a method of treating a fibrotic pulmonary disease in a subject in need thereof is provided. The method includes administering to the subject an effective amount of an modulator of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B, thereby treating a fibrotic pulmonary disease in the subject.

[0028] In another aspect, a method of treating a fibrotic pulmonary disease in a subject in need thereof is provided. The method includes (i) determining whether a subject expresses an elevated level of a fibrotic pulmonary disease marker protein as set forth in Table 3A or a decreased level of a fibrotic pulmonary disease marker protein as set forth in Table 3B relative to a standard control, and (ii) when an elevated expression level of the fibrotic pulmonary disease marker protein of Table 3A or a decreased expression level of the fibrotic pulmonary disease marker protein of Table 3B is found relative to the standard control, administering to the subject a fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker

protein set forth in Table 3A or an agonist of a fibrotic pulmonary disease marker protein set forth in Table 3B, thereby treating the subject.

[0029] In one aspect a kit is provided. The kit includes a marker protein binding agent capable of binding to a substance within a biological sample from a human subject having or at risk of developing a fibrotic pulmonary disease, wherein the substance is a fibrotic pulmonary disease marker protein or fragment thereof set forth in the tables provided herein (e.g., Table 1A, IB, 2A, 2B, 3A, 3B, 4A, or 4B); and a detecting reagent or a detecting apparatus capable of indicating binding of the marker protein binding agent to the substance.

[0030] In one aspect a kit is provided. The kit includes a marker protein binding agent capable of binding to a substance within a biological sample from a human subject having a fibrotic pulmonary disease, wherein the substance is a fibrotic pulmonary disease marker protein or fragment thereof set forth in the tables provided herein (e.g., Table 1A, IB, 2A, 2B, 3A, 3B, 4A, or 4B); and a detecting reagent or a detecting apparatus capable of indicating binding of the marker protein binding agent to the substance.

[0031] In another aspect, a complex in vitro is provided. The complex includes a marker protein binding agent bound to a fibrotic pulmonary disease marker protein or fragment thereof set forth in the tables provided herein (e.g., Table 1A, IB, 2A, 2B, 3A, 3B, 4A, or 4B), wherein the fibrotic pulmonary disease marker protein is extracted from a human subject having or at risk of developing a fibrotic pulmonary disease.

[0032] In another aspect, a complex in vitro is provided. The complex includes a marker protein binding agent bound to a fibrotic pulmonary disease marker protein or fragment thereof set forth in the tables provided herein (e.g., Table 1A, IB, 2A, 2B, 3A, 3B, 4A, or 4B), wherein the fibrotic pulmonary disease marker protein is extracted from a human subject having a fibrotic pulmonary disease.

BRIEF DESCRIPTION OF THE DRAWINGS

[0033] Fig.1. Overview of the proteomics assay. (Step 1) The specific protein to be measured (P_i) binds tightly to its cognate SOMAmer binding molecule (S), which includes a photo-cleavable biotin (PCB) and fluorescent label (L) at the 5' end. **(Step 2)** Bound protein-SOMAmer complexes are captured onto streptavidin coated beads (SA) by photo-cleavable biotin on the SOMAmer. Unbound proteins are washed away. Bound proteins are tagged with biotin (B). **(Step 3)** The photo-cleavable biotin is cleaved by UV light (hn) and the protein-SOMAmer complexes are released into solution. **(Step 4)** The protein-SOMAmer complexes are

captured onto SA and the SOMAmers are eluted into solution (Step 5) and recovered for quantification in **Step 6**, hybridization to a custom DNA microarray. Each probe spot contains DNA with sequence complementary to a specific SOMAmer, and the fluorescent intensity of each probe spot is proportional to the amount of SOMAmer recovered, which is proportional to the amount of protein present in the original sample. PCB: Photo-cleavable biotin; SOMAmer: Slow off-rate modified aptamer. (reference: High-content affinity-based proteomics: unlocking protein biomarker discovery, Edward N Brody¹, Larry Gold^{†1,2}, Richard M Lawn¹, Jeffrey J Walker¹ and Dom Zichil Expert Review of Molecular Diagnostics, November 2010, Vol. 10, No. 8, Pages 1013-1022 , DOI 10.1586/erm.10.89).

10 **[0034] Fig.2:** Comparison of relative rank of pathways defined by differentially expressed proteins to those defined by the proteins on the panel as a whole.

[0035] Fig.3: Hierarchical Clustering of IPF patients.

[0036] Fig.4: Hierarchical Clustering of IPF patients.

[0037] Fig.5 - Fig.47: Fibrotic pulmonary disease biomarker proteins that predict disease progression in patient cohort #1 using aptamer based proteomic platform for determination of protein levels. The following description applies to the histograms shown in Fig.5-Fig.47:

15 **[0038]** (Upper left histogram) Kaplan-Meier curve for Progression Free Survival (PFS). X-axis indicates days, y-axis indicates percent survival. Subjects are grouped based on a protein level cut-off suggested by CART analysis. Number of subjects above and below cut-off are indicated (i.e. "high N" and "low N"). Unadjusted p-value for significance is shown.

[0039] (Upper right histogram) Kaplan-Meier curve for Progression Free Survival (PFS). X-axis indicates days, y-axis indicates percent survival. Subjects are grouped based on quartiles of protein levels (i.e. Q1 indicates the lowest quartile). Unadjusted p-values for significance compared to Q1 are shown.

25 **[0040]** (Lower left histogram) Kaplan-Meier curve for All-Cause Mortality. X-axis indicates days, y-axis indicates percent survival. Subjects are grouped based on a protein level cut-off suggested by CART analysis. Number of subjects above and below cut-off are indicated (i.e. "high N" and "low N"). Unadjusted p-value for significance is shown.

30 **[0041]** (Lower right histogram) Kaplan- Meier curve for All-Cause Mortality. X-axis indicates days, y-axis indicates percent survival. Subjects are grouped based on quartiles of protein levels

(i.e. Q1 indicates the lowest quartile). Unadjusted p-values for significance compared to Q1 are shown.

[0042] Fig.48 - Fig.57: Selected fibrotic pulmonary disease biomarker proteins that predict disease progression in two independent patient cohorts. Plasma protein level was quantitated by ELISA. The following description applies to the plots shown in Fig.48 - Fig.57:

[0043] (Upper left plot) Kaplan-Meier Curve for Progression Free Survival (PFS). X-axis indicates days, Y-axis indicates proportion of survival. Subjects are grouped based on median (Fig. 48-Fig. 53 and Fig. 56-57) or tertiles (Fig. 54 and 55) of protein levels. p-values of Log-rank test are shown.

[0044] (Upper right plot) Kaplan-Meier Curve for Progression Free Survival (PFS). X-axis indicates days, Y-axis indicates proportion of survival. Subjects are grouped based on a protein level cut-off determined by CART analysis. Number of subjects above and below cut-off are indicated (i.e. "High N" and "Low N"). p-values of Log-rank test are shown.

[0045] (Lower left plot) Kaplan-Meier Curve for All-Cause Mortality. X-axis indicates days, Y-axis indicates proportion of survival. Subjects are grouped based on median (Fig. 48- Fig. 53 and Fig. 56-Fig. 57) or tertiles (Fig. 54 and Fig. 55) of protein levels. p-values of Log-rank test are shown.

[0046] (Lower right plot) Kaplan-Meier Curve for All-Cause Mortality. X-axis indicates days, Y-axis indicates proportion of survival. Subjects are grouped based on a protein level cut-off determined by CART analysis. Number of subjects above and below cut-off are indicated (i.e. "High N" and "Low N"). p-values of Log-rank test are shown.

DETAILED DESCRIPTION OF THE INVENTION

I. DEFINITIONS

[0047] The term "disease" refers to any deviation from the normal health of a mammal and includes a state when disease symptoms are present, as well as conditions in which a deviation (e.g., infection, gene mutation, genetic defect, etc.) has occurred, but symptoms are not yet manifested. According to the present invention, the methods disclosed herein are suitable for use in a patient that is a member of the Vertebrate class, Mammalia, including, without limitation, primates, livestock and domestic pets (e.g., a companion animal). Typically, a patient will be a human patient.

[0048] The terms "pulmonary disease," "pulmonary disorder," "lung disease," etc. are used interchangeably herein. The term is used to broadly refer to lung disorders characterized by difficulty breathing, coughing, airway discomfort and inflammation, increased mucus, and/or pulmonary fibrosis.

5 [0049] An "airway mucosal sample" can be obtained using methods known in the art, *e.g.*, a bronchial epithelial brush as described herein. Additional methods include endobronchial biopsy, bronchial wash, bronchoalveolar lavage, whole lung lavage, transendoscopic biopsy, and transtracheal wash.

10 [0050] The terms "subject," "patient," "individual," and the like as used herein are not intended to be limiting and can be generally interchanged. That is, an individual described as a "patient" does not necessarily have a given disease, but may be merely seeking medical advice.

[0051] The term "subject" as used herein includes all members of the animal kingdom prone to suffering from the indicated disorder. In some aspects, the subject is a mammal, and in some aspects, the subject is a human.

15 [0052] A "control" sample or value refers to a sample that serves as a reference, usually a known reference, for comparison to a test sample. For example, a test sample can be taken from a patient suspected of having a given pulmonary disease and compared to samples from a known pulmonary disease patient, or a known normal (non-disease) individual. A control can also represent an average value gathered from a population of similar individuals, *e.g.*, pulmonary
20 disease patients or healthy individuals with a similar medical background, same age, weight, etc. A control value can also be obtained from the same individual, *e.g.*, from an earlier-obtained sample, prior to disease, or prior to treatment. One of skill will recognize that controls can be designed for assessment of any number of parameters.

25 [0053] One of skill in the art will understand which controls are valuable in a given situation and be able to analyze data based on comparisons to control values. Controls are also valuable for determining the significance of data. For example, if values for a given parameter are widely variant in controls, variation in test samples will not be considered as significant.

[0054] As used herein, the terms "pharmaceutically" acceptable is used synonymously with physiologically acceptable and pharmacologically acceptable. A pharmaceutical composition
30 will generally comprise agents for buffering and preservation in storage, and can include buffers and carriers for appropriate delivery, depending on the route of administration.

[0055] The terms "dose" and "dosage" are used interchangeably herein. A dose refers to the amount of active ingredient given to an individual at each administration. For the present invention, the dose will generally refer to the amount of pulmonary disease treatment, anti-inflammatory agent, agonist or antagonist. The dose will vary depending on a number of factors, including the range of normal doses for a given therapy, frequency of administration; size and tolerance of the individual; severity of the condition; risk of side effects; and the route of administration. One of skill will recognize that the dose can be modified depending on the above factors or based on therapeutic progress. The term "dosage form" refers to the particular format of the pharmaceutical, and depends on the route of administration. For example, a dosage form can be in a liquid form for nebulization, *e.g.*, for inhalants, in a tablet or liquid, *e.g.*, for oral delivery, or a saline solution, *e.g.*, for injection.

[0056] As used herein, the terms "treat" and "prevent" are not intended to be absolute terms. Treatment can refer to any delay in onset, reduction in the frequency or severity of symptoms, amelioration of symptoms, improvement in patient comfort and/or respiratory function, etc. The effect of treatment can be compared to an individual or pool of individuals not receiving a given treatment, or to the same patient prior to, or after cessation of, treatment.

[0057] "Treating" or "treatment" as used herein (and as well-understood in the art) also broadly includes any approach for obtaining beneficial or desired results in a subject's condition, including clinical results. Beneficial or desired clinical results can include, but are not limited to, alleviation or amelioration of one or more symptoms or conditions, diminishment of the extent of a disease, stabilizing (*i.e.*, not worsening) the state of disease, prevention of a disease's transmission or spread, delay or slowing of disease progression, amelioration or palliation of the disease state, diminishment of the reoccurrence of disease, and remission, whether partial or total and whether detectable or undetectable. In other words, "treatment" as used herein includes any cure, amelioration, or prevention of a disease. Treatment may prevent the disease from occurring; inhibit the disease's spread; relieve the disease's symptoms (*e.g.*, ocular pain, seeing halos around lights, red eye, very high intraocular pressure), fully or partially remove the disease's underlying cause, shorten a disease's duration, or do a combination of these things.

[0058] "Treating" and "treatment" as used herein include prophylactic treatment. Treatment methods include administering to a subject a therapeutically effective amount of an active agent. The administering step may consist of a single administration or may include a series of administrations. The length of the treatment period depends on a variety of factors, such as the

severity of the condition, the age of the patient, the concentration of active agent, the activity of the compositions used in the treatment, or a combination thereof. It will also be appreciated that the effective dosage of an agent used for the treatment or prophylaxis may increase or decrease over the course of a particular treatment or prophylaxis regime. Changes in dosage may result and become apparent by standard diagnostic assays known in the art. In some instances, chronic administration may be required. For example, the compositions are administered to the subject in an amount and for a duration sufficient to treat the patient.

[0059] The term "prevent" refers to a decrease in the occurrence of pulmonary disease symptoms in a patient. As indicated above, the prevention may be complete (no detectable symptoms) or partial, such that fewer symptoms are observed than would likely occur absent treatment.

[0060] The term "therapeutically effective amount," as used herein, refers to that amount of the therapeutic agent sufficient to ameliorate the disorder, as described above. For example, for the given parameter, a therapeutically effective amount will show an increase or decrease of at least 5%, 10%, 15%, 20%, 25%, 40%, 50%, 60%, 75%, 80%, 90%, or at least 100%.

Therapeutic efficacy can also be expressed as "-fold" increase or decrease. For example, a therapeutically effective amount can have at least a 1.2-fold, 1.5-fold, 2-fold, 5-fold, or more effect over a control.

[0061] The term "diagnosis" refers to a relative probability that a pulmonary disease is present in the subject. Similarly, the term "prognosis" refers to a relative probability that a certain future outcome may occur in the subject. For example, in the context of the present invention, prognosis can refer to the likelihood that an individual will develop a pulmonary disease, or the likely severity of the disease (*e.g.*, severity of symptoms, rate of functional decline, survival, etc.). The terms are not intended to be absolute, as will be appreciated by any one of skill in the field of medical diagnostics.

[0062] The terms "correlating" and "associated," in reference to determination of a pulmonary disease risk factor, refers to comparing the presence or amount of the risk factor (*e.g.*, decreased or increased expression of an IPF biomarker protein) in an individual to its presence or amount in persons known to suffer from, or known to be at risk of, the pulmonary disease, or in persons known to be free of pulmonary disease, and assigning an increased or decreased probability of having/ developing the pulmonary disease to an individual based on the assay result(s).

[0063] "Nucleic acid" or "oligonucleotide" or "polynucleotide" or grammatical equivalents used herein means at least two nucleotides covalently linked together. Oligonucleotides are typically from about 5, 6, 7, 8, 9, 10, 12, 15, 25, 30, 40, 50 or more nucleotides in length, up to about 100 nucleotides in length. Nucleic acids and polynucleotides are a polymers of any length, including longer lengths, *e.g.*, 200, 300, 500, 1000, 2000, 3000, 5000, 7000, 10,000, etc. A nucleic acid of the present invention will generally contain phosphodiester bonds, although in some cases, nucleic acid analogs are included that may have alternate backbones, comprising, *e.g.*, phosphoramidate, phosphorothioate, phosphorodithioate, or O-methylphosphoroamidite linkages (see Eckstein, *Oligonucleotides and Analogues: A Practical Approach*, Oxford University Press); and peptide nucleic acid backbones and linkages. Other analog nucleic acids include those with positive backbones; non-ionic backbones, and non-ribose backbones, including those described in U.S. Patent Nos. 5,235,033 and 5,034,506, and Chapters 6 and 7, ASC Symposium Series 580, *Carbohydrate Modifications in Antisense Research*, Sanghui & Cook, eds.. Nucleic acids containing one or more carbocyclic sugars are also included within one definition of nucleic acids. Modifications of the ribose-phosphate backbone may be done for a variety of reasons, *e.g.*, to increase the stability and half-life of such molecules in physiological environments or as probes on a biochip. Mixtures of naturally occurring nucleic acids and analogs can be made; alternatively, mixtures of different nucleic acid analogs, and mixtures of naturally occurring nucleic acids and analogs may be made.

[0064] The terms "identical" or percent "identity," in the context of two or more nucleic acids (*e.g.*, genomic sequences or subsequences or coding sequences) or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (*i.e.*, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more identity over a specified region), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Such sequences are then said to be "substantially identical." This definition also refers to the compliment of a test sequence. Optionally, the identity exists over a region that is at least about 10 to about 100, about 20 to about 75, about 30 to about 50 amino acids or nucleotides in length.

[0065] An example of algorithms suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al*, *Nuc. Acids Res.* 25:3389-3402 (1977) and Altschul *et al*, *J. Mol. Biol.* 215:403-410

(1990), respectively. As will be appreciated by one of skill in the art, the software for performing BLAST analyses is publicly available through the website of the National Center for Biotechnology Information (ncbi.nlm.nih.gov).

[0066] The terms "polypeptide," "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers, those containing modified residues, and non-naturally occurring amino acid polymer.

[0067] The term "amino acid" refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function similarly to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, *e.g.*, hydroxyproline, γ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, *e.g.*, an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, *e.g.*, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs may have modified R groups (*e.g.*, norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions similarly to a naturally occurring amino acid.

[0068] Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes.

[0069] "Conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical or associated, *e.g.*, naturally contiguous, sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode most proteins. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to another of

the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes silent variations of the nucleic acid. One of skill will recognize that in certain contexts each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is

5 ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, often silent variations of a nucleic acid which encodes a polypeptide is implicit in a described sequence with respect to the expression product, but not with respect to actual probe sequences.

10 **[0070]** As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing

15 functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants, interspecies homologs, and alleles of the invention. Typically conservative substitutions for one another: 1) Alanine (A), Glycine (G); 2) Aspartic acid (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); 6)

20 Phenylalanine (F), Tyrosine (Y), Tryptophan (W); 7) Serine (S), Threonine (T); and 8) Cysteine (C), Methionine (M) (*see, e.g.,* Creighton, *Proteins* (1984)).

[0071] A "label" or a "detectable moiety" is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, chemical, or other physical means. For example, useful labels include ³²P, fluorescent dyes, electron-dense reagents, enzymes (*e.g.,* as commonly

25 used in an ELISA), biotin, digoxigenin, or haptens and proteins or other entities which can be made detectable, *e.g.,* by incorporating a radiolabel into a peptide or antibody specifically reactive with a target peptide. Any method known in the art for conjugating an antibody to the label may be employed, *e.g.,* using methods described in Hermanson, Bioconjugate Techniques 1996, Academic Press, Inc., San Diego.

30 **[0072]** A "labeled protein or polypeptide" is one that is bound, either covalently, through a linker or a chemical bond, or noncovalently, through ionic, van der Waals, electrostatic, or hydrogen bonds to a label such that the presence of the labeled protein or polypeptide may be

detected by detecting the presence of the label bound to the labeled protein or polypeptide. Alternatively, methods using high affinity interactions may achieve the same results where one of a pair of binding partners binds to the other, *e.g.*, biotin, streptavidin.

[0073] "Antibody" refers to a polypeptide comprising a framework region from an immunoglobulin gene or fragments thereof that specifically binds and recognizes an antigen, *e.g.*, a specific bacterial antigen. Typically, the "variable region" contains the antigen-binding region of the antibody (or its functional equivalent) and is most critical in specificity and affinity of binding. See Paul, *Fundamental Immunology* (2003).

[0074] An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kD) and one "heavy" chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (VL) and variable heavy chain (VH) refer to these light and heavy chains respectively.

[0075] Antibodies can exist as intact immunoglobulins or as any of a number of well-characterized fragments that include specific antigen-binding activity. Such fragments can be produced by digestion with various peptidases. Pepsin digests an antibody below the disulfide linkages in the hinge region to produce F(ab)'₂, a dimer of Fab which itself is a light chain joined to VH-CH₁ by a disulfide bond. The F(ab)'₂ may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the F(ab)'₂ dimer into an Fab' monomer. The Fab' monomer is essentially Fab with part of the hinge region (*see Fundamental Immunology* (Paul ed., 3d ed. 1993). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized *de novo* either chemically or by using recombinant DNA methodology. Thus, the term antibody, as used herein, also includes antibody fragments either produced by the modification of whole antibodies, or those synthesized *de novo* using recombinant DNA methodologies (*e.g.*, single chain Fv) or those identified using phage display libraries (*see, e.g.*, McCafferty *et al*, *Nature* 348:552-554 (1990)).

[0076] The term "aptamer" as provided herein refers to short oligonucleotides (*e.g.* deoxyribonucleotides), which fold into diverse and intricate molecular structures that bind with high affinity and specificity to proteins, peptides, and small molecules in a non-Watson Crick manner. An aptamer can thus be used to detect or otherwise target nearly any molecule of

interest, including a fibrotic pulmonary disease marker protein. Methods of constructing and determining the binding characteristics of aptamers are well known in the art. For example, such techniques are described in U.S. Patent Nos. 5,582,981, 5,595,877 and 5,637,459. Aptamers are typically at least 5 nucleotides, 10, 20, 30 or 40 nucleotides in length, and can be composed of modified nucleic acids to improve stability. Flanking sequences can be added for structural stability, *e.g.*, to form 3-dimensional structures in the aptamer. Aptamers can be selected in vitro from very large libraries of randomized sequences by the process of systemic evolution of ligands by exponential enrichment (SELEX as described in Ellington AD, Szostak JW (1990) In vitro selection of RNA molecules that bind specific ligands. *Nature* 346:818-822; Tuerk C, Gold L (1990) Systematic evolution of ligands by exponential enrichment: RNA ligands to bacteriophage T4 DNA polymerase. *Science* 249:505-510) or by developing SOMAmers (slow off-rate modified aptamers) (Gold L et al. (2010) Aptamer-based multiplexed proteomic technology for biomarker discovery. *PLoS ONE* 5(12):e15004). Applying the SELEX and the SOMAmer technology includes for instance adding functional groups that mimic amino acid side chains to expand the aptamer's chemical diversity. As a result high affinity aptamers for almost any protein target are enriched and identified.

[0077] A "biomarker", "marker protein" or "biomarker protein" as provided herein refers to any assayable characteristics or compositions that is used to identify, predict, or monitor a condition (*e.g.*, a fibrotic pulmonary disease or lack thereof) or a therapy for said condition in a subject or sample. A biomarker is, for example, a protein or combination of proteins whose presence, absence, or relative amount is used to identify a condition (*e.g.* IPF) or status of a condition (*e.g.* IPF) in a subject or sample. Biomarkers identified herein are measured to determine levels, expression, activity, or to detect fragments, variants or homologs of said biomarkers. Variants include amino acid or nucleic acid variants or post translationally modified variants. In embodiments, the marker protein is a protein or fragment thereof as set forth in Table 1A, 1B, 2A, 2B, 3A, 3B, 4A, or 4B. In embodiments, the marker protein is a homolog of the protein listed in Table 1A, 1B, 2A, 2B, 3A, 3B, 4A, or 4B. The marker proteins provided herein are identified by accession numbers referring to the corresponding amino acid and/or nucleic acid sequence of the individual marker proteins. Therefore, a person of ordinary skill in the art will immediately recognize the sequences of the marker proteins provided herein.

[0078] In some examples of the disclosed methods, when the expression level of a biomarker(s) is assessed, the level is compared with control expression level of the biomarker(s). By control level is meant the expression level of a particular biomarker(s) from a sample or

subject lacking a disease (e.g. IPF), at a selected stage of a disease or disease state, or in the absence of a particular variable such as a therapeutic agent. Alternatively, the control level comprises a known amount of biomarker. Such a known amount correlates with an average level of subjects lacking a disease, at a selected stage of a disease or disease state, or in the absence of a particular variable such as a therapeutic agent. A control level also includes the expression level of one or more biomarkers from one or more selected samples or subjects as described herein. For example, a control level includes an assessment of the expression level of one or more biomarkers in a sample from a subject that does not have a disease (e.g. IPF), is at a selected stage of progression of a disease (e.g. IPF), or has not received treatment for a disease. Another exemplary control level includes an assessment of the expression level of one or more biomarkers in samples taken from multiple subjects that do not have a disease, are at a selected stage of progression of a disease, or have not received treatment for a disease.

[0079] When the control level includes the expression level of one or more biomarkers in a sample or subject in the absence of a therapeutic agent, the control sample or subject is optionally the same sample or subject to be tested before or after treatment with a therapeutic agent or is a selected sample or subject in the absence of the therapeutic agent. Alternatively, a control level is an average expression level calculated from a number of subjects without a particular disease. A control level also includes a known control level or value known in the art.

[0080] In one particular example, a biomarker is a protein or combination of proteins whose expression level in a subject or sample is indicative of a fibrotic pulmonary disease or a fibrotic pulmonary disease activity. The expression level of a biomarker or a combination of a plurality of biomarkers may be increased or decreased compared to a control level. For example, the expression level of a biomarker or a combination of a plurality of biomarkers as provided herein may be increased or decreased in a subject compared to the expression level of the same subject at an earlier time point. Therefore, the expression level of a biomarker as provided herein may be indicative of a specific disease stage. Alternatively, the biomarker may be indicative of the efficacy of treatment. In other words, the expression level of a biomarker may be indicative of whether a patient is responsive to a treatment. The biomarker may further be indicative of the activity of a disease, wherein the activity of a disease refers to the change of one or more biomarker expression levels over the course of the disease.

[0081] The terms "agonist," "activator," "upregulator," etc. refer to a substance capable of detectably increasing the expression or activity of a given gene or protein. The agonist can

increase expression or activity 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in comparison to a control in the absence of the agonist. In certain instances, expression or activity is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold or higher than the expression or activity in the absence of the agonist.

5 **[0082]** The terms "inhibitor," "repressor" or "antagonist" or "downregulator" interchangeably refer to a substance capable of detectably decreasing the expression or activity of a given gene or protein. The antagonist can decrease expression or activity 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in comparison to a control in the absence of the antagonist. In certain instances, expression or activity is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold or lower than
10 the expression or activity in the absence of the antagonist.

[0083] The pulmonary diseases contemplated herein can include any pulmonary disorders, lung fibrosis diseases, interstitial lung diseases, idiopathic interstitial pneumonias (IIP), idiopathic pulmonary fibrosis, familial interstitial pneumonia (FIP), non-specific interstitial pneumonia (NSIP), hypersensitivity pneumonitis, acute respiratory distress syndrome (ARDS),
15 scleroderma associated interstitial lung disease (SSc-ILD), Sarcoidosis, Beryllium disease, rheumatoid arthritis associated lung disorder, collagen vascular associated lung disorder, cigarette smoke associated lung disorders, Sjogren's syndrome, mixed connective tissue disease, etc.

[0084] Pulmonary fibrotic conditions, *e.g.*, interstitial lung diseases (ILD) are characterized by
20 shortness of breath, chronic coughing, fatigue and weakness, loss of appetite, and rapid weight loss. Pulmonary fibrosis is commonly linked to interstitial lung diseases (*e.g.*, autoimmune disorders, viral infections or other microscopic injuries), but can be idiopathic. Fibrosis involves exchange of normal lung tissue with fibrotic tissue (scar tissue) that leads to reduced oxygen capacity.

25 **[0085]** Idiopathic interstitial pneumonias (IIP) are a subset of diffuse interstitial lung diseases of unknown etiology (the term "idiopathic" indicates unknown origin). IIPs are characterized by expansion of the interstitial compartment (*i.e.*, that portion of the lung parenchyma sandwiched between the epithelial and endothelial basement membranes) with an infiltrate of inflammatory cells. The inflammatory infiltrate is sometimes accompanied by fibrosis, either in the form of
30 abnormal collagen deposition or proliferation of fibroblasts capable of collagen synthesis.

[0086] Idiopathic Pulmonary Fibrosis (IPF) occurs in thousands of people worldwide with a doubling of prevalence over the past 10 years. Onset of IPF occurs around 50 to 70 years of age

and starts with progressive shortness of breath and hypoxemia. IPF median survival is around 3-5 years and is to date untreatable. The etiology and pathogenesis of the condition is not well understood. About 5-20 percent of all cases of IPF have a family history and inheritance appears to be autosomal dominant. The clinical course of IPF is highly variable. Individuals diagnosed with IPF can experience a slow and steady decline, whereas others may decline more rapidly, thereby exhibiting a progressive form of idiopathic pulmonary fibrosis. Patients may also experience periods of relative stability interrupted by periods of rapid decline (known as acute exacerbations). A progressive form of idiopathic pulmonary fibrosis is characterized by the rapid decline of clinical and physical markers (e.g., FEV1, VC, FVC, DL_CO as defined below).

[0087] Additional fibrotic pulmonary diseases include Acute Interstitial Pneumonia (AIP), Respiratory Bronchiolitis-associated Interstitial Lung Disease (RBILD), Desquamative Interstitial Pneumonia (DIP), Non-Specific Interstitial Pneumonia (NSIP), Bronchiolitis obliterans, with Organizing Pneumonia (BOOP).

[0088] AIP is a rapidly progressive and histologically distinct form of interstitial pneumonia.

The pathological pattern is an organizing form of diffuse alveolar damage (DAD) that is also found in acute respiratory distress syndrome (ARDS) and other acute interstitial pneumonias of known causes (see Clinical Atlas of Interstitial Lung Disease (2006 ed.) pp61-63).

[0089] RBILD is characterized by inflammatory lesions of the respiratory bronchioles in cigarette smokers. The histologic appearance of RBILD is characterized by the accumulation of pigmented macrophages within the respiratory bronchioles and the surrounding airspaces, variably, peribronchial fibrotic alveolar septal thickening, and minimal associated mural inflammation (see Wells *et al.* (2003) *Sem Respir. Crit. Care Med.* vol. 24).

[0090] DIP is a rare interstitial lung disease characterized by the accumulation of macrophages in large numbers in the alveolar spaces associated with interstitial inflammation and/or fibrosis.

The macrophages frequently contain light brown pigment. Lymphoid nodules are common, as is a sparse but distinct eosinophil infiltrate. DIP is most common in smokers (see Tazelaar *et al.* (Sep. 21, 2010) *Histopathology*).

[0091] NSIP is characterized pathologically by uniform interstitial inflammation and fibrosis appearing over a short period of time. NSIP differs from other interstitial lung diseases in that it has a generally good prognosis. In addition, the temporal uniformity of the parenchymal changes seen in NSIP contrasts greatly with the temporal heterogeneity of usual interstitial pneumonia (see Coche *et al.* (2001) *Brit J Radiol* 74: 189).

[0092] BOOP, unlike NSIP, can be fatal within days of first acute symptoms. It is characterized by rapid onset of acute respiratory distress syndrome; therefore, clinically, rapidly progressive BOOP can be indistinguishable from acute interstitial pneumonia. Histological features include clusters of mononuclear inflammatory cells that form granulation tissue and plug the distal airways and alveolar spaces. These plugs of granulation tissue may form polyps that migrate within the alveolar ducts or may be focally attached to the wall. (see White & Ruth-Saad (2007) *Crit. Care Nurse* 27:53).

[0093] Further details about the characteristics and therapies available for these diseases can be found, *e.g.*, on the website of the American Lung Association at lungusa.org/lung-disease/pulmonary-fibrosis.

[0094] Diagnostic indicators of pulmonary disorders include biopsy (*e.g.*, VATS or surgical lung biopsy), high resolution computed tomography (HRCT) or breathing metrics, such as forced expiratory volume (FEV1), vital capacity (VC), forced vital capacity (FVC), and FEV1/FVC.

II. METHODS

[0095] Provided herein, *inter alia*, are biomarkers (*e.g.* protein biomarkers) for diagnosing fibrotic pulmonary diseases as well as evaluation of progression, activity and treatment of fibrotic pulmonary diseases. Further provided herein are therapeutic targets for ameliorating fibrotic pulmonary diseases.

[0096] In one aspect, a method of determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB in a subject that has or is at risk for developing a fibrotic pulmonary disease is provided. The method includes (i) obtaining a biological sample from the subject and (ii) determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB in the biological sample. In some embodiments, the method includes selecting a subject that has or is at risk for developing a fibrotic pulmonary disease. The selected subject may be treated for fibrotic pulmonary disease. In some embodiments, the subject is not treated for fibrotic pulmonary disease. The subject may be part of a plurality of subjects participating in a clinical trial. Wherein the subject is part of a clinical trial, the selecting is at least part based on the determining of an expression level as provided herein.

[0097] A fibrotic pulmonary disease marker protein is a biomarker protein useful to identify, predict, or monitor a fibrotic pulmonary disease or lack thereof or a therapy for fibrotic pulmonary disease in a subject or sample. A person of ordinary skill in the art will immediately

recognize that determining an expression level of a fibrotic pulmonary disease marker protein described herein includes determining the level of one or more fibrotic pulmonary disease marker proteins in a sample (e.g. patient biological sample such as a blood-derived biological sample). Thus in some embodiments, the expression level of a plurality of fibrotic pulmonary disease marker proteins is determined. Wherein the expression level of a plurality of fibrotic pulmonary disease marker proteins is determined, the level of at least two (e.g. 3, 4, 5, 6, 7, 8, 9, 10 etc.) fibrotic pulmonary disease marker proteins is determined and the at least two fibrotic pulmonary disease marker proteins are independently different.

[0098] In some embodiments, the fibrotic pulmonary disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia. In embodiments, the fibrotic pulmonary disease is a progressive form of idiopathic pulmonary fibrosis. In other embodiments, the fibrotic pulmonary disease marker protein is a progressive fibrotic pulmonary disease marker protein. A progressive fibrotic pulmonary disease marker protein is a biomarker protein indicative of a fibrotic pulmonary disease patient having or being at risk of developing progressive fibrotic pulmonary disease (e.g., a progressive form of a fibrotic pulmonary disease). Thus, in some embodiments, the subject has or is at risk for developing a progressive fibrotic pulmonary disease. In other embodiments, the progressive fibrotic pulmonary disease is idiopathic pulmonary fibrosis. A progressive fibrotic pulmonary disease is a disease wherein certain clinical or physiological parameters decline over the course of the disease. Commonly used parameters to determine fibrotic pulmonary disease progression include for example, breathing metrics, such as forced expiratory volume (FEV1), vital capacity (VC), forced vital capacity (FVC), FEV1/FVC and diffusing capacity of carbon monoxide (**DLco**)- For example, a fibrotic pulmonary disease patient showing a 5% decline in FVC as compared to a control level may be considered a progressive fibrotic pulmonary disease patient. A control level may be the FVC of the same patient measured at an earlier stage of the fibrotic pulmonary disease or the FVC calculated from a number of subjects lacking the fibrotic pulmonary disease. Thus, in some embodiments, the FVC of a progressive fibrotic pulmonary disease patient is at least 5% less than a control level.

[0099] In other embodiments, the FVC of a progressive fibrotic pulmonary disease patient is about 5% less than a control level. In some embodiments, the FVC of a progressive fibrotic pulmonary disease patient is at least 10% less than a control level. In other embodiments, the FVC of a progressive fibrotic pulmonary disease patient is about 10% less than a control level.

[0100] Another parameter of progressive fibrotic pulmonary disease is a decline in the 6-minute walk test distance (6MWD). For example, a fibrotic pulmonary disease patient showing a decline of more than 50 meters in the 6MWD compared to a control distance may be considered a progressive fibrotic pulmonary disease patient. A control distance refers to the 6MWD of the same patient measured at an earlier stage of the fibrotic pulmonary disease or the 6MWD calculated from a number of subjects lacking the fibrotic pulmonary disease. In some embodiments, the decline in the 6MWD is at least 30 meters. In other embodiments, the decline in the 6MWD is about 50 meters.

[0101] Yet, another parameter of progressive fibrotic pulmonary disease is a decline in diffusing capacity of carbon monoxide (DL_{CO}). For example, a fibrotic pulmonary disease patient showing a decline of more than 15% in DL_{CO} compared to a control level may be considered a progressive fibrotic pulmonary disease patient. A control level refers to the DL_{CO} of the same patient measured at an earlier stage of the fibrotic pulmonary disease or the DL_{CO} calculated from a number of subjects lacking the fibrotic pulmonary disease. Thus, in some embodiments, the DL_{CO} of a progressive fibrotic pulmonary disease patient is at least 10% less compared to a control level. In other embodiments, the DL_{CO} of a progressive fibrotic pulmonary disease patient is about 15% less than a control level.

[0102] In other embodiments, the biological sample is a blood-derived biological sample of the subject. In some further embodiments, the blood-derived biological sample is whole blood, serum or plasma. In some embodiments, the biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0103] The methods provided herein including embodiments thereof further include treating a subject for fibrotic pulmonary diseases. As described above the expression level of one or more fibrotic pulmonary disease marker proteins is determined. In some embodiments, an expression level of a fibrotic pulmonary disease marker protein set forth in Table 1A is determined. In other embodiments, an expression level of a fibrotic pulmonary disease marker protein set forth in Table IB is determined. In some embodiments, an expression level of a fibrotic pulmonary disease marker protein set forth in Table 2A is determined. In other embodiments, an expression level of a fibrotic pulmonary disease marker protein set forth in Table 2B is determined. In some embodiments, an expression level of a fibrotic pulmonary disease marker protein set forth in Table 4A is determined. In other embodiments, an expression level of a fibrotic pulmonary disease marker protein set forth in Table 4B is determined.

[0104] The expression levels of the one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table 1B may be increased or decreased in a fibrotic pulmonary disease patient. The expression levels of the one or more fibrotic pulmonary disease marker proteins set forth in Table 2A or Table 2B may be increased or decreased in a fibrotic pulmonary disease patient. The expression levels of the one or more fibrotic pulmonary disease marker proteins set forth in Table 4A or Table 4B may be increased or decreased in a fibrotic pulmonary disease patient.

[0105] In some embodiments, the one or more fibrotic pulmonary disease marker proteins set forth in Table 1A, 2A or 4A is increased. In other embodiments, the one or more fibrotic pulmonary disease marker proteins set forth in Table 1A, 2A or 4A is decreased. In some embodiments, the one or more fibrotic pulmonary disease marker proteins set forth in Table 1B, 2B or 4B is increased. In other embodiments, the one or more fibrotic pulmonary disease marker proteins set forth in Table 1B, 2B or 4B is decreased.

[0106] In some embodiments, the expression level of the fibrotic pulmonary disease marker protein set forth in Table 1A is elevated relative to a standard control. In other embodiments, the expression level of the fibrotic pulmonary disease marker protein set forth in Table 1B is decreased relative to a standard control. In some embodiments, the expression level of the fibrotic pulmonary disease marker protein set forth in Table 2A is elevated relative to a standard control. In other embodiments, the expression level of the fibrotic pulmonary disease marker protein set forth in Table 2B is decreased relative to a standard control. In some embodiments, the expression level of the fibrotic pulmonary disease marker protein set forth in Table 4A is elevated relative to a standard control. In other embodiments, the expression level of the fibrotic pulmonary disease marker protein set forth in Table 4B is decreased relative to a standard control.

[0107] A treatment regimen for a fibrotic pulmonary disease patient with modulated expression levels of one or more fibrotic pulmonary disease marker proteins as disclosed herein (e.g. in Table 1A, 1B, 2A, 2B, 4A or 4B) may be administering to the patient an effective amount of a modulator affecting the one or more increased or decreased biomarker protein expression levels. Therefore, in some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 1A or 1B. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in

Table 2A or 2B. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 4A or 4B.

[0108] In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 1A. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 2A. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 4A. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 1B. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 2B. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 4B. In some embodiments, the modulator is an antagonist. In other embodiments, the antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer. In other embodiments, the modulator is an agonist. In other embodiments, the agonist is a peptide, small molecule, nucleic acid, antibody or aptamer. In embodiments, the agonist is a fibrotic pulmonary disease marker protein set forth in Table 1A or 1B. In embodiments, the agonist is a fibrotic pulmonary disease marker protein set forth in Table 2A or 2B. In embodiments, the agonist is a fibrotic pulmonary disease marker protein set forth in Table 4A or 4B.

[0109] The methods provided herein may include combinatorial treatment of a fibrotic pulmonary disease patient with a modulator of one or more fibrotic pulmonary disease marker proteins and a fibrotic pulmonary disease treatment. Thus, the method includes administering to the subject an effective amount of a further therapeutic agent. Exemplary therapeutic agents include, but are not limited to, the agents selected from the group consisting of steroids (including but not limited to prednisolone), cytotoxic agents (including but not limited to azathioprine and cyclophosphamide), bardoxolone, LPA agonists (including but not limited to AM152); Torisel (temsirolimus); PI3K inhibitors; pentraxin or serum amyloid P (including but not limited to Pentraxin-2 (PTX-2 or PRM-151)); MEK inhibitors (including but not limited to ARRY-162 and ARRY-300); p38 inhibitors; PAI-1 inhibitors (including but not limited to Tiplaxtinin); agents that reduce the activity of transforming growth factor-beta (TGF- β)

(including but not limited to GC-1008 (Genzyme/MedImmune); lerdelimumab (CAT-152; Trabio, Cambridge Antibody); metelimumab (CAT-192, Cambridge Antibody); LY-2157299 (Eli Lilly); ACU-HTR-028 (Opko Health)) including antibodies that target one or more TGF- β isoforms, inhibitors of TGF- β receptor kinases TGFBR1 (ALK5) and TGFBR2, and modulators of post-receptor signaling pathways; chemokine receptor signaling; endothelin receptor antagonists including inhibitors that target both endothelin receptor A and B and those that selectively target endothelin receptor A (including but not limited to ambrisentan; avosentan; bosentan; clazosentan; darusentan; BQ-153; FR-139317, L-744453; macitentan; PD-145065; PD-156252; PD163610; PS-433540; S-0139; sitaxentan sodium; TBC-3711; zibotentan); agents that reduce the activity of connective tissue growth factor (CTGF) (including but not limited to FG-3019, FibroGen), and including other CTGF-neutralizing antibodies; matrix metalloproteinase (MMP) inhibitors (including but not limited to MMPI-12, PUP-1 and tigapotide trifluate); agents that reduce the activity of epidermal growth factor receptor (EGFR) including but not limited to erlotinib, gefitinib, BMS-690514, cetuximab, antibodies targeting EGF receptor, inhibitors of EGF receptor kinase, and modulators of post-receptor signaling pathways; agents that reduce the activity of platelet derived growth factor (PDGF) (including but not limited to Imatinib mesylate (Novartis)) and also including PDGF neutralizing antibodies, antibodies targeting PDGF receptor (PDGFR), inhibitors of PDGFR kinase activity, and post-receptor signaling pathways; agents that reduce the activity of vascular endothelial growth factor (VEGF) (including but not limited to axitinib, bevacizumab, BIBF-1 120, CDP-791, CT-322, IMC-18F1, PTC-299, and ramucirumab) and also including VEGF-neutralizing antibodies, antibodies targeting the VEGF receptor 1 (VEGFR1, Flt-1) and VEGF receptor 2 (VEGFR2, KDR), the soluble form of VEGFR1 (sFlt) and derivatives thereof which neutralize VEGF, and inhibitors of VEGF receptor kinase activity; inhibitors of multiple receptor kinases such as BIBF-1 120 which inhibits receptor kinases for vascular endothelial growth factor, fibroblast growth factor, and platelet derived growth factor; agents that interfere with integrin function (including but not limited to STX-100 and IMG-388) and also including integrin targeted antibodies; agents that interfere with the pro-fibrotic activities of IL-4 (including but not limited to AER-001, AMG-317, APG-201, and sIL-4Ra) and IL-13 (including but not limited to AER-001, AMG-3 17, anrukinzumab, CAT-354, cintredekin besudotox, MK-6105, QAX-576, SB-3 13, SL-102, and TNX-650) and also including neutralizing anti-bodies to either cytokine, antibodies that target IL-4 receptor or IL-13 receptor, the soluble form of IL-4 receptor or derivatives thereof that is reported to bind and neutralize both IL-4 and IL-13, chimeric proteins including all or part of IL-13 and a toxin particularly pseudomonas endotoxin, signaling through the JAK-

STAT kinase pathway; agents that interfere with epithelial mesenchymal transition including inhibitors of mTor (including but not limited to AP-23573 or rapamycin); agents that reduce levels of copper such as tetrathiomolybdate; agents that reduce oxidative stress including N-acetyl cysteine and tetrathiomolybdate; and interferon gamma, inhibitors of phosphodiesterase 4 (PDE4) (including but not limited to Roflumilast); inhibitors of phosphodiesterase 5 (PDE5) (including but not limited to mirodenafil, PF-4480682, sildenafil citrate, SLx-2101, tadalafil, udenafil, UK-369003, vardenafil, and zaprinast); or modifiers of the arachidonic acid pathway including cyclooxygenase and 5-lipoxygenase inhibitors (including but not limited to Zileuton), compounds that reduce tissue remodeling or fibrosis including prolyl hydrolase inhibitors (including but not limited to 1016548, CG-0089, FG-2216, FG-4497, FG-5615, FG-6513, fibrostatin A (Takeda), lufironil, P-1894B, and safironil) and peroxisome proliferator-activated receptor (PPAR)-gamma agonists (including but not limited to pioglitazone and rosiglitazone), and combinations thereof. In some embodiments, the therapeutic agent is an anti-fibrotic drug. In some embodiments, the therapeutic agent is an idiopathic pulmonary fibrosis drug. In other embodiment, the idiopathic pulmonary fibrosis drug is a mucolytic drug.

[0110] In another aspect, a method of determining whether a subject has or is at risk of developing a fibrotic pulmonary disease is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB in a subject; (ii) determining whether the expression level is increased or decreased relative to a standard control, wherein an elevated expression level of a fibrotic pulmonary disease marker protein in Table 1A or a decreased expression level of a fibrotic pulmonary disease marker protein in Table IB relative to the standard control indicates that the subject has or is at risk of developing a fibrotic pulmonary disease; and (iii) based at least in part on the expression level in step (ii), determining whether the subject has or is at risk for developing a fibrotic pulmonary disease.

[0111] In some embodiments, the method includes selecting a subject that has or is at risk for developing a fibrotic pulmonary disease. In other embodiments, the fibrotic pulmonary disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia. In some embodiments, the one or more fibrotic pulmonary disease marker proteins is a progressive fibrotic pulmonary disease marker protein. In other embodiments, the subject has or is at risk for developing a progressive fibrotic pulmonary disease. In some embodiments, the progressive fibrotic pulmonary disease is idiopathic pulmonary fibrosis.

[0112] The expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB is detected from a biological sample of the subject. In some embodiments, the biological sample is a blood-derived biological sample of the subject. In other embodiments, the blood-derived biological sample is whole blood, serum or plasma. In some
5 embodiments, the biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0113] As described above the methods provided herein include treating for fibrotic pulmonary disease. Thus, in some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in
10 Table 1A or IB. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 1A. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table IB. In some embodiments, the method includes administering to the subject an effective amount of a
15 modulator of the fibrotic pulmonary disease marker protein set forth in Table 2A. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 2B. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 4A. In some
20 embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 4B. In some embodiments, the modulator is an antagonist. In other embodiments, the antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer. In one embodiment, the modulator is an agonist. In one embodiment, the agonist is a peptide, small molecule, nucleic acid, antibody or
25 aptamer.

[0114] As described above the methods provided herein may further include administering to the subject an effective amount of a further therapeutic agent (e.g., as described above). In some further embodiments, the therapeutic agent is an idiopathic pulmonary fibrosis drug. In some other further embodiments, the idiopathic pulmonary fibrosis drug is a mucolytic drug.

30 [0115] In another aspect, a method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker

protein set forth in Table 1A or Table IB in a fibrotic pulmonary disease patient (ii) determining whether the expression level is modulated relative to a standard control, wherein a modulated expression level of a fibrotic pulmonary disease marker protein in Table 1A or Table IB relative to the standard control indicates that the fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, and (iii) based at least in part on the expression level in step (ii), determining whether the fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease. In some embodiments, the determining whether the expression level is modulated relative to a standard control comprises determining whether the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients. In embodiments, the determining whether the expression level is modulated relative to a standard control includes determining whether the expression level is elevated or suppressed relative to an expression level of the same fibrotic pulmonary disease marker protein in other fibrotic pulmonary disease patients. In embodiments, the determining whether the expression level is modulated relative to a standard control includes determining whether the expression level is elevated or suppressed relative to an expression level of a different fibrotic pulmonary disease marker protein in the patient. In some embodiments, the fibrotic pulmonary disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia. In other embodiments, the fibrotic pulmonary disease marker protein is a progressive fibrotic pulmonary disease marker protein.

[0116] In embodiments, an increased expression level of a fibrotic pulmonary disease marker protein in Table 1A relative to a standard control indicates an increased risk for progression of the fibrotic pulmonary disease. In embodiments, a decreased expression level of a fibrotic pulmonary disease marker protein in Table 1A relative to a standard control indicates an decreased risk for progression of the fibrotic pulmonary disease. In embodiments, an increased expression level of a fibrotic pulmonary disease marker protein in Table IB relative to a standard control indicates an increased risk for progression of the fibrotic pulmonary disease. In embodiments, a decreased expression level of a fibrotic pulmonary disease marker protein in Table IB relative to a standard control indicates an decreased risk for progression of the fibrotic pulmonary disease.

[0117] In some embodiments, the subject has or is at risk for developing a progressive fibrotic pulmonary disease. In other embodiments, the progressive fibrotic pulmonary disease is idiopathic pulmonary fibrosis. In other embodiments, the biological sample is a blood-derived biological sample of the subject. In some further embodiments, the blood-derived biological

sample is whole blood, serum or plasma. In some embodiments, the biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0118] In another aspect, a method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease is provided. The method includes (i) determining a first expression level of a protein set forth in Table 1A or Table IB in the patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A or Table IB in the patient at a second time point; and (iii) comparing the second expression level of a protein set forth in Table 1A or Table IB to the first expression level of a protein set forth in Table 1A or Table IB, wherein when the second expression level of a protein set forth in Table 1A is greater than the first level of a protein set forth in Table 1A; or wherein when the second expression level of a protein set forth in Table IB is smaller than the first level of a protein set forth in Table IB, the patient is at risk for progression of the fibrotic pulmonary disease. The patient at risk for progression of the fibrotic pulmonary disease may have received treatment for the fibrotic pulmonary disease prior to the determining in step (i). Where the patient at risk for progression of the fibrotic pulmonary disease has received treatment for the fibrotic pulmonary disease prior to the determining in step (i) the treatment may be altered after the determining in step (i) and before the determining in step (ii). Alternatively, the patient at risk for progression of the fibrotic pulmonary disease may not have received treatment for the fibrotic pulmonary disease prior to the determining in step (i). Where the patient at risk for progression of the fibrotic pulmonary disease has not received treatment for the fibrotic pulmonary disease prior to the determining in step (i) the patient may receive treatment after the determining in step (i). Thus, in some embodiments, the method includes administering a fibrotic pulmonary disease treatment after the determining in step (i). In some embodiments, the method further includes determining a rate of progression of the fibrotic pulmonary disease in the patient based on the comparing. In some embodiments, the method further includes predicting a rate of progression of the fibrotic pulmonary disease in the patient based on the comparing.

[0119] In other embodiments, the determining the first expression level of a protein set forth in Table 1A or Table IB and the second expression level of a protein set forth in Table 1A or Table IB includes normalizing the first expression level of a protein set forth in Table 1A or Table IB and the second expression level of a protein set forth in Table 1A or Table IB to a protein expressed from a standard gene in the patient. In some further embodiments, the standard gene is a so-called housekeeping gene, as is commonly known in the art, such as GAPDH or beta-actin. In embodiments, the standard gene is non-differentially expressed. Where the standard

gene is non-differentially expressed, the expression level of the standard gene remains unchanged over the time course of the disease. In some embodiment, the first expression level is detected from a first biological sample of the subject and the second expression level is detected from a second biological sample of the subject. In some embodiment, the first biological sample is a first bodily fluid sample and the second biological sample is a second bodily fluid sample.

[0120] In another aspect, a method of determining a fibrotic pulmonary disease activity in a patient is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB in a subject; (ii) determining whether the expression level is modulated relative to a standard control, thereby determining a fibrotic pulmonary disease activity in the patient; and (iii) based at least in part on the expression level in step (ii), determining the fibrotic pulmonary disease activity in the patient. In embodiments, the expression level of a fibrotic pulmonary disease marker protein in Table 1A is increased relative to the standard control. In embodiments, the expression level of a fibrotic pulmonary disease marker protein in Table IB is decreased relative to the standard control.

[0121] In another aspect, a method of determining a fibrotic pulmonary disease activity in a patient is provided. The method includes (i) determining a first expression level of a protein set forth in Table 1A or Table IB in the patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A or Table IB in the patient at a second time point; and (iii) comparing the second expression level of a protein set forth in Table 1A or Table IB to the first expression level of a protein set forth in Table 1A or Table IB, thereby determining the fibrotic pulmonary disease activity in the patient. In some embodiments, the method includes administering a fibrotic pulmonary disease treatment after the determining in step (i). In some embodiment, the method includes determining a change in a fibrotic pulmonary disease activity.

[0122] In another aspect, a method of treating a fibrotic pulmonary disease in a subject in need thereof is provided. The method includes administering to the subject an effective amount of a modulator of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB, thereby treating a fibrotic pulmonary disease in the subject. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 1A. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic

pulmonary disease marker protein set forth in Table 2A. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic

pulmonary disease marker protein set forth in Table 4A. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic

5 pulmonary disease marker protein set forth in Table IB. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic

pulmonary disease marker protein set forth in Table 2B. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic

10 pulmonary disease marker protein set forth in Table 4B. In other embodiments, the modulator is an antagonist. In some further embodiments, the antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer. In some embodiments, the modulator is an agonist. In some other further embodiment, the agonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0123] In one aspect, a method of treating a fibrotic pulmonary disease in a subject in need thereof is provided. The method includes (i) determining whether a subject expresses an

15 elevated level of a fibrotic pulmonary disease marker protein as set forth in Table 1A or a decreased level of a fibrotic pulmonary disease marker protein as set forth in Table IB relative to a standard control; and (ii) when an elevated expression level of the fibrotic pulmonary disease marker protein of Table 1A or a decreased expression level of the fibrotic pulmonary disease marker protein of Table IB is found relative to the standard control, administering to the subject

20 a fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker protein set forth in Table 1A or an agonist of a fibrotic pulmonary disease marker protein set forth in Table IB, thereby treating the subject. In some embodiment, the fibrotic pulmonary disease treatment is a mucolytic drug.

[0124] The aspects provided herein including embodiments refer to one or more fibrotic

25 pulmonary disease marker protein set forth in the tables provided herein. In some aspects, one or more fibrotic pulmonary disease marker proteins are set forth in Table 1A or Table IB. In

embodiments, the one or more fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB is a protein set forth in Table 2A or Table 2B. In embodiments, the one or more

fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB is a protein set forth

30 in Table 4A or Table 4B. In some embodiments, the one or more fibrotic pulmonary disease marker protein is set forth in Table 2A. In some embodiments, the one or more fibrotic pulmonary disease marker protein is set forth in Table 2B. In some embodiments, the one or

more fibrotic pulmonary disease marker protein is set forth in Table 4A. In some embodiments, the one or more fibrotic pulmonary disease marker protein is set forth in Table 4B.

[0125] In some embodiments, the subject has or is at risk for developing a progressive fibrotic pulmonary disease. In other embodiments, the progressive fibrotic pulmonary disease is idiopathic pulmonary fibrosis. In embodiments, the progressive fibrotic pulmonary disease is a progressive form of idiopathic pulmonary fibrosis. In other embodiments, the biological sample is a blood-derived biological sample of the subject. In some further embodiments, the blood-derived biological sample is whole blood, serum or plasma. In some embodiments, the biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0126] In some embodiments, an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A is determined. In other embodiments, an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3B is determined. In embodiments, the risk for developing a progressive fibrotic pulmonary disease is increased relative to a standard control. In embodiments, the risk for developing a progressive fibrotic pulmonary disease is decreased relative to a standard control.

[0127] The expression levels of the one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B may be increased or decreased in a fibrotic pulmonary disease patient. In some embodiments, the expression level of the fibrotic pulmonary disease marker protein set forth in Table 3A is elevated relative to a standard control. In other embodiments, the expression level of the fibrotic pulmonary disease marker protein set forth in Table 3B is decreased relative to a standard control.

[0128] In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 3A or 3B. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 3A. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 3B. Thus, the method includes administering to the subject an effective amount of a further therapeutic agent. In some embodiments, the therapeutic agent is an idiopathic pulmonary fibrosis drug. In other embodiment, the idiopathic pulmonary fibrosis drug is a mucolytic drug.

[0129] In another aspect, a method of determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B in a subject that has or is at risk for developing a fibrotic pulmonary disease is provided. The method includes (i) obtaining a biological sample from the subject, and (ii) determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B in the biological sample.

[0130] In some embodiments, the method includes selecting a subject that has or is at risk for developing a fibrotic pulmonary disease. In other embodiments, the fibrotic pulmonary disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia. In some embodiments, the one or more fibrotic pulmonary disease marker proteins is a progressive fibrotic pulmonary disease marker protein. In other embodiments, the subject has or is at risk for developing a progressive fibrotic pulmonary disease. In some embodiments, the progressive fibrotic pulmonary disease is idiopathic pulmonary fibrosis. In embodiments, the progressive fibrotic pulmonary disease is a progressive form of idiopathic pulmonary fibrosis.

[0131] The expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B is detected from a biological sample of the subject. In some embodiments, the biological sample is a blood-derived biological sample of the subject. In other embodiments, the blood-derived biological sample is whole blood, serum or plasma. In some embodiments, the biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0132] As described above the methods provided herein include treating for fibrotic pulmonary disease. Thus, in some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 3A or 3B. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 3A. In some embodiments, the method includes administering to the subject an effective amount of a modulator of the fibrotic pulmonary disease marker protein set forth in Table 3B.

[0133] As described above the methods provided herein may further include administering to the subject an effective amount of a further therapeutic agent. In some further embodiments, the therapeutic agent is an idiopathic pulmonary fibrosis drug. In some other further embodiments, the idiopathic pulmonary fibrosis drug is a mucolytic drug.

[0134] In another aspect, a method of determining whether a subject has or is at risk of developing a fibrotic pulmonary disease is provided. The method includes (i) detecting an

expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject, (ii) determining whether said expression level is increased or decreased relative to a standard control, wherein an elevated expression level of a fibrotic pulmonary disease marker protein in Table 3A or a decreased expression level of a fibrotic pulmonary disease marker protein in Table 3B relative to the standard control indicates that the subject has or is at risk of developing a fibrotic pulmonary disease, and (iii) based at least in part on the expression level in step (ii), determining whether the subject has or is at risk for developing a fibrotic pulmonary disease.

[0135] In other embodiments, the determining the first expression level of a protein set forth in Table 3A or Table 3B and the second expression level of a protein set forth in Table 3A or Table 3B includes normalizing the first expression level of a protein set forth in Table 3A or Table 3B and the second expression level of a protein set forth in Table 3A or Table 3B to a protein expressed from a standard gene in the patient. In some further embodiments, the standard gene is a so-called housekeeping gene, as is commonly known in the art, such as GAPDH or beta-actin. In embodiments, the standard gene is non-differentially expressed. Where the standard gene is non-differentially expressed, the expression level of the standard gene remains unchanged over the time course of the disease. In some embodiment, the first expression level is detected from a first biological sample of the subject and the second expression level is detected from a second biological sample of the subject. In some embodiment, the first biological sample is a first bodily fluid sample and the second biological sample is a second bodily fluid sample.

[0136] In another aspect, a method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker protein set forth in Table 3A or 3B in a fibrotic pulmonary disease patient, (ii) determining whether the expression level is modulated relative to a standard control, wherein a modulated expression level of a fibrotic pulmonary disease marker protein in Table 3A or 3B relative to the standard control indicates that the fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, and (iii) based at least in part on the expression level in step (ii), determining whether the fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease. In some embodiments, the determining whether the expression level is modulated relative to a standard control comprises determining whether the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients. In embodiments, the determining whether the expression level is modulated relative to a standard control includes

determining whether the expression level is elevated or suppressed relative to an expression level of the same fibrotic pulmonary disease marker protein in other fibrotic pulmonary disease patients. In embodiments, the determining whether the expression level is modulated relative to a standard control includes determining whether the expression level is elevated or suppressed relative to an expression level of a different fibrotic pulmonary disease marker protein in the patient. In some embodiments, the fibrotic pulmonary disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia. In other embodiments, the fibrotic pulmonary disease marker protein is a progressive fibrotic pulmonary disease marker protein.

[0137] In embodiments, an increased expression level of a fibrotic pulmonary disease marker protein in Table 3A relative to a standard control indicates an increased risk for progression of the fibrotic pulmonary disease. In embodiments, a decreased expression level of a fibrotic pulmonary disease marker protein in Table 3A relative to a standard control indicates an decreased risk for progression of the fibrotic pulmonary disease. In embodiments, an increased expression level of a fibrotic pulmonary disease marker protein in Table 3B relative to a standard control indicates an increased risk for progression of the fibrotic pulmonary disease. In embodiments, a decreased expression level of a fibrotic pulmonary disease marker protein in Table 3B relative to a standard control indicates an decreased risk for progression of the fibrotic pulmonary disease.

[0138] In some embodiments, the subject has or is at risk for developing a progressive fibrotic pulmonary disease. In other embodiments, the progressive fibrotic pulmonary disease is idiopathic pulmonary fibrosis. In embodiments, the progressive fibrotic pulmonary disease is a progressive form of idiopathic pulmonary fibrosis. In other embodiments, the biological sample is a blood-derived biological sample of the subject. In some further embodiments, the blood-derived biological sample is whole blood, serum or plasma. In some embodiments, the biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0139] In one aspect, a method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease is provided. The method includes (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in the patient at a first time point, (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in the patient at a second time point, (iii) comparing the second expression level of a protein set forth in Table 3A or Table 3B to the first expression level of a protein set forth in

Table 3A or Table 3B, wherein when the second expression level of a protein set forth in Table 3A is greater than the first level of a protein set forth in Table 3A, or wherein when the second expression level of a protein set forth in Table 3B is smaller than the first level of a protein set forth in Table 3B, the patient is at risk for progression of the fibrotic pulmonary disease. The patient at risk for progression of the fibrotic pulmonary disease may have received treatment for the fibrotic pulmonary disease prior to the determining in step (i). Where the patient at risk for progression of the fibrotic pulmonary disease has received treatment for the fibrotic pulmonary disease prior to the determining in step (i) the treatment may be altered after the determining in step (i) and before the determining in step (ii). Alternatively, the patient at risk for progression of the fibrotic pulmonary disease may not have received treatment for the fibrotic pulmonary disease prior to the determining in step (i). Where the patient at risk for progression of the fibrotic pulmonary disease has not received treatment for the fibrotic pulmonary disease prior to the determining in step (i) the patient may receive treatment after the determining in step (i). Thus, in some embodiments, the method includes administering a fibrotic pulmonary disease treatment after the determining in step (i). In some embodiments, the method further includes determining a rate of progression of the fibrotic pulmonary disease in the patient based on the comparing.

[0140] In other embodiments, the determining the first expression level of a protein set forth in Table 3A or Table 3B and the second expression level of a protein set forth in Table 3A or Table 3B includes normalizing the first expression level of a protein set forth in Table 3A or Table 3B and the second expression level of a protein set forth in Table 3A or Table 3B to a protein expressed from a standard gene in the patient. In some further embodiments, the standard gene is a so-called housekeeping gene, as is commonly known in the art, such as GAPDH or beta-actin. In embodiments, the standard gene is non-differentially expressed. Where the standard gene is non-differentially expressed, the expression level of the standard gene remains unchanged over the time course of the disease. In some embodiment, the first expression level is detected from a first biological sample of the subject and the second expression level is detected from a second biological sample of the subject. In some embodiment, the first biological sample is a first bodily fluid sample and the second biological sample is a second bodily fluid sample.

[0141] In another aspect, a method of determining a fibrotic pulmonary disease activity in a patient is provided. The method includes (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject; (ii) determining whether the expression level is modulated relative to a standard control, thereby

determining a fibrotic pulmonary disease activity in the patient; and (iii) based at least in part on the expression level in step (ii), determining the fibrotic pulmonary disease activity in the patient. In embodiments, the expression level of a fibrotic pulmonary disease marker protein in Table 3A is increased relative to the standard control. In embodiments, the expression level of a
5 fibrotic pulmonary disease marker protein in Table 3B is decreased relative to the standard control.

[0142] In another aspect, a method of determining a fibrotic pulmonary disease activity in a patient is provided. The method includes (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in the patient at a first time point, (ii) determining a second
10 expression level of a protein set forth in Table 3A or Table 3B in the patient at a second time point, (iii) comparing the second expression level of a protein set forth in Table 3A or Table 3B to the first expression level of a protein set forth in Table 3A or Table 3B, thereby determining the fibrotic pulmonary disease activity in the patient. In embodiments, the method includes determining a change in a fibrotic pulmonary disease activity.

[0143] In another aspect, a method of treating a fibrotic pulmonary disease in a subject in need thereof is provided. The method includes administering to the subject an effective amount of an modulator of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B, thereby treating a fibrotic pulmonary disease in the subject.

[0144] In another aspect, a method of treating a fibrotic pulmonary disease in a subject in need
20 thereof is provided. The method includes (i) determining whether a subject expresses an elevated level of a fibrotic pulmonary disease marker protein as set forth in Table 3A or a decreased level of a fibrotic pulmonary disease marker protein as set forth in Table 3B relative to a standard control, and (ii) when an elevated expression level of the fibrotic pulmonary disease marker protein of Table 3A or a decreased expression level of the fibrotic pulmonary disease
25 marker protein of Table 3B is found relative to the standard control, administering to the subject a fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker protein set forth in Table 3A or an agonist of a fibrotic pulmonary disease marker protein set forth in Table 3B, thereby treating the subject. In some embodiments, the determining whether the expression level is modulated relative to a standard control includes determining whether the
30 expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients. Therefore a standard control as referred to herein may include or may be an average value

gathered from a population of pulmonary disease patients. In other embodiments, a standard control is an average value gathered from a population of normal patients.

III. KITS AND COMPOSITIONS

[0148] The invention provides kits for detection of fibrotic pulmonary disease marker proteins or fragments thereof in a subject. The kit can be for personal use or provided to medical professionals. The kit can be a kit for diagnosing or prognosing a fibrotic pulmonary disease, or for monitoring the progression of disease or the efficacy of treatment.

[0145] In one aspect a kit is provided. The kit includes a marker protein binding agent (e.g., an aptamer, optionally labeled) capable of binding to a substance within a biological sample (e.g., whole blood, serum or plasma) from a human subject having or at risk of developing a fibrotic pulmonary disease, wherein the substance is a fibrotic pulmonary disease marker protein or fragment thereof set forth in the tables provided herein (e.g., Table 1A, 1B, 2A, 2B, 3A, 3B, 4A, or 4B); and a detecting reagent or a detecting apparatus capable of indicating binding of the marker protein binding agent to the substance. The kit can further include assay containers (tubes), buffers, or enzymes necessary for carrying out the detection assay. In embodiments, the kit further includes a sample collection device for collecting a sample from a subject. In embodiments, the human subject has a fibrotic pulmonary disease.

[0146] In another aspect, a kit is provided. The kit includes a marker protein binding agent capable of binding to a substance within a biological sample from a human subject having a fibrotic pulmonary disease, wherein the substance is a fibrotic pulmonary disease marker protein or fragment thereof set forth in the tables provided herein (e.g., Table 1A, 1B, 2A, 2B, 3A, 3B, 4A, or 4B); and a detecting reagent or a detecting apparatus capable of indicating binding of the marker protein binding agent to the substance.

[0147] In some embodiments, the kit includes components to examine more than one fibrotic pulmonary disease marker protein or fragment thereof. For example, the kit can include more than one marker protein binding agent capable of binding to one or more fibrotic pulmonary disease marker proteins or fragments thereof set forth in Table 1A, 1B, 2A, 2B, 3A, 3B, 4A, or 4B. In embodiments, the kit includes a plurality of marker protein binding agents. Where the kit includes a plurality of marker protein binding agents a plurality of fibrotic pulmonary disease marker proteins or fragments thereof (e.g., as set forth in Table 1A, 1B, 2A, 2B, 3A, 3B, 4A, or 4B) are detected.

[0148] The kit will generally include at least one vial, test tube, flask, bottle, syringe or other container means, into which the testing agent, can be suitably reacted or aliquoted. Kits can also include components for comparing results such as a suitable control sample, for example a positive and/or negative control. The kit can also include a collection device for collecting and/or holding the sample from the subject. The collection device can include a sterile swab or needle (for collecting blood), and/or a sterile tube (*e.g.*, for holding the swab or a bodily fluid sample).

[0149] In another aspect, a complex in vitro is provided. The complex includes a marker protein binding agent bound to a fibrotic pulmonary disease marker protein or fragment thereof set forth in the tables provided herein (*e.g.*, Table 1A, 1B, 2A, 2B, 3A, 3B, 4A, or 4B), wherein the fibrotic pulmonary disease marker protein is extracted from a human subject having or at risk of developing a fibrotic pulmonary disease. In embodiments, the subject has a fibrotic pulmonary disease.

[0150] In another aspect, a complex in vitro is provided. The complex includes a marker protein binding agent bound to a fibrotic pulmonary disease marker protein or fragment thereof set forth in the tables provided herein (*e.g.*, Table 1A, 1B, 2A, 2B, 3A, 3B, 4A, or 4B), wherein the fibrotic pulmonary disease marker protein is extracted from a human subject having a fibrotic pulmonary disease.

IV. METHODS OF DETECTION

[0151] The methods provided herein include the step of determining (detecting) an expression level of a fibrotic pulmonary disease marker protein or fragment thereof. Methods for detecting and identifying proteins and their interactions with other proteins or nucleic acid molecules involve conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature (*see, e.g.*, Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition 1989, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.; *Animal Cell Culture*, R. I. Freshney, ed., 1986).

[0152] "Determining an expression level of a protein" or "determining a level of expression of a protein" as provided herein includes methods and technologies well known in the art. For example, capture arrays for expression profiling may be used to determine an expression level of a protein. Capture arrays employ high affinity capture reagents, such as conventional antibodies, single domains, engineered scaffolds, peptides or nucleic acid aptamers, to bind and detect

specific target ligands in high throughput manner. Antibody arrays have the required properties of specificity and acceptable background, and some are available commercially (BD Biosciences, San Jose, CA; Clontech, Mountain View, CA; BioRad; Sigma, St. Louis, MO). Antibodies for capture arrays are made either by conventional immunization (polyclonal sera and hybridomas), or as recombinant fragments, usually expressed in *E. coli*, after selection from phage or ribosome display libraries (Cambridge Antibody Technology, Cambridge, UK; Biolnvent, Lund, Sweden; Affitech, Walnut Creek, CA; Biosite, San Diego, CA). In addition to the conventional antibodies, Fab and scFv fragments, single V-domains from camelids or engineered human equivalents (Domantis, Waltham, MA) are optionally useful in arrays.

[0153] The term scaffold refers to ligand-binding domains of proteins, which are engineered into multiple variants capable of binding diverse target molecules with antibody-like properties of specificity and affinity. The variants are produced in a genetic library format and selected against individual targets by phage, bacterial or ribosome display. Such ligand-binding scaffolds or frameworks include Affibodies based on *S. aureus* protein A (Affibody, Bromma, Sweden), Trinetins based on fibronectins (Phylos, Lexington, MA) and Anticalins based on the lipocalin structure (Pieris Proteolab, Freising-Weihestephane, Germany). These are used on capture arrays in a similar fashion to antibodies and have advantages of robustness and ease of production.

[0154] Nonprotein capture molecules, notably the nucleic acid aptamers which bind protein ligands with high specificity and affinity, are also used in arrays (SomaLogic, Boulder, CO). Aptamers are selected from libraries of oligonucleotides by the Selex™ procedure (SomaLogic, Boulder, CO) and their interaction with protein is enhanced by covalent attachment, through incorporation of brominated deoxyuridine and UV-activated crosslinking (photoaptamers). Photocrosslinking to ligand reduces the cross reactivity of aptamers due to the specific steric requirements. Aptamers have the advantages of ease of production by automated oligonucleotide synthesis and the stability and robustness of DNA; on photoaptamer arrays, universal fluorescent protein stains are used to detect binding.

[0155] Protein analytes binding to antibody arrays are detected directly or indirectly, for example, via a secondary antibody. Direct labeling is used for comparison of different samples with different colors. Where pairs of antibodies directed at the same protein ligand are available, sandwich immunoassays provide high specificity and sensitivity and are therefore the method of choice for low abundance proteins such as cytokines; they also give the possibility of detection

of protein modifications. Label-free detection methods, including mass spectrometry, surface plasmon resonance and atomic force microscopy, avoid alteration of ligand. What is required from any method is optimal sensitivity and specificity, with low background to give high signal to noise. Since analyte concentrations cover a wide range, sensitivity has to be tailored appropriately. Serial dilution of the sample or use of antibodies of different affinities are solutions to this problem. Proteins of interest are frequently those in low concentration in body fluids and extracts, requiring detection in the pg range or lower, such as cytokines or the low expression products in cells.

[0156] An alternative to an array of capture molecules is one made through molecular imprinting technology, in which peptides (e.g., from the C-terminal regions of proteins) are used as templates to generate structurally complementary, sequence-specific cavities in a polymerizable matrix; the cavities can then specifically capture (denatured) proteins that have the appropriate primary amino acid sequence (ProteinPrint™, Aspira Biosystems, Burlingame, CA).

[0157] Another methodology which is useful diagnostically and in expression profiling is the ProteinChip® array (CIPHERgen, Fremont, CA), in which solid phase chromatographic surfaces bind proteins with similar characteristics of charge or hydrophobicity from mixtures such as plasma or tumor extracts, and SELDI-TOF mass spectrometry is used to detect the retained proteins.

[0158] Large-scale functional chips have been constructed by immobilizing large numbers of purified proteins and are used to assay a wide range of biochemical functions, such as protein interactions with other proteins, drug-target interactions, enzyme-substrates, etc. Generally they require an expression library, cloned into E. coli, yeast or similar from which the expressed proteins are then purified, e.g., via a His tag and immobilized. Cell free protein transcription/translation is a viable alternative for synthesis of proteins which do not express well in bacterial or other in vivo systems.

[0159] In embodiments, determining (detecting) an expression level of a fibrotic pulmonary disease marker protein or fragment thereof as provided herein includes contacting a fibrotic pulmonary disease marker protein with a marker protein binding agent. A "marker protein binding agent" as provided herein refers to a substance capable of binding a fibrotic pulmonary disease marker protein. The marker protein binding agent may be a nucleic acid or a protein. In embodiments, the marker protein binding agent is an aptamer. In embodiments, the marker protein binding agent is peptide. In embodiments, the marker protein binding agent is a small

molecule. In embodiments, the marker protein binding agent is an antibody. In embodiments, the fibrotic pulmonary disease marker protein or fragment thereof is contacted with a marker protein binding agent in a biological sample (e.g., whole blood, serum or plasma). In embodiments, the marker protein binding agent includes a detectable moiety. In embodiments, the detectable moiety is a fluorescent moiety. In embodiments, the marker protein binding agent includes a capturing moiety. A "capturing moiety" refers to a protein or nucleic acid, which is covalently, through a linker or a chemical bond, or noncovalently attached to the marker protein binding agent and is capable of interacting with a capturing agent. An example of a capturing moiety useful for the methods provided herein is biotin. In embodiments, the capturing moiety is biotin. In embodiments, the capturing moiety is a cleavable capturing moiety. In embodiments, the capturing moiety is photocleavable biotin.

[0160] A "capturing agent" as provided herein refers to an agent capable of binding a capturing moiety. The interaction between the capturing moiety and the capturing agent may be a high affinity interaction, wherein the capturing moiety and the capturing agent bind to each other (e.g., biotin, streptavidin). An example of a capturing agent useful for the methods provided herein are streptavidin coated beads. In embodiments, the capturing agent is a streptavidin coated bead. Without limitation any suitable affinity binding pairs known in the art may be used as capturing moiety and capturing agent in the methods provided herein. For example, the capturing moiety may be an antibody and the capturing agent may be an antigen-coated bead. In embodiments, the capturing moiety is biotin and the capturing agent is a streptavidin coated bead.

[0161] The marker protein binding agent may bind non-covalently to the fibrotic pulmonary disease marker protein through ionic, van der Waals, electrostatic or hydrogen bonds. Upon binding of the marker protein binding agent to the fibrotic pulmonary disease marker protein a disease marker-protein binding agent complex is formed. The methods provided herein including embodiments thereof include detecting the disease marker-protein binding agent complex, thereby determining the expression level of a fibrotic pulmonary disease marker protein or fragment thereof in a biological sample. Thus, in embodiments, the determining includes (a) contacting a fibrotic pulmonary disease marker protein with a marker protein binding agent in the biological sample, thereby forming a disease marker protein-binding agent complex; and (b) detecting the disease marker protein-binding agent complex. The disease marker protein-binding agent complex may be separated from the sample and unbound components contained therein by contacting the disease marker protein-binding agent complex

with a capturing agent as described above (e.g., streptavidin-coated beads). Thus, in
embodiments, the detecting includes contacting the disease marker protein-binding agent
complex with a capturing agent, thereby forming a captured disease marker protein-binding
agent complex. The captured disease marker protein-binding agent complex may be washed to
5 remove any unbound components.

[0162] For the methods provided herein, the fibrotic pulmonary disease marker protein or
fragment thereof may be contacted with a tagging moiety. A "tagging moiety" as provided
herein is a composition capable of non-covalently binding to the fibrotic pulmonary disease
marker protein or fragment thereof. In embodiments, the tagging moiety is biotin. Upon binding
10 to the fibrotic pulmonary disease marker protein or fragment thereof, the tagging moiety may
bind through high affinity interaction with a tagging agent (e.g., streptavidin). In embodiments,
the fibrotic pulmonary disease marker protein or fragment thereof is contacted with a tagging
moiety after the formation of a captured disease marker protein-binding agent complex. In
embodiments, the fibrotic pulmonary disease marker protein or fragment thereof is contacted
15 with a tagging moiety before the formation of a captured disease marker protein-binding agent
complex. In embodiments, the fibrotic pulmonary disease marker protein or fragment thereof is
contacted with a tagging moiety at the same time as the formation of a captured disease marker
protein-binding agent complex. In embodiments, the detecting further includes (1) contacting
the captured disease marker protein-binding agent complex with a tagging moiety; thereby
20 forming a tagged disease marker protein-binding agent complex; and (2) separating the tagged
disease marker protein-binding agent complex from the biological sample.

[0163] Once the tagged disease marker protein-binding agent complex is separated from the
biological sample, the interaction between capturing moiety (e.g., photocleavable biotin) and
capturing agent (e.g., streptavidin-coated beads) is reversed (e.g. through cleavage of the
25 photocleavable biotin) and a cleaved disease marker protein-binding agent complex is formed.
The cleaved disease marker protein-binding agent complex includes a fibrotic pulmonary disease
marker protein or fragment thereof bound to a marker protein binding agent and a tagging
moiety. Thus, in embodiments, the detecting further includes after the separating of step (2)
separating the capturing binding agent from the tagged disease marker protein-binding agent
30 complex, thereby forming a cleaved disease marker protein-binding agent complex.

[0164] The cleaved disease marker protein-binding agent complex may be contacted with a
tagging agent (e.g., streptavidin-coated beads) and the tagging moiety (e.g., biotin) bound to the

fibrotic pulmonary disease marker protein or fragment thereof may form a high affinity interaction with the tagging agent. The cleaved disease marker protein-binding agent complex may be captured by a tagging agent (e.g., streptavidin-coated beads) and the marker protein binding agent may be subsequently separated, (e.g., eluted by affinity chromatography) from the
5 cleaved disease marker protein-binding agent complex. Once the marker protein binding agent (e.g., aptamer) is released (separated) from the cleaved disease marker protein-binding agent complex it may be quantified using standard techniques known in the art to quantify labeled nucleic acids molecules (e.g., hybridization to a custom DNA microarray). Thus, in
embodiments, the detecting further includes (3) separating the marker protein binding agent from
10 the cleaved disease marker protein-binding agent complex; thereby forming a released marker protein binding agent; and (4) determining an amount of released marker protein binding agent.

V. MARKER PROTEINS

[0165] The fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 1A, 1B, 2A, 2B, 3A, 3B, 4A, or 4B) are applicable to all methods, kits and compositions
15 described herein. For the methods, kits and compositions described herein the expression level of one or more fibrotic pulmonary disease marker proteins may be determined (detected). In embodiments, the expression level of at least one fibrotic pulmonary disease marker protein is determined (detected). In embodiments, the expression level of a plurality (e.g., 2, 3, 4, 5, 6, 7,
8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20) of fibrotic pulmonary disease marker proteins is
20 determined (detected). Where the expression level of a plurality of fibrotic pulmonary disease marker proteins is determined (detected), the expression level of a combination of any one of the fibrotic pulmonary disease marker proteins provided herein is determined (detected).

[0166] For example, the expression level of a combination of fibrotic pulmonary disease marker proteins set forth in Table 4A and/or Table 4B may be determined (detected) in the
25 methods, kits or compositions provided herein. In embodiments, the fibrotic pulmonary disease marker protein is a2-Macroglobulin, PUR8, iC3b, C4b, CAPG, PARC, Cathepsin H, Discoidin domain receptor 2, sICAM-5, LKHA4, MMP-7, MMP-9, NXPH1, OLR1, PTK6, or calgranulin B. In embodiments, the fibrotic pulmonary disease marker protein is a2-Macroglobulin, PUR8,
iC3b, C4b, CAPG, PARC, Cathepsin H, Discoidin domain receptor 2, sICAM-5, LKHA4,
30 MMP-7, MMP-9, NXPH1, OLR1, PTK6, and calgranulin B. In embodiments, the fibrotic pulmonary disease marker protein is a2-Macroglobulin. In embodiments, the fibrotic pulmonary disease marker protein is PUR8. In embodiments, the fibrotic pulmonary disease marker protein is iC3b. In embodiments, the fibrotic pulmonary disease marker protein is C4b. In

embodiments, the fibrotic pulmonary disease marker protein is CAPG. In embodiments, the fibrotic pulmonary disease marker protein is PARC. In embodiments, the fibrotic pulmonary disease marker protein is Cathepsin H. In embodiments, the fibrotic pulmonary disease marker protein is Discoidin domain receptor 2. In embodiments, the fibrotic pulmonary disease marker protein is sICAM-5. In embodiments, the fibrotic pulmonary disease marker protein is LKHA4. In embodiments, the fibrotic pulmonary disease marker protein is MMP-7. In embodiments, the fibrotic pulmonary disease marker protein is MMP-9. In embodiments, the fibrotic pulmonary disease marker protein is NXPH1. In embodiments, the fibrotic pulmonary disease marker protein is OLR1. In embodiments, the fibrotic pulmonary disease marker protein is PTK6. In embodiments, the fibrotic pulmonary disease marker protein is calgranulin B. In embodiments, the expression level of a2-Macroglobulin, PUR8, iC3b, C4b, CAPG, PARC, Cathepsin H, Discoidin domain receptor 2, sICAM-5, LKHA4, MMP-7, MMP-9, NXPH1, OLR1, PTK6, or calgranulin B is determined. In embodiments, the expression level of a2-Macroglobulin, PUR8, iC3b, C4b, CAPG, PARC, Cathepsin H, Discoidin domain receptor 2, sICAM-5, LKHA4, MMP-7, MMP-9, NXPH1, OLR1, PTK6, and calgranulin B is determined.

[0167] In embodiments, the expression level of a2-Macroglobulin and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of PUR8 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of iC3b and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of C4b and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of CAPG and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of PARC and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of Cathepsin H and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of Discoidin domain receptor 2 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of sICAM-5 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in

Table 4A or 4B) is determined (detected). In embodiments, the expression level of LKHA4 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in

Table 4A or 4B) is determined (detected). In embodiments, the expression level of MMP-7 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in

5 Table 4A or 4B) is determined (detected). In embodiments, the expression level of MMP-9 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in

Table 4A or 4B) is determined (detected). In embodiments, the expression level of NXPH1 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in

Table 4A or 4B) is determined (detected). In embodiments, the expression level of OLR1 and

10 one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in

Table 4A or 4B) is determined (detected). In embodiments, the expression level of PTK6 and

one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in

Table 4A or 4B) is determined (detected). In embodiments, the expression level of calgranulin B

and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in

15 Table 4A or 4B) is determined (detected).

[0168] In embodiments, the fibrotic pulmonary disease marker protein is AMPK a2b2gl,

CAMK2A, CAMK2B, CAMK2D, COMMD7, FER, FYN, Integrin albl, LYNB, METAP1,

MMP-3, PDPK1, phosphoglycerate kinase 1, SHC1, or SRCN1. In embodiments, the fibrotic pulmonary disease marker protein is AMPK a2b2gl, CAMK2A, CAMK2B, CAMK2D,

20 COMMD7, FER, FYN, Integrin albl, LYNB, METAP1, MMP-3, PDPK1, phosphoglycerate

kinase 1, SHC1, and SRCN1. In embodiments, the fibrotic pulmonary disease marker protein is

AMPK a2b2gl. In embodiments, the fibrotic pulmonary disease marker protein is CAMK2A.

In embodiments, the fibrotic pulmonary disease marker protein is CAMK2B. In embodiments,

the fibrotic pulmonary disease marker protein is CAMK2D. In embodiments, the fibrotic

25 pulmonary disease marker protein is COMMD7. In embodiments, the fibrotic pulmonary

disease marker protein is FER. In embodiments, the fibrotic pulmonary disease marker protein is

FYN. In embodiments, the fibrotic pulmonary disease marker protein is Integrin albl. In

embodiments, the fibrotic pulmonary disease marker protein is LYNB. In embodiments, the

fibrotic pulmonary disease marker protein is METAP1. In embodiments, the fibrotic pulmonary

30 disease marker protein is MMP-3. In embodiments, the fibrotic pulmonary disease marker

protein is PDPK1. In embodiments, the fibrotic pulmonary disease marker protein is

phosphoglycerate kinase 1. In embodiments, the fibrotic pulmonary disease marker protein is

SHC1. In embodiments, the fibrotic pulmonary disease marker protein is SRCN1. In

embodiments, the expression level of AMPK $\alpha 2$, CAMK2A, CAMK2B, CAMK2D, COMMD7, FER, FYN, Integrin $\alpha 1$, LYNB, METAP1, MMP-3, PDPK1, phosphoglycerate kinase 1, SHC1, or SRCN1 is determined. In embodiments, the expression level of AMPK $\alpha 2$, CAMK2A, CAMK2B, CAMK2D, COMMD7, FER, FYN, Integrin $\alpha 1$, LYNB, METAP1, MMP-3, PDPK1, phosphoglycerate kinase 1, SHC1, and SRCN1 is determined.

[0169] In embodiments, the expression level of AMPK $\alpha 2$ and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of CAMK2A and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of CAMK2B and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of CAMK2D and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of COMMD7 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of FER and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of FYN and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of Integrin $\alpha 1$ and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of LYNB and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of METAP1 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of MMP-3 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of PDPK1 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of phosphoglycerate kinase 1 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected). In embodiments, the expression level of SHC1 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in

Table 4A or 4B) is determined (detected). In embodiments, the expression level of SRCN1 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 4A or 4B) is determined (detected).

[0170] As provided herein the methods, kits and compositions may include determining (detecting) the expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B. Thus, in embodiments, the fibrotic pulmonary disease marker protein is TIMP-1, MMP-7, PTN, Activin A, HGF, Midkine, VEGF121, PDE3A, SBDS, Dkk-4, sICAM-5, SREC-I, ERK-1, DKK1, a2-Macroglobulin, NAGK, UFC1, or SGTA. In embodiments, the fibrotic pulmonary disease marker protein is TIMP-1, MMP-7, PTN, Activin A, HGF, Midkine, VEGF121, PDE3A, SBDS, Dkk-4, sICAM-5, SREC-I, ERK-1, DKK1, a2-Macroglobulin, NAGK, UFC1, and SGTA. In embodiments, the fibrotic pulmonary disease marker protein is TIMP-1. In embodiments, the fibrotic pulmonary disease marker protein is MMP-7. In embodiments, the fibrotic pulmonary disease marker protein is PTN. In embodiments, the fibrotic pulmonary disease marker protein is Activin A. In embodiments, the fibrotic pulmonary disease marker protein is HGF. In embodiments, the fibrotic pulmonary disease marker protein is Midkine. In embodiments, the fibrotic pulmonary disease marker protein is VEGF121. In embodiments, the fibrotic pulmonary disease marker protein is PDE3A. In embodiments, the fibrotic pulmonary disease marker protein is SBDS. In embodiments, the fibrotic pulmonary disease marker protein is Dkk-4. In embodiments, the fibrotic pulmonary disease marker protein is sICAM-5. In embodiments, the fibrotic pulmonary disease marker protein is SREC-I. In embodiments, the fibrotic pulmonary disease marker protein is ERK-1. In embodiments, the fibrotic pulmonary disease marker protein is DKK1. In embodiments, the fibrotic pulmonary disease marker protein is a2-Macroglobulin. In embodiments, the fibrotic pulmonary disease marker protein is NAGK. In embodiments, the fibrotic pulmonary disease marker protein is UFC1. In embodiments, the fibrotic pulmonary disease marker protein is SGTA. In embodiments, the expression level of TIMP-1, MMP-7, PTN, Activin A, HGF, Midkine, VEGF121, PDE3A, SBDS, Dkk-4, sICAM-5, SREC-I, ERK-1, DKK1, a2-Macroglobulin, NAGK, UFC1, or SGTA is determined. In embodiments, the expression level of TIMP-1, MMP-7, PTN, Activin A, HGF, Midkine, VEGF121, PDE3A, SBDS, Dkk-4, sICAM-5, SREC-I, ERK-1, DKK1, a2-Macroglobulin, NAGK, UFC1, and SGTA is determined.

[0171] In embodiments, the expression level of CCL18 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of YKL40 and one or more fibrotic pulmonary

disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of TIMP-1 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of MMP-7 and one or more fibrotic pulmonary

5 disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of PTN and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of Activin A and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is

10 determined (detected). In embodiments, the expression level of HGF and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of Midkine and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of VEGF121 and one or more

15 fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of PDE3A and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of SBDS and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B)

20 is determined (detected). In embodiments, the expression level of Dkk-4 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of sICAM-5 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of SREC-I and one or more

25 fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of ERK-1 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of DKK1 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B)

30 is determined (detected). In embodiments, the expression level of α 2-Macroglobulin and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of NAGK and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of UFCL and one or more

fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of SGTA and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected).

- 5 **[0172]** In embodiments, the fibrotic pulmonary disease marker protein is PGRP-S, sFRP-3, P-Cadherin, SSRP1, a2-HS-Glycoprotein, Persephin, ABL2, SCF sR, EMR2, MSP, Marapsin, NXPH1, ZAP70, CHL1, Rb, CAD 15, CK-MB, IL-1 sR9, Lymphotoxin a2/bl, PH, Survivin, IL-6 sRa, CLC7A, 17-beta-HSD 1, or NCAM-120. In embodiments, the fibrotic pulmonary disease marker protein is PGRP-S, sFRP-3, P-Cadherin, SSRP1, a2-HS-Glycoprotein, Persephin, ABL2, SCF sR, EMR2, MSP, Marapsin, NXPH1, ZAP70, CHL1, Rb, CAD 15, CK-MB, IL-1 sR9, Lymphotoxin a2/bl, PH, Survivin, IL-6 sRa, CLC7A, 17-beta-HSD 1, and NCAM-120. In
- 10 embodiments, the fibrotic pulmonary disease marker protein is PGRP-S. In embodiments, the fibrotic pulmonary disease marker protein is sFRP-3. In embodiments, the fibrotic pulmonary disease marker protein is P-Cadherin. In embodiments, the fibrotic pulmonary disease marker protein is SSRP1. In embodiments, the fibrotic pulmonary disease marker protein is a2-HS-Glycoprotein. In embodiments, the fibrotic pulmonary disease marker protein is Persephin. In
- 15 embodiments, the fibrotic pulmonary disease marker protein is ABL2. In embodiments, the fibrotic pulmonary disease marker protein is SCF sR. In embodiments, the fibrotic pulmonary disease marker protein is EMR2. In embodiments, the fibrotic pulmonary disease marker protein is MSP. In embodiments, the fibrotic pulmonary disease marker protein is Marapsin. In
- 20 embodiments, the fibrotic pulmonary disease marker protein is NXPH. In embodiments, the fibrotic pulmonary disease marker protein is ZAP70. In embodiments, the fibrotic pulmonary disease marker protein is CHL1. In embodiments, the fibrotic pulmonary disease marker protein is Rb. In embodiments, the fibrotic pulmonary disease marker protein is CAD 15. In
- 25 embodiments, the fibrotic pulmonary disease marker protein is CK-MB. In embodiments, the fibrotic pulmonary disease marker protein is IL-1 sR9. In embodiments, the fibrotic pulmonary disease marker protein is Lymphotoxin a2/bl. In embodiments, the fibrotic pulmonary disease marker protein is PH. In embodiments, the fibrotic pulmonary disease marker protein is Survivin. In embodiments, the fibrotic pulmonary disease marker protein is IL-6 sRa. In
- 30 embodiments, the fibrotic pulmonary disease marker protein is CLC7A. In embodiments, the fibrotic pulmonary disease marker protein is 17-beta-HSD 1. In embodiments, the fibrotic pulmonary disease marker protein is NCAM-120. In embodiments, the expression level of PGRP-S, sFRP-3, P-Cadherin, SSRP1, a2-HS-Glycoprotein, Persephin, ABL2, SCF sR, EMR2,

MSP, Marapsin, NXPH1, ZAP70, CHL1, Rb, CAD 15, CK-MB, IL-1 sR9, Lymphotoxin a2/bl, PH, Survivin, IL-6 sRa, CLC7A, 17-beta-HSD 1, or NCAM-120 is determined. In

embodiments, the expression level of PGRP-S, sFRP-3, P-Cadherin, SSRP1, a2-HS-

Glycoprotein, Persephin, ABL2, SCF sR, EMR2, MSP, Marapsin, NXPH1, ZAP70, CHL1, Rb,

5 CAD 15, CK-MB, IL-1 sR9, Lymphotoxin a2/bl, PH, Survivin, IL-6 sRa, CLC7A, 17-beta-HSD 1, and NCAM-120 is determined.

[0173] In embodiments, the expression level of PGRP-S and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined

(detected) (detected). In embodiments, the expression level of sFRP-3 and one or more fibrotic

10 pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is

determined (detected). In embodiments, the expression level of P-Cadherin and one or more

fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B)

is determined (detected). In embodiments, the expression level of SSRP1 and one or more

15 fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B)

is determined (detected). In embodiments, the expression level of a2-HS-Glycoprotein and one

or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table

3A or 3B) is determined (detected). In embodiments, the expression level of Persephin and one

or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table

3A or 3B) is determined (detected). In embodiments, the expression level of ABL2 and one or

20 more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A

or 3B) is determined (detected). In embodiments, the expression level of SCF sR and one or

more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A

or 3B) is determined (detected). In embodiments, the expression level of EMR2 and one or more

fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B)

25 is determined (detected). In embodiments, the expression level of MSP and one or more fibrotic

pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is

determined (detected). In embodiments, the expression level of Marapsin and one or more

fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B)

is determined (detected). In embodiments, the expression level of NXPH1 and one or more

30 fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B)

is determined (detected). In embodiments, the expression level of ZAP70 and one or more

fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B)

is determined (detected). In embodiments, the expression level of CHL1 and one or more

fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of Rb and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of CAD 15 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of CK-MB and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of IL-1 sR9 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of Lymphotoxin a2/bl and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of PH and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of Survivin and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of IL-6 sRa and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of CLC7A and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of 17-beta-HSD 1 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected). In embodiments, the expression level of NCAM-120 and one or more fibrotic pulmonary disease marker proteins as provided herein (e.g., set forth in Table 3A or 3B) is determined (detected).

[0174] For the methods, kits or compositions provided herein the expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A, IB, 2A, 2B, 3A, 3B, 4A and/or Table 4B may be determined (detected) in combination with the expression level of known markers associated with fibrotic pulmonary diseases. In embodiments, the expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A, IB, 2A, 2B, 3A, 3B, 4A or Table 4B is determined in combination with the expression level of MMP7, PARC/CCL18, YKL-40(Chitinase-Like-3), MMP3, SP-D, KL-6/Mucl, or Osteopontin (OPN). In embodiments, the expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A, IB, 2A, 2B, 3A, 3B, 4A or Table 4B is determined in combination with the

expression level of MMP7, PARC/CCL18, YKL-40(Chitinase-Like-3), MMP3, SP-D, KL-6/Mucl, and Osteopontin (OPN). In embodiments, the expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A, IB, 2A, 2B, 3A, 3B, 4A or Table 4B is determined in combination with the expression level of MMP1, SP-A, S100a2 (enrage),

- 5 Periostin, ICAM1, VCAM, IL8, VEGF, CD28, Hsp47, IL6, LDH, PAI-1, Protein C, Thrombomodulin, vonWillebrand Factor (vWF), MMP8, IL2Ralpha, IL2Rgamma, TGFB1, TGFB2, TGFB3, or TNFalpha. In embodiments, the expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A, IB, 2A, 2B, 3A, 3B, 4A or Table 4B is determined in combination with the expression level of MMP1, SP-A, S100a2 (enrage),
- 10 Periostin, ICAM1, VCAM, IL8, VEGF, CD28, Hsp47, IL6, LDH, PAI-1, Protein C, Thrombomodulin, vonWillebrand Factor (vWF), MMP8, IL2Ralpha, IL2Rgamma, TGFB1, TGFB2, TGFB3, and TNFalpha. In embodiments, the expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A, IB, 2A, 2B, 3A, 3B, 4A or Table 4B is determined in combination with the expression level of BNP, albumin, COMP, CCL1 1, CCL13,
- 15 CCL17, Serum amyloid A, or CXCL13. In embodiments, the expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A, IB, 2A, 2B, 3A, 3B, 4A or Table 4B is determined in combination with the expression level of BNP, albumin, COMP, CCL1 1, CCL13, CCL17, Serum amyloid A, and CXCL13.

- [0175] The following discussion of the invention is for the purposes of illustration and description, and is not intended to limit the invention to the form or forms disclosed herein.
- 20 Although the description of the invention has included description of one or more embodiments and certain variations and modifications, other variations and modifications are within the scope of the invention, *e.g.*, as may be within the skill and knowledge of those in the art, after understanding the present disclosure. All publications, patents, patent applications, Genbank
- 25 numbers, UniProt numbers, and websites cited herein are hereby incorporated by reference in their entireties for all purposes.

VI. SPECIFIC EMBODIMENTS

[0176] Samples and Sample Selection

- [0177] Plasma samples were obtained from IPF patients. Plasma samples for demographically matched healthy control subjects with a summary of health conditions and information regarding medication use were obtained from a separate collection protocol. All samples were obtained
- 30 under appropriate written Informed Consent. Histories were reviewed to provide the best

possible match of IPF and control groups. This excluded significant pulmonary disease from the healthy control group and recent use of prednisolone or other drugs (other than proton pump inhibitors) used off label in the treatment of IPF from the IPF cohort.

[0178] IPF patient samples were selected to include all patients that died. In addition, IPF patient samples were selected to include patients that experienced a broad range of subsequent FVC change. The ratio of male and female IPF patients (2.5: 1) is similar to that in the IPF population. Control samples were selected for a balanced design across groups. Normal Healthy control subjects were selected to be demographically consistent with the IPF patient samples for consistency with the clinical trial population (age/gender). Normal Healthy control samples were collected into comparable collection vials (i.e. with matched preservatives).. Both IPF and control samples were collected at multiple sites.

[0179] Table I. Summary demographics for IPF patients and healthy controls are shown below.

	Patients (n)	All		Male		Female	
		IPF	Controls	IPF	Controls	IPF	Controls
Gender		140	70	100	50	40	20
	Male	100 (71.4%)	50 (71.4%)	100%	100%		
	Females	40 (28.6%)	20 (28.6%)			100%	100%
Ethnic origin							
	White	100%	100%	100%	100%	100%	100%
Smoking status							
	Never	65 (32.1%)	22 (31.4)	31 (31%)	16 (32%)	14 (35%)	6 (30%)
	Previous	92 (65.7%)	46 (65.7)	67 (67%)	33 (66%)	25 (62.5%)	13 (65%)
	Current	3 (2.1%)	2 (2.9%)	2 (2%)	1 (2%)	1 (2.5%)	1 (5%)
Region							
	United States	98 (70.0%)	100%				
	Canada	10 (7.1%)					
	Europe	32 (22.9%)					
Age (year)		66.9 (7.2)	66.9 (7.3)	66.9 (7.2)	66.9 (7.7)	67.0 (7.3)	67.0 (6.5)
Weight (kg)		87.4 (18.2)	84.8 (17.1)	91.1 (17.7)	88.0 (15.7)	78.4 (16.4)	77.0 (18.2)
BMI		30.1 (5.2)		30.1 (5.0)		30.1 (5.4)	
Baseline							
	FVC (% predicted)	72.4 (13.2)		72.4 (12.5)		72.3 (14.0)	
	DLCO (% predicted, High-corrected)	45.9 (8.5)		45.4 (8.3)		46.8 (9.1)	
	6MWT (m)	395.2 (95.1)		408.7 (90.4)		362.4 (59.3)	
Outcome							
	Death	25 (17.9%)		16 (16%)		9 (22.5%)	
	All patients that died were selected						
	Δ FVC 24 weeks (% predicted)	-6.9		-6.7		-7.2	
	Δ FVC 48 weeks (% predicted)	-13.7		-12.7		-16.0	

[0180] From plasma of IPF patients and healthy controls, 1129 proteins were profiled using a broad aptamer-based proteomic platform (SomaLogic, Colorado, US). Briefly, the technology uses DNA aptamers containing chemically modified nucleotides as highly specific protein binding reagents in a multiplexed assay that transforms the quantity of each targeted protein into

a corresponding quantify of the aptamer (Fig.I). Protein quantities are then detected on a microarray platform and recorded as Relative Fluorescence Units (RFU). Data meeting quality control standards were obtained for 130/140 IPF patients and 69/70 controls.

[0181] From plasma of IPF patients and controls, proteins were profiled in commercially available sandwich immunoassays (ELISAs) from various companies (R&D Systems Inc., Minneapolis, MN; Abeam Inc., Cambridge, UK). In brief, the assay involves binding of the protein of interest to a capture matrix on a specific antibody and detection using a second enzymatically conjugated antibody, usually horse radish peroxidase. The HRP reacts with a chromogenic substrate to create a colorimetric reading which is read by a microplate reader and expressed as an OD. A standard curve is used to convert this to a defined concentration of protein. Data meeting quality control standards ranged from 191 to 213 in patient cohort #1 to 466 to 613 patients patient cohort #2. Survival and Progression Free Survival (PFS) Kaplan Meier Plots were generated using median and Rpart analysis.

[0182] Statistical analysis: Relative protein concentrations (measured in relative fluorescence units; RFU) were normalized against hybridization controls and median normalized (calibrated) across all groups. Normalized protein concentrations were log transformed for better fidelity with the Gaussian assumptions in the statistical models. Differential protein regulation was evaluated based on univariate tests (per protein). Differentially expressed proteins were identified by ANOVA with corrections for multiple comparisons (FDR <0.01 and fold change >1.2x). This resulted in the identification of 262 differentially expressed proteins of which 174 are up-regulated in IPF (Table 1A) relative to healthy controls and 88 are down-regulated (Table 1B). This analysis confirmed several previously known proteins that are differentially expressed in IPF including matrix metalloproteinase 7 (MMP7), chemokine (C-C motif) ligand (CCL18), and surfactant protein D (SFTPD).

[0183] In order to identify biological processes that are enriched in IPF, the pathways associated with differentially expressed proteins in IPF were compared to those associated with those on proteins on the 1129 protein panel as a whole. In each case, relevant processes were identified using MetaCore (Thomson Reuters, New York, NY). To identify processes that are overrepresented by the differentially expressed proteins in IPF relative to the protein panel, the rank order of processes associated with each case were compared. Fig.2 shows a plot of the highest ranked processes associated with differentially expressed proteins in IPF (Y axis) compared to the processes associated with the proteins on the panel as a whole (X axis). The

processes highlighted in the red box (lower right section) are enriched in IPF. Table II lists the associated processes that show a shift of at least 10 positions between IPF and the panel as a whole (with processes associated with the greatest change in rank at the top). Among these processes are those involved in TGF- β signaling, cell adhesion, and epithelial mesenchymal transition (EMT). These processes are known to be associated with IPF and fibrotic conditions and suggest that the plasma proteomic signature obtained is consistent with disease state.

[0184] Table II. The IPF plasma protein profile shows activation of pathways involved in TGF- β signaling, cell adhesion, and EMT.

Rank increase (X-Y)	Rank in IPF (Y-axis)	Process Network
66	17	Cardiac development BMP TGF beta signaling
65	22	Cell adhesion Synaptic contact
50	30	Cytoskeleton Macropinocytosis and its regulation
47	13	Apoptosis Anti-Apoptosis mediated by external signals by Estrogen
34	35	Reproduction Spermatogenesis, motility and copulation
32	44	Reproduction GnRH signaling pathway
28	49	Reproduction Gonadotropin regulation
25	16	Reproduction Male sex differentiation
25	38	Signal Transduction TGF-beta, GDF and Activin signaling
23	7	Cell cycle G1-S Growth factor regulation
22	14	Development Neurogenesis Axonal guidance
19	5	Development EMT Regulation of epithelial-to-mesenchymal transition
19	42	Inflammation MIF signaling
17	12	Cell adhesion Attractive and repulsive receptors
17	40	Development Synaptogenesis
17	41	Cell adhesion Cadherins
16	18	Inflammation Amphotericin signaling
15	39	Immune response Phagosome in antigen presentation
14	6	Apoptosis Anti-Apoptosis mediated by external signals via MAPK and JAK/STAT
14	36	Cell adhesion Cell-matrix interactions
13	27	Signal transduction WNT signaling

[0185] Identification of proteins that predict disease progression: Protein concentrations and patient outcomes (particularly time to death or disease progression) were analyzed using R. For this analysis, disease progression was defined as death or a >10% absolute decrease in percent predicted FVC or a >15% absolute decrease in percent predicted DLCO (corrected for Hb) at any time compared to baseline, documented by at 2 consecutive assessments performed at least 6 weeks apart. Patients were followed from Baseline to last known survival date or completion of study, whichever is shortest. Patients undergoing lung transplant were censored at the time of transplant.

[0186] For each of 1129 proteins, an initial analysis using a method tree-based model (CART, R package), partitioned patients into a high and low risk range of protein concentration that would maximize the difference in Hazard Ratios (HR) for mortality or progression free survival.

The output from this analysis was a protein concentration threshold (in RFU) for further evaluation. In a further analysis, the HR and p value associated with this threshold were determined and reported using a Cox proportional hazard (PH) model (survival package of R). An additional analysis stratified the patients by protein expression (in quartiles) and reported associated differences in either survival or PFS compared to the patients in the lowest quartile. The resultant Kaplan-Meier curves were manually evaluated resulting in the identification of 43 proteins that predict mortality, progression free survival, or both (Table 3; Fig.5- Fig.47). This list includes MMP-7, a previously reported marker of disease progression in IPF.

[0187] Unsupervised Hierarchical Clustering: Unsupervised hierarchical clustering was performed based on a subset of 150 differentially expressed proteins that met more stringent inclusion criteria (FDR<0.001 and FC>1.25). The Partek Genomics Suite was used for this analysis (Partek, St. Louis, MO). In this analysis, the majority IPF patients cluster together (Fig.3 branches 5-7). Most of the remaining patients form a secondary cluster has increased similarity to healthy controls (Fig.3; branch 1). One cluster of upregulated proteins and one cluster of downregulated proteins are consistent across the majority of IPF patients. The consistently upregulated proteins (Fig.4) includes the previously reported IPF plasma protein markers MMP-7 and CCL-18. Seven of the 17 protein in this cluster are differentially upregulated at the mRNA level in IPF lung tissue (InterMune analysis of data in the GEO database).

[0188] In addition, several differentially expressed proteins correspond to mRNAs previously shown to be differentially expressed in IPF lung tissue

p-value (FDR) threshold	0.01	0.01	0.01	0.01	0.01
Fold Change threshold	none	1.15	1.2	1.5	2
number of proteins above the threshold	351	311	262	94	20

[0189] Table IIIA and IIIB. Bivariate analysis of the association between mortality and plasma protein levels (i.e. MMP-7, CCL-18, YKL-40, Activin A and A2M) by Cox proportional hazards regression model. Protein levels were quantitated by ELISA in two independent patient cohorts (i.e. Cohort 1 in Table IIIA and Cohort 2 in Table IIIB).

[0190] Table IIIA:

	Cohort 1					
	N	Death	N High Conc.	HR	95% CI	p-value
MMP-7	221	25	74	2.3	1.0-5.0	0.041
CCL-18	213	24	140	2.8	1.0-8.1	0.062
YKL-40	191	20	105	2.0	0.8-5.1	0.165
Activin A	213	24	102	2.4	1.0-5.6	0.043
A2-macroglobulin	211	23	33	2.1	0.8-5.2	0.126

[0191] Table IIIB.

	Cohort 2					
	N	Death	N High Conc.	HR	95% CI	p-value
MMP-7	613	97	205	1.8	1.2-2.6	0.006
CCL-18	480	77	223	1.9	1.2-3.0	0.005
YKL-40	466	75	203	2.0	1.3-3.2	0.003
Activin A	468	76	176	2.6	1.7-4.2	3.23e-05
A2-macroglobulin	478	77	258	1.7	1.1-2.8	0.02

[0192] Table IV. Mortality risk of CCL18 and YKL40 determined by ELISA with 2 separate patient cohorts.

EntrezGeneID	EntrezGeneSymbol	UniProt	Target	Correlation with risk
6362	CCL18	P55774	PARC	positive
1116	CHI3L1	P36222	YKL40	positive

VII. TABLES

[0193] TABLE 1A. Fibrotic pulmonary disease biomarker proteins (upregulated).

Marker proteins have been identified with $p(\text{FDR}) < 0.01$, $\text{FC} > 1.2x$, $n=262$; 174 proteins upregulated in IPF vs. control are shown.

Target	Entrez GeneID	Entrez GeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
sICAM-5	7087	ICAM5	Q9UMF0	5.36E-16	1.6161	IPF-Baseline up vs Control
PUR8	158	ADSL	P30566	3.05E-14	2.87734	IPF-Baseline up vs Control

Target	Entrez GeneID	Entrez GeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
Cathepsin H	1512	CTSH	P09668	5.68E-13	1.30088	IPF-Baseline up vs Control
STRATIFIN	2810	SFN	P31947	1.15E-12	1.62451	IPF-Baseline up vs Control
iC3b	718	C3	P01024	6.55E-10	1.73599	IPF-Baseline up vs Control
MMP-7	4316	MMP7	P09237	2.06E-09	1.5265	IPF-Baseline up vs Control
LKHA4	4048	LTA4H	P09960	3.44E-09	8.53381	IPF-Baseline up vs Control
Soggy- 1	27120	DKKL1	Q9UK85	9.10E-09	1.50067	IPF-Baseline up vs Control
C3b	718	C3	P01024	2.16E-08	2.31267	IPF-Baseline up vs Control
CLC7A	64581	CLEC7A	Q9BXN2	3.87E-08	1.48678	IPF-Baseline up vs Control
BASI	682	BSG	P35613	4.77E-08	1.76885	IPF-Baseline up vs Control
ER	2099	ESR1	P03372	4.82E-08	1.48871	IPF-Baseline up vs Control
calgranulin B	6280	S100A9	P06702	6.55E-08	1.83576	IPF-Baseline up vs Control
SOD	6647	SOD1	P00441	7.89E-08	1.45524	IPF-Baseline up vs Control
Cathepsin A	5476	CTSA	P10619	1.34E-07	1.3084	IPF-Baseline up vs Control
cGMP-stimulated PDE	5138	PDE2A	000408	1.54E-07	1.42056	IPF-Baseline up vs Control
PARC	6362	CCL18	P55774	1.84E-07	1.40298	IPF-Baseline up vs Control
NID2	22795	NID2	Q14112	2.97E-07	1.44597	IPF-Baseline up vs Control
Cadherin-6	1004	CDH6	P55285	3.74E-07	1.6071	IPF-Baseline up vs Control
LGMN	5641	LGMN	Q99538	4.82E-07	1.28215	IPF-Baseline up vs Control
WISP-1	8840	WISP1	095388	5.92E-07	1.39604	IPF-Baseline up vs Control
tau	4137	MAPT	P10636	6.96E-07	1.83756	IPF-Baseline up vs Control
41	2035	EPB41	P11171	8.08E-07	1.53461	IPF-Baseline up vs Control
RANTES	6352	CCL5	P13501	9.62E-07	1.55401	IPF-Baseline up vs Control
Histone H1.2	3006	HIST1H1C	P16403	9.96E-07	3.83001	IPF-Baseline up vs Control
Bcl-2	596	BCL2	P10415	1.08E-06	1.60709	IPF-Baseline up vs Control

Target	Entrez GeneID	Entrez GeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
PRL	5617	PRL	P01236	1.25E-06	1.42039	IPF-Baseline up vs Control
Cadherin-12	1010	CDH 12	P55289	1.29E-06	1.88342	IPF-Baseline up vs Control
PKC-Z	5590	PRKCZ	Q05513	1.45E-06	1.90214	IPF-Baseline up vs Control
PSA1	5682	PSMA1	P25786	1.47E-06	1.22511	IPF-Baseline up vs Control
BMP10	27302	BMP10	O95393	1.56E-06	1.52213	IPF-Baseline up vs Control
Ephrin-B3	1949	EFNB3	Q15768	1.64E-06	1.58277	IPF-Baseline up vs Control
EphB4	2050	EPHB4	P54760	1.73E-06	1.47915	IPF-Baseline up vs Control
TGF-b3	7043	TGFB3	P10600	2.09E-06	1.40029	IPF-Baseline up vs Control
GFRa-3	2676	GFRA3	O60609	2.25E-06	1.29682	IPF-Baseline up vs Control
MMP-9	4318	MMP9	P14780	2.58E-06	1.48952	IPF-Baseline up vs Control
Protease nexin 1	5270	SERPINE2	P07093	3.09E-06	1.98673	IPF-Baseline up vs Control
Calcineurin B a	5534	PPP3R1	P63098	3.23E-06	1.39475	IPF-Baseline up vs Control
PTP-1C	5777	PTPN6	P29350	3.91E-06	1.90799	IPF-Baseline up vs Control
MMP-17	4326	MMP17	Q9ULZ9	4.58E-06	1.92995	IPF-Baseline up vs Control
HMG-1	3146	HMGB1	P09429	5.23E-06	1.97342	IPF-Baseline up vs Control
NCC27	1192	CLIC1	O00299	5.67E-06	2.043	IPF-Baseline up vs Control
TYK2	7297	TYK2	P29597	6.51E-06	1.51262	IPF-Baseline up vs Control
IL22RA1	58985	IL22RA1	Q8N6P7	7.73E-06	1.70245	IPF-Baseline up vs Control
RBM39	9584	RBM39	Q14498	7.90E-06	1.95172	IPF-Baseline up vs Control
Peroxisredoxin-6	9588	PRDX6	P30041	8.15E-06	1.74142	IPF-Baseline up vs Control
FCGR1	2209	FCGR1A	P12314	8.71E-06	1.59004	IPF-Baseline up vs Control
MOZ	7994	KAT6A	Q92794	8.86E-06	1.28366	IPF-Baseline up vs Control
SP-D	6441	SFTPD	P35247	9.08E-06	4.00143	IPF-Baseline up vs Control
Ubiquitin+1	6233	RPS27A	P62979	9.17E-06	1.32953	IPF-Baseline up vs Control

Target	Entrez GeneID	Entrez GeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
Histone H2A.Z	3015	H2AFZ	P0C0S5	1.04E-05	2.20185	IPF-Baseline up vs Control
a2-Macroglobulin	2	A2M	P01023	1.19E-05	1.60132	IPF-Baseline up vs Control
PECAM-1	5175	PECAM1	P16284	1.21E-05	1.23637	IPF-Baseline up vs Control
ON	6678	SPARC	P09486	1.25E-05	1.38543	IPF-Baseline up vs Control
MIF	4282	MIF	P14174	1.37E-05	1.62735	IPF-Baseline up vs Control
CaMKK alpha	84254	CAMKK1	Q8N5S9	1.69E-05	1.684	IPF-Baseline up vs Control
PSME1	5720	PSME1	Q06323	1.75E-05	1.7017	IPF-Baseline up vs Control
LY86	9450	LY86	095711	2.03E-05	1.25277	IPF-Baseline up vs Control
HPG-	3248	HPGD	P15428	2.61E-05	1.36537	IPF-Baseline up vs Control
TrkA	4914	NTRK1	P04629	2.64E-05	1.20709	IPF-Baseline up vs Control
Rab GDP dissociation inhibitor beta	2665	GDI2	P50395	3.61E-05	1.36366	IPF-Baseline up vs Control
PBEF	10135	NAMPT	P43490	3.63E-05	2.23087	IPF-Baseline up vs Control
CAPG	822	CAPG	P40121	3.69E-05	1.90263	IPF-Baseline up vs Control
PAI-1	5054	SERPINE1	P05121	3.80E-05	1.52823	IPF-Baseline up vs Control
BDNF	627	BDNF	P23560	3.90E-05	1.6203	IPF-Baseline up vs Control
ANGL3	27329	ANGPTL3	Q9Y5C1	4.07E-05	1.30295	IPF-Baseline up vs Control
CD30	943	TNFRSF8	P28908	4.28E-05	1.25432	IPF-Baseline up vs Control
FIt-3	2322	FLT3	P36888	4.47E-05	1.45412	IPF-Baseline up vs Control
G-CSF-R	1441	CSF3R	Q99062	4.55E-05	1.25163	IPF-Baseline up vs Control
tPA	5327	PLAT	P00750	4.88E-05	1.2056	IPF-Baseline up vs Control
IL-12	3592 3593	IL12A IL12B	P29459 P29460	5.19E-05	1.24153	IPF-Baseline up vs Control
XPNPEP1	7511	XPNPEP1	Q9NQW7	6.32E-05	1.57494	IPF-Baseline up vs Control
PAK6	56924	PAK6	Q9NQU5	6.40E-05	1.7035	IPF-Baseline up vs Control
Gro-b/g	None	CXCL3	P19876	6.56E-05	1.61602	IPF-Baseline up

Target	Entrez GeneID	Entrez GeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
		CXCL2	P19875			vs Control
HDGR2	84717	HDGFRP2	Q7Z4V5	6.57E-05	1.34838	IPF-Baseline up vs Control
PTK6	5753	PTK6	Q13882	6.80E-05	1.54905	IPF-Baseline up vs Control
NET4	59277	NTN4	Q9HB63	7.40E-05	1.27615	IPF-Baseline up vs Control
C4b	720 721	C4A C4B	P0C0L4 P0C0L5	8.41E-05	2.21215	IPF-Baseline up vs Control
al-Antitrypsin	5265	SERPINA1	P01009	8.42E-05	1.21183	IPF-Baseline up vs Control
prostatic binding protein	5037	PEBP1	P30086	8.56E-05	1.4349	IPF-Baseline up vs Control
OLR1	4973	OLR1	P78380	9.39E-05	1.77154	IPF-Baseline up vs Control
GHC2	83733	SLC25A18	Q9H1K4	9.52E-05	1.28086	IPF-Baseline up vs Control
GAS1	2619	GAS1	P54826	9.60E-05	1.58615	IPF-Baseline up vs Control
Nucleoside diphosphate kinase A	4830	NME1	P15531	9.93E-05	1.31321	IPF-Baseline up vs Control
IMDH2	3615	IMPDH2	P12268	9.96E-05	1.37417	IPF-Baseline up vs Control
PGP9.5	7345	UCHL1	P09936	0.00010213 8	1.44393	IPF-Baseline up vs Control
4EBP2	1979	EIF4EBP2	Q13542	0.00010254 8	1.27799	IPF-Baseline up vs Control
Discoidin domain receptor 2	4921	DDR2	Q16832	0.00010501 3	1.25463	IPF-Baseline up vs Control
hnRNP A2/B1	3181	HNRNPA2B 1	P22626	0.00010549 6	3.10531	IPF-Baseline up vs Control
TARC	6361	CCL17	Q92583	0.00011108 8	1.62767	IPF-Baseline up vs Control
Peroxiredoxin-1	5052	PRDX1	Q06830	0.00011153 3	1.4596	IPF-Baseline up vs Control
IL-12 Rb1	3594	IL12RB1	P42701	0.00011327	1.23088	IPF-Baseline up vs Control
MDHC	4190	MDH 1	P40925	0.00011599 4	1.32468	IPF-Baseline up vs Control
HO-2	3163	HMOX2	P30519	0.00012411 6	1.51311	IPF-Baseline up vs Control
IFN-lambda 2	282616	IL28A	Q8IZJ0	0.00012691 4	1.36754	IPF-Baseline up vs Control
RAP	4043	LRPAP1	P30533	0.00012715 8	1.4188	IPF-Baseline up vs Control
LDH-H 1	3945	LDHB	P07195	0.00013364 1	1.42348	IPF-Baseline up vs Control

Target	Entrez GeneID	Entrez GeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
MFRP	83552	MFRP	Q9BY79	0.000139768	2.08405	IPF-Baseline up vs Control
Epo	2056	EPO	P01588	0.00014028	1.65765	IPF-Baseline up vs Control
Myokinase human	203	AK1	P00568	0.000141109	1.31277	IPF-Baseline up vs Control
AK1A1	10327	AKR1A1	P14550	0.000147383	1.43383	IPF-Baseline up vs Control
GREM1	26585	GREM1	O60565	0.00015057	1.57243	IPF-Baseline up vs Control
NXPH1	30010	NXPH1	P58417	0.000152003	1.40755	IPF-Baseline up vs Control
CAMK1	8536	CAMK1	Q14012	0.000155007	1.23113	IPF-Baseline up vs Control
C3	718	C3	P01024	0.000157238	1.23997	IPF-Baseline up vs Control
PSA6	5687	PSMA6	P60900	0.000194759	1.44708	IPF-Baseline up vs Control
17-beta-HSD 1	3292	HSD17B1	P14061	0.000214675	1.27653	IPF-Baseline up vs Control
STX1a	6804	STX1A	Q16623	0.000239348	1.31737	IPF-Baseline up vs Control
DnaJ homolog	131118	DNAJC19	Q96DA6	0.000247922	1.21217	IPF-Baseline up vs Control
UBE2N	7334	UBE2N	P61088	0.00028064	1.51364	IPF-Baseline up vs Control
P-Selectin	6403	SELP	P16109	0.00028186	1.25965	IPF-Baseline up vs Control
RS7	6201	RPS7	P62081	0.000283023	2.12908	IPF-Baseline up vs Control
IDUA	3425	IDUA	P35475	0.000297778	1.20044	IPF-Baseline up vs Control
Triosephosphate isomerase	7167	TPI1	P60174	0.000298866	1.48785	IPF-Baseline up vs Control
AIF1	199	AIF1	P55008	0.000300015	1.39219	IPF-Baseline up vs Control
Topoisomerase 1	7150	TOPI	P11387	0.000302245	2.53892	IPF-Baseline up vs Control
UBC9	7329	UBE2I	P63279	0.000323054	1.61829	IPF-Baseline up vs Control
TSLP R	64109	CRLF2	Q9HC73	0.000341473	1.27071	IPF-Baseline up vs Control
Cytochrome P450 3A4	1576	CYP3A4	P08684	0.000343888	1.38319	IPF-Baseline up vs Control
eIF-5	1983	EIF5	P55010	0.000349155	1.36514	IPF-Baseline up vs Control
PIGR	5284	PIGR	P01833	0.000373112	1.49802	IPF-Baseline up vs Control

Target	Entrez GeneID	Entrez GeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
complement factor H-related 5	81494	CFHR5	Q9BXR6	0.00046835 1	1.45947	IPF-Baseline up vs Control
Ubiquitin	6233	RPS27A	P62979	0.00049552	1.82873	IPF-Baseline up vs Control
kallikrein 14	43847	KLK14	Q9P0G3	0.00050058 6	1.22009	IPF-Baseline up vs Control
HPV E7 Type 16	148907 9	Human- virus	P03129	0.00051024 7	1.23204	IPF-Baseline up vs Control
amyloid precursor protein	351	APP	P05067	0.00057953 3	1.33454	IPF-Baseline up vs Control
TCTP	7178	TPT1	P13693	0.00063108 1	1.51891	IPF-Baseline up vs Control
IL-1b	3553	IL1B	P01584	0.00063598 9	1.33065	IPF-Baseline up vs Control
PPIE	10450	PPIE	Q9UNP9	0.00064706 6	1.46298	IPF-Baseline up vs Control
NAGK	55577	NAGK	Q9UJ70	0.00065310 8	1.28239	IPF-Baseline up vs Control
RBP	5950	RBP4	P02753	0.00066015 4	1.52463	IPF-Baseline up vs Control
PA2G4	5036	PA2G4	Q9UQ80	0.00069210 9	1.50882	IPF-Baseline up vs Control
NAP-2	5473	PPBP	P02775	0.00071171 6	1.52551	IPF-Baseline up vs Control
KREM2	79412	KREMEN2	Q8NCW0	0.00082447 2	1.54929	IPF-Baseline up vs Control
TGF-b1	7040	TGFB1	P01137	0.00087606	1.2611	IPF-Baseline up vs Control
Angiopoietin-1	284	ANGPT1	Q15389	0.00089519 3	1.35839	IPF-Baseline up vs Control
PSA2	5683	PSMA2	P25787	0.00089601 9	1.30114	IPF-Baseline up vs Control
PF-4	5196	PF4	P02776	0.00090741 6	1.52612	IPF-Baseline up vs Control
YES	7525	YES1	P07947	0.00092361 2	1.24438	IPF-Baseline up vs Control
CTAP-III	5473	PPBP	P02775	0.00096789 1	1.42696	IPF-Baseline up vs Control
TWEAK	8742	TNFSF12	043508	0.00098247	1.38582	IPF-Baseline up vs Control
CSF-1	1435	CSF1	P09603	0.00099998 8	1.66978	IPF-Baseline up vs Control
FLRT1	23769	FLRT1	Q9NZU 1	0.00109578	1.28199	IPF-Baseline up vs Control
EP15R	58513	EPS15L1	Q9UBC2	0.00110187	1.27645	IPF-Baseline up vs Control
Azurocidin	566	AZU1	P20160	0.0011022	1.33249	IPF-Baseline up vs Control

Target	Entrez GeneID	Entrez GeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
COX-2	5743	PTGS2	P35354	0.00112433	1.51298	IPF-Baseline up vs Control
ASM 3A	10924	SMPDL3A	Q92484	0.00113407	1.45068	IPF-Baseline up vs Control
MCP-4	6357	CCL13	Q99616	0.00123469	1.53294	IPF-Baseline up vs Control
Myeloperoxidase	4353	MPO	P05164	0.00127381	1.43084	IPF-Baseline up vs Control
SKP1	6500	SKP1	P63208	0.00128357	1.29436	IPF-Baseline up vs Control
MMP-1	4312	MMP1	P03956	0.00137452	1.50942	IPF-Baseline up vs Control
RAN	5901	RAN	P62826	0.00138722	1.35935	IPF-Baseline up vs Control
PACAP-27	116	ADCYAP1	P18509	0.00147735	1.26374	IPF-Baseline up vs Control
NG36	10919	EHMT2	Q96KQ7	0.00155681	1.21559	IPF-Baseline up vs Control
MBD4	8930	MBD4	095243	0.00157949	1.46935	IPF-Baseline up vs Control
PSA-ACT	354 12	KLK3 SERPINA3	P07288 P01011	0.00175059	1.49302	IPF-Baseline up vs Control
PAFAH beta subunit	5049	PAFAH1B2	P68402	0.00176448	1.26625	IPF-Baseline up vs Control
OBCAM	4978	OPCML	Q14982	0.00177672	1.23158	IPF-Baseline up vs Control
BMPER	168667	BMPER	Q8N8U9	0.00180385	1.26938	IPF-Baseline up vs Control
BPI	671	BPI	P17213	0.00183319	4.13979	IPF-Baseline up vs Control
MATK	4145	MATK	P42679	0.00204296	1.57543	IPF-Baseline up vs Control
CATE	1510	CTSE	P14091	0.00209243	1.55577	IPF-Baseline up vs Control
Plasmin	5340	PLG	P00747	0.00216666	1.33246	IPF-Baseline up vs Control
CPNE1	8904	CPNE1	Q99829	0.00217877	1.55166	IPF-Baseline up vs Control
ULBP-3	79465	ULBP3	Q9BZM4	0.00223735	1.27446	IPF-Baseline up vs Control
Cripto	6997	TDGF1	P13385	0.00225129	1.30062	IPF-Baseline up vs Control
Calpain 1	823 826	CAPN 1 CAPNS1	P07384 P04632	0.00234514	1.27918	IPF-Baseline up vs Control
Thrombospondin-1	7057	THBS1	P07996	0.00248811	1.46794	IPF-Baseline up vs Control
CHST6	4166	CHST6	Q9GZX3	0.00249809	1.23918	IPF-Baseline up vs Control

Target	Entrez GeneID	Entrez GeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
Testican-1	6695	SPOCK1	Q08629	0.00253377	1.26887	IPF-Baseline up vs Control
TIMP-3	7078	TIMP3	P35625	0.00261046	1.42678	IPF-Baseline up vs Control
GX	8399	PLA2G10	015496	0.00279047	1.23456	IPF-Baseline up vs Control
PDE7A	5150	PDE7A	Q13946	0.00292702	1.23995	IPF-Baseline up vs Control
SSRP1	6749	SSRP1	Q08945	0.00300883	2.08277	IPF-Baseline up vs Control

[0194] TABLE IB. Fibrotic pulmonary disease biomarker proteins (downregulated). Marker proteins have been identified with $p(\text{FDR}) < 0.01$, $\text{FC} > 1.2x$, $n=262$; 88 proteins downregulated in IPF vs. control are shown.

Study 2 differentially expressed protein						
Target	Entrez GeneID	Entrez GeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
METAP1	23173	METAP1	P53582	9.21E-22	-3.52411	IPF-Baseline down vs Control
SLIK5	26050	SLITRK5	O94991	1.26E-16	-1.53867	IPF-Baseline down vs Control
sRAGE	177	AGER	Q15109	1.96E-15	-1.69619	IPF-Baseline down vs Control
TGF- β R III	7049	TGFBR3	Q03167	4.54E-14	-1.30835	IPF-Baseline down vs Control
RGM-C	148738	HFE2	Q6ZVN8	5.48E-13	-1.31264	IPF-Baseline down vs Control
NCAM-120	4684	NCAM1	P13591	6.20E-13	-1.30594	IPF-Baseline down vs Control
NCAM-L1	3897	L1CAM	P32004	1.37E-12	-1.33744	IPF-Baseline down vs Control
phosphoglycerate kinase 1	5230	PGK1	P00558	1.45E-12	-1.85791	IPF-Baseline down vs Control
CNTRF alpha	1271	CNTRF	P26992	2.18E-12	-1.49834	IPF-Baseline down vs Control
IL-1 sRI	3554	IL1R1	P14778	3.36E-12	-1.44229	IPF-Baseline down vs Control
Lymphotoxin $\alpha 1/\beta 2$	4049 4050	LTA LTB	P01374 Q06643	4.79E-12	-1.28121	IPF-Baseline down vs Control
bFGF-R	2260	FGFR1	P11362	4.84E-12	-1.23515	IPF-Baseline down vs Control
Endoglin	2022	ENG	P17813	3.75E-11	-1.44841	IPF-Baseline down vs Control
ATS13	11093	ADAMTS13	Q76LX8	3.76E-11	-1.34398	IPF-Baseline down vs Control
UNC5H4	137970	UNC5D	Q6UXZ4	3.77E-11	-1.50039	IPF-Baseline down vs Control
MATN2	4147	MATN2	O00339	4.08E-11	-1.2313	IPF-Baseline down vs Control
LRIG3	121227	LRIG3	Q6UXM1	6.28E-11	-1.38206	IPF-Baseline down vs Control
DAF	1604	CD55	P08174	1.59E-10	-1.23646	IPF-Baseline down vs Control
Sonic Hedgehog	6469	SHH	Q15465	2.58E-10	-1.52834	IPF-Baseline down vs Control

Study 2 differentially expressed protein							
CAMK2D	817	CAMK2D	Q13557	3.80E-10	-1.77385	IPF-Baseline down vs Control	
LYNB	4067	LYN	P07948	1.87E-09	-1.70397	IPF-Baseline down vs Control	
Semaphorin-6A	57556	SEMA6A	Q9H2E6	2.14E-09	-1.34688	IPF-Baseline down vs Control	
AMPK a2b2g1	5563 5565 5571	PRKAA2 PRKAB2 PRKAG1	P54646 O43741 P54619	2.59E-09	-1.84555	IPF-Baseline down vs Control	
FER	2241	FER	P16591	2.94E-09	-2.17179	IPF-Baseline down vs Control	
CAMK2B	816	CAMK2B	Q13554	4.33E-09	-1.74339	IPF-Baseline down vs Control	
MMP-3	4314	MMP3	P08254	8.42E-09	-2.73309	IPF-Baseline down vs Control	
Carbonic anhydrase 6	765	CA6	P23280	1.00E-08	-1.6323	IPF-Baseline down vs Control	
EphB6	2051	EPHB6	O15197	1.10E-08	-1.20157	IPF-Baseline down vs Control	
EphA1	2041	EPHA1	P21709	1.67E-08	-1.37492	IPF-Baseline down vs Control	
ERBB1	1956	EGFR	P00533	1.74E-08	-1.21591	IPF-Baseline down vs Control	
Ephrin-A5	1946	EFNA5	P52803	2.03E-08	-1.36768	IPF-Baseline down vs Control	
CAMK2A	815	CAMK2A	Q9UQM7	2.05E-08	-1.66666	IPF-Baseline down vs Control	
TrkC	4916	NTRK3	Q16288	2.39E-08	-1.24118	IPF-Baseline down vs Control	
WFKN2	124857	WFIKN2	Q8TEU8	3.40E-08	-1.34719	IPF-Baseline down vs Control	
RGMA	56963	RGMA	Q96B86	4.41E-08	-1.35993	IPF-Baseline down vs Control	
IDS	3423	IDS	P22304	5.26E-08	-1.3268	IPF-Baseline down vs Control	
TrkB	4915	NTRK2	Q16620	6.60E-08	-1.2045	IPF-Baseline down vs Control	
Cathepsin V	1515	CTSL2	O60911	8.41E-08	-1.42082	IPF-Baseline down vs Control	
Mn SOD	6648	SOD2	P04179	9.06E-08	-1.22193	IPF-Baseline down vs Control	
MP2K4	6416	MAP2K4	P45985	9.30E-08	-1.30101	IPF-Baseline down vs Control	
Cadherin-5	1003	CDH5	P33151	1.23E-07	-1.24254	IPF-Baseline down vs Control	
PCSK7	9159	PCSK7	Q16549	3.02E-07	-1.39555	IPF-Baseline down vs Control	
IL-1 R AcP	3556	IL1RAP	Q9NPH3	1.42E-06	-1.28716	IPF-Baseline down vs Control	
EDA	1896	EDA	Q92838	1.45E-06	-1.40884	IPF-Baseline down vs Control	
BOC	91653	BOC	Q9BWW1	1.78E-06	-1.30703	IPF-Baseline down vs Control	

Study 2 differentially expressed protein						
COMMD7	149951	COMMD7	Q86VX2	5.50E-06	-1.3293	IPF-Baseline down vs Control
FYN	2534	FYN	P06241	6.42E-06	-2.37353	IPF-Baseline down vs Control
SHC1	6464	SHC1	P29353	9.32E-06	-1.41344	IPF-Baseline down vs Control
PDGF Rb	5159	PDGFRB	P09619	1.47E-05	-1.40369	IPF-Baseline down vs Control
IGF-I sR	3480	IGF1R	P08069	1.63E-05	-1.25281	IPF-Baseline down vs Control
IGFBP-2	3485	IGFBP2	P18065	1.79E-05	-1.27751	IPF-Baseline down vs Control
sTie-2	7010	TEK	Q02763	2.29E-05	-1.2446	IPF-Baseline down vs Control
LY9	4063	LY9	Q9HBG7	2.42E-05	-1.30885	IPF-Baseline down vs Control
CD109	135228	CD109	Q6YHK3	2.86E-05	-1.42742	IPF-Baseline down vs Control
CYTT	1470	CST2	P09228	3.39E-05	-1.36018	IPF-Baseline down vs Control
LYN	4067	LYN	P07948	3.60E-05	-1.51471	IPF-Baseline down vs Control
XTP3A	79077	DCTPP1	Q9H773	4.51E-05	-1.23348	IPF-Baseline down vs Control
BGN	633	BGN	P21810	4.93E-05	-1.2413	IPF-Baseline down vs Control
ART	181	AGRP	O00253	5.06E-05	-1.31411	IPF-Baseline down vs Control
Elafin	5266	PI3	P19957	5.83E-05	-1.40487	IPF-Baseline down vs Control
ADAM 9	8754	ADAM9	Q13443	7.47E-05	-1.65831	IPF-Baseline down vs Control
PDPK1	5170	PDPK1	O15530	8.30E-05	-1.81173	IPF-Baseline down vs Control
sLeptin R	3953	LEPR	P48357	9.45E-05	-1.27922	IPF-Baseline down vs Control
BMP-7	655	BMP7	P18075	9.59E-05	-1.2765	IPF-Baseline down vs Control
Integrin a1b1	3672 3688	ITGA1 ITGB1	P56199 P05556	0.000113895	-1.54663	IPF-Baseline down vs Control
SRCN1	6714	SRC	P12931	0.000147453	-1.41024	IPF-Baseline down vs Control
JNK2	5601	MAPK9	P45984	0.000168267	-1.21905	IPF-Baseline down vs Control
IR	3643	INSR	P06213	0.000171593	-1.24842	IPF-Baseline down vs Control
GASP-2	114928	GPRASP2	Q96D09	0.000215307	-1.24857	IPF-Baseline down vs Control
b-ECGF	2246	FGF1	P05230	0.000217493	-1.20768	IPF-Baseline down vs Control
Siglec-3	945	CD33	P20138	0.000285117	-1.33873	IPF-Baseline down vs Control
CYTN	1469	CST1	P01037	0.000526955	-1.25847	IPF-Baseline down vs Control
LAG-3	3902	LAG3	P18627	0.000543879	-1.28161	IPF-Baseline down vs Control

Study 2 differentially expressed protein						
Cystatin M	1474	CST6	Q15828	0.000602595	-1.23413	IPF-Baseline down vs Control
Adiponectin	9370	ADIPOQ	Q15848	0.000685824	-1.24709	IPF-Baseline down vs Control
DC-SIGN	30835	CD209	Q9NNX6	0.000768396	-1.20751	IPF-Baseline down vs Control
PDK1	5163	PDK1	Q15118	0.00105885	-2.05488	IPF-Baseline down vs Control
PERL	4025	LPO	P22079	0.00108209	-1.23178	IPF-Baseline down vs Control
Nectin-like protein 2	23705	CADM1	Q9BY67	0.00130427	-1.20929	IPF-Baseline down vs Control
TEC	7006	TEC	P42680	0.00144851	-1.30303	IPF-Baseline down vs Control
Ck-b-8-1	6368	CCL23	P55773	0.00156062	-1.20917	IPF-Baseline down vs Control
Activin A	3624	INHBA	P08476	0.001905	-1.20745	IPF-Baseline down vs Control
Cofilin-1	1072	CFL1	P23528	0.00192739	-1.22991	IPF-Baseline down vs Control
PKC-A	5578	PRKCA	P17252	0.00215465	-1.31513	IPF-Baseline down vs Control
KPCT	5588	PRKCQ	Q04759	0.00227261	-1.79846	IPF-Baseline down vs Control
IL-5 Ra	3568	IL5RA	Q01344	0.00245521	-1.27362	IPF-Baseline down vs Control
BST1	683	BST1	Q10588	0.00295217	-1.22839	IPF-Baseline down vs Control
FGF-19	9965	FGF19	O95750	0.00303264	-1.42081	IPF-Baseline down vs Control

[0195] TABLE 2A. Fibrotic pulmonary disease biomarker proteins overlapping between study 1 and study 2 (upregulated). Marker proteins (n=8) that have been identified with $p(\text{FDR}) < 0.01$, FOL.2, up-regulated in study 2 (Table 1A, n=174) and that have been identified with $p(\text{FDR}) < 0.01$, FOL.15, up-regulated in study 1 (n=125); and are upregulated in IPF vs.

5 control are shown.

Consistent differentially expressed proteins in Study 1 and 2					Study 2 data		Study 1 data	
Entrez GeneID	Entrez GeneSymbol	UniProt	Target	Fold-Change	p-value	Fold Change	p-value	Fold Change
7087	ICAM5	Q9UMF0	sICAM-5	IPF1 up vs control	5.36E-16	1.6161	2.42E-08	2.07
1512	CTSH	P09668	Cathepsin H	IPF1 up vs control	5.68E-13	1.30088	1.37E-06	1.45
718	C3	P01024	iC3b	IPF1 up vs control	6.55E-10	1.73599	3.15E-03	1.44
4316	MMP7	P09237	MMP-7	IPF1 up vs control	2.06E-09	1.5265	2.38E-07	2.71
4048	LTA4H	P09960	LKHA4	IPF1 up vs control	3.44E-09	8.53381	4.48E-03	3.24
6362	CCL18	P55774	PARC	IPF1 up vs control	1.84E-07	1.40298	3.15E-03	1.5
2	A2M	P01023	a2-Macroglobulin	IPF1 up vs control	1.19E-05	1.60132	1.76E-03	1.36
3425	IDUA	P35475	IDUA	IPF1 up vs control	0.000297778	1.20044	7.00E-03	1.36

[0196] TABLE 2B. Fibrotic pulmonary disease biomarker proteins overlapping between study 1 and study 2 (downregulated). Marker proteins (n=20) that have been identified with $p(\text{FDR}) < 0.01$, FOL.2, down-regulated in study 2 (Table 1A, n=88) and that have been identified with $p(\text{FDR}) < 0.01$, FOL.15, down-regulated in study 1 (n=175); and are downregulated in IPF vs. control are shown.

10

Consistent differentially expressed proteins in Study 1 and 2					Study 2 data		Study 1 data	
Entrez GeneID	EntrezGene Symbol	UniProt	Target	Fold-Change	p-value	Fold Change	p-value	Fold Change
5230	PGK1	P00558	phosphoglycerate kinase 1	IPF1 down vs control	1.45E-12	-1.85791	9.50E-05	-1.42
817	CAMK2D	Q13557	CAMK2D	IPF1 down vs control	3.80E-10	-1.77385	2.38E-07	-3.21
4067	LYN	P07948	LYN	IPF1 down vs control	1.87E-09	-1.70397	1.75E-04	-2.08
5563 5565 5571	PRKAA2 PRKAB2 PRKAG1	P54646 O43741 P54619	AMPK a2b2g1	IPF1 down vs control	2.59E-09	-1.84555		
2241	FER	P16591	FER	IPF1 down vs control	2.94E-09	-2.17179	4.56E-06	-3.5
816	CAMK2B	Q13554	CAMK2B	IPF1 down vs control	4.33E-09	-1.74339	3.13E-05	-2.8
815	CAMK2A	Q9UQM7	CAMK2A	IPF1 down vs control	2.05E-08	-1.66666	2.38E-07	-3.79
4916	NTRK3	Q16288	TrkC	IPF1 down vs control	2.39E-08	-1.24118	1.38E-07	-3.43
3423	IDS	P22304	IDS	IPF1 down vs control	5.26E-08	-1.3268	2.71E-02	-1.24
149951	COMMMD7	Q86VX2	COMMMD7	IPF1 down vs control	5.50E-06	-1.3293	4.46E-02	-1.5
2534	FYN	P06241	FYN	IPF1 down vs control	6.42E-06	-2.37353	1.48E-08	-2.09
6464	SHC1	P29353	SHC1	IPF1 down vs control	9.32E-06	-1.41344	5.08E-07	-2.71
4067	LYN	P07948	LYN	IPF1 down vs control	3.60E-05	-1.51471	6.01E-08	-2.11
633	BGN	P21810	BGN	IPF1 down vs control	4.93E-05	-1.2413	3.21E-03	-2.01
5170	PDPK1	O15530	PDPK1	IPF1 down vs control	8.30E-05	-1.81173	3.71E-05	-1.57
3672 3688	ITGA1 ITGB1	P56199 P05556	Integrin a1b1	IPF1 down vs control	0.000113895	-1.54663	7.91E-05	-3.74
7006	TEC	P42680	TEC	IPF1 down vs control	0.00144851	-1.30303	3.30E-04	-1.95
1072	CFL1	P23528	Cofilin-1	IPF1 down vs control	0.00192739	-1.22991	4.48E-03	-1.25
5578	PRKCA	P17252	PKC-A	IPF1 down vs control	0.00215465	-1.31513	2.42E-08	-2.34
5588	PRKCQ	Q04759	KPCT	IPF1 down vs control	0.00227261	-1.79846	7.99E-03	-1.65
							4.48E-03	-1.53

[0197] TABLE 3A. Fibrotic pulmonary disease biomarker proteins that predict disease progression (positively correlated with risk).

Entrez GeneID	EntrezGene Symbol	UniProt	Target	Mortality high risk	PFS high risk	correlation with risk	Mortality p-value	PFS p-value
7076	TIMP1	P01033	TIMP-1	High 82	High 88	positive	0.0286	0.00251
4316	MMP7	P09237	MMP-7	High 78	High 79	positive	0.0142	0.000637
5764	PTN	P21246	PTN	High 49	High 60	positive	0.00235	0.00537
3624	INHBA	P08476	Activin A	High 31	High 35	positive	0.099	0.00987
3082	HGF	P14210	HGF	High 109	High 109	positive	0.0255	0.00583
4192	MDK	P21741	Midkine	High 67	High 74	positive	0.023	0.00489
7422	VEGFA	P15692	VEGF121	High 94	High 43	positive	0.00664	0.00254
5139	PDE3A	Q14432	PDE3A	High 45	High 34	positive	0.0214	0.0215
51119	SBDS	Q9Y3A5	SBDS	High 31	High 32	positive	0.00365	0.0427
27121	DKK4	Q9UBT3	Dkk-4	High 51	High 64	positive	0.00529	0.0625
7087	ICAM5	Q9UMF0	sICAM-5	High 123	High 102	positive	0.229	0.00223
8578	SCARF1	Q14162	SREC-I	High 36	High 24	positive	0.00057	0.000365
5595	MAPK3	P27361	ERK-1	High 53	High 53	positive	0.00218	0.0681
22943	DKK1	O94907	DKK1	High 78	High 78	positive	0.0154	0.0157
2	A2M	P01023	a2-Macroglobulin	High 30	Low 121	positive	0.0294	0.174
55577	NAGK	Q9UJ70	NAGK	High 87	Low 118	positive	0.0081	0.166
51506	UFC1	Q9Y3C8	UFC1	High 63	High 52	positive	0.00892	0.135
6449	SGTA	O43765	SGTA	High 52	High 40	positive	0.00721	0.0566

[0198] TABLE 3B. Fibrotic pulmonary disease biomarker proteins that predict disease progression (negatively correlated with risk).

Entrez GeneID	EntrezGene Symbol	UniProt	Target	Mortality high risk	PFS high risk	correlation with risk	Mortality p-value	PFS p-value	Rank
8993	PGLYRP1	075594	PGRP-S	Low 31	Low 34	negative	0.000324	0.000662	3
2487	FRZB	Q92765	sFRP-3	Low 53	Low 80	negative	0.00825	0.0058	3
1001	CDH3	P22223	P-Cadherin	Low 45	Low 63	negative	0.00171	0.00217	4
6749	SSRP1	Q08945	SSRP1	Low 31	Low 31	negative	0.0105	0.00317	5
197	AHSG	P02765	a2-HS-Glycoprotein	Low 36	Low 36	negative	5.27e-05	0.000759	6
5623	PSPN	060542	Persephin	Low 34	Low 43	negative	0.000166	0.0206	6
27	ABL2	P42684	ABL2	Low 21	Low 39	negative	0.00648	0.00353	7
3815	KIT	P10721	SCF sr	Low 27	Low 76	negative	0.00122	0.0343	7
30817	EMR2	Q9UHX3	EMR2	Low 64	Low 69	negative	0.00183	0.044	9
4485	MST1	P26927	MSP	Low 88	Low 90	negative	0.00839	0.0153	9
83886	PRSS27	Q9BQR3	Marapsin	Low 59	Low 10	negative	0.00187	0.0087	10
30010	NXPH 1	P58417	NXPH 1	Low 34	Low 34	negative	0.00843	0.0226	10
7535	ZAP 70	P43403	ZAP70	Low 60	Low 26	negative	0.0399	0.000135	10
10752	CHL1	000533	CHL1	Low 47	Low 50	negative	0.000716	0.0835	11
5925	RBI	P06400	Rb	Low 97	Low 87	negative	0.0133	0.0466	11
1013	CDH15	P55291	CAD15	Low 65	Low 10	negative	0.0292	0.000523	12
1152 1158	CKB CKM	P12277	CK-MB	Low 83	Low 91	negative	0.0122	0.0328	12
26280	IL1RAPL2	Q9NP60	IL-1 sR9	Low 73	Low 52	negative	0.00508	0.0669	12
4049 4050	LTA LTB	P01374	Lymphotoxin a2/b1	Low 8	Low 66	negative	0.000341	0.00625	12
5539	PPY	P01298	PH	Low 41	Low 68	negative	0.000555	0.00305	12
332	BIRC5	015392	Survivin	Low 71	Low 34	negative	0.0456	0.016	12
3570	IL6R	P08887	IL-6 sRa	Low 94	Low 8	negative	0.00141	0.0889	13
64581	CLEC7A	Q9BXN2	CLC7A	Low 72	Low 76	negative	0.0474	0.00921	13

Entrez GeneID	EntrezGene Symbol	UniProt	Target	Mortality high risk	PFS high risk	correlation with risk	Mortality p-value	PFS p-value	Rank
3292	HSD17B1	P14061	17-beta-HSD 1	Low 87	Low 83	negative	0.0242	0.00483	14
4684	NCAM1	P13591	NCAM-120	Low 51	Low 54	negative	0.0043	0.0522	14

[0199] **TABLE 4A. Fibrotic pulmonary disease biomarker proteins (upregulated).** Marker proteins have been identified with $p(\text{FDR}) < 0.01$, $\text{FC} > 1.2x$, $n = 262$; 16 proteins upregulated in IPF vs. control are shown.

Target	EntrezGeneID	EntrezGeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
a2- Macroglobulin	2	A2M	P01023	1.19E-05	1.60132	IPF-Baseline up vs Control
PUR8	158	ADSL	P30566	3.05E-14	2.87734	IPF-Baseline up vs Control
iC3b	718	C3	P01024	6.55E-10	1.73599	IPF-Baseline up vs Control
C4b	720 721	C4A C4B	P0C0L4 P0C0L5	8.41E-05	2.21215	IPF-Baseline up vs Control
CAPG	822	CAPG	P40121	3.69E-05	1.90263	IPF-Baseline up vs Control
PARC	6362	CCL18	P55774	1.84E-07	1.40298	IPF-Baseline up vs Control
Cathepsin H	1512	CTSH	P09668	5.68E-13	1.30088	IPF-Baseline up vs Control
Discoidin domain receptor 2	4921	DDR2	Q16832	0.000105	1.25463	IPF-Baseline up vs Control
sICAM-5	7087	ICAM5	Q9UMF0	5.36E-16	1.6161	IPF-Baseline up vs Control

Target	EntrezGeneID	EntrezGeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
LKHA4	4048	LTA4H	P09960	3.44E-09	8.53381	IPF-Baseline up vs Control
MMP-7	4316	MMP7	P09237	2.06E-09	1.5265	IPF-Baseline up vs Control
MMP-9	4318	MMP9	P14780	2.58E-06	1.48952	IPF-Baseline up vs Control
NXPH1	30010	NXPH1	P58417	0.000152	1.40755	IPF-Baseline up vs Control
OLR1	4973	OLR1	P78380	9.39E-05	1.77154	IPF-Baseline up vs Control
PTK6	5753	PTK6	Q13882	6.80E-05	1.54905	IPF-Baseline up vs Control
calgranulin B	6280	S100A9	P06702	6.55E-08	1.83576	IPF-Baseline up vs Control

[0200] TABLE 4B. Fibrotic pulmonary disease biomarker proteins (downregulated). Marker proteins have been identified with $p(\text{FDR}) < 0.01$, $\text{FC} > 1.2x$, $n=262$; 15 proteins downregulated in IPF vs. control are shown.

Target	EntrezGeneID	EntrezGeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
AMPK a2b2g1	5563 5565	PRKAA2 PRKAB2	P54646			IPF-Baseline
	5571	PRKAG1	O43741	2.59E-09	-1.84555	down vs Control
			P54619			IPF-Baseline
CAMK2A	815	CAMK2A	Q9UQM7	2.05E-08	-1.666666	down vs Control
CAMK2B	816	CAMK2B	Q13554	4.33E-09	-1.74339	IPF-Baseline
CAMK2D						down vs Control
	817	CAMK2D	Q13557	3.80E-10	-1.77385	IPF-Baseline
COMMD7						down vs Control
	149951	COMMD7	Q86VX2	5.50E-06	-1.3293	IPF-Baseline
FER						down vs Control
	2241	FER	P16591	2.94E-09	-2.17179	IPF-Baseline
FYN						down vs Control
	2534	FYN	P06241	6.42E-06	-2.37353	IPF-Baseline
Integrin a1b1						down vs Control
	3672 3688	ITGA1 ITGB1	P56199 P05556	0.000114	-1.54663	IPF-Baseline

Target	EntrezGeneID	EntrezGeneSymbol	UniProt	p-value	FoldChange (IPF vs. Control)	FoldChange (Description)
LYNB	4067	LYN	P07948	1.87E-09	-1.70397	IPF-Baseline down vs Control
METAP1	23173	METAP1	P53582	9.21E-22	-3.52411	IPF-Baseline down vs Control
MMP-3	4314	MMP3	P08254	8.42E-09	-2.73309	IPF-Baseline down vs Control
PDPK1	5170	PDPK1	O15530	8.30E-05	-1.81173	IPF-Baseline down vs Control
phosphoglycerate kinase 1	5230	PGK1	P00558	1.45E-12	-1.85791	IPF-Baseline down vs Control
SHC1	6464	SHC1	P29353	9.32E-06	-1.41344	IPF-Baseline down vs Control
SRCN1	6714	SRC	P12931	0.000147	-1.41024	IPF-Baseline down vs Control

VIII. EMBODIMENTS

[0201] Embodiment 1. A method of determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB in a subject that has or is at risk for developing a fibrotic pulmonary disease, said method comprising: (i) obtaining a biological sample from said subject; (ii) determining an expression level of a fibrotic pulmonary disease marker protein or fragment thereof set forth in Table 1A or Table IB in said biological sample.

[0202] Embodiment 2. The method of embodiment 1, wherein said determining comprises: (a) contacting a fibrotic pulmonary disease marker protein with a marker protein binding agent in said biological sample, thereby forming a disease marker protein-binding agent complex; and (b) detecting said disease marker protein-binding agent complex.

[0203] Embodiment 3. The method of embodiment 2, wherein said marker protein binding agent comprises a detectable moiety.

[0204] Embodiment 4. The method of embodiment 3, wherein said marker protein binding agent comprises a capturing moiety.

[0205] Embodiment 5. The method of embodiment 4, wherein said capturing moiety is a cleavable capturing moiety.

[0206] Embodiment 6. The method of any one of embodiments 2-5, wherein said detecting comprises contacting said disease marker protein-binding agent complex with a capturing agent, thereby forming a captured disease marker protein-binding agent complex.

[0207] Embodiment 7. The method of embodiment 6, wherein said detecting further comprises: (1) contacting said captured disease marker protein-binding agent complex with a tagging moiety; thereby forming a tagged disease marker protein-binding agent complex; and (2) separating said tagged disease marker protein-binding agent complex from said biological sample.

[0208] Embodiment 8. The method of embodiment 7, wherein said detecting further comprises after said separating of step (2) separating said capturing binding agent from said tagged disease marker protein-binding agent complex, thereby forming a cleaved disease marker protein-binding agent complex.

[0209] Embodiment 9. The method of embodiment 8, wherein said detecting further comprises: (3) separating said marker protein binding agent from said cleaved disease marker

protein-binding agent complex; thereby forming a released marker protein binding agent; and (4) determining an amount of released marker protein binding agent.

[0210] Embodiment 10. The method of any one of embodiments 1-9, further comprising selecting a subject that has or is at risk for developing a fibrotic pulmonary disease.

5 [0211] Embodiment 11. The method of any one of embodiments 1-9, wherein said fibrotic pulmonary disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia.

[0212] Embodiment 12. The method of any one of embodiments 1-9, wherein said fibrotic pulmonary disease marker protein is a progressive fibrotic pulmonary disease marker protein.

10 [0213] Embodiment 13. The method of embodiment 12, wherein said subject has or is at risk for developing a progressive fibrotic pulmonary disease.

[0214] Embodiment 14. The method of embodiment 13, wherein said progressive fibrotic pulmonary disease is idiopathic pulmonary fibrosis.

[0215] Embodiment 15. The method of embodiment 1, wherein said biological sample is a blood-derived biological sample of said subject.

15 [0216] Embodiment 16. The method of embodiment 15, wherein said blood-derived biological sample is whole blood, serum or plasma.

[0217] Embodiment 17. The method of embodiment 1, wherein said biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

20 [0218] Embodiment 18. The method of embodiment 1, wherein an expression level of a fibrotic pulmonary disease marker protein set forth in Table 1A is determined.

[0219] Embodiment 19. The method of embodiment 18, further comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 1A.

25 [0220] Embodiment 20. The method of embodiment 19, wherein said modulator is an antagonist.

[0221] Embodiment 21. The method of embodiment 20, wherein said antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0222] Embodiment 22. The method of embodiment 20, wherein said expression level of said fibrotic pulmonary disease marker protein set forth in Table 1A is elevated relative to a standard control.

5 [0223] Embodiment 23. The method of embodiment 1, wherein an expression level of a fibrotic pulmonary disease marker protein set forth in Table IB is determined.

[0224] Embodiment 24. The method of embodiment 23, further comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table IB.

10 [0225] Embodiment 25. The method of embodiment 24, wherein said modulator is an agonist.

[0226] Embodiment 26. The method of embodiment 24, wherein said agonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

15 [0227] Embodiment 27. The method of embodiment 25, wherein said expression level of said fibrotic pulmonary disease marker protein set forth in Table IB is decreased relative to a standard control.

[0228] Embodiment 28. The method of embodiment 19 or 24, further comprising administering to said subject an effective amount of a further therapeutic agent.

[0229] Embodiment 29. The method of embodiment 28, wherein said therapeutic agent is an idiopathic pulmonary fibrosis drug or an anti-fibrotic drug.

20 [0230] Embodiment 30. The method of embodiment 28, wherein said idiopathic pulmonary fibrosis drug is a mucolytic drug.

[0231] Embodiment 31. A method of determining whether a subject has or is at risk of developing a fibrotic pulmonary disease, said method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table
25 IB in a subject; (ii) determining whether said expression level is increased or decreased relative to a standard control, wherein an elevated expression level of a fibrotic pulmonary disease marker protein in Table 1A or a decreased expression level of a fibrotic pulmonary disease marker protein in Table IB relative to said standard control indicates that said subject has or is at risk of developing a fibrotic pulmonary disease; and (iii) based at least in part on said expression

level in step (ii), determining whether said subject has or is at risk for developing a fibrotic pulmonary disease.

[0232] Embodiment 32. The method of embodiment 31, further comprising selecting a subject that has or is at risk for developing a fibrotic pulmonary disease.

5 [0233] Embodiment 33. The method of embodiment 31, wherein said fibrotic pulmonary disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia.

[0234] Embodiment 34. The method of embodiment 31, wherein said one or more fibrotic pulmonary disease marker proteins is a progressive fibrotic pulmonary disease marker protein.

10 [0235] Embodiment 35. The method of embodiment 34, wherein said subject has or is at risk for developing a progressive fibrotic pulmonary disease.

[0236] Embodiment 36. The method of embodiment 35, wherein said progressive fibrotic pulmonary disease is idiopathic pulmonary fibrosis.

15 [0237] Embodiment 37. The method of embodiment 31, wherein said expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB is detected from a biological sample of said subject.

[0238] Embodiment 38. The method of embodiment 37, wherein said biological sample is a blood-derived biological sample of said subject.

[0239] Embodiment 39. The method of embodiment 38, wherein said blood-derived biological sample is whole blood, serum or plasma.

20 [0240] Embodiment 40. The method embodiment 37, wherein said biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0241] Embodiment 41. The method of embodiment 31, further comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 1A.

25 [0242] Embodiment 42. The method of embodiment 41, wherein said modulator is an antagonist.

[0243] Embodiment 43. The method of embodiment 42, wherein said antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0244] Embodiment 44. The method of embodiment 31, further comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table IB.

5 [0245] Embodiment 45. The method of embodiment 44, wherein said modulator is an agonist.

[0246] Embodiment 46. The method of embodiment 45, wherein said agonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0247] Embodiment 47. The method of embodiments 41 or 44, further comprising administering to said subject an effective amount of a further therapeutic agent.

10 [0248] Embodiment 48. The method of embodiment 47, wherein said therapeutic agent is an idiopathic pulmonary fibrosis drug or an anti-fibrotic drug.

[0249] Embodiment 49. The method of embodiment 48, wherein said idiopathic pulmonary fibrosis drug is a mucolytic drug.

15 [0250] Embodiment 50. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, the method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB in a fibrotic pulmonary disease patient; (ii) determining whether said expression level is modulated relative to a standard control, wherein a modulated expression level of a fibrotic pulmonary disease marker protein in Table 1A or Table IB relative to said
20 standard control indicates that said fibrotic pulmonary disease patient is at risk for progression of said fibrotic pulmonary disease; and (iii) based at least in part on said expression level in step (ii), determining whether said fibrotic pulmonary disease patient is at risk for progression of said fibrotic pulmonary disease.

25 [0251] Embodiment 51. The method of embodiment 50, wherein said determining whether the expression level is modulated relative to a standard control comprises determining whether the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients.

[0252] Embodiment 52. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, said method comprising: (i)
30 determining a first expression level of a protein set forth in Table 1A or Table IB in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A

or Table IB in said patient at a second time point; (iii) comparing said second expression level of a protein set forth in Table 1A or Table IB to said first expression level of a protein set forth in Table 1A or Table IB, wherein when said second expression level of a protein set forth in Table 1A is greater than said first level of a protein set forth in Table 1A, or wherein when said second expression level of a protein set forth in Table IB is smaller than said first level of a protein set forth in Table IB, the patient is at risk for progression of the fibrotic pulmonary disease.

[0253] Embodiment 53. The method of embodiment 52 further comprising administering a fibrotic pulmonary disease treatment after said determining in step (i).

[0254] Embodiment 54. The method of embodiment 52, further comprising predicting a rate of progression of said fibrotic pulmonary disease in said patient based on said comparing.

[0255] Embodiment 55. The method of embodiment 52, wherein said determining said first expression level of a protein set forth in Table 1A or Table IB and said second expression level of a protein set forth in Table 1A or Table IB comprises normalizing said first expression level of a protein set forth in Table 1A or Table IB and said second expression level of a protein set forth in Table 1A or Table IB to a protein expressed from a standard gene in said patient.

[0256] Embodiment 56. The method of embodiment 55, wherein said first expression level is detected from a first biological sample of said subject and said second expression level is detected from a second biological sample of said subject.

[0257] Embodiment 57. The method of embodiment 56, wherein said first biological sample is a first bodily fluid sample and said second biological sample is a second bodily fluid sample.

[0258] Embodiment 58. A method of determining a fibrotic pulmonary disease activity in a patient, said method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB in a subject; (ii) determining whether said expression level is modulated relative to a standard control, thereby determining a fibrotic pulmonary disease activity in said patient; and (iii) based at least in part on said expression level in step (ii), determining said fibrotic pulmonary disease activity in said patient.

[0259] Embodiment 59. The method of embodiment 58, wherein said expression level of a fibrotic pulmonary disease marker protein in Table 1A is increased relative to said standard control.

[0260] Embodiment 60. The method of embodiment 58, wherein said expression level of a fibrotic pulmonary disease marker protein in Table IB is decreased relative to said standard control.

5 [0261] Embodiment 61. A method of determining a fibrotic pulmonary disease activity in a patient, said method comprising: (i) determining a first expression level of a protein set forth in Table 1A or Table IB in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A or Table IB in said patient at a second time point; (iii) comparing said second expression level of a protein set forth in Table 1A or Table IB to said first expression level of a protein set forth in Table 1A or Table IB, thereby determining said
10 fibrotic pulmonary disease activity in said patient.

[0262] Embodiment 62. The method of embodiment 61, further comprising administering a fibrotic pulmonary disease treatment after said determining in step (i).

[0263] Embodiment 63. A method of determining a change in fibrotic pulmonary disease activity in a patient, said comprising: (i) determining a first expression level of a protein set forth
15 in Table 1A or Table IB in a patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A or Table IB in said patient at a second time point; and (iii) comparing said second expression level of a protein set forth in Table 1A or Table IB to said first expression level of a protein set forth in Table 1A or Table IB, thereby determining said change in fibrotic pulmonary disease activity in the patient.

20 [0264] Embodiment 64. A method of determining a change in fibrotic pulmonary disease activity in a patient, said comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB in a subject; (ii) determining whether said expression level is modulated relative to a standard control, thereby determining a change in fibrotic pulmonary disease activity in said patient; and (iii) based at least
25 in part on said expression level in step (ii), determining said change in fibrotic pulmonary disease activity in said patient.

[0265] Embodiment 65. A method of monitoring the effect of treatment for a fibrotic pulmonary disease in a patient undergoing fibrotic pulmonary disease therapy or a patient that has received fibrotic pulmonary disease therapy comprising: (i) determining a first expression
30 level of a protein set forth in Table 1A or Table IB in the patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A or Table IB in the patient at a second time point; and (iii) comparing the second expression level of a protein set

forth in Table 1A or Table IB to the first expression level of a protein set forth in Table 1A or Table IB, thereby determining the effect of treatment for a fibrotic pulmonary disease in the patient.

[0266] Embodiment 66. A method of monitoring the effect of treatment for a fibrotic pulmonary disease in a patient undergoing fibrotic pulmonary disease therapy or a patient that has received fibrotic pulmonary disease therapy comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table IB in a subject; (ii) determining whether the expression level is modulated relative to a standard control, thereby monitoring the effect of treatment for a fibrotic pulmonary disease in the patient; and (iii) based at least in part on the expression level in step (ii), monitoring the effect of treatment for a fibrotic pulmonary disease in the patient.

[0267] Embodiment 67. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising administering to said subject an effective amount of an modulator of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB, thereby treating a fibrotic pulmonary disease in said subject.

[0268] Embodiment 68. The method of embodiment 67, comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 1A.

[0269] Embodiment 69. The method of embodiment 68, wherein said modulator is an antagonist.

[0270] Embodiment 70. The method of embodiment 69, wherein said antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0271] Embodiment 71. The method of embodiment 67, comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table IB.

[0272] Embodiment 72. The method of embodiment 71, wherein said modulator is an agonist.

[0273] Embodiment 73. The method of embodiment 72, wherein said agonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0274] Embodiment 74. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising: (i) determining whether a subject expresses an elevated level of a fibrotic pulmonary disease marker protein as set forth in Table 1A or a decreased level of a fibrotic pulmonary disease marker protein as set forth in Table 1B relative to a standard control; and (ii) when an elevated expression level of said fibrotic pulmonary disease marker protein of Table 1A or a decreased expression level of said fibrotic pulmonary disease marker protein of Table 1B is found relative to said standard control, administering to said subject a fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker protein set forth in Table 1A or an agonist of a fibrotic pulmonary disease marker protein set forth in Table 1B, thereby treating said subject.

[0275] Embodiment 75. The method of embodiment 74, wherein said fibrotic pulmonary disease treatment is a mucolytic drug.

[0276] Embodiment 76. A method of determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B in a subject that has or is at risk for developing a fibrotic pulmonary disease, said method comprising: (i) obtaining a biological sample from said subject; and (ii) determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B in said biological sample.

[0277] Embodiment 77. A method of determining whether a subject has or is at risk of developing a fibrotic pulmonary disease, said method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject; (ii) determining whether said expression level is increased or decreased relative to a standard control, wherein an elevated expression level of a fibrotic pulmonary disease marker protein in Table 3A or a decreased expression level of a fibrotic pulmonary disease marker protein in Table 3B relative to said standard control indicates that said subject has or is at risk of developing a fibrotic pulmonary disease; and (iii) based at least in part on said expression level in step (ii), determining whether said subject has or is at risk for developing a fibrotic pulmonary disease.

[0278] Embodiment 78. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, the method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker protein set forth in Table 3A or 3B in a fibrotic pulmonary disease patient; (ii) determining whether said expression level is modulated relative to a standard control, wherein a modulated expression

level of a fibrotic pulmonary disease marker protein in Table 3A or 3B relative to said standard control indicates that said fibrotic pulmonary disease patient is at risk for progression of said fibrotic pulmonary disease; and (iii) based at least in part on said expression level in step (ii), determining whether said fibrotic pulmonary disease patient is at risk for progression of said fibrotic pulmonary disease.

[0279] Embodiment 79. The method of embodiment 78, wherein said expression level of a fibrotic pulmonary disease marker protein in Table 3A is increased relative to said standard control.

[0280] Embodiment 80. The method of embodiment 78, wherein said expression level of a fibrotic pulmonary disease marker protein in Table 3B is decreased relative to said standard control.

[0281] Embodiment 81. The method of embodiment 78, wherein said determining whether the expression level is modulated relative to a standard control comprises determining whether the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients.

[0282] Embodiment 82. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, said method comprising: (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in said patient at a second time point; (iii) comparing said second expression level of a protein set forth in Table 3A or Table 3B to said first expression level of a protein set forth in Table 3A or Table 3B, wherein when said second expression level of a protein set forth in Table 3A or Table 3B is greater than said first level of a protein set forth in Table 3A or Table 3B, or wherein when said second expression level of a protein set forth in Table 3A or Table 3B is smaller than said first level of a protein set forth in Table 3A or Table 3B, the patient is at risk for progression of the fibrotic pulmonary disease.

[0283] Embodiment 83. A method of determining a fibrotic pulmonary disease activity in a patient, said method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject; (ii) determining whether said expression level is modulated relative to a standard control, thereby determining a fibrotic pulmonary disease activity in said patient; and (iii) based at least in part on

said expression level in step (ii), determining said fibrotic pulmonary disease activity in said patient.

[0284] Embodiment 84. The method of embodiment 83, wherein said expression level of a fibrotic pulmonary disease marker protein in Table 3A is increased relative to said standard control.

[0285] Embodiment 85. The method of embodiment 83, wherein said expression level of a fibrotic pulmonary disease marker protein in Table 3B is decreased relative to said standard control.

[0286] Embodiment 86. A method of determining a fibrotic pulmonary disease activity in a patient, said method comprising: (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in said patient at a second time point; (iii) comparing said second expression level of a protein set forth in Table 3A or Table 3B to said first expression level of a protein set forth in Table 3A or Table 3B, thereby determining said fibrotic pulmonary disease activity in said patient.

[0287] Embodiment 87. A method of determining a change in fibrotic pulmonary disease activity in a patient, said comprising: (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in a patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in said patient at a second time point; and (iii) comparing said second expression level of a protein set forth in Table 3A or Table 3B to said first expression level of a protein set forth in Table 3A or Table 3B, thereby determining said change in fibrotic pulmonary disease activity in the patient.

[0288] Embodiment 88. A method of determining a change in fibrotic pulmonary disease activity in a patient, said comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject; (ii) determining whether said expression level is modulated relative to a standard control, thereby determining a change in fibrotic pulmonary disease activity in said patient; and (iii) based at least in part on said expression level in step (ii), determining said change in fibrotic pulmonary disease activity in said patient.

[0289] Embodiment 89. A method of monitoring the effect of treatment for a fibrotic pulmonary disease in a patient undergoing fibrotic pulmonary disease therapy or a patient that

has received fibrotic pulmonary disease therapy comprising: (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in the patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in the patient at a second time point; and (iii) comparing the second expression level of a protein set forth in Table 3A or Table 3B to the first expression level of a protein set forth in Table 3A or Table 3B, thereby determining the effect of treatment for a fibrotic pulmonary disease in the patient.

[0290] Embodiment 90. A method of monitoring the effect of treatment for a fibrotic pulmonary disease in a patient undergoing fibrotic pulmonary disease therapy or a patient that has received fibrotic pulmonary disease therapy comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject; (ii) determining whether the expression level is modulated relative to a standard control, thereby monitoring the effect of treatment for a fibrotic pulmonary disease in the patient; and (iii) based at least in part on the expression level in step (ii), monitoring the effect of treatment for a fibrotic pulmonary disease in the patient.

[0291] Embodiment 91. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising administering to said subject an effective amount of an modulator of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B, thereby treating a fibrotic pulmonary disease in said subject.

[0292] Embodiment 92. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising: (i) determining whether a subject expresses an elevated level of a fibrotic pulmonary disease marker protein as set forth in Table 3A or a decreased level of a fibrotic pulmonary disease marker protein as set forth in Table 3B relative to a standard control; and (ii) when an elevated expression level of said fibrotic pulmonary disease marker protein of Table 3A or a decreased expression level of said fibrotic pulmonary disease marker protein of Table 3B is found relative to said standard control, administering to said subject a fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker protein set forth in Table 3A or an agonist of a fibrotic pulmonary disease marker protein set forth in Table 3B, thereby treating said subject.

[0293] Embodiment 93. The method of embodiment 92, wherein said determining whether the expression level is modulated relative to a standard control comprises determining whether

the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients.

[0294] Embodiment 94. A kit comprising: (a) a marker protein binding agent capable of binding to a substance within a biological sample from a human subject having or at risk of developing a fibrotic pulmonary disease; wherein said substance is a fibrotic pulmonary disease marker protein or fragment thereof set forth in Table 1A, IB, 2A, 2B, 3A, 3B, 4A, or 4B; (b) a detecting reagent or a detecting apparatus capable of indicating binding of said marker protein binding agent to said substance.

[0295] Embodiment 95. The kit of embodiment 94, further comprising:

c) a sample collection device for collecting a sample from a subject.

[0296] Embodiment 96. The kit of embodiment 94 or 95, wherein said subject has a fibrotic pulmonary disease.

[0297] Embodiment 97. A complex in vitro comprising a marker protein binding agent bound to a fibrotic pulmonary disease marker protein or fragment thereof set forth in Table 1A, IB, 2A, 2B, 3A, 3B, 4A, or 4B, wherein said fibrotic pulmonary disease marker protein is extracted from a human subject having or at risk of developing a fibrotic pulmonary disease.

[0298] Embodiment 98. The complex of embodiment 97, wherein said subject has a fibrotic pulmonary disease.

[0299] Embodiment 99. A method of determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB in a subject that has or is at risk for developing a fibrotic pulmonary disease, said method comprising: (i) obtaining a biological sample from said subject; and (ii) determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table IB in said biological sample.

[0300] Embodiment 100. The method of embodiment 99, further comprising selecting a subject that has or is at risk for developing a fibrotic pulmonary disease.

[0301] Embodiment 101. The method of embodiment 99, wherein said fibrotic pulmonary disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia.

[0302] Embodiment 102. The method of embodiment 99, wherein said fibrotic pulmonary disease marker protein is a progressive fibrotic pulmonary disease marker protein.

[0303] Embodiment 103. The method of embodiment 102, wherein said subject has or is at risk for developing a progressive fibrotic pulmonary disease.

[0304] Embodiment 104. The method of embodiment 103, wherein said progressive fibrotic pulmonary disease is progressive idiopathic pulmonary fibrosis.

5 [0305] Embodiment 105. The method of embodiment 99, wherein said biological sample is a blood-derived biological sample of said subject.

[0306] Embodiment 106. The method of embodiment 105, wherein said blood-derived biological sample is whole blood, serum or plasma.

10 [0307] Embodiment 107. The method of embodiment 99, wherein said biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0308] Embodiment 108. The method of embodiment 99, wherein an expression level of a fibrotic pulmonary disease marker protein set forth in Table 1A is determined.

15 [0309] Embodiment 109. The method of embodiment 108, further comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 1A.

[0310] Embodiment 110. The method of embodiment 109, wherein said modulator is an antagonist.

[0311] Embodiment 111. The method of embodiment 110, wherein said antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

20 [0312] Embodiment 112. The method of embodiment 110, wherein said expression level of said fibrotic pulmonary disease marker protein set forth in Table 1A is elevated relative to a standard control.

[0313] Embodiment 113. The method of embodiment 99, wherein an expression level of a fibrotic pulmonary disease marker protein set forth in Table IB is determined.

25 [0314] Embodiment 114. The method of embodiment 113, further comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table IB.

[0315] Embodiment 115. The method of embodiment 114, wherein said modulator is an agonist.

[0316] Embodiment 116. The method of embodiment 114, wherein said agonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0317] Embodiment 117. The method of embodiment 115, wherein said expression level of said fibrotic pulmonary disease marker protein set forth in Table IB is decreased relative to a standard control.

[0318] Embodiment 118. The method of embodiment 109 or 114, further comprising administering to said subject an effective amount of a further therapeutic agent.

[0319] Embodiment 119. The method of embodiment 118, wherein said therapeutic agent is an idiopathic pulmonary fibrosis drug.

10 [0320] Embodiment 120. The method of embodiment 118, wherein said idiopathic pulmonary fibrosis drug is a mucolytic drug.

[0321] Embodiment 121. A method of determining whether a subject has or is at risk of developing a fibrotic pulmonary disease, said method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table
15 IB in a subject; (ii) determining whether said expression level is increased or decreased relative to a standard control, wherein an elevated expression level of a fibrotic pulmonary disease marker protein in Table 1A or a decreased expression level of a fibrotic pulmonary disease marker protein in Table IB relative to said standard control indicates that said subject has or is at risk of developing a fibrotic pulmonary disease; and (iii) based at least in part on said expression
20 level in step (ii), determining whether said subject has or is at risk for developing a fibrotic pulmonary disease.

[0322] Embodiment 122. The method of embodiment 121, further comprising selecting a subject that has or is at risk for developing a fibrotic pulmonary disease.

[0323] Embodiment 123. The method of embodiment 121, wherein said fibrotic pulmonary
25 disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia.

[0324] Embodiment 124. The method of embodiment 121, wherein said one or more fibrotic pulmonary disease marker proteins is a progressive fibrotic pulmonary disease marker protein.

[0325] Embodiment 125. The method of embodiment 124, wherein said subject has or is at risk for developing a progressive fibrotic pulmonary disease.

[0326] Embodiment 126. The method of embodiment 125, wherein said progressive fibrotic pulmonary disease is progressive idiopathic pulmonary fibrosis.

[0327] Embodiment 127. The method of embodiment 121, wherein said expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 1A or Table 1B is
5 detected from a biological sample of said subject.

[0328] Embodiment 128. The method of embodiment 127, wherein said biological sample is a blood-derived biological sample of said subject.

[0329] Embodiment 129. The method of embodiment 128, wherein said blood-derived biological sample is whole blood, serum or plasma.

10 [0330] Embodiment 130. The method embodiment 127, wherein said biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0331] Embodiment 131. The method of embodiment 121, further comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 1A.

15 [0332] Embodiment 132. The method of embodiment 131, wherein said modulator is an antagonist.

[0333] Embodiment 133. The method of embodiment 132, wherein said antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0334] Embodiment 134. The method of embodiment 121, further comprising administering
20 to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 1B.

[0335] Embodiment 135. The method of embodiment 134, wherein said modulator is an agonist.

[0336] Embodiment 136. The method of embodiment 135, wherein said agonist is a peptide,
25 small molecule, nucleic acid, antibody or aptamer.

[0337] Embodiment 137. The method of embodiments 131 or 134, further comprising administering to said subject an effective amount of a further therapeutic agent.

[0338] Embodiment 138. The method of embodiment 137, wherein said therapeutic agent is an idiopathic pulmonary fibrosis drug.

[0339] Embodiment 139. The method of embodiment 138, wherein said idiopathic pulmonary fibrosis drug is a mucolytic drug.

[0340] Embodiment 140. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, said method comprising: (i) determining a first expression level of a protein set forth in Table 1A or Table IB in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A or Table IB in said patient at a second time point; (iii) comparing said second expression level of a protein set forth in Table 1A or Table IB to said first expression level of a protein set forth in Table 1A or Table IB, wherein when said second expression level of a protein set forth in Table 1A is greater than said first level of a protein set forth in Table 1A, or wherein when said second expression level of a protein set forth in Table IB is smaller than said first level of a protein set forth in Table IB, the patient is at risk for progression of the fibrotic pulmonary disease.

[0341] Embodiment 141. The method of embodiment 140 further comprising administering a fibrotic pulmonary disease treatment after said determining in step (i).

[0342] Embodiment 142. The method of embodiment 140, further comprising determining a rate of progression of said fibrotic pulmonary disease in said patient based on said comparing.

[0343] Embodiment 143. The method of embodiment 140, wherein said determining said first expression level of a protein set forth in Table 1A or Table IB and said second expression level of a protein set forth in Table 1A or Table IB comprises normalizing said first expression level of a protein set forth in Table 1A or Table IB and said second expression level of a protein set forth in Table 1A or Table IB to a protein expressed from a standard gene in said patient.

[0344] Embodiment 144. The method of embodiment 143, wherein said standard gene is a GAPDH or beta-actin.

[0345] Embodiment 145. The method of embodiment 144, wherein said first expression level is detected from a first biological sample of said subject and said second expression level is detected from a second biological sample of said subject.

[0346] Embodiment 146. The method of embodiment 145, wherein said first biological sample is a first bodily fluid sample and said second biological sample is a second bodily fluid sample.

[0347] Embodiment 147. A method of determining a fibrotic pulmonary disease activity in a patient, said method comprising: (i) determining a first expression level of a protein set forth in

Table 1A or Table 1B in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 1A or Table 1B in said patient at a second time point; (iii) comparing said second expression level of a protein set forth in Table 1A or Table 1B to said first expression level of a protein set forth in Table 1A or Table 1B, thereby determining said fibrotic pulmonary disease activity in said patient.

[0348] Embodiment 148. The method of embodiment 147, further comprising administering a fibrotic pulmonary disease treatment after said determining in step (i).

[0349] Embodiment 149. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising administering to said subject an effective amount of an modulator of a fibrotic pulmonary disease marker protein set forth in Table 1A or Table 1B, thereby treating a fibrotic pulmonary disease in said subject.

[0350] Embodiment 150. The method of embodiment 149, comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 1A.

[0351] Embodiment 151. The method of embodiment 150, wherein said modulator is an antagonist.

[0352] Embodiment 152. The method of embodiment 151, wherein said antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0353] Embodiment 153. The method of embodiment 149, comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 1B.

[0354] Embodiment 154. The method of embodiment 153, wherein said modulator is an agonist.

[0355] Embodiment 155. The method of embodiment 154, wherein said agonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0356] Embodiment 156. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising: (i) determining whether a subject expresses an elevated level of a fibrotic pulmonary disease marker protein as set forth in Table 1A or a decreased level of a fibrotic pulmonary disease marker protein as set forth in Table 1B relative to a standard control; and (ii) when an elevated expression level of said fibrotic pulmonary disease marker

protein of Table 1A or a decreased expression level of said fibrotic pulmonary disease marker protein of Table 1B is found relative to said standard control, administering to said subject a fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker protein set forth in Table 1A or an agonist of a fibrotic pulmonary disease marker protein set forth in Table 1B, thereby treating said subject.

[0357] Embodiment 157. The method of embodiment 156, wherein said fibrotic pulmonary disease treatment is a mucolytic drug.

[0358] Embodiment 158. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, the method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker protein set forth in Table 1A or 1B in a fibrotic pulmonary disease patient; (ii) determining whether said expression level is modulated relative to a standard control, wherein a modulated expression level of a fibrotic pulmonary disease marker protein in Table 1A or 1B relative to said standard control indicates that said fibrotic pulmonary disease patient is at risk for progression of said fibrotic pulmonary disease; and (iii) based at least in part on said expression level in step (ii), determining whether said fibrotic pulmonary disease patient is at risk for progression of said fibrotic pulmonary disease.

[0359] Embodiment 159. The method of embodiment 158, wherein said determining whether the expression level is modulated relative to a standard control comprises determining whether the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients.

[0360] Embodiment 160. A method of determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B in a subject that has or is at risk for developing a fibrotic pulmonary disease, said method comprising: (i) obtaining a biological sample from said subject; and (ii) determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B in said biological sample.

[0361] Embodiment 161. A method of determining whether a subject has or is at risk of developing a fibrotic pulmonary disease, said method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject; (ii) determining whether said expression level is increased or decreased relative to a standard control, wherein an elevated expression level of a fibrotic pulmonary disease marker protein in Table 3A or a decreased expression level of a fibrotic pulmonary disease

marker protein in Table 3B relative to said standard control indicates that said subject has or is at risk of developing a fibrotic pulmonary disease; and (iii) based at least in part on said expression level in step (ii), determining whether said subject has or is at risk for developing a fibrotic pulmonary disease.

5 **[0362]** Embodiment 162. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, said method comprising: (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in said patient at a second time point; (iii) comparing said second expression level of
10 a protein set forth in Table 3A or Table 3B to said first expression level of a protein set forth in Table 3A or Table 3B, wherein when said second expression level of a protein set forth in Table 3A is greater than said first level of a protein set forth in Table 3A, or wherein when said second expression level of a protein set forth in Table 1B is smaller than said first level of a protein set forth in Table 1B, the patient is at risk for progression of the fibrotic pulmonary disease.

15 **[0363]** Embodiment 163. A method of determining a fibrotic pulmonary disease activity in a patient, said method comprising: (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in said patient at a second time point; (iii)
20 comparing said second expression level of a protein set forth in Table 3A or Table 3B to said first expression level of a protein set forth in Table 3A or Table 3B, thereby determining said fibrotic pulmonary disease activity in said patient.

[0364] Embodiment 164. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising administering to said subject an effective amount of an modulator of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B,
25 thereby treating a fibrotic pulmonary disease in said subject.

[0365] Embodiment 165. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising: (i) determining whether a subject expresses an elevated level of a fibrotic pulmonary disease marker protein as set forth in Table 3A or a decreased level of a fibrotic pulmonary disease marker protein as set forth in Table 3B relative to a standard
30 control; and (ii) when an elevated expression level of said fibrotic pulmonary disease marker protein of Table 3A or a decreased expression level of said fibrotic pulmonary disease marker protein of Table 3B is found relative to said standard control, administering to said subject a

fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker protein set forth in Table 3A or an agonist of a fibrotic pulmonary disease marker protein set forth in Table 3B, thereby treating said subject.

[0366] Embodiment 166. The method of embodiment 165, wherein said determining whether the expression level is modulated relative to a standard control comprises determining whether the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients.

[0367] Embodiment 167. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, the method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker protein set forth in Table 3A or 3B in a fibrotic pulmonary disease patient; (ii) determining whether the expression level is modulated relative to a standard control, wherein a modulated expression level of a fibrotic pulmonary disease marker protein in Table 3A or 3B relative to said standard control indicates that said fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease; and (iii) based at least in part on the expression level in step (ii), determining whether said fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease.

[0368] Embodiment 168. The method of embodiment 167, wherein said determining whether the expression level is modulated relative to a standard control comprises determining whether the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients.

[0369] Embodiment 169. A method of determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 2A or Table 2B in a subject that has or is at risk for developing a fibrotic pulmonary disease, said method comprising: (i) obtaining a biological sample from said subject; and (ii) determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 2A or Table 2B in said biological sample.

[0370] Embodiment 170. The method of embodiment 169, further comprising selecting a subject that has or is at risk for developing a fibrotic pulmonary disease.

[0371] Embodiment 171. The method of embodiment 169, wherein said fibrotic pulmonary disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia.

[0372] Embodiment 172. The method of embodiment 169, wherein said fibrotic pulmonary disease marker protein is a progressive fibrotic pulmonary disease marker protein.

[0373] Embodiment 173. The method of embodiment 172, wherein said subject has or is at risk for developing a progressive fibrotic pulmonary disease.

5 [0374] Embodiment 174. The method of embodiment 173, wherein said progressive fibrotic pulmonary disease is progressive idiopathic pulmonary fibrosis.

[0375] Embodiment 175. The method of embodiment 169, wherein said biological sample is a blood-derived biological sample of said subject.

10 [0376] Embodiment 176. The method of embodiment 175, wherein said blood-derived biological sample is whole blood, serum or plasma.

[0377] Embodiment 177. The method of embodiment 169, wherein said biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0378] Embodiment 178. The method of embodiment 169, wherein an expression level of a fibrotic pulmonary disease marker protein set forth in Table 2A is determined.

15 [0379] Embodiment 179. The method of embodiment 178, further comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 2A.

[0380] Embodiment 180. The method of embodiment 179, wherein said modulator is an antagonist.

20 [0381] Embodiment 181. The method of embodiment 180, wherein said antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0382] Embodiment 182. The method of embodiment 180, wherein said expression level of said fibrotic pulmonary disease marker protein set forth in Table 2A is elevated relative to a standard control.

25 [0383] Embodiment 183. The method of embodiment 169, wherein an expression level of a fibrotic pulmonary disease marker protein set forth in Table 2B is determined.

[0384] Embodiment 184. The method of embodiment 183, further comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 2B.

[0385] Embodiment 185. The method of embodiment 184, wherein said modulator is an agonist.

[0386] Embodiment 186. The method of embodiment 184, wherein said agonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

5 [0387] Embodiment 187. The method of embodiment 185, wherein said expression level of said fibrotic pulmonary disease marker protein set forth in Table 2B is decreased relative to a standard control.

[0388] Embodiment 188. The method of embodiment 179 or 184, further comprising administering to said subject an effective amount of a further therapeutic agent.

10 [0389] Embodiment 189. The method of embodiment 188, wherein said therapeutic agent is an idiopathic pulmonary fibrosis drug.

[0390] Embodiment 190. The method of embodiment 188, wherein said idiopathic pulmonary fibrosis drug is a mucolytic drug.

[0391] Embodiment 191. A method of determining whether a subject has or is at risk of
15 developing a fibrotic pulmonary disease, said method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 2A or Table 2B in a subject; (ii) determining whether said expression level is increased or decreased relative to a standard control, wherein an elevated expression level of a fibrotic pulmonary disease marker protein in Table 2A or a decreased expression level of a fibrotic pulmonary disease
20 marker protein in Table 2B relative to said standard control indicates that said subject has or is at risk of developing a fibrotic pulmonary disease; and (iii) based at least in part on said expression level in step (ii), determining whether said subject has or is at risk for developing a fibrotic pulmonary disease.

[0392] Embodiment 192. The method of embodiment 191, further comprising selecting a
25 subject that has or is at risk for developing a fibrotic pulmonary disease.

[0393] Embodiment 193. The method of embodiment 191, wherein said fibrotic pulmonary disease is idiopathic pulmonary fibrosis or familial interstitial pneumonia.

[0394] Embodiment 194. The method of embodiment 191, wherein said one or more fibrotic pulmonary disease marker proteins is a progressive fibrotic pulmonary disease marker protein.

[0395] Embodiment 195. The method of embodiment 194, wherein said subject has or is at risk for developing a progressive fibrotic pulmonary disease.

[0396] Embodiment 196. The method of embodiment 195, wherein said progressive fibrotic pulmonary disease is progressive idiopathic pulmonary fibrosis.

5 [0397] Embodiment 197. The method of embodiment 191, wherein said expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 2A or Table 2B is detected from a biological sample of said subject.

[0398] Embodiment 198. The method of embodiment 197, wherein said biological sample is a blood-derived biological sample of said subject.

10 [0399] Embodiment 199. The method of embodiment 198, wherein said blood-derived biological sample is whole blood, serum or plasma.

[0400] Embodiment 200. The method embodiment 197, wherein said biological sample is urine, saliva, a pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

[0401] Embodiment 201. The method of embodiment 191, further comprising administering
15 to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 2A.

[0402] Embodiment 202. The method of embodiment 201, wherein said modulator is an antagonist.

[0403] Embodiment 203. The method of embodiment 202, wherein said antagonist is a
20 peptide, small molecule, nucleic acid, antibody or aptamer.

[0404] Embodiment 204. The method of embodiment 191, further comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 2B.

[0405] Embodiment 205. The method of embodiment 204, wherein said modulator is an
25 agonist.

[0406] Embodiment 206. The method of embodiment 205, wherein said agonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0407] Embodiment 207. The method of embodiments 201 or 204, further comprising administering to said subject an effective amount of a further therapeutic agent.

[0408] Embodiment 208. The method of embodiment 207, wherein said therapeutic agent is an idiopathic pulmonary fibrosis drug.

[0409] Embodiment 209. The method of embodiment 208, wherein said idiopathic pulmonary fibrosis drug is a mucolytic drug.

5 [0410] Embodiment 210. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, said method comprising: (i) determining a first expression level of a protein set forth in Table 2A or Table 2B in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 2A or Table 2B in said patient at a second time point; (iii) comparing said second expression level of
10 a protein set forth in Table 2A or Table 2B to said first expression level of a protein set forth in Table 2A or Table 2B, wherein when said second expression level of a protein set forth in Table 2A is greater than said first level of a protein set forth in Table 2A, or wherein when said second expression level of a protein set forth in Table 2B is smaller than said first level of a protein set forth in Table 2B, the patient is at risk for progression of the fibrotic pulmonary disease.

15 [0411] Embodiment 211. The method of embodiment 210 further comprising administering a fibrotic pulmonary disease treatment after said determining in step (i).

[0412] Embodiment 212. The method of embodiment 210, further comprising determining a rate of progression of said fibrotic pulmonary disease in said patient based on said comparing.

[0413] Embodiment 213. The method of embodiment 210, wherein said determining said
20 first expression level of a protein set forth in Table 2A or Table 2B and said second expression level of a protein set forth in Table 2A or Table 2B comprises normalizing said first expression level of a protein set forth in Table 2A or Table 2B and said second expression level of a protein set forth in Table 2A or Table 2B to a protein expressed from a standard gene in said patient.

[0414] Embodiment 214. The method of embodiment 213, wherein said standard gene is a
25 GAPDH or beta-actin.

[0415] Embodiment 215. The method of embodiment 214, wherein said first expression level is detected from a first biological sample of said subject and said second expression level is detected from a second biological sample of said subject.

[0416] Embodiment 216. The method of embodiment 215, wherein said first biological
30 sample is a first bodily fluid sample and said second biological sample is a second bodily fluid sample.

[0417] Embodiment 217. A method of determining a fibrotic pulmonary disease activity in a patient, said method comprising: (i) determining a first expression level of a protein set forth in Table 2A or Table 2B in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 2A or Table 2B in said patient at a second time point; (iii) comparing said second expression level of a protein set forth in Table 2A or Table 2B to said first expression level of a protein set forth in Table 2A or Table 2B, thereby determining said fibrotic pulmonary disease activity in said patient.

[0418] Embodiment 218. The method of embodiment 217, further comprising administering a fibrotic pulmonary disease treatment after said determining in step (i).

[0419] Embodiment 219. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising administering to said subject an effective amount of an modulator of a fibrotic pulmonary disease marker protein set forth in Table 2A or Table 2B, thereby treating a fibrotic pulmonary disease in said subject.

[0420] Embodiment 220. The method of embodiment 219, comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 2A.

[0421] Embodiment 221. The method of embodiment 220, wherein said modulator is an antagonist.

[0422] Embodiment 222. The method of embodiment 221, wherein said antagonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0423] Embodiment 223. The method of embodiment 219, comprising administering to said subject an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in Table 2B.

[0424] Embodiment 224. The method of embodiment 223, wherein said modulator is an agonist.

[0425] Embodiment 225. The method of embodiment 224, wherein said agonist is a peptide, small molecule, nucleic acid, antibody or aptamer.

[0426] Embodiment 226. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising: (i) determining whether a subject expresses an elevated level of a fibrotic pulmonary disease marker protein as set forth in Table 2A or a decreased level

of a fibrotic pulmonary disease marker protein as set forth in Table 2B relative to a standard control; and (ii) when an elevated expression level of said fibrotic pulmonary disease marker protein of Table 2A or a decreased expression level of said fibrotic pulmonary disease marker protein of Table 2B is found relative to said standard control, administering to said subject a fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker protein set forth in Table 2A or an agonist of a fibrotic pulmonary disease marker protein set forth in Table 2B, thereby treating said subject.

[0427] Embodiment 227. The method of embodiment 226, wherein said fibrotic pulmonary disease treatment is a mucolytic drug.

[0428] Embodiment 228. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, the method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker protein set forth in Table 2A or 2B in a fibrotic pulmonary disease patient; (ii) determining whether said expression level is modulated relative to a standard control, wherein a modulated expression level of a fibrotic pulmonary disease marker protein in Table 2A or 2B relative to said standard control indicates that said fibrotic pulmonary disease patient is at risk for progression of said fibrotic pulmonary disease; and (iii) based at least in part on said expression level in step (ii), determining whether said fibrotic pulmonary disease patient is at risk for progression of said fibrotic pulmonary disease.

[0429] Embodiment 229. The method of embodiment 228, wherein said determining whether the expression level is modulated relative to a standard control comprises determining whether the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients.

[0430] Embodiment 230. A method of determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B in a subject that has or is at risk for developing a fibrotic pulmonary disease, said method comprising: (i) obtaining a biological sample from said subject; and (ii) determining an expression level of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B in said biological sample.

[0431] Embodiment 231. A method of determining whether a subject has or is at risk of developing a fibrotic pulmonary disease, said method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker proteins set forth in Table 3A or Table 3B in a subject; (ii) determining whether said expression level is increased or decreased relative

to a standard control, wherein an elevated expression level of a fibrotic pulmonary disease marker protein in Table 3A or a decreased expression level of a fibrotic pulmonary disease marker protein in Table 3B relative to said standard control indicates that said subject has or is at risk of developing a fibrotic pulmonary disease; and (iii) based at least in part on said expression level in step (ii), determining whether said subject has or is at risk for developing a fibrotic pulmonary disease.

[0432] Embodiment 232. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, said method comprising: (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in said patient at a second time point; (iii) comparing said second expression level of a protein set forth in Table 3A or Table 3B to said first expression level of a protein set forth in Table 3A or Table 3B, wherein when said second expression level of a protein set forth in Table 3A is greater than said first level of a protein set forth in Table 3A, or wherein when said second expression level of a protein set forth in Table 3B is smaller than said first level of a protein set forth in Table 3B, the patient is at risk for progression of the fibrotic pulmonary disease.

[0433] Embodiment 233. A method of determining a fibrotic pulmonary disease activity in a patient, said method comprising: (i) determining a first expression level of a protein set forth in Table 3A or Table 3B in said patient at a first time point; (ii) determining a second expression level of a protein set forth in Table 3A or Table 3B in said patient at a second time point; (iii) comparing said second expression level of a protein set forth in Table 3A or Table 3B to said first expression level of a protein set forth in Table 3A or Table 3B, thereby determining said fibrotic pulmonary disease activity in said patient.

[0434] Embodiment 234. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising administering to said subject an effective amount of an modulator of a fibrotic pulmonary disease marker protein set forth in Table 3A or Table 3B, thereby treating a fibrotic pulmonary disease in said subject.

[0435] Embodiment 235. A method of treating a fibrotic pulmonary disease in a subject in need thereof, said method comprising: (i) determining whether a subject expresses an elevated level of a fibrotic pulmonary disease marker protein as set forth in Table 3A or a decreased level of a fibrotic pulmonary disease marker protein as set forth in Table 3B relative to a standard control; and (ii) when an elevated expression level of said fibrotic pulmonary disease marker

protein of Table 3A or a decreased expression level of said fibrotic pulmonary disease marker protein of Table 3B is found relative to said standard control, administering to said subject a fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker protein set forth in Table 3A or an agonist of a fibrotic pulmonary disease marker protein set forth in Table 3B, thereby treating said subject.

[0436] Embodiment 236. The method of embodiment 235, wherein said determining whether the expression level is modulated relative to a standard control comprises determining whether the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients.

[0437] Embodiment 237. A method of determining whether a fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease, the method comprising: (i) detecting an expression level of one or more fibrotic pulmonary disease marker protein set forth in Table 3A or 3B in a fibrotic pulmonary disease patient; (ii) determining whether the expression level is modulated relative to a standard control, wherein a modulated expression level of a fibrotic pulmonary disease marker protein in Table 3A or 3B relative to said standard control indicates that said fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease; and (iii) based at least in part on the expression level in step (ii), determining whether said fibrotic pulmonary disease patient is at risk for progression of the fibrotic pulmonary disease.

[0438] Embodiment 238. The method of embodiment 237, wherein said determining whether the expression level is modulated relative to a standard control comprises determining whether the expression level is elevated or suppressed relative to other fibrotic pulmonary disease patients.

WHAT IS CLAIMED IS:

1 1. A method of determining an expression level of a fibrotic pulmonary
2 disease marker protein set forth in Table 2A or Table 2B in a subject that has or is at risk for
3 developing a fibrotic pulmonary disease, said method comprising:

4 (i) obtaining a biological sample from said subject; and

5 (ii) determining an expression level of a fibrotic pulmonary disease marker
6 protein set forth in Table 2A or Table 2B in said biological sample.

1 2. The method of claim 1, further comprising selecting a subject that has or
2 is at risk for developing a fibrotic pulmonary disease.

1 3. The method of claim 1, wherein said fibrotic pulmonary disease is
2 idiopathic pulmonary fibrosis or familial interstitial pneumonia.

1 4. The method of claim 1, wherein said fibrotic pulmonary disease marker
2 protein is a progressive fibrotic pulmonary disease marker protein.

1 5. The method of claim 4, wherein said subject has or is at risk for
2 developing a progressive fibrotic pulmonary disease.

1 6. The method of claim 5, wherein said progressive fibrotic pulmonary
2 disease is progressive idiopathic pulmonary fibrosis.

1 7. The method of claim 1, wherein said biological sample is a blood-derived
2 biological sample of said subject.

1 8. The method of claim 7, wherein said blood-derived biological sample is
2 whole blood, serum or plasma.

1 9. The method of claim 1, wherein said biological sample is urine, saliva, a
2 pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

1 10. The method of claim 1, wherein an expression level of a fibrotic
2 pulmonary disease marker protein set forth in Table 2A is determined.

1 11. The method of claim 10, further comprising administering to said subject
2 an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in
3 Table 2A.

1 12. The method of claim 11, wherein said modulator is an antagonist.

1 13. The method of claim 12, wherein said antagonist is a peptide, small
2 molecule, nucleic acid, antibody or aptamer.

1 14. The method of claim 12, wherein said expression level of said fibrotic
2 pulmonary disease marker protein set forth in Table 2A is elevated relative to a standard control.

1 15. The method of claim 1, wherein an expression level of a fibrotic
2 pulmonary disease marker protein set forth in Table 2B is determined.

1 16. The method of claim 15, further comprising administering to said subject
2 an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in
3 Table 2B.

1 17. The method of claim 16, wherein said modulator is an agonist.

1 18. The method of claim 16, wherein said agonist is a peptide, small molecule,
2 nucleic acid, antibody or aptamer.

1 19. The method of claim 17, wherein said expression level of said fibrotic
2 pulmonary disease marker protein set forth in Table 2B is decreased relative to a standard
3 control.

1 20. The method of claim 11 or 16, further comprising administering to said
2 subject an effective amount of a further therapeutic agent.

1 21. The method of claim 20, wherein said therapeutic agent is an idiopathic
2 pulmonary fibrosis drug.

1 22. The method of claim 20, wherein said idiopathic pulmonary fibrosis drug
2 is a mucolytic drug.

1 23. A method of determining whether a subject has or is at risk of developing
2 a fibrotic pulmonary disease, said method comprising:

3 (i) detecting an expression level of one or more fibrotic pulmonary disease marker
4 proteins set forth in Table 2A or Table 2B in a subject;

5 (ii) determining whether said expression level is increased or decreased relative to
6 a standard control, wherein an elevated expression level of a fibrotic pulmonary disease marker
7 protein in Table 2A or a decreased expression level of a fibrotic pulmonary disease marker
8 protein in Table 2B relative to said standard control indicates that said subject has or is at risk of
9 developing a fibrotic pulmonary disease; and

10 (iii) based at least in part on said expression level in step (ii), determining whether
11 said subject has or is at risk for developing a fibrotic pulmonary disease.

1 24. The method of claim 23, further comprising selecting a subject that has or
2 is at risk for developing a fibrotic pulmonary disease.

1 25. The method of claim 23, wherein said fibrotic pulmonary disease is
2 idiopathic pulmonary fibrosis or familial interstitial pneumonia.

1 26. The method of claim 23, wherein said one or more fibrotic pulmonary
2 disease marker proteins is a progressive fibrotic pulmonary disease marker protein.

1 27. The method of claim 26, wherein said subject has or is at risk for
2 developing a progressive fibrotic pulmonary disease.

1 28. The method of claim 27, wherein said progressive fibrotic pulmonary
2 disease is progressive idiopathic pulmonary fibrosis.

1 29. The method of claim 23, wherein said expression level of one or more
2 fibrotic pulmonary disease marker proteins set forth in Table 2A or Table 2B is detected from a
3 biological sample of said subject.

1 30. The method of claim 29, wherein said biological sample is a blood-
2 derived biological sample of said subject.

1 31. The method of claim 30, wherein said blood-derived biological sample is
2 whole blood, serum or plasma.

1 32. The method claim 29, wherein said biological sample is urine, saliva, a
2 pulmonary tissue, bronchoalveolar lavage sample, or exhaled breath condensate.

1 33. The method of claim 23, further comprising administering to said subject
2 an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in
3 Table 2A.

1 34. The method of claim 33, wherein said modulator is an antagonist.

1 35. The method of claim 34, wherein said antagonist is a peptide, small
2 molecule, nucleic acid, antibody or aptamer.

1 36. The method of claim 23, further comprising administering to said subject
2 an effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in
3 Table 2B.

1 37. The method of claim 36, wherein said modulator is an agonist.

1 38. The method of claim 37, wherein said agonist is a peptide, small molecule,
2 nucleic acid, antibody or aptamer.

1 39. The method of claims 33 or 36, further comprising administering to said
2 subject an effective amount of a further therapeutic agent.

1 40. The method of claim 39, wherein said therapeutic agent is an idiopathic
2 pulmonary fibrosis drug.

1 41. The method of claim 40, wherein said idiopathic pulmonary fibrosis drug
2 is a mucolytic drug.

1 42. A method of determining whether a fibrotic pulmonary disease patient is
2 at risk for progression of the fibrotic pulmonary disease, said method comprising:

3 (i) determining a first expression level of a protein set forth in Table 2A or Table
4 2B in said patient at a first time point;

5 (ii) determining a second expression level of a protein set forth in Table 2A or
6 Table 2B in said patient at a second time point;

(iii) comparing said second expression level of a protein set forth in Table 2A or Table 2B to said first expression level of a protein set forth in Table 2A or Table 2B, wherein when said second expression level of a protein set forth in Table 2A is greater than said first level of a protein set forth in Table 2A, or wherein when said second expression level of a protein set forth in Table 2B is smaller than said first level of a protein set forth in Table 2B, the patient is at risk for progression of the fibrotic pulmonary disease.

43. The method of claim 42 further comprising administering a fibrotic pulmonary disease treatment after said determining in step (i).

44. The method of claim 42, further comprising determining a rate of progression of said fibrotic pulmonary disease in said patient based on said comparing.

45. The method of claim 42, wherein said determining said first expression level of a protein set forth in Table 2A or Table 2B and said second expression level of a protein set forth in Table 2A or Table 2B comprises normalizing said first expression level of a protein set forth in Table 2A or Table 2B and said second expression level of a protein set forth in Table 2A or Table 2B to a protein expressed from a standard gene in said patient.

46. The method of claim 45, wherein said standard gene is a GAPDH or beta-actin.

47. The method of claim 46, wherein said first expression level is detected from a first biological sample of said subject and said second expression level is detected from a second biological sample of said subject.

48. The method of claim 47, wherein said first biological sample is a first bodily fluid sample and said second biological sample is a second bodily fluid sample.

49. A method of determining a fibrotic pulmonary disease activity in a patient, said method comprising:

(i) determining a first expression level of a protein set forth in Table 2A or Table 2B in said patient at a first time point;

(ii) determining a second expression level of a protein set forth in Table 2A or Table 2B in said patient at a second time point;

7 (iii) comparing said second expression level of a protein set forth in Table 2A or
8 Table 2B to said first expression level of a protein set forth in Table 2A or Table 2B, thereby
9 determining said fibrotic pulmonary disease activity in said patient.

1 50. The method of claim 49, further comprising administering a fibrotic
2 pulmonary disease treatment after said determining in step (i).

1 51. A method of treating a fibrotic pulmonary disease in a subject in need
2 thereof, said method comprising administering to said subject an effective amount of an
3 modulator of a fibrotic pulmonary disease marker protein set forth in Table 2A or Table 2B,
4 thereby treating a fibrotic pulmonary disease in said subject.

1 52. The method of claim 51, comprising administering to said subject an
2 effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in
3 Table 2A.

1 53. The method of claim 52, wherein said modulator is an antagonist.

1 54. The method of claim 53, wherein said antagonist is a peptide, small
2 molecule, nucleic acid, antibody or aptamer.

1 55. The method of claim 51, comprising administering to said subject an
2 effective amount of a modulator of said fibrotic pulmonary disease marker protein set forth in
3 Table 2B.

1 56. The method of claim 55, wherein said modulator is an agonist.

1 57. The method of claim 56, wherein said agonist is a peptide, small molecule,
2 nucleic acid, antibody or aptamer.

1 58. A method of treating a fibrotic pulmonary disease in a subject in need
2 thereof, said method comprising:

- 3 (i) determining whether a subject expresses an elevated level of a fibrotic
4 pulmonary disease marker protein as set forth in Table 2A or a decreased level of a fibrotic
5 pulmonary disease marker protein as set forth in Table 2B relative to a standard control; and
6 (ii) when an elevated expression level of said fibrotic pulmonary disease marker
7 protein of Table 2A or a decreased expression level of said fibrotic pulmonary disease marker

8 protein of Table 2B is found relative to said standard control, administering to said subject a
9 fibrotic pulmonary disease treatment, an antagonist of a fibrotic pulmonary disease marker
10 protein set forth in Table 2A or an agonist of a fibrotic pulmonary disease marker protein set
11 forth in Table 2B, thereby treating said subject.

1 59. The method of claim 58, wherein said fibrotic pulmonary disease
2 treatment is a mucolytic drug.

1 60. A method of determining whether a fibrotic pulmonary disease patient is
2 at risk for progression of the fibrotic pulmonary disease, the method comprising:

3 (i) detecting an expression level of one or more fibrotic pulmonary disease marker
4 protein set forth in Table 2A or 2B in a fibrotic pulmonary disease patient;

5 (ii) determining whether said expression level is modulated relative to a standard
6 control,

7 wherein a modulated expression level of a fibrotic pulmonary disease marker
8 protein in Table 2A or 2B relative to said standard control indicates that said fibrotic pulmonary
9 disease patient is at risk for progression of said fibrotic pulmonary disease; and (iii) based at least
10 in part on said expression level in step (ii), determining whether said fibrotic pulmonary disease
11 patient is at risk for progression of said fibrotic pulmonary disease.

1 61. The method of claim 60, wherein said determining whether the expression
2 level is modulated relative to a standard control comprises determining whether the expression
3 level is elevated or suppressed relative to other fibrotic pulmonary disease patients.

1

FIG. 1




FIG. 2




FIG. 3





FIG.4



*mRNAs encoding the proteins were upregulated in IPF Lung

FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37




FIG. 38




FIG. 39




FIG. 40




FIG. 41




FIG. 42




FIG. 43




FIG. 44




FIG. 45




FIG. 46




FIG. 47




FIG. 48




FIG. 49




FIG. 50




FIG. 51




FIG. 52




FIG. 53




FIG. 54




FIG. 55




FIG. 56





FIG. 57



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2014/029392**A. CLASSIFICATION OF SUBJECT MATTER****G01N 33/68(2006.01)i, G01N 33/53(2006.01)i, C12Q 1/68(2006.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

G01N 33/68; C40B 30/06; A61K 39/395; C40B 40/06; G01N 33/53; C12Q 1/06; C12Q 1/68; C40B 30/04

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models

Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: fibrotic pulmonary disease, marker, idiopathic pulmonary fibrosis(IPF), biological sample

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2012-0035067 AI (KAMINSKI et al.) 09 February 2012 See abstract, claims 1 and 2 and paragraph [0108].	1-10, 49
A		15
A	MA et al., 'Quantitation of desmosine and isodesmosine in urine, plasma, and sputum by LC-MS/MS as biomarkers for elastin degradation' Journal of Chromatography B, Vol.879, No.21, pp.1893-1898 (2011) See abstract and page 1893.	1-10, 15, 49
A	US 2011-0250589 AI (LAMA) 13 October 2011 See abstract and claim 1.	1-10, 15, 49
A	US 2012-0282276 AI (HOGABOAM et al.) 08 November 2012 See abstract and claim 1.	1-10, 15, 49
A	US 2011-0280865 AI (MARSH et al.) 17 November 2011 See abstract and claim 1.	1-10, 15, 49

I Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 July 2014 (10.07.2014)

Date of mailing of the international search report

17 July 2014 (17.07.2014)

Name and mailing address of the ISA/KR



International Application Division
Korean Intellectual Property Office
 189 Cheongsa-ro, Seo-gu, Daejeon Metropolitan City, 302-701,
 Republic of Korea

Facsimile No. +82-42-472-7140

Authorized officer

KIM, Seung Beom

Telephone No. +82-42-481-3371



Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 11-14, 16-48, 50-61
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 11-14, 16-48 and 50-61 pertain to methods for treatment of the human body by therapy, as well as diagnostic methods, and thus relate to a subject matter which this International Searching Authority is not required, under PCT Article 17(2)(a)(i) and PCT Rule 39.1(iv), to search.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically :
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of any additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos. :
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2014/029392

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2012-0035067 A1	09/02/2012	WO 2010-028274 A1	11/03/2010
US 2011-0250589 A1	13/10/2011	US 8637232 B2	28/01/2014
		WO 2011-091270 A2	28/07/2011
		WO 2011-091270 A3	08/12/2011
US 2012-0282276 A1	08/11/2012	CN 102640001 A	15/08/2012
		EP 2496944 A2	12/09/2012
		JP 2013-509869 A	21/03/2013
		WO 2011-054893 A2	12/05/2011
		WO 2011-054893 A3	21/07/2011
US 2011-0280865 A1	17/11/2011	CA 2749821 A1	22/07/2010
		EP 2387716 A2	23/11/2011
		EP 2387716 A4	17/10/2012
		WO 2010-083392 A2	22/07/2010
		WO 2010-083392 A3	07/10/2010

专利名称(译)	蛋白质组学ipf标记		
公开(公告)号	EP2972392A1	公开(公告)日	2016-01-20
申请号	EP2014762377	申请日	2014-03-14
申请(专利权)人(译)	InterMune公司 , INC.		
当前申请(专利权)人(译)	InterMune公司 , INC.		
[标]发明人	KOSSEN KARL QIN XIAOLI LIM SHARLEEN R SEIWERT SCOTT D RUHRMUND DONALD		
发明人	KOSSEN, KARL QIN, XIAOLI LIM, SHARLEEN, R. SEIWERT, SCOTT, D. RUHRMUND, DONALD		
IPC分类号	G01N33/68 G01N33/53 C12Q1/68		
CPC分类号	A61P11/00 C12Q1/6883 C12Q2600/118 C12Q2600/158 G01N33/6854 G01N33/6893 G01N2800/12		
代理机构(译)	HGF LIMITED		
优先权	61/801290 2013-03-15 US 61/801476 2013-03-15 US 61/874947 2013-09-06 US		
其他公开文献	EP2972392A4		
外部链接	Espacenet		

摘要(译)

本文提供了用于鉴定，预测或监测纤维化肺病（例如特发性肺纤维化（IPF））的新型生物标志物。本文提供的生物标志物可进一步用于IPF的治疗性治疗。