



(11) **EP 2 748 337 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
07.08.2019 Bulletin 2019/32

(51) Int Cl.:
C12Q 1/68 (2018.01) G01N 33/53 (2006.01)
G01N 33/574 (2006.01)

(21) Application number: **12833913.2**

(86) International application number:
PCT/US2012/056668

(22) Date of filing: **21.09.2012**

(87) International publication number:
WO 2013/044099 (28.03.2013 Gazette 2013/13)

(54) **CELL RESPONSE ASSAY FOR CANCER AND METHODS OF PRODUCING AND USING SAME**

ZELLREAKTIONSTEST FÜR KREBS SOWIE VERFAHREN ZU SEINER HERSTELLUNG UND ANWENDUNG

DOSAGE DE SENSIBILITÉ CELLULAIRE POUR UN CANCER ET PROCÉDÉS DE PRODUCTION ET D'UTILISATION ASSOCIÉS

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

US-A1- 2003 129 665 US-A1- 2008 038 738
US-A1- 2008 038 738

(30) Priority: **23.09.2011 US 201161538302 P**

(43) Date of publication of application:
02.07.2014 Bulletin 2014/27

(73) Proprietor: **Siemens Healthcare Diagnostics Inc. Tarrytown, NY 10591 (US)**

(72) Inventor: **PUGIA, Michael Granger, IN 46530 (US)**

(74) Representative: **Schweitzer, Klaus et al Plate Schweitzer Zounek Patentanwälte Rheingaustrasse 196 65203 Wiesbaden (DE)**

(56) References cited:
WO-A1-2009/124997 WO-A1-2009/124997
WO-A1-2011/109440 WO-A1-2011/109440
WO-A2-2006/060561 US-A1- 2003 129 665

- **MARILEILA VARELLA-GARCIA ET AL: "The UroVysion fluorescence in situ hybridization assay is an effective tool for monitoring recurrence of bladder cancer", UROLOGIC ONCOLOGY: SEMINARS AND ORIGINAL INVESTIGATIONS, vol. 22, no. 1, 1 January 2004 (2004-01-01), pages 16-19, XP055178124, ISSN: 1078-1439, DOI: 10.1016/S1078-1439(03)00098-X**
- **WYFFELS: 'Synthesis and preliminary evaluation of radiolabeled bis(zinc(II)-dipicolylamine) coordination complexes as cell death imaging agents' BIOORG. MED. CHEM. vol. 19, 01 June 2011, pages 3425 - 3433, XP028216865**
- **Veronique Neumeister ET AL: "In Situ Identification of Putative Cancer Stem Cells by Multiplexing ALDH1, CD44, and Cytokeratin Identifies Breast Cancer Patients with Poor Prognosis", AMERICAN JOURNAL OF PATHOLOGY., vol. 176, no. 5, 1 May 2010 (2010-05-01), pages 2131-2138, XP055388459, US ISSN: 0002-9440, DOI: 10.2353/ajpath.2010.090712**
- **Yanjie Hu ET AL: "Detection of circulating tumor cells in breast cancer patients utilizing multiparameter flow cytometry and assessment of the prognosis of patients in different CTCs levels", CYTOMETRY. PART A, vol. 77A, no. 3, 1 March 2010 (2010-03-01), pages 213-219, XP055222260, US ISSN: 1552-4922, DOI: 10.1002/cyto.a.20838**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 748 337 B1

Description

CROSS REFERENCE TO RELATED APPLICATION

[0001] The present application is based on provisional application Serial No. 61/538,302, filed September 23, 2011, the entire contents of which are herein incorporated by reference.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0002] Not Applicable.

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

[0003] The presently disclosed and claimed inventive concept(s) generally relates to cellular assays. More particularly, but not by way of limitation, the presently disclosed and claimed inventive concept(s) relates to methods of assaying biological samples to determine the cellular response to treatment/therapy and/or monitor progression of a disease state.

DESCRIPTION OF THE BACKGROUND ART

[0004] Affinity assays using fluorescence compounds are commonly used for cell and tissue analysis for biomarkers. The fluorescent compounds (labels) are attached (conjugated) to affinity molecules. Said affinity molecules may be, for example but not by way of limitation, antibodies, antigens, nucleic acid or other molecules that associate with the biomarkers of interest. This direct affinity method is a commonly used method to stain cells and tissues (See Horan and <http://www.crm.ed.ac.uk/facilities/flow-cytometry/protocols/immunofluorescence-staining>). The cells or tissues are incubated with conjugates, and unbound materials are washed away from cell bound materials. The conjugates are read cytometrically with a flow cytometer, scanning microscope or other device capable of detecting the fluorescence emission of the cellular material (see for example, US2008/0187198; Krivacic, 2004; and Herzenberg, 2002). Protocols vary and may or may not require cell or tissue fixation, depending on the conjugate used.

[0005] In practice, cell counts are measured using flow cytometry or scanning microscopy technologies which capture signals of cells at the excitation and emission wavelengths required to detect the dye. Different conjugates can be simultaneously determined (multiplexed) by using fluorescent dyes with distinct excitation wavelengths and emission wavelengths. Multiplexing of fluorescence signals is commonly used to detect as many properties as possible at one time. For example, biomarkers conjugated to phycoerythrin (ex 592, em 614nm) can be measured independent from biomarkers conju-

gated to fluorescein-5-isothiocyanate (FITC) (ex 488, em 525nm) using different excitation and emission filters. A list of known fluorophores is commonly available to researchers in the field (see <http://www.fluorophores.org/>).

[0006] Cell and tissue assays are typically performed for various types of cancer assays. Both cell and tissue assays use affinity reagents to detect biomarkers on or inside the cell, such as peptides, proteins, nucleic acid and modifications thereof (see, for example, Punnoose, 2010). Nucleic acids are measured as messenger RNA, Micro RNA and/or DNA prior to or after PCR amplification. Antibodies are also used for detecting protein biomarkers such as, but not limited to, EGFR, IGFRE, ERBB2, PSA, PL2L, KRAS, EPCAM, CK, and CD. A variety of tumor markers have detected breast, colon, prostate and melanoma cancer cells (see, for example, Mocellin, 2006). The use of biomarkers for monitoring cancer progression and/or treatment is also common practice. For example, the use of the plasminogen activator system (uPA) is monitored by measuring uPA, the plasminogen inhibitor PAI-1, and the complex of both (see Carney et al., 2009).

[0007] Cell assays are often looking for rare but clinically significant events. For example, circulating tumor cells (CTC) are significant at 5 cancer cells in 80,000,000 normal blood cells. The CTC must be isolated by enrichment or depletion to eliminate interference from normal cells (see, for example, Pantel et al., 2008). The isolated rare cells of interest then have to be assayed by methods that can detect each individual cell. Additionally the biomarkers expressed on these cells can be as low as 100 molecules per cell. Overall high sensitivity detection methods are required.

[0008] The sensitivity of traditional fluorescent labels for detecting biomarkers is generally poor for rare cell analysis. Excess affinity molecules must be used to increase the number of probes associated with the cell or tissue. This problem can be overcome partially using enzymes to catalytically increase the number of fluorescent labels. For example, Tyramide Signal Amplification (TSA) uses peroxidase enzyme to catalyze the deposition of multiple fluorophores on tyrosines of proteins by formation of tyramide with phenolic labeled fluorophores. The enzyme is associated with an affinity label to direct which cells are reacted (see, for example, Bobrow et al., 1993). However, enzymatic amplification involves multiple lengthy steps that increase the difficulty of the measurements.

[0009] Nanoparticles can be used to increase the number of fluorescent labels. However, multiple affinity molecules must also be attached to the nanoparticles, and adsorption of organic compounds onto nanoparticles also decreases the fluorescence signals. As a result, the nanoparticles with fluorescent labels (such as FITC) are no more sensitive than directly conjugated fluorescent labels. Additional problems with nanoparticles are that the size and coating must be carefully controlled for the particles to cross the cell walls (see, for example, Good-

man et al., 2007).

[0010] Rare earth elements, like europium (Eu), when chelated, or in oxides or silicate lattices, have fluorescent or luminescent behaviors that allow detection thereof in a method with high sensitivity. These phosphors have been used as high sensitivity signals for detection of material. For example, Ryan et al. (1973) first explained the use of rare-earth metal activated phosphors to detect explosive material. Nanoparticles containing the europium phosphor can be fluorescent in a 500 to 700 nm range and thus can be useful as a tracer for explosives.

[0011] Chelation of europium is critical to stability and strength of the fluorescence signal. Europium has long been shown to form a strong chelate with beta diketones. Shepard and Serigne (1934) have shown thenoyltrifluoracetone (TTA) has an association constant (log Ka) of 8.0 when converted to enolate. Stary (1959) demonstrated that benzoylacetone (HBA) has a log K of 18.9. Haar and Umland (1962) showed 8-hydroxyquinoline (oxine) forms an oxinate with europium. Dyrssen (1956) describes Beta-isopropyltropolone (HIPT) log K - 6.24 dissociation constant. Eu^{2+} resembles Ba^{2+} and can be complexed by various amine poly carboxylate ions such as DTPA (log Ka 23), DCTA (log Ka 19), EDTA (log Ka 17), HEDTA (log Ka 15), HTA (log Ka 11), and IMDA (log Ka 6). Matsumoto et al. (2007) demonstrated that europium with a variety of chelates in silicate nanoparticles can be a fluorophor in the range of 500 to 700 nm. Ullman et al. (1994) and Kraus et al. (2001) demonstrate europium with a variety of chelates in polystyrene nanoparticles. In particular, N,N,N,N'-[2,6-bis(3'-aminomethyl-1"-pyrazolyl)-4-phenylpyridine} tetra kis (acetic acid) (BB-TA) and 3-(2-thienoyl)-1,1,1-trifluoroacetone (TTA) are useful. In general, a wide variety of compounds has been shown to complex with lanthanide elements (for example but not by way of limitation, europium (Eu)), and this complex provides stability and strength to the fluorescence signals.

[0012] In practice, rare earth metal labels (such as but not limited to, lanthanides like Eu, Sm, Tm, Pr) or others like Tb have been applied as high sensitivity fluorescents or luminescent detection technology in nanoparticles (Ullman et al. 1994; Kraus et al., 2001; and Harma et al., 2001). These labels have been applied to various analytical methods for clinical diagnostics using nanoparticles. Ullman et al. demonstrated that nanoparticles containing europium chelate were useful in diagnostics as biomarker labels that generate luminance. Kraus et al. also showed nanoparticles with Europium chelate (such as but not limited to Eu TTA_3) were detectable above 500 nm and useful as diagnostic labels for biomarkers. Harma et al. showed the advantages of time resolved fluorescence using europium-label detection technology. In general europium fluorescence is either an acceptable high sensitivity label when the signal is used directly or amplified with subsequent reactions.

[0013] Another type of fluorescent label is a molecule which binds directly to the biomarker or a probe. These

labels do not need affinity molecules. For example, 4', 6'-diamidino-2'-phenylindole, dihydrochloride (DAPI) is a probe that fluoresces blue (455 nm) when bound to double stranded DNA and excited by exposure to light at 345 nm (see, for example, Morikawa et al., 1981). The detection of DNA in cells is an indication of a living cell nucleus. Other probes containing bis(zinc²⁺ dipicolylamine) groups bind to surfaces enriched with anionic phospholipids, especially phosphatidylserine (PS) exposed on cell membranes (see, for example, US Patent No. 7,179,616). The appearance of phosphatidylserine (PS) on the cell surface indicates cell apoptosis, prior to DNA fragmentation, morphological changes, and plasma membrane permeabilization.

[0014] Overall it is known that cell and tissue assays can be done by affinity assays whether using affinity reagents with fluorescent labels or fluorescent probes. It is also known that these signals can be multiplexed, and high sensitivity labels and probes are needed to detect rare events. Further, it is known that biomarkers can be selected for assay applications in the field of cancer.

[0015] Cell and tissues assays in the field of cancer are used to direct treatment. With the entry of more antibody-based chemotherapies, a patient's cancer must be typed to assure the cancer antigen is present (personalized medicine). The number of ever increasing cancer antigens is becoming increasingly complex (see, for example, Cheever et al., 2009). As a result, the selection of biomarkers to test is a difficult choice; the expression level, number of epitopes, expression on cancer stem cells, cellular location of expression, and the number of patients with antigen positive cancers all must be considered. Antigens with therapeutic function, immunogenicity (T cell expression), oncogenicity (association with the oncogenic process) and specificity rank highest in value. Even with a ranking, selection is not straight forward and is in need of a simplification; picking the wrong antigen for a patient is still likely, and all antigens cannot be measured at one time.

[0016] Chemo resistance is another important parameter that cell and tissue assays try to address by measuring cells and tissue during the treatment course to determine if cancer antigens are still present. Another method for determining chemo resistance is to culture or grow the cancer cells and see if the cancer antigens are expressed by living cells (see, for example, the so called EPISPOT method of Alix-anabieres et al., 2009). Another method is to measure the cancer stem cell antigens as a measure of chemo resistance (see for example, Pardalet al., 2003; and Gao et al., 2010). However, all three of these methods also rely on "picking" the right antigen and fail to give good information for all patients.

[0017] Thus the current field lacks a comprehensive measure of the metastatic potential of a cancer cell/tissue. The presently disclosed and claimed inventive concept(s) provide a cell based response assay that overcomes the defects and disadvantages of the prior art.

BRIEF DESCRIPTION OF THE DRAWINGS

[0018] FIGURE 1 graphically depicts a cell response assay constructed in accordance with the presently disclosed and claimed inventive concept(s).

DETAILED DESCRIPTION OF THE INVENTION

[0019] Before explaining at least one embodiment of the inventive concept(s) in detail by way of exemplary drawings, experimentation, results, and laboratory procedures, it is to be understood that the inventive concept(s) is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the drawings, experimentation and/or results. The inventive concept(s) is capable of other embodiments or of being practiced or carried out in various ways. As such, the language used herein is intended to be given the broadest possible scope and meaning; and the embodiments are meant to be exemplary - not exhaustive. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

[0020] Unless otherwise defined herein, scientific and technical terms used in connection with the presently disclosed and claimed inventive concept(s) shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures utilized in connection with, and techniques of, cell and tissue culture, molecular biology, and protein and oligo- or polynucleotide chemistry and hybridization described herein are those well known and commonly used in the art. Standard techniques are used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (e.g., electroporation, lipofection). Enzymatic reactions and purification techniques are performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The foregoing techniques and procedures are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. See e.g., Sambrook et al. *Molecular Cloning: A Laboratory Manual* (2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) and Coligan et al. *Current Protocols in Immunology* (Current Protocols, Wiley Interscience (1994)). The nomenclatures utilized in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[0021] All patents, published patent applications and non-patent publications mentioned in the specification are indicative of the level of skill of those skilled in the art to which this presently disclosed and claimed inventive concept(s) pertains.

[0022] All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this presently disclosed and claimed inventive concept(s) have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the presently disclosed and claimed inventive concept(s). All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the inventive concept(s) as defined by the appended claims.

[0023] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

The use of the word "a" or "an" when used in conjunction with the term "comprising" in the claims and/or the specification may mean "one," but it is also consistent with the meaning of "one or more," "at least one," and "one or more than one." The use of the term "or" in the claims is used to mean "and/or" unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and "and/or." Throughout this application, the term "about" is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects. The use of the term "at least one" will be understood to include one as well as any quantity more than one, including but not limited to, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, 100, etc. The term "at least one" may extend up to 100 or 1000 or more, depending on the term to which it is attached; in addition, the quantities of 100/1000 are not to be considered limiting, as higher limits may also produce satisfactory results.

[0024] The term "about" is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value and/or the variation that exists among study subjects.

[0025] As used in this specification and claim(s), the words "comprising" (and any form of comprising, such as "comprise" and "comprises"), "having" (and any form of having, such as "have" and "has"), "including" (and any form of including, such as "includes" and "include") or "containing" (and any form of containing, such as "contains" and "contain") are inclusive or open-ended and do not exclude additional, unrecited elements or method steps.

[0026] The term "or combinations thereof" as used

herein refers to all permutations and combinations of the listed items preceding the term. For example, "A, B, C, or combinations thereof" is intended to include at least one of: A, B, C, AB, AC, BC, or ABC, and if order is important in a particular context, also BA, CA, CB, CBA, BCA, ACB, BAC, or CAB. Continuing with this example, expressly included are combinations that contain repeats of one or more item or term, such as BB, AAA, MB, BBC, AAABCCCC, CBBAAA, CABABB, and so forth. The skilled artisan will understand that typically there is no limit on the number of items or terms in any combination, unless otherwise apparent from the context.

[0027] The term "probe" as used herein will be understood to refer to any type of affinity reagent that binds to a specific biomarker as described herein. Examples of probes include, but are not limited to, antibodies (or binding fragments or derivatives thereof), receptors, organic molecules, inorganic molecules, ligands, nucleic acids (including but not limited to, DNA, RNA, microRNA, mRNA, siRNA, etc.), peptides, polypeptides, proteins, epitopes, antigens, ligands, receptors, complexes, lipids, glycoproteins, glycolipids, glycosaminoglycans, carbohydrates, polycarbohydrates, glycoconjugates, and any combination or derivative thereof.

[0028] The term "biomarker" as used herein will be understood to refer to any target site on the surface of or inside of a cell that a probe can have affinity therefor and thus can bind to said moiety. The "biomarker" may be, for example but not by way of limitation, a nucleic acid, peptide, polypeptide, protein, epitope, antigen, ligand, receptor, complex (i.e., an MHC-peptide complex), lipid, glycoprotein, glycolipid, glycosaminoglycan, carbohydrate, polycarbohydrate, glycoconjugate, and any combination or derivative thereof.

[0029] The terms "peptide", "polypeptide" and "protein" are used herein to refer to a polymer of amino acid residues. The term "polypeptide" as used herein is a generic term to refer to native protein, protein fragments, or analogs of a polypeptide sequence. Hence, native protein, protein fragments, and analogs are species of the polypeptide genus.

[0030] The terms "polynucleotide", and "nucleic acid" are used interchangeably. They refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. The following are non-limiting examples of polynucleotides: coding or non-coding regions of a gene or gene fragment, loci (locus) defined from linkage analysis, exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. If present, modifications to the nucleotide structure may be imparted before or after assembly of the polymer. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynu-

cleotide may be further modified, such as by conjugation with a labeling component. The terms "isolated nucleic acid" and "isolated polynucleotide" are used interchangeably; a nucleic acid or polynucleotide is considered "isolated" if it: (1) is not associated with all or a portion of a polynucleotide in which the "isolated polynucleotide" is found in nature, (2) is linked to a polynucleotide to which it is not linked in nature, or (3) does not occur in nature as part of a larger sequence.

[0031] The term "antibody" is used in the broadest sense, and specifically covers monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments so long as they exhibit the desired biological activity. Thus, the terms "Antibody" or "antibody peptide(s)" refer to a full length immunoglobulin molecule (i.e., an intact antibody), or a binding fragment thereof that competes with the intact antibody for specific antigen binding. Binding fragments may be produced by recombinant DNA techniques, or by enzymatic or chemical cleavage of intact antibodies. Binding fragments include Fab, Fab', F(ab')₂, Fv, scFv, disulfide linked Fv, Fd, diabodies, single-chain antibodies, single domain antibodies (such as but not limited to, NANOBOODIES®) and other antibody fragments that retain at least a portion of the variable region of an intact antibody. See, e.g., Hudson et al. (Nature Med., 9:129-134 (2003)).

[0032] The term "antigen binding fragment" or "antigen-binding portion" of an antibody, as used herein, refers to one or more fragments of an antibody that retain the ability to bind to an antigen. The antigen-binding function of an antibody can be performed by fragments of an intact antibody. Examples of binding fragments encompassed within the term "antigen-binding fragment" of an antibody include but are not limited to, Fab, Fab', F(ab')₂, Fv, scFv, disulfide linked Fv, Fd, diabodies, single-chain antibodies, single domain antibodies (such as but not limited to, NANOBOODIES®), isolated CDRH3, and other antibody fragments that retain at least a portion of the variable region of an intact antibody. These antibody fragments are obtained using conventional recombinant and/or enzymatic techniques and are screened for antigen binding in the same manner as intact antibodies.

[0033] The term "epitope" includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor. In certain embodiments, an epitope is a region of an antigen that is specifically bound by an antibody. Epitopic determinants usually include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl, or sulfonyl groups. In certain embodiments, an epitope may have specific three dimensional structural characteristics (e.g., a "conformational epitope"), as well as specific charge characteristics.

[0034] The term "nanoparticle" as used herein refers to a particle having dimensions of from about 1 to about 5000 nanometers, and having any size, shape or morphology. The nanoparticles utilized in accordance with

the present invention may be naturally occurring, commercially available nanoparticles, or the nanoparticles may be synthesized for use in accordance with the present invention, as described herein below and as known in the art. Particular examples of nanoparticles that may be utilized in accordance with the present invention include, but are not limited to, poly(lactic-co-glycolic) acid (PLGA) nanoparticles, poly lactic acid (PLA) nanoparticles, Chitosen nanoparticles, liposomes, and derivatives or combinations thereof.

[0035] As used herein, the terms "label" or "labeled" refers to incorporation of a detectable marker, e.g., by attachment of a fluorescent, enzymatic or colorimetric label or incorporation of a radiolabeled amino acid. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ³H, ¹⁴C, ¹⁵N, ³⁵S, ⁹⁰Y, ⁹⁹Tc, ¹¹¹In, ¹²⁵I, ¹³¹I), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic labels (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

[0036] The terms "label", "detectable marker" and "detection moiety" are used interchangeably herein.

[0037] A fluorophore may be employed in the methods of the present invention and detected via any of numerous colorimetric and fluorescence detection methods. Depending on the application and purpose, such methods include, but are not limited to, absorbance spectroscopy, fluorescence spectroscopy, fluorescence activated cytometry (FACS), fluorescence microscopy, fluorescence resonance energy transfer (FRET), and the like.

[0038] Various types of fluorophores, depending on the application and purpose, may be employed in accordance with the present invention. Examples of suitable fluorophores are described herein below. Other examples are given in US Patent Nos. 7,465,747 and 7,955,859, issued to Matsumoto et al. on December 16, 2008 and June 7, 2011, respectively; and US Publication No. US2007/0026407, published February 1, 2007 (the entire contents of which are expressly incorporated herein by reference in their entirety).

[0039] Ample guidance regarding fluorophore selection, methods of linking fluorophores to various types of molecules, and methods of use thereof is available in the literature of the art [for example, refer to: Richard P. Haugland, "Molecular Probes: Handbook of Fluorescent Probes and Research Chemicals 1992-1994", 5th ed., Molecular Probes, Inc. (1994); U.S. Pat. No. 6,037,137 to Oncoimmunin Inc.; Hermanson, "Bioconjugate Techniques", Academic Press New York, N.Y. (1995); Kay M.

et al., 1995. *Biochemistry* 34:293; Stubbs et al., 1996. *Biochemistry* 35:937; Gakamsky D. et al., "Evaluating Receptor Stoichiometry by Fluorescence Resonance Energy Transfer," in "Receptors: A Practical Approach," 2nd ed., Stanford C. and Horton R. (eds.), Oxford University Press, UK. (2001); U.S. Pat. No. 6,350,466 to Targesome, Inc.]. Therefore, no further description is considered necessary.

[0040] The terms "substantial increase" and "substantial decrease", as well as grammatical equivalents thereof, will be understood herein to refer to at least a 12% increase or decrease, such as at least a 30% increase or decrease, at least a 50% increase or decrease, at least a 75% increase or decrease, or at least a 90% increase or decrease.

[0041] The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, and sarcoma. More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial carcinoma, salivary gland carcinoma, kidney cancer, renal cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma and various types of head and neck cancer.

[0042] The term "metastasis" as used herein will be understood to refer to the spread of cancer from a primary tumor to other parts of the body. Metastasis is a sequential, multistep process in which tumor cells detach from a primary tumor, migrate through the basement membrane and extracellular matrix, and invade the lymphatic and/or blood systems. This is followed by the establishment of secondary tumors at distant sites.

[0043] "Mammal" for purposes of treatment refers to any animal classified as a mammal, including human, domestic and farm animals, nonhuman primates, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, etc.

[0044] The term "patient" includes human and veterinary subjects. In certain embodiments, a patient is a mammal. In certain other embodiments, the patient is a human.

[0045] The term "biological sample" as used herein will be understood to refer to a sample of biological fluid. Biological samples include, but are not limited to, blood, plasma, serum, sputum, cerebrospinal fluid (CSF), tears, mucus, urine, tissue, other types of specimens, and the like.

[0046] The phrase "providing a biological sample" as used herein refers to obtaining a biological sample for use in methods described and claimed herein. Most often, this will be done by removing a sample of cells from an animal, but can also be accomplished by using previously isolated cells (e.g., isolated by another person,

at another time and/or for another purpose). The step of "providing a biological sample" may further include various isolation and/or purification steps known in the art for providing a specific component of a biological sample for use in the methods described and claimed herein.

[0047] Numerous aspects and advantages of the inventive concept(s) will be apparent to those skilled in the art upon consideration of the following detailed description which provides illumination of the practice of the presently disclosed and claimed inventive concept(s).

[0048] The presently disclosed and claimed inventive concept(s) generally relates to multiplexed direct affinity conjugate assays of cells using probes able to determine a cellular response to treatments; this type of assay is herein termed a "cell response assay".

[0049] The presently disclosed and claimed inventive concept(s) generally relates to the use of a combination of biomarkers for cancer cell/tissue type, metastatic potential and chemo resistance in a multiplexed affinity assay for cancer cells and tissue. Fluorescent labels either conjugated to affinity molecules/reagents or capable of directly binding to the biomarker are used and are referred to herein as labeled probes. This combination of labeled probes for biomarkers allows determination of the cellular response to treatments by simultaneously measuring cell type, chemo resistance nature, and metastatic potential and/or cell viability, and allows simple multiplexed signals with high sensitivity.

[0050] Surprisingly, the biomarker affinity markers selected in this order solves the problem of "picking" the right antigen for cancer antigen type and also provides accurate information on the response to therapy. More particularly, the presently disclosed and claimed inventive concept(s) relates to methods of using said affinity molecules and molecular probes as compositions, dosage forms, and kits.

[0051] The presently disclosed and claimed inventive concept(s) combines cell biomarker(s) for tissue type with biomarker(s) for metastatic potential and biomarker(s) for cell viability in an affinity assay for cancer cells and tissue. In Figure 1, the biomarker for cancer cell type is represented as biomarker A and is always present in the disease state. The biomarker for metastatic potential is represented as biomarker B and is present in the disease state with metastatic invasiveness potential. The biomarker for chemo resistance is represented as biomarker C and is present in unresponsive cells. Fluorescent labels either conjugated to the affinity reagents or capable of directly binding the biomarker are used to simultaneously detect biomarkers A, B and C; simultaneous detection is obtained by using fluorescent labels that are detectable at different excitations and/or emission wavelengths.

[0052] The desired response to the treatment would be first for the number of cancer cells to decrease overall. However, the unexpected response identified by the presently disclosed and claimed inventive concept(s) is for biomarker B (i.e., metastatic potential) to increase or

become present in the course of treatment in a greater proportion of cancer cells detected by biomarker A, and for biomarker C (i.e. chemo resistance) to decrease or become absent in the course of treatment in a greater proportion of cancer cells detected by biomarker A. In order for this combination of cell biomarkers to correctly predict the response, the biomarker for tissue type, metastatic potential and cell viability must be biochemically linked in a process. Biomarkers A, B and C must co-exist in the disease state and be fundamental to progression or relapse of the disease. Further, the presence of biomarkers B and C must be linked such that the presence of one facilitates the absence of the other. This combination of biomarkers allows the determination of the cell response to treatments; by simultaneously measuring cell type, chemo resistance nature, and metastatic potential (and possibly also cell viability), the "cell based response assay" can accurately monitor cancer treatment and/or monitor disease progression/relapse.

[0053] Turning now to the particular embodiments of the presently disclosed and claimed inventive concept(s), one embodiment thereof is directed to a multiplexed assay for cancer. Said assay involves simultaneously measuring: (1) at least one cancer cell type biomarker (also referred to herein as Biomarker A) in a biological sample of a cancer patient utilizing a first labeled probe that binds to said cancer cell type biomarker; (2) at least one chemo resistance biomarker (also referred to herein as Biomarker C) in the biological sample utilizing a second labeled probe that binds to said chemo resistance biomarker; and (3) at least one metastatic potential biomarker (also referred to herein as Biomarker B) in the biological sample utilizing a third labeled probe that binds to said metastatic potential biomarker. The at least three labeled probes are measured at different excitations and emission wavelengths.

[0054] The assays/methods described herein may be utilized for a specific cancer, or the assays/methods may be general assays for all types of cancers. Likewise, the cancer cell type biomarker utilized in accordance with the presently disclosed and claimed inventive concept(s) may be a non-specific cancer cell biomarker, or may be specific for a certain type of cancer. Examples of cancers that may be detected/monitored by the currently disclosed and claimed inventive concept(s) include, but are not limited to, lung, bronchus, colon, rectum, pancreas, prostate, breast, liver, bile duct, bladder, ovary, brain, central nervous system (CNS), kidney, pelvis, uterine corpus, oral cavity or pharynx or melanoma cancers.

[0055] Non-limiting examples of non-specific cancer cell type biomarkers include, but are not limited to, cytokeratins (CK), EpCAM, N-cadherin, E-cadherin and vimentin.

[0056] For example but not by way of limitation, the carcinoma cells may be indicated by cytokeratin (CK) as a biomarker A, and the first labeled probe may comprise multiple labeled antibodies for separate cytokeratins (such as but not limited to, cytokeratins 8/18/19. Cytok-

erations (CK) are proteins of keratin-containing intermediate filaments found in the intracytoplasmic cytoskeleton of epithelial tissue and refer to a family of fibrous structural proteins. There are two fundamental types of cytokeratins: the acidic type I cytokeratins and the basic or neutral type II cytokeratins. Cytokeratins are usually found in pairs comprising a type I cytokeratin and a type II cytokeratin. Basic or neutral cytokeratins include (but are not limited to) CK1, CK2, CK3, CK4, CK5, CK6, CK7, CK8 and CK9. Acidic cytokeratins include (but are not limited to) CK10, CK12, CK 13, CK14, CK16, CK17, CK18, CK19 and CK20.

[0057] In other instances the, Epithelial cell adhesion molecule (EpCAM) is a protein that signifies the presence of epithelial cells in a carcinoma. EpCAM has also been designated as TACSTD1 (tumor-associated calcium signal transducer 1) and CD326 (cluster of differentiation 326). Thus, the terms "EpCAM", "TACSTD1" and "CD326" are used herein interchangeably.

[0058] In other instances, some cancer cells undergo an Epithelial to Mesenchymal transition (EMT) that may be indicated by an increase in cancer cell biomarkers such as but not limited to, Vimentin and Galectin-3, which signify cells with a loss of anchorage. Alternatively, certain other cancer cells undergo a Mesenchymal to Epithelial transition (MET) indicated by an increase in cancer cell biomarkers such as but not limited to, N-Cadherin, and E-Cadherin. The EMT and MET stages for cancer cells signify a change in sub type of cells. It is important to detect as many cancer cells as possible and to additionally know their sub types to measure the entire tumorigenicity.

[0059] The term "cancer cell type biomarker" also includes markers that identify specific genetic mutations that cause oncoproteins or oncogenes to be regulated (whether tumor promoting or tumor suppressing). These types of cancer cell type biomarkers may be utilized alone or in combination with any of the other cancer cell type biomarkers described herein above. Examples of these types of cancer cell type biomarkers include, but are not limited to, HER2/neu, VEGF-165, KRAS, EGFR, WAF, BAX-1, PDGF, Rb, Jagged 1, Notch, VEGF, VEGFR, k-Ras, CAIX, MIB1, MDM, PR, ER, SEL5, SEM1, PI3K, Akt2, twist 1, EML-4, ALK, Braf, DRAFF, c-met, and combinations thereof. These oncoproteins and oncogenes are used to direct targeted therapies based on their presence in/on cells. For example, the presence of HER2/neu is used to prescribe Herceptin therapy; however, HER2/neu, as with other oncoprotein/oncogene markers, is often only expressed on a fraction of cancer cells, and not all patients have the specific genetic mutation leading to the biomarker. HER2/neu-positive expression is present in only 10-20% of breast cancer patients, and even in the HER2/neu-positive patients, the marker is only present in approximately 30% of their circulating tumor cells. Therefore these markers are often measured in combination with another cancer cell type biomarker.

[0060] In other instances, the carcinoma cells may be

indicated by a tissue specific marker. For example, Prostate-Specific Antigen (PSA) is a protein that signifies the presence of cells of the prostate gland. Non-limiting examples of specific cancer cell type biomarkers include, but are not limited to: prostate specific antigen (PSA) or prostate specific membrane antigen (PSMA) can be used to detect prostate cancer cells; MUC1, CA 15-3 and CA 27-29 can be used to detect breast cancer cells; Carcinoembryonic Antigen (CEA), CA19-9 or Galactosyl Transferase II can be used to detect colon cancer cells; MSLN (mesothelin) can be used to detect pancreatic cancer cells; CA 125 or Follicle-Stimulating Hormone (FSH) receptor can be used to detect ovarian cancer cells; Alpha-Fetoprotein can be used to detect liver cancer cells; Melan-A (MLANA), Tyrosinase (TYR), CSPG4, or MITF can be used to detect melanoma cancer cells; and Parathyroid related protein (PTHrP) or TSHR can be used to detect thyroid cancer cells.

[0061] Any metastatic potential biomarkers known in the art or otherwise contemplated herein may be utilized in accordance with the presently disclosed and claimed inventive concept(s). The third labeled probe may be any probe capable of detecting said metastatic potential biomarker; for example but not by way of limitation, the third labeled probe may be one or more labeled antibodies against any of the markers of tumor cell invasiveness listed below. In some instances, the metastatic potential biomarker may be measured as a ratio of one of the below-listed biomarkers to the cancer cell type biomarker. Particular non-limiting examples of biomarkers of metastatic potential that may be utilized in accordance with the presently disclosed and claimed inventive concept(s) include a marker(s) of tumor cell invasiveness where the marker measures cancer cell invasion into the extracellular membrane through proteolytic events, such as urokinase plasminogen activator (uPA), plasminogen activator inhibitor (PAI-1), CD95, serine proteases such as plasmin, ADAM, and others; serine protease inhibitors such as Bikunin; matrix metalloproteinases such as MMP9; matrix metalloproteinase inhibitors such as TIMP-1; and combinations thereof. For example but not by way of limitation, a prostate cancer assay may measure PSA as a cancer cell type biomarker, and measure a ratio of PAI-1/PSA for metastatic potential.

[0062] Any chemo resistance biomarkers known in the art or otherwise contemplated herein may be utilized in accordance with the presently disclosed and claimed inventive concept(s). In certain embodiments, the chemo resistance biomarker is a biomarker that detects the prevention of cell death (apoptosis), for example but not by way of limitation, a detection of the presence of a cancer stem cell biomarker, such as but not limited to, PL2L piwi like, ADLH, β -integrin, α 6 integrin, c-kit, c-met, LIF-R, CXCR4, ESA, CD 20, CD44, CD133, CK5, TRAF2 and ABC transporters. In another non-limiting example, cancer cells that contain CD44 but lack CD24 are indicative of a cancer stem cell phenotype. Also, cancer cells that lack CD45 and CD31 but contain CD34 are indicative of

a cancer stem cell. Also, cancer cells that contain CD44 and CD24 as well as ESA are indicative of a cancer stem cell. Further, the presence of both CD24 and ESA are indicative of a cancer stem cell. Somatic stem cells in solid tumor carcinomas can become cancer stem cells. Other cancer cells resistant to CD95 induced-apoptosis are chemo resistant. Cancer stem cells are chemo resistant in nature, and contain a capacity for self renewal (asymmetric divisions) as well as being capable of differentiation into a hierarchy of progeny cells to form a tumor. Cancer stem cell self renewal is activated by the stem cell signaling pathways (Wnt, Sonic Hedge hog and Notch) and at the epigenetic level by Polycomb gener (BMI-1 and EZH2). This self renewal occurs at the expense of apoptosis signaling pathways (Caspase 3, 5, 8, p53). Measuring cancer stem cell biomarkers and combinations thereof as an indication of chemo resistance provides a measure of (a) drug resistance, (b) the inability to activate apoptosis, and/or (c) the inability to shutdown self renewal.

[0063] The probes described and claimed herein may be labeled by any methods known in the art or otherwise contemplated herein. For example, but not by way of limitation, the labels of the first, second and third (and any additional fourth, fifth, sixth, etc.) labeled probes may be selected from a comprehensive catalogue of fluorescent (luminescent) dyes. For example but not by way of limitation, a comprehensive catalogue exists online at <http://www.fluorophores.org> (the entire contents of which are expressly incorporated herein by reference). This catalogue includes commonly used labels such as fluorescein-5-isothiocyanate (FITC), phycoerythrin, sulforhodamine 101 (Texas Red), 2-[4-(aminoiminomethyl)phenyl]-1H-Indole-6-carboximidamide (DAPI), 3H-Indolium (Cy5), 1H-benz[e]indolium (Cy 5.5), 3H-Indolium (Cy 7), ALEXA FLUOR@488, ALEXA FLUOR@555, ALEXA FLUOR@647, and combinations and derivatives thereof that have been synthesized in order to provide better reagents. Additionally, rare earth metals and rare earth element-containing nanoparticles can be used as labels.

[0064] The biological samples utilized in the methods of the presently disclosed and claimed inventive concept(s) may be utilized in the form they are obtained (i.e., tissue sample), or they may be exposed to one or more isolating steps (i.e., isolation of cancer cells therefrom).

[0065] While the assays described herein above describe the measurement of three biomarkers with three labeled probes, it is to be understood that one or more additional biomarker(s) may also be measured in the assay (utilizing additional one or more additional labeled probe(s)). For example, but not by way of limitation, the assay may further include measuring at least one additional biomarker in the biological sample utilizing a fourth labeled probe that binds to said additional biomarker, measuring a second additional biomarker in the biological sample utilizing a fifth labeled probe that binds to said additional biomarker, measuring a third additional biomarker in the biological sample utilizing a sixth labeled

probe that binds to said additional biomarker, etc. Ideally, a maximum of six to eight labeled probes are utilized; otherwise, it is difficult to detect all of the labeled probes at different, non-overlapping excitations and emission wavelengths.

[0066] These additional biomarkers may include biomarkers that detect other cellular properties, specifically the presence of cell nuclei or of intact cell membranes, for the cells that are positive for the cancer cell type biomarker. For example but not by way of limitation, the presence of cell nuclei may be detected utilizing a DNA binding probe such as but not limited to, DAPI. In another non-limiting example, the detection of intact cell membranes may involve the detection of anionic phospholipids (such as but not limited to, phosphatidylserine) on the surface of cells, and the second labeled probe may be bis(zinc2+dipicolylamine) and/or PSVue™.

[0067] The one or more additional biomarker(s)/labeled probe(s) may also be utilized to reduce any false positive results obtained with the cancer cell type biomarker; for example but not by way of limitation, white blood cells may be excluded from the assay by excluding cells positive for one or more markers of cluster of differentiation (also referred to as "cluster of designation"; often abbreviated as CD). For example but not by way of limitation, markers such as CD45, CTLA-4, CD4, CD68 and/or CD8 present on white blood cells can be used to indicate that a cell is not a cancer cell. In a particular non-limiting example, CD45 antigen (also known as PTPRC, Protein tyrosine phosphatase receptor type C, and originally called leukocyte common antigen (all terms used herein interchangeably)) is useful in detecting all white blood cells. Additionally, CD45 can be used to differentiate the different types of white blood cells when combined with other CD markers. For example, granulocytes are indicated by CD45+, CD15+; monocytes are indicated by CD45+, CD14+; T lymphocytes are indicated by CD45+, CD3+; T helper cells are indicated by CD45+, CD3+, CD4+; cytotoxic T cells are indicated by CD45+, CD3+, CD8+; B lymphocytes are indicated by CD45+, CD19+ or CD45+, CD20+; thrombocytes are indicated by CD45+, CD61+; and natural killer cells are indicated by CD16+, CD56+, CD3-. Additionally, two commonly used CD molecules are CD4 and CD8, which are, in general, used as markers for helper and cytotoxic T cells, respectively. These molecules are defined in combination with CD3+, as some other leukocytes also express these CD molecules (some macrophages express low levels of CD4; dendritic cells express high levels of CD8).

[0068] The presently disclosed and claimed inventive concept(s) is also directed to a kit that includes the first, second and third labeled probes (that bind to a cancer cell type biomarker, a chemo resistance biomarker, and a metastatic potential biomarker, respectively) as described in detail herein above. Additional labeled probes for detecting other biomarkers as further described herein above may also be included in the kit. All of the labeled

probes present in the kit are measured at different excitations and emission wavelengths.

[0069] The kit may also include means for isolating cancer cells from a biological sample. Said means are well known in the art (see for example, Lianidou and Markou, 2011; the entire contents of which are hereby expressly incorporated herein by reference); therefore, no additional discussion thereof is deemed necessary.

[0070] The presently disclosed and claimed inventive concept(s) is also directed to methods of (1) monitoring cancer treatment in a cancer patient undergoing said treatment, and (2) monitoring progression/relapse of the disease in the cancer patient. In the method of (1) monitoring cancer treatment, a first biological sample is obtained from the patient prior to exposure to a cancer treatment, and a second biological sample is obtained from the patient following exposure to the cancer treatment. In the method of (2) monitoring progression/relapse of disease, the first biological sample is obtained from the patient at a first time point, and the second biological sample is obtained from the patient at a subsequent time point. In both methods, the levels of a cancer cell type biomarker (as described herein above), a chemo resistance biomarker (as described herein above), and a metastatic potential biomarker (as described herein above) are then measured utilizing first, second and third labeled probes, respectively (as described in detail herein above), in the first and second biological samples. The cells to which the first labeled probe is bound are identified in the first and second biological samples, and then the levels of (i) the chemo resistance biomarker and (ii) the metastatic potential marker in the cells to which the first labeled probe is bound are compared in the first and second biological samples. It is then determined in the method of (1) that the cancer treatment is effective if the level of the chemo resistance biomarker is decreased and the level of the metastatic potential biomarker is increased in the second biological sample when compared to the first biological sample; alternatively, it is then determined in the method of (2) that the cancer has progressed/relapsed if the level of the chemo resistance biomarker is increased and the level of the metastatic potential biomarker has decreased in the second biological sample when compared to the first biological sample.

[0071] The biological samples utilized in the methods of the presently disclosed and claimed inventive concept(s) may be utilized in the form they are obtained (i.e., tissue sample), or they may be exposed to one or more isolating steps (i.e., isolation of cancer cells therefrom).

[0072] While the methods described herein above describe the measurement of three biomarkers with three labeled probes, it is to be understood that additional biomarkers may also be included in the methods (utilizing additional labeled probes). For example, but not by way of limitation, the method may further include measuring at least one additional biomarker in the first and second biological samples utilizing a fourth labeled probe that binds to said additional biomarker, measuring a second

additional biomarker in the first and second biological samples utilizing a fifth labeled probe that binds to said additional biomarker, measuring a third additional biomarker in the first and second biological samples utilizing a sixth labeled probe that binds to said additional biomarker, etc. Ideally, a maximum of six to eight labeled probes are utilized; otherwise, it is difficult to detect all of the labeled probes at different, non-overlapping excitations and emission wavelengths. When additional biomarkers are measured, the methods may also include the step of comparing the levels of the additional biomarker(s) in the cells to which the first labeled probe is bound in the first and second biological samples.

[0073] These additional biomarkers may include biomarkers that detect other cellular properties, specifically the presence of cell nuclei or of intact cell membranes, for the cells that are positive for the cancer cell type biomarker. For example but not by way of limitation, the presence of cell nuclei may be detected utilizing a DNA binding probe such as but not limited to, DAPI. In another non-limiting example, the detection of intact cell membranes may involve the detection of anionic phospholipids (such as but not limited to, phosphatidylserine) on the surface of cells, and the second labeled probe may be bis(zinc2+dipicolylamine) and/or PSVue™.

[0074] The one or more additional biomarker(s)/labeled probe(s) may also be utilized to reduce any false positive results obtained with the cancer cell type biomarker; for example but not by way of limitation, white blood cells may be excluded from the methods by excluding cells positive for one or more markers of cluster of differentiation (as described in detail herein above).

EXAMPLES

[0075] Examples are provided hereinbelow. However, the present invention is to be understood to not be limited in its application to the specific experimentation, results and laboratory procedures. Rather, the Examples are simply provided as one of various embodiments and are meant to be exemplary, not exhaustive.

Example 1 - Multiplexed Assay for Breast Cancer

[0076] The following procedure shows the multiplexing of Biomarkers A (cancer cell type), B (metastatic potential) and C (chemo resistance) using a unique combination of affinity reagents with fluorescent labels and an affinity label to provide a multiplexed assay for breast cancer. This allows the simultaneous measurement of cancer cell type, chemo resistance nature, and metastatic potential, as described herein above, in combination with a marker indicating the presence of cell nucleus (utilizing a nucleic acid binding probe). The data demonstrated that four simultaneous signals can be measured with this combination but not with the traditional combinations of the prior art. The presently disclosed and claimed inventive concept(s) has the potential to measure up to

eight signals utilizing various fluorophores taught herein.

[0077] The procedure to test the method was to collect 4 mL normal whole blood fresh into a VACUTAINER® tube (BD 6 mL VACUTAINER® K2EDTA tube, 10.8 mg, ref 367863, Becton Dickinson, Oakville, ON) and transfer the blood into a round centrifuge tube sealed with top. Cancer cells were added by 40 μ L of breast cell cancer suspension to obtain a range of 3000 to 8000 cells/mL (8 cells/ μ L or 1-2 per high powered filed HRP (HRP=0.2 μ L)). The blood was lysed by adding 3 mL of lysis buffer (155mM NH_4Cl , 20mM KHCO_3 , 0.1 mM Na_2EDTA , pH 7.4). The tubes were rolled tubes to assure that the cells were evenly distributed and that lysis was complete, and then the sample was spun down for 15 minutes at 12,000 rpm. Next, the plasma was decanted as waste supernatant, and 4.0 mL of TES buffer (50 mM N-Tris [Hydroxymethyl]-2-aminoethane-sulfonic acid: TES, pH 7.4, 2150mM NaCl , 2 mM MgSO_4 and 1 mM KCO_3) and 1 % albumin was added, and the tube was rocked to mix the cells. The tube was spun down again for 15 minutes at 12,000 rpm, and the supernatant was decanted as waste. A final 2.0 mL of TES wash buffer was added with rocking to mix cells.

[0078] The cells were then reacted with the reagents by taking 0.5 mL of washed blood to the centrifuge tube, and 40 μ L of antibodies for HER2/neu labeled with Cy5 and 40 μ L of antibodies for urokinase like plasminogen activator (uPA) or plasminogen activator inhibitor (PAI-1) labeled with Texas Red (at working stock of 0.001 to 0.1 mg/mL) and PL2L piwi like labeled with FITC (at working stock of 0.001 to 0.1 mg/mL) were added. Next, 10 μ L of a 100 μ g/ml 4,6 Diamidino-2-phenylindole dihydrochloride (DAPI; a nucleic acid binding probe) was added from a solution in TES buffer. This was followed by incubation for 15 minutes at 37°C.

[0079] The sample was centrifuged down for 5 minutes @ 7500 rpm, and the supernatant was decanted off as waste (~450 μ L). 500 μ L of TES buffer was then added, and the sample was vortexed and centrifuged down at 5 minutes @ 7500 rpm. Again, the supernatant was decanted off by tapping as waste (~450 μ L). A final 500 μ L of TES wash buffer was added for final imaging, and 5 μ L of the solution was placed on a slide with a long cover slip of glass. Phase contrast and fluorescence microscopy was conducted with the Leica DM5000 (Leica Microsystems, Buffalo Grove, IL). A sample of cells was stained with the PSS-550 probe (ex 553, em 615nm) and measured with an excitation band pass filter at 540-580 nm and an emission band pass filter at 610-680 nm to determine cell membrane flopping as a measure of cell death in addition to cell counts .

[0080] The Texas Red label (ex 592, em 614 nm) was measured with an excitation band pass filter at 540-580 nm and an emission band pass filter at 610-680 nm. The FITC label (ex 488, em 525nm) was measured with an excitation band pass filter at 460 - 500 nm and an emission band pass filter at 512 -543 nm. The DAPI probe (ex 355, em 460 nm) was measured with an excitation

band pass filter at 340-380 nm and an emission band pass filter at 450-490 nm. The Cy5 label (ex 646, em 676 nm) was measured with an excitation band pass filter at 590-650 nm and an emission band pass filter at 665-735 nm.

[0081] The ideal multiplexing was demonstrated by combination of the rare earth nanoparticles with FITC, Texas Red, Cy5 and DAPI. The europium nanoparticle (ex 355, em 617 nm) was measured at an excitation filter band pass at 340-380 nm and emission long pass filter of > 590 nm. Simultaneously measuring biomarkers for cancer cell type, metastatic potential and chemoresistance, along with a biomarker indicating the presence of cell nucleus was possible, along with the removal of false positives (i.e., white blood cells) by detecting the presence of the biomarker CD45. Current data allows for five signals. However, the assays/methods described and claimed herein have the potential for detection and measurement of six signals/biomarkers, if the rare earth nanoparticles can be multiplexed into three different emission signals. The europium label greatly increased signal over FITC and Texas Red by 2-3 orders of magnitude.

Example 2 - Cell Response Assay for Breast Cancer

[0082] A novel cell response assay for breast cancer was developed by using HER2/neu as a tissue type marker for carcinoma breast cancer (Biomarker A) that is always present in the disease state. Biomarker B for metastatic potential utilized uPA and PAI antigen; these are markers of tumor invasiveness and are present in aggressive cancer cell differentiation and growth and increase with metastatic invasiveness potential of the cells. Biomarker C for chemo resistance utilized PL2L piwi like antigen (see for example, Gao et al., 2008).

[0083] The procedure shown in Example 1 was used to measure cancer cells before and after treatment with and without camptothecin. Camptothecin is known to activate apoptosis in cancer cells and kill cancer cells like a chemotherapy agent (See Gupta, 1997). Cells were tested by the cell response assay and by a traditional prior art assay. The novel "cell response assay" utilized HER2/neu as Biomarker A for cell type, the tumor invasiveness markers uPA & PAI-1 as Biomarker B for metastatic potential, and the cancer stem cell marker PL2L piwi like as Biomarker C for chemo resistance nature. The traditional assay used EpCAM as a marker for epithelial tissue type and cytokeratin (CK) as a marker of cells of cancer origin. Additionally, CD45, a marker for white blood cells, was used to reduce false results, with CD45 positive results being excluded from the analysis. Both the traditional and cell response assays used DAPI to determine the presence of cell nuclei.

[0084] The camptothecin concentration could be adjusted to be sufficient to kill >80%, 60% and 40% of the cancer cells in cultures. Cellular analysis showed cancer cell count to decrease as the cell death % increased and phosphatidylserine flopping increased. Cellular analysis

showed biomarker B, namely uPA and PA1, to increase (or was detected in a greater number of cancer cells) when the cell death % was highest. Cellular analysis showed biomarker C, namely PL2L piwi like to decrease (or was detected in a lesser number of cancer cells) when the cell death % was highest. All three biomarkers co-exist in the disease state and were fundamental to progression or relapse of the disease. This analysis of the specific combination of biomarkers described and claimed herein allows the determination of the cellular response to treatments by simultaneously measuring cancer cell type, chemo resistance nature, and metastatic potential (with or without the additional measurement of cell viability), and is herein termed a "cell based response assay".

[0085] In contrast, in the traditional assay, the CD45 biomarker does not co-exist with EpCAM and CK. Both EpCAM and CK were present in all cells independent of the % of cells that were killed.

EXAMPLE 3 - MULTIPLEXED ASSAY FOR PROSTATE CANCER

[0086] The following procedure shows the multiplexing of Biomarkers A (cancer cell type), B (metastatic potential) and C (chemo resistance) using a unique combination of affinity reagents with fluorescent labels and an affinity label to provide a multiplexed assay for prostate cancer. This allows the simultaneous measurement of cancer cell type, chemo resistance nature, and metastatic potential, as described herein above, in combination with a marker indicating the presence of cell nucleus (utilizing a nucleic acid binding probe). The data demonstrated that four simultaneous signals can be measured with this combination but not with the traditional combinations of the prior art. The presently disclosed and claimed inventive concept(s) has the potential to measure up to eight signals utilizing various fluorophores taught herein.

[0087] The procedure to test the method was to collect 4 mL normal whole blood fresh into a VACUTAINER® tube (BD 6 mL VACUTAINER® K2EDTA tube, 10.8 mg, ref 367863, Becton Dickinson, Oakville, ON) and transfer the blood into a round centrifuge tube sealed with top. Cancer cells were added by 40 µL of prostate cell cancer suspension to obtain a range of 3000 to 8000 cells/mL (8 cells/µL or 1-2 per high powered filed HRP (HRP=0.2 µL)). The blood was lysed by adding 3 mL of lysis buffer (155mM NH₄Cl, 20mM KHCO₃, 0.1 mM Na₂EDTA, pH 7.4). The tubes were rolled tubes to assure that the cells were evenly distributed and that lysis was complete, and then the sample was spun down for 15 minutes at 12,000 rpm. Next, the plasma was decanted as waste supernatant, and 4.0 mL of TES buffer (50 mM N-Tris [Hydroxymethyl]-2-aminoethane-sulfonic acid: TES, pH 7.4, 2150mM NaCl, 2 mM MgSO₄ and 1 mM KCO₃) and 1 % albumin was added, and the tube was rocked to mix the cells. The tube was spun down again for 15 minutes at 12,000 rpm, and the supernatant was decanted as waste.

A final 2.0 mL of TES wash buffer was added with rocking to mix cells.

[0088] The cells were then reacted with the reagents by taking 0.5 mL of washed blood to the centrifuge tube, and 40 µL of antibodies for PSA labeled with Cy5 and 40 µL of antibodies for urokinase like plasminogen activator (uPA) or plasminogen activator inhibitor (PAI-1) labeled with Texas Red (at working stock of 0.001 to 0.1 mg/mL) and PL2L piwi like labeled with FITC (at working stock of 0.001 to 0.1 mg/mL) were added. Next, 10 µl of a 100 µg/ml 4,6 Diamidino-2-phenylindole dihydrochloride (DAPI; a nucleic acid binding probe) was added from a solution in TES buffer. This was followed by incubation for 15 minutes at 37°C.

[0089] The sample was centrifuged down for 5 minutes @ 7500 rpm, and the supernatant was decanted off as waste (~450 µL). 500 µL of TES buffer was then added, and the sample was vortexed and centrifuged down at 5 minutes @ 7500 rpm. Again, the supernatant was decanted off by tapping as waste (~450 µL). A final 500 µL of TES wash buffer was added for final imaging, and 5 µL of the solution was placed on a slide with a long cover slip of glass. Phase contrast and fluorescence microscopy was conducted with the Leica DM5000 (Leica Microsystems, Buffalo Grove, IL).

[0090] The Texas Red label (ex 592, em 614 nm) was measured with a excitation band pass filter at 540-580 nm and emission band pass filter at 610-680 nm. The FITC label (ex 488, em 525nm) was measured with an excitation band pass filter at 460 - 500 nm and an emission band pass filter at 512 -543 nm. The DAPI probe (ex 355, em 460 nm) was measured with an excitation band pass filter at 340-380 nm and an emission band pass filter at 450-490 nm. The Cy5 label (ex 646, em 676 nm) was measured with an excitation band pass filter at 590-650 nm and an emission band pass filter at 665-735 nm.

[0091] The ideal multiplexing was demonstrated by combination of the rare earth nanoparticles with FITC, Texas Red, Cy5 and DAPI. The europium nanoparticle (ex 355, em 617 nm) was measured at an excitation filter band pass at 340-380 nm and emission long pass filter of > 590 nm. Simultaneously measuring biomarkers for cancer cell type, metastatic potential and chemoresistance, along with a biomarker indicating the presence of cell nucleus was possible, along with the removal of false positives (i.e., white blood cells) by detecting the presence of the biomarker CD45. Current data allows for five signals. However, the assays/methods described and claimed herein have the potential for detection and measurement of six signals/biomarkers, if the rare earth nanoparticles can be multiplexed into three different emission signals. The europium label greatly increased signal over FITC and Texas Red by 2-3 orders of magnitude.

EXAMPLE 4 - CELL RESPONSE ASSAY FOR PROSTATE CANCER

[0092] Protease, like plasmin, is expressed during metastasis of malignant cells as part of the tissue regeneration and fibrinolysis process, indirectly promoting cell proliferation. Cancer cells use cell-bound plasmin to activate the plasminogen signaling for urokinase. Inhibitors of plasmin such as Bikunin and inhibitors of plasminogen activation such as PAI-1 prevent cell-bound plasmin activation and suppress tumor invasiveness.

[0093] A novel cell response assay for prostate cancer was developed by using PSA as a tissue type marker for prostate carcinoma (Biomarker A) that is always present in the disease state. The markers used in this example for metastatic potential (Biomarker B) were urokinase plasminogen activator (uPA) and plasminogen activator inhibitor (PAI-1), which is a measure of tissue invasiveness and is present in an aggressive cancer cell differentiation and growth with increased metastatic invasiveness potential. Chemo resistance nature (Biomarker C) was determined using PL2L piwi like antigen to determine the inhibitory response to tissue invasiveness.

[0094] The procedure shown in Example 3 was used to measure prostate cancer cells before and after treatment with and without camptothecin. Camptothecin is known to activate apoptosis in cancer cells and kill cancer cells like a chemotherapy agent (See Gupta, 1997). Cells were tested by the cell response assay and by a traditional prior art assay. The novel "cell response assay" utilized PSA as Biomarker A for cell type, the tumor invasiveness markers uPA & PAI-1 as Biomarker B for metastatic potential, and the cancer stem cell marker PL2L piwi like as Biomarker C for chemo resistance nature. The traditional assay used EpCAM as a marker for epithelial tissue type and cytokeratin (CK) as a marker of cells of cancer origin. Additionally, CD45, a marker for white blood cells, was used to reduce false results, with CD45 positive results being excluded from the analysis. Both the traditional and cell response assays used DAPI to determine the presence of cell nuclei.

[0095] The camptothecin concentration could be adjusted to be sufficient to kill >80%, 60% and 40% of the cancer cells in cultures. Cellular analysis showed cancer cell count to decrease as the cell death % increased and phosphatidylserine flopping increased. Cellular analysis showed biomarker B, namely uPA and PA1, to increase (or was detected in a greater number of cancer cells) when the cell death % was highest. Cellular analysis showed biomarker C, namely PL2L piwi like to decrease (or was detected in a lesser number of cancer cells) when the cell death % was highest. All three biomarkers co-exist in the disease state and were fundamental to progression or relapse of the disease. This analysis of the specific combination of biomarkers described and claimed herein allows the determination of the cellular response to treatments by simultaneously measuring cancer cell type, chemo resistance nature, and metastat-

ic potential (with or without the additional measurement of cell viability), and is herein termed a "cell based response assay".

[0096] In contrast, in the traditional assay, the CD45 biomarker does not co-exist with EpCAM and CK. Both EpCAM and CK were present in all cells independent of the % of cells that were killed.

EXAMPLE 5

[0097] The following procedure demonstrates the multiplexing of Biomarkers A, B and C as described herein previously to provide a general assay for all types of cancers. This procedure is performed as described herein above in Examples 1/3, except as follows: (a) the cancer suspension added to the procedure is selected from one of the following cell lines - SKBR, MCF or MDA (breast cancer), UC3 or T24 (bladder cancer), PC3-9 (prostate cancer), and A549 (lung cancer); and (b) the non-specific cancer cell biomarker cytokeratin is utilized for Biomarker A. In this particular example, antibodies to the cytokeratins 8/18/19 (wherein said antibodies are labeled with Cy5) are utilized as the first labeled probe that binds to Biomarker A. As described in Examples 1 and 3, antibodies to uPA or PAI-1 labeled with Texas Ted are utilized as the second labeled probe that binds to Biomarker B, and antibodies to PL2L piwi like labeled with FITC is utilized as the third labeled probe that binds to Biomarker C. The nucleic acid binding probe DAPI is also utilized as described in Examples 1 and 3.

[0098] This multiplexed assay utilizing a non-specific cancer cell biomarker allows the assays/methods described herein to be adapted for use with all types of cancers.

[0099] Thus, in accordance with the present invention, there has been provided a cell response assay, as well as methods of producing and using same, that fully satisfies the objectives and advantages set forth hereinabove. Although the invention has been described in conjunction with the specific drawings, experimentation, results and language set forth hereinabove, it is evident that many alternatives, modifications, and variations will be apparent to those skilled in the art.

Claims

1. A cell response assay for cancer, comprising the steps of:
 - isolating cancer cells from a biological sample of a cancer patient;
 - measuring at least one cancer cell type biomarker on the surface of or inside of the cell utilizing a first labeled probe that binds to said cancer cell type biomarker, wherein the at least one cancer cell type biomarker comprises at least one of:

- (a) a combination of at least one type I cyto-keratin and at least one type II cyto-keratin;
- (b) a combination of vimentin and galectin-3; and
- (c) a combination of N-cadherin and E-cad-herin; or
- (d) an oncoprotein/oncogene selected from the group consisting of HER2/neu, VEGF-165, KRAS, EGFr, WAF, BAX-1, PDGF, Rb, Jagged 1, Notch, VEGF, VEGHR, k-Ras, CAIX, MIB1, MDM, PR, ER, SEL5, SEM1, PI3K, Akt2, twist 1, EML-4, ALK, Braf, DRAFF, c-met, and combinations thereof;

- measuring at least one chemo resistance bi-omarker on the surface of or inside of the cell utilizing a second labeled probe that binds to said chemo resistance biomarker, wherein the at least one chemo resistance biomarker is se-lected from the group consisting of PL2L piwi like, ADLH, b-integrin, a6 integrin, c-kit, c-met, LIF-R, CXCR4, ESA, CD 20, CD44, CD133, CK5, TRAF2, ABC transporters and combina-tions thereof;

- measuring at least one metastatic poten-tial bi-omarker on the surface of or inside of the cell utilizing a third labeled probe that binds to said metastatic potential biomarker, wherein the at least one metastatic potential biomarker is se-lected from the group consisting of

- urokinase plasminogen activator (uPA), plasminogen activator inhibitor (PAI-1), CD95, a serine protease, a serine protease inhibitor, a matrix metalloproteinase, a ma-trix metalloproteinase inhibitor, and combi-nations thereof; and

wherein the at least three labeled probes are meas-ured at different excitations and emission wave-lengths.

2. The assay of claim 1, wherein the assay is for a can-cer selected from the group consisting of lung, bron-chus, colon, rectum, pancreas, prostate, breast, liv-er, bile duct, bladder, ovary, brain, central nervous system (CNS), kidney, pelvis, uterine corpus, oral cavity, pharynx, melanoma, and combinations there-of.
3. The assay of claim 1, wherein at least one of:
 - (a) the assay is for prostate cancer, and the at least one cancer cell type biomarker comprises at least one of prostate specific antigen (PSA), prostate specific membrane antigen (PSMA) and combinations thereof;
 - (b) the assay is for breast cancer, and the at

- least one cancer cell type biomarker comprises at least one of MUC1, CA 15-3, CA 27-29 and combinations thereof;
- (c) the assay is for colon cancer, and the at least one cancer cell type biomarker comprises at least one of Carcinoembryonic Antigen (CEA), CA19-9, Galactosyl Transferase II and combi-nations thereof;
- (d) the assay is for pancreatic cancer, and the at least one cancer cell type biomarker compris-es MSLN (mesothelin);
- (e) the assay is for ovarian cancer, and the at least one cancer cell type biomarker comprises at least one of CA 125, Follicle-Stimulating Hor-mone (FSH) receptor and combinations thereof;
- (f) the assay is for liver cancer, and the at least one cancer cell type biomarker comprises Al-pha-Fetoprotein;
- (g) the assay is for melanoma, and the at least one cancer cell type biomarker comprises at least one of Melan-A (MLANA), Tyrosinase (TYR), CSPG4, MITF and combinations thereof; and
- (h) the assay is for thyroid cancer, and the at least one cancer cell type biomarker comprises at least one of Parathyroid related protein (PTHrP), TSHR and combinations thereof.

4. The assay of claim 1, wherein the at least one chemo resistance biomarker comprises at least one of:

- (a) presence of CD44 and absence of CD24;
- (b) presence of CD34 and absence of CD45 and CD31;
- (c) presence of CD44, CD24 and ESA; and
- (d) presence of CD24 and ESA.

5. The assay of claim 1, wherein at least one of:

- (a) the serine protease is selected from the group consisting of plasmin, ADAM, and combi-nations thereof;
- (b) the serine protease inhibitor comprises Bikunin;
- (c) the matrix metalloproteinase comprises MMP9; and
- (d) the matrix metalloproteinase inhibitors com-prises TIMP-1.

6. The assay of claim 1, wherein at least one of the first, second and third labeled probes comprise at least one labeled antibody to a biomarker.

7. The assay of claim 1, wherein the labels of the first, second and third labeled probes are selected from the group consisting of

- fluorescein-5-isothiocyanate (FITC), phyco-

erythrin, sulforhodamine 101 (Texas Red), 2-[4-(aminoiminomethyl)phenyl]-1H-Indole-6-carboximidamide (DAPI), 3H-Indolium (Cy5), 1H-benz[e]indolium (Cy 5.5), 3H-Indolium (Cy 7), ALEXA FLUOR®488, ALEXA FLUOR®555, ALEXA FLUOR®647, rare earth metals, rare earth element-containing nanoparticles, and combinations and derivatives thereof.

8. The assay of claim 1, wherein the biological sample is further defined as a tissue sample.

9. The assay of claim 1, further comprising measuring at least one additional biomarker in the biological sample utilizing a fourth labeled probe that binds to said additional biomarker.

10. The assay of claim 9, wherein the at least one additional biomarker comprises at least one white blood cell biomarker selected from the group consisting of

- CD45, CTLA-4, CD4, CD68, CD8, and combinations thereof,

and wherein the assay further comprises the step of excluding cells positive for at least one white blood cell biomarker.

11. The assay of claim 9, wherein the at least one additional biomarker comprises a biomarker indicating the presence of cell nuclei, and wherein the fourth labeled probe comprises 4',6-diamidino-2'-phenylindole, dihydrochloride (DAPI).

12. The assay of claim 9, wherein the at least one additional biomarker comprises phosphatidylserine, and wherein the fourth labeled probe comprises at least one of bis(zinc²⁺dipicolylamine) and PSVue™.

13. A method of monitoring cancer treatment in a cancer patient undergoing said treatment, the method comprising the steps of :

- measuring the levels of a cancer cell type biomarker in a first biological sample and in a second biological sample utilizing a first labeled probe that binds to said cancer cell type biomarker, wherein the first biological sample is obtained from the patient prior to exposure to a cancer treatment, and wherein the second biological sample is obtained from the patient following exposure to the cancer treatment, wherein the at least one cancer cell type biomarker comprises at least one of:

- (a) a combination of at least one type I cytokeratin and at least one type II cytokeratin;
- (b) a combination of vimentin and galectin-

- 3; and
- (c) a combination of N-cadherin and E-cadherin; or
- (d) an oncoprotein/oncogene selected from the group consisting of HER2/neu, VEGF-165, KRAS, EGFr, WAF, BAX-1, PDGF, Rb, Jagged 1, Notch, VEGF, VEGHR, k-Ras, CAIX, MIB1, MDM, PR, ER, SEL5, SEM1, PI3K, Akt2, twist 1, EML-4, ALK, Braf, DRAFF, c-met, and combinations thereof;

- measuring the levels of a chemo resistance biomarker in the first and second biological samples utilizing a second labeled probe that binds to said chemo resistance biomarker, wherein the at least one chemo resistance biomarker is selected from the group consisting of PL2L piwi like, ADLH, b-integrin, a6 integrin, c-kit, c-met, LIF-R, CXCR4, ESA, CD 20, CD44, CD133, CK5, TRAF2, ABC transporters and combinations thereof;

- measuring the levels of a metastatic potential biomarker in the first and second biological samples utilizing a third labeled probe that binds to said metastatic potential biomarker, wherein the at least one metastatic potential biomarker is selected from the group consisting of

- urokinase plasminogen activator (uPA), plasminogen activator inhibitor (PAI-1), CD95, a serine protease, a serine protease inhibitor, a matrix metalloproteinase, a matrix metalloproteinase inhibitor, and combinations thereof

and wherein the at least three labeled probes are measured at different excitations and emission wavelengths;

- comparing the levels of the chemo resistance biomarker in the cells to which the second labeled probe is bound in the first and second biological samples;

- comparing the levels of the metastatic potential biomarker in the cells to which the third labeled probe is bound in the first and second biological samples; and

- determining that the cancer treatment is effective if the level of the chemo resistance biomarker is decreased and the level of the metastatic potential biomarker is increased in the second biological sample when compared to the first biological sample.

14. A method of monitoring progression/relapse of cancer in a cancer patient, the method comprising the steps of :

- measuring the levels of a cancer cell type biomarker in a first biological sample and in a second biological sample utilizing a first labeled probe that binds to said cancer cell type biomarker, wherein the first biological sample is obtained from the patient at a first time point, and the second biological sample is obtained from the patient at a subsequent time point, wherein the at least one cancer cell type biomarker comprises at least one of:

- (a) a combination of at least one type I cytokeratin and at least one type II cytokeratin;
- (b) a combination of vimentin and galectin-3; and
- (c) a combination of N-cadherin and E-cadherin; or
- (d) an oncoprotein/oncogene selected from the group consisting of HER2/neu, VEGF-165, KRAS, EGFr, WAF, BAX-1, PDGF, Rb, Jagged 1, Notch, VEGF, VEGHR, k-Ras, CAIX, MIB1, MDM, PR, ER, SEL5, SEM1, PI3K, Akt2, twist 1, EML-4, ALK, Braf, DRAFF, c-met, and combinations thereof;

- measuring the levels of a chemo resistance biomarker in the first and second biological samples utilizing a second labeled probe that binds to said chemo resistance biomarker, wherein the at least one chemo resistance biomarker is selected from the group consisting of PL2L piwi like, ADLH, b-integrin, a6 integrin, c-kit, c-met, LIF-R, CXCR4, ESA, CD 20, CD44, CD133, CK5, TRAF2, ABC transporters and combinations thereof;

- measuring the levels of a metastatic potential biomarker in the first and second biological samples utilizing a third labeled probe that binds to said metastatic potential biomarker, wherein the at least one metastatic potential biomarker is selected from the group consisting of

- urokinase plasminogen activator (uPA), plasminogen activator inhibitor (PAI-1), CD95, a serine protease, a serine protease inhibitor, a matrix metalloproteinase, a matrix metalloproteinase inhibitor, and combinations thereof

and wherein the at least three labeled probes are measured at different excitations and emission wavelengths;

- comparing the levels of the chemo resistance biomarker in the cells to which the second labeled probe is bound in the first and second biological samples;
- comparing the levels of the metastatic potential

biomarker in the cells to which the third labeled probe is bound in the first and second biological samples; and

- determining that the cancer has progressed/relapsed if the level of the chemo resistance biomarker is increased and the level of the metastatic potential biomarker is decreased in the second biological sample when compared to the first biological sample.

Patentansprüche

1. Zellreaktionstest für Krebs, umfassend die Schritte:

- Isolieren von Krebszellen aus einer biologischen Probe eines Krebspatienten;
- Messen mindestens eines Biomarkers für einen Krebszelltyp auf der Oberfläche von oder innerhalb der Zelle unter Verwendung einer ersten markierten Sonde, die an den Biomarker für einen Krebszelltyp bindet, wobei der mindestens eine Biomarker für einen Krebszelltyp mindestens einen aus:

- (a) einer Kombination aus mindestens einem Typ-I-Zytokeratin und mindestens einem Typ-II-Zytokeratin;
- (b) einer Kombination aus Vimentin und Galectin-3; und
- (c) einer Kombination aus N-Cadherin und E-Cadherin; oder
- (d) einem Onkoprotein/Onkogen, ausgewählt aus der Gruppe bestehend aus HER2/neu, VEGF-165, KRAS, EGFr, WAF, BAX-1, PDGF, Rb, Jagged 1, Notch, VEGF, VEGHR, k-Ras, CAIX, MIB1, MDM, PR, ER, SEL5, SEM1, PI3K, Akt2, twist 1, EML-4, ALK, Braf, DRAFF, c-met und Kombinationen davon umfasst;

- Messen mindestens eines Biomarkers für Chemoresistenz auf der Oberfläche von oder innerhalb der Zelle unter Verwendung einer zweiten markierten Sonde, die an den Biomarker für Chemoresistenz bindet, wobei der mindestens eine Biomarker für Chemoresistenz ausgewählt ist aus der Gruppe bestehend aus PL2L-piwi-like, ADLH, b-Integrin, a6-Integrin, c-kit, c-met, LIF-R, CXCR4, ESA, CD 20, CD44, CD133, CK5, TRAF2, ABC-Transportern und Kombinationen davon;

- Messen des mindestens einen Biomarkers für Metastasierungspotential auf der Oberfläche von oder innerhalb der Zelle unter Verwendung einer dritten markierten Sonde, die an den Biomarker für Metastasierungspotential bindet, wobei der mindestens eine Biomarker für Metasta-

sierungspotential ausgewählt ist aus der Gruppe bestehend aus

- Urokinase-Plasminogen-Aktivator (uPA), Plasminogen-Aktivator-Inhibitor (PAI-1), CD95, einer Serinprotease, einem Serinproteaseinhibitor, einer Matrixmetalloproteinase, einem Matrixmetalloproteinaseinhibitor und Kombinationen davon; und

wobei die mindestens drei markierten Sonden bei verschiedenen Anregungs- und Emissionswellenlängen gemessen werden.

2. Test nach Anspruch 1, wobei der Test für einen Krebs ist, ausgewählt aus der Gruppe bestehend aus Lungen-, Bronchial-, Darm-, Mastdarm-, Bauchspeicheldrüsen-, Prostata-, Brust-, Leber-, Gallengang-, Blasen-, Eiserstock-, Gehirn-, Zentralnervensystem(ZNS)-, Nieren, Becken-, Gebärmutter-, Mundhöhlen-, Rachenkrebs, Melanom und Kombinationen davon.

3. Test nach Anspruch 1, wobei mindestens einer:

(a) der Tests für Prostatakrebs ist, und der mindestens eine Biomarker für einen Krebszelltyp mindestens einen aus einem prostataspezifischen Antigen (PSA), prostataspezifischen Membranantigen (PSMA) und Kombinationen davon umfasst;

(b) der Tests für Brustkrebs ist, und der mindestens eine Biomarker für einen Krebszelltyp mindestens einen aus MUC1, CA 15-3, CA 27-29 und Kombinationen davon umfasst;

(c) der Test für Darmkrebs ist, und der mindestens eine Biomarker für einen Krebszelltyp mindestens einen aus karzinoembryonalem Antigen (CEA), CA19-9, Galactosyltransferase II und Kombinationen davon umfasst;

(d) der Test für Bauchspeicheldrüsenkrebs ist, und der mindestens eine Biomarker für einen Krebszelltyp MSLN (Mesothelin) umfasst;

(e) der Test für Eierstockkrebs ist, und der mindestens eine Biomarker für einen Krebszelltyp mindestens einen aus CA 125, follikelstimulierendes-Hormon(FSH)-Rezeptor und Kombinationen davon umfasst;

(f) der Test für Leberkrebs ist, und der mindestens eine Biomarker für einen Krebszelltyp Alpha-Fetoprotein umfasst;

(g) der Test für Melanom ist, und der mindestens eine Biomarker für einen Krebszelltyp mindestens einen aus Melan-A (MLANA), Tyrosinase (TYR), CSPG4, MITF und Kombinationen davon umfasst; und

(h) der Test für Schilddrüsenkrebs ist, und der mindestens eine Biomarker für einen Krebszell-

typ mindestens einen aus Parathyroid-related-Protein (PTHrP), TSHR und Kombinationen davon umfasst.

4. Test nach Anspruch 1, wobei der mindestens eine Biomarker für Chemoresistenz mindestens einen aus:

(a) Gegenwart von CD44 und Abwesenheit von CD24;

(b) Gegenwart von CD34 und Abwesenheit von CD45 und CD31;

(c) Gegenwart von CD44, CD24 und ESA; und

(d) Gegenwart von CD24 und ESA umfasst.

5. Test nach Anspruch 1, wobei mindestens eine/einer:

(a) der Serinproteasen ausgewählt ist aus der Gruppe bestehend aus Plasmin, ADAM und Kombinationen davon;

(b) der Serinproteaseinhibitoren Bikunin umfasst;

(c) der Matrixmetalloproteinasen MMP9 umfasst; und

(d) der Matrixmetalloproteinaseinhibitoren TIMP-1 umfasst.

6. Test nach Anspruch 1, wobei mindestens eine der ersten, zweiten und dritten markierten Sonden mindestens einen markierten Antikörper gegen einen Biomarker umfassen.

7. Test nach Anspruch 1, wobei die Marker der ersten, zweiten und dritten markierten Sonden ausgewählt sind aus der Gruppe bestehend aus

F fluorescein-5-isothiocyanat (FITC), Phycoerythrin, Sulforhodamin 101 (Texasrot), 2-[4-(Aminoiminomethyl)phenyl]-1H-Indol-6-carboximidamid (DAPI), 3H-Indol (Cy5), 1H-Benz[e]indol (Cy 5.5), 3H-Indol (Cy7), ALEXA FLUOR®488, ALEXA FLUOR®555, ALEXA FLUOR®647, Seltenerdmetalle, Seltenerdelemententhaltende Nanopartikel und Kombinationen und Derivate davon.

8. Test nach Anspruch 1, wobei die biologische Probe ferner als Gewebeprobe definiert ist.

9. Test nach Anspruch 1, ferner umfassend das Messen mindestens eines zusätzlichen Biomarkers in der biologischen Probe unter Verwendung einer vierten markierten Sonde, die an den zusätzlichen Biomarker bindet.

10. Test nach Anspruch 9, wobei der mindestens eine zusätzliche Biomarker mindestens einen Biomarker für weiße Blutzellen umfasst, ausgewählt aus der

Gruppe bestehend aus

- CD45, CTLA-4, CD4, CD68, CD8 und Kombinationen davon,

und wobei der Test ferner den Schritt zum Ausschließen von Zellen umfasst, die für mindestens einen Biomarker für weiße Blutzellen positiv sind.

11. Test nach Anspruch 9, wobei der mindestens eine zusätzliche Biomarker einen Biomarker umfasst, der die Gegenwart von Zellkernen anzeigt, und wobei die vierte markierte Sonde 4',6-Diamidino-2'-phenylindol-Dihydrochlorid (DAPI) umfasst.

12. Test nach Anspruch 9, wobei der mindestens eine zusätzliche Biomarker Phosphatidylserin umfasst, und wobei die vierte markierte Sonde mindestens eine aus Bis (zink²⁺dipicolylamin) und PSVue™ umfasst.

13. Verfahren zum Überwachen der Krebsbehandlung bei einem Krebspatienten, der sich der Behandlung unterzieht, wobei das Verfahren die folgenden Schritte umfasst:

- Messen der Konzentrationen eines Biomarkers für einen Krebszelltyp in einer ersten biologischen Probe und in einer zweiten biologischen Probe unter Verwendung einer ersten markierten Sonde, die an den Biomarker für einen Krebszelltyp bindet, wobei die erste biologische Probe von dem Patienten erhalten wird, bevor er sich einer Krebsbehandlung unterzieht, wobei die zweite biologische Probe von dem Patienten erhalten wird, nachdem er sich der Krebsbehandlung unterzogen hat, wobei der mindestens eine Biomarker für einen Krebszelltyp mindestens einen aus:

(a) einer Kombination aus mindestens einem Typ-I-Zytokeratin und mindestens einem Typ-II-Zytokeratin;

(b) einer Kombination aus Vimentin und Galectin-3; und

(c) einer Kombination aus N-Cadherin und E-Cadherin; oder

(d) einem Onkoprotein/Onkogen, ausgewählt aus der Gruppe bestehend aus HER2/neu, VEGF-165, KRAS, EGFr, WAF, BAX-1, PDGF, Rb, Jagged 1, Notch, VEGF, VEGHR, k-Ras, CAIX, MIB1, MDM, PR, ER, SEL5, SEM1, PI3K, Akt2, twist 1, EML-4, ALK, Braf, DRAFF, c-met und Kombinationen davon umfasst;

- Messen der Konzentrationen eines Biomarkers für Chemoresistenz in der ersten und der

zweiten biologischen Probe unter Verwendung einer zweiten Sonde, die an den Biomarker für Chemoresistenz bindet, wobei der mindestens eine Biomarker für Chemoresistenz ausgewählt ist aus der Gruppe bestehend aus PL2L-piwi-like, ADLH, b-Integrin, a6-Integrin, c-kit, c-met, LIF-R, CXCR4, ESA, CD 20, CD44, CD133, CK5, TRAF2, ABC-Transportern und Kombinationen davon;

- Messen der Konzentrationen eines Biomarkers für Metastasierungspotential in der ersten und der zweiten biologischen Probe unter Verwendung einer dritten Sonde, die an den Biomarker für Metastasierungspotential bindet, wobei der mindestens eine Biomarker für Metastasierungspotential ausgewählt ist aus der Gruppe bestehend aus

- Urokinase-Plasminogen-Aktivator (uPA), Plasminogen-Aktivator-Inhibitor (PAI-1), CD95, einer Serinprotease, einem Serinproteaseinhibitor, einer Matrixmetalloproteinase, einem Matrixmetalloproteinaseinhibitor und Kombinationen davon,

und wobei die mindestens drei markierten Sonden bei verschiedenen Anregungs- und Emissionswellenlängen gemessen werden;

- Vergleichen der Konzentrationen des Biomarkers für Chemoresistenz in den Zellen, an den die zweite markierte Sonde in der ersten und der zweiten biologischen Probe gebunden ist;

- Vergleichen der Konzentrationen des Biomarkers für Metastasierungspotential in den Zellen, an den die dritte markierte Sonde in der ersten und der zweiten biologischen Probe gebunden ist; und

- Bestimmen, dass die Krebsbehandlung wirksam ist, wenn die Konzentration des Biomarkers für Chemoresistenz verringert ist und die Konzentration des Biomarkers für Metastasierungspotential in der zweiten biologischen Probe im Vergleich zur ersten biologischen Probe erhöht ist.

14. Verfahren zum Überwachen des Fortschreitens/der Wiedererkrankung bei einem Krebspatienten, wobei das Verfahren die folgenden Schritte umfasst:

- Messen der Konzentrationen eines Biomarkers für einen Krebszelltyp in einer ersten biologischen Probe und in einer zweiten biologischen Probe unter Verwendung einer ersten markierten Sonde, die an den Biomarker für einen Krebszelltyp bindet, wobei die erste biologische Probe zu einem ersten Zeitpunkt von dem Patienten erhalten wird und die zweite bi-

ologische Probe zu einem nachfolgenden Zeitpunkt von dem Patienten erhalten wird, wobei der mindestens eine Biomarker für einen Krebszelltyp mindestens einen aus:

- (a) einer Kombination aus mindestens einem Typ-I-Zytokeratin und mindestens einem Typ-II-Zytokeratin;
- (b) einer Kombination aus Vimentin und Galectin-3; und
- (c) einer Kombination aus N-Cadherin und E-Cadherin; oder
- (d) einem Onkoprotein/Onkogen, ausgewählt aus der Gruppe bestehend aus HER2/neu, VEGF-165, KRAS, EGFr, WAF, BAX-1, PDGF, Rb, Jagged 1, Notch, VEGF, VEGHR, k-Ras, CAIX, MIB1, MDM, PR, ER, SEL5, SEM1, PI3K, Akt2, twist 1, EML-4, ALK, Braf, DRAFF, c-met und Kombinationen davon umfasst;

- Messen der Konzentrationen eines Biomarkers für Chemoresistenz in der ersten und der zweiten biologischen Probe unter Verwendung einer zweiten Sonde, die an den Biomarker für Chemoresistenz bindet, wobei der mindestens eine Biomarker für Chemoresistenz ausgewählt ist aus der Gruppe bestehend aus PL2L-piwi-like, ADLH, b-Integrin, a6-Integrin, c-kit, c-met, LIF-R, CXCR4, ESA, CD 20, CD44, CD133, CK5, TRAF2, ABC-Transportern und Kombinationen davon;

- Messen der Konzentrationen eines Biomarkers für Metastasierungspotential in der ersten und der zweiten biologischen Probe unter Verwendung einer dritten Sonde, die an den Biomarker für Metastasierungspotential bindet, wobei der mindestens eine Biomarker für Metastasierungspotential ausgewählt ist aus der Gruppe bestehend aus

- Urokinase-Plasminogen-Aktivator (uPA), Plasminogen-Aktivator-Inhibitor (PAI-1), CD95, einer Serinprotease, einem Serinproteaseinhibitor, einer Matrixmetalloproteinase, einem Matrixmetalloproteinaseinhibitor und Kombinationen davon,

und wobei die mindestens drei markierten Sonden bei verschiedenen Anregungs- und Emissionswellenlängen gemessen werden;

- Vergleichen der Konzentrationen des Biomarkers für Chemoresistenz in den Zellen, an den die erste markierte Sonde in der ersten und der zweiten biologischen Probe gebunden ist;
- Vergleichen der Konzentrationen des Biomarkers für Metastasierungspotential in den Zellen,

an den die dritte markierte Sonde in der ersten und der zweiten biologischen Probe gebunden ist; und

- Bestimmen, dass der Krebs fortgeschritten/zurückgekehrt ist, wenn die Konzentration des Biomarkers für Chemoresistenz erhöht ist und die Konzentration des Biomarkers für Metastasierungspotential in der zweiten biologischen Probe im Vergleich zur ersten biologischen Probe verringert ist.

Revendications

1. Dosage de sensibilité cellulaire pour un cancer, comprenant les étapes consistant à :

- isoler des cellules cancéreuses d'un échantillon biologique d'un patient cancéreux;
- mesurer au moins un biomarqueur de type cellule cancéreuse à la surface ou à l'intérieur de la cellule en utilisant une première sonde marquée qui se lie audit biomarqueur de type cellule cancéreuse, où ledit au moins un biomarqueur de type cellule cancéreuse comprend au moins l'un des éléments suivants :

- (a) une combinaison d'au moins une cyto-kératine de type I et d'au moins une cyto-kératine de type II ;
- (b) une combinaison de vimentine et de galectine-3 ; et
- (c) une combinaison de N-cadhérine et de E-cadhérine ; ou
- (d) une oncoprotéine/oncogène choisie dans le groupe constitué par HER2/neu, VEGF-165, KRAS, EGFr, WAF, BAX-1, PDGF, Rb, Jagged 1, Notch, VEGF, VEGHR, k-Ras, CAIX, MIB1, MDM, PR, ER, SEL5, SEM1, PI3K, Akt2, twist 1, EML-4, ALK, Braf, DRAFF, c-met, et des combinaisons de celles-ci ;

- mesurer au moins un biomarqueur de chimio-résistance sur la surface de ou à l'intérieur de la cellule en utilisant une deuxième sonde marquée qui se lie audit biomarqueur de chimio-résistance, dans lequel ledit moins un biomarqueur de chimio-résistance est choisi dans le groupe constitué de PL2L de type piwi, ADLH, b-intégrin, a6 intégrin, c-kit, c-met, LIF-R, CXCR4, ESA, CD 20, CD44, CD133, CK5, TRAF2, transporteurs ABC et de leurs combinaisons ;

- mesurer au moins un biomarqueur de potentiel métastatique à la surface ou à l'intérieur de la cellule en utilisant une troisième sonde marquée qui se lie audit biomarqueur de potentiel métas-

tatique, dans lequel ledit au moins un biomarqueur de potentiel métastatique est choisi dans le groupe constitué de :

- activateur plasminogène urokinase (uPA),
- inhibiteur activateur de plasminogène (PAI-1),
- CD95, une protéase à sérine, un inhibiteur de protéase à sérine, une métalloprotéinase matricielle, un inhibiteur de métalloprotéinase matricielle et leurs combinaisons ; et

où lesdites au moins trois sondes marquées sont mesurées à des des longueurs d'onde d'émission et d'excitation différentes.

2. Dosage selon la revendication 1, où le dosage porte sur un cancer choisi dans le groupe constitué de poumon, bronche, côlon, rectum, pancréas, prostate, sein, foie, canal biliaire, vessie, ovaire, cerveau, système nerveux central (CNS), rein, bassin, corps utérin, cavité orale, pharynx, mélanome, et de leurs combinaisons.

3. Dosage selon la revendication 1, où au moins :

- (a) un dosage est destiné au cancer de la prostate et ledit au moins un biomarqueur de type de cellule cancéreuse comprend au moins un élément parmi un antigène spécifique de prostate (PSA), un antigène de membrane spécifique de prostate (PSMA) et leur combinaisons ;
- (b) un dosage est destiné au cancer du sein et ledit au moins un biomarqueur de type de cellule cancéreuse comprend au moins un élément parmi MUC1, CA 15-3, CA 27-29 et leurs combinaisons ;
- (c) un dosage porte sur le cancer du côlon et ledit au moins un biomarqueur de type de cellules cancéreuses comprend au moins un élément parmi un antigènes Carcinoembryonnaire (CEA), un CA19-9, un Galactosyl Transferase II et leurs combinaisons ;
- (d) un dosage est destiné au cancer du pancréas et ledit au moins un biomarqueur de type de cellule cancéreuse comprend le MSLN (mésothéline) ;
- (e) un dosage est destiné au cancer de l'ovaire et ledit au moins un biomarqueur de type de cellules cancéreuse comprend au moins un élément parmi le CA 125, le récepteur d'Hormone Folliculo-Stimulante (FSH) et leurs combinaisons ;
- (f) un dosage porte sur le cancer du foie et ledit au moins un biomarqueur de type de cellule cancéreuse comprend l'Alpha-Foetoprotéine ;
- (g) un dosage porte sur le mélanome et ledit au moins un biomarqueur de type de cellule can-

céreuse comprend au moins un élément parmi le Melan-A (MLANA), le Tyrosinase (TYR), le CSPG4, le MITF et leurs combinaisons ; et
(h) un dosage est destiné au cancer de la thyroïde et ledit au moins un biomarqueur type de cellule cancéreuse comprend au moins un élément parmi une protéine apparentée à la parathyroïde (PTHrP), un TSHR et leurs combinaisons.

4. Dosage selon la revendication 1, où au moins un biomarqueur de chimio-résistance comprend au moins l'une des caractéristiques suivantes :

- (a) présence de CD44 et absence de CD24 ;
- (b) présence de CD34 et absence de CD45 et de CD31 ;
- (c) présence de CD44, CD24 et ESA ; et
- (d) présence de CD24 et de ESA.

5. Dosage selon la revendication 1, où au moins le dosage comprend l'une des caractéristiques suivantes :

- (a) la protéase à sérine est choisie dans le groupe constitué par la plasmine, l'ADAM et leurs combinaisons ;
- (b) l'inhibiteur de protéase à sérine comprend de la Bikunine ;
- (c) la métalloprotéinase matricielle comprend la MMP9 ; et
- (d) les inhibiteurs de métalloprotéinase matricielle comprennent du TIMP-1.

6. Dosage selon la revendication 1, où au moins une des première, deuxième et troisième sondes marquées comprend au moins un anticorps marqué contre un biomarqueur.

7. Dosage selon la revendication 1, où les étiquettes des première, deuxième et troisième sondes marquées sont choisies dans le groupe constitué de

- fluorescéine-5-isothiocyanate (FITC), phycoérythrine, sulforhodamine 101 (Texas Red), 2-[4-(aminoiminométhyl)phényl]-1H-Indole-6-carboximidamide (DAPI), 3H-Indolium (Cy5), 1H-benz[e]indolium (Cy 5.5), 3H-Indolium (Cy 7), ALEXA FLUOR®488, ALEXA FLUOR®555, ALEXA FLUOR®647, métaux des terres rares et nanoparticules contenant des éléments de terres rares et leurs combinaisons et dérivés.

8. Dosage selon la revendication 1, où l'échantillon biologique est en outre défini comme un échantillon de tissu.

9. Dosage selon la revendication 1, comprenant en

- outre la mesure d'au moins un biomarqueur supplémentaire dans l'échantillon biologique en utilisant une quatrième sonde marquée qui se lie audit biomarqueur supplémentaire.
- 5
10. Dosage selon la revendication 9, où au moins un biomarqueur supplémentaire comprend au moins un biomarqueur de globules blancs choisi dans le groupe constitué de
- 10
- CD45, CTLA-4, CD4, CD68, CD8 et leurs combinaisons,
- et où le dosage comprend en outre l'étape d'exclusion des cellules positives pour au moins un biomarqueur de globules blancs.
- 15
11. Dosage selon la revendication 9, où ledit au moins un biomarqueur supplémentaire comprend un biomarqueur indiquant la présence de noyaux cellulaires, et où la quatrième sonde marquée comprend du 4',6-diamidino-2'-phénylindole, dihydrochlorure (DAPI)
- 20
12. Dosage selon la revendication 9, où au moins un biomarqueur supplémentaire comprend de la phosphatidylsérine, et où la quatrième sonde marquée comprend au moins l'un de bis(zinc²⁺dipicolylamine) et PSVue™.
- 25
- 30
13. Procédé de contrôle du traitement du cancer chez un patient cancéreux subissant ledit traitement, le procédé comprenant les étapes consistant à :
- 35
- mesurer les niveaux d'un biomarqueur de type cellule cancéreuse dans un premier échantillon biologique et dans un deuxième échantillon biologique en utilisant une première sonde marquée qui se lie audit biomarqueur de type cellule cancéreuse, où ledit premier échantillon biologique est obtenu du patient avant exposition à un traitement du cancer, et où le deuxième échantillon biologique est obtenu du patient à la suite de l'exposition au traitement du cancer, où ledit au moins un biomarqueur de type cellule cancéreuse comprend au moins l'un des éléments suivants :
- 40
- (a) une combinaison d'au moins une cyto-kératine de type I et d'au moins une cyto-kératine de type II ;
- 45
- (b) une combinaison de vimentine et de galectine-3 ; et
- (c) une combinaison de N-cadhérine et de E-cadhérine ; ou
- 50
- (d) une oncoprotéine/oncogène choisie dans le groupe constitué par HER2/neu, VEGF-165, KRAS, EGFr, WAF, BAX-1, PDGF, Rb, Jagged 1, Notch, VEGF, VEGHR, k-Ras, CAIX, MIB1, MDM, PR, ER, SELS, SEM1, PI3K, Akt2, twist 1, EML-4, ALK, Braf, DRAFF, c-met, et des combinaisons de celles-ci ;
- mesurer les niveaux d'un biomarqueur de chimio-résistance sur le premier et le deuxième échantillons biologiques en utilisant une deuxième sonde marquée qui se lie audit biomarqueur de chimio-résistance, dans lequel ledit moins un biomarqueur de chimio-résistance est choisi dans le groupe constitué de PL2L de type piwi, ADLH, b-integrin, a6 integrin, c-kit, c-met, LIF-R, CXCR4, ESA, CD 20, CD44, CD133, CK5, TRAF2, transporteurs ABC et de leurs combinaisons ;
- mesurer les niveaux d'un biomarqueur de potentiel métastatique sur le premier et deuxième échantillon biologique en utilisant une troisième sonde marquée qui se lie audit biomarqueur de potentiel métastatique, dans lequel ledit au moins un biomarqueur de potentiel métastatique est choisi dans le groupe constitué de :
- activateur plasminogène urokinase (uPA), inhibiteur activateur de plasminogène (PAI-1), CD95, une protéase à sérine, un inhibiteur de protéase à sérine, une métalloprotéinase matricielle, un inhibiteur de métalloprotéinase matricielle et leurs combinaisons ;
- et où lesdites au moins trois sondes marquées sont mesurées à des longueurs d'onde d'émission et d'excitation différentes
- 55
- comparer les niveaux du biomarqueur de chimio-résistance dans les cellules auxquelles la deuxième sonde marquée est liée dans les premier et deuxième échantillons biologiques ;
- comparer les niveaux du biomarqueur de potentiel métastatique dans les cellules auxquelles la troisième sonde marquée est liée dans les premier et deuxième échantillons biologiques ; et
- déterminer que le traitement du cancer est efficace si le niveau du biomarqueur de chimio-résistance est réduit et que le niveau du biomarqueur potentiel métastatique est augmenté dans le deuxième échantillon biologique par rapport au premier échantillon biologique.
14. Procédé de contrôle du progrès/rechute du cancer dans un patient cancéreux, le procédé comprenant les étapes consistant à :
- mesurer les niveaux d'un biomarqueur de type cellule cancéreuse dans un premier échantillon biologique et dans un deuxième échantillon bio-

logique en utilisant une première sonde marquée qui se lie audit biomarqueur de type cellule cancéreuse, où ledit premier échantillon biologique est obtenu du patient à un premier instant, et le deuxième échantillon biologique est obtenu du patient à instant ultérieur, où ledit au moins un biomarqueur de type cellule cancéreuse comprend au moins l'un des éléments suivants :

- (a) une combinaison d'au moins une cyto-kératine de type I et d'au moins une cyto-kératine de type II ;
- (b) une combinaison de vimentine et de galectine-3 ; et
- (c) une combinaison de N-cadhérine et de E-cadhérine ; ou
- (d) une oncoprotéine/oncogène choisie dans le groupe constitué par HER2/neu, VEGF-165, KRAS, EGFr, WAF, BAX-1, PDGF, Rb, Jagged 1, Notch, VEGF, VEGHR, k-Ras, CAIX, MIB1, MDM, PR, ER, SELS, SEM1, PI3K, Akt2, twist 1, EML-4, ALK, Braf, DRAFF, c-met, et des combinaisons de celles-ci ;

- mesurer les niveaux d'un biomarqueur de chimio-résistance sur le premier et le deuxième échantillons biologiques en utilisant une deuxième sonde marquée qui se lie audit biomarqueur de chimio-résistance, dans lequel ledit moins un biomarqueur de chimio-résistance est choisi dans le groupe constitué de PL2L de type piwi, ADLH, b-integrin, a6 integrin, c-kit, c-met, LIF-R, CXCR4, ESA, CD 20, CD44, CD133, CK5, TRAF2, transporteurs ABC et de leurs combinaisons;

- mesurer les niveaux d'un biomarqueur de potentiel métastatique sur le premier et deuxième échantillon biologique en utilisant une troisième sonde marquée qui se lie audit biomarqueur de potentiel métastatique, dans lequel ledit au moins un biomarqueur de potentiel métastatique est choisi dans le groupe constitué de :

- activateur plasminogène urokinase (uPA), inhibiteur activateur de plasminogène (PAI-1), CD95, une protéase à sérine, un inhibiteur de protéase à sérine, une métalloprotéinase matricielle, un inhibiteur de métalloprotéinase matricielle et leurs combinaisons;

et où lesdites au moins trois sondes marquées sont mesurées à des longueurs d'onde d'émission et d'excitation différentes


- comparer les niveaux du biomarqueur de chimio-résistance dans les cellules auxquelles la

deuxième sonde marquée est liée dans les premier et deuxième échantillons biologiques ;

- comparer les niveaux du biomarqueur de potentiel métastatique dans les cellules auxquelles la troisième sonde marquée est liée dans les premier et deuxième échantillons biologiques ; et

- déterminer que le traitement du cancer a progressé/rechuté si le niveau du biomarqueur de chimio-résistance est augmenté et que le niveau du biomarqueur potentiel métastatique est réduit dans le deuxième échantillon biologique par rapport au premier échantillon biologique.

FIGURE 1
Cell Response Assay



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 61538302 A [0001]
- US 20080187198 A [0004]
- US 7179616 B [0013]
- US 7465747 B [0038]
- US 7955859 B, Matsumoto [0038]
- US 20070026407 A [0038]
- US 6037137 A [0039]
- US 6350466 B [0039]

Non-patent literature cited in the description

- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1989 [0020]
- **COLIGAN et al.** Current Protocols in Immunology. Current Protocols, Wiley Interscience, 1994 [0020]
- **HUDSON et al.** *Nature Med.*, 2003, vol. 9, 129-134 [0031]
- **RICHARD P. HAUGLAND.** Molecular Probes: Handbook of Fluorescent Probes and Research Chemicals 1992-1994. Molecular Probes, Inc, 1994 [0039]
- **HERMANSON.** Bioconjugate Techniques. Academic Press, 1995 [0039]
- **KAY M. et al.** *Biochemistry*, 1995, vol. 34, 293 [0039]
- **STUBBS et al.** *Biochemistry*, 1996, vol. 35, 937 [0039]
- Evaluating Receptor Stoichiometry by Fluorescence Resonance Energy Transfer. **GAKAMSKY D. et al.** Receptors: A Practical Approach. Oxford University Press, 2001 [0039]

专利名称(译)	癌症的细胞反应测定法及其制备和使用方法		
公开(公告)号	EP2748337B1	公开(公告)日	2019-08-07
申请号	EP2012833913	申请日	2012-09-21
[标]申请(专利权)人(译)	西门子医疗保健诊断公司		
申请(专利权)人(译)	西门子医疗诊断INC.		
当前申请(专利权)人(译)	西门子医疗诊断INC.		
[标]发明人	PUGIA MICHAEL		
发明人	PUGIA, MICHAEL		
IPC分类号	C12Q1/68 G01N33/53 G01N33/574		
CPC分类号	A61K31/4745 C12Q1/6886 C12Q2537/143 C12Q2565/102 C12Q2600/158 G01N33/57407 G01N2800/60 G01N33/57415 G01N33/57434 G01N33/5748 G01N33/57484 G01N2333/8132 G01N2333/82 G01N2333/91205 G01N2333/96463		
代理机构(译)	施韦泽KLAUS		
优先权	61/538302 2011-09-23 US		
其他公开文献	EP2748337A4 EP2748337A1		
外部链接	Espacenet		

摘要(译)

提供了癌症的细胞应答测定。在该测定中，在生物样品中同时测量癌细胞类型生物标志物，化学抗性生物标志物和转移潜在生物标志物的水平。

