



(11) **EP 2 305 807 A1**

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 153(4) EPC

(43) Date of publication:
06.04.2011 Bulletin 2011/14

(51) Int Cl.:
C12N 15/09 ^(2006.01) **C12Q 1/25** ^(2006.01)
C12Q 1/68 ^(2006.01) **G01N 33/53** ^(2006.01)

(21) Application number: **09762576.8**

(86) International application number:
PCT/JP2009/061068

(22) Date of filing: **11.06.2009**

(87) International publication number:
WO 2009/151150 (17.12.2009 Gazette 2009/51)

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK TR
Designated Extension States:
AL BA RS

- **SATOH, Hideo**
Ibaraki-shi
Osaka 567-0826 (JP)
- **TARUI, Hirokazu**
Toyonaka-shi
Osaka 561-0802 (JP)

(30) Priority: **11.06.2008 JP 2008152619**

(74) Representative: **Duckworth, Timothy John**
J.A. Kemp & Co.
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

(71) Applicant: **Sumitomo Chemical Company, Limited**
Tokyo 104-8260 (JP)

(72) Inventors:
• **TOMIGAHARA, Yoshitaka**
Toyonaka-shi
Osaka 560-0013 (JP)

(54) **METHOD FOR DETECTING OR QUANTIFYING DNA**

(57) The present invention relates to a method for quantifying or detecting DNA having a target DNA region, and so on.

EP 2 305 807 A1

Description

TECHNICAL FIELD

5 **[0001]** The present invention relates to a method for quantifying or detecting DNA having a target DNA region, and so on.

BACKGROUND ART

10 **[0002]** Known as a method for quantifying or detecting DNA having a target DNA region contained in a specimen are, for example, a method of detecting DNA having a target DNA region amplified by a chain reaction of DNA synthesis by DNA polymerase (Polymerase Chain Reaction; hereinafter, sometimes referred to as PCR) after extraction of DNA from a specimen, a method of detecting DNA by hybridization of a fluorescent-labeled oligonucleotide with a target DNA region possessed by DNA in a specimen, and so on (see, for example, J. Cataract. Refract. Surg., 2007;33(4):635-641, Environ. Mol. Mutagen., 1991;18(4):259-262).

15

DISCLOSURE OF THE INVENTION

[0003] It is an object of the present invention to provide a method for quantifying or detecting DNA having a target DNA region in a simple and convenient manner.

20

[0004] Specifically, the present invention provides:

[Invention 1]

[0005] A method for quantifying or detecting DNA comprising a target DNA region contained in a specimen comprising:

25

- (1) First step of preparing from the specimen DNA for which the target DNA region is to be detected;
- (2) Second step of treating the DNA prepared in First step with a DNA methylation enzyme;
- (3) Third step of preparing single-stranded methylated DNA from the DNA treated in Second step, and making a detection oligonucleotide bind with the single-stranded methylated DNA to obtain a test DNA complex;
- 30 (4) Fourth step of making an immobilized methylated-DNA antibody bind with the test DNA complex obtained in Third step to obtain a detection complex; and
- (5) Fifth step of quantifying or detecting DNA comprising a target DNA region in the single-stranded DNA by quantifying or detecting the detection oligonucleotide contained in the detection complex obtained in Fourth step by its identification function (hereinafter, sometimes referred to as the present method);

35

[Invention 2]

[0006] The method according to Invention 1, wherein a counter oligonucleotide is added in obtaining a test DNA complex in Third step;

40

[Invention 3]

[0007] The method according to Invention 1 or 2, wherein the immobilized methylated-DNA antibody is a methylcytosine antibody;

45

[Invention 4]

[0008] The method according to any one of Inventions 1 to 3, wherein the DNA methylation enzyme is a cytosine methylation enzyme or Sssl methylase;

50

[Invention 5]

[0009] The method according to any one of Inventions 1 to 4, wherein the DNA comprising a target DNA region contained in the specimen is DNA comprising a target DNA region in DNA generated from RNA by a reverse transcriptase;

55

[Invention 6]

[0010] The method according to any one of Inventions 1 to 5, wherein the specimen is any of the following biological

specimen:

- 5 (a) mammalian blood, body fluid, excreta, body secretion, cell lysate, or tissue lysate,
(b) DNA extracted from one selected from the group consisting of mammalian blood, body fluid, excreta, body secretion, cell lysate, and tissue lysate,
(c) DNA prepared by using as a template RNA extracted from one selected from the group consisting of mammalian tissue, cell, tissue lysate and cell lysate,
(e) DNA extracted from bacterium, fungus or virus, or
10 (f) DNA prepared by using as a template RNA extracted from cell, fungus or virus;

[Invention 7]

15 **[0011]** The method according to any one of Inventions 1 to 6, wherein the DNA comprising a target DNA region obtained in First step is DNA digested in advance with a restriction enzyme recognition cleavage site for which is not present in the target DNA region, a synthesized oligonucleotide, or DNA purified in advance;

[Invention 8]

20 **[0012]** The method according to any one of Inventions 1 to 7, wherein the identification function of the detection oligonucleotide is any of the following identification function:

- (a) fluorescence detection of FITC, or
(b) detection by FITC antibody;

25 [Invention 9]

30 **[0013]** The method according to any one of Inventions 1 to 8, wherein the detection oligonucleotide comprises a repetitive sequence or a nucleotide sequence of an overlapping gene or a pseudo gene in human genome or a nucleotide sequence capable of complementarily binding with a part thereof;

[Invention 10]

[0014] The method according to Invention 9, wherein the repetitive sequence in human genome is LINE or SINE;

35 [Invention 11]

[0015] The method according to any one of Inventions 1 to 10, wherein the detection oligonucleotide comprises a nucleotide sequence capable of complementarily binding with any one of the following nucleotide sequences:

- 40 (1) the nucleotide sequence of SEQ ID NO: 37 or a nucleotide sequence having 80% or more sequence identity to the same,
(2) a complementary sequence of the nucleotide sequence of SEQ ID NO: 37 or a nucleotide sequence having 80% or more sequence identity to the same,
(3) the nucleotide sequence of SEQ ID NO: 39 or a nucleotide sequence having 80% or more sequence identity to the same, or
45 (4) a complementary sequence of the nucleotide sequence of SEQ ID NO: 39 or a nucleotide sequence having 80% or more sequence identity to the same;

50 [Invention 12]

[0016] The method according to any one of Inventions 1 to 10, wherein the detection oligonucleotide comprises any one of the following nucleotide sequences:

- 55 (1) the nucleotide sequence of SEQ ID NO: 38 or a nucleotide sequence having 80% or more sequence identity to the same,
(2) a complementary sequence of the nucleotide sequence of SEQ ID NO: 38 or a nucleotide sequence having 80% or more sequence identity to the same,
(3) the nucleotide sequence of SEQ ID NO: 40 or a nucleotide sequence having 80% or more sequence identity to

the same, or

(4) a complementary sequence of the nucleotide sequence of SEQ ID NO: 40 or a nucleotide sequence having 80% or more sequence identity to the same;

5 [Invention 13]

[0017] The method according to any one of Inventions 1 to 12, wherein concentration of a sodium salt in a solution used in a DNA extracting operation for preparing DNA from a specimen in First step is 100 mM or more and 1000 mM or less;

10 [Invention 14]

[0018] The method according to any one of Inventions 1 to 12, wherein concentration of a sodium salt in a solution used in a DNA extracting operation for preparing DNA from a specimen in First step is 100 mM or more and 200 mM or less;

15 [Invention 15]

[0019] A method for selecting a specimen from a cancer patient comprising the step of evaluating that a specimen from a test subject is a specimen from a cancer patient when there is significant difference between a quantification result or a detection result of DNA quantified or detected by using the specimen from the test subject according to the method of any one of Inventions 1 to 14 and a quantification result or a detection result of DNA quantified or detected by using a specimen from a healthy subject according to the same method, and identifying a specimen from a cancer patient based on a result of the evaluation;

25 [Invention 16]

[0020] The method according to Invention 15, wherein the specimen is mammalian serum; and

30 [Invention 17]

[0021] The method according to Invention 15 or 16, wherein DNA comprising a target DNA region is free DNA comprising the target DNA region in a mammalian serum; and so on.

BRIEF DESCRIPTION OF THE DRAWINGS

35 **[0022]**

Fig. 1 is a drawing showing results obtained by using 0.5 $\mu\text{g/mL}$ of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F1 in Example 1. DNA amounts measured by absorbance (450nm) are shown for Solution A (10 ng/10 μL TE buffer solution), Solution B (1 ng/10 μL TE buffer solution), Solution C (0.1 ng/10 μL TE buffer solution), and Solution D (0 ng/10 μL TE buffer solution (negative control solution)), respectively in this order from the right.

Fig. 2 is a drawing showing results obtained by using 0.5 $\mu\text{g/mL}$ of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F1 in Example 2. DNA amounts measured by absorbance (450nm) are shown for Solution MA (10 ng each/20 μL TE buffer solution), Solution MB (1 ng each/20 μL TE buffer solution), Solution MC (0.1 ng each/20 μL TE buffer solution), and Solution MD (0 ng each/20 μL TE buffer solution (negative control solution)), respectively in this order from the right.

Fig. 3 is a drawing showing results obtained by using 0.5 $\mu\text{g/mL}$ of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F2 in Example 3. DNA amounts measured by absorbance (450nm) are shown for Solution MA (10 ng each/20 μL TE buffer solution), Solution MB (1 ng each/20 μL TE buffer solution), Solution MC (0.1 ng each/20 μL TE buffer solution), and Solution MD (0 ng each/20 μL TE buffer solution (negative control solution)), respectively in this order from the right.

Fig. 4 is a drawing showing results obtained by using 0.5 $\mu\text{g/mL}$ of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotides F1 and F2 in Example 4. DNA amounts measured by absorbance (450nm) are shown for Solution MA (10 ng each/20 μL TE buffer solution), Solution MB (1 ng each/20 μL TE buffer solution), Solution MC (0.1 ng each/20 μL TE buffer solution), and Solution MD (0 ng each/20 μL TE buffer solution (negative control solution)), respectively in this order from the right.

Fig. 5 is a drawing showing results obtained by using 0.5 $\mu\text{g/mL}$ of a biotin-labeled methylcytosine antibody and 5'-

end FITC-labeled oligonucleotide F3 in Example 5. DNA amounts measured by absorbance (450nm) are shown for Solution A (10 ng/10 μ L TE buffer solution), Solution B (1 ng/10 μ L TE buffer solution), Solution C (0.1 ng/10 μ L TE buffer solution), and Solution D (0 ng/10 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

5 Fig. 6 is a drawing showing results obtained by using 0.5 μ g/mL of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F3 in Example 6. DNA amounts measured by absorbance (450nm) are shown for Solution MA (10 ng each/20 μ L TE buffer solution), Solution MB (1 ng each/20 μ L TE buffer solution), and Solution MC (0 ng each/20 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

10 Fig. 7 is a drawing showing results obtained by using 0.5 μ g/mL of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotides F3 and F4 in Example 7. DNA amounts measured by absorbance (450nm) are shown for Solution MA (10 ng each/20 μ L TE buffer solution), Solution MB (1 ng each/20 μ L TE buffer solution), and Solution MC (0 ng each/20 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

15 Fig. 8 is a drawing showing results obtained by using 0.5 μ g/mL of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F5 in Example 8. DNA amounts measured by absorbance (450nm) are shown for Solution A (100 ng/5 μ L TE buffer solution), Solution B (10 ng/5 μ L TE buffer solution), Solution C (1 ng/5 μ L TE buffer solution), and Solution D (0 ng/5 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

20 Fig. 9 is a drawing showing results obtained by using 0.5 μ g/mL of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F6 in Example 9. DNA amounts measured by absorbance (450nm) are shown for Solution A (100 ng/5 μ L TE buffer solution), Solution B (10 ng/5 μ L TE buffer solution), Solution C (1 ng/5 μ L TE buffer solution), and Solution D (0 ng/5 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

25 Fig. 10 is a drawing showing results obtained by using 0.5 μ g/mL of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F3 in Example 10. DNA amounts measured by absorbance (450nm) are shown for Solution A (10 ng/10 μ L TE buffer solution), Solution B (1 ng/10 μ L TE buffer solution), Solution C (0.1 ng/10 μ L TE buffer solution), and Solution D (0 ng/10 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

30 Fig. 11 is a drawing showing results obtained by using 0.5 μ g/mL of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F3 in Example 11. DNA amounts measured by absorbance (450nm) are shown for Solution MA (10 ng each/20 μ L TE buffer solution), Solution MB (1 ng each/20 μ L TE buffer solution), and Solution MC (0 ng each/20 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

35 Fig. 12 is a drawing showing results obtained by using 0.5 μ g/mL of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F4 in Example 12. DNA amounts measured by absorbance (450nm) are shown for Solution MA (10 ng each/20 μ L TE buffer solution), Solution MB (1 ng each/20 μ L TE buffer solution), Solution MC (0 ng each/20 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

40 Fig. 13 is a drawing showing results obtained by using 0.5 μ g/mL of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotides F3 and F4 in Example 13. DNA amounts measured by absorbance (450nm) are shown for Solution MA (10 ng each/20 μ L TE buffer solution), Solution MB (1 ng each/20 μ L TE buffer solution), and Solution MC (0 ng each/20 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

45 Fig. 14 is a drawing showing results obtained by using 0.5 μ g/mL of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F5 in Example 14. DNA amounts measured by absorbance (450nm) are shown for Solution A (100 ng/5 μ L TE buffer solution), Solution B (10 ng/5 μ L TE buffer solution), Solution C (1 ng/5 μ L TE buffer solution), and Solution D (0 ng/5 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

50 Fig. 15 is a drawing showing results obtained by using 0.5 μ g/mL of a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F6 in Example 15. DNA amounts measured by absorbance (450nm) are shown for Solution A (100 ng/5 μ L TE buffer solution), Solution B (10 ng/5 μ L TE buffer solution), Solution C (1 ng/5 μ L TE buffer solution), and Solution D (0 ng/5 μ L TE buffer solution (negative control solution)), respectively in this order from the right.

55 Fig. 16 is a drawing showing results obtained by conducting Treatment 1 and detecting DNA using a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F6 in Example 16. Measured absorbances (450nm) are shown for Solution A (10 ng/10 μ L rat serum solution), Solution B (1 ng/10 μ L rat serum solution), Solution C (0.1 ng/10 μ L rat serum solution), and Solution D (rat serum (negative control solution)), respectively in this order from the right.

Fig. 17 is a drawing showing results obtained by conducting Treatment 2 and detecting DNA using a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F6 in Example 16. Measured absorbances (450nm)

are shown for Solution A (10 ng/10 μ L rat serum solution), Solution B (1 ng/10 μ L rat serum solution), Solution C (0.1 ng/10 μ L rat serum solution), and Solution D (rat serum (negative control solution)), respectively in this order from the right.

Fig. 18 is a drawing showing results obtained by detecting DNA using a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F6 in Example 17. Measured absorbances (450nm) are shown for Solution A (4 ng/40 μ L human serum solution), Solution B (2 ng/40 μ L human serum solution), Solution C (1 ng/40 μ L human serum solution), and Solution D (human serum (negative control solution)), respectively in this order from the right.

Fig. 19 is a drawing showing a comparison between a result detected by using a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F6 and a result quantified by real time PCR in Example 18. In Fig. 19, the detection result is plotted on the vertical axis, and the quantification result by real-time PCR is plotted on the horizontal axis. The straight lines in the graph represent regression line (thick line) and standard error range (thin line).

Fig. 20 shows DNA detection results using a biotin-labeled methylcytosine antibody and 5'-end FITC-labeled oligonucleotide F6 for human serum samples aged 57 or younger in Example 18, which are plotted separately for cancer patients and healthy subjects, together with respective mean values and standard deviations.

MODE FOR CARRYING OUT THE INVENTION

[0023] Examples of the "specimen" in the present method include (a) mammalian blood, body fluid, excreta, body secretion, cell lysate, or tissue lysate, (b) DNA extracted from one selected from the group consisting of mammalian blood, body fluid, excreta, body secretion, cell lysate and tissue lysate, (c) DNA prepared by using as a template RNA extracted from one selected from the group consisting of mammalian tissue, cell, tissue lysate and cell lysate, (e) DNA extracted from cell, fungus or virus, and (f) DNA prepared by using as a template RNA extracted from cell, fungus or virus. The term "tissue" means broadly including blood and lymph node.

[0024] The term "mammal" means animals classified into animal kingdom, Chordata, Chordate subphylum, and Mammalia, and concrete examples include human being, monkey, marmoset, guinea pig, rat, mouse, cattle, sheep, dog, and cat.

[0025] The term "body fluid" means a liquid existing between cells constituting an individual body, and concretely, plasma and interstitial fluid are recited, and it often functions to maintain homeostasis of an individual body. More concrete examples include lymph, tissue fluid (interinstitutional fluid, intercellular fluid, interstitial fluid), celomic fluid, serous cavity fluid, pleural effusion, ascetic fluid, pericardial fluid, cerebral fluid (spinal fluid), joint fluid (spinal fluid), eye aqueous fluid (aqueous fluid), and cerebrospinal fluid.

[0026] The term "body secretion" is a secretion from an exocrine gland, and concrete examples include saliva, gastric juice, bile, pancreatic juice, intestinal juice, sweat, tear, runny nose, semen, vaginal lubricant, amniotic fluid, and milk.

[0027] When the specimen is blood, body fluid or body secretion of a human being, a sample collected for a clinical test in a regular health check of human may be utilized.

[0028] Examples of the "cell lysate" include lysates containing intracellular fluids obtained by grinding cells, such as cell strains, primary cultured cells or blood cells, cultured in a plate for cell culture. As a method of grinding cells, a method based on sonication, a method using a surfactant, a method of using an alkaline solution and the like are recited. For lysing cells, a commercially available kit and the like may be used.

[0029] For example, after culturing cells to be confluent in a 10 cm plate, the culture solution is removed, and 0.6 mL of a RIPA buffer (1x TBS, 1% nonidet P-40, 0.5% sodium deoxysholate, 0.1% SDS, 0.004% sodium azide) is added to the plate. After shaking slowly the plate at 4°C for 15 minutes, cells adhered on the plate are removed by using a scraper or the like, and the liquid on the plate is transferred to a microtube. After adding 10 mg/mL PMSF in an amount of 1/10 volume of the liquid, the tube is left still on ice for 30 to 60 minutes, the solution is centrifuged at 4°C for 10 minutes at 10,000xg, to obtain the supernatant as a cell lysate.

[0030] As the "tissue lysate", lysates containing intracellular fluids obtained by grinding cells in tissues collected from animals such as mammals can be recited.

[0031] Concretely, after measuring the weight of a tissue obtained from an animal, the tissue is cut into small pieces with the use of a razor or the like. When a frozen tissue is used, it is necessary to make a smaller piece. After cutting, an ice-cooled RIPA buffer is added in a rate of 3 mL per 1 g of tissue, and homogenized at 4°C. Here, as the RIPA buffer, a protease inhibitor, a phosphatase inhibitor and the like may be added, and for example, 10 mg/mL PMSF in an amount of 1/10 volume of the RIPA buffer may be added. For homogenization, a sonicator or a pressurized cell grinder is used. In an operation of homogenization, a homogenized liquid is constantly kept at 4°C for preventing heat generation. The homogenized liquid is transferred to a microtube, and centrifuged at 4°C for 10 minutes at 10,000xg, and the supernatant is obtained as a tissue lysate.

[0032] Examples of the "specimen" in the present method include samples and surface adhered matters collected from foods, rivers, soils or general commercial products, and microorganisms such as fungi, cells, viruses and nucleic

acids thereof can be contained.

[0033] Examples of the DNA used as a specimen include genomic DNA obtained by extraction from the biological sample or the microorganism, and DNA fragment or RNA derived from genomic DNA. For obtaining genomic DNA from a sample derived from a mammal, for example, a commercially available DNA extraction kit and the like may be used.

[0034] The term "target DNA region" (hereinafter, sometimes referred to as a target region) in the present method means a DNA region intended to be detected or quantified by the present method in DNA contained in a specimen. The target DNA region is represented by a nucleotide sequence on DNA when the specimen is DNA. When the specimen is RNA, the target DNA region is represented by a nucleotide sequence on DNA prepared from RNA by a reverse transcriptase, and is a complementary nucleotide sequence of a prescribed nucleotide sequence to be detected on RNA. In the present method, when cytosine is methylated and detected or quantified, a target region desirably contains a region abundantly containing cytosine or CpG as will be described later.

[0035] First step is a step of preparing from a specimen DNA for which a target DNA region is to be detected.

[0036] Examples of DNA prepared in First step include a DNA sample digested in advance with a restriction enzyme recognition cleavage site for which is not present in the target DNA region possessed by the DNA, a DNA sample purified in advance, free DNA in blood, DNA derived from microbial genome, and DNA prepared from RNA in a specimen by a reverse transcriptase. As DNA prepared in First step, for example, DNA that is designed based on gene information of the specimen and artificially synthesized may be recited.

[0037] When blood is used as a specimen, plasma or serum is prepared from blood by a routine method, and the prepared plasma or serum is used as a specimen, and free DNA (containing DNA derived from cancer cells such as gastric cancer cells) contained therein is analyzed, and thus DNA derived from cancer cells such as gastric cancer cells can be analyzed away from DNA derived from hemocytes, and sensitivity of detecting cancer cells such as gastric cancer cells, and tissues containing the same can be improved.

[0038] Examples of DNA prepared in First step include DNA derived from microorganisms such as gram-positive bacteria, gram-negative bacteria, fungi, viruses and pathogenic protozoans, and DNA obtained from RNA derived from such microorganisms by a reverse transcriptase. For example, genomic DNA or DNA prepared by a reverse transcriptase from RNA of *Mycoplasma genitalium*, *Mycoplasma pneumoniae*, *Borrelia burgdorferi* B31, *Rickettsia prowazekii*, *Treponema pallidum*, *Chlamydia pneumoniae*, *Chlamydia trachomatis*, *Helicobacter pylori* J99, *Helicobacter pylori* 26695, *Haemophilus influenzae* Rd, *Mycobacterium tuberculosis* H37Rv, *Pseudomonas aeruginosa*, *Legionella pneumophila*, *Serratia marcescens*, *Escherichia coli*, *Listeria monocytogenes*, *Salmonella enterica*, *Campylobacter jejuni* subsp. *Jejuni*, *Staphylococcus aureus*, *Vibrio parahaemolyticus*, *Bacillus cereus*, *Clostridium botulinum*, *Clostridium perfringens*, *Yersinia enterocolitica*, *Yersinia pseudotuberculosis*, *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Candida albicans*, *Cryptococcus neoformans*, *Aspergillus fumigatus*, *Pneumocystis carinii*, *Coccidioides immitis*, *Cytomegalovirus*, *human herpesvirus 5*, *Epstein-Barr virus*, *Human Immunodeficiency Virus*, *Human Papilloma Virus*, *Enterovirus*, *Norovirus Influenza Virus*, *Toxoplasma gondii*, *Cryptosporidium parvum*, or *Entamoeba histolytica* may be used for detection of a microorganism responsible for an infection in a specimen, or a microorganism responsible for a food poisoning in food.

[0039] For preparing genomic DNA, for example, when the specimen is a sample derived from a mammal, a commercially available DNA extraction kit (Genfind v2 Kit (available from BECKMAN COULTER), FastPure DNA Kit (available from TAKARA BIO INC.)) and the like may be used.

[0040] When the specimen is a microorganism such as fungus, genomic DNA may be prepared by a general preparation method of yeast genome or the like as described in *Methods in Yeast Genetics* (Cold Spring Harbor Laboratory Press), and when the specimen is a prokaryote such as *Escherichia coli*, a general preparation method of microorganism genome or the like as described in *Molecular Cloning -A Laboratory Manual-* (Cold Spring Harbor Laboratory Press) may be used.

[0041] When the specimen is a food sample, DNA may be prepared after separating a microorganism or the like from the food, and genomic DNA of non-microorganism and genome derived from a microorganism contained in the food may be obtained at the same time. When the specimen is a tissue derived from a mammal, and the target DNA region is DNA derived from a virus, RNA may be extracted from the tissue using such as a commercially available RNA extraction kit (ISOGEN(311-02501)(available from NIPPON GENE CO., LTD.), or FastRNA Pro Green Kit (available from Funakoshi Corporation), FastRNA Pro Blue Kit (available from Funakoshi Corporation), FastRNA Pro Red Kit (available from Funakoshi Corporation), and the like), and DNA may be obtained by a reverse transcriptase. When the specimen is a specimen derived from a mammal, viral DNA may be extracted after extracting virus particles, or after extracting virus particles, viral RNA may be extracted using a commercially available kit (QuickGene RNA tissue kit SII, available FUJIFILM Corporation) or the like, and DNA derived from the virus may be obtained by a reverse transcriptase. RNA may be extracted from a tissue infected by a virus, and DNA derived from the virus may be obtained by a reverse transcriptase, or DNA may be obtained from a tissue infected by a virus, and DNA derived from the virus may be obtained. When DNA is obtained from RNA by a reverse transcriptase, a commercially available kit (Transcripser high fidelity cDNA synthesis

kit, available from Roche Diagnostics K.K.) and the like may be used.

[0042] In "methylated DNA", any of four kinds of bases constituting gene (genomic DNA) is methylated. For example, in a mammal is known a phenomenon that only cytosine in a nucleotide sequence represented by 5'-CG-3' (C represents cytosine, and G represents guanine. Hereinafter, the nucleotide sequence is occasionally denoted by "CpG") is methylated. A methylation site of cytosine is position 5. In DNA duplication antecedent to cell division, only cytosine in "CpG" in a template chain derived from a parent cell is methylated in nascent double-stranded DNA, and cytosine in "CpG" in a nascent DNA chain is also methylated rapidly by the action of a methyltransferase. In this cytosine methylation is methylated cytosine in CpG in a nascent DNA chain that complementarily binds to CpG containing methylated cytosine in a DNA chain derived from a parent cell. Therefore, the methylation condition of DNA of the parent cell is taken over as it is to new two sets of DNA after DNA duplication. The term "CpG pair" means a double-stranded DNA in which a nucleotide sequence represented by CpG binds to CpG complementary to the sequence.

[0043] The term "single-stranded methylated DNA" means single-stranded DNA in which is methylated cytosine at a position 5 in a nucleotide sequence represented by 5'-CG-3' in a nucleotide sequence of the single-stranded DNA.

[0044] Examples of the "target DNA region" include promoter regions, untranslated regions or translated regions (coding regions) of useful protein genes such as Lysyl oxidase, HRAS-like suppressor, bA305P22.2.1, Gamma filamin, HAND1, Homologue of RIKEN 2210016F16, FLJ32130, PPARG angiopoietin-related protein, Thrombomodulin, p53-responsive gene 2, Fibrillin 2, Neurofilament 3, disintegrin and metalloproteinase domain 23, G protein-coupled receptor 7, G-protein coupled somatostatin and angiotensin-like peptide receptor, and Solute carrier family 6 neurotransmitter transporter noradrenalin member 2, and preferably include DNA regions containing one or more CpG present in these nucleotide sequences. In the present method, methylated DNA of "target DNA region" may be detected or quantified individually, and, for example, when more methylated DNA of "target DNA region" is detected in one detection system, the quantification accuracy and detection sensitivity are improved correspondingly.

[0045] To be more specific, when the useful protein gene is a Lysyl oxidase gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a Lysyl oxidase gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 1 (corresponding to a nucleotide sequence represented by base No. 16001 to 18661 in the nucleotide sequence described in Genbank Accession No. AF270645) can be recited. In the nucleotide sequence of SEQ ID NO: 1, ATG codon encoding methionine at amino terminal of Lysyl oxidase protein derived from human is represented in base No. 2031 to 2033, and a nucleotide sequence of the above exon 1 is represented in base No. 1957 to 2661.

[0046] To be more specific, when the useful protein gene is a HRAS-like suppressor gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a HRAS-like suppressor gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 2 (corresponding to a nucleotide sequence represented by base No. 172001 to 173953 in the nucleotide sequence described in Genbank Accession No. AC068162) can be recited. In the nucleotide sequence of SEQ ID NO: 2, the nucleotide sequence of exon 1 of a HRAS-like suppressor gene derived from human is represented in base No. 1743 to 1953.

[0047] To be more specific, when the useful protein gene is a bA305P22.2.1 gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a bA305P22.2.1 gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 3 (corresponding to a nucleotide sequence represented by base No. 13001 to 13889 in the nucleotide sequence described in Genbank Accession No. AL121673) can be recited. In the nucleotide sequence of SEQ ID NO: 3, ATG codon encoding methionine at amino terminal of bA305P22.2.1 protein derived from human is represented in base No. 849 to 851, and a nucleotide sequence of the above exon 1 is represented in base No. 663 to 889.

[0048] To be more specific, when the useful protein gene is a Gamma filamin gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a Gamma filamin gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 4 (corresponding to a complementary sequence to a nucleotide sequence represented by base No. 63528 to 64390 in the nucleotide sequence described in Genbank Accession No. AC074373) can be recited. In the nucleotide sequence of SEQ ID NO: 4, ATG codon encoding methionine at amino terminal of Gamma filamin protein derived from human is represented in base No. 572 to 574, and a nucleotide sequence of the above exon 1 is represented in base No. 463 to 863.

[0049] To be more specific, when the useful protein gene is a HAND1 gene, as a nucleotide sequence that includes

at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a HAND1 gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 5 (corresponding to a complementary sequence to a nucleotide sequence represented by base No. 24303 to 26500 in the nucleotide sequence described in Genbank Accession No. AC026688) can be recited. In the nucleotide sequence of SEQ ID NO: 5, ATG codon encoding methionine at amino terminal of HAND1 protein derived from human is represented in base No. 1656 to 1658, and a nucleotide sequence of the above exon 1 is represented in base No. 1400 to 2198.

[0050] To be more specific, when the useful protein gene is a Homologue of RIKEN 2210016F16 gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a Homologue of RIKEN 2210016F16 gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 6 (corresponding to a complementary nucleotide sequence to a nucleotide sequence represented by base No. 157056 to 159000 in the nucleotide sequence described in Genbank Accession No. AL354733) can be recited. In the nucleotide sequence of SEQ ID NO: 6, a nucleotide sequence of exon 1 of a Homologue of a RIKEN 2210016F16 gene derived from human is represented in base No. 1392 to 1945.

[0051] To be more specific, when the useful protein gene is a FLJ32130 gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a FLJ32130 gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 7 (corresponding to a complementary nucleotide sequence to a nucleotide sequence represented by base No. 1 to 2379 in the nucleotide sequence described in Genbank Accession No. AC002310) can be recited. In the nucleotide sequence of SEQ ID NO: 7, ATG codon encoding methionine at amino terminal of FLJ32130 protein derived from human is represented in base No. 2136 to 2138, and a nucleotide sequence assumed to be the above exon 1 is represented in base No. 2136 to 2379.

[0052] To be more specific, when the useful protein gene is a PPARG angiopoietin-related protein gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a PPARG angiopoietin-related protein gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 8 can be recited. In the nucleotide sequence of SEQ ID NO: 8, ATG codon encoding methionine at amino terminal of PPARG angiopoietin-related protein derived from human is represented in base No. 717 to 719, and a nucleotide sequence of the 5' side part of the above exon 1 is represented in base No. 1957 to 2661.

[0053] To be more specific, when the useful protein gene is a Thrombomodulin gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a Thrombomodulin gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 9 (corresponding to a nucleotide sequence represented by base No. 1 to 6096 in the nucleotide sequence described in Genbank Accession No. AF495471) can be recited. In the nucleotide sequence of SEQ ID NO: 9, ATG codon encoding methionine at amino terminal of Thrombomodulin protein derived from human is represented in base No. 2590 to 2592, and a nucleotide sequence of the above exon 1 is represented in base No. 2048 to 6096.

[0054] To be more specific, when the useful protein gene is a p53-responsive gene 2 gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a p53-responsive gene 2 gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 10 (corresponding to a complementary sequence to a nucleotide sequence represented by base No. 113501 to 116000 in the nucleotide sequence described in Genbank Accession No. AC009471) can be recited. In the nucleotide sequence of SEQ ID NO: 10, a nucleotide sequence of exon 1 of a p53-responsive gene 2 gene derived from human is represented in base No. 1558 to 1808.

[0055] To be more specific, when the useful protein gene is a Fibrillin2 gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a Fibrillin2 gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 11 (corresponding to a complementary sequence to a nucleotide sequence represented by base No. 118801 to 121000 in the nucleotide sequence described in Genbank Accession No. AC113387) can be recited. In the nucleotide sequence of SEQ ID NO: 11, a nucleotide sequence of exon 1 of a Fibrillin2

gene derived from human is represented in base No. 1091 to 1345.

[0056] To be more specific, when the useful protein gene is a Neurofilament3 gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a Neurofilament3 gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 12 (corresponding to a complementary sequence to a nucleotide sequence represented by base No. 28001 to 30000 in the nucleotide sequence described in Genbank Accession No. AF106564) can be recited. In the nucleotide sequence of SEQ ID NO: 12, a nucleotide sequence of exon 1 of a Neurofilament3 gene derived from human is represented in base No. 614 to 1694.

[0057] To be more specific, when the useful protein gene is a disintegrin and metalloproteinase domain 23 gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a disintegrin and metalloproteinase domain 23 gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 13 (corresponding to a nucleotide sequence represented by base No. 21001 to 23300 in the nucleotide sequence described in Genbank Accession No. AC009225) can be recited. In the nucleotide sequence of SEQ ID NO: 13, a nucleotide sequence of exon 1 of a disintegrin and metalloproteinase domain 23 gene derived from human is represented in base No. 1194 to 1630.

[0058] To be more specific, when the useful protein gene is a G protein-coupled receptor 7 gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a G protein-coupled receptor 7 gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 14 (corresponding to a nucleotide sequence represented by base No. 75001 to 78000 in the nucleotide sequence described in Genbank Accession No. AC009800) can be recited. In the nucleotide sequence of SEQ ID NO: 14, a nucleotide sequence of exon 1 of a G protein-coupled receptor 7 gene derived from human is represented in base No. 1666 to 2652.

[0059] To be more specific, when the useful protein gene is a G-protein coupled somatostatin and angiotensin-like peptide receptor gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a G-protein coupled somatostatin and angiotensin-like peptide receptor gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 15 (corresponding to a complementary sequence to a nucleotide sequence represented by base No. 57001 to 60000 in the nucleotide sequence described in Genbank Accession No. AC008971) can be recited. In the nucleotide sequence of SEQ ID NO: 15, a nucleotide sequence of exon 1 of a G-protein coupled somatostatin and angiotensin-like peptide receptor gene derived from human is represented in base No. 776 to 2632.

[0060] To be more specific, when the useful protein gene is a Solute carrier family 6 neurotransmitter transporter noradrenalin member 2 gene, as a nucleotide sequence that includes at least one nucleotide sequence represented by CpG present in a nucleotide sequence of its promoter region, untranslated region or translated region (coding region), a nucleotide sequence of a genomic DNA containing exon 1 of a Solute carrier family 6 neurotransmitter transporter noradrenalin member 2 gene derived from human, and a promoter region located 5' upstream of the same can be recited, and more concretely, the nucleotide sequence of SEQ ID NO: 16 (corresponding to a complementary sequence to a nucleotide sequence represented by base No. 78801 to 81000 in the nucleotide sequence described in Genbank Accession No. AC026802) can be recited. In the nucleotide sequence of SEQ ID NO: 16, a nucleotide sequence of exon 1 of a Solute carrier family 6 neurotransmitter transporter noradrenalin member 2 gene derived from human is represented in base No. 1479 to 1804.

[0061] Second step is a step of treating the DNA prepared in First step with a DNA methylation enzyme.

[0062] The "DNA methylation enzyme" means an enzyme that methylates a base in DNA, and various kinds DNA methylation enzymes are isolated from mammalian cells, bacteria and the like. DNA methylation enzymes are classified into several kinds such as adenine methylation enzymes, and cytosine methylation enzymes according to the kind of the base of a substrate. A cytosine methylation enzyme is an enzyme that recognizes a specific sequence in a DNA nucleotide sequence, and methylates cytosine near the sequence, and different cytosine methylation enzymes are known according to the recognized nucleotide sequences.

[0063] A number of methylation reactions of DNA catalyzed by a DNA methylation enzyme are found from a primitive immune system called a restriction-modification system. The restriction-modification system is a function that digests foreign DNA (in particular, bacteriophage) with a restriction enzyme after regularly methylating the entire genome functioning in bacteria to protect it from being digested by a restriction enzyme (restriction endonuclease) that recognizes a specific sequence, and is a system for protecting a microbial genome from bacteriophage infection. Enzymes functioning

in methylation of genome are known to methylate cytosine or adenine, and often known to methylate nitrogen at position 6 (N6) or carbon at position 5 (C5) of a purine residue. Among these enzymes, known as a cytosine methylation enzyme that methylates C5 of cytosine are SssI (M.SssI) methylase, AluI methylase, HhaI methylase, HpaII methylase, MspI methylase, HaeIII methylase, and so on. These enzymes that methylate position C5 of cytosine recognize different nucleotide sequences, and a cytosine methylation enzyme that recognizes CpG is only SssI.

[0064] As a methylation reaction of DNA in human genome, methylation at position 5 (C5) of cytosine in CpG is known as epigenetics (the mechanism generating diversity of gene expression independent of gene sequence), and as such a cytosine methylation enzyme, DNA methyltransferase is known. As a DNA methyltransferase, Dnmt1 methyltransferase is known.

[0065] In human cells, since position C5 of cytosine in a CpG sequence is methylated, for methylating genome artificially, the same position of the same cytosine in the same sequence (CpG) with methylation in a human cell can be methylated by using SssI.

[0066] For methylating DNA by a cytosine methylation enzyme, concretely, for example, a DNA sample is added with 5 μ L of an optimum 10x buffer (NEBuffer2 (available from NEB Inc.)), 0.5 μ L of S-adenosyl methionine (3.2 mM, available from NEB Inc.) and 0.5 μ L of cytosine methylation enzyme SssI(available from NEB Inc.) and then the resultant mixture is added with sterilized ultrapure water to make the liquid amount 50 μ L, and then incubated at 37°C for 30 minutes. Since methylation in Second step is executed for making the target DNA region bind with a support by making it bind with the methylated DNA antibody, Second step is not necessarily executed insofar as the target DNA region of the extracted DNA sample is methylated.

[0067] In Second step, when a methylated base constituting the methylated DNA modified with the DNA methylation enzyme differs from the methylated base possessed by the later-described detection oligonucleotide, the methylated base on the detection oligonucleotide can be used as an identification function. For example, when the detection oligonucleotide is 6-methyladenine, and cytosine can be modified into 5-methylcytosine in Second step, a 6-methyladenine antibody may be used as the identification function of the detection oligonucleotide, and immobilization to a support can be achieved using a methylcytosine antibody. Concretely, when modification is achieved by SssI methylase in Second step, immobilization to a support may be achieved by using a methylcytosine antibody. That is, when the detection oligonucleotide has methyladenine, and the detection oligonucleotide is detected by a methyladenine antibody, it is possible to detect only methyladenine of the detection oligonucleotide without detecting methylcytosine of DNA having a target DNA region.

[0068] Third step is a step of preparing single-stranded methylated DNA from the DNA treated with a DNA methylation enzyme obtained in Second step, and making a detection oligonucleotide bind with the single-stranded methylated DNA having a target DNA region to obtain a test DNA complex.

[0069] The "detection oligonucleotide" in the present method is such an oligonucleotide that bases of nucleotides constituting the oligonucleotide has an identification function for detecting or quantifying the detection oligonucleotide, and an adhesion sequence for detection which is a nucleotide sequence for binding by complementation with the DNA having a target DNA region.

[0070] The "detection oligonucleotide" in the present invention may comprise a repetitive sequence in human genome as will be described later, a nucleotide sequence of an overlapping gene or a pseudo gene or a nucleotide sequence capable of complementarily binding with a part of the same. Concretely, for example, the nucleotide sequences of SEQ ID NOs: 38 and 40, nucleotide sequences having a complementary sequence therewith and the like are recited.

[0071] The adhesion sequence for detection is a nucleotide sequence required for forming a conjugate (double strand) with the single-stranded DNA containing a target DNA region, namely, a nucleotide sequence containing a sequence capable of binding with a part of the nucleotide sequence of the target DNA region by complementary base pairing, or a nucleotide sequence containing a nucleotide sequence that is complementary to a part of a nucleotide sequence in a further 5'-end side DNA region than 5'-end of the target DNA region or a nucleotide sequence containing a nucleotide sequence that is complementary to a part of a nucleotide sequence in a further 3'-end side DNA region than 3' -end of the target DNA region, and desirably not inhibiting binding of the methylated DNA antibody and the methylated DNA in the target DNA region.

[0072] The expression "not inhibiting binding of the methylated DNA antibody and the methylated DNA in the target DNA region" means that complementary binding between the present oligonucleotide and the single-stranded DNA does not occur in the occupied space required for the methylated DNA antibody to bind with the methylated single-stranded DNA. That is, it is supposed that the methylated DNA antibody occupies not only the methylated base (cytosine) to which the methylated DNA antibody directly binds, but also the peripheral space where the methylated base (cytosine) exists for binding with the methylated base (cytosine). Therefore, it suffices that the adhesion sequence for detection fails to complementary bind with the single-stranded DNA in the occupied space required for the methylated DNA antibody to bind with the methylated DNA. The adhesion sequence for detection to be bound with the single-stranded DNA is not necessarily one kind, and two or more kinds may be used unless binding of the methylated DNA antibody is inhibited. By using a plurality of the present oligonucleotides, it is possible to improve the quantity accuracy and detection sensitivity.

[0073] The expression "obtaining single-stranded methylated DNA having a target DNA region, and making a detection oligonucleotide bind with the single-stranded methylated DNA to obtain a test DNA complex" in Third step means forming a test DNA complex made up of methylated DNA having a target DNA region (hereinafter, also sometimes referred to as single-stranded methylated DNA) and the detection oligonucleotide that are bound complementarily. Concretely, the single-stranded methylated DNA methylated by a DNA methylation enzyme in Second step is prepared into a 1 ng/ μL solution in Tris-HCl buffer (10 mM), and a detection oligonucleotide capable of binding with the single-stranded methylated DNA is prepared into a 0.02 μM solution in Tris-HCl buffer (10 mM), and 10 μL of respective oligonucleotide solutions and a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μL of a 100 mM MgCl₂ solution, and 10 μL of a 1 mg/mL BSA solution are mixed, and the resultant mixture is further added with sterilized ultra pure water to make the liquid amount 100 μL . Then the mixture is heated at 95°C for 10 minutes, rapidly cooled to 70°C and retained at this temperature for 10 minutes, then cooled to 50°C and retained at this temperature for 10 minutes, retained at 37°C for 10 minutes, and then returned to room temperature, to promote formation of the test DNA complex of the single-stranded methylated DNA and the detection oligonucleotide.

[0074] The term "complementarily bind" means that double-stranded DNA is formed by base-pairing through a hydrogen bond between bases. For example, it means that bases that constitute respective single-stranded DNAs of a double strand forming DNA form a double strand by base-pairing between purine and pyrimidine, and more concretely, double-stranded DNA is formed by base-pairing through plural sequential hydrogen bonds between thymine and adenine, and guanine and cytosine. Binding based on complementation may be sometimes referred to as "complementarily binding". The term "complementarily binding" may be sometimes expressed by "capable of complementary binding", "capable of complementary base-pairing", "bind by complementation" or "bind (complementary bind (by base-pairing)) by complementation". Nucleotide sequences that are capable of complementarily binding may be sometimes expressed by "having complementation" or "complementary" to each other. It also means binding of inosine contained in an artificially prepared oligonucleotide with cytosine, adenine or thymine through hydrogen bonding.

[0075] In the present method, when the single-stranded methylated DNA and the detection oligonucleotide "bind by complementation", it is also included the case where a part of the nucleotide sequence constituting the adhesion sequence for detection of the detection oligonucleotide fails to base-pair with single-stranded methylated DNA. For example, the case where at least 75%, preferably 80% or more bases of the bases constituting the adhesion sequence for detection base-pair with the single-stranded methylated DNA, and the adhesion sequence for detection is able to bind with an oligonucleotide having at least 75% or more, preferably 80% or more homology with the test oligonucleotide is included.

[0076] In Third step of the present method, as a preferred aspect in forming the test DNA complex of the single-stranded methylated DNA and the detection oligonucleotide, for example, addition of a counter oligonucleotide is recited.

[0077] The "counter oligonucleotide" is obtained by dividing a polynucleotide having the same nucleotide sequence as the target DNA region into "short" oligonucleotides. Here, "short" means length of 10 to 100 bases, and more preferably 20 to 50 bases. The counter oligonucleotide is not designed on the nucleotide sequence where the detection oligonucleotide and the single-stranded methylated DNA bind. The counter oligonucleotide is added excessively compared to the target DNA region. When the target DNA region is a plus strand, it is added for preventing a complementary strand (minus strand) of the target DNA region and the target DNA region (plus strand) from rebinding by complementation when the target DNA region (plus strand) is made into a single-strand and caused to bind with the later-described immobilized methylated DNA antibody. This is because in making the methylated DNA antibody bind with the methylated target DNA region, and measuring an amount of target DNA or an index value correlated therewith, the methylated target region in a single strand state is more likely to bind with the methylated DNA antibody. The counter oligonucleotide is preferably added in an amount of at least 10 times, preferably 100 times or more compared to the target DNA region.

[0078] When the target DNA region in the DNA sample prepared in First step is single-stranded DNA, single-stranded methylated DNA can be obtained as the DNA sample without conducting any special operation for "preparing single-stranded methylated DNA" in Third step. Concretely, as the DNA sample, single-stranded DNA synthesized from RNA by a reverse transcriptase treatment, and single-stranded DNA obtained from a virus whose genome is single-stranded DNA, among the DNAs extracted from the virus are recited. When the target DNA region in the DNA sample prepared in First step is double-stranded DNA, an operation for making double-stranded DNA into single strands may be conducted for "preparing single-stranded methylated DNA" in Third step. In this case, concretely, the operation may include heating at 95°C for several minutes, followed by rapid cooling to 4°C or less.

[0079] The term "adhesion sequence for detection" in the detection oligonucleotide is an oligonucleotide comprising a nucleotide sequence complementary to a nucleotide sequence possessed by the target DNA region, and means a sequence having a homology of 75% or more, more preferably 90% or more with the nucleotide sequence of the target DNA region capable of pairing with the adhesion sequence for detection. It suffices that the adhesion sequence for detection does not inhibit binding between the later-described immobilized methylated DNA antibody and the test DNA complex. Further, "adhesion sequence for detection" is designed to have a nucleotide sequence that binds with the target DNA region or the vicinity of the target DNA region, and to be able to form a detection complex in Fourth step as will be described later. Only one or two or more adhesion sequences for detection may be designed in the same repetitive

sequence (in the target DNA region) as will be described later. When two or more adhesion sequences are designed, it suffices that the plural adhesion sequences for detection and a specific adhesion sequence as will be described later do not mutually inhibit binding with the single-stranded methylated DNA.

5 **[0080]** Fourth step is a step of making an immobilized methylated DNA antibody bind with the test DNA complex obtained in Third step to obtain a detection complex.

[0081] The "immobilized methylated DNA antibody" is a methylated DNA antibody binding with a methylated base in DNA as an antigen, which is immobilized to a "support". As the antibody, preferably, an antibody having the property of recognizing and binding with the cytosine that is methylated at position 5 in single-stranded DNA, and more preferably a methylcytosine antibody may be used. Also a commercially available methylated DNA antibody capable of specifically recognizing DNA in a methylated state described in the specification and specifically binding with the same may be used.

10 **[0082]** As the "support", the material and form thereof are not particularly limited as far as methylated DNA antibody can bind thereto. For example, any form suited for use purpose may be employed, including the forms of tube, test plate, filter, disc, bead and the like. As the material, those used as supports for a usual immune measuring method, for example, synthetic resins such as polystyrene, polypropylene, polyacrylamide, polymethylmethacrylate, polysulfone, polyacrylonitrile and nylon, or those obtained by incorporating a reactive functional group such as sulfonic group, amino group or the like to the synthetic resins can be recited. Also, glass, polysaccharides or derivatives thereof (cellulose, nitrocellulose and the like), silica gel, porous ceramics, metal oxides and the like may be used.

[0083] The "support" may be a microparticle, and a microparticle as same as the support may be bound to the detection oligonucleotide. As the microparticles, latex beads, gold colloids (gold nanoparticles) and the like are recited.

20 **[0084]** When the same kinds of microparticles are bound to the support and the detection oligonucleotide, the microparticle serving as the support, and the microparticle bound to the detection oligonucleotide are able to concurrently bind and aggregate on the methylated DNA having a target DNA region. This means that as a result of formation of a detection complex by binding of the immobilized methylated DNA antibody, the detection oligonucleotide, and the methylated single-stranded DNA, the microparticle serving as the support and the microparticle bound to the detection oligonucleotide form an aggregate. In this case, when the microparticle is a latex bead, the aggregate can be detected by change in turbidity. When the microparticle is a gold colloid (gold nanoparticle), the aggregate can be detected by change in color tone (pink to purple).

25 **[0085]** When a plurality of methylated DNA antibodies immobilized to supports are bound concurrently on DNA having one target DNA region, aggregation of microparticles can be detected by binding of the methylated DNA antibodies immobilized on microparticles, with plural methylated DNAs on the DNA having a target DNA region. For example, when the support is a latex bead, an aggregate of microparticles can be detected by change in turbidity. When the support is a gold colloid (gold nanoparticle), an aggregate of microparticles can be detected by color tone change (from pink to purple). In this case, an equivalent result to that obtained by adding the detection oligonucleotide can be obtained even when the detection oligonucleotide is not added.

30 **[0086]** An amount detected by aggregation of microparticles is correlated with a sum of DNA methylated by Second step. When DNA obtained in First step is DNA contained in a cell of a tissue, the degree of methylation of DNA contained in the cell of the tissue differs depending on the tissue, so that the degree of methylation of DNA obtained in Second step includes both the degree of methylation in the cell of the tissue, and the degree of methylation in Second step. That is, an amount detected by an aggregate of microparticles when a DNA methylation enzyme treatment is not executed in Second step is a value correlated with the degree of methylation in the cell of the tissue.

35 **[0087]** "Methylated DNA antibody" is an antibody that binds with a methylated base in DNA as an antigen. Concretely, methylcytosine antibody can be recited, and an antibody having the property of recognizing cytosine whose position 5 is methylated in single-stranded DNA and binding thereto can be recited. Commercially available methylated DNA antibodies may also be used as far as they specifically recognize DNA in a methylation state described in the present specification, and are capable of specifically binding thereto. Such a methylated DNA antibody can be prepared in a conventional method using a methylated base, methylated DNA or the like as an antigen. For concretely preparing a methylcytosine antibody, selection is made according to specific binding to methylcytosine in DNA as an indicator from antibodies prepared using DNA containing 5-methylcytosine, 5-methylcytosine, or 5-methylcytosine as an antigen. Considering the property of such immobilized methylated-DNA antibody (the fact that one antibody binds to one methylated base (cytosine)), improvements in quantification accuracy and detection sensitivity are expected by selecting the region where a number of methylated bases (cytosine), namely CpG, are present, as the target DNA region.

40 **[0088]** Known as antibodies that can be obtained by immunizing an animal with an antigen are an antibody of IgG fraction (polyclonal antibody), an antibody producing a single clone (monoclonal antibody) and the like that can be obtained by immunizing an animal with an antigen. In the present invention, since an antibody capable of specifically recognizing methylated DNA or methylcytosine is preferred, use of a monoclonal antibody is advisable.

45 **[0089]** As a method of preparing a monoclonal antibody, a procedure based on a cell fusion method can be recited. For example, in the cell fusion method, a hybridoma is prepared by allowing cell fusion between a spleen cell (B cell) derived from an immunized mouse and a myeloma cell, and an antibody produced by the hybridoma is selected for

preparation of a methyl cytosine antibody (monoclonal antibody). When a monoclonal antibody is prepared by a cell fusion method, it is not necessary to purify an antigen, and for example, a mixture of 5-methyl cytidine, 5-methyl cytosine or DNA or the like containing 5-methyl cytosine may be administered as an antigen to an animal used for immunization. As an administration method, 5-methyl cytidine, 5-methyl cytosine or DNA or the like containing 5-methyl cytosine is directly administered to a mouse for production of an antibody. When an antibody is difficult to be produced, an antigen bound to a support may be used for immunization. Also, by thoroughly mixing an adjuvant solution (prepared, for example, by mixing liquid paraffin and Aracel A, and mixing killed tubercle bacilli as an adjuvant) and an antigen and administering the same, or immunizing via liposome incorporating the same, immunity of an antigen can be improved. Also a method involving adding equivalent amounts of a solution containing an antigen and an adjuvant solution, fully emulsifying them, and subcutaneously or intraperitoneally injecting the resultant mixture to a mouse, and a method of adding killed Bordetella pertussis as an adjuvant after mixing well with alum water are known. A mouse may be boosted intraperitoneally or intravenously after an appropriate term from initial immunization. When the amount of an antigen is small, a solution in which the antigen is suspended may be directly injected into a mouse spleen to effect immunization.

[0090] After exenterating a spleen and peeling an adipose tissue off after several days from the final immunization, a spleen cell suspension is prepared. The spleen cell is fused, for example, with an HGPRT-deficient myeloma cell to prepare a hybridoma. As a cell fusion agent, any means capable of efficiently fusing a spleen cell (B cell) and a myeloma cell is applicable, and for example, a method of using a hemagglutinating virus of Japan (HVJ), polyethyleneglycol (PEG) and the like are recited. Cell fusion may be conducted by a method using a high voltage pulse.

[0091] After the cell fusion operation, cells are cultured in an HAT medium, a clone of a hybridoma in which a spleen cell and a myeloma cell are fused is selected, and the cell is allowed to grow until screening becomes possible. In a method of detecting an antibody for selecting a hybridoma that produces an intended antibody, or a method of measuring a titer of an antibody, an antigen-antibody reaction system may be used. Concretely, as a method of measuring an antibody against a soluble antigen, a radioisotope immune assay (RIA), an enzyme-linked immunosorbent assay (ELISA) and the like can be recited.

[0092] Single-stranded DNA is able to bind with an anti-methylation antibody as far as CpG existing therein is methylated at least at one site. Therefore, the term "methylated" in the present method means DNA in which CpG existing therein is methylated at least at one site, and is not limited to DNA in which every CpG existing therein is methylated.

[0093] The "detection complex" in Fourth step means a complex made up of the test DNA complex obtained in Third step and the immobilized methylated DNA antibody bound to each other. Since the methylated DNA antibody is used as an entity that can be immobilized to a support, it suffices that the methylated DNA antibody can be eventually immobilized to the support in the condition that the test DNA complex of the single-stranded DNA containing methylated DNA in the target DNA region and the detection oligonucleotide is formed, and

- (1) the methylated DNA antibody may be immobilized to the support in the stage prior to binding of the test DNA complex and the methylated DNA antibody, or
- (2) the methylated DNA antibody may be immobilized to the support in the stage posterior to binding of the test DNA complex and the methylated DNA antibody.

[0094] One exemplary concrete method for immobilizing a methylated DNA antibody to a support involves immobilizing a biotinylated methylated DNA antibody obtained by biotinylating a methylated DNA antibody to a streptavidin-coated support (for example, a PCR tube coated with streptavidin, magnetic beads coated with streptavidin, a chromatostrip partially coated with streptavidin and so on).

[0095] Also there is a method of letting a molecule having an active functional group such as an amino group, a thiol group, or an aldehyde group covalently bind to a methylated DNA antibody, and letting the resultant conjugate covalently bind to a support made of glass, a polysaccharide derivative, silica gel or the synthetic resin or thermostable plastic whose surface is activated by a silane coupling agent or the like. The above described covalent bonding may be achieved, for example, by covalently coupling the molecule having an active functional group to a methylated DNA antibody using a spacer formed by serially connecting five triglycerides, a cross linker or the like.

[0096] A methylated DNA antibody may be directly immobilized to a support, or an antibody against a methylated DNA antibody (secondary antibody) may be immobilized to a support, and a methylated antibody may be bound to the secondary antibody to achieve immobilization to a support.

[0097] The expression "making the immobilized methylated DNA antibody bind with a test DNA complex to obtain a detection complex" which is Fourth step means making the single-stranded methylated DNA contained in the test DNA complex obtained in Third step bind with the immobilized methylated DNA antibody to immobilize it to the support. For example, when "the methylated DNA antibody is immobilized to the support in the stage prior to binding of the test DNA complex and the methylated DNA antibody", concretely, for example, as the methylated DNA antibody that can be immobilized to the support, a "biotin-labeled biotinylated methylated DNA antibody" is used, and when the DNA sample derived from a specimen is a DNA sample derived from genome contained in a specimen derived from a biological

specimen, the practice may be executed in the following manner.

(a) An appropriate amount (for example, 4 $\mu\text{g}/\text{mL}$ solution, 100 $\mu\text{L}/\text{well}$) of a biotinylated methylated DNA antibody is added to an avidin-coated plate, and then left still at room temperature, for example, for about 2 hours to promote immobilization of the biotinylated methylated DNA antibody and streptavidin. Thereafter, the remaining solution is removed and washed. A washing buffer (for example, 0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)) is added in an amount of, for example, 300 $\mu\text{L}/\text{well}$, and the solution is removed. This operation is repeated several times and the biotinylated methylcytosine antibody that is immobilized to the support is left on the well.

(b) A DNA sample derived from genome contained in a specimen derived from a biological specimen is added with 5 μL of NEBuffer2 (available from NEB Inc.), 0.5 μL of S-adenosyl methionine (3.2 mM, available from NEB Inc.), and 0.5 μL of cytosine methylation enzyme SssI (available from NEB Inc.), and then the mixture is added with sterilized ultra pure water to make the liquid amount 50 μL , and incubated at 37°C for 30 minutes. Then methylated single-stranded DNA obtained by separating double-stranded DNA, a detection oligonucleotide, and an annealing buffer (for example, 33 mM Tris-Acetate pH 7.9, 66 mM KOAc, 10 mM MgOAc_2 , 0.5 mM Dithiothreitol) are mixed, and heated at 95°C, for example, for several minutes. Then for forming a conjugate of single-stranded DNA containing a target DNA region and a detection oligonucleotide, the mixture is rapidly cooled to the temperature lower than T_m value of the detection oligonucleotide by about 10 to 20°C, and retained at this temperature for example, for several minutes, and then returned to room temperature (the conjugate formed in this stage includes a conjugate of single-stranded DNA not containing DNA methylated in a target region and a detection oligonucleotide, as well as a conjugate of methylated single-stranded DNA containing DNA methylated in a target DNA region and a detection oligonucleotide).

(c) The formed conjugate of methylated single-stranded DNA and a detection oligonucleotide is added to an avidin-coated plate to which a biotinylated methylated DNA antibody is immobilized, and then left still at room temperature for about 3 hours to promote formation of a complex of a biotinylated methylated DNA antibody, methylated single-stranded DNA containing DNA methylated in a target DNA region, among the methylated single-stranded DNAs, and the detection oligonucleotide (formation of complex) (in this stage, if the DNA sample containing a target DNA region is not methylated, the conjugate of single-stranded DNA and a detection oligonucleotide will not form a complex with a methylated DNA antibody). Thereafter, the remaining solution is removed and washed. A washing buffer (for example, 0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)) is added in an amount of, for example, 300 $\mu\text{L}/\text{well}$, and the solution is removed. This washing operation is repeated several times to leave the complex on the well (separation of complex).

[0098] As the annealing buffer used in (b), those suited for binding the detection oligonucleotide and methylated single-stranded DNA containing a target DNA region may be used without limited to the aforementioned annealing buffer. Stability of binding increases when an annealing buffer in which a divalent ion, preferably a magnesium ion is dissolved at a concentration of 1 to 600 mM is used.

[0099] The washing operation in (a) and (c) is important for removing an unimmobilized methylated DNA antibody suspended in the solution, unmethylated single-stranded DNA that is not bound to a methylated DNA antibody and suspended in the solution, methylated single-stranded DNA in a region other than the target region that does not form a conjugate with the detection oligonucleotide and the like, from the reaction solution. The washing buffers may be those suited for removing the free methylated DNA antibody, methylated single-stranded DNA suspended in the solution and the like, and a DELFIA buffer (available from Perkin Elmer, Tris-HCl pH 7.8 with Tween 80), a TE buffer and the like may also be used without limited to the aforementioned washing buffer.

[0100] In the above (a) to (c), after binding of the methylated single-stranded DNA obtained by methylating the target DNA region, and the detection oligonucleotide, a biotinylated methylated DNA antibody is caused to bind with the resultant conjugate to form a complex, however, the order is not limited thereto. That is, after causing the methylated single-stranded DNA containing a target DNA region to bind with the biotinylated methylated DNA antibody, the detection oligonucleotide may be bound to the resultant conjugate to form a complex. For example, a biotinylated methylated DNA antibody immobilized to a support coated with streptavidin may be added with a DNA sample derived from genome, 5 μL of NEBuffer2 (available from NEB Inc.), 0.5 μL of S-adenosyl methionine (3.2 mM, available from NEB Inc.), and 0.5 μL of cytosine methylation enzyme SssI (available from NEB Inc.), and then the resultant mixture is added with sterilized ultra pure water to make the liquid amount 50 μL , and incubated at 37°C for 30 minutes, and then added with methylated single-stranded DNA obtained by separation from double-stranded DNA, to thereby form a conjugate of the biotinylated methylated DNA antibody immobilized to the support and the methylated single-stranded DNA (the conjugate formed in this stage includes a conjugate of methylated single-stranded DNA methylated in a region other than the target DNA region and a methylated antibody, as well as a conjugate of methylated single-stranded DNA containing DNA methylated in a target DNA region and the methylated antibody). Thereafter, the detection oligonucleotide may be added thereto to

form a complex with the conjugate having methylated single-stranded DNA containing DNA methylated in a target DNA region, among the aforementioned bound bodies, and the complex may be separated (in this stage, the conjugate of the methylated single-stranded DNA containing DNA methylated in a region other than the target DNA region and the methylated antibody will not form a complex).

5 **[0101]** The operations of the above (a) to (c) may be conducted using a chromatostrip. In this case, the operations are concretely conducted as follows. For example, first, an appropriate amount of a biotinylated methylated DNA antibody is developed by a chromatostrip that is partly coated with streptavidin. By this operation, the biotinylated methylated DNA antibody is immobilized to the part coated with streptavidin. Then the conjugate of the methylated single-stranded DNA and the detection oligonucleotide obtained in (b) (the conjugate formed in this stage includes a conjugate of single-
10 single-stranded DNA not containing DNA methylated in a target region and a detection oligonucleotide, as well as a conjugate of methylated single-stranded DNA containing DNA methylated in a target DNA region and a detection oligonucleotide) is developed by the chromatostrip. Through these operations, a complex made up of the methylated single-stranded DNA obtained by adding a DNA sample derived from genome DNA, 5 μ L of NEBuffer2 (available from NEB Inc.), 0.5 μ L of S-adenosyl methionine (3.2 mM, available from NEB Inc.), and 0.5 μ L of cytosine methylation enzyme Sssl (available from NEB Inc.), and adding the resultant mixture with sterilized ultra pure water to make the liquid amount 50 μ L, and incubating at 37°C for 30 minutes, followed by separation from double-stranded DNA, the detection oligonucleotide, and the biotinylated methylated DNA antibody is immobilized to the part coated with streptavidin (formation of complex and immobilization to support)(in this stage, the conjugate of single-stranded DNA not containing DNA methylated in a target DNA region and a detection oligonucleotide fails to form a complex). The order of operations for
20 formation of a complex is not limited to this order of operations. For example, after forming a complex made up of the single-stranded DNA containing DNA methylated in a target DNA region, the detection oligonucleotide, and the biotinylated methylated DNA antibody, this may be developed by a chromatostrip, and the complex may be immobilized to the part coated with streptavidin. In these operations, unnecessary components can be removed and a washing operation can be omitted by developing the solution by the chromatostrip. Between these operations, a washing operation (development of the chromatostrip by a washing buffer (for example, 0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)) may be executed.

25 **[0102]** Fifth step is a step of quantifying or detecting DNA having a target DNA region in the single-stranded DNA by quantifying or detecting the detection oligonucleotide contained in the detection complex obtained in Fourth step according to its identification function.

30 **[0103]** The term "detection" used herein means determining whether the degree of methylation of DNA methylated up to Second step is a certain degree or more according to the identification function of the detection oligonucleotide. That is, it usually means that a significant value is obtained by the identification function as will be described later.

35 **[0104]** The term "quantification" means quantification of an amount detected by the identification function determined in Fifth step. That is, it means that a value correlated with a total amount of the DNA having a target DNA region obtained in First step, and the methylated DNA methylated by DNA methylation enzyme treatment in Second step is obtained. For example, the quantified value of the amount of the methylated DNA antibody or the present oligonucleotide as the identification function as will be described later is a value correlated with the amount of DNA in the target DNA region in the specimen, and, for example, when the specimen is 1 mL of serum, it means acquiring a value correlated with a total amount of the DNA having a target DNA region obtained in First step and the methylated DNA methylated by DNA
40 methylation enzyme treatment in Second step, among the DNAs of the target region contained in 1 mL of serum.

45 **[0105]** The term "identification function" in Fifth step is a function capable of detecting or quantifying the detection oligonucleotide. The identification function may be any function possessed by the detection oligonucleotide, and for example, an identification function based on labeling of the detection oligonucleotide, and an identification function imparted to the detection oligonucleotide by a detection molecule binding to the detection oligonucleotide can be recited. Concretely, a fluorescent or a chromogenic property of the detection oligonucleotide labeled with europium, gold colloid, latex bead, radioisotope, fluorescent substance (FITC or the like), horseradish Peroxidase (HRP), alkaline phosphatase, or the like at its 5'-end or 3'-end are recited.

50 **[0106]** For detection of europium, after adding and mixing Enhancement Solution (available from PerkinElmer), and keeping still at room temperature for about 45 minutes, fluorescence (excitation 340nm/fluorescence 612 nm) may be measured by a fluorescent detector. When the detection oligonucleotide is a methylated oligonucleotide, as a detection molecule, concretely, a methylated DNA antibody, an osmium complex (J. Am. Chem. Soc., 2007;129:5612-5620) and the like can be recited. The term "methylated oligonucleotide" means an oligonucleotide wherein at least one of bases of nucleotides constituting the oligonucleotide is methylated, and more concretely, means oligonucleotides including 5-methylcytosine, 6-methyladenine and the like. Further, when the detection oligonucleotide is labeled with FITC, a FITC
55 antibody can be recited as a detection molecule.

[0107] When the detection molecule is a methylated DNA antibody, the function capable of quantifying or detecting by being bound to the antibody can impart an identification function to the antibody. However, a methylated DNA antibody having substrate cross reactivity with an immobilized methylated DNA antibody cannot be used. For example, when an

immobilized methylated DNA antibody uses a methylcytosine antibody, a methylated DNA antibody capable of binding with methylcytosine cannot be used. Concretely, labels such as europium label, gold colloid label, latex bead label, radioisotope label, fluorescent substance (FITC or the like) label, horseradish Peroxidase (HRP) label, alkaline phosphatase label, biotin label and the like are fluorescent, chromogenic and the like functions. As a method of imparting an identification function to the antibody as a detection molecule, the identification function may be directly bound to an antibody which is a detection molecule, or a secondary antibody or a tertiary antibody having an identification function may be bound to an antibody which is a detection molecule. Concretely, since an antibody labeled with a fluorescent substance, an antibody labeled with horseradish Peroxidase (HRP), an antibody labeled with alkaline phosphatase, an antibody labeled with biotin, and an antibody labeled with europium are commercially available, they may be used as a secondary antibody or a tertiary antibody. Also an antibody to which a substrate detectable by an enzyme cycle method is bound may be used. As a means for quantifying or detecting these functions, for example, measurement by a radiation detector, a spectrophotometer and the like, or visual observation and the like are recited. For example, as a case of detecting or quantifying a detection oligonucleotide according to its identification function, when a secondary antibody to which europium is added as a concretely detectable or quantifiable function, Enhancement Solution (available from PerkinElmer) is added and mixed after the detection complex is bound to the secondary antibody, and left still at room temperature for about 45 minutes. Thereafter, fluorescence (excitation 340 nm/fluorescence 612 nm) may be measured by a fluorescent detector.

[0108] When a methylated DNA antibody (methylated DNA antibody having substrate specificity different from that of an immobilized methylated DNA antibody) is caused to bind with methylated DNA on the detection oligonucleotide and detection or quantification is executed according to its function, concretely, when the own characteristic of the antibody is used as the function, the operation may be executed in the following manner. After causing the methylated DNA antibody to bind with the complex, a secondary antibody against the methylated DNA antibody (for example, Eu-N1-labeled mouse IgG antibody: available from PerkinElmer) is added, and left still at room temperature for about 1 hour, to thereby promote binding of the secondary antibody to the complex. Thereafter, Enhancement Solution (available from PerkinElmer) is added and mixed, and left still at room temperature, for example, for about 45 minutes. Then, by measuring fluorescence (excitation 340 nm/fluorescence 612 nm) by a fluorescent detector, detection or quantification is conducted.

[0109] When the methylated DNA antibody binding to methylated DNA on the detection oligonucleotide is labeled with FITC, an antibody to which FITC is bound may also be used as a secondary antibody. In this case, fluorescence of FITC may be measured by a known method to achieve detection or quantification, or detection or quantification may be achieved by using an anti-FITC antibody as a secondary antibody. Further, when FITC is directly bound to the detection oligonucleotide, FITC may be used as an identification function, or labeling function may be imparted by a horseradish Peroxidase (HRP)-labeled FITC antibody, an alkaline phosphatase-labeled FITC antibody, a biotin-labeled FITC antibody, an europium-labeled FITC antibody and the like. Concretely, as the detection oligonucleotide, when a FITC-labeled oligonucleotide is used as the detection oligonucleotide, a detection complex immobilized to a support containing the detection oligonucleotide is added with an antibody labeled with horseradish Peroxidase (HRP) (for example, HRP-labeled FITC antibody (available from Jackson ImmunoResearch Laboratories)), and left still at room temperature for about 1 to 2 hours, to promote binding of the FITC antibody to the detection complex. Then after washing and removing the FITC antibody solution, an appropriate substrate (for example, Substrate Reagent Pack #DY999: available from R&D SYSTEMS) is added and mixed. After leaving still at room temperature for about 5 to 60 minutes, a stop solution (2N H₂SO₄ aqueous solution) is added to stop the reaction of horseradish Peroxidase (HRP), and absorbance at 450 nm may be measured within 30 minutes after stopping of the reaction.

[0110] When biotin is not used for immobilization of the methylated DNA antibody, a biotinylated detection oligonucleotide can be used for detection or quantification. When a biotinylated detection oligonucleotide is detected or quantified, for example, HRP-labeled streptavidin is added and mixed with a immobilized detection complex to form and separate a conjugate of the biotinylated detection oligonucleotide and HRP-labeled streptavidin, and then activity of HRP is measured by a known method to thereby detect or quantify the biotinylated methylated DNA antibody.

[0111] The identification function may utilize a substrate used in a high sensitivity detection method such as an enzyme cycle method and the like. Concretely, an antibody to which an enzyme used in an enzyme cycle method may be bound as a detection molecule to the detection complex. The identification function imparted to the detection molecule in the present method is not limited to the methods as described above.

[0112] It suffices that the "detection molecule" has a property of detecting or quantifying a detection oligonucleotide. The detection molecule may recognize a detection sequence of a detection oligonucleotide, or may be preliminarily bound to a detection oligonucleotide. That is, the detection molecule has a property of specifically binding to a detection oligonucleotide and has an "identification function" which is the function or characteristic that is utilized for quantification or detection or is a molecule capable of being provided with the identification function. Concretely, it suffices that the detection molecule is able to bind with a methylated oligonucleotide and detect the methylated oligonucleotide, and specifically binds with the methylated oligonucleotide to exhibit the identification function when the detection sequence is the methylated oligonucleotide (a methylated DNA antibody having substrate cross reactivity with the immobilized

methylated DNA antibody cannot be used. For example, when the immobilized methylated DNA antibody uses a methylcytosine antibody, a methylcytosine antibody cannot be used as a detection molecule). As others, for example, the detection molecule may be a methylated DNA antibody. However, a methylated DNA antibody having substrate cross reactivity with the immobilized methylated DNA antibody cannot be used. For example, when the immobilized methylated DNA antibody uses a methylcytosine antibody, the methylcytosine antibody cannot be used as a detection molecule. When the detection sequence is a detection molecule itself, it is not necessary to add a new detection molecule for detecting the detection oligonucleotide, and by detecting the detection molecule incorporated into the detection oligonucleotide, it becomes possible to detect the detection oligonucleotide.

[0113] As a method of quantifying or detecting minor substances contained in a biological sample such as blood or urine, immunological measuring methods are generally used. Among such methods, what is called immuno chromatography using chromatography is currently widely used in various situations including, for example, clinical examinations in hospitals, assays in laboratories and the like because of its simple operation and short time required for assay. In recent years, what is called hybrid chromatography has been utilized in which labeled DNA (gene) is developed on a chromatostrip, and target DNA (gene) is detected by hybridization using a probe capable of capturing the target DNA (gene). Also this method is now coming to be widely used in situations including, for example, clinical examinations in hospitals, assays in laboratories and the like because of its simple operation and short required time for assay. The present method conceptually enables a combined method of the immuno chromatography and the hybrid chromatography. In the present method, since the order of formation of a complex and selection of a complex is not particularly limited, various methods are possible. Concretely, such methods may be executed in the following manner.

[0114] Method 1: A sample directly after end of Second step is added with a detection oligonucleotide having an identification function to form a conjugate of methylated single-stranded DNA containing a target DNA region and the detection oligonucleotide having an identification function (the conjugate formed in this stage includes a conjugate of single-stranded DNA not containing DNA methylated in a target DNA region and a detection oligonucleotide, as well as a conjugate of single-stranded DNA containing DNA methylated in a target DNA region and a detection oligonucleotide), and then added with a biotinylated methylated antibody, to form a complex in which a single-stranded DNA containing DNA methylated in a target DNA region, the detection oligonucleotide having an identification function, and a biotinylated methylated DNA antibody are bound. As the obtained sample is dropped (introduced) into an introduction part of a chromatostrip, the complex migrates in a development part by a capillary phenomenon, and is trapped in the part preliminarily coated with streptavidin. Thereafter, by detecting or quantifying the detection oligonucleotide contained in the obtained complex according to its identification function, DNA having a target DNA region can be detected or quantified.

[0115] Method 2: A sample directly after end of Second step is added with a biotinylated methylated DNA antibody, to form a conjugate of methylated single-stranded DNA having methylated cytosine (there exist single-stranded DNA containing a target DNA region and single-stranded DNA other than the target) and a biotinylated methylated DNA antibody (the conjugate formed in this stage includes a conjugate of methylated single-stranded DNA other than the target DNA region and a methylated antibody, as well as a conjugate of single-stranded DNA containing DNA methylated in a target DNA region and a methylated antibody). As the obtained sample is dropped (introduced) into an introduction part of a chromatostrip, the conjugate migrates in a development part by a capillary phenomenon, and is trapped in the part preliminarily coated with streptavidin (also in this stage, the conjugate includes the conjugate of methylated single-stranded DNA other than the target DNA region and a methylated antibody, as well as the conjugate of single-stranded DNA containing DNA methylated in a target DNA region and a methylated antibody). Thereafter, as the detection oligonucleotide having an identification function is dropped (introduced) into the introduction part, it migrates in the development part, and binds only to methylated single-stranded DNA containing a target DNA region, among the aforementioned bound bodies, to form a complex in which single-stranded DNA containing DNA methylated in a target DNA region, a detection oligonucleotide having an identification function, and a biotinylated methylated DNA antibody are bound. By detecting or quantifying the detection oligonucleotide contained in the obtained complex according to its identification function, it is possible to detect or quantify DNA having a target DNA region.

[0116] Method 3: As a biotinylated methylated DNA antibody is dropped (introduced) into an introduction part of a chromatostrip, the methylated DNA antibody migrates in a development part by a capillary phenomenon, and is trapped in the part preliminarily coated with streptavidin. Then as a sample directly after end of Second step is dropped (introduced) into the introduction part, it migrates in the development part, and single-stranded DNA having methylated cytosine is trapped by the already trapped methylated DNA antibody as the conjugate (the conjugate formed in this stage includes a conjugate of methylated single-stranded DNA other than the target DNA region and a methylated antibody, as well as a conjugate of single-stranded DNA containing DNA methylated in a target DNA region and a methylated antibody). Thereafter, as the detection oligonucleotide having an identification function is dropped (introduced) into the introduction part, it migrates in the development part, and binds only to methylated single-stranded DNA containing a target DNA region, among the aforementioned bound bodies, to form a detection complex in which single-stranded DNA containing DNA methylated in a target DNA region, a detection oligonucleotide having an identification function, and a biotinylated

methylated DNA antibody are bound. By detecting or quantifying the detection oligonucleotide contained in the obtained complex according to its identification function, it is possible to detect or quantify DNA having a target DNA region.

5 [0117] Method 4: As a biotinylated methylated DNA antibody is dropped (introduced) into an introduction part of a chromatostrip, the methylated DNA antibody migrates in a development part by a capillary phenomenon, and is trapped
10 in the part preliminarily coated with streptavidin. A sample directly after end of Second step is added with a detection oligonucleotide having an identification function, to form a test DNA complex which is a conjugate of methylated single-stranded DNA containing a target DNA region and a detection oligonucleotide having an identification function. Then as the obtained conjugate is dropped (introduced) into the introduction part, it migrates in the development part, and only single-stranded DNA containing DNA methylated in a target DNA region, among the aforementioned bound bodies,
15 binds to the methylated DNA antibody that is already trapped, to form a detection complex in which single-stranded DNA containing DNA methylated in a target DNA region, a detection oligonucleotide having an identification function, and a biotinylated methylated DNA antibody are bound. By detecting or quantifying the detection oligonucleotide contained in the obtained complex according to its identification function, it is possible to detect or quantify DNA having a target DNA region.

20 [0118] Also a plurality of detection sites may be provided in the target DNA region (using detection oligonucleotides respectively capable of complementarily binding to different target DNA regions), and each target DNA region may be sequentially detected or quantified, and also by using a detection oligonucleotide capable of complementarily binding to a plurality of target DNA regions such that a repetitive sequence in genome or an overlapping gene or a plurality of different genes are concurrently detected for enabling formation of complexes with a plurality of target DNA regions, the detection sensitivity can be dramatically improved. Also by designing a number of detection oligonucleotides in a single target region, and using them on the support side or on the detection side, the detection sensitivity can be dramatically improved.

25 [0119] As a method of executing the process of forming a complex of the detection oligonucleotide, the biotinylated methylated antibody, and the single-stranded DNA methylated in a target DNA region and binding it to the support, a method of using an immunological antibody method may be applied without limited to Methods 1 to 4 as described above. For example, the ELISA method allows execution of the process of forming a complex and binding it to a support in the order described in Methods 1 to 4 because it uses the principle similar to that of the chromatostrip method.

30 [0120] The DNA methylation enzyme, the detection oligonucleotide, or the methylated DNA antibody that can be used in the present method are useful as reagents for a detection kit. The present method also provides a detection kit containing such a DNA methylation enzyme, a specific oligonucleotide, or a methylated DNA antibody as a reagent, and a detection chip comprising such a present oligonucleotide, or a methylated DNA antibody immobilized on a support, and the scope of the present method includes use in the form of a detection kit or a detection chip as described above using the substantial principle of the present invention.

35 [0121] Generally, for examining presence or absence of a pathogenic microorganism contained in a biopsy sample or food, presence or absence of such a pathogenic microorganism is examined, or such a pathogenic microorganism is identified by a test based on immunization method for each microbial antigen. However, preparation of an antibody used for such a immunization method is not easy, and for detecting a plurality of pathogenic microorganisms, it is necessary to prepare antibodies against respective antigens of the pathogenic microorganisms. By using the present method, it is possible to realize a simple test-four pathogenic microorganisms without conducting such complicated
40 antibody preparation. Further, according to the present method, since nucleotide sequences of different pathogenic microorganisms can be tested at the same time, it is possible to detect several kinds of pathogenic microorganisms contained in one specimen at the same time. Concretely, food poisoning bacteria such as *Listeria monocytogenes*, *Salmonella enterica*, *Campylobacter jejuni* subsp. *Jejuni*, *Staphylococcus aureus*, *Vibrio parahaemolyticus*, *Bacillus cereus*, *Clostridium botulinum*, *Yersinia enterocolitica*, *Yersinia pseudotuberculosis* and *Clostridium perfringens* are known, however, a technique of detecting several kinds of these food poisoning bacteria at the same time is not known. However, by using the present method, it is possible to detect nucleotide sequences of several kinds of food poisoning bacteria at the same time. Further, when a nucleotide sequence found plurally in genome such as CRISPR (Clustered regularly interspaced short palindromic repeats) region is selected as a nucleotide sequence to be detected by a specific oligonucleotide, detection at higher sensitivity is realized compared to the case of detecting one gene in one genome.
45 Such a technique is useful also for diagnosis of infection and rapid detection of food poisoning bacteria. Further, the present method may be used for identification of an industrially useful bacterium, or for a simple test of a microbial community in soil, river or lake sediments and the like by detecting genomes of microorganisms in such environments. Of the microorganisms in such environments, inhabitation of, for example, *Methanococcus jannaschii*, *Methanobacterium thermoautotrophicum deltaH*, *Aquifex aeolicus*, *Pyrococcus horikoshii* OT3, *Archaeoglobus fulgidus*, *Thermotoga maritima* MSB8, *Aeropyrum pernix* K1, and *Haloferax mediterranei* can be verified. It is also possible to detect and identify industrially available bacteria such as *Geobacter sulfurreducens* and microorganisms used for fermentation such as *Streptococcus thermophilus*.

55 [0122] When the target DNA region in the present method is a nucleotide sequence derived from a microorganism, it

means genomic DNA or a DNA fragment extracted from a specimen, or a nucleotide sequence of DNA obtained from RNA extracted from a specimen by a reverse transcriptase. Therefore, as a nucleotide sequence capable of complementarily binding with a detection oligonucleotide, a region specific to the microorganism may be selected. For example, when the target DNA region in the present invention is a microbial nucleotide sequence, for selectively extracting the target region from the specimen, a nucleotide sequence that is peculiar to the microorganism in the vicinity of the target region may be selected as a nucleotide sequence that specifically binds with the specific oligonucleotide, from microbial genomic DNA, or nucleotide sequences of DNA that are prepared from RNA extracted from the specimen by a reverse transcriptase.

[0123] For example, applicable as a region where the target region for detecting genome in a microorganism and the specific oligonucleotide complementarily bind is, concretely, a nucleotide sequence not encoding a gene such as a region corresponding to the nucleotide number 271743-272083 of yeast chromosome VII (SEQ ID NO: 29) shown, for example, in Genbank Accession No. NC_001139, or the nucleotide number 384569-384685 of yeast chromosome VII (SEQ ID NO: 34) shown, for example, in Genbank Accession No. NC_001139. It is also useful to detect a nucleotide sequence conserved among pathogenic microorganisms in a characteristic gene that is common in various pathogenic microorganisms because a method of detecting a plurality of pathogenic microorganisms at the same time can be provided. Concretely, since mce-family gene (Mycobacterium tuberculosis), tRNA-Tyrnucleotide sequence on 13th chromosome (Cryptococcus neoformans), and chitin synthase activator (Chs3) have a nucleotide sequence peculiar to *Aspergillus fumigatus* and genus *Neosartorya*, they can be used for assay of infection by a microorganism, by assaying whether DNA derived from these microorganisms is contained in DNA extracted from a biopsy sample of human expectoration or lung. Further, since actA (*Listeria monocytogenes*), pyrG (NC 002163, *Campylobacter jejuni* subsp. *jejuni*) and the like are common genes peculiar to food poisoning bacteria, these genes may be used for a microbial assay in food poisoning. ThrA has a sequence that is conserved among *Salmonella enterica*, *Yersinia enterocolitica*, and *Escherichia coli*, so that a plurality of microorganisms can be detected by one gene.

[0124] Of nucleotide sequences published on a database, a nucleotide sequence peculiar to a microorganism may be retrieved, and a nucleotide sequence peculiar to a microorganism may be searched. For example, a nucleotide sequence on a published database such as PubMed may be obtained through regular procedure, and the obtained nucleotide sequence can be examined whether it is a peculiar nucleotide sequence by Blast search through regular procedure. The peculiar nucleotide sequence means that a nucleotide sequence in a detection object does not include a nucleotide sequence having homology with a nucleotide sequence derived from an organism other than the genomic nucleotide sequence of the microorganism to be detected.

[0125] In particular, when the specimen is a human biopsy sample, it is important to design a specific oligonucleotide that would not complementarily bind with human genes. Similarly, when the specimen is food, it is important to design a specific oligonucleotide that would not complementarily bind with a nucleotide sequence derived from an organism other than the detection object contained in the food.

[0126] For detecting free DNA in blood, any region correlated with an amount of free DNA may be used, and when it is intended to quantify or detect free DNA, what is called repetitive sequence where the same sequence in genome appears repetitively, several or more times, is preferred, and a simple repetitive sequence (called tandem repetitive sequence, or tandem repeat), and an interspersed repeat sequence are more preferred.

[0127] The simple repetitive sequence is **characterized in that** the same sequences neighbor in the same orientation, and a series of nucleotide sequences such as satellite DNA, minisatellite, microsatellite, centromere, telomere, kinetochore, and ribosome group genes are known.

[0128] The interspersed repetitive sequence is **characterized in that** the same sequences are interspersed without neighboring each other, and is believed to be DNA derived from retrotransposon. Interspersed repetitive sequences are classified into SINE (Short Interspersed Repetitive Element: short chain interspersed repetitive sequence) and LINE (Long Interspersed Elements: long chain interspersed repetitive sequence) depending on the length of the nucleotide sequence, and as a human nucleotide sequence, Alu sequence and LINE-1 sequence are respectively known as representative repetitive sequences. Also an inactive processed pseudogene that is reverse transcribed from RNA or protein, and a gene sequence amplified by gene duplication are also known.

[0129] The term duplicated gene indicates the case where a plurality of genes having high homology exist on one genome, and is, in many cases, a nucleotide sequence that exists in tandem near one gene. It is often the case where a pseudogene is one of duplicated genes.

[0130] As concrete examples of the repetitive sequence, such sequences as (A)_n, (T)_n, (GA)_n, (CA)_n, (TAA)_n, (GGA)_n, (CAGC)_n, (CATA)_n, (GAAA)_n, (TATG)_n, (TTTG)_n, (TTTA)_n, (TTTC)_n, (TAAA)_n, (TTCA)_n, (TATAA)_n, (TCTCC)_n, (TTTCC)_n, (TTTAA)_n, (TTTTTC)_n, (TTTTA)_n, (TTTTG)_n, (CAAAA)_n, (CACCC)_n, (TATATG)_n, (CATATA)_n, (TCTCTG)_n, (AGGGGG)_n, (CCCCCA)_n, and (TGGGGG)_n (n means a number of repetition) are known as repetition comprising a relatively short nucleotide sequence, and as a sequence derived from a transcription factor, MER1-Charlie, and Zaphod of hAT group, and MER2-Tigger, Tc-1, and Mariner of Tc-1 group can be recited. As others, concretely, Tigger1, Tigger2a, Tigger5, Charlie4a, Charlie7 and the like are known. These sequences are generally short and simple nucleotide se-

quences, and are difficult to set the specific adhesion sequence and a detection adhesion sequence as will be described later, however, these sequences can be used in the present method as far as they have a sequence that can be set into setting objects of the specific adhesion sequence and a detection adhesion sequence as will be described later. Therefore, it is not necessarily excluded as an object of the present method. Further, satellite DNA, minisatellite, microsatellite and the like are repetitive sequences classified into simple repetitive sequences.

[0131] Further, as a sequence having multi-copies in gene, ALR6 as a sequence existing in centromere, U2 and U6 as snRNA, as well as the genes such as tRNA and rRNA that are generally known to have multi-copies in genome, and the genes that have plural copies in genome as a result of gene duplication are recited.

[0132] It is also known that a retrovirus, a retrotransposon having LTR (Long terminal repeat) in its terminal, an endogenous sequence such as MaLRs (Mammalian apparent LTR-Retrotransposons) considered to be derived from viruses, and LTR derived from a retrovirus exist in multicopy in one genome.

[0133] For example, as the LTR derived from a retrovirus, concretely, subfamilies such as LTR1, LTR1B, LTR5, LTR7, LTR8, LTR16A1, LTR16A1, LTR16C, LTR26, LTR26E, MER48, and MLT2CB are known. The LTRs derived from a retrotransposon are classified into classes of ERV, ERVK and ERVL, and concrete examples include subfamilies such as LTR8A, LTR28, MER21B, MER83, MER31B, MER49, MER66B, HERVH, ERVL, LTR16A1, LTR33A, LTR50, LTR52, MLT2A1, MLT2E, MER11C, and MER11C. Further, MaLRs indicate DNA factors including LTRs in both ends likewise a typical retrotransposon, wherein an internal sequence sandwiched between LTRs is not derived from a retrovirus. For example, subfamilies such as MLT1A1, MLT1A2, MLT1B, MLT1C, MLT1D, MLT1F, MLT1G, MLT1H, MLT1J, MLT1K, MLT1I, MLT2CB, MSTA, MSTA-int, MSTB, THE1A, THE1B, THE1B-internal, and THE1 can be recited.

[0134] The interspersed repetitive sequences are **characterized in that** the same sequences are interspersed without neighboring each other, and are considered to be derived from a retrotransposon. Further, the interspersed repetitive sequences are classified into SINE (Short Interspersed Repetitive Element: short chain interspersed repetitive sequences) and LINE (Long Interspersed Elements: long-chain interspersed repetitive sequences) according to the length. Most of SINEs are sequences belonging to the Alu family. A common feature is that it has a sequence of 3'-side or a sequence of 5'-side of 7SL RNA, and that it has an AT-Rich region sandwiched between a Left-monomer and a Right-monomer. As subfamilies of the Alu family, Alu, AluJb, AluJo, AluSc, AluSg, AluSp, AluSq, AluSx, AluY, and FAM (Fossil Alu Monomer), FLAM (Free Left Alu Monomer) having a sequence of FAM, and FRAM (Free Right Alu Monomer) can be recited. As SINEs other than the Alu family, MIR, and Ther/MIR3 are known, and MIR and MIR3 are known as respective subfamilies. As subfamilies of the Alu family including other biological species, B1, B2, B4, PB1, PB1D and so on are known. As LINEs, subfamilies of LINE1 to Line23 are reported, and it is known that subfamilies such as LINE-1, LINE2, and LINE3 broadly exist in a genome. As for LINE-1, for example, L1M1, L1M2, L1M3, L1M3d, L1M4, L1M4c, L1MA2, L1MA7, L1MA8, L1MA9, L1MB1, L1MB1, L1MB3, L1MB4, L1MB5, L1MB6, L1MB7, L1MCa, L1MCb, L1MC2, L1MC3, L1MC4, L1MC4a, L1MC5, L1MDa, L1ME, L1MEc, L1MEd, L1MEg, L1ME1, L1ME2, L1ME3, L1ME3A, L1ME3B, L1ME4a, L1PB3, L1P4, L1PA2, L1PA3, L1PA4, L1PA5, L1PA6, L1PA7, L1PA10, L1PA12, L1PA13, L1PA14, L1PA16, L1PB1, L1PB3, L1PB4, L1PREC2, and HAL1 are known, and as LINE-2, subfamilies such as L2 and L2c are known. For example, if the later-described specific adhesion sequence and the detection adhesion sequence can be set, for a sequence common to the Alu family or subfamilies of Alu, or the LINE-1 family or subfamilies of LINE-1, a plurality of detection objects can be set in one genome, so that sensitivity of genome detection can be improved.

[0135] As a target DNA region, concretely, for example, a partial sequence of LINE-1 (the nucleotide sequence of SEQ ID NO: 37), a partial sequence of Alu (the nucleotide sequence of SEQ ID NO: 39) or nucleotide sequences having homology to these sequences can be recited.

[0136] For example, when a repetitive sequence in a certain region needs to be examined, databases such as Repbase (<http://www.girinst.org/repbase/>) and RepeatMasker (<http://www.repeatmasker.org/>) may be used because it is difficult to retrieve a general sequence retrieving database such as PubMed. If a specific adhesion sequence of the present method can be set, the detection sensitivity can be improved. Measuring these repetitive sequences can be treated, for example, as a surrogate marker of a free DNA amount in blood, and can be utilized for identification of an organism species when an organism species-specific repetitive sequence is noted.

[0137] In the present method, by measuring a repetitive sequence, a nucleotide sequence existing plurally in one genome can be measured concurrently. For example, a nucleotide sequence having a sequence homology of 80% or higher with the nucleotide sequence of SEQ ID NO: 37 has about 280 copies in a human genome, and a nucleotide sequence having a sequence identity of 80% or higher with the nucleotide sequence of SEQ ID NO: 39 has about 820 copies in a human genome. Therefore, if a specific adhesion sequence can be set in each nucleotide sequence, the detection sensitivity of one genome can be improved to 280 to 820 folds theoretically, compared to the case where a specific adhesion sequence is set for a sequence having just one kind in genome.

[0138] A duplicated gene means a gene or a gene fragment that is generated by doubling of a specific gene or gene fragment in genome due to gene duplication. Gene duplication is a phenomenon that a certain region of DNA including a gene is overlapped. As a cause of gene duplication, abnormality of gene recombination, translocation of retrotransposon,

duplication of the entire chromosome and the like are recited. For example, it means that one gene is copied and inserted into genomic DNA, and the copy is inserted to a different chromosome site in some cases, and inserted near the original gene in the other cases. The site where copied genes are aligned as a result of insertion near the original gene is called a tandem repeat, and a group of genes generated by gene duplication is called a gene family.

5 **[0139]** A pseudogene means a gene having a characteristic nucleotide sequence that is assumable to have encoded a gene product (particularly protein) in a sequence of DNA, but currently losing the function. It is assumed that it is generated as a result of mutation of the original functioning sequence. For example, there is the case where a stop codon arises by mutation and a peptide chain of a protein is shortened, so that the function as a protein is no longer effective, and there is the case where a function of a regulatory sequence required for normal transcription is impaired due to mutation such as single nucleotide substitution. In many pseudogenes, the original normal genes are remained separately, however, those becoming pseudogenes by themselves are also known.

10 **[0140]** Pseudogenes are classified into three types according to the characteristic of the gene sequence. There are known the case where DNA prepared from mRNA by a reverse transcriptase of retrotransposon is inserted into genome (processed pseudogene), the case where an original gene sequence is duplicated in genome, and a part of the copies loses the function due to mutation or the like to become a pseudogene (duplicated pseudogene or non-processed pseudogene), and the case where gene in genome (in the condition of single gene with no duplicated gene) loses the function to become a pseudogene.

15 **[0141]** Currently, among the genes known as pseudogenes, transcribed examples, examples having a gene function (whether it is called a pseudogene is not determined) and the like also have been known, so that the term "pseudogene" in the present method means the "processed pseudogene" or "duplicated pseudogene (non-processed pseudogene)" rather than presence or absence of gene function or whether it is transcribed or not.

20 **[0142]** When DNA having a target DNA region is a repetitive sequence in genome, the repetitive sequence is a group of nucleotide sequences having homology, so that there is the possibility that in complementary base pairing between DNA having a target DNA region and an adhesion sequence for detection, every nucleotide sequence fails to base-pair with DNA having a target DNA region. Concretely, as the LINE sequence, there are about 280 copies of nucleotide sequences having a homology of 80% or more with respect to SEQ ID NO: 37 in genome, and as the SINE (Alu) sequence, there are found about 820 copies of nucleotide sequences having a homology of 80% or more with respect to SEQ ID NO: 39 in genome. However, these nucleotide sequences having homology found in genome include those having one or several different bases with respect to SEQ ID NO: 37 and SEQ ID NO: 39.

25 **[0143]** In First step of the present method, it is preferred to extract DNA from a specimen by a system containing a sodium salt at high concentration. Concretely, as a concentration of sodium salt in a solution (for example, buffer) used in a DNA extraction operation for obtaining DNA from a specimen in First step of the present method, at least 50 mM or more, and preferably 100 mM or more can be recited. More concretely, 50 mM or more and 1000 mM or less, preferably 100 mM or more and 1000 mM or less, more preferably 100 mM or more and 200 mM or less can be recited. Any salts including NaCl, NaCO₃, Na₂SO₄ and the like are applied as far as it is a salt containing a sodium ion, and preferably means NaCl.

30 **[0144]** The present invention is a method of selecting a specimen derived from a cancer patient, and includes the steps of evaluating a specimen derived from a test subject as a specimen derived from a cancer patient when there is a significant difference between a DNA quantification result or detection result quantified or detected using a specimen derived from a test subject by the method according to any one of Inventions 1 to 13, and a DNA quantification result or detection result quantified or detected using a specimen derived from a healthy subject by the method, and identifying the specimen derived from a cancer patient based on the evaluation result. As a preferred aspect of the present invention, the invention in which the specimen is a serum derived from a mammal, and the invention in which the DNA comprising a target DNA region is free DNA comprising a target DNA region in serum derived from a mammal can be recited. Use of these inventions will make it possible to identify a cancer patient in a simple and convenient manner by a blood test.

35 **[0145]** Here, the "cancer patient" is a test subject developing a cancer, and as the cancer, solid cancers developing in organs of human and mammals, and non-solid cancers developing in blood of human and mammals such as lung cancer (non-small-cell lung cancer, small-cell lung cancer), esophageal cancer, gastric cancer, duodenal cancer, colon cancer, rectal cancer, hepatic cancer (hepatocarcinoma, cholangiocellular carcinoma), gallbladder cancer, bile duct cancer, pancreatic cancer, colon cancer, anal cancer, breast cancer, cervical cancer, uterine cancer, ovarian cancer, vulvar cancer, vaginal cancer, prostate cancer, kidney cancer, ureter cancer, bladder cancer, prostate cancer, penile cancer, testicular (testis) cancer, maxillary cancer, tongue cancer, (nasal-, oro-, hypo-) pharyngeal cancer, acute myeloid leukemia, chronic myelogenous leukemia, acute lymphoblastic leukemia, chronic lymphoblastic leukemia, malignant lymphoma, myelodysplastic syndrome, thyroid cancer, brain tumor, osteosarcoma and skin cancer (basal cell cancer, squamous cell cancer) are included.

EXAMPLES

[0146] In the following, the present invention will be described in detail by way of examples, however, the present invention is not limited to these examples.

Example 1

[0147] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 μg/μL solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4).

[0148] A 0.5 μg/mL solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0149] For genomic DNA derived from human blood purchased from Clontech, a DNA fragment (X, SEQ ID NO: 19, the region corresponding to the nucleotide numbers 25687390-25687775 shown in Genbank Accession No. NT_029419) used as a test sample was amplified by conducting PCR using the following oligonucleotide primers (PF1 and PR1) designed for PCR of SEQ ID NO: 17 and SEQ ID NO: 18 and the following reaction condition.

<Oligonucleotide primer designed for PCR>

[0150]

PF1:5'- CTCAGCACCCAGGCGGCC -3' (SEQ ID NO: 17)

PR1:5'- CTGGCCAAACTGGAGATCGC -3' (SEQ ID NO: 18)

<DNA fragment>

[0151]

X: 5' -

CTCAGCACCCAGGCGGCCGCGATCATGAGGCGCGAGCGGCGCGCGGGCTGTTGCAGAGTCTT

GAGCGGGTGGCACACCGCGATGTAGCGGTGGCTGTCATGACTACCAGCATGTAGGCCGACG

CAAACATGCCGAACACCTGCAGGTGCTTCACCACGCGGCACAGCCAGTCGGGGCCGCGGAAG

CGGTAGGTGATGTCCCAGCACATTTGCGGCAGCACCTGGAAGAATGCCACGGCCAGGTCCGGC

CAGGCTGAGGTGTCGGATGAAGAGGTGCATGCGGGACGTCTTGCGCGGCGTCCGGTGCAGAG

CCAGCAGTACGCTGCTGTTGCCAGCACGGCCACCGCGAAAGTCACCGCCAGCACGGCGATC

TCCAGTTTGGCCAG -3' (SEQ ID NO: 19)

[0152] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μL of 5 μM of the above primer solutions, 5 μL of each 2 mM dNTP, and 5 μL of 10xbuffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μL of 5U/μL thermostable DNA polymerase (AmpliTaQ Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 30 seconds at 95°C, 30 seconds at 61°C and 45

EP 2 305 807 A1

seconds at 72°C.

[0153] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment X was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0154] For the DNA fragment X, the following solutions were prepared respectively in duplicate.

- 5
10
- Solution A: DNA fragment X 10ng/10 μ L TE buffer solution
 - Solution B: DNA fragment X 1ng/10 μ L TE buffer solution
 - Solution C: DNA fragment X 0.1ng/10 μ L TE buffer solution
 - Solution D: TE buffer solution (negative control solution)

15 **[0155]** Ten (10) μ L of each obtained solution, 0.5 μ L of SssI methylase (available from NEB Inc.), 5 μ L of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μ L of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

20 **[0156]** Synthesized was 5'-end FITC-labeled oligonucleotide F1 having the nucleotide sequence of SEQ ID NO: 21 capable of binding by complementation with oligonucleotide X' comprising a target DNA region of SEQ ID NO: 20, and a 0.02 μ M solution in Tris-HCl buffer (10 mM) was prepared.

<Oligonucleotide comprising target DNA region>

25 **[0157]**

X' : 5' -

30
35
40

```
CTCAGCACCCAGGCGGCCGCGATCATGAGGCGCGAGCGGCGCGGGCTGTTGCAGAGTCTT
GAGCGGGTGGCACACCGCGATGTAGCGGTTCGGCTGTCATGACTACCAGCATGTAGGCCGACG
CAAACATGCCGAACACCTGCAGGTGCTTCACCACGCGGCACAGCCAGTCGGGGCCGCGGAAG
CGGTAGGTGATGTCCCAGCACATTTGCGGCAGCACCTGGAAGAATGCCACGGCCAGGTCCGGC
CAGGCTGAGGTGTCGGATGAAGAGGTGCATGCGGGACGTCTTGCGCGGGCGTCCGGTGCAGAG
CCAGCAGTACGCTGCTGTTGCCAGCACGGCCACCGCGAAAGTCACCGCCAGCACGGCGATC
```

45 TCCAGTTTGGCCAG -3' (SEQ ID NO: 20)

<5'-end FITC-labeled oligonucleotide>

50 **[0158]**

F1: 5'- CTGGCCAAACTGGAGAT -3' (SEQ ID NO: 21)

[0159] Each obtained reaction liquid was subjected to the following treatments.

55 **[0160]** In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KoAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0161] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immo-

bilized, 100 μL of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of

5 the present method).
[0162] Then 100 μL of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 $\mu\text{g}/100 \mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.
 10 **[0163]** One hundred (100) μL of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 15 minutes, 50 μL of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

15 **[0164]** The result is shown in Fig. 1. In Solution A, Solution B, and Solution C, increase in absorbance was observed compared to Solution D. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

20 Example 2

[0165] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4).

25 **[0166]** A 0.5 $\mu\text{g}/\text{mL}$ solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

30 **[0167]** For genomic DNA derived from human blood purchased from Clontech, a DNA fragment (X, SEQ ID NO: 19, the region corresponding to the nucleotide numbers 25687390-25687775 shown in Genbank Accession No. NT_029419) used as a test sample was amplified by conducting PCR using the following oligonucleotide primers (PF1 and PR1) designed for PCR of SEQ ID NO: 17 and SEQ ID NO: 18 and the following reaction condition.

35 <Oligonucleotide primer designed for PCR>

40 **[0168]**

PF1:5'- CTCAGCACCCAGGCGGCC -3' (SEQ ID NO: 17)
 PR1:5'- CTGCCAAACTGGAGATCGC -3' (SEQ ID NO: 18)

<DNA fragment>

45 **[0169]**

X: 5' -

5 CTCAGCACCCAGGCGGCCGCGATCATGAGGCGCGAGCGGGCGCGGGGCTGTTGCAGAGTCTT
 GAGCGGGTGGCACACCGCGATGTAGCGGTTCGGCTGTCATGACTACCAGCATGTAGGCCGACG
 10 CAAACATGCCGAACACCTGCAGGTGCTTCACCACGCGGCACAGCCAGTCGGGGCCGCGGAAG
 CGGTAGGTGATGTCCCAGCACATTTGCGGCAGCACCTGGAAGAATGCCACGGCCAGGTTCGGC
 CAGGCTGAGGTGTCGGATGAAGAGGTGCATGCGGGACGTCTTGCGCGGCGTCCGGTGCAGAG
 15 CCAGCAGTACGCTGCTGTTGCCCAGCACGGCCACCGCGAAAGTCACCGCCAGCACGGCGATC
 TCCAGTTTGGCCAG -3' (SEQ ID NO: 19)

20 **[0170]** As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10xbuffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 30 seconds at 95°C, 30 seconds at 61°C and 45 seconds at 72°C.

25 **[0171]** After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment X was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0172] For the DNA fragment X, the following solutions were prepared respectively in duplicate.

30 Solution A: DNA fragment X 10ng/10 μ L TE buffer solution
 Solution B: DNA fragment X 1ng/10 μ L TE buffer solution
 Solution C: DNA fragment X 0.1ng/10 μ L TE buffer solution
 Solution D: TE buffer solution (negative control solution)

35 **[0173]** For genomic DNA derived from human blood purchased from Clontech, a DNA fragment (Y, SEQ ID NO: 24, the region corresponding to the nucleotide numbers 76606-76726 shown in Genbank Accession No. AC009800) used as a test sample was amplified by conducting PCR using the following oligonucleotide primers (PF2 and PR2) designed for PCR of SEQ ID NO: 22 and SEQ ID NO: 23 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

40 **[0174]**

PF2:5'- TGAGCTCCGTAGGGCGTCC -3' (SEQ ID NO: 22)
 PR2:5'- GCGCCGGGTCCGGGCC -3' (SEQ ID NO: 23)

45 <DNA fragment>

[0175]

50 Y: 5' -

GCGCCGGGTCCGGGCCCGATGCGTTGGCGGGCCAGGGCTCCGAGAACGAGGCGTTGTCCATC
 55 TCAACGAGGGCAGAGGAGCCGGCGACCTGGCGTCCCCAAGGACGCCCTACGGAGCTCA -3'
 (SEQ ID NO: 24)

EP 2 305 807 A1

[0176] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μL of 5 μM of the above primer solutions, 5 μL of each 2 mM dNTP, and 5 μL of 10xbuffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl_2 , 0.01% Gelatin) were mixed with 0.25 μL of 5U/ μL thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL . After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 50 cycles each consisting of 30 seconds at 95°C, 30 seconds at 60°C and 45 seconds at 72°C.

[0177] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment Y was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0178] For the DNA fragment Y, the following solutions were prepared respectively.

Solution A: DNA fragment Y 10ng/10 μL TE buffer solution
Solution: DNA fragment Y 1ng/10 μL TE buffer solution
Solution C: DNA fragment Y 0.1ng/10 μL TE buffer solution
Solution D: TE buffer solution (negative control solution)

[0179] Solution A of DNA fragment X and Solution A of DNA fragment Y, Solution B of DNA fragment X and Solution B of DNA fragment Y, Solution C of DNA fragment X and Solution C of DNA fragment Y, and Solution D of DNA fragment X and Solution D of DNA fragment Y prepared in the above were respectively mixed, to prepare the following DNA fragment-mixed Solutions MA to MD respectively in duplicate.

Solution MA: 10 ng/20 μL TE buffer solution
Solution MB: 1 ng/20 μL TE buffer solution
Solution MC: 0.1 ng/20 μL TE buffer solution
Solution MD: TE buffer solution (negative control solution)

[0180] Twenty (20) μL of each obtained solution, 0.5 μL of Sssl methylase (available from NEB Inc.), 5 μL of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μL of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL . The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

[0181] Synthesized was 5'-end FITC-labeled oligonucleotide F1 having the nucleotide sequence of SEQ ID NO: 21 capable of binding by complementation with oligonucleotide X' comprising a target DNA region of SEQ ID NO: 20, and a 0.02 μM solution in Tris-HCl buffer (10 mM) was prepared.

<Oligonucleotide comprising target DNA region>

[0182]

X' : 5' -

CTCAGCACCCAGGCGGCCGCGATCATGAGGCGCGAGCGGCCGCGGGCTGTTGCAGAGTCTT
GAGCGGGTGGCACACCGCGATGTAGCGGTTCGGCTGTCATGACTACCAGCATGTAGGCCGACG
CAAACATGCCGAACACCTGCAGGTGCTTCACCACGCGGCACAGCCAGTCGGGGCCGCGGAAG
CGGTAGGTGATGTCCCAGCACATTTGCGGCAGCACCTGGAAGAATGCCACGGCCAGGTCGGC
CAGGCTGAGGTGTCGGATGAAGAGGTGCATGCGGGACGTCTTGCGCGGCGTCCGGTGCAGAG
CCAGCAGTACGCTGCTGTTGCCAGCACGGCCACCGCGAAAGTCACCGCCAGCACGGCGATC
TCCAGTTTGGCCAG -3' (SEQ ID NO: 20)

<5'-end FITC-labeled oligonucleotide>

[0183]

EP 2 305 807 A1

F1:5'- CTGGCCAAACTGGAGAT -3' (SEQ ID NO: 21)

[0184] Each obtained reaction liquid was subjected to the following treatments.

[0185] In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc2, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0186] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μ L of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

[0187] Then 100 μ L of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 μ g/100 μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0188] One hundred (100) μ L of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 30 minutes, 50 μ L of a stop solution (2N H₂ SO₄ aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0189] The result is shown in Fig. 2. In Solution MA, Solution MB, and Solution MC, increase in absorbance was observed compared to Solution MD. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 3

[0190] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 μ g/ μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4).

[0191] A 0.5 μ g/mL solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH 7.4) was prepared, and each 100 μ L of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0192] For genomic DNA derived from human blood purchased from Clontech, a DNA fragment (X, SEQ ID NO: 19, the region corresponding to the nucleotide numbers 25687390-25687775 shown in Genbank Accession No. NT_029419) used as a test sample was amplified by conducting PCR using the following oligonucleotide primers (PF1 and PR1) designed for PCR of SEQ ID NO: 17 and SEQ ID NO: 18 and the following reaction condition.

<Oligonucleotide primer designed for PCR>

[0193]

PF1:5'- CTCAGCACCCAGGCGGCC -3' (SEQ ID NO: 17)

PR1:5'- CTGGCCAAACTGGAGATCGC -3' (SEQ ID NO: 18)

EP 2 305 807 A1

<DNA fragment>

[0194]

5

X: 5' -

CTCAGCACCCAGGCGGCCGCGATCATGAGGCGCGAGCGGCGCGGGCTGTTGCAGAGTCTT

10

GAGCGGGTGGCACACCGCGATGTAGCGGTTCGGCTGTCATGACTACCAGCATGTAGGCCGACG

CAAACATGCCGAACACCTGCAGGTGCTTCACCACGCGGCACAGCCAGTCGGGGCCGCGGAAG

15

CGGTAGGTGATGTCCCAGCACATTTGCGGCAGCACCTGGAAGAATGCCACGGCCAGGTCCGGC

CAGGCTGAGGTGTCGGATGAAGAGGTGCATGCGGGACGTCTTGCGCGGGCGTCCGGTGCAGAG

CCAGCAGTACGCTGCTGTTGCCAGCACGGCCACCGCGAAAGTCACCGCCAGCACGGCGATC

20

TCCAGTTTGGCCAG -3' (SEQ ID NO: 19)

[0195] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10xbuffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 30 seconds at 95°C, 30 seconds at 61°C and 45 seconds at 72°C.

[0196] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment X was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0197] For the DNA fragment X, the following solutions were prepared respectively in duplicate.

Solution A: DNA fragment X 10ng/10 μ L TE buffer solution

Solution B: DNA fragment X 1ng/10 μ L TE buffer solution

35

Solution C: DNA fragment X 0.1ng/10 μ L TE buffer solution

Solution D: TE buffer solution (negative control solution)

[0198] For genomic DNA derived from human blood purchased from Clontech, a DNA fragment (Y, SEQ ID NO: 24, the region corresponding to the nucleotide numbers 76606-76726 shown in Genbank Accession No. AC009800) used as a test sample was amplified by conducting PCR using the following oligonucleotide primers (PF2 and PR2) designed for PCR of SEQ ID NO: 22 and SEQ ID NO: 23 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

45

[0199]

PF2:5'- TGAGCTCCGTAGGGCGTCC -3' (SEQ ID NO: 22)

PR2:5'- GCGCCGGTCCGGGCC -3' (SEQ ID NO: 23)

50

<DNA fragment>

[0200]

55

Y : 5' -

GCGCCGGGTCCGGGCCCCGATGCGTTGGCGGGCCAGGGCTCCGAGAACGAGGCGTTGTCCATC
TCAACGAGGGCAGAGGAGCCGGCGACCTGGCGTCCCCAAGGACGCCCTACGGAGCTCA - 3'

(SEQ ID NO: 24)

[0201] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10xbuffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 50 cycles each consisting of 30 seconds at 95°C, 30 seconds at 60°C and 45 seconds at 72°C.

[0202] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment Y was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0203] For the DNA fragment Y, the following solutions were prepared respectively.

Solution A: DNA fragment Y 10ng/10 μ L TE buffer solution
Solution B: DNA fragment Y 1ng/10 μ L TE buffer solution
Solution C: DNA fragment Y 0.1ng/10 μ L TE buffer solution
Solution D: TE buffer solution (negative control solution)

[0204] Solution A of DNA fragment X and Solution A of DNA fragment Y, Solution B of DNA fragment X and Solution B of DNA fragment Y, Solution C of DNA fragment X and Solution C of DNA fragment Y, and Solution D of DNA fragment X and Solution D of DNA fragment Y prepared in the above were respectively mixed, to prepare the following DNA fragment-mixed Solutions MA to MD respectively in duplicate.

Solution MA: 10 ng/20 μ L TE buffer solution
Solution MB: 1 ng/20 μ L TE buffer solution
SolutionMC: 0.1 ng/20 μ L TE buffer solution
Solution MD: TE buffer solution (negative control solution)

[0205] Twenty (20) μ L of each obtained solution, 0.5 μ L of Sssl methylase (available from NEB Inc.), 5 μ L of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μ L of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

[0206] Synthesized was 5'-end FITC-labeled oligonucleotide F2 having the nucleotide sequence of SEQ ID NO: 26 capable of binding by complementation with oligonucleotide X' comprising a target DNA region of SEQ ID NO: 25, and a 0.02 μ M solution in Tris-HCl buffer (10 mM) was prepared.

<Oligonucleotide comprising target DNA region>

[0207]

Y' : 5' -

GCGCCGGGTCCGGGCCCCGATGCGTTGGCGGGCCAGGGCTCCGAGAACGAGGCGTTGTCCATC
TCAACGAGGGCAGAGGAGCCGGCGACCTGGCGTCCCCAAGGACGCCCTACGGAGCTCA - 3'

(SEQ ID NO: 25)

<5'-end FITC-labeled oligonucleotide>

[0208]

5 F2:5'- GACAACGCCTCGTTCTCGG -3' (SEQ ID NO: 26)

[0209] Each obtained reaction liquid was subjected to the following treatments.

[0210] In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0211] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μ L of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

[0212] Then 100 μ L of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 μ g/100 μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0213] One hundred (100) μ L of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 30 minutes, 50 μ L of a stop solution (2N H₂ SO₄ aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0214] The result is shown in Fig. 3. In Solution MA and Solution MB, increase in absorbance was observed compared to Solution MD. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 4

[0215] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 μ g/ μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4).

[0216] A 0.5 μ g/mL solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH 7. 4) was prepared, and each 100 μ L of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0217] For genomic DNA derived from human blood purchased from Clontech, a DNA fragment (X, SEQ ID NO: 19, the region corresponding to the nucleotide numbers 25687390-25687775 shown in Genbank Accession No. NT_029419) used as a test sample was amplified by conducting PCR using the following oligonucleotide primers (PF1 and PR1) designed for PCR of SEQ ID NO: 17 and SEQ ID NO: 18 and the following reaction condition.

<Oligonucleotide primer designed for PCR>

[0218]

EP 2 305 807 A1

PF1:5'- CTCAGCACCCAGGCGGCC -3' (SEQ ID NO: 17)
PR1:5'- CTGGCCAAACTGGAGATCGC -3' (SEQ ID NO: 18)

<DNA fragment>

5

[0219]

X: 5' -

10

CTCAGCACCCAGGCGGCCGCGATCATGAGGCGCGAGCGGCGCGGGCTGTTGCAGAGTCTT

GAGCGGGTGGCACACCGCGATGTAGCGGTCGGCTGTCATGACTACCAGCATGTAGGCCGACG

15

CAAACATGCCGAACACCTGCAGGTGCTTACCACGCGGCACAGCCAGTCGGGGCCGCGGAAG

CGGTAGGTGATGTCCCAGCACATTTGCGGCAGCACCTGGAAGAATGCCACGGCCAGGTCGGC

20

CAGGCTGAGGTGTTCGGATGAAGAGGTGCATGCGGGACGTCCTTTCGCGGGCGTCCGGTGCAGAG

CCAGCAGTACGCTGCTGTTGCCAGCACGGCCACCGCGAAAGTCACCGCCAGCACGGCGATC

TCCAGTTTGGCCAG -3' (SEQ ID NO: 19)

25

[0220] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10xbuffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 30 seconds at 95°C, 30 seconds at 61°C and 45 seconds at 72°C.

30

[0221] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment X was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0222] For the DNA fragment X, the following solutions were prepared respectively in duplicate.

35

Solution A: DNA fragment X 10ng/10 μ L TE buffer solution
Solution B: DNA fragment X 1ng/10 μ L TE buffer solution
Solution C: DNA fragment X 0.1ng/10 μ L TE buffer solution
Solution D: TE buffer solution (negative control solution)

40

[0223] For genomic DNA derived from human blood purchased from Clontech, a DNA fragment (Y, SEQ ID NO: 24, the region corresponding to the nucleotide numbers 76606-76726 shown in Genbank Accession No. AC009800) used as a test sample was amplified by conducting PCR using the following oligonucleotide primers (PF2 and PR2) designed for PCR of SEQ ID NO: 22 and SEQ ID NO: 23 and the following reaction condition.

45

<Oligonucleotide primers designed for PCR>

[0224]

50

PF2:5'- TGAGCTCCGTAGGGCGTCC -3' (SEQ ID NO: 22)
PR2:5'- GCGCCGGTCCGGGCC -3' (SEQ ID NO: 23)

<DNA fragment>

55

[0225]

Y: 5' -

GCGCCGGGTCCGGGCCCCGATGCGTTGGCGGGCCAGGGCTCCGAGAACGAGGCGTTGTCCATC
 TCAACGAGGGCAGAGGAGCCGGCGACCTGGCGTCCCCCAAGGACGCCCTACGGAGCTCA - 3'

(SEQ ID NO: 24)

[0226] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10xbuffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 50 cycles each consisting of 30 seconds at 95°C, 30 seconds at 60°C and 45 seconds at 72°C.

[0227] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment Y was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0228] For the DNA fragment Y, the following solutions were prepared respectively.

- Solution A: DNA fragment Y 10ng/10 μ L TE buffer solution
- Solution B: DNA fragment Y 1ng/10 μ L TE buffer solution
- Solution C: DNA fragment Y 0.1ng/10 μ L TE buffer solution
- Solution D: TE buffer solution (negative control solution)

[0229] Solution A of DNA fragment X and Solution A of DNA fragment Y, Solution B of DNA fragment X and Solution B of DNA fragment Y, Solution C of DNA fragment X and Solution C of DNA fragment Y, and Solution D of DNA fragment X and Solution D of DNA fragment Y prepared in the above were respectively mixed, to prepare the following DNA fragment-mixed Solutions MA to MD respectively in duplicate.

- Solution MA: 10 ng/20 μ L TE buffer solution
- Solution MB: 1 ng/20 μ L TE buffer solution
- Solution MC: 0.1 ng/20 μ L TE buffer solution
- Solution MD: TE buffer solution (negative control solution)

[0230] Ten (10) μ L of each obtained solution, 0.5 μ L of SssI methylase (available from NEB Inc.), 5 μ L of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μ L of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

[0231] Synthesized were 5'-end FITC-labeled oligonucleotide F1 having the nucleotide sequence of SEQ ID NO: 21 capable of binding by complementation with oligonucleotide X' comprising a target DNA region of SEQ ID NO: 20 and 5'-end FITC-labeled oligonucleotide F2 having the nucleotide sequence of SEQ ID NO: 26 capable of binding by complementation with oligonucleotide X' comprising a target DNA region of SEQ ID NO: 25, and respective 0.02 μ M solutions in Tris-HCl buffer (10 mM) were prepared.

<Oligonucleotide comprising target DNA region>

[0232]

X' : 5' -

5 CTCAGCACCCAGGCGGCCGCGATCATGAGGCGCGAGCGGCCGCGGGCTGTTGCAGAGTCTT
 GAGCGGGTGGCACACCGCGATGTAGCGGTTCGGCTGTCATGACTACCAGCATGTAGGCCGACG
 CAAACATGCCGAACACCTGCAGGTGCTTCACCACGCGGCACAGCCAGTCGGGGCCGCGGAAG
 10 CGGTAGGTGATGTCCCAGCACATTTGCGGCAGCACCTGGAAGAATGCCACGGCCAGGTTCGGC
 CAGGCTGAGGTGTCGGATGAAGAGGTGCATGCGGGACGTCTTGCGCGGCGTCCGGTGCAGAG
 CCAGCAGTACGCTGCTGTTGCCAGCACGGCCACCGCGAAAGTCACCGCCAGCACGGCGATC
 15 TCCAGTTTGGCCAG -3' (SEQ ID NO: 20)

Y' : 5' -

20 GCGCCGGGTCCGGGCCCCGATGCGTTGGCGGGCCAGGGCTCCGAGAACGAGGCGTTGTCCATC
 TCAACGAGGGCAGAGGAGCCGGCGACCTGGCGTCCCCAAGGACGCCCTACGGAGCTCA -3'
 25 (SEQ ID NO: 25)

<5'-end FITC-labeled oligonucleotide>

30 **[0233]**

F1:5'- CTGGCCAAACTGGAGAT -3' (SEQ ID NO: 21)
 F2:5'- GACAACGCCTCGTTCTCGG -3' (SEQ ID NO: 26)

35 **[0234]** Each obtained reaction liquid was subjected to the following treatments.

[0235] In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0236] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μ L of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

[0237] Then 100 μ L of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 μ g/100 μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0238] One hundred (100) μ L of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 30 minutes, 50 μ L of a stop solution (2N H₂ SO₄ aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0239] The result is shown in Fig. 4. In Solution MA, Solution MB, and Solution MC, increase in absorbance was observed compared to Solution MD. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 5

[0240] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylation kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 µg/ µL solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4).

[0241] A 0.5 µg/mL solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH 7.4) was prepared, and each 100 µL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 µL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0242] Yeast strain X2180-1A of baker's yeast was cultured in a YPD medium (1% Yeast extract, 2% Peptone, 2% Glucose, pH 5.6 to 6.0) to a turbidity of OD₆₀₀ 0.6 to 1.0, and centrifuged at 10,000 g for 10 minutes, to prepare 1 × 10⁷ of yeast cells. From the prepared yeast cells, a yeast genome was acquired using a generally used preparation method of a yeast genome as described in Methods in Yeast Genetics (Cold Spring Harbor Laboratory).

[0243] The prepared yeast cells were suspended in Buffer A (1 M sorbitol, 0.1 M EDTA, pH 7.4), added with 2-mercaptoethanol (final concentration 14 mM) and 100 U zymolase (10 mg/ml), and incubated under stirring at 30°C for an hour until the solution became clear. After collecting a protoplast by centrifugation at 550 g for 10 minutes, it was suspended in Buffer B (50 mM Tris-HCl, pH 7.4, 20 mM EDTA), added with sodium dodecyl sulfate in 1% (w/v), and then incubated at 65°C for 30 minutes. Sequentially, 5 M CH₃COOK was added and mingled in a volume ratio of 2/5, and the mixture was cooled on ice for 30 minutes, and then centrifuged at 15,000 g for 30 minutes to collect the supernatant. The collected supernatant was added with 3 M CH₃COONa in a volume ratio of 1/10 and an equal amount of isopropanol and mingled well, and the precipitate obtained by centrifugation at 15,000 g at 4°C for 30 minutes was rinsed with 70% ethanol and collected. After drying, the precipitate was dissolved in 1 mL of TE buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA), and added with RNase A (available from Sigma) in a concentration of 40 µg/ml, incubated at 37°C for an hour, and then the mixture was added with proteinase K (available from Sigma) and sodium dodecyl sulfate in a concentrations of 500 µg/mL and 1% (w/v), respectively, and shaken at 55°C for about 16 hours. After end of the shaking, the mixture was extracted with phenol [saturated with 1 M Tris-HCl (pH 8.0)]-chloroform. An aqueous layer was collected, added with NaCl in a concentration of 0.5 N, and allowed to precipitate from ethanol, and the generated precipitate was collected. The collected precipitate was rinsed with 70% ethanol, to obtain genomic DNA.

[0244] From the obtained genomic DNA, a DNA fragment to be used as a test sample (S, SEQ ID NO: 29, the region corresponding to the nucleotide numbers 271743-272083 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF3 and PR3) designed for PCR of SEQ ID NO: 27 and SEQ ID NO: 28 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

[0245]

PF3: 5'-AGGTGAGCTACGTGTGTTTGG-3' (SEQ ID NO: 27)

PR3: 5'-AGACATGTGCTCACGTACGGT-3' (SEQ ID NO: 28)

<DNA fragment>

[0246]

EP 2 305 807 A1

S:

5' -AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGC
5 AGCTTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGT
GGCGGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCGATAGTCGAGCCGCCTGTCAC
10 GTGCGCGGCCGCCCTGCTCCGTTTGACGCGATGCATAGCATGCGACCACCCAGTAATCATA
TGCTGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTC
15 CACACCGTACGTGAGCACATGTCT-3' (SEQ ID NO: 29)

20 **[0247]** As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10xbuffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

25 **[0248]** After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0249] For the DNA fragment S, the following solutions were prepared respectively in duplicate.

30 Solution A: DNA fragment S 10ng/20 μ L TE buffer solution
Solution B: DNA fragment S 1ng/20 μ L TE buffer solution
Solution C: DNA fragment S 0.1ng/20 μ L TE buffer solution
Solution D: TE buffer solution (negative control solution)

35 **[0250]** Twenty (20) μ L of each obtained solution, 0.5 μ L of SssI methylase (available from NEB Inc.), 5 μ L of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μ L of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

40 **[0251]** Synthesized was 5'-end FITC-labeled oligonucleotide F3 having the nucleotide sequence of SEQ ID NO: 31 capable of binding by complementation with oligonucleotide S' comprising a target DNA region of SEQ ID NO: 30, and a 0.02 μ M solution in Tris-HCl buffer (10 mM) was prepared.

<Oligonucleotide comprising target DNA region>

45 **[0252]**

S' : 5' -

5 AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGCAGC
 TTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGTGGC
 GGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGGCAGTGTGCGATAGTCGAGCCGCCTGTCACGTG
 10 CGCGGCCGCCCTGCTCCGTTTGTGACGCGATGCATAGCATGCGACCACCCAGTAATCATACTGC
 TGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTCCAC
 ACCGTACGTGAGCACATGTCT -3' (SEQ ID NO: 30)

<5'-end FITC-labeled oligonucleotide>

[0253]

F3:5'- CTGGCCAAACTGGAGAT -3' (SEQ ID NO: 31)

[0254] Each obtained reaction liquid was subjected to the following treatments.

[0255] In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0256] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μ L of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

[0257] Then 100 μ L of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 μ g/100 μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0258] One hundred (100) μ L of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 30 minutes, 50 μ L of a stop solution (2N H₂SO₄ aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0259] The result is shown in Fig. 5. In Solution A, Solution B, and Solution C, increase in absorbance was observed compared to Solution D. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 6

[0260] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 μ g/ μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O,

154 mM NaCl pH7.4).

[0261] A 0.5 $\mu\text{g}/\text{mL}$ solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0262] Yeast strain X2180-1A of baker's yeast was cultured in a YPD medium (1% Yeast extract, 2% Peptone, 2% Glucose, pH 5.6 to 6.0) to a turbidity of OD_{600} 0.6 to 1.0, and centrifuged at 10,000 g for 10 minutes, to prepare 1×10^7 of yeast cells. From the prepared yeast cells, a yeast genome was acquired using a generally used preparation method of a yeast genome as described in Methods in Yeast Genetics (Cold Spring Harbor Laboratory).

[0263] The prepared yeast cells were suspended in Buffer A (1 M sorbitol, 0.1 M EDTA, pH 7.4), added with 2-mercaptoethanol (final concentration 14 mM) and 100 U zymolase (10 mg/ml), and incubated under stirring at 30°C for an hour until the solution became clear. After collecting a protoplast by centrifugation at 550 g for 10 minutes, it was suspended in Buffer B (50 mM Tris-HCl, pH 7.4, 20 mM EDTA), added with sodium dodecyl sulfate in 1% (w/v), and then incubated at 65°C for 30 minutes. Sequentially, 5 M CH_3COOK was added and mingled in a volume ratio of 2/5, and the mixture was cooled on ice for 30 minutes, and then centrifuged at 15,000 g for 30 minutes to collect the supernatant. The collected supernatant was added with 3 M CH_3COONa in a volume ratio of 1/10 and an equal amount of isopropanol and mingled well, and the precipitate obtained by centrifugation at 15,000 g at 4°C for 30 minutes was rinsed with 70% ethanol and collected. After drying, the precipitate was dissolved in 1 mL of TE buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA), and added with RNase A (available from Sigma) in a concentration of 40 $\mu\text{g}/\text{ml}$, incubated at 37°C for an hour, and then the mixture was added with proteinase K (available from Sigma) and sodium dodecyl sulfate in concentrations of 500 $\mu\text{g}/\text{mL}$ and 1% (w/v), respectively, and shaken at 55°C for about 16 hours. After end of the shaking, the mixture was extracted with phenol [saturated with 1 M Tris-HCl (pH 8.0)]-chloroform. An aqueous layer was collected, added with NaCl in a concentration of 0.5 N, and allowed to precipitate from ethanol, and the generated precipitate was collected. The collected precipitate was rinsed with 70% ethanol, to obtain genomic DNA.

[0264] From the obtained genomic DNA, a DNA fragment to be used as a test sample (S, SEQ ID NO: 29, the region corresponding to the nucleotide numbers 271743-272083 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF3 and PR3) designed for PCR of SEQ ID NO: 27 and SEQ ID NO: 28 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

[0265]

PF3: 5'-AGGTGAGCTACGTGTGTTTGG-3' (SEQ ID NO: 27)

PR3: 5'-AGACATGTGCTCACGTACGGT-3' (SEQ ID NO: 28)

<DNA fragment>

[0266]

S:

5' -AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGC

EP 2 305 807 A1

AGCTTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGT
GGCGGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCGATAGTCGAGCCGCCTGTCAC
GTGCGCGGCCGCCCTGCTCCGTTTGACGCGATGCATAGCATGCGACCACCCAGTAATCATA
TGCTGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTC
CACACCGTACGTGAGCACATGTCT-3' (SEQ ID NO: 29)

[0267] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10 \times buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

[0268] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0269] For the DNA fragment S, the following solutions were prepared respectively in duplicate.

Solution A: DNA fragment S 10ng/10 μ L TE buffer solution

Solution B: DNA fragment S 1ng/10 μ L TE buffer solution

Solution C: TE buffer solution (negative control solution)

[0270] From the obtained genomic DNA, a DNA fragment to be used as a test sample (T, SEQ ID NO: 34, the region corresponding to the nucleotide numbers 384569-384685 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF4 and PR4) designed for PCR of SEQ ID NO: 32 and SEQ ID NO: 33 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

[0271]

PF4: 5'-GGACCTGTGTTTGACGGGTAT-3' (SEQ ID NO: 32)

PR4: 5'-AGTACAGATCTGGCGTTCTCG-3' (SEQ ID NO: 33)

<DNA fragment>

[0272]

T:

5' -GGACCTGTGTTTGACGGGTATAACACTAAGTTGCGCAATTTGCTGTATTGCGAAATCCG
CCCGGACGATATCACTCTTGAGCGCATGTGCCGTTTCCGAGAACGCCAGATCTGTACT-3'

(SEQ ID NO: 34)

[0273] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10 \times buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

[0274] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment

EP 2 305 807 A1

S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0275] For the DNA fragment T, the following solutions were prepared respectively in duplicate.

5 Solution A: DNA fragment T 10ng/10 μ L TE buffer solution

Solution B: DNA fragment T 1ng/10 μ L TE buffer solution

Solution C: TE buffer solution (negative control solution)

[0276] Solution A of DNA fragment S and Solution A of DNA fragment T, Solution B of DNA fragment S and Solution B of DNA fragment T, and Solution C of DNA fragment S and Solution C of DNA fragment T prepared in the above were respectively mixed, to prepare the following DNA fragment-mixed Solutions MA to MC respectively in duplicate.

15 Solution MA: 10 ng/20 μ L TE buffer solution

Solution MB: 1 ng/20 μ L TE buffer solution

Solution MC: TE buffer solution (negative control solution)

[0277] Twenty (20) μ L of each obtained solution, 0.5 μ L of SssI methylase (available from NEB Inc.), 5 μ L of 10 \times NEBuffer2 (available from NEB Inc.), and 0.5 μ L of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

[0278] Synthesized was 5'-end FITC-labeled oligonucleotide F3 having the nucleotide sequence of SEQ ID NO: 31 capable of binding by complementation with oligonucleotide S' comprising a target DNA region of SEQ ID NO: 30, and a 0.02 μ M solution in Tris-HCl buffer (10 mM) was prepared.

<Oligonucleotide comprising target DNA region>

[0279]

30 S' : 5' -

AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGCAGC

35 TTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGTGGC

GGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCGATAGTCGAGCCGCCTGTACAGTG

CGCGGCCCGCCCTGCTCCGTTTGACGCGATGCATAGCATGCGACCACCCAGTAATCATACTGC

40 TGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTCCAC

ACCGTACGTGAGCACATGTCT -3' (SEQ ID NO: 30)

<5'-end FITC-labeled oligonucleotide>

[0280]

50 F3:5'- AGACATGTGCTCACGTACGGT -3' (SEQ ID NO: 31)

[0281] Each obtained reaction liquid was subjected to the following treatments.

[0282] In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room

temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0283] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μ L of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

[0284] Then 100 μ L of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 μ g/100 μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0285] One hundred (100) μ L of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 60 minutes, 50 μ L of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0286] The result is shown in Fig. 6. In Solution MA and Solution MB, increase in absorbance was observed compared to Solution MC. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 7

[0287] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 μ g/ μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4).

[0288] A 0.5 μ g/mL solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4) was prepared, and each 100 μ L of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0289] Yeast strain X2180-1A of baker's yeast was cultured in a YPD medium (1% Yeast extract, 2% Peptone, 2% Glucose, pH 5.6 to 6.0) to a turbidity of OD₆₀₀ 0.6 to 1.0, and centrifuged at 10,000 g for 10 minutes, to prepare 1×10^7 of yeast cells. From the prepared yeast cells, a yeast genome was acquired using a generally used preparation method of a yeast genome as described in Methods in Yeast Genetics (Cold Spring Harbor Laboratory).

[0290] The prepared yeast cells were suspended in Buffer A (1 M sorbitol, 0.1 M EDTA, pH 7.4), added with 2-mercaptoethanol (final concentration 14 mM) and 100 U zymolase (10 mg/ml), and incubated under stirring at 30°C for an hour until the solution became clear. After collecting a protoplast by centrifugation at 550 g for 10 minutes, it was suspended in Buffer B (50 mM Tris-HCl, pH 7.4, 20 mM EDTA), added with sodium dodecyl sulfate in 1% (w/v), and then incubated at 65°C for 30 minutes. Sequentially, 5 M CH_3COOK was added and mingled in a volume ratio of 2/5, and the mixture was cooled on ice for 30 minutes, and then centrifuged at 15,000 g for 30 minutes to collect the supernatant. The collected supernatant was added with 3 M CH_3COONa in a volume ratio of 1/10 and an equal amount of isopropanol and mingled well, and the precipitate obtained by centrifugation at 15,000 g at 4°C for 30 minutes was rinsed with 70% ethanol and collected. After drying, the precipitate was dissolved in 1 mL of TE buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA), and added with RNase A (available from Sigma) in a concentration of 40 μ g/ml, incubated at 37°C for an hour, and then the mixture was added with proteinase K (available from Sigma) and sodium dodecyl sulfate in concentrations of 500 μ g/mL and 1% (w/v), respectively, and shaken at 55°C for about 16 hours. After end of the shaking, the mixture was extracted with phenol [saturated with 1 M Tris-HCl (pH 8.0)]-chloroform. An aqueous layer was collected, added with NaCl in a concentration of 0.5 N, and allowed to precipitate from ethanol, and the generated precipitate was collected. The collected precipitate was rinsed with 70% ethanol, to obtain genomic DNA.

[0291] From the obtained genomic DNA, a DNA fragment to be used as a test sample (S, SEQ ID NO: 29, the region corresponding to the nucleotide numbers 271743-272083 of yeast chromosome VII shown in Genbank Accession No.

EP 2 305 807 A1

NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF3 and PR3) designed for PCR of SEQ ID NO: 27 and SEQ ID NO: 28 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

[0292]

PF3: 5'-AGGTGAGCTACGTGTGTTTGG-3' (SEQ ID NO: 27)

PR3: 5'-AGACATGTGCTCACGTACGGT-3' (SEQ ID NO: 28)

<DNA fragment>

[0293]

S:

5' -AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGC
AGCTTGTACGATTGGTGACCCGCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGT
GGCGGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCGATAGTCGAGCCGCTGTCAC
GTGCGCGGCCGCCCTGCTCCGTTTGACGCGATGCATAGCATGCGACCACCCAGTAATCATAC
TGCTGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTC
CACACCGTACGTGAGCACATGTCT-3' (SEQ ID NO: 29)

[0294] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10 \times buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

[0295] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0296] For the DNA fragment S, the following solutions were prepared respectively in duplicate.

Solution A: DNA fragment S 10ng/10 μ L TE buffer solution

Solution B: DNA fragment S 1ng/10 μ L TE buffer solution

Solution C: TE buffer solution (negative control solution)

[0297] From the obtained genomic DNA, a DNA fragment to be used as a test sample (T, SEQ ID NO: 34, the region corresponding to the nucleotide numbers 384569-384685 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF4 and PR4) designed for PCR of SEQ ID NO: 32 and SEQ ID NO: 33 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

[0298]

PF4: 5'-GGACCTGTGTTTGACGGGTAT-3' (SEQ ID NO: 32)

PR4: 5'-AGTACAGATCTGGCGTTCTCG-3' (SEQ ID NO: 33)

EP 2 305 807 A1

<DNA fragment>

[0299]

5 T:
5' -GGACCTGTGTTTGGACGGGTATAACACTAAGTTGCGCAATTTGCTGTATTGCGAAATCCG
10 CCCGGACGATATCACTCTTGAGCGCATGTGCCGTTTCCGAGAACGCCAGATCTGTACT-3'
(SEQ ID NO: 34)

15 [0300] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10 \times buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

20 [0301] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0302] For the DNA fragment T, the following solutions were prepared respectively in duplicate.

25 Solution A: DNA fragment T 10ng/10 μ L TE buffer solution
Solution B: DNA fragment T 1ng/10 μ L TE buffer solution
Solution C: TE buffer solution (negative control solution)

30 [0303] Solution A of DNA fragment S and Solution A of DNA fragment T, Solution B of DNA fragment S and Solution B of DNA fragment T, and Solution C of DNA fragment S and Solution C of DNA fragment T prepared in the above were respectively mixed, to prepare the following DNA fragment-mixed Solutions MA to MC respectively in duplicate.

Solution MA: 10 ng/20 μ L TE buffer solution
Solution MB: 1 ng/20 μ L TE buffer solution
35 Solution MC: TE buffer solution (negative control solution)

[0304] Twenty (20) μ L of each obtained solution, 0.5 μ L of SssI methylase (available from NEB Inc.), 5 μ L of 10 \times NEBuffer2 (available from NEB Inc.), and 0.5 μ L of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

40 [0305] Synthesized were 5'-end FITC-labeled oligonucleotide F3 having the nucleotide sequence of SEQ ID NO: 31 capable of binding by complementation with oligonucleotide S' comprising a target DNA region of SEQ ID NO: 30 and 5'-end FITC-labeled oligonucleotide F4 having the nucleotide sequence of SEQ ID NO: 36 capable of binding by complementation with oligonucleotide T' comprising a target DNA region of SEQ ID NO: 35, and respective 0.02 μ M solutions in Tris-HCl buffer (10 mM) were prepared.

<Oligonucleotide comprising target DNA region>

[0306]

50

55

S' : 5' -

5 AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGCAGC
 TTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGTGGC
 GGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCATAGTCGAGCCGCCTGTCACGTG
 10 CGCGGCCGCCCTGCTCCGTTTGACGCGATGCATAGCATGCGACCACCCAGTAATCATACTGC
 TGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTCCAC
 15 ACCGTACGTGAGCACATGTCT -3' (SEQ ID NO: 30)

T' : 5' -

20 GGACCTGTGTTTGACGGGTATAACACTAAGTTGCGCAATTTGCTGTATTGCGAAATCCGCCC
 25 GGACGATATCACTCTTGAGCGCATGTGCCGTTTCCGAGAACGCCAGATCTGTACT -3'
 (SEQ ID NO: 35)

30 <5'-end FITC-labeled oligonucleotide>

[0307]

35 F3:5'- AGACATGTGCTCACGTACGGT -3' (SEQ ID NO: 31)
 F4:5'- AGTACAGATCTGGCGTTCTCG -3' (SEQ ID NO: 36)

[0308] Each obtained reaction liquid was subjected to the following treatments.

40 **[0309]** In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

45 **[0310]** To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μ L of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

50 **[0311]** Then 100 μ L of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 μ g/100 μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

55 **[0312]** One hundred (100) μ L of a substrate (available from R&D, #DY999) was added to each well and mixed, to

start the reaction. After leaving still at room temperature for about 60 minutes, 50 μL of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0313] The result is shown in Fig. 7. In Solution MA and Solution MB, increase in absorbance was observed compared to Solution MC. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 8

[0314] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylation kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4).

[0315] A 0.5 $\mu\text{g}/\text{mL}$ solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0316] Using genomic DNA derived from human blood purchased from Clontech, the following solutions were prepared respectively in duplicate.

[0317] Solution A: Genomic DNA derived from human blood 100 ng/5 μL TE buffer solution

Solution B: Genomic DNA derived from human blood 10 ng/5 μL TE buffer solution

Solution C: Genomic DNA derived from human blood 1 ng/5 μL TE buffer solution

Solution D: TE buffer solution (negative control solution)

[0318] Five (5) μL of each obtained solution, 10 U of restriction enzyme XspI, and 2 μL of 10x buffer optimum for XspI (200 mM Tris-HCl pH8.5, 100 mM MgCl_2 , 10 mM Dithiothreitol, 1000 mM KCl) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 20 μL . The reaction liquid was incubated at 37°C for 1 hour.

[0319] Twenty (20) μL of each obtained solution, 0.5 μL of SssI methylase (available from NEB Inc.), 5 μL of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μL of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL . The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

[0320] Synthesized was 5'-end FITC-labeled oligonucleotide F5 having the nucleotide sequence of SEQ ID NO: 38 capable of binding by complementation with oligonucleotide Z (region corresponding to the nucleotide numbers 115-386 shown in Genbank Accession No. M80340) comprising a target DNA region of SEQ ID NO: 37 designed in LINE1 region known as human transposon, and a 0.02 μM solution in Tris-HCl buffer (10 mM) was prepared.

<Oligonucleotide comprising target DNA region>

[0321]

Z: 5' -

TAGGGAGTGCCAGACAGTGGGCGCAGGCCAGTGTGTGTGCGCACCGTGCGCGAGCCGAAGCA
 GGGCGAGGCATTGCCTCACCTGGGAAGCGCAAGGGGTGAGGGAGTTCCTTTCTGAGTCAAA
 GAAAGGGGTGACGGTCGCACCTGGAAAATCGGGTCACTCCCACCCGAATATTGCGCTTTTCA
 GACCGGCTTAAGAAACGGCGCACCCACGAGACTATATCCCACACCTGGCTCGGAGGGTCCTAC
 GCCCACGGAATCTCGCTGATTGC -3' (SEQ ID NO: 37)

<5'-end FITC-labeled oligonucleotide>

[0322]

F5:5'- CTGGCCAAACTGGAGAT -3' (SEQ ID NO: 38)

[0323] Each obtained reaction liquid was subjected to the following treatments.

[0324] In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0325] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μ L of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

[0326] Then 100 μ L of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 μ g/100 μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0327] One hundred (100) μ L of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 10 minutes, 50 μ L of a stop solution (2N H₂SO₄ aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0328] The result is shown in Fig. 8. In Solution A, Solution B, and Solution C, increase in absorbance was observed compared to Solution D. The intensity increased depending on the concentration of genomic DNA. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 9

[0329] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 μ g/ μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4).

[0330] A 0.5 μ g/mL solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-con-

EP 2 305 807 A1

taining phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0331] Using genomic DNA derived from human blood purchased from Clontech, the following solutions were prepared respectively in duplicate.

Solution A: Genomic DNA derived from human blood 100 ng/5 μL TE buffer solution
Solution B: Genomic DNA derived from human blood 10 ng/5 μL TE buffer solution
Solution C: Genomic DNA derived from human blood 1 ng/5 μL TE buffer solution
Solution D: TE buffer solution (negative control solution)

[0332] Five (5) μL of each obtained solution, 4 U of restriction enzyme MspI, and 2 μL of 10x buffer optimum for MspI (100 mM Tris-HCl pH 8.5, 100 mM MgCl_2 , 10 mM Dithiothreitol, 500 mM NaCl) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 20 μL . The reaction liquid was incubated at 37°C for 1 hour.

[0333] Twenty (20) μL of each obtained solution, 0.5 μL of SssI methylase (available from NEB Inc.), 5 μL of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μL of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL . The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

[0334] Synthesized was 5'-end FITC-labeled oligonucleotide F6 having the nucleotide sequence of SEQ ID NO: 40 capable of binding by complementation with oligonucleotide W (region corresponding to the nucleotide numbers 178-262 shown in Genbank Accession No. AF458110) comprising a target DNA region of SEQ ID NO: 39 designed in Alu region known as human transposon, and a 0.02 μM solution in Tris-HCl buffer (10 mM) was prepared.

<Oligonucleotide comprising target DNA region>

[0335]

W: 5' -

CGGGCGCGGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCCGAGGTGGGCGGATCACG

AGGTCAGGAGATCGAGACCATCC -3' (SEQ ID NO: 39)

<5'-end FITC-labeled oligonucleotide>

[0336]

F6:5'- GGATGGTCTCGATCTCCTGAC -3' (SEQ ID NO: 40)

[0337] Each obtained reaction liquid was subjected to the following treatments.

[0338] In a PCR tube, 40 μL of the reaction liquid prepared in the above, 10 μL of the 5'-end FITC-labeled oligonucleotide solution, 10 μL of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc_2 , 5 mM Dithiothreitol), 10 μL of a 100 mM MgCl_2 solution, 10 μL of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μL and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0339] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μL of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was

removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

[0340] Then 100 μL of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 $\mu\text{g}/100 \mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0341] One hundred (100) μL of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 8 minutes, 50 μL of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0342] The result is shown in Fig. 9. In Solution A, Solution B, and Solution C, increase in absorbance was observed compared to Solution D. The intensity increased depending on the concentration of genomic DNA. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 10

[0343] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylation kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4).

[0344] A 0.5 $\mu\text{g}/\text{mL}$ solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0345] Yeast strain X2180-1A of baker's yeast was cultured in a YPD medium (1% Yeast extract, 2% Peptone, 2% Glucose, pH 5.6 to 6.0) to a turbidity of OD_{600} 0.6 to 1.0, and centrifuged at 10,000 g for 10 minutes, to prepare 1×10^7 of yeast cells. From the prepared yeast cells, a yeast genome was acquired using a generally used preparation method of a yeast genome as described in Methods in Yeast Genetics (Cold Spring Harbor Laboratory).

[0346] The prepared yeast cells were suspended in Buffer A (1 M sorbitol, 0.1 M EDTA, pH 7.4), added with 2-mercaptoethanol (final concentration 14 mM) and 100 U zymolase (10 mg/ml), and incubated under stirring at 30°C for an hour until the solution became clear. After collecting a protoplast by centrifugation at 550 g for 10 minutes, it was suspended in Buffer B (50 mM Tris-HCl, pH 7.4, 20 mM EDTA), added with sodium dodecyl sulfate in 1% (w/v), and then incubated at 65°C for 30 minutes. Sequentially, 5 M CH_3COOK was added and mingled in a volume ratio of 2/5, and the mixture was cooled on ice for 30 minutes, and then centrifuged at 15,000 g for 30 minutes to collect the supernatant. The collected supernatant was added with 3 M CH_3COONa in a volume ratio of 1/10 and an equal amount of isopropanol and mingled well, and the precipitate obtained by centrifugation at 15,000 g at 4°C for 30 minutes was rinsed with 70% ethanol and collected. After drying, the precipitate was dissolved in 1 mL of TE buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA), and added with RNase A (available from Sigma) in a concentration of 40 $\mu\text{g}/\text{mL}$, incubated at 37°C for an hour, and then the mixture was added with proteinase K (available from Sigma) and sodium dodecyl sulfate in concentrations of 500 $\mu\text{g}/\text{mL}$ and 1% (w/v), respectively, and shaken at 55°C for about 16 hours. After end of the shaking, the mixture was extracted with phenol [saturated with 1 M Tris-HCl (pH 8.0)]-chloroform. An aqueous layer was collected, added with NaCl in a concentration of 0.5 N, and allowed to precipitate from ethanol, and the generated precipitate was collected. The collected precipitate was rinsed with 70% ethanol, to obtain genomic DNA.

[0347] From the obtained genomic DNA, a DNA fragment to be used as a test sample (S, SEQ ID NO: 29, the region corresponding to the nucleotide numbers 271743-272083 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF3 and PR3) designed for PCR of SEQ ID NO: 27 and SEQ ID NO: 28 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

[0348]

EP 2 305 807 A1

PF3: 5'-AGGTGAGCTACGTGTGTTTGG-3' (SEQ ID NO: 27)

PR3: 5'-AGACATGTGCTCACGTACGGT-3' (SEQ ID NO: 28)

<DNA fragment>

5

[0349]

S:

10

5' -AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGC

AGCTTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGT

15

GGCGGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCGATAGTCGAGCCGCCTGTCAC

GTGCGCGGCCGCCCTGCTCCGTTTGTGACGCGATGCATAGCATGCGACCACCCAGTAATCATAC

20

TGCTGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTC

CACACCGTACGTGAGCACATGTCT-3' (SEQ ID NO: 29)

25

[0350] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10 \times buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

30

[0351] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0352] For the DNA fragment S, the following solutions were prepared respectively in duplicate.

35

- Solution A: DNA fragment S 10ng/20 μ L TE buffer solution
- Solution B: DNA fragment S 1ng/20 μ L TE buffer solution
- Solution C: DNA fragment S 0.1ng/20 μ L TE buffer solution
- Solution D: TE buffer solution (negative control solution)

40

[0353] Twenty (20) μ L of each obtained solution, 0.5 μ L of SssI methylase (available from NEB Inc.), 5 μ L of 10 \times NEBuffer2 (available from NEB Inc.), and 0.5 μ L of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

45

[0354] Synthesized was 5'-end FITC-labeled oligonucleotide F3 having the nucleotide sequence of SEQ ID NO: 31 capable of binding by complementation with oligonucleotide S' comprising a target DNA region of SEQ ID NO: 30, and a 0.02 μ M solution in Tris-HCl buffer (10 mM) was prepared.

<Oligonucleotide comprising target DNA region>

[0355]

50

55

S' : 5' -

5 AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGCAGC
 TTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGTGGC
 10 GCGGGCTCAATGACCTGTGGCGCCCGT TTTGTGGCGTGCATAGTCGAGCCGCCTGTCACGTG
 CGCGGCCGCCCTGCTCCGTTTACGCGATGCATAGCATGCGACCACCCAGTAATCATACTGC
 TGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTCCAC
 15 ACCGTACGTGAGCACATGTCT -3' (SEQ ID NO: 30)

<5'-end FITC-labeled oligonucleotide>

[0356]

F3:5'- CTGGCCAAACTGGAGAT -3' (SEQ ID NO: 31)

20 **[0357]** Synthesized were counter oligonucleotides C1, C2, C3, C4, C5, C6, C7, C8, C9 and C10 having the nucleotide sequences of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48 and SEQ ID NO: 49 and SEQ ID NO: 50, respectively capable of binding by complementation with a minus strand of DNA fragment S' comprising the target DNA region of SEQ ID NO: 30, and respective 0.01 μM TE buffer solutions were prepared.

<Counter oligonucleotides>

[0358]

30 C1:5'- AGGTGAGCTACGTGTGTTTGG -3' (SEQ ID NO: 41)
 C2:5'- GCGTCGTGCACTGGCTCACTTGTACGCGCA -3' (SEQ ID NO: 42)
 35 C3:5'- CTTGTACGATTGGTGACCCGCCTTTTCGAC -3' (SEQ ID NO: 43)
 C4:5'- ACTGGACCGCTATGGACGTGGCGGCGGTGT -3' (SEQ ID NO: 44)
 C5:5'- GCGGGCGGCTCAATGACCTGTGGCGCCCGT -3' (SEQ ID NO: 45)
 C6:5'- TTGTGGCGTGCATAGTCGAGCCGCCTGTC -3' (SEQ ID NO: 46)
 C7:5'- ACGTGC GCGCCGCCCTGCTCCGTT -3' (SEQ ID NO: 47)
 40 C8:5'- TGACGCGATGCATAGCATGCGACCACCCAG -3' (SEQ ID NO: 48)
 C9:5'- ACTGCTGACGCTATTGGTCACGTGGTTATG -3' (SEQ ID NO: 49)
 C10:5'- CTGCTGTTGACTGCGGTGGCGTCCCGTTTC -3' (SEQ ID NO: 50)

[0359] Each obtained reaction liquid was subjected to the following treatments.

45 **[0360]** In a PCR tube, 40 μL of the reaction liquid prepared in the above, 10 μL of the 5'-end FITC-labeled oligonucleotide solution, 10 μL of the counter oligonucleotide solution, 10 μL of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μL of a 100 mM MgCl₂ solution, 10 μL of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μL and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

50 **[0361]** To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μL of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO₄ 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

[0362] Then 100 μL of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 $\mu\text{g}/100 \mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0363] One hundred (100) μL of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 30 minutes, 50 μL of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0364] The result is shown in Fig. 10. In Solution A, Solution B, and Solution C, increase in absorbance was observed compared to Solution D. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 11

[0365] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4).

[0366] A 0.5 $\mu\text{g}/\mu\text{L}$ solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0367] Yeast strain X2180-1A of baker's yeast was cultured in a YPD medium (1% Yeast extract, 2% Peptone, 2% Glucose, pH 5.6 to 6.0) to a turbidity of OD_{600} 0.6 to 1.0, and centrifuged at 10,000 g for 10 minutes, to prepare 1×10^7 of yeast cells. From the prepared yeast cells, a yeast genome was acquired using a generally used preparation method of a yeast genome as described in Methods in Yeast Genetics (Cold Spring Harbor Laboratory).

[0368] The prepared yeast cells were suspended in Buffer A (1 M sorbitol, 0.1 M EDTA, pH 7.4), added with 2-mercaptoethanol (final concentration 14 mM) and 100 U zymolase (10 mg/ml), and incubated under stirring at 30°C for an hour until the solution became clear. After collecting a protoplast by centrifugation at 550 g for 10 minutes, it was suspended in Buffer B (50 mM Tris-HCl, pH 7.4, 20 mM EDTA), added with sodium dodecyl sulfate in 1% (w/v), and then incubated at 65°C for 30 minutes. Sequentially, 5 M CH_3COOK was added and mingled in a volume ratio of 2/5, and the mixture was cooled on ice for 30 minutes, and then centrifuged at 15,000 g for 30 minutes to collect the supernatant. The collected supernatant was added with 3 M CH_3COONa in a volume ratio of 1/10 and an equal amount of isopropanol and mingled well, and the precipitate obtained by centrifugation at 15,000 g at 4°C for 30 minutes was rinsed with 70% ethanol and collected. After drying, the precipitate was dissolved in 1 mL of TE buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA), and added with RNase A (available from Sigma) in a concentration of 40 $\mu\text{g}/\text{ml}$, incubated at 37°C for an hour, and then the mixture was added with proteinase K (available from Sigma) and sodium dodecyl sulfate in concentrations of 500 $\mu\text{g}/\text{mL}$ and 1% (w/v), respectively, and shaken at 55°C for about 16 hours. After end of the shaking, the mixture was extracted with phenol [saturated with 1 M Tris-HCl (pH 8.0)]-chloroform. An aqueous layer was collected, added with NaCl in a concentration of 0.5 N, and allowed to precipitate from ethanol, and the generated precipitate was collected. The collected precipitate was rinsed with 70% ethanol, to obtain genomic DNA.

[0369] From the obtained genomic DNA, a DNA fragment to be used as a test sample (S, SEQ ID NO: 29, the region corresponding to the nucleotide numbers 271743-272083 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF3 and PR3) designed for PCR of SEQ ID NO: 27 and SEQ ID NO: 28 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

[0370]

PF3: 5'-AGGTGAGCTACGTGTGTTTGG-3' (SEQ ID NO: 27)

EP 2 305 807 A1

PR3: 5'-AGACATGTGCTCACGTACGGT-3' (SEQ ID NO: 28)

<DNA fragment>

5 **[0371]**

S:

10 5' -AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGC
AGCTTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGT
15 GGCGGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCGATAGTCGAGCCGCCTGTCAC
GTGCGCGGCCGCCCTGCTCCGTTTGACGCGATGCATAGCATGCGACCACCCAGTAATCATA
TGCTGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTC
20 CACACCGTACGTGAGCACATGTCT-3' (SEQ ID NO: 29)

25 **[0372]** As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10 \times buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

30 **[0373]** After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0374] For the DNA fragment S, the following solutions were prepared respectively in duplicate.

35 Solution A: DNA fragment S 10ng/10 μ L TE buffer solution
Solution B: DNA fragment S 1ng/10 μ L TE buffer solution
Solution C: TE buffer solution (negative control solution)

40 **[0375]** From the obtained genomic DNA, a DNA fragment to be used as a test sample (T, SEQ ID NO: 34, the region corresponding to the nucleotide numbers 384569-384685 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF4 and PR4) designed for PCR of SEQ ID NO: 32 and SEQ ID NO: 33 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

45 **[0376]**

PF4: 5'-GGACCTGTGTTTGACGGGTAT-3' (SEQ ID NO: 32)
PR4: 5'-AGTACAGATCTGGCGTTCTCG-3' (SEQ ID NO: 33)

<DNA fragment>

50 **[0377]**

55

T:

5' -GGACCTGTGTTTGGACGGGTATAACACTAAGTTGCGCAATTTGCTGTATTGCGAAATCCG
 5
 CCCGGACGATATCACTCTTGAGCGCATGTGCCGTTTCCGAGAACGCCAGATCTGTACT-3'

(SEQ ID NO: 34)

10
[0378] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μL of 5 μM of the above primer solutions, 5 μL of each 2 mM dNTP, and 5 μL of 10×buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μL of 5U/μL thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

15
[0379] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

20
[0380] For the DNA fragment T, the following solutions were prepared respectively in duplicate.

Solution A: DNA fragment T 10ng/10 μL TE buffer solution
 Solution B: DNA fragment T 1ng/10 μL TE buffer solution
 Solution C: TE buffer solution (negative control solution)

25
[0381] Solution A of DNA fragment S and Solution A of DNA fragment T, Solution B of DNA fragment S and Solution B of DNA fragment T, and Solution C of DNA fragment S and Solution C of DNA fragment T prepared in the above were respectively mixed, to prepare the following DNA fragment-mixed Solutions MA to MC respectively in duplicate.

30
 Solution MA: 10 ng/20 μL TE buffer solution
 Solution MB: 1 ng/20 μL TE buffer solution
 Solution MC: TE buffer solution (negative control solution)

35
[0382] Twenty (20) μL of each obtained solution, 0.5 μL of SssI methylase (available from NEB Inc.), 5 μL of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μL of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL. The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

40
[0383] Synthesized was 5'-end FITC-labeled oligonucleotide F3 having the nucleotide sequence of SEQ ID NO: 31 capable of binding by complementation with oligonucleotide S' comprising a target DNA region of SEQ ID NO: 30, and a 0.02 μM solution in Tris-HCl buffer (10 mM) was prepared.

<Oligonucleotide comprising target DNA region>

[0384]

45
 S' : 5' -

50
 AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGCAGC
 TTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGTGGC
 GGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCATAGTCGAGCCGCCTGTCACGTG
 CGCGGCCGCCCTGCTCCGTTTGGACGCGATGCATAGCATGCGACCACCCAGTAATCATACTGC
 55
 TGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTCCAC
 ACCGTACGTGAGCACATGTCT -3' (SEQ ID NO: 30)

<5'-end FITC-labeled oligonucleotide>

[0385]

5 F3:5'- AGACATGTGCTCACGTACGGT -3' (SEQ ID NO: 31)

[0386] Synthesized were counter oligonucleotides C1, C2, C3, C4, C5, C6, C7, C8, C9 and C10 having the nucleotide sequences of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48 and SEQ ID NO: 49 and SEQ ID NO: 50, respectively capable of binding by complementation with a minus strand of DNA fragment S' comprising the target DNA region of SEQ ID NO: 30, and respective 0.01 μ M TE buffer solutions were prepared.

<Counter oligonucleotides>

15 **[0387]**

C1:5'- AGGTGAGCTACGTGTGTTTGG -3' (SEQ ID NO: 41)
 C2:5'- GCGTCGTGCACTGGCTCACTTGTACGCGCA -3' (SEQ ID NO: 42)
 C3:5'- CTTGTACGATTGGTGACCCGCCTTTTCGAC -3' (SEQ ID NO: 43)
 20 C4:5'- ACTGGACCGCTATGGACGTGGCGGCGGTGT -3' (SEQ ID NO: 44)
 C5:5'- GGCGGCGGCTCAATGACCTGTGGCGCCCGT -3' (SEQ ID NO: 45)
 C6:5'- TTGTGGCGTGCGATAGTCGAGCCGCCTGTC -3' (SEQ ID NO: 46)
 C7:5'- ACGTGCGCGGCCGCCCTGCTCCGTT -3' (SEQ ID NO: 47)
 C8:5'- TGACGCGATGCATAGCATGCGACCACCCAG -3' (SEQ ID NO: 48)
 25 C9:5'- ACTGCTGACGCTATTGGTCACGTGGTTATG -3' (SEQ ID NO: 49)
 C10:5'- CTGCTGTTGACTGCGGTGGCGTCCCGTTTC -3' (SEQ ID NO: 50)

[0388] Each obtained reaction liquid was subjected to the following treatments.

30 **[0389]** In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of the counter oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

35 **[0390]** To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μ L of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

40 **[0391]** Then 100 μ L of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 μ g/100 μ L solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

45 **[0392]** One hundred (100) μ L of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 60 minutes, 50 μ L of a stop solution (2N H₂SO₄ aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

50 **[0393]** The result is shown in Fig. 11. In Solution MA and Solution MB, increase in absorbance was observed compared to Solution MC. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 12

[0394] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 μg/ μL solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4).

[0395] A 0.5 μg/mL solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0396] Yeast strain X2180-1A of baker's yeast was cultured in a YPD medium (1% Yeast extract, 2% Peptone, 2% Glucose, pH 5.6 to 6.0) to a turbidity of OD₆₀₀ 0.6 to 1.0, and centrifuged at 10,000 g for 10 minutes, to prepare 1 × 10⁷ of yeast cells. From the prepared yeast cells, a yeast genome was acquired using a generally used preparation method of a yeast genome as described in Methods in Yeast Genetics (Cold Spring Harbor Laboratory).

[0397] The prepared yeast cells were suspended in Buffer A (1 M sorbitol, 0.1 M EDTA, pH 7.4), added with 2-mercaptoethanol (final concentration 14 mM) and 100 U zymolase (10 mg/ml), and incubated under stirring at 30°C for an hour until the solution became clear. After collecting a protoplast by centrifugation at 550 g for 10 minutes, it was suspended in Buffer B (50 mM Tris-HCl, pH 7.4, 20 mM EDTA), added with sodium dodecyl sulfate in 1% (w/v), and then incubated at 65°C for 30 minutes. Sequentially, 5 M CH₃COOK was added and mingled in a volume ratio of 2/5, and the mixture was cooled on ice for 30 minutes, and then centrifuged at 15,000 g for 30 minutes to collect the supernatant. The collected supernatant was added with 3 M CH₃COONa in a volume ratio of 1/10 and an equal amount of isopropanol and mingled well, and the precipitate obtained by centrifugation at 15,000 g at 4°C for 30 minutes was rinsed with 70% ethanol and collected. After drying, the precipitate was dissolved in 1 mL of TE buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA), and added with RNase A (available from Sigma) in a concentration of 40 μg/ml, incubated at 37°C for an hour, and then the mixture was added with proteinase K (available from Sigma) and sodium dodecyl sulfate in concentrations of 500 μg/mL and 1% (w/v), respectively, and shaken at 55°C for about 16 hours. After end of the shaking, the mixture was extracted with phenol [saturated with 1 M Tris-HCl (pH 8.0)]-chloroform. An aqueous layer was collected, added with NaCl in a concentration of 0.5 N, and allowed to precipitate from ethanol, and the generated precipitate was collected. The collected precipitate was rinsed with 70% ethanol, to obtain genomic DNA.

[0398] From the obtained genomic DNA, a DNA fragment to be used as a test sample (S, SEQ ID NO: 29, the region corresponding to the nucleotide numbers 271743-272083 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF3 and PR3) designed for PCR of SEQ ID NO: 27 and SEQ ID NO: 28 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

[0399]

PF3: 5'-AGGTGAGCTACGTGTGTTTGG-3' (SEQ ID NO: 27)

PR3: 5'-AGACATGTGCTCACGTACGGT-3' (SEQ ID NO: 28)

<DNA fragment>

[0400]

S:

5' -AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGC

AGCTTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGT

GGCGGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCGATAGTCGAGCCGCCTGTCAC
 5 GTGCGCGGCCGCCCTGCTCCGTTTGACGCGATGCATAGCATGCGACCACCCAGTAATCATAC
 TGCTGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTC
 10 CACACCGTACGTGAGCACATGTCT-3' (SEQ ID NO: 29)

[0401] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10 \times buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

[0402] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0403] For the DNA fragment S, the following solutions were prepared respectively in duplicate.

20 Solution A: DNA fragment S 10ng/10 μ L TE buffer solution
 Solution B: DNA fragment S 1ng/10 μ L TE buffer solution
 Solution C: TE buffer solution (negative control solution)

25 [0404] From the obtained genomic DNA, a DNA fragment to be used as a test sample (T, SEQ ID NO: 34, the region corresponding to the nucleotide numbers 384569-384685 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF4 and PR4) designed for PCR of SEQ ID NO: 32 and SEQ ID NO: 33 and the following reaction condition.

30 <Oligonucleotide primers designed for PCR>

[0405]

35 PF4: 5'-GGACCTGTGTTTGACGGGTAT-3' (SEQ ID NO: 32)
 PR4: 5'-AGTACAGATCTGGCGTTCTCG-3' (SEQ ID NO: 33)

<DNA fragment>

[0406]

T:

40 5' -GGACCTGTGTTTGACGGGTATAACACTAAGTTGCGCAATTTGCTGTATTGCGAAATCCG
 45 CCCGGACGATATCACTCTTGAGCGCATGTGCCGTTTCCGAGAACGCCAGATCTGTACT-3'
 (SEQ ID NO: 34)

50 [0407] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10 \times buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

[0408] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0409] For the DNA fragment T, the following solutions were prepared respectively in duplicate.

EP 2 305 807 A1

Solution A: DNA fragment T 10ng/10 μ L TE buffer solution

Solution B: DNA fragment T 1ng/10 μ L TE buffer solution

Solution C: TE buffer solution (negative control solution)

5 **[0410]** Solution A of DNA fragment S and Solution A of DNA fragment T, Solution B of DNA fragment S and Solution B of DNA fragment T, and Solution C of DNA fragment S and Solution C of DNA fragment T prepared in the above were respectively mixed, to prepare the following DNA fragment-mixed Solutions MA to MC respectively in duplicate.

Solution MA: 10 ng/20 μ L TE buffer solution

10 Solution MB: 1 ng/20 μ L TE buffer solution

Solution MC: TE buffer solution (negative control solution)

15 **[0411]** Twenty (20) μ L of each obtained solution, 0.5 μ L of SssI methylase (available from NEB Inc.), 5 μ L of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μ L of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

20 **[0412]** Synthesized was 5'-end FITC-labeled oligonucleotide F4 having the nucleotide sequence of SEQ ID NO: 36 capable of binding by complementation with oligonucleotide T' comprising a target DNA region of SEQ ID NO: 35, and a 0.02 μ M solution in Tris-HCl buffer (10 mM) was prepared.

20

<Oligonucleotide comprising target DNA region>

[0413]

25

T' : 5' -

GGACCTGTGTTTGACGGGTATAACACTAAGTTGCGCAATTTGCTGTATTGCGAAATCCGCC

30

GGACGATATCACTCTTGAGCGCATGTGCCGTTTCCGAGAACGCCAGATCTGTACT - 3'

35

(SEQ ID NO: 35)

<5'-end FITC-labeled oligonucleotide>

[0414]

40

F4:5'- AGTACAGATCTGGCGTTCTCG -3' (SEQ ID NO: 36)

45 **[0415]** Synthesized were counter oligonucleotides C11, C12, C13 and C14 having the nucleotide sequences of SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53 and SEQ ID NO: 54, respectively capable of binding by complementation with a minus strand of DNA fragment T' comprising the target DNA region of SEQ ID NO: 35, and respective 0.01 μ M TE buffer solutions were prepared.

<Counter oligonucleotides>

50 **[0416]**

C11:5'- GGACCTGTGTTTGACGGGTAT -3' (SEQ ID NO: 51)

C12:5'- AACACTAAGTTGCGCAATTTGCTGT -3' (SEQ ID NO: 52)

C13:5'- ATTGCGAAATCCGCCCGACGATAT -3' (SEQ ID NO: 53)

55

C14:5'- CACTCTTGAGCGCATGTGCCGTTTC -3' (SEQ ID NO: 54)

[0417] Each obtained reaction liquid was subjected to the following treatments.

[0418] In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide

solution, 10 μL of the counter oligonucleotide solution, 10 μL of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc_2 , 5 mM Dithiothreitol), 10 μL of a 100 mM MgCl_2 solution, 10 μL of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μL and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0419] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μL of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

[0420] Then 100 μL of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 $\mu\text{g}/100 \mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0421] One hundred (100) μL of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 60 minutes, 50 μL of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0422] The result is shown in Fig. 12. In Solution MA and Solution MB, increase in absorbance was observed compared to Solution MC. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 13

[0423] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4).

[0424] A 0.5 $\mu\text{g}/\text{mL}$ solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0425] Yeast strain X2180-1A of baker's yeast was cultured in a YPD medium (1% Yeast extract, 2% Peptone, 2% Glucose, pH 5.6 to 6.0) to a turbidity of OD_{600} 0.6 to 1.0, and centrifuged at 10,000 g for 10 minutes, to prepare 1×10^7 of yeast cells. From the prepared yeast cells, a yeast genome was acquired using a generally used preparation method of a yeast genome as described in Methods in Yeast Genetics (Cold Spring Harbor Laboratory).

[0426] The prepared yeast cells were suspended in Buffer A (1 M sorbitol, 0.1 M EDTA, pH 7.4), added with 2-mercaptoethanol (final concentration 14 mM) and 100 U zymolase (10 mg/ml), and incubated under stirring at 30°C for an hour until the solution became clear. After collecting a protoplast by centrifugation at 550 g for 10 minutes, it was suspended in Buffer B (50 mM Tris-HCl, pH 7.4, 20 mM EDTA), added with sodium dodecyl sulfate in 1% (w/v), and then incubated at 65°C for 30 minutes. Sequentially, 5 M CH_3COOK was added and mingled in a volume ratio of 2/5, and the mixture was cooled on ice for 30 minutes, and then centrifuged at 15,000 g for 30 minutes to collect the supernatant. The collected supernatant was added with 3 M CH_3COONa in a volume ratio of 1/10 and an equal amount of isopropanol and mingled well, and the precipitate obtained by centrifugation at 15,000 g at 4°C for 30 minutes was rinsed with 70% ethanol and collected. After drying, the precipitate was dissolved in 1 mL of TE buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA), and added with RNase A (available from Sigma) in a concentration of 40 $\mu\text{g}/\text{ml}$, incubated at 37°C for an hour, and then the mixture was added with proteinase K (available from Sigma) and sodium dodecyl sulfate in a concentrations of 500 $\mu\text{g}/\text{mL}$ and 1% (w/v), respectively, and shaken at 55°C for about 16 hours. After end of the shaking,

EP 2 305 807 A1

the mixture was extracted with phenol [saturated with 1 M Tris-HCl (pH 8.0)]-chloroform. An aqueous layer was collected, added with NaCl in a concentration of 0.5 N, and allowed to precipitate from ethanol, and the generated precipitate was collected. The collected precipitate was rinsed with 70% ethanol, to obtain genomic DNA.

[0427] From the obtained genomic DNA, a DNA fragment to be used as a test sample (S, SEQ ID NO: 29, the region corresponding to the nucleotide numbers 271743-272083 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF3 and PR3) designed for PCR of SEQ ID NO: 27 and SEQ ID NO: 28 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

[0428]

PF3: 5'-AGGTGAGCTACGTGTGTTTGG-3' (SEQ ID NO: 27)

PR3: 5'-AGACATGTGCTCACGTACGGT-3' (SEQ ID NO: 28)

<DNA fragment>

[0429]

S :

5' -AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGC

AGCTTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGT

GGCGGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCGATAGTCGAGCCGCCTGTCAC

GTGCGCGGCCCGCCCTGCTCCGTTTGGACGCGATGCATAGCATGCGACCACCCAGTAATCATAAC

TGCTGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTC

CACACCGTACGTGAGCACATGTCT-3' (SEQ ID NO: 29)

[0430] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μ L of 5 μ M of the above primer solutions, 5 μ L of each 2 mM dNTP, and 5 μ L of 10 \times buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μ L of 5U/ μ L thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μ L. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

[0431] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

[0432] For the DNA fragment S, the following solutions were prepared respectively in duplicate.

Solution A: DNA fragment S 10ng/10 μ L TE buffer solution

Solution B: DNA fragment S 1ng/10 μ L TE buffer solution

Solution C: TE buffer solution (negative control solution)

[0433] From the obtained genomic DNA, a DNA fragment to be used as a test sample (T, SEQ ID NO: 34, the region corresponding to the nucleotide numbers 384569-384685 of yeast chromosome VII shown in Genbank Accession No. NC_001139) was amplified by conducting PCR using oligonucleotide primers (PF4 and PR4) designed for PCR of SEQ ID NO: 32 and SEQ ID NO: 33 and the following reaction condition.

<Oligonucleotide primers designed for PCR>

[0434]

5 PF4: 5'-GGACCTGTGTTTGACGGGTAT-3' (SEQ ID NO: 32)
 PR4: 5'-AGTACAGATCTGGCGTTCTCG-3' (SEQ ID NO: 33)

<DNA fragment>

10 **[0435]**

T:

15 5' -GGACCTGTGTTTGACGGGTATAACACTAAGTTGCGCAATTTGCTGTATTGCGAAATCCG
 CCCGGACGATATCACTCTTGAGCGCATGTGCCGTTTCCGAGAACGCCAGATCTGTACT-3'
 20 (SEQ ID NO: 34)

[0436] As a reaction liquid of PCR, 10 ng of genomic DNA as a template, each 3 μL of 5 μM of the above primer solutions, 5 μL of each 2 mM dNTP, and 5 μL of 10×buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin) were mixed with 0.25 μL of 5U/μL thermostable DNA polymerase (AmpliTaq Gold), and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL. After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 20 seconds at 95°C, 30 seconds at 58°C and 30 seconds at 72°C.

[0437] After conducting PCR, amplification was checked by 2% agarose gel electrophoresis, and the DNA fragment S was purified by Wizard SV Gel/PCR Kit (PROMEGA Corporation).

30 **[0438]** For the DNA fragment T, the following solutions were prepared respectively in duplicate.

- Solution A: DNA fragment T 10ng/10 μL TE buffer solution
- Solution B: DNA fragment T 1ng/10 μL TE buffer solution
- Solution C: TE buffer solution (negative control solution)

35 **[0439]** Solution A of DNA fragment S and Solution A of DNA fragment T, Solution B of DNA fragment S and Solution B of DNA fragment T, and Solution C of DNA fragment S and Solution C of DNA fragment T prepared in the above were respectively mixed, to prepare the following DNA fragment-mixed Solutions MA to MC respectively in duplicate.

- 40 Solution MA: 10 ng/20 μL TE buffer solution
- Solution MB: 1 ng/20 μL TE buffer solution
- Solution MC: TE buffer solution (negative control solution)

[0440] Twenty (20) μL of each obtained solution, 0.5 μL of Sssl methylase (available from NEB Inc.), 5 μL of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μL of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL. The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

[0441] Synthesized were 5'-end FITC-labeled oligonucleotide F3 having the nucleotide sequence of SEQ ID NO: 31 capable of binding by complementation with oligonucleotide S' comprising a target DNA region of SEQ ID NO: 30 and 5'-end FITC-labeled oligonucleotide F4 having the nucleotide sequence of SEQ ID NO: 36 capable of binding by complementation with oligonucleotide T' comprising a target DNA region of SEQ ID NO: 35, and respective 0.02 μM solutions in Tris-HCl buffer (10 mM) were prepared.

<Oligonucleotide comprising target DNA region>

55 **[0442]**

EP 2 305 807 A1

S' : 5' -

5 AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGCAGC
TTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGTGGC
10 GGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCGATAGTCGAGCCGCCTGTCACGTG
CGCGGCCGCCCTGCTCCGTTTGACGCGATGCATAGCATGCGACCACCCAGTAATCATACTGC
TGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTCCAC
15 ACCGTACGTGAGCACATGTCT -3' (SEQ ID NO: 30)

T' : 5' -

20 GGACCTGTGTTTGACGGGTATAACACTAAGTTGCGCAATTTGCTGTATTGCGAAATCCGCCC
GGACGATATCACTCTTGAGCGCATGTGCCGTTTCCGAGAACGCCAGATCTGTACT -3'
(SEQ ID NO: 35)

25 <5'-end FITC-labeled oligonucleotide>

[0443]

30 F3:5'- AGACATGTGCTCACGTACGGT -3' (SEQ ID NO: 31)
F4:5'- AGTACAGATCTGGCGTTCTCG -3' (SEQ ID NO: 36)

35 **[0444]** Synthesized were counter oligonucleotides C1, C2, C3, C4, C5, C6, C7, C8, C9 and C10 having the nucleotide sequences of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48 and SEQ ID NO: 49 and SEQ ID NO: 50, respectively capable of binding by complementation with a minus strand of DNA fragment S' comprising the target DNA region of SEQ ID NO: 30, and respective 0.01 μ M TE buffer solutions were prepared.

40 **[0445]** Synthesized were counter oligonucleotides C11, C12, C13 and C14 having the nucleotide sequences of SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53 and SEQ ID NO: 54, respectively capable of binding by complementation with a minus strand of DNA fragment T' comprising the target DNA region of SEQ ID NO: 35, and respective 0.01 μ M TE buffer solutions were prepared.

<Oligonucleotide comprising target DNA region>

45 **[0446]**

EP 2 305 807 A1

S' : 5' -

AGGTGAGCTACGTGTGTTTGGGCGTCGTGCACTGGCTCACTTGTACGCGCAGAAATGGCAGC
5 TTGTACGATTGGTGACCCGCCTTTTCGACACTGGACCGCTATGGACGTGGCGGCGGTGTGGC
GGCGGCTCAATGACCTGTGGCGCCCGTTTGTGGCGTGCGATAGTCGAGCCGCCTGTCACGTG
10 CGCGGCCCGCCCTGCTCCGTTTGACGCGATGCATAGCATGCGACCACCCAGTAATCATACTGC
TGACGCTATTGGTCACGTGGTTATGGCAGCTGCTGTTGACTGCGGTGGCGTCCCGTTTCCAC
ACCGTACGTGAGCACATGTCT -3' (SEQ ID NO: 30)

T' : 5' -

20 GGACCTGTGTTTGACGGGTATAACACTAAGTTGCGCAATTTGCTGTATTGCGAAATCCGCCC
GGACGATATCACTCTTGAGCGCATGTGCCGTTTCCGAGAACGCCAGATCTGTACT -3'
(SEQ ID NO: 35)

<Counter oligonucleotides>

[0447]

30 C1:5'- AGGTGAGCTACGTGTGTTTGG -3' (SEQ ID NO: 41)
C2:5'- GCGTCGTGCACTGGCTCACTTGTACGCGCA -3' (SEQ ID NO: 42)
C3:5'- CTTGTACGATTGGTGACCCGCCTTTTCGAC -3' (SEQ ID NO: 43)
C4:5'- ACTGGACCGCTATGGACGTGGCGGCGGTGT -3' (SEQ ID NO: 44)
C5:5'- GGC GGCGGCTCAATGACCTGTGGCGCCCGT -3' (SEQ ID NO: 45)
35 C6:5'- TTGTGGCGTGCGATAGTCGAGCCGCCTGTC -3' (SEQ ID NO: 46)
C7:5'- ACGTGCGCGGCCCGCCCTGCTCCGTT -3' (SEQ ID NO: 47)
C8:5'- TGACGCGATGCATAGCATGCGACCACCCAG -3' (SEQ ID NO: 48)
C9:5'- ACTGCTGACGCTATTGGTCACGTGGTTATG -3' (SEQ ID NO: 49)
C10:5'- CTGCTGTTGACTGCGGTGGCGTCCCGTTTC -3' (SEQ ID NO: 50)
40 C11:5'- GGACCTGTGTTTGACGGGTAT -3' (SEQ ID NO: 51)
C12:5'- AACACTAAGTTGCGCAATTTGCTGT -3' (SEQ ID NO: 52)
C13:5'- ATTGCGAAATCCGCCCCGACGATAT -3' (SEQ ID NO: 53)
C14:5'- CACTCTTGAGCGCATGTGCCGTTTC -3' (SEQ ID NO: 54)

45 [0448] Each obtained reaction liquid was subjected to the following treatments.

[0449] In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of the counter oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed.
50 Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0450] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μ L of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of

the present method).

[0451] Then 100 μL of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 $\mu\text{g}/100 \mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was

added with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0452] One hundred (100) μL of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 60 minutes, 50 μL of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0453] The result is shown in Fig. 13. In Solution MA and Solution MB, increase in absorbance was observed compared to Solution MC. The intensity increased depending on the concentration of the DNA fragment. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 14

[0454] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylation kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4).

[0455] A 0.5 $\mu\text{g}/\text{mL}$ solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0456] Using genomic DNA derived from human blood purchased from Clontech, the following solutions were prepared respectively in duplicate.

Solution A: Genomic DNA derived from human blood 100 ng/5 μL TE buffer solution

Solution B: Genomic DNA derived from human blood 10 ng/5 μL TE buffer solution

Solution C: Genomic DNA derived from human blood 1 ng/5 μL TE buffer solution

Solution D: TE buffer solution (negative control solution)

[0457] Five (5) μL of each obtained solution, 10 U of restriction enzyme Xspl, and 2 μL of 10x buffer optimum for Xspl (200 mM Tris-HCl pH 8.5, 100 mM MgCl_2 , 10 mM Dithiothreitol, 1000 mM KCl) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 20 μL . The reaction liquid was incubated at 37°C for 1 hour.

[0458] Twenty (20) μL of each obtained solution, 0.5 μL of Sssl methylase (available from NEB Inc.), 5 μL of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μL of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL . The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

[0459] Synthesized was 5'-end FITC-labeled oligonucleotide F5 having the nucleotide sequence of SEQ ID NO: 38 capable of binding by complementation with oligonucleotide Z (region corresponding to the nucleotide numbers 115-386 shown in Genbank Accession No. M80340) comprising a target DNA region of SEQ ID NO: 37 designed in LINE1 region known as human transposon, and a 0.02 μM solution in Tris-HCl buffer (10 mM) was prepared.

<Oligonucleotide comprising target DNA region>

[0460]

Z: 5' -

EP 2 305 807 A1

TAGGGAGTGCCAGACAGTGGGCGCAGGCCAGTGTGTGTGCGCACCGTGCGCGAGCCGAAGCA
5 GGGCGAGGCATTGCCTCACCTGGGAAGCGCAAGGGGTCAGGGAGTTCCTTTCTGAGTCAA
GAAAGGGGTGACGGTCGCACCTGGAAAATCGGGTCACTCCCACCCGAATATTGCGCTTTTCA
GACCGGCTTAAGAAACGGCGCACCCACGAGACTATATCCCACACCTGGCTCGGAGGGTCCTAC
10 GCCCACGGAATCTCGCTGATTGC -3' (SEQ ID NO: 37)

<5'-end FITC-labeled oligonucleotide>

[0461]

F5:5'- ATAGTCTCGTGGTGCGCCGT -3' (SEQ ID NO: 38)

[0462] Synthesized were counter oligonucleotides C15, C16, C17, C18 and C19 having the nucleotide sequences of
20 SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58 and SEQ ID NO: 59, respectively capable of binding
by complementation with a minus strand of DNA fragment W comprising the target DNA region of SEQ ID NO: 39, and
respective 0.01 μ M TE buffer solutions were prepared.

<Oligonucleotide comprising target DNA region>

[0463]

Z: 5' -

30 TAGGGAGTGCCAGACAGTGGGCGCAGGCCAGTGTGTGTGCGCACCGTGCGCGAGCCGAAGCA
GGGCGAGGCATTGCCTCACCTGGGAAGCGCAAGGGGTCAGGGAGTTCCTTTCTGAGTCAA
35 GAAAGGGGTGACGGTCGCACCTGGAAAATCGGGTCACTCCCACCCGAATATTGCGCTTTTCA
GACCGGCTTAAGAAACGGCGCACCCACGAGACTATATCCCACACCTGGCTCGGAGGGTCCTAC
GCCCACGGAATCTCGCTGATTGC -3' (SEQ ID NO: 37)

<Counter oligonucleotides>

[0464]

45 C15:5'- CAGTGTGTGTGCGCACCGTGCGCGAGCCGA -3' (SEQ ID NO: 55)
C16:5'- GGCGAGGCATTGCCTCACCTGGGAAGCGCA -3' (SEQ ID NO: 56)
C17:5'- GGTGACGGTCGCACCTGGAAAATCGGGTCA -3' (SEQ ID NO: 57)
C18:5'- ACCCGAATATTGCGCTTTTCAGACCGGCTT -3' (SEQ ID NO: 58)
50 C19:5'- TCGGAGGGTCCTACGCCACGGAATCTCGC -3' (SEQ ID NO: 59)

[0465] Each obtained reaction liquid was subjected to the following treatments.

[0466] In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide
solution, 10 μ L of the counter oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc,
100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were
55 added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed.
Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this
temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained
at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-

labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0467] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μL of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature for 1 hour. Then the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

[0468] Then 100 μL of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 $\mu\text{g}/100 \mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0469] One hundred (100) μL of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 9 minutes, 50 μL of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0470] The result is shown in Fig. 14. In Solution A, Solution B, and Solution C, increase in absorbance was observed compared to Solution D. The intensity increased depending on the concentration of genomic DNA. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 15

[0471] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin using a commercially available biotinylation kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the catalogue. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4).

[0472] A 0.5 $\mu\text{g}/\text{mL}$ solution of the synthetically obtained biotin-labeled methylcytosine antibody in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4) was prepared, and each 100 μL of this was added to an 8-well strip coated with streptavidin (available from PerkinElmer), and left still at room temperature for about 1 hour to immobilize it to the wells. Thereafter, the solution was removed by pipetting, and 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by decantation. This operation was repeated two more times (the above corresponds to preparation of an immobilized methylated DNA antibody used in the present method).

[0473] Using genomic DNA derived from human blood purchased from Clontech, the following solutions were prepared respectively in duplicate.

- Solution A: Genomic DNA derived from human blood 100 ng/5 μL TE buffer solution
- Solution B: Genomic DNA derived from human blood 10 ng/5 μL TE buffer solution
- Solution C: Genomic DNA derived from human blood 1 ng/5 μL TE buffer solution
- Solution D: TE buffer solution (negative control solution)

[0474] Five (5) μL of each obtained solution, 4 U of restriction enzyme MspI, and 2 μL of 10x buffer optimum for MspI (100 mM Tris-HCl pH 8.5, 100 mM MgCl_2 , 10 mM Dithiothreitol, 500 mM NaCl) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 20 μL . The reaction liquid was incubated at 37°C for 1 hour.

[0475] Twenty (20) μL of each obtained solution, 0.5 μL of SssI methylase (available from NEB Inc.), 5 μL of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μL of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL . The reaction liquid was incubated at 37°C for 30 minutes (the foregoing process corresponds to First step of the present method).

[0476] Synthesized was 5'-end FITC-labeled oligonucleotide F6 having the nucleotide sequence of SEQ ID NO: 40 capable of binding by complementation with oligonucleotide W (region corresponding to the nucleotide numbers 178-262 shown in Genbank Accession No. AF458110) comprising a target DNA region of SEQ ID NO: 39 designed in Alu region known as human transposon, and a 0.02 μM solution in Tris-HCl buffer (10 mM) was prepared.

EP 2 305 807 A1

<Oligonucleotide comprising target DNA region>

[0477]

5

W: 5' -

CGGGCGCGGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCCGAGGTGGGCGGATCACG

10

AGGTCAGGAGATCGAGACCATCC -3' (SEQ ID NO: 39)

15 <5'-end FITC-labeled oligonucleotide>

[0478]

F6:5'- GGATGGTCTCGATCTCCTGAC -3' (SEQ ID NO: 40)

20

[0479] Synthesized were counter oligonucleotides C20 and C21 having the nucleotide sequences of SEQ ID NO: 60 and SEQ ID NO: 61, respectively capable of binding by complementation with a minus strand of DNA fragment W comprising the target DNA region of SEQ ID NO: 39, and respective 0.01 μ M TE buffer solutions were prepared.

25 <Oligonucleotide comprising target DNA region>

[0480]

30

W: 5' -

CGGGCGCGGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCCGAGGTGGGCGGATCACG

AGGTCAGGAGATCGAGACCATCC -3' (SEQ ID NO: 39)

35

<Counter oligonucleotides>

[0481]

40

C20:5'- CGGGCGCGGTGGCTCACGCCTGTAATCCCA -3' (SEQ ID NO: 60)

C21:5'- TTTGGGAGGCCGAGGTGGGCGGATCACGAG -3' (SEQ ID NO: 61)

[0482] Each obtained reaction liquid was subjected to the following treatments.

45 **[0483]** In a PCR tube, 40 μ L of the reaction liquid prepared in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of the counter oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc₂, 5 mM Dithiothreitol), 10 μ L of a 100 mM MgCl₂ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the resultant mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L and mixed. Then the PCR tube was heated at 95°C for 10 minutes, cooled rapidly to 70°C, and retained for 10 minutes at this

50

temperature. Then the mixture was cooled to 50°C and retained at the temperature for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature, to promote formation of a conjugate of the 5'-end FITC-labeled oligonucleotide and the DNA fragment (the foregoing process corresponds to Second step of the present method).

[0484] To an 8-well strip coated with streptavidin to which the biotin-labeled methylcytosine antibody has been immobilized, 100 μ L of the reaction liquid of the DNA fragment prepared above was added, and left still at room temperature

55 for 1 hour. Then the solution was removed by pipetting, and 200 μ L of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂ PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH7.4)] was added, and then the buffer was removed by pipetting. This operation was repeated two more times (the foregoing process corresponds to Third step of the present method).

EP 2 305 807 A1

[0485] Then 100 μL of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 $\mu\text{g}/100 \mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was added with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0486] One hundred (100) μL of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction. After leaving still at room temperature for about 8 minutes, 50 μL of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured (the foregoing process corresponds to Fourth step of the present method).

[0487] The result is shown in Fig. 15. In Solution A, Solution B, and Solution C, increase in absorbance was observed compared to Solution D. The intensity increased depending on the concentration of genomic DNA. In this experiment, it was revealed that the DNA fragment can be detected and quantified by forming and selecting a complex of the methylcytosine antibody, the methylated DNA fragment, and the immobilized 5'-end biotin-labeled oligonucleotide, and quantifying or detecting FITC in the complex by its function.

Example 16

[0488] As a serum sample, mixed liquids of a TE buffer solution of genomic DNA derived from human blood DNA (Human Genomic DNA, #636401, Clontech) and serum collected from rat (Wistar Hannover) were prepared respectively in quadruplicate as follows.

[0489] Serum sample A: Genomic DNA derived from human blood 10 ng/10 μL TE buffer solution + rat serum 10 μL

[0490] Serum sample B: Genomic DNA derived from human blood 1 ng/10 μL TE buffer solution + rat serum 10 μL

[0491] Serum sample C: Genomic DNA derived from human blood 0.1 ng/10 μL TE buffer solution + rat serum 10 μL

[0492] Serum sample D: 0 ng/10 μL TE buffer solution + rat serum 10 μL (negative control)

[0493] For Serum samples A to D prepared in the above, Treatment 1 or Treatment 2 was conducted respectively in duplicate. Treatment 1:

[0494] Twenty (20) μL of a serum sample and 4 μL of a buffer (500 mM Tris-HCl (pH 7.5), 100 mM MgCl_2 , 10 mM DTT, 1000 mM NaCl) were mixed, and the mixture was added with sterilized ultrapure water to make a liquid amount 40 μL , and mixed. Then, the PCR tube was retained at 95°C for 10 minutes, retained at 4°C for 10 minutes, and then returned to room temperature. After centrifugation at 9100xg for 10 minutes, the supernatant was collected.

Treatment 2:

[0495] Twenty (20) μL of a serum sample and 4 μL of a buffer (330 mM Tris-Acetate (pH 7.9), 100 mM $\text{Mg}(\text{OAc})_2$, 5 mM DTT, 660 mM KOAc) were mixed, and the mixture was added with sterilized ultrapure water to make a liquid amount 40 μL , and mixed. Then, the PCR tube was retained at 95°C for 10 minutes, retained at 4°C for 10 minutes, and then returned to room temperature. After centrifugation at 9100xg for 10 minutes, the supernatant was collected.

[0496] Twenty (20) μL of each solution prepared by Treatment 1 or Treatment 2, 2U of restriction enzyme MspI, and 5 μL of 10x buffer optimum for MspI (100 mM Tris-HCl pH 7.5, 100 mM MgCl_2 , 10 mM Dithiothreitol, 500 mM NaCl) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL . The reaction liquid was incubated at 37°C for 1 hour.

[0497] Thirty (30) μL of the solution obtained by the above enzyme treatment, 0.5 μL of SssI methylase (available from NEB Inc.), 5 μL of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μL of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultrapure water to prepare a reaction liquid having a liquid amount of 50 μL . The reaction liquid was incubated at 37°C for 30 minutes.

[0498] As a specific oligonucleotide used for obtaining a target DNA region (W, SEQ ID NO: 39, region corresponding to the nucleotide number 178-262 shown in Genbank Accession No. AF458110) designed in Alu region known as human transposon and having the nucleotide sequence of SEQ ID NO: 39, was synthesized 5'-end biotin-labeled oligonucleotide F1 comprising the nucleotide sequence of SEQ ID NO: 40 that binds by complementation with a plus strand of the target DNA region W, a 0.02 μM solution in Tris-HCl buffer (10 mM) was prepared

<Target DNA region>

[0499]

W: 5' -

CGGGCGCGGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCCGAGGTGGGCGGATCACG
 AGGTCAGGAGATCGAGACCATCC -3' (SEQ ID NO: 39)

<5'-end FITC-labeled oligonucleotide>

[0500]

F1:5'- GGATGGTCTCGATCTCCTGAC -3' (SEQ ID NO: 40)

[0501] Fifty (50) μL of the reaction liquid obtained in the above, 10 μL of the 5'-end FITC-labeled oligonucleotide solution, 10 μL of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc_2 , 5 mM Dithiothreitol), 10 μL of a 100 mM MgCl_2 solution, 10 μL of a 1 mg/mL BSA solution were added, and the mixture was further added with sterilized ultrapure water to make a liquid amount 100 μL , and mixed. Then for forming a double strand between the target DNA region and the 5'-end FITC-labeled oligonucleotide, the PCR tube was retained at 95°C for 10 minutes, rapidly cooled to 70°C, and retained at this temperature for 10 minutes. Then the PCR tube was cooled to 50°C and retained for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature.

[0502] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin by using a commercially available biotinylating kit (Biotin Labeling Kit-NH2, available from DOJINDO LABORATORIES) according to the method described in the protocol. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4).

[0503] 100 μL of the reaction liquid obtained by the above heat treatment was added with 1 μL of a x5 diluted solution of the biotin-labeled methylcytosine antibody (0.05 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4)), and left still at room temperature for 1 hour, to form a detection complex comprising the target DNA region, the 5'-end FITC-labeled oligonucleotide, and the biotin-labeled methylcytosine antibody.

[0504] The reaction liquid obtained in the above was transferred to an 8-well strip coated with streptavidin (StreptaWell, #11645692001, available from Roche), and left still at room temperature for about 60 minutes, to immobilize the detection complex comprising the target DNA region, the 5'-end FITC-labeled oligonucleotide and the biotin-labeled methylcytosine antibody via a biotin-streptavidin bond. Thereafter, the solution was removed by decantation, and each well was washed three times with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4)].

[0505] Thereafter, 100 μL of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 $\mu\text{g}/100 \mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was washed with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4)], and the buffer was removed by decantation. This operation was repeated two more time.

[0506] One hundred (100) μL of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction.

[0507] The reaction was left still at room temperature for about 30 minutes, and 50 μL of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured, and an average value was calculated in duplicate for the obtained measured values.

[0508] The results are shown in Fig. 16 and Fig. 17. In Treatment 1, absorbance increased depending on the concentration in Solution A (10 ng), Solution B (1 ng), and Solution C (0.1 ng) of genomic DNA derived from human blood, compared to Solution D (0 ng: control solution) (Fig. 16). On the other hand, in Treatment 2, absorbance increased in Solution A (10 ng) of genomic DNA derived from human blood, compared to Solution D (0 ng: control solution), however, in Solution B (1 ng) and Solution C (0.1 ng), increase in absorbance was not observed (Fig. 17).

[0509] In the present experiment, it was revealed that human genomic DNA in serum can be detected and quantified with high sensitivity by forming and selecting a complex of an immobilized biotin-labeled methylcytosine antibody, a methylated DNA fragment, and a 5'-end FITC-labeled oligonucleotide, and quantifying and detecting FITC in the complex according to its function. In Treatment 1, human genomic DNA in serum was detected with better sensitivity than in Treatment 2.

EP 2 305 807 A1

Example 17

[0510] As a serum sample, mixed liquids of a TE buffer solution of genomic DNA derived from human blood (Human Genomic DNA, #636401, Clontech) and a human serum purchased from Kohjin Bio Co., Ltd (individual human serum) were prepared respectively in duplicate as follows.

Serum sample A: Genomic DNA derived from human blood 4 ng/10 μ L TE buffer solution + human serum 40 μ L
Serum sample B: Genomic DNA derived from human blood 2 ng/10 μ L TE buffer solution + human serum 40 μ L
Serum sample C: Genomic DNA derived from human blood 1 ng/10 μ L TE buffer solution + human serum 40 μ L
Serum sample D: Genomic DNA derived from human blood 0 ng/10 μ L TE buffer solution + human serum 40 μ L (negative control solution)

For Serum samples A to D prepared above, the following treatments were conducted respectively in duplicate.

[0511] In a PCR tube, 50 μ L of a serum sample, and 20 μ L of a buffer (500 mM Tris-HCl (pH 7.5), 100 mM $MgCl_2$, 10 mM DTT, 1000 mM NaCl) were charged, and the result mixture was added with sterilized ultra pure water to make the liquid amount 100 μ L, and mixed. Then the PCR tube was retained at 95°C for 10 minutes, and cooled to 4°C, and then returned to room temperature. After centrifugation at 9100xg for 10 minutes, 20 μ L of the supernatant was collected.

[0512] Twenty (20) μ L of the solution prepared by the above treatment, 2 U of restriction enzyme MspI, and 5 μ L of a 10xbuffer optimum for MspI (100 mM Tris-HCl pH 7.5, 100 mM $MgCl_2$, 10 mM Dithiothreitol, 500 mM NaCl) were mixed, and further added with sterilized ultra pure water to make the liquid amount 50 μ L. The reaction liquid was incubated at 37°C for 1 hour.

[0513] Thirty (30) μ L of the solution obtained by the above enzyme treatment, 0.5 μ L of SssI methylase (available from NEB Inc.), 5 μ L of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μ L of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultra pure water to make the liquid amount 50 μ L. The reaction liquid was incubated at 37°C for 30 minutes.

[0514] Synthesized was 5'-end FITC-labeled oligonucleotide F1 comprising the nucleotide sequence of SEQ ID NO: 40 capable of binding by complementation with the target DNA region (W, SEQ ID NO: 39, the region corresponding to base number 178-262 shown in Genbank Accession No. AF458110) designed in an Alu region known as human transposon, and a 0.02 μ M solution in Tris-HCl buffer (10 mM) was prepared.

<Target DNA region>

[0515]

W: 5' -

CGGGCGCGGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCCGAGGTGGGCGGATCACG

AGGTCAGGAGATCGAGACCATCC -3' (SEQ ID NO: 39)

<5'-end biotin-labeled oligonucleotide>

[0516]

F6:5'- GGATGGTCTCGATCTCCTGAC -3' (SEQ ID NO: 40)

[0517] Fifty (50) μ L of the reaction liquid obtained in the above, 10 μ L of the 5'-end FITC-labeled oligonucleotide solution, 10 μ L of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM $MgOAc_2$, 5 mM Dithiothreitol), 10 μ L of a 100 mM $MgCl_2$ solution, 10 μ L of a 1 mg/mL BSA solution were added, and the mixture was further added with sterilized ultrapure water to make a liquid amount 100 μ L, and mixed. Then for forming a double strand between the target DNA region and the 5'-end FITC-labeled oligonucleotide, the PCR tube was retained at 95°C for 10 minutes, rapidly cooled to 70°C, and retained at this temperature for 10 minutes. Then the PCR tube was cooled to 50°C and retained for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature.

[0518] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with

EP 2 305 807 A1

biotin by using a commercially available biotinylating kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the protocol. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 µg/µL solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH 7.4).

5 **[0519]** 100 µL of the reaction liquid obtained by the above heat treatment was added with 1 µL of a x5 diluted solution of the biotin-labeled methylcytosine antibody (0.05 µg/µL solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH 7.4)), and left still at room temperature for 1 hour, to form a detection complex comprising the target DNA region, the 5'-end FITC-labeled oligonucleotide, and the biotin-labeled methylcytosine antibody.

10 **[0520]** The reaction liquid obtained in the above was transferred to an 8-well strip coated with streptavidin (StreptaWell, #11645692001, available from Roche), and left still at room temperature for about 60 minutes, to immobilize the detection complex comprising the target DNA region, the 5'-end FITC-labeled oligonucleotide and the biotin-labeled methylcytosine antibody via a biotin-streptavidin bond. Thereafter, the solution was removed by decantation, and each well was washed three times with 200 µL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO₄·7H₂O, 154 mM NaCl pH 7.4)].

15 **[0521]** Thereafter, 100 µL of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 µg/100 µL solution in 0.1% BSA-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH 7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was washed with 200 µL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH₂PO₄, 3 mM Na₂HPO 7H₂O, 154 mM NaCl pH 7.4)], and the buffer was removed by decantation. This operation was repeated two more time.

20 **[0522]** One hundred (100) µL of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction.

25 **[0523]** The reaction was left still at room temperature for about 20 minutes, and 50 µL of a stop solution (2N H₂SO₄ aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured.

[0524] The results are shown in Fig. 18. In Solution A (4 ng), Solution B (2 ng), and Solution C (1 ng) of genomic DNA derived from human blood, absorbance increased depending on the concentration, compared to Solution D (0 ng: control solution).

30 **[0525]** In the present experiment, it was revealed that human genomic DNA in human serum can be detected and quantified by forming a detection complex made up of a target DNA region, a 5'-end FITC-labeled oligonucleotide, and a biotin-labeled methylcytosine antibody, using the DNA sample extracted by the present inventive method, and selecting by immobilization via a biotin-streptavidin bond, and detecting FITC in the complex according to its function.

35 Example 18

[0526] As a serum sample, the following human serums were used. Human serums purchased from Kohjin Bio Co., Ltd (individual human serums)

Lot No.:

40

N51438 (healthy subject)
N51439 (healthy subject)
N51441 (healthy subject)

45 Human serums purchased from ProMedDx (individual human serums) Lot No.:

11171268 (healthy subject, age 56, male)
11171292 (healthy subject, age 62, male)
11171297 (healthy subject, age 67, male)
50 11202510 (healthy subject, age 67, female)
11202522 (healthy subject, age 64, female)
11202527 (healthy subject, age 52, female)
11202615 (healthy subject, age 75, female)
11202618 (healthy subject, age 78, female)
55 10958886 (healthy subject, age 56, male)
10958979 (healthy subject, age 39, male)
10958980 (healthy subject, age 45, male)
10960268 (healthy subject, age 37, male)

EP 2 305 807 A1

10960272 (healthy subject, age 50, male)
10960276 (healthy subject, age 30, male)
10960285 (healthy subject, age 39, male)
11003457 (healthy subject, age 38, male)
5 11003479 (healthy subject, age 51, male)
11003480 (healthy subject, age 48, male)
11324997 (healthy subject, age 59, male)
11325001 (healthy subject, age 61, male)
10325022 (healthy subject, age 61, male)
10 10870623 (breast cancer patient, age 33, female)
10929521 (breast cancer patient, age 55, female)
10989644 (breast cancer patient, age 45, female)
11209430 (breast cancer patient, age 80, female)
10929514 (breast cancer patient, age 57, female)
15 10843055 (breast cancer patient, age 59, female)
10984680 (breast cancer patient, age 64, female)
10840414 (lung cancer patient, age 54, female)
10929506 (lung cancer patient, age 55, male)
11091955 (lung cancer patient, age 76, female)
20 11103346 (lung cancer patient, age 66, female)
11142322 (lung cancer patient, age 62, female)
11152564 (lung cancer patient, age 67, male)
11152571 (lung cancer patient, age 67, male)
11153198 (lung cancer patient, age 69, female)
25 11209435 (lung cancer patient, age 61, male)
11230621 (lung cancer patient, age 71, female)
11153192 (lung cancer patient, age 59, male)
10715942 (lung cancer patient, age 64, male)
10840422 (lung cancer patient, age 78, female)
30 10935547 (prostate cancer patient, age 83, male)
11000243 (prostate cancer patient, age 78, male)
11071226 (prostate cancer patient, age 84, male)

35 **[0527]** For the above Serum samples, the following treatments were conducted respectively in duplicate.

Treatment 1:

40 **[0528]** In a PCR tube, 40 μL of a serum sample, and 20 μL of a buffer (500 mM Tris-HCl (pH 7.5), 100 mM MgCl_2 , 10 mM DTT, 1000 mM NaCl) were charged, and the result mixture was added with sterilized ultra pure water to make the liquid amount 100 μL , and mixed. Then the PCR tube was retained at 95°C for 10 minutes, and cooled to 4°C, and then returned to room temperature. After centrifugation at 9100xg for 10 minutes, 20 μL of the supernatant was collected.

45 **[0529]** Twenty (20) μL of the solution prepared by the above treatment, 2 U of restriction enzyme MspI, and 5 μL of a 10xbuffer optimum for MspI (100 mM Tris-HCl pH 7.5, 100 mM MgCl_2 , 10 mM Dithiothreitol, 500 mM NaCl) were mixed, and further added with sterilized ultra pure water to make the liquid amount 50 μL . The reaction liquid was incubated at 37°C for 1 hour.

[0530] Thirty (30) μL of the solution obtained by the above enzyme treatment, 0.5 μL of SssI methylase (available from NEB Inc.), 5 μL of 10xNEBuffer2 (available from NEB Inc.), and 0.5 μL of 3.2 mM S-adenosyl methionine (available from NEB Inc.) were mixed, and added with sterilized ultra pure water to make the liquid amount 50 μL . The reaction liquid was incubated at 37°C for 30 minutes.

50 **[0531]** Synthesized was 5'-end FITC-labeled oligonucleotide F6 comprising the nucleotide sequence of SEQ ID NO: 40 capable of binding by complementation with the target DNA region (W, SEQ ID NO: 39, the region corresponding to base number 178-262 shown in Genbank Accession No. AF458110) designed in an Alu region known as human transposon, and a 0.02 μM solution in Tris-HCl buffer (10 mM) was prepared.

55 <Target DNA region>

[0532]

W: 5' -

CGGGCGCGGTGGCTCACGCCTGTAATCCAGCACTTTGGGAGGCCGAGGTGGGCGGATCACG
 AGGTCAGGAGATCGAGACCATCC -3' (SEQ ID NO: 39)

<5'-end biotin-labeled oligonucleotide>

[0533]

F6:5'- GGATGGTCTCGATCTCCTGAC -3' (SEQ ID NO: 40)

[0534] Fifty (50) μL of the reaction liquid obtained in the above, 10 μL of the 5'-end FITC-labeled oligonucleotide solution, 10 μL of a buffer (330 mM Tris-Acetate pH 7.9, 660 mM KOAc, 100 mM MgOAc_2 , 5 mM Dithiothreitol), 10 μL of a 100 mM MgCl_2 solution, 10 μL of a 1 mg/mL BSA solution were added, and the mixture was further added with sterilized ultrapure water to make a liquid amount 100 μL , and mixed. Then for forming a double strand between the target DNA region and the 5'-end FITC-labeled oligonucleotide, the PCR tube was retained at 95°C for 10 minutes, rapidly cooled to 70°C, and retained at this temperature for 10 minutes. Then the PCR tube was cooled to 50°C and retained for 10 minutes, and further retained at 37°C for 10 minutes, and returned to room temperature.

[0535] A commercially available methylcytosine antibody (available from Aviva Systems Biology) was labeled with biotin by using a commercially available biotinylation kit (Biotin Labeling Kit-NH₂, available from DOJINDO LABORATORIES) according to the method described in the protocol. The obtained biotin-labeled methylcytosine antibody was refrigerated as a 0.25 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4).

[0536] 100 μL of the reaction liquid obtained by the above heat treatment was added with 1 μL of a x5 diluted solution of the biotin-labeled methylcytosine antibody (0.05 $\mu\text{g}/\mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4)), and left still at room temperature for 1 hour, to form a detection complex comprising the target DNA region, the 5'-end FITC-labeled oligonucleotide, and the biotin-labeled methylcytosine antibody.

[0537] The reaction liquid obtained in the above was transferred to an 8-well strip coated with streptavidin (StreptaWell, #11645692001, available from Roche), and left still at room temperature for about 60 minutes, to immobilize the detection complex comprising the target DNA region, the 5'-end FITC-labeled oligonucleotide and the biotin-labeled methylcytosine antibody via a biotin-streptavidin bond. Thereafter, the solution was removed by decantation, and each well was washed three times with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4)].

[0538] Thereafter, 100 μL of a HRP-labeled FITC antibody solution [available from Jackson ImmunoResearch Laboratories, 0.005 $\mu\text{g}/100 \mu\text{L}$ solution in 0.1% BSA-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4)] was added to each well, and left still at room temperature for 1 hour. After leaving still, each well was washed with 200 μL of a washing buffer [0.05% Tween20-containing phosphate buffer (1 mM KH_2PO_4 , 3 mM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 154 mM NaCl pH 7.4)], and the buffer was removed by decantation. This operation was repeated two more times.

[0539] One hundred (100) μL of a substrate (available from R&D, #DY999) was added to each well and mixed, to start the reaction.

[0540] The reaction was left still at room temperature for about 25 minutes, and 50 μL of a stop solution (2N H_2SO_4 aqueous solution) was added to each well to stop the reaction. Within 30 minutes after stopping of the reaction, absorbance at 450 nm was measured, and an average value was calculated in duplicate for the obtained measured values.

[0541] On the other hand, DNA in the solution obtained by the above enzyme treatment (MspI treatment) was quantified by real time PCR.

[0542] As a standard sample for measuring concentration, an MspI-treated human genomic DNA solution was prepared in the following manner. A 5 ng/ μL TE buffer solution of genomic DNA derived from human blood (Human Genomic DNA, #636401, available from Clontech) in a TE buffer solution was prepared, and 20 μL of this solution, 2 U of restriction enzyme MspI, and 5 μL of a 10xbuffer optimum for MspI (100 mM Tris-HCl pH 7.5, 100 mM MgCl_2 , 10 mM Dithiothreitol, 500 mM NaCl) were mixed, and added with sterilized ultra pure water to make the liquid amount 50 μL . This was prepared for each treatment. The reaction liquid was incubated at 37°C for 1 hour. For the obtained reaction liquid, 10^{-5} , 10^{-4} , 10^{-1} , 10^{-2} , 10^{-1} , 1, 10 ng/5 μL solutions were prepared by dilution with TE buffer.

[0543] For amplifying the target DNA region (W, SEQ ID NO: 39, the region corresponding to base number 178-262

EP 2 305 807 A1

shown in Genbank Accession No. AF458110) designed in an Alu region known as human transposon and quantifying it by real time PCR, Forward primer (F) comprising the nucleotide sequence of SEQ ID NO: 62 and Reverse primer (R) comprising the nucleotide sequence of SEQ ID NO: 63 were designed.

5 <Target DNA region>

[0544]

10 W: 5' -

CGGGCGCGGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCCGAGGTGGGCGGATCACG

AGGTCAGGAGATCGAGACCATCC -3' (SEQ ID NO: 39)

15

<Forward primer>

[0545]

20

F:5'- GGTGGCTCACGCCTGTAATC -3' (SEQ ID NO: 62)

<Reverse primer>

25

[0546]

R:5'- GGATGGTCTCGATCTCCTGAC -3' (SEQ ID NO: 63)

30

[0547] As a reaction liquid of PCR, a liquid prepared by mixing 5 μ L of an MspI-treated human genomic DNA solution prepared in the above which is to be a template, or a standard sample for concentration measurement prepared in the above, and each 1.5 μ L of 5 μ M solutions of primers comprising the nucleotide sequences of SEQ ID NO: 62 and SEQ ID NO: 63, 0.1x amount of SYBR® Green I (available from Lonza), each 2.5 μ L of 2 mM dNTPs, 2.5 μ L of a 10xPCR buffer (100 mM Tris-HCl pH 8.3, 500 mM KCl, 15 mM MgCl₂, 0.01% Gelatin), and 0.125 μ L of heat-resistant DNA polymerase (AmpliTaq Gold, 5 U/ μ L, available from ABI) and adding with sterilized ultra pure water to make the liquid amount 25 μ L was used. Real-time PCR was conducted using Mx3005P (Stratagene). After retaining the reaction liquid at 95°C for 10 minutes, PCR was conducted by 40 cycles each consisting of 30 seconds at 95°C, 30 seconds at 61°C and 45 seconds at 72°C, to amplify the target DNA region. According to a result of the real-time PCR, DNA in a serum sample was quantified.

35

40

[0548] The results are shown in Fig. 19 and Fig. 20. A measured value by the present method, and a value quantified by the real-time PCR were compared, to reveal that there is a correlation (coefficient of correlation: R = 0.62)(Fig. 19). A comparison between cancer patients and healthy subjects was made for the result of quantification for human serum samples aged 57 or younger, to reveal that serum DNA concentration increased in cancer patient compared to healthy subjects (Fig. 20).

45

[0549] In the present experiment, it was revealed that free DNA in human serum can be detected and quantified with high sensitivity by forming and selecting a complex of a methylcytosine antibody, a methylated DNA fragment, and a 5'-end biotin-labeled oligonucleotide, and detecting the methylcytosine antibody in the complex according to its function. It was also revealed that serum DNA concentration differs between cancer patients and healthy subjects at age 57 or younger, and the difference can be detected in a simple manner by the present method.

50

INDUSTRIAL APPLICABILITY

55

[0550] According to the present invention, it becomes possible to provide a method for quantifying or detecting DNA having a target DNA region in a simple and convenient manner. Further, it becomes possible to provide a method for selecting a specimen derived from a cancer patient by using a specimen derived from a test subject (preferably serum) and comparing a result in the specimen and a result in a specimen derived from a healthy subject, and so on.

Free Text in Sequence Listing

SEQ ID NOs:17 to 63

5 Designed oligonucleotide

[0551]

10

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

<110> Sumitomo Chemical Co., Ltd.

<120> Method for determining or detecting DNA

<130> S21066W001

<150> JP2008-152619

<151> 2008-06-11

<160> 63

<210> 1

<211> 2661

<212> DNA

<213> Homo sapiens

<400> 1

acagacatgt	gccaccatgc	ccagctaatt	ttttgtttgt	ttgtttgttt	gtttgtattt	60
ttagtagaga	tggggttttg	ccatgttggc	caggctggtc	tcgaactcct	gacctcgaat	120
gataatgac	cgccgcttgg	cctccaaagt	gctaggatta	caggtgtgag	ccactgcgcc	180
aggcctgggc	actttcttta	gtagtttgag	gagcaacatt	tttgacagtg	tccttctgct	240
caagattcaa	gatcccagat	aaaattaaac	catctagaga	gatggcttga	ttggccaaac	300
ctggatctca	tgaccacttc	ttgaagtggg	taagtctcat	aaatgctcag	tccttccact	360
atgcaactga	gtggggtggg	tgggaagccc	ctcaaaggaa	aatccggttg	ttcttactag	420
aaagaaaagg	aaaaatggatg	tgaggcagtc	aaaatcagca	gaggtccacc	acaccaccaa	480
aatgtgggtga	ttaaataatgg	agagacagag	actaacagag	gtatgtgaat	attgaagtat	540
gtctggacaa	tagcccaatg	atgagaccaa	taaaatggtt	accaaactct	ggttttgagt	600
agtagtggtta	aatcagacca	tttagtaacc	atTTTTTgtt	gcaaagtttc	tagcactgcc	660
caaaccctga	gtggtatatg	aataactcgt	ccattatgta	tctctttcca	gtcagcataa	720
tttatccccc	acctatattc	ttttctgacc	actcctactt	ccttctcttt	accaaactct	780
aaactctaag	gctgtttctt	cagcaacttc	tttgtttaga	ttggaagata	aattaaacag	840
catgcatgtt	tttactgact	ttcagtattt	aacagagggt	atttaatttt	tttttaaatc	900
caaagtcaaa	cttctttata	agatgaagga	gaaaaatgtc	ttataaaatg	catatgtgaa	960
gatgcctttc	gagtgccttc	tcatgcagac	ttgttctagt	ctttaatgaa	tcttctttgt	1020
agacactgtg	gagatgaaag	atggttctcc	acttctactc	aaagtacaaa	tcaggccggc	1080
atTTTgaaaa	agagacaggt	ttattcatag	ctgcagcgtt	agctggcttt	gttccctgta	1140
caatttcact	tttggttatt	aaaatattca	ctgtaggaaa	taaatttgta	accattttct	1200
catattacct	acacacagaa	aaacaaaatt	tgatatcctg	gggtttatTT	gctgagggcg	1260
cttcccataa	aagcgagaga	gtgtgcgttg	ggaaatgtgt	ctggttaact	cttttatgga	1320
taaacttttag	tcacaatcct	cccccgcccc	cctctcacc	ccagcaccct	cccaacctcc	1380
cgacttcccc	cctctcaagg	gctggtgacc	taatagcatt	tttcttctgt	catatTTTgg	1440
cgtcgcccc	tggcctggct	gccttcgcct	gtctgagttt	tttgaattc	ctgcatgttc	1500
gccccagatt	aagccagtgt	gtctcaggat	gtgtgtttcc	ttttgttctt	tccccttaac	1560
gctccctgtg	caacgtgtct	ggggggagga	gggcagggac	gggagagagg	gaggggcaga	1620
ggcggaggagc	tgtccgcctt	gcacgtttcc	aatcgcatta	cgtgaacaaa	tagctgaggg	1680
gcgccggggc	cagaacggct	tgtgtaactt	tgcaaacgtg	ccagaaagtt	taaactctct	1740
ctccttccct	cactccagac	actgcccgtc	ctccgggact	gccgcgcggc	tccccgttgc	1800
cttccaggac	tgagaaaggg	gaaagggaa	ggtgccacgt	ccgagcagcc	gccttgactg	1860
gggaagggtc	tgaatcccac	ccttggcatt	gcttggtgga	gactgagata	cccgtgctcc	1920
gctcgcctcc	ttggttgaag	atTTTctcct	ccctcacgtg	atTTTgagccc	cgTTTTtatt	1980
ttctgtgagc	cacgtcctcc	tcgagcgggg	tcaatctggc	aaaaggagtg	atgctgcttc	2040
cctggaccgt	gctcctgctc	gggccttttg	agctctgctc	gctagtgcac	tgcgccccct	2100
ccgcccggcg	ccaacagcag	ccccgcgcg	agccgcgggc	ggctccgggc	gcctggcgcc	2160
agcagatcca	atgggagaac	aacggggcagg	tgttcagctt	gctgagcctg	ggctcacagt	2220
accagcctca	gcgccgccgg	gacccggggc	ccgccgtccc	tggtgcagcc	aacgcctccg	2280
cccagcagcc	ccgcaactcc	atcctgctga	tccgcgacaa	ccgcaccgcc	gcggcgcgaa	2340
cgcggaacggc	cggtcatctc	ggagtcaccg	ctggccgccc	caggcccacc	gcccgtcact	2400
ggttccaagc	tggctactcg	acatctagag	cccgcgaacc	tggcgcctcg	cgcgcggaga	2460
accagacagc	gcccggagaa	gttctctgct	tcagtaacct	gcggccgccc	agccgcgtgg	2520
acggcatggt	gggcgacgac	ccttacaacc	cctacaagta	ctctgacgac	aacccttatt	2580
acaactacta	cgatacttat	gaaaggccca	gacctggggg	caggtaccgg	cccggatacg	2640
gcactggcta	cttccagtac	g				2661

<210> 2

<211> 1953

<212> DNA

<213> Homo sapiens

EP 2 305 807 A1

	<400> 2						
	tataaattcc	acgcaggcat	tgaattgaat	ttgttcttaa	ccaaatgcgt	tttatctata	60
	cctggcagga	atctagaagt	gaaattacaa	gatttatttc	attttaattc	tattatgaag	120
	catttaatca	caaataccct	gaaaatgaaa	agataattta	tcattttacc	ttgactgagc	180
5	aactctcctc	acttcacatt	catgaatcca	taacgcagag	aggagactgg	atgattaagt	240
	gtttgattag	agaaaacaga	ttaacctagc	aaacataata	aatttggctc	ataagcagga	300
	tggctttata	aatgctcaca	atacctctcc	tgtataaaaat	catgaaccac	ttcctacagt	360
	gatgactcca	tcgaaatagt	tgagaaacat	aaagcaaatg	catgtttatg	gctttctctt	420
	tgagacatta	aaaggtatt	gaaaggcata	tctgattcag	cttataactc	tgatataata	480
	ttaaggaaca	tgtaagaaaa	tattaatgca	taaaaaaagc	tacaacttct	caagtgttct	540
	agtttccact	ttgtcaataa	ttacgttttc	aatgtccttc	tgtggactgt	ttccaaaggt	600
10	gccaatccag	acccaaagtt	tcagatcact	cagattcacc	cttaaccttc	ataacacaac	660
	ccaatagcct	tacgaaaaaa	gttgcatatt	taggtagtgt	ttatcccatt	atgacaaaaat	720
	acataaaatt	agcgagatat	tttttagcct	tcaaataagt	gggaaaaaat	ccttttagct	780
	gagattccat	ttacatcaga	ataaaaaatct	aagttatgac	taggttgaag	caacgtcctg	840
	tgcagcgctc	cataaagttc	acttagtctt	caagggttcc	ttacttagct	aggttagtat	900
	tcctggcctc	tttttttagc	agtgagaaaa	aggatactct	ccctgcccc	gctttatttt	960
15	taaactcaca	gccatattct	ggaggtctct	gctggctatt	tggcgcgtgg	gggaggaggg	1020
	gggcccgggg	agggggcg	ggcggggtct	ggaggtctgt	gctggctatc	tggcgtgtgt	1080
	gtgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	aggtctctgc	tggctatctg	gcgtgtgtgt	1140
	gtgtgtgtgt	tgggtgtgt	aagcagtgag	gtgttttag	ggccagtcct	tcctccgcca	1200
	ctttgctgac	tcaaagaccc	agaggctttc	ttgggtgca	ggtaccatga	ttccttgggc	1260
	cctaagggaa	ttttgttag	gctagaagag	tgggtgact	catgatgggt	gtacccgaac	1320
20	attcctgggc	tcaacaaaac	cgattatctt	tataaccgcg	gcgcctagca	cagcgcctgg	1380
	tgccctaaac	gttggctgcg	ggaacgtccg	agacgcgggt	gcggagccgg	gggaggaaata	1440
	actggttgcg	cggcgtttg	accgtaggcg	ctggagcgcg	tgcgttgcgt	gcgcgcgcgg	1500
	aggcggctgc	gtcggggcgc	gagaaggtgc	agttccccgg	cgggcgggcg	ggcgggaggg	1560
	cgaagctggg	ctcggggcca	agcgaggtct	agccggagcg	actgtgcccc	gcctcctggg	1620
	cggagcgggc	ggctccccat	ggtcagagcc	tcgtgcccgc	tcggcagcgc	ccggacgccc	1680
	agcccagcgc	gtcggcccc	cgggtgccc	gcgtctcaga	gccgcggagg	ggcggcggg	1740
25	accgtttcag	cgtggcggcg	ctggtgctgg	cgttggccct	ggaggacggc	cccagtgat	1800
	ggctggcgcc	tgccctcccg	gtgtctcccc	ggtacagatg	gagtcgtccc	gcggccgccc	1860
	gcggcaaggt	cggcagctgc	gaggccaaga	gagaccccag	gacacacaca	gctgcctccc	1920
	ggtgcgagaa	gaagaccccg	gcttgagagt	gag			1953
	<210> 3						
30	<211> 889						
	<212> DNA						
	<213> Homo sapiens						
	<400> 3						
	cggccccatg	gctccgtgtc	gtgtccaagg	gatgggctgg	cacctcttgg	accaggctta	60
	ccaccagggc	ccttctctga	agccccagtc	tgaccggcct	gctgctggga	atccccctct	120
35	gccccacac	taacctctgc	tggggctgag	ccaggggcgcg	tcggacagtc	agggcgacc	180
	agccagggcg	accgttggcc	ccgctcctat	ggggcagcag	ggaccgacgt	cagcaggggtg	240
	gggaggggcac	ccgagtggta	tgccccgccc	tgccccgctt	gcccgcctctg	gtggccgtct	300
	ggggggcgaca	agtccctgaga	gaaccagacg	gaagcgcgct	gggactgaca	cgtggacttg	360
	ggcgggtgctg	cccgggtggg	tcagcctggg	ctgggagcca	gccccgggac	acagctgtgc	420
	ccacgcgctg	tgagcacc	aagcccgatg	cagccacccc	cagacgaggc	ccgagggac	480
40	atggccgggg	acaccagtg	gtccaggtgt	ggcgggggtg	aggggagggg	gggtgggagc	540
	ggtggagatg	gggcccgtgg	gagggagctg	agatactgcc	acgtgggacg	atgctaggtg	600
	gggagggctg	agctgggagg	gctcctctgg	ctgtggggcc	ccctgtgttc	cttgtgggag	660
	gtggaaggaa	gtgagtgcc	tgtccttctt	ccctgccaatg	agattccagg	accggacctg	720
	gcaagtgtccc	tatcccagcc	agtgttctctg	gggctcttcc	aggcagggct	atgttcccca	780
	ggccaggggc	attgtcctgg	acagtcagga	ggcatacccc	tcgccaggtg	gaaccaccct	840
45	gtgtatgcat	gaccctgaca	agcaggcgcc	aggacagtca	ggaggccag		889
	<210> 4						
	<211> 863						
	<212> DNA						
	<213> Homo sapiens						
50	<400> 4						
	gttgttgggt	gtgaatggag	aactgtgggc	cctccccgac	accttccagc	gggacggcaa	60
	cggggggcca	gggggtgggc	gccatcaacc	ccgtcccacc	gccaggacgg	cgcgggggag	120
	ggccggcggg	ggcggggcgt	cctgtaagcc	gcgcccccca	cccgcgggcg	gggaggcatt	180
	cctggggaggc	ggcgctctg	acgtggaccc	ggggggcgcg	ggcacggcgg	ggggggcggc	240
	gtccgggggc	ttcttaaacc	ccccgccccg	gcccagcccc	cacttcccga	gcaccgctcc	300
55	gaccctggag	ggagagagag	ccagagagcg	gcccagcggc	taggagggcc	gcccagcctc	360
	gccgagcccc	gccagccccg	gcgcgagaga	agttggagag	gagagcagcg	cagcgcagcg	420

EP 2 305 807 A1

5
 agtcccgtgg tcgcgcccca acagcgcccc acagccccg atagcccaaa ccgcgccct 480
 agccccggcc gcacccccag cccgcgccag catgatgaac aacagcggct actcagacgc 540
 cggcctcggc ctgggcgatg agacagacga gatgccgtcc acggagaagg acctggcgga 600
 ggacgcgccg tggaaagaaga tccagcagaa cacattcacg cgctggtgca atgagcacct 660
 caagtgcgtg ggcaagcgcc tgaccgacct gcagcgcgac ctcagcgacg ggctccggct 720
 catcgcgtg ctcgaggtgc tcagccagaa gcgcatgtac cgcaagttcc atccgcgcc 780
 caacttccgc caaatgaagc tggagaacgt gtccgtggcc ctcgagttcc tcgagcgcga 840
 gcacatcaag ctcgtgtcca tag 863

10
 <210> 5
 <211> 2198
 <212> DNA
 <213> Homo sapiens

15
 <400> 5
 aagagaggca cactccctct accacaccga gggagggggc gttgagctga gaaaggttga 60
 gagaatgagg gaccaggtgta ggtggacatc ggccaagaaa ggaaccacag cgggaggtaa 120
 gaccgagagt ccccagcttg aagcgtcacc actccgggat tcccagattc caacgcgagc 180
 ctggggaaag cccacagtgg agagagtccg gctggcaggg aatggcccta ccccgggggt 240
 gaaatctcgg aggtcctgctc agccgagtcg cgcctctgctg ctgatgcgtg agagatgccg 300
 gacgtcgcgt ttgcctgtgc gaggcctcgcg gatgtgtgc agtcttggtc ccctctgct 360
 gtgtctaacc ccgaatgctg gtgtctcgag gtgtgagctt cggggccgggt gtctttaaag 420
 aaccaaagat tcttaaggag tgatgatctg ggtagagcgg cccgacgtag ccgctccc 480
 aggtctcggg gcgagtcctg cggacagacc agaggagacc tgctggccag atgccccggg 540
 cccaaggcgg acgccagact gtctctgcgc cagccgggct ggcttcgga atggatcagg 600
 caccggggag gccggagtgg atctcagacc ctcaagccgg gaacaaacc gtcgatgcc 660
 gtgggctcgg agtccgcctc ctctctccc ccccaccct accctgctt ccgaaaggct 720
 tcttcgctgg tcagtagctg cgtgcccgct tgcctgaggc tgggtcagaa ttggcgggct 780
 ggtaacgacc ccgtgcacaa gcggctccca gtctctccag aaagggccga tgactaaggg 840
 gtgggggtgg gggcggaggg ctggaaggtg ttaggaaaga acgttagcgg cctatctgt 900
 cttcagcagc gccctctcat cttctagctc tgacgccgag cagagcagtt ggagctcggg 960
 actgggaact gctggaattc ctatcttagac ttctagacag tctagaaaca agaactttc 1020
 tttccctggg cctcagtttc cttgtctgta aatcaaaaag gcgggctcta ggtgtaggcc 1080
 ttcttttcag ttggtgattc tggattcctt tccttgatc cgtggggagg ggggtgcagc 1140
 aacagctccag ggcgttgccc gtcctgtgcc tcaagtacgt agtccccgtg cccgccccct 1200
 caacaccccc agcagcccg cccctaagc ccgcagagca gggagctgag tgggaggggc 1260
 agaggcgggg ccggttcca gtccctgctg gcggactaga gtggcgcggg ctgagcgtaa 1320
 aacctgggat agccactccc ctttttctt atccccgcc ccctgccatt ggctccccgg 1380
 agaggttgac atcaaaaggc cggctttata taagccagat ccgcagggga gtccgcagaa 1440
 gggttaaaca ggtctttggg cttcggcgac ctgcgccgag gcagaaaccg gtaagaagac 1500
 agtgggctgc gcgtctcatt ttcagccttg cccggactct cccaaagccg gcgcccagta 1560
 gtggctccag agcccaacag tggcccccg cagtctctgg ggcgcatgga gcggcgtaa 1620
 tagggctggc ggcgcaggcc agtagccgct ccaacatgaa cctcgtgggc agctacgcac 1680
 accatcacca ccataccac cgcaccctg cgcacccccat cctccacgaa cccttctct 1740
 tcggtccggc ctgcgcgtgt catcaggaaa ggccctactt ccagagctgg ctgctgagcc 1800
 cggctgacgc tgccccggac ttccctgcgg gcgggcccgc gcccgcgcc gctgcagccg 1860
 ccaccgccta tggctctgac gccaggcctg ggcagagccc cgggcggctg gaggcgctt 1920
 gcggccgtct tggccggcgg aaaggctcag gaccaaagaa ggagcggaga gccactgaga 1980
 gcattaacag gcatttcgcg gagttgcgag agtgcattcc caacgtgccg gccgacacca 2040
 agctctccaa gatcaagact ctgcgcctag ccaccagcta catgcctac ctgatggacg 2100
 tgctggccaa ggatgcacag tctggcgatc ccgaggcctt caaggctgaa ctcaagaagg 2160
 cggatggcgg ccgtgagagc aagcggaaaa gggagctg 2198

45
 <210> 6
 <211> 1945
 <212> DNA
 <213> Homo sapiens

50
 <400> 6
 ctggatgaca gactgagact ccgtctcaaa aaaaagctc catttgggag gccgaggagg 60
 gtggattacc tgaggtcagg agtttgagac cagcctggcc cacatagga aacccatct 120
 ctactaaaaa tacaaaaatt agtcagggtg ggtggctgac acctataatc ccagctactt 180
 gggaaagctga ggcagggaga atcacttgaa ccggggagggt ggaggttga gtgagctgag 240
 atcatgccac tgcactccag cctgggcgac aggggtgagat tctgtctcaa acaacaaat 300
 ttaaagctc cgaatcctcc aaaaatacca agattttctt gtcggtaact agagatgggt 360
 actgatgatt attttaata ggtgattttc aaagatgtga acgttatcca tggagattta 420
 agtctcaaaa aggaaaaaaa atgcatacct ttatactaaa acttcatcac cagtcaaatt 480
 tggatcatca ctaaattggc ttctacacct ctctcctaat ataaggtaact tgtgtaagtt 540
 tgcagttgtg agacacttat ttctcattt ttaatgtctt ctcagtaggg ccactgatat 600
 agtcactatt tgactgacca gaatggttgg cactggtgat tggctcataa agtgcctcgc 660

EP 2 305 807 A1

	atthagggg	ctcaattatc	aaaggtttaa	atcctagccc	aaaccattgc	tgtgatgggg	720
	gtaaataaat	gaaccactca	gcttcacttg	caaaaagcggg	atcaccaatag	ccgctttcgt	780
	catgaccag	cctaggtgag	athtagtact	taagtacact	gccaggcaca	caaggttaat	840
	ttaacaat	aacacatttg	tttctcatc	catcttcca	aaccttcaa	ctaatectaa	900
5	cgttcgttcg	gccaaatggg	ccaggaattc	acttaaacaa	aaacaaaaa	caaacaaac	960
	aaaaaacac	tccctggggc	ttggggaagg	aggcaccgcc	gcccattgctg	cagtctgggg	1020
	gtggctcagt	cctcagcacc	cagatctacg	gccataatgc	tcttcgaggc	caaggagccc	1080
	ggatgcgggg	cggtgcgaa	ggcgtcttgc	tcaggctgcg	ggaaaggaga	ggggtgggag	1140
	cggggtgggg	gcatcgcgac	ccagggcaag	gcggcagtc	gccgtcttcg	agtcccacct	1200
	gtccgaagcg	gggtgagaaa	aggcaaaaaca	tggcaaagcc	atgcacctcc	cagggtgggc	1260
	aactcacggc	cggtgaacgc	cggaccctta	gcagtttcca	gacctttgga	accggaagcg	1320
10	gagcctgaga	gcgcgcccga	gagggcgtga	acgggaccgc	tttcccggaa	gtgcttgccg	1380
	ccctgcccga	gcgagctgcc	ccggggcttc	tctggtttcc	taatcagggc	aacgcgcgcg	1440
	gagagaacct	ttaccttggc	tgcactaagt	tctcggtgcc	actccctggc	agggcgggac	1500
	cttgtttagg	ccctgtgatc	gcgcggttcg	tagtagcgca	aggcgcagag	tggaccttga	1560
	cccgcctagg	gcgggaagag	tttggcccgc	cgggtcccga	agggcagaat	ggacgggctc	1620
	ctaaatccca	gggaatcctc	taaattcatt	gcgaaaaaca	gtcgggatgt	gtttattgac	1680
15	agcggagcg	tacggaagg	ggcagagctg	ctcgtggcca	aggcggcggg	gccagagctg	1740
	cgctgagg	ggtggaagc	ccttcatgag	ctgaaccca	gggcccga	cgaggccgcg	1800
	gtcaactggg	tgttcgtgac	agacacgctc	aacttctcct	tttggctcga	gcaggacgag	1860
	cacaagtgtg	tggtgaggtg	cagagggaaa	acatacagtg	ggtactggtc	cctgtgcgcc	1920
	gccgtcaaca	gagccctcga	cgaag				1945
	<210>	7					
20	<211>	2379					
	<212>	DNA					
	<213>	Homo sapiens					
	<400>	7					
25	aagcttgtgg	tttacttggg	cctctgcctc	atctttcttc	ttttgcgctt	cagcctgcgc	60
	attcgcttcc	tccactaggg	tctcatgggtg	cagaggtttc	caagaagatg	gtgtgaaggc	120
	cgagatcatt	tggttatatt	ataaaataga	atgcaaattc	acacaagttt	ttgtttttta	180
	tttatttatt	tttttagaga	tgagggtcttg	ctatgttggt	tagtctggtc	tcgaactcct	240
	ggcctcgtga	tcctcccacc	ttgacctccc	aaagtgtctg	gattacaggc	ctgaggctg	300
	agccacttca	cccaactgaa	ttcacatttt	ttttttctt	ttctgagacg	gagtctcact	360
	ctgtcaccca	gtatggagtg	cagtggcgcg	actgcggctc	actgcaagct	ccgtctctcg	420
30	ggttcaagtg	attctcatgc	ctcagcccc	caagtagctg	gaattacagg	ggtgcactac	480
	cacacctggc	taatttttct	gttttagtag	agatggggtt	tcaccatggt	gcctggctc	540
	aaactcctga	ctttaagtga	tccacacacc	tcagcctccc	aaagtgtctg	gattacaggt	600
	gtgagcctcc	acaccggcc	gaattcacat	gaattttaa	gtgatgtctt	caaagtgggt	660
	tcactgtggg	gatgggcagc	ttttgttat	acatctagaa	cgttcctctt	ctgtttctat	720
	gaatactcgg	ttgaaagg	ctgaaaaacg	gtcttaagag	attatctgat	tcgtttccca	780
35	gttttattac	tcacatatca	gctgtaattt	gagcacgttt	tctgattgag	acaagactca	840
	gatggtatta	aacattacta	caacacatcc	gggcacgggtg	gctcacgcct	gtaatcccag	900
	cactttggga	ggccgaggcg	ggcggatcac	gaggtcagga	gatcgagacc	atcctggcta	960
	acacggtgaa	gccctgtctc	tactaaaaat	acaaaaaatt	aggcgggcat	ggtggcgggc	1020
	gcctgtagtc	ccagctactc	gggaggctga	ggcaggagaa	tggcgtgaac	ccgggaggcg	1080
	gagcttgcaag	tgagccgaga	ctcgcgccact	gcactccagc	ctgggcgaca	gagcaagact	1140
	ccatctcaaa	aaaaaaaaaa	aaaaaaaaaaa	actacaacac	tataaattca	tatctattat	1200
40	aatagtactt	tgtgcagggc	cctaccctaa	gtccttaacc	gaaccggaa	gcgagaagat	1260
	gacttttggt	tgtttttaga	gatgggcgcc	tggctctgtc	gccagcctgg	agtgtggggg	1320
	cgcgatctcg	actcacagca	gcctccacct	cccaggttca	ggcgatcttc	ctgcctcagc	1380
	ccctcgagga	gctgggacca	ccggcgcgct	ccatcgcgct	cggctaggag	ctgactttga	1440
	atccgggctc	tgcgctggc	cttctgcatc	tctataaggg	aagacatctg	tgacctcggg	1500
	gcaaagggtca	aattagatcc	tgggtagat	cctgttcccg	ctgcccctcg	ggctggcact	1560
45	gccaggagta	ctcagagctc	aaagctggga	tctgcagctc	cttaccact	cagtgcacgc	1620
	cgctaaggc	tttgcgcttc	acctttactc	acctcgaagc	cctggacatc	cgcatctgcc	1680
	ctaagacttc	tcacctcagt	agcagaagga	agtcgcgctc	gctggccaca	gcctctctcc	1740
	taggagaccg	tccgggaaaa	gcgagtcagg	gtagaccctg	aggcccctca	gctccggcta	1800
	ttttcagatc	tgctcgtcct	tcacctcag	cctttcaaac	aggccactcc	aaaaaaaaagc	1860
	ccaatcacag	ccttctctct	tctcctggcc	ttccggcact	gtccaatcaa	cgtacgccat	1920
50	ctatcggtta	gtggtgttgc	ggggccacc	ttcccgtgg	tttccctcgt	ggtgtgtaaa	1980
	ggcagagagg	aaaggcgagg	ggtgttgacg	ccaggaaggt	tccatcttgg	ttaagggcag	2040
	gagtccctta	cggacttgtc	tgaggaaaga	caggaaagcg	ccagcatctc	caccttcccc	2100
	ggaagcctcc	ctttgccagg	cagaaaaggg	ttcccattgg	gccgcccctg	gcgcccgcgc	2160
	cggccacgtg	acccggggag	gccgggcccc	ggaggacgag	ggaaagcagg	ccggggcggc	2220
	tgagcttcgc	ggacgtggcc	gtgtacttct	ctcccagga	gtgggaatgc	ctgcggccag	2280
	cgagagggc	cctgtaccgg	gacgtgatgc	gggagacctt	cgccacctg	ggcgcgctgg	2340
55	gtgaggccgg	gccctccggc	cgggaccccc	agtccgtcg			2379

EP 2 305 807 A1

<210> 8
 <211> 933
 <212> DNA
 <213> Homo sapiens

5
 <400> 8
 gagacgtact ctggctctgt cgcccaggct ggagcgcaat ggcgccatct cggcgcactg 60
 caacctccac ctcccgggtt caagcgattc tactgcctca gcctcccagag tagctgggac 120
 tacaggcgcg cactaccaag cccgggtaat ttcttttgta ttttttagtag agactggggtt 180
 tcacgatgtt ggccgggctg gtctggaagt cttgacctca agcgtgccc ctctccgcc 240
 ctgggtaagg cggggggcgg aatagggggc ttgcaatttc aactagagg cgggcgccgt 300
 10 ggggaaaga agagtcacgt ctcccacggt tcgtagagga agcctgcct gagcctggag 360
 cggggcggg agagccacag tttggcatcc ccagggcatc ccccagccc cagactacca 420
 ggctccaga ggacaggacc ccacccccgg ccacaggccc tgccccagc actccccgca 480
 ccccgctcc aagactcctc cgcccactcc gcacccaact tataaaaacc gtccctgggc 540
 gggcgggga gaagccgagc tgagcggatc ctcacacgac tgtgatccga ttctttccag 600
 cgacttctgc aaccaagcgg gtcttaccct cggctcctcg cgtctccagt cctcgacct 660
 15 ggaacccaa cgtccccgag agtccccgaa tccccgctcc caggctacct aagaggatga 720
 gcggtgctcc gacggccggg gcagccctga tgctctgccc cgccaccgcc gtgctactga 780
 gcgctcaggg cggaccgctg cagtccaagt cgccgcgctt tgcgtcctgg gacgagatga 840
 atgtcctggc gcacggactc ctgcccgtcg gccaggggct gcgcgaacac gcggagcgca 900
 cccgcagcca gctgagcgcg ctggagcggc gcc 933

20
 <210> 9
 <211> 6096
 <212> DNA
 <213> Homo sapiens

<400> 9
 25 atctgcacct cctcatatag ggttgatcca agtttcacag acatcactga gttcttagtg 60
 gactcagcta ttggggctgt tctcacactt ttttttctt tgcaagaatc agcaatgggt 120
 gcaagtggac ctgtgtagga cgtccagtga aacattgtgt tggatcaatc gctagaatcc 180
 atccaagaac tcagccagcc tgggtgtagg tgagatctga tccttgaatg tccctcagt 240
 gcttttaggg ctggcaggtt cagaagggcc ctctcatcac cccccaggg cctcattcct 300
 tgtttaacac tttgctatca cagtcttgaa tccttgtaat tgaacaatgg accccacatt 360
 30 ttacttttgc actggtttct gattctgtaa ccgatcctgt ccccctctt tgtctcattc 420
 actctgggaa ttgtccccac attctgagac ctttcagcag tgccccaac aggttcctgc 480
 ccttatctga agtccacc caccccccat ggcggcacc caggcagccc tgcttttgcg 540
 tcccgcgtag gcagctgtg caccggagtc acgaccccc gattcagcct aggcagccc 600
 agcttgactg ctcccggcgg acaagcccta ctgtgctatc tgccgctctt ccttcctct 660
 tcccaggggg tccgcgtcag gggaggcgca gctgtgtgca tccggggagc ttcagacccc 720
 cggtgcccagc agctccttcg tttcctgggt gctggggcgg ccttcccagc gaagagctca 780
 35 actcagcggg acgtttggag gctctctgcc ccaaggcgt ggggagtgt cggcgggaca 840
 gtcgtgcttg cctttttcac tttcagagt tccacgccc acccgtttg tccactgagg 900
 tcagtccagt ccagcccggc ccaccccacc ggtgctgtc tgtcgcactg ggcagacgcc 960
 atactctctg ttcttgttta aagcccagga tctactgggc cctggaggca agaggtgaac 1020
 gcagcggaa ccacgctgag ctgcccggga acggagctc caaccccaga aggaggactc 1080
 tgtgtccta cacctaac ctitttagcc cgaacttct ccaacttct tggctttgtt 1140
 tagagctcga cagcgcggcc ccctggcgt cgttgtgagg acagtagagg agagaggcaa 1200
 40 ggggtgtttt aaacagttt cctctacca ttatggggg gacccgagg ggagaccac 1260
 tcttccgcat tcccggtaag tgaaccacc gaagaggtc aaagtgacg attcccattg 1320
 cctcctccag cccccccc accctgccc tccacaggac ggtggctctt cagtgccctt 1380
 tgccgagcaa gtggcgttt tatgcacgt ggtatcaatt cggactctgg acgaaatgga 1440
 aacctcctta gccgacccgg gtgggatcag ctgggatcct gcgcgctccc ctgggggggt 1500
 45 gccagccact ctgttggggt gcaagaagca ccacctctc gaagctgggc cgaactggc 1560
 caggctgact cgctcccacg cgcccggccc taccggcgc cctgacagt tccccgcgac tccccacc 1620
 cgctctgag cgggtccgg acttgcagtg cctgacagt tccccgcgac tccccacc 1680
 gatgagatgg ggtctggcgt tggccagtgc gtgtccagg actcgcgggt ccttggccag 1740
 ccatggggca gagggcgctg gtgttaggccc agtcttccc accctgccc gtcaccccag 1800
 ccacaccac tgtcctgtga ggccaagcgc gctccgctgg tttcctgagc caggcacctt 1860
 ggccgaggac aggatccagc tgtctctct tgcgatcctg tcttgggga agtccagtc 1920
 ctaggcagg cctcccaaag tgcccttggg gccgatcacc cctcccagc tcttgcagg 1980
 50 cctgtgcacc acctccccc ctccccattc aaagccctt tctctgaagt ctccggttcc 2040
 cagagctctt gcaatccagg ctttcttgg aagtggctgt aacatgtatg aaaagaaaga 2100
 aaggaggacc aagagatgaa agagggctgc acgcgtggg gcccgagtgg tgggagggga 2160
 cagtactctt gttacagggg tgcctggcct ccttggcgc tggcccctgtc ggcccggccc 2220
 gagaacctcc ctgcccagg gcagggttta ctcatcccgg cgaggatgac ccatgcccga 2280
 55 gggcggggc aagggcgcc agagaaccca gcaatccgag tatgcccag cagcccttcc 2340
 caccaggcac ttccttctt tcccgaacg tccagggagg gagggccggg cacttataaa 2400
 ctcgagccct ggccgatccg catgtcagag gctgcctcgc aggggctgag cgcagcggca 2460

EP 2 305 807 A1

	agaagtgtct	gggctgggac	ggacaggaga	ggctgtcgcc	atcggcgctcc	tgtgcccctc	2520
	tgctccggca	cggccctgtc	gcagtgcccc	cgctttcccc	ggcgcctgca	cgcggcgccg	2580
	ctgggtaaca	tgcttggggt	cctgggtcctt	ggcgcgctgg	ccctggccgg	cttgggggtc	2640
	cccgcaccgg	cagagccgca	gccgggtggc	agccagtgcg	tcgagcacga	ctgcttcgcg	2700
5	ctctacccgg	gccccgcgac	cttctctcaat	gccagtcaga	tctgcgacgg	actgcgggggc	2760
	cacctaataga	cagtgcgctc	ctcgggtggct	gccgatgtca	tttctttgct	actgaacggc	2820
	gacggcgggc	ttggccgccc	gcgccctctgg	atcggcctgc	agctgccacc	cggctgcggc	2880
	gaccccaagc	gcctcggggc	cctgcgcggc	ttccagtggg	ttacgggaga	caacaacacc	2940
	agctatagca	ggtgggcacg	gctcgacctc	aatggggctc	ccctctgagg	cccgttgtgc	3000
	gtcgtgtctt	ccgctgctga	ggccactgtg	cccagcgagc	cgatctggga	ggagcagcag	3060
	tgcaagtga	aggccgatgg	cttctctctgc	gagttccact	tcccagccac	ctgcaggcca	3120
10	ctggctgtgg	agcccggcgc	cgcggtctgc	cccgtctcga	tcacctacgg	caccccgttc	3180
	gcgcccgccg	gagcggactt	ccaggcgctg	ccggtgggca	gctccgcccg	ggtggctccc	3240
	ctcggcttac	agctaatgtg	caccgcgccg	cccggagcgg	tccaggggca	ctgggcccagg	3300
	gaggcgccgg	gcgcttggga	ctgcagcgtg	gagaacggcg	gctgcgagca	cgcgtgcaat	3360
	gcatcccttg	gggtcccccg	ctgccagtgc	ccagccggcg	ccgccctgca	ggcagacggg	3420
	cgctctctga	ccgcatccgc	gacgcagctc	tgcaacgacc	tctgcgagca	cttctgctgt	3480
15	cccaaccctg	accagccggg	ctcctactcg	tgcagtgtcg	agaccggcta	ccggctggcg	3540
	gcccaccaac	accggtgcga	ggacgtggat	gactgcatac	tggagcccag	tccgtgtccg	3600
	cagcgtgtg	tcaaacacaca	gggtggcttc	gagtgccact	gctaccctaa	ctacgacctg	3660
	gtggacggcg	agtgtgtgga	gcccgtggac	ccgtgttca	gagccaactg	cgagtaccag	3720
	tgccagcccc	tgaaccacaac	tagctacctc	tgcgtctgcg	ccgagggctt	cgcgccattt	3780
	ccccacgagc	cgcacaggtg	ccagatgttt	tgcaaccaga	ctgcctgtcc	agccgactgc	3840
20	gaccccaaca	cccaggctag	ctgtgagtg	cctgaaggct	acatccttga	cgacggtttc	3900
	atctgcacgg	acatcgacga	gtgcgaaaac	ggcgcttct	gctccggggg	gtgccacaac	3960
	ctcccgggta	cccttcgagtg	catctgcggg	cccagctcgg	cccttgcccc	ccacattggc	4020
	accgactgtg	actccggcaa	ggtggacggg	ggcgacagcg	gctctggcga	gcccccgccc	4080
	agcccgacgc	ccggctccac	cttgactcct	ccggccgtgg	ggctcgtgca	ttcgggcttg	4140
	ctcataggca	tctccatcgc	gagcctgtgc	ctgggtgggg	cgcttttggc	gctcctctgc	4200
25	cacctgcgca	agaagcaggg	cgccgcccagg	gccaagatgg	agtacaagtg	gctgcccctt	4260
	tccaaggagg	tagtgcctgca	gcacgtgccc	accgagcggg	cgccgcagag	actctgagcg	4320
	gcctccgtcc	aggagcctgg	ctccgtccag	gagcctgtgc	ctcctcacc	ccagctttgc	4380
	taccaaaagca	cccttagctgg	cattacagct	ggagaagacc	ctccccgcac	cccccaagct	4440
	gttttctctt	attccatggc	taactggcga	gggggtgatt	agagggagga	gaatgagcct	4500
	cggcctcttc	cgtagcttca	ctggaccact	gggcaatgat	ggcaattttg	taacgaagac	4560
30	acagactgcg	atttgtccca	ggtcctcact	accgggcgca	ggagggtgag	cgttattggg	4620
	cggcagcctt	ctgggcagac	cttgacctcg	tgggctaggg	atgactaaaa	tatttatttt	4680
	ttttaagtat	ttaggttttt	gtttgttttc	tttgttctta	cctgtatgtc	tccagtatcc	4740
	actttgcaca	gcttccgggt	ctctctctct	ctacaactc	ccacttgtca	tgtgacaggt	4800
	aaactatctt	ggtgaatttt	ttttctctag	ccctctcaca	tttatgaagc	aagccccact	4860
	tattcccat	tcttctctag	tttctctctc	caggaactgg	gccaactcac	ctgagtcacc	4920
	ctacctgtgc	ctgaccctac	ttcttttggc	cttagctgtc	tgctcagaca	gaacccttac	4980
35	atgaaacaga	aacaaaaaca	ctaaaaataa	aaatggccat	ttgctttttc	accagatttg	5040
	ctaatttatc	ctgaaatttc	agattcccag	agcaaaaata	ttttaaacaa	aggttgagat	5100
	gtaaaaggta	ttaaattgat	gttgctggac	tgtcatagaa	attacacca	aagaggtatt	5160
	tatctttact	tttaaacagt	gagcctgaat	tttgttgctg	ttttgatttg	tactgaaaaa	5220
	tggtaatgtg	tgctaattct	cttatgcaat	ttcctttttt	gttattatta	cttatttttg	5280
	acagtgttga	aaatgttcag	aaggttgctc	tagattgaga	gaagagacaa	acacctcca	5340
40	ggagacagtt	caagaaagct	tcaaaactga	tgattcatgc	caattagcaa	ttgactgtca	5400
	ctgttccttg	tcaactgtag	acaaaaataa	aaccagctct	actggtcttg	tggaattggg	5460
	agcttgggaa	tggatccctg	aggatgcccc	attagggcct	agccttaatc	aggtcctcag	5520
	agaatttcta	ccatctcaga	gaggcctttt	ggaatgtggc	ccctgaacaa	gaattggaag	5580
	ctgcctctgc	ggttagaagt	ggtagaatct	cgagaatcct	aggctccacc	ccatccagtt	5640
	catgagaatc	tatatttaac	aagatctgca	gggggtgtgt	ctgctcagta	atctgaggac	5700
	aaccattcca	gactgcttcc	aattttctgg	aatacatgaa	atatagatca	gttataagta	5760
45	gcaggccaag	tcaggccctt	attttcaaga	aactgaggaa	ttttctttgt	gtagctttgc	5820
	ictttggtag	aaaaggctag	gtacacagct	ctagacactg	ccacacaggg	tctgcaaggt	5880
	ctttggttca	gctaagctag	gaatgaaatc	ctgcttcagt	gtatggaaat	aaatgtatca	5940
	tagaaatgta	acttttghta	gacaaaggtt	ttcctcttct	attttgtaa	ctcaaaatat	6000
	ttgtacatag	ttatattttt	attggagata	atctagaaca	caggcaaaat	ccttgcttat	6060
	gacatcactt	gtacaaaata	aacaaataac	aatgtg			6096
50	<210>	10					
	<211>	2500					
	<212>	DNA					
	<213>	Homo sapiens					
55	<400>	10					
	accacttct	gtgtgtggat	agtatcctgc	aggagagatg	ttgtctgcag	tgtgagctgg	60
	gccaccgga	gtgtgtgaat	aggatcctgc	aggagaaatg	gaatccggag	tgtgagctgc	120

	atccgctgta	gaggggtggat	aaaatcctgc	aggaaagatg	gcattctggaa	tgctcagcggg	180
	agccaccgac	ctctgaggat	gcacccccga	gggtgtgatgc	ggggccagtt	ccaaggctgg	240
	gtaggtttt	accctggctt	ctgtgttga	ctctcattct	cttcctcttt	cttctaatac	300
	ctgctctggg	agggcatcagg	ccatgtccag	tgtgcaggcc	atggagacc	acacggcaag	360
5	gaactggaac	ccccgtccag	cagcctcggg	ggctccagtc	ttagatgggtg	ccctgtggtc	420
	agcaatgcac	ctgtgacctc	cgggctatgt	ctcgtggtag	ttgcttttgt	gttttaacat	480
	agcaacagga	aactagccta	ttaccacca	atcccattcc	aggctgcttt	caaacgcagc	540
	tcaggctaga	acaccagcac	ggggacacag	ctgagacttg	gggtttgca	cgggaacag	600
	cccatgctgt	gcctctgaat	ctggcaccgt	cacctgtgg	cctgggttca	gcaactggc	660
	ctcaccttcc	ttgtctgtga	aattcagact	gggtccttgt	gagatgattg	gagagaatgt	720
10	atgaactatg	tgagaacgcc	acctttgtgc	gtatctcacg	cagtgtcttc	cctccttcc	780
	aaagtcttct	gctgtctcta	gacacaccgg	acgtgggggg	gggggggttc	gtgggtctcc	840
	tcctaggtct	gtcccaggag	ggcacgcact	gaaggccgcg	agaatcccgg	gggctgcatt	900
	gcgccgcgcc	aaggactcca	cacaggacct	ttcattttcc	caactgtgct	gagccaggcg	960
	gccggcagag	agcaggtggc	tgacaggccc	cggggagccg	gaccgcctgg	gtctaattct	1020
	cccgcagact	cccttgctgt	gcgctttggg	gcttgggctt	cagtttcttc	aaaaggaatg	1080
	aggggctttt	ttggaacggt	aaataatttc	ctacgtggtt	cgcggtaggg	agaaggagaa	1140
15	agagaggagc	gcgcctgagc	gcctggaatc	gtgccggat	cagagcaagc	gctctaaaag	1200
	tgttacaaac	attaagggcg	caactaaaaa	accgtagtg	agcgcaggca	gaaaccacgg	1260
	gtaagagaag	tggaagaagc	tcgcgtaggc	cccagggctc	cgagccccga	gtctcagcgc	1320
	cagaatcagc	ggtgccaatg	ctctcctccg	cgccccgag	cgctcgcctt	ggcctagcgg	1380
	gcccggccac	cgggatgagg	gcgctcaggc	cggacgctgg	ggccccgggt	tctcgccccg	1440
	ccccgcccct	ggggattcag	aggggcccgg	aggagcctcg	cgcatgtgca	cagctggcgc	1500
20	ccccgcccc	ccgcgcacag	ctgggacgtg	ggccgcggcc	gggcgggccc	agtcgggagc	1560
	cggccgtggt	ggctccgtgc	gtccgagcgt	ccgtccgcgc	cgctcggccat	ggccaagcgc	1620
	tccaggggcc	ccgggcgccc	ctgcctgttg	gcgctcgtgc	tgttctgctc	ctgggggacg	1680
	ctggccgtgg	tgcccagaaa	gcccggcgca	gggtgtccga	gccgctgcct	gtgcttccgc	1740
	accaccgtgc	gctgcatgca	tctgctgctg	gaggccgtgc	ccgccgtggc	gccgcagacc	1800
	tccatcctgt	gagtgcggcg	ggggacgccc	ggggcgcggg	gtcccggggct	tcgtggagat	1860
	ccgggagcgc	aggggtgatc	ggaggtgggg	ggcgcggagg	gtggaggggg	gtgggggacg	1920
25	gcggggggcc	tggggacttg	ggagcagaaa	gggaacctcc	gaagggggac	gtggggggac	1980
	ctgggcgccc	ggacccgctg	ggcctttggt	cgccctgcgg	gagacgccga	ggggcggaac	2040
	agagcgctgt	gcgcgcggcc	ttcgtagccc	cctttgttcg	gaactcggaa	tccccgagc	2100
	actgggaagt	tgttggagcc	tccggggctc	ccccgcctcg	cctcccgcgg	ccccctctca	2160
	tgctccgccc	gcctcccgtc	tccccctggt	tcgcggcccc	tctctcgcct	accttcccc	2220
	cgctcaggac	ccctcggctc	ccctcgcctc	cccagcgcgc	gcgcagcccc	ctccgtctct	2280
30	ccagccccct	ccgcccgtt	cctcgtcctg	ttcgtcctcc	tctcgcctc	ctctctctcc	2340
	tcccccttct	ctctctctc	cccttctctc	tctctctccc	cttctctctc	ctctctctct	2400
	ccccctctcc	tcccccttcc	ccttctctcc	ccccagcctc	cgccctctcc	ccctcccccg	2460
	ccctttggag	cgcagtgc	acccatccc	cccgcgcccg			2500

<210> 11
 <211> 2200
 <212> DNA
 <213> Homo sapiens

	<400> 11						
	cctcggcccc	tccagccggc	ccccggggcc	cctcctctcg	gcgcccggac	cttggccctc	60
	ctctctcttt	cccacttctc	tctttgccc	aacttcgccc	ccatcccccg	ctcatttctt	120
40	ctcgcacccg	ggctcgccaa	tccctctttc	caagtccttc	ttccagcccc	gccttctctt	180
	cgggttcgcc	ccccttctcc	ccaatctccg	tcctcttccc	tcccttcgcc	ctccccctt	240
	tccttctctt	tcccctcacc	caaccctggt	tccccctggt	cctcagtcct	gatctctccc	300
	ttactctgtc	cccgccact	ctgcgcccgc	ctctcagtc	gggttgagcc	ccacgtgtgg	360
	acggccgcgc	ccccactgac	agccgcccgc	cgccggcccc	ccccgcgccc	cgccgggctt	420
	ctaaaacccc	cgcgccgcgc	cctccaccgc	cgcatcttct	ccagcgcca	gcctcccgcc	480
45	ctctctcttt	ctggccgcac	gccccggccc	cgcgaccctc	cgcccggtc	cgcagccgct	540
	acccgcgctt	cgttgccctg	tgggactccg	agcgagcccc	gagggaaacc	tcctcttctt	600
	ctggggggca	ctttgtttg	cttgccctgtt	tctttctggt	gacttttgca	gctttccaat	660
	atccgtcttc	ggagcgcacg	ggaatccgcc	gagctctgcg	tgaggccct	ttttctttt	720
	gaggttcaca	tttttgaaa	ttttacgcca	gggcttttgt	aatttctctc	cccggccgct	780
	gacggctcctg	gagtcgctcg	gggctttagg	ccggttatgc	aacgtgtacc	gctcggggct	840
50	gccggctgca	cctccgcccgc	gcctcgcgcg	tcactgcctc	agacccggcg	ccccgcgtct	900
	cgcttcgccc	gcagtcaggg	ggccggcgct	ctgtcgaggt	ctccagctag	agcagggagc	960
	ccgagcccga	gggagtcctc	ggagccgacg	aagggttat	tagaccctga	ctctttctg	1020
	aggcgcgcag	atthtgcctt	tgatcactcc	ctctccgccc	gtctacggcc	gcgcgctttc	1080
	ggcggggccc	atggggagaa	gacggaggct	gtgtctccag	ctctacttcc	tgtgtgctgg	1140
	ctgtgtgggtg	ctctgggcgc	agggcacggc	cgccagcctt	cagcctctct	cgcccaagcc	1200
	gccccggccc	cagccgcccgc	cgcaacaggt	tcggctccgct	acagcaggct	ctgaaggcgg	1260
55	gtttctagcg	cccagtatc	gcgaggaggg	tgccgcagtg	gccagccgcg	tccgcccggc	1320
	aggacagcag	gacgtgctcc	gagggtaagt	gggcaagcgg	ctccgcacct	agggctccgg	1380

EP 2 305 807 A1

	cttgggggag	gggggaatcc	tcagtttggc	ggctttctgg	cccactccgt	cccagaccct	1440
	ttagctggag	cctagagctg	cagccccctt	tgccagaata	tccaaagacc	cccaggagcg	1500
	cgccccctt	ttccttccca	acccccgagc	tcagcgggcg	gaaagccctc	tctccggggg	1560
	ttggcggcg	ggtggttagg	gggtccaggg	gtgccgatcg	cagagcgtgt	gcagagctcg	1620
5	cgctgcggga	acaggttctg	aatgtccggc	ggcaggcggg	cctgggtccg	cctgctgcag	1680
	gggccagaga	agcctgcttg	ctccccacgt	cggggccgcc	gctcgtgagc	ctttgtttg	1740
	aggacgtgtg	cagggttcac	agctcacctt	ctcatcgta	acccgagcgc	tccaccttgc	1800
	gacgcgcttt	ccttgacacg	tcggggccaa	agtaacagtt	gaccaaggag	gaatggattt	1860
	gggaaggagg	gcaaggattc	tttggaacgg	aatgggtccct	ttgttctctg	catctggaag	1920
	ctagaatagt	agcaattat	atgtttccat	gcctcttttc	gccctttaa	aaggcaggca	1980
	agggacgaca	gatgaaaggc	agtgtttaga	catttctgac	cctcctgcat	tccagcatct	2040
10	agctcttttg	cttccacgtc	tgcttccccg	tctccaataa	tttgaagtgt	aattttgatt	2100
	tgtttgttgt	cctgaaatct	actcgtcgg	ggcattgctt	acgaagaccg	tttataatgtt	2160
	gtgtcatccc	tctacctatc	tgttacgtga	ccgcgcttgt			2200

<210> 12

<211> 2000

<212> DNA

<213> Homo sapiens

<400> 12

	ttggaagaaa	aggatctccg	aggaaggggc	tgagagaagg	gcaggggtgaa	ctggactaaa	60
	ggccagagta	ggaaggagaa	gagggggccaa	aaaagaaggg	gatgaaatta	agcacagaag	120
	atgggtaaa	aaaaaagtat	caggggaaagg	gcaaaataag	agaaagcctt	gaggataaga	180
	gggtagaagg	ctaaagaaca	agggggaccac	tgggtcgggg	aagcgctgcc	tgaacggcgg	240
	gacagtgaca	aagaaagggc	gctggcgata	ttcgacccaa	gggtgcgaaa	cgcaatcggg	300
	aggtgagaaa	tggaagaag	gcgaatgcc	ggctacaagt	agcctgggac	tgaaggggga	360
	cctgggggag	gggctgggcc	cagggcagaa	aagtccaggt	tcccatgagg	cctgggcccc	420
	cgtggagcgg	gcgctgaatc	accgttcagc	cgccccctc	ccctcctccc	cgaccggtgc	480
	ccgcagctcc	cgctcctcg	gccgcccgtc	ccacggggcg	gggccctggc	ccgggaccag	540
	cgcccggtct	ataaatgggc	tgccggcagg	ccggcagaac	gctgtgacag	ccacacgccc	600
	caaggcctcc	aagatgagct	acacgttgga	ctcgtgggc	aaccctgccc	cctaccggcg	660
	ggtaaccgag	accgctcga	gcttcagccg	cgtagcggc	tccccgtcca	gtggcttccg	720
	ctcgcagctc	tggtcccggc	gctcggccc	caccgtgtcc	tcctcctata	agcgcagcat	780
	gctcgcctcc	gcctcgtctt	acagctcggc	catgctcagc	tccgcccaga	gcagccttga	840
	cttcagccag	tcctcgtccc	tgctcaacgg	cggctccgga	cccggcggcg	actacaagct	900
	gtcccgtctc	aacgagaagg	agcagctgca	ggggctgaac	gaccgctttg	ccggctacat	960
	agagaaggtg	cactacctgg	agcagcagaa	taaggagatt	gaggcggaga	tccaggcgct	1020
	gcggcagaag	caggcctcgc	acgcccagct	gggcgacgag	tacgaccagg	agatccgga	1080
	gctgcgcgcc	accctggaga	tggtgaacca	cgagaaggct	caggtgcagc	tggactcggg	1140
	ccacctggag	gaagacatcc	accggctcaa	ggagcgcttt	gaggaggagg	cgcggttgcg	1200
	cgacgacact	gaggcggcca	tccgcgcgct	gcgcaaagac	atcgaggagg	cgtagctggt	1260
	caaggtggag	ctggacaaga	agggtgcagtc	gctgcaggat	gaggtggcct	tcctgcggag	1320
	caaccacgag	gaggaggtgg	ccgaccttct	ggcccagatc	caggcatcgc	acatcacggt	1380
	ggagcgcaaa	gactacctga	agacagacat	ctgcagggcg	ctgaaggaaa	tccgctccca	1440
	gctcgaaagc	cactcagacc	agaatatgca	ccaggccgaa	gagtggttca	aatgccgcta	1500
	cgccaagctc	accgagcggg	ccgagcagaa	caaggaggcc	atccgctccg	ccaaggaaga	1560
	gatcgcggag	taccggcgcc	agctgcagtc	caagagcatc	gagctagagt	cggtgcgagg	1620
	caccaaggag	tccttggagc	ggcagctcag	gcacatcgag	gagcggcaca	accacgacct	1680
	cagcagctac	caggtaggaa	ccgcggctgc	gcgccagcc	tgccgagcgg	ccagcggcgc	1740
	gcgcccccca	cacttgggct	cgtagcccagg	cgccctctcc	gccgcgctcc	ctgggtggccg	1800
	ctcgtctagag	cacgcgcgcc	gcagaccctag	ggtatttggc	gatcagcgtc	ctcggccatc	1860
	tcatcctcca	cactccgccc	ccaccacct	gccccagctg	ctaaggggtc	tgaccttttt	1920
	cagaaacgtg	catcttttcc	agttctaatt	ttgcacgctt	gcacgtttaa	agcaggaggg	1980
	atgaattcgg	tagtgataaa					2000

<210> 13

<211> 2300

<212> DNA

<213> Homo sapiens

<400> 13

	tcagattgtc	attgggaggg	tgaataaatg	aatgcttgca	ttatgagagt	ttgggggag	60
	aaatatgcc	cagactctta	tctgaagcca	tcagatttag	tggtgagaa	cccaccgaag	120
	tcagggattt	acatttttta	cagcaacgag	agaaaacttc	ccctttctc	tgcagaagtc	180
	aggactggat	ctcaaaaata	gaaatgtgtc	ctcctaaatg	tgtgcccac	ccgtgggtt	240
	acaaacaacg	gatttcccaa	gatagctgcc	acacacttgg	tttctaact	ctgtattgt	300
	tccccgccag	aatgtcgaag	tccttcccga	atatgcccag	tcatacttt	tgaactttt	360
	agcaaacacc	gtccggcttc	ttgtgctttc	ctcaaagacc	ccaggcaccg	gcagggagg	420
	cacaggccgg	ggcagagcgc	ccctgcgcgg	gggattcctg	ccactccgcg	ccagcctgcg	480

EP 2 305 807 A1

5
10
15
20
25

```

gcgcaaacgc tcttctcagc cgcagtccca cccgctgctg gcaatctgaa tgaggagccg 540
cgctatTTTT acctccccgg ctgcaatcct ttatatTTac atgcaggaag caaatatata 600
agggattaag aaggagatgc gtggccttag tttatccaga gcaggaagag gttggaatag 660
gagagggtat gtgaagtctg gggTggTgga aaaggcaggt ggacttcggc TggtTgTtt 720
ctccccgatca tcccTgtctc tggcctggaa acccccgtac tctctttctt ctggcttatc 780
cgtgactgcc ggctccccct ccaccgcccc catcttttga ggtaccaccg gtcacctccg 840
atgctgcttg ggctgctgca tcaactctgt gctttaccctt cttccccgcc cccaacaaa 900
gcatgCGcag TgcgtTccgg gccaggcaac agcagcagca cagcatccag caacagcatc 960
agcaccCGaa gccccgctcg ggCGcGctct cggggggcgg ggCGcAcgCC cgctccgCGc 1020
gtccccgCGc cgctcGctcc cGcGcGtccc cGcGcGctc gctccCGcGc gccGcctcag 1080
catcctcagG cccggCGgca gccccCGcag tCGctgaaGc ggccGcGccc gccgggggag 1140
ggagtagccG ctggggagGc tccaagTtgg cggagcggcg aggaccctg gactcctctg 1200
cgtcccGccc cgggagTggc Tgcgaggcta ggCGagccGg gaaaggggGc gccGcccagc 1260
ccccgagcccc gcgcccGtg ccccgagccc ggagccccct gccCGccGcg gaccatGcg 1320
cGccgagccG gcgtgaccGg ctccGcccGc ggccGccccg cagctagccc ggCGctctcG 1380
ccggccacac ggagCGcGc cCGggagcta tgagcattga agccGccccg cagcagctcG 1440
cggcagccGc ccctggcggg ctgcagcctt gccgCGcctt cctgCGgccc ccaacGcggc 1500
ccccCGgct cggtGcctGc cagcGccccg gcccGcAcGc cgccctGccg ctgcttctc 1560
gtccttctcc TgctGcctcc gctcGccGcc tCGtcccGgc cccGcGcctg gggggctgct 1620
gCGcccagcG gtgggtatgg ccccgTgccc tttGcgtTgg ctttcccGcg gggccctGca 1680
gaggaaaGcg aaggGcGcGc gggTccGtGt gctccggGct TgtcccGgc tcggcctttc 1740
cttcccctcc tccaccctt cGaaaaacc attccaattc ttgtTggtc 1800
acttttggaa gtccatttct gttgcattcG cctcagcGat ctctgTgcaa cttgCGgagG 1860
cactgggagG TgtttagTgg atcctggGtc gagcaagTtG ctactTgcGt ggtTgaaCct 1920
ggcaaccagT ggatgggaaa tacagcGagG atagTgTaa tgggctgTtt cctcaaattc 2040
taatgtgaat gcggggagga TgtagTgata cggTtggaaT taaggTggGg gaggcacact tcggggacca 2100
cgtatccGgc gcattcagTg catacttcat gcacgacctt Tggttctgat ttctcaaagT 2160
aagaattaag gtgctgaaga ataataataa taccatcttc tatatattga tgattggaag 2220
gctTgtcatt ataatgaaCa attaatataa gaaaataaaa agaaattggg aagctgccag 2280
tactgaaag cagaaagctg gctTgtcag gaaaataaaa agaaattggg aagctgccag 2300
catctgtatc cctacatggc

```

<210> 14
<211> 3000
<212> DNA
<213> Homo sapiens

30
35
40
45
50
55

```

<400> 14
tactgCCgac tttaggtctc tctggatctc aggccccctt ctctaagatg catcctagag 60
gaccaaaaat acactttatt tgggcttcGc ctgcttttGt ggaagggtag tttactagag 120
gatataatct cgtgttttaa tttgctctct ctccTaaagg aaatgtggag aaaaaaaaaa 180
agcagaaatt ggaaataacc aatatttagt ttatttcatt cgattcttag gggaaactggt 240
gaggagccta agatgatttt cccttcttag agaaagaatc caaagtccag ggaaatagcg 300
acaggggagT tcaagactGc cctctctagT ccttccttgg ctactctccg ctGcGatcGc 360
aggatagctc tcattagcag gagaatcggg caagtgtgtg gataagtaga gagtgtgttG 420
aacaacttGt aacgTtttat gaaatacGca ttgtcatggT tccTaaaag gctttgCGga 480
agccgTttGt ctttactaat caagtcttta cttacacaaa agtagaagta gaagtagttt 540
tagaaaacat actaacaatc ttctatcccc ttgaagacca gagtagcaga aaacaggtga 600
tttgattatT aaaattGcac tcactttttc ctccTttcag atttcacatt acattagccc 660
atTtGtGtta cggTgtataa aaaatggaaC aggcGcctcc actgcatTgt tctcctttaa 720
aaatagatca cttacaccct aactttgTtt tcctTaaatt cgattctTaa caggagagct 780
ttctattatt tcagatggag TgaggtTgca cGactgggat ggaagaaagg aatccctTaa 840
atttggggga atttctgttc tctgttccaa gaccatttta cttgggggtgT ggggggtggc 900
gcggcggtca gggcagTgga acgCagTcGc ggctgCGcCa tccctGcact tccaggcGcg 960
cgggagggac cggcggggac gcgagctGcg gactctggcg aactcggggg aggcagacag 1020
ggggagggcGg acaccagcc ggCagGcGtC tcagcctccc cGcagccGgc gggcttttct 1080
cctgacagct ccaggaagg cagaccctt cccagccag cCaggtAagg taaagactGc 1140
TgtTgagctt gctgttactg agggcGcaca gaccctgggg agaccgaagc ttgccactGc 1200
gggattctgt ggggtaacct gggTctacGg aagTttcctg aaagagggga gaagggTttg 1260
catttttctt atggaggatt ctctctctc tagcatttcg tttgatgtat tcaactggta 1320
gaagtGagat ttcaacaggt agcagagagc gctcagctgg aggaggtttg gggcGcGcg 1380
gcGccacccc caccctctc cgggaccGcg cctatttcta aagttacacg tcGacgaact 1440
aacctatgct ttaaattcct ctTccagcc cGgtgagTcc gcggcGacat tgggcccgtg 1500
ggTggctggg aacggTcccc tctccGgaa aaaccagaga acggctTgga gagctgaaac 1560
gagcgtccGc gagcaggtcc gtgcagaacc gggcttcagG accGctgagc tccgtagggc 1620
gtcctTgggg gacGccaggt cGccggctcc tctGccctcG ttgagatgga caacGcctcG 1680
ttctcggagc cctggcccGc caacGcatcG gcccCGgacc cGcGcGctgag ctgctccaac 1740
gcgtcGactc tggcGccGct gccggcGccg ctggcggTgg ctgtaccagT TgtctacGcg 1800
gtgatctGcg ccgtgggtct ggCGggCaac tccGccGtGc Tgtacgtgtt gctgCGgGcg 1860
ccccgatga agaccGtcac caacctgttc atcctcaacc TggccatcGc cGacgagctc 1920

```

EP 2 305 807 A1

	ttcacgctgg	tgctgccc	caacatcgcc	gacttcctgc	tgccgagctg	gcccttcggg	1980
	gagctcatgt	gcaagctcat	cgtdggctatc	gaccagtaca	acaccttctc	cagcctctac	2040
	ttcctcaccg	tcatgagcgc	cgaccgctac	ctgggtgggt	tggccactgc	ggagtcgcgc	2100
	cggttgcccg	cccgcaccta	cagcgcccg	cgcgcggtga	gcctggccgt	gtgggggatc	2160
5	gtcacactcg	tcgtgctgcc	cttcgcagtc	ttcgcccg	tagacgacga	gcagggcccg	2220
	cgccagtgcg	tgctagtctt	tccgcagccc	gaggccttct	ggtggcgcg	gagccgcctc	2280
	tacacgctcg	tgctgggctt	cgccatcccc	gtgtccacca	tctgtgtcct	ctataccacc	2340
	ctgtgtgccc	ggctgcatgc	catgcggtcg	gacagccacg	ccaaggccct	ggagcgcgc	2400
	aagaagcggg	tgaccttctc	ggtgggtggca	atcctggcgg	tgctgcctct	ctgctggacg	2460
	ccctaccacc	tgagcaccgt	ggtggcgctc	accaccgacc	tcccgcagac	gccgctggtc	2520
10	atcgctatct	cctacttcat	caccagcctg	agctacgcca	acagctgcct	caacccttct	2580
	ctctacgcct	tcctggacgc	cagcttccgc	aggaacctcc	gccagctgat	aacttgccgc	2640
	gcggcagcct	gactccccca	ggtcccggct	ccgcaactgc	ccgccaactc	tgccagcga	2700
	gggaggagcc	ggcgccagag	tgcgggacca	gacagggcgc	ctaggcctcc	tggggaaacc	2760
	gactcgcgcc	ccatacccga	cctagcagat	cggaagcgt	gagactgtgc	ccgaggttg	2820
	accttgccaa	gccctccagg	tgatgcgcgg	ccatgcccgg	tgaggagaac	tgaggctgag	2880
	atcgccacac	tgagggtctc	ctaaagccga	ggtggaggaa	gaggagggta	gaggaggag	2940
15	gcggtattgc	tgggaaaccg	cccctcctg	ccctgctccc	tgctgcccc	cccagccct	3000

<210> 15
 <211> 3000
 <212> DNA
 <213> Homo sapiens

20	<400> 15						
	gaatacatta	aagtagggg	aacccttgag	cccagacttc	tgccatgtga	agaccctttg	60
	aaaatcctga	caaacaacag	tactgctgaa	gtggtcagct	aattaaagag	gggaggtgga	120
	gctgtccttt	gtgtatccaa	taagtaccca	ttatctcatt	tgagcatgaa	aagaggccac	180
	tgattact	ttcaagaagg	aaagtaagca	ggatagctca	tatttttaga	accattcctc	240
	accaaattgga	ataattccgg	tgaaaagtgg	gagtgaggaa	gaaagaaaaa	aaaaacttct	300
25	aatcataatg	tttgggaata	agaaaggaag	aagaaaactca	cgcaaaagcc	gactttctcc	360
	tgagctgta	aaataaactc	ttaagacctt	tcctgctgaa	actctggaga	ggaaaactgg	420
	agtggcgggt	gggctttgcc	tgagctcaa	ctctccctcg	cgccgcgggc	gcggctgggt	480
	tcagcactc	ggaaagcgc	cctcgccggc	ccccgggatt	acgatgctc	cttggggccc	540
	gcccctttgg	cctgcaagt	gccaccgtaa	ctggtgagag	ccgctggcaa	cccaccocga	600
	ggtgacaacc	gcggagagac	gcagacacc	actgacctcc	aggaagctga	gcgtggtgga	660
30	tggaactcta	cgatctcttt	ctctccaagg	acggaacct	catccaagca	gtcccagagg	720
	aaacggataa	aggtatttga	aagggagcga	gcggcccaa	atcgcaaat	tgagcggctg	780
	ggggagttat	gcgccagtgc	cccagtgacc	gcgggacacg	gagaggggaa	gtctgcgttg	840
	tacataagga	cctagggact	ccgagcttgg	cttgagaacc	cttggacgcc	gagtgcttgc	900
	cttacgggct	gcactcctca	actctgctcc	aaagcagccg	ctgagctcaa	ctcctgcgtc	960
	cagggcgctt	gctgcgcgcc	aggacgcgct	tagtaccag	ttcctgggct	ctctcttcag	1020
35	tagctgcttt	gaaagctccc	acgcagctcc	cgaggctag	cctggcaaca	aaactggggt	1080
	aaaccgtgtt	atcttaggtc	ttgtccccca	gaacatgacc	tagaggtacc	tgcgcatgca	1140
	gatggccgat	gcagccacga	tagccaccat	gaataaggca	gcaggcgggg	acaagctagc	1200
	agaactcttc	agtctggtcc	cggaccttct	ggaggcggcc	aacacgagtg	gtaacgcgtc	1260
	gctgcagctt	ccggacttgt	ggtgggagct	ggggctggag	ttgccggacg	gcgcgcgcc	1320
	aggacatccc	ccgggcagcg	gcggggcaga	gagcgcggac	acagaggccc	gggtgcggtg	1380
	tctcatcagc	gggtgtact	gggtgggtgtg	gccttgggg	ttggcgggca	acctgtggt	1440
40	tctctacctg	atgaagagca	tgagggctg	gcgcaagtcc	tctatcaacc	tcttctgtac	1500
	caacctggcg	ctgacggact	ttcagtttgt	gctcaccctg	cccttctggg	cggtggagaa	1560
	cgctcttgac	ttcaaatggc	ccttcggcaa	ggccatgtgt	aagatcgtgt	ccatggtgac	1620
	gtccatgaac	atgtacgcca	gcgtgttctt	cctcactgcc	atgagtgatga	gcgctacca	1680
	ttcgggtggc	tcggctctga	agagccaccg	gaccgagga	cacggccggg	gcgactgctg	1740
	cggccggagc	ctgggggaca	gctgctgctt	ctcggccaag	gcgctgtgtg	tgtggatctg	1800
45	ggctttggcc	gcgctggcct	cgctgcccag	tgccatttct	tccaccacgg	tcaaggtgat	1860
	gggcgaggag	ctgtgcctgg	tgcgitttcc	ggacaagtgt	ctgggcccgc	acagggcgtt	1920
	ctggctgggc	ctctaccact	gcgagaaggt	gctgctgggc	ttcgtgctgc	cgctgggcat	1980
	cattatcttg	tgctacctgc	tgctggtgcg	cttcatcgcc	gaccgcccgc	cgccggggac	2040
	caaaggaggg	gcccgcgtag	ccggaggagc	cccagccgga	gccagcgcct	ggagactgtc	2100
	gaaggtcacc	aaatcagtg	ccatcgttgt	cctgtccttc	ttcctgtgtt	ggctgcccga	2160
50	ccaggcctc	accacctgga	gcacctctat	caagtcaac	gcggtgccct	tcagccagga	2220
	gtatttctctg	tgccagggtat	acgcgttccc	tgtgagcgtg	tgcttagcgc	actccaacag	2280
	ctgcctcaac	cccgtcctct	actgcctcgt	gcgcccgcgag	ttccgcaagg	cgctcaagag	2340
	cctgctgtgg	cgatcgcgt	ctccttcgat	caccagcatg	cgccccctca	ccgccaactac	2400
	caagccggag	cacgagatc	aggggctgca	ggccccggcg	ccgccccacg	cgcccgcca	2460
	gccggacctg	ctctactacc	cacctggcgt	cggtgtctac	agcggggggc	gctacgacct	2520
	gctgcccagc	agctctgcct	actgacgcag	gcctcaggcc	cagggcgcgc	cgctggggca	2580
55	aggtggcctt	ccccggcgg	taaagaggtg	aaaggatgaa	ggagggctgg	ggggggcccc	2640
	atttaagaag	taggtgggag	gaggatggg	agagcatgga	ggaggagcct	gtggataggg	2700

EP 2 305 807 A1

5 cgaggacctt ctctggagag gagatgcttc gaaatcaggt ggagagagga aattggcaaa 2760
 gggatagaga cgagccccac gggccagaca gccaacctcc gctccgcacc ccacagcctc 2820
 tccttactct tcccacgctg agtagtgctg gggcgccag aagcgaagac aagcagcaaa 2880
 aatgtagaga aattggcacg gggagcgggg cttagccaaa tgatgcacag acaattgtgc 2940
 ccgtttattc cagcgacttc tgcggagagg gcagccgtcg gcacaaacac tcctttgcgt 3000

<210> 16
 <211> 2200
 <212> DNA
 <213> Homo sapiens

10 <400> 16
 gtccccgat tccctcacc atcatataac gtgtgtattt attatgtttc ccgtttcctc 60
 tgtctccgcc agcagaatgt aaactccatg aggtcaggaa tctccgagtt atgttgccgc 120
 agtghtaatcc aagagcccgg aacagtgcct ggacacagc gggcatatgg aagaacaaat 180
 gtgtgaaggt gtgaatgaat gaataattga aagaataaat agtagttctc agcctcacag 240
 aacacgggtc acaacctcaa atgacctgct accctgccca taaataacag agatgcagga 300
 15 gtaagtgtcg ggctgtgacc tgtcaacatg ctaagccgct caaacaacac tgcccaacag 360
 cccgctggcc gcctatattg agcactgggc cctgagccgc acattcccat ttcgttgata 420
 aagaaactga ccagatagtt taagtggcct gctgcggaag acagagctgg tgctgcaccg 480
 gtcgctgctt ccccagtcct tttttggcct cttttctgac gcgacgcaga cccagttct 540
 ggagagtctg tcaactcgctc cccgtggtgg gagatcagag gcctgggtgc cttgggagcg 600
 gcgagcggtg ctccggcgag gatagaaagg gagtgccgag ccgagtcacc cagatccctg 660
 20 ggaacccgcg ccaccctccc gccctgccc atccccggcc gcgctgtcag tctccattag 720
 cgtaaacagw ctccagacgg agcggggccgg gcgctggggt aatgcaatcg gcgctgtacc 780
 tggggcgcag gctacattac cagcccggcc cccgctaggc acgcccagaa cccagtcagc 840
 cgcccccctgc cggccgcccc gcgcctccag ctcttccccg gccccgcccg aacgccacac 900
 ggcggagccc agccccagcc cgcgccctag agcctgcca ggcgcccggc gtcgggggcc 960
 ggcagggcgc aaggcaccag ggatcccctc gccgcccggc acgtgagtg gcctgagcg 1020
 cgggacaggg ctaggtctgc ctgggaggcc cgggcccaga cgcgccagca gagggctagc 1080
 25 gagttttagt tgcaagtacg ttaagtgtcc gagaaggctc ctgtggctgt tgaagtgtcg 1140
 cggacctgag ctggggaggg ggtcggcacg ctgccctcag cctcggtgag ttcaatccca 1200
 gccatttggg gcagggcaga gtgggtgaac gaggaaaagt gctgcagggg cttcagccgc 1260
 ccccagaggg ctgtcagaag tctccaactc ttgagttccg gcgtgcccc aacctgttt 1320
 ccaaattttt ccagcggacg cgcgctcttt tctgggaacc ctgctgccc tcagcgcgcg 1380
 ctcatcccag tgtctaaggc gctcccgggt ggtcttggga gttgcaagta gggaggaacg 1440
 30 gccgggtaac cacctctttt ccttttatcc aagcagagcc tcggcgtgcc cccaggaccg 1500
 gtaaagtccc tctcgccagc cgcattccatg cttctggcgc ggatgaacc gcaggtgcag 1560
 cccgagaaca acggggcgga cacgggtcca gacgagcccc ttcgggcgcg caaaactgcg 1620
 gagctgctgg tgggaaagga gcgcaacggc gtccagtgcc tgctggcgcc ccgagcggc 1680
 gacgcgcagc cccgggagac ctggggcaag aagatcgact tcctgctgtc cgtagtcggc 1740
 ttcgcagtgg acctggccaa cgtgtggcgc ttcccctacc tctgctacaa gaacggcggc 1800
 35 ggtgagcgtg gggctgggct gggaaattga atctgggagg tccactgtct gcagcgggtg 1860
 ctgggacagg agctggaata cacacggaag ggagggcagg agacaggggc aaatctgggg 1920
 cgcagaagaa actggacagg gctaaccggga aaaaaaaaaag attggagtcc tctggaaggt 1980
 cattttccca ggctctttgc agagtacctc gagctcattc cagcggaggt gtcaggattg 2040
 ggcaccctgg aagcaaaaca gcagaagagt gaaatcgagt catgacccta aagtcatggt 2100
 aggggatgag atggaagga cagaatctgg ggtgccaggt tgggtggggg agcctgacct 2160
 40 tttgatggtc tgctggaagg gaggtggaga ttccaagagc

<210> 17
 <211> 18
 <212> DNA
 <213> Artificial sequence

45 <220>
 <223> Designed oligonucleotide

<400> 17
 ctcagcacc aggcggcc 18

50 <210> 18
 <211> 20
 <212> DNA
 <213> Artificial sequence

55 <220>
 <223> Designed oligonucleotide

<400> 18

ctggccaaac tggagatcgc 20

<210> 19
<211> 386
<212> DNA
5 <213> Artificial Sequence

<220>
<223> Designed oligonucleotide

<400> 19
10 ctcagcacc aggcgccgc gatcatgagg cgcgagcggc gcgcgggctg ttgcagagtc 60
ttgagcgggt ggcacaccgc gatgtagcgg tcggctgtca tgactaccag catgtaggcc 120
gacgcaaaca tgccgaacac ctgcaggtgc ttcaccacgc ggcacagcca gtcggggccg 180
cgggaagcggg aggtgatgtc ccagcacatt tgcggcagca cctggaagaa tgccacggcc 240
aggtcggcca ggctgaggtg tcggatgaag aggtgcatgc gggacgtctt gcgcggcgtc 300
cgggtgcagag ccagcagtac gctgctgttg cccagcacgg ccaccgcgaa agtcaccgcc 360
15 agcacggcga tctccagttt ggcag 386

<210> 20
<211> 386
<212> DNA
20 <213> Artificial Sequence

<220>
<223> Designed oligonucleotide

<400> 20
25 ctcagcacc aggcgccgc gatcatgagg cgcgagcggc gcgcgggctg ttgcagagtc 60
ttgagcgggt ggcacaccgc gatgtagcgg tcggctgtca tgactaccag catgtaggcc 120
gacgcaaaca tgccgaacac ctgcaggtgc ttcaccacgc ggcacagcca gtcggggccg 180
cgggaagcggg aggtgatgtc ccagcacatt tgcggcagca cctggaagaa tgccacggcc 240
aggtcggcca ggctgaggtg tcggatgaag aggtgcatgc gggacgtctt gcgcggcgtc 300
cgggtgcagag ccagcagtac gctgctgttg cccagcacgg ccaccgcgaa agtcaccgcc 360
30 agcacggcga tctccagttt ggcag 386

<210> 21
<211> 17
<212> DNA
35 <213> Artificial Sequence

<220>
<223> Designed oligonucleotide

<400> 21
ctggccaaac tggagat 17

<210> 22
<211> 19
<212> DNA
40 <213> Artificial Sequence

<220>
<223> Designed oligonucleotide

<400> 22
45 tgagctccgt agggcgtcc 19

<210> 23
<211> 17
<212> DNA
50 <213> Artificial Sequence

<220>
<223> Designed oligonucleotide

<400> 23
55 gcgccgggtc cgggcc 17

EP 2 305 807 A1

<210> 24
 <211> 121
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> Designed oligonucleotide
 <400> 24
 gcgccgggtc cgggccgat gcgttggcgg gccagggctc cgagaacgag gcgttgtcca 60
 10 tctcaacgag ggcagaggag ccggcgacct ggcgtcccc aaggacgcc tacggagctc 120
 a 121
 <210> 25
 <211> 121
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 <223> Designed oligonucleotide
 <400> 25
 gcgccgggtc cgggccgat gcgttggcgg gccagggctc cgagaacgag gcgttgtcca 60
 20 tctcaacgag ggcagaggag ccggcgacct ggcgtcccc aaggacgcc tacggagctc 120
 a 121
 <210> 26
 <211> 19
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> Designed oligonucleotide
 <400> 26
 30 gacaacgcct cgttctcgg 19
 <210> 27
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 <223> Designed oligonucleotide
 <400> 27
 aggtgagcta cgtgtgtttg g 21
 40
 <210> 28
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> Designed oligonucleotide
 <400> 28
 agacatgtgc tcacgtacgg t 21
 50
 <210> 29
 <211> 391
 <212> DNA
 <213> Artificial Sequence
 55
 <220>
 <223> Designed oligonucleotide
 <400> 29
 aggtgagcta cgtgtgtttg ggcgtcgtgc actggctcac ttgtacgcg agaaatggca 60
 gcttgtagca ttggtgacct gccitttcga cactggacct ctatggacgt ggcggcggtg 120

EP 2 305 807 A1

tggcggcggc tcaatgacct gtggcgcccg tttgtggcgt gcgatagtcg agccgcctgt 180
 cacgtgcgcg gccgccctgc tccgtttgac gcgatgcata gcatgcgacc acccagtaat 240
 catactgctg acgctattgg tcacgtggtt atggcagctg ctggtgactg cgggtggcgtc 360
 ccgtttccac accgtacgtg agcacatgct t 391

5

<210> 30
 <211> 391
 <212> DNA
 <213> Artificial Sequence

10

<220>
 <223> Designed oligonucleotide

<400> 30
 aggtgagcta cgtgtgtttg ggcgtcgtgc actggctcac ttgtacgcgc agaaatggca 60
 gcttgtagca ttggtgacct gccttttcga cactggaccg ctatggacgt ggcggcgggtg 120
 tggcggcggc tcaatgacct gtggcgcccg tttgtggcgt gcgatagtcg agccgcctgt 180
 cacgtgcgcg gccgccctgc tccgtttgac gcgatgcata gcatgcgacc acccagtaat 240
 catactgctg acgctattgg tcacgtggtt atggcagctg ctggtgactg cgggtggcgtc 360
 ccgtttccac accgtacgtg agcacatgct t 391

15

<210> 31
 <211> 17
 <212> DNA
 <213> Artificial Sequence

20

<220>
 <223> Designed oligonucleotide

25

<400> 31
 ctggccaaac tggagat 17

<210> 32
 <211> 21
 <212> DNA
 <213> Artificial Sequence

30

<220>
 <223> Designed oligonucleotide

<400> 32
 ggacctgtgt ttgacgggta t 21

35

<210> 33
 <211> 21
 <212> DNA
 <213> Artificial Sequence

40

<220>
 <223> Designed oligonucleotide

<400> 33
 agtacagatc tggcgttctc g 21

45

<210> 34
 <211> 117
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed oligonucleotide

50

<400> 34
 ggacctgtgt ttgacgggta taactaag ttgcgcaatt tgctgtattg cgaaatccgc 60
 ccggacgata tcaacttga gcgcatgtgc cgtttccgag aacgccagat ctgtact 117

55

<210> 35
 <211> 117
 <212> DNA
 <213> Artificial Sequence

EP 2 305 807 A1

<220>
 <223> Designed oligonucleotide
 5 <400> 35
 ggacctgtgt ttgacgggta taacactaag ttgcgcaatt tgctgtattg cgaatccgc 60
 ccggacgata tcaactttga gcgcatgtgc cgtttccgag aacgccagat ctgtact 117
 <210> 36
 <211> 21
 <212> DNA
 10 <213> Artificial Sequence
 <220>
 <223> Designed oligonucleotide
 15 <400> 36
 agtacagatc tggcgttctc g 21
 <210> 37
 <211> 271
 <212> DNA
 <213> Artificial Sequence
 20 <220>
 <223> Designed oligonucleotide
 <400> 37
 tagggagtgc cagacagtgg gcgcaggcca gtgtgtgtgc gcaccgtgcg cgagccgaag 60
 25 cagggcgagg cattgcctca cctgggaagc gcaaggggtc agggagtcc ctttctgagt 120
 caaagaaagg ggtgacggtc gcacctggaa aatcgggtca ctcccacccg aatattgctc 180
 ttttcagacc ggcttaagaa acggcgcacc acgagactat atcccacacc tggctcggag 240
 ggtcctacgc ccacggaatc tcgctgattg c 271
 <210> 38
 <211> 17
 <212> DNA
 30 <213> Artificial Sequence
 <220>
 <223> Designed oligonucleotide
 35 <400> 38
 ctggcctaac tggagat 17
 <210> 39
 <211> 85
 <212> DNA
 <213> Artificial Sequence
 40 <220>
 <223> Designed oligonucleotide
 <400> 39
 45 cgggcgcggt ggctcacgcc tgtaatccca gcactttggg aggccgaggt gggcggatca 60
 cgaggtcagg agatcgagac catcc 85
 <210> 40
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 50 <220>
 <223> Designed oligonucleotide
 <400> 40
 ggatggtctc gatctcctga c 21
 55 <210> 41
 <211> 21

<212> DNA
 <213> Artificial Sequence

 <220>
 <223> Designed oligonucleotide
 5

 <400> 41
 aggtgagcta cgtgtgttg g 21

 <210> 42
 <211> 30
 10 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Designed oligonucleotide

 <400> 42
 15 gcgctgtgca ctggctcact tgtacgcgca 30

 <210> 43
 <211> 30
 20 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Designed oligonucleotide

 <400> 43
 25 cttgtacgat tggtagcccg ccttttcgac 30

 <210> 44
 <211> 30
 <212> DNA
 <213> Artificial Sequence
 30

 <220>
 <223> Designed oligonucleotide

 <400> 44
 35 actggaccgc tatggacgtg gcggcgggtg 30

 <210> 45
 <211> 30
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Designed oligonucleotide
 40

 <400> 45
 ggcggcggct caatgacctg tggcgcccgt 30

 <210> 46
 <211> 30
 45 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Designed oligonucleotide

 <400> 46
 50 ttgtggcgtg cgatagtcga gccgcctgct 30

 <210> 47
 <211> 25
 <212> DNA
 <213> Artificial Sequence
 55

 <220>

	<223> Designed oligonucleotide	
	<400> 47	
	acgtgcgcgg ccgccctgct ccgtt	25
5	<210> 48	
	<211> 30	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
10	<223> Designed oligonucleotide	
	<400> 48	
	tgacgcgatg catagcatgc gaccaccag	30
	<210> 49	
15	<211> 30	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Designed oligonucleotide	
20	<400> 49	
	actgctgacg ctattggtca cgtggttatg	30
	<210> 50	
	<211> 30	
25	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Designed oligonucleotide	
30	<400> 50	
	ctgctgttga ctgcggtggc gtcccgtttc	30
	<210> 51	
	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Designed oligonucleotide	
	<400> 51	
	ggacctgtgt ttgacgggta t	21
40	<210> 52	
	<211> 25	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
45	<223> Designed oligonucleotide	
	<400> 52	
	aacactaagt tgcgcaattt gctgt	25
	<210> 53	
50	<211> 25	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Designed oligonucleotide	
55	<400> 53	
	attgcgaaat ccgcccggac gatat	25

5
 <210> 54
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed oligonucleotide

10
 <400> 54
 cactcttgag cgcatgtgcc gtttc 25

<210> 55
 <211> 30
 <212> DNA
 <213> Artificial Sequence

15
 <220>
 <223> Designed oligonucleotide

<400> 55
 cagtgtgtgt gcgcaccgtg cgcgagccga 30

20
 <210> 56
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed oligonucleotide

25
 <400> 56
 ggcgagggcat tgcctcacct gggaagcgca 30

<210> 57
 <211> 30
 <212> DNA
 <213> Artificial Sequence

30
 <220>
 <223> Designed oligonucleotide

<400> 57
 ggtgacggtc gcacctgaa aatcgggtca 30

<210> 58
 <211> 30
 <212> DNA
 <213> Artificial Sequence

40
 <220>
 <223> Designed oligonucleotide

<400> 58
 acccgaatat tgcgcttttc agaccggctt 30

45
 <210> 59
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Designed oligonucleotide

50
 <400> 59
 tcggagggtc ctacgcccac ggaatctcgc 30

55
 <210> 60
 <211> 30
 <212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide

<400> 60

cgggcgcggt ggctcacgcc tgtaatcca

30

<210> 61

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide

<400> 61

tttgggaggc cgaggtgggc ggatcacgag

30

<210> 62

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide

<400> 62

ggtggctcac gcctgtaatc

20

<210> 63

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide

<400> 63

ggatggcttc gatctcctga c

21

Claims

1. A method for quantifying or detecting DNA comprising a target DNA region contained in a specimen comprising:

(1) First step of preparing from the specimen DNA for which the target DNA region is to be detected;

(2) Second step of treating the DNA prepared in First step with a DNA methylation enzyme;

(3) Third step of preparing single-stranded methylated DNA from the DNA treated in Second step, and making a detection oligonucleotide bind with the single-stranded methylated DNA to obtain a test DNA complex;

(4) Fourth step of making an immobilized methylated-DNA antibody bind with the test DNA complex obtained in Third step to obtain a detection complex; and

(5) Fifth step of quantifying or detecting DNA comprising a target DNA region in the single-stranded DNA by quantifying or detecting the detection oligonucleotide contained in the detection complex obtained in Fourth step by its identification function.

2. The method according to claim 1, wherein a counter oligonucleotide is added in obtaining a test DNA complex in Third step.

3. The method according to claim 1 or 2, wherein the immobilized methylated-DNA antibody is a methylcytosine antibody.

EP 2 305 807 A1

4. The method according to any one of claims 1 to 3, wherein the DNA methylation enzyme is a cytosine methylation enzyme or Sssl methylase.
5. The method according to any one of claims 1 to 4, wherein the DNA comprising a target DNA region contained in the specimen is DNA comprising a target DNA region in DNA generated from RNA by a reverse transcriptase.
6. The method according to any one of claims 1 to 5, wherein the specimen is any of the following biological specimen:
- (a) mammalian blood, body fluid, excreta, body secretion, cell lysate, or tissue lysate,
 - (b) DNA extracted from one selected from the group consisting of mammalian blood, body fluid, excreta, body secretion, cell lysate, and tissue lysate,
 - (c) DNA prepared by using as a template RNA extracted from one selected from the group consisting of mammalian tissue, cell, tissue lysate and cell lysate,
 - (e) DNA extracted from bacterium, fungus or virus, or
 - (f) DNA prepared by using as a template RNA extracted from cell, fungus or virus.
7. The method according to any one of claims 1 to 6, wherein the DNA comprising a target DNA region obtained in First step is DNA digested in advance with a restriction enzyme recognition cleavage site for which is not present in the target DNA region, a synthesized oligonucleotide, or DNA purified in advance.
8. The method according to any one of claims 1 to 7, wherein the identification function of the detection oligonucleotide is any of the following identification function:
- (a) fluorescence detection of FITC, or
 - (b) detection by FITC antibody.
9. The method according to any one of claims 1 to 8, wherein the detection oligonucleotide comprises a repetitive sequence or a nucleotide sequence of an overlapping gene or a pseudo gene in human genome or a nucleotide sequence capable of complementarily binding with a part thereof.
10. The method according to claim 9, wherein the repetitive sequence in human genome is LINE or SINE;
11. The method according to any one of claims 1 to 10, wherein the detection oligonucleotide comprises a nucleotide sequence capable of complementarily binding with any one of the following nucleotide sequences:
- (1) the nucleotide sequence of SEQ ID NO: 37 or a nucleotide sequence having 80% or more sequence identity to the same,
 - (2) a complementary sequence of the nucleotide sequence of SEQ ID NO: 37 or a nucleotide sequence having 80% or more sequence identity to the same,
 - (3) the nucleotide sequence of SEQ ID NO: 39 or a nucleotide sequence having 80% or more sequence identity to the same, or
 - (4) a complementary sequence of the nucleotide sequence of SEQ ID NO: 39 or a nucleotide sequence having 80% or more sequence identity to the same.
12. The method according to any one of claims 1 to 10, wherein the detection oligonucleotide comprises any one of the following nucleotide sequences:
- (1) the nucleotide sequence of SEQ ID NO: 38 or a nucleotide sequence having 80% or more sequence identity to the same,
 - (2) a complementary sequence of the nucleotide sequence of SEQ ID NO: 38 or a nucleotide sequence having 80% or more sequence identity to the same,
 - (3) the nucleotide sequence of SEQ ID NO: 40 or a nucleotide sequence having 80% or more sequence identity to the same, or
 - (4) a complementary sequence of the nucleotide sequence of SEQ ID NO: 40 or a nucleotide sequence having 80% or more sequence identity to the same.
13. The method according to any one of claims 1 to 12, wherein concentration of a sodium salt in a solution used in a DNA extracting operation for preparing DNA from a specimen in First step is 100 mM or more and 1000 mM or less.

EP 2 305 807 A1

14. The method according to any one of claims 1 to 12, wherein concentration of a sodium salt in a solution used in a DNA extracting operation for preparing DNA from a specimen in First step is 100 mM or more and 200 mM or less.

5
15. A method for selecting a specimen from a cancer patient comprising the step of evaluating that a specimen from a test subject is a specimen from a cancer patient when there is significant difference between a quantification result or a detection result of DNA quantified or detected by using the specimen from the test subject according to the method of any one of claims 1 to 14 and a quantification result or a detection result of DNA quantified or detected by using a specimen from a healthy subject according to the same method, and identifying a specimen from a cancer patient based on a result of the evaluation.

10
16. The method according to claim 15, wherein the specimen is mammalian serum.

17. The method according to claim 15 or 16, wherein DNA comprising a target DNA region is free DNA comprising the target DNA region in a mammalian serum.

15

20

25

30

35

40

45

50

55

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig.19





Fig.20



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2009/061068

A. CLASSIFICATION OF SUBJECT MATTER C12N15/09(2006.01)i, C12Q1/25(2006.01)i, C12Q1/68(2006.01)i, G01N33/53(2006.01)i According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C12N15/09, C12Q1/25, C12Q1/68, G01N33/53 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Jitsuyo Shinan Koho 1922-1996 Jitsuyo Shinan Toroku Koho 1996-2009 Kokai Jitsuyo Shinan Koho 1971-2009 Toroku Jitsuyo Shinan Koho 1994-2009 Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CA/BIOSIS/MEDLINE/WPIDS (STN), JSTPlus/JMEDPlus/JST7580 (JDreamII)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BAEK, T.J., et al., Development of a photodiode array biochip using a bipolar semiconductor and its application to detection of human papilloma virus. Anal. Bioanal. Chem., Mar 2008, Vol.390, No.5, p.1373-1378	1-17
A	JP 2004-536596 A (U.S. Genomics, Inc.), 09 December, 2004 (09.12.04), & US 2002/0197639 A1 & EP 1402071 A & WO 2002/101353 A2	1-17
A	JP 2004-347508 A (Japan Science and Technology Agency), 09 December, 2004 (09.12.04), & WO 2004/104582 A1	1-17
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 19 August, 2009 (19.08.09)		Date of mailing of the international search report 01 September, 2009 (01.09.09)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/JP2009/061068

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2005/080565 A1 (Japan Science and Technology Agency), 01 September, 2005 (01.09.05), & US 2008/0227652 A1 & EP 1717312 A1	1-17
A	GAO, L., et al., DNA microarray: a high throughput approach for methylation detection. Colloids Surf. B: Biointerfaces., 2005, Vol.40, No.3-4, p.127-131	1-17

Form PCT/ISA/210 (continuation of second sheet) (April 2007)

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- *J. Cataract. Refract. Surg.*, 2007, vol. 33 (4), 635-641 [0002]
- *Environ. Mol. Mutagen.*, 1991, vol. 18 (4), 259-262 [0002]
- *Methods in Yeast Genetics*. Cold Spring Harbor Laboratory Press [0040]
- *Molecular Cloning-A Laboratory Manual*. Cold Spring Harbor Laboratory Press [0040]
- *J. Am. Chem. Soc.*, 2007, vol. 129, 5612-5620 [0106]
- *Methods in Yeast Genetics*. Cold Spring Harbor Laboratory [0242] [0262] [0289] [0345] [0367] [0396] [0425]

专利名称(译)	检测或定量DNA的方法		
公开(公告)号	EP2305807A1	公开(公告)日	2011-04-06
申请号	EP2009762576	申请日	2009-06-11
[标]申请(专利权)人(译)	住友化学有限公司		
申请(专利权)人(译)	住友化学公司		
当前申请(专利权)人(译)	住友化学公司		
[标]发明人	TOMIGAHARA YOSHITAKA SATOH HIDEO TARUI HIROKAZU		
发明人	TOMIGAHARA, YOSHITAKA SATOH, HIDEO TARUI, HIROKAZU		
IPC分类号	C12N15/09 C12Q1/25 C12Q1/68 G01N33/53		
CPC分类号	C12Q1/6816 C12Q1/6804 C12Q1/6827 C12Q1/6834 C12Q1/6886		
优先权	2008152619 2008-06-11 JP		
其他公开文献	EP2305807A4		
外部链接	Espacenet		

摘要(译)

本发明涉及定量或检测具有靶DNA区域的DNA的方法，等等。

X:5'-

CTCAGCACCCAGGCGCGCATGAGGCGAGCGCGCGGGCTGTTGCAGAGTCTT
GAGCGGGTGGCACACCGCATGTAGCGGTGGCTGTCATGACTACCAGCATGTAGGCCGACG
CAAACATGCCGAACACCTGCAGGTGCTTACCACGCGGCACAGCCAGTGGGGCCGCGGAAG
CGGTAGGTGATGTCCAGCACATTTGCGGCAGCACCTGGAAGAATGCCACGGCCAGGTGGC
CAGGCTGAGGTGTCGGATGAAGAGGTGCATGCGGGAGCTCTTGCAGGGCTCCGGTGCAGAG
CCAGCAGTACGCTGCTGTTGCCAGCACGGCCACCGGAAAGTCACCGCCAGCACGGCGATC
TCCAGTTTGGCCAG -3' (SEQ ID NO: 19)