



(11)

**EP 2 576 832 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:  
**04.05.2016 Bulletin 2016/18**

(51) Int Cl.:  
**C12Q 1/68 (2006.01) C12N 15/12 (2006.01)**  
**G01N 33/53 (2006.01) G01N 33/574 (2006.01)**

(21) Application number: **11787338.0**

(86) International application number:  
**PCT/US2011/037926**

(22) Date of filing: **25.05.2011**

(87) International publication number:  
**WO 2011/150075 (01.12.2011 Gazette 2011/48)**

(54) **COMPOSITIONS AND METHODS FOR DETECTING A NEOPLASIA**

ZUSAMMENSETZUNGEN UND VERFAHREN ZUM NACHWEIS EINER NEOPLASIE

COMPOSITIONS ET MÉTHODES POUR DÉTECTER UNE NÉOPLASIE

(84) Designated Contracting States:  
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**

(30) Priority: **25.05.2010 US 348010 P**

(43) Date of publication of application:  
**10.04.2013 Bulletin 2013/15**

(73) Proprietor: **The Johns Hopkins University**  
**Baltimore, MD 21218 (US)**

(72) Inventors:  
• **AHUJA, Nita**  
**Lutherville, MD 21093 (US)**  
• **BAYLIN, Stephen**  
**Baltimore, MD 21230 (US)**  
• **HERMAN, James, Gordon**  
**Lutherville, MD 21093 (US)**  
• **WANG, Jeff**  
**Timonium, MD 21093 (US)**  
• **BAILEY, Vasudev**  
**Baltimore, MD 21210 (US)**  
• **YI, Joo-Mi**  
**Laurel, MD 20723 (US)**

(74) Representative: **Hiebl, Inge Elisabeth**  
**Kraus & Weisert**  
**Patentanwälte PartGmbB**  
**Thomas-Wimmer-Ring 15**  
**80539 München (DE)**

(56) References cited:  
• **SHAMES DAVID S ET AL: "A genome-wide screen for promoter methylation in lung cancer identifies novel methylation markers for multiple malignancies", PLOS MEDICINE, PUBLIC LIBRARY OF SCIENCE, US, vol. 3, no. 12, 1 December 2006 (2006-12-01), XP002527542, ISSN: 1549-1676, DOI: 10.1371/JOURNAL.PMED.0030486**  
• **CHOI J E ET AL: "Aberrant methylation of ADAMTS1 in non-small cell lung cancer", CANCER GENETICS AND CYTOGENETICS, ELSEVIER SCIENCE PUBLISHING, NEW YORK, NY, US, vol. 187, no. 2, 1 December 2008 (2008-12-01), pages 80-84, XP026067163, ISSN: 0165-4608, DOI: 10.1016/J.CANCERGENCYTO.2008.08.001 [retrieved on 2008-11-21]**  
• **PHILIP J. JOHNSON ET AL: "Plasma Nucleic Acids in the Diagnosis and Management of Malignant Disease", CLINICAL CHEMISTRY, vol. 48, no. 8, 1 August 2002 (2002-08-01), pages 1186-1193, XP055042293, ISSN: 0009-9147**  
• **ESTELLER M: "Relevance of DNA methylation in the management of cancer", LANCET ONCOLOGY, LANCET PUBLISHING GROUP, LONDON, GB, vol. 4, no. 6, 1 June 2003 (2003-06-01), pages 351-358, XP004809686, ISSN: 1470-2045, DOI: 10.1016/S1470-2045(03)01115-X**  
• **G STRATHDEE ET AL: "Aberrant DNA methylation in cancer: potential clinical interventions", EXPERT REVIEWS IN MOLECULAR MEDICINE, vol. 4, 1 March 2002 (2002-03-01), pages 1-17, XP055087285,**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 2 576 832 B1**

- SEO D ET AL: "Genomic medicine: bringing biomarkers to clinical medicine", CURRENT OPINION IN CHEMICAL BIOLOGY, CURRENT BIOLOGY LTD, LONDON, GB, vol. 9, no. 4, 1 August 2005 (2005-08-01), pages 381-386, XP027847778, ISSN: 1367-5931 [retrieved on 2005-08-01]
- SHAMES, D. S. ET AL.: 'A genome-wide screen for promoter methylation in lung cancer identifies novel methylation markers for multiple malignancies' PLOS MEDICINE. vol. 3, no. 12, December 2006, pages 2244 - 2263, XP002527542
- CHOI, J. E. ET AL.: 'Aberrant methylation of ADAMTS1 in non-small cell lung cancer' CANCER GENETICS AND CYTOGENETICS vol. 187, no. 2, December 2008, pages 80 - 84, XP026067163
- JOHNSON, P. J. ET AL.: 'Plasma nucleic acids in the diagnosis and management of malignant disease' CLINICAL CHEMISTRY vol. 48, no. 8, August 2002, pages 1186 - 1193, XP055042293
- AHLQUIST, T. ET AL.: 'Gene methylation profiles of normal mucosa, and benign and malignant colorectal tumors identify early onset markers' MOLECULAR CANCER. vol. 7, no. 94, 31 December 2008, XP021050163
- MORRIS, M. R. ET AL. IDENTIFICATION OF CANDIDATE TUMOUR SUPPRESSOR GENES FREQUENTLY METHYLATED IN RENAL CELL CARCINOMA pages 2104 - 2117, XP055100662
- AHLQUIST TERJE ET AL: "Gene methylation profiles of normal mucosa, and benign and malignant colorectal tumors identify early onset markers", MOLECULAR CANCER, BIOMED CENTRAL, LONDON, GB, vol. 7, no. 1, 31 December 2008 (2008-12-31), page 94, XP021050163, ISSN: 1476-4598, DOI: 10.1186/1476-4598-7-94

**Description****BACKGROUND OF THE INVENTION**

5 **[0001]** Pancreatic cancer is the fourth leading cause of cancer mortality with a death rate nearly equal to the incidence of this disease. Detection of cancer specific, abnormally DNA methylated gene promoter sequences has emerged as one of the leading tumor biomarker detection strategies. Recently, there have been successful reports of DNA methylation screening using various body fluids, such as stool for detection of colorectal cancer, sputum for lung cancer, and urine for prostate cancer.

10 **[0002]** However, in pancreatic cancer no screening tool is currently available for early detection. This is particularly relevant as pancreatic cancer is often found once it is already metastatic or locally advanced and the diagnosis is often delayed because patients present with nonspecific gastrointestinal symptoms. The development of a screening modality for pancreatic cancer which identifies early stage cancers amenable to surgical curative resection would then have a potential impact in reducing mortality from this currently lethal disease.

15 **[0003]** SHAMES DAVID S ET AL: "A genome-wide screen for promoter methylation in lung cancer identifies novel methylation markers for multiple malignancies", PLOS MEDICINE, PUBLIC LIBRARY OF SCIENCE, US, vol. 3, no. 12, 1 December 2006 (2006-12-01), ISSN: 1549-1676, DOI: 10.1371/JOURNAL.PMED.0030486 discloses a high-throughput global expression profiling approach in lung cancer cells to identify new cancer-specific methylation markers. 132 genes were identified that have 5' CpG islands that are induced from undetectable levels by 5-aza-2'-deoxycytidine in multiple non-small cell lung cancer cells and are expressed in immortalized human bronchial epithelial cells.

20 **[0004]** CHOI J E ET AL: "Aberrant methylation of ADAMTS1 in non-small cell lung cancer", CANCER GENETICS AND CYTOGENETICS, ELSEVIER SCIENCE PUBLISHING, NEW YORK, NY, US, vol. 187, no. 2, 1 December 2008 (2008-12-01), pages 80-84, ISSN: 0165-4608, DOI: 10.1016/J.CANCERGENCYT0.2008.08.001 discloses the expression of ADAMTS1 in non-small cell lung cancer (NSCLC) and the relationship between ADAMTS1 methylation and clinicopathologic features in NSCLC patients. The results suggest that aberrant methylation of ADAMTS1 frequency occurs in NSCLCs and may play a role in the pathogenesis of NSCLC.

25 **[0005]** PHILIP J. JOHNSON ET AL: "Plasma Nucleic Acids in the Diagnosis and Management of Malignant Disease", CLINICAL CHEMISTRY, vol. 48, no. 8, 1 August 2002 (2002-08-01), pages 1186-1193, ISSN: 0009-9147 discloses circulating nucleic acids as an emerging class of molecular tumor markers such as plasma DNA, microsatellite analysis, chromosomal translocations, epigenetic alterations and viral DNA.

30 **[0006]** ESTELLER M: "Relevance of DNA methylation in the management of cancer", LANCET ONCOLOGY, LANCET PUBLISHING GROUP, LONDON, GB, vol. 4, no. 6, 1 June 2003 (2003-06-01), pages 351-358, ISSN: 1470-2045, DOI: 10.1016/S1470-2045(03) 01115-X generally discloses DNA methylation as a therapeutic target.

35 **[0007]** G STRATHDEE ET AL: "Aberrant DNA methylation in cancer: potential clinical interventions", EXPERT REVIEWS IN MOLECULAR MEDICINE, vol. 4, 1 March 2002 (2002-03-01), pages 1-17, presents studies of DNA methylation in cancer opening up new opportunities for diagnosis, prognosis and treatment of human tumors.

40 **[0008]** AHLQUIST TERJE ET AL: "Gene methylation profiles of normal mucosa, and benign and malignant colorectal tumors identify early onset markers", MOLECULAR CANCER, BIOMED CENTRAL, LONDON, GB, vol. 7, no. 1, 31 December 2008 (2008-12-31), page 94, ISSN: 1476-4598, DOI: 10.1186/1476-4598-7-94 discloses epigenetic markers that can discriminate between non-malignant and malignant tissue from the large bowel, i.e. markers with diagnostic potential.

**SUMMARY OF THE INVENTION**

45 **[0009]** As described below, the present invention features methods for diagnosing, detecting or characterizing a neoplasia (pancreatic cancer, colon cancer, lung cancer) featuring *BNC1* and *ADAMTS1*. Advantageously, the method provides for screening serum for increased promoter methylation of *BNC1* and *ADAMTS1* to identify early stage neoplasia or a propensity to develop a neoplasia.

50 **[0010]** In one aspect, the invention provides an in vitro method for detecting or characterizing a neoplasia in a biologic sample of a subject, the method comprising detecting the methylation of a *BNC1* and *ADAMTS1* gene, wherein detection of methylation detects or characterizes the presence of a neoplasia in the sample, the method optionally further comprising the step of detecting an alteration in the sequence or expression level of a *Brca1*, *Brca2*, *p16*, *K-ras*, *APC*, *PalB2*, *DPC4*, *EGFR*, and/or *EML-ALK4* gene or polypeptide, wherein the neoplasia is a cancer selected from colon, lung or pancreatic cancer and wherein the level of methylation present at *BNC1* and *ADAMTS1* gene promoter is compared to a reference selected from the level of methylation present in a control sample obtained from a patient that does not have a neoplasia, a baseline level of methylation present in a biologic sample derived from a patient, prior to, during or after, treatment for a neoplasia or a standardized curve.

55 **[0011]** In another aspect, the invention provides an in vitro method for detecting or characterizing lung or colon cancer

in a sample derived from a subject, the method comprising detecting methylation of a *BNC1* and *ADAMTS1* gene, wherein detection of methylation detects or characterizes lung or colon cancer in the subject and further comprising detecting an alteration in the sequence or expression level of a *Brca1*, *Brca2*, *p16*, *K-ras*, *APC*, *EGFR* (*lung*), and/or *EML-ALK4* gene or polypeptide, wherein the level of methylation present at *BNC1* and *ADAMTS1* gene promoter is compared to a reference selected from the level of methylation present in a control sample obtained from a patient that does not have a neoplasia, a baseline level of methylation present in a biologic sample derived from a patient, prior to, during or after, treatment for a neoplasia or a standardized curve. In another embodiment, the subject is identified as having a propensity to develop a neoplasia (e.g., is identified as a smoker, having colon polyps or adenomas, or a family history of cancer).

**[0012]** In another aspect, the invention provides an in vitro method for detecting or characterizing pancreatic cancer in a serum or plasma sample derived from a subject, the method comprising detecting the methylation of *BNC1* and *ADAMTS1* gene, wherein detection of methylation detects or characterizes pancreatic cancer in the subject, further comprising the step of detecting an alteration in the sequence or expression of a *Brca1*, *Brca2*, *p16*, *K-ras*, *APC*, *PalB2*, and/or *DPC4* gene or polypeptide relative to a reference, wherein the level of methylation present at *BNC1* and *ADAMTS1* gene promoter is compared to a reference selected from the level of methylation present in a control sample obtained from a patient that does not have a neoplasia, a baseline level of methylation present in a biologic sample derived from a patient, prior to, during or after, treatment for a neoplasia or a standardized curve. In another embodiment, the method further involves imaging the subject, and localizing the cancer. In one embodiment, the subject is identified as having a propensity to develop a pancreatic cancer (e.g., is identified as a smoker, has a *Brca1* or *Brca2* mutation, pancreatic cyst, chronic pancreatitis, or a family history of cancer).

**[0013]** Further described is a method of monitoring a subject diagnosed as having a neoplasia, the method involving detecting an alteration in promoter methylation level in a *BNC1* and *ADAMTS1* gene in a subject sample relative to a reference, where an altered level indicates an altered severity of neoplasia in the subject. In one embodiment, the reference is the level of methylation present in a sample previously obtained from the subject; is a baseline level of methylation present in a sample from the subject obtained prior to therapy; or is the level of methylation present in a normal patient sample. In another embodiment, a decreased level of methylation relative to a reference indicates a reduced severity of the neoplasia, and an increased level of methylation relative to a reference indicates an increased severity of neoplasia.

**[0014]** Further described is a method for selecting a treatment for a subject diagnosed as having a neoplasia, the method involving detecting methylation of a *BCN1* and *ADAMTS1* gene, where detection of methylation indicates that epigenetic therapy should be selected for treatment of said subject. In one embodiment, the epigenetic therapy is selected from the group consisting of entinostat, SAHA (suberoylanilide hydroxamic acid), depsipeptide, azacytidine, and deazacytidine.

**[0015]** Further described is a kit for the analysis of promoter methylation, the kit involving at least one primer capable of distinguishing between methylated and unmethylated *BNC1* and *ADAMTS1* promoter sequences. In one embodiment, the kit further contains a pair of primers for amplifying the promoter sequence of a reference gene. In another embodiment, the kit further contains a detectable probe, where the probe is capable of binding to the promoter sequence. In yet another embodiment, the probe is detected by fluorescence, by autoradiography, by an immunoassay, by an enzymatic assay, or by a colorimetric assay.

**[0016]** In other embodiments of the above aspects, the biologic sample is a tissue or biologic fluid sample (e.g., of blood, serum, plasma, urine, pancreatic juice, pancreatic cyst fluid, or lung lavage). The reference is the level of methylation present at the promoter in a control sample. In other embodiments, the control sample is derived from a healthy subject. In still other embodiments, the methylation is detected by quantitative methylation-specific PCR (QMSP). In other embodiments, the level of methylation is quantified or the frequency of methylation is quantified. In still other embodiments, the methylation levels of the *BNC1* and *ADAMTS1* promoters are quantified. In still other embodiments, the method results in at least 50% - 100% (e.g., 50%, 60%, 70%, 80%, 90%, 100%) sensitivity. In still other embodiments, the subject is identified as having a propensity to develop a neoplasia (e.g., is a smoker, has a *Brca1* or *Brca2* mutation, pancreatic cyst, chronic pancreatitis, presence of colon polyps or adenomas, or a family history of cancer). In various embodiments, the method further involves detecting an alteration in the sequence or expression level of a *Brca1*, *Brca2*, *p16*, *K-ras*, *APC*, *PalB2*, *DPC4*, *EGFR*, and/or *EML-ALK4* gene or polypeptide. In still other embodiments, the alteration is a sequence alteration or alteration in expression level. In still other embodiments, the subject is a human patient. In various embodiments of the above aspects, the methylation is detected or quantified by methylation on beads or quantitative methylation-specific PCR.

**[0017]** The invention provides methods for detecting or characterizing a neoplasia in a biologic sample of a subject, the method comprising detecting promoter methylation of *BNC1* and *ADAMTS1* genes.

## Definitions

**[0018]** Unless defined otherwise, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art to which this invention belongs. The following references provide one of skill with a general definition of many of the terms used in this invention: Singleton et al., Dictionary of Microbiology and Molecular Biology (2nd ed. 1994); The Cambridge Dictionary of Science and Technology (Walker ed., 1988); The Glossary of Genetics, 5th Ed., R. Rieger et al. (eds.), Springer Verlag (1991); and Hale & Marham, The Harper Collins Dictionary of Biology (1991). As used herein, the following terms have the meanings ascribed to them below, unless specified otherwise.

**[0019]** By "BNC1 gene" is meant a polynucleotide encoding a BNC1 polypeptide. An exemplary BNC1 gene nucleotide sequence is publically available at GeneBank No.: NM\_001717. An exemplary sequence is provided below.

```

1  ggggcggggg  cggccatcgt  gctgcgcagc  ctgggcgctt  ggggagccgc  ccacttcgcc
61  gggtcgcgcc  ccgacggccg  gagcgtggat  gcggcggcgc  ccgccgagcc  ggggcggacg
15  121  cggggcggcc  cgggcccggg  agacgcgccg  gcagccccgg  caccgcagcg  gtcgcaggat
181  ggccgaggct  atcagctgta  ctctgaactg  tagttgccaa  agtttcaaac  ccgggaaaat
241  aaaccaccgt  cagtgtgacc  aatgcaagca  tggatgggtg  gcccacgctc  taagtaagct
301  aaggatcccc  cccatgtatc  caacaagcca  ggtggagatt  gtccagtcca  atgtagtgtt
361  tgatattagc  agcctcatgc  tctatgggac  ccaggccatc  cccgttcgcc  taaaaatcct
20  421  actggaccgg  ctcttcagtg  tgttgaagca  agatgagggt  ctccagatcc  tccatgcctt
481  ggactggaca  cttcaggatt  atatccgtgg  atacgtactg  caggatgcat  caggaaaggt

```

25

30

35

40

45

50

55

EP 2 576 832 B1

541 gttggatcac tggagcatca tgaccagtga ggaagaagtg gccaccttgc agcagttcct  
 601 tcgttttgga gagaccaaat ctatagttga actcatggca attcaagaga aagaagagca  
 661 atccatcatc ataccacctt ccacagcaaa tgtagatc agggctttca tcgagagctg  
 721 cagtcacagg agttctagcc tccccactcc tgtggacaaa gaaacccca gcagtataca  
 781 cccctttgag aacctcataa gcaacatgac tttcatgctg cttttccagt tcttcaacc  
 841 tctgcctcct gcactgatag ggtcattgcc cgaacaatat atgttggagc agggctatga  
 901 ccaaagtcag gaccccaaac aggaagtcca tgggcccttc cctgacagca gcttctaac  
 961 ttccagttcc acaccatttc aggttgaaaa agatcagtg ttaaactgtc cggatgctat  
 1021 tactaaaaaa gaagacagca cccatttaag tgactccagc tcatacaaca ttgtcactaa  
 1081 gtttgaaagg acacagttat ccctgaggc caaagtgaag cctgagagga atagccttgg  
 1141 tacaagaag ggccgggtgt tctgcactgc atgtgagaag accttctatg acaaaggcac  
 1201 cctcaaaatc cactacaatg ccgtccactt gaagatcaag cataagtgca ccatcgaagg  
 1261 gtgtaacatg gtgttcagct ccctaaggag ccggaatcgc catagcgcca accccaacc  
 1321 tcggctgcac atgccaatga acagaaataa ccgggacaaa gacctcagga acagcctgaa  
 1381 cctggccagc tctgagaact acaagtgcc aggtttcaca gtgacgtccc cagactgtag  
 1441 gcctcctccc agctaccctg gttcaggaga ggattccaaa ggccaaccag ctttcccaa  
 1501 cattgggcaa aatggtgtgc tttttccaa cctaaagaca gtccagccag tcttctctt  
 1561 ctaccgcagt ccagccacgc ctgccgaggt agcaaacacg cctgggatac tcccttccct  
 1621 cccgctgttg tctcttcaa tcccagaaca gctcatttca aacgaaatgc cttttagatg  
 1681 ccttcccaag aagaaatcca ggaagtccag tatgcctatc aaaatagaga aagaagctgt  
 1741 ggaatatagc aatgagaaaa gacacaacct cagctcagat gaagacatgc ccctacaggt  
 1801 ggtcagtgaa gatgagcagg aggcctgcag tccctcagtca cacagagtat ctgaggagca  
 1861 gcatgtacag tcaggaggct tagggaagcc tttccctgaa ggggagaggc cctgccatcg  
 1921 tgaatcagta attgagtcca gtggagccat cagccaaacc cctgagcagg ccacacacaa  
 1981 ttcagagagg gagactgagc agacaccagc attgatcatg gtgccaaggg aggtcgagga  
 2041 tgggtggccat gaacactact tcacacctgg gatggaacc caagttcctt tttctgacta  
 2101 catggaactg cagcagcgc tgctggctgg gggactctc agtgctttgt ccaacagggg  
 2161 aatggctttt ccttgtcttg aagattctaa agaactggag cacgtgggtc agcatgcatt  
 2221 agcaaggcag atagaagaaa atcgttcca gtgtgacatc tgcaagaaga ctttaaaaa  
 2281 tgctttagt gtgaaaattc atcacaagaa tatgcatgtc aaagaaatgc acacatgcac  
 2341 agtggagggc tgtaatgcta ctttccctc ccgcaggagc agagacagac acagctcaaa  
 2401 cctaaacctc caccaaaaag cattgagcca ggaagcattg gagagtagtg aagatcattt  
 2461 ccgtgcagct taccttctga aagatgtggc taaggaagcc tatcaggatg tggcttttac  
 2521 acagcaagcc tcccagacat ctgtcatctt caaaggaaca agtcgaatgg gcagtctggt  
 2581 ttaccaata acgcaagtc acagtgccag cctggagagc tacaactctg gcccttgag  
 2641 cgagggcacc atcctggatt tgagcactac ctgagcatg aagtcagaga gtgagcaca  
 2701 ttcttctggt gactctgagc ggtgagtgga ggaaggcact gtgcttatgg aggcagtgga  
 2761 tgggaactgt gaagggtcga gccttgtccc tggggaagat gagtaccca tctgtgtcct  
 2821 gatggagaag gctgaccaga gccttgttag cctgccttct gggttgccca taacctgtca  
 2881 tctctgcaa aagacataca gtaacaaagg gacctttagg gccactaca aaactgtgca  
 2941 cctccggcag ctccacaaat gcaaagtacc aggtgcaac accatgtttt cgtctgttcg  
 3001 cagtcgaaac agacacagcc agaatccaa cctgcacaaa agcctggcct catctccaag  
 3061 tcacctccag taacaagatg gcaaaccaag tatgctcaga taagctttt tcataattca  
 3121 ggaataaagt agtccataga aatgtttctg tttcatatca tttggggcga gtcaggcaaa  
 3181 agtatttgat ttgactttat agttttccac agcacaatga gcaaaagaca aacctcgtgg  
 3241 gaagatgaca ctggggcagc ctttccattt attttctta gcccaagagg tctttcactg  
 3301 atacaaggaa aacttgacga aatgtgattt tttccagatt tgtttacatg ttccctggga  
 3361 cagatccagg tctgcagatc gacaccagtg ggcccaggac ctgggggtg ctttaaatga  
 3421 ggcttgagc gttaaaggctc ttggataaga agggctctgg ggaagaagac tctgtggaca  
 3481 agataccagt ccccaaaaaca gcattttcag ttccttctc aattagtttg aaatccagac  
 3541 ctgagtttg aagactgatt ttttgagacc atccctgtgt ttggagtgga taattgtccc  
 3601 tcccctcagc cctgcaccag aggtctcata tgttaccaca gggagttctc agaggattgg  
 3661 gttggcctct aacatgttcc ttgttaattc ttgttctgta acatgcattc aagaagctag  
 3721 gggaaaaata tctcatgcac ttaataatg gtcttcaatt taatttaaaa atatttgac  
 3781 aatatttaat ttgtgcttat gtggtgtttg gtgtgagtg agatattgca ctgtgtcacc  
 3841 tctggatctc tgctcagaag cagaacaagt gatgacctaa atgtcaaat cactgctcgt  
 3901 tttcatttgg tgaacttcaa actctgttct ttttggtcac ctgtggaatg aatgcaagca

EP 2 576 832 B1

3961 tgatTTTTggc agaacattt gtacatatc tgccgtagat aatgtgggtc tgatggttgt  
 4021 tgtgtatTTT cagtatcact ggatccctca gtcttcaccg ttttataaac gtataagatt  
 4081 aggatgaact tttgaattta cttggtagga aaaaaagtag gacattattg ccatattgta  
 5 4141 tgtcttaata ttttaacttat tcggaaatat attccacact gttacataca ttttccatgg  
 4201 tagaaaggaa gttcagtcag tcctgtggaa tgaaaccatc tcctaaaatt cagcatttgc  
 4261 agcattctaa aagcctgtgt aggtacaagg acattgattt tgtattcaga attcaagtta  
 4321 actatctttt aaattcgtgg ttgatgtaag taataaaaaa cattcttaaa gttgaggggt  
 4381 ataagagaga ttatttctgt ggtctaaagg ttaaaaaagcc aacaacctgt taccaattat  
 10 4441 ttcagctttt tttgttttaa taagtgtgac aacttaaaac ttgtttctat ttaaagtgaa  
 4501 atgtatcttt caactgttta gttaccagc tgtttaatat tccagtcttc ccaaagtgaa  
 4561 aagatttgta tacaaatgtt ttctatgatt taataaaaaat atatggcaca ccaaaaaaaa  
 4621 aaaaaaa

15 gtttccttttcaaggatccccaccagcagcaaggcaggtggaagaaactg  
 agggaaagaggaagcaggacccccgcccagctctagcctttacag  
 actccttttaaggagctgaagaggttgggagaggcttgccctgccccggg  
 gcagcagaaacccgggcgctgctgcagcagctgtgggcgatctccgtcag  
 ctectgggctccaggatcctgcccctcagcccctccgcgtctactctcg  
 20 gggccaccacctccagttggttctccagggcacttcatttcattccgtc  
 cctagcatccaataagacgcgaaaaacttaatcccattttacaactgaag  
 ccactgagggcccagagagcaaagccatttgtcctgggaaagtggcgaagc  
 tgggactttgactcttcagaccaccaaggggaaggggtgtgtgtgtgt  
 gtgtgtgtgtgtgcgtgtgcggtggggggcgctgttttgagttcttaag  
 25 aaaacctcctggcgaccccccttctccacatcccaagacgctcgtcccgc  
 actttctcgggaatgaggtttctgcagggcagggcgcgctgecttcttc  
 ctccgcggcagtgagacccccgagggcgccccagggtaggaggggaggccg  
 aatcatctcctgagaagagcgcagagaacttcagagcgtttcgcccttc  
 cccgggagagggcaaacaccgacacgtctgtgtctttaccaacaagtgc  
 ttcaagcccggcgggggcagacacctccgcgcggccgcccggcgaggtct  
 30 ccgcggtctgcgggggcccagggcctcgcctcagctgcgctgatttagggc  
 gttatccggtcccggggcgggagggcgcctcccggggcgcgaaagcagcgc  
 ccgcggcgtggggcgaccgcgggtgggcgagggggcagggggaggggcg  
 gagagggcgtccccggggcgagggggcgggcgtgcgggcacacgcgggtgc

35 GCGGCGGGGGCGGCCATCGTGCTGCGCAGCCTGGGCGCTTGGGGAGCCGC  
 CCACTTCGCCGGGTCGCGCCCCGACGGCCGGAGCGTGATGCGGCGGCGC  
 CCGCCGAGCCGGGGCGGACGCGGGGCGGCCCGGGCCGGGAGACGCGCCG  
 GCAGCCCCGGCACCGCAGCGGTCGACAGGATGGCCGAGgt aagcgcggcgc

40  
 1 gcactcgtg gaaagcggct ccgagccagg ggctattgca aagccagggt gcgctaccgg  
 61 acggagaggg gagagccctg agcagagtga gcaacatcgc agccaaggcg gaggccgaag  
 121 aggggcgcca ggcaccaatc tccgcgttgc ctacgccccg gaggcgcccc agagcgttc

[0020] By "BNC1 promoter" is meant a polynucleotide sequence sufficient to direct expression of a BNC1 coding sequence. The sequence of an exemplary BNC1 promoter (BNC1 at chr15:81715659-8174472) is provided below:

**BNC1 promoter region upstream 1kb from Exon 1**

**BNC1 Exon 1**

[0021] By "ADAMTS1 gene" is meant a polynucleotide sequence encoding a ADAMTS1 protein. An exemplary ADAMTS1 gene sequence is publically available at GeneBank No.: NM\_006988. An exemplary ADAMTS1 gene sequence is provided below:

EP 2 576 832 B1

181 ttgtcccagc agagccactc tgccctgcgcc tgccctctcag tgtctccaac tttgcgctgg  
 241 aagaaaaact tcccgcgcgc cggcagaact gcagcgcctc cttttagtga ctccgggagc  
 301 ttcggctgta gccggctctg cgcgccttc caacgaataa tagaaattgt taattttaac  
 361 aatccagagc aggccaaacga ggctttgctc tcccgaaccg aactaaaggt ccctcgctcc  
 421 gtgcgctgct acgagcgggtg tctcctgggg ctccaatgca gcgagctgtg cccgaggggt  
 481 tcggaaggcg caagctgggc agcgacatgg ggaacgcgga gcgggctccg gggctcggga  
 541 gctttgggcc cgtaccacag ctgctgctgc tcgccgcggc gctactggcc gtgtcggacg  
 601 cactcgggcg cccctccgag gaggacgagg agctagtggg gccggagctg gagcgcgcc  
 661 cgggacacgg gaccacgcgc ctccgcctgc acgcctttga ccagcagctg gatctggagc  
 721 tgcggcccga cagcagcttt ttggcgcccg gcttcacgct ccagaacgtg gggcgcaaat  
 781 ccgggtccga gacgcgcctt ccggaaccg acctggcgca ctgcttctac tccggcaccg  
 841 tgaatggcga tcccagctcg gctgccgccc tcagcctctg cgagggcgtg cgcggcgctt  
 901 tctacctgct gggggaggcg tatttcatcc agccgctgcc cgcgccagc gagcgcctcg  
 961 ccaccgccgc cccaggggag aagccgcggc caccactaca gttccacctc ctgcgggga  
 1021 atcggcaggg cgacgtcggc ggacgtgcg gggctcgtgga cgacgagccc cggccgactg  
 1081 gaaagcggga gaccgaagac gaggacgaag ggactgaggg cgaggacgaa ggggctcagt  
 1141 ggtccgcgca ggaccgggca ctgcaaggcg taggacagcc cacaggaact ggaagcataa  
 1201 gaaagaagcg atttgtgtcc agtcaccgct atgtggaac catgcttgtg gcagaccagt  
 1261 cgatggcaga attccacggc agtggctctaa agcattacct tctcacgttg tttcggtg  
 1321 cagccagatt gtacaaacac cccagcattc gtaattcagt tagcctgggtg gtggtgaaga  
 1381 tcttggcat ccacgatgaa cagaaggggc cggaaagtgc ctccaatgct gccctcactc  
 1441 tgcggaactt ttgcaactgg cagaagcagc acaaccacc cagtgaccgg gatgcagagc  
 1501 actatgacac agcaattctt ttaccagac aggacttgtg tgggtcccag acatgtgata  
 1561 ctcttgggat ggctgatggt ggaactgtgt gtgatccgag cagaagctgc tccgtcatag  
 1621 aagatgatgg tttacaagct gccttaccac cagcccatga attaggccac gtgtttaaca  
 1681 tgccacatga tgatgcaaag cagtgtgcc gccttaatgg tgtgaaccag gattcccaca  
 1741 tgatggcgtc aatgctttcc aacctggacc acagccagcc ttggtctcct tgcagtgcct  
 1801 acatgattac atcatttctg gataatggtc atggggaatg tttgatggac aagcctcaga  
 1861 atcccataca gctcccaggc gatctccctg gcacctcgta cgatgccaac cggcagtgcc  
 1921 agtttacatt tggggaggac tccaaacact gccccgatgc agccagcaca tgtagcacct  
 1981 tgtggtgtac cggcacctct ggtggggtgc tgggtgtgtca aaccaaacac ttcccgtggg  
 2041 cggatggcac cagctgtgga gaagggaat ggtgtatcaa cggcaagtgt gtgaacaaaa  
 2101 ccgacagaaa gcattttgat acgccttttc atggaagctg ggaatgtgg gggccttggg  
 2161 gagacttcc gagaacgtgc ggtggaggag tccagtacac gatgagggaa tgtgacaacc  
 2221 cagtccaaa gaatggaggg aagtactgtg aaggcaaacg agtgcgctac agtccctgta  
 2281 accttgagga ctgtccagac aataatggaa aaaccttag agaggaacaa tgtgaagcac  
 2341 acaacgagtt ttcaaaagct tcctttggga gtgggcctgc ggtggaatgg attcccaagt  
 2401 acgctggcgt ctaccaaag gacaggtgca agctcatctg ccaagccaaa ggcattggct  
 2461 acttcttctg tttgcagccc aagggtgtag atggtactcc atgtagccca gattccacct  
 2521 ctgtctgtgt gcaaggacag tgtgtaaaag ctggttgtga tcgcatcata gactccaaaa  
 2581 agaagtttga taaatgtggt gtttgcgggg gaaatggatc tacttgtaaa aaaatatcag  
 2641 gatcagttac tagtgcaaaa cctggatctc atgatatcat cacaattcca actggagcca  
 2701 ccaacatcga agtgaacag cgaaccaga ggggatccag gaacaatggc agctttcttg  
 2761 ccatcaaagc tgctgatggc acatatattc ttaatggtga ctacactttg tccaccttag  
 2821 agcaagacat tatgtacaaa ggtgttgtct tgaggtacag cggctcctct gcggcattgg  
 2881 aaagaattcg cagcttttag cctctcaaag agcccttgac catccaggtt ctactgtgg  
 2941 gcaatgcctc tcgacctaaa attaaataca cctactcgt aaagaagaag aaggaatctt  
 3001 tcaatgctat ccccactttt tcagcatggg tcattgaaga gtggggcgaa tgttctaagt  
 3061 catgtgaatt gggttggcag agaagactgg tagaatgccg agacattaat ggacagcctg  
 3121 cttccgagtg tgcaaaggaa gtgaagccag ccagcaccag acctgtgca gaccatcctt  
 3181 gccccagtg gcagctgggg gagtgtcat catgttctaa gacctgtgg aagggttaca  
 3241 aaaaagaag cttgaagtgt ctgtccatg atggaggggt gttatctcat gagagctgtg  
 3301 atcctttaa gaaacctaaa cattcatag actttgac aatggcagaa tgcagttaag  
 3361 tggtttaagt ggtgttagct ttgagggcaa ggcaaagtga ggaagggctg gtgcagggaa  
 3421 agcaagaagg ctggagggat ccagcgtatc ttgccagtaa ccagtgaggt gtatcagtaa  
 3481 ggtgggatta tgggggtaga tagaaaagga gttgaatcat cagagtaaac tgccagttgc  
 3541 aaatttgata ggatagttag tgaggattat taacctctga gcagtgatat agcataataa

EP 2 576 832 B1

3601 agccccgggc attattatta ttatttcttt tgttacatct attacaagtt tagaaaaaac  
 3661 aaagcaattg tcaaaaaaag ttagaactat tacaaccct gtttcctggg acttatcaaa  
 3721 tacttagtat catgggggtt gggaaatgaa aagtaggaga aaagtgagat tttactaaga  
 5 3781 cctgttttac tttacctcac taacaatggg gggagaaagg agtacaataa ggatctttga  
 3841 ccagcactgt ttatggctgc tatggtttca gagaatgttt atacattatt tctaccgaga  
 3901 attaaaactt cagattgttc aacatgagag aaaggctcag caacgtgaaa taacgcaaat  
 3961 ggcttcctct ttcctttttt ggaccatctc agtctttatt tgtgtaattc attttgagga  
 4021 aaaaacaact ccatgtattt attcaagtgc attaaagtct acaatggaaa aaaagcagtg  
 4081 aagcattaga tgctggtaaa agctagagga gacacaatga gcttagtacc tccaacttcc  
 10 4141 tttctttcct accatgtaac cctgctttgg gaatatggat gtaaagaagt aacttgtgtc  
 4201 tcatgaaaat cagtacaatc acacaaggag gatgaaacgc cggaacaaaa atgaggtgtg  
 4261 tagaacaggg tcccacaggt ttggggacat tgagatcact tgtcttggg tggggaggct  
 4321 gctgaggggt agcaggtcca tctccagcag ctggtccaac agtcgtatcc tgggtgaatgt  
 4381 ctgttcagct cttctgtgag aatatgattt tttccatatg tatatagtaa aatatgttac  
 15 4441 tataaattac atgtacttta taagtattgg tttgggtgtt cttccaaga aggactatag  
 4501 ttagtaataa atgcctataa taacatattt atttttatac atttatttct aatgaaaaaa  
 4561 acttttaaat tataatcgctt ttgtggaagt gcatataaaa tagagtattt atacaatata  
 4621 tgttactaga aataaaaaga cacttttggg aaaaaaaaaa aaaaaaaaaa

20 By "ADAMTS1 promoter" is meant is meant a polynucleotide sequence sufficient to direct expression of a BNC1 coding sequence. The sequence of an exemplary *AdamTS1* (ADAMTS1 at chr21:27130477-27139599) promoter is provided below:

ADAMTS1 promoter region upstream 1kb from Exon 1

25  
 30  
 35  
 40  
 45  
 atttccttctttccccctctgcacgcttgetagccccagcgatcgctgct  
 ggccccgggtaggaaagtggggttctctggcogtttctgacgctggcc  
 tagggcttgcagctgctgttgagtgaaagcacgcagactggcgggagccg  
 atcatttctcgaatgaagaagaaaaagcgaattccctccttatgctcta  
 gggaattgagccgcgtcccagatcaccattccagaaatgtgaaaccggg  
 ccctcacaagtctctctggtgaagaggtggcgtgaggggtggggtt  
 gtggaggggtgaaggcataagcaaacatattttaaaatccagatcgtagga  
 agtgtcacctggcccctcaccagcatgctttctgggggaagcgcaggg  
 ccaagcttccctagaaaagctggggcgaagagagagcaggcggcggcta  
 35 aggagctcctggcaggctgggaaggtggagaagtggggtgaggtatttt  
 ctagaaagtgtagccctagctcatctcctagattggggaagagggactg  
 agggaggagggaaaggagaccagggcagctccaggatagggaaatgttga  
 agaagggactgcttctccaaccgaaccctccctcctgggaaccgcagcc  
 cagcgcggttaactgagttaccgcaaccgggcggtgggaggaaggggtgt  
 40 ccaggaaacccggcgagggagaaaagcgggtggaagggagagtcttctccct  
 ggagcggccccagcagtagcaagtgtggtcacagcggccccttccgcccc  
 tagattgacgagcagtagcgtggagccagcggaggctgccccctccc  
 ctcccagccccgcagcgggagcgggttagcaaccaacggagccggggg  
 cggcgtctttgggatggaaaagggccaaaggggaggagtggggtgggggt  
 45 gggggtttcactggtccactataaaaggaccgctcggctgcccggttctt

ADAMTS1 Exon 1

50  
 55  
 GCACTCGCTGGAAAGCGGCTCCGAGCCAGGGGCTATTGCAAAGCCAGGGT  
 GCGCTACCGGACGGAGAGGGGAGAGCCCTGAGCAGAGTGAGCAACATCGC  
 AGCCAAGGCGGAGGCCGAAGAGGGGCGCCAGGCACCAATCTCCGCGTTGC  
 CTCAGCCCCGGAGGCGCCCCAGAGCGCTTCTTGTCCCAGCAGAGCCACTC  
 IGCCGTGCGCCTGCCTCTCAGTGTCTCCAACCTTTGCGCTGGAAGAAAACT  
 TCCCGCGCGCCGGCAGAAGTGCAGCGCCTCCTTTTAGTGACTCCGGGAGC  
 TTCGGCTGTAGCCGGCTCTGCGCGCCCTTCCAACGAATAATAGAAATTGT

TAATTTTAACAATCCAGAGCAGGCCAACGAGGCTTTGCTCTCCCGACCCG  
 AACTAAAGGTCCCTCGCTCCGTGCGCTGCTACGAGCGGTGTCTCCTGGGG  
 CTCCAATGCAGCGAGCTGTGCCCGAGGGGTTTCGGAAGGCGCAAGCTGGGC  
 5 AGCGACATGGGGAACGCGGAGCGGGCTCCGGGGTCTCGGAGCTTTGGGCC  
 AGTACCCACGCTGCTGCTGCTCGCCGCGGCGCTACTGGCCGTGTGGACG  
 CACTCGGGCGCCCCCTCCGAGGAGGACGAGGAGCTAGTGGTGCCGGAGCTG  
 GAGCGCGCCCCGGGACACGGGACCACGCGCCTCCGCCTGCACGCCTTTGA  
 CCAGCAGCTGGATCTGGAGCTGCGGCCGACAGCAGCTTTTTGGCGCCCG  
 10 GCTTCACGCTCCAGAACGTGGGGCGCAAATCCGGGTCCGAGACGCCGCTT

**[0022]** By "alteration" is meant an increase or decrease. An alteration may be by as little as 1%, 2%, 3%, 4%, 5%, 10%, 20%, 30%, or by 40%, 50%, 60%, or even by as much as 75%, 80%, 90%, or 100%. An alteration may be a change in sequence relative to a reference sequence or a change in expression level, activity, or epigenetic marker (e.g., promoter methylation or histone alterations).

**[0023]** By "biologic sample" is meant any tissue, cell, fluid, or other material derived from an organism.

**[0024]** In this disclosure, "comprises," "comprising," "containing" and "having" and the like can have the meaning ascribed to them in U.S. Patent law and can mean "includes," "including," and the like; "consisting essentially of" or "consists essentially" likewise has the meaning ascribed in U.S. Patent law and the term is open-ended, allowing for the presence of more than that which is recited so long as basic or novel characteristics of that which is recited is not changed by the presence of more than that which is recited, but excludes prior art embodiments.

**[0025]** "Detect" refers to identifying the presence, absence or amount of the analyte to be detected.

**[0026]** By "clinical aggressiveness" is meant the severity of the neoplasia. Aggressive neoplasias are more likely to metastasize than less aggressive neoplasias. While conservative methods of treatment are appropriate for less aggressive neoplasias, more aggressive neoplasias require more aggressive therapeutic regimens.

**[0027]** By "control" is meant a standard of comparison. For example, the methylation level present at a promoter in a neoplasia may be compared to the level of methylation present at that promoter in a corresponding normal tissue.

**[0028]** By "diagnostic" is meant any method that identifies the presence of a pathologic condition or characterizes the nature of a pathologic condition (e.g., a neoplasia). Diagnostic methods differ in their sensitivity and specificity. While a particular diagnostic method may not provide a definitive diagnosis of a condition, it suffices if the method provides a positive indication that aids in diagnosis.

**[0029]** By "frequency of methylation" is meant the number of times a specific promoter is methylated in a number of samples.

**[0030]** By "increased methylation" is meant a detectable positive change in the level, frequency, or amount of methylation. Such an increase may be by 5%, 10%, 20%, 30%, or by as much as 40%, 50%, 60%, or even by as much as 75%, 80%, 90%, or 100%. In certain embodiments, the detection of any methylation in an *AdamTS1* and *BNC* promoter in a subject sample is sufficient to identify the subject as having a neoplasia, a pre-cancerous lesion, or the propensity to develop a neoplasia.

**[0031]** By "isolated polynucleotide" is meant a nucleic acid (e.g., a DNA) that is free of the genes which, in the naturally-occurring genome of the organism from which the nucleic acid molecule of the invention is derived, flank the gene. The term therefore includes, for example, a recombinant DNA that is incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of a prokaryote or eukaryote; or that exists as a separate molecule (for example, a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. In addition, the term includes an RNA molecule that is transcribed from a DNA molecule, as well as a recombinant DNA that is part of a hybrid gene encoding additional polypeptide sequence.

**[0032]** By "methylation level" is meant the number of methylated alleles. Methylation level can be represented as the methylation present at a target gene/reference gene x 100. Any ratio that allows the skilled artisan to distinguish neoplastic tissue from normal tissue is useful in the methods of the invention. One skilled in the art appreciates that the cutoff value is selected to optimize both the sensitivity and the specificity of the assay. In certain embodiments, merely detecting promoter methylation of *BNC1* and *AdamTS1* genes in a biological sample of a subject is sufficient to identify the subject as having cancer, a pre-cancerous lesion, or having a propensity to develop cancer.

**[0033]** By "tumor marker profile" is meant an alteration present in a subject sample relative to a reference. In one embodiment, a tumor marker profile includes promoter methylation of *BNC1* and *AdamTS1* genes, as well as mutations present in *Brca1*, *p16*, *K-ras*, *APC*, *PalB2*, *DPC4*, EGFR, EML- ALK4 or other marker known in the art.

**[0034]** By "sensitivity" is meant the percentage of subjects with a particular disease that are correctly detected as having the disease. For example, an assay that detects 98/100 of carcinomas has 98% sensitivity.

**[0035]** By "severity of neoplasia" is meant the degree of pathology. The severity of a neoplasia increases, for example, as the stage or grade of the neoplasia increases.

**[0036]** By "specificity" is meant the percentage of subjects without a particular disease who test negative.

**[0037]** By "neoplasia" is meant any disease that is caused by or results in inappropriately high levels of cell division, inappropriately low levels of apoptosis, or both. For example, cancer is an example of a neoplasia. Examples of cancers include, without limitation, pancreatic cancer, including islet cell and adenocarcinomas), duodenal cancers, cholangiocarcinomas, ampullary tumors, leukemia's (e.g., acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, chronic leukemia, chronic myelocytic leukemia, chronic lymphocytic leukemia), polycythemia vera, lymphoma (Hodgkin's disease, non-Hodgkin's disease), Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors such as sarcomas and carcinomas (e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colorectal carcinoma, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, neuroendocrine carcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, nile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, uterine cancer, testicular cancer, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, schwannoma, meningioma, melanoma, neuroblastoma, and retinoblastoma). Lymphoproliferative disorders are also considered to be proliferative diseases.

**[0038]** By "periodic" is meant at regular intervals. Periodic patient monitoring includes, for example, a schedule of tests that are administered daily, bi-weekly, bi-monthly, monthly, bi-annually, or annually.

**[0039]** By "promoter" is meant a nucleic acid sequence sufficient to direct transcription. In general, a promoter includes, at least, 50, 75, 100, 125, 150, 175, 200, 250, 300, 400, 500, 750, 1000, 1500, or 2000 nucleotides upstream of a given coding sequence (i.e upstream of the coding sequence for BNC1 and ADAMTS1 polypeptides).

**[0040]** By "marker" is meant any protein or polynucleotide having an alteration in methylation, expression level or activity that is associated with a disease or disorder.

**[0041]** By "reduces" is meant a negative alteration of at least 10%, 25%, 50%, 75%, or 100%.

**[0042]** By "reference" is meant a standard or control condition. Exemplary references include a baseline of methylation present in a healthy control subject or a standardized curve.

**[0043]** A "reference sequence" is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset of or the entirety of a specified sequence; for example, a segment of a full-length cDNA or gene sequence, or the complete cDNA or gene sequence. For polypeptides, the length of the reference polypeptide sequence will generally be at least about 16 amino acids, preferably at least about 20 amino acids, more preferably at least about 25 amino acids, and even more preferably about 35 amino acids, about 50 amino acids, or about 100 amino acids. For nucleic acids, the length of the reference nucleic acid sequence will generally be at least about 50 nucleotides, preferably at least about 60 nucleotides, more preferably at least about 75 nucleotides, and even more preferably about 100 nucleotides or about 300 nucleotides or any integer thereabout or therebetween.

**[0044]** By "subject" is meant a mammal, including, but not limited to, a human or non-human mammal, such as a bovine, equine, canine, ovine, or feline.

**[0045]** Ranges provided herein are understood to be shorthand for all of the values within the range. For example, a range of 1 to 50 is understood to include any number, combination of numbers, or sub-range from the group consisting 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42 43, 44, 45, 46, 47, 48, 49, or 50.

**[0046]** As used herein, the terms "treat," treating," "treatment," and the like refer to reducing or ameliorating a disorder and/or symptoms associated therewith. It will be appreciated that, although not precluded, treating a disorder or condition does not require that the disorder, condition or symptoms associated therewith be completely eliminated.

**[0047]** Unless specifically stated or obvious from context, as used herein, the term "or" is understood to be inclusive. Unless specifically stated or obvious from context, as used herein, the terms "a", "an", and "the" are understood to be singular or plural.

**[0048]** Unless specifically stated or obvious from context, as used herein, the term "about" is understood as within a range of normal tolerance in the art, for example within 2 standard deviations of the mean. About can be understood as within 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.1%, 0.05%, or 0.01% of the stated value. Unless otherwise clear from context, all numerical values provided herein are modified by the term about.

**[0049]** Any methods provided herein can be combined with one or more of any of the other methods provided herein.

## BRIEF DESCRIPTION OF THE DRAWINGS

**[0050]**

Figures 1A, 1B, and 1C show a schematic diagram, a gel, and a bar graph, respectively. Figure 1A depicts a schematic that is a step-wise representation of the screening method used to identify epigenetic biomarkers. Figure 1B depicts the results of methylation analysis of 8 cancer-specific genes in pancreatic cancer cell lines. M=methylation signal; U=unmethylated signal. IVD= *in vitro* methylated DNA. ddH<sub>2</sub>O=water control adding no DNA. Figure 1C is a bar graph that displays the methylation frequencies of the 8 genes in a series of pancreatic cancer patients from JHU (stage 1-4; n=143), along with a series of normal pancreas from non cancerous patients (n=4), and a series of pancreatic intraductal neoplasia (PanIN) (n=20 ranging from Pan IN 1-3).

Figures 2A, 2B, and 2C depict promoter methylation gels and corresponding bar graphs, a CpG methylation diagram, and a schematic depicting methylation-on-bead (MOB) technology, respectively. Figure 2A shows silencing of *BNC1* and *ADAMTS1* genes in pancreatic cancer cell lines by a methylation specific PCR (MSP) analysis of *BNC1* and *ADAMTS1* gene promoter regions (upper panel depicting gel bands), and the correlation of promoter methylation with gene expression by quantitative RT-PCR in pancreatic cancer cell lines (lower panel showing data in bar graph format. M=methylation signal; U=unmethylated signal. IVD= *in vitro* methylated DNA. Real-time RT-PCR expression is shown as fold change  $\pm$  SE relative to mock-treated cells during 5uM 5-Aza-deoxycytidine (DAC) and 300 nM Trichostatin A (TSA) treatments. Figure 2B shows the results of bisulfite sequencing analysis of CpG islands in the *BNC1* gene promoter region. The *BNC1* promoter regions is depicted as a line, with CpG regions designated as vertical slashes. TIS indicates the transcriptional start site. Location of CpG sites (bisulfite primers (BST): upstream region from -331 to +36 relative to transcriptional start site). Bisulfite sequencing analysis of the *BNC1* gene in Panc1 (pancreatic cancer cell line), normal pancreas tissue, pancreatic primary tumors and HCT116 (DNMT1/3b<sup>-/-</sup>) as a negative control. Open and filled circles represent unmethylated and methylated CpG sites, respectively, and each row represents a single clone. Figure 2C shows step-wise illustration of the nanoassay used to detect methylation in serum. DNA extraction and bisulfite conversion are carried out using methylation-on-beads (MOB). DNA is eluted using PCR buffer, and amplified using MSP modified primers and fluorophore-labeled NTPs. Samples are analyzed using MS-qFRET or gel electrophoresis.

Figure 3 depicts plots showing the results of quantitative MSP (qMSP) analysis of *BNC1* and *ADAMTS1* using real-time PCR. Figure 3 shows a statistically increased frequency of methylation when normal pancreas tissue is compared to invasive cancers. Figure 3 also shows that *BNC1* methylation could be detected at the earliest stages of pancreatic carcinogenesis (e.g. PanIN's) as compared to *ADAMTS1* methylation. Normal pancreas (n=4 normal tissues, n=10 surrounding normal tissues) and Pancreatitis samples (n=30), PanIN (n=20), and tumor (stage 2; n=12), respectively. Real-time MSP is shown as fold change for Methylated signal relative to Unmethylated signal. In MSP analysis, signals for unmethylated (U) and methylated (M) DNA are shown for each sample. Horizontal bar indicates mean. Figures 4A, 4B, and 4C show colony formation assays, Panc1 charts and bar graphs, and MIA-PaCa2 charts and bar graphs, respectively. Figure 4A shows colony formation by Panc1 and MIA-PaCa2 cells transfected with pcDNA3.1 or *BNC1*-pcDNA3.1 and grown for 2 weeks in medium containing G418. Results are plotted as the mean colony numbers relative to pcDNA3.1 transfectants in three independent experiments (Panc1; \*p=0.0275, MIA-PaCa2; \*p=0.0294). Figure 4B shows graphs that summarize cell proliferation assay in Panc1 cells, while Figure 4C shows analogous data for MIA-PaCa cells. Figures 4A and 4B are each divided into a right panel, a middle panel, and a left panel, which depict relevant data for Panc1 and MIA-PaCa cells, respectively. Right panel: *BNC1* transfected Panc1 cells (*BNC1*-pcDNA3.1) were compared with control cells transfected with empty vector (pcDNA3.1). Results are plotted as the mean cells number in three different independent experiments (Panel; \*p=0.0469; 72 hrs, \*\*p=0.0086; 96hrs: MIA-PaCa2, \*p=0.0469; 48hrs, \*p=0.0318; 72hrs, \*p=0.0389; 96hrs). Middle panel: Cell proliferation measured by 3H-thymidine incorporation (Panc1; \*p=0.0286, MIA-PaCa2; \*p=0.0256). Left panel: Invasion of Panc1 and MIA-PaCa2 cells through matrigel-coated transwells relative to control cells transfected with empty vector in three independent experiments (NS: not statistically significant).

## DETAILED DESCRIPTION OF THE INVENTION

**[0051]** The invention features methods that are useful for identifying a subject as having or having a propensity to develop neoplasia (pancreatic cancer, colon cancer, lung cancer).

**[0052]** The invention is based, at least in part, on the discovery that detection of *BNC1* and *ADAMTS1* promoter methylation was useful for identifying with high sensitivity the earliest stages of pancreas cancers. Notably, detection of *BNC1* and *ADAMTS1* promoter methylation in circulating DNA in serum was useful for the early detection of lung, colon and pancreatic cancer, especially in high risk individuals. Moreover, *BNC1* is a tumor suppressor gene in pancreatic cancer that is inactivated by promoter methylation.

**[0053]** A genome-wide transcriptome approach was used to identify new cancer specific DNA methylation alterations in pancreatic carcinoma. Methylation frequencies were analyzed for genes, *BNC1* and *ADAMTS1*, by methylation specific PCR and quantitative methylation specific PCR, as well as expression analysis by real-time PCR and immunohistochemistry. A novel nanoparticle-enabled Methylation On Beads technology was used to detect very early stage pancreatic

cancers. The biological role of *BNC1* gene was examined by colony formation, cell proliferation, and invasion assays in pancreatic cancer cell lines. This analysis led to the identification of *BNC1* (91.8%) and *ADAMTS1* (66.7%) as genes that showed a high frequency of methylation in pancreas cancer tissues (n=143). *BNC1* was frequently methylated in the earliest stages of pancreas carcinogenesis including carcinoma *in situ* or pancreatic intraepithelial neoplasia PanIN3 (100%) and Stage 1 invasive cancers (97.4%). Using the ultrasensitive nanoparticle-enabled MOB assay, these alterations were detected in serum samples from patients with pancreas cancer, with a sensitivity for *BNC1* of 79% (95% CI: 0.6-0.8) and for *ADAMTS1* of 48% (95% CI: 0.3-0.6) (n=42 cancers, Stages 1-4), while specificity was 88% for *BNC1* (95% CI: 0.6-0.9) and 92% for *ADAMTS1* (95% CI: 0.7-0.9) among 26 individuals without cancer. *BNC1* overexpression in pancreatic cancer cell lines showed suppressive effect by colony formation, and cell proliferation, but not invasion.

### Pancreatic Cancer

**[0054]** Pancreatic cancer is a deadly cancer with an overall 5-year survival rate of less than 5% and no improvements in survival over the last 3 decades. Pancreatic cancer currently ranks as the fourth leading cause of cancer related death in United States with an estimated 42,470 new cases and 35,240 deaths in 2009 and its incidence is rising. One of the major factors attributed to the dismal prognosis of pancreas cancer is the delayed diagnosis of the disease. Only about 10% of cases are amenable to potential curative surgical resection. However, long term 5-year survival is attainable in selected patients with early-stage pancreatic cancer who can undergo curative surgical resection. Early detection of pancreatic cancer is, therefore, thought to be the best modality for improving survival in this lethal disease. However, no screening test is available for detection of pancreatic cancer.

**[0055]** Pancreatic cancer is characterized by multiple genetic and epigenetic changes. In recent years, it has become apparent that pancreatic cancer is as much a disease of mis-regulated epigenetics, as it is a disease of genetic mutation. In particular, changes in DNA promoter methylation patterns likely play a crucial role in tumorigenesis and cancer progression. In order to address the need for both clinical diagnostics as well as therapeutics, many studies have employed DNA methylation of specific genes for application in diagnostics of multiple cancers. Such diagnostic tests can in principle be used for early detection of cancers, for assessing prognosis, and for therapeutics as predictors of response to therapy. Early detection of disease results in an improved clinical outcome for most types of cancer (Jemal A et al. (2011) Cancer Statistics, 2010. CA Cancer J Clin. Mar-Apr;61(2):133-4). Therefore, much effort has been invested in developing efficient screening technologies for early detection strategies.

### Cancer Screening

**[0056]** Ideally, methods for cancer screening should be easy to perform, cost-effective, noninvasive, and provide a benefit to patients. Current methods of screening for pancreatic cancer are inadequate. For example, endoscopic ultrasound has shown promise for identifying high risk patients, but requires access to specialized centers and is an expensive and invasive modality that needs to be repeated at frequent intervals. The present invention provides significant advantages over existing screening technologies for pancreatic and other cancers. Significantly, the invention provides methods for detecting *BNC1* and *ADAMTS1* methylation. Increased methylation in *BNC1* and *ADAMTS1* is associated with all stages of pancreatic cancer. *BNC1* gene was methylated at 100% frequency in PanIN3, which is felt to be the penultimate step prior to development of invasive carcinoma.

**[0057]** Interestingly, *BNC1* showed a potential tumor suppressive role as measured by suppression of colony formation, cell proliferation, and invasion in pancreatic cancer cells. Methylation of *BNC1* and *ADAMTS1* showed high sensitivity and specificity for detecting pancreatic cancer in sera using nano-enabled methylation based technology termed Methylation-on-Beads (MOB). The results reported herein below indicate that methylation-based screening is useful for identifying subjects at risk of not only pancreatic cancer, but also, lung and colon cancers using a non-invasive inexpensive modality.

**[0058]** While the methods of the invention are suitable for screening the population at large, it is particularly useful for screening subjects identified as at increased risk for having a neoplasia. Subjects identified as having an increased risk of neoplasia (pancreatic cancer, lung cancer, colon cancer) include but are not limited to smokers and subjects having a *Brca1* or *Brca2* mutation. Subjects identified as having an increased risk of pancreatic cancer include subjects identified as having pancreatic cysts or pancreatitis, as well as patients that have a family history of pancreatic cancer, particularly in 1 or 2 relatives.

### Types of biological samples

**[0059]** The level of promoter methylation in each of the genes identified herein (i.e *BNC1* and *ADAMTS1*) can be measured in different types of biologic samples. In one embodiment, the biologic sample is a blood, plasma, or serum sample. In another embodiment, the sample is a tissue sample that includes cells of a tissue or organ (e.g., pancreatic

cells, cells of a pancreatic cyst or pancreas lesion, lung cells, and colon cells). Pancreatic tissue is obtained, for example, from a biopsy of the pancreas. In another embodiment, the biologic sample is a biologic fluid sample (e.g., blood, blood serum, plasma, urine, stool, pancreatic cyst fluid, fluid from the major/minor pancreatic duct (i.e., "pancreatic juice") lung lavage, stool, sputum, or any other biological fluid useful in the methods of the invention).

### Methylation-on-Beads

**[0060]** In brief, Methylation-on-Beads is a single-tube method for polynucleotide extraction and bisulfite conversion that provides a rapid and highly efficient method for DNA extraction, bisulfite treatment and detection of DNA methylation using silica superparamagnetic particles (SSP). All steps are implemented without centrifugation or air drying that provides superior yields relative to conventional methods for DNA extraction and bisulfite conversion. SSP serve as solid substrate for DNA binding throughout the multiple stages of each process. Specifically, SSP are first used to capture genomic DNA from raw biological samples, processed biological samples or cultured cells. Sodium bisulfite treatment is then carried out in the presence of SSP without tube transfers. Finally, the bisulfite treated DNA is analyzed to determine the methylation status. DNA extraction yield was found to be 5-20 times the yield from conventional extraction. 90% of the input DNA was recovered after bisulfite treatment. In addition, Methylation-on-Beads total process time was completed in less than 6 hours when compared to 3 days for conventional methods. Hence, Methylation-on-Beads allows for convenient, efficient and contamination-resistant methylation detection in a single tube or other reaction platform. Methods for carrying out methylation-on-beads are known in the art, and described, for example, in PCT/US2009/000039.

### Methylation-specific quantum dot FRET

**[0061]** If desired, methods of the invention may be advantageously combined with methylation-specific quantum dot fluorescence resonance energy transfer (MS-qFRET). See, for example, PCT/US2009/000039. MS-qFRET provides for the qualitative and quantitative detection of methylated DNA, as well as for the detection of low-abundance methylated DNA. In this technique, quantum dots are used to capture methylation-specific PCR (MSP) amplicons and to determine the methylation status via fluorescence resonance energy transfer (FRET). Desirably, MS-qFRET has low intrinsic background noise, high resolution and high sensitivity. MS-qFRET detects as little as 15 pg of methylated DNA in the presence of a 10,000-fold excess of unmethylated alleles, enables reduced use of PCR (8 cycles), and allows for multiplexed analyses.

**[0062]** More specifically, bisulfite-treated DNA is amplified through PCR, wherein the forward primer is biotinylated and the reverse primer is labeled with an organic fluorophore. Next, streptavidin-conjugated quantum dots (QDs) are introduced to capture the labeled PCR products via streptavidin-biotin binding, bringing the QDs (serving as donors) and fluorophores (serving as acceptors) in close proximity allowing FRET to occur. Finally, PCR products are detected by emissions of fluorophores accompanied by quenching of QDs. Spectral information is processed to determine the level of DNA methylation. Fluorescence responses are measured using a fluorospectrometer.

**[0063]** PCR with labeled primers is run. Products are then subject to PCR purification (Qiagen Corporation) in order to recover PCR product that is free of primers, primer-dimers, Taq and dNTPs. For conjugating with quantum dots (Invitrogen Corporation), 1  $\mu$ L of 100 mM NaCl is mixed with 7  $\mu$ L PCR mix. 1  $\mu$ L of deionized (DI) H<sub>2</sub>O is added to this mix. Finally, 1  $\mu$ L of 1 nM QD is added and the mixture is left undisturbed for 15 minutes.

**[0064]** Mixtures of defined methylation levels ranging from 100%, 75%, 50%, 25%, and 1% of the total 150 ng input DNA are obtained. To quantify the level of methylation, a "q-score": a score that is based on the normalized FRET efficiencies of acceptor and donor emission in MS-qFRET is defined. In any FRET process, as the level of the acceptor emission increases, the decay of donor emission increases as well. The FRET efficiency can then be calculated based

on the proximity ratio formalism,  $E = \frac{I_A}{I_A + I_D}$  ( $I_D$  and  $I_A$  corresponding to donor and acceptor intensity). Further, the

q-score was determined by normalizing the calculated E for the DNA mixture to an appropriate concentration of IVD only as a methylated control (q-score = 1) and NL only and as an unmethylated control (q-score = 0). By including positive and negative controls in every assay a standard curve is created in order to quantify and compare methylation levels of unknown samples using low-amplification cycles.

### Diagnostic assays

**[0065]** The present invention provides a number of diagnostic assays that are useful for the identification or characterization of a neoplasia (pancreatic cancer, lung cancer, colon cancer). In one embodiment, the neoplasia is characterized by quantifying or determining the methylation level of the following promoters: *BNC1* and *ADAMTS1* in the neoplasia. In one embodiment, methylation levels are determined using quantitative methylation specific PCR (QMSP) to detect

CpG methylation in genomic DNA. QMSP uses sodium bisulfate to convert unmethylated cytosine to uracil. A comparison of sodium bisulfate treated and untreated DNA provides for the detection of methylated cytosines.

**[0066]** While the examples provided below describe methods of detecting methylation levels using QMSP, the skilled artisan appreciates that the invention is not limited to such methods. Methylation levels are quantifiable by any standard method, such methods include, but are not limited to real-time PCR, Southern blot, bisulfite genomic DNA sequencing, restriction enzyme-PCR, MSP (methylation-specific PCR), methylation-sensitive single nucleotide primer extension (MS-SNuPE) (see, for example, Kuppuswamy et al., Proc. Natl Acad. Sci. USA, 88, 1143-1147, 1991), DNA microarray based on fluorescence or isotope labeling (see, for example, Adorján Nucleic Acids Res., 30: e21 and Hou Clin. Biochem., 36:197-202, 2003), mass spectroscopy, methyl accepting capacity assays, and methylation specific antibody binding. See also U.S. Patent Nos.: 5,786,146, 6,017,704, 6,300,756, and 6,265,171.

**[0067]** The primers used for amplification of the CpG-containing nucleic acid in the specimen, after bisulfite modification, specifically distinguish between untreated or unmodified DNA, methylated, and non-methylated DNA. Methylation specific primers for the non-methylated DNA preferably have a T in the 3' CG pair to distinguish it from the C retained in methylated DNA, and the complement is designed for the antisense primer. Methylation specific primers usually contain relatively few Cs or Gs in the sequence since the Cs will be absent in the sense primer and the Gs absent in the antisense primer (C becomes modified to U(uracil) which is amplified as T(thymidine) in the amplification product).

**[0068]** The primers embrace oligonucleotides of sufficient length and appropriate sequence so as to provide specific initiation of polymerization on a significant number of nucleic acids in the polymorphic locus. Specifically, the term "primer" as used herein refers to a sequence comprising two or more deoxyribonucleotides or ribonucleotides, preferably more than three, and most preferably more than 8, which sequence is capable of initiating synthesis of a primer extension product, which is substantially complementary to a polymorphic locus strand. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the inducing agent for polymerization. The exact length of primer will depend on many factors, including temperature, buffer, and nucleotide composition. The oligonucleotide primer typically contains between 12 and 27 or more nucleotides, although it may contain fewer nucleotides. Primers are designed to be "substantially" complementary to each strand of the genomic locus to be amplified and include the appropriate G or C nucleotides as discussed above. This means that the primers must be sufficiently complementary to hybridize with their respective strands under conditions that allow the agent for polymerization to perform. In other words, the primers should have sufficient complementarity with the 5' and 3' flanking sequences to hybridize therewith and permit amplification of the genomic locus. While exemplary primers are provided herein, it is understood that any primer that hybridizes with the target sequences of the invention are useful in the method of the invention for detecting methylated nucleic acid.

**[0069]** In one embodiment, methylation specific primers amplify a desired genomic target using the polymerase chain reaction (PCR). The amplified product is then detected using standard methods known in the art. In one embodiment, a PCR product (*i.e.*, amplicon) or real-time PCR product is detected by probe binding. In one embodiment, probe binding generates a fluorescent signal, for example, by coupling a fluorogenic dye molecule and a quencher moiety to the same or different oligonucleotide substrates (*e.g.*, TaqMan® (Applied Biosystems, Foster City, CA, USA), Molecular Beacons (see, for example, Tyagi et al., Nature Biotechnology 14(3):303-8, 1996), Scorpions® (Molecular Probes Inc., Eugene, OR, USA)). In another example, a PCR product is detected by the binding of a fluorogenic dye that emits a fluorescent signal upon binding (*e.g.*, SYBR® Green (Molecular Probes)). Such detection methods are useful for the detection of a methylation specific PCR product.

**[0070]** The methylation level of *BNC1* and *ADAMTS1* promoters described herein defines the methylation profile of a neoplasia. The level of methylation present at any particular promoter is compared to a reference. In one embodiment, the reference is the level of methylation present in a control sample obtained from a patient that does not have a neoplasia. In another embodiment, the reference is a baseline level of methylation present in a biologic sample derived from a patient prior to, during, or after treatment for a neoplasia. In yet another embodiment, the reference is a standardized curve.

**[0071]** The methylation level of the promoters described herein (*BNC1* and *ADAMTS1*) is used, alone or in combination with other standard methods, to characterize the neoplasia. For example, methods for detecting *BNC1* and *ADAMTS1* promoter methylation may be carried out prior to or concurrently with testing for alterations in other biomarkers, such as *BRCA1*, *p16*, *K-ras*, *APC*, *PalB2*, *DPC4*, *EGFR*, *EML- ALK4* gene or polypeptide. In one embodiment, the alteration in a *Brca1*, *p16*, *K-ras*, *APC*, *PalB2*, *DPC4*, *EGFR*, *EML- ALK4* gene is a mutation in the sequence of the gene relative to a reference. In other embodiments, the alteration is in the level of expression or activity of the *Brca1*, *p16*, *K-ras*, *APC*, *PalB2*, *DPC4*, *EGFR*, or *EML- ALK4* gene or polypeptide. Once a subject is identified as having increased *BNC1* and *ADAMTS1* promoter methylation, further diagnostic testing may be carried out to localize the pathology to a particular organ or organ system. Typically, after the subject has been identified as having increased promoter methylation in *BNC1* and *ADAMTS1*, imaging studies are carried out. Such studies include, but are not limited to, endoscopic ultrasound, MRI, CT scan, and PET scan.

**[0072]** If desired, diagnostic methods of the invention can be combined with conventional diagnostic methods used to determine the stage or grade of a neoplasia. Grading is used to describe how abnormal or aggressive the neoplastic

cells appear, while staging is used to describe the extent of the neoplasia. The grade and stage of the neoplasia is indicative of the patient's long-term prognosis (*i.e.*, probable response to treatment and survival). Thus, the methods of the invention are useful for predicting a patient's prognosis, and for selecting a course of treatment.

5 [0073] In conventional diagnostic methods, a pathologist will view a tissue sample from the tumor and determine the grade based on the degree of pathology observed. Typically, pancreatic tumors are graded from 1-4. For a grade I tumor, cells present in the sample are most similar to normal pancreatic cells. Grade 4 samples contain cells that are most dissimilar to normal pancreatic cells. High-grade neoplasias are the most deadly because they are most aggressive and fast growing. High-grade neoplasias typically move rapidly into surrounding tissues, such as lymph nodes and bones.

10 [0074] Stage refers to the extent of a cancer. In pancreatic cancer, for example, one staging method divides the cancer into four categories, stage I pancreatic cancer is found only in the pancreas itself or has started to spread to the tissues next to the pancreas (such as the small intestine, the stomach, or the bile duct). Stage II pancreatic cancer has spread to nearby organs such as the stomach, spleen, or colon, but has not entered the lymph nodes. Stage III pancreatic cancer has spread to the lymph nodes near the pancreas. Cancer may have spread to nearby organs. Stage IV pancreatic cancer has spread to organs near the pancreas (stage IVA) or to organs far away from the pancreas (stage IVB). Stage 15 IVA pancreatic cancer has spread to organs that are near the pancreas (such as the stomach, spleen, or colon) but has not spread to distant organs (such as the liver or lungs). Stage IVB pancreatic cancer has spread to distant organs (such as the liver or lungs).

### 20 Selection of a treatment method

[0075] Identifying the presence of increased promoter methylation in *BNC1* and *ADAMTS1*, indicates that the subject likely has an invasive cancer or a pre-cancerous lesion (*e.g.*, cancer-in-situ). Further diagnostic testing may be carried out to localize the pathology to a particular organ or organ system. For example, endoscopic ultrasound, MRI, CT scan, PET scan,, bronchoscopy, colonoscopy, esophagogastroduodenoscopy, laparoscopic surgery to localize a lesion or 25 any other modality known in the art may be used to characterize the neoplasia or pre-cancerous lesion (*e.g.*, PanIN's, or precursor lung conditions such as dysplasia, carcinoma in situ, atypical adenomatous hyperplasia or precursor colonic lesions such as adenomas). Pre-cancerous lesions are likely to be susceptible to conservative treatment methods. Conservative treatment methods include, for example, cancer surveillance, which involves periodic patient monitoring using diagnostic assays of the invention, alone or in combination, with diagnostic imaging or chemoprevention.

30 [0076] More aggressive neoplasias are less susceptible to conservative treatment methods. For aggressive neoplasias, an aggressive method of treatment should be selected. Aggressive therapeutic regimens typically include one or more of the following therapies: surgical resection, radiation therapy, and chemotherapy. After a subject is diagnosed as having a neoplasia (*e.g.*, prostate cancer, lung cancer, colon cancer) a method of treatment is selected. Where the methods of the invention identify a subject as having a neoplasia associated with increased promoter methylation in *BNC1* and 35 *ADAMTS1*, therapy to reduce this promoter methylation may be selected. For example, epigenetic therapy may be selected to relieve promoter methylation in *BNC1* and *ADAMTS1*. Such epigenetic therapy may involve the use of histone deacetylase inhibitors or methylation inhibitors. Exemplary agents include, but are not limited to, entinostat, SAHA (suberoylanilide hydroxamic acid), depsipeptide (Fujisawa Pharmaceuticals), azacytidine, deazacytidine, romidepsin (Istodax), Vorinostat, polyamine analogues, zebularine and other novel emerging drugs targeting methylation, histone 40 changes or the entire polycomb complex. Epigenetic therapy may be combined with more conventional therapies.

### Patient monitoring

45 [0077] The diagnostic methods of the invention are also useful for monitoring the course of a pre-cancerous lesion in a patient or for assessing the efficacy of a therapeutic regimen. In one embodiment, the diagnostic methods of the invention are used periodically to monitor the methylation levels of *BNC1* and *ADAMTS1*. In one example, the neoplasia is characterized using a diagnostic assay of the invention prior to administering therapy. This assay provides a baseline that describes the methylation level of one or more promoters or the methylation profile of the neoplasia prior to treatment. Additional diagnostic assays are administered during the course of therapy to monitor the efficacy of a selected therapeutic 50 regimen. A therapy is identified as efficacious when a diagnostic assay of the invention detects a decrease in methylation levels at one or more promoters relative to the baseline level of methylation.

### Kits

55 [0078] Described are kits for the diagnosis or monitoring of a neoplasia in a biological sample obtained from a subject. In various embodiments, the kit includes at least one primer or probe whose binding distinguishes between a methylated and an unmethylated *BNC1* and *ADAMTS1* promoter sequence, together with instructions for using the primer or probe to identify a neoplasia. In another embodiment, the kit further comprises a pair of primers suitable for use in a polymerase

chain reaction (PCR). In yet another embodiment, the kit further comprises a detectable probe. In yet another embodiment, the kit further comprises a pair of primers capable of binding to and amplifying a reference sequence. In yet other embodiments, the kit comprises a sterile container which contains the primer or probe; such containers can be boxes, ampoules, bottles, vials, tubes, bags, pouches, blister-packs, or other suitable container form known in the art. Such containers can be made of plastic, glass, laminated paper, metal foil, or other materials suitable for holding nucleic acids. The instructions will generally include information about the use of the primers or probes described herein and their use in diagnosing a neoplasia. Preferably, the kit further comprises any one or more of the reagents described in the diagnostic assays described herein. In other embodiments, the instructions include at least one of the following: description of the primer or probe; methods for using the enclosed materials for the diagnosis of a neoplasia; precautions; warnings; indications; clinical or research studies; and/or references. The instructions may be printed directly on the container (when present), or as a label applied to the container, or as a separate sheet, pamphlet, card, or folder supplied in or with the container.

**[0079]** The following examples are offered by way of illustration, not by way of limitation.

**[0080]** The practice of the present invention employs, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry and immunology, which are well within the purview of the skilled artisan. Such techniques are explained fully in the literature, such as, "Molecular Cloning: A Laboratory Manual", second edition (Sambrook, 1989); "Oligonucleotide Synthesis" (Gait, 1984); "Animal Cell Culture" (Freshney, 1987); "Methods in Enzymology" "Handbook of Experimental Immunology" (Weir, 1996); "Gene Transfer Vectors for Mammalian Cells" (Miller and Calos, 1987); "Current Protocols in Molecular Biology" (Ausubel, 1987); "PCR: The Polymerase Chain Reaction", (Mullis, 1994); "Current Protocols in Immunology" (Coligan, 1991). These techniques are applicable to the production of the polynucleotides and polypeptides of the invention, and, as such, may be considered in making and practicing the invention. Particularly useful techniques for particular embodiments will be discussed in the sections that follow.

## EXAMPLES

### Example 1: Identification of cancer-specific promoter methylation candidate genes in pancreas hypermethylome

**[0081]** In order to identify new DNA methylation biomarkers for pancreatic cancer, previously established whole human transcriptome microarray screening was used to identify genes silenced by promoter hypermethylation. To identify global hypermethylation-dependent gene expression changes in pancreatic cancer, a genome-wide expression array-based approach was performed in four different human pancreatic cancer cell lines (PL45, MIA-PaCa2, Panc1, and Capan 1) with a pharmacologic strategy using 5-aza-deoxycytidine and Trichostatin A (TSA) using standard array-based methodology (e.g. Agilent Technologies 44K). As seen in Figure 1A, a total of 1,427 unique genes were initially identified in four cell lines, which met the criteria for candidate hypermethylated genes in the pancreatic cancer cell lines.

**[0082]** In order to identify genes that only showed cancer specific methylation, the following experimental validation criteria was used: 1) the gene was expressed in normal pancreas tissue, 2) the gene displayed no/low methylation in normal pancreatic tissues (cancer-free tissues), and 3) the gene was frequently methylated in pancreas cancer cell lines, and methylation was also seen in primary tumors. Based on these criteria, 8 genes (*TFPI2*, *ASCL2*, *BNC1*, *TWIST1*, *BNIP3*, *ADAMTS1*, *PNMT*, and *EVL*) were identified that displayed cancer-specific methylation in pancreatic cancers.

### Example 2: *BNC1* and *ADAMTS1*-DNA methylation biomarkers for early detection of pancreatic cancer

**[0083]** Next, the methylation status of these 8 genes (*TFPI2*, *ASCL2*, *BNC1*, *TWIST1*, *BNIP3*, *ADAMTS1*, *PNMT*, and *EVL*) was examined in a large series of primary pancreatic tumor samples (n=143; Stages 1-4) using methylation specific PCR (see Table 1).

**Table 1.** Clinical information of pancreas cancer patient primary tumor samples in this study

	Pancreas Primary tumors					
	Normal (n=4)	PanINs (N=20)	Stage 1 (N=38)	Stage 2 (N=78)	Stage 3 (N=5)	Stage 4 (N=2)
<b>Median Survival (months)</b>		Not Met	43.0	18.0	14.1	19.6
<b>Gender</b>						
<b>Male</b>	3(75%)	9 (45%)	9(23.7%)	44 (57%)	1 (20%)	2 (100%)
<b>Femal</b>	1(25%)	11 (55%)	29(76.3%)	34 (43%)	4 (80%)	0.0%

(continued)

		Pancreas Primary tumors					
<u>Grade</u>		Normal (n=4)	PanINs (N=20)	Stage 1 (N=38)	Stage 2 (N=78)	Stage 3 (N=5)	Stage 4 (N=2)
5	Well-Moderately		9 (45%)	3 (7.9%)	4 (5.1%)	1 (20%)	0(0%)
	Poorly		8 (40%)	23 (60.5%)	41 (52%)	4 (40%)	1 (50%)
10			3 (15%)	12 (31.6%)	33 (42.9%)	4 (40%)	1 (50%)

[0084] As shown in Figure 1B, the most frequently methylated gene was *BNC1* (90.5%), followed by *ADAMTS1* (66.7%), *TWIST1* (66.7%), *ASCL2* (53.79%), *BNIP3* (49%), *TFPI2* (53.7%), *EVL* (46.3%), and *PNMT* (26.5%). Interestingly, two of the genes that demonstrated frequent methylation in this cohort of primary pancreatic cancer samples, *BNC1* (90.5%) and *ADAMTS1* (66.7%), also showed frequent methylation in the precursor lesions of pancreatic cancer, pancreatic intraepithelial neoplasia (PanIN) lesion. Specifically, *BNC1* methylation frequency was 70% (n=14/20) and *ADAMTS1* methylation frequency was 25% (n=5/20).

[0085] The ability to identify high grade precursor lesions such as PanIN3 (or pancreatic carcinoma *in situ*) represents one of the best means for achieving early detection of pancreatic cancer, and facilitating the implementation of chemoprevention strategies (see Figure 1C). *BNC1* was methylated in PanIN 1 (56%), PanIN 2 (75%), and PanIN 3 (100%) during PanIN progression. This data indicated that DNA methylation of *BNC1* gene is an early event in pancreatic carcinogenesis and that the prevalence of methylation increases with pancreatic cancer progression.

[0086] This genome-wide DNA hypermethylome screening is based on a pharmacological approach using 5-deoxy-azacytidine and TSA, which has shown an excellent correlation between promoter DNA methylation and inhibition of gene expression and re-expression after treatment with a DNA demethylating drug, such as 5-deoxy-azacytidine, in prior studies on breast and colorectal cancer. Here, promoter methylation of *BNC1* and *ADAMTS1* was tested, and correlated with a lack of gene expression in pancreatic cancer cell lines (Panc1, MIA-PaCa2, Capan1, and PL45) using quantitative RT-PCR (qRT-PCR) (Figure 2A, upper panel). Both genes showed virtually undetectable (or no/low) endogenous gene expression, and significant re-expression after 5-deoxy-azacytidine treatment. As shown in Figure 2A, lower panel, treatment with a histone deacetylase inhibitor, such as TSA, resulted in minimal re-expression, except in the case of PL45, which may be regulated both by promoter DNA methylation and histone modifications. Promoter-associated CpG island methylation in the *BNC1* promoter was detected by bisulfite sequencing analysis in pancreatic cancer cell line (Panc1), a primary pancreas cancer sample (stage 2), and normal pancreatic tissue. As shown in Figure 2B, bisulfite sequencing analysis showed dense methylation in the pancreatic cancer cell line and the primary pancreatic cancer and minimal/no methylation in normal pancreas samples consistent with the methylation specific PCR analysis.

### Example 3: Detection of DNA methylation in pancreatic cancer patient sera using methylation on beads technology

[0087] Early detection of disease results in an improved clinical outcome for most types of cancer. This is particularly relevant for pancreatic cancer, which is poorly accessible and difficult to diagnose. Consequently, 90% of patients with pancreatic cancer present with metastatic or advanced regional disease. Based on the methylation studies, *BNC1* and *ADAMTS1* have emerged as important biomarkers for the early detection of pancreatic cancer. To determine whether *BNC1* and *ADAMTS1* promoter DNA methylation could be detected in pancreatic cancer patient sera, a highly-sensitive nanoenabled assay, termed methylation on Beads was used, which was developed to improve sensitivity of methylation detection (see, e.g., Figure 2C). To test the utility of methylation on beads as a tool for early detection of pancreatic cancers, serum samples from a series of pancreatic cancer patients (n=42; stages 1-4), as well as a panel of normal healthy serum samples (n=26) was tested. Sensitivity was determined based on the assumption that all patients with cancer would be methylated for both genes, while all healthy normal volunteers would be unmethylated for both genes. Methylation was analyzed in 42 serum samples from pancreatic cancer patients and 26 normal volunteers. 33 of the 42 pancreatic cancer patients showed methylation for *BNC1*, while 20 of the 42 showed methylation for *ADAMTS1* (see Table 2).

[0088] Overall, for all stages included, the sensitivity for *BNC1* and *ADAMTS1* was determined to be 78.6% and 47.6%, respectively. Sensitivity of detection of stage I pancreatic cancers was 90% for both genes. Amongst the 26 normal serum samples, 3 of the normal volunteers showed methylation for *BNC1* while 2 showed methylation for *ADAMTS1*. Specificity of detection was determined to be 88.4% for *BNC1* and 92.4% for *ADAMTS1*. Computed results have been presented using a 95% confidence interval and are presented in detail in Table 2. Results in the corresponding tumor

were higher for both genes, with 100% sensitivity and specificity for *BNC1* and 79.2% sensitivity and 100% specificity for *ADAMTS1*.

[0089] Additionally, the sensitivity and specificity of *BNC1* and *ADAMTS1* was also tested in other types of cancer, including both colon cancer and lung cancer (Table 2). This data showed that *BNC1* was also a useful biomarker for the detection of early stages of colon cancer. For example, *BNC1* showed a sensitivity of 84.6% (n=11/13) for stage 1 colon cancers and advanced adenomas. Similarly, *BNC1* also showed good sensitivity for the detection of early stage lung cancer as stage 1 lung cancer was detected with a sensitivity of 75% (n=12/16).

**Table 2. Sensitivity and specificity of *BNC1* and *ADAMTS1* genes in various cancer patient sera samples**

		<i>BNC1</i>		<i>ADAMTS1</i>	
<b>Cancer Types:</b>					
<b><i>Pancreatic Cancer</i></b>					
		<b>Sensitivity(%)</b>			
Stage	N	Estimated Value	95% CI	Estimated Value	95% CI
I	10	90.0% (9/10)		90.0% (9/10)	
II-IV	32	75.0% (24/32)		34.4% (11/32)	
Total	42	78.6% (33/42)	60-90%	47.6% (20/42)	40-70%
<b><i>Colon Cancer</i></b>					
		<b>Sensitivity(%)</b>			
Stage		Estimated Value	95% CI	Estimated Value	95% CI
Adenomas, stage I	13	84.6% (11/13)		61.5% (8/13)	
II-III	15	66.7% (10/15)		73.3% (11/15)	
Other cancer*	2	100.0% (2/2)		50.0% (1/2)	
Total	30	76.7% (23/30)	70-90%	66.7% (20/30)	40-70%
<b><i>Lung Cancer</i></b>					
		<b>Sensitivity (%)</b>			
Stage		Estimated Value	95% CI	Estimated Value	95% CI
I	16	75.0 % (12/16)		62.5% (10/16)	
II-III	4	100.0% (4/4)		100.0% (4/4)	
Total	20	80.0 % (16/20)	60-90%	70.0% (14/20)	50-90%
<b>Normal</b>					
<b>Specificity</b>	26	88.4%	70-90%	92.4%	70-90%

CI:

Confidential Interval.

\*other colorectal malignancies.

[0090] A subset of tissue samples from patients diagnosed with non-cancerous pancreatic diseases, such as pancreatitis, was collected and analyzed. Patients with chronic pancreatitis maybe at increased risk for pancreatic cancer and amongst patients with pancreatic cancer, 5% or more of patients have underlying chronic pancreatitis. Chronic inflammatory conditions likely increases the frequency of methylation as a field defect which may then increase risk of subsequent malignancy. The methylation of *BNC1* and *ADMATS1* DNA was compared between different conditions (normal, pancreatitis, PanIN, and invasive cancers) using quantitative methylation specific PCR analysis. As shown in Figure 3, *BNC1* and *ADAMTS1* showed statistically increased frequency of methylation when comparing normal pancreas tissues and invasive cancers (p<0.0001; both *BNC1* and *ADAMTS1*), as well as chronic pancreatitis and invasive cancers (p<0.0001, both *BNC1* and *ADAMTS1*). There was a low frequency of *BNC1* and *ADAMTS1* methylation present in non-cancerous disease, such as pancreatitis. More interestingly, there was a significant quantitative difference between PanINs and invasive cancers on both genes (p<0.001 both *BNC1* and *ADAMTS1*). However, *BNC1* methylation could be detected in the earliest stages of pancreatic carcinogenesis such as PanIN's, compared to *ADAMTS1* methylation (see Figure 3), indicating that *BNC1* methylation is a very early event, and that *ADMATS1* methylation is a later event during pancreas cancer development.

**Example 4: Tumor suppressive effects of *BNC1* gene over-expression in pancreatic cancer cells**

[0091] According to the methylation data in pancreatic cancer patients, *BNC1* is a useful DNA methylation biomarker

in pancreatic cancer patients. To determine whether *BNC1* played a role as a tumor suppressor in pancreatic cancer cells, *in vitro* colony formation assays were performed to determine the effects of full-length *BNC1* transfected into Panc1 and MIA-PaCa2 cells lacking *BNC1* expression. Compared with control cells transfected with empty vector, over-expression of a full-length *BNC1* gene induced a nearly 2.2-fold (Panc1) and 9-fold (MIA-PaCa2) reduction of G418-resistant colonies (see Figure 4A). In addition, there was a 75% (Panc1) and 81% (MIA-PaCa2) decrease in cell proliferation as measured by [<sup>3</sup>H] thymidine activity (Figure 4B and 4C; left and middle panel). However, overexpression of *BNC1* gene had no effect on Panc1 and MIA-PaCa2 cells migration and invasion through matrigel-coated transwell membranes compared with control cells transfected with empty vector (Figure 4B and 4C; right panel). Taken together, these data indicated that *BNC1* gene has tumor-suppressive effects in human pancreatic cancer cells.

[0092] In this study, genome-wide gene expression profiling using a pharmacological approach (5-deoxy-azacytidine and TSA) was used in pancreatic cancer cell lines, a platform which has been used to identify an early detection marker for colorectal cancer. Defining the DNA hypermethylome has been useful as it has allowed the identification of not only novel DNA methylation biomarker candidates, but also of tumor suppressor gene candidates in many types of cancers (colon, breast, ovarian, etc). The identified DNA methylation biomarkers will facilitate diagnostics, and also contribute to therapeutics as predictors of response to therapy.

[0093] The poor accessibility of the pancreas along with the late presentation of symptoms has thwarted attempts at timely detection of malignancy, and contributed to the high mortality rates of pancreatic cancer, which is the fourth leading cause of cancer death in both men and women. Therefore, the development of cancer biomarkers of pancreatic cancer is the best hope for early detection.

[0094] As reported herein above, the promoter methylation of *BNC1* and *ADAMTS1* is an excellent early detection biomarker for pancreatic cancer. *BNC1* and *ADAMTS1* have not been described as a DNA methylation biomarker in pancreatic cancers, although they have been reported methylated in lung cancers.

[0095] Screening with a nanobased high sensitive technology allowed the sensitive and specific detection of pancreatic cancer in its earliest stages. Using *BNC1* and *ADAMTS1*, provided for the detection of very early stages of pancreatic carcinoma, with an overall sensitivity of 83.7% (95%CI; 74.2- 90.3%) and a specificity of 84.6% (95%CI; 64.3-95.0%). Significantly, the sensitivity achieved in this study is higher than previously reported for serum hypermethylation markers. The present invention provides a cost-effective approach for screening individuals identified as having risk factors for pancreatic cancer in the general population. Additionally, the nano-based methylation on beads detection method significantly reduced the quantity of serum necessary for analysis. The invention provides a paradigm for widespread screening for pancreatic cancers using a simple blood test.

[0096] In conclusion, this is the first study to describe the utility of *BNC1* and *ADAMTS1* promoter methylation as biomarkers in pancreatic cancer patient serum using nanoparticle-enabled technology. In addition, this is the first report to demonstrate a tumor suppressor role for *BNC1* in pancreatic cancer. These data indicate that *BNC1* promoter methylation is useful as a sensitive and specific noninvasive pre-selection modality for diagnosing subjects as having cancer and identifying individuals at risk for pancreatic cancer.

[0097] The results presented hereinabove were carried out using the following methods and materials.

#### Gene Expression Microarray Analysis

[0098] Total RNA was harvested from log phase cells using standard methods (e.g. with TRIzol (Invitrogen™) and the RNeasy kit (Qiagen™) according to the manufacturer's instructions), including a DNase digestion step. RNA was quantified using a spectrophotometer (e.g. the NanoDrop™ ND-100 (<http://www.nanodrop.com/>)) followed by quality assessment with a microfluidics analysis platform (e.g. the Agilent™ 2100 Bioanalyzer (Agilent Technologies, <http://www.agilent.com/>)). RNA concentrations for individual samples were greater than 200ng/11, with 28S/18S ratios greater than 2.2 and RNA integrity of 10 (10 scored as the highest). Sample amplification and labeling procedures were carried out using standard methods (e.g. Low RNA Input Fluorescent Linear Amplification Kit (Agilent Technologies™) according to the manufacturer's instructions). The labeled cRNA was purified using spin columns (e.g. the RNeasy mini kit (Qiagen™)) and quantified. RNA spike-in controls (Agilent Technologies™) were added to RNA samples before amplification. Samples (0.75 1g) labeled with Cy3 or Cy5 were mixed with control targets (Agilent Technologies™), assembled on Oligo Microarray, hybridized, and processed according to the Agilent microarray protocol. Scanning was performed with a microarray scanner (Agilent™ G2565BA, using recommended settings). Data analysis was conducted as previously reported (Schuebel K, Chen W, Cope L, Glöckner SC, Suzuki H, et al., Comparing the DNA hypermethylome with gene mutations in human colorectal cancer.2007, PLOS Genetics, 3(9):1709-1723..

#### Patient samples and study population

[0099] Patient samples were prepared from formalin-fixed, paraffin-embedded (FFPE) tissue samples from patients with pancreatic cancer procured from the pathology archives of the Johns Hopkins Hospital in accordance with all rules

and regulations of the Institutional Review Board (IRB) and as per HIPAA compliance. A total of 167 patients were analyzed in the current study. The Johns Hopkins cohort consisted of 143 tissue samples from pancreatic cancer patients with stage 1 through stage 4 disease who underwent primary surgery and other treatment for pancreatic cancer at the JHH from 1998 to 2009 (Median follow up of 6.4 years). Patients in this cohort were similar by stage with respect to gender, proportion of cases with lymphovascular invasion, and pathologic grade (Table 1). An additional 20 FFPE tissue samples were obtained from patients who had undergone surgical resection but diagnosed with precancerous lesions, pancreatic intraepithelial neoplasias (PanINs). Finally, 4 FFPE tissue samples were obtained from pathology archives from patients who had undergone pancreatic resection with no abnormality noted on gross or microscopic examination.

#### Pancreatic cancer patient serum samples

[0100] Patient serum samples were obtained from individuals with pancreatic cancer prior to undergoing surgical treatment at the JHH after obtaining informed consent from 2007 to 2009. Matching tumor samples were drawn from the pathology archives of the Johns Hopkins Hospital in accordance with all rules and regulations of the Institutional Review Board (IRB) and as per HIPAA compliance. A total of 36 serum samples were tested, 23 of which had matching FFPE tissue available. Patient serum samples were also obtained from individuals with pancreas (42 samples, Stages 1-4), colon (30 samples, Stages 1-3 and adenomas) and lung cancers (20 samples, Stages 1-3) prior to undergoing surgical treatment at the JHH after obtaining informed consent. Additionally, 26 serum samples were obtained from normal, healthy volunteers to serve as controls.

#### Cell culture and treatment

[0101] Cancer cell lines (Pancreas cancer cell lines; Panc1, MIA-PaCa2, PL45, Capan1, Bxpc3, Capan2, PK9, Aspc1, PK8, and PL3) were obtained from ATCC and cultured in appropriate media and under conditions described by ATCC, with media obtained from InVitrogen™, supplemented with 10% fetal bovine serum (Gemini Bio-Products) and 1% penicillin/streptomycin (InVitrogen™). DKO cells (HCT116 colon cancer cells with genetic disruption of *DNMT1* and *DNMT3b*) were cultured as described previously (Rhee et al., 2002, Nature).

#### DNA Methylation analyses and RT-PCR for expression analysis

[0102] Primer pairs were preferentially designed near the putative transcriptional start site (TSS) in the 5' CpG islands of the genes. Primer sequences for methylation specific PCR analysis were designed using MSPPrimer (<http://www.mspprimer.org>). All primer sequences are listed in Table 3. For expression studies using RT-PCR, primers were designed using the open access program Primer3 (<http://frodo.wi.mit.edu/cgi-bin/primer3/primer3 www.cgi>).

Table 3: Primers used for MSP, RT-PCR, Bisulfite sequencing analyses

Genes	Primers	S(5-3)	AS(5-3)
ADAMTS1	Unmeth	tTATTGtAAAGtAGGGTgTtAtttGGAtG	aaaaCTaAaaCAACaCaaAaATTaaTaCCTaaCa
	Meth	tTAGGGTGCGtTAICGGAC	TaAaaCAACCGCGaAaATTaaTaCCTaaCG
	RT	tcgaagtgaaacacagcggaac	gcgctgtaccctcaagacaa
ASCL2	Unmeth	tGtTAGGGGAAGtGtGttAGttAGTtGtG	CaTACaCaTaaCaCaTaaCTCACaaaaAaCATCa
	Meth	GAtAGCGCGtttAGttAGTCCG	TaaCGCGTaaCTCACGaaaAaCATCG
BNC1	Unmeth	TTtGtttTTTTGGGAGAG GtAAAAtttGAtAtG	aaAaACCTCaCCaaCaaCCaaCa
	Meth	GtttTTtttCCGGGAGAGGtAAAAttCGAtAC	CCGaaCGaCCGaaCG
	RT	acaaaagccctggcctcatct	tcgccccaaatgatatagaaa
	Bisulfite		
	Sequencing	AATtA TtTtT GAGAAAGAGY GtAGAGAAtT	CCCCAAaCRCCCCAaaCTaC
BNI/P3	Unmeth	tAtGtttTGtGtAtGtAGGtttAAAGTtGtG	aCTCCaaACTaAaCaaAaCCCCa
	Meth	GtACGCGtAG GtttAAAGTCCG	CCGaaACTaAaCGaaAaCCCCCG
EVL	Unmeth	GTGTGTTTTTTTTGAGGATTTGGAGTTGTTTG	ACCACCAAAAAATAAAAAAACAAAAAACCA
	Meth	GAGGATTCGGAGTCGTTT	CCGAAAAATAAAAAAACGAAAAACGAAACCG
PNMT	Unmeth	TGGGGAtGATTGttGtAGttG	CaCCaAaCCaCCCaAAAAaCCATaTaCa
	Meth	GGGGACGATTGtCGtGtAGtC	AaCCGCCCGaaaCCATaTaCG
TFPI2	Unmeth	CCCACATAAAACAAACACCCAAACCA	TGGTTTGTGGGTAAGGTGTTTG
	Meth	GTTTCGTTGGGTAAGCGGTTT	CATAAAACGAAACACCCGAAACCG
TWIST1	Unmeth	tAGAtTtGtGGGtTtGtAGAtttGGAtttG	aCaaaAaaaaAaaaaAcaaTaTaaTaaCCCCa
	Meth	CGGGtTtGtAGAtCGGtAtC	CGAAaaaaAaaaaACGaaTaTaaTaaCCCCCG

[0103] For methylation-specific PCR analysis, DNA was extracted using the standard phenolchloroform extraction method. Bisulfite modification of genomic DNA was carried out using standard protocols (e.g. with the EZ DNA methylation Kit (Zymo Research™)). Methylation analysis of CAN gene (Cancer gene, as defined by Wood LD, Parsons DW, Jones S et al., The genomic landscapes of human breast and colorectal cancers, Science. 2007, 318(5853):1108-1113.) promoters was performed using MSP primer pairs located close to the putative transcriptional start site in the 5' CpG island with 2 $\mu$ l of bisulfite-treated DNA as template and JumpStart Red Taq DNA Polymerase (Sigma™) for amplification as previously described (Herman JG, Graff JR, Myöhänen S, Nelkin BD, Baylin SB., Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. Proc Natl Acad Sci USA. 1996 Sep 3;93(18):9821-9826).

[0104] Total RNA was extracted from cell lines using standard methods (e.g. the RNeasy Mini Kit (Qiagen™)), and treated with DNase. For reverse transcription (RT) reaction, 1 $\mu$ g of total RNA was subjected to the first strand cDNA synthesis using standard cDNA synthesis protocols (e.g. Superscript III first strand cDNA synthesis kit (Invitrogen™) used according to the manufacturer's instructions). Expression analysis was performed by RT-PCR using 1 $\mu$ l of cDNA as template and standard amplification protocols (e.g. JumpStart Red Taq DNA Polymerase (Sigma™)).

#### Quantitative methylation-specific PCR using real-time PCR

[0105] Sodium bisulfite modification, which converts unmethylated cytosine residues to uracil residues, was first carried out on lug genomic DNA isolated from the paraffin-embedded tissue sections using standard methylation procedures (e.g. EZ DNA methylation kit (Zymo Research Co) according to the manufacturer's instructions). For quantitative real-time analyses, standard procedures were used. For example, the Power SYBR Green PCR kit (Applied Biosystems) was used and the amplification conditions consisted of an initial 10 minute denaturation step at 95°C, followed by 40 cycles of denaturation at 95°C for 15 seconds and annealing and extension for 30 seconds and 60 seconds, respectively. An ABI StepOnePlusReal-Time PCR System was used (Applied Biosystems), and for quantification the comparative cycle threshold (Ct) method was used, normalizing the Ct values for the indicated gene to the Ct values of Unmethylated reaction relative to a methylated reaction sample.

#### Methylation on Beads (MOB) method

[0106] Methods and buffers used for methylation on beads extraction are as described previously (Bailey VJ, Zhang Y, Keeley BP, Yin C, Pelosky KL, Brock M, Baylin SB, Herman JG, Wang TH. Single-tube analysis of DNA methylation with silica superparamagnetic beads. Clin Chem. 56(6):1022-1025). Briefly, 50 $\mu$ L of Protease K was added into the bottom of an eppendorf tube, and 200 $\mu$ L methylation on beads Binding Buffer 1 and 100 $\mu$ L Lysing Buffer to 200  $\mu$ L was added to the serum sample. The sample was incubated at 55°C for 15 minutes (cell lines, whole blood), and 200 $\mu$ L 100% Isopropyl Alcohol was added, and the sample was mixed. 35 $\mu$ L of Magnetic Beads was then added to the sample and mixed. The tube was then placed on the magnetic holder to capture the beads, and the supernatant was discarded. The tube was removed from the magnetic holder, and the beads were washed with 500 $\mu$ L of Wash Buffer 1. The beads were once again captured on the magnetic holder, and then washed a second time in 500 $\mu$ L of Wash buffer 2. This step was then repeated one time, and the beads were again captured, and the supernatant was discarded. 50 $\mu$ L of Elution Buffer was added to the eppendorf tube containing the washed beads. To denature the DNA for bisulfite treatment, 6 $\mu$ L 2M NaOH was added to the tube, mixed, and incubated at 70°C for 10-15 minutes. 12 $\mu$ L of freshly prepared hydroquinone solution was then added and mixed. 200 $\mu$ L of prepared NaBisulfite solution was added and mixed. The NaBisulfite solution was vortexed to insure that it was fully saturated. The sample was incubated at 65°C for 4-6 hours, and then 120 $\mu$ L of 10M NaOH was added, mixed, and incubated at 50°C for an additional 10 minutes. 350 $\mu$ L of methylation on beads Binding Buffer 2, 100 $\mu$ L of IPA was added to the sample and mixed. The beads were then captured and washed once in Wash Buffer 1 and twice in Wash Buffer 2 as described above. The beads were then captured and the supernatant discarded. The DNA was then eluted from the magnetic beads with 100 $\mu$ L of PCR buffer. The sample was then divided as desired for subsequent PCR amplification.

#### Claims

1. An in vitro method for detecting or characterizing a neoplasia in a biologic sample of a subject, the method comprising detecting the methylation of a *BNC1* and *ADAMTS1* gene, wherein detection of methylation detects or characterizes the presence of a neoplasia in the sample, the method optionally further comprising the step of detecting an alteration in the | sequence or expression level of a | *Brcal*, *Brca2*, *p16*, *K-ras*, *APC*, *PalB2*, *DPC4*, *EGFR*, and/or *EML- ALK4* gene or polypeptide, wherein the neoplasia is a cancer selected from colon, lung or pancreatic cancer and wherein the level of methylation present at *BNC1* and *ADAMTS1* gene promoter is compared to a reference selected from the level of methylation present in a control sample obtained from a patient that does not have a neoplasia, a baseline

level of methylation present in a biologic sample derived from a patient, prior to, during or after, treatment for a neoplasia or a standardized curve.

2. The method of claim 1, wherein the method detects an increase in methylation relative to a reference.
3. The method of claim 1, wherein the biologic sample is a tissue or biologic fluid sample.
4. The method of claim 2, wherein the reference is the level of methylation present at the promoter in a control sample preferably derived from a healthy subject.
5. The method of any one of claims 1-4, wherein the methylation is detected by quantitative methylation-specific PCR (QMSP).
6. The method of claim 5, wherein the level or the frequency of methylation is quantified.
7. The method of any one of claims 1-6, wherein the methylation levels of the *BNC1* and *ADAMTS1* promoters are quantified.
8. The method of claim 1, wherein the subject is a smoker, has a *Brcal* or *Brca2* mutation, pancreatic cyst, chronic pancreatitis, presence of colon polyps or adenomas, or a family history of cancer.
9. An in vitro method according to claim 1 for detecting or characterizing lung or colon cancer in a sample derived from a subject, the method comprising detecting methylation of a *BNC1* and *ADAMTS1* gene, wherein detection of methylation detects or characterizes lung or colon cancer in the subject and further comprising detecting an alteration in the sequence or expression level of a *Brca1*, *Brca2*, *p16*, *K-ras*, *APC*, *EGFR (lung)*, and/or *EML-ALK4* gene or polypeptide, wherein the level of methylation present at *BNC1* and *AD1MTS1* gene promoter is compared to a reference selected from the level of methylation present in a control sample obtained from a patient that does not have a neoplasia, a baseline level of methylation present in a biologic sample derived from a patient, prior to, during or after, treatment for a neoplasia or a standardized curve.
10. The method of claim 9, wherein the subject is a smoker, has colon polyps or adenomas, or a family history of cancer.
11. An in vitro method according to claim 1 for detecting or characterizing pancreatic cancer in a serum or plasma sample derived from a subject, the method comprising detecting the methylation of *BNC1* and *ADAMTS1* gene, wherein detection of methylation detects or characterizes pancreatic cancer in the subject, further comprising the step of detecting an alteration in the sequence or expression of a *Brca1*, *Brca2*, *p16*, *K-ras*, *APC*, *PalB2*, and/or *DPC4* gene or polypeptide relative to a reference, wherein the level of methylation present at *BNC1* and *ADAMTS1* gene promoter is compared to a reference selected from the level of methylation present in a control sample obtained from a patient that does not have a neoplasia, a baseline level of methylation present in a biologic sample derived from a patient, prior to, during or after, treatment for a neoplasia or a standardized curve. |
12. The method of claim 11, wherein the method detects an increase in methylation relative to a reference.
13. The method of claim 11, wherein the subject is identified as a smoker, has a *Brcal* or *Brca2* mutation, pancreatic cyst, chronic pancreatitis, or a family history of cancer.

#### Patentansprüche

1. In-vitro-Verfahren zur Detektion oder Charakterisierung einer Neoplasie in einer biologischen Probe eines Subjekts; das Verfahren umfasst die Detektion der Methylierung eines *BNC1*- und *ADAMTS1*-Gens, wobei die Detektion der Methylierung das Vorliegen einer Neoplasie in der Probe detektiert oder charakterisiert; das Verfahren umfasst optional ferner den Schritt der Detektion einer Veränderung auf dem Sequenz- oder Expressionslevel eines *Brca1*-, *Brca2*-, *p16*-, *K-ras*-, *APC*-, *PalB2*-, *DPC4*-, *EGFR*- und/oder *EML-ALK4*-Gens oder -polypeptids, wobei die Neoplasie ein Krebs ausgewählt aus Dickdarm-, Lungen- oder Bauchspeicheldrüsenkrebs ist und wobei das Methylierungslevel eines *BNC1*- und *ADAMTS1*-Genpromotors mit einer Referenz verglichen wird, die ausgewählt ist aus dem Methylierungslevel einer Kontrollprobe, die von einem Patienten, der keine Neoplasie hat, erhalten wurde, einem Methylierungslevel-Ausgangswert von einer biologischen Probe, die von einem Patient vor, während oder

## EP 2 576 832 B1

nach der Behandlung einer Neoplasie stammt, oder einer standardisierten Kurve.

2. Verfahren gemäß Anspruch 1, wobei das Verfahren eine Erhöhung der Methylierung in Relation zu einer Referenz detektiert.
3. Verfahren gemäß Anspruch 1, wobei die biologische Probe eine Gewebe- oder biologische Flüssigkeitsprobe ist.
4. Verfahren gemäß Anspruch 2, wobei die Referenz das Methylierungslevel des Promotors in einer Kontrollprobe, die vorzugsweise von einem gesunden Subjekt stammt, ist.
5. Verfahren nach einem der Ansprüche 1 bis 4, wobei die Methylierung durch quantitative methylierungsspezifische PCR (QMSP) detektiert wird.
6. Verfahren gemäß Anspruch 5, wobei das Methylierungslevel oder die Methylierungsfrequenz quantifiziert wird.
7. Verfahren gemäß einem der Ansprüche 1 bis 6, wobei die Methylierungslevel der *BNC1*- und *ADAMTS1*-Promotoren quantifiziert werden.
8. Verfahren gemäß Anspruch 1, wobei das Subjekt ein Raucher ist, eine *Brca1*- oder *Brca2*-Mutation, eine Pankreaszyste, chronische Pankreatitis, das Vorliegen von Dickdarpolyphen oder -adenomen oder eine Familienanamnese mit Krebs aufweist.
9. In-vitro-Verfahren gemäß Anspruch 1 zur Detektion oder Charakterisierung von Lungen- oder Dickdarmkrebs in einer Probe, die von einem Subjekt stammt; das Verfahren umfasst die Detektion der Methylierung eines *BNC1*- und *ADAMTS1*-Gens, wobei die Detektion der Methylierung Lungen- oder Dickdarmkrebs in dem Subjekt detektiert oder charakterisiert, und welches ferner die Detektion einer Veränderung auf dem Sequenz- oder Expressionslevel eines *Brca1*-, *Brca2*-, *p16*-, *K-ras*-, *APC*-, *EGFR* (*Lungen*)- und/oder *EML- ALK4*-Gens oder - polypeptids umfasst, wobei das Methylierungslevel eines *BNC1*- und *ADAMTS1*-Genpromotors mit einer Referenz verglichen wird, die ausgewählt ist aus dem Methylierungslevel einer Kontrollprobe, die von einem Patient, der keine Neoplasie hat, erhalten wurde, einem Methylierungslevel-Ausgangswert von einer biologischen Probe, die von einem Patienten vor, während oder nach der Behandlung einer Neoplasie stammt, oder einer standardisierten Kurve.
10. Verfahren gemäß Anspruch 9, wobei das Subjekt ein Raucher ist, Dickdarpolyphen oder -adenome oder eine Familienanamnese mit Krebs aufweist.
11. In-vitro-Verfahren gemäß Anspruch 1 zur Detektion oder Charakterisierung von Bauchspeicheldrüsenkrebs in einer Serum- oder Plasmaprobe, die von einem Subjekt stammt; das Verfahren umfasst die Detektion der Methylierung eines *BNC1*- und *ADAMTS1*-Gens, wobei die Detektion der Methylierung Bauchspeicheldrüsenkrebs in dem Subjekt detektiert oder charakterisiert, und welches ferner den Schritt der Detektion einer Veränderung der Sequenz oder Expression eines *Brca1*-, *Brca2*-, *p16*-, *K-ras*-, *APC*-, *PalB2*- und/oder *DPC4*-Gens oder -polypeptids in Relation zu einer Referenz umfasst, wobei das Methylierungslevel eines *BNC1*- und *ADAMTS1*-Genpromotors mit einer Referenz verglichen wird, die ausgewählt ist aus dem Methylierungslevel einer Kontrollprobe, die von einem Patienten der keine Neoplasie hat, erhalten wurde, einem Methylierungslevel-Ausgangswert von einer biologischen Probe, die von einem Patienten vor, während oder nach der Behandlung einer Neoplasie stammt, oder einer standardisierten Kurve.
12. Verfahren gemäß Anspruch 11, wobei das Verfahren einen Anstieg der Methylierung in Relation zu einer Referenz detektiert.
13. Verfahren gemäß Anspruch 11, wobei das Subjekt als Raucher identifiziert ist, eine *Brca1*- oder *Brca2*-Mutation, Pankreaszyste, chronische Pankreatitis oder eine Familienanamnese mit Krebs aufweist.

### Revendications

1. Procédé in vitro pour détecter ou caractériser une néoplasie dans un échantillon biologique d'un sujet, le procédé comprenant la détection de la méthylation du gène *BNC1* et *ADAMTS1*, dans lequel la détection de la méthylation détecte ou caractérise la présence d'une néoplasie dans l'échantillon, le procédé comprenant en outre éventuelle-

ment l'étape de détection d'une altération de la séquence ou du niveau d'expression d'un gène *Brca1*, *Brca2*, *p16*, *K-ras*, *APC*, *PalB2*, *DPC4*, *EGFR* et/ou *EML-ALK4* ou d'un polypeptide, dans lequel la néoplasie est un cancer choisi parmi le cancer du colon, du poumon ou du pancréas et dans lequel le niveau de méthylation présent au niveau d'un promoteur du gène *BANC1* et *ADAMTS1* est comparé à une référence choisie parmi le niveau de méthylation présent dans un échantillon témoin obtenu auprès d'un patient qui ne présente pas de néoplasie, un niveau de méthylation de base présent dans un échantillon biologique dérivé d'un patient, avant ou pendant ou après un traitement d'une néoplasie ou d'une courbe normalisée.

- 5
  - 10
  - 15
  - 20
  - 25
  - 30
  - 35
  - 40
  - 45
  - 50
  - 55
2. Procédé selon la revendication 1, dans lequel le procédé détecte une augmentation de la méthylation par rapport à une référence.
  3. Procédé selon la revendication 1, dans lequel l'échantillon biologique est un tissu ou un échantillon de fluide biologique.
  4. Procédé selon la revendication 2, dans lequel la référence est le niveau de méthylation présent au niveau du promoteur dans un échantillon témoin dérivé de préférence d'un sujet en bonne santé.
  5. Procédé selon l'une quelconque des revendications 1 à 4, dans lequel la méthylation est détectée par une PCR quantitative spécifique d'une méthylation (QMSP).
  6. Procédé selon la revendication 5, dans lequel le niveau ou la fréquence de méthylation est quantifié.
  7. Procédé selon l'une quelconque des revendications 1 à 6, dans lequel les niveaux de méthylation des promoteurs de *BNC1* et *ADANTS1* sont quantifiés.
  8. Procédé selon la revendication 1, dans lequel le sujet est un fumeur, présente une mutation de *Brcal* ou de *Brca2*, un kyste pancréatique, une pancréatite chronique, la présence de polypes ou d'adénomes dans le colon ou des antécédents familiaux de cancer.
  9. Procédé in vitro selon la revendication 1, pour détecter ou caractériser un cancer du poumon ou du colon dans un échantillon dérivé d'un sujet, le procédé comprenant la détection de la méthylation du gène *BNC1* et *ADAMTS1*, dans lequel la détection de la méthylation détecte ou caractérise le cancer du poumon ou du colon chez le sujet et comprenant en outre la détection d'une altération de la séquence ou du niveau d'expression d'un gène *Brca1*, *Brca2*, *p16*, *K-ras*, *APC*, *EGFR* (poumon) et/ou *EML-ALK4* ou d'un polypeptide, dans lequel le niveau de méthylation présent au niveau d'un promoteur du gène *BNC1* et *ADAMTS1* est comparé à une référence choisie parmi le niveau de méthylation présent dans un échantillon témoin obtenu auprès d'un patient qui ne présente pas de néoplasie, un niveau de méthylation de base présent dans un échantillon biologique dérivé d'un patient, avant ou pendant ou après un traitement d'une néoplasie ou d'une courbe normalisée.
  10. Procédé selon la revendication 9, dans lequel le sujet est un fumeur, présente des polypes ou des adénomes dans le colon ou des antécédents familiaux de cancer.
  11. Procédé in vitro selon la revendication 1, pour détecter ou caractériser un cancer du pancréas dans un échantillon de sérum ou de plasma dérivé d'un sujet, le procédé comprenant la détection de la méthylation du gène *BNC1* et *ADAMTS1*, dans lequel la détection de la méthylation détecte ou caractérise le cancer du pancréas chez le sujet, comprenant en outre l'étape de détection d'une altération de la séquence ou de l'expression d'un gène *Brca1*, *Brca2*, *p16*, *K-ras*, *APC*, *PalB2* et/ou *DPC4* ou d'un polypeptide, par rapport à une référence, dans lequel le niveau de méthylation présent au niveau du promoteur est comparé à une référence choisie parmi le niveau de méthylation présent dans un échantillon témoin obtenu auprès d'un patient qui ne présente pas de néoplasie, un niveau de méthylation de base présent dans un échantillon biologique dérivé d'un patient, avant ou pendant ou après un traitement d'une néoplasie ou d'une courbe normalisée.
  12. Procédé selon la revendication 11, dans lequel le procédé détecte une augmentation de la méthylation par rapport à une référence.
  13. Procédé selon la revendication 11, dans lequel le sujet est identifié comme étant un fumeur, présentant une mutation de *Brcal* ou de *Brca2*, un kyste pancréatique, une pancréatite chronique, des antécédents familiaux de cancer.







Figure 1A



Figures 1B and 1C






Figure 2C





Figure 3



## REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

## Patent documents cited in the description

- US 2009000039 W [0060] [0061]
- US 5786146 A [0066]
- US 6017704 A [0066]
- US 6300756 B [0066]
- US 6265171 B [0066]

## Non-patent literature cited in the description

- **SHAMES DAVID S et al.** A genome-wide screen for promoter methylation in lung cancer identifies novel methylation markers for multiple malignancies. *PLOS MEDICINE, PUBLIC LIBRARY OF SCIENCE, US*, 01 December 2006, vol. 3 (12), ISSN 1549-1676 [0003]
- Aberrant methylation of ADAMTS1 in non-small cell lung cancer. **CHOI J E et al.** CANCER GENETICS AND CYTOGENETICS. ELSEVIER SCIENCE PUBLISHING, 01 December 2008, vol. 187, 80-84 [0004]
- **PHILIP J. JOHNSON et al.** Plasma Nucleic Acids in the Diagnosis and Management of Malignant Disease. *CLINICAL CHEMISTRY*, 01 August 2002, vol. 48, ISSN 0009-9147, 1186-1193 [0005]
- Relevance of DNA methylation in the management of cancer. **ESTELLER M.** LANCET ONCOLOGY. LANCET PUBLISHING GROUP, 01 June 2003, vol. 4, 351-358 [0006]
- **G STRATHDEE et al.** Aberrant DNA methylation in cancer: potential clinical interventions. *EXPERT REVIEWS IN MOLECULAR MEDICINE*, 01 March 2002, vol. 4, 1-17 [0007]
- **AHLQUIST TERJE et al.** Gene methylation profiles of normal mucosa, and benign and malignant colorectal tumors identify early onset markers. *MOLECULAR CANCER, BIOMED CENTRAL*, 31 December 2008, vol. 7 (1), ISSN 1476-4598, 94 [0008]
- **SINGLETON et al.** Dictionary of Microbiology and Molecular Biology. 1994 [0018]
- The Cambridge Dictionary of Science and Technology. 1988 [0018]
- The Glossary of Genetics. Springer Verlag, 1991 [0018]
- **HALE ; MARHAM.** *The Harper Collins Dictionary of Biology*, 1991 [0018]
- **JEMAL A et al.** *Cancer Statistics*, 2011, 2010 [0055]
- *CA Cancer J Clin.*, vol. 61 (2), 133-4 [0055]
- **KUPPUSWAMY et al.** *Proc. Natl Acad. Sci. USA*, 1991, vol. 88, 1143-1147 [0066]
- *Adorján Nucleic Acids Res.*, vol. 30, e21 [0066]
- *Hou Clin. Biochem.*, 2003, vol. 36, 197-202 [0066]
- **TYAGI et al.** *Nature Biotechnology*, 1996, vol. 14 (3), 303-8 [0069]
- **SAMBROOK.** *Molecular Cloning: A Laboratory Manual*. 1989 [0080]
- **GAIT.** *Oligonucleotide Synthesis*, 1984 [0080]
- **FRESHNEY.** *Animal Cell Culture*, 1987 [0080]
- *Methods in Enzymology.* **WEIR.** *Handbook of Experimental Immunology*. 1996 [0080]
- **MILLER ; CALOS.** *Gene Transfer Vectors for Mammalian Cells*, 1987 [0080]
- **AUSUBEL.** *Current Protocols in Molecular Biology*, 1987 [0080]
- **MULLIS.** *PCR: The Polymerase Chain Reaction*, 1994 [0080]
- **COLIGAN.** *Current Protocols in Immunology*, 1991 [0080]
- **SCHUEBEL K ; CHEN W ; COPE L ; GLÖCKNER SC ; SUZUKI H et al.** Comparing the DNA hypermethylome with gene mutations in human colorectal cancer. *PLOS Genetics*, 2007, vol. 3 (9), 1709-1723 [0098]
- **RHEE et al.** *Nature*, 2002 [0101]
- **WOOD LD ; PARSONS DW ; JONES S et al.** The genomic landscapes of human breast and colorectal cancers. *Science*, 2007, vol. 318 (5853), 1108-1113 [0103]
- *Proc Natl Acad Sci USA.*, 03 September 1996, vol. 93 (18), 9821-9826 [0103]
- *Clin Chem.*, vol. 56 (6), 1022-1025 [0106]

专利名称(译)	用于检测瘤形成的组合物和方法		
公开(公告)号	<a href="#">EP2576832B1</a>	公开(公告)日	2016-05-04
申请号	EP2011787338	申请日	2011-05-25
[标]申请(专利权)人(译)	约翰霍普金斯大学		
申请(专利权)人(译)	约翰·霍普金斯大学		
当前申请(专利权)人(译)	约翰·霍普金斯大学		
[标]发明人	AHUJA NITA BAYLIN STEPHEN HERMAN JAMES GORDON WANG JEFF BAILEY VASUDEV YI JOO MI		
发明人	AHUJA, NITA BAYLIN, STEPHEN HERMAN, JAMES, GORDON WANG, JEFF BAILEY, VASUDEV YI, JOO-MI		
IPC分类号	C12Q1/68 C12N15/12 G01N33/53 G01N33/574		
CPC分类号	C12Q1/6886 C12Q1/686 C12Q2600/154 G01N33/57407 G01N33/57419 G01N33/57423 G01N33/57438		
优先权	61/348010 2010-05-25 US		
其他公开文献	EP2576832A4 EP2576832A2		
外部链接	<a href="#">Espacenet</a>		

#### 摘要(译)

本发明提供了用于检测受试者样品(例如,血清,血液,血浆,组织)中的瘤形成(例如,胰腺癌,肺癌,结肠癌)的组合物和方法。在特定的实施方案中,本发明提供了检测血清中循环DNA中BNC1和ADAMTS1启动子甲基化的方法。

```

1  gcggcggggg cggccatcgt gctgcgcagc ctgggcgctt ggggagccgc ccacttcgcc
61  gggtcgcgcc ccgacggccg gagcgtggat gcggcgggc ccgccgagcc ggggcggacg
121  cggggcggcc cgggccggg agacgcggcg gcagccccc caccgcagcg gtcgcaggat
181  ggccgaggct atcagctgta ctctgaactg tagttgcaa agtttcaaac ccgggaaaat
241  aaaccaccgt cagtgtgacc aatgcaagca tggatgggtg gcccacgctc taagtaagct
301  aaggatcccc cccatgtatc caacaagcca ggtggagatt gtccagtcga atgtagtgtt
361  tgatattagc agcctcatgc tctatggac ccaggccatc cccgttcgcc taaaaatcct
421  actggaccgg ctcttcagtg tgttgaagca agatgaggtt ctccagatcc tccatgcctt
481  ggactggaca ctcaagatt atatccgtgg atacgtactg caggatgcac caggaaaggt

```