



(11) **EP 2 643 694 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:  
**03.01.2018 Bulletin 2018/01**

(51) Int Cl.:  
**G01N 33/50 (2006.01) C12Q 1/68 (2018.01)**

(21) Application number: **11842900.0**

(86) International application number:  
**PCT/US2011/062125**

(22) Date of filing: **23.11.2011**

(87) International publication number:  
**WO 2012/071557 (31.05.2012 Gazette 2012/22)**

(54) **RAPID IN VIVO GENE MUTATION ASSAY BASED ON THE PIG-A GENE**

SCHNELLER IN-VIVO-GENMUTATIONS-ASSAY AUF BASIS DES PIG-A-GENS

ÉVALUATION DE MUTATION GÉNÉTIQUE IN VIVO RAPIDE SUR LA BASE DU GÈNE PIG-A

(84) Designated Contracting States:  
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**

(30) Priority: **24.11.2010 US 417104 P**

(43) Date of publication of application:  
**02.10.2013 Bulletin 2013/40**

(73) Proprietor: **Litron Laboratories Ltd. Rochester, NY 14623 (US)**

(72) Inventors:  
• **DERTINGER, Stephen, D. Webster, NY 14580 (US)**  
• **BRYCE, Steven, M. Rochester, NY 14609 (US)**

(74) Representative: **Zaccaro, Elisabetta et al Notarbartolo & Gervasi S.p.A. Corso di Porta Vittoria, 9 20122 Milano (IT)**

(56) References cited:  
**WO-A1-99/36778 US-A1- 2007 274 919**  
**US-A1- 2009 311 706 US-B2- 7 824 874**

- **D. J. ARATEN ET AL: "Clonal populations of hematopoietic cells with paroxysmal nocturnal hemoglobinuria genotype and phenotype are present in normal individuals", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 96, no. 9, 27 April 1999 (1999-04-27) , pages 5209-5214, XP055106018, ISSN: 0027-8424, DOI: 10.1073/pnas.96.9.5209**

- **STEPHEN D DERTINGER ET AL: "When pigs fly: Immunomagnetic separation facilitates rapid determination of mutant frequency by flow cytometric analysis", MUTATION RESEARCH. GENETIC TOXICOLOGY AND ENVIRONMENTAL MUTAGENESIS, ELSEVIER, AMSTERDAM, NL, vol. 721, no. 2, 20 January 2011 (2011-01-20), pages 163-170, XP028369770, ISSN: 1383-5718, DOI: 10.1016/J.MRGENTOX.2011.01.009 [retrieved on 2011-01-28]**
- **BRYCE STEVEN MET AL: "In vivo mutation assay based on the endogenous Pig-a locus", ENVIRONMENTAL AND MOLECULAR MUTAGENESIS, WILEY-LISS, INC. CHICHESTER, GB, vol. 49, no. 4, 1 May 2008 (2008-05-01), pages 256-264, XP002622290, ISSN: 0893-6692, DOI: 10.1002/EM.20379 [retrieved on 2008-02-20]**
- **PHONETHEPSWATH ET AL.: 'Erythrocyte-based Pig-a gene mutation assay: demonstration of cross- species potential.' MUTAT RES. vol. 657, no. 2, 2008, pages 122 - 126, XP025679907**
- **"Lympholyte -M", , 1 November 2009 (2009-11-01), pages 325-334259, XP055257187, Retrieved from the Internet: URL: [http://www.tebu-bio.com/Product/007CL5\\_031/LYMPHOLYTE\\_M.html](http://www.tebu-bio.com/Product/007CL5_031/LYMPHOLYTE_M.html) [retrieved on 2016-03-10]**

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

**EP 2 643 694 B1**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**Description****FIELD OF THE INVENTION**

5 [0001] The present invention relates to methods and materials for the quantitative analysis of *in vivo* mutant cell frequencies in individuals, particularly using peripheral blood samples of vertebrates.

**BACKGROUND OF THE INVENTION**

10 [0002] Several authors have articulated the possibility of using the endogenous *Pig-A* gene as a reporter of somatic mutation (Araten et al., "Clonal Populations of Hematopoietic Cells with Paroxysmal Nocturnal Hemoglobinuria Genotype and Phenotype are Present in Normal Individuals," Proc Natl Acad Sci USA 96:5209-5214 (1999); Chen et al., "Glyco-phosphatidylinositol-anchored Protein Deficiency as a Marker of Mutator Phenotypes in Cancer," Cancer Res. 61:654-658 (2001)). As with the *HPRT* locus, *Pig-A* is located on the X-chromosome. Iida and colleagues isolated the human genomic gene, and found that it contains six exons over its 17 kb length (Iida et al., "Characterization of Genomic PIG-A Gene: A Gene for GPI Anchor Biosynthesis and Paroxysmal Nocturnal Hemoglobinuria," Blood 83:3126-3131 (1994)). As demonstrated by Kawagoe et al., "Molecular cloning of Murine pig-a, a Gene for GPI-anchor Biosynthesis, and Demonstration of Interspecies Conservation of its Structure, Function, and Genetic Locus," Genomics 23:566-574 (1994), there is a high degree of interspecies conservation of the gene's structure, function, and locus. The *Pig-A* gene product acts in the first step in glycosylphosphatidylinositol (GPI) anchor biosynthesis, and the entire process is thought to require at least 12 genes. Mutation of any one of these could theoretically result in GPI anchor deficiency. However, all other genes involved in GPI anchor synthesis are autosomal. Mutations on both alleles would have to occur to ablate expression of GPI anchors, and this is expected to be a very rare event. Thus, an inability to anchor GPI-linked proteins in the outer membrane is believed to be virtually equivalent to *Pig-A* mutation.

25 [0003] This key assumption, as well as practical aspects of assay development, greatly benefit from research on Paroxysmal Nocturnal Hemoglobinuria (PNH). PNH is a genetic disorder that affects 1 to 10 per million individuals, and is caused by a somatic *Pig-A* gene mutation within a bone marrow stem cell (Norris et al., "The Defect in Glycosylphosphatidylinositol Anchor Synthesis in Paroxysmal Nocturnal Hemoglobinuria," Blood 83:816-821 (1994)). Since bone marrow stem cells are the precursors for the entire hematopoietic system, the gene mutation tends to affect numerous lineages. Erythrocytes, granulocytes and monocytes are typically affected. In a minority of cases, however, lymphocytes are also affected. A key finding is that all PNH clones to date exhibit mutation at the *Pig-A* locus (Nishimura et al., "Paroxysmal Nocturnal Hemoglobinuria: An Acquired Genetic Disease," Am J Hematol 62:175-182 (1999)). Furthermore, an analysis of 146 PNH patients by Nishimura and colleagues provides important examples of the types of mutations that lead to GPI anchor deficiency. Single-base substitutions and frame-shift events are the most highly represented classes of mutation observed. Even so, there are three examples of large deletions (entire gene, 4 kb, and 737 base pairs), as well as a large insertion (88 base pairs). The mutations are widely distributed in the coding regions and splice sites, although others have found a somewhat higher frequency of missense mutations in exon 2 relative to other exons (Nafa et al., "The Spectrum of Somatic Mutations in the PIG-A Gene in Paroxysmal Nocturnal Hemoglobinuria Includes Large Deletions and Small Duplications," Blood Cells Mol Dis 24:370-384 (1998)). Taken together, the PNH literature provides strong evidence that an *in vivo* assay based on the *Pig-A* gene would be sensitive to each important class of mutation.

35 [0004] In a report by Miura et al., "Development of an In Vivo Gene Mutation Assay Using the Endogenous Pig-A Gene: I. Flow Cytometric Detection of CD59-Negative Peripheral Red Blood Cells and CD48-Negative Spleen T-Cells From the Rat," Environ. Molec. Mutagen. 49:614-621 (2008), a method for quantifying the frequency of mutant phenotype erythrocytes was identified. In that flow cytometry-based assay, anti-CD45 antibody was used to differentiate leukocytes from erythrocytes, and anti-CD59-FITC was used to distinguish mutant phenotype erythrocytes from wild-type erythrocytes. The authors also described a second approach whereby the fluorescent reagent FLAER and flow cytometry could be used to quantify the frequency of mutant phenotype erythrocytes. However, these approaches did not differentiate mature erythrocytes from the immature fraction of erythrocytes (reticulocytes). This is a significant disadvantage of the approach of Miura *et al.*, because differential staining of mature and immature erythrocytes allows one to determine the percentage of reticulocytes among total erythrocytes simultaneously with *Pig-A* mutation measurements. These percent reticulocyte values provide important information regarding bone marrow toxicity, a parameter that is valuable for interpreting any genotoxicity endpoint that is based on hematopoietic cells. Differentially staining reticulocytes and mature erythrocytes also allows one to measure *Pig-A* mutation frequency in both the total RBC cohort as well as the reticulocyte fraction. The latter measurement is valuable for some experimental designs, since maximal mutagenic responses are obtained in this fraction of cells sooner than those observed in the total erythrocyte pool. Furthermore, the approach of Miura *et al.* for distinguishing erythrocytes from leukocytes was less than ideal. Namely, in their hands, anti-CD45 did not afford clear resolution of nucleated cells from erythrocytes. Rather than distinct populations, a continuum of CD45-

associated fluorescent events was observed. The consequence of this is contamination of the erythrocyte analyses with leukocytes that failed to exhibit sufficient differential fluorescent resolution. This likely contributed to the high and variable baseline mutation frequencies that were reported by these investigators. Arten et al. in PNAS 96: 5209-5214 (1999), analyze the hypothesis that PIG-A gene mutations are not the only culprits for the development of PNH and uses a method of magnetic bead enrichment for PNH granulocytes whereby part of a cellular sample was depleted of GPI-positive white cells by incubating with a murine (IgM) anti-CD16 antibody and separating with rat anti-mouse IgM magnetic Dynabeads.

[0005] In U.S. Patent No. 7,824,874 to Dertinger, a method of enumerating *Pig-A* mutant cell frequency from peripheral blood samples is identified. The described methodology uses a three-color labeling approach to distinguish GPI anchor-deficient cells from GPI anchor-expressing cells, platelets from other blood cells, and reticulocytes from erythrocytes. In U.S. Patent Application Publ. No. US20090311706 to Dertinger and Phonethepswath et al., "Erythrocyte-based *Pig-A* Gene Mutation Assay: Demonstration of Cross-Species Potential," *Mutat. Res.* 657:122-126 (2008), another method of enumerating *Pig-A* mutation frequency from peripheral blood samples is identified. In this method, the peripheral blood sample is treated in a manner to substantially separate RBCs from platelets and leukocytes, thereby enriching the sample for RBCs and making the method less susceptible to spuriously high readings. However, in both cases, the time required to evaluate millions of cells, especially reticulocytes, for the *Pig-A* mutant phenotype is a very time-consuming process, and most analyses are therefore based on suboptimal numbers of total cells interrogated for the mutant phenotype. This situation leads to less reliable estimates of mutation frequency, especially in situations when mutation frequency is low, as is typically the case in individuals that have not been exposed to potent mutagen(s). It would be desirable to obtain an assay that can identify and quantify many more *Pig-A* mutant phenotype cells per unit time. Such an assay would be more practical to perform, in terms of efficiently studying large numbers of specimens, and it would be endowed with greater reliability and sensitivity, *i.e.*, greater power to detect modest changes to the frequency of mutant phenotype erythrocytes and/or mutant phenotype reticulocytes.

[0006] The present invention is directed to overcoming these and other deficiencies in the prior art.

## DEFINITIONS

[0007] For purposes of the present invention, the following terms are defined as follows:

"*PIG-A*" and "*Pig-A*" are intended to mean the phosphatidylinositol glycan complementation group A gene. When referring to the human gene, the convention is to capitalize all letters, that is, *PIG-A*. When referring to other species, the convention is to use lower case letters, that is, *Pig-a*. However, for the purposes of the present invention, the term *Pig-A* is meant to refer to any vertebrate species, including man.

[0008] "GPI" is intended to mean glycosylphosphatidylinositol, a glycolipid that is attached to the C-terminus of certain proteins during posttranslational modification. Certain GPI-anchored proteins such as CD24, CD59 and CD55 normally appear on the cell surface of erythrocytes.

[0009] "Erythrocytes" is intended to mean enucleated red blood cells, regardless of RNA content. Erythrocytes is abbreviated RBCs.

[0010] "Normochromatic erythrocytes" is intended to mean enucleated red blood cells that have matured to the point that RNA content is negligible. Normochromatic erythrocytes is abbreviated NCEs.

[0011] "Reticulocytes" is intended to mean recently formed enucleated red blood cells that are characterized by the presence of cytoplasmic RNA. Reticulocytes is abbreviated RETs.

[0012] "*Pig-A* mutant cells" is intended to mean erythrocytes and/or reticulocytes with altered *Pig-A* DNA sequence, such that transcription of the *Pig-A* gene is affected, resulting in a phenotype that is distinguishable by either a lack of or significant deficiency of GPI-anchored proteins on the cell surface.

[0013] "Wild-type cells" is intended to mean erythrocytes and/or reticulocytes with a normal complement of GPI-anchored proteins on the cell surface.

[0014] "Paramagnetic beads" is intended to mean paramagnetic particles, typically though not exclusively nanoparticles, coated with antibodies or other ligands that bind to a specific surface antigen. This causes the cells expressing this antigen to attach to the paramagnetic beads (or vice versa). Afterwards the cell solution is transferred on a column placed in a strong magnetic field. In this step, the magnetic field should be strong enough such that the cells attached to the paramagnetic beads (expressing the antigen) stay on the column, while other cells (not expressing the antigen) flow through. With this process, the cells can be physically separated based on antigen expression profiles.

[0015] "Counting beads" is intended to mean latex particles or other flow cytometry-compatible particles that can be resolved from cells based on light scatter and/or fluorescence emission that are used to generate cell to Counting Bead ratios.

[0016] "Expression time" is intended to mean the period of time following exposure of a DNA damaging event until

the time that a mutated cell both expresses the GPI-anchor deficient phenotype and also appears in the tissue compartment under consideration (for example, in peripheral blood circulation).

## SUMMARY OF THE INVENTION

5

[0017] A first aspect of the present invention relates to methods for the enumeration of in vivo *Pig-A* gene locus mutation comprising the steps of: providing an enriched erythrocyte sample obtained from a mammal exposed to an exogenous agent, said sample comprising normochromatic erythrocytes and reticulocytes, and having a reduction in the frequency of platelets and leukocytes; contacting the enriched erythrocyte sample with a first fluorescent reagent that labels GPI anchor-expressing erythrocytes, but not GPI anchor-deficient erythrocytes, and a second fluorescent reagent that specifically labels platelets; separating the enriched erythrocyte sample into a first portion comprising platelets and GPI anchor-expressing erythrocytes and a second portion comprising GPI anchor-deficient erythrocytes; contacting the enriched erythrocyte sample or the second portion with a third fluorescent reagent that differentially labels normochromatic erythrocytes from reticulocytes and leukocytes; exciting the first, second, and third fluorescent reagents in the second portion with light of appropriate excitation wavelength, wherein the third fluorescent reagent has a fluorescent emission spectrum that does not substantially overlap with the fluorescent emission spectra of the first and/or second fluorescent reagents; and detecting the fluorescent emission and light scatter produced by the GPI anchor-deficient erythrocytes of the second portion labeled with the third fluorescent reagent, while excluding contaminating GPI anchor-expressing erythrocytes and reticulocytes, platelets, and leukocytes, and counting the number of GPI anchor-deficient erythrocytes and reticulocytes.

20

[0018] Further disclosed herein is a method that includes the steps of: providing an enriched erythrocyte sample obtained from a mammal exposed to an exogenous agent, said sample comprising normochromatic erythrocytes and reticulocytes, and having a reduction in the frequency of platelets and leukocytes; contacting the enriched erythrocyte sample with a first fluorescent reagent that labels GPI anchor-expressing erythrocytes, but not GPI anchor-deficient erythrocytes, and a second fluorescent reagent that specifically labels platelets; obtaining a quantitative measure of the total number of erythrocytes and/or reticulocytes in the enriched erythrocyte sample; separating the enriched erythrocyte sample into a first portion comprising platelets and GPI anchor-expressing erythrocytes and a second portion comprising GPI anchor-deficient erythrocytes; contacting the enriched erythrocyte sample or the second portion with a third fluorescent reagent that differentially labels normochromatic erythrocytes from reticulocytes and leukocytes; exciting the first, second, and third fluorescent reagents in the second portion with light of appropriate excitation wavelength, wherein the third fluorescent reagent has a fluorescent emission spectrum that does not substantially overlap with the fluorescent emission spectra of the first and/or second fluorescent reagents; detecting the fluorescent emission and light scatter produced by the GPI anchor-deficient erythrocytes of the second portion labeled with the third fluorescent reagent, while excluding contaminating GPI anchor-expressing erythrocytes and reticulocytes, platelets, and leukocytes, and counting the number of GPI anchor-deficient erythrocytes and reticulocytes; and calculating the frequency of GPI anchor-deficient erythrocytes and/or reticulocytes relative to total erythrocytes and/or reticulocytes present in the sample.

25

30

35

[0019] A second aspect of the present disclosure relates to a method of assessing the DNA-damaging potential of an exogenous chemical or physical agent. This method is carried out by exposing a mammal to an exogenous chemical or physical agent, and then performing the method according to the present disclosure, wherein a significant deviation in the frequency of mutant phenotype RBCs or RETs from a baseline mutant phenotype RBC or RET frequency in unexposed or vehicle control mammals indicates the genotoxic potential of the exogenous chemical or physical agent.

40

[0020] A third aspect of the present disclosure relates to a method of evaluating the effects of an exogenous agent that can modify endogenously-induced DNA damage. This method is carried out by administering to a mammal an exogenous agent that may modify endogenously-induced genetic damage; and then performing the flow cytometric method according to the present disclosure, wherein a significant deviation in the frequency of mutant phenotype RBCs or RETs from a baseline mutant phenotype RBC or RET frequency indicates that the exogenous agent can modify endogenous DNA damage.

45

[0021] A fourth aspect of the present disclosure relates to a method of evaluating the effects of an exogenous agent that can modify exogenously-induced DNA damage. This method is carried out by administering to a mammal a first exogenous agent that may modify exogenously-induced genetic damage; exposing the mammals to a second exogenous agent that causes genetic damage; and then performing the flow cytometric method according to the present disclosure, wherein a significant deviation in the frequency of mutant phenotype RBCs or RETs for genotoxicant-exposed mammals indicates that the first exogenous agent can modify exogenously-induced DNA damage.

50

[0022] The invention further relates to a kit comprising:

55

- a first fluorescent reagent that binds GPI anchor-expressing cells, but not GPI anchor-deficient cells;
- a second fluorescent reagent that specifically binds platelets;
- a third fluorescent reagent that differentially labels normochromatic erythrocytes, reticulocytes, and leukocytes, the

third fluorescent reagent having a fluorescent emission spectrum that does not substantially overlap with the fluorescent emission spectra of the first or second fluorescent reagent;  
 a fourth reagent suitable for separating erythrocytes from platelets and leukocytes, the fourth reagent comprising a density gradient medium;  
 5 a fifth reagent comprising paramagnetic particles, the fifth reagent specifically binding to GPI anchor-expressing cells or at least a portion of the first fluorescent reagent bound thereon;  
 and instructions for using the reagents and cell separation products, and for detecting and calculating the frequency of GPI anchor-deficient erythrocytes and/or reticulocytes relative to total erythrocytes and/or reticulocytes in the sample.

[0023] As demonstrated herein, improved methods for enumerating *Pig-A* mutation frequency in blood specimens are described. The experimental results presented in the accompanying examples were conducted with Sprague Dawley rats. Two peripheral blood erythrocyte populations were evaluated for the GPI anchor-deficient phenotype: RBCs, and the newly formed immature fraction, RETs. As demonstrated by the accompanying examples, the present invention can achieve *in vivo* mutant cell frequency measurements with previously unattainable precision and rates of data acquisition. Moreover, the experimental results confirm that the invention can be practiced with a variety of genotoxicants that cause genetic damage via different modes.

## BRIEF DESCRIPTION OF THE DRAWINGS

### [0024]

Figure 1 is a schematic that illustrates the overall strategy used for one embodiment of the present invention to rapidly determine *Pig-A* mutant cell frequency in peripheral blood samples. Separation of RBCs from leukocytes and platelets of the sample to form an enriched erythrocyte sample can be achieved using Lympholyte®-Mammal or other similar reagent. A phycoerythrin (PE) conjugated anti-CD59 antibody (anti-CD59-PE) represents a preferred first fluorescent reagent for labeling GPI anchor-expressing erythrocytes, but not GPI anchor-deficient erythrocytes, and a PE conjugated anti-CD61 antibody (anti-61-PE) represents a preferred second fluorescent reagent that specifically labels platelets. Anti-PE magnetic particles are preferred reagents for separating the enriched erythrocyte sample into a platelet, GPI anchor-expressing portion and a GPI anchor-deficient portion. The reagents identified in Figure 1 are exemplary, and the invention is not limited to practice with the reagents illustrated in Figure 1.

Figure 2 is a schematic that highlights the differential staining and also the relative depletion/enrichment status for individual cell types with one embodiment of the present invention. As indicated above, anti-CD59-PE and anti-CD61-PE represent preferred first and second fluorescent reagents of the present invention for labeling GPI anchor-expressing erythrocytes and platelets, respectively. Anti-PE magnetic beads are preferred reagents for separating platelets and GPI anchor-expressing erythrocytes from the GPI anchor-deficient erythrocytes, and SYTO®13 represents a preferred third fluorescent reagent for differentially labeling normochromatic erythrocytes from reticulocytes and leukocytes. The reagents identified in Figure 2 are exemplary, and the invention is not limited to practice with the reagents illustrated in Figure 2.

Figure 3 shows flow cytometric bivariate plots (SYTO®13 fluorescence vs. anti-CD59-PE fluorescence) for a blood specimen from an N-ethyl-N-nitrosourea-treated rat that was processed according to one embodiment of the present invention. The plot to the left shows the results from a pre-magnetic field analysis, whereas the plot to the right shows a post-magnetic field analysis. As shown in the pre-magnetic field analysis (left plot), the wild-type cells (high PE fluorescence signal) vastly outnumber the mutant phenotype cells (low PE fluorescence signal), despite exposure to a potent mutagen. In contrast, as shown in the post-magnetic field analysis (right plot), the wild-type cells are depleted and the mutant phenotype cells (low PE fluorescence signal) are enriched. From these two analyses mutant phenotype RBC and mutant phenotype RET frequencies can be calculated as described herein. The pre-magnetic field analysis is used to determine the total percentage of RETs, the total RBC to counting bead ratio, and the total RET to counting bead ratio. The post-magnetic field analyses are used to determine the mutant phenotype RBC to counting bead ratio and the total mutant phenotype RET to counting bead ratio.

Figure 4 shows observed mutant phenotype cell frequencies ( $\times 10^{-6}$ ) versus known, relative mutant cell frequencies. Data points depicted by squares represent pre-magnetic field analyses (on average 1,323,693 RBCs interrogated per sample). Data points depicted by circles represent post-magnetic field analyses (on average 112,632,575 RBCs interrogated per sample). The  $R^2$  values are linear correlation coefficients.

Figures 5A-5C show longitudinal data from a 28-day 1,3-propane sultone (1,3-PS) study. Figure 5A shows the mean percentage of reticulocytes (%RETs) versus time. Figure 5B shows the mean mutant reticulocyte (RET<sup>CD59-</sup>) frequencies versus time. Figure 5C shows the mean mutant erythrocyte (RBC<sup>CD59-</sup>) frequencies versus time. Error bars signify standard error of the mean. Asterisks indicate statistical significance compared to same-day vehicle

control values (Dunnett's t-test,  $p < 0.05$ ; note that analyses were performed on log-transformed data, therefore statistical comparisons were based on log means rather than arithmetic means).

## DETAILED DESCRIPTION OF THE INVENTION

5

**[0025]** The present disclosure relates to methods and kits for the measurement of RET frequency, *Pig-A* mutant phenotype RBC frequency, and *Pig-A* mutant phenotype RET frequency. These methods and kits can be used with an optical device designed for illumination and analysis of cell specimens.

10

**[0026]** Preferably, these measurements are accomplished using flow cytometry technology. The advantageous characteristics of this assay relative to other *in vivo* mutation assays which have been reported to date are that it does not require transgenic animals, special breeding programs, or cell culture; and it is compatible with all mammalian species. Other advantages will become apparent in the discussion of the various embodiments.

15

**[0027]** With this method, blood specimens are obtained from mammals. If the exposure that one wishes to evaluate is acute, then a period of expression time is allowed to occur before samples are collected into an anticoagulant solution according to standard practices. If the exposure that one wishes to evaluate has been protracted, for instance as occurs for subchronic or chronic toxicity tests, then it will usually not be necessary to allow for additional expression time before blood samples are collected. Alternatively, multiple samples can be collected over a period of time to monitor such long-term exposure.

20

**[0028]** Peripheral blood samples obtained from a subject are preferably treated in a manner effective substantially to separate RBCs from platelets and leukocytes. This is illustrated in Figure 1, step A. In one embodiment, the sample is enriched for RBCs such that leukocytes and platelets amount to less than about 0.17% of events per sample, more preferably less than about 0.01% of events per sample. One approach for achieving this degree of separation involves separation via centrifugation in an appropriate density gradient, e.g., Lympholyte®-Mammal (Cedarlane Laboratories, Burlington, NC), which affords a cell pellet that can be resuspended, e.g., in balanced salt solution, to form an enriched RBC sample. Other commercially available density gradient products that can be used for this purpose exist, for instance Ficoll™ PM400 has a more than 30 year track record for blood cell separation procedures, and is based on a polysucrose and sodium diatrizoate formulation. Additional products include Ficoll-Paque PLUS™, Ficoll-Paque PREMIUM™, Percoll™, and Percoll PLUS™. Whereas some of these products can be purchased at densities that are optimized for use with human blood (i.e., 1.077 g/mL), others have been optimized for mouse and rat blood (1.084 g/mL). Whatever reagent(s) are used for this purpose, the enriched RBC sample can then be treated in the manner described herein.

25

30

**[0029]** To achieve *Pig-A* mutant cell scoring according to the present invention, the enriched RBC sample is incubated with a first fluorescent reagent that binds to GPI-anchor competent (wild-type) RBCs, but not GPI-anchor deficient (*Pig-A* mutant) RBCs. This is illustrated in Figure 1, step B. By using a first fluorescent reagent that is either directly conjugated to a fluorochrome, or else one that can be readily bound to a subsequently applied fluorochrome, secondary antibody, or other tag (e.g., biotin), differential labeling of wild-type and mutant phenotype cells can be achieved. Ideally, the specificity of the interaction is high, for instance that which is typically found with antibody-antigen interactions.

35

**[0030]** Preferred GPI-anchor specific antibodies include, without limitation, fluorochrome-conjugated anti-CD59 and/or anti-CD24 and/or anti-CD55, as well as mixtures thereof. Alternatively, these antibodies can be used in combination with secondary antibodies labeled with a fluorochrome.

40

**[0031]** Subsequent to or concurrently with differential labeling of mutant phenotype versus wild-type RBCs, the enriched blood sample is contacted with a second fluorescent reagent that specifically binds to platelets (although platelets have largely been eliminated with the reagent used in a previous step to leuko- and platelet-deplete blood samples, some platelets can remain and interfere with analysis). This is also illustrated in Figure 1 at step B. The second fluorescent reagent can be conjugated directly to a fluorochrome or can be readily bound to a subsequently applied fluorochrome. By using a fluorescent reagent with a different emission spectrum as the first reagent fluorochrome, it is possible to specifically label platelets and thereby exclude them from mutant cell frequency measurements based on this fluorescence signal.

45

**[0032]** Alternately, it is possible to use the same fluorochrome to label platelets as was used to label wild-type erythrocytes. This one "shared" fluorochrome is appropriate when paramagnetic particles directed against the shared fluorochrome are used to deplete samples of wild-type cells. In this case, platelets are retained along with the wild-type RBCs in the magnetic field and are prevented from interfering with mutant cell scoring. Of the rare platelets that pass through the initial and subsequent physical depletions steps, light scatter is able to provide further differentiation from RBCs. Of the rare platelets that pass through the physical depletions steps as well as light scatter gating criteria, their first fluorochrome-positive phenotype gives them a wild-type RBC characteristic, a situation that does not appreciably effect the rare mutant phenotype RBC or mutant phenotype RET frequency measurements.

50

55

**[0033]** Preferred platelet-specific antibodies include, without limitation, anti-CD61 and/or anti-CD42b, as well as mixtures thereof. These antibodies can be directly conjugated to a fluorochrome or can be readily bound to a subsequently applied fluorochrome

**[0034]** Subsequent to differential labeling of mutant phenotype versus wild-type RBCs, the enriched blood sample is contacted with a reagent that includes paramagnetic particles and specifically recognizes wild-type RBCs or the fluorochrome, secondary antibody, or other tag associated with the first fluorescent reagent bound to wild-type RBCs. This is illustrated in Figure 1 at step C. The paramagnetic particles facilitate the separation of the wild-type and mutant RBCs. The paramagnetic particles may also specifically recognize platelets, or the fluorochrome or antibody associated with the second fluorescent reagent bound to the labeled platelets. This facilitates the removal or separation of platelets from the sample.

**[0035]** One example of paramagnetic bead-containing reagent includes anti-first fluorochrome beads, for instance Anti-Phycoerythrin (PE) or Anti-Fluorescein (FITC) MicroBeads from Miltenyi Biotec (Germany). Figure 3 illustrates how effective these particular Anti-PE MicroBeads can be. Here, flow cytometric bivariate plots (SYTO®13 fluorescence vs. anti-CD59-PE fluorescence) are presented for a blood specimen from an N-ethyl-N-nitrosourea-treated rat that was processed according to one embodiment of the present invention. The plot to the left shows the results from a pre-magnetic field analysis, whereas the plot to the right shows a post-magnetic field analysis. As shown in the pre-magnetic field analysis (left plot), wild-type cells (high PE fluorescence signal) ordinarily vastly outnumber mutant phenotype cells (low PE fluorescence signal). In contrast, as shown in the post-magnetic field analysis, the effective use of Anti-PE MicroBeads can severely deplete samples of wild-type cells and thereby greatly enrich samples for mutant phenotype cells (low PE fluorescence signal).

**[0036]** Following sufficient exposure of the enriched sample to the paramagnetic particles, typically on the order of 15 to 30 minutes, the sample is then placed in a magnetic field that is sufficiently strong enough to hold the majority of the wild-type RBCs in the magnetic field while the eluate becomes enriched for mutant phenotype RBCs. This is illustrated in Figure 1 at step E. This may be accomplished with the aid of a column, for instance one that is packed with ferromagnetic spheres. The eluate, now depleted (or very nearly depleted) of wild-type RBCs and enriched for mutant phenotype RBCs, is then concentrated.

**[0037]** The sample is contacted with a third fluorescent reagent that specifically binds to nucleic acids. This is also illustrated in Figure 1 at step E. By using a third fluorescent reagent containing a fluorochrome, differential labeling of NCEs, RETs, and any remaining leukocytes is achieved. In one embodiment of the present invention, the concentrated eluate, depleted of wildtype RBCs and enriched for mutant phenotype RBCs is contacted with the third fluorescent reagent that specifically binds to nucleic acids. Exemplary third fluorescent reagents include, without limitation, cyanine dyes, such as those in the SYTO® family of nucleic acid dyes, especially SYTO®13, SYTO®59, SYTO®83, SYTO® RNA Select™ (all available from Invitrogen, USA). Other suitable third fluorescent reagents include thiazole dyes (e.g., thiazole orange) and acridine orange. Of these, SYTO®13 dye is preferred.

**[0038]** In an alternative aspect of the disclosure, the enriched erythrocyte sample is contacted with the third fluorescent reagent prior to separation or enrichment for the mutant RBC phenotype. Exemplary third fluorescent reagents for use in this aspect include reagents that permeate the cells and covalently bind to DNA following photo-induction, e.g., ethidium monoazide (EMA) and propidium monoazide (PMA). In accordance with this aspect, the cells are fixed or otherwise permeabilized using known procedures and reagents prior to exposure to the third fluorescent reagent.

**[0039]** The labeled samples are then subjected to optical detection to enumerate *Pig-A* mutant cells using any suitable optical detection system. Preferred optical detection systems have one or more light sources, preferably in the form of one or more amplified or collimated beams of light, that are able to excite the fluorescent reagents. Exemplary optical detection systems include, without limitation, single-laser flow cytometers and dual- or multiple-laser flow cytometers.

**[0040]** Single-laser flow cytometric analysis uses a single focused laser beam with an appropriate emission band to excite the several fluorescent reagents. Dual- or multiple-laser flow cytometric analysis use two or more focused laser beams with appropriate emission bands, in much the same manner as a single-laser flow cytometer. Different emission bands afforded by the two or more lasers allow for additional combinations of fluorescent dyes or immunochemical-conjugated fluorochromes to be employed.

**[0041]** As labeled cells pass through the focused laser beam, they exhibit a fluorescent emission maxima characteristic of the fluorochromes or dyes associated therewith. The flow cytometer is equipped with appropriate detection devices to enable detection of the fluorescent emissions and light scatter produced by the cells. In this way, mutant cell populations (i.e., mutant erythrocytes and/or reticulocytes) are counted.

**[0042]** Cell population counts can be expressed as cells per unit volume of sample (i.e., cell density measurement) or per unit time (based on the fluidic rate and the time taken to analyze the sample). Alternatively, counting beads can be added to the sample and the fluorescent emission and light scatter of the counting beads is detected and counted along with the enumerated cell populations to obtain a cell-to-bead ratio. The counting beads can be a suspension of latex particles or similar uniform particle that can be readily differentiated from the cells. Preferred latex particles include, without limitation, CountBright™ Absolute Counting Beads from Invitrogen. In one embodiment of the present invention, such counting beads are added after erythrocytes have been labeled with antibodies and contacted with paramagnetic beads that recognize GPI-anchor-expressing cells. This is schematically illustrated in Figure 1 at step D<sub>1</sub>, and Figure 2. However, it will be appreciated by those knowledgeable in the art that there are alternate and equally acceptable times

during the labeling and cell separation procedures when counting beads can be added and used effectively to obtain the desired cell population counts. For instance, in one alternative embodiment of the present invention, counting beads are added to an isotonic buffer solution that includes the third fluorescent reagent used to label nucleic acids. In this case, the dilution factor used to create pre-column samples, and the volume of post-column eluates are important for

determining the pre- and post-column cell-to-Counting Bead ratios that are used to calculate mutant cell frequencies. **[0043]** In order to calculate the frequency of mutant phenotype erythrocytes and/or mutant phenotype reticulocytes relative to total erythrocytes and/or reticulocytes, a quantitative measure of the total number of erythrocytes and reticulocytes in the enriched erythrocyte sample is obtained from the pre-separation sample. This is illustrated in Figure 1 at step D<sub>2</sub>. The quantitative measurement can be a cell per volume unit measurement, a cell per unit time measurement, or cell-to-bead ratio as described above. To obtain an accurate frequency calculation, both cell counts (i.e., the mutant phenotype cell counts and the overall total sample cell counts) should be obtained and expressed in the same manner.

**[0044]** While immunodetection reagents are described for use in the methods of the present invention, it should be appreciated that any suitable immunolabel can be used, including without limitation monoclonal antibodies, polyclonal antibodies, mono-specific polyclonal antibody preparations, chimeric antibodies, single chain antibodies, synthetic antibodies, and any antibody fragments, e.g., Fab fragments, Fab' fragments, F(ab)<sub>2</sub> fragments, F(ab')<sub>2</sub> fragments, Fd fragments, Fv fragments, dAb fragments, and isolated complementarity determining regions ("CDRs") (see U.S. Patent Nos. 7,037,498, 7,034,121, 7,041,870, and 7,074,405). These antibody fragments can be made by conventional procedures, such as proteolytic fragmentation procedures, as described in J. Goding, *Monoclonal Antibodies: Principles and Practice*, pp. 98-118 (N.Y. Academic Press 1983). Methods for preparing antibodies that are specific to an analyte of interest are well known in the art. Conjugation of desired fluorochromes to antibodies is also well known in the art, and such immunoreagents are commercially available.

**[0045]** Whereas the fluorochrome associated with the first and second fluorescent reagents may be the same, the nucleic acid dye and counting beads, if used, should be selected such that they can all be excited by the one or more light sources, yet their emission spectra are sufficiently distinct.

**[0046]** Samples that have been contacted with three fluorescent reagents as described should be stored at about 4°C and protected from light until ready for analysis. In a preferred embodiment of the present disclosure, samples are leuko- and platelet-depleted, stained, and analyzed on the same day of harvest. Alternately, the samples are maintained in a suitable anticoagulant or else leuko- and platelet-depleted, and then stored refrigerated until they can be stained and analyzed, preferably within approximately five days of harvest.

**[0047]** The low frequency of *Pig-A* mutant cells in healthy mammals, for instance those that are untreated, sham-exposed or vehicle-treated, demands that a sensitive mutation scoring system be capable of interrogating at least several hundred cells per individual, but more ideally greater than 10<sup>6</sup> cells per individual. Unless considerably more than 10<sup>6</sup> cells per individual are interrogated per individual, the mutation frequency measurement is a rough approximation, since the spontaneous frequency is thought to be on the order of 1 x 10<sup>6</sup>. Therefore, to enhance the precision of mutation frequency measurements, especially at or near spontaneous levels, it is important to interrogate many millions of cells for the mutant phenotype. These numbers of cells have been difficult to acquire with current methods, especially for the RET subpopulation of cells. For instance, in Dertinger et al., "Integration of Mutation and Chromosomal Damage Endpoints into 28-Day Repeat Dose Toxicology Studies," *Toxicol. Sci.* 115, 401 - 411 (2010), the investigators stopped data acquisition once 10<sup>6</sup> total RBCs were acquired, and in a separate analysis that employed SYTO®13 thresholding, when approximately 0.3 x 10<sup>6</sup> RETs were acquired (per sample). These stop modes were dictated by the practical consideration of time spent on the flow cytometer—twenty plus minutes per sample. These stop modes led to a majority of vehicle control analyses returning zero values, a less than ideal situation that diminishes the ability of the assay to detect modest changes to mutation frequency.

**[0048]** While it requires less time to acquire millions of RBCs as opposed to RETs, there are at least two reasons why there is merit in performing *Pig-A* mutation measurements in the RET fraction in addition to the total RBC pool.

**[0049]** One advantage of RET-based measurements is demonstrated by work with the peripheral blood cells of PNH patients. It has been shown that the number of circulating GPI-deficient neutrophils is typically higher than the number of affected RBCs. One explanation for this finding is that PNH RBCs are subject to complement-mediated intravascular lysis. Therefore, it is possible that an analytical system that focuses on the newly formed RETs would provide a more accurate mutation frequency. At least one previous report supports this view insofar as staining to identify RETs in the blood of PNH patients showed that the percentage of abnormal (GPI-deficient) RETs was similar to the percentage of affected neutrophils (Ware et al., "Immunophenotypic Analysis of Reticulocytes in Paroxysmal Nocturnal Hemoglobinuria," *Blood* 86:1586-1589 (1995)). These data indicate that premature destruction of PNH RBCs may account for the fact that most patients have more GPI-deficient neutrophils than RBCs. Thus, a mutation assay based on RETs may be more sensitive than one based on total RBCs, especially if mutant RBCs exhibit a shortened lifespan.

**[0050]** It is important to note, however, that the specimens analyzed by Ware *et al.* were from PNH patients that expressed high frequencies of mutant RBCs, in the range of approximately 20 to 95%. These high frequencies are not found in non-PNH mammals, even following exposure to potent mutagens. Therefore, Ware *et al.* were not required to

utilize a methodology that was capable of accurately enumerating GPI anchor-deficient cells at baseline and near-baseline frequencies, as is the case for a mutation assay system as described herein.

5 [0051] A second reason for studying *Pig-A* mutation in RETs is that this subpopulation would be expected to reflect genotoxicant-induced mutation more rapidly than the total RBC pool (Phonethepswath et al., "Pig-a Mutation Kinetics in Rat Erythrocytes Following Exposure to Five Prototypical Mutagens," *Toxicol. Sci.* 114, 59 - 70 (2010)). That is, mutation frequency in the RET population would be expected to be a "leading indicator" of genotoxicant exposure. This temporal relationship reflects the fact that the spontaneous mutant frequency can only be affected by genotoxicant exposure after a sufficient length of time has elapsed, one that allows for a significant fraction of pre-existing (low mutation frequency) cells to be replaced by cells that exhibit an elevated mutation frequency. The time-frame for turning over blood RETs (days) is considerably shorter than for the total RBC pool (several weeks to about 4 months, depending on species), and is consequently expected to provide a quickly responding cohort, versus one that lags in time. Thus, shortening of mutant cell expression time is therefore a major advantage for studying *Pig-A* mutation in blood RETs.

10 [0052] While rapid interrogation of erythrocytes for the extremely rare *Pig-A* mutation phenotype is clearly desirable, it is not easily achieved with current methods, especially for the RET subpopulation. In particular, the low incidence of RETs relative to total RBCs poses technical challenges. Despite the high throughput capacity of modern flow cytometers (often on the order of 7000 events per second), the interrogation of approximately  $10^6$  RETs for mutation is time consuming. At this rate, a whole blood specimen with 3% RETs would require approximately 80 minutes to interrogate  $10^6$  RETs. The method reported herein overcomes the low throughput capacity of previously described approaches, enabling researchers to interrogate many times more RBCs and RETs than was previously possible, and in a fraction of the time. A second consideration is that data file sizes are extremely large when flow cytometric data for several hundred or more RETs per sample are acquired, and all the mature RBCs that accompany them are also saved to the same data file.

15 [0053] One significant use of the present invention relates to genotoxicity assessment. In this case, an exogenous test agent is applied over a range of doses or intensities to mammals of interest. This test agent exposure may occur one or several times as is the case in acute or subacute toxicity tests, or repeatedly as is the case in subchronic and chronic toxicity tests. The test agent may be a chemical or formulation, or it could be a physical entity, such as energy. Chemicals which are known to damage DNA include, but are not limited to: certain metals (e.g., arsenic, cadmium and nickel), alkylating agents (e.g., N-ethyl-N-nitrosourea, methyl methanesulfonate, etc.), intercalating agents (e.g., proflavin), anti-metabolites (e.g., 5-fluorouracil), organic genotoxicants that are generated by combustion processes (e.g., polycyclic aromatic hydrocarbons such as benzo(a)pyrene), as well as organic genotoxicants that are found in nature (e.g., aflatoxins such as aflatoxin B1). Examples of physical agents that are known to damage DNA include, but are not limited to: X-rays, gamma radiation, neutron radiation, beta radiation, and UV radiation.

20 [0054] After an appropriate length of expression time which allows mutated bone marrow stem cells or erythroid progenitor cells to appear in peripheral blood as GPI-anchor deficient RETs or RBCs (i.e., from several days to several weeks post exposure), blood is harvested and prepared for flow cytometric enumeration of *Pig-A* mutants according to procedures outlined above and described in detail in the following examples.

25 [0055] Certain agents may offer protection from DNA damage, while others may magnify risk of damage. The present invention can also be used to evaluate the effects of an agent which can modify (i.e., enhance or suppress) such damage. To assess the suspected protective effects of an agent, mammals can be exposed to the putative protective agent either prior to, concurrently, or soon after exposure to a known genotoxicant. Any protective effect afforded by the agent can be measured relative to damage caused by the genotoxicant alone. Putative protective agents can be vitamins, bioflavonoids and anti-oxidants, dietary supplements (e.g., herbal supplements), or any other protective agent, whether naturally occurring or synthesized by man.

30 [0056] To assess the ability of an agent to synergistically or additively enhance genotoxicity, mammals can be exposed to the agent prior to, concurrently, or shortly after exposure to a known genotoxicant. Any additive or synergistic effect caused by the agent can be measured relative to damage caused by the genotoxicant alone.

35 [0057] The assays of the present invention can likewise be used to monitor chronic exposure to genotoxicant agents, for example, in individuals that work in environmental remediation, manufacturing industries that involve exposure to such agents, agricultural environments that involve pesticide or insecticide usage, etc. For these uses, period testing can be performed on a regular basis (e.g., weekly, monthly, quarterly, seasonally, biannually, etc.).

## 50 EXAMPLES

[0058] The examples below are intended to exemplify the practice of the present invention but are by no means intended to limit the scope thereof.

### 55 Example 1-Reproducibility of Technical Replicates

[0059] One male Sprague Dawley rat (8 - 10 weeks old) was treated on three consecutive days (i.e., days 1, 2, and

EP 2 643 694 B1

3) with either vehicle (water) or else 60 mg 1,3-propane sultone (PS) per kilogram body weight per day *via* oral gavage. Cardiac puncture blood specimens were collected on day 29, that is 26 days after the last administration. Each blood sample was processed (i.e., stained and analyzed *via* flow cytometry) three separate times to evaluate the reproducibility of cell handling procedures and the analytical platform.

**[0060]** Figure 1 and 2 illustrate the cell processing procedures used in this study. Specimens of approximately 80  $\mu$ l whole blood were first debulked of leukocytes and platelets *via* centrifugation through Lympholyte®-Mammal. Incubation with anti-CD59-PE provided differential labeling of wild-type and mutant phenotype RBCs, while concurrent exposure to anti-CD61-PE endowed rare contaminating platelets with PE fluorescence. Samples were subsequently contacted with Anti-PE MicroBeads from Miltenyi, and after a washing step *via* centrifugation, latex particles (CountBright™, from Invitrogen) were added in phosphate buffered saline and represented Counting Beads. One fraction of each blood sample was then stained with SYTO®13 and analyzed on a FACSCanto II flow cytometer running FACS Diva software for 3 minutes each. These "pre-magnetic field" specimens were used to generate the data that appear in Table I, which were used to calculate RBC: Counting Bead and RET:Counting Bead ratios.

**[0061]** After adding Counting Beads to the samples, the majority of each specimen was applied to LC Separation Columns (Miltenyi) that were suspended in a magnetic field (MidiMACS™ Separator, Miltenyi). 5 mL of phosphate buffered saline was used to elute mutant phenotype cells and Counting Beads, while the column held the vast majority of PE-labeled particles (i.e., wild-type RBCs and contaminating platelets). The eluate was pelleted *via* centrifugation and resuspended in 300  $\mu$ l phosphate buffered saline with SYTO®13. These samples were analyzed on the flow cytometer for 4 minutes each. These "post-magnetic field" specimens were used to generate the data that appear in Table II, which were used to calculate mutant phenotype RBC: Counting Bead and mutant phenotype RET:Counting Bead ratios.

**[0062]** Table III shows calculated RET frequencies (percentage) based on the pre-magnetic field analyses, as well as the incidence of mutant phenotype RBCs per 10<sup>6</sup> RBCs and the incidence of mutant phenotype RETs per 10<sup>6</sup> RETs that were calculated based on ratios determined in the pre- and post-magnetic field analyses. As these data demonstrate, treatment with the mutagenic chemical PS clearly led to elevated frequencies of mutant phenotype RBCs and RETs, and the results are reproducible across technical replicates.

**Table I. Flow Cytometric Analyses, Pre-magnetic Field, Technical Replicates**

PRE-Magnetic Field Analyses							
Treatment (Replicate)	Mutant RETs	Wt RETs	Mutant NCEs	Wt NCEs	Counting Beads	RBC:Bead Ratio	RET: Bead Ratio
Vehicle (1)	0	30,267	7	1,339,253	156	8,779.02	194.02
Vehicle (2)	0	27,387	9	1,207,862	144	8,578.18	190.19
Vehicle (3)	0	31,093	8	1,354,878	173	8,011.44	179.73
PS (1)	1	46,060	83	1,098,308	169	6,771.91	272.55
PS (2)	1	46,655	75	1,098,129	182	6,290.44	256.35
PS (3)	2	43,955	64	1,032,849	158	6,815.63	278.21

**Table II. Flow Cytometric Analyses, Post-magnetic Field, Technical Replicates**

POST-Magnetic Field Analyses							
Treatment (Replicate)	Mutant RETs	Mutant NCEs	Counting Beads	Mutant RBC:Bead Ratio	Mutant RET: Bead Ratio	Total RBC Equivalentents	Total RET Equivalentents
Vehicle (1)	1	84	16,106	0.00528	0.00006	141,394,884	3,124,874
Vehicle (2)	1	81	16,681	0.00492	0.00006	143,092,630	3,172,518
Vehicle (3)	1	92	12,803	0.00726	0.00008	102,570,457	2,301,062
PS (1)	95	6,466	15,856	0.41379	0.00599	107,375,331	4,321,557
PS (2)	79	7,590	18,558	0.41324	0.00426	116,737,977	4,757,374
PS (3)	94	7,054	19,070	0.41324	0.37483	129,974,120	5,305,443

Table III. Calculated Frequencies Derived From Pre- and Post-Magnetic Field Analyses, Technical Replicates

Treatment (Replicate)	%RET	Calculated Frequencies	
		Mutant RBCs per 10 <sup>6</sup> Total RBCs	Mutant RETs per 10 <sup>6</sup> Total RETs
Vehicle (1)	2.21	0.6	0.3
Vehicle (2)	2.22	0.6	0.3
Vehicle (3)	2.24	0.9	0.4
PS (1)	4.02	61.1	22
PS (2)	4.08	65.7	16.6
PS (3)	4.08	55	17.7

### Example 2-Spiking Experiment

**[0063]** One male Sprague Dawley rat (7 - 8 weeks old) was treated on fourteen consecutive days (i.e., days 1 - 14) with either vehicle (water) or 40 mg 1,3-propane sultone (1,3-PS) per kilogram body weight per day *via* oral gavage. Cardiac puncture blood specimens were collected on day 35 (i.e., 21 days after the last administration). The blood sample from the vehicle control rat was processed (i.e., stained and analyzed *via* flow cytometry) two separate times and is referred to herein as the "0" sample; blood from the 1,3-PS-treated rat was processed two separate times and is referred to herein as the "100" sample. To construct a specimen with an intermediate frequency of mutant cells, eight parts vehicle control blood were combined with two parts 1,3-PS blood and this spiked sample (referred to herein as the "20" sample) was processed two separate times. Another intermediate frequency specimen was prepared by combining nine parts vehicle control blood with one part 1,3-PS blood and this spiked sample (referred to herein as the "10" sample) was processed two separate times.

**[0064]** The eight specimens described above were processed as follows: 80  $\mu$ l whole blood were first debulked of leukocytes and platelets *via* centrifugation through Lympholyte®-Mammal. Incubation with anti-CD59-PE provided differential labeling of wild-type and mutant phenotype RBCs, while concurrent exposure to anti-CD61-PE endowed rare contaminating platelets with PE fluorescence. Samples were subsequently contacted with anti-PE MicroBeads from Miltenyi, and after a washing step *via* centrifugation, latex particles (CountBright™, from Invitrogen) were added in phosphate buffered saline and represented Counting Beads. One fraction of each blood sample was then stained with SYTO®13 and analyzed on a FACSCanto II flow cytometer running FACS Diva software for 3 minutes each. These pre-magnetic field specimens were used to generate the data that appear in Table IV, which were used to calculate RBC: Counting Bead and RET: Counting Bead ratios.

**[0065]** After adding Counting Beads to the samples, the majority of each specimen was applied to LC Separation Columns (Miltenyi) that were suspended in a magnetic field (MidiMACS™ Separator, Miltenyi). 5 mL of phosphate buffered saline was used to elute mutant phenotype cells and Counting Beads, while the column held the vast majority of PE-labeled particles (i.e., wild-type RBCs and contaminating platelets). The eluate was pelleted *via* centrifugation and resuspended in 300  $\mu$ l phosphate buffered saline with SYTO®13. These samples were analyzed on the flow cytometer for 4 minutes each. These post-magnetic field specimens were used to generate the data that appear in Table V, which were used to calculate mutant phenotype RBC:Counting Bead and mutant phenotype RET:Counting Bead ratios.

**[0066]** Table VI shows calculated RET frequencies (percentage) based on the pre-magnetic field analyses, as well as the incidence of mutant phenotype RBCs per 10<sup>6</sup> RBCs and the incidence of mutant phenotype RETs per 10<sup>6</sup> RETs that were calculated based on ratios determined in the pre- and post-magnetic field analyses.

**[0067]** Table VII shows mutant phenotype RBCs frequencies (per 10<sup>6</sup> RBCs) based on pre-magnetic field analyses as well as post-magnetic field analyses. Whereas the pre-magnetic field frequency determinations are based on an average of 1,323,693 RBCs per specimen, the post-magnetic field frequency measurements are based on an average of 112,632,575 RBCs per specimen. Figure 4 is a graph that depicts each observed mutant phenotype cell frequency against the known, relative mutant cell frequency. From these data it is clear that the invention described herein, which includes wild-type cell depletion steps, provides more precise enumeration of rare GPI-anchor deficient cells relative to analyses based on total erythrocytes that do not benefit from depletion step(s).

EP 2 643 694 B1

Table IV. Flow Cytometric Analyses, Pre-magnetic Field, Spiking Experiment

PRE-Magnetic Field Analyses							
Sample ID (Replicate)	Mutant RETs	Wt RETs	Mutant NCEs	Wt NCEs	Counting Beads	RBC:Bead Ratio	RET: Bead Ratio
0 (1)	0	37,666	7	1,206,954	186	6,691.54	202.51
0 (2)	0	42,322	10	1,356,514	193	7,247.91	219.28
10 (1)	0	45,106	5	1,290,058	151	8,842.18	298.72
10 (2)	0	44,928	8	1,336,885	182	7,592.42	246.86
20 (1)	0	47,520	11	1,292,199	151	8,872.38	314.70
20 (2)	0	48,649	9	1,308,497	159	8,535.57	305.97
100 (1)	1	79,656	36	1,265,814	168	8,008.97	474.15
100 (2)	1	70,002	31	1,116,652	162	7,325.22	432.12

Table V. Flow Cytometric Analyses, Post-magnetic Field, Spiking Experiment

POST-Magnetic Field Analyses							
Treatment (Replicate)	Mutant RETs	Mutant NCEs	Counting Beads	Mutant RBC:Bead Ratio	Mutant RET: Bead Ratio	Total RBC Equivalents	Total RET Equivalents
0 (1)	2	130	16,503	0.00800	0.00012	110,430,534	3,341,946
0 (2)	0	150	16,041	0.00935	0.00000	116,263,672	3,517,550
10 (1)	20	449	13,819	0.03394	0.00145	122,190,069	4,127,946
10 (2)	18	502	13,998	0.03715	0.00129	106,278,738	3,455,506
20 (1)	50	720	13,036	0.05907	0.00384	115,660,399	4,102,455
20 (2)	35	765	13,509	0.05922	0.00259	115,306,962	4,133,329
100 (1)	173	3,304	14,475	0.24021	0.01195	115,929,844	6,863,304
100 (2)	182	3,192	13,515	0.24965	0.01347	99,000,378	5,840,065

Table VI. Calculated Frequencies Derived From Pre- and Post-Magnetic Field Analyses, Spiking Experiment

Calculated Frequencies			
Treatment (Replicate)	%RET	Mutant RBCs per 10 <sup>6</sup> Total RBCs	Mutant RETs per 10 <sup>6</sup> Total RETs
0 (1)	3.03	1.2	0.6
0 (2)	3.03	1.3	0
10 (1)	3.38	3.8	4.8
10 (2)	3.25	4.9	5.2
20 (1)	3.55	6.7	12.2
20 (2)	3.58	6.9	8.5
100 (1)	5.92	30	25.2
100 (2)	5.9	34.1	31.2

Table VII. Calculated Frequencies, Pre- versus Post-Magnetic Field Analyses, Spiking Experiment

		Calculated Frequencies	
5	Treatment (Replicate)	Mutant RBCs per 10 <sup>6</sup> Total RBCs (Pre-Column)*	Mutant RBCs per 10 <sup>6</sup> Total RBCs (Post-Column)**
	0 (1)	6	1.2
	0 (2)	7	1.3
10	10 (1)	4	3.8
	10 (2)	6	4.9
	20 (1)	8	6.7
	20 (2)	7	6.9
	100 (1)	27	30
15	100 (2)	27	34.1

\* Avg. number of RBCs interrogated per sample = 1,323,693  
\*\* Avg. number of RBCs interrogated per sample = 112,632,575

### 20 Example 3-*In Vivo* Responses to the Mutagenic Chemical 1,3-Propane Sultone

**[0068]** Groups consisting of six male Sprague Dawley rats (7 - 8 weeks old) were treated on twenty-eight consecutive days (i.e., days 1 through 28) with either water (vehicle), 12.5 mg 1,3-PS per kilogram body weight per day, 25 mg 1,3-PS per kilogram body weight per day, or 50 mg 1,3-PS per kilogram body weight per day *via* oral gavage. Note that one rat in the high dose group died on day 11 and others in this group continued to lose weight, therefore after 14 days of administering 50 mg/kg/day, the top dose was reduced to 37.5 mg/kg/day for the remaining 14 days of administration. Tail vein blood specimens were collected on day -1 (i.e., one day before the start of treatment), and again on days 15, 29, and 42.

**[0069]** Each blood sample was processed according to procedures outlined in Figures 1 and 2. More specifically, specimens consisting of 80  $\mu$ l whole blood each were first debulked of leukocytes and platelets *via* centrifugation through Lympholyte®-Mammal. Incubation with anti-CD59-PE provided differential labeling of wild-type and mutant phenotype RBCs, while concurrent exposure to anti-CD61-PE endowed rare contaminating platelets with PE fluorescence. Cells were subsequently contacted with Anti-PE MicroBeads from Miltenyi, and after a washing step *via* centrifugation, latex particles (CountBright™, from Invitrogen) were added in phosphate buffered saline and represented Counting Beads. One fraction of each blood sample was then stained with SYTO®13 and analyzed on a FACSCalibur flow cytometer running CellQuest Pro software for 3 minutes each. These pre-magnetic field specimens were used to generate the data used to calculate RBC:Counting Bead and RET:Counting Bead ratios.

**[0070]** After adding Counting Beads to the samples, the majority of each specimen was applied to LC Separation Columns (Miltenyi) that were suspended in a magnetic field (QuadroMACS™ Separator, Miltenyi). 5 mL of phosphate buffered saline was used to elute mutant phenotype cells and Counting Beads, while the column held the vast majority of PE-labeled particles (i.e., wild-type RBCs and contaminating platelets). The eluate was pelleted *via* centrifugation and resuspended in 300  $\mu$ l phosphate buffered saline with SYTO®13. These samples were analyzed on the flow cytometer for 3 - 3.5 minutes each. These post-magnetic field specimens were used to generate the data used to calculate mutant phenotype RBC:Counting Bead and mutant phenotype RET:Counting Bead ratios.

**[0071]** Longitudinal %RET data are shown in Figure 5A. Day 15 %RET were reduced in a dose-dependent manner, and a rebound effect was evident at the top two dose levels at later time points with %RET increasing and ultimately exceeding vehicle control values.

**[0072]** As shown by Figure 5B, elevated mutant phenotype cells were first observed in the RET population. These increases occurred in a dose-dependent manner, and each dose group was significantly elevated relative to vehicle controls at every time point studied. While a modest reduction to the top dose group's mean mutant RET frequency was observed between days 29 and 42, the other dose groups exhibited similar or slightly elevated values over this same timeframe.

**[0073]** Mutant RBC frequencies increased at a slower rate compared to mutant RETs, with maximal observed values occurring on day 42 (Figure 5C). Each of the 1,3-PS dose groups exhibited elevated frequencies that were significantly higher than those observed in vehicle-treated rats from Day 15 through Day 42, even at Day 15 where the increases over the baseline were relatively small.

**[0074]** The immunomagnetic separation procedure proved capable of analyzing large numbers of RETs and RBCs for the mutant phenotype. For example, considering the day 29 time point, the average number of cell equivalents

interrogated for CD59 expression was  $1.18 \times 10^8$  RBCs and  $2.93 \times 10^6$  RETs per sample. As these values are on the order of 100x and 10x higher than previous work with these endpoints (Phonethepswath et al., "Pig-a Mutation: Kinetics in Rat Erythrocytes Following Exposure to Five Prototypical Mutagens," *Toxicol. Sci.* 114:59-70 (2010); Dertinger et al., "Integration of Mutation and Chromosomal Damage Endpoints into 28-Day Repeat Dose Toxicology Studies," *Toxicol. Sci.* 115:401-411 (2010)), it was of interest to evaluate the statistical power of this new methodology. As shown in Table VIII, given a group size of 6 rats, 3-fold increases in mutant RETs and 2-fold increases in mutant RBCs were detectable with greater than 80% power.

Table VIII. Power Analyses, Based on Day 29 Data

Parameter	Increase Over Baseline	Chance of Detecting
Mutant RETs	2x	49%
	2.5x	69%
	3x	82%
	3.5x	90%
	4x	95%
Mutant RBCs	2x	90%
	2.5x	99%
	3x	99.9%
	3.5x	> 99.9%
	4x	> 99.9%

Example 4- In Vivo Responses to Diverse Mutagenic Chemicals

[0075] The same experimental design used in Example 3 was used to study several additional mutagenic chemicals agents. That is, groups of six male Sprague Dawley rats (age 7 - 8 weeks) were treated for twenty-eight consecutive days with vehicle, low, mid, and high dose levels of each test article via oral gavage. High dose levels approximated maximum tolerated doses, mid dose was one-half of the high dose, and low dose was one-quarter of the high dose. The identity of these chemicals and their primary mode of action are presented in Table IX.

[0076] Tail vein blood samples were collected from each animal over time, specifically on day -1 (i.e., one day before the start of treatment), and again on days 15, 29, and 42. Each blood sample was processed according to procedures outlined in Figures 1 and 2. More specifically, specimens consisting of 80 µl whole blood each were first debulked of leukocytes and platelets via centrifugation through Lympholyte®-Mammal. Incubation with anti-CD59-PE provided differential labeling of wild-type and mutant phenotype RBCs, while concurrent exposure to anti-CD61-PE endowed rare contaminating platelets with PE fluorescence. Cells were subsequently contacted with Anti-PE MicroBeads from Miltenyi, and after a washing step via centrifugation, latex particles (CountBright™, from Invitrogen) were added in phosphate buffered saline and represented Counting Beads. One fraction of each blood sample was then stained with SYTO®13 and analyzed on a FACSCalibur flow cytometer running CellQuest Pro software for 3 minutes each. These pre-magnetic field specimens were used to generate the data used to calculate RBC:Counting Bead and RET:Counting Bead ratios.

[0077] After adding Counting Beads to the samples, the majority of each specimen was applied to LC Separation Columns (Miltenyi) that were suspended in a magnetic field (QuadroMACS™ Separator, Miltenyi). Five milliliters of phosphate buffered saline was used to elute mutant phenotype cells and Counting Beads, while the column held the vast majority of PE-labeled particles (i.e., wild-type RBCs and contaminating platelets). The eluate was pelleted via centrifugation and resuspended in 300 µl phosphate buffered saline with SYTO®13. These samples were analyzed on the flow cytometer for 3 - 3.5 minutes each. These post-magnetic field specimens were used to generate the data used to calculate mutant phenotype RBC:Counting Bead and mutant phenotype RET:Counting Bead ratios.

[0078] Each of these diverse genotoxic chemicals was observed to significantly increase the frequency of circulating mutant RETs and RBCs. The data are summarized in Table IX, which provides information about the time point at which the maximal effect was observed for both the mutant RET and mutant RBC endpoints.

Table IX. Maximum Mutant Cell Responses for a Series of Diverse Genotoxicants

Chemical	Mode of Genotoxic Action	Maximum Avg. Mutant Cell Freq. x 10 <sup>-6</sup> (Day Maximum Value was Observed)
Thiotepa	Alkylating agent, organophosphorus compound	79 Mutant RETs (d29) ; 69 Mutant RBCs (d42)

(continued)

Chemical	Mode of Genotoxic Action	Maximum Avg. Mutant Cell Freq. x 10 <sup>-6</sup> (Day Maximum Value was Observed)
5 2-Acetylaminofluorene	Aromatic amine	50 Mutant RETs (d42) ; 58 Mutant RBCs (d29)
Chlorambucil	Alkylating agent, nitrogen mustard	40 Mutant RETs (d42) ; 21 Mutant RBCs (d42)
Melphalan	Alkylating agent, nitrogen mustard	33 Mutant RETs (d42) ; 23 Mutant RBCs (d42)
Cyclophosphamide	Alkylating agent, nitrogen mustard	5 Mutant RETs (d29) ; 3 Mutant RBCs (d42)
10 Azathioprine	Purine analog	13 Mutant RETs (d29) ; 4 Mutant RBCs (d42)
1,3-Propane Sultone (see Example 3 above)	Alkylating agent, cyclic sulfonate ester	106 Mutant RETs (d29) ; 72 Mutant RBCs (d42)

Note that untreated male Sprague Dawley rats' historical control mean mutant RET and mutant RBC frequencies are < 1 x 10<sup>-6</sup>.

**[0079]** All of the features described herein (including any accompanying claims, abstract and drawings), and/or all of the steps of any method or process so disclosed, may be combined with any of the above aspects in any combination, except combinations where at least some of such features and/or steps are mutually exclusive.

### Claims

1. A method for the enumeration of in vivo *Pig-A* gene locus mutation, the method comprising:

25 providing an enriched erythrocyte sample obtained from a mammal exposed to an exogenous agent, said sample comprising normochromatic erythrocytes and reticulocytes, and having a reduction in the frequency of platelets and leukocytes;  
 30 contacting the enriched erythrocyte sample with a first fluorescent reagent that labels GPI anchor-expressing erythrocytes, but not GPI anchor-deficient erythrocytes, and a second fluorescent reagent that specifically labels platelets;  
 separating the enriched erythrocyte sample into a first portion comprising platelets and GPI anchor-expressing erythrocytes and a second portion comprising GPI anchor-deficient erythrocytes;  
 35 contacting the second portion with a third fluorescent reagent that differentially labels normochromatic erythrocytes from reticulocytes and leukocytes;  
 exciting the first, second, and third fluorescent reagents in the second portion with light of appropriate excitation wavelength, wherein the third fluorescent reagent has a fluorescent emission spectrum that does not substantially overlap with the fluorescent emission spectra of the first or second fluorescent reagents; and  
 40 detecting the fluorescent emission and light scatter produced by the GPI anchor-deficient erythrocytes of the second portion labeled with the third fluorescent reagent, while excluding contaminating GPI anchor-expressing erythrocytes and reticulocytes, platelets, and leukocytes, and counting the number of GPI anchor-deficient erythrocytes and reticulocytes.

2. The method according to claim 1, wherein the first fluorescent reagent comprises an anti-CD59, anti-CD24, or anti-CD55 antibody, or a combination thereof.

3. The method according to claim 1 or 2, wherein the second fluorescent reagent comprises an anti-CD61 antibody or anti-CD42b antibody, or a combination thereof.

4. The method according to one of claims 1-3, wherein the third fluorescent reagent is a nucleic acid dye.

5. The method according to one of claims 1-4, wherein said separating comprises:

55 contacting the enriched erythrocyte sample with paramagnetic particles that bind specifically to the labeled GPI anchor-expressing erythrocytes;  
 subjecting the enriched erythrocyte sample containing the paramagnetic particles to a magnetic field under conditions effective to retain the labeled GPI anchor-expressing erythrocytes, but not GPI anchor-deficient erythrocytes in the magnetic field; and

collecting the GPI anchor-deficient erythrocytes not retained in the magnetic field.

6. The method according to claim 5 further comprising concentrating the collected GPI anchor-deficient erythrocytes.

5 7. The method according to claim 5, wherein the paramagnetic particles also bind to the second fluorescent reagent-labeled platelets, and said subjecting retains labeled platelets in the magnetic field.

8. The method according to one of claims 1-7 further comprising:

10 obtaining a quantitative measure of the total number of erythrocytes and/or reticulocytes in the enriched erythrocyte sample; and  
calculating the frequency of GPI anchor-deficient erythrocytes and/or reticulocytes relative to total erythrocytes and/or reticulocytes present in the sample.

15 9. The method according to claim 8, wherein said obtaining comprises:

contacting the enriched erythrocyte sample with a fluorescent reagent that differentially labels normochromatic erythrocytes from reticulocytes and leukocytes;  
exciting the fluorescent reagent with light of appropriate excitation wavelength;  
20 detecting the fluorescent emission and light scatter produced by the fluorescent reagent, and counting the number of erythrocytes and/or reticulocytes per unit volume of sample.

10. The method according to claim 8, further comprising:

25 adding counting beads to the enriched erythrocyte sample prior to said separating;  
counting, during said detecting step, the number of counting beads; and  
calculating a GPI anchor-deficient erythrocyte to bead ratio and/or GPI anchor-deficient reticulocyte to bead ratio.

11. The method according to claim 10, wherein said obtaining comprises:

30 contacting the enriched erythrocyte sample containing counting beads with a fluorescent reagent that differentially labels normochromatic erythrocytes from reticulocytes and leukocytes;  
exciting the fluorescent reagent and counting beads with light of appropriate excitation wavelength;  
35 detecting the fluorescent emission and light scatter produced by the fluorescent reagent and counting beads, and counting the number of erythrocytes and/or reticulocytes and counting beads in the sample; and  
calculating an erythrocyte to bead ratio and/or reticulocyte to bead ratio.

12. The method according to claim 8 further comprising:

40 providing a second enriched erythrocyte sample, wherein said second sample is obtained from the mammal prior to exposure to the exogenous agent;  
performing said contacting with the first and second fluorescent reagents, said separating, said contacting with the third fluorescent reagent, said exciting, and said detecting on the second enriched erythrocyte sample; and  
45 comparing the results obtained from the first and second samples, wherein a statistically significant deviation in the frequency of GPI anchor-deficient erythrocytes or reticulocytes between the first and second samples indicates the genotoxic potential of the exogenous agent.

13. The method according to claim 8, wherein

50 the exogenous agent is a chemical or physical agent and a statistically significant deviation in the frequency of GPI anchor-deficient erythrocytes or reticulocytes from a baseline GPI-anchor-deficient erythrocytes or reticulocyte frequency in unexposed or vehicle control mammals indicates the genotoxic potential of the exogenous chemical or physical agent; or  
the exogenous agent is an agent may modify endogenously-induced genetic damage and a statistically significant deviation in the frequency of GPI anchor-deficient erythrocytes or reticulocytes from a baseline GPI anchor-deficient  
55 erythrocytes or reticulocyte frequency indicates that the exogenous agent can modify endogenous DNA damage; or  
the exogenous agent comprises a first exogenous agent that may modify exogenously-induced genetic damage and a second exogenous agent that causes genetic damage, and a statistically significant deviation in the frequency of GPI anchor-deficient erythrocytes or reticulocytes for the genotoxicant-exposed mammal, as compared to a

mammal receiving only the second exogenous agent, indicates that the first exogenous agent can modify exogenously-induced DNA damage.

14. A kit comprising:

- 5 a first fluorescent reagent that binds GPI anchor-expressing cells, but not GPI anchor-deficient cells;  
 a second fluorescent reagent that specifically binds platelets;  
 a third fluorescent reagent that differentially labels normochromatic erythrocytes, reticulocytes, and leukocytes,  
 10 the third fluorescent reagent having a fluorescent emission spectrum that does not substantially overlap with the fluorescent emission spectra of the first or second fluorescent reagent;  
 a fourth reagent suitable for separating erythrocytes from platelets and leukocytes, the fourth reagent comprising a density gradient medium;  
 a fifth reagent comprising paramagnetic particles, the fifth reagent specifically binding to GPI anchor-expressing cells or at least a portion of the first fluorescent reagent bound thereon; and  
 15 instructions for using the reagents and cell separation products, and for detecting and calculating the frequency of GPI anchor-deficient erythrocytes and/or reticulocytes relative to total erythrocytes and/or reticulocytes in the sample.

15. The kit according to claim 14, wherein

- 20 the first fluorescent reagent comprises an anti-CD59, anti-CD24, or anti-CD55 antibody, or a combination thereof; the second fluorescent reagent comprises an anti-CD61 antibody or anti-CD42b antibody, or a combination thereof; and/or  
 the third fluorescent reagent is a nucleic acid dye.

16. The kit according to one of claims 14 or 15 further comprising:

- 25 a column suitable for use with paramagnetic particles and magnetic fields;  
 a computer readable storage medium that contains a cytometry data acquisition template for flow cytometric mutant erythrocyte and mutant reticulocyte scoring;  
 30 one or more of balanced salt solutions and an anticoagulant solution; and/or  
 counting beads.

Patentansprüche

1. Verfahren zur Zählung von in-vivo-Pig-A-Gen-Locus-Mutation, wobei das Verfahren umfasst:

- 35 Bereitstellen einer angereicherten Erythrozyten-Probe, die von einem Säuger erhalten wurde, welcher einem exogenen Agens ausgesetzt war, wobei die Probe normochromatische Erythrozyten und Retikulozyten umfasst und eine Reduktion der Häufigkeit von Thrombozyten und Leukozyten hat;  
 40 In-Kontakt-Bringen der angereicherten Erythrozyten-Probe mit einem ersten Fluoreszenzreagens, das GPI-Anker-exprimierende Erythrozyten, nicht aber GPI-Anker-defiziente Erythrozyten markiert, und einem zweiten Fluoreszenzreagens, das spezifisch Thrombozyten markiert;  
 Trennen der angereicherten Erythrozyten-Probe in einen ersten Teil, der Thrombozyten und GPI-Anker-exprimierende Erythrozyten umfasst, und einen zweiten Teil, der GPI-Anker-defiziente Erythrozyten umfasst;  
 45 In-Kontakt-Bringen des zweiten Teils mit einem dritten Fluoreszenzreagens, das normochromatische Erythrozyten unterschiedlich zu Retikulozyten und Leukozyten markiert;  
 Anregen des ersten, zweiten und dritten Fluoreszenzreagens im zweiten Teil mit Licht geeigneter Anregungswellenlänge, wobei das dritte Fluoreszenzreagens ein Fluoreszenz-Emissionsspektrum hat, das im Wesentlichen nicht mit den Fluoreszenz-Emissionsspektren des ersten oder zweiten Fluoreszenzreagens überlappt, und  
 50 Detektieren der Fluoreszenzemission und der Lichtstreuung, die durch die GPI-Anker-defizienten Erythrozyten des zweiten Teils, die mit dem dritten Fluoreszenzreagens markiert sind, produziert wurde, während kontaminierende GPI-Anker-exprimierende Erythrozyten und Retikulozyten, Thrombozyten und Leukozyten ausgeschlossen werden, und Zählen der Anzahl von GPI-Anker-defizienten Erythrozyten und Retikulozyten.

2. Verfahren nach Anspruch 1, wobei das erste Fluoreszenzreagens einen Anti-CD59, Anti-CD24- oder Anti-CD55-Antikörper oder eine Kombination davon umfasst.

## EP 2 643 694 B1

3. Verfahren nach Anspruch 1 oder 2, wobei das zweite Fluoreszenzreagens einen Anti-CD61-Antikörper oder Anti-CD42b-Antikörper oder eine Kombination davon umfasst.

4. Verfahren nach einem der Ansprüche 1 bis 3, wobei das dritte Fluoreszenzreagens ein Nucleinsäurefarbstoff ist.

5. Verfahren nach einem der Ansprüche 1 bis 4, wobei das Trennen umfasst:

In-Kontakt-Bringen der angereicherten Erythrozyten-Probe mit paramagnetischen Partikeln, welche spezifisch an die markierten GPI-Anker-exprimierenden Erythrozyten binden;

Unterwerfen der angereicherten Erythrozyten-Probe, die die paramagnetischen Partikel enthält, einem Magnetfeld unter Bedingungen, die wirksam sind, um die markierten GPI-Anker-exprimierenden Erythrozyten, nicht aber GPI-Anker-defiziente Erythrozyten in dem Magnetfeld zurückzuhalten, und Sammeln der GPI-Anker-defizienten Erythrozyten, die nicht im Magnetfeld zurückgehalten werden.

6. Verfahren nach Anspruch 5, das außerdem Konzentrieren der gesammelten GPI-Anker-defizienten Erythrozyten umfasst.

7. Verfahren nach Anspruch 5, wobei die paramagnetischen Partikel auch an die mit dem zweiten Fluoreszenzreagens markierten Thrombozyten binden und das Unterwerfen die markierten Thrombozyten in dem Magnetfeld zurückhält.

8. Verfahren nach einem der Ansprüche 1 bis 7, außerdem umfassend:

Erhalten einer quantitativen Messgröße der Gesamtzahl von Erythrozyten und/oder Retikulozyten in der angereicherten Erythrozyten-Probe und

Berechnen der Häufigkeit von GPI-Anker-defizienten Erythrozyten und/oder Retikulozyten relativ zu den gesamten Erythrozyten und/oder Retikulozyten, die in der Probe vorliegen.

9. Verfahren nach Anspruch 8, wobei das Erhalten umfasst:

In-Kontakt-Bringen der angereicherten Erythrozyten-Probe mit einem Fluoreszenzreagens, das normochromatische Erythrozyten unterschiedlich zu Retikulozyten und Leukozyten markiert;  
Anregen des Fluoreszenzreagens mit Licht geeigneter Anregungswellenlänge;  
Detektieren der Fluoreszenzemission und der Lichtstreuung, die durch das Fluoreszenzreagens produziert werden, und Zählen der Anzahl von Erythrozyten und/oder Retikulozyten pro Probeneinheitvolumen.

10. Verfahren nach Anspruch 8, außerdem umfassend:

Zugeben von Zähl-Beads zu der angereicherten Erythrozyten-Probe vor dem Trennen;

Zählen, während des Schrittes des Detektierens, der Anzahl von Zähl-Beads und

Berechnen eines GPI-Anker-defiziente Erythrozyten-zu-Beads-Verhältnisses und/oder eines GPI-Anker-defiziente Retikulozyten-zu-Beads-Verhältnisses.

11. Verfahren nach Anspruch 10, wobei das Erhalten umfasst:

In-Kontakt-Bringen der angereicherten Erythrozyten-Probe, die Zähl-Beads enthält, mit einem Fluoreszenzreagens, das normochromatische Erythrozyten unterschiedlich zu Retikulozyten und Leukozyten markiert;

Anregen des Fluoreszenzreagens und der Zähl-Beads mit Licht geeigneter Anregungswellenlänge;

Detektieren der Fluoreszenzemission und der Lichtstreuung, die durch das Fluoreszenzreagens und die Zähl-Beads produziert wurden, und Zählen der Anzahl von Erythrozyten und/oder Retikulozyten und Zähl-Beads in der Probe und Berechnen eines Erythrozyten-zu-Beads-Verhältnisses und/oder Retikulozyten-zu-Beads-Verhältnisses.

12. Verfahren nach Anspruch 8, außerdem umfassend:

Bereitstellen einer zweiten angereicherten Erythrozyten-Probe, wobei die zweite Probe von dem Säuger vor Exposition gegenüber dem exogenen Agens erhalten wurde;

Durchführen des In-Kontakt-Bringens mit dem ersten und dem zweiten Fluoreszenzreagens, des Trennens, des In-Kontakt-Bringens mit dem dritten Fluoreszenzreagens, des Anregens und des Detektierens mit der

## EP 2 643 694 B1

zweiten angereicherten Erythrozyten-Probe und  
Vergleichen der Resultate, die mit der ersten und der zweiten Probe erhalten worden waren, wobei eine statistisch signifikante Abweichung in der Häufigkeit von GPI-Anker-defizienten Erythrozyten oder Retikulozyten zwischen der ersten und der zweiten Probe das genotoxische Potential des exogenen Agenses anzeigt.

5

### 13. Verfahren nach Anspruch 8, wobei

das exogene Agens ein chemisches oder physikalisches Agens ist und eine statistisch signifikante Abweichung der Häufigkeit von GPI-Anker-defizienten Erythrozyten oder Retikulozyten ab einer Basislinien-Häufigkeit von GPI-Anker-defizienten Erythrozyten oder Retikulozyten bei nicht ausgesetzten oder Vehikel-Kontroll-Säugern das genotoxische Potential des exogenen chemischen oder physikalischen Agenses anzeigt, oder

10

das exogene Agens ein Agens ist, das eine endogen induzierte genetische Schädigung modifizieren kann und eine statistisch signifikante Abweichung der Häufigkeit von GPI-Anker-defizienten Erythrozyten oder Retikulozyten ab einer Basislinien-Häufigkeit von GPI-Anker-defizienten Erythrozyten oder Retikulozyten anzeigt, dass das exogene Agens endogene DNA-Schädigung modifizieren kann, oder

15

das exogene Agens ein erstes exogenes Agens, das eine exogen induzierte genetische Schädigung modifizieren kann, und ein zweites exogenes Agens, das eine genetische Schädigung verursacht, umfasst, und eine statistisch signifikante Abweichung der Häufigkeit von GPI-Anker-defizienten Erythrozyten oder Retikulozyten für den dem Genotoxikans ausgesetzten Säuger im Vergleich zu einem Säuger, der nur das zweite exogene Agens erhält, anzeigt, dass das erste exogene Agens exogen eine induzierte DNA-Schädigung modifizieren kann.

20

### 14. Kit, umfassend:

ein erstes Fluoreszenzreagens, das GPI-Anker-exprimierende Zellen, nicht aber GPI-Anker-defiziente Zellen bindet;

25

ein zweites Fluoreszenzreagens, das spezifisch Thrombozyten bindet;

ein drittes Fluoreszenzreagens, das normochromatische Erythrozyten, Retikulozyten und Leukozyten unterschiedlich markiert, wobei das dritte Fluoreszenzreagens ein Fluoreszenz-Emissionsspektrum hat, das mit den Fluoreszenz-Emissionsspektren des ersten oder zweiten Fluoreszenzreagens nicht wesentlich überlappt;

30

ein viertes Reagens, das zum Trennen von Erythrozyten von Thrombozyten und Leukozyten geeignet ist, wobei das vierte Reagens ein Dichtegradientenmedium umfasst;

ein fünftes Reagens, das paramagnetische Partikel umfasst, wobei das fünfte Reagens spezifisch an GPI-Anker-exprimierende Zellen oder wenigstens einen Teil des ersten daran gebundenen Fluoreszenzreagens bindet, und

35

Anleitungen zum Verwenden der Reagenzien und Zelltrennungsprodukte und zum Detektieren und Berechnen der Häufigkeit von GPI-Anker-defizienten Erythrozyten und/oder Retikulozyten relativ zu den gesamten Erythrozyten und/oder Retikulozyten in der Probe.

### 15. Kit nach Anspruch 14, wobei

das erste Fluoreszenzreagens einen Anti-CD59-, Anti-CD24- oder Anti-CD55-Antikörper oder eine Kombination davon umfasst;

40

das zweite Fluoreszenzreagens einen Anti-CD61-Antikörper oder Anti-CD42b-Antikörper oder eine Kombination davon umfasst und/oder

das dritte Fluoreszenzreagens ein Nucleinsäurefarbstoff ist.

45

### 16. Kit nach einem der Ansprüche 14 oder 15, außerdem umfassend:

eine Säule, die zur Verwendung mit paramagnetischen Partikeln und Magnetfeldern geeignet ist;

ein Computer-lesbares Speichermedium, das eine Zytometriedaten-Erfassungsmatrize zum Scoring von mutanten Erythrozyten und mutanten Retikulozyten mittels Durchflusszytometrie enthält;

50

eine oder mehrere ausgewogene Salzlösungen und eine Antikoagulans-Lösung und/oder Zähl-Beads.

## Revendications

55

### 1. Procédé de dénombrement d'une mutation du locus du gène *Pig-A* *in vivo*, le procédé comprenant :

la mise à disposition d'un échantillon d'érythrocytes enrichi obtenu à partir d'un mammifère exposé à un agent exogène, ledit échantillon comprenant des érythrocytes normochromatiques et des réticulocytes, et présentant

## EP 2 643 694 B1

une réduction de la fréquence des plaquettes et des leucocytes ;  
la mise en contact de l'échantillon d'érythrocytes enrichi avec un premier réactif fluorescent qui marque les érythrocytes exprimant une ancre GPI, mais pas les érythrocytes déficients en ancre GPI, et un deuxième réactif fluorescent qui marque spécifiquement les plaquettes ;

5 la séparation de l'échantillon d'érythrocytes enrichi en une première partie comprenant les plaquettes et les érythrocytes exprimant une ancre GPI et une seconde partie comprenant les érythrocytes déficients en ancre GPI ;

la mise en contact de la seconde partie avec un troisième réactif fluorescent qui marque les érythrocytes normochromatiques de manière différentielle par rapport aux réticulocytes et aux leucocytes ;

10 l'excitation du premier, deuxième, et troisième réactif fluorescent dans la seconde partie avec une lumière ayant une longueur d'onde d'excitation appropriée, où le troisième réactif fluorescent possède un spectre d'émission de fluorescence qui ne chevauche pas essentiellement les spectres d'émission de fluorescence du premier ou du deuxième réactif fluorescent ; et

15 la détection de l'émission de fluorescence et de la diffusion de la lumière produites par les érythrocytes déficients en ancre GPI de la seconde partie marquée avec le troisième réactif fluorescent, tout en excluant les érythrocytes et les réticulocytes exprimant une ancre GPI contaminants, les plaquettes, et les leucocytes, et en comptant le nombre d'érythrocytes et de réticulocytes déficients en ancre GPI.

20 **2.** Procédé selon la revendication 1, dans lequel le premier réactif fluorescent comprend un anticorps anti-CD59, anti-CD24, ou anti-CD55, ou une combinaison de ceux-ci.

**3.** Procédé selon la revendication 1 ou 2, dans lequel le deuxième réactif fluorescent comprend un anticorps anti-CD61 ou un anticorps anti-CD42b, ou une combinaison de ceux-ci.

25 **4.** Procédé selon l'une quelconque des revendications 1-3, dans lequel le troisième réactif fluorescent est un colorant des acides nucléiques.

**5.** Procédé selon l'une quelconque des revendications 1-4, dans lequel ladite séparation comprend :

30 la mise en contact de l'échantillon d'érythrocytes enrichi avec des particules paramagnétiques qui se lient spécifiquement aux érythrocytes exprimant une ancre GPI marqués ;

la soumission de l'échantillon d'érythrocytes enrichi contenant les particules paramagnétiques à un champ magnétique dans des conditions efficaces pour retenir les érythrocytes exprimant une ancre GPI marqués, mais pas les érythrocytes déficients en ancre GPI, dans le champ magnétique ; et

35 la collecte des érythrocytes déficients en ancre GPI non retenus dans le champ magnétique.

**6.** Procédé selon la revendication 5, comprenant en outre la concentration des érythrocytes déficients en ancre GPI collectés.

40 **7.** Procédé selon la revendication 5, dans lequel les particules paramagnétiques se lient également aux plaquettes marquées par le deuxième réactif fluorescent, et ladite soumission retient les plaquettes marquées dans le champ magnétique.

**8.** Procédé selon l'une quelconque des revendications 1-7, comprenant en outre :

45 l'obtention d'une mesure quantitative du nombre total d'érythrocytes et/ou de réticulocytes dans l'échantillon d'érythrocytes enrichi ; et

le calcul de la fréquence des érythrocytes et/ou réticulocytes déficients en ancre GPI par rapport aux érythrocytes et/ou réticulocytes totaux présents dans l'échantillon.

50 **9.** Procédé selon la revendication 8, dans lequel ladite obtention comprend :

la mise en contact de l'échantillon d'érythrocytes enrichi avec un réactif fluorescent qui marque les érythrocytes normochromatiques de manière différentielle par rapport aux réticulocytes et aux leucocytes ;

55 l'excitation du réactif fluorescent avec une lumière ayant une longueur d'onde d'excitation appropriée ;  
la détection de l'émission de fluorescence et de la diffusion de la lumière produites par le réactif fluorescent, et le comptage du nombre d'érythrocytes et/ou de réticulocytes par unité de volume d'échantillon.

10. Procédé selon la revendication 8, comprenant en outre :

l'ajout de billes de comptage à l'échantillon d'érythrocytes enrichi avant ladite séparation ;  
le comptage, pendant ladite étape de détection, du nombre de billes de comptage ; et  
le calcul d'un rapport érythrocyte déficient en ancre GPI/bille et/ou d'un rapport réticulocyte déficient en ancre GPI/bille.

11. Procédé selon la revendication 10, dans lequel ladite obtention comprend :

la mise en contact de l'échantillon d'érythrocytes enrichi contenant les billes de comptage avec un réactif fluorescent qui marque les érythrocytes normochromatiques de manière différentielle par rapport aux réticulocytes et aux leucocytes ;  
l'excitation du réactif fluorescent et des billes de comptage avec une lumière ayant une longueur d'onde d'excitation appropriée ;  
la détection de l'émission de fluorescence et de la diffusion de la lumière produites par le réactif fluorescent et les billes de comptage, et le comptage du nombre d'érythrocytes et/ou de réticulocytes et de billes de comptage dans l'échantillon ; et  
le calcul d'un rapport érythrocyte/bille et/ou d'un rapport réticulocyte/bille.

12. Procédé selon la revendication 8, comprenant en outre :

la mise à disposition d'un second échantillon d'érythrocytes enrichi, où ledit second échantillon est obtenu à partir du mammifère avant l'exposition à l'agent exogène ;  
la réalisation de ladite mise en contact avec le premier et le deuxième réactif fluorescent, ladite séparation, ladite mise en contact avec le troisième réactif fluorescent, ladite excitation, et ladite détection sur le second échantillon d'érythrocytes enrichi ; et  
la comparaison des résultats obtenus à partir du premier et du second échantillon, où une déviation statistiquement significative de la fréquence des érythrocytes ou des réticulocytes déficients en ancre GPI entre le premier et le second échantillon indique le potentiel génotoxique de l'agent exogène.

13. Procédé selon la revendication 8, dans lequel

l'agent exogène est un agent chimique ou physique et une déviation statistiquement significative de la fréquence des érythrocytes ou des réticulocytes déficients en ancre GPI par rapport à une fréquence d'érythrocytes ou de réticulocytes déficients en ancre GPI de référence chez des mammifères de contrôle véhicules ou non exposés indique le potentiel génotoxique de l'agent chimique ou physique exogène ; ou

l'agent exogène est un agent qui peut modifier une lésion génétique induite de manière endogène et une déviation statistiquement significative de la fréquence des érythrocytes ou des réticulocytes déficients en ancre GPI par rapport à une fréquence d'érythrocytes ou de réticulocytes déficients en ancre GPI de référence indique que l'agent exogène peut modifier un endommagement de l'ADN endogène ; ou

l'agent exogène comprend un premier agent exogène qui peut modifier une lésion génétique induite de manière exogène et un second agent exogène qui provoque des lésions génétiques, et une déviation statistiquement significative de la fréquence des érythrocytes ou des réticulocytes déficients en ancre GPI pour le mammifère exposé à l'agent génotoxique, par comparaison à un mammifère recevant uniquement le second agent exogène, indique que le premier agent exogène peut modifier un endommagement de l'ADN induit de manière exogène.

14. Kit comprenant :

un premier réactif fluorescent qui se lie à des cellules exprimant une ancre GPI, mais pas à des cellules déficientes en ancre GPI ;

un deuxième réactif fluorescent qui se lie spécifiquement aux plaquettes ;

un troisième réactif fluorescent qui marque de manière différentielle les érythrocytes normochromatiques, les réticulocytes, et les leucocytes, le troisième réactif fluorescent possédant un spectre d'émission de fluorescence qui ne chevauche pas essentiellement les spectres d'émission de fluorescence du premier ou du deuxième réactif fluorescent ;

un quatrième réactif approprié pour séparer les érythrocytes des plaquettes et des leucocytes, le quatrième réactif comprenant un milieu à gradient de densité ;

un cinquième réactif comprenant des particules paramagnétiques, le cinquième réactif se liant spécifiquement à des cellules exprimant une ancre GPI ou à au moins une partie du premier réactif fluorescent lié sur celles-ci ; et

## EP 2 643 694 B1

des instructions pour utiliser les réactifs et les produits de séparation cellulaire, et pour détecter et calculer la fréquence des érythrocytes et/ou réticulocytes déficients en ancre GPI par rapport aux érythrocytes et/ou réticulocytes totaux dans l'échantillon.

- 5    **15.** Kit selon la revendication 14, dans lequel  
le premier réactif fluorescent comprend un anticorps anti-CD59, anti-CD24, ou anti-CD55, ou une combinaison de ceux-ci ;  
le deuxième réactif fluorescent comprend un anticorps anti-CD61 ou un anticorps anti-CD42b, ou une combinaison de ceux-ci ; et/ou  
10    le troisième réactif fluorescent est un colorant des acides nucléiques.

- 16.** Kit selon l'une quelconque des revendications 14 ou 15, comprenant en outre :

15    une colonne appropriée pour être utilisée avec des particules paramagnétiques et des champs magnétiques ;  
un support de stockage pouvant être lu par ordinateur qui contient un modèle d'acquisition de données de cytométrie pour l'évaluation d'érythrocytes mutants et de réticulocytes mutants par cytométrie en flux ;  
l'une ou plusieurs de solutions salines équilibrées et d'une solution anticoagulante ; et/ou  
des billes de comptage.

20

25

30


35

40


45

50

55




**FIG. 1**



**FIG. 2**

Blood from a Mutagenized Rat,  
Pre-Magnetic Field



Blood from a Mutagenized Rat,  
Post-Magnetic Field






FIG. 3



**FIG. 4**



FIGS. 5A-C

## REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

## Patent documents cited in the description

- US 7824874 B, Dertinger [0005]
- US 20090311706 A, Dertinger [0005]
- US 7037498 B [0044]
- US 7034121 B [0044]
- US 7041870 B [0044]
- US 7074405 B [0044]

## Non-patent literature cited in the description

- **ARATEN et al.** Clonal Populations of Hematopoietic Cells with Paroxysmal Nocturnal Hemoglobinuria Genotype and Phenotype are Present in Normal Individuals. *Proc Natl Acad Sci USA*, 1999, vol. 96, 5209-5214 [0002]
- **CHEN et al.** Glycophosphatidylinositol-anchored Protein Deficiency as a Marker of Mutator Phenotypes in Cancer. *Cancer Res.*, 2001, vol. 61, 654-658 [0002]
- **IIDA et al.** Characterization of Genomic PIG-A Gene: A Gene for GPI Anchor Biosynthesis and Paroxysmal Nocturnal Hemoglobinuria. *Blood*, 1994, vol. 83, 3126-3131 [0002]
- **KAWAGOE et al.** Molecular cloning of Murine pig-a, a Gene for GPI-anchor Biosynthesis, and Demonstration of Interspecies Conservation of its Structure, Function, and Genetic Locus. *Genomics*, 1994, vol. 23, 566-574 [0002]
- **NORRIS et al.** The Defect in Glycosylphosphatidylinositol Anchor Synthesis in Paroxysmal Nocturnal Hemoglobinuria. *Blood*, 1994, vol. 83, 816-821 [0003]
- **NISHIMURA et al.** Paroxysmal Nocturnal Hemoglobinuria: An Acquired Genetic Disease. *Am J Hematol*, 1999, vol. 62, 175-182 [0003]
- **NAFA et al.** The Spectrum of Somatic Mutations in the PIG-A Gene in Paroxysmal Nocturnal Hemoglobinuria Includes Large Deletions and Small Duplications. *Blood Cells Mol Dis*, 1998, vol. 24, 370-384 [0003]
- **MIURA et al.** Development of an In Vivo Gene Mutation Assay Using the Endogenous Pig-A Gene: I. Flow Cytometric Detection of CD59-Negative Peripheral Red Blood Cells and CD48-Negative Spleen T-Cells From the Rat. *Environ. Molec. Mutagen.*, 2008, vol. 49, 614-621 [0004]
- **ARTEN et al.** *PNAS*, 1999, vol. 96, 5209-5214 [0004]
- **PHONETHEPSWATH et al.** Erythrocyte-based Pig-a Gene Mutation Assay: Demonstration of Cross-Species Potential. *Mutat. Res.*, 2008, vol. 657, 122-126 [0005]
- **J. GODING.** Monoclonal Antibodies: Principles and Practice. Academic Press, 1983, 98-118 [0044]
- **DETINGER et al.** Integration of Mutation and Chromosomal Damage Endpoints into 28-Day Repeat Dose Toxicology Studies. *Toxicol. Sci.*, 2010, vol. 115, 401-411 [0047]
- **WARE et al.** Immunophenotypic Analysis of Reticulocytes in Paroxysmal Nocturnal Hemoglobinuria. *Blood*, 1995, vol. 86, 1586-1589 [0049]
- **PHONETHEPSWATH et al.** Pig-a Mutation Kinetics in Rat Erythrocytes Following Exposure to Five Prototypical Mutagens. *Toxicol. Sci.*, 2010, vol. 114, 59-70 [0051]
- **PHONETHEPSWATH et al.** Pig-a Mutation: Kinetics in Rat Erythrocytes Following Exposure to Five Prototypical Mutagens. *Toxicol. Sci.*, 2010, vol. 114, 59-70 [0074]
- **DETINGER et al.** Integration of Mutation and Chromosomal Damage Endpoints into 28-Day Repeat Dose Toxicology Studies. *Toxicol. Sci.*, 2010, vol. 115, 401-411 [0074]

专利名称(译)	基于Pig-a基因的体内快速基因突变检测		
公开(公告)号	<a href="#">EP2643694A1</a>	公开(公告)日	2013-10-02
申请号	EP2011842900	申请日	2011-11-23
[标]申请(专利权)人(译)	LITRON LAB		
申请(专利权)人(译)	LITRON LABORATORIES LTD.		
当前申请(专利权)人(译)	LITRON LABORATORIES LTD.		
[标]发明人	DERTINGER STEPHEN D BRYCE STEVEN M		
发明人	DERTINGER, STEPHEN, D. BRYCE, STEVEN, M.		
IPC分类号	G01N33/53 G01N33/00 C12Q1/68		
CPC分类号	C12Q1/6827 G01N33/5002 G01N33/5014 G01N33/54326 C12Q2563/131		
优先权	61/417104 2010-11-24 US		
其他公开文献	EP2643694B1 EP2643694A4		
外部链接	<a href="#">Espacenet</a>		

#### 摘要(译)

本发明涉及用于定量分析个体中猪-A基因的体内突变频率的方法和试剂盒，特别是使用脊椎动物的外周血样品。