



(11) **EP 1 754 783 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
**21.02.2007 Bulletin 2007/08**

(51) Int Cl.:  
**C12N 5/06 (2006.01)**

(21) Application number: **05025978.7**

(22) Date of filing: **28.04.2004**

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
HU IE IT LI LU MC NL PL PT RO SE SI SK TR**

(30) Priority: **02.05.2003 US 467051 P**  
**25.07.2003 US 628783**

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
**04760595.1 / 1 624 758**

(71) Applicant: **THE SCRIPPS RESEARCH INSTITUTE**  
**La Jolla, CA 92037 (US)**

(72) Inventors:  
• **Friedlander, Martin**  
**Del Mar, CA 92014 (US)**

- **Otani, Atsushi**  
**San Diego, CA 92122 (US)**
- **Dasilva, Karen**  
**Irvine, CA 92612 (US)**
- **Hanekamp, Stacey**  
**Spring Valley, CA 91977 (US)**

(74) Representative: **Schnappauf, Georg et al**  
**Dr. Volker Vossius**  
**Patent- und Rechtsanwaltskanzlei**  
**Geibelstrasse 6**  
**81679 München (DE)**

Remarks:

This application was filed on 29 - 11 - 2005 as a  
divisional application to the application mentioned  
under INID code 62.

(54) **Transfected hematopoietic stem cells and methods of treatment of neovascular eye diseases therewith**

(57) Isolated, mammalian, adult bone marrow-derived, lineage negative hematopoietic stem cell populations (Lin<sup>-</sup> HSCs) containing endothelial progenitor cells (EPCs) have been transfected with therapeutically useful genes. The transfected isolated stem cell populations of the invention are useful for delivering genes to the eye

for cell-based gene therapy. Methods of preparing transfected isolated stem cell populations of the invention, and methods of treating ocular diseases and injury are also described.

**EP 1 754 783 A1**

**Description****Cross-Reference to Related Applications**

5 **[0001]** This application is a continuation-in-part of U.S. Application for Patent Serial No. 10/628,783, filed on July 25, 2003, which claims the benefit of Provisional Application for Patent Serial No. 60/398,522, filed on July 25, 2002, and Provisional Application for Patent Serial No. 60/467,051, filed on May 2, 2003, the entire disclosures of which are incorporated herein by reference

**Statement of Government Interest**

10 **[0002]** A portion of the work described herein was supported by grant number CA92577 from the National Cancer Institute and by grants number EY11254, EY12598 and EY125998 from the National Institutes of Health. The United States Government has certain rights in this invention.

15

**Field of the Invention**

20 **[0003]** This invention relates to isolated, mammalian, lineage negative hematopoietic stem cell (Lin<sup>-</sup> HSC) populations derived from bone marrow and their uses. More particularly, the invention relates to isolated, mammalian, lineage negative hematopoietic stem cell (Lin<sup>-</sup> HSC) populations containing endothelial progenitor cells (EPC). The invention also relates to treatment of vascular diseases of the eye by administering Lin<sup>-</sup> HSC and transfected Lin<sup>-</sup> HSC populations to the eye.

**Background of the Invention**

25

30 **[0004]** Inherited degenerations of the retina affect as many as 1 in 3500 individuals and are characterized by progressive night blindness, visual field loss, optic nerve atrophy, arteriolar attenuation, altered vascular permeability and central loss of vision often progressing to complete blindness (Heckenlively, J. R., editor, 1988; *Retinitis Pigmentosa*, Philadelphia: JB Lippincott Co.). Molecular genetic analysis of these diseases has identified mutations in over 110 different genes accounting for only a relatively small percentage of the known affected individuals (Humphries *et al.*, 1992, *Science* 256:804-808; Farrar *et al.* 2002, *EMBO J.* 21:857-864.). Many of these mutations are associated with enzymatic and structural components of the phototransduction machinery including rhodopsin, cGMP phosphodiesterase, *rd*s peripherin, and RPE65. Despite these observations, there are still no effective treatments to slow or reverse the progression of these retinal degenerative diseases. Recent advances in gene therapy have led to successful reversal of the *rd*s (Ali *et al.* 2000, *Nat. Genet.* 25:306-310) and *rd* (Takahashi *et al.* 1999, *J. Virol.* 73:7812-7816) phenotypes in mice and the RPE65 phenotype in dogs (Acland *et al.* 2001, *Nat. Genet.* 28:92-95) when the wild type transgene is delivered to photoreceptors or the retinal pigmented epithelium (RPE) in animals with a specific mutation.

35

40 **[0005]** Age related macular degeneration (ARMD) and diabetic retinopathy (DR) are the leading causes of visual loss in industrialized nations and do so as a result of abnormal retinal neovascularization. Since the retina consists of well-defined layers of neuronal, glial, and vascular elements, relatively small disturbances such as those seen in vascular proliferation or edema can lead to significant loss of visual function. Inherited retinal degenerations, such as retinitis pigmentosa (RP), are also associated with vascular abnormalities, such as arteriolar narrowing and vascular atrophy. While significant progress has been made in identifying factors that promote and inhibit angiogenesis, no treatment is currently available to specifically treat ocular vascular disease.

45

50 **[0006]** For many years it has been known that a population of stem cells exists in the normal adult circulation and bone marrow. Different sub-populations of these cells can differentiate along hematopoietic lineage positive (Lin<sup>+</sup>) or lineage negative (Lin<sup>-</sup>) lineages. Furthermore, the lineage negative hematopoietic stem cell (HSC) population has recently been shown to contain endothelial progenitor cells (EPC) capable of forming blood vessels *in vitro* and *in vivo* (See Asahara *et al.* 1997, *Science* 275: 964-7). These cells can participate in normal and pathological postnatal angiogenesis (See Lyden *et al.* 2001 *Nat. Med.* 7, 1194-201; Kalka *et al.* 2000, *Proc. Natl. Acad. Sci. U. S. A.* 97:3422-7; and Kocher *et al.* 2001, *Nat. Med.* 7: 430-6) as well as differentiate into a variety of non-endothelial cell types including hepatocytes (See Lagasse *et al.* 2000, *Nat. Med.* 6:1229-34), microglia (See Priller *et al.* 2002 *Nat. Med.* 7:1356-61), cardiomyocytes (See Orlic *et al.* 2001, *Proc. Natl. Acad. Sci. U. S. A.* 98:10344-9) and epithelium (See Lyden *et al.* 2001, *Nat. Med.* 7: 1194-1201). Although these cells have been used in several experimental models of angiogenesis, the mechanism of EPC targeting to neovasculature is not known and no strategy has been identified that will effectively increase the number of cells that contribute to a particular vasculature.

55

**[0007]** Hematopoietic stem cells from bone marrow are currently the only type of stem cell commonly used for therapeutic applications. Bone marrow HSC's have been used in transplants for over 40 years. Currently, advanced methods

of harvesting purified stem cells are being investigated to develop therapies for treatment of leukemia, lymphoma, and inherited blood disorders. Clinical applications of stem cells in humans have been investigated for the treatment of diabetes and advanced kidney cancer in limited numbers of human patients.

## 5 Summary of the Invention

[0008] The present invention provides isolated, mammalian, population of hematopoietic stem cells (HSCs) that do not express lineage surface antigens (Lin) on their cell surface, i.e, lineage negative hematopoietic stem cells (Lin<sup>-</sup> HSCs). The Lin<sup>-</sup> HSC populations of the present invention include endothelial progenitor cells (EPC), also known as endothelial precursor cells, that selectively target activated retinal astrocytes when intravitreally injected into the eye. The Lin<sup>-</sup> HSCs of the present invention preferably are derived from adult mammalian bone marrow, more preferably from adult human bone marrow.

[0009] In a preferred embodiment the Lin<sup>-</sup> HSC populations of the present invention are isolated by extracting bone marrow from an adult mammal; separating a plurality of monocytes from the bone marrow; labeling the monocytes with biotin-conjugated lineage panel antibodies to one or more lineage surface antigens, removing monocytes that are positive for the lineage surface antigens and then recovering a Lin<sup>-</sup> HSC population containing EPCs. Preferably the monocytes are labeled with biotin-conjugated lineage panel antibodies to one or more lineage surface antigen selected from the group consisting of CD2, CD3, CD4, CD11, CD11a, Mac-1, CD14, CD16, CD19, CD24, CD33, CD36, CD38, CD45, Ly-6G, TER-119, CD45RA, CD56, CD64, CD68, CD86, CD66b, HLA-DR, and CD235a (Glycophorin A). Preferably, at least about 20% of the cells of the isolated Lin<sup>-</sup> HSC population of the present invention express the surface antigen CD31.

[0010] The EPC's within the population of Lin<sup>-</sup> HSCs of the present invention extensively incorporate into developing retinal vessels and remain stably incorporated into neovasculature of the eye. The isolated, Lin<sup>-</sup> HSC populations of the present invention can be used to rescue and stabilize degenerating retinal vasculature in mammals, to rescue neuronal networks, and to facilitate repair of ischemic tissue.

[0011] In one preferred embodiment, the cells of the isolated Lin<sup>-</sup> HSC populations are transfected with a therapeutically useful gene. For example, the cells can be transfected with polynucleotides that operably encode for neurotrophic agents or anti-angiogenic agents that selectively target neovasculature and inhibit new vessel formation without affecting already established vessels through a form of cell-based gene therapy. In one embodiment, the isolated, Lin<sup>-</sup> HSC populations of the present invention include a gene encoding an angiogenesis inhibiting peptide. The angiogenesis inhibiting Lin<sup>-</sup> HSCs are useful for modulating abnormal blood vessel growth in diseases such as ARMD, DR and certain retinal degenerations associated with abnormal vasculature. In another preferred embodiment, the isolated, Lin<sup>-</sup> HSCs of the present invention include a gene encoding a neurotrophic peptide. The neurotrophic Lin<sup>-</sup> HSCs are useful for promoting neuronal rescue in ocular diseases involving retinal neural degeneration, such as glaucoma, retinitis pigmentosa, and the like.

[0012] A particular advantage of ocular treatments with the isolated Lin<sup>-</sup> HSC populations of the present invention is a vasculotropic and neurotrophic rescue effect observed in eyes intravitreally treated with the Lin<sup>-</sup> HSCs. Retinal neurons and photoreceptors are preserved and visual function is maintained in eyes treated with the isolated Lin<sup>-</sup> HSCs of the invention. The present invention provides a method for treating retinal degeneration comprising administering isolated Lin<sup>-</sup> HSC cells derived from bone marrow, which contain endothelial progenitor cells that selectively target activated retinal astrocytes, wherein at least about 50 % the isolated Lin<sup>-</sup> HSCs express the surface antigen CD31 and at least about 50 % the isolated Lin<sup>-</sup> HSCs express the surface antigen CD117 (c-kit).

[0013] The present invention also provides a method of isolating lineage negative hematopoietic stem cell populations containing endothelial progenitor cells from adult mammalian bone marrow, preferably from adult human bone marrow. In addition, a line of genetically identical cells (i.e., clones) can be generated from human Lin<sup>-</sup> HSCs that are useful in regenerative or reparative treatment of retinal vasculature, as well as for treatment or amelioration of retinal neuronal tissue degeneration.

## Brief Description of the Drawings

### 50 [0014]

Figure 1 (a and b) depicts schematic diagrams of developing mouse retina. (a) Development of primary plexus. (b) The second phase of retinal vessel formation. GCL, ganglion cell layer; IPL, inner plexus layer; INL, inner nuclear layer; OPL, outer plexus layer; ONL, outer nuclear layer; RPE, retinal pigment epithelium; ON, optic nerve; P, periphery.

Figure 1c depicts flow cytometric characterization of bone marrow-derived Lin<sup>+</sup> HSC and Lin<sup>-</sup>HSC separated cells. Top row: Dot plot distribution of non-antibody labeled cells, in which R1 defines the quantifiable-gated area of positive PE-staining; R2 indicates GFP-positive; Middle row: Lin<sup>-</sup>HSC (C57B/6) and Bottom row: Lin<sup>+</sup> HSC (C57B/6) cells,

each cell line labeled with the PE-conjugated antibodies for Sca-1, c-kit, Flk-1/KDR, CD31. Tie-2 data was obtained from Tie-2-GFP mice. Percentages indicate percent of positive-labeled cells out of total Lin<sup>-</sup>HSC or Lin<sup>+</sup> HSC population.

Figure 2 depicts engraftment of Lin<sup>-</sup> HSCs into developing mouse retina. (a) At four days post-injection (P6) intravitreally injected eGFP<sup>+</sup> Lin<sup>-</sup>HSC cells attach and differentiate on the retina (b) Lin<sup>-</sup>HSC (B6.129S7-Gtrosa26 mice, stained with  $\beta$ -gal antibody) establish themselves ahead of the vasculature stained with collagen IV antibody (asterisk indicates tip of vasculature). (c) Most of Lin<sup>+</sup>HSC cells (eGFP<sup>+</sup>) at four days post-injection (P6) were unable to differentiate. (d) Mesenteric eGFP<sup>+</sup> murine EC four days post-injection (P6). (e) Lin<sup>-</sup> HSCs (eGFP<sup>+</sup>) injected into adult mouse eyes. (f) Low magnification of eGFP<sup>+</sup> Lin<sup>-</sup> HSCs (arrows) homing to and differentiating along the pre-existing astrocytic template in the GFAP-GFP transgenic mouse. (g) Higher magnification of association between Lin<sup>-</sup> cells (eGFP) and underlying astrocyte (arrows). (h) Non-injected GFAP-GFP transgenic control. (i) Four days post-injection (P6), eGFP<sup>+</sup> Lin<sup>-</sup> HSCs migrate to and undergo differentiation in the area of the future deep plexus. Left figure captures Lin<sup>-</sup> HSC activity in a whole mounted retina; right figure indicates location of the Lin<sup>-</sup> cells (arrows) in the retina (top is vitreal side, bottom is scleral side). (j) Double labeling with  $\alpha$ -CD31-PE and  $\alpha$ -GFP-alexa 488 antibodies. Seven days after injection, the injected Lin<sup>-</sup>HSCs (eGFP), red were incorporated into the vasculature (CD31). Arrowheads indicate the incorporated areas. (k) eGFP<sup>+</sup> Lin<sup>-</sup> HSC cells form vessels fourteen days post-injection (P17). (l and m) Intra-cardiac injection of rhodamine-dextran indicates that the vessels are intact and functional in both the primary (l) and deep plexus (m).

Figure 3 (a and b) shows that eGFP<sup>+</sup> Lin<sup>-</sup> HSC cells home to the gliosis (indicated by GFAP expressing-astrocytes, far left image) induced by both laser (a) and mechanical (b) induced injury in the adult retina (asterisk indicates injured site). Far right images are a higher magnification, demonstrating the close association of the Lin<sup>-</sup>HSCs and astrocytes. Calibration bar= 20 $\mu$ M.

Figure 4 shows that Lin<sup>-</sup>HSC cells rescue the vasculature of the retinal degeneration mouse. (a-d) Retinas at 27 days post-injection (P33) with collagen IV staining; (a) and (b), retinas injected with Lin<sup>+</sup> HSC cells (Balb/c) showed no difference in vasculature from normal FVB mice; (c) and (d) retinas injected with Lin<sup>-</sup> HSCs (Balb/c), exhibited a rich vascular network analogous to a wild-type mouse; (a) and (c), frozen sections of whole retina (top is vitreal side, bottom is scleral side) with DAPI staining; (b) and (d), deep plexus of retinal whole amount; (e) bar graph illustrating the increase in vascularity of the deep vascular plexus formed in the Lin<sup>-</sup>HSC cell-injected retinas (n=6). The extent of deep retinal vascularization was quantified by calculating the total length of vessels within each image. Average total length of vessels/high power field (in microns) for Lin<sup>-</sup> HSC, Lin<sup>+</sup>HSC or control retinas were compared. (f) Comparison of the length of deep vascular plexus after injection with Lin<sup>-</sup>HSC (R, right eye) or Lin<sup>+</sup>HSC (L, left eye) cells from *rd/rd* mouse. The results of six independent mice are shown (each color represents each mouse). (g) and (h) Lin<sup>-</sup> HSC cells also (Balb/c) rescued the *rd/rd* vasculature when injected into P15 eyes. The intermediate and deep vascular plexus of Lin<sup>-</sup>HSC (G) or Lin<sup>+</sup>HSC (H) cell injected retinas (one month after injection) are shown.

Figure 5 depicts photomicrographs of mouse retinal tissue: (a) deep layer of retinal whole mount (*rd/rd* mouse), five days post-injection (P11) with eGFP<sup>+</sup> Lin<sup>-</sup>HSCs visible (gray). (b) and (c) P60 retinal vasculature of Tie-2-GFP (*rd/rd*) mice that received Balb/c Lin<sup>-</sup> cells (b) or Lin<sup>+</sup>HSC cell (c) injection at P6. Only endogenous endothelial cells (GFP-stained) are visible in the left panels of (b) and (c). The middle panels of (b) and (c) are stained with CD31 antibody; arrows indicate the vessels stained with CD31 but not with GFP, the right panels of (b) and (c) show staining with both GFP and CD31. (d)  $\alpha$ -SMA staining of Lin<sup>-</sup> HSC injected (left panel) and control retina (right panel).

Figure 6 shows that T2-TrpRS-transfected Lin<sup>-</sup> HSCs inhibit the development of mouse retinal vasculature. (a) Schematic representation of human TrpRS, T2-TrpRS and T2-TrpRS with an Igk signal sequence at the amino terminus. (b) T2-TrpRS transfected Lin<sup>-</sup> HSC-injected retinas express T2-TrpRS protein *in vivo*. (1) Recombinant T2-TrpRS produced in *E. coli*; (2) Recombinant T2-TrpRS produced in *E. coli*; (3) Recombinant T2-TrpRS produced in *E. coli*; (4) control retina; (5) Lin<sup>-</sup>HSC + pSecTag2A (vector only) injected retina; (6) Lin<sup>-</sup>HSC + pKLe135 (Igk-T2-TrpRS in pSecTag) injected retina. (a) Endogenous TrpRS. (b) Recombinant T2-TrpRS. (c) T2-TrpRS of Lin<sup>-</sup>HSC injected retina. (c-f) Representative primary (superficial) and secondary (deep) plexuses of injected retinas, seven days post-injection; (c) and (d) Eyes injected with empty plasmid-transfected Lin<sup>-</sup>HSC developed normally; (e) and (f) the majority of T2-TrpRS-transfected Lin<sup>-</sup>HSC injected eyes exhibited inhibition of deep plexus; (c) and (e) primary (superficial) plexus; (d) and (f) secondary (deep) plexus. Faint outline of vessels observed in (f) are "bleed-through" images of primary network vessels shown in (e).

Figure 7 shows the DNA sequence encoding His<sub>6</sub>-tagged T2-TrpRS, SEQ ID NO: 1.

Figure 8 shows the amino acid sequence of His<sub>6</sub>-tagged T2-TrpRS, SEQ ID NO: 2.

Figure 9 illustrates photomicrographs and electroretinograms (ERG) of retinas from mice whose eyes were injected with the Lin<sup>-</sup> HSC of the present invention and with Lin<sup>+</sup> HSC (controls).

Figure 10 depicts statistical plots showing a correlation between neuronal rescue (y-axis) and vascular rescue (x-axis) for both the intermediate (Int.) and deep vascular layers of *rd/rd* mouse eyes treated with Lin<sup>-</sup>HSC.

Figure 11 depicts statistical plots showing no correlation between neuronal rescue (y-axis) and vascular rescue (x-

axis) for *rd/rd* mouse eyes that were treated with Lin<sup>+</sup>HSC.

Figure 12 is a bar graph of vascular length (y-axis) in arbitrary relative units for *rd/rd* mouse eyes treated with the Lin<sup>-</sup>HSC (dark bars) and untreated (light bars) *rd/rd* mouse eyes at time points of 1 month (1M), 2 months (2M), and 6 months (6M) post-injection.

Figure 13 includes three bar graphs of the number of nuclei in the outer neural layer (ONR) of *rd/rd* mice at 1 month (1M), 2 months (2M) and 6 months (6M), post-injection, and demonstrates a significant increase in the number of nuclei for eyes treated with Lin<sup>-</sup> HSC (dark bars) relative to control eyes treated with Lin<sup>+</sup> HSC (light bars).

Figure 14 depicts plots of the number of nuclei in the outer neural layer for individual *rd/rd* mice, comparing the right eye (R, treated with Lin<sup>-</sup> HSC) relative to the left eye (L, control eye treated with Lin<sup>+</sup> HSC) at time points (post injection) of 1 month (1M), 2 months (2M), and 6 months (6M); each line in a given plot compares the eyes of an individual mouse.

Figure 15 depicts retinal vasculature and neural cell changes in *rd1/rd1* (C3H/HeJ, left panels) or wild type mice (C57BL/6, right panels). Retinal vasculature of intermediate (upper panels) or deep (middle panels) vascular plexuses in whole-mounted retinas (red: collagen IV, green: CD31) and sections (red: DAPI, green: CD31, lower panels) of the same retinas are shown (P: postnatal day). (GCL: ganglion cell layer, INL: inter nuclear layer, ONL: outer nuclear layer).

Figure 16 shows that Lin<sup>-</sup> HSC injection rescues the degeneration of neural cells in *rd1/rd1* mice. A, B and C, retinal vasculature of intermediate (int.) or deep plexus and sections of Lin<sup>-</sup>HSC injected eye (right panels) and contralateral control cell (CD31<sup>-</sup>) injected eye (left panels) at P30 (A), P60 (B), and P180 (C). D, the average total length of vasculature (+ or - standard error of the mean) in Lin<sup>-</sup> HSC injected or control cell (CD31<sup>-</sup>) injected retinas at P30 (left, n=10), P60 (middle, n=10), and P180 (right, n=6). Data of intermediate (Int.) and deep vascular plexus are shown separately (Y axis: relative length of vasculature). E, the average numbers of cell nuclei in the ONL at P30 (left, n=10), P60 (middle, n=10), or P180 (right, n=6) of control cell (CD31<sup>-</sup>) or Lin<sup>-</sup>HSC injected retinas (Y axis: relative number of cell nuclei in the ONL). F, Linear correlations between the length of vasculature (X axis) and the number of cell nuclei in the ONL (Y axis) at P30 (left), P60 (middle), and P180 (right) of Lin<sup>-</sup>HSC or control cell injected retinas.

Figure 17 demonstrates that retinal function is rescued by Lin<sup>-</sup> HSC injection. Electroretinographic (ERG) recordings were used to measure the function of Lin<sup>-</sup> HSC or control cell (CD31<sup>-</sup>) injected retinas. A and B, Representative cases of rescued and non-rescued retinas 2 months after injection. Retinal section of Lin<sup>-</sup> HSC injected right eye (A) and CD31<sup>-</sup> control cell injected left eye (B) of the same animal are shown (green: CD31 stained vasculature, red: DAPI stained nuclei). C, ERG results from the same animal shown in A & B.

Figure 18 shows that a population of human bone marrow cells can rescue degenerating retinas in the *rd1* mouse (A-C). The rescue is also observed in another model of retinal degeneration, *rd10* (D-K). A, human Lin<sup>-</sup> HSCs (hLin<sup>-</sup>HSCs) labeled with green dye can differentiate into retinal vascular cells after intravitreal injection into C3SnSmn. *CB17-Prkdc SCID* mice. B and C, Retinal vasculature (left panels; upper: intermediate plexus, lower: deep plexus) and neural cells (right panel) in hLin<sup>-</sup> HSC injected eye (B) or contralateral control eye (C) 1.5 months after injection. D-K, Rescue of *rd10* mice by Lin<sup>-</sup>HSCs (injected at P6). Representative retinas at P21 (D: Lin<sup>-</sup>HSCs, H: control cells), P30 (E: Lin<sup>-</sup>HSCs, I: control cells), P60 (F: Lin<sup>-</sup>HSCs, J: control cells), and P105 (G: Lin<sup>-</sup>HSCs, K: control cells) are shown (treated and control eyes are from the same animal at each time point). Retinal vasculature (upper image in each panel is the intermediate plexus; the middle image in each panel is the deep plexus) was stained with CD31 (green) and Collagen IV (red). The lower image in each panel shows a cross section made from the same retina (red: DAPI, green: CD31).

Figure 19 demonstrates that crystallin  $\alpha A$  is up regulated in rescued outer nuclear layer cells after treatment with Lin<sup>-</sup> HSCs but not in contralateral eyes treated with control cells. Left panel; IgG control in rescued retina, Middle panel; crystallin  $\alpha A$  in rescued retina, Right panel; crystallin  $\alpha A$  in non-rescued retina.

Figure 20 includes tables of genes that are upregulated in murine retinas that have been treated with the Lin<sup>-</sup> HSCs of the present invention. (A) Genes whose expression is increased 3-fold in mouse retinas treated with murine Lin<sup>-</sup> HSCs. (B) Crystallin genes that are upregulated in mouse retinas treated with murine Lin<sup>-</sup> HSC. (C) Genes whose expression is increased 2-fold in mouse retinas treated with human Lin<sup>-</sup> HSCs. (D) Genes for neurotrophic factors or growth factors whose expression is upregulated in mouse retinas treated with human Lin<sup>-</sup> HSCs.

Figure 21 illustrates the distribution of CD31 and integrin  $\alpha 6$  surface antigens on CD133 positive (CD133<sup>+</sup>) and CD133 negative (CD133<sup>-</sup>) human Lin<sup>-</sup> HSC populations of the present invention.

Figure 22 illustrates postnatal retinal development in wild-type C57/B16 mice raised in normal oxygen levels (normoxia), at post natal days P0 through P30.

Figure 23 illustrates oxygen-induced retinopathy model in C57/B16 mice raised in high oxygen levels (hyperoxia; 75% oxygen) between P7 and P12, followed by normoxia from P12-P17.

Figure 24 demonstrates vascular rescue by treatment with the Lin<sup>-</sup> HSC populations of the present invention in the oxygen -induced retinopathy model.

**Detailed Description of Preferred Embodiments**

**[0015]** Stem cells are typically identified by the distribution of antigens on the surface of the cells (for a detailed discussion see *Stem Cells: Scientific Progress and Future Directions*, a report prepared by the National Institutes of Health, Office of Science Policy, June 2001, Appendix E: Stem Cell Markers, which is incorporated herein by reference to the extent pertinent).

**[0016]** Hematopoietic stem cells are that stem cells that are capable of developing into various blood cell types e.g., B cells, T cells, granulocytes, platelets, and erythrocytes. The lineage surface antigens are a group of cell-surface proteins that are markers of mature blood cell lineages, including CD2, CD3, CD11, CD11a, Mac-1 (CD11b:CD18), CD14, CD16, CD19, CD24, CD33, CD36, CD38, CD45, CD45RA, murine Ly-6G, murine TER-119, CD56, CD64, CD68, CD86 (B7.2), CD66b, human leucocyte antigen DR (HLA-DR), and CD235a (Glycophorin A). Hematopoietic stem cells that do not express significant levels of these antigens are commonly referred to a lineage negative (Lin<sup>-</sup>). Human hematopoietic stem cells commonly express other surface antigens such as CD31, CD34, CD117 (c-kit) and/or CD133. Murine hematopoietic stem cells commonly express other surface antigens such as CD34, CD117 (c-kit), Thy-1, and/or Sca-1.

**[0017]** The present invention provides isolated hematopoietic stem cells that do not express significant levels of a "lineage surface antigen" (Lin) on their cell surfaces. Such cells are referred to herein as "lineage negative" or "Lin<sup>-</sup>" hematopoietic stem cells. In particular this invention provides a population of Lin<sup>-</sup> hematopoietic stem cells (Lin<sup>-</sup> HSCs) that include endothelial progenitor cells (EPCs), which are capable of incorporating into developing vasculature and then differentiating to become vascular endothelial cells. Preferably the isolated Lin<sup>-</sup> HSC populations are present in a culture medium such as phosphate buffered saline (PBS).

**[0018]** As used herein and in the appended claims, the phrase "adult" in reference to bone marrow, includes bone marrow isolated postnatally, i.e., from juvenile and adult individuals, as opposed to embryos. The term "adult mammal" refers to both juvenile and fully mature mammals.

**[0019]** The present invention provides isolated, mammalian, lineage negative hematopoietic stem cell (Lin<sup>-</sup> HSC) populations containing endothelial progenitor cells (EPCs). The isolated Lin<sup>-</sup> HSC populations of the present invention preferably comprise mammalian cells in which at least about 20% of the cells express the surface antigen CD31, which is commonly present on endothelial cells. In other embodiment, at least about 50 % of the cells express CD31, more preferably at least about 65%, most preferably at least about 75%. Preferably at least about 50% of the cells of the Lin<sup>-</sup> HSC populations of the present invention preferably express the integrin  $\alpha$ 6 antigen.

**[0020]** In one preferred murine Lin<sup>-</sup> HSC population embodiment, at least about 50 % of the cells express CD31 antigen and at least about 50 % of the cells express the CD117 (c-kit) antigen. Preferably, at least about 75% of the Lin<sup>-</sup> HSC cells express the surface antigen CD31, more preferably about 81 % of the cells. In another preferred murine embodiment, at least about 65 % of the cells express the surface antigen CD117, more preferably about 70 % of the cells. A particularly preferred embodiment of the present invention is a population of murine Lin<sup>-</sup> HSCs in which about 50 % to about 85% of the cells express the surface antigen CD31 and about 70 % to about 75% of the cells express the surface antigen CD 117.

**[0021]** Another preferred embodiment is a human Lin<sup>-</sup> HSC population in which the cells are CD 133 negative, in which at least about 50 % of the cells express the CD31 surface antigen and at least about 50 % of the cells express the integrin  $\alpha$ 6 antigen. Yet another preferred embodiment is a human Lin<sup>-</sup> HSC population in which the cells are CD133 positive, in which at less than about 30% of the cells express the CD31 surface antigen and less than about 30% of the cells express the integrin  $\alpha$ 6 antigen.

**[0022]** The isolated Lin<sup>-</sup> HSC populations of the present invention selectively target astrocytes and incorporate into the retinal neovasculature when intravitreally injected into the eye of the mammalian species, such as a mouse or a human, from which the cells were isolated.

**[0023]** The isolated Lin<sup>-</sup> HSC populations of the present invention include endothelial progenitor cells that differentiate to endothelial cells and generate vascular structures within the retina. In particular, the Lin<sup>-</sup> HSC populations of the present invention are useful for the treatment of retinal neovascular and retinal vascular degenerative diseases, and for repair of retinal vascular injury. The Lin<sup>-</sup> HSC cells of the present invention promote neuronal rescue in the retina and promote upregulation of anti-apoptotic genes. It has surprisingly been found that adult human Lin<sup>-</sup> HSC cells of the present invention can inhibit retinal degeneration even in severe combined immunodeficient (SCID) mice suffering from retinal degeneration. Additionally, the Lin<sup>-</sup> HSC populations can be utilized to treat retinal defects in the eyes of neonatal mammals, such as mammals suffering from oxygen induced retinopathy or retinopathy of prematurity.

**[0024]** The present invention also provides a method of treating ocular diseases in a mammal comprising isolating from the bone marrow of the mammal a lineage negative hematopoietic stem cell population that includes endothelial progenitor cells, and intravitreally injecting the isolated stem cells into an eye of the mammal in a number sufficient to arrest the disease. The present method can be utilized to treat ocular diseases such as retinal degenerative diseases, retinal vascular degenerative diseases, ischemic retinopathies, vascular hemorrhages, vascular leakage, and choroidopathies in neonatal, juvenile or fully mature mammals. Examples of such diseases include age related macular

degeneration (ARMD), diabetic retinopathy (DR), presumed ocular histoplasmosis (POHS), retinopathy of prematurity (ROP), sickle cell anemia, and retinitis pigmentosa, as well as retinal injuries.

**[0025]** The number of stem cells injected into the eye is sufficient for arresting the disease state of the eye. For example, the number of cells can be effective for repairing retinal damage of the eye, stabilizing retinal neovasculature, maturing retinal neovasculature, and preventing or repairing vascular leakage and vascular hemorrhage.

**[0026]** Cells of the Lin<sup>-</sup> HSC populations of the present invention can be transfected with therapeutically useful genes, such as genes encoding antiangiogenic proteins for use in ocular, cell-based gene therapy and genes encoding neurotrophic agents to enhance neuronal rescue effects.

**[0027]** The transfected cells can include any gene which is therapeutically useful for treatment of retinal disorders. In one preferred embodiment, the transfected Lin<sup>-</sup> HSCs of the present invention include a gene operably encoding an antiangiogenic peptide, including proteins, or protein fragments such as TrpRS or antiangiogenic fragments thereof, e.g., the T1 and T2 fragments of TrpRS, which are described in detail in co-owned, co-pending U.S. patent application Serial No. 10/080,839, the disclosure of which is incorporated herein by reference. The transfected Lin<sup>-</sup> HSCs encoding an antiangiogenic peptide of the present invention are useful for treatment of retinal diseases involving abnormal vascular development, such as diabetic retinopathy, and like diseases. Preferably the Lin<sup>-</sup> HSCs are human cells.

**[0028]** In another preferred embodiment, the transfected Lin<sup>-</sup> HSCs of the present invention include a gene operably encoding a neurotrophic agent such as nerve growth factor, neurotrophin-3, neurotrophin-4, neurotrophin-5, ciliary neurotrophic factor, retinal pigmented epithelium-derived neurotrophic factor, insulin-like growth factor, glial cell line-derived neurotrophic factor, brain-derived neurotrophic factor, and the like. Such neurotrophic Lin<sup>-</sup> HSCs are useful for promoting neuronal rescue in retinal neuronal degenerative diseases such as glaucoma and retinitis pigmentosa, in treatment of injuries to the retinal nerves, and the like. Implants of ciliary neurotrophic factor have been reported as useful for the treatment of retinitis pigmentosa (see Kirby *et al.* 2001, *Mol Ther.* 3(2):241-8; Farrar *et al.* 2002, *EMBO Journal* 21:857-864). Brain-derived neurotrophic factor reportedly modulates growth associated genes in injured retinal ganglia (see Fournier, *et al.*, 1997, *J. Neurosci. Res.* 47:561-572). Glial cell line derived neurotrophic factor reportedly delays photoreceptor degeneration in retinitis pigmentosa (see McGee *et al.* 2001, *Mol Ther.* 4(6):622-9).

**[0029]** The present invention also provides a method of isolating a lineage negative hematopoietic stem cells comprising endothelial progenitor cells from bone marrow of a mammal. The method entails the steps of (a) extracting bone marrow from an adult mammal; (b) separating a plurality of monocytes from the bone marrow; (c) labeling the monocytes with biotin-conjugated lineage panel antibodies to one or more lineage surface antigens, preferably lineage surface antigens selected from the group consisting of CD2, CD3, CD4, CD11, CD11a, Mac-1, CD14, CD16, CD19, CD24, CD33, CD36, CD38, CD45, Ly-6G (murine), TER-119 (murine), CD45RA, CD56, CD64, CD68, CD86 (B7.2), CD66b, human leucocyte antigen DR (HLA-DR), and CD235a (Glycophorin A); (d) removing monocytes that are positive for said one or more lineage surface antigens from the plurality of monocytes and recovering a population of lineage negative hematopoietic stem cells containing endothelial progenitor cells, preferably in which at least about 20% of the cells express CD31.

**[0030]** When the Lin<sup>-</sup> HSC are isolated from adult human bone marrow, preferably the monocytes are labeled with biotin-conjugated lineage panel antibodies to lineage surface antigens CD2, CD3, CD4, CD11a, Mac-1, CD14, CD16, CD19, CD33, CD38, CD45RA, CD64, CD68, CD86 (B7.2), and CD235a. When the Lin<sup>-</sup> HSC are isolated from adult murine bone marrow, preferably the monocytes are labeled with biotin-conjugated lineage panel antibodies to lineage surface antigens CD3, CD11, CD45, Ly-6G, and TER-119.

**[0031]** In a preferred method, the cells are isolated from adult human bone marrow and are further separated by CD133 lineage. One preferred method of isolating human Lin<sup>-</sup> HSCs includes the additional steps of labeling the monocytes with a biotin-conjugated CD133 antibody and recovering a CD133 positive, Lin<sup>-</sup> HSC population. Typically, less than about 30 % of such cells express CD31 and less than about 30% of such cell express integrin  $\alpha 6$ . The human Cd133 positive, Lin<sup>-</sup> HSC populations of the present invention can target sites of peripheral ischemia-driven neovascularization when injected into eyes that are not undergoing angiogenesis.

**[0032]** Another preferred method of isolating human Lin<sup>-</sup> HSCs includes the additional steps of labeling the monocytes with a biotin-conjugated CD133 antibody, removing CD133 positive cells, and recovering a CD133 negative, Lin<sup>-</sup> HSC population. Typically, at least about 50 % of such cells express CD31 and at least about 50% of such cell express integrin  $\alpha 6$ . The human CD 133 negative, Lin<sup>-</sup> HSC populations of the present invention can incorporate into developing vasculature when injected into eyes that are undergoing angiogenesis.

**[0033]** The present invention also provides methods for treating ocular angiogenic diseases by administering transfected Lin<sup>-</sup> HSC cells of the present invention by intravitreal injection of the cells into the eye. Such transfected Lin<sup>-</sup> HSC cells comprise Lin<sup>-</sup> HSC transfected with a therapeutically useful gene, such as a gene encoding antiangiogenic or neurotrophic gene product. Preferably the transfected Lin<sup>-</sup> HSC cells are human cells.

**[0034]** Preferably, at least about  $1 \times 10^5$  Lin<sup>-</sup> HSC cells or transfected Lin<sup>-</sup> HSC cells are administered by intravitreal injection to a mammalian eye suffering from a retinal degenerative disease. The number of cells to be injected may depend upon the severity of the retinal degeneration, the age of the mammal and other factors that will be readily apparent to one of ordinary skill in the art of treating retinal diseases. The Lin<sup>-</sup> HSC may be administered in a single dose or by

multiple dose administration over a period of time, as determined by the clinician in charge of the treatment.

**[0035]** The Lin<sup>-</sup> HSCs of the present invention are useful for the treatment of retinal injuries and retinal defects involving an interruption in or degradation of the retinal vasculature or retinal neuronal degeneration. Human Lin<sup>-</sup> HSCs also can be used to generate a line of genetically identical cells, i.e., clones, for use in regenerative or reparative treatment of retinal vasculature, as well as for treatment or amelioration of retinal neuronal degeneration.

## METHODS

### Example 1. Cell Isolation and Enrichment; Preparation of Murine Lin<sup>-</sup> HSC Populations A and B.

**[0036] General Procedure.** All *in vivo* evaluations were performed in accordance with the NIH Guide for the Care and Use of Laboratory Animals, and all evaluation procedures were approved by The Scripps Research Institute (TSRI, La Jolla, CA) Animal Care and Use Committee. Bone marrow cells were extracted from B6.129S7-Gtosa26, Tie-2GFP, ACTbEGFP, FVB/NJ (*rd/rd* mice) or Balb/cBYJ adult mice (The Jackson Laboratory, ME).

**[0037]** Monocytes were then separated by density gradient separation using HISTOPAQUE® polysucrose gradient (Sigma, St. Louis, MO) and labeled with biotin conjugated lineage panel antibodies (CD45, CD3, Ly-6G, CD11, TER-119, Pharmingen, San Diego, CA) for Lin<sup>-</sup> selection in mice. Lineage positive (Lin<sup>+</sup>) cells were separated and removed from Lin<sup>-</sup> HSC using a magnetic separation device (AUTOMACS™ sorter, Miltenyi Biotech, Auburn, CA). The resulting Lin<sup>-</sup> HSC population, containing endothelial progenitor cells was further characterized using a FACS™ Calibur flow cytometer (Becton Dickinson, Franklin Lakes, NJ) using following antibodies: PE-conjugated-Sca-1, c-kit, KDR, and CD31 (Pharmingen, San Diego, CA). Tie-2-GFP bone marrow cells were used for characterization of Tie-2.

**[0038]** To harvest adult mouse endothelial cells, mesenteric tissue was surgically removed from ACTbEGFP mouse and placed in collagenase (Worthington, Lakewood, NJ) to digest the tissue, followed by filtration using a 45µm filter. Flow-through was collected and incubated with Endothelial Growth Media (Clonetics, San Diego, CA). Endothelial characteristics were confirmed by observing morphological cobblestone appearance, staining with CD31 mAb (Pharmingen) and examining cultures for the formation of tube-like structures in MATRIGEL™ matrix (Beckton Dickinson, Franklin Lakes, NJ).

**[0039] Murine Lin<sup>-</sup> HSC Population A.** Bone marrow cells were extracted from ACTbEGFP mice by the General Procedure described above. The Lin<sup>-</sup> HSC cells were characterized by FACS flow cytometry for CD31, c-kit, Sca-1, Flk-1, and Tie-2 cell surface antigen markers. The results are shown in FIG. 1c. About 81 % of the Lin<sup>-</sup> HSC exhibited the CD31 marker, about 70.5 % of the Lin<sup>-</sup> HSC exhibited the c-kit marker, about 4% of the Lin<sup>-</sup> HSC exhibited the Sca-1 marker, about 2.2% of the Lin<sup>-</sup> HSC exhibited the Flk-1 marker and about 0.91 % of the Lin<sup>-</sup> HSC cell exhibited the Tie-2 marker. In contrast, the Lin<sup>+</sup> HSC that were isolated from these bone marrow cells had a significantly different cell marker profile (i.e., CD31: 37.4%; c-kit: 20%; Sca-1: 2.8%; Flk-: 0.05 %).

**[0040] Murine Lin<sup>-</sup> HSC Population B.** Bone marrow cells were extracted from Ba1b/C, ACTbEGFP, and C3H mice by the General Procedure described above. The Lin<sup>-</sup> HSC cells were analyzed for the presence of cell surface markers (Sca-1, KDR, c-kit, CD34, CD31 and various integrins:  $\alpha$ 1,  $\alpha$ 2,  $\alpha$ 3,  $\alpha$ 4,  $\alpha$ 5,  $\alpha$ 6,  $\alpha$ M,  $\alpha$ V,  $\alpha$ X,  $\alpha$ IIb,  $\beta$ 1,  $\beta$ 2,  $\beta$ 3,  $\beta$ 4,  $\beta$ 5 and  $\beta$ 7). The results are shown in Table 1.

**Table 1. Characterization of Lin<sup>-</sup> HSC Population B.**

Cell Marker	Lin <sup>-</sup> HSC
$\alpha$ 1	0.10
$\alpha$ 2	17.57
$\alpha$ 3	0.22
$\alpha$ 4	89.39
$\alpha$ 5	82.47
$\alpha$ 6	77.70
$\alpha$ L	62.69
$\alpha$ M	35.84
$\alpha$ X	3.98
$\alpha$ V	33.64
$\alpha$ IIb	0.25
$\beta$ 1	86.26
$\beta$ 2	49.07
$\beta$ 3	45.70

## EP 1 754 783 A1

(continued)

	Cell Marker	Lin <sup>-</sup> HSC
5	p4	0.68
	β5	9.44
	β7	11.25
	CD31	51.76
	CD34	55.83
10	Flk-1/KDR	2.95
	c-kit (CD117)	74.42
	Sca-1	7.54

### Example 2. Intravitreal Administration of Cells in a Murine Model.

15 [0041] An eyelid fissure was created in a mouse eyelid with a fine blade to expose the P2 to P6 eyeball. Lineage negative HSC Population A of the present invention (approximately 10<sup>5</sup> cells in about 0.5 μl to about 1 μl of cell culture medium) was then injected intravitreally using a 33-gauge (Hamilton, Reno, NV) needled-syringe.

### Example 3. EPC Transfection.

20 [0042] Murine Lin<sup>-</sup> HSC (Population A) were transfected with DNA encoding the T2 fragment of TrpRS also enclosing a His<sub>6</sub> tag (SEQ ID NO:1, FIG. 7) using FuGENE™6 Transfection Reagent (Roche, Indianapolis, IN) according to manufacturer's protocol. Lin<sup>-</sup> HSC cells (about 10<sup>6</sup> cell per ml) were suspended in opti-MEM® medium (Invitrogen, Carlsbad, CA) containing stem cell factor (PeproTech, Rocky Hill, NJ). DNA (about 1 μg) and FuGENE reagent (about 25 3 μl) mixture was then added, and the mixtures were incubated at about 37 °C for about 18 hours. After incubation, cells were washed and collected. The transfection rate of this system was approximately 17 % that was confirmed by FACS analysis. T2 production was confirmed by western blotting. The amino acid sequence of His<sub>6</sub>-tagged T2-TrpRS is shown as SEQ ID NO: 2, FIG. 8.

### Example 4. Immunohistochemistry and Confocal Analysis.

30 [0043] Mouse retinas were harvested at various time points and were prepared for either whole mounting or frozen sectioning. For whole mounts, retinas were fixed with 4 % paraformaldehyde, and blocked in 50 % fetal bovine serum (FBS) and 20% normal goat serum for one hour at ambient room temperature. Retinas were processed for primary 35 antibodies and detected with secondary antibodies. The primaries used were: anti-Collagen IV (Chemicon, Temecula, CA), anti-β-gal (Promega, Madison, WI), anti-GFAP (Dako Cytomation, Carpinteria, CA), anti-α-smooth muscle actin (α-SMA, Dako Cytomation). Secondary antibodies used were conjugated either to Alexa 488 or 594 fluorescent markers (Molecular Probes, Eugene, OR). Images were taken using an MRC 1024 Confocal microscope (Bio-Rad, Hercules, CA). Three-dimensional images were created using LASERSHARP® software (Bio-Rad) to examine the three different 40 layers of vascular development in the whole mount retina. The difference in GFP pixel intensity between enhanced GFP (eGFP) mice and GFAP/wtGFP mice, distinguished by confocal microscopy, was utilized to create the 3D images.

### Example 5. *In vivo* Retinal Angiogenesis Quantification Assay in Mice.

45 [0044] For T2-TrpRS analysis, the primary and deep plexus were reconstructed from the three dimensional images of mouse retinas. The primary plexus was divided into two categories: normal development, or halted vascular progression. The categories of inhibition of deep vascular development were construed based upon the percentage of vascular inhibition including the following criteria: complete inhibition of deep plexus formation was labeled "Complete", normal 50 vascular development (including less than 25 % inhibition) was labeled "Normal" and the remainder labeled "Partial." For the *rd/rd* mouse rescue data, four separate areas of the deeper plexus in each whole mounted retina were captured using a 10x lens. The total length of vasculature was calculated for each image, summarized and compared between the groups. To acquire accurate information, Lin<sup>-</sup> HSC were injected into one eye and Lin<sup>+</sup> HSC into another eye of the same mouse. Non-injected control retinas were taken from the same litter.

### Example 6. Adult Retinal Injury Murine Models.

55 [0045] Laser and scar models were created using either a diode laser (150 mW, 1 second, 50 mm) or mechanically

by puncturing the mouse retina with a 27 gauge needle. Five days after injury, cells were injected using the intravitreal method. Eyes were harvested from the mice five days later.

#### Example 7. Neurotrophic Rescue of Retinal Regeneration.

**[0046]** Adult murine bone marrow derived lineage negative hematopoietic stem cells (Lin<sup>-</sup> HSC) have a vasculotropic and neurotrophic rescue effect in a mouse model of retinal degeneration. Right eyes of 10-day old mice were injected intravitreally with about 0.5 microliters containing about 10<sup>5</sup> Lin<sup>-</sup>HSC of the present invention and evaluated 2 months later for the presence of retinal vasculature and neuronal layer nuclear count. The left eyes of the same mice were injected with about the same number of Lin<sup>+</sup>HSC as a control, and were similarly evaluated. As shown in FIG. 9, in the Lin<sup>-</sup>HSC treated eyes, the retinal vasculature appeared nearly normal, the inner nuclear layer was nearly normal and the outer nuclear layer (ONL) had about 3 to about 4 layers of nuclei. In contrast, the contralateral Lin<sup>+</sup>HSC treated eye had a markedly atrophic middle retinal vascular layer, a completely atrophic outer retinal vascular layer; the inner nuclear layer was markedly atrophic and the outer nuclear layer was completely gone. This was dramatically illustrated in Mouse 3 and Mouse 5. In Mouse 1, there was no rescue effect and this was true for approximately 15% of the injected mice.

**[0047]** When visual function was assessed with electroretinograms (ERG), the restoration of a positive ERG was observed when both the vascular and neuronal rescue was observed (Mice 3 and 5). Positive ERG was not observed when there was no vascular or neuronal rescue (Mouse 1). This correlation between vascular and neurotrophic rescue of the *rd/rd* mouse eyes by the Lin<sup>-</sup> HSC of the present invention is illustrated by a regression analysis plot shown in FIG. 10. A correlation between neuronal (y-axis) and vascular (x-axis) recovery was observed for the intermediate vasculature type ( $r=0.45$ ) and for the deep vasculature ( $r=0.67$ ).

**[0048]** FIG. 11 shows the absence of any statistically significant correlation between vascular and neuronal rescue by Lin<sup>+</sup> HSC. The vascular rescue was quantified and the data are presented in Figure 12. Data for mice at 1 month (1M), 2 months (2M), and 6 months (6M), post-injection shown in FIG. 12, demonstrate that vascular length was significantly increased in eyes treated with the Lin<sup>-</sup> HSC of the present invention (dark bars) relative to the vascular length in untreated eyes from the same mouse (light bars), particularly at 1 month and 2 months, post-injection. The neurotrophic rescue effect was quantified by counting nuclei in the inner and outer nuclear layers about two months after injection of Lin<sup>-</sup> HSC or Lin<sup>+</sup>HSC. The results are presented in Figures 13 and 14.

#### Example 8. Human Lin<sup>-</sup> HSC Population.

**[0049]** Bone marrow cells were extracted from healthy adult human volunteers by the General Procedure described above. Monocytes were then separated by density gradient separation using HISTOPAQUE® polysucrose gradient (Sigma, St. Louis, MO). To isolate the Lin<sup>-</sup> HSC population from human bone marrow mononuclear cells the following biotin conjugated lineage panel antibodies were used with the magnetic separation system (AUTOMACS™ sorter, Miltenyi Biotech, Auburn, CA): CD2, CD3, CD4, CD11a, Mac-1, CD14, CD16, CD19, CD33, CD38, CD45RA, CD64, CD68, CD86, CD235a (Pharmingen).

**[0050]** The human Lin<sup>-</sup> HSC population was further separated into two sub-populations based on CD133 expression. The cells were labeled with biotin-conjugated CD133 antibodies and separated into CD133 positive and CD133 negative sub-populations.

#### Example 9. Intravitreal Administration of Human and Murine Cells in Murine Models for Retinal Degeneration.

**[0051]** C3H/HeJ, C3SnSmn.CB17-*Prkdc* *SCID*, and *rd10* mouse strains were used as retinal degeneration models. C3H/HeJ and C3SnSmn.CB17-*Prkdc* *SCID* mice (The Jackson Laboratory, Maine) were homozygous for the retinal degeneration 1 (*rd1*) mutation, a mutation that causes early onset severe retinal degeneration. The mutation is located in exon 7 of the *Pde6b* gene encoding the rod photoreceptor cGMP phosphodiesterase  $\beta$  subunit. The mutation in this gene has been found in human patients with autosomal recessive retinitis pigmentosa (RP). C3SnSmn.CB17-*Prkdc* *SCID* mice are also homozygous for the severe combined immune deficiency spontaneous mutation (*Prkdc* *SCID*) and were used for human cell transfer experiments. Retinal degeneration in *rd10* mice is caused by a mutation in exon 13 of *Pde6b* gene. This is also a clinically relevant RP model with later onset and milder retinal degeneration than *rd1/rd1*. All evaluations were performed in accordance with the NIH Guide for the Care and Use of Laboratory Animals, and all procedures were approved by The Scripps Research Institute Animal Care and Use Committee.

**[0052]** An eyelid fissure was created in a mouse eyelid with a fine blade to expose the P2 to P6 eyeball. Lineage negative HSC cells for murine population A or human population C (approximately 10<sup>5</sup> cells in about 0.5  $\mu$ l to about 1  $\mu$ l of cell culture medium) were then injected in the mouse eye intravitreally using a 33-gauge (Hamilton, Reno, NV) needled-syringe. To visualize the injected human cells, cells were labeled with dye (Cell tracker green CMFDA, Molecular Probes) before injection.

**[0053]** Retinas were harvested at various time points and fixed with 4% paraformaldehyde (PFA) and methanol followed by blocking in 50% FBS/20% NGS for one hour at room temperature. To stain retinal vasculature, retinas were incubated with anti-CD31 (PharMingen) and anti-collagen IV (Chemicon) antibodies followed by Alexa 488 or 594 conjugated secondary antibodies (Molecular Probes, Eugene, Oregon). The retinas were laid flat with four radial relaxing incisions to obtain a whole mount preparation. Images of vasculature in intermediate or deep retinal vascular plexuses (see Dorrell *et al.* 2002 *Invest Ophthalmol. Vis. Sci.* 43:3500-3510) were obtained using a Radiance MP2100 confocal microscope and LASERSHARP® software (Biorad, Hercules, California). For quantification of vasculature, four independent fields (900  $\mu\text{m}$  x 900  $\mu\text{m}$ ) were chosen randomly from the mid portion of the intermediate or deep vascular layer and the total length of vasculature was measured using LASERPIX® analyzing software (Biorad). The total lengths of these four fields in the same plexus were used for further analysis.

**[0054]** The flat-mounted retinas were re-embedded for cryostat sections. Retinas were placed in 4 % PFA overnight followed by incubation with 20 % sucrose. The retinas were embedded in optimal cutting temperature compound (OCT: Tissue-Tek; Sakura FineTech, Torrance, CA). Cryostat sections (10  $\mu\text{m}$ ) were re-hydrated in PBS containing the nuclear dye DAPI (Sigma-Aldrich, St. Louis, Missouri). DAPI-labeled nuclear images of three different areas (280  $\mu\text{m}$  width, unbiased sampling) in a single section that contained optic nerve head and the entire peripheral retina were taken by confocal microscope. The numbers of the nuclei located in ONL of the three independent fields in one section were counted and summed up for analysis. Simple linear-regression analysis was performed to examine the relationship between the length of vasculature in the deep plexus and the number of cell nuclei in the ONL.

**[0055]** Following overnight dark-adaptation, mice were anesthetized by intraperitoneal injection of 15  $\mu\text{g}/\text{gm}$  ketamine and 7  $\mu\text{g}/\text{gm}$  xylazine. Electroretinograms (ERGs) were recorded from the corneal surface of each eye after pupil dilation (1 % atropine sulfate) using a gold loop corneal electrode together with a mouth reference and tail ground electrode. Stimuli were produced with a Grass Photoc Stimulator (PS33 Plus, Grass Instruments, Quincy, MA) affixed to the outside of a highly reflective Ganzfeld dome. Rod responses were recorded to short-wavelength (Wratten 47A;  $\lambda_{\text{max}} = 470 \text{ nm}$ ) flashes of light over a range of intensities up to the maximum allowable by the photic stimulator (0.668  $\text{cd}\cdot\text{s}/\text{m}^2$ ). Response signals were amplified (CP511 AC amplifier, Grass Instruments), digitized (PCI-1200, National Instruments, Austin, TX) and computer-analyzed. Each mouse served as its own internal control with ERGs recorded from both the treated and untreated eyes. Up to 100 sweeps were averaged for the weakest signals. The averaged responses from the untreated eye were digitally subtracted from the responses from the treated eye and this difference in signal was used to index functional rescue.

**[0056]** Microarray analysis was used for evaluation of Lin<sup>-</sup> HSC-targeted retinal gene expression. P6 *rd/rd* mice were injected with either Lin<sup>-</sup> or CD31<sup>-</sup> HSCs. The retinas of these mice were dissected 40 days post-injection in RNase free medium (rescue of the retinal vasculature and the photoreceptor layer is obvious at this time point after injection). One quadrant from each retina was analyzed by whole mount to ensure that normal HSC targeting as well as vasculature and neural protection had been achieved. RNA from retinas with successful injections was purified using a TRIzol (Life Technologies, Rockville, MD), phenol/chloroform RNA isolation protocol. RNA was hybridized to Affymetrix Mu74Av2 chips and gene expression was analyzed using GENESPRING® software (SiliconGenetics, Redwood City, CA). Purified human or mouse HSCs were injected intravitreally into P6 mice. At P45 the retinas were dissected and pooled into fractions of 1) human HSC-injected, rescued mouse retinas, 2) human HSC-injected, non-rescued mouse retinas, and 3) mouse HSC-injected, rescued mouse retinas for purification of RNA and hybridization to human-specific U133A Affymetrix chips. GENESPRING® software was used to identify genes that were expressed above background and with higher expression in the human HSC-rescued retinas. The probe-pair expression profiles for each of these genes were then individually analyzed and compared to a model of normal human U133A microarray experiments using dChip to determine human species specific hybridization and to eliminate false positives due to cross-species hybridization.

**[0057]** When the CD133 positive and CD133 negative Lin<sup>-</sup> HSC sub-population was intravitreally injected into the eyes of neonatal SCID mice, the greatest extent of incorporation into the developing vasculature was observed for the CD133 negative sub-population, which expresses both CD31 and integrin  $\alpha 6$  surface antigens (see FIG. 21, bottom). The CD133 positive sub-population, which does not express CD31 or integrin  $\alpha 6$  (FIG. 21, top) appears to target sites of peripheral ischemia-driven neovascularization, but not when injected into eyes undergoing angiogenesis.

#### **Example 10. Intravitreal Administration of Murine Cells in Murine Models for Oxygen Induced Retinal Degeneration.**

**[0058]** New born wild-type C57B16 mice were exposed to hyperoxia (75 % oxygen) between postnatal days P7 to P12 in an oxygen-induced retinal degeneration (OIR) model. FIG 22 illustrates normal postnatal vascular development in C57B16 mice from P0 to P30. At P0 only budding superficial vessels can be observed around the optic disc. Over the next few days, the primary superficial network extends toward the periphery, reaching the far periphery by day P10. Between P7 and P12, the secondary (deep) plexus develops. By P17, an extensive superficial and deep network of vessels is present (FIG. 22, insets). In the ensuing days, remodeling occurs along with development of the tertiary

(intermediate) layer of vessels until the adult structure is reached approximately at P21.

[0059] In contrast, in the OIR model, following exposure to 75 % oxygen at P7-P12, the normal sequence of events is severely disrupted (FIG. 23). Adult murine Lin<sup>-</sup> HSC populations of the invention were intravitreally injected at P3 in an eye of a mouse that was subsequently subjected to OIR, the other eye was injected with PBS or CD31 negative cells as a control. FIG. 24 illustrates that the Lin<sup>-</sup> HSC populations of the present invention can reverse the degenerative effects of high oxygen levels in the developing mouse retina. Fully developed superficial and deep retinal vasculature was observed at P17 in the treated eyes, whereas in the control eyes showed large avascular areas with virtually no deep vessels (FIG. 24). Approximately 100 eyes of mice in the OIR model were observed. Normal vascularization was observed in 58% of the eyes treated with the Lin<sup>-</sup> HSC populations of the invention, compared to 12 % of the control eyes treated with CD31- cells and 3 % of the control eyes treated with PBS.

## RESULTS

### Murine Retinal Vascular Development; A Model for Ocular Angiogenesis

[0060] The mouse eye provides a recognized model for the study of mammalian retinal vascular development, such as human retinal vascular development. During development of the murine retinal vasculature, ischemia-driven retinal blood vessels develop in close association with astrocytes. These glial elements migrate onto the third trimester human fetus, or the neonatal rodent, retina from the optic disc along the ganglion cell layer and spread radially. As the murine retinal vasculature develops, endothelial cells utilize this already established astrocytic template to determine the retinal vascular pattern (See FIG. 1a and b). FIG. 1 (a and b) depicts schematic diagrams of developing mouse retina. FIG. 1a depicts development of the primary plexus (dark lines at upper left of the diagram) superimposed over the astrocyte template (light lines) whereas, FIG. 1b depicts the second phase of retinal vessel formation. In the Figures, GCL stands for ganglion cell layer; IPL stands for inner plexus layer; INL stands for inner nuclear layer; OPL stands for outer plexus layer; ONL stands for outer nuclear layer; RPE stands for retinal pigment epithelium; ON stands for optic nerve; and P stands for periphery.

[0061] At birth, retinal vasculature is virtually absent. By postnatal day 14 (P14) the retina has developed complex primary (superficial) and secondary (deep) layers of retinal vessels coincident with the onset of vision. Initially, spoke-like peripapillary vessels grow radially over the pre-existing astrocytic network towards the periphery, becoming progressively interconnected by capillary plexus formation. These vessels grow as a monolayer within the nerve fiber through P10 (FIG. 1a). Between P7-P8 collateral branches begin to sprout from this primary plexus and penetrate into the retina to the outer plexiform layer where they form the secondary, or deep, retinal plexus. By P21, the entire network undergoes extensive remodeling and a tertiary, or intermediate, plexus forms at the inner surface of inner nuclear layer (FIG. 1b).

[0062] The neonatal mouse retinal angiogenesis model is useful for studying the role of HSC during ocular angiogenesis for several reasons. In this physiologically relevant model, a large astrocytic template exists prior to the appearance of endogenous blood vessels, permitting an evaluation of the role for cell-cell targeting during a neovascular process. In addition, this consistent and reproducible neonatal retinal vascular process is known to be hypoxia-driven, in this respect having similarities to many retinal diseases in which ischemia is known to play a role.

### Enrichment of Endothelial Progenitor Cells (EPC) From Bone Marrow

[0063] Although cell surface marker expression has been extensively evaluated on the EPC population found in preparations of HSC, markers that uniquely identify EPC are still poorly defined. To enrich for EPC, hematopoietic lineage marker positive cells (Lin<sup>+</sup>), i.e., B lymphocytes (CD45), T lymphocytes (CD3), granulocytes (Ly-6G), monocytes (CD11), and erythrocytes (TER-119), were depleted from bone marrow mononuclear cells of mice. Sca-1 antigen was used to further enrich for EPC. A comparison of results obtained after intravitreal injection of identical numbers of either Lin<sup>-</sup> Sca-1<sup>+</sup> cells or Lin<sup>-</sup> cells, no difference was detected between the two groups. In fact, when only Lin<sup>-</sup> Sca-1<sup>-</sup> cells were injected, far greater incorporation into developing blood vessels was observed.

[0064] The Lin<sup>-</sup>HSC populations of the present invention are enriched with EPCs, based on functional assays. Furthermore, Lin<sup>+</sup>HSC populations functionally behave quite differently from the Lin<sup>-</sup>HSC populations. Epitopes commonly used to identify EPC for each fraction (based on previously reported *in vitro* characterization studies) were also evaluated. While none of these markers were exclusively associated with the Lin<sup>-</sup> fraction, all were increased about 70 to about 1800% in the Lin<sup>-</sup> HSC, compared to the Lin<sup>+</sup>HSC fraction (FIG. 1c). FIG. 1c illustrates flow cytometric characterization of bone marrow-derived Lin<sup>+</sup> HSC and Lin<sup>-</sup> HSC separated cells. The top row of FIG. 1c shows a hematopoietic stem cell dot plot distribution of non-antibody labeled cells. R1 defines the quantifiable-gated area of positive PE-staining; R2 indicates GFP-positive. Dot plots of Lin<sup>-</sup> HSC are shown in the middle row and dot plots of Lin<sup>+</sup> HSC are shown in the bottom row. The C57B/6 cells were labeled with the PE-conjugated antibodies for Sca-1, c-kit, Flk-1/KDR, CD31. Tie-2 data was obtained from Tie-2-GFP mice. The percentages in the corners of the dot plots indicate the percent of positive-

labeled cells out of total Lin<sup>-</sup> or Lin<sup>+</sup> HSC population. Interestingly, accepted EPC markers like Flk-1/KDR, Tie-2, and Sca-1 were poorly expressed and, thus, not used for further fractionation.

### Intravitreally Injected HSC Lin<sup>-</sup> Cells Contain EPC That Target Astrocytes and Incorporate into Developing Retinal Vasculature

**[0065]** To determine whether intravitreally injected Lin<sup>-</sup> HSC can target specific cell types of the retina, utilize the astrocytic template and participate in retinal angiogenesis, approximately 10<sup>5</sup> cells from a Lin<sup>-</sup> HSC composition of the present invention or Lin<sup>+</sup> HSC cells (control, about 10<sup>5</sup> cells) isolated from the bone marrow of adult (GFP or LacZ transgenic) mice were injected into postnatal day 2 (P2) mouse eyes. Four days after injection (P6), many cells from the Lin<sup>-</sup> HSC composition of the present invention, derived from GFP or LacZ transgenic mice were adherent to the retina and had the characteristic elongated appearance of endothelial cells (FIG. 2a). FIG. 2 illustrates engraftment of Lin<sup>-</sup> cells into developing mouse retina. As shown in FIG. 2a, the four days post-injection (P6) intravitreally injected eGFP<sup>+</sup> Lin<sup>-</sup> HSC attach and differentiate on the retina.

**[0066]** In many areas of the retinas, the GFP-expressing cells were arranged in a pattern conforming to underlying astrocytes and resembled blood vessels. These fluorescent cells were observed ahead of the endogenous, developing vascular network (FIG. 2b). Conversely, only a small number of Lin<sup>+</sup>HSC (FIG. 2c), or adult mouse mesenteric endothelial cells (FIG. 2d) attached to the retinal surface. In order to determine whether cells from an injected Lin<sup>-</sup> HSC population could also attach to retinas with already established vessels, we injected a Lin<sup>-</sup> HSC composition into adult eyes. Interestingly, no cells were observed to attach to the retina or incorporate into established, normal retinal blood vessels (FIG. 2e). This indicates that the Lin<sup>-</sup> HSC compositions of the present invention do not disrupt a normally developed vasculature and will not initiate abnormal vascularization in normally developed retinas.

**[0067]** In order to determine the relationship between an injected Lin<sup>-</sup> HSC compositions of the present invention and retinal astrocytes, a transgenic mouse was used, which expressed glial fibrillary acidic protein (GFAP, a marker of astrocytes) and promoter-driven green fluorescent protein (GFP). Examination of retinas of these GFAP-GFP transgenic mice injected with Lin<sup>-</sup> HSC from eGFP transgenic mice demonstrated co-localization of the injected eGFP EPC and existing astrocytes (FIG. 2f-h, arrows). Processes of eGFP+Lin<sup>-</sup> HSC were observed to conform to the underlying astrocytic network (arrows, FIG. 2g). Examination of these eyes demonstrated that the injected, labeled cells only attached to astrocytes; in P6 mouse retinas, where the retinal periphery does not yet have endogenous vessels, injected cells were observed adherent to astrocytes in these not yet vascularized areas. Surprisingly, injected, labeled cells were observed in the deeper layers of the retina at the precise location where normal retinal vessels will subsequently develop (FIG. 2i, arrows).

**[0068]** To determine whether injected Lin<sup>-</sup> HSC of the present invention are stably incorporated into the developing retinal vasculature, retinal vessels at several later time points were examined. As early as P9 (seven days after injection), Lin<sup>-</sup> HSC incorporated into CD31<sup>+</sup> structures (FIG. 2j). By P16 (14 days after injection), the cells were already extensively incorporated into retinal vascular-like structures (FIG. 2k). When rhodamine-dextran was injected intravascularly (to identify functional retinal blood vessels) prior to sacrificing the animals, the majority of Lin<sup>-</sup> HSC were aligned with patent vessels (FIG. 2l). Two patterns of labeled cell distribution were observed: (1) in one pattern, cells were interspersed along vessels in between unlabeled endothelial cells; and (2) the other pattern showed that vessels were composed entirely of labeled cells. Injected cells were also incorporated into vessels of the deep vascular plexus (FIG. 2m). While sporadic incorporation of Lin<sup>-</sup> HSC-derived EPC into neovasculature has been previously reported, this is the first report of vascular networks being entirely composed of these cells. This demonstrates that cells from a population of bone marrow-derived Lin<sup>-</sup> HSC of the present invention injected intravitreally can efficiently incorporate into any layer of the forming retinal vascular plexus.

**[0069]** Histological examination of non-retinal tissues (e.g., brain, liver, heart, lung, bone marrow) did not demonstrate the presence of any GFP positive cells when examined up to 5 or 10 days after intravitreal injection. This indicates that a sub-population of cells within the Lin<sup>-</sup> HSC fraction selectively target to retinal astrocytes and stably incorporate into developing retinal vasculature. Since these cells have many characteristics of endothelial cells (association with retinal astrocytes, elongate morphology, stable incorporation into patent vessels and not present in extravascular locations), these cells represent EPC present in the Lin<sup>-</sup> HSC population. The targeted astrocytes are of the same type observed in many of the hypoxic retinopathies. It is well known that glial cells are a prominent component of neovascular fronds observed in DR and other forms of retinal injury. Under conditions of reactive gliosis and ischemia-induced neovascularization, activated astrocytes proliferate, produce cytokines, and up-regulate GFAP, similar to that observed during neonatal retinal vascular template formation in many mammalian species including humans.

**[0070]** Lin<sup>-</sup> HSC populations of the present invention will target activated astrocytes in adult mouse eyes as they do in neonatal eyes, Lin<sup>-</sup> HSC cells were injected into adult eyes with retinas injured by photo-coagulation (FIG. 3a) or needle tip (FIG. 3b). In both models, a population of cells with prominent GFAP staining was observed only around the injury site (FIG. 3a and b). Cells from injected Lin<sup>-</sup> HSC compositions localized to the injury site and remained specifically

associated with GFAP-positive astrocytes (FIG. 3a and b). At these sites, Lin<sup>-</sup> HSC cells were also observed to migrate into the deeper layer of retina at a level similar to that observed during neonatal formation of the deep retinal vasculature. Uninjured portions of retina contained no Lin<sup>-</sup> HSC cells, identical to that observed when Lin<sup>-</sup> HSC were injected into normal, uninjured adult retinas (FIG. 2e). These data indicate that Lin<sup>-</sup> HSC compositions can selectively target activated glial cells in injured adult retinas with gliosis as well as neonatal retinas undergoing vascularization.

### Intravitreally Injected Lin<sup>-</sup> HSC Can Rescue and Stabilize Degenerating Vasculature

**[0071]** Since intravitreally injected Lin<sup>-</sup> HSC compositions target astrocytes and incorporate into the normal retinal vasculature, these cells also stabilize degenerating vasculature in ischemic or degenerative retinal diseases associated with gliosis and vascular degeneration. The *rd/rd* mouse is a model for retinal degeneration that exhibits profound degeneration of photoreceptor and retinal vascular layers by one month after birth. The retinal vasculature in these mice develops normally until P16 at which time the deeper vascular plexus regresses; in most mice the deep and intermediate plexuses have nearly completely degenerated by P30.

**[0072]** To determine whether HSC can rescue the regressing vessels, Lin<sup>+</sup> or Lin<sup>-</sup> HSC (from Balb/c mice) were injected into *rd/rd* mice intravitreally at P6. By P33, after injection with Lin<sup>+</sup> cells, vessels of the deepest retinal layer were nearly completely absent (FIG. 4a and b). In contrast, most Lin<sup>-</sup> HSC-injected retinas by P33 had a nearly normal retinal vasculature with three parallel, well-formed vascular layers (FIG. 4a and 4d). Quantification of this effect demonstrated that the average length of vessels in the deep vascular plexus of Lin<sup>-</sup> injected *rd/rd* eyes was nearly three times greater than untreated or Lin<sup>+</sup> cell-treated eyes (FIG. 4e). Surprisingly, injection of a Lin<sup>-</sup> HSC composition derived from *rd/rd* adult mouse (FVB/N) bone marrow also rescued degenerating *rd/rd* neonatal mouse retinal vasculature (FIG. 4f). Degeneration of the vasculature in *rd/rd* mouse eyes is observed as early as 2-3 weeks post-natally. Injection of Lin<sup>-</sup> HSC as late as P15 also resulted in partial stabilization of the degenerating vasculature in the *rd/rd* mice for at least one month (FIG. 4g and 4h).

**[0073]** A Lin<sup>-</sup> HSC composition injected into younger (e.g., P2) *rd/rd* mice also incorporated into the developing superficial vasculature. By P11, these cells were observed to migrate to the level of the deep vascular plexus and form a pattern identical to that observed in the wild type outer retinal vascular layer (FIG. 5a). In order to more clearly describe the manner in which cells from injected Lin<sup>-</sup> HSC compositions incorporate into, and stabilize, degenerating retinal vasculature in the *rd/rd* mice, a Lin<sup>-</sup> HSC composition derived from Balb/c mice was injected into Tie-2-GFP FVB mouse eyes. The FVB mice have the *rd/rd* genotype and because they express the fusion protein Tie-2-GFP, all endogenous blood vessels are fluorescent.

**[0074]** When non-labeled cells from a Lin<sup>-</sup> HSC composition are injected into neonatal Tie-2-GFP FVB eyes and are subsequently incorporated into the developing vasculature, there should be non-labeled gaps in the endogenous, Tie-2-GFP labeled vessels that correspond to the incorporated, non-labeled Lin<sup>-</sup> HSC that were injected. Subsequent staining with another vascular marker (e.g., CD-31) then delineates the entire vessel, permitting determination as to whether non-endogenous endothelial cells are part of the vasculature. Two months after injection, CD31-positive, Tie-2-GFP negative, vessels were observed in the retinas of eyes injected with the Lin<sup>-</sup> HSC composition (FIG. 5b). Interestingly, the majority of rescued vessels contained Tie-2-GFP positive cells (FIG. 5c). The distribution of pericytes, as determined by staining for smooth muscle actin, was not changed by Lin<sup>-</sup> HSC injection, regardless of whether there was vascular rescue (FIG. 5d). These data clearly demonstrate that intravitreally injected Lin<sup>-</sup> HSC compositions of the present invention migrate into the retina, participate in the formation of normal retinal blood vessels, and stabilize endogenous degenerating vasculature in a genetically defective mouse.

### Inhibition of Retinal Angiogenesis by Transfected Cells from Lin<sup>-</sup> HSC

**[0075]** The majority of retinal vascular diseases involve abnormal vascular proliferation rather than degeneration. Transgenic cells targeted to astrocytes can be used to deliver an anti-angiogenic protein and inhibit angiogenesis. Cells from Lin<sup>-</sup> HSC compositions were transfected with T2-tryptophanyl-tRNA synthetase (T2-TrpRS). T2-TrpRS is a 43 kD fragment of TrpRS that potently inhibits retinal angiogenesis (FIG. 6a). On P12, retinas of eyes injected with a control plasmid-transfected Lin<sup>-</sup> HSC composition (no T2-TrpRS gene) on P2 had normal primary (FIG. 6c) and secondary (FIG. 6d) retinal vascular plexuses. When the T2-TrpRS transfected Lin<sup>-</sup> HSC composition of the present invention was injected into P2 eyes and evaluated 10 days later, the primary network had significant abnormalities (FIG. 6e) and formation of the deep retinal vasculature was nearly completely inhibited (FIG. 6f). The few vessels observed in these eyes were markedly attenuated with large gaps between vessels. The extent of inhibition by T2-TrpRS-secreting Lin<sup>-</sup> HSCs is detailed in Table 2.

**[0076]** T2-TrpRS is produced and secreted by cells in the Lin<sup>-</sup> HSC composition *in vitro* and after injection of these transfected cells into the vitreous, a 30 kD fragment of T2-TrpRS in the retina (FIG. 6b) was observed. This 30 kD fragment was specifically observed only in retinas injected with transfected Lin<sup>-</sup> HSC of the present invention and this

decrease in apparent molecular weight compared to the recombinant or *in vitro*-synthesized protein may be due to processing or degradation of the T2-TrpRS *in vivo*. These data indicate that Lin<sup>-</sup> HSC compositions can be used to deliver functionally active genes, such as genes expressing angiostatic molecules, to the retinal vasculature by targeting to activated astrocytes. While it is possible that the observed angiostatic effect is due to cell-mediated activity this is very unlikely since eyes treated with identical, but non-T2-transfected Lin<sup>-</sup> HSC compositions had normal retinal vasculature.

**Table 2. Vascular Inhibition by T2-TrpRS-secreting Lin<sup>-</sup> HSCs**

	Primary Plexus		Deep Plexus		
	Inhibited	Normal	Complete	Partial	Normal
TsTrpRs (15 eyes)	60% (9 eyes)	40% (6 eyes)	33.3% (5 eyes)	60% (9 eyes)	6.7% (1 eye)
Control (13 eyes)	0% (0 eyes)	100% (13 eyes)	0% (0 eyes)	38.5% (5 eyes)	61.5% (8 eyes)

**[0077]** Intravitreally injected Lin<sup>-</sup> HSC populations localize to retinal astrocytes, incorporate into vessels, and can be useful in treating many retinal diseases. While most cells from injected HSC compositions adhere to the astrocytic template, small numbers migrate deep into the retina, homing to regions where the deep vascular network will subsequently develop. Even though no GFAP-positive astrocytes were observed in this area prior to 42 days postnatally, this does not rule out the possibility that GFAP-negative glial cells are already present to provide a signal for Lin<sup>-</sup> HSC localization. Previous studies have shown that many diseases are associated with reactive gliosis. In DR, in particular, glial cells and their extracellular matrix are associated with pathological angiogenesis.

**[0078]** Since cells from injected Lin<sup>-</sup> HSC compositions specifically attached to GFAP-expressing glial cells, regardless of the type of injury, Lin<sup>-</sup> HSC compositions of the present invention can be used to target pre-angiogenic lesions in the retina. For example, in the ischemic retinopathies such as diabetes, neovascularization is a response to hypoxia. By targeting Lin<sup>-</sup> HSC compositions to sites of pathological neovascularization, developing neovasculature can be stabilized preventing abnormalities of neovasculature such as hemorrhage or edema (the causes of vision loss associated with DR) and can potentially alleviate the hypoxia that originally stimulated the neovascularization. Abnormal blood vessels can be restored to normal condition. Furthermore, angiostatic proteins, such as T2-TrpRS can be delivered to sites of pathological angiogenesis by using transfected Lin<sup>-</sup> HSC compositions and laser-induced activation of astrocytes. Since laser photocoagulation is a commonly used in clinical ophthalmology, this approach has application for many retinal diseases. While such cell-based approaches have been explored in cancer therapy, their use for eye diseases is more advantageous since intraocular injection makes it possible to deliver large numbers of cells directly to the site of disease.

### Neurotrophic and Vasculotropic Rescue by Lin<sup>-</sup>HSC

**[0079]** MACS was used to separate Lin<sup>-</sup> HSC from bone marrow of enhanced green fluorescent protein (eGFP), C3H (*rd/rd*), FVB (*rd/rd*) mice as described above. Lin<sup>-</sup> HSC containing EPC from these mice were injected intravitreally into P6 C3H or FVB mouse eyes. The retinas were collected at various time points (1 month, 2 months, and 6 months) after injection. The vasculature was analyzed by scanning laser confocal microscope after staining with antibodies to CD31 and retinal histology after nuclear staining with DAPI. Microarray gene expression analysis of mRNA from retinas at varying time points was also used to identify genes potentially involved in the effect.

**[0080]** Eyes of *rd/rd* mice had profound degeneration of both neurosensory retina and retinal vasculature by P21. Eyes of *rd/rd* mice treated with Lin<sup>-</sup> HSC on P6 maintained a normal retinal vasculature for as long as 6 months; both deep and intermediate layers were significantly improved when compared to the controls at all timepoints (1M, 2M, and 6M) (see FIG. 12). In addition, we observed that retinas treated with Lin<sup>-</sup>HSC were also thicker (1M; 1.2-fold, 2M; 1.3-fold, 6M; 1.4-fold) and had greater numbers of cells in the outer nuclear layer (1M; 2.2-fold, 2M; 3.7-fold, 6M; 5.7-fold) relative to eyes treated with Lin<sup>+</sup> HSC as a control. Large scale genomic analysis of "rescued" (e.g., Lin<sup>-</sup> HSC) compared to control (untreated or non-Lin<sup>-</sup> treated) *rd/rd* retinas demonstrated a significant upregulation of genes encoding sHSPs (small heat shock proteins) and specific growth factors that correlated with vascular and neural rescue, including genes encoding the proteins listed in Figure 20, panels A and B.

**[0081]** The bone marrow derived Lin<sup>-</sup> HSC populations of the present invention significantly and reproducibly induced maintenance of a normal vasculature and dramatically increased photoreceptor and other neuronal cell layers in the *rd/rd* mouse. This neurotrophic rescue effect correlated with significant upregulation of small heat shock proteins and growth factors and provides insights into therapeutic approaches to currently untreatable retinal degenerative disorders.

***Rd1/rd1* Mouse Retinas Exhibit Profound Vascular and Neuronal Degeneration.**

**[0082]** Normal postnatal retinal vascular and neuronal development in mice has been well described and is analogous to changes observed in the third trimester human fetus (Dorrell *et al.*, 2002, *Invest. Ophthalmol. Vis. Sci.* 43:3500-3510). Mice homozygous for the *rd1* gene share many characteristics of human retinal degeneration (Frasson *et al.*, 1999, *Nat. Med.* 5:1183-1187) and exhibit rapid photoreceptor (PR) loss accompanied by severe vascular atrophy as the result of a mutation in the gene encoding PR cGMP phosphodiesterase (Bowes *et al.* 1990, *Nature* 347:677-680). To examine the vasculature during retinal development and its subsequent degeneration, antibodies against collagen IV (CIV), an extracellular matrix (ECM) protein of mature vasculature, and CD31 (PECAM-1), a marker for endothelial cells were used (Figure 15). Retinas of *rd1/rd1* (C3H/HeJ) developed normally until approximately postnatal day (P) 8 when degeneration of the photoreceptor-containing outer nuclear layer (ONL) began. The ONL rapidly degenerated and cells died by apoptosis such that only a single layer of nuclei remained by P20. Double staining of the whole-mounted retinas with antibodies to both CIV and CD31 revealed details of the vascular degeneration in *rd1/rd1* mice similar to that described by others (Blanks *et al.*, 1986, *J. Comp. Neurol.* 254:543-553). The primary and deep retinal vascular layers appeared to develop normally though P12 after which there is a rapid loss of endothelial cells as evidenced by the absence of CD31 staining. CD31 positive endothelial cells were present in a normal distribution through P12 but rapidly disappeared after that. Interestingly, CIV positive staining remained present throughout the time points examined, suggesting that the vessels and associated ECM formed normally, but only the matrix remained after P13 by which time no CD31 positive cells were observed. (Figure 15, middle panels). The intermediate vascular plexus also degenerates after P21, but the progression is slower than that observed in the deep plexus (Figure 15, upper panel). Retinal vascular and neural cell layers of a normal mouse are shown for comparison to the *rd1/rd1* mouse (right panels, Figure 15).

**Neuroprotective Effect of Bone Marrow-Derived Lin<sup>-</sup> HSCs in *rd1/rd1* Mice.**

**[0083]** Intravitreally injected Lin<sup>-</sup> HSCs incorporate into endogenous retinal vasculature in all three vascular plexuses and prevent the vessels from degenerating. Interestingly, the injected cells are virtually never observed in the outer nuclear layer. These cells either incorporate into the forming retinal vessels or are observed in close proximity to these vessels. Murine Lin<sup>-</sup> HSCs (from C3H/HeJ) were intravitreally injected into C3H/HeJ (*rd1/rd1*) mouse eyes at P6, just prior to the onset of degeneration. By P30, control cell (CD31<sup>-</sup>)-injected eyes exhibited the typical *rd1/rd1* phenotype, i.e., nearly complete degeneration of the deep vascular plexus and ONL was observed in every retina examined. Eyes injected with Lin<sup>-</sup> HSCs maintained normal-appearing intermediate and deep vascular plexuses. Surprisingly, significantly more cells were observed in the internuclear layer (INL) and ONL of Lin<sup>-</sup> HSC-injected eyes than in control cell-injected eyes (Figure 16A). This rescue effect of Lin<sup>-</sup> HSCs could be observed at 2 months (Figure 16B) and for as long as 6 months after injection (Figure 16C). Differences in the vasculature of the intermediate and deep plexuses of Lin<sup>-</sup> HSC-injected eyes, as well as the neuronal cell-containing INL and ONL, were significant at all time points measured when rescued and non-rescued eyes were compared (Figure 16B and C). This effect was quantified by measuring the total length of the vasculature (Figure 16D) and counting the number of DAPI-positive cell nuclei observed in the ONL (Figure 16E). Simple linear-regression analysis was applied to the data at all time points.

**[0084]** A statistically significant correlation was observed between vascular rescue and neuronal (e.g., ONL thickness) rescue at P30 ( $p < 0.024$ ) and P60 ( $p < 0.034$ ) in the Lin<sup>-</sup> HSC-injected eyes (Figure 16F). The correlation remained high, although not statistically significant ( $p < 0.14$ ) at P180 when comparing Lin<sup>-</sup> HSC-injected retinas to control cell-injected retinas (Figure 16F). In contrast, control cell-injected retinas showed no significant correlation between the preservation of vasculature and ONL at any time point (Figure 16F). These data demonstrate that intravitreal injection of Lin<sup>-</sup> HSCs results in concomitant retinal vascular and neuronal rescue in retinas of *rd1/rd1* mice. Injected cells were not observed in the ONL or any place other than within, or in close proximity to, retinal blood vessels.

**Functional Rescue of Lin<sup>-</sup> HSC-injected *rd1/rd1* Retinas**

**[0085]** Electroretinograms (ERGs) were performed on mice 2 months after injection of control cells or murine Lin<sup>-</sup> HSCs (Figure 17). Immunohistochemical and microscopic analysis was done with each eye following ERG recordings to confirm that vascular and neuronal rescue had occurred. Representative ERG recordings from treated, rescued and control, non-rescued eyes show that in the rescued eyes, the digitally subtracted signal (treated minus untreated eyes) produced a clearly detectable signal with an amplitude on the order of 8-10 microvolts (Figure 17). Clearly, the signals from both eyes are severely abnormal. However, consistent and detectable ERGs were recordable from the Lin<sup>-</sup> HSC-treated eyes. In all cases the ERG from the control eye was non-detectable. While the amplitudes of the signals in rescued eyes were considerably lower than normal, the signals were consistently observed whenever there was histological rescue and were on the order of magnitude of those reported by other, gene based, rescue studies. Overall these results demonstrate of some degree of functional rescue in the eyes treated with the Lin<sup>-</sup> HSCs of the invention.

### Human bone marrow (hBM)-derived Lin<sup>-</sup> HSCs also Rescue Degenerating Retinas

[0086] Lin<sup>-</sup> HSCs isolated from human bone marrow behave similarly to murine Lin<sup>-</sup> HSCs. Bone marrow was collected from human donors and the Lin<sup>+</sup> HSCs were depleted, producing a population of human Lin<sup>-</sup> HSCs (hLin<sup>-</sup> HSCs). These cells were labeled ex-vivo with fluorescent dye and injected into C3SnSmn.CB17-*Prkdc* SCID mouse eyes. The injected hLin<sup>-</sup> HSCs migrated to, and targeted, sites of retinal angiogenesis in a fashion identical to that observed when murine Lin<sup>-</sup> HSCs were injected (Figure 18A). In addition to the vascular targeting, the human Lin<sup>-</sup> HSCs also provided a robust rescue effect on both the vascular and neuronal cell layers of the *rd1/rd1* mice (Figure 18B and 18C). This observation confirms the presence of cells in human bone marrow that target retinal vasculature and can prevent retinal degeneration.

### Lin<sup>-</sup> HSCs have Vasculo- and Neurotrophic Effects in the *rd10/rd10* Mouse

[0087] While the *rd1/rd1* mouse is the most widely used and best characterized model for retinal degeneration (Chang *et al.* 2002, *Vision Res.* 42:517-525), the degeneration is very rapid and in this regard differs from the usual, slower time course observed in the human disease. In this strain, photoreceptor cell degeneration begins around P8, a time when the retinal vasculature is still rapidly expanding (Figure 15). Subsequent degeneration of the deep retinal vasculature occurs even while the intermediate plexus is still forming and, thus, the retinas of *rd1/rd1* mice never completely develops, unlike that observed in most humans with this disease. An *rd10* mouse model, which has a slower time course of degeneration and more closely resembles the human retinal degenerative condition, was used to investigate Lin<sup>-</sup> HSC-mediated vascular rescue. In the *rd10* mouse, photoreceptor cell degeneration begins around P21 and vascular degeneration begins shortly thereafter.

[0088] Since normal neurosensory retinal development is largely complete by P21, the degeneration is observed to start after the retina has completed differentiation and in this way is more analogous to human retinal degenerations than the *rd1/rd1* mouse model. Lin<sup>-</sup> HSCs or control cells from *rd10* mice were injected into P6 eyes and the retinas were evaluated at varying time points. At P21 the retinas from both Lin<sup>-</sup> HSC and control cell-injected eyes appeared normal with complete development of all vascular layers and normal development of the INL and ONL (Figure 18D and 18H). At approximately P21 the retinal degeneration began and progressed with age. By P30, the control cell-injected retinas exhibited severe vascular and neuronal degeneration (Figure 18I), while the Lin<sup>-</sup> HSC-injected retinas maintained nearly normal vascular layers and photoreceptor cells (Figure 18E). The difference between the rescued and non-rescued eyes was more pronounced at later time points (compare Figure 18F and 18G to 18J and 18K). In the control treated eyes, the progression of vascular degeneration was very clearly observed by immunohistochemical staining for CD31 and collagen IV (Figure 18I-K). The control-treated eyes were nearly completely negative for CD31, whereas collagen IV-positive vascular "tracks" remained evident, indicating that vascular regression, rather than incomplete vascular formation, had occurred. In contrast, Lin<sup>-</sup> HSC-treated eyes had both CD31 and collagen IV-positive vessels that appeared very similar to normal, wild-type eyes (compare Figure 18F and 18I).

### Gene Expression Analysis of *rd1/rd1* Mouse Retinas after Lin<sup>-</sup> HSC Treatment

[0089] Large scale genomics (microarray analysis) was used to analyze rescued and non-rescued retinas to identify putative mediators of neurotrophic rescue. Gene expression in *rd1/rd1* mouse retinas treated with Lin<sup>-</sup> HSCs was compared to uninjected retinas as well as retinas injected with control cells (CD31<sup>-</sup>). These comparisons were each done in triplicate. To be considered present, genes were required to have expression levels at least 2-fold higher than background levels in all three triplicates. Genes that were upregulated 3-fold in Lin<sup>-</sup> HSC-protected retinas compared to control cell-injected and non-injected *rd1/rd1* mouse retinas are shown in FIG. 20, panels A and B. Many of the significantly upregulated genes, including MAD and Ying Yang-1 (YY-1), encode proteins with functions involving the protection of cells from apoptosis. A number of crystallin genes, which have sequence homology and similar functions to known heat-shock proteins involving protection of cells from stress, were also upregulated by Lin<sup>-</sup> HSC treated retinas. Expression of  $\alpha$ -crystallin was localized to the ONL by immunohistochemical analysis (FIG. 19).

[0090] Messenger RNA from *rd1/rd1* mouse retinas rescued with human Lin<sup>-</sup> HSCs were hybridized to human specific Affymetrix U133A microarray chips. After stringent analysis, a number of genes were found whose mRNA expression was human specific, above background, and significantly higher in the human Lin<sup>-</sup> HSC rescued retinas compared to the murine Lin<sup>-</sup> HSC rescued retinas and the human control cell-injected non-rescued retinas (FIG. 20, panel C). CD6, a cell adhesion molecule expressed at the surface of primitive and newly differentiated CD34<sup>+</sup> hematopoietic stem cells, and interferon alpha 13, another gene expressed by hematopoietic stem cells, were both found by the microarray bioinformatics technique, validating the evaluation protocol. In addition, several growth factors and neurotrophic factors were expressed above background by human Lin<sup>-</sup> HSC rescued mouse retina samples (FIG. 20, panel D).

## DISCUSSION

5 [0091] Markers for lineage-committed hematopoietic cells were used to negatively select a population of bone marrow-derived Lin<sup>-</sup> HSC containing EPC. While the sub-population of bone marrow-derived Lin<sup>-</sup> HSC that can serve as EPC is not characterized by commonly used cell surface markers, the behavior of these cells in developing or injured retinal vasculature is entirely different than that observed for Lin<sup>+</sup> or adult endothelial cell populations. These cells selectively target to sites of retinal angiogenesis and participate in the formation of patent blood vessels.

10 [0092] Inherited retinal degenerative diseases are often accompanied by loss of retinal vasculature. Effective treatment of such diseases requires restoration of function as well as maintenance of complex tissue architecture. While several recent studies have explored the use of cell-based delivery of trophic factors or stem cells themselves, some combination of both may be necessary. For example, use of growth factor therapy to treat retinal degenerative disease resulted in unregulated overgrowth of blood vessels resulting in severe disruption of the normal retinal tissue architecture. The use of neural or retinal stem cells to treat retinal degenerative disease may reconstitute neuronal function, but a functional vasculature will also be necessary to maintain retinal functional integrity. Incorporation of cells from a Lin<sup>-</sup> HSCs of the present invention into the retinal vessels of *rd/rd* mice stabilized the degenerative vasculature without disrupting retinal structure. This rescue effect was also observed when the cells were injected into P15 *rd/rd* mice. Since vascular degeneration begins on P16 in *rd/rd* mice, this observation expands the therapeutic window for effective Lin<sup>-</sup> HSC treatment. Retinal neurons and photoreceptors are preserved and visual function is maintained in eyes injected with the Lin<sup>-</sup> HSC of the present invention.

20 [0093] Adult bone marrow-derived Lin<sup>-</sup> HSCs exert profound vasculo- and neurotrophic effects when injected intraventrically into mice with retinal degenerative disease. This rescue effect persists for up to 6 months after treatment and is most efficacious when the Lin<sup>-</sup> HSCs are injected prior to complete retinal degeneration (up to 16 days after birth in mice that ordinarily exhibit complete retinal degeneration by 30 days postnatally). This rescue is observed in 2 mouse models of retinal, degeneration and, remarkably, can be accomplished with adult human bone marrow-derived HSCs when the recipient is an immunodeficient rodent with retinal degeneration (e.g., the SCID mouse) or when the donor is a mouse with retinal degeneration. While several recent reports have described a partial phenotypic rescue in mice or dogs with retinal degeneration after viral based gene rescue with the wild type gene (Ali, *et al.* 2000, *Nat Genet* 25:306-310; Takahashi *et al.* 1999, *J. Virol.* 73:7812-7816; Acland *et al.* 2001, *Nat. Genet.* 28:92-95.), the present invention is the first generic cell-based therapeutic effect achieved by vascular rescue. Thus, the potential utility of such an approach in treating a group of diseases (e.g., retinitis pigmentosa) with over 100 known associated mutations is more practical than creating individual gene therapies to treat each known mutation.

25 [0094] The precise molecular basis of the neurotrophic rescue effect remains unknown, but is observed only when there is concomitant vascular stabilization/rescue. The presence of injected stem cells, per se, is not sufficient to generate a neurotrophic rescue and the clear absence of stem cell-derived neurons in the outer nuclear layer rules out the possibility that the injected cells are transforming into photoreceptors. Data obtained by microarray gene expression analysis demonstrated a significant up-regulation of genes known to have anti-apoptotic effects. Since most neuronal death observed in retinal degenerations is by apoptosis, such protection may be of great therapeutic benefit in prolonging the life of photoreceptors and other neurons critical to visual function in these diseases. C-myc is a transcription factor that participates in apoptosis by upregulation various downstream apoptosis-inducing factors. C-myc expression was increased 4.5 fold in *rd/rd* mice over wild-type indicating potential involvement in the photoreceptor degeneration observed in the *rd1/rd1* mouse. Mad1 and YY-1, two genes dramatically upregulated in Lin<sup>-</sup> HSC-protected retinas (Fig. 20, panel A), are known to suppress the activity of c-myc, thus inhibiting c-myc induced apoptosis. Overexpression of Mad1 has also been shown to suppress Fas-induced activation of caspase-8, another critical component of the apoptotic pathway. Upregulation of these two molecules may play a role in protection of the retina from vascular and neural degeneration by preventing the initiation of apoptosis that normally leads to degeneration in *rd/rd* mice.

35 [0095] Another set of genes that were greatly upregulated in Lin<sup>-</sup>HSC protected retinas includes members of the crystallin family (Fig 20, panel B). Similar to heat-shock and other stress-induced proteins, crystallins may be activated by retinal stress and provide a protective effect against apoptosis. Abnormally low expression of  $\alpha$ A-crystallins is correlated with photoreceptor loss in a rat model of retinal dystrophy and a recent proteomic analysis of the retina in the *rd/rd* mouse demonstrated induction of crystalline up-regulation in response to retinal degeneration. Based on our microarray data of EPC-rescued *rd/rd* mouse retinas, upregulation of crystallins appear to play a key role in EPC mediated retinal neuroprotection.

40 [0096] Genes such as c-myc, Mad1, Yx-1 and the crystallins are likely to be downstream mediators of neuronal rescue. Neurotrophic agents can regulate anti-apoptotic gene expression, although our microarray analysis of retinas rescued with mouse stem cells did not demonstrate induction of increased levels of known neurotrophic factors. Analysis of human bone marrow-derived stem cell-mediated rescue with human specific chips did, on the other hand, demonstrate low, but significant increases in the expression of multiple growth factor genes.

45 [0097] The upregulated genes include several members of the fibroblast growth factor family and otoferlin. Mutations

in the otoferlin gene are associated with genetic disorders leading to deafness due to auditory neuropathy. It is possible that otoferlin production by injected Lin<sup>-</sup> HSCs contributes to the prevention of retinal neuropathy as well. Historically, it has long been assumed that vascular changes observed in patients and animals with retinal degeneration were secondary to decreased metabolic demand as the photoreceptors die. The present data indicate that, at least for mice with inherited

5 retinal degeneration, preserving normal vasculature can help maintain components of the outer nuclear layer as well. Recent reports in the literature would support the concept that tissue-specific vasculature has trophic effects that go beyond that expected from simply providing vascular "nourishment." For example, liver endothelial cells can be induced to produce, after VEGFR1 activation, growth factors critical to hepatocyte regeneration and maintenance in the face of hepatic injury (LeCouter *et al.* 2003, *Science* 299:890-893).

10 **[0098]** Similar indicative interactions between vascular endothelial cells and adjacent hepatic parenchymal cells are reportedly involved in liver organogenesis, well before the formation of functional blood vessels. Endogenous retinal vasculature in individuals with retinal degeneration may not facilitate so dramatic a rescue, but if this vasculature is buttressed with endothelial progenitors derived from bone marrow hematopoietic stem cell populations, they may make the vasculature more resistant to degeneration and at the same time facilitate retinal neuronal, as well as vascular,

15 survival. In humans with retinal degeneration, delaying the onset of complete retinal degeneration may provide years of additional sight. The animals treated with the Lin<sup>-</sup> HSCs of the present invention had significant preservation of an ERG, which may be sufficient to support vision.

**[0099]** Clinically, it is widely appreciated that there can be substantial loss of photoreceptors and other neurons while still preserving functional vision. At some point, the critical threshold is crossed and vision is lost. Since nearly all of the

20 human inherited retinal degenerations are of early, but slow, onset, it may be possible to identify an individual with retinal degeneration, treat them intravitreally with an autologous bone marrow stem cell graft and delay retinal degeneration with concomitant loss of vision. To enhance targeting and incorporation of these stem cells, the presence of activated astrocytes would be desirable. This can be accomplished by early treatment when there is an associated gliosis or by using a laser to stimulate local proliferation of activated astrocytes.

25 **[0100]** The Lin<sup>-</sup> HSC populations of the present invention contain a population of EPC that can promote angiogenesis by targeting reactive astrocytes and incorporate into an established template without disrupting retinal structure. The Lin<sup>-</sup> HSC of the present invention also provide a surprising long-term neurotrophic rescue effect in eyes suffering from retinal degeneration. In addition, genetically modified, autologous Lin<sup>-</sup> HSC compositions containing EPC can be trans-

30 planted into ischemic or abnormally vascularized eyes and can stably incorporate into new vessels and continuously deliver therapeutic molecules locally for prolonged periods of time. Such local delivery of genes that express pharmacological agents in physiologically meaningful doses represents a new paradigm for treating currently untreatable ocular diseases.

35 **[0101]** Numerous variations and modifications of the embodiments described above may be effected without departing from the spirit and scope of the novel features of the invention. No limitations with respect to the specific embodiments illustrated herein are intended or should be inferred.

40

45

50

55

## SEQUENCE LISTING

<110> The Scripps Research Institute

<120> TRANSFECTED HEMATOPOIETIC STEM CELLS AND METHODS OF TREATMENT OF NEOVASCULAR EYE DISEASES THEREWITH

<130> 18-197 T1

<140> Divisional application of EP 04 760 595.1

<141> 2004-04-28

<150> US 60/467,051

<151> 2003-05-02

<150> US 10/628,783

<151> 2003-07-25

<160> 5

<170> FastSEQ for windows Version 4.0

<210> 1

<211> 4742

<212> DNA

<213> Artificial sequence

<220>

<223> DNA encoding His-tagged human T2-TrpRS

<400> 1

tggcgaatgg	gacgcgccct	gtagcggcgc	attaagcgcg	gcgggtgtgg	tggttacgcg	60
cagcgtgacc	gctacacttg	ccagcgcgcc	agcgcgccgt	cctttcgcct	tcttcccttc	120
ctttctcgcc	acgttcgccc	gctttccccg	tcaagctcta	aatcgggggc	tccctttagg	180
gttccgattt	agtgttttac	ggcacctcga	ccccaaaaaa	cttgattagg	gtgatgggtc	240
acgtagtggt	ccatcgccct	gatagacggt	ttttcgccct	ttgacgttgg	agtccacggt	300
ctttaatagt	ggactcttgt	tccaaactgg	aacaacactc	aaccctatct	cggtctattc	360
ttttgattta	taaggggattt	tgccgatttc	ggcctattgg	ttaaaaaatg	agctgattta	420
acaaaaattt	aacgcgaatt	taaacaaaa	attaacggtt	acaatttcag	gtggcacttt	480
tcggggaaat	gtgcgcggaa	cccctatttg	tttatttttc	taaatacatt	caaatatgta	540
tccgctcatg	agacaataac	cctgataaat	gcttcaataa	tattgaaaaa	ggaagagtat	600
gagtattcaa	catttccggtg	tcgcccctat	tccctttttt	gcggcatttt	gccttccctg	660
ttttgctcac	ccagaaacgc	tggtgaaagt	aaaagatgct	gaagatcagt	tggtgtcacg	720
agtgggttac	atcgaactgg	atctcaacag	cggtaaagatc	cttgagagtt	ttcgccccga	780
agaagctttt	ccaatgatga	gcacttttaa	agttctgcta	tgtggcgcg	tattatcccc	840
tattgacgcc	gggcaagagc	aactcggctg	ccgcatacac	tattctcaga	atgacttggg	900
tgagtactca	ccagtcacag	aaaagcatct	tacggatggc	atgacagtaa	gagaattatg	960
cagtgtgccc	ataacatga	gtgataaac	tcggccaac	ttacttctga	caacgatcgg	1020
aggaccgaag	gagctaaccg	cttttttgca	caacatgggg	gatcatgtaa	ctcgccttga	1080
tcgttgggaa	ccggagctga	atgaagccat	accaaacgac	gagcgtgaca	ccacgatgcc	1140
tgagcaatg	gcaacaacgt	tcgcaaac	attaactggc	gaactactta	ctctagcttc	1200
ccggcaacaa	taatatagact	ggatggaggc	ggataaagtt	gcaggaccac	ttctgcgctc	1260
ggccttccg	tgtggctggt	ttattgctga	taaacttggg	gccgggtgagc	gtgggtctcg	1320
cggtatcatt	gcagcactgg	ggccagatgg	taagccctcc	cgatcgtag	ttatctacac	1380
gacggggagt	caggcaacta	tggatgaacg	aatagacag	atcgctgaga	taggtgcctc	1440
actgattaag	cattggtaac	tgtcagacca	agtttactca	tatatacttt	agattgattt	1500
aaaacttcat	ttttaattta	aaaggatcta	ggtgaagatc	ctttttgata	atctcatgac	1560
caaaatccct	taacgtgagt	tttcgttcca	ctgagcgtca	gaccccgtag	aaaagatcaa	1620
aggatcttct	tgagatcctt	tttttctg	cgtaactctg	tgcttgcaaa	caaaaaaacc	1680
accgctacca	gcggtgggtt	gtttgccgga	tcaagagcta	ccaactcttt	ttccgaaggt	1740
aactggcttc	agcagagcgc	agataccaaa	tactgtcctt	ctagtgtagc	cgtagttagg	1800
ccaccacttc	aagaactctg	tagcaccgcc	tacatacctc	gctctgctaa	tcctgttacc	1860
agtggctgct	gccagtggcg	ataagctgtg	tcttaccggg	ttggactcaa	gacgatagtt	1920
accggataag	gcgcagcgg	cgggctgaac	ggggggttcg	tgcacacagc	ccagcttggg	1980
gcgaacgacc	tacaccgaac	tgagatacct	acagcgtgag	ctatgagaaa	gcgccacgct	2040

tcccaaggga agaaaggcgg acaggtatcc ggtaagcggc agggtcggaa caggagagcg 2100  
 caccagggag cttccagggg gaaacgcctg gtatctttat agtcctgtcg ggtttcgcca 2160  
 cctctgactt gagcgtcgat ttttgtgatg ctctgcaggg gggcggagcc tatggaaaaa 2220  
 cgccagcaac gcggcctttt tacggttcct ggcccttttg tggccttttg ctcacatggt 2280  
 5 ctttcctgcg ttatcccctg attctgtgga taaccgtatt accgcctttg agtgagctga 2340  
 taccgctcgc cgcagccgaa cgaccgagcg cagcgagtca gtgagcgagg aagcgggaaga 2400  
 gcgcctgatg cggatatttc tccttacgca tctgtgaggg atttcacacc gcataatagg 2460  
 tgcactctca gtacaatctg ctctgatgcc gcatagttaa gccagttatac actccgctat 2520  
 cgctacgtga ctgggtcatg gctgcgcccc gacacccgcc aacacccgct gacgcgcctt 2580  
 gacgggcttg tctgctcccc gcacccgctt acagacaagc tgtgaccgtc tccgggagct 2640  
 10 gcatgtgtca gaggttttca ccgtcatcac cgaaacgcgc gaggcagctg cggtaaagct 2700  
 catcagcgtg gtcgtgaagc gattcacaga tgtctgcctg ttcacccgcg tccagctcgt 2760  
 tgagtttctc cagaagcgtt aatgtctggc ttctgataaa gcggggccatg ttaagggcgg 2820  
 ttttttctctg tttggctact gatgcctccg tgtaaggggg atttctgttc atgggggtaa 2880  
 tgataccgat gaaacgagag aggatgtcta cgatacgggt tactgatgat gaacatgccc 2940  
 ggttactgga acgttgtgag ggtaaacaac tggcgggatg gatgcggcgg gaccagagaa 3000  
 15 aaatcactca gggcaatgc cagcgtctcg ttaatacaga ttaggtgtt ccacagggta 3060  
 gccagcagca tcctgcatg cagatccgga acataatgt gcagggcgct gacttccgcg 3120  
 tttccagact ttacgaaaca cggaaaccga agaccattca tgttgttgc cagtcgca 3180  
 acgttttgca gcagcagctg cttcacgttc gctcgcgtat cggtgattca ttctgtaac 3240  
 cagtaaggca accccgccag cctagccggg tcctcaacga caggagcacg atcatgcgca 3300  
 cccgtggcca ggacccaacg ctgcccgaga tctcgatccc gcgaaattaa tacgactcac 3360  
 tataaggaga ccacaacggt ttccctctag aaataathtt gtttaacttt aagaaggaga 3420  
 20 tatacatatg agtgcaaaag gcatagacta cgataagctc attgttcggt ttggaagtag 3480  
 taaaattgac aaagagctaa taaaccgaat agagagagcc accggccaaa gaccacacca 3540  
 cttcctgcmc agaggcatct tcttctcaca cagagatatg aatcaggttc ttgatgccta 3600  
 tgaaaataag aagccathtt atctgtacac gggccggggc ccctcttctg aagcaatgca 3660  
 tgtaggtcac ctcatccat ttathttcac aaagtggctc caggatgtat ttaacgtgcc 3720  
 25 cttggctcatc cagatgacgg atgacgagaa gtatctgtgg aaggacctga ccctggacca 3780  
 ggccataggc gatgctgttg agaatgcaa ggacatcatc gcctgtggct ttgacatcaa 3840  
 caagactttc atattctctg acctggacta catggggatg agctcagggt tctacaaaaa 3900  
 tgtggtgaaag attcaaaagc atgttacctt caaccaagtg aaaggcattt tcggcttcac 3960  
 tgacagcagc tgcattggga agatcagttt tcttgccatc caggctgctc cctccttcag 4020  
 caactcattc ccacagatct tccgagacag gacggatata cagtgcctta tcccatgtgc 4080  
 cattgaccag gatccttact ttagaatgac aaggacgct gcccccagga tcggctatcc 4140  
 30 taaaccagcc ctgtttgact ccaccttctt cccagccctg cagggcgccc agacaaaaat 4200  
 gaagtgccagc gacccaaact cctccatctt cctcaccgac acggccaagc agatcaaaaac 4260  
 aaggttcaat aagcatgctg tttctggagg gagagacacc atcgaggagc acaggcagtt 4320  
 tgggggcaac tgtgatgtgg acgtgtcttt catgtacctg accttcttcc tcgaggacga 4380  
 cgacaagctc gagcagatca ggaaggatta caccagcggg gccatgctca ccggtgagct 4440  
 caagaaggca ctcatagagg ttctgcagcc cttgatcgca gagcaccagg cccggcgcaa 4500  
 35 ggaggtcacg gatgagatag tgaagagtt cactactccc cggaagctgt ccttcgactt 4560  
 tcagaagctt gcggccgcac tcgagcacca ccaccaccac cactgagatc cggctgctaa 4620  
 caaagcccga aaggaagctg agttggctgc tgccaccgct gagcaataac tagcataacc 4680  
 ccttggggcc tctaaacggg tcttgagggg ttttttgctg aaaggaggaa ctatatccgg 4740  
 at 4742

<210> 2  
 <211> 392  
 <212> PRT  
 <213> Artificial sequence

<220>  
 <223> His-tagged human T2-TrpRS

<400> 2  
 Met Ser Ala Lys Gly Ile Asp Tyr Asp Lys Leu Ile Val Arg Phe Gly  
 1 5 10 15  
 Ser Ser Lys Ile Asp Lys Glu Leu Ile Asn Arg Ile Glu Arg Ala Thr  
 20 25 30  
 50 Gly Gln Arg Pro His His Phe Leu Arg Arg Gly Ile Phe Phe Ser His  
 35 40 45  
 Arg Asp Met Asn Gln Val Leu Asp Ala Tyr Glu Asn Lys Lys Pro Phe  
 50 55 60  
 Tyr Leu Tyr Thr Gly Arg Gly Pro Ser Ser Glu Ala Met His Val Gly  
 65 70 75 80  
 55 His Leu Ile Pro Phe Ile Phe Thr Lys Trp Leu Gln Asp Val Phe Asn  
 85 90 95

EP 1 754 783 A1

Val Pro Leu Val Ile Gln Met Thr Asp Asp Glu Lys Tyr Leu Trp Lys  
 100  
 Asp Leu Thr Leu Asp Gln Ala Tyr Gly Asp Ala Val Glu Asn Ala Lys  
 115  
 5 Asp Ile Ile Ala Cys Gly Phe Asp Ile Asn Lys Thr Phe Ile Phe Ser  
 130  
 Asp Leu Asp Tyr Met Gly Met Ser Ser Gly Phe Tyr Lys Asn Val Val  
 145 150 155  
 Lys Ile Gln Lys His Val Thr Phe Asn Gln Val Lys Gly Ile Phe Gly  
 165 175  
 10 Phe Thr Asp Ser Asp Cys Ile Gly Lys Ile Ser Phe Pro Ala Ile Gln  
 180 185  
 Ala Ala Pro Ser Phe Ser Asn Ser Phe Pro Gln Ile Phe Arg Asp Arg  
 195 200  
 Thr Asp Ile Gln Cys Leu Ile Pro Cys Ala Ile Asp Gln Asp Pro Tyr  
 210 215 220  
 15 Phe Arg Met Thr Arg Asp Val Ala Pro Arg Ile Gly Tyr Pro Lys Pro  
 225 230 235  
 Ala Leu Leu His Ser Thr Phe Phe Pro Ala Leu Gln Gly Ala Gln Thr  
 245 250 255  
 Lys Met Ser Ala Ser Asp Pro Asn Ser Ser Ile Phe Leu Thr Asp Thr  
 260 265 270  
 20 Ala Lys Gln Ile Lys Thr Lys Val Asn Lys His Ala Phe Ser Gly Gly  
 275 280 285  
 Arg Asp Thr Ile Glu Glu His Arg Gln Phe Gly Gly Asn Cys Asp Val  
 290 295 300  
 Asp Val Ser Phe Met Tyr Leu Thr Phe Phe Leu Glu Asp Asp Asp Lys  
 305 310 315  
 25 Leu Glu Gln Ile Arg Lys Asp Tyr Thr Ser Gly Ala Met Leu Thr Gly  
 325 330 335  
 Glu Leu Lys Lys Ala Leu Ile Glu Val Leu Gln Pro Leu Ile Ala Glu  
 340 345 350  
 His Gln Ala Arg Arg Lys Glu Val Thr Asp Glu Ile Val Lys Glu Phe  
 355 360 365  
 30 Met Thr Pro Arg Lys Leu Ser Phe Asp Phe Gln Lys Leu Ala Ala Ala  
 370 375 380  
 Leu Glu His His His His His His

<210> 3  
 <211> 379  
 <212> PRT  
 <213> Artificial sequence

<220>  
 <223> Human T2-TrpRS

<400> 3  
 Met Ser Ala Lys Gly Ile Asp Tyr Asp Lys Leu Ile Val Arg Phe Gly  
 1 5 10 15  
 Ser Ser Lys Ile Asp Lys Glu Leu Ile Asn Arg Ile Glu Arg Ala Thr  
 20 25 30  
 35 Gly Gln Arg Pro His His Phe Leu Arg Arg Gly Ile Phe Phe Ser His  
 40 45  
 Arg Asp Met Asn Gln Val Leu Asp Ala Tyr Glu Asn Lys Lys Pro Phe  
 50 55 60  
 Tyr Leu Tyr Thr Gly Arg Gly Pro Ser Ser Glu Ala Met His Val Gly  
 65 70 75 80  
 His Leu Ile Pro Phe Ile Phe Thr Lys Trp Leu Gln Asp Val Phe Asn  
 85 90 95  
 50 Val Pro Leu Val Ile Gln Met Thr Asp Asp Glu Lys Tyr Leu Trp Lys  
 100 105 110  
 Asp Leu Thr Leu Asp Gln Ala Tyr Ser Tyr Ala Val Glu Asn Ala Lys  
 115 120 125  
 Asp Ile Ile Ala Cys Gly Phe Asp Ile Asn Lys Thr Phe Ile Phe Ser  
 130 135 140  
 55 Asp Leu Asp Tyr Met Gly Met Ser Ser Gly Phe Tyr Lys Asn Val Val



EP 1 754 783 A1

210 215 220  
 Ser Asp Cys Ile Gly Lys Ile Ser Phe Pro Ala Ile Gln Ala Ala Pro  
 225 230 235 240  
 Ser Phe Ser Asn Ser Phe Pro Gln Ile Phe Arg Asp Arg Thr Asp Ile  
 5 Gln Cys Leu Ile Pro Cys Ala Ile Asp Gln Asp Pro Tyr Phe Arg Met  
 260 265 270  
 Thr Arg Asp Val Ala Pro Arg Ile Gly Tyr Pro Lys Pro Ala Leu Leu  
 275 280  
 His Ser Thr Phe Phe Pro Ala Leu Gln Gly Ala Gln Thr Lys Met Ser  
 290 295 300  
 Ala Ser Asp Pro Asn Ser Ser Ile Phe Leu Thr Asp Thr Ala Lys Gln  
 305 310 315 320  
 Ile Lys Thr Lys Val Asn Lys His Ala Phe Ser Gly Gly Arg Asp Thr  
 325 330 335  
 Ile Glu Glu His Arg Gln Phe Gly Gly Asn Cys Asp Val Asp Val Ser  
 340 345 350  
 15 Phe Met Tyr Leu Thr Phe Phe Leu Glu Asp Asp Asp Lys Leu Glu Gln  
 355 360 365  
 Ile Arg Lys Asp Tyr Thr Ser Gly Ala Met Leu Thr Gly Glu Leu Lys  
 370 375 380  
 20 Lys Ala Leu Ile Glu Val Leu Gln Pro Leu Ile Ala Gln His Gln Ala  
 385 390 395 400  
 Arg Arg Lys Glu Val Thr Asp Glu Ile Val Lys Glu Phe Met Thr Pro  
 405 410 415  
 Arg Lys Leu Ser Phe Asp Phe Gln  
 420

25 <210> 5  
 <211> 401  
 <212> PRT  
 <213> Artificial sequence

30 <220>  
 <223> Human T1-TrpRS

<400> 5  
 Ser Asn His Gly Pro Asp Ala Thr Glu Ala Glu Glu Asp Phe Val Asp  
 1 5 10 15  
 Pro Trp Thr Val Gln Thr Ser Ser Ala Lys Gly Ile Asp Tyr Asp Lys  
 20 25 30  
 35 Leu Ile Val Arg Phe Gly Ser Ser Lys Ile Asp Lys Glu Leu Ile Asn  
 35 40 45  
 Arg Ile Glu Arg Ala Thr Gly Gln Arg Pro His His Phe Leu Arg Arg  
 50 55 60  
 Gly Ile Phe Phe Ser His Arg Asp Met Asn Gln Val Leu Asp Ala Tyr  
 65 70 75 80  
 40 Glu Asn Lys Lys Pro Phe Tyr Leu Tyr Thr Gly Arg Gly Pro Ser Ser  
 85 90 95  
 Glu Ala Met His Val Gly His Leu Ile Pro Phe Ile Phe Thr Lys Trp  
 100 105 110  
 Leu Gln Asp Val Phe Asn Val Pro Leu Val Ile Gln Met Thr Asp Asp  
 115 120 125  
 45 Glu Lys Tyr Leu Trp Lys Asp Leu Thr Leu Asp Gln Ala Tyr Ser Tyr  
 130 135 140  
 Ala Val Glu Asn Ala Lys Asp Ile Ile Ala Cys Gly Phe Asp Ile Asn  
 145 150 155 160  
 Lys Thr Phe Ile Phe Ser Asp Leu Asp Tyr Met Gly Met Ser Ser Gly  
 165 170 175  
 50 Phe Tyr Lys Asn Val Val Lys Ile Gln Lys His Val Thr Phe Asn Gln  
 180 185 190  
 Val Lys Gly Ile Phe Gly Phe Thr Asp Ser Asp Cys Ile Gly Lys Ile  
 195 200 205  
 Ser Phe Pro Ala Ile Gln Ala Ala Pro Ser Phe Ser Asn Ser Phe Pro  
 210 215 220  
 55 Gln Ile Phe Arg Asp Arg Thr Asp Ile Gln Cys Leu Ile Pro Cys Ala  
 225 230 235 240

Ile Asp Gln Asp Pro Tyr Phe Arg Met Thr Arg Asp Val Ala Pro Arg  
 Ile Gly Tyr Pro Lys Pro Ala Leu Leu His Ser Thr Phe Phe Pro Ala  
 5 Leu Gln Gly Ala Gln Thr Lys Met Ser Ala Ser Asp Pro Asn Ser Ser  
 Ile Phe Leu Thr Asp Thr Ala Lys Gln Ile Lys Thr Lys Val Asn Lys  
 His Ala Phe Ser Gly Gly Arg Asp Thr Ile Glu Glu His Arg Gln Phe  
 10 305 310 315 320  
 Gly Gly Asn Cys Asp Val Asp Val Ser Phe Met Tyr Leu Thr Phe Phe  
 325 330 335  
 Leu Glu Asp Asp Asp Lys Leu Glu Gln Ile Arg Lys Asp Tyr Thr Ser  
 340 345  
 Gly Ala Met Leu Thr Gly Glu Leu Lys Lys Ala Leu Ile Glu Val Leu  
 15 355 360 365  
 Gln Pro Leu Ile Ala Glu His Gln Ala Arg Arg Lys Glu Val Thr Asp  
 370 375 380  
 Glu Ile Val Lys Glu Phe Met Thr Pro Arg Lys Leu Ser Phe Asp Phe  
 385 390 395 400  
 Gln

**Claims**

1. A transfected mammalian lineage negative hematopoietic stem cell population comprising a stem cell population of hematopoietic stem cells and endothelial progenitor cells, wherein cells from the stem cell population express a therapeutically useful anti-angiogenic protein fragment of human tryptophanyl tRNA synthetase (TrpRS).
2. The transfected stem cell population of claim 1 wherein at least about 20% of the cells express the surface antigen CD31.
3. The transfected stem cell population of claim 1 wherein at least about 50% of the cells express the surface antigen CD31.
4. The transfected stem cell population of claim 1 wherein at least about 75% of the cells express the surface antigen CD31.
5. The transfected stem cell population of claim 1 wherein at least about 50% of the cells express the surface antigen for integrin  $\alpha_6$ .
6. The transfected stem cell population of claim 1 wherein the cells are derived from adult bone marrow.
7. The transfected stem cell population of claim 1 wherein the cells are murine cells.
8. The transfected stem cell population of claim 7 wherein at least about 50% of the cells express the surface antigen CD31 and at least about 50% of the cells express the surface antigen CD 117.
9. The transfected stem cell population of claim 7 wherein at least about 65% of the cells express the surface antigen CD117.
10. The transfected stem cell population of claim 7 wherein at least about 80% of the cells express the surface antigen CD31 and at least about 70% of the cells express the surface antigen CD 117.
11. The transfected stem cell population of claim 1 wherein the cells are human cells.
12. The transfected stem cell population of claim 11 wherein the cells are CD 133 negative, at least about 50% of the cells express the surface antigen for integrin  $\alpha_6$ , and at least about 50% of the cells express the surface antigen CD31.

## EP 1 754 783 A1

13. The transfected stem cell population of claim 11 wherein the cells are CD 133 positive, less than about 30% of the cells express the surface antigen for integrin  $\alpha_6$ , and less than about 30% of the cells express the surface antigen CD31.
- 5 14. The transfected stem cell population of claim 1 further including a cell culture medium.
15. The transfected stem cell population of claim 1 wherein the cell population is further transfected with a DNA operably encoding a neurotrophic agent.
- 10 16. The transfected stem cell population of claim 15 wherein the neurotrophic agent is selected from the group consisting of nerve growth factor, neurotrophin-3, neurotrophin-4, neurotrophin-5, ciliary neurotrophic factor, retinal pigmented epithelium-derived neurotrophic factor, insulin-like growth factor, glial cell line-derived neurotrophic factor, and brain-derived neurotrophic factor.
- 15 17. The transfected stem cell population of claim 1 for delivering transgenes to the retinal vasculature of a mammal by intravitreally injecting the transfected stem cell population into the eye of the mammal.
18. The transfected stem cell population of claim 1 for targeted delivery of an anti-angiogenic protein fragment of TrpRS to astrocytes in the eye of a mammal by intravitreally injecting the transfected stem cell population into the eye of  
20 the mammal.
19. The transfected stem cell population of any one of claims 1, 17 and 18 wherein the transfected stem cell population expresses a fragment of TrpRS which is T2-TrpRS (SEQ ID NO: 3).
- 25 20. Use of a transfected stem cell population of claim 1 for the preparation of a pharmaceutical composition for treating a vascular disease of the eye in a mammal wherein the pharmaceutical composition is for intravitreally injecting the transfected stem cell population into an eye of the mammal.
- 30 21. The use of claim 20 wherein the disease is selected from the group consisting of a retinal degenerative disease, a retinal vascular degenerative disease, an ischemic retinopathy, a vascular hemorrhage, a vascular leakage, a choroidopathy, age related macular degeneration, diabetic retinopathy, presumed ocular histoplasmosis, retinopathy of prematurity, sickle cell anemia, and retinitis pigmentosa.
- 35 22. Use of a transfected stem cell population of claim 1 for the preparation of a pharmaceutical composition for rescuing neuronal networks in the eye of a mammal wherein the pharmaceutical composition is for intravitreally injecting the transfected stem cell population into the eye of the mammal.
- 40 23. Use of a transfected stem cell population of claim 1 for the preparation of a pharmaceutical composition for stimulating upregulation of anti-apoptotic genes in the eye of a mammal wherein the pharmaceutical composition is for intravitreally injecting the transfected stem cell population into the eye of the mammal.
24. The use of any one of claims 20, 22 and 23 wherein the transfected stem cell population expresses a fragment of TrpRS which is T2-TrpRS (SEQ ID NO: 3).
- 45 25. Use of a transfected stem cell population of claim 1 for the preparation of a pharmaceutical composition for rescuing blood vessels in the eye of a mammal wherein the pharmaceutical composition is for intravitreally injecting the transfected stem cell population into the eye of the mammal.
- 50 26. The use of claim 25 wherein the transfected stem cell population expresses a fragment of TrpRS selected from the group consisting of T2-TrpRS (SEQ ID NO: 3), mini-TrpRS (SEQ ID NO: 4), and T1-TrpRS (SEQ ID NO: 5).

55




FIG. 1

C




FIG. 1 Cont.




FIG. 2




FIG. 2 Cont.




FIG. 3




FIG. 4




FIG. 5




FIG. 6

## DNA encoding His-tagged T2 fragment of human TrpRS

```

tggcgaatgg gacgcgccct gtagcggcgc attaagcgcg gcgggtgtgg tggttacgcg 60
cagcgtgacc gctacacttg ccagcgcctc agcgcgccgt cctttcgctt tcttcccttc 120
ctttctcgcc acgttcgcgc gctttccccc tcaagctcta aatcgggggc tccctttagg 180
gttccgattt agtgctttac ggcacctoga ccccaaaaaa cttgattagg gtgatggttc 240
acgtagtggg ccatcgccct gatagacggt ttttcgccct ttgacgttgg agtccacggt 300
ctttaatagt ggactcttgt tccaaactgg aacaacactc aaccctatct cggctctattc 360
ttttgattta taagggattt tgccgatttc ggcctattgg ttaaaaaatg agctgattta 420
acaaaaatth aacgcgaatt ttaacaaaat attaacgttt acaatttcag gtggcacttt 480
tcggggaaat gtgcgcgaa cccctatttg tttatttttc taaatacatt caaatatgta 540
tccgctcatg agacaataac cctgataaat gcttcaataa tattgaaaaa ggaagagtat 600
gagtattcaa catttccgtg tcgcccttat tccctttttt gcggcatttt gccttctctg 660
ttttgctcac ccagaaacgc tgggtgaaagt aaaagatgct gaagatcagt tgggtgcacg 720
agtgggttac atcgaactgg atctcaacag cggtaagatc cttgagagtt ttcgccccga 780
agaacgtttt ccaatgatga gcacttttaa agttctgcta tgtggcgcgg tattatcccg 840
tattgacgcc gggcaagagc aactcggtcg ccgcatacac tattctcaga atgacttggg 900
tgagtactca ccagtcacag aaaagcatct tacggatggc atgacagtaa gagaattatg 960
cagtgtctgcc ataaccatga gtgataaac tgccggccaac ttacttctga caacgatcgg 1020
aggaccgaag gagctaaccg cttttttgca caacatgggg gatcatgtaa ctgccttga 1080
tcgttgggaa ccggagctga atgaagccat accaaacgac gagcgtgaca ccacgatgcc 1140
tcgagcaatg gcaacaacgt tcgcgaaact attaactggc gaactactta ctctagcttc 1200
ccggcaacaa ttaatagact ggatggaggc ggataaagtt gcaggaccac ttctgcgctc 1260
ggcccttccg gctggctggt ttattgctga taaatctgga gccgggtgagc gtgggtctcg 1320
cggtatcatt gcagcactgg ggccagatgg taagccctcc cgtatcgtag ttatctacac 1380
gacggggagt caggcaacta tggatgaacg aaatagacag atcgtgaga taggtgcctc 1440
actgattaag catttgtaac tgtcagacca agtttactca tataactttt agattgattt 1500
aaaacttcat ttttaattta aaaggatcta ggtgaagatc ctttttgata atctcatgac 1560
caaaatccct taacgtgagt tttcgttcca ctgagcgtca gaccccgtag aaaagatcaa 1620
aggatcttct tgagatcctt tttttctgcg cgtaatctgc tgcttgcaaa caaaaaaac 1680
accgctacca ccggtggttt gtttgccgga tcaagagcta ccaactcttt tccgaaggt 1740
aactggcttc agcagagcgc agataccaaa tactgtcctt ctagtgtagc cgtagttagg 1800
ccaccacttc aagaactctg tagcacgccc tacatacctc gctctgctaa tctgtttacc 1860
agtggctgct gccagtgccg cggctgaac ggggggttcg tgacacagc ccagcttggg 1920
accgataag gcgcagcggg tgagatacct acagcgtgag ctatgagaaa gcgccacgct 2040
gccaagagg agaaaggcgg acaggatcc ggtaagcggc aggttcggaa caggagagcg 2100
cagcagggag cttccagggg gaaacgcctg gtatctttat agtcctgtcg ggtttcgcca 2160
cctctgactt gagcgtcgat ttttgtgatg ctgctcaggg gggcggagcc tatggaaaaa 2220
cgccagcaac gcggcctttt tacggttctt ggccttttgc tgcacatggt tggccttttg 2280
ctttcctgcg ttatcccctg attctgtgga taaccgtatt accgcctttg agtgagctga 2340
taccgctcgc cgcagccgaa cgaccgagcg cagcaggtca gtgagcaggg aagcgggaaga 2400
gcgcctgatg cggtatthttc tccctacgca tctgtgcggg atttcacacc gcatatatgg 2460
tgactcttca gtacaatctg ctctgatgcc gcatagttaa gccagtatac actccgctat 2520
cgctacgtga ctgggtcatg gctgcccccc gacacccgcc aacacccgct gacgcgccct 2580
gacgggcttg tctgctcccg gcatccgctt acagacaagc tgtgaccgtc tccgggagct 2640
gcatgtgtca gaggttttca ccgtcatcac cgaaacgcgc gaggcagctg cggtaaagct 2700
catcagcgtg gtcgtgaagc gattcacaga tgtctgcctg ttcacccgcg tccagctcgt 2760
tgagtttctc cagaagcgtt aatgtctggc ttctgataaa gcgggccaatg ttaagggcgg 2820
tttttctctg tttggtcact gatgcctccg tgtaaggggg atttctgttc atgggggtaa 2880
tgataccgat gaaacgagag aggatgctca cgatacgggt tactgatgat gaacatgccc 2940
ggttactgga acgttgtgag ggtaaacaaac tggcggtatg gatgcggcgg gaccagagaa 3000
aaatcactca gggtaaatgc cagcgtctcg ttaatacaga tgtaggtggt ccacagggta 3060
gccagcagca tcttgcgatg cagatccgga acataatggt gcagggcgct gacttccgcg 3120
ttccagact ttacgaaaca cggaaaccga agaccattca tgtgtgtgct caggtcgcag 3180
acgttttgca gcagcagtcg cttcacgttc gctcgcgtat cggtgattca ttctgctaac 3240
cagtaaggca accccgccag cctagccggg tcccaacga caggagcacg atcatgcgca 3300

```

FIG. 7

```

cccgtggcca ggaccaacg ctgcccgaga tctcgatccc gcgaaattaa tacgactcac 3360
tatagggaga ccacaacggt ttccctctag aaataatfff gtttaacttt aagaaggaga 3420
tatacat atg agt gca aaa ggc ata gac tac gat aag ctc att gtt cgg 3469
ttt gga agt agt aaa att gac aaa gag cta ata aac cga ata gag aga 3517
gcc acc ggc caa aga cca cac cac ttc ctg cgc aga ggc atc ttc ttc 3565
tca cac aga gat atg aat cag gtt ctt gat gcc tat gaa aat aag aag 3613
cca ttt tat ctg tac acg ggc cgg ggc ccc tct tct gaa gca atg cat 3661
gta ggt cac ctc att cca ttt att ttc aca aag tgg ctc cag gat gta 3709
ttt aac gtg ccc ttg gtc atc cag atg acg gat gac gag aag tat ctg 3757
tgg aag gac ctg acc ctg gac cag gcc tat ggc gat gct gtt gag aat 3805
gcc aag gac atc atc gcc tgt ggc ttt gac atc aac aag act ttc ata 3853
ttc tct gac ctg gac tac atg ggg atg agc tca ggt ttc tac aaa aat 3901
gtg gtg aag att caa aag cat gtt acc ttc aac caa gtg aaa ggc att 3949
ttc ggc ttc act gac agc gac tgc att ggg aag atc agt ttt cct gcc 3997
atc cag gct gct ccc tcc ttc agc aac tca ttc cca cag atc ttc cga 4045
gac agg acg gat atc cag tgc ctt atc cca tgt gcc att gac cag gat 4093
cct tac ttt aga atg aca agg gac gtc gcc ccc agg atc ggc tat cct 4141
aaa cca gcc ctg ttg cac tcc acc ttc ttc cca gcc ctg cag ggc gcc 4189
cag acc aaa atg agt gcc agc gac cca aac tcc tcc atc ttc ctc acc 4237
gac acg gcc aag cag atc aaa acc aag gtc aat aag cat gcg ttt tct 4285
gga ggg aga gac acc atc gag gag cac agg cag ttt ggg ggc aac tgt 4333
gat gtg gac gtg tct ttc atg tac ctg acc ttc ttc ctc gag gac gac 4381
gac aag ctc gag cag atc agg aag gat tac acc agc gga gcc atg ctc 4429
acc ggt gag ctc aag aag gca ctc ata gag gtt ctg cag ccc ttg atc 4477
gca gag cac cag gcc cgg cgc aag gag gtc acg gat gag ata gtg aaa 4525
gag ttc atg act ccc cgg aag ctg tcc ttc gac ttt cag aag ctt gcg 4573
gcc gca ctc gag cac cac cac cac cac cac tgagatccgg ctgctaacia 4623
agcccgaaag gaagctgagt tggctgctgc caccgctgag caataactag cataaccct 4683
tggggcctct aaacgggtct tgaggggttt tttgctgaaa ggaggaacta tatccggat 4742

```

FIG. 7 Cont.

## His-tagged T2 fragment of human TrpRS

Met Ser Ala Lys Gly Ile Asp Tyr Asp Lys Leu Ile Val Arg Phe Gly  
 1 5 10 15  
 Ser Ser Lys Ile Asp Lys Glu Leu Ile Asn Arg Ile Glu Arg Ala Thr  
 20 25 30  
 Gly Gln Arg Pro His His Phe Leu Arg Arg Gly Ile Phe Phe Ser His  
 35 40 45  
 Arg Asp Met Asn Gln Val Leu Asp Ala Tyr Glu Asn Lys Lys Pro Phe  
 50 55 60  
 Tyr Leu Tyr Thr Gly Arg Gly Pro Ser Ser Glu Ala Met His Val Gly  
 65 70 75 80  
 His Leu Ile Pro Phe Ile Phe Thr Lys Trp Leu Gln Asp Val Phe Asn  
 85 90 95  
 Val Pro Leu Val Ile Gln Met Thr Asp Asp Glu Lys Tyr Leu Trp Lys  
 100 105 110  
 Asp Leu Thr Leu Asp Gln Ala Tyr Gly Asp Ala Val Glu Asn Ala Lys  
 115 120 125  
 Asp Ile Ile Ala Cys Gly Phe Asp Ile Asn Lys Thr Phe Ile Phe Ser  
 130 135 140  
 Asp Leu Asp Tyr Met Gly Met Ser Ser Gly Phe Tyr Lys Asn Val Val  
 145 150 155 160  
 Lys Ile Gln Lys His Val Thr Phe Asn Gln Val Lys Gly Ile Phe Gly  
 165 170 175  
 Phe Thr Asp Ser Asp Cys Ile Gly Lys Ile Ser Phe Pro Ala Ile Gln  
 180 185 190  
 Ala Ala Pro Ser Phe Ser Asn Ser Phe Pro Gln Ile Phe Arg Asp Arg  
 195 200 205  
 Thr Asp Ile Gln Cys Leu Ile Pro Cys Ala Ile Asp Gln Asp Pro Tyr  
 210 215 220  
 Phe Arg Met Thr Arg Asp Val Ala Pro Arg Ile Gly Tyr Pro Lys Pro  
 225 230 235 240  
 Ala Leu Leu His Ser Thr Phe Phe Pro Ala Leu Gln Gly Ala Gln Thr  
 245 250 255  
 Lys Met Ser Ala Ser Asp Pro Asn Ser Ser Ile Phe Leu Thr Asp Thr  
 260 265 270  
 Ala Lys Gln Ile Lys Thr Lys Val Asn Lys His Ala Phe Ser Gly Gly  
 275 280 285  
 Arg Asp Thr Ile Glu Glu His Arg Gln Phe Gly Gly Asn Cys Asp Val  
 290 295 300  
 Asp Val Ser Phe Met Tyr Leu Thr Phe Phe Leu Glu Asp Asp Asp Lys  
 305 310 315 320  
 Leu Glu Gln Ile Arg Lys Asp Tyr Thr Ser Gly Ala Met Leu Thr Gly  
 325 330 335  
 Glu Leu Lys Lys Ala Leu Ile Glu Val Leu Gln Pro Leu Ile Ala Glu  
 340 345 350  
 His Gln Ala Arg Arg Lys Glu Val Thr Asp Glu Ile Val Lys Glu Phe  
 355 360 365  
 Met Thr Pro Arg Lys Leu Ser Phe Asp Phe Gln Lys Leu Ala Ala Ala  
 370 375 380  
 Leu Glu His His His His His His  
 385 390

FIG. 8






FIG. 9

Simple Regression Analysis -2M



Int. Vasculature

$$y=0.005328x+86.228974$$
$$r=0.452836$$
$$p=0.1888$$




Deep Vasculature

$$y=0.008873x+142.327$$
$$r=0.669651$$
$$p=0.0342$$


FIG. 10

Simple Regression Analysis – 2M Control



Deep Vasculature

$$y=0.003517x+47.465646$$
$$r=0.349191$$
$$p=0.3227$$



Int. Vasculature

$$y=0.000056x+52.905069$$
$$r=0.019418$$
$$p=0.9575$$

**FIG. 11**




FIG. 12

Number of Nuclei in ONL




FIG. 13




FIG. 14

FIG. 15




FIG. 16




FIG. 17





FIG. 18



$\alpha$ A-crystallin expression in rescued *rd/rd* retina



green:  $\alpha$ A-crystallin, blue: DAPI

**FIG. 19**

**(A) Genes whose expression is increased 3-fold in Lin- HSC injected retinas**

Common Name	(normalized data)			Genbank	Keywords
	Lin (-)	CD31 (-)	NI rd/rd		
Tgfp	11.86	0.53	0.66	L38444	T-cell-specific protein
H-2D4(q)	7.09	0.92	0.69	X52914	transplantation antigen
H2-K2; H-2K2	4.51	0.70	0.55	M27134	cell surface glycoprotein
Lzp-s	6.51	0.65	0.99	X51547	lysozyme; lysozyme P
Kcnj5	4.50	0.85	0.72	U33631	G-protein gated K+ channel
Scya8	5.19	0.47	1.00	AB023418	MCP-2 precursor
Ly6a	4.02	0.98	0.79	X04653	Ly-6 alloantigen
Anxa1	2.49	0.60	0.51	AV003419	EST
Pip5k1c	3.40	0.94	0.78	AB006916	phosphatidylinositolkinase
MAD	3.76	0.56	0.89	X83106	MAX dimerization protein
Cxadr	3.98	0.81	1.00	U90715	coxsackie-adenovirus receptor
Isg15	2.22	0.64	0.45	X56602	interferon inducible protein
EST	3.51	0.90	0.98	AA790936	EST
Tm4sf1	3.02	0.49	0.70	AV087000	EST
IgG VH-II	2.64	0.95	0.91	X02463	Ig heavy chain; variable region
YY1	2.97	0.85	0.87	M74590	delta-transcription factor
Psmb9	3.29	0.49	0.97	D44456	polypeptide complex subunit 2
H2-Aa	2.63	0.88	0.94	X52643	I-E alpha NON, MHC

**(B) Crystallin genes**

Common Name	(normalized data)			Genbank	Keywords
	Lin (-)	CD31 (-)	NI rd/rd		
Crybb2	8.73	0.55	0.83	M60559	beta-B2-crystallin
Cryaa	4.00	0.57	1.00	J00376	alpha-A-crystallin
CrygD	2.09	0.74	0.97	AJ224342	gamma-D-crystallin
Cryba1	6.52	0.93	0.60	AJ239052	beta-A3/A1-crystallin
Crygs	2.89	0.97	0.85	AF032995	gamma-S-crystallin
CrygC	5.07	1.00	0.83	Z22574	gamma-C-crystallin
CrygF	1.94	1.00	0.69	AJ224343	gamma-F-crystallin

**(C) Genes upregulated at least 2-fold in human stem cell rescued retinas**

Common Name	(raw data)			Genbank	Keywords
	MR	HR	HNR		
EST	397	1885	489	AK024177	EST
CD6	960	2155	840	NM_006725	CD6 antigen
IFN $\alpha$ 13	241	506	220	NM_006900	interferon, alpha 13
TRAG3	269	543	229	NM_004909	taxol resistance associated gene 3
ELA2	229	442	187	NM_001972	elastase 2, neutrophil
PP32D	179	345	146	NM_012404	acidic nuclear phosphoprotein 32D
otoferlin	232	485	239	NM_004802	otoferlin
acyl oxidase	175	332	164	NM_004035	acyl-Coenzyme A oxidase
HS6ST1	204	385	210	NM_004807	heparan sulfate 6-O-sulfotransferase
ACCN3	187	337	181	NM_004769	testis amiloride-sensitive cation channel 3

**(D) Neurotrophic factors or growth factor genes upregulated in human stem cell rescued retinas**

Common Name	(raw data)			Genbank	Keywords
	MR	HR	HNR		
IGF2	743	1245	839	NM_016412	insulin-like growth factor 2
SCGF	474	547	402	NM_002975	stem cell growth factor
TGF $\beta$ 1	841	1011	819	NM_000660	transforming growth factor beta 1
EGFR S8	499	593	491	A1343292	epidermal growth factor receptor pathway substrate 8-related prot. 1
FGF-6	524	611	526	NM_020996	fibroblast growth factor 6
IGFBP	927	991	895	AW338791	insulin-like growth factor binding prot.
FGF-12	471	523	507	AL119322	fibroblast growth factor 12
EGF-like	991	1092	1088	M60278	diphtheria toxin receptor, EGF-like
FGF-21	592	615	590	NM_019113	fibroblast growth factor 21

FIG. 20






FIG. 21




Post natal development of retinal vasculature in wild type C57/Bl6 mice raised in normoxia.

**FIG. 22**

**FIG. 23**



**Oxygen-induced retinopathy model: Post natal development of retinal vasculature in C57/Bl6 mice raised in 75% oxygen between P7 and P12, returned to normoxia from P12-P17 and analyzed with fluorescein dextran imaging in the confocal microscope at P17.**



Rescue effect of LinHSC in OIR model. CD31-CD34-CD11b<sup>+</sup> HSC (left panels, "Cntrl") or LinHSC (right panels) were injected intravitreally on P3 into the left or right eyes, respectively, in the same animal. On P7 the animal was placed into 75% oxygen for 5 days, normoxia on P12 and analyzed by confocal microscopy on P17. Red is lectin and blue is DAPI staining.

**FIG. 24**



European Patent Office

**PARTIAL EUROPEAN SEARCH REPORT**

Application Number

which under Rule 45 of the European Patent Convention EP 05 02 5978 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	OTANI ATSUSHI ET AL: "Bone marrow-derived stem cells target retinal astrocytes and can promote or inhibit retinal angiogenesis." NATURE MEDICINE. SEP 2002, vol. 8, no. 9, September 2002 (2002-09), pages 1004-1010, XP002988691 ISSN: 1078-8956 * the whole document * -----	1-10, 12-14, 17,18, 20-23,25	INV. C12N5/06
P,L, X	WO 2004/010959 A2 (SCRIPPS RESEARCH INST [US]) 5 February 2004 (2004-02-05)  invalidates priority * page 17, paragraph 2 - page 27, paragraph 2 *  ----- -/--	1-14,17, 18, 20-23,25	
			TECHNICAL FIELDS SEARCHED (IPC)
			C12N
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search: see sheet C</p>			
Place of search		Date of completion of the search	Examiner
The Hague		10 January 2007	Manu, Dominique
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

7  
EPO FORM 1503 03.02 (P04C07)



DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (IPC)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	<p>YIN A H ET AL: "AC133, a novel marker for human hematopoietic stem and progenitor cells."                      BLOOD. 15 DEC 1997,                      vol. 90, no. 12,                      15 December 1997 (1997-12-15), pages                      5002-5012, XP002264586                      ISSN: 0006-4971                      * page 5005, right-hand column, paragraph 3 *                      * figure 3b *</p> <p style="text-align: center;">-----</p>	12,13	
			TECHNICAL FIELDS SEARCHED (IPC)



Claim(s) not searched:  
15,16,19,24,26

Reason for the limitation of the search:

The amendments filed with the present divisional application add subject-matter which extends beyond the content of the earlier application EP04760595, contrary to the requirements of Article 76(1) EPC. No basis was found in the earlier application EP04760595 for the subject-matter of claims 15, 16, 19, 24 and 26.

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 05 02 5978

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-01-2007

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2004010959 A2	05-02-2004	AU 2003259687 A1	16-02-2004
		CA 2493122 A1	05-02-2004
		EP 1542536 A2	22-06-2005
		JP 2005538742 T	22-12-2005
		KR 20050027132 A	17-03-2005
		MX PA05000972 A	12-09-2005
		US 2006104963 A1	18-05-2006
		US 2004197902 A1	07-10-2004
		US 2005129665 A1	16-06-2005
		WO 2004098499 A2	18-11-2004
		WO 2006031467 A2	23-03-2006
-----			

专利名称(译)	转染的造血干细胞和治疗新生血管性眼病的方法		
公开(公告)号	<a href="#">EP1754783A1</a>	公开(公告)日	2007-02-21
申请号	EP2005025978	申请日	2004-04-28
[标]申请(专利权)人(译)	斯克里普斯研究学院		
申请(专利权)人(译)	斯克里普斯研究所		
当前申请(专利权)人(译)	斯克里普斯研究所		
[标]发明人	FRIEDLANDER MARTIN OTANI ATSUSHI DASILVA KAREN HANEKAMP STACEY		
发明人	FRIEDLANDER, MARTIN OTANI, ATSUSHI DASILVA, KAREN HANEKAMP, STACEY		
IPC分类号	C12N5/06 C12N5/078 A61K35/12 A61K35/28 A61K48/00 A61P9/10 A61P27/02 A61P27/12 C12N5/071 C12N5/074 C12N5/0789 C12N5/10 C12N15/09 G01N33/48 G01N33/53		
CPC分类号	A61K35/12 A61K48/00 A61K2035/124 A61P7/02 A61P7/06 A61P9/10 A61P9/14 A61P25/00 A61P27/02 A61P27/12 A61P41/00 A61P43/00 C12N5/0647 C12N5/0692 C12N5/10		
优先权	60/467051 2003-05-02 US 10/628783 2003-07-25 US		
外部链接	<a href="#">Espacenet</a>		

### 摘要(译)

已经用治疗上有用的基因转染分离的，哺乳动物的，成体骨髓衍生的，含有内皮祖细胞 (EPC) 的谱系阴性造血干细胞群 (Lin-HSC)。本发明的转染的分离的干细胞群可用于将基因递送至眼睛以进行基于细胞的基因治疗。还描述了制备本发明的转染的分离的干细胞群的方法，以及治疗眼病和损伤的方法。

SEQUENCE LISTING

<110> The Scripps Research Institute  
<120> TRANSFECTED HEMATOPOIETIC STEM CELLS AND METHODS OF TREATMENT OF NEOVASCULAR EYE DISEASES THEREWITH  
<130> 18-197 T1  
<140> Divisional application of EP 04 760 595.1  
<141> 2004-04-28  
<150> US 60/467,051  
<151> 2003-05-02  
<150> US 10/628,783  
<151> 2003-07-25  
<160> 5  
<170> FastSEQ for Windows Version 4.0  
<210> 1742  
<211> DNA  
<212> Artificial sequence  
<220> DNA encoding His-tagged human T2-TrpRS  
<400> 1  
TGGCGAATGG AAGCCGCTCT GTAACGGGCG ATTAGCGCG GCGGGTGTGG TGGTACTCG 60  
CAGCGTGACC GCTACACTTG CAGCGCCCTC AGCGCCGCTC CCTTCGCTT TCTTCCCTC 120  
CTTCTGCC ACGTTCGCG GGTCTTCCCA TCGAGCTCA ATCGGGGCG TCCCTTAGG 180  
GTTCCGATTT AGTCTTTC AGGACCTCGA CCCCAGAAA CTGATTTAGG GTGATGTTT 240  
ACGTAATGGG GCACTGCCCT GATAGACGTC TTTCCGCTT TGGCGTTG AGTCCACTT 300  
CTTAATAGT GGACTCTTGT TCCAACTGG AACAACTCT AACCCATCT CCGTCACTC 360  
CTCGATTA CAGGAGATTT TCCGATTTT GGCCTATGG GGCCTATGG TAAAAAATG AGTCACTT 420  
ACAAAAATTT AAGCGAATT TCAACAAAT ATTAACGTT ACAATTCAG GTGGCACTT 480  
TGGGAAAT AGCGGAA CCGCTATTTG TTATTTTC AAAAAAT CAATAGTA 540  
TCCGCTCTG GGCATATC CTGTAATA GTTCAAAA AATLAAAA GAAAGATA 600  
GAGATTCAG CATTCCTGT TCGCCTTAT TCCCTTTT GCGGATTT GCTTCTCT 660  
TTTGTCTC CCAAGATG TGTGAAAT AAAAAATG GAGATCTGT TGGTCTCT 720  
AGTGGTTC ATCAACTGG ATCTCAACG CGTAAAGTC CTTGAGGTT TCTGCCCG 780  
AGAACTTT CCAATGATA GCACTTTAA AGTACTCA TGTGCGTA TATTTCCG 840  
TATTGACGC GGCAGAGC AACTCGGTC CCGATACAC TATTTCTAGA ATGACTTGT 900  
GAGGACTCA CAGTCCAG AAAAGACTT TCGGCAAC TATTTCTGA CAAAGATG 960  
CAGTCTGC ATAACATGA GTGATAAC TCGGCAAC TATTTCTGA CAAAGATG 1020  
AGGCGAAG GCGTAACT CTATTTC AACAATGGG GATCATGTA CTTCTCTG 1080  
TCTTGGAA CCGAGTTPA AAGAACT ATAACTGAG GAGTTTGA CCAAGTCT 1140  
TGCAGCAAT GCAACAAGT TCGCAACT ATAACTGG GAACTACTA CTCCTGCT 1200  
GGCCTTCC GCTGCTGT TATTGCTGA TAACTCGA GCGGTGAG CAGGACAC TCTGCTC 1260  
CGATCATC CAGCACTGG GCGCAATGG TAACTCTC GATCCCTC CATTCTAG 1320  
GACGGGAGT CAGCAACTA TGGATGAG AATAGACG ATCGTGAAG TAGGTGCTC 1380  
AAACTTCA TTTCACTA AAAGGATCTA GGTGAGATC CTTTGTGA ATCCATGA 1440  
GAAATCTT TAAATGAT TTTTCTCA CTAAGTCA GACCCCGTAG AAAGATGA 1500  
ACCGTACCA GGGTGGTCTT GTTTGCGGA TCAAGAGCTA CCAACTCTT TCCGAAAT 1560  
AGATCTCT TGAATCTT TTTTCTCA CTAAGTCA TGTGTAGC CAGGATAG 1620  
CCACCTCT AAGACTCTG TGGCCTCC TACAACCT GCTCTGTA TCTGTATC 1680  
AGTGGTCT GCGGATCG ATAGTCTG TCTATCGT TGTGACTCA GCAATGAT 1740  
ACGGAAGG GGCAGCGT CGGGCTGAC GCGGGGCTC TGCACACG CAGCTTGA 1800  
GCGAGCAC TACACGAC TGAATACC ACGGATCG CATAGAAA GGCACACT 1860