



(11) **EP 1 393 078 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:  
**08.07.2009 Bulletin 2009/28**

(21) Application number: **02743017.2**

(22) Date of filing: **15.05.2002**

(51) Int Cl.:  
**G01N 33/68 (2006.01)**

(86) International application number:  
**PCT/EP2002/005350**

(87) International publication number:  
**WO 2002/093175 (21.11.2002 Gazette 2002/47)**

(54) **METHODS FOR THE IN VITRO DIAGNOSTIC OR FOR THE MONITORING OF A DISEASE INVOLVING AN INFLAMMATORY REACTION**

VERFAHREN FÜR DIE IN-VITRO-DIAGNOSTIK ODER FÜR DIE ÜBERWACHUNG EINER KRANKHEIT MIT ENTZÜNDUNGSREAKTION

PROCEDES POUR LE DIAGNOSTIC IN VITRO OU POUR LA SURVEILLANCE D'UNE MALADIE IMPLIQUANT UNE REACTION INFLAMMATOIRE

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR**

(30) Priority: **16.05.2001 US 290973 P**

(43) Date of publication of application:  
**03.03.2004 Bulletin 2004/10**

(73) Proprietor: **Forceville, Xavier**  
**77860 Saint Germain sur Morin (FR)**

(72) Inventor: **Forceville, Xavier**  
**77860 Saint Germain sur Morin (FR)**

(74) Representative: **Catherine, Alain et al**  
**Cabinet Harlé & Phélip**  
**7 rue de Madrid**  
**75008 Paris (FR)**

(56) References cited:  
• **PERSSON-MOSCHOS M ET AL:**  
**"SELENOPROTEIN P IN SERUM AS A BIOCHEMICAL MARKER OF SELENIUM STATUS" ANALYST, LONDON, GB, vol. 120, no. 3, March 1995 (1995-03), pages 833-836, XP002924559**

- **NICHOL COLIN ET AL:** "Changes in the concentrations of plasma selenium and selenoproteins after minor elective surgery: Further evidence for a negative acute phase response?" **CLINICAL CHEMISTRY**, vol. 44, no. 8 PART 1, August 1998 (1998-08), pages 1764-1766, XP001084140 ISSN: 0009-9147
- **P. GALLOWAY ET AL.:** "Effect of the inflammatory response of trace element and vitamin status" **ANN. CLIN. BIOCHEM.**, [Online] 2000, pages 289-297, XP002206057 Retrieved from the Internet: <URL:www.leeds.ac.uk/acb/annals/annals\_pdf /May00/ACB289.PDF> [retrieved on 2002-07-04]
- **GÄRTNER R ET AL:** "[Significance of selenium in intensive care medicine. Clinical studies of patients with SIRS/sepsis syndrome]" **MEDIZINISCHE KLINIK (MUNICH, GERMANY: 1983) GERMANY 15 OCT 1999**, vol. 94 Suppl 3, 15 October 1999 (1999-10-15), pages 54-57, XP001088107 ISSN: 0723-5003
- **FORCEVILLE X ET AL:** "Selenium, systemic immune response syndrome, sepsis, and outcome in critically ill patients." **CRITICAL CARE MEDICINE. UNITED STATES SEP 1998**, vol. 26, no. 9, September 1998 (1998-09), pages 1536-1544, XP001088109 ISSN: 0090-3493
- **Forceville, X. et al (2000) Intensive Care Medicine 26 (Supplement issue) S228**
- **Mostert, V. et al (1998) Arch. Biochem. Biophys. 357:326-330**
- **Harrison, I. et al (1996) Analyst 121:189-194**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 1 393 078 B1**

**Description****FIELD OF THE INVENTION**

5 **[0001]** The present invention relates to the field of the diagnostic and of the monitoring of several diseases involving an inflammatory reaction, especially those which are associated with damages to the vascular endothelium in human and animals.

**Background of the invention**

10 **[0002]** Severe sepsis and more particularly at a further stage, septic shock, leading to multiple organ failure are major medical problems in critically ill patients and are the most common cause of death in medical and surgical intensive care unit (ICU) (Definition and epidemiology This LG, Dhainaut JF ; in Septic Shock, Dhainaut JF, This LG, G Parked. ;  
 15 **Sunders 2000**). Septic shock is a public health problem because of its frequency (500 000 to 700 000 new cases per year in the United States), and of its mortality rate (around 45%). The systemic repercussion of septic shock is considered as the result of an uncontrolled and generalized immuno-inflammatory reaction. In the case of septic shock, that reaction results from a systemic host response to invasive infection. (From Celsus to Galen to Bone: The Illnesses, Syndromes, and Diseases of Acute Inflammation, Marshall JC, Aarts MA, in Yearbook of Intensive Care and Emergency Medicine, Vincent JL ed.; Springer Verlag, 2001; 3-12). This reaction leads to organ dysfunction and, at a further stage, organ failure.

20 **[0003]** The physiopathology of septic shock could be considered as a consequence of an overproduction of free radicals (Novelli, G. P., Role of free radicals in septic shock, J Physiol Pharmacol; 1997,48 (4): 517-527) (Zimmerman, J. J., Defining the role of oxyradicals in the pathogenesis of sepsis, Critical Care Medicine; 1995 , 23, (4): 616-617) This does not contradict the usual diagrams concerning septic shock but points to free radicals as being directly responsible for cellular damages. The main origin of this deleterious overproduction of free radicals would be the activated polynuclear  
 25 neutrophils adhering to the venous postcapillary segment (Bast, A.; Haenen, G. R.; Doelman, C. J., Oxidants and antioxidants: state of the art, Am J Med; 1991, 91, (3C): 2S-13S). In septic shock, nitric oxide is known to have a marked role. It appears early in septic shock. Its deleterious effects on the organism seems to be essentially mediated by peroxynitrite (ONOO-), formed during the simultaneous synthesis of NO by endothelium cells and superoxide anion by phagocytic cells, such as polymorphonuclears. Moreover, the endothelium is a particular target for peroxynitrite, Selenium may be an efficient treatment of septic shock in particular for increasing selenoprotein P concentration (Selenium and the "free" electron" -Selenium a trace to be followed in septic or inflammatory ICU patients, Forceville X, Intensive Care  
 30 Medicine; 2001, 27: 16-18).

35 **[0004]** Forceville et al. (2000, Intensive Care Medicine, Vol, 26 : 5228) have observed a decrease in selenoprotein P plasma concentration in patients undergoing a septic shock. Further, Forceville et al. (1998, Critical Care Medicine, Vol. 26: 1536-1544) have shown a decrease in plasma selenium concentrations in ICU patients affected with a severe inflammatory response syndrome (SIRS).

40 **[0005]** The early diagnosis and treatment of septic shock have to be conducted in a state of emergency. In man, at the present time, in the only phase III positive therapeutical trial in the treatment of the inflammatory reaction of septic shock, patients had to begin treatment within the first 24 hours of septic shock (Efficacy and Safety of Recombinant Human Activated Protein C for Severe Sepsis, Bernard G.R. , Vincent JL, Laterre PF, New England Journal of Medicine; 2001). In fact, microcirculation endothelium is rapidly damaged during septic shock. The massive alterations of endothelium have a major physiopathological role in the consequences of septic shock and lead to organ failure (Sepsis/septic shock: Participation of the microcirculation: An abbreviated review, Hinshaw LB Critical Care Medicine 1996, 24 (6), 1072-1078).

45 **[0006]** There is a need in the art to have a relatively simple and reproducible biological marker to precisely identify septic shock patients in order to have reproducible epidemiological data, to define appropriate populations of patients to be included in sepsis trials, to identify populations of patients who might benefit from those therapies, and to monitor the evolution of septic shock under those therapies. Moreover that marker should ideally target the, or a, biological process responsible for the manifestation of septic shock. "The search for such (a) marker(s) is an important priority for  
 50 ongoing investigations" (From Celsus to Galen to Bone: The Illnesses, Syndromes, and Diseases of Acute Inflammation, Marshall JC, Aarts MA, in Yearbook of Intensive Care and Emergency Medicine, Vincent JL ed.; Springer Verlag, 2001; 3-12).

55 **[0007]** However, the diagnostic of sepsis and septic shock remains presently complex and depends on definitions that are based on clinical manifestations (Groeneveld, This, in Dhainaut 2000, Pg. 355). They characterize a subgroup of patients with systemic inflammatory response syndrome not only due to infection but also to diseases such as pancreatitis, extended burns, and polytrauma. For further details, see the American College of Chest Physicians and the Society of Critical Care Medicine, conference consensus, Chicago 1991 (The ACCP-SCCM consensus conference on sepsis and organ failure, Chest 1992; 101:1644-55), and the article by Marshall and Aarts (From Celsus to Galen to

Bone: The Illnesses, Syndromes, and Diseases of Acute Inflammation, Marshall JC, Aarts MA, in Yearbook of Intensive Care and Emergency Medicine, Vincent JL ed.; Springer Verlag, 2001; 3-12). Sepsis, severe sepsis and at a further stage septic shock may be due to infection associated or not with detected bacteremia due to bacteria, but also fungi or virus.

**[0008]** Septic shock is defined as a sepsis-induced hypoperfusion (i.e. systolic blood pressure < 90 mmHg or a reduction of  $\geq 40$  mmHg from baseline) despite adequate fluid resuscitation along with the presence of perfusion abnormality that may include -but are not limited to- lactic acidosis, oliguria or an acute alteration in mental state. Patients who are receiving inotropic or vasopressor agents may not be hypotensive at the time that perfusion abnormalities are measured (Bone, 1992, ref. 2925) (Definition and epidemiology This LG, Dhainaut JF ; in Septic Shock, Dhainaut JF, This LG, G Park ed. ; Sunders 2000). This clinical definition established in the ACCP-SCCM consensus conference on sepsis and organ failure, Chicago, August 1991 remains the only valid one (The ACCP-SCCM consensus conference on sepsis and organ failure, Chest 1992; 101:1644-55).

**[0009]** Different markers of septic shock, and especially of its initial manifestation, have been proposed, but are non-specific and insensitive (Diagnosis : from clinical signs to haemodynamic evaluation, Groeneveld AB J, This GT ; in Septic Shock, Dhainaut JF, This LG, G Park ed. ; Sunders 2000). As activation of inflammatory pathways, including the cytokine network, is considered to play a major role in the pathogenesis of septic shock, cytokines have been extensively studied as a potential marker of septic shock (Cytokines and Anticytokines in the Pathogenesis of sepsis, van der Poll T, van Deventer S JH, infectious disease Clinics of North America; 13 (2) 1999; 413-426). In comparison with other cytokines, IL-6 (a mixed pro- and anti-inflammatory cytokine) has been reported most consistently in the circulation of septic patients. However, IL-6 level shows considerable variation, and it is not possible to define a cut-off value to identify septic shock patients (Cytokines and Anticytokines in the Pathogenesis of sepsis, van der Poll T, van Deventer S JH, infectious disease Clinics of North America; 13 (2) 1999; 413-426) Procalcitonin (PCT), C-reactive protein (CRP), leucocytosis, lactate concentration, coagulation parameters, and other parameters have also been proposed to identify septic shock patients (Diagnosis of sepsis: Novel and Conventional Parameters, Rheinhardt K, Meisner M and Hartog C, Advances in sepsis, 2001; 42-51). However, though they seem to be useful parameters to improve the diagnosis and monitoring of septic shock, they cannot clearly identify septic shock patients due to their lack of specificity and the diagnosis of septic shock remains based on clinical parameters. (From Celsus to Galen to Bone: The Illnesses, Syndromes, and Diseases of Acute Inflammation, Marshall JC, Aarts MA, in Yearbook of Intensive Care and Emergency Medicine, Vincent JL ed.; Springer Verlag, 2001; 3-12). Presently, laboratory abnormalities can only be supportive for the diagnosis of sepsis and cannot prove the presence of septic shock. Taken together, abnormalities in hematological and chemical laboratory parameters may only constitute indicators of septic shock (Diagnosis : from clinical signs to haemodynamic evaluation, Groeneveld AB J, This GT ; in Septic Shock, Dhainaut JF, This LG, G Park ed. ; Sunders 2000). Each alone of this parameters carries little meaning. (Defining a Clinical Syndrome of Systemic inflammation, Vincent JL, Byl B, Sepsis; 200; 4:15-19).

**[0010]** There is also an urgent need in the art to find a specific and reliable parameter to define septic shock patients. A precise definition of septic shock patients could allow an early diagnosis and a follow-up of septic shock. This parameter should ideally be closely related to the physiopathological process of septic shock (From Celsus to Galen to Bone: The Illnesses, Syndromes, and Diseases of Acute Inflammation, Marshall JC, Aarts MA, in Yearbook of Intensive Care and Emergency Medicine, Vincent JL ed.; Springer Verlag, 2001; 3-12).

#### **SUMMARY OF THE INVENTION**

**[0011]** The present invention is based on the observation that selenoprotein P can be used as a biochemical marker of septic shock, both in man and in animals, and more generally as a biochemical marker of the severity of the systemic inflammation response syndrome (SIRS).

**[0012]** Thus, a first object of the present invention consists of a method for the in vitro diagnostic or for the monitoring of a disease involving an inflammatory reaction within a patient as defined in claim 1.

**[0013]** By « a disease involving an inflammatory reaction », it is intended herein a disease wherein the inflammatory reaction is associated with an activation of the peripheral blood phagocytic cells, namely monocytes/macrophages and most importantly polymorphonuclear cells and more specifically neutrophils. Most preferably, a « disease involving an inflammatory reaction » as defined above is also associated with a damage to the vascular endothelial cells.

**[0014]** By « amount » of selenoprotein P, it is intended herein any parameter or unit of measure which allows the quantification of selenoprotein P in a sample. An amount of selenoprotein P may be expressed as an absolute quantity in a sample (e.g. number of moles of selenoprotein P) or as a concentration (e.g. nmol/L of selenoprotein P in the sample).

**[0015]** The damage to the vascular endothelial cells may be present at the onset of the disease or alternatively is a consequence of the onset of the disease and thus consists of a secondary damage to the vascular endothelial cells.

**[0016]** In a first preferred embodiment, the disease consists of a systemic inflammation response syndrome (SIRS), and most preferably of a septic shock.

**[0017]** In a second preferred embodiment, the disease consists of a chronic inflammatory disease. Said chronic inflammatory disease may be selected from the group consisting of a systemic disease such as polyarteritis nodosa. The chronic inflammatory disease may also be a disease related to a bacterial, a fungal or a viral infection or may be an eclampsia, therefore the follow up of selenoprotein P during pregnancy may be done during routine pregnancy follow up.

**[0018]** In a further preferred embodiment, the occurrence or the aggravation of the disease is diagnosed in a further step d) when the amount of Selenoprotein P measured in the biological sample is lower than the normal amount which is contained within non-affected individuals or when the amount of selenoprotein P measured in the biological sample is lower than that measured in a sample from the same patient in a former test.

**[0019]** In another preferred embodiment, the persistence of the disease is diagnosed in a further step d) when the amount of Selenoprotein P measured in the biological sample is lower than the normal amount which is contained within non-affected individuals or when the amount of selenoprotein P measured in the biological sample is lower than or not higher than, or equal to, the amount measured in a sample from the same patient in a former test.

**[0020]** In a still further preferred embodiment of the method disclosed above, the recovering from the disease is diagnosed in a further step d) when the amount of selenoprotein P measured in the biological sample is at least equal to the normal amount which is contained within non-affected individuals or when the amount of selenoprotein P measured in the biological sample is higher to that measured in a sample from the same patient in a former test.

**[0021]** In a further preferred embodiment of the method above, mortality is predicted when the amount of selenoprotein P which is measured within a serial of biological samples collected during the time period of the therapeutical treatment is always lower than 20nmol/L, preferably lower than 10 nmol/L.

**[0022]** It is hereby also described a kit for the in vitro diagnostic or for the monitoring of a disease involving an inflammatory reaction associated with a damage of the vascular endothelial cells within a patient, wherein said kit comprises (i) means for assaying the amount of Selenoprotein P contained in a biological sample.

**[0023]** In a first preferred instance, the kit further comprises (ii) a sample or a plurality of samples containing each respectively a predetermined amount of Selenoprotein P to be used as a standard reference.

**[0024]** In a second preferred instance of the kit, the means for assaying the amount of Selenoprotein P contained in the biological sample consist of antibodies directed against Selenoprotein P.

#### **DETAILED DESCRIPTION OF THE INVENTION**

**[0025]** It has now been found according to the invention that a marked decrease in the blood concentration of selenoprotein P is measured in patients suffering from diseases that involve an inflammatory reaction.

**[0026]** The inventor has now shown that a marked decrease in the blood concentration of selenoprotein P occur in patients affected with severe sepsis, particularly in patients which are affected with a systemic inflammation response syndrome (SIRS) and that a more further decrease in the blood selenoprotein P level is measured in patients which are undergoing a septic shock.

**[0027]** Selenoprotein P has been known since 1971. It is the only selenoprotein to contain more than one selenium atom per polypeptide chain in the form of 10 selenocysteine residues per polypeptide (Orphan selenoproteins Burk RF, Hill KE BioEssays 21 (3): 231-237, 1999) (Selenoprotein P: Properties, Functions, and Regulation Mostert V, Archives of Biochemistry and Biophysics; 376 (2): 433-438, 2000). Selenoproteins are defined as proteins containing selenocysteine. Selenocysteine is known to be the 21st amino acid by the replacement of sulfur to selenium in the cysteine. The comparison of deduced amino acid sequences of selenoprotein P from different species indicates almost complete conservation of cysteine and selenocysteine residues when these amino acids are considered together. Selenoprotein P is mainly synthesized in the liver. There are at least two distinct isoforms of selenoprotein P in human plasma (A Novel Method for the Purification of Selenoprotein P from Human Plasma, Mostert V, Lombeck I, Abel J, Archives of Biochemistry and Biophysics; 357 (2): 326-330, 1998).

**[0028]** Selenoprotein P is present in large amounts of the surface of the endothelial cells. Through its heparin-binding properties it could bind to the inflamed endothelium (Burk, R. F.; Hill, K. E.; Boeglin, M. E.; Ebner, F. F.; Chittum, H. S., Selenoprotein P associates with endothelial cells in rat tissues, Histochem Cell Biol ; 1997, 108 (1), 11-15).

**[0029]** Because of the high stability of the selenoprotein P plasma concentration in healthy individuals as well as in individuals affected with diverse pathologies which are not involving an inflammatory reaction, selenoprotein P consists of a highly reliable marker both for diagnosing inflammatory diseases and for monitoring the recovering of patients from these inflammatory diseases, for example during the whole time period of treatment and even after the end of the treatment in order to ensure the total recovery of the patient from the disease.

**[0030]** In anorexic, malnutrition, selenium deficiency, in particular in some countries of China, severe alcoholic disease i.e. cirrhosis, dialyzed or apheresis patients as in inflammatory bowel diseases patients, seleno protein P plasma concentration may be expected to be low, reflecting the low selenium status of these patients and perhaps in case of inflammatory diseases the chronic consumption and/or reduced synthesis for severe liver diseases i.e. cirrhosis.

**[0031]** Thus, the present invention firstly relates to a method for the in vitro diagnostic or for the monitoring of a disease

## EP 1 393 078 B1

involving an inflammatory reaction within a patient as defined in claim 1.

**[0032]** By a « biological sample », it is herein intended a blood sample, such as a whole blood sample. Most preferably, the biological sample consists of a blood plasma sample.

**[0033]** According to a first aspect of the method, the disease involving an inflammatory reaction involving an inflammatory reaction consists of a Systemic Inflammation Response Syndrome (SIRS).

**[0034]** According to a second aspect, the disease involving an inflammatory reaction consists of a septic shock.

**[0035]** According to a third aspect, the disease involving an inflammatory reaction associated with a damage of the vascular endothelial cells consists of a chronic inflammatory disease.

**[0036]** According to a fourth aspect, the chronic inflammatory disease is selected from the group consisting of the inflammatory bowel disease or a systemic disease such as polyarteritis nodosa.

**[0037]** In a fifth aspect, the disease involving an inflammatory reaction consists of a chronic inflammatory disease related to a bacterial, a fungal or a viral infection. The diagnostic or the monitoring of the evolution of the inflammatory reactions occurring during a HIV infections is an illustrative example of the use of the diagnostic method disclosed herein.

**[0038]** Most preferably, the method according to the invention is highly useful for diagnosing as well as monitoring septic shock disorders which require a treatment based on a high level of catecholamines such as norepinephrine and epinephrine.

**[0039]** More preferably, the method according to the invention is performed for the diagnosis, the monitoring or for pronostic purposes for the following pathologies involving inflammatory reactions:

a/ Main target diseases (for diagnostic, severity, pronostic)

⇒ septic shock associated or not with bacteremia, funguemia, parasitemia, viremia, associated or not with multiple organ failure,

b/ Associated target diseases (for pronostic and severity, also for diagnostic)- SIRS :

- infections without septic shock ie. with severe sepsis, (ie. peritonitis, pneumonia, meningitidis, catheter related infection, urinay, biliary tract infection, attack of malaria, especially cerebral malaria) non infectious origin : pancreatitis, burns, polytrauma associated of not with fat embolism, massive transfusion, adult respiratory distress syndrom (ARDS), or similar diseases especially in newborn acute renal failure (ARF) related to inflammatory disease, major surgery.

- shock whatever the origin especially with multiple organ dysfunction and severe hypoperfusion : i.e. hemorrhagic, cardiogenic, anaphylactic.

- ischemia-reperfusion :

- crush syndrom, aortic cross clamping, ischemia with reperfusion (ie. mesenteric, lower limb ischemia).

- prolonged status epilepticus, acute rhabdomyolysis (ie. exertionnal rabdomyolysis) malignant hyperthermia.

- eclampsia

- severe status asthmaticus

- tumoral lysis syndrom, exacerbation of systemic diseases. (Acute) hemolytic syndrom

- Metals intoxication i.e. iron (Fe), copper (Cu) ; all intoxication or poisoning involving vascular oxidative stress (i.e. paraquat, diquat) transplantation (especially if associated with graft reaction, second set, graft versus host reaction).

c/ All diseases involving oxidative vascular damages, for the folow-up especially in cases of exacerbation and the monitoring of treatment involving or not selenium compound.

⇒ Chronic infection :

i.e. HIV, endocarditis, (infection related arteritis i.e. syphilis)

⇒ Systemic diseases : (especially those associated with elevated ANCA) :

⇒ i.e necrotizing angeitis, Wegner's syndrome, polyarteritis nodosa, allergic granulomatosis, temporal arteritis, nephroangiosclerosis, Takayasu's disease, Buerger's disease

• Autoimmune diseases : i.e. rheumatoid arthritis, systemic lupus erythematosus (LES), acute rheumatic fever, dermatomyositis, systemic scleroderma thrombotic thrombocytopenic purpura

⇒ hematologic and cancerologic diseases -associated with elevated cytokines (i.e. TNF- $\alpha$ ) or tumoral lysis, or endothelium aggression

• Kahler's disease, Lymphoma especially T cell Lymphoma,  
• tumoral lysis syndrom either spontaneous or induced by anti carcinologic treatment in hematologic or tumoral diseases (i.e. acute leukemia, solid tumor especially with elevated cytokines) .

• Immun complex diseases (i.e. Berger's disease)

⇒ Atherosclerosis (especially for follow-up and the monitoring of treatment prevention or for acute exacerbation, including or not selenium)

Post-operative vascular restenosis following vascular angioplasty Diabetes (diabetic angiitis)

⇒ Pregnancy follow-up, monitoring of treatment (especially for the early detection of preeclampsia).

⇒ Severe liver diseases, inflammatory bowel disease.

**[0040]** In general, the method of the invention enables the one skilled in the art to diagnostic and monitor patients with the inflammatory diseases selected from the group of inflammatory diseases which are defined in the ACCD-ISCM Conference consensus (from Celsus to Galen to Bone: The illnesses syndromes and diseases of acute inflammation, Marshall JC, Aarts MA, in Yearbook of Intensive Care and Emergency Medicine, Vincent JL ed.; Springer Verlag, 2001; 3-12).

**[0041]** In the general population, selenoprotein P plasma concentration is stable, though it could be somewhat affected by the selenium nutritional status of the population. In a survey of 414 people from various European countries, selenoprotein P level varies from 100% for Spain, to 76 % for Greece (Variation in selenoprotein P concentration in serum from different European regions, Marchaluk E, Persson-Moschos M, Thorling EB, Akesson B, Eur J Clin Nutr, 1995; 49 (1): 42-48). Selenoprotein P appears to be an index of selenium nutritional status especially in the case of low selenium nutritional status. Indeed, in the Finland studies, a baseline selenium intake of about 100  $\mu$ g/day abolishes the increase of selenoprotein P under oral supplements of 200  $\mu$ g/day selenium, contrary to what was observed in the case of a baseline selenium intake of about 40  $\mu$ g/day (Plasma selenoprotein P levels of healthy males in different selenium status after oral supplementation with different form of selenium, Persson-Moschos M, Alftan G, Akesson B, Eur J Clin Nutr; 1998, 52 (5): 363-367). Also, In the United States selenium nutritional status is known to be higher than in Europe (Rayman, M. P., The importance of selenium to human health, Lancet; 2000, 356: 233-41). Compared to other selenoprotein, such as glutathion peroxidase, selenoprotein P concentration is better preserved during the decrease of selenium intake (Selenoprotein P in serum as a biochemical marker of selenium status, Persson-Moschos M, Huang W, Srikumar TS, Akesson B, Lindeberg S, Analyst, 1995; 120 (3): 833-836).

**[0042]** As it is detailed hereabove, although the selenoprotein P plasma concentration is highly stable, some differences in the normal selenoprotein P plasma concentration were observed when individuals from various geographical origins were compared (see Marchaluk et al., 1995, Persson-Moschos et al., 1998; Rayman et al., 2000; *OP. Cit.*).

**[0043]** The discrepancies in selenoprotein P concentration which are observed in individuals belonging to distinct countries of origin do not require an adaptation of the method described above depending of the country of birth or of the genealogy of the patient tested, since these discrepancies do not significantly affect the statistical significance of the marked decrease in the selenoprotein P concentration, except in countries with very low selenium intake i.e. part of China, Finland without supplementation, New Zeland.

**[0044]** Nevertheless, in a most preferred embodiment of the method described wherein, the amount of selenoprotein P which is measured at step a) is compared, at step b), with the amount of selenoprotein P which would be expected within a group of non-affected individuals from the same geographical origin than that of patient tested.

**[0045]** Consequently, when the patient originates from Europe, United States of America, Australia, or an asian country, the amount of selenoprotein P which is measured at step a) will be preferably compared, at step b) of the method, with the mean amount of selenoprotein P which is contained in the same biological sample of a group of non-infected individuals originating respectively from Europe, United States of America, Australia or an asian country.

[0046] In a further most preferred embodiment of the method, the country of origin may be taken into account, such as France, Spain or Greece, for example.

[0047] In a specific embodiment of the invention, the method described above allows the diagnosis of the occurrence of the disease in a further step c), when the amount of selenoprotein P which is measured in the biological sample of the patient tested is statistically lower than the normal amount which is contained within non-affected individuals.

[0048] In a further particular embodiment of the invention, the method described above allows to diagnose the persistence of the disease in a further step c), when the amount of selenoprotein P which is measured in the biological sample of the patient tested is lower than the normal amount which is contained within non-affected individuals, or alternatively when the amount of selenoprotein P is lower or not significantly higher than the amount found for the same patient in a former test.

[0049] In still a further embodiment of the invention, the method allows the diagnosis of the recovering of the patient from the disease in a further step c), when the amount of selenoprotein P which is measured in the biological sample of the patient tested is at least equal to the normal amount which is contained within the non-affected individuals, or alternatively when the amount of selenoprotein P is higher than the amount found for the same patient in a former test.

[0050] By « at least identical » as used herein, is intended an amount of selenoprotein P which is of a value which is at least not statistically distinct from the mean amount of selenoprotein P which is contained in the biological sample which is compared.

[0051] Without wishing to be bound by any particular theory, the inventor believes that the treatment of patients affected with a disease involving an inflammatory reaction associated with a damage to the vascular endothelial cells, particularly patients affected from SIRS or a septic shock, may need to obtain selenoprotein P plasma concentration values which are higher than those found in non-affected individuals. This situation may be the consequence of the therapeutical treatment based mainly on the administration of selenium. A supra-normal value of selenoprotein P plasma concentration may indicate a patient status wherein the endothelium is protected against endothelium oxydative aggression, whether it be acute or chronic.

[0052] Thus, in another particular embodiment of the invention, the method disclosed herein allows to diagnose the recovering from the disease in a further c), when the amount of selenoprotein P which is measured in the biological sample of the patient tested is higher than the normal amount which is contained within the non-affected individuals.

[0053] Preferably, the standard level of selenoprotein P plasma concentration which is contained in a biological sample from an individual which is not affected with the disease involving an inflammatory reaction associated with a damage to the vascular endothelial cells, at step b) of the method, is of at least 50 nmol/L. Most preferably, this selenoprotein P plasma concentration value is comprised between 40 and 80 nmol/L, advantageously between 40 and 70 nmol/L and most preferably between 40 and 60 nmol/L.

[0054] Most preferably, the selenoprotein P plasma concentration is indicative of a disease involving an inflammatory reaction associated with a damage to the vascular endothelial cells when the amount of selenoprotein P which is measured within the biological sample of the patient tested is lower than 30 nmol/L.

[0055] In a particular embodiment, the diagnostic of a selenoprotein P plasma concentration of less than 20 nmol/L is indicative of patients with septic shock requiring high level of catecholamines such as norepinephrine and epinephrine.

[0056] In a further embodiment, the diagnostic of a selenoprotein P plasma concentration of less than 15 nmol/L and most preferably of less than 10 nmol/L is predictive of mortality, especially when there is a persistence of a weak selenoprotein P amount during a selenium-based treatment of the patient.

[0057] However, at the stage of patients recovering from the disease, the inventor has observed supranormal values of selenoprotein P plasma concentration up to 106 nmol/L in recovering septic shock patients who survived, and supra normal values up to 181 nmol/L in non SIRS agressed patients.

[0058] In a still further embodiment, the methods of the invention allows to predict mortality when the amount of selenoprotein P which is measured within a serial of biological samples collected during the time period of the therapeutical treatment is always lower than 20nmol/L, preferably lower than 10 nmol/L.

[0059] Without wishing to be bound by any particular theory, the inventor believes that the increase of selenoprotein P may be an interesting defense mechanism of the endothelium in situation of aggression. This reinforces the interest of the monitoring of selenoprotein P plasma concentration in all situations of generalized blood inflammation associated with immediate or secondary endothelium oxydative damages, whether or not it is associated with a selenium-based treatment or supplementation.

[0060] According to a specific embodiment of the method disclosed herein, the comparison step c) might take into account previously known chronic deficiencies in selenoprotein P of the tested patient, in order to avoid interpreting a weak decrease of selenoprotein P plasma concentration as regards the selenoprotein P standard concentration as indicative of an inflammatory disease.

[0061] As it is disclosed herein, the method of the invention allows the use of a novel biochemical marker, namely plasma concentration of selenoprotein P, in order to diagnose septic shock, and even the severity of the septic shock in man and in animal.

[0062] Moreover, the follow up of the selenoprotein P plasma concentration in a patient permits the monitoring of the evolution of the disease, for example septic shock. A persistent low level of selenoprotein P plasma concentration less than 20 nmol/L or 30 nmol/L, especially if abundant selenium administration is provided during the therapeutical treatment, will correspond to the persistence of the systemic inflammation which was diagnosed according to the method of the invention.

[0063] Indeed, the method of the invention, by following up the plasma concentration of selenoprotein P, is very useful to monitor the benefit to the patient of a selenium-based treatment.

[0064] Further, the persistence of a low level of selenoprotein P consists of a marker of the gravity and the mortality of the disease in a septic shock patients.

[0065] In a most preferred embodiment, the measure of the amount of selenoprotein P which is contained in the biological sample of the patient tested, in step a) of the disclosed method, consists of an immuno assay wherein antibodies directed against selenoprotein P are used.

[0066] For performing such an immunoassay, the one skilled in the art will advantageously refers to the article of Mostert et al. (1998, Archives of Biochemistry and Biophysics, vol. 357 (2): 326-330).

[0067] When performing the method of the invention for monitoring purposes, said method will be performed at least daily during the period of time starting from the admission of the patients in an emergency care unit of an hospital. After this critical period, the method of the invention may be performed every two-days time period during the therapeutical treatment in order to monitor the recovering of the patients from the disease.

[0068] A daily dosage of selenoprotein P is preferred for acute diseases, especially if selenium-based treatment - e.g. by selenoprotein P - in order to achieve the optimal selenoprotein P plasma concentration adapted to the disease and its severity. Then, a dosage every two days periods or a weekly dosage is advantageous during the recovering period and then monthly after recovering of the normal selenoprotein P plasma concentration.

[0069] For chronic diseases in the absence of an acute disorder, or in cases of pregnancy, selenoprotein P dosages may be part of the routine biological examination.

[0070] Monitoring of selenium-based treatment can be achieved with selenoprotein P plasma concentration monitoring in order to obtain optimal selenoprotein P concentration according to the gravity of the disease and the importance of oxidative stress in the blood compartment.

[0071] In cases of septic shock with high level of oxidative stress in the blood compartment, it could be appropriate to obtain and maintain very high level of selenoprotein P much higher than the reference values.

[0072] It is hereby also described a kit for the *in vitro* diagnostic or for the monitoring of a disease involving an inflammatory reaction within a patient, wherein said kit comprises (i) means for assaying the amount of selenoprotein P contained in a biological sample.

[0073] In a preferred instance, said kit further comprises (ii) a sample or a plurality of samples containing each respectively a predetermined amount of selenoprotein P to be used as a standard reference.

[0074] Most preferably, said kit comprises one standard sample containing selenoprotein P at a determined concentration, said concentration being comprised between 50 and 80 nmol/L, and most preferably about 50 nmol/L selenoprotein P.

[0075] In the instance wherein said kit comprises a plurality of standard samples containing selenoprotein P, the respective concentration are preferably 200, 100, 50, 30, 20 and 10 nmol/L selenoprotein P which are used to generate a standard curve allowing a precised determination of the selenoprotein P plasma concentration within the biological sample of the patient tested.

[0076] In a most preferred instance, the standard samples contain a selenoprotein P in a liquid solution, either a physiological saline buffer (e.g. minimum essential medium - MEM) or a plasma sample containing a predetermined concentration of selenoprotein P.

[0077] Standard samples of selenoprotein P may be contained in lyophilized form in a vial that will be added with the required volume of a suitable buffer at the time of use of the kit.

[0078] In a specific instance of the kit above, the means for assaying the amount of selenoprotein P contained in the biological sample consist of antibodies directed against selenoprotein P. Most preferably, said antibodies are those disclosed by MOSTERT et al. (1998).

[0079] Thus, the kit may also contain a vial containing the antibodies directed against selenoprotein P, either in suspension in a suitable buffer or alternatively under a lyophilized form.

[0080] By the determination of selenoprotein P plasma concentration by immunoblot analysis using purified selenoprotein P as a standard (A Novel Method for the Purification of selenoprotein P from Human Plasma, Mostert V, Lombeck I, Abel J, Archives of Biochemistry and Biophysics; 357 (2): 326-330, 1998), the reference level of selenoprotein P in the general population is about 50 nmol/L and more than 40 nmol/L. Selenoprotein P plasma concentration could be considered as low when it is less than 30 nmol/L and very low when it is less than 10 nmol/L.

[0081] The present invention will be further disclosed by, Without in any case being limited to, the following examples.

**EXAMPLES**

**[0082]** In a preliminary study on 21 ICU patient and 7 healthy volunteers, selenoprotein P plasma concentration has been measured by immunoblot analysis using purified selenoprotein P as a standard (A Novel Method for the Purification of selenoprotein P from Human Plasma, Mostert V, Lombeck I, Abel J, Archives of Biochemistry and Biophysics; 357 (2): 326-330, 1998). In this study, 7 septic shock patients were included among the 21 ICU patients. Dosages have been realized at admission in ICU and later on at weekly intervals during their ICU length of stay. At admission, a very low selenoprotein P plasma concentration was observed in septic shock ICU patients (see Table 1 below) compared to non systemic inflammatory ICU patients. Extremely low level of selenoprotein P plasma concentration, even undetectable, was observed only in septic shock patients during their length of stay. Decrease of selenoprotein P was associated with severe inflammatory adverse events such as severe infections or acute respiratory distress syndrome. Last measurement before death was very low in the 3 septic shock patients who died compared to the value of the 3 defined as non severe SIRS patients who died. These selenoprotein P value was significantly lower than the one observed in the 3 patients who died in the non inflammatory group ( $p < 0.05$ ; Mann-Whitney test). It is to underline that this difference is significative despite the small number of patients involved .

**[0083]** The level of selenoprotein P at admission was poorly correlated ( $r$  about 0.3) with gravity index as measured to the Simplified Acute Physiology Score II (SAPS II) (A new simplifies acute physiology score (SAPS II) based on European/North American multicenter study, Le Gall JR, Lemeshow S, Saulnier F, JAMA 1993; 270: 2957-2963).

**[0084]** This indicates that selenoprotein P doesn't assess the global gravity of the patients, but rather measures septic shock's (and SIRS's) gravity.

**Table 1**

<b>Selenoprotein P Plasma concentration at admission in septic shock ICU patients</b>		
	<b>septic shock patients (all intubated)</b>	<b>Including septic shock patients with grade 4 in the cardiovascular function scale of the SOFA score</b>
number of patients	(7 patients) (measurement at admission on only 6 patients)	(4 patients) (measurement at admission on only 3 patients)
Selenoprotein P	$15 \pm 11$ nmol/L (1 - 28)	$16 \pm 9$ nmol/L (7 - 28)
Mortality	(3 patients)	(3 patients)
SOFA (The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure, Vincent JL, Moreno R, Takala J and al, Intensive Care Medicine 1996, 22:707-710).		

**[0085]** Results are given in mean  $\pm$  standard deviation; followed by minima and maxima in brackets.

**EXAMPLE 2:**

**Selenoprotein P measurement in septic shock patients and comparison with healthy- volunteers and with other ICU patients.**

**[0086]** The selenoprotein P plasma concentration at admission was significantly lower in the septic shock patient group compared to the non severe SIRS (see below for definition, and Table 2) patient group ( $p = 0,0015$  ; Mann-Whitney test ), and healthy volunteers ( $p = 0.003$ ). It is to underline that this difference is significative despite the small number of patients involved, supporting the marked difference between the two groups.

**[0087]** As well, The selenoprotein P plasma concentration at admission was significantly lower in the severe SIRS patient group compared to the non severe SIRS ( $p < 0.05$ ) and to the healthy volunteer group ( $p < 0.01$ ) (see below Table 2).

**[0088]** Non severe SIRS patients were defined as patients without suspected or proven severe infection or septic shock, severe acute pancreatitis, extended burns; without post operative event with a SAPS II of more than 30; and without exposition to major ischemia-reperfusion process such as aortic cross clamping, inferior leg ischemia-reperfusion, mesenteric vascular ischemia.

**[0089]** Mean gravity index score, as defined by the SAPS II value, was lower in the non SIRS patient group than in the severe SIRS patient group : SAPS II  $36 \pm 11$  vs. SAPS II  $47 \pm 14$ . But there were also patients in the non SIRS

group with very high values of SAPS II (6 patients with SAPS II values superior to 35).

**Table 2**

<b>Selenoprotein P plasma concentration at admission in ICU patients</b>			
<b>Groups</b>	<b>Septic shock or severe SIRS patients</b>	<b>Non severe SIRS patients</b>	<b>Healthy volunteers</b>
number of patients	(9 patients) (measurement at admission on only 8 patients)	(12 patients)	(7 volunteers)
Selenoprotein P	17 ± 12 nmol/L (1 - 38)	55 ± 45 nmol/L (20 - 181)	49 ± 8 nmol/L (39 - 65)
(p < 0.01 vs. healthy volunteers) (p < 0.05 vs. non severe SIRS patients) SIRS : Systemic Inflammatory Response Syndrome			

**[0090]** initial dosages of the 7 patients admitted for septic shock were all inferior to 30 nmol/L. In this group, 4 patients required epinephrine or norepinephrine at doses superior to 0.1 µg/kg.min, corresponding to a cardiovascular scale value in the SOFA score of 4, which is the maximum cardiovascular dysfunction. In these 4 patients, 3 had a selenoprotein P value less than 15 nmol/L at their initial measurement. Three of these 4 patients died as a consequence of their septic shock. All of these three patients had had initially, at admission, a plasma selenoprotein P concentrations inferior to 15 nmol/L.

**[0091]** Moreover, selenoprotein P plasma concentration decreased further between admission and the first weekly measurement. Their values were inferior to 10 nmol/L in 3 out of the 7 patients in septic shock, and inferior to 16 nmol/L for 6 out of the 7 septic shock patients. The only septic shock patient who had a persistent selenoprotein P plasma concentration at 28 nmol/L at the first weekly measurement survived.

### **EXAMPLE 3:**

#### **Other SIRS patients**

**[0092]** In the severe SIRS group, one patient had at admission only a severe sepsis, retrospectively diagnosed as *Bordetella pertussis*, associated with severe asthma. Her selenoprotein P plasma concentration at admission was the highest observed in the severe SIRS group : 38 nmol/L. A rapid decrease of the selenoprotein P plasma concentration to a concentration of 7.5 nmol/L was observed, simultaneously as she developed a very severe Adult Respiratory Distress Syndrome (ARDS), requiring 100 % FiO<sub>2</sub> ventilation, with high level of positive end expiratory pressure, nitric oxide and prone-supine positions.

**[0093]** In the severe SIRS group, another patient was a severe polytrauma patient. This 34 year old man had severe thoracic and abdominal injuries requiring emergency surgery. In addition, his both femurs were broken and he had also a broken left humerus. He was initially in shock, and required abundant transfusion. He developed a severe rhabdomyolysis, and a very convincing clinical and biological picture of severe fat embolism, and later, infection was suspected. In this patient, the selenoprotein P plasma concentration was low at admission (selenoprotein P : 10 nmol/L) and remained low during his ICU length of stay. He was discharged from ICU to the orthopedic ward in less than three weeks.

**[0094]** As selenoprotein P is stable in the general population and as this young polytrauma patient was healthy before his accident, such a low initial plasma selenoprotein P concentration indicates that the marked decrease of selenoprotein P observed in the severe SIRS group rapidly appears in the case of a severe SIRS.

### **EXAMPLE 4:**

#### **Non severe SIRS patients**

**[0095]** In the non severe SIRS group, four patients out of 12 had a selenoprotein P plasma concentration between 20 and 30 nmol/L at their initial measurement. One of them belongs to a selenium deficient group (chronic dialysis); and two were misclassified as they had ischemia-reperfusion injuries (one with a status epilepticus well known to have transient lactate acidosis, and the other an inferior leg vascular cross clamping followed by hemorrhage and myocardial infarcts).

[0096] On the opposite, high level of selenoprotein P was observed in the non severe ICU SIRS patients; in this group, 6 patients had a selenoprotein P plasma concentration superior to 60 nmol/L, even in ICU patients with high SAPS II value.

[0097] Indeed, there is little correlation between SAPS II gravity index score, and the selenoprotein P plasma concentration for the all patients (r=0.33). There was also little correlation between SAPS II gravity index score, and the selenoprotein P plasma concentration in the severe SIRS patient group (r = 0,25). One should underlined that fact, as SAPS II gravity index score reflects the probability of death of ICU patients, but not the specific gravity of the SIRS or septic shock (From Celsus to Galen to Bone: The Illnesses, Syndromes, and Diseases of Acute Inflammation, Marshall JC, Aarts MA, in Yearbook of Intensive Care and Emergency Medicine, Vincent JL ed.; Springer Verlag, 2001; 3-12). Mean SAPS II gravity index score was lower in the non severe SIRS patients,  $47 \pm 14$  vs.  $36 \pm 11$  (p =0.055) than in the severe SIRS patients. However, the mortality rate of the two group was close with 3 deceased patients out of 9 in the severe SIRS patient group (all of them septic shock patients) and 3 deceased patients out of 12 in the non severe SIRS patient group.

**EXAMPLE 5:**

**Gravity index, Follow-up**

[0098] One should underlined that the last selenoprotein P plasma concentration observed before death were significantly lower (p < 0.05) in the 3 deceased septic shock patients, compared to the 3 deceased patients of the non SIRS group (see Table 3 below).

[0099] Moreover, the surviving patient with a very low selenoprotein P plasma concentration of 3 nmol/L at his last dosage before discharge from ICU was an anorexic patient, with therefore an initial selenium deficiency, which had made during his ICU length of stay several severe nosocomial infections, including 2 nosocomial peritonitis, following the initial peritonitis. This patient had at his discharge a persistent abundant enteric fistula.

[0100] The mean selenoprotein P plasma concentration of the surviving severe SIRS patients observed before discharge was 23 nmol/L compared to 10 nmol/L for the deceased severe SIRS patients. However, it is to be noticed that selenium supplementation was only 40 µg per day, according to the daily recommended intake. This supplementation is probably very insufficient, in such inflammatory patients. (Selenium and the "free" electron" -Selenium a trace to be followed in septic or inflammatory ICU patients, Forceville X, Intensive Care Medicine; 2001, 27: 16-18). Indeed, as for all selenoprotein, the syntheses of selenoprotein P requires available selenium.

[0101] Selenoprotein P is a good marker for the follow-up of selenium supplementation or treatment in inflammatory ICU patients.

**Table 3**

<b>Selenoprotein P plasma concentration at the latest value before ICU discharge</b>			
<b>Groups</b>	<b>Septic shock or severe SIRS patients</b>	<b>Non severe SIRS patients</b>	
deceased patients	(3 patients) (all septic shock patients)	(3 patients) (non SIRS diseased patients)	
Selenoprotein P	10 nmol/L (0-24)	49 (27-84)	P<0.05
ICU surviving patients	(6 patients)	(9 patients)	
Selenoprotein P	23 nmol/L (3-60)	41 (20-77)	
Results are given in mean ± standard deviation; followed by minima and maxima in brackets SIRS : Systemic Inflammatory Response Syndrome.			

**Claims**

1. A method for the in vitro diagnostic or for the monitoring of a disease involving an inflammatory reaction within a patient, wherein said method comprises the steps of

a) measuring the amount of selenoprotein P which is contained in a biological sample from said patient;

b) comparing the amount of selenoprotein P measured at step a)

i) with the amount of selenoprotein P which is contained in a biological sample from an individual which is not affected with the disease ; or

ii) with the amount of selenoprotein P which was contained in a biological sample from the same patient in a former test; and

c) diagnosing a septic shock when the amount of selenoprotein P which is measured within the biological sample is lower than 30 nmol/L.

2. The method of claim 1, wherein the biological sample consists of a blood sample.
3. The method of claim 1, wherein the biological sample consists of a blood plasma sample.
4. The method of claim 1, wherein said disease involving an inflammatory reaction consists of a Systemic Inflammation Response Syndrome (SIRS).
5. The method of claim 1, wherein said disease involving an inflammatory reaction consists of a septic shock.
6. The method of claim 1, wherein said disease involving an inflammatory reaction consists of a chronic inflammatory disease.
7. The method of claim 6, wherein said chronic inflammatory disease is selected from the group consisting of a systemic disease such as polyarteritis nodosa.
8. The method of claim 6, wherein said disease involving an inflammatory reaction consists of a chronic inflammatory disease related to a bacterial, a fungal or a viral infection.
9. The method of claim 1, wherein at step b), the amount of selenoprotein P which is measured is compared with the amount of selenoprotein P which would be expected within a group of non affected individuals from the same geographical origin than that of the patient tested.
10. The method of claim 1, wherein the occurrence of the disease is diagnosed in a further step d) when the amount of selenoprotein P measured in the biological sample is lower than the normal amount which is contained within non-affected individuals.
11. The method of claim 1, wherein the persistence of the disease is diagnosed in a further step d) when the amount of selenoprotein P measured in the biological sample is lower than the normal amount which is contained within non-affected individuals or when the amount of selenoprotein P measured in the biological sample is lower or not higher than, or equal to, the amount measured in a sample from the same patient in a former test.
12. The method of claim 1, wherein the recovering from the disease is diagnosed in a further step d) when the amount of selenoprotein P measured in the biological sample is at least equal to the normal amount which is contained within non-affected individuals or the amount of selenoprotein P measured in the biological sample is higher than the amount measured in a sample from the same patient in a former test.
13. The method of claim 1, wherein the amount which is contained in a biological sample from an individual which is not affected with the disease at step b) is of at least 40 nmol/L.
14. The method of claim 1, wherein mortality is predicted when the amount of selenoprotein P which is measured within a serial of biological samples collected during the time period of the therapeutical treatment is always lower than 20nmol/L, preferably lower than 10 nmol/L.
15. The method of claim 1, wherein step a) consists of an immunoassay wherein antibodies directed against selenoprotein P are used.

**Patentansprüche**

- 5
1. Verfahren zur in-vitro-Diagnose oder zur Überwachung einer Krankheit, die eine Entzündungsreaktion innerhalb eines Patienten beinhaltet, wobei das Verfahren die folgenden Schritte umfasst:
- a) Messen der Menge an Selenoprotein P, die in einer von dem Patienten stammenden biologischen Probe enthalten ist;
- b) Vergleichen der in Schritt a) gemessenen Menge an Selenoprotein P:
- 10
- i) mit der Menge an Selenoprotein P, die in einer biologischen Probe enthalten ist, die von einem Individuum stammt, das nicht an der Krankheit leidet; oder
- ii) mit der Menge an Selenoprotein P, die bei einem früheren Test in einer von demselben Patienten stammenden biologischen Probe enthalten war; und
- 15
- c) Diagnostizieren eines septischen Schocks, wenn die Menge an Selenoprotein P, die in der biologischen Probe gemessen wird, kleiner als 30 nmol/l ist.
2. Verfahren gemäß Anspruch 1, wobei die biologische Probe aus einer Blutprobe besteht.
- 20
3. Verfahren gemäß Anspruch 1, wobei die biologische Probe aus einer Blutplasmaprobe besteht.
4. Verfahren gemäß Anspruch 1, wobei die Krankheit, die eine Entzündungsreaktion beinhaltet, aus einem systemischen inflammatorischen Response-Syndrom (SIRS) besteht.
- 25
5. Verfahren gemäß Anspruch 1, wobei die Krankheit, die eine Entzündungsreaktion beinhaltet, aus einem septischen Schock besteht.
6. Verfahren gemäß Anspruch 1, wobei die Krankheit, die eine Entzündungsreaktion beinhaltet, aus einer chronischen Entzündungskrankheit besteht.
- 30
7. Verfahren gemäß Anspruch 6, wobei die chronische Entzündungskrankheit aus der Gruppe ausgewählt ist, die aus einer systemischen Krankheit, wie Polyarteriitis nodosa, besteht.
8. Verfahren gemäß Anspruch 6, wobei die Krankheit, die eine Entzündungsreaktion beinhaltet, aus einer chronischen Entzündungskrankheit besteht, die mit einer Bakterien-, Pilz- oder Virusinfektion zusammenhängt.
- 35
9. Verfahren gemäß Anspruch 1, wobei in Schritt b) die Menge an Selenoprotein P, die gemessen wird, mit der Menge an Selenoprotein P verglichen wird, die man innerhalb einer Gruppe von nicht betroffenen Individuen derselben geographischen Herkunft wie der getestete Patient erwarten würde.
- 40
10. Verfahren gemäß Anspruch 1, wobei das Auftreten der Krankheit in einem weiteren Schritt d) diagnostiziert wird, wenn die Menge an Selenoprotein P, die in der biologischen Probe gemessen wird, kleiner ist als die normale Menge, die in nicht betroffenen Individuen enthalten ist.
- 45
11. Verfahren gemäß Anspruch 1, wobei das Fortbestehen der Krankheit in einem weiteren Schritt d) diagnostiziert wird, wenn die Menge an Selenoprotein P, die in der biologischen Probe gemessen wird, kleiner ist als die normale Menge, die in nicht betroffenen Individuen enthalten ist, oder wenn die Menge an Selenoprotein P, die in der biologischen Probe gemessen wird, kleiner oder nicht größer oder gleich groß ist wie die Menge, die bei einem früheren Test in einer von demselben Patienten stammenden Probe gemessen wurde.
- 50
12. Verfahren gemäß Anspruch 1, wobei die Genesung von der Krankheit in einem weiteren Schritt d) diagnostiziert wird, wenn die Menge an Selenoprotein P, die in der biologischen Probe gemessen wird, wenigstens gleich der normalen Menge ist, die in nicht betroffenen Individuen enthalten ist, oder wenn die Menge an Selenoprotein P, die in der biologischen Probe gemessen wird, größer ist als die Menge, die bei einem früheren Test in einer von demselben Patienten stammenden Probe gemessen wurde.
- 55
13. Verfahren gemäß Anspruch 1, wobei die Menge, die in einer biologischen Probe enthalten ist, die von einem Individuum stammt, das nicht an der Krankheit leidet, in Schritt b) wenigstens 40 nmol/l beträgt.

14. Verfahren gemäß Anspruch 1, wobei eine Sterbewahrscheinlichkeit vorhergesagt wird, wenn die Menge an Selenoprotein P, welche in einer Reihe von biologischen Proben gemessen wird, die während des Zeitraums der therapeutischen Behandlung gewonnen wurden, stets kleiner als 20 nmol/l, vorzugsweise kleiner als 10 nmol/l, ist.

5 15. Verfahren gemäß Anspruch 1, wobei Schritt a) aus einem Immunoassay besteht, wobei gegen Selenoprotein P gerichtete Antikörper verwendet werden.

## Revendications

10 1. Procédé pour le diagnostic in vitro ou pour la surveillance d'une maladie impliquant une réaction inflammatoire chez un patient, ledit procédé comprenant les étapes de :

15 a) mesurer la quantité de sélénoprotéine P qui est présente dans un échantillon biologique provenant dudit patient;

b) comparer la quantité de sélénoprotéine P mesurée au niveau de l'étape a)

20 i) avec la quantité de sélénoprotéine P qui est présente dans un échantillon biologique provenant d'un individu non affecté par la maladie; ou

ii) avec la quantité de sélénoprotéine P qui était présente dans un échantillon biologique provenant du même patient dans un test antérieur; et

25 c) diagnostiquer un choc septique lorsque la quantité de sélénoprotéine P qui est mesurée dans l'échantillon biologique est inférieure à 30 nmol/L.

2. Procédé selon la revendication 1, dans lequel l'échantillon biologique consiste en un échantillon de sang.

3. Procédé selon la revendication 1, dans lequel l'échantillon biologique consiste en un échantillon de plasma sanguin.

30 4. Procédé selon la revendication 1, dans lequel ladite maladie impliquant une réaction inflammatoire consiste en un Syndrome de Réponse Inflammatoire Systémique (SIRS).

5. Procédé selon la revendication 1, dans lequel ladite maladie impliquant une réaction inflammatoire consiste en un choc septique.

35 6. Procédé selon la revendication 1, dans lequel ladite maladie impliquant une réaction inflammatoire consiste en une maladie inflammatoire chronique.

40 7. Procédé selon la revendication 6, dans lequel ladite maladie inflammatoire chronique est choisie dans le groupe constitué par une maladie systémique telle que la polyartérite noueuse.

8. Procédé selon la revendication 6, dans lequel ladite maladie impliquant une réaction inflammatoire consiste en une maladie inflammatoire chronique associée à une infection bactérienne, fongique ou virale.

45 9. Procédé selon la revendication 1, dans lequel au niveau de l'étape b), la quantité de sélénoprotéine P qui est mesurée est comparée à la quantité de sélénoprotéine P à laquelle on devrait s'attendre au sein d'un groupe d'individus non affectés, issu de la même origine géographique que le patient testé.

50 10. Procédé selon la revendication 1, dans lequel l'apparition de la maladie est diagnostiquée dans une étape ultérieure d) lorsque la quantité de sélénoprotéine P mesurée dans l'échantillon biologique est inférieure à la quantité normale qui est présente chez les individus non affectés.

55 11. Procédé selon la revendication 1, dans lequel la persistance de la maladie est diagnostiquée dans une étape ultérieure d) lorsque la quantité de sélénoprotéine P mesurée dans l'échantillon biologique est inférieure à la quantité normale qui est présente chez les individus non affectés ou lorsque la quantité de sélénoprotéine P mesurée dans l'échantillon biologique est inférieure ou non supérieure ou égale à la quantité mesurée dans un échantillon provenant du même patient dans un test antérieur,

## EP 1 393 078 B1

- 5
12. Procédé selon la revendication 1, dans lequel la guérison de la maladie est diagnostiquée dans une étape ultérieure d) lorsque la quantité de sélénoprotéine P mesurée dans l'échantillon biologique est au moins égale à la quantité normale qui est présente chez les individus non affectés ou lorsque la quantité de sélénoprotéine P mesurée dans l'échantillon biologique est supérieure à la quantité mesurée dans un échantillon provenant du même patient dans un test antérieur.
- 10
13. Procédé selon la revendication 1, dans lequel la quantité qui est présente dans un échantillon biologique provenant d'un individu non affecté par la maladie au niveau de l'étape b) est d'au moins 40 nmol/L.
14. Procédé selon la revendication 1, dans lequel la mortalité est pronostiquée lorsque la quantité de sélénoprotéine P qui est mesurée dans une série d'échantillons biologiques prélevés durant la période de temps du traitement thérapeutique est toujours inférieure à 20 nmol/L, de préférence inférieure à 10 nmol/L.
- 15
15. Procédé selon la revendication 1, dans lequel l'étape a) consiste en un dosage immunologique dans lequel on utilise des anticorps dirigés contre la sélénoprotéine P.
- 20
- 25
- 30
- 35
- 40
- 45
- 50
- 55

## REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

## Non-patent literature cited in the description

- Definition and epidemiology This LG. **Dhainaut JF ; Dhainaut JF ; This LG ; G Parked**. Septic Shock. *Sunders*, 2000 [0002]
- From Celsus to Galen to Bone: The Illnesses, Syndromes, and Diseases of Acute Inflammation. **Marshall JC ; Aarts MA**. Yearbook of Intensive Care and Emergency Medicine. Springer Verlag, 2001, 3-12 [0002] [0006] [0007] [0009] [0010]
- **Novelli, G. P.** Role of free radicals in septic shock. *J Physiol Pharmacol*, 1997, vol. 48 (4), 517-527 [0003]
- **Zimmerman, J. J.** Defining the role of oxyradicals in the pathogenesis of sepsis. *Critical Care Medicine*, 1995, vol. 23 (4), 616-617 [0003]
- **Bast, A. ; Haenen, G. R. ; Doelman, C. J.** Oxidants and antioxidants: state of the art. *Am J Med*, 1991, vol. 91 (3C), 2S-13S [0003]
- **Forceville X.** Selenium and the "free" electron" -Selenium a trace to be followed in septic or inflammatory ICU patients. *Intensive Care Medicine*, 2001, vol. 27, 16-18 [0003] [0100]
- **Forceville et al.** *Intensive Care Medicine*, 2000, vol. 26, 5228 [0004]
- **Forceville et al.** *Critical Care Medicine*, 1998, vol. 26, 1536-1544 [0004]
- **Bernard G.R. ; Vincent JL ; Laterre PF.** Efficacy and Safety of Recombinant Human Activated Protein C for Severe Sepsis. *New England Journal of Medicine*, 2001 [0005]
- Sepsis/septic shock: Participation of the microcirculation: An abbreviated review. *Hinshaw LB Critical Care Medicine*, 1996, vol. 24 (6), 1072-1078 [0005]
- The ACCP-SCCM consensus conference on sepsis and organ failure. *Chest*, 1992, vol. 101, 1644-55 [0007] [0008]
- Definition and epidemiology This LG. **Dhainaut JF**. Septic Shock. *Sunders*, 2000 [0008]
- Diagnosis : from clinical signs to haemodynamic evaluation. **Groeneveld AB J ; This GT**. Septic Shock. *Sunders*, 2000 [0009] [0009]
- **van der Poll T ; van Deventer S JH.** Cytokines and Anticytokines in the Pathogenesis of sepsis. *infectious disease Clinics of North America*, 1999, vol. 13 (2), 413-426 [0009] [0009]
- **Rheinart K ; Meisner M ; Hartog C.** Diagnosis of sepsis: Novel and Conventional Parameters. *Advances in sepsis*, 2001, 42-51 [0009]
- **Vincent JL ; Byl B.** Defining a Clinical Syndrome of Systemic inflammation. *Sepsis*, 2000, vol. 4, 15-19 [0009]
- **Burk RF ; Hill KE.** Orphan selenoproteins. *BioEssays*, 1999, vol. 21 (3), 231-237 [0027]
- Selenoprotein P: Properties, Functions, and Regulation Mostert V. *Archives of Biochemistry and Biophysics*, 2000, vol. 376 (2), 433-438 [0027]
- **Mostert V ; Lombeck I ; Abel J.** A Novel Method for the Purification of Selenoprotein P from Human Plasma. *Archives of Biochemistry and Biophysics*, 1998, vol. 357 (2), 326-330 [0027]
- **Burk, R. F. ; Hill, K. E. ; Boeglin, M. E. ; Ebner, F. F. ; Chittum, H. S.** Selenoprotein P associates with endothelial cells in rat tissues. *Histochem Cell Biol*, 1997, vol. 108 (1), 11-15 [0028]
- Celsus to Galen to Bone: The illnesses syndromes and diseases of acute inflammation. **Marshall JC ; Aarts MA**. Yearbook of Intensive Care and Emergency Medicine. Springer Verlag, 2001, 3-12 [0040]
- **Marchaluk E ; Persson-Moschos M ; Thorling EB ; Akesson B.** Variation in selenoprotein P concentration in serum from different European regions. *Eur J Clin Nutr*, 1995, vol. 49 (1), 42-48 [0041]
- **Persson-Moschos M ; Aiftan G ; Akesson B.** Plasma selenoprotein P levels of healthy males in different selenium status after oral supplementation with different form of selenium. *Eur J Clin Nutr*, 1998, vol. 52 (5), 363-367 [0041]
- **Rayman, M. P.** The importance of selenium to human health. *Lancet*, 2000, vol. 356, 233-41 [0041]
- **Persson-Moschos M ; Huang W ; Srikumar TS ; Akesson B ; Lindeberg S.** Selenoprotein P in serum as a biochemical marker of selenium status. *Analyst*, 1995, vol. 120 (3), 833-836 [0041]
- **Mostert et al.** *Archives of Biochemistry and Biophysics*, 1998, vol. 357 (2), 326-330 [0066]
- **Mostert V ; Lombeck I ; Abel J.** A Novel Method for the Purification of selenoprotein P from Human Plasma. *Archives of Biochemistry and Biophysics*, 1998, vol. 357 (2), 326-330 [0080] [0082]
- **Le Gall JR ; Lemeshow S ; Saulnier F.** European/North American multicenter study. *JAMA*, 1993, vol. 270, 2957-2963 [0083]
- **Vincent JL ; Moreno R ; Takala J.** Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. *Intensive Care Medicine*, 1996, vol. 22, 707-710 [0084]

专利名称(译)	用于体外诊断或用于监测涉及炎症反应的疾病的方法		
公开(公告)号	<a href="#">EP1393078B1</a>	公开(公告)日	2009-07-08
申请号	EP2002743017	申请日	2002-05-15
[标]申请(专利权)人(译)	福塞维尔XAVIER		
申请(专利权)人(译)	福塞维尔, XAVIER		
当前申请(专利权)人(译)	福塞维尔, XAVIER		
[标]发明人	FORCEVILLE XAVIER		
发明人	FORCEVILLE, XAVIER		
IPC分类号	G01N33/68 G01N33/53		
CPC分类号	G01N33/6893 G01N2333/47 Y10S436/801 Y10S436/811		
优先权	60/290973 2001-05-16 US		
其他公开文献	EP1393078A1		
外部链接	<a href="#">Espacenet</a>		

#### 摘要(译)

本发明涉及体外诊断或监测涉及患者体内炎症反应的疾病的方法，其中所述方法包括以下步骤：a) 从所述患者提供生物样品；b) 测量所述生物样品中含有的硒蛋白P的量；c) 将步骤b) i) 测量的硒蛋白P的量与来自未受疾病影响的个体的生物样品中含有的硒蛋白P的量进行比较；或ii) 含有来自同一患者的生物样品中的硒蛋白P的量。

**Table 2**

Selenoprotein P plasma concentration at admission in ICU patients			
Groups	Septic shock or severe SIRS patients	Non severe SIRS patients	Healthy volunteers
number of patients	(9 patients) (measurement at admission on only 8 patients)	(12 patients)	(7 volunteers)
Selenoprotein P	17 ± 12 nmol/L (1 - 38)	55 ± 45 nmol/L (20 - 181)	49 ± 8 nmol/L (39 - 65)
<p>(p &lt; 0.01 vs. healthy volunteers)            (p &lt; 0.05 vs. non severe SIRS patients)            SIRS : Systemic Inflammatory Response Syndrome</p>			