

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number
WO 01/83519 A1

(51) International Patent Classification⁷: C07K 7/06, 7/08,
14/195, 16/12, A61K 39/02, G01N 33/53, 33/551, 33/68

S.A. 5000 (AU). HEUZENROEDER, Michael, W.
[AU/AU]; Infectious Diseases Laboratories, Institute of
Medical and Veterinary Sciences, Frome Road, Adelaide,
S.A. 5000 (AU).

(21) International Application Number: PCT/AU01/00478

(22) International Filing Date: 27 April 2001 (27.04.2001)

(74) Agent: A.P.T. PATENT AND TRADE MARK ATTOR-
NEYS; G.P.O. Box 772, Adelaide, S.A. 5001 (AU).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PQ 7120 27 April 2000 (27.04.2000) AU

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID,
IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT,
LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL,
PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ,
UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicants (*for all designated States except US*): UNI-
VERSITY OF SOUTH AUSTRALIA [AU/AU]; North
Terrace, Adelaide, S.A. 5000 (AU). MEDVET SCIENCE
PTY LTD [AU/AU]; 20 Dalgleish Street, Thebarton, S.A.
5031 (AU). RURAL INDUSTRIES RESEARCH AND
DEVELOPMENT CORPORATION [AU/AU]; P.O. Box
4776, Kingston, ACT 2604 (AU).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): VANNIASINKAM,
Thiru [AU/AU]; School of Pharmacy and Medical Labora-
tory Science, University of South Australia North Terrace,
Adelaide, S.A. 5000 (AU). BARTON, Mary [AU/AU];
School of Pharmacy and Medical Laboratory Science,
University of South Australia North Terrace, Adelaide,

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTIGENIC PEPTIDE FRAGMENTS OF VapA PROTEIN, AND USES THEREOF

(57) Abstract: An isolated peptide fragment of the Vap A protein that binds antibodies specific for *Rhodococcus equi* and the VapA protein. In a preferred form the peptide contains an amino acid sequence of 5 or more amino acid residues that is identical to or homologous to the amino acid sequence of at least one region of the VapA protein that is responsible for immunological recognition. Methods of diagnosing a vertebrate for the presence of *R. equi* using the peptide and methods of vaccinating a vertebrate against *R. equi* using the peptide are also claimed.



WO 01/83519 A1

ANTIGENIC PEPTIDE FRAGMENTS OF VapA PROTEIN, AND USES THEREOF

5 FIELD OF THE INVENTION

The present invention relates to peptide fragments that mimic immunogenic properties specific for the VapA protein and can be used as a basis for diagnoses and vaccinations of vertebrates against *Rhodococcus equi*.

10 BACKGROUND OF THE INVENTION

Rhodococcus equi is an encapsulated and rod shaped, Gram positive bacterium that is considered to be a soil saprophyte that survives well in the environment. *R. equi* has long been considered a pathogen in horses principally in foals fewer than 6 months old (particularly 1-3 months old). Infection by the organism is
15 accompanied by extra-pulmonary manifestations, causes a pyogranulomatous pneumonia, often such as bacteraemia, lymphadenitis, meningitis and enteritis (Barton and Hughes, 1980; Giguere and Prescott, 1997; Takai, 1997). Infections are often fatal if untreated. Apart from causing disease in horses, *R. equi* also causes infections in cattle, pigs and goats (Barton, 1992). *R. equi* is also known
20 to cause severe pulmonary and disseminated disease in immuno-compromised humans, particularly AIDS patients (Capdevila *et al.*, 1997).

In Australia most equine *R. equi* infections occur in summer (December to February) when the age of the foals as well as the warm and dry environmental
25 conditions make the animals more susceptible to infection (Barton and Hughes, 1984).

R. equi produces a range of putative virulence factors such as cholesterol oxidase, phospholipase C and lecithinase (Smola *et al.* 1994). However one of the more
30 important putative virulence factors is considered to be a 17kDa virulence associated protein (VapA) which is plasmid encoded. This protein is known to be produced by up to 90% of equine clinical isolates of *R. equi*. Although VapA producing strains are widespread among disease causing isolates, recent work has shown that VapA protein alone is not sufficient to cause disease in foals and that
35 other as yet unknown plasmid borne factors are likely to be involved (Giguere *et al.*, 1999). The role of VapA in virulence is yet to be elucidated, although there is strong evidence to suggest that the plasmid encoding the protein may play an

important part in the survival of the organism within macrophages (Hondalus and Mosser, 1994).

5 Current techniques for the identification of *R. equi* such as culture and phenotypic testing are often not successful due to the slow growing nature of the organism. It is also often difficult to differentiate *R. equi* from closely related organisms based on biochemical tests alone. It has been shown that the detection of high levels of IgG antibodies to the VapA protein is an indicator of *R. equi* infection in horses (Prescott *et al.*, 1996; Higuchi *et al.*, 1997). Of the immunodiagnostic tests developed to date, none have been found suitable for use on a routine basis mainly due to the use of laborious protein extraction techniques (Higuchi *et al.*, 10 1997; Takai, 1997). Thus an improved immunodiagnostic test based on the VapA protein would be of great benefit as a routine diagnostic test.

15 Further, attempts at immunisation against Rhodococcus infection have met with limited success to date. The use of antibiotics has been found to be only partially effective and vaccines developed over the years for the prevention of *R. equi* infection in horses have not been particularly effective.

20 Mosser in WO 99/05304 has found that an avirulent strain transformed with a VapA expressing plasmid is avirulent but is also protective and describes a vaccine comprising DNA encoding the VapA protein or fragment thereof. Mosser's results are based on the immunogenicity of the whole VapA protein. However, Mosser has found that VapA is poorly immunogenic in mice (personal communication).
25

Peptides, and particularly relatively small peptides, have an advantage over whole proteins in diagnostics and therapeutics in that they are more readily produced than the whole protein, and they generate a population of homogenous molecules, i.e. single peptides composed of the same amino acids. Further it may not be economically viable to synthesise large proteins and therefore native proteins are often obtained by extraction. However native proteins derived from natural sources may contain other proteins or peptides of the same origin as the target proteins and as such the complexity and variability of mixtures of native antigen
30 proteins can persist even after fractionation and purification and this can be a barrier to their use in immunoassays and vaccines.
35

For these reasons it may be more preferable if a vaccine were based on one or more small peptides that have the immunogenic properties of the whole VapA protein. However, Mosser does not identify any peptide fragments that provide for recognition of antibodies specific for VapA. Further, it is often difficult to identify peptide fragments that mimic immunogenic properties specific for a whole protein because in many cases the peptide fragments do not take up the three dimensional structure necessary for immunological recognition.

OBJECT OF THE INVENTION

The object of one aspect of this invention is to provide one or more peptide fragments that mimic immunogenic properties specific for the VapA protein and can be used as a basis for diagnoses and vaccinations of vertebrates against *R. equi* that ameliorate one or more of the problems associated with existing diagnoses and vaccinations, or at least to provide the public with a useful choice.

For the purpose of this specification the word "comprising" means "including but not limited to", and the word "comprise" has a corresponding meaning.

SUMMARY OF THE INVENTION

The present invention arises out of experiments designed to map the antigenic epitopes of the VapA protein that are reactive with sera from infected horses. The inventors have discovered peptide fragments of the VapA protein that are recognised by the sera of horses that have been infected by *R. equi* and are unrecognised by control horses.

As a result the invention provides peptide targets for use in assays for the early and rapid diagnosis of infection in vertebrates. Further, the invention provides peptide vaccines for protecting vertebrates against infection by *R. equi*.

Therefore, in one form of a first aspect the invention could be said to reside in an isolated peptide fragment of the VapA protein that is capable of binding antibodies specific for *R. equi* and the VapA protein.

The peptide preferably includes an amino acid sequence that is identical to or homologous to an amino acid sequence of at least one region of the VapA protein that is responsible for immunological recognition. The peptide preferably contains an amino acid sequence of 5 or more amino acid residues that is identical

to or homologous to the amino acid sequence of at least one region of the VapA protein that is responsible for immunological recognition. The amino acid sequence may be part of a larger peptide. The amino acid sequence of the VapA protein has been previously published (Sekizaki *et al.* 1995; GenBank database
5 Accession No: D21236) and therefore the larger peptide sequence may contain the 5 or more amino acid sequence of the present invention plus other segments of the VapA protein.

A putative 20 amino acid region of the VapA protein that is recognised by
10 antibodies in the sera of horses infected with *R. equi* has been identified as TSLNLQKDEPNGRASDTAGQ [Seq I.D. No. 1], although it will be understood that the minimal region for antigenic recognition may be further defined within the identified sequence, or additionally the identified sequence may contain two or more separate adjacent epitopes. Thus the peptide may be any peptide that is
15 capable of mimicking this region in so far as providing VapA specific immunogenicity. Therefore the peptide may be part of a larger peptide that contains the amino acid sequence TSLNLQKDEPNGRASDTAGQ [Seq I.D. No. 1] of the present invention, as well as one or more of the amino acids either side of that sequence in the native VapA protein.

20 Therefore in one form of the first aspect the peptide has 5 or more amino acid residues and contains all or part of the sequence TSLNLQKDEPNGRASDTAGQ [Seq I.D. No. 1], or immunologically active derivative or analogue thereof. Preferably the peptide contains 7 to 30 amino acid residues, and more preferably
25 10 to 12 amino acid residues.

The peptide may be selected from the list including
TSLNLQKDEPNGRASDTAGQ [Seq I.D. No. 1],
SLNLQKDEPNGRASDTAGQ [Seq I.D. No. 2], TSLNLQKDEPNGRASDTAG
30 [Seq I.D. No. 3], LNLQKDEPNGRASDTAGQ [Seq I.D. No. 4],
TSLNLQKDEPNGRASDTA [Seq I.D. No. 5], NLQKDEPNGRASDTAGQ [Seq I.D. No. 6], TSLNLQKDEPNGRASDT [Seq I.D. No. 7],
LQKDEPNGRASDTAGQ [Seq I.D. No. 8], TSLNLQKDEPNGRASD [Seq I.D. No. 9], SLNLQKDEPNGRASDT [Seq I.D. No. 10], LNLQKDEPNGRASDTA
35 [Seq I.D. No. 11], NLQKDEPNGRASDTAG [Seq I.D. No. 12],
QKDEPNGRASDTAGQ [Seq I.D. No. 13], TSLNLQKDEPNGRAS [Seq I.D. No. 14], SLNLQKDEPNGRASD [Seq I.D. No. 15], LNLQKDEPNGRASDT

[Seq I.D. No. 16], NLQKDEPNGRASDTA [Seq I.D. No. 17],
LQKDEPNGRASDTAG [Seq I.D. No. 18], KDEPNGRASDTAGQ [Seq I.D. No.
19], TSLNLQKDEPNGRA [Seq I.D. No. 20], SLNLQKDEPNGRAS [Seq I.D.
No. 21], LNLQKDEPNGRASD [Seq I.D. No. 22], NLQKDEPNGRASDT [Seq
5 I.D. No. 23], LQKDEPNGRASDTA [Seq I.D. No. 24], QKDEPNGRASDTAG
[Seq I.D. No. 25], DEPNGRASDTAGQ [Seq I.D. No. 26], TSLNLQKDEPNGR
[Seq I.D. No. 27], SLNLQKDEPNGRA [Seq I.D. No. 28], LNLQKDEPNGRAS
[Seq I.D. No. 29], NLQKDEPNGRASD [Seq I.D. No. 30], LQKDEPNGRASDT
[Seq I.D. No. 31], QKDEPNGRASDTA [Seq I.D. No. 32], KDEPNGRASDTAG
10 [Seq I.D. No. 33], EPNGRASDTAGQ [Seq I.D. No. 34], TSLNLQKDEPNG
[Seq I.D. No. 35], SLNLQKDEPNGR [Seq I.D. No. 36], LNLQKDEPNGRAS
[Seq I.D. No. 37], NLQKDEPNGRAS [Seq I.D. No. 38], LQKDEPNGRASD
[Seq I.D. No. 39], QKDEPNGRASDT [Seq I.D. No. 40], KDEPNGRASDTA
[Seq I.D. No. 41], DEPNGRASDTAG [Seq I.D. No. 42], PNGRASDTAGQ [Seq
15 I.D. No. 43], TSLNLQKDEPN [Seq I.D. No. 44], SLNLQKDEPNG [Seq I.D.
No. 45], LNLQKDEPNGR [Seq I.D. No. 46], NLQKDEPNGRA [Seq I.D. No.
47], LQKDEPNGRAS [Seq I.D. No. 48], QKDEPNGRASD [Seq I.D. No. 49],
KDEPNGRASDT [Seq I.D. No. 50], DEPNGRASDTA [Seq I.D. No. 51],
EPNGRASDTAG [Seq I.D. No. 52], NGRASDTAGQ [Seq I.D. No. 53],
20 TSLNLQKDEP [Seq I.D. No. 54], SLNLQKDEPN [Seq I.D. No. 55],
LNLQKDEPNGR [Seq I.D. No. 56], NLQKDEPNGRA [Seq I.D. No. 57],
LQKDEPNGRA [Seq I.D. No. 58], QKDEPNGRAS [Seq I.D. No. 59],
KDEPNGRASD [Seq I.D. No. 60], DEPNGRASDT [Seq I.D. No. 61],
EPNGRASDTA [Seq I.D. No. 62], PNGRASDTAG [Seq I.D. No. 63],
25 GRASDTAGQ [Seq I.D. No. 64], TSLNLQKDE [Seq I.D. No. 65],
SLNLQKDEP [Seq I.D. No. 66], LNLQKDEPN [Seq I.D. No. 67],
NLQKDEPNG [Seq I.D. No. 68], LQKDEPNGR [Seq I.D. No. 69],
QKDEPNGRA [Seq I.D. No. 70], KDEPNGRAS [Seq I.D. No. 71],
DEPNGRASD [Seq I.D. No. 72], EPNGRASDT [Seq I.D. No. 73],
30 PNGRASDTA [Seq I.D. No. 74], NGRASDTAG [Seq I.D. No. 75],
RASDTAGQ [Seq I.D. No. 76], TSLNLQKD [Seq I.D. No. 77], SLNLQKDE
[Seq I.D. No. 78], LNLQKDEP [Seq I.D. No. 79], NLQKDEPN [Seq I.D. No.
80], LQKDEPNG [Seq I.D. No. 81], QKDEPNGR [Seq I.D. No. 82],
KDEPNGRA [Seq I.D. No. 83], DEPNGRAS [Seq I.D. No. 84], EPNGRASD
35 [Seq I.D. No. 85], PNGRASDT [Seq I.D. No. 86], NGRASDTA [Seq I.D. No.
87], GRASDTAG [Seq I.D. No. 88], ASDTAGQ [Seq I.D. No. 89], TSLNLQK
[Seq I.D. No. 90], SLNLQKD [Seq I.D. No. 91], LNLQKDE [Seq I.D. No. 92],

NLQKDEP [Seq I.D. No. 93], LQKDEPN [Seq I.D. No. 94], QKDEPNG [Seq I.D. No. 95], KDEPNGR [Seq I.D. No. 96], DEPNGRA [Seq I.D. No. 97], EPNGRAS [Seq I.D. No. 98], PNGRASD [Seq I.D. No. 99], NGRASDT [Seq I.D. No. 100], GRASDTA [Seq I.D. No. 101], RASDTAG [Seq I.D. No. 102],
5 TSLNLQ [Seq I.D. No. 103], SLNLQK [Seq I.D. No. 104], LNLQKD [Seq I.D. No. 105], NLQKDE [Seq I.D. No. 106], LQKDEP [Seq I.D. No. 107], KDEPNG [Seq I.D. No. 108], DEPNGR [Seq I.D. No. 109], EPNGRA [Seq I.D. No. 110], PNGRAS [Seq I.D. No. 111], NGRASD [Seq I.D. No. 112], GRASDT [Seq I.D. No. 113], RASDTA [Seq I.D. No. 114], ASDTAG [Seq I.D. No. 115], SDTAGQ
10 [Seq I.D. No. 116], TSLNL [Seq I.D. No. 117], SLNLQ [Seq I.D. No. 118], LNLQK [Seq I.D. No. 119], NLQKD [Seq I.D. No. 120], LQKDE [Seq I.D. No. 121], QKDEP [Seq I.D. No. 122], KDEPN [Seq I.D. No. 123], DEPNG [Seq I.D. No. 124], EPNGR [Seq I.D. No. 125], PNGRA [Seq I.D. No. 126], NGRAS [Seq I.D. No. 127], GRASD [Seq I.D. No. 128], RASDT [Seq I.D. No. 129], ASDTA
15 [Seq I.D. No. 130], SDTAG [Seq I.D. No. 131], DTAGQ [Seq I.D. No. 132].

Preferably the peptide is selected from the list including TSLNLQKDEPN [Seq I.D. No. 44], NLQKDEPNGRA [Seq I.D. No. 47], KDEPNGRASDT [Seq I.D. No. 50], and PNGRASDTAGQ [Seq I.D. No. 43]. Most preferably the peptide is
20 NLQKDEPNGRA [Seq I.D. No. 47].

Whether a peptide of the present invention provides for VapA specific immunogenicity can be determined routinely by following the procedures outlined herein.

25

The peptide may also be homologous to any of the abovementioned peptides provided that the peptide provides for VapA specific immunogenicity. In this context, a peptide is considered homologous to a peptide of the present invention when it is immuno cross-reactive with antibodies specific for the *R. equi* VapA
30 protein. It will be recognised by those skilled in the art that some amino acid sequences within the peptide can be varied without significant effect on the structure or function of the peptide. Thus for instance it is anticipated that 'type' amino acid substitutions still retain immuno cross reactivity and as such a neutral amino acid may be conservatively substituted with another neutral natural or non-
35 natural amino acid, an acidic amino acid may be conservatively substituted with a natural or non-natural acidic amino acid, a hydrophilic amino acid may be

substituted with another hydrophilic amino acid, and so on, provided that the immunological function of the peptide is not altered by the substitution.

Typically seen as conservative substitutions are the replacement of one for
5 another among the aliphatic amino acids Ala, Val, Leu and Ile; interchange of the
hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu;
substitutions between the amide residues Asn and Gln; exchange of the basic
residues Lys and Arg; and replacements among the aromatic residues Phe and
Tyr. Preferably the homologous peptide shares 50% homology with a peptide of
10 the present invention, more preferably shares 70% homology, and most preferably
shares 90% homology.

The peptide may also be a non-native polypeptide which contains the epitope
embodied in any one or more of the abovementioned peptides. Thus the
15 polypeptide may contain any one or more of the abovementioned peptides, and
variations thereof, as part of a larger peptide or protein. In order to present the
peptide in a form more suitable for eliciting an immune reaction to raise
antibodies thereagainst, for example, the peptide may be part of a chimeric
protein.

20 The invention may also be said to reside in a polynucleotide that encodes any one
or more of the abovementioned peptides or partial VapA proteins, or a cell
transformed with a recombinant plasmid that expresses any one or more of the
abovementioned peptides or partial VapA proteins.

25 The peptides of the present invention may be used in diagnosing a vertebrate for
the presence of *R. equi*. Thus the peptides may be used to directly or indirectly
detect the presence of antibodies to the VapA protein present in the vertebrate.
Alternatively, the peptides of the present invention may be used to produce
30 antibodies that can in turn be used to detect the presence of VapA antigens in the
vertebrate.

Thus, in one form of a second aspect the invention could be said to reside in a
method of diagnosing a vertebrate for the presence of *R. equi* by detecting the
35 presence of antibodies to *R. equi* in the vertebrate, the method including the steps
of:

obtaining a sample of antibody containing fluid from the vertebrate,

contacting the sample with at least one peptide fragment of a region for
specific immunogenic recognition of the VapA protein, or
derivative thereof, and
assaying for the formation of the antigen:antibody complex to detect the
presence of antibodies to the VapA protein.

5

The antibody containing fluid may be any biological fluid in the vertebrate that
contains antibodies including, but not limited to, serum, plasma, whole blood,
cerebro spinal fluid, amniotic fluid, and synovial fluid.

10

The peptide fragment may be any of the peptides of the first aspect of the
invention including derivatives, variants and chimeric peptides and proteins
containing the peptides.

15 In contacting the biological sample with the peptide, the peptide fragment may be
attached or conjugated to a carrier molecule or attached or conjugated to a solid
support. A solid support in the present invention means any solid material to
which the peptide can be complexed or attached. Examples of such solid supports
include, but are not limited to, microtitre plates, petri dishes, bottles, slides, and
20 other such containers made of plastic, glass, polyvinyl, polystyrene, and other
solid materials which allow detection of labelled antibodies. Other suitable
carriers for binding the peptide exist or will be able to be ascertained by routine
experimentation.

25 After contacting the peptides of the present invention under conditions suitable
for formation of an antigen:antibody complex any component of the biological
sample that is not bound to the peptide fragment on the solid support may be
washed or otherwise removed from the bound complex.

30 In a preferred form of the invention the peptide(s) are covalently or non-
covalently bound to the surface of a microtitre well. A serum sample suspected
of containing antibody may then be added, unbound sample washed away and the
level of antibodies bound in the antigen:antibody complex assayed.

35 The assay for the formation of antigen:antibody compounds preferably involves
adding a compound that enables detection of the antibodies which are specifically
bound to the peptide. These assays employ a wide variety of labels and provide

for a varying range of sensitivity and susceptibility to interference. Labels include radionuclides, enzymes, fluorescers, chemiluminescers, particles, ligands, enzyme substrates, enzyme cofactors, enzyme inhibitors, light emitter-quencher combinations and the like. The immunoassays may be homogenous or
5 heterogenous, where the distinction relates to the use of a separation step for separating uncomplexed label from complexed label.

In a preferred form of the invention the assay is an ELISA and labelled anti-antibodies that are specific for the VapA antibodies may be added to bind the
10 VapA antibodies, thereby allowing detection and quantification of the VapA antibodies. Thus, for example, in the case of an equine diagnostic test, the VapA antibodies may be detected with labelled goat anti-horse IgG.

It will be appreciated that the antibodies being assayed may be members of any of
15 the five major classes of antibodies and therefore the diagnostic method encompasses IgA, IgD, IgE, IgG and IgM antibodies and the labelled anti-antibodies for use in an ELISA may be IgA, IgD, IgE, IgG, and IgM anti-antibodies, respectively.

20 Alternatively, VapA antibodies may be detected by binding with anti-antibodies specific for the VapA antibodies, and then a second labelled antibody specific for the anti-antibody may be used to detect the anti-antibodies. Thus for instance in the case of an equine diagnostic test, the VapA antibodies may be detected with
25 labelled goat anti-horse IgG and then labelled anti-goat IgG may be added to bind and detect the anti-horse IgG. This latter form of assay is believed to give better specificity in that it tends to give less false positives.

Alternatively, a peptide:antibody complex may be used to detect the presence of
30 VapA antibody in the sample. Preferably the antibody that forms part of the complex is labelled with a suitable label as discussed.

The labelled antibodies specific for VapA antibodies may be labelled with a radioisotope, which can then be determined by such means as the use of a gamma counter or a scintillation counter.

35

Another way in which the antibodies specific for VapA antibodies can be detectably labelled is by linking to an enzyme. This enzyme, in turn, when later

exposed to its substrate, will react with the substrate in such a manner as to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorometric or visual means. Examples of enzymes which can be used to detectably label the antibody of the present invention include

5 malate dehydrogenase, staphylococcal nuclease, delta-V-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-VI-phosphate dehydrogenase, glucoamylase and acetylcholine esterase. Avidin-biotin binding

10 may be used to facilitate the enzyme labelling.

It is also possible to label the antibodies specific for VapA antibodies with a fluorescent compound. When the fluorescently labelled antibody is exposed to light of the proper wavelength, its presence can then be detected due to the

15 fluorescence of the dye. Among the most commonly used fluorescent labelling compounds are fluorescein isothiocyanate, rhodamine, phycoerytherin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine.

The antibodies specific for VapA antibodies can also be detectably labelled using

20 fluorescent emitting metals such as ^{152}Eu , or others of the lanthanide series. These metals can be attached to the antibody molecule using such metal chelating groups as diethylenetriaminepentacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA).

25 The antibodies specific for VapA antibodies can also be detectably labelled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescent labelling compounds are luminol,

30 isoluminol, therromatic acridinium ester, imidazole, acridinium salt and oxalate ester.

Likewise, a bioluminescent compound may be used to label the antibodies specific for VapA antibodies. Bioluminescence is a type of chemiluminescence

35 found in biological systems in which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent antibody is

determined by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labelling are luciferin, luciferase and aequorin.

5 Another technique which may also result in greater sensitivity when used in conjunction with the present invention consists of coupling the antibodies specific for VapA antibodies to low molecular weight haptens. The haptens can then be specifically detected by means of a second reaction. For example, it is common to use such haptens as biotin (reacting with avidin) or dinitrophenyl, pyridoxal and fluorescamine (reacting with specific anti-hapten antibodies) in this manner.

10

In addition, the sensitivity of the assay may be increased by use of amplification strategies including substrate cycling and enzyme channelling as described by Lindbladh *et al.* (1993).

15 Preferably the vertebrate is selected from the list including horses, cattle, pigs, goats and humans. Most preferably the vertebrate is a horse.

In one form of a third aspect the invention could be said to reside in a method of diagnosing a vertebrate for the presence of *R. equi* by detecting the presence of VapA antigens in the vertebrate, the method including the steps of:

20

producing an antibody specific to at least one peptide fragment of a region for specific immunogenic recognition of the VapA protein, or derivative thereof, obtaining a putative VapA antigen containing biological sample from the vertebrate,

25

contacting the sample with the raised antibodies under conditions for formation of an antibody:antigen complex, and

assaying for the formation of the antibody:antigen complex to detect the presence of antibodies to the VapA protein.

30 The antibodies can be, for example, polyclonal or monoclonal antibodies as well as chimeric, single chain, equinised antibodies, and Fab fragments. Various procedures known in the art may used for the production of such antibodies and fragments.

35 The antibody may be produced by immunising a suitable animal such as mice, guinea pigs, rabbits, goats, sheep, horses, with one or more peptides of the present invention and isolating antibody producing cells from the immunised animal.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, 1975), the trioma
5 technique, the human B-cell hybridoma technique (Kozbor *et al.*, 1983). In a preferred method, the antibody producing cells may be fused with a myeloma cell to produce a pool of hybridoma cells which can then be screened for cells that produce the monoclonal antibody.

10 Monoclonal antibody fragments may also be used in the above method. Thus, the VapA antigen containing biological sample may be contacted with a fragment of a monoclonal antibody specific for a peptide of the present invention. It is to be understood that where reference is made to a fragment of a monoclonal antibody the term includes, but is not limited to, Fab, Fv and peptide fragments of the
15 monoclonal antibody, and it may also include such fragments when made as part of a different larger peptide or protein, which may be the product of a recombinant vector.

Alternatively, additional antibodies capable of binding to the VapA protein may
20 be produced in a two step procedure through the use of anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and that, therefore, it is possible to obtain an antibody which binds to a second antibody. Thus, VapA protein specific antibodies can be used to immunise an animal and the splenocytes of the animal are used to produce hybridoma cells,
25 and the hybridoma cells are screened to identify clones which produce an antibody whose ability to block the VapA protein specific antibody can be blocked by the VapA protein. Such antibodies comprise anti-idiotypic antibodies to the VapA protein specific antibody.

30 The step of assaying for formation of the antibody:antigen complex may include the step of separating the bound antibody:antigen complex from unbound antibody. Thus, the raised antibodies, or fragments thereof may be labelled as discussed in the second aspect of this invention. Any antibody:antigen complex formed by contacting the sample with the antibody may be separated from
35 unbound antibody using suitable techniques such as immunoprecipitation or techniques for separation based on size. For example, a mixture obtained after contacting the sample with antibody may be filtered through a suitable membrane

so that antibody:antigen complex is retained on the membrane and unbound antibody passes through the membrane. The labelled antibody:antigen complex can then be quantitatively assayed using standard techniques for the label used.

5 Alternatively the raised antibodies may be used in an ELISA based assay to bind the VapA protein in the sample, and a labelled antibody specific for another immuno-recognition site on VapA can be used to assay for bound VapA protein. Thus, the raised antibodies or fragments thereof may be attached or conjugated to a solid support. After contacting the raised antibodies or fragments thereof under
10 conditions suitable for formation of an antibody:antigen complex any component of the biological sample that is not bound to the antibodies or antibody fragments on the support may be washed or otherwise separated from the bound complex.

The assay for the formation of antibody:antigen complex preferably involves
15 adding a labelled antibody specific for the VapA protein and assaying for the presence of the labelled antibody as discussed with respect to the second aspect of this invention. These assays employ a wide variety of labels including radionuclides, enzymes, fluorescers, chemilumescers, particles, ligands, enzyme substrates, enzyme cofactors, enzyme inhibitors, light emitter-quencher
20 combinations and the like as described for the second aspect of this invention.

Other known methods may also be used to assay for levels of VapA antigen in the biological sample. For example VapA protein expression in tissues can be studied with classical immunohistological methods. In these, specific recognition
25 is provided by the primary antibody but the secondary detection systems can utilise fluorescent, enzyme, or other conjugated secondary antibodies. As a result an immunological staining of tissue section for pathological examination is obtained. Tissues can also be extracted, for example with urea and neutral detergent, for the liberation of the VapA protein for Western-blot or dot/slot assay
30 (Jalkanen *et al.*, 1987).

The invention may also be said to reside in a diagnostic kit containing one or more of the peptides of the present invention, in conjunction with appropriate reagents. In addition a control lymphocyte extract obtained from a healthy
35 vertebrate may be included, as well as serum from the healthy vertebrate.

Alternatively the invention may be said to reside in a diagnostic kit containing one or more antibodies, or fragments thereof, specific for the peptide fragments of the present invention, in conjunction with appropriate reagents.

- 5 The invention may also reside in a reagent for use in the above diagnostic kit. Thus in one form the reagent may be a peptide of the present invention. In another form the reagent may be an antibody, or fragment thereof, of the present invention.
- 10 In a fourth aspect, the invention could be said to reside in a method of vaccinating a vertebrate against *Rhodococcus equi*, the method including the step of administering a peptide fragment of a region for specific immunogenic recognition of VapA, or an immunogenically active derivative or variant thereof to thereby induce an immune reaction. This might be especially useful in
- 15 vaccinating a pregnant mare to potentially provide protection for the progeny.

The peptide may form part of a larger peptide or protein by cross linking to a suitable carrier. The carrier may be a carrier protein including but not limited to bovine serum albumin, tetanus toxoid, cholera toxin and deactivated diphtheria

20 toxin. For example, the peptide may be chemically coupled to the β -subunit of the heat labile (LT) enterotoxigenic *E. coli* (ETEC) toxin which has previously been used as a carrier protein for other bacterial epitopes.

Alternatively the carrier may be an organic molecule, or a solid phase carrier. In one

25 preferred form the carrier is a Ni-NTA (nickel-nitriloacetic acid) agarose to which the peptide is bound through a (Histidine)₆-tag at either the N or the C terminus. In another form the peptide may be displayed on bacteriophage particles. Thus nucleic acid vectors are used wherein an oligonucleotide encoding the peptide is fused to a portion of a gene encoding the transmembrane portion of an integral

30 protein. Upon expression of the fusion protein it is embedded in the outer cell membrane with the peptide portion of the protein facing outward. Thus the peptide is linked to a solid support which is the bacterial cell itself.

Preferably the peptide is administered in pharmaceutical dosage form as a

35 composition or formulation comprising an immunogenically effective amount of the peptide. The amount of peptide administered will vary depending on the pharmacokinetic parameters, severity of the disease treated or immunogenic

response desired. Doses may be set by a physician or veterinarian considering relevant factors including the age, weight and condition of the vertebrate including, in the case of immunogenic dosage forms, whether the vertebrate has been previously exposed to the microorganism responsible for the disease to be vaccinated against as well as the release characteristics of the peptide from pharmaceutical dosage forms of the present invention.

The vertebrate itself may be vaccinated directly according to the method of the invention, or alternatively a progeny may be passively vaccinated by maternal vaccination.

In a fifth aspect the invention could be said to reside in a method of vaccinating a vertebrate against *R. equi*, the method including the step of administering a nucleic acid sequence that encodes a peptide fragment of a region for specific immunogenic recognition of VapA so that a host cell expresses the peptide fragment to thereby induce an immune reaction.

The techniques for cloning and expressing nucleic acid sequences that encode the amino acid sequences corresponding to the peptide fragments of the present invention, e.g. synthesis of oligonucleotides, PCR, transforming cells, constructing vectors, expression systems and the like are well known in the art and standard reference materials may be consulted for specific conditions and procedures (Sambrook *et al.*, eds., Molecular Cloning, A Laboratory Manual, 2nd Edition, Cold Spring Harbour Laboratory Press, 1989).

The nucleic acid sequence may encode a chimeric peptide or protein in which a peptide fragment of the present invention is tagged into the protein, wherein the protein is expressed by a suitable host.

In one specific example, a mutagenised gene encoding the chimeric protein and expressing the peptide may be inserted into a suitable DNA vaccine vector such as pcDNA3 which is a mammalian expression vector often used in DNA vaccines. In one particular form of the invention the *groEL* gene may be used as the vaccine candidate. The *groEL* gene found in both eukaryotic and prokaryotic organisms encodes a heat shock protein (hsp60 or hsp65). This heat inducible protein is a chaperonin or molecular chaperone that acts together with other

similar proteins to help in the folding and assembly of various oligomeric proteins in the cell (Craig *et al.*, 1993).

5 Immunity to *R. equi* has been shown to depend primarily on Th1 type cell mediated immune response and studies have shown that DNA vaccines may be used to elicit a strong Th1 type response in the host unlike protein vaccines which predominantly elicit a Th2 type response (Leclerc *et al.*, 1997; Robinson, 1997).

10 It has previously been shown that DNA vaccines based on the *groEL* gene are effective in *R. equi* related organisms such as *M. tuberculosis* (Lowrie *et al.*, 1997; Lowrie *et al.*, 1999). The reason why this gene is so effective is not known but it is thought that the immunogenicity of the GroEL protein may contribute to its effectiveness. Since GroEL is a chaperonin protein (60-65 kDa) it is synthesised in large amounts by organisms when infecting a host therefore if the
15 host were to produce high levels of antibodies to the GroEL (as the result of the *in vivo* production of GroEL brought about by a DNA vaccine) these antibodies would help in resisting onset of disease caused by natural infection.

20 The method may also include the step of assaying for the presence of antibodies to VapA to ensure that there is an immune reaction to the expressed peptide. The assay for antibodies may be any of the assays discussed with respect to the second aspect of this invention.

25 In a sixth aspect the invention may be said to reside in a composition for vaccinating a vertebrate against *Rhodococcus equi*, the composition comprising at least one peptide fragment of the region for specific immunogenic recognition of VapA according to the first and fourth aspects of the invention.

30 The composition may be injected or may be added to a pharmaceutically acceptable carrier as will be apparent to those skilled in the art and as set out in "Remington's Pharmaceutical Sciences", Sixteenth Edition, Mack Publishing Co, 1980, and include water and other polar substances, including lower molecular weight alkanes, polyalkanols such as ethylene glycol, polyethylene glycol and propylene glycol as well as non-polar carriers.

35

The method of administering the vaccine may vary and could include intravenous, buccal, oral, transdermal and nasal as well as intramuscular or subcutaneous

administration. Preferably, the vaccine is administered by inhalation which may then set up local immunity. Alternatively the vaccine may be administered using other forms of mucosal priming.

- 5 In a seventh aspect the invention could be said to reside in a method for treating a vertebrate infected with *Rhodococcus equi*, the method including the step of administering to the vertebrate a therapeutic agent capable of binding VapA protein, wherein the agent is an antibody or other small molecule capable of binding any one or more of the peptides of the present invention.

10

The antibody may be produced by immunising a suitable animal with one or more peptides of the present invention and isolating antibody producing cells from the immunised animal. The antibody producing cells may be fused with a myeloma cell to produce a pool of hybridoma cells which can then be screened for cells that

15 produce the monoclonal antibody.

Whilst monoclonal antibodies may be successfully used in therapy, they are large proteins that may invoke an immune response in the recipient and therefore it may be preferable that monoclonal antibody fragments be used in therapy.

- 20 Therefore the therapeutic agent may be a Fab, Fv or peptide fragment of a monoclonal antibody directed to a peptide of the present invention.

Other compounds capable of binding the specific immunogenic region of the VapA protein may be isolated by screening for binding to peptides of the present

25 invention. For example, a scramble of randomly synthesised compounds could be passed through a solid matrix to which a peptide of the present invention is bound. Following washing the strongly binding compounds remain and can be eluted and characterised using standard techniques. The screening may also be a competitive binding screen used to identify compounds that bind the peptide in

30 preference to a monoclonal antibody specific for that peptide.

The nature of the compounds obtained by screening is not limited and may include, but is not limited to, peptides, oligonucleotides, amino acids, nucleic acids or sugars. The methods used for the binding assay can be any one of the

35 many common techniques known to those skilled in the art. Such methods may include affinity selection chromatography, ultrafiltration assays, the scintillation proximity assay, interfacial optical techniques, the quartz crystal microbalance,

the jet ring cell, interferometric assays using porous silicon to immobilise the receptor. Reference to such techniques can be found in Woodbury *et al.*, 1999. By way of example, a scramble of randomly synthesised oligonucleotides could be passed through a solid matrix to which a peptide of the present invention is
5 bound. Following washing the strongly binding oligonucleotides remain and can be eluted under different conditions (salt, pH etc). The sequence can be determined by PCR and tested for inhibition of VapA.

It is to be understood that where reference is made to a fragment of a monoclonal
10 antibody the term includes, but is not limited to, Fab, Fv and peptide fragments of the monoclonal antibody, and it may also include such fragments when made as part of a different larger peptide or protein, which may be the product of a recombinant vector. Thus the variable region of the respective monoclonal antibody may be cloned and be made part of a hybrid protein with properties
15 appropriate for the therapeutic purposes of the respective agent. Thus for example the monoclonal antibody may be "equinised" by recombining nucleic acid encoding the variable region of the monoclonal antibody with nucleic acid encoding non-variable regions of equine origin in an appropriate expression vector.

20 The invention may also reside in a pharmaceutical preparation including the therapeutic agent, antibody or antibody fragment defined above in a pharmaceutically effective carrier. The formulation and preparation of any of these pharmaceutical compositions using antibodies, antibody fragments or other
25 compounds is well known to those skilled in the art of pharmaceutical formulation (see e.g. "Remington's Pharmaceutical Sciences", Sixteenth Edition, Mack Publishing Co, 1980).

By way of a shorthand notation the following three and one letter abbreviations
30 for amino acid residues are used in the specification as defined in Table 1.

Where a specific amino acid residue is referred to by its position in the polypeptide of an protein, the amino acid abbreviation is used with the residue number given in superscript (i.e. Xaaⁿ)

35

TABLE 1

	Amino Acid	Three-letter Abbreviation	One letter Abbreviation
	Alanine	Ala	A
5	Arginine	Arg	R
	Asparagine	Asn	N
	Aspartic Acid	Asp	D
	Cysteine	Cys	C
	Glutamine	Gln	Q
10	Glutamic acid	Glu	E
	Glycine	Gly	G
	Histidine	His	H
	Isoleucine	Ile	I
	Leucine	Leu	L
15	Lysine	Lys	K
	Methionine	Met	M
	Phenylalanine	Phe	F
	Proline	Pro	P
	Serine	Ser	S
20	Threonine	Thr	T
	Tryptophan	Trp	W
	Tyrosine	Tyr	Y
	Valine	Val	V

25 BRIEF DESCRIPTION OF THE DRAWINGS

For a better understanding the invention will now be described with reference to an illustrated embodiment. The drawings describe an illustrated embodiment wherein,

30 Figure 1 is a plot of the optical density for a number of peptides against positive sera. Fifty-one positive sera were used to screen fifty overlapping peptide fragments from the mature form of VapA in an ELISA. A positive result was considered to be at least twice the background OD at 450nm (background OD range 0.04 - 0.3).
35 Numbers above columns indicate the actual number of sera reacting with a given peptide; and

Figure 2 is a plot of the optical density for a number of peptides against positive sera and shows peptides used for further definition of the B-cell epitope within LQKDEPNGRASD [Seq I.D. No. 39], using peptides derived from that sequence. Positive peptides were defined as having an OD of at least 3 times the standard deviation above the lowest 25% of all OD values at 450nm wavelength (OD range 0.13-0.78). Numbers above columns indicate actual numbers of sera reacting with a given peptide. NLQKDEPNGRA [Seq I.D. No. 47] is included for comparison.

DETAILED DESCRIPTION OF THE INVENTION

Example 1 - Peptide based assay for R. equi

A potential 20 amino acid region of the VapA protein that appears to be recognised by the sera of horses that have been infected with *R. equi* and is unrecognised by control horses which have not been infected by the organism was identified.

Biotinylated peptides synthesized by Mimotopes, Victoria, Australia were used in all assays. The peptide bank used in the initial screening of sera was designed based upon the published sequence of VapA (Genbank accession No: D21236). A total of 50 overlapping peptides, each 11 amino acid residues in length (offset by 3 residues at a time) were synthesized beginning from the predicted signal peptide cleavage site between amino acids 31 and 32 up to and including the C terminus.

A second set of peptides was used to further define the region between peptides 11 to 14 (LQKDEPNGRASD [Seq I.D. No. 39]) of the VapA protein. A total of nineteen peptides were designed based upon this region, twelve were truncated peptides and contained single stepwise amino acid deletions starting from either the N or C terminal. Six peptides were overlapping 6-mers and covered the sequence offset by one residue at a time beginning at the N-terminal. A final peptide KDEPNGR [Seq I.D. No. 96] was designed based upon the core sequence of the B-cell epitope identified in assays using the previous 18 peptides.

A total of seventy foal sera, most of these from animals aged between 4 to 12 weeks, were used to screen the peptides. Fifty-one sera were from foals with

current *R. equi* disease (positive sera). Sixteen sera were from foals with no known history of *R. equi* infection and three sera were from foals that had recovered from *R. equi* infection 10 months previously (negative sera). Thirty-nine of the positive sera and all negative sera were obtained from studs in South Australia and New South Wales, Australia.

Biotin-SGSG(spacer) labelled peptide units were used as antigen targets. Neutravidin bound to the surface of a micro-titre plate was then used to bind the biotin labelled peptide to the plate. The procedure is as follows:

A solution of neutravidin was diluted 1 in 300 in sterile distilled water to a final concentration of 3.3 µg/ml, and 100 µl per well was added to a micro-titre plate and the plate was left overnight at 37°C. 200 µl 1% casein /1xPBS/ .05% Tween 20 was then added to each well and the plate was left for at least 1 hour at room temperature (or at 4° C if left longer) before the solution in the plate was flicked out. The plate was then washed five times with 1xPBS/.05%Tween 20 using an automatic plate washer and the plates were dried by slapping against absorbent paper.

The peptides were diluted to a concentration of 5 nmol/ml in 1xPBS/.05%Tween 20 and 100 µl of each peptide was added to different wells. The plate was incubated with shaking at room temperature for 1 hour before the solution in the plate was flicked out. The plate was then washed five times with 1xPBS/.05%Tween 20 using an automatic plate washer and the plates were dried by slapping against absorbent paper.

A sera samples were then diluted 1 in 250 in 1xPBS/.05%Tween20 and 100 µl was added to each well. The plate was left overnight at 4°C before the solution in the plate was flicked out. The plate was then washed five times with 1xPBS/.05%Tween 20 using an automatic plate washer and the plates were dried by slapping against absorbent paper.

Horseradish peroxidase labelled goat anti-horse IgG was diluted 1 in 5000 in 1% casein / 1xPBS/ .05% Tween 20 and 100 µl was added per well. The plate was incubated at room temperature for 1 hour before the solution in the plate was flicked out. The plate was then washed five times with 1xPBS/.05%Tween 20

using an automatic plate washer and the plates were dried by slapping against absorbent paper.

5 The plate was then washed twice with 1xPBS and 1 TMB (tetramethyl benzidine) tablet was dissolved in 10 ml phosphate citrate buffer. 100 μ l of the solution was added per well and the plate was incubated at room temperature in the dark for 15 minutes before 100 μ l 1N H₂SO₄ was added per well to stop reaction.

10 The plate was then read using an ELISA plate reader at 450nm.

Interpretation of data. In the initial assay using the fifty peptides in the overlapping bank of the entire VapA protein, a positive result was assigned by using a cut off value of twice the background OD. The background OD was the mean of the lowest 50% of all OD values obtained with that particular serum and all OD readings that were twice this value were considered positive. The background ranged from 0.04 to 0.3, indicating a high degree of variability in the reactivity of sera with the peptide bank.

20 In the assay to identify the most reactive peptides containing elements of region LQKDEPNGRASD, the cut off OD value for a positive result was determined using the mean value of the lowest 25% of all OD values obtained with that serum (range 0.13 - 0.78) plus 3 times their standard deviation. All OD readings above the cut off were considered positive.

25 A positive result in the *R. equi* whole cell ELISA was based upon twice the background OD. The background OD (0.07 to 0.37) was the OD value of the well containing all reagents and sera used in the corresponding test assay without the whole cell antigen preparation. All OD values above the background OD were considered positive.

30 The fifty-one positive sera screened against fifty peptides recognized an epitope between amino acids 62 to 81 of the VapA sequence corresponding to peptides 11 to 14 (OD values between 0.25 and 1.5). The amino acid sequences of these peptides are TSLNLQKDEPN (P 11; [Seq I.D. No. 44]), NLQKDEPNGRA (P 12; [Seq I.D. No. 47]), KDEPNGRASDT (P 13; [Seq I.D. No. 50]) and PNGRASDTAGQ (P 14; [Seq I.D. No. 43]). Peptide [Seq I.D. No. 47] was 35 universally recognized by all fifty-one sera associated with current *R. equi*

infection. Forty-nine of these sera recognized at least two peptides in this region and two sera were positive with only peptide 12. Thirteen sera were positive with all four peptides 11-14. Four sera were positive with peptides 15 or 16 in addition to reacting with at least one peptide in the region 11 to 14.

5

In addition, eleven of the positive sera also reacted positively with one or two of peptides 41 to 43 (region between amino acids 152 and 168 of VapA). The sequence of this secondary epitope did not have any similarity to the sequence encompassed by peptides 11-14. Ten of these sera gave a positive result with peptide 41 which corresponded to sequence YLNINFFDSSG [Seq I.D. No. 133] (Fig. 1).

10

All sera from animals with no known history of *R. equi* infection gave a negative result with peptides P11-P14.

15

These assays show that a major linear epitope of VapA lies in the region between peptides 11 to 14 corresponding to amino acids 62 to 81 of the VapA precursor protein sequence. Based on the universal reactivity of peptide NLQKDEPNGRA [Seq I.D. No. 47] it is likely that a B-cell epitope is in this region of VapA.

20

The region between peptides 11-14 of VapA contains predominantly hydrophilic residues and analysis of the precursor VapA sequence using the Hopp and Woods hydrophobicity algorithm indicated that the region corresponding to P11-P14 was the most hydrophilic region of the entire protein (Hopp and Woods, 1981).

25

Studies have shown VapA to be a lipid modified, hydrophobic, surface-expressed protein (Tanet *al.* 1995). Therefore it would be expected that the hydrophilic region of this protein would lie on the cell surface and consequently be more likely to interact with the host immune system. Interestingly, the minor epitope identified between peptides 41 to 43 was within the hydrophobic region of the VapA protein. This may mean that occasionally non cell-surface exposed regions of VapA do interact with the host immune system, although to a much lesser extent than the major cell surface domain of the protein.

30

Example 2 - Vaccine Development

35

The antigenic region or derivative peptide identified was synthesised with a (Histidine)₆-tag at either the N or C terminii. The 6xHis tag allows the peptide to

bind strongly with Ni-NTA (nickelnitriloacetic acid) agarose thereby greatly increasing its immunogenicity. The peptide Ni-NTA agarose was then used directly with a suitable adjuvant (Freund's) as a sub-cutaneous or intra muscular vaccine (Sheibani and Frazier, 1998).

5

This vaccine was then used to determine whether it can elicit anti VapA antibodies in mice. This was tested by use of the mouse sera in an ELISA against the peptide epitope, and in Western blots to determine whether the sera can detect the VapA protein. Challenge studies then followed using the immunised mice.

10 This entailed either intra peritoneal or aerosol inoculation of the mice with virulent *R. equi*. Protection against fatal infection was taken as good evidence that VapA is protective.

DNA Vaccine

15

pcDNA3 vectors containing *groEL* and *vapA* were used to transiently transfect Cos7 cells in order to check for expression of chimeric protein(s) in a mammalian system using Western blot analysis.

20 The DNA vaccines were then tested for antibody production in BALB/c mice (50µg DNA in 50 µl saline injected into the quadriceps muscle using gene gun delivery).

Purified GroEL and VapA proteins can also be used to vaccinate mice in the
25 murine model. Purified protein can be produced in a bacterial system using (Hisidine)₆ tagged protein and NiNTA agarose purification system. The efficacy of the vaccines can be tested by subjecting the immunized animals to an intraperitoneal challenge of 10⁷ *R. equi* organisms (clinical isolates). Appropriate controls can be used.

30

The epitope or a derivative can then be fused by epitope tagging using the modified inverse PCR mutagenesis (IPCRM) procedure as described by Gama and Breitweiser (Gama and Breitweiser, 1999). The epitope can be tagged into a hydrophilic or antibody accessible region of the GroEL protein, these regions can
35 be chosen using hydropathy plots of the protein. The mutagenised *groEL* gene can then be inserted into a pcDNA3 vaccine vector. After construction of the DNA vaccine using the mutagenised *groEL* it can be tested as described above.

It will be appreciated that this disclosure is not intended to limit the invention to the preferred embodiment or details thereof. It is intended to give an overview of the invention as conceived and other embodiments will be apparent to the skilled
5 addressee all of which fall within the spirit of the invention. In addition, reference in this specification to a document is not to be taken as an admission that the disclosure therein constitutes common general knowledge in Australia.

REFERENCES

- Babuik, L. Al. *et al.* 1998 *Curr. Top. Microbiol. Immunol.* **226**: 90-106.
- Barton, M.D. and Hughes, K.L. 1980 *Vet Bull.* **50**: 65-80.
- 5 Barton, M.D. and Hughes, K.L. 1984 *Vet Microbiol.* **9**: 65-76.
- Barton, M.D. 1992 *Equine Infectious Diseases VI* ed. Plowright W, Rossdale P.D. and Wade J.F R&W publications (Newmarket) Ltd. p77.
- Capdevila, J.A. *et al.* 1997 *J. Infect. Dis.* **29**: 535-541.
- Craig, E.A. *et al.* 1993 *Microbiol. Rev.* **57**: 402-414.
- 10 Gama, L. and Breitweiser, G.E. 1999 *BioTechniques* **26**: 814-816.
- Giguere, S. and Prescott, J.F. 1997 *Vet Microbiol* **56**: 313-334.
- Giguere, S. *et al.* 1999 *Infect Immun* **67**: 3548-3557.
- Higuchi, T. *et al.* 1997 *Equine Vet. J.* **29**: 274-278.
- Hondalus, M.K and Mosser, D.M. 1994 *Infect. Immun.* 4167-4175.
- 15 Hopp, T.P. and Woods, K.R. 1981 *Proc. Natl. Acad. Sci. USA* **78**: 3824-3828.
- Kohler and Milstein 1975 *Nature* **256**:495-497
- Kozbor *et al.* 1983 *Immunology Today* **4**:72.
- Leclerc, C. *et al.* 1997 *Cell Immunol.* **179**: 97-106.
- Lindbladh *et al.* 1993 *Trends in Biochem. Sci.* **18**:279-283.
- 20 Lowrie, D.B. *et al.* 1997 *Vaccine* **15**: 834-838.
- Lowrie, D.B. *et al.* 1999 *Nature* **400**: 269-271.
- Montgomery, D.L. *et al.* 1997 *DNA Vaccines* **74**: 195-205.
- Mustafa, A.S. *et al.* 1993 *Infect. Immun.* **61**: 5294-5301.
- Prescott, J.F. *et al.* 1996 *Equine Vet.* **28**: 344-349.
- 25 Ramsay, A.J. *et al.* 1997 *Cell Biol.* **75**: 360-363.
- Robinson, H.L. 1997 *Vaccine* **15**: 785-787.
- Sekizaki, T. *et al.* 1995 *Gene* **155**: 135-136.
- Sheibani, N. and Frazier W.A. 1998 *BioTechniques* **25**: 28-32.
- Smola, J. *et al.* 1994 *J. Appl. Bacteriol.* **77**: 325-333.
- 30 Stugnell, R.A. *et al.* 1997 *Immunol. Cell Biol.* **75**: 364-369.
- Takai, S. 1997 *Vet Microbiol* **56**: 167-176.
- Tan, C. *et al.* 1995 *Can. J. Vet. Res.* **59**: 51-59
- Tascon, R.E. *et al.* 1996 *Nature Med.* **2**: 888-898.
- Woodbury *et al.* 1999 *J Chromatogr B Biomed Sci Appl* **725**:113-37

CLAIMS

1. An isolated peptide fragment of the Vap A protein, or immunologically active homolog, derivative or analogue thereof, said peptide fragment binding antibodies specific for *R. equi* and the VapA protein.
5
2. An isolated peptide fragment as in claim 1 wherein the peptide contains an amino acid sequence of 5 or more amino acid residues that is identical to or homologous to the amino acid sequence of at least one region of the VapA protein that is responsible for immunological recognition.
10
3. An isolated peptide fragment as in claim 2 wherein the amino acid sequence includes the said 5 or more amino acid sequence plus other segments of the VapA protein or sequences homologous to the other segments of the VapA protein.
15
4. An isolated peptide fragment as in claim 2 wherein the peptide fragment has 5 or more amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof.
20
5. An isolated peptide fragment as in claim 4 wherein the peptide is part of a larger peptide that contains 5 or more amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof, as well as one or more of the amino acids either side of that sequence in the native VapA protein.
25
6. An isolated peptide fragment as in claim 4 wherein the peptide contains 7 to 30 amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof.
30
7. An isolated peptide fragment as in claim 6 wherein the peptide contains 10 to 12 amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof.
- 35 8. An isolated peptide fragment as in claim 2 wherein the peptide contains an amino acid sequence selected from the list including any one of Seq I.D. No. 1 through to Seq I.D. No. 132 inclusive.

9. An isolated peptide fragment as in claim 8 wherein the peptide contains an amino acid sequence selected from the list including Seq I.D. No. 44, Seq I.D. No. 47, Seq I.D. No. 50, and Seq I.D. No. 43.

5

10. An isolated peptide fragment as in claim 9 wherein the peptide contains an amino acid sequence of Seq I.D. No. 47.

11. An isolated peptide fragment as in claim 4 wherein the homologous peptide shares 50% homology with the peptide fragment having 5 or more amino acid residues and containing all or part of the sequence of Seq I.D. No. 1.

10

12. An isolated peptide fragment as in claim 4 wherein the homologous peptide shares 70% homology with the peptide fragment having 5 or more amino acid residues and containing all or part of the sequence of Seq I.D. No. 1.

15

13. An isolated peptide fragment as in claim 4 wherein the homologous peptide shares 90% homology with the peptide fragment having 5 or more amino acid residues and containing all or part of the sequence of Seq I.D. No. 1.

20

14. A polynucleotide that encodes a peptide fragment of the Vap A protein that binds antibodies specific for *R. equi* and the VapA protein.

15. A polynucleotide as in claim 13 wherein the peptide fragment has 5 or more amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof.

25

16. A cell transformed with a recombinant plasmid that expresses a peptide fragment of the Vap A protein that binds antibodies specific for *R. equi* and the VapA protein.

30

17. A cell as in claim 16 wherein the peptide fragment has 5 or more amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof.

35

18. A method of diagnosing a vertebrate for the presence of *R. equi* by detecting the presence of antibodies to *R. equi* in the vertebrate, the method including the steps of:
- 5 obtaining a sample of antibody containing fluid from the vertebrate,
contacting the sample with at least one peptide fragment of a region for
specific immunogenic recognition of the VapA protein, or
derivative thereof, and
assaying for the formation of the antigen:antibody complex to detect the
presence of antibodies to the VapA protein.
- 10
19. A method of diagnosing a vertebrate as in claim 18 wherein the peptide contains an amino acid sequence of 5 or more amino acid residues that is identical to or homologous to the amino acid sequence of at least one region of the VapA protein that is responsible for immunological recognition.
- 15
20. A method of diagnosing a vertebrate as in claim 19 wherein the amino acid sequence includes the said 5 or more amino acid sequence plus other segments of the VapA protein or sequences homologous to the other segments of the VapA protein.
- 20
21. A method of diagnosing a vertebrate as in claim 19 wherein the peptide fragment has 5 or more amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof.
- 25
22. A method of diagnosing a vertebrate as in claim 21 wherein the peptide is part of a larger peptide that contains 5 or more amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof, as well as one or more of the amino acids either
30 side of that sequence in the native VapA protein.
23. A method of diagnosing a vertebrate as in claim 21 wherein the peptide contains 7 to 30 amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue
35 thereof.

24. A method of diagnosing a vertebrate as in claim 23 wherein the peptide contains 10 to 12 amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof.
- 5
25. A method of diagnosing a vertebrate as in claim 19 wherein the peptide contains an amino acid sequence selected from the list including any one of Seq I.D. No. 1 through to Seq I.D. No. 132 inclusive.
- 10
26. A method of diagnosing a vertebrate as in claim 25 wherein the peptide contains an amino acid sequence selected from the list including Seq I.D. No. 44, Seq I.D. No. 47, Seq I.D. No. 50, and Seq I.D. No. 43.
- 15
27. A method of diagnosing a vertebrate as in claim 26 wherein the peptide contains an amino acid sequence of Seq I.D. No. 47.
- 20
28. A method of diagnosing a vertebrate as in claim 19 wherein the antibody containing fluid is selected from the list including serum, plasma, whole blood, cerebro spinal fluid, amniotic fluid, and synovial fluid.
29. A method of diagnosing a vertebrate as in claim 19 wherein the peptide fragment is attached or conjugated to a carrier molecule.
- 25
30. A method of diagnosing a vertebrate as in claim 19 wherein the peptide fragment is attached or conjugated to a solid support.
- 30
31. A method of diagnosing a vertebrate as in claim 30 wherein the sample is contacted with the the peptide fragment under conditions suitable for formation of an antigen:antibody complex and any component of the biological sample that is not bound to the peptide fragment on the solid support is washed or otherwise removed from the bound complex.
- 35
32. A method of diagnosing a vertebrate as in claim 31 wherein the peptide fragment is bound to the surface of a microtitre well, a serum sample suspected of containing antibody is then added, unbound sample washed away and the level of antibodies bound in the antigen:antibody complex assayed.

33. A method of diagnosing a vertebrate as in claim 19 wherein the assay for the formation of antigen:antibody complex involves adding a compound that enables detection of the antibodies which are specifically bound to the peptide.
- 5 34. A method of diagnosing a vertebrate as in claim 33 wherein the assay is an ELISA and labelled anti-antibodies that are specific for the VapA antibodies are added to bind the VapA antibodies to allow detection and quantification of the VapA antibodies.
- 10 35. A method of diagnosing a vertebrate as in claim 34 wherein the vertebrate is equine and the VapA antibodies are detected with labelled goat anti-horse IgG.
36. A method of diagnosing a vertebrate as in claim 33 wherein VapA antibodies are detected by binding with anti-antibodies specific for the VapA antibodies, and then a second labelled antibody specific for the anti-antibody is used to detect the anti-antibodies.
- 15 37. A method of diagnosing a vertebrate as in claim 36 wherein the vertebrate is equine and the VapA antibodies are detected with labelled goat anti-horse IgG and then labelled anti-goat IgG is added to bind and detect the anti-horse IgG.
- 20 38. A method of diagnosing a vertebrate as in claim 33 wherein a peptide:antibody complex is used to detect the presence of VapA antibodies.
- 25 39. A method of diagnosing a vertebrate as in claim 38 wherein the antibody that forms part of the peptide:antibody complex is labelled.
40. A method of diagnosing a vertebrate as in claim 36 wherein the anti-antibodies specific for VapA antibodies are detectably labelled by linking to an enzyme which is later exposed to its substrate so as to react with the substrate to produce a chemical moiety which can be detected.
- 30 41. A method of diagnosing a vertebrate as in claim 40 wherein the enzyme is selected from the list including malate dehydrogenase, staphylococcal nuclease, delta-V-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease,
- 35

urease, catalase, glucose-VI-phosphate dehydrogenase, glucoamylase and acetylcholine esterase.

42. A method of diagnosing a vertebrate as in claim 34 wherein the sensitivity of the method is increased by coupling the anti-antibodies specific for VapA antibodies to low molecular weight haptens and detecting the haptens by means of a second reaction.
44. A method of diagnosing a vertebrate as in claim 42 wherein the hapten is biotin which is reacted with avidin.
45. A method of diagnosing a vertebrate as in claim 34 wherein the sensitivity of the assay is increased by use of amplification strategies including substrate cycling and enzyme channelling.
46. A method of diagnosing a vertebrate as in claim 34 wherein the vertebrate is selected from the list including horses, cattle, pigs, goats and humans.
47. A method of diagnosing a vertebrate as in claim 32 wherein the vertebrate is a horse.
48. A method of diagnosing a vertebrate for the presence of *R. equi* by detecting the presence of VapA antigens in the vertebrate, the method including the steps of:
- raising an antibody specific to at least one peptide fragment of a region for specific immunogenic recognition of the VapA protein, or derivative thereof, obtaining a putative VapA antigen containing biological sample from the vertebrate,
- contacting the sample with the raised antibodies under conditions for formation of an antibody:antigen complex, and assaying for the formation of the antibody:antigen complex to detect the presence of antibodies to the VapA protein.
49. A method of diagnosing a vertebrate as in claim 48 wherein the raised antibodies are any one of polyclonal antibodies, monoclonal antibodies, chimeric antibodies, single chain antibodies, equinised antibodies or Fab fragments.

50. A method of diagnosing a vertebrate as in claim 48 wherein the antibody is raised by immunising an animal with one or more peptide fragment of a region for specific immunogenic recognition of the VapA protein, or derivative thereof, and isolating antibody producing cells from the immunised animal.
- 5
51. A method of diagnosing a vertebrate as in claim 48 wherein additional antibodies capable of binding to the VapA protein are produced in a two step procedure through the use of anti-idiotypic antibodies.
- 10 52. A method of diagnosing a vertebrate as in claim 48 wherein the step of assaying for formation of the antibody:antigen complex includes the step of separating the bound antibody:antigen complex from unbound antibody.
- 15 53. A method of diagnosing a vertebrate as in claim 52 wherein the raised antibodies, or fragments thereof are labelled, any antibody:antigen complex formed by contacting the sample with the antibody is separated from unbound antibody and the labelled antibody:antigen complex is then quantitatively assayed for the label used.
- 20 54. A method of diagnosing a vertebrate as in claim 53 wherein the raised antibodies are used in an ELISA based assay to bind the VapA protein in the sample, and a labelled antibody specific for another immuno-recognition site on VapA is used to assay for bound VapA protein.
- 25 55. A method of diagnosing a vertebrate as in claim 54 wherein the raised antibodies or fragments thereof are attached or conjugated to a solid support and after contacting the raised antibodies or fragments thereof under conditions suitable for formation of an antibody:antigen complex any component of the biological sample that is not bound to the antibodies or antibody fragments on the support is washed or otherwise separated from the bound complex.
- 30
- 35 56. A method of diagnosing a vertebrate as in claim 53 wherein the assay for the formation of antibody:antigen complex involves adding a labelled antibody specific for the VapA protein and assaying for the presence of the labelled antibody.

57. A method of diagnosing a vertebrate as in claim 56 wherein the antibodies specific for the VapA protein are detectably labelled by linking to an enzyme which is later exposed to its substrate so as to react with the substrate to produce a chemical moiety which can be detected.

5

58. A method of diagnosing a vertebrate as in claim 57 wherein the enzyme is selected from the list including malate dehydrogenase, staphylococcal nuclease, delta-V-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-VI-phosphate dehydrogenase, glucoamylase and acetylcholine esterase.

10

59. A method of diagnosing a vertebrate as in claim 56 wherein the sensitivity of the method is increased by coupling the antibodies specific for the VapA protein to low molecular weight haptens and detecting the haptens by means of a second reaction.

15

60. A method of diagnosing a vertebrate as in claim 59 wherein the hapten is biotin which is reacted with avidin.

20

61. A diagnostic kit containing one or more peptide fragment of a region for specific immunogenic recognition of the VapA protein, or derivative thereof, in conjunction with appropriate reagents.

25

62. A diagnostic kit as in claim 61 wherein a control lymphocyte extract obtained from a healthy vertebrate is included, as well as serum from the healthy vertebrate.

30

63. A diagnostic kit containing one or more antibodies, or fragments thereof, specific for one or more peptide fragment of a region for specific immunogenic recognition of the VapA protein, or derivatives thereof, in conjunction with appropriate reagents.

35

64. A method of vaccinating a vertebrate against *Rhodococcus equi*, the method including the step of administering a peptide fragment of a region for

specific immunogenic recognition of VapA, or an immunogenically active derivative or variant thereof to thereby induce an immune reaction.

65. A method of vaccinating a vertebrate as in claim 64 wherein the peptide
5 contains an amino acid sequence of 5 or more amino acid residues that is identical to or homologous to the amino acid sequence of at least one region of the VapA protein that is responsible for immunological recognition.
66. A method of vaccinating a vertebrate as in claim 65 wherein wherein the
10 amino acid sequence includes the said 5 or more amino acid sequence plus other segments of the VapA protein or sequences homologous to the other segments of the VapA protein.
67. A method of vaccinating a vertebrate as in claim 65 wherein the peptide
15 fragment has 5 or more amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof.
68. A method of vaccinating a vertebrate as in claim 67 wherein the peptide is
20 part of a larger peptide that contains 5 or more amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof, as well as one or more of the amino acids either side of that sequence in the native VapA protein.
69. A method of vaccinating a vertebrate as in claim 67 wherein the peptide
25 contains 7 to 30 amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof.
70. A method of vaccinating a vertebrate as in claim 69 wherein the peptide
30 contains 10 to 12 amino acid residues and contains all or part of the sequence of Seq I.D. No. 1, or immunologically active homolog, derivative or analogue thereof.
71. A method of vaccinating a vertebrate as in claim 65 wherein the peptide
35 contains an amino acid sequence selected from the list including any one of Seq I.D. No. 1 through to Seq I.D. No. 132 inclusive.

72. A method of vaccinating a vertebrate as in claim 71 wherein the peptide contains an amino acid sequence selected from the list including Seq I.D. No. 44, Seq I.D. No. 47, Seq I.D. No. 50, and Seq I.D. No. 42.
- 5
73. A method of vaccinating a vertebrate as in claim 72 wherein the peptide contains an amino acid sequence of Seq I.D. No. 47.
74. A method of vaccinating a vertebrate as in claim 65 wherein the peptide is cross linked to a suitable carrier to form a larger peptide or protein.
- 10
75. A method of vaccinating a vertebrate as in claim 71 wherein the carrier is a carrier protein selected from the list including bovine serum albumin, tetanus toxoid, cholera toxin and deactivated diphtheria toxin.
- 15
76. A method of vaccinating a vertebrate as in claim 71 wherein the carrier is a solid phase carrier.
77. A method of vaccinating a vertebrate as in claim 76 wherein the carrier is a Ni-NTA (nickel-nitriloacetic acid) agarose to which the peptide is bound through a (Histidine)₆-tag at either the N or the C terminus.
- 20
78. A method of vaccinating a vertebrate as in claim 76 wherein the peptide is displayed on bacteriophage particles.
- 25
79. A method of vaccinating a vertebrate as in claim 65 wherein the peptide is administered in pharmaceutical dosage form as a composition or formulation comprising an immunogenically effective amount of the peptide.
- 30
80. A method of vaccinating a vertebrate as in claim 65 wherein a progeny is passively vaccinated by maternal vaccination.
81. A method of vaccinating a vertebrate against *R. equi*, the method including the step of administering a nucleic acid sequence that encodes a peptide fragment of a region for specific immunogenic recognition of VapA so that a host cell expresses the peptide fragment to thereby induce an immune reaction.
- 35

82. A method of vaccinating a vertebrate as in claim 81 wherein the nucleic acid sequence encodes a chimeric peptide or protein in which a peptide fragment of the present invention is tagged into the protein, wherein the protein is expressed by a suitable host.

5

83. A method of vaccinating a vertebrate as in claim 82 wherein a mutagenised gene encoding the chimeric protein and expressing the peptide is inserted into a DNA vaccine vector.

10 84. A method of vaccinating a vertebrate as in claim 83 wherein the *groEL* gene is used as the vaccine candidate.

85. A method of vaccinating a vertebrate as in claim 81 wherein the method also includes the step of assaying for the presence of antibodies to VapA to ensure that there is an immune reaction to the expressed peptide.

15

86. A composition for vaccinating a vertebrate against *Rhodococcus equi*, the composition comprising at least one peptide fragment of the region for specific immunogenic recognition of VapA, and a pharmaceutically acceptable carrier

20

87. A method for treating a vertebrate infected with *Rhodococcus equi*, the method including the step of administering to the vertebrate a therapeutic agent capable of binding VapA protein, wherein the agent is an antibody or other small molecule capable of binding at least one peptide fragment of the region for specific immunogenic recognition of VapA.

25

88. A method for treating a vertebrate as in claim 87 wherein the antibody is produced by immunising a suitable animal with one or more peptides of the present invention, isolating antibody producing cells from the immunised animal, fusing the antibody producing cells with a myeloma cell to produce a pool of hybridoma cells, and screening for cells that produce the monoclonal antibody.

30

89. A method for treating a vertebrate as in claim 87 wherein the therapeutic agent is a Fab, Fv or peptide fragment of a monoclonal antibody directed to at least one peptide fragment of the region for specific immunogenic recognition of VapA.

35

90. A method of screening for compounds capable of binding at least one specific immunogenic region of the VapA protein, the method including the step of screening for compounds that bind at least one peptide fragment of the region for specific immunogenic recognition of VapA.

5

91. A method of screening as in claim 90 wherein the method includes the steps of passing a scramble of randomly synthesised compounds through a solid matrix to which at least one peptide fragment of the region for specific immunogenic recognition of VapA is bound, washing so that only the strongly
10 binding compounds remain, eluting and characterising the strongly binding compounds.

92. A method of screening as in claim 91 wherein the screening is a competitive binding screen used to identify compounds that bind the peptide
15 fragment in preference to a monoclonal antibody specific for that peptide fragment.

1/1




FIGURE 1




FIGURE 2

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU01/00478

A. CLASSIFICATION OF SUBJECT MATTER		
Int. Cl. ⁷ : C07K 7/06, 7/08, 14/195, 16/12; A61K 39/02; G01N 33/53, 33/551, 33/68		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
CA: Seq. Id's 117-132 and keywords: Rhodococcus, equine, vap a, virulence associated		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CA 2,125,426 A1 (PRESCOTT, John F. <i>et al.</i>) 9 December 1995 see whole document	1-92
X	TAN, C. <i>et al.</i> "Molecular Characterization of a Lipid-modified Virulence-associated Protein of <i>Rhodococcus equi</i> and its potential in Protective Immunity", <i>Can. J. Vet Res</i> 1995; 59 pp. 51-59 see whole document	1-92
X	SEKIZAKI, Tsutomu <i>et al.</i> "Sequence of the <i>Rhodococcus equi</i> gene encoding the virulence-associated 15—17-kDa antigens", <i>Gene</i> , 155 (1995); 59 pp. 135-136 see whole document	1-92
<input type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex		
* Special categories of cited documents:		
"A"	document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed	
Date of the actual completion of the international search	Date of mailing of the international search report	
13 June 2001	15 June 2001	
Name and mailing address of the ISA/AU	Authorized officer	
AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaaustralia.gov.au Facsimile No. (02) 6285 3929	DAVID GRIFFITHS Telephone No : (02) 6283 2628	

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/AU01/00478

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report	Patent Family Member
CA 2,125,426	NONE

END OF ANNEX

专利名称(译)	VapA蛋白的抗原肽片段及其用途		
公开(公告)号	EP1276757A4	公开(公告)日	2004-11-24
申请号	EP2001925213	申请日	2001-04-27
[标]申请(专利权)人(译)	南澳大学 梅德维特科学股份有限公司 农村INDS RES & DEV		
申请(专利权)人(译)	南澳大利亚大学 MEDVET科学PTY. LTD. 农业工业研究与发展公司		
当前申请(专利权)人(译)	南澳大利亚大学 MEDVET科学PTY. LTD. 农业工业研究与发展公司		
[标]发明人	HEUZENROEDER MICHAEL W		
发明人	VANNIASINKAM, THIRUSCHOOL PHARMACY MED. LAB. SCI. BARTON, MARYSCHOOL OF PHARMACY AND MED. LAB. SCI. HEUZENROEDER, MICHAEL W.		
IPC分类号	A61K39/00 A61K39/05 C07K14/34 C07K16/12 G01N33/569 C07K7/06 C07K7/08 C07K14/195 A61K39/02 G01N33/53 G01N33/551 G01N33/68		
CPC分类号	A61K39/00 A61K39/05 A61K2039/505 A61K2039/53 A61K2039/6087 C07K14/34 C07K16/1267 C07K2317/34 G01N33/56911		
优先权	2000PQ7120 2000-04-27 AU		
其他公开文献	EP1276757A1		
外部链接	Espacenet		

摘要(译)

VapA蛋白的分离肽片段，该片段结合对马红球菌和VapA蛋白具有特异性的抗体。在优选的形式中，该肽包含具有5个或更多个氨基酸残基的氨基酸序列，其与负责免疫识别的VapA蛋白的至少一个区域的氨基酸序列相同或同源。还要求保护使用该肽诊断脊椎动物的马鞭毛虫的存在的方法，以及使用该肽对脊椎动物进行针对马鞭毛虫的疫苗接种的方法。