



(11) **EP 1 664 325 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
08.10.2014 Bulletin 2014/41

(51) Int Cl.:
C12Q 1/00 (2006.01) **C12Q 1/34** (2006.01)
G01N 33/53 (2006.01) **G01N 33/68** (2006.01)

(21) Application number: **04782672.2**

(86) International application number:
PCT/US2004/028238

(22) Date of filing: **30.08.2004**

(87) International publication number:
WO 2005/021779 (10.03.2005 Gazette 2005/10)

(54) **MASS SPECTROMETRY METHODS FOR SIMULTANEOUS DETECTION OF METABOLIC ENZYME ACTIVITY AND METABOLITE LEVELS**

MASSENSPEKTROMETRIEVERFAHREN ZUM GLEICHZEITIGEN NACHWEIS VON
METABOLISCHER ENZYMAKTIVITÄT UND METABOLITENSPIEGELN

PROCEDES DE SPECTROMETRIE DE MASSE DESTINES A LA DETECTION SIMULTANEE DE
L'ACTIVITE METABOLIQUE D'ENZYMES ET DE NIVEAUX DE METABOLITES

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PL PT RO SE SI SK TR**

(30) Priority: **29.08.2003 US 652732**

(43) Date of publication of application:
07.06.2006 Bulletin 2006/23

(73) Proprietor: **PerkinElmer LAS, Inc.**
Boston, MA 02118 (US)

(72) Inventor: **CERDA, Blas**
Milford, MA 01757 (US)

(74) Representative: **Høiberg A/S**
St. Kongensgade 59 A
1264 Copenhagen K (DK)

(56) References cited:
US-A- 5 629 210 US-A- 5 719 035
US-B1- 6 258 605 US-B1- 6 455 321
US-B1- 6 670 194

- **RASHED M S:** "Clinical applications of tandem mass spectrometry: ten years of diagnosis and screening for inherited metabolic diseases" **JOURNAL OF CHROMATOGRAPHY, BIOMEDICAL APPLICATIONS, ELSEVIER, AMSTERDAM, NL, vol. 758, no. 1, 5 July 2001 (2001-07-05), pages 27-48, XP004248035 ISSN: 0378-4347**
- **IM W.B. ET AL.:** 'Bacterial degradation of biotin' **J. BIOL. CHEM. vol. 248, no. 22, pages 7798 - 7805, XP002983412**
- **GALLIKIENLE M ET AL:** "Evaluation of enzyme activities by gas chromatography-mass spectrometry: HMGCoA reductase and cholesterol 7alpha-hydroxylase", **27 April 1984 (1984-04-27), JOURNAL OF CHROMATOGRAPHY, ELSEVIER SCIENCE PUBLISHERS B.V, NL, PAGE(S) 267 - 276, XP026475009, ISSN: 0021-9673 [retrieved on 1984-04-27]**
- **L.M. BENSON ET AL.:** "Simultaneous structure-activity determination of disulfiram photolysis products by on-line continuous-flow liquid secondary ion mass spectrometry and enzyme inhibition assay", **JOURNAL OF CHROMATOGRAPHY A, vol. 693, 1995, pages 162-166, Amsterdam**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 664 325 B1

Description

BACKGROUND OF THE INVENTION

5 [0001] The present Invention relates generally to diagnostic medicine and more specifically to methods for diagnosing metabolic disorders.

[0002] Metabolism is the process of building the body's molecular structures from nutrients (anabolism) and breaking them down for energy (catabolism). Metabolic processes result in growth, produce energy, eliminate wastes, and control distribution of nutrients in the body. Homeostasis, or a steady state, in the body is a result of normal metabolism.

10 [0003] Enzymes play an important role in metabolism because they catalyze conversion of one molecule to another during metabolic processes. When a metabolic enzyme is faulty or present in an abnormal amount in an individual, a metabolic disorder can result. Metabolic disorders are frequently due to genetic inheritance that leads to absence or overproduction of an enzyme or production of a faulty enzyme.

15 [0004] As an example, deficiencies produced by an inactive gene can prevent the body from making an enzyme (or enzymes) needed to break down certain amino acids or types of fats. Such an enzyme deficiency typically prevents normal metabolism of a nutrient, causing the nutrient or its metabolite to build up in the body to toxic levels. An enzyme deficiency can also cause nutrient deficiencies when an enzyme is unable to make a normal end-product of a metabolic process. Nutrient deficiencies resulting from metabolic disorders can lead to impaired growth and development, and other severe health problems.

20 [0005] Various analysis and screening methods for detecting diseases are known.

[0006] Clinical screening applications for inherited diseases using tandem mass spectrometry are described by Rashed M. S. in "Clinical applications of tandem mass spectrometry: ten years of diagnosis and screening for inherited metabolic diseases", Journal of Chromatography B, vol. 758, 2001, pages 27-48. The document does not disclose the simultaneous detection of metabolic enzyme activity and analyte level.

25 [0007] Methods for screening newborns using electrospray tandem mass spectrometry for the detection of metabolites are disclosed in US 6,258,605. The document does not disclose the simultaneous detection of metabolic enzyme activity and analyte level.

[0008] Methods for assaying enzyme activity and detecting fluorescent readout are disclosed in US 5,719,035. The simultaneous detection of metabolic enzyme activity and analyte level is not described therein.

30 [0009] Various analyses of biotin metabolites are disclosed by IM W. B., et al. in "Bacterial Degradation of Biotin", J. Biol. Chem., vol. 248, no. 22, 1973, pages 7798-7805. The disclosure is limited to showing 'mass spectra of methyl esters of Catabolite 2-B-1 and d-bisnorbiotin' in Figure 6.

[0010] Methods of screening newborns using electrospray tandem mass spectrometry for the detection of metabolites are disclosed in US 6,455,321. The document does not disclose the simultaneous detection of metabolic enzyme activity and analyte level.

35 [0011] A rapid screening test for Smith-Lemli-Opitz Syndrome is disclosed in US 5,629,210. The document does not disclose the simultaneous detection of metabolic enzyme activity and analyte level.

[0012] A rapid quantitative analysis of proteins or protein function in complex mixtures is described in US 6,670,194. The document does not disclose the simultaneous detection of metabolic enzyme activity and analyte level.

40 [0013] SIM assays specifically designed for HMGCoA reductase in purified rat liver microsomes are disclosed by M. Galli-Kienle et al. in "Evaluation of enzyme activities by gas chromatography-mass spectrometry: HMGCoA reductase and cholesterol 7 α -hydrolase" Journal of Chromatography, vol. 289, 1984, pages 267-276. The document does not disclose the simultaneous detection of metabolic enzyme activity and analyte level.

45 [0014] Assays of purified yeast aldehyde dehydrogenase activity are disclosed by L. M. Benson et al. in "Simultaneous structure-activity determination of disulfiram photolysis products by on-line continuous-flow liquid secondary ion mass spectrometry and enzyme inhibition assay", Journal of chromatography, vol. 693, 1995, pages 162-166. The document does not disclose the simultaneous detection of metabolic enzyme activity and analyte level.

50 [0015] Many inherited metabolic disorders are fatal in the first weeks or months of postnatal life, for example, severe defects in the conversion of pyruvate to acetyl coenzyme A (CoA), some urea cycle defects, and severe defects in the processing of fructose. Infants and children with a treatable metabolic disorder are often identified by newborn screening in developed countries. For example, testing for phenylketonurea is routine practice in developed countries. Phenylketonurea is an example of a relatively common inherited metabolic disorder that occurs in about one out of 16,000 live births in the United States. Individuals having phenylketonuria do not produce the enzyme necessary to break down phenylalanine (an amino acid). Fortunately, when recognized, this metabolic disorder can be successfully treated by dietary restriction.

55 [0016] Unfortunately, newborns are not typically screened for other metabolic disorders such as homocysteinuria, maple syrup urine disease, organic acid disorders, and disorders of fatty acid oxidation. As a result, these disorders are often detected in infants and children after damage has occurred and effects such as developmental delay and mental

retardation become apparent. However, such effects often can be reduced or avoided with early detection and sustained dietary restriction. Such early detection involves examining a blood sample for an enzyme activity or a metabolic marker (a metabolite).

[0017] Wide-spread newborn testing for multiple inherited metabolic disorders has been unavailable due, in part, to cost associated with each test. One approach for reducing costs is to examine multiple enzyme activities and metabolites and in a single test. However, development of multi-enzyme/metabolite testing has been hindered by the different conditions needed for enzyme tests in comparison to metabolite tests. Currently, enzyme and metabolite tests cannot be carried out simultaneously from a single patient sample.

[0018] Thus, there exists a need for methods for efficiently diagnosing a metabolic disorder in an individual. The present invention satisfies this need and provides related advantages as well.

SUMMARY OF THE INVENTION

[0019] The present invention provides a method for detecting a metabolic disorder in an individual.

[0020] The method involves (a) contacting a sample to determine the presence of (i) one or more metabolically indicative enzymes which have an altered activity as a result of a metabolic disorder and (ii) one or more metabolic analytes, which have an altered amount as a result of a metabolic disorder, with one or more substrates for said one or more enzymes to produce a reaction admixture, under conditions in which at least one of said enzymes is capable of acting on a corresponding substrate to generate at least one product; (b) contacting said reaction admixture with a reagent that inhibits the ability of said one or more enzymes to act on a corresponding substrate, wherein said one or more metabolic analytes and said at least one product are soluble in said reagent; to produce a test sample and (c) determining the amount or presence of said one or more metabolic analytes and said at least one product contained in said test sample using mass spectrometry, wherein a determined amount or presence of said one or more metabolic analytes and said at least one product correlates with presence or absence of said metabolic disorder. A sample used in a method of the invention can be, for example, a body fluid sample, such as blood, and can be obtained from a neonate, newborn, child or adult. In one embodiment, the method can include, in step (a) contacting the sample with one or more reference substrates. In another embodiment, the method can include, in step (b), contacting the sample with one or more reference products. In a further embodiment, the method can include, in step (d), prior to determining, adding one or more reference products corresponding to the at least one product. In yet another embodiment, the method can include, in step (d), prior to determining, adding one or more reference analytes corresponding to the one or more metabolic analytes contained in the sample.

[0021] The invention provides another method for detecting a metabolic disorder in an individual where the one or more substrates are present in dried form in a vessel.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022] Figure 1 shows simultaneous detection of biotinidase activity and metabolic analytes (amino acids and carnitines) using a method of the invention.

DETAILED DESCRIPTION OF THE INVENTION

[0023] This invention relates to the discovery of metabolic disorder diagnostic tests that allow detection of at least two relevant markers simultaneously and from a single sample. In particular, the inventor has discovered that one sample can be used to simultaneously detect a metabolic analyte, as well as activity of a metabolic enzyme, using mass spectrometry.

[0024] The present invention involves detecting at least two markers or indicia of one or more metabolic disorders from a single sample. At least one index corresponds to the activity of a metabolically indicative enzyme in the sample, and at least one index corresponds to the presence or amount of a metabolic analyte endogenously contained in the sample. Previous clinical methods for genetic screening using mass spectrometry detected only metabolic analytes in a single sample. In particular, Chace (US 6,258,605 and US 6,455,321) describes preparing a patient sample (dried blood spot) by extraction in methanol prior to analysis. Under such conditions, enzymes are inactive such that measuring enzyme activity is unfeasible, moreover, un contemplated.

[0025] Conversely, well-known methods for determining enzymatic activity do not include detection of metabolic analytes but instead measure a tag on the enzymatic product, such as a fluorescent tag. Modes of detection for tagged products cannot simultaneously detect metabolic analytes present in a patient sample. Another disadvantage of fluorimetric methods with respect to enzymatic assays is that artificial substrates tagged with a fluorescent group are different enough from their natural counterparts that enzyme activity toward such substrates is inherently reduced and thus sensitivity of the assays is reduced. The methods of the invention advantageously can involve detection of metabolically

indicative enzyme activity toward natural substrate.

[0026] As described in Example I, the activity of metabolically indicative enzyme biotinidase, as well as the concentration of alpha amino acids and carnitines, were simultaneously determined from a single sample. As is shown in Figure I, in spite of its multiplex character, the sensitivity of the method for detecting biotinidase (Fig. 1) is comparable to a well-known fluorimetric enzyme assay method. Similarly, as shown in Tables 3 and 4, a method of the invention has sensitivity comparable to well-known mass spectrometry methods that measure analytes without regard for enzyme activities. Based on this discovery of simultaneous detection of metabolic analytes and enzyme activity from a single sample, the invention provides methods for detecting metabolic disorders.

[0027] Detecting of a metabolic disorder can be used in diagnosing or predicting susceptibility to a metabolic disorder because biochemical indicators examined using the methods of the invention are indicative of a metabolic disorder, whether or not physiologic or behavioral symptoms of the disorder have become apparent.

[0028] The methods of the invention also are useful for monitoring the metabolism of an individual, such as one undergoing treatment for a metabolic disorder. As a non-limiting example, the methods can be used for determining therapeutic efficacy of a particular treatment. Based on this determination, the individual can be offered additional or alternative therapeutic options. The methods of the invention also can be useful for assessing patient compliance with a particular treatment modality, such as dietary restriction. Therefore, the invention is applicable to screening, diagnosis, prognosis, monitoring therapy and compliance, and any other application in which determining the presence or amount of one or more metabolic analytes and at least one metabolically indicative enzyme activity is useful.

[0029] The methods of the invention involve simultaneously detecting one or more analytes and one or more metabolically indicative enzymes. By providing a multiplexed format, the invention advantageously reduces technical performance errors because the number of manipulations is fewer than tests performed separately. Moreover, assay costs associated with simultaneous detection of multiple analytes and enzyme activities is lower than those associated with individual tests because fewer materials (assay vessels, liquid transfer materials, and the like) are used and less operator time is required.

[0030] As used herein, the term "metabolic disorder" means any abnormal condition in the body that causes loss of metabolic control of the body's steady state. Metabolism is the physiological and biochemical process by which food is converted in the body to forms that provide energy for bodily activities. Metabolism includes creation of specific molecules that the body uses for its myriad activities. Such molecules include hormones, neurotransmitters, proteins such as enzymes, and membrane constituents. Metabolism also includes degradation processes that enable cells to excrete waste products. Exemplary metabolic processes include processes for absorption and modification of vitamins and minerals; for degrading molecules to provide energy or to be excreted; processes for making acetyl-coenzyme A, non-essential amino acids, cholesterol, long-chain fatty acids, prostaglandins, complex lipids and proteins; and for neutralizing toxins. Therefore, a metabolic disorder is a condition that interferes with normal creation or destruction of biological molecules that regulate health. A metabolic disorder arising early in life and resulting from genetic inheritance is generally referred to as an "inborn error of metabolism", whereas a metabolic disorder that arises as a result of disease or lifestyle is generally referred to as an "acquired metabolic disorder", although genetic predisposition generally exists.

[0031] Elevated amino acids, free carnitine and acylcarnitine levels are examples of metabolites that can be indicative of one or more of several metabolic disorders. Free carnitine and acylcarnitines are markers for disorders that are classified as fatty acid oxidation (FAO) disorders and organic aciduria disorders (OAD). Similarly, amino acids are used as markers for several metabolic disorders collectively known as amino acidopathies.

[0032] In Fatty Acid Oxidation (FAO) Disorders disorders, enzymes necessary for fatty acid breakdown are unavailable or have reduced activity. Breakdown, or oxidation, of fatty acids is necessary for energy production when glucose levels, the body's main source of energy, are low. Without this energy supply some individuals may have recurring incidences of low blood sugar levels. In cases of fasting, often caused by illnesses such as ear infections or flu, there may be metabolic crisis. Affected individuals may show vomiting, diarrhea, lethargy, seizures and coma. Failure to diagnose FAO disorders may result in excessive fat buildup in the liver, heart and kidneys. This buildup can cause a variety of symptoms, ranging from hepatic failure, encephalopathy, and heart and eye complications to general problems with muscle development. Many of these clinical symptoms can lead to death. Many deaths due to FAO disorders have been misdiagnosed as SIDS or Reye's Syndrome.

[0033] In Organic Aciduria (OA) Disorders, the metabolic pathways of organic acids are disrupted in OA disorders and thus accumulation of the acids in blood and urine alters the acid-base balance of the body. Resulting modifications or adaptations to intermediary metabolic pathways may cause numerous clinical symptoms, including metabolic acidosis, ketosis, hyperammonemia, failure to thrive, sepsis or coma.

[0034] In disorders of amino acid metabolism (amino acidopathies), enzymes necessary for the metabolism of certain amino acids are unavailable or have reduced activity. As a result, the concentration of the affected amino acids and alternate metabolites increases in the infant's body. These excesses can have deleterious effects on the infant's health including death. Some commonly studied amino acidopathies are:

[0035] Phenylketonuria (PKU) is a disorder of aromatic amino acid metabolism in which phenylalanine cannot be

converted to tyrosine. If untreated, PKU leads to various degrees of mental retardation. Hyperphenylalaninemia leads to mental retardation and muscular rigidity. Homocystinuria leads to vascular occlusive disease, osteoporosis, accumulation of homocystine and methionine, and variable developmental delays. Maple Syrup Urine Disease (MSUD) is caused by a disorder of branched-chain amino acid metabolism resulting in elevated levels of leucine, isoleucine and valine in the blood. If untreated, lethargy progressive to coma, developmental delay, and convulsions will develop. Tyrosinemia type I (hereditary tyrosinemia), leads to acute hepatic failure or chronic cirrhosis and hepatocellular carcinoma. Citrullinemia leads to convulsions, anorexia, vomiting and lethargy, followed rapidly by potentially lethal coma.

[0036] As used herein, the term "metabolically indicative" when used in reference to an enzyme means that the amount or activity of the enzyme is altered as a result of a metabolic disorder in an individual. Therefore, the amount or activity of a metabolically indicative enzyme can be increased or decreased in an individual having a metabolic disorder in comparison to an individual free of the metabolic disorder. Although the activity of a metabolically indicative enzyme can be altered in an individual having a metabolic disorder, it is understood that the individual can be free of, or can have outward signs or symptoms of, the disorder.

[0037] As used herein, the term "metabolic analyte" means a substance that is altered in amount, activity, or physical characteristic in the body of an individual experiencing a metabolic disorder in comparison to an individual not experiencing a metabolic disorder. Exemplary metabolic analytes include an amino acid, fatty acid and organic acid, and other small biological molecules, as well as a macromolecule such as a nucleic acid, polypeptide or carbohydrate, and the like.

[0038] The invention provides methods for detecting a metabolic disorder in an individual.

[0039] One method involves (a) contacting a sample to determine the presence of (i) one or more metabolically indicative enzymes which have an altered activity as a result of a metabolic disorder and (ii) one or more metabolic analytes, which have an altered amount as a result of a metabolic disorder, with one or more substrates for said one or more enzymes to produce a reaction admixture, under conditions in which at least one of said enzymes is capable of acting on a corresponding substrate to generate at least one product; (b) contacting said reaction admixture with a reagent that inhibits the ability of said one or more enzymes to act on a corresponding substrate, wherein said one or more metabolic analytes and said at least one product are soluble in said reagent; to produce a test sample and (c) determining the amount or presence of said one or more metabolic analytes and said at least one product contained in said test sample using mass spectrometry, wherein a determined amount or presence of said one or more metabolic analytes and said at least one product correlates with presence or absence of said metabolic disorder.

[0040] In some embodiments, the invention provides a method for detecting a metabolic disorder, wherein the one or more substrates are present in dried form in a vessel.

[0041] In some embodiments, the invention provides a method for detecting a metabolic disorder, wherein the reaction admixture is contacted with one or more reference products and one or more reference analytes and the amount of said one or more metabolic analytes and said at least one product contained in said test samples is determined with respect to said one or more reference products and said one or more reference analytes, using mass spectrometry.

[0042] In some embodiments, the invention provides a method for detecting a metabolic disorder, wherein said vessel is contacted with a solution to generate said conditions in which at least one of said enzymes is capable of acting on a corresponding substrate to generate at least one product.

[0043] In some embodiments, the invention provides a method for detecting a metabolic disorder, wherein the sample is separated into two sample portions and each sample portion is subjected to steps (a) and (b); combining the two reaction admixtures to produce a test sample; and subjecting the combined reaction admixtures to step (c).

[0044] In some embodiments, the invention provides a method for detecting a metabolic disorder, wherein the sample is separated into two sample portions, wherein step (a) comprises contacting each sample portion with an individual vessel, each vessel containing said one or more substrates for said one or more enzymes in dried form; subjecting each sample portion to step (b); combining the two reaction admixtures to produce a test sample; and subjecting the combined reaction admixtures to step (c).

[0045] In any of the above methods for detecting a metabolic disorder, one or more reference products and/or reference analytes can be added to a test sample prior to mass spectrometry. Similarly, one or more reference substrates can be added to a reaction admixture or solution before an enzymatic reaction is allowed to proceed. Such additions can be useful when the methods are carried out in a quantitative format, as is described in more detail herein below.

[0046] The methods of the invention can be carried out in a variety of formats. As an example of a typical procedure, a sample from a normal individual or one suspected of having a metabolic disorder is placed in a vessel. The type of the sample (for example, whole blood, serum, tissue) is chosen based on the particular metabolic disorder, and any sample pre-treatment to release enzymes is performed based on the sample type. An enzyme that correlates with the test subject metabolic disorder is known, as is the enzyme substrate. At least one metabolic analyte known to correlate with the metabolic disorder, or a related or different metabolic disorder, is also known.

[0047] To the sample is added an aqueous solution containing a substrate for the enzyme indicative of the metabolic disease. The enzymatic reaction is allowed to proceed for a time period sufficient for the enzyme to produce a detectable or desired amount of product. The reaction is then stopped by adding a stop reagent that includes a reference analyte

to be detected. The reagent denatures the enzyme, but maintains the product and reference form of metabolic analyte in solution (reference analyte). The sample is subjected to tandem mass spectrometry to detect the product, metabolic analyte, and reference analyte. The amount of metabolic analyte is determined based on the reference analyte, and the presence of the product is detected.

[0048] As another example, a sample from a normal individual or one suspected of having a metabolic disorder is added to a vessel containing the metabolically indicative enzyme substrate. If the sample is contained in a suitable solution, the enzymatic reaction is allowed to proceed. If the sample is in dried form, such as on membrane or paper, a suitable solution is applied to the sample when in the vessel. After a time period sufficient to solubilize the enzyme or allow enzyme to be released from the material on which it is dried (if present), the enzymatic reaction is allowed to proceed. As above, the enzymatic reaction is allowed to proceed for a time period sufficient for the enzyme to produce a detectable or desired amount of product. The reaction is then stopped by adding a stop reagent that includes a reference analyte and a reference product. The amounts of metabolic analyte and enzymatic product are determined based on the reference analyte and product.

[0049] As a further example of a procedure for performing a method of the invention, after addition of the stop reagent, the test sample can be processed to separate soluble from insoluble material. Separation can be performed, for example, by centrifugation, filtration and the like. The resultant clarified test sample can then be subjected to mass spectrometry. Another optional step is to add a detergent or other solubilizing agent to the test sample, for example in the case that the reaction admixture and reagent are immiscible.

[0050] Although not required, the test sample can be derivatized prior to mass spectrometry to enhance detection of the amount of metabolic analyte, enzymatic product and/or reference substance. An exemplary well-known derivatization method is reaction of sample with acid and butanol.

[0051] A method of the invention can be used to simultaneously determine presence or amount of one or more metabolic analytes and at least one enzymatic product. As such, the methods are applicable to simultaneously determining presence or amount of two or more metabolic analytes and at least one enzymatic product, three or more metabolic analytes and at least one enzymatic product, four or more metabolic analytes and at least one enzymatic product, including at least five metabolic analytes, at least five metabolic analytes, at least ten metabolic analytes and at least twenty metabolic analytes, together with a at least one enzymatic product. Similarly, the methods are applicable to simultaneously determining presence of amount of one or more metabolic analytes and two or more enzymatic products, one or more metabolic analytes and three or more enzymatic products, one or more metabolic analytes and four or more enzymatic products, one or more metabolic analytes and five more enzymatic products, as well as one or more metabolic analytes and more than five enzymatic products.

[0052] The methods of the invention involve detecting a metabolic disorder by testing a "sample." As used herein, the term "sample" means any biological fluid, cell, tissue, or fraction thereof, that includes one or more enzymes having an endogenous enzyme indicative of a metabolic state, and one or more endogenous metabolic analytes. A sample can be, for example, a specimen obtained from an individual or can be derived from such a specimen. For example, a sample can be a tissue section obtained by biopsy, or cells that are placed in or adapted to tissue culture. Exemplary samples therefore include cultured fibroblasts, cultured amniotic fluid cells, and chorionic villus sample. A sample can also be a biological fluid specimen such as urine, blood, plasma, serum, saliva, semen, sputum, cerebral spinal fluid, tears, mucus, and the like. A sample can be further fractionated, if desired, to a fraction containing particular cell types. For example, a blood sample can be fractionated into serum or into fractions containing particular types of blood cells such as red blood cells or white blood cells (leukocytes). If desired, a sample can be a combination of samples from an individual such as a combination of a tissue and fluid sample, and the like.

[0053] For used in the methods of the invention, a sample contains one or more metabolically indicative enzymes and one or more endogenous metabolic analytes. To allow enzyme activity to occur in solution while practicing the methods of the invention, the sample can be suitably processed to release metabolically indicative enzyme(s) and/or endogenous metabolic analytes into solution. Well known methods for releasing or solubilizing enzymes in samples containing cells include treatment with detergent, freeze/thaw treatment, drying and rehydrating, and the like.

[0054] The origin of a sample used in a method of the invention will depend upon the metabolically indicative enzyme(s) and metabolic analyte(s) to be detected. The body fluids and tissues in which metabolically indicative enzymes and metabolic analytes are contained are well known to those skilled in the art. As non-limiting examples, a whole blood sample can be used when biotinidase is the metabolically indicative enzyme; a liver biopsy can be used when carbamyl-phosphate synthetase is the metabolically indicative enzyme; urine can be used when branched-chain ketoacid dehydrogenase is the metabolically indicative enzyme and a skin biopsy can be used when alpha-L-fucosidase is the metabolically indicative enzyme.

[0055] Methods for obtaining samples that preserve the activity or integrity of molecules in the sample are well known to those skilled in the art. Such methods include the use of appropriate buffers and/or inhibitors, including nuclease, protease and phosphatase inhibitors, that preserve or minimize changes in the molecules in the sample. Such inhibitors include, for example, chelators such as ethylenediamine tetraacetic acid (EDTA), ethylene glycol bis(P-aminoethyl

ether)N,N,N1,N1-tetraacetic acid (EGTA), protease inhibitors such as phenylmethylsulfonyl fluoride (PMSF), aprotinin, leupeptin, antipain and the like, and phosphatase inhibitors such as phosphate, sodium fluoride, vanadate and the like. Appropriate buffers and conditions for isolating molecules are well known to those skilled in the art and can be varied depending, for example, on the type of molecule in the sample to be characterized (see, for example, Ausubel et al.

[0056] Various methods to fractionate a fluid sample or cell extract are well known to those skilled in the art, including subcellular fractionation or chromatographic techniques such as ion exchange, hydrophobic and reverse phase, size exclusion, affinity, hydrophobic charge-induction chromatography, and the like (Ausubel et al. supra., 1999; Scopes, Protein Purification: Principles and Practice, third edition; Springer-Verlag, New York (1993); Burton and Harding, J. Chromatogr. A 814:71-81 (1998)). As an example, a serum sample from an individual containing leukocytes can be fractionated to isolate leukocytes, if desired, or subfractionated, for example, into macrophages, T cells, B cells, granulocytes, monocytes, neutrophils, eosinophils, basophils, mast cells, and the like.

[0057] For use in a method of the invention, a sample can be in a variety of physical states. For example, a sample can be a liquid or solid, can be dissolved or suspended in a liquid, can be in an emulsion or gel, and can be absorbed onto a material. As a non-limited example, a sample can be a liquid blood sample, liquid serum sample, liquid white blood cell sample, dried blood, serum, or white cell sample, or such a sample absorbed onto a paper or polymer substrate. Example 1 describes use of a dried blood sample in a method of the invention.

[0058] A sample used in a method of the invention can be separated prior to use. Separation can be desirable if particular metabolically indicative enzymatic activities present in the sample require distinct assay conditions, or if particular metabolic analytes endogenously present in the sample are soluble in different solution environments. Separation also can be desirable to reduce the presence of substances that interfere with detection of a metabolically indicative enzyme or metabolic analyte in a sample, and to retain that interfering portion of the sample in order to detect one or more distinct metabolically indicative enzymes or metabolic analytes in the portion.

[0059] Individuals of all ages can be affected by a metabolic disorder. Therefore, a sample used in a method of the invention can be obtained from an individual of any age, including a neonate, newborn, baby, child, and adult. Exemplary metabolic disorders that can be characterized by adult onset include Schindler disease (lysosomal enzyme alpha-N-acetylgalactosaminidase deficiency), urea cycle disorders and sphingolipidoses. The methods of the invention are applicable to human and non-human individuals. As such, the methods can be used on a sample obtained, for example, from any veterinary or research subject for which a metabolically indicative enzyme and endogenous metabolic analyte are known. Exemplary non-human animals include a horse, dog, cat, rabbit, rat, mouse, fish, turtle and lizard. The methods of the invention also are applicable to lower organisms, such as yeast, archeobacteria and bacteria, and plants, so long as a sample containing an endogenous metabolically indicative enzyme and endogenous metabolic analyte can be obtained.

[0060] The methods of the invention involve detecting the enzymatic product of at least one "metabolically indicative enzyme." Any metabolically indicative enzyme can be assayed using the methods of the invention, so long as a substrate is known for the enzyme, which substrate is soluble in a selected "reagent," and which substrate can be modified to a form distinctly detectable by mass spectrometry, if desired, for use as an internal standard. Metabolically indicative enzymes are represented in every enzyme class, including oxidoreductases, hydrolases, lyases, transferases, ligases and isomerases. As a non-limiting example, well-known metabolically indicative enzymes and their associated metabolic disorders are shown below in Table 1. A specific example of a metabolically indicative enzyme is biotinidase.

Table 1:

Metabolic disorder	Metabolically indicative enzyme	Exemplary metabolic analyte
3-methylglutaconic aciduria	3-methylglutaconic hydratase	3-methylglutaconic acid
Smith-Lemli Opitz Syndrome		7-dehydrocholesterol
4-hydroxybutyric aciduria	Succinic semialdehyde dehydrogenase	
Adenylosuccinate lyase deficiency	Adenylosuccinate lyase	succinylAICA nucleoside and succinyladenosine

EP 1 664 325 B1

(continued)

	Metabolic disorder	Metabolically indicative enzyme	Exemplary metabolic analyte
5	Beta-Ketothiolase Deficiency	3 oxoacid CoA transferase and beta-ketothiolase	
	Biotinidase deficiency	Biotinidase	Biotin
10	Carbamylphosphate synthetase deficiency	Carbamylphosphate synthetase	
	Duarte Galactosemia	Gal-1-P-uridylyltransferase	Gal 1 P
15	Ehlers-Danlos Syndrome, Type VI	Lysyl hydroxylase	pyridinoline (Pyr), and deoxypyridinoline, (Dpyr) and their ratio
	Fucosidosis	alpha-L-fucosidase	
20	Guanidinoacetate Methyltransferase Deficiency	Guanidinoacetate methyltransferase	Guanidinoacetate
	Krabbe Disease	Beta-Galactocerebrosidase	
25	Long Chain Acyl-CoA Dehydrogenase Deficiency	Acyl-CoA Dehydrogenases Short-, Medium- and Long Chain	
	Maple Syrup Unne Disease	Branched-Chain Ketoacid Dehydrogenase	branched chain amino acids (valine, isoleucine and leucine)
30	Methyltetrahydrofolate Reductase Deficiency	Methyltetrahydrofolate Reductase	
	Phenylketonurea	phenylalanine hydroxylase	Biopterin, Pteridine, Phenylalanine
35	Tay-Sachs Disease	Beta-N-Acetylglucosaminidase, Hexosaminidase (Beta-Hexosaminidase-A)	Gangliosides
	Urea Cycle Disorder	Argininase, Argininosuccinate Lyase, Carbamyl Phosphate Synthetase	Orotic acid
40	Zellweger Syndrome	Dihydroxyacetonephosphate acyltransferase, Phytanic Acid Oxidase	Pipecolic Acid, Very Long Chain Fatty Acids
	Classic galactosemia	Galactose-1-phosphate undyl transferase (GALT)	
45	Galactosemia	Galactokinase	
	Galactosemia	Undine diphosphate galactose-4-epimerase	
	G6PD deficiency	Glucose-6-phosphate dehydrogenase (G6PD)	
50	Sialidosis type II	Neuraminidase	
	Hurler disease	α -Iduronidase	
	Hunter disease	Iduronate sulfatase	
55	Sanfilipo disease type A (MPS IIIA)	Heparan sulphamidase	
	Sanfilipo disease type B (MPS IIIB)	α -N-acetyl glucosaminidase	
	Sanfilipo disease type C (MPS IIIC)	Glucosamine N-acetyl transferase	

(continued)

Metabolic disorder	Metabolically indicative enzyme	Exemplary metabolic analyte
Morquio disease type A (MPS IVA)	N-acetyl galactosamine-6-sulphatase	
Morquio disease type B (MPS NB)	β -Galactosidase	
Maroteaux-Lamy disease (MPS VI)	Arylsulphatase B	
Sly disease (MPS VII)	β -Glucuronidase	
α -Mannosidosis	α -Mannosidase	
β -Mannosidosis	β -Mannosidase	
α -Fucosidosis	α -Fucosidase	
Sialidosis	α -N-acetylneuraminidase	
Galactosialidosis	α -N-acetylneuraminidase, p-Galactosidase	
Aspartylglucosaminuria	Aspartylglucosaminidase	
Schidler disease	α -N-acetylgalactosaminidase,	
Pompe disease	α -Glucosidase	
Tay-Sachs	α -Hexosaminidase	
Sandhoff disease	α -Hexosaminidase	
GM-1 gangliosidosis	β -Galactosidase	
Gaucher disease	β -Glucosidase, Chitotriosidase	
Krabbe disease	Galactocerebrosidase	
Metachromatic leucodystrophy	Arylsulphatase A	
Fabry disease	α -Galactosidase A	
Farber disease	Ceramidase	
Niemann-Pick disease	Sphingomyelinase	

[0061] Substrates for metabolically indicative enzymes are well known to those skilled in the art. As used herein, the term "substrate" means a molecule or complex acted upon by a metabolically indicative enzyme. In particular, a substrate has a physiological affinity for and must be able to be acted upon by a corresponding enzyme. A substrate useful in the methods of the invention can be native or modified. Modified substrates useful in the invention have a mass different from the native substrate and retain the ability to be acted upon by the corresponding enzyme. A variety of chemical modification reactions useful for increasing the mass of a substrate are well known to those skilled in the art. Exemplary modifications suitable for substrates include isotopic labeling and any chemical modification that alters the mass or mass-to-charge ratio of the substrate. Once a substrate is modified, activity can be confirmed using routine methods, such as comparison of enzyme activity using native substrate vs. modified substrate, or by quantitation of the product via internal standardization using one or more reference products of known concentration. As a non-limiting example, when the metabolically indicative enzyme is biotinidase, a substrate can be native biocytin or modified biocytin.

[0062] As used herein, the term "product" when used in reference to enzymatic activity means a molecular entity that is produced as a result of enzyme acting on substrate. Therefore, the presence or amount of a product in a sample correlates with the presence or amount of enzymatic activity in the sample. As such, an amount of product determined using a method of the invention can be expressed as a level of enzymatic activity using routine calculations, if desired, for example, when assessing differences between test sample and reference level enzymatic activity.

[0063] The methods of the invention involve use of a reagent that inhibits the ability of one or more metabolically indicative enzymes to act on a corresponding substrate. As used herein, the term "inhibit" means to reduce or block enzyme activity such that product formation is undetectable or insignificant.

[0064] As used herein, a "reagent" for use in the methods of the invention is or contains a chemical agent that specifically or non-specifically inhibits the ability of a metabolically indicative enzyme to act on a corresponding substrate, and does not substantially interfere with mass spectrometry analysis. Exemplary non-specific reagents that inhibit enzymatic reactions are chaotropes, which are chemical agents that denature proteins, and solutions containing chaotropes. Non-limiting examples of chaotropes are alcohols, such as methanol, ethanol and isobutyl alcohol; urea and guanidine salts: Exemplary specific reagents that inhibit enzymatic reactions include enzyme inhibitors effective for a selected enzyme. A large number of enzyme inhibitors are widely available and well-known to those skilled in the art. A reagent for use in the methods of the invention can be selected based on, for example, suitability for inhibiting activity of metabolically indicative enzyme(s), and ability to maintain solubility of metabolic analyte(s) and enzymatic product(s). A selected reagent can be selected to have a minimal number of components in order to minimize background for mass spectrometry. A selected reagent can be miscible or immiscible with a particular reaction admixture or solution, with rapid mixing being one method for ensuring that enzymatic reaction is inhibited. Examples of reagents useful in the invention include organic solvents and solutions thereof, organic acids and solutions thereof, alcohols and solutions thereof, reagents that are non-polar relative to the reaction admixture or solution, and reagents that have a pH distinct from the reaction admixture or solution, the pH creating an unsuitable environment for enzymatic activity, and the like. Non-limiting specific reagents include methanol and solutions thereof; trifluoroacetic acid and solutions thereof; and acetonitrile and solutions thereof. Those skilled in the art of mass spectrometry will be able to identify reagents that do not interfere with mass spectrometry determinations.

[0065] The ability of a selected reagent to inhibit an enzymatic reaction can be determined by routine methods. For example, the effectiveness of a candidate "reagent" can be tested by adding the reagent or a known inhibitor of the enzyme to an enzymatic reaction in progress. The amount of enzymatic activity under each condition can then be compared to confirm if a selected reagent is suitable for use in the invention methods.

[0066] The methods of the invention involve detecting at least one "metabolic analyte." A variety of metabolic analytes can be detected using the methods of the invention, so long as the analyte is soluble in a selected reagent, has a known mass, and can be modified, if desired, for use as an internal standard. Metabolic analytes associated with particular metabolic disorders are well known to those skilled in the art. As non-limiting examples, well-known metabolic analytes and their associated metabolic disorders are shown above in Table 1. Amino acid and carnitine metabolic analytes are shown in Table 2, and can correspond with common metabolic disorders shown in Table 3. Amino acid disorders and corresponding amino acid metabolic analytes are also shown in Table 4. Fatty acid disorders and corresponding fatty acid metabolic analytes are shown in Table 5; additional fatty acid disorders shown in Table 3.

Table 2

ANALYTE NAME	ABBREVIATION
Amino Acids	
Alanine	Ala
Arginine	Arg
Citrulline	Cit
Glycine	Gly
Leucine	Leu
Methionine	Met
Ornithine	Orn
5-Oxoproline	5-Oxo Pro
Phenylalanine	Phe
Tyrosine	Tyr
Valine	Val
Proline	Pro
Carnitines	
Free carnitine	C0
Acetylcarnitine	C2
Propionylcarnitine	C3

EP 1 664 325 B1

(continued)

5
10
15
20
25
30
35
40
45
50

Carnitines	
Malonylcarnitine	C3DC
Butyrylcarnitine	C4
3-Hydroxy-butyrylcarnitine	C4OH
Isovalerylcarnitine	C5
Tiglylcarnitine	C5:1
Glutarylcamitine	C5DC
3-Hydroxy-isovalerylcarnitine	C5OH
Hexanoylcarnitine	C6
Adipylcarnitine	C6DC
Octanoylcarnitine	C8
Octenoylcarnitine	C8:1
Decanoylcarnitine	C10
Decenoylcarnitine	C10:1
ANALYTE NAME	ABBREVIATION
Carnitines	
Decadienoylcarnitine	C10:2
Dodecanoylcarnitine	C12
Dodecenoylcarnitine	C12:1
Tetradecanoylcarnitine (Myristoylcarnitine)	C14
Tetradecenoylcarnitine	C14:1
T etradecadienoylcarnitine	C14:2
3-Hydroxy-tetradecanoylcarnitine	C14OH
Hexadecanoylcarnitine (palmitoylcarnitine)	C16
Hexadecenoylcarnitine	C16:1
3-Hydroxy-hexadecanoylcarnitine	C16OH
3-Hydroxy-hexadecenoylcarnitine	C16:1OH
Octadecanoylcarnitine (Stearoylcarnitine)	C18
Octadecenoylcarnitine (Oleylcarnitine)	C18:1
Octadecadienoylcarnitine (Linoleylcarnitine)	C18:2
3-Hydroxy-octadecanoylcarnitine	C18OH
3-Hydroxy-octadecenoylcarnitine	C18:1OH

Table 3

Amino Acid Disorders	
ARGD	Argininemia
ASA, ASL, ALD	Argininosuccinic aciduria
	(Argininosuccinate lyase deficiency, Argininosuccinase deficiency)
ASD, ASS	Citrullinemia

EP 1 664 325 B1

(continued)

Amino Acid Disorders		
	(Argininosuccinic acid synthetase deficiency, Argininosuccinate synthetase deficiency)	
5	HCU, HCYS	Homocystinuria
		(Cystathione synthase deficiency)
	Hmet	Hypermethioninemia
	HHH	Hyperornithinemia, Hyperammonemia, Hyperhomocitrullinuria syndrome (Omithine translocase deficiency)
10	PRO	Hyperprolinemia
	MSUD, BCKA	Maple Syrup Urine Disease (Branched chain ketoaciduria)
	NKG, NKHI	Nonketotic hyperglycinemia
	PKU	Phenylketonuria
15	PYP/PIP	Pyroglutamic/pipecolic acidemia
		Tyrosenemia, Type I
		Tyrosenemia, Type II
		5-oxoprolinuria
20		(Pyroglutamic aciduria)
Fatty Acid and Organic Acid Disorders		
	2-MBCD	2-methylbutyryl CoA dehydrogenase deficiency
		2,4-Dienoyl-CoA reductase deficiency
25	HMG	3 -hydroxy-3-methylglutaryl CoA lyase deficiency
		(hydroxymethylglutaric acidemia)
	3MCC, 3-MMC	3-methylcrotonyl CoA carboxylase deficiency
		(3-methylcrotonylglycinemia)
	CPTI	Carnitine palmitoyltransferase, type I deficiency
30	CPT II	Carnitine palmitoyltransferase, type II deficiency
	CTD	Carnitine transporter defect
	CATR, CACT	Carnitine/acylcarnitine translocase defect
	EMA	Ethylmalonic acidemia
35	GAI	Glutaric acidemia, type I
		(Glutaryl CoA dehydrogenase deficiency)
	IBCD	Isobutyryl CoA dehydrogenase deficiency
	IVA	Isovaleric acidemia
	LCAD	Long-chain acyl-CoA dehydrogenase deficiency
40	LCHAD	Long-chain hydroxyacyl-CoA dehydrogenase deficiency
	MA	Malonic aciduria
	MCAD	Medium-chain acyl-CoA dehydrogenase deficiency
	MMA	Methylmalonic acedemia
45	BKT	Mitochondrial acetoacetyl CoA thiolase deficiency
		(Beta-Ketothiolase deficiency)
	MADD. GA II	Multiple acyl-CoA dehydrogenase deficiency
		(Glutaric acidemia, type II)
	MCD	Multiple Co-A carboxylase deficiency
50		(Holocarboxylase synthetase deficiency)
	PA, PPA	Propionic acidemia
	SCAD	Short-chain acyl-CoA dehydrogenase deficiency
	SCHAD	Short-chain hydroxyacyl-CoA dhydrogenase deficiency
	TFP	Trifunctional protein deficiency
55	VLCAD	Very-long-chain acyl-CoA dehydrogenase deficiency

Aminoacidopathies

Phenylketonuria Phe ^{2,4,6} Phe, Phe/Tyr ⁷ Phe, Tyr ^{1,5} Maple syrup urine disease Leu+Ile ¹ Leu+Ile, Val ^{2,4,6} Leu, Val ⁵ Leu, Leu/Phe ⁷ Homocystinuria Met ^{1,2,4,6} Met, Met/Phe, Met/Leu ⁵ Met, Met/Phe ⁷ Hypermethioninemia Met ² Met, Met/Phe ⁷ Citrullinemia Cit ^{1,2,6} Cit/Phe, Cit/Tyr ⁴ Cit, Orn/Cit, Cit/Arg ⁵ Cit, Cit/Arg ⁷ Nonketotic hyperglycinemia Gly ^{2,4,5}	Tyrosinemia Type I ^{1,4} Met, Tyr ^{2,6} Tyr ⁷ Tyr, Tyr/Phe Tyrosinemia II ² Tyr ⁷ Argininemia Arg ² Arg, Arg/Orn PRO Hyperprolinemia ⁴ Pro/Phe HHH ² Orn, HomoCit ⁷ Orn, Orn/Cit 5-oxoprolinuria ² 5-Oxopro ⁴ pyroGlu/Phe Argininosuccinic aciduria ^{1,6} Cit ⁴ Cit/Tyr ⁵ Asa ⁷ Cit, Cit/Arg
--	--

Table 5

Fatty Acid Oxidation Disorders	
SCAD	LCAD
C4, C6 ¹	⁴ C14:2 C14 1. C14 1/C16
C4 ^{2,5,7}	
C4 C5 C4/C2 C5/C2 ⁴	VLCAD
C4, C4/C2, C4/C3 ⁶	¹ C14, C14.1. C16. C18
MCAD	² C14:1, C14. C16
C8, C10:1 ¹	³ C14.1, C14, C16:1, C16, C18:1, C18, C14:1/C12:1
C8, C10 C10:1, C6 ²	⁴ C14:2, C14:1, C14:1/C16
C6, C8, C10, C10:1, C8/C10 ³	⁵ C14.1 or C14
C6, C8, C10:1 ^{4,7}	⁶ C14, C14.1 C14:2, C16.1 C14:1/C16
C6, C8, C10:1, C10, C8/C2, C8/C10, C8/C12 ⁵	⁷ C14 1
C6, C8, C10:1, C8/C10 ⁶	LCHAD
	¹ C14, C14:1, C16, C18, C14OH, C14:1OH, C16OH C18OH

(continued)

Fatty Acid Oxidation Disorders			
5 10	Glutaric acidemia Type II		
	C5DC	1	2 C16OH, C18:1OH, C18OH
	C4, C5, C8;1, C8, C12 C14, C16, C5DC	2	3 C16OH, C18:1OH, C18OH, C16OH/C16
	C4, C5DC, C5, C6 C8, C10	3	5 C14OH, C16:1OH, C16OH, C18:1OH, C18:OH, C14:1, C14
	C6 C8, C10 C5/C2	4	6 C16OH, C18:1, C18:1OH, C18:2, C18:2OH
	C4 C5, C6, C8, C10	6	7 C16OH
	C8, C10, C5DC	7	
15	Carnitine transporter defect		Trifunctional protein deficiency
	C0, C2	6	2 C16OH, C18:1OH, C18OH
			6 C16OH, C18:1OH, C18OH, C16OH/C16
20	CPT I		5 C14OH, C16:1OH, C16OH C18:1OH, C18:OH, C14:1, C14
	C16, C18:1 C18	2	
	C0, C16, C18, C0/(C16+C18)	5	Carnitine/acylcarnitine translocase deficiency
25	Ethylmalonic acidemia		2 C16, C18:1, C18
	C4, C5 C4/C2, C5/C2	4	5 C0, C16, C18, C0/(C16+C18)
			7 C16
30	2,4 Dienoyl CoA reductase deficiency	6	CPT II
	C10.2		1 C14, C14:1, C16, C16:1
			3 C16 C18 C18:1, C16/C14:1
			6 C16, C18:1, C18:2
			7 C16

[0067] In the methods of the invention, the presence or amount of one or more metabolic analytes can be determined. An amount of a metabolic analyte refers to an absolute amount of a molecule in a sample or to a relative amount of the molecule, including amounts determined under steady-state or non-steady-state conditions. The amount of a metabolic analyte can be determined relative to a reference molecule in a sample.

[0068] The methods of the invention involve determining the presence or amount of at least one metabolic analyte and/or product, where the presence or amount of each metabolic analyte and/or product correlates the presence or absence of a metabolic disorder. To quantitate the amount of a metabolic analyte or product in a test sample, an internal standard can be used, as is described below. The methods of the invention can be used quantitatively, if desired, to allow comparison of test sample results with known or pre-determined standard amounts of particular analyte and/or product amounts. The methods of the invention can be used qualitatively when the presence of the metabolic analyte or enzyme in the sample is indicative of a metabolic disorder, for example, when the metabolic analyte detected is the result of abnormal metabolic processes, or when enzyme activity is not detected in a normal sample. The methods can also be used qualitatively when a test sample is compared with a reference sample, which can be either a normal reference or metabolic disorder reference. In this format, the relative amount of analyte or product can be indicative of a metabolic disorder.

[0069] An internal standard, or reference analyte, for a metabolic analyte useful in a method of the invention can be any modification or analog of the analyte that is detectable by mass spectrometry. A reference analyte is separately detectable from the metabolic analyte based on a unique physical characteristic, such as a unique mass or charge-to-mass ratio. In addition, a suitable generic reference standard can be used. Such an internal standard will, for example, co-elute with the analyte or product if a separation method such as chromatography is used prior to mass spectrometric analysis. A commonly used internal standard for mass spectrometry is a stable isotopically labeled form or chemical derivative of the analyte or product. Non-limiting examples of internal standards for metabolic analytes that are amino acids include isotopically labeled amino acids such as ¹⁵N,²-¹³C-Glycine, ²H₄-Alanine, ²H₈-Valine, ²H₃-Leucine, ²H₃-Methionine, ²H₅-Phenylalanine, ²H₄-Tyrosine, ²H₃-Aspartate, ²H₃-Glutamate, ²H₂-Ornithine ²H Cl, ²H₂-Citrulline, and ²H₄-¹³C-Arginine-HCl. Non-limiting examples of internal standards for metabolic analytes that are carnitines include isotopically labeled carnitines such as ²H₉-carnitine, ²H₃-acetylcarnitine, ²H₃-propionylcarnitine, ²H₃-butyrylcarnitine,

^2H 9-isovalerylcarnitine, ^2H 3-octanoylcarnitine, ^2H 9-myristoylcarnitine, and ^2H 3-palmitoylcarnitine. An internal standard for an enzymatic product, or reference product, can be any modification or analog of the product that is identifiable by mass spectrometry. A reference product is separately detectable from the product based on a unique physical characteristic, such as a unique mass or charge-to-mass ratio. A non-limiting example of an internal standard for an enzymatic product is an isotopically labeled form of the product, such as a deuterated form of the product.

[0070] Generally, a cut-off value for each metabolic analyte or enzyme activity will be known in the art for commonly tested analytes and enzymes. A cut-off value is typically an analyte concentration or enzyme activity, or ratio thereof, above or below which is considered indicative of a metabolic disorder or cause for retest. Thus, in accordance with the present invention a reference level of a metabolic analyte or enzymatic activity (as indicated by presence or amount of enzymatic product) in a particular sample type is identified as a cut-off value, above which there is a significant correlation between the presence of the analyte or enzymatic activity and presence or absence of a metabolic disorder. Exemplary cut-off values for amino acid and carnitine/acylcarnitine butyl esters are described in US 6,455,321. Also described are procedures for determining significance of an elevated amount of a metabolic analyte in a blood sample from a newborn. This method can be applied to a variety of types of samples (for example, serum and tissue biopsy) from a subject of any age.

[0071] Those of skill in the art will recognize that some cut-off values are not absolute in that clinical correlations are still significant over a range of values on either side of the cutoff; however, it is possible to select an optimal cut-off value (e.g. varying H-scores, and the like) of analyte, enzyme activity for a particular sample types. Cut-off values determined for use in a method of the invention generally are compared with published ranges but can be individualized to the methodology used and patient population. It is understood that improvements in optimal cut-off values could be determined depending on the sophistication of statistical methods used and on the number and source of samples used to determine reference level values for the different analyte, enzyme activity and sample types. Therefore, established cut-off values can be adjusted up or down, on the basis of periodic re-evaluations or changes in methodology or population distribution. In addition, instrument-specific cut-off values can be used, if desired, for example such as when inter-instrument performance comparability is <10%.

[0072] The reference level can be determined by a plurality of methods, provided that the resulting reference level accurately provides an amount of metabolic analyte or enzyme activity above which exists a first group of individuals having a different probability of metabolic disorder than that of a second group of individuals having metabolic analyte or enzyme activity amount below the reference level. The reference level can be determined by comparison of metabolic analyte or enzyme activity amount in populations of patients having the same metabolic disorder. This can be accomplished, for example, by histogram analysis, in which an entire cohort of patients are graphically presented, wherein a first axis represents the amount of metabolic analyte or enzyme activity and a second axis represents the number of individuals in the cohort whose sample contain metabolic analyte or enzyme activity at a given amount. Two or more separate groups of individuals can be determined by identification of subsets populations of the cohort which have the same or similar levels of metabolic analyte or enzyme activity. Determination of the reference level can then be made based on an amount which best distinguishes these separate groups. The reference level can be a single number, equally applicable to every individual, or the reference level can vary, according to specific subpopulations of individuals. For example, older individuals might have a different reference level than younger individuals for the same metabolic disorder.

[0073] The methods of the invention use mass spectrometry for determining the amount or presence of one more metabolic analytes and at least one metabolically indicative enzyme product. A variety of configurations of mass spectrometers can be used in a method of the invention. Several types of mass spectrometers are available or can be produced with various configurations. In general, a mass spectrometer has the following major components: a sample inlet, an ion source, a mass analyzer, a detector, a vacuum system, and instrument-control system, and a data system. Difference in the sample inlet, ion source, and mass analyzer generally define the type of instrument and its capabilities. For example, an inlet can be a capillary-column liquid chromatography source or can be a direct probe or stage such as used in matrix-assisted laser desorption. Common ion sources are, for example, electrospray, including nanospray and microspray or matrix-assisted laser desorption. Common mass analyzers include a quadrupole mass filter, ion trap mass analyzer and time-of-flight mass analyzer.

[0074] The ion formation process is a starting point for mass spectrum analysis. Several ionization methods are available and the choice of ionization method depends on the sample to be analyzed. For example, for the analysis of amino acids a relatively gentle ionization procedure such as electrospray ionization (ESI) can be desirable. For ESI, a solution containing the sample is passed through a fine needle at high potential which creates a strong electrical field resulting in a fine spray of highly charged droplets that is directed into the mass spectrometer. Other ionization procedures include, for example, fast-atom bombardment (FAB) which uses a high-energy beam of neutral atoms to strike a solid sample causing desorption and ionization. Matrix-assisted laser desorption ionization (MALDI) is a method in which a laser pulse is used to strike a sample that has been crystallized in an UV-absorbing compound matrix. Other ionization procedures known in the art include, for example, plasma and glow discharge, plasma desorption ionization, resonance

ionization, and secondary ionization.

[0075] Electrospray ionization (ESI) has several properties that are useful for the invention described herein. For example, ESI can be used for biological molecules such as amino acids and carnitines that are difficult to ionize or vaporize. In addition, the efficiency of ESI can be very high which provides the basis for highly sensitive measurements. Furthermore, ESI produces charged molecules from solution, which is convenient for analyzing metabolic analytes, products and reference molecules that are in solution. In contrast, ionization procedures such as MALDI require crystallization of the sample prior to ionization.

[0076] Since ESI can produce charged molecules directly from solution, it is compatible with samples from liquid chromatography systems. For example, a mass spectrometer can have an inlet for a liquid chromatography system, such as an HPLC, so that fractions flow from the chromatography column into the mass spectrometer. This in-line arrangement of a liquid chromatography system and mass spectrometer is sometimes referred to as LC-MS. A LC-MS system can be used, for example, to separate metabolic analytes, products and reference molecules from complex mixtures before mass spectrometry analysis. In addition, chromatography can be used to remove salts or other buffer components from the sample before mass spectrometry analysis. For example, desalting of a sample using a reversed-phase HPLC column, in-line or off-line, can be used to increase the efficiency of the ionization process and thus improve sensitivity of detection by mass spectrometry.

[0077] A variety of mass analyzers are available that can be paired with different ion sources. Different mass analyzers have different advantages as known to one skilled in the art and as described herein. The mass spectrometer and methods chosen for detection depends on the particular assay, for example, a more sensitive mass analyzer can be used when a small amount of ions are generated for detection. Several types of mass analyzers and mass spectrometry methods are described below.

[0078] Quadrupole mass spectrometry utilizes a quadrupole mass filter or analyzer. This type of mass analyzer is composed of four rods arranged as two sets of two electrically connected rods. A combination of rf and dc voltages are applied to each pair of rods which produces fields that cause an oscillating movement of the ions as they move from the beginning of the mass filter to the end. The result of these fields is the production of a high-pass mass filter in one pair of rods and a low-pass filter in the other pair of rods. Overlap between the high-pass and low-pass filter leaves a defined m/z that can pass both filters and traverse the length of the quadrupole. This m/z is selected and remains stable in the quadrupole mass filter while all other m/z have unstable trajectories and do not remain in the mass filter. A mass spectrum results by ramping the applied fields such that an increasing m/z is selected to pass through the mass filter and reach the detector. In addition, quadrupoles can also be set up to contain and transmit ions of all m/z by applying a rf-only field. This allows quadrupoles to function as a lens or focusing system in regions of the mass spectrometer where ion transmission is needed without mass filtering. This will be of use in tandem mass spectrometry as described further below.

[0079] A quadrupole mass analyzer, as well as the other mass analyzers described herein, can be programmed to analyze a defined m/z or mass range. This property of mass spectrometers is useful for the invention described herein. Since the mass range of metabolic analytes, enzymatic products and reference molecules will be known prior to an assay, a mass spectrometer can be programmed to transmit ions of the projected correct mass range while excluding ions of a higher or lower mass range. The ability to select a mass range can decrease the background noise in the assay and thus increase the signal-to-noise ratio as well as increasing the specificity of the assay. Therefore, the mass spectrometer can accomplish an inherent separation step as well as detection and identification of metabolic analytes, enzymatic products and reference molecules.

[0080] Ion trap mass spectrometry utilizes an ion trap mass analyzer. In these mass analyzers, fields are applied so that ions of all m/z are initially trapped and oscillate in the mass analyzer. Ions enter the ion trap from the ion source through a focusing device such as an octapole lens system. Ion trapping takes place in the trapping region before excitation and ejection through an electrode to the detector. Mass analysis is accomplished by sequentially applying voltages that increase the amplitude of the oscillations in a way that ejects ions of increasing m/z out of the trap and into the detector. In contrast to quadrupole mass spectrometry, all ions are retained in the fields of the mass analyzer except those with the selected m/z . One advantage to ion traps is that they have very high sensitivity, as long as one is careful to limit the number of ions being trapped at one time. Control of the number of ions can be accomplished by varying the time over which ions are injected into the trap. The mass resolution of ion traps is similar to that of quadrupole mass filters, although ion traps do have low m/z limitations.

[0081] Time-of-flight mass spectrometry utilizes a time-of-flight mass analyzer. For this method of m/z analysis, an ion is first given a fixed amount of kinetic energy by acceleration in an electric field (generated by high voltage). Following acceleration, the ion enters a field-free or "drift" region where it travels at a velocity that is inversely proportional to its m/z . Therefore, ions with low m/z travel more rapidly than ions with high m/z . The time required for ions to travel the length of the field-free region is measured and used to calculate the m/z of the ion.

[0082] One consideration in this type of mass analysis is that the set of ions being studied be introduced into the analyzer at the same time. For example, this type of mass analysis is well suited to ionization techniques like MALDI which produces ions in short well-defined pulses. Another consideration is to control velocity spread produced by ions

that have variations in their amounts of kinetic energy. The use of longer flight tubes, ion reflectors, or higher accelerating voltages can help minimize the effects of velocity spread. Time-of-flight mass analyzers have a high level of sensitivity and a wider m/z range than quadrupole or ion trap mass analyzers. Also data can be acquired quickly with this type of mass analyzer because no scanning of the mass analyzer is necessary.

[0083] Tandem mass spectrometry can utilize combinations of the mass analyzers described above. Tandem mass spectrometers can use a first mass analyzer to separate ions according to their m/z in order to isolate an ion of interest for further analysis. The isolated ion of interest is then broken into fragment ions (called collisionally activated dissociation or collisionally induced dissociation) and the fragment ions are analyzed by the second mass analyzer. These types of tandem mass spectrometer systems are called tandem in space systems because the two mass analyzers are separated in space, usually by a collision cell. Tandem mass spectrometer systems also include tandem in time systems where one mass analyzer is used, however the mass analyzer is used sequentially to isolate an ion, induce fragmentation, and then perform mass analysis.

[0084] Mass spectrometers in the tandem in space category have more than one mass analyzer. For example, a tandem quadrupole mass spectrometer system can have a first quadrupole mass filter, followed by a collision cell, followed by a second quadrupole mass filter and then the detector. Another arrangement is to use a quadrupole mass filter for the first mass analyzer and a time-of-flight mass analyzer for the second mass analyzer with a collision cell separating the two mass analyzers. Other tandem systems are known in the art including reflectron-time-of-flight, tandem sector and sector-quadrupole mass spectrometry.

[0085] Mass spectrometers in the tandem in time category have one mass analyzer that performs different functions at different times. For example, an ion trap mass spectrometer can be used to trap ions of all m/z . A series of rf scan functions are applied which ejects ions of all m/z from the trap except the m/z of ions of interest. After the m/z of interest has been isolated, an rf pulse is applied to produce collisions with gas molecules in the trap to induce fragmentation of the ions. Then the m/z values of the fragmented ions are measured by the mass analyzer. Ion cyclotron resonance instruments, also known as Fourier transform mass spectrometers, are an example of tandem-in-time systems.

[0086] Several types of tandem mass spectrometry experiments can be performed by controlling the ions that are selected in each stage of the experiment. The different types of experiments utilize different modes of operation, sometimes called "scans," of the mass analyzers. In a first example, called a mass spectrum scan, the first mass analyzer and the collision cell transmit all ions for mass analysis into the second mass analyzer. In a second example, called a product ion scan, the ions of interest are mass-selected in the first mass analyzer and then fragmented in the collision cell. The ions formed are then mass analyzed by scanning the second mass analyzer. In a third example, called a precursor ion scan, the first mass analyzer is scanned to sequentially transmit the mass analyzed ions into the collision cell for fragmentation. The second mass analyzer mass-selects the product ion of interest for transmission to the detector. Therefore, the detector signal is the result of all precursor ions that can be fragmented into a common product ion. Other experimental formats include neutral loss scans where a constant mass difference is accounted for in the mass scans. The use of these different tandem mass spectrometry scan procedures can be advantageous when large sets of analytes are measured in a single experiment as with multiplex experiments.

[0087] In view of the above, those skilled in the art recognize that different mass spectrometry methods, for example, quadrupole mass spectrometry, ion trap mass spectrometry, time-of-flight mass spectrometry and tandem mass spectrometry, can use various combinations of ion sources and mass analyzers which allows for flexibility in designing customized detection protocols. In addition, mass spectrometers can be programmed to transmit all ions from the ion source into the mass spectrometer either sequentially or at the same time. Furthermore, a mass spectrometer can be programmed to select ions of a particular mass for transmission into the mass spectrometer while blocking other ions. The ability to precisely control the movement of ions in a mass spectrometer allows for greater options in detection protocols which can be advantageous when a large number of analytes, for example, from a multiplex experiment, are being analyzed.

[0088] Different mass spectrometers have different levels of resolution, that is, the ability to resolve peaks between ions closely related in mass. The resolution is defined as $R=m/\Delta m$, where m is the ion mass and Δm is the difference in mass between two peaks in a mass spectrum. For example, a mass spectrometer with a resolution of 1000 can resolve an ion with a m/z of 100.0 from an ion with a m/z of 100.1. Those skilled in the art will therefore select a mass spectrometer having a resolution appropriate for the analyte(s) to be detected.

[0089] Mass spectrometers can resolve ions with small mass differences and measure the mass of ions with a high degree of accuracy. Therefore, analytes of similar masses can be used together in the same experiment since the mass spectrometer can differentiate the mass of even closely related molecules. The high degree of resolution and mass accuracy achieved using mass spectrometry methods allows the use of large sets of analytes because they can be distinguished from each other.

[0090] Additional mass spectrometry methods are well known in the art (see Burlingame et al. Anal. Chem. 70:647R-716R (1998); Kinter and Sherman, New York (2000)). Exemplary descriptions of mass spectrometry methods for detecting metabolic analytes include Chace DH, Hillman SL, Van Hove JLK, Naylor EW. Clin Chem 1997;43:210613; Rashed

MS, Bucknall MP, Little D, et al. Clin Chem 1997;43:112941; Matern D, Strauss AW, Hillman SL, Mayatepek E, Millington DS, Trefz FK. Pediatr Res 1999;46:459, and Millington DS, Kodo N, Terada N, Roe D, Chace DH. International Journal of Mass Spectrometry and Ion Processes 1991;111:21128.

[0091] One or more steps of a method of the invention can be automated, if desired. Those skilled in the art will know how to use autosamplers, liquid handling devices, robots, computers and other routine laboratory instruments for automation.

[0092] All journal articles, reference, and patent citations provided above, in parentheses or otherwise, whether previously stated or not, are incorporated herein by reference.

[0093] It is understood that modifications that do not substantially affect the function of the various embodiments of this invention are also included within the definition of the invention provided herein: Accordingly, the following examples are intended to illustrate but not limit the present invention.

EXAMPLE 1

[0094] This example describes simultaneous determination of the amount of a metabolic analyte and an metabolically indicative enzyme product from a patient sample.

[0095] The enzyme biotinidase is a known metabolically indicative enzyme for the metabolic disorder termed biotinidase deficiency. In the untreated state, profound biotinidase deficiency in newborns typically is characterized by neurological and skin-related symptoms, such as seizures, hypotonia, and rash, often accompanied by hyperventilation, laryngeal stridor, and apnea. Older children may also have alopecia, ataxia, developmental delay, neurosensory hearing loss, optic atrophy, and recurrent infections. Individuals with partial biotinidase deficiency may have hypotonia, skin rash, and hair loss, particularly during times of stress. Once detected, biotinidase deficiency is often treatable by administration of biotin.

[0096] It has been previously established that individuals with profound biotinidase deficiency have <10% of mean normal serum biotinidase activity. Individuals with partial biotinidase deficiency have 10-30% of mean normal serum biotinidase activity. Therefore, it is understood that a decrease in biotinidase activity compared to normal reference level is indicative of biotinidase deficiency.

[0097] Acylcarnitines and α -amino acid also were assayed. Different patterns of the plasma acylcarnitine profile can indicate the diagnosis of fatty acid oxidation disorders (MCAD, VLCAD, SCAD, MAD, LCHAD, and CPTII), as well as some organic acidemias (propionic acidemia, methylmalonic acidemia, isovaleric acidemia, glutaric acidemia type I, 3-methylcrotonyl CoA carboxylase deficiency, B-ketothiolase deficiency, etc.). Plasma amino acids analysis will detect specific amino acid disorders such as phenylketonuria (PKU), Maple Syrup Urine Disease (MSUD), urea cycle defects, non-ketotic hyperglycinemia and homocystinuria. This analysis is also indicated in patients with general failure to thrive as an initial screen. For assessment of a metabolic disorder, plasma amino acids are compared to age-matched normal values

[0098] Activity of the enzyme biotinidase and the amount of the α -amino acid and acylcarnitine analytes were measured simultaneously as follows:

[0099] Dried blood samples on paper were each punched into a well of a microwell plate. Each sample was reconstituted with 50 microliter of buffer solution containing isotopically labeled biocytin (the substrate for biotinidase), and the mixture is incubated for 17 hours at 42°C. 140 microliter of methanol containing isotopically labeled internal standards for α -amino acids and acylcarnitines and for the product of the biotinidase reaction were then added, and the plate was shaken for 30 minutes at 27°C. The plate was then covered with aluminum foil and placed-on an autosampler. Each processed sample was analyzed by tandem mass spectrometry, and concentrations of biotinidase product, endogenous α -amino acids and endogenous acylcarnitines were quantified based on internal standards.

[0100] Tables 6 and 7 show data obtained using the above method, as compared to data obtained using well known mass spectrometry screening methods for metabolic analytes without simultaneously measuring enzyme activity. A slope of one indicates identical results between the compared methods.

Table 6: α -Amino acid (AA) comparison:

AA	SLOPE	R2
ALA	1.04	0.9916
ARG	1.11	0.9975
CIT	1.13	0.9912
GLY	1.36	0.9836
LEU	1.14	0.9987

(continued)

AA	SLOPE	R2
MET	1.00	0.9989
ORN	1.02	0.9951
PHE	0.99	0.9949
TYR	1.10	0.9917
VAL	1.04	0.9836
MEAN	1.09	0.9927

Table 7: Acylcarnitine comparison:

AC	SLOPE	R2
C0	1.05	0.9994
C2	1.12	0.9902
C3	1.11	0.9915
C4	1.10	0.9946
C5DC	0.89	0.9641
C6	0.99	0.9973
C8	1.00	0.9987
C10	1.05	0.9987
C12	1.01	0.9968
C14	1.06	0.9952
C16	1.21	0.9837
C18	1.16	0.9844
MEAN	1.06	0.9912

[0101] These results show that the above method produces data that correlate with well known mass spectrometry screening methods in which α -amino acids and acylcarnitines are extracted and measured without measuring enzyme activity simultaneously. These results also show that the above method produces data that correlate with known enzyme activity screening methods that measure fluorescence of a tagged product without measuring simultaneously concentrations of α -amino acids and acylcarnitines.

Claims

1. A method for detecting a metabolic disorder, which is a Fatty Acid Oxidation Disorder, Organic Aciduria Disorder, Amino Acidopathy or a disorder listed in Table 1 in an individual, comprising the simultaneous detection of metabolic enzyme activity and metabolite levels **characterized by**:

(a) contacting a sample to determine the presence of

- (i) one or more metabolically indicative enzymes which have an altered activity as a result of a metabolic disorder and
- (ii) one or more metabolic analytes, which have an altered amount as a result of a metabolic disorder,

with one or more substrates for said one or more enzymes to produce a reaction admixture, under conditions in which at least one of said enzymes is capable of acting on a corresponding substrate to generate at least

one product;

(b) contacting said reaction admixture with a reagent that inhibits the ability of said one or more enzymes to act on a corresponding substrate, wherein said one or more metabolic analytes and said at least one product are soluble in said reagent; to produce a test sample and

(c) determining the amount or presence of said one or more metabolic analytes and said at least one product contained in said test sample using mass spectrometry,

wherein a determined amount or presence of said one or more metabolic analytes and said at least one product correlates with presence or absence of said metabolic disorder.

2. The method of claim 1, wherein said sample is a body fluid sample.

3. The method of claim 2, wherein said body fluid sample is blood.

4. The method of claim 1, wherein said sample is dried.

5. The method of claim 1, wherein said individual is a human suspected of having a metabolic disorder.

6. The method of claim 1, wherein said individual is a neonate.

7. The method of claim 1, wherein said individual is a newborn.

8. The method of claim 1, wherein said individual is a child.

9. The method of claim 1, wherein said individual is an adult.

10. The method of claim 1, wherein said metabolic disorder is an inborn error of metabolism.

11. The method of claim 1, wherein said metabolic disorder is an acquired metabolic disorder.

12. The method of claim 1, wherein said metabolically indicative enzyme is selected from the group consisting of oxidoreductase, hydrolase, lyase, transferase, ligase and isomerase.

13. The method of claim 12, wherein the enzyme is a hydrolase.

14. The method of claim 13, wherein the enzyme is a biotinidase.

15. The method of claim 14, wherein said substrate is biocytin.

16. The method of claim 1, wherein said metabolic analyte is one or more amino acids.

17. The method of claim 1, wherein said metabolic analyte is an acylcarnitine or plurality of acylcarnitines.

18. The method of claim 1, wherein step (a) further comprises contacting said sample with one or more reference substrates.

19. The method of claim 1, wherein step (b) further comprises contacting said sample with one or more reference products.

20. The method of claim 1, wherein step (d) further comprises, prior to determining, adding one or more reference products corresponding to the at least one product.

21. The method of claim 1, wherein step (d) further comprises, prior to determining, adding one or more reference analytes corresponding to the one or more metabolic analytes contained in said sample.

22. The method of claim 1, wherein said reaction admixture is aqueous.

23. The method of claim 22, wherein said reagent is non-aqueous.

24. The method of claim 23, wherein said reagent comprises an organic solvent.
25. The method of claim 23, wherein said reagent comprises an alcohol.
- 5 26. The method of claim 25, wherein said reagent is methanol.
27. The method of claim 1, wherein said mass spectrometry is tandem mass spectrometry.
28. The method of claim 1, wherein the one or more substrates are present in dried form in a vessel.
- 10 29. The method of claim 1, wherein the reaction admixture is contacted with one or more reference products and one or more reference analytes and the amount of said one or more metabolic analytes and said at least one product contained in said test samples is determined with respect to said one or more reference products and said one or more reference analytes, using mass spectrometry.
- 15 30. The method of claim 1, wherein the sample is separated into two sample portions and each sample portion is subjected to (a) and (b); combining the two reaction admixtures to produce a test sample; and subjecting the combined reaction admixtures to (c).
- 20 31. The method of claim 1, wherein the sample is separated into two sample portions, wherein (a) comprises contacting each sample portion with an individual vessel, each vessel containing said one or more substrates for said one or more enzymes in dried form; subjecting each sample portion to (b); combining the two reaction admixtures to produce a test sample; and subjecting the combined reaction admixtures to (c).
- 25 32. The method of claim 28, wherein said vessel is contacted with a solution to generate said conditions in which at least one of said enzymes is capable of acting on a corresponding substrate to generate at least one product.
- 30

Patentansprüche

- 35 1. Verfahren zum Nachweis einer metabolischen Störung, die eine Fettsäureoxidationsstörung, eine organische Azidurie, eine Aminoazidopathie oder eine der in Tabelle 1 aufgelisteten Störungen bei einem Individuum sein kann, aufweisend den gleichzeitigen Nachweis von metabolischer Enzymaktivität und Metabolitenspiegeln, **gekennzeichnet durch:**
- 40 (a) Kontaktieren einer Probe zum Bestimmen des Vorhandenseins
- (i) eines oder mehrerer metabolisch indikativer Enzyme, das bzw. die eine veränderte Aktivität infolge einer metabolischen Störung aufweisen, und
- (ii) eines oder mehrerer metabolischer Analyte, das bzw. die infolge einer metabolischen Störung in veränderter Menge vorliegen,
- 45 mit einem oder mehreren Substraten für das eine oder die mehreren Enzyme, um einen Reaktionszusatz herzustellen, unter Bedingungen, bei denen wenigstens eines der Enzyme in der Lage ist, auf ein entsprechendes Substrat einzuwirken, um wenigstens ein Produkt zu erzeugen,
- 50 (b) Kontaktieren des Reaktionszusatzes mit einem Reagens, das die Eigenschaft des einen oder der mehreren Enzyme, auf ein entsprechendes Substrat einzuwirken, hemmt, wobei das eine oder die mehreren metabolischen Analyte und das wenigstens eine Produkt in dem Reagens lösbar sind, um eine Testprobe herzustellen, und
- (c) Bestimmen der Menge oder des Vorhandenseins des einen oder der mehreren metabolischen Analyte und des wenigstens einen Produktes, das in der Testprobe enthalten ist, unter Anwendung der Massenspektrometrie,
- 55 wobei eine bestimmte Menge oder das Vorhandensein des einen oder der mehreren metabolischen Analyte und des wenigstens einen Produktes mit dem Vorliegen oder dem Nichtvorliegen der metabolischen Störung korreliert.

EP 1 664 325 B1

2. Verfahren nach Anspruch 1, bei dem die Probe eine Körperflüssigkeitsprobe ist.
3. Verfahren nach Anspruch 2, bei dem die Körperflüssigkeitsprobe Blut ist.
- 5 4. Verfahren nach Anspruch 1, bei dem die Probe getrocknet ist.
5. Verfahren nach Anspruch 1, bei dem das Individuum ein humanes Lebewesen mit Verdacht auf eine metabolische Störung ist.
- 10 6. Verfahren nach Anspruch 1, bei dem das Individuum ein Neugeborenes ist.
7. Verfahren nach Anspruch 1, bei dem das Individuum ein reifes Neugeborenes ist.
8. Verfahren nach Anspruch 1, bei dem das Individuum ein Kind ist.
- 15 9. Verfahren nach Anspruch 1, bei dem das Individuum ein Erwachsener ist.
10. Verfahren nach Anspruch 1, bei dem die metabolische Störung eine angeborene Stoffwechseldefunktion ist.
- 20 11. Verfahren nach Anspruch 1, bei dem die metabolische Störung eine erworbene metabolische Störung ist.
12. Verfahren nach Anspruch 1, bei dem die metabolisch indikativen Enzyme ausgewählt werden aus einer Gruppe bestehend aus Oxidoreduktasen, Hydrolasen, Lyasen, Transferasen, Ligasen und Isomerasen.
- 25 13. Verfahren nach Anspruch 12, bei dem das Enzym eine Hydrolase ist.
14. Verfahren nach Anspruch 13, bei dem das Enzym eine Biotinidase ist.
15. Verfahren nach Anspruch 14, bei dem das Substrat Biocytin ist.
- 30 16. Verfahren nach Anspruch 1, bei dem das metabolische Analyt aus einer oder mehreren Aminosäuren besteht.
17. Verfahren nach Anspruch 1, bei dem das metabolische Analyt aus einem Acylcarnitin oder einer Vielzahl von Acylcarnitinen besteht.
- 35 18. Verfahren nach Anspruch 1, bei dem Schritt (a) weiterhin das Kontaktieren der Probe mit einer oder mit mehreren Referenzsubstraten umfasst.
19. Verfahren nach Anspruch 1, bei dem Schritt (b) weiterhin das Kontaktieren der Probe mit einer oder mit mehreren Referenzprodukten umfasst.
- 40 20. Verfahren nach Anspruch 1, bei dem Schritt (d), vor dem Bestimmen, weiterhin das Beigeben eines oder mehrerer Referenzprodukte entsprechend dem wenigstens einen Produkt umfasst.
- 45 21. Verfahren nach Anspruch 1, bei dem Schritt (d), vor dem Bestimmen, weiterhin das Beigeben eines oder mehrerer Referenzanalyte entsprechend dem einen oder der mehreren metabolischen Analyte, die in der Probe enthalten sind, umfasst.
22. Verfahren nach Anspruch 1, bei dem der Reaktionszusatz wässrig ist.
- 50 23. Verfahren nach Anspruch 22, bei dem das Reagens nichtwässrig ist.
24. Verfahren nach Anspruch 23, bei dem das Reagens eine organische Lösung aufweist.
- 55 25. Verfahren nach Anspruch 23, bei dem das Reagens Alkohol aufweist.
26. Verfahren nach Anspruch 25, bei dem das Reagens Methanol ist.

27. Verfahren nach Anspruch 1, bei dem es sich bei der Massenspektrometrie um die Tandem-Massenspektrometrie handelt.
28. Verfahren nach Anspruch 1, bei dem das eine oder die mehreren Substrate in getrockneter Form in einem Behältnis vorliegen.
29. Verfahren nach Anspruch 1, bei dem der Reaktionszusatz mit einem oder mit mehreren Referenzprodukten und mit einem oder mehreren Referenzanalyten in Kontakt gebracht wird und die Menge des einen oder der mehreren metabolischen Analyte und des wenigstens einen Produktes, die in den Testproben enthalten sind, bezüglich des einen oder der mehreren Referenzprodukte und des einen oder der mehreren Referenzanalyte bestimmt werden, und zwar unter Anwendung der Massenspektrometrie.
30. Verfahren nach Anspruch 1, bei dem die Probe in zwei Probenteile getrennt wird und jeder Probenteil den Schritten (a) und (b) unterzogen wird;
Kombinieren der beiden Reaktionszusätze, um eine Testprobe herzustellen, und
Unterziehen der kombinierten Reaktionszusätze dem Schritt (c).
31. Verfahren nach Anspruch 1, bei dem die Probe in zwei Probenteile getrennt wird, wobei Schritt (a) das Kontaktieren jedes Probenteils mit einem individuellen Behältnis umfasst, wobei jedes Behältnis das eine oder die mehreren Substrate für das eine oder die mehreren Enzyme in getrockneter Form enthält,
Unterziehen jedes Probenteils dem Schritt (b),
Kombinieren der zwei Reaktionszusätze, um eine Testprobe herzustellen, und
Unterziehen des kombinierten Reaktionszusatzes dem Schritt (c).
32. Verfahren nach Anspruch 28, bei dem das Behältnis mit einer Lösung in Kontakt gebracht wird, um die Bedingungen zu erzeugen, unter denen wenigstens eines der Enzyme in der Lage ist, auf ein entsprechendes Substrat einzuwirken, um wenigstens ein Produkt zu erzeugen.

Revendications

1. Procédé de détection d'un trouble métabolique, qui est un trouble de l'oxydation des acides gras, une acidurie organique, une aminoacidopathie ou un trouble proposé dans la liste du tableau I chez un individu, comprenant la détection simultanée de l'activité métabolique des enzymes et des niveaux de métabolites, **caractérisé par** :
 - (a) la mise en contact d'un échantillon, pour déterminer la présence de :
 - (i) une ou plusieurs enzymes donnant des indications sur le métabolisme dont l'activité est modifiée en résultat d'un trouble métabolique et
 - (ii) un ou plusieurs analytes métaboliques, dont la quantité est modifiée en résultat d'un trouble métabolique,avec un ou plusieurs substrats pour ladite ou lesdites enzymes afin de produire un mélange réactionnel, dans des conditions dans lesquelles au moins l'une desdites enzymes est capable d'agir sur un substrat correspondant pour générer au moins un produit ;
 - (b) la mise en contact dudit mélange réactionnel avec un réactif qui inhibe la capacité de ladite ou desdites enzymes à agir sur un substrat correspondant, dans lequel ledit ou lesdits analytes métaboliques et ledit ou lesdits produits sont solubles dans ledit réactif ; pour produire un échantillon de test et
 - (c) la détermination de la quantité ou de la présence dudit ou desdits analytes métaboliques et dudit ou desdits produits contenus dans ledit échantillon de test par spectrométrie de masse,dans lequel une quantité déterminée ou une présence dudit ou desdits analytes métaboliques et dudit ou desdits produits est corrélée à la présence ou à l'absence dudit trouble métabolique.
2. Procédé selon la revendication 1, dans lequel ledit échantillon est un échantillon de fluide corporel.
3. Procédé selon la revendication 2, dans lequel ledit échantillon de fluide corporel est du sang.
4. Procédé selon la revendication 1, dans lequel ledit échantillon est sec.

EP 1 664 325 B1

5. Procédé selon la revendication 1, dans lequel ledit individu est un humain dont on soupçonne qu'il souffre d'un trouble métabolique.
- 5 6. Procédé selon la revendication 1, dans lequel ledit individu est un nouveau-né.
7. Procédé selon la revendication 1, dans lequel ledit individu est un nourrisson.
8. Procédé selon la revendication 1, dans lequel ledit individu est un enfant.
- 10 9. Procédé selon la revendication 1, dans lequel ledit individu est un adulte.
10. Procédé selon la revendication 1, dans lequel ledit trouble métabolique est une erreur innée du métabolisme.
11. Procédé selon la revendication 1, dans lequel ledit trouble métabolique est un trouble acquis du métabolisme.
- 15 12. Procédé selon la revendication 1, dans lequel ladite enzyme donnant des indications sur le métabolisme est sélectionnée dans le groupe constitué de l'oxydoréductase, de l'hydrolase, de la lyase, de la transférase, de la ligase et de l'isomérase.
- 20 13. Procédé selon la revendication 12, dans lequel l'enzyme est une hydrolase.
14. Procédé selon la revendication 13, dans lequel l'enzyme est une biotinidase.
15. Procédé selon la revendication 14, dans lequel ledit substrat est la biocytine.
- 25 16. Procédé selon la revendication 1, dans lequel ledit analyte métabolique est un ou plusieurs acides aminés.
17. Procédé selon la revendication 1, dans lequel ledit analyte métabolique est une acylcarnitine ou une pluralité d'acylcarnitines.
- 30 18. Procédé selon la revendication 1, dans lequel, l'étape (a) comprend en outre la mise en contact dudit échantillon avec un ou plusieurs substrats de référence.
19. Procédé selon la revendication 1, l'étape (b) comprend en outre la mise en contact dudit échantillon avec un ou plusieurs produits de référence.
- 35 20. Procédé selon la revendication 1, dans lequel l'étape (d) comprend en outre, avant la détermination, l'ajout d'un ou de plusieurs produits de référence correspondant au ou aux produits.
- 40 21. Procédé selon la revendication 1, dans lequel l'étape (d) comprend en outre, avant la détermination, l'ajout d'un ou plusieurs analytes de référence correspondant au ou aux analytes métaboliques contenus dans ledit échantillon.
22. Procédé selon la revendication 1, dans lequel ledit mélange réactionnel est aqueux.
- 45 23. Procédé selon la revendication 22, dans lequel ledit réactif est non aqueux.
24. Procédé selon la revendication 23, dans lequel ledit réactif comprend un solvant organique.
25. Procédé selon la revendication 23, dans lequel ledit réactif comprend un alcool.
- 50 26. Procédé selon la revendication 25, dans lequel ledit réactif est le méthanol.
27. Procédé selon la revendication 1, dans lequel ladite spectrométrie de masse est la spectrométrie de masse en tandem.
- 55 28. Procédé selon la revendication 1, dans lequel le ou les substrats sont présents sous une forme sèche dans un récipient.

29. Procédé selon la revendication 1, dans lequel le mélange réactionnel est mis en contact avec un ou plusieurs produits de référence et un ou plusieurs analytes de référence et la quantité dudit ou desdits analytes métaboliques et dudit ou desdits produits contenus dans lesdits échantillons de test est déterminée relativement au dit ou auxdits produits de référence et au dit ou auxdits analytes de référence, par spectrométrie de masse.

5

30. Procédé selon la revendication 1, dans lequel l'échantillon est séparé en deux parties d'échantillon et chaque partie d'échantillon est soumise à (a) et (b) ;
la combinaison des deux mélanges réactionnels pour produire un échantillon de test; et
la soumission des mélanges réactionnels combinés à (c).

10

31. Procédé selon la revendication 1, dans lequel l'échantillon est séparé en deux parties d'échantillons, dans lequel (a) comprend la mise en contact de chaque partie d'échantillon avec un récipient individuel, chaque récipient contenant ledit ou lesdits substrats pour ladite ou lesdites enzymes sous une forme sèche;
la soumission de chaque partie d'échantillon à (b) ;
la combinaison des deux mélanges réactionnels pour produire un échantillon de test; et
la soumission des mélanges réactionnels combinés à (c).

15

32. Procédé selon la revendication 28, dans lequel ledit récipient est mis en contact avec une solution pour générer lesdites conditions dans lesquelles au moins l'une desdites enzymes est capable d'agir sur un substrat correspondant pour générer au moins un produit.

20

25

30


35

40

45

50

55



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 6258605 B [0007] [0024]
- US 5719035 A [0008]
- US 6455321 B [0010] [0024] [0070]
- US 5629210 A [0011]
- US 6670194 B [0012]

Non-patent literature cited in the description

- **RASHED M. S.** Clinical applications of tandem mass spectrometry: ten years of diagnosis and screening for inherited metabolic diseases. *Journal of Chromatography B*, 2001, vol. 758, 27-48 [0006]
- **IM W. B. et al.** Bacterial Degradation of Biotin. *J. Biol. Chem.*, 1973, vol. 248 (22), 7798-7805 [0009]
- **M. GALLI-KIENLE et al.** Evaluation of enzyme activities by gas chromatography-mass spectrometry: HMGCoA reductase and cholesterol 7 α -hydrolase. *Journal of Chromatography*, 1984, vol. 289, 267-276 [0013]
- **L. M. BENSON et al.** Simultaneous structure-activity determination of disulfiram photolysis products by on-line continuous-flow liquid secondary ion mass spectrometry and enzyme inhibition assay. *Journal of chromatography*, 1995, vol. 693, 162-166 [0014]
- **AUSUBEL et al.** Current Protocols in Molecular Biology. John Wiley & Sons, 1999, vol. 47 [0055]
- **HARLOW ; LANE.** Antibodies: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1988 [0055]
- **HARLOW ; LANE.** Using Antibodies: A Laboratory Manual. Cold Spring Harbor Press, 1999 [0055]
- Tietz Textbook of Clinical Chemistry. W.B. Saunders, 1999 [0055]
- **SCOPES.** Protein Purification: Principles and Practice. Springer-Verlag, 1993 [0056]
- **BURTON ; HARDING.** *J. Chromatoqr. A*, 1998, vol. 814, 71-81 [0056]
- **BURLINGAME et al.** *Anal. Chem.*, 1998, vol. 70, 647R-716R [0090]
- **CHACE DH ; HILLMAN SL ; VAN HOVE JLK.** *Naylor EW. Clin Chem*, 1997, vol. 43, 210613 [0090]
- **RASHED MS ; BUCKNALL MP ; LITTLE D et al.** *Clin Chem*, 1997, vol. 43, 112941 [0090]
- **MATERN D ; STRAUSS AW ; HILLMAN SL ; MAY-ATEPEK E ; MILLINGTON DS ; TREFZ FK.** *Pediatr Res*, 1999, vol. 46, 459 [0090]
- **MILLINGTON DS ; KODO N ; TERADA N ; ROE D ; CHACE DH.** *International Journal of Mass Spectrometry and Ion Processes*, 1991, vol. 111, 21128 [0090]

专利名称(译)	用于同时检测代谢酶活性和代谢物水平的质谱法		
公开(公告)号	EP1664325B1	公开(公告)日	2014-10-08
申请号	EP2004782672	申请日	2004-08-30
申请(专利权)人(译)	PERKINELMER LAS , INC.		
当前申请(专利权)人(译)	PERKINELMER医学版 , INC.		
[标]发明人	CERDA BLAS		
发明人	CERDA, BLAS		
IPC分类号	C12Q1/00 C12Q1/34 G01N33/53 G01N33/68 C12Q1/68		
CPC分类号	C12Q1/00 G01N33/6893 G01N2800/04		
优先权	10/652732 2003-08-29 US		
其他公开文献	EP1664325A4 EP1664325A1		
外部链接	Espacenet		

摘要(译)

本发明提供了一种检测个体代谢紊乱的方法。该方法包括 (a) 使含有 (i) 一种或多种代谢指示酶和 (ii) 一种或多种代谢分析物的样品与一种或多种酶的一种或多种底物接触以产生反应混合物, 条件是其中至少一种所述酶能够作用于相应的底物上以产生至少一种产物; (b) 使所述反应混合物与抑制所述一种或多种酶作用于相应底物的能力的试剂接触, 其中所述一种或多种代谢分析物和所述至少一种产物可溶于所述试剂中; 产生测试样品和 (c) 使用质谱法测定所述一种或多种代谢分析物和所述测试样品中包含的所述至少一种产物的存在或量, 其中确定所述一种或多种代谢分析物的存在或量和所述至少一种产品与所述代谢紊乱的存在或不存在相关。

Aminoacidopathies	
Phenylketonuria Phe ^{2,4,6} Phe, Phe/Tyr ⁷ Phe, Tyr ^{1,5}	Tyrosinemia Type I ^{1,4} Met, Tyr ^{2,6} Tyr ⁷ Tyr, Tyr/Phe
Maple syrup urine disease Leu+Ile ¹ Leu+Ile, Val ^{2,4,6} Leu, Val ⁵ Leu, Leu/Phe ⁷	Tyrosinemia II ² Tyr ⁷
Homocystinuria Met ^{1,2,4,6} Met, Met/Phe, Met/Leu ⁵ Met, Met/Phe ⁷	Argininemia Arg ² Arg, Arg/Orn
Hypermethioninemia Met ² Met, Met/Phe ⁷	PRO Hyperprolinemia ⁴ Pro/Phe
Citrullinemia Cit ^{1,2,6} Cit/Phe, Cit/Tyr ⁴ Cit, Orn/Cit, Cit/Arg ⁵ Cit, Cit/Arg ⁷	HHH ² Orn, HomoCit ⁷ Orn, Orn/Cit
Nonketotic hyperglycinemia Gly ^{2,4,5}	5-oxoprolinuria ² 5-Oxopro ⁴ pyroGlu/Phe
	Argininosuccinic aciduria ^{1,6} Cit ⁴ Cit/Tyr ⁵ Asa ⁷ Cit, Cit/Arg