



(11) **EP 2 267 449 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication: **29.12.2010 Bulletin 2010/52** (51) Int Cl.: **G01N 33/53 (2006.01)**

(21) Application number: **10009115.6**

(22) Date of filing: **11.04.2006**

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR**  
Designated Extension States:  
**AL BA HR MK YU**

(30) Priority: **12.04.2005 JP 2005114350**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:  
**06731578.8 / 1 870 710**

(71) Applicant: **Matsumori, Akira**  
**Minoh-shi,**  
**Osaka 562-0045 (JP)**

(72) Inventor: **Matsumori, Akira**  
**Minoh-shi,**  
**Osaka 562-0045 (JP)**

(74) Representative: **Blodig, Wolfgang**  
**Wächtershäuser & Hartz**  
**Patentanwaltpartnerschaft**  
**Weinstraße 8**  
**80333 München (DE)**

Remarks:

This application was filed on 01-09-2010 as a divisional application to the application mentioned under INID code 62.

(54) **Biomarker for diagnosing heart failure and the use thereof**

(57) The invention relates to the following method and to a kit for carrying out the method:  
a method for detecting heart failure characterized in that the method comprises determining a free immunoglobulin kappa light chain level and/or a free immunoglobulin

lambda light chain level of a specimen and comparing at least one value of the kappa chain level, the lambda chain level, and a kappa/lambda ratio with that of a healthy subject.

**EP 2 267 449 A1**

**Description**

## Technical Field

5 **[0001]** The present invention relates to a novel biomarker for diagnosis of heart disease and to use thereof.

## Background Art

10 **[0002]** Generally, heart failure is defined as a pathological condition in which heart function is deteriorated by various conditions such as an ischemic heart disease (e.g., myocardial infarction or angina pectoris), cardiomyopathy, or hypertension. Currently available biomarkers for diagnosing such a heart failure include atrial natriuretic peptide (ANP), brain natriuretic peptide (BNP), and N-terminal proBNP (NT-proBNP). Assaying such a marker can determine whether or not a patient suffers from heart failure, and how severe the heart failure is.

15 **[0003]** However, since the cause of many diseases including heart failure cannot be determined as a single factor, in actual clinical settings a specific disease cannot be definitely diagnosed through assaying only one type of biomarker. In addition, since they are found in blood, urine, etc. in response to progress of pathological conditions, biomarkers may fail to specifically diagnose the target pathological conditions when the progress of the conditions varies among individual patients. Therefore, currently, many diseases are generally diagnosed by use of a variety of markers in combination.

## 20 Disclosure of the Invention

## Problems to be Solved by the Invention

25 **[0004]** Thus, for the diagnosis of heart diseases such as heart failure, keen demand exists for development of a novel marker other than known markers such as ANP and BNP.

## Means for Solving the Problems

30 **[0005]** The present inventor has carried out extensive studies in an attempt to develop a novel biomarker for diagnosing heart disease, and has paid attention to a free immunoglobulin light chain dissociated from an immunoglobulin heavy chain (i.e., free light chain (FLC)), which has never been employed for the diagnosis of heart disease.

35 **[0006]** Hitherto, FLC assay has been employed for the diagnosis of blood diseases such as myeloma and amyloidosis. In myeloma, FLC assay results vary depending on the type of tumor cells. Specifically, when the tumor cells produce kappa ( $\kappa$ ) chains, the kappa chain level and the kappa/lambda ratio increase, whereas when the tumor cells produce lambda ( $\lambda$ ) chains, the lambda chain level increases and the kappa/lambda ratio decreases. The type of myeloma has been diagnosed on the basis of the assay characteristics.

40 **[0007]** The aforementioned FLC is a type of pathological condition markers and is conceived to be a marker which indicates the activation of B lymphocytes. Therefore, if FLC can be employed for the diagnosis of heart failure, activation of B lymphocytes can possibly be detected in the diagnosis of heart failure. If the activation can be detected, screening and development of a novel heart failure therapeutic drug can be attained through regulating activation of B lymphocytes. Thus, FLC is a promising candidate for a useful marker, which differs from conventional markers.

45 **[0008]** The present inventor has further carried out extensive studies on application of FLC to the diagnosis of heart disease, and have found that FLC, which has never been employed for the diagnosis of heart disease, can be applied to the diagnosis of heart disease including heart failure. The present invention has been accomplished on the basis of this finding. Accordingly, the present invention is directed to the following.

**[0009]** [1] A method for detecting heart disease, **characterized in that** the method comprises determining a free kappa light chain level and/or a free lambda light chain level of a specimen and comparing at least one value of the kappa chain level, the lambda chain level, and a kappa/lambda ratio with that of a healthy subject.

50 **[0010]** [2] A method for detecting a complication of viral infection in a patient with heart disease, **characterized in that** the method comprises determining a free kappa light chain level and/or a free lambda light chain level of a specimen and comparing at least one value of the kappa chain level, the lambda chain level, and a kappa/lambda ratio with that of a healthy subject.

55 **[0011]** [3] A method for detecting severity of heart disease, **characterized in that** the method comprises determining a free kappa light chain level and/or a free lambda light chain level of a specimen and comparing at least one value of the kappa chain level, the lambda chain level, and a kappa/lambda ratio with that of a healthy subject or a patient suffering slight heart disease.

**[0012]** [4] A method as described in [1], [2], or [3] above, wherein the heart disease is heart failure, cardiomyopathy, myocarditis, or acute myocardial infarction.

**[0013]** [5] A method as described in [1], [2], or [3] above, wherein the free light chain levels are determined through immunoassay.

**[0014]** [6] A kit for diagnosing heart disease and for carrying out a method as recited in [1], [2], or [3] above, the kit comprising at least one reagent selected from the group consisting of a) to d):

- a) an immobilized FLC-specific antibody,
- b) a labeled FLC-specific antibody,
- c) an FLC-specific antibody, and
- d) a secondary antibody which can specifically bound to an antibody as recited in c) above.

**[0015]** [7] A kit as described in [6] above, wherein the heart disease is heart failure, cardiomyopathy, myocarditis, or acute myocardial infarction.

**[0016]** [8] A kit as described in [6] above, wherein the FLC levels are determined through immunoassay.

#### Effects of the Invention

**[0017]** The present invention has first elucidated that FLC is useful for the diagnosis of heart diseases such as heart failure, cardiomyopathy, myocarditis, and acute myocardial infarction. In addition, the invention has first elucidated that FLC is useful for the diagnosis of a complication of viral infection (e.g., infection with hepatitis C virus) in a patient with cardiomyopathy or myocarditis. Therefore, the method for detecting heart disease and the kit for diagnosing heart disease are remarkably useful for the diagnosis of heart disease as well as a complication of viral infection in a patient with heart disease.

**[0018]** In addition, since the kappa level and the lambda level correlate to the NT-proBNP level, which is useful for determination of severity and prediction of prognosis of heart diseases such as heart failure, FLC assay is useful for the determination of severity and prediction of prognosis of heart diseases such as heart failure.

**[0019]** Furthermore, the present invention realizes evaluation of the effects of a therapeutic drug for heart disease and that of an antiviral drug. Therefore, the invention is of great value in development of therapeutic agents for heart disease and viral infections.

#### Brief Description of the Drawing

##### **[0020]**

[Fig. 1] Images showing detection of kappa chains or lambda chains in the cardiac muscular tissue suffering heart failure (hematoxylin-eosin stain, FLC kappa (kappa chain detection, and FLC lambda (lambda chain detection).

#### Best Modes for Carrying Out the Invention

##### 1. Method for diagnosis of heart disease

**[0021]** As mentioned above, at least one FLC-related value of kappa chain level, lambda chain level, and kappa/lambda ratio reflects conditions and prognosis of heart disease. Therefore, heart disease can be diagnosed through quantitation of the free light chains (kappa chain and lambda chain) in specimens.

The specimen to be analyzed is blood, body fluid, tissue, excretions, etc. separated from subjects, clinical specimens, etc. Among these specimens, blood-originating specimens such as blood, serum, and plasma, and urine specimens are particularly preferred.

**[0022]** FLC assay may be performed through a known immunoassay technique (ELISA, RIA, etc.).

Specifically, in a preferred immunoassay procedure, a microamount of FLC present in a specimen can be determined at sufficient sensitivity, and no cross-reactivity to intact immunoglobulin is observed (e.g.,  $\leq 1\%$ ), whereby effects of components other than FLC can be minimized. Preferably, an enzymatic immunoassay method developed by Nakano *et al.* is employed (J. Immunol. Methods, 2003, 275, 9-17).

**[0023]** In order to determine FLC at higher sensitivity, preferably employed is an immunoassay method in which an FLC-specific polyclonal antibody is used as an antibody for trapping FLC in a specimen, and an FLC-specific monoclonal antibody is used as an antibody for detecting the trapped FLC.

**[0024]** The two types of antibodies employed in the invention are selected from FLC-specific monoclonal antibodies and FLC-specific polyclonal antibodies. Such antibodies may be antibodies *per se* or active fragments thereof [F(ab')<sub>2</sub>, Fab', etc.].

**[0025]** The FLC-specific antibody refers to an antibody which exhibits very weak reactivity to a light chain bound to

the heavy chain present in intact immunoglobulin or which does not recognize the light chain, but which strongly reacts with a light chain not bound to the heavy chain.

**[0026]** These FLC-specific monoclonal antibodies and polyclonal antibodies may be produced through a conventional method. For example, monoclonal antibodies are prepared by use of a free kappa chain or a free lambda chain as an antigen, and an antibody which does not bind to intact immunoglobulin and which specifically binds to FLC is selected from the thus-produced antibodies.

**[0027]** The FLC-specific polyclonal antibody may be produced through a conventional method. For example, polyclonal-antibodies are prepared by use of a free kappa chain or a free lambda chain, and an antibody which binds to intact immunoglobulin is absorbed and removed from the thus-produced polyclonal-antibodies. Such antibodies, which are now commercially available, may also be employed. However, FLC specificity of such commercial products must be checked in advance.

**[0028]** The immunoassay method employed in the invention is the same as conventionally employed method in terms of technique, procedure, etc. Specifically, examples of the antibody for trapping FLC in a specimen include an immobilized antibody fixed on a support.

**[0029]** Examples of the support employed for preparing such an immobilized antibody include generally employed supports, including synthetic organic polymer compounds such as polyvinyl chloride, polystyrene, styrene-divinylbenzene copolymers, styrene-maleic anhydride copolymers, nylon, polyvinyl alcohol, polyacrylamide, polyacrylonitrile, polypropylene, and polymethylene methacrylate; polysaccharides such as dextran derivatives (e.g., Sephadex), agarose gel (e.g., Sepharose or Biogel), and cellulose (e.g., paper disk or filter paper); and inorganic polymer compounds such as glass, silica gel, and silicone. Such a support may have an introduced functional group (e.g., an amino group, a carboxyl group, a carbonyl group, a hydroxyl group, or a sulfhydryl group).

**[0030]** Such a support may assume any form, such as a plate form (e.g., microtiter plate or disk), a particulate form (e.g., beads), a tubular form (e.g., test tube), a fibrous form, a membrane form, a microparticulate form (e.g., latex particles), a capsule form, or an endoplasmic reticulum form. A support of suitable form may be appropriately selected in consideration of the assay method employed.

**[0031]** Binding of an antibody to a support may be carried out through a known technique such as physical adsorption, ionic binding, covalent binding, or entrapment [see, for example, "Koteika Koso" ("Immobilized Enzyme") (edited by Ichiro Senhata, published by Kodansha Ltd. on March 20, 1975)]. Particularly, physical adsorption is preferred, from the viewpoint of convenience. An antibody may be bound directly to a support, or a linker may be provided between an antibody and a support.

In order to suppress non-specific binding, the thus-prepared immobilized reagent may be subjected to blocking treatment by use of a generally employed blocking agent such as gelatin or BSA.

**[0032]** Examples of the antibody for detecting trapped FLCs include an antibody labeled with a labeling agent. Examples of labeling agents employed include radioisotopes such as  $^{32}\text{P}$ ,  $^3\text{H}$ ,  $^{14}\text{C}$ , and  $^{125}\text{I}$ ; enzymes such as  $\beta$ -galactosidase, peroxidase, alkaline phosphatase, glucose-6-phosphate dehydrogenase, catalase, glucose oxidase, lactate oxidase, alcohol oxidase, and monoamine oxidase; coenzymes and prosthetic groups such as FAD, FMN, ATP, biotin, and heme; fluorescein derivatives such as fluorescein isothiocyanate and fluorescein thiofurbamyl; rhodamine derivatives such as tetramethylrhodamine B isothiocyanate; fluorescent dyes such as umbelliferone and 1-anilino-8-naphthalenesulfonate; and luminol derivatives such as luminol, isoluminol, and N-(6-aminoethyl)-N-ethylisoluminol.

**[0033]** Binding of an antibody to a labeling agent may be carried out through a method appropriately selected from among known methods described in the literature [for example, "Zoku Seikagaku Jikken Koza 5, Men-eki Seikagaku Kenkyu-ho" ("Sequel to Biochemical Experiments 5, Immunobiochemical Studies"), published by Tokyo Kagaku Dojin Co., Ltd. (1986), pp. 102-112].

**[0034]** Assay employing such an immobilized antibody and labeled antibody may employ a typical immunoassay procedure as it is. Specifically, an immobilized antibody is reacted with a test sample, followed by optional BF separation, and the resultant product is reacted with a labeled antibody (two-step method), or an immobilized antibody, a test sample, and a labeled antibody are reacted simultaneously (one-step method); and subsequently, FLCs contained in the sample are detected or quantitatively determined through a method which is known *per se*.

**[0035]** Details of immunoassay are described in, for example, the following references.

(1) "Zoku Rajio Imunoassei" ("Sequel to Radioimmunoassay") edited by Hiroshi Irie (published by Kodansha Ltd. on May 1, 1979)

(2) "Koso Men-eki Sokutei-ho" ("Enzyme Immunoassay") (2nd Edition) edited by Eiji Ishikawa, et al. (published by Igaku-Shoin Ltd. on December 15, 1982)

(3) Rinsho Byori Extra Edition No. 53 "Rinsho Kensa no tameno Imunoassei -Gijutsu to Oyo- ("Immunoassay for Clinical Tests -Techniques and Applications-") (published by The Clinical Pathology Press, 1983)

(4) "Biotechnology Encyclopedia" (published by CMC Publishing Co., Ltd. on October 9, 1986)

[0036]

- (5) "Methods in ENZYMOLOGY Vol. 70" (Immunochemical techniques (Part A))  
 (6) "Methods in ENZYMOLOGY Vol. 73" (Immunochemical techniques (Part B))  
 (7) "Methods in ENZYMOLOGY Vol. 74" (Immunochemical techniques (Part C))  
 (8) "Methods in ENZYMOLOGY Vol. 84" (Immunochemical techniques (Part D: Selected Immunoassay))  
 (9) "Methods in ENZYMOLOGY Vol. 92" (Immunochemical techniques (Part E: Monoclonal Antibodies and General Immunoassay Methods))

[(5) to (9), published by Academic Press]

**[0037]** The pathological condition and prognosis of heart disease is diagnosed on the basis of at least one value of the thus-determined free kappa light chain level, free lambda light chain level, and kappa/lambda ratio.

Examples of the target heart disease include heart failure, cardiomyopathy, myocarditis, acute myocardial infarction, and these heart diseases concomitant with viral infection.

**[0038]** The diagnosis is performed by comparing at least one value of the free kappa light chain level, free lambda light chain level, and kappa/lambda ratio of a specimen of a patient with that of a healthy subject. The diagnosis realizes detection of heart disease, a complication of viral infection in a patient with heart disease, or severity of heart disease, as well as discrimination between acute myocardial infarction and myocarditis or heart failure.

**[0039]** As mentioned in Example 1 hereinbelow, the kappa chain level, the lambda chain level, and the kappa/lambda ratio vary in response to the pathological conditions and correlate to the response of conventional markers and conventional heart function indices.

Therefore, on the basis of the aforementioned values, heart disease, a complication of viral infection in a patient with heart disease, or severity of heart disease can be detected. The detection of heart disease includes differential diagnosis of heart failure, cardiomyopathy, myocarditis, and acute myocardial infarction.

## 2. Kit for diagnosing heart disease

**[0040]** The kit of the present invention, for carrying out the aforementioned heart disease diagnosing method, includes, for example, the following reagents:

- 1) an immobilized FLC-specific antibody and
- 2) a labeled FLC-specific antibody.

**[0041]** The FLC-specific antibody for forming the immobilized FLC-specific antibody and the labeled FLC-specific antibody may be an FLC-specific monoclonal antibody or an FLC-specific polyclonal antibody. Two FLC-specific monoclonal antibodies may be used in combination, or an FLC-specific polyclonal antibody and an FLC-specific monoclonal antibody may be used in combination. Among them, a combination of an immobilized FLC-specific polyclonal antibody and a labeled FLC-specific monoclonal antibody is preferred, from the viewpoint of sensitivity.

**[0042]** Instead of a labeled FLC-specific antibody, a non-labeled FLC-specific antibody and a secondary antibody may be used. In this case, the kit includes the following reagents:

- 1) an immobilized FLC-specific antibody,
- 2) an FLC-specific antibody, and
- 3) a labeled anti-immunoglobulin antibody.

**[0043]** As is the above case, the immobilized FLC-specific antibody or the FLC-specific antibody may be a monoclonal antibody or a polyclonal antibody. A combination of an immobilized FLC-specific polyclonal antibody and an FLC-specific monoclonal antibody is preferred, from the viewpoint of sensitivity.

**[0044]** If required, in addition to the aforementioned reagents, the kit of the present invention includes an appropriate reagent, depending on the assay method, selected from among a color developing reagent, a reaction-terminating reagent, a standard antigen reagent, a sample pre-treating reagent, and other reagents.

## Examples

**[0045]** The present invention will next be described by way of examples, which should not be construed as limiting the invention thereto.

## EP 2 267 449 A1

### (1) Serum specimens

**[0046]** Serum specimens tested in the experiment were collected from healthy volunteers, heart failure patients, cardiomyopathy patients, and myocarditis patients.

### (2) FLC-specific monoclonal antibody

**[0047]** An FLC-specific monoclonal antibody was produced through immunizing BALB/c mice with FLC (product of Bethyl laboratories, Montgomery, TX) (J. Immunol. Methods, 275, 9-17. (2003)). An F(ab')<sub>2</sub> fragment of the produced antibody was labeled with horseradish peroxidase (HRP) (J. Histochem. Cytochem., 22, 1084-91. (1974)).

### (3) FLC assay

**[0048]** FLC assay was performed according to a description (J. Immunol. Methods, 2003, 275, 9-17). Specifically, a specimen or a standard concentration solution (each 100 μL) was added to a 96-well microplate (Nunc) coated with the FLC-specific monoclonal antibody, followed by reaction at room temperature for two hours. After washing, an HRP labeled anti-kappa light chain antibody or an HRP-labeled anti-lambda light chain antibody (each 100 μL) serving as secondary antibodies were added to the microplate, followed by reaction at 37°C for 30 minutes. After reaction, the microplate was washed with PBS containing 0.05% Tween 20, and the reactions were allowed to develop color by dispensing to each well a color developing liquid (OPD, product of Sigma) (each 100 μL). The color development was terminated with 1N sulfuric acid, and absorbance at 492 nm was measured. In the assay, 50mM Tris-HCl (pH 7.5) containing 1% bovine serum albumin was employed as a buffer.

### Example 1

**[0049]** Heart failure patients (63) (from Dec. 2002 to Nov. 2004) were investigated as follows. Among the patients, 17 patients were investigated in terms of NT-proBNP level and stem cell factor (SCF) level, with correlation to FLC levels. The results are shown in Table 1.

### **[0050]**

[Table 1]

X	NT-proBNP(y)			SCF(y)		
		r <sup>2</sup>	P		r <sup>2</sup>	p
Kappa	y=280x-4920	0.511	0.0013	y=12.9x+381	0.3518	0.012
Lambda	y=51.6x-455	0.806	<0.0001	y=1.72x+642	0.289	0.0259
Kappa/lambda	y=-32600x+18900	0.564	0.0005		0.0641	0.327

### Example 2

**[0051]** Serum FLC levels determined in a multi-center clinical trial on myocarditis treatment are shown below. (1) Table 2 shows serum FLC levels of all myocarditis patients.

### **[0052]**

[Table 2]

	n	kappa (mg/L)	lambda (mg/L)	kappa/lambda
Myocarditis	1,318	37.6±19.2**	72.4±124.2*	0.64±0.32
Healthy subjects	17	27.9±5.3	43.6±8.8	0.66±0.16
Mean ± standard deviation, *: p<0.0005, **p<0.0001				

**[0053]** (2) Table 3 shows the effect of infection with hepatitis C virus (HCV) on FLC levels.

### **[0054]**

EP 2 267 449 A1

[Table 3]

	n	kappa (mg/L)	lambda (mg/L)	kappa/lambda
HCV (+)	42	30.6±14.8*	173.1±37.1*	0.27±0.22*
HCV (-)	1,276	37.9±19.3*	69.1±81.7*	0.66±0.31
Healthy subjects	17	27.9±5.3	43.6±8.8	0.66±0.16

Mean ± standard deviation, \*p<0.0001 with respect to healthy subject, HCV (+,-), p<0.0001

[0055] (3) Table 4 shows FLC levels of hepatitis C virus antibody-positive cardiomyopathy patients.  
[0056]

[Table 4]

	n	kappa	lambda	kappa/lambda
Hepatitis C virus 5 antibody (+)		42.5±27.2*	79.9±74.0*	0.558±0.338
Healthy subjects	17	27.9±5.3	43.6±8.8	0.661±0.162

\*p<0.05

(4) Relationship between FLC levels and severity of heart failure

[0057] The tested subjects were classified into two categories in terms of severity (slight (grades I + II) and severe (grades III + IV)). Each grade denotes a heart function status on the basis of subjective symptoms graded by the New York Heart Association (NYHA). The results are shown in Table 5.

[0058]

[Table 5]

	NYHA heart function scale			p
	Slight (grades I + II) (n=17)	Severe (grades III + IV) (n=63)		
Kappa	32.0±15.4	40.8±39.2		NS
Lambda	55.0±44.0	93.2±67.9		0.03
Kappa/lambda	0.67±0.31	0.46±0.14		0.0001

(5) Results of Examples 1 and 2

[0059] (5-1) The kappa and lambda levels of heart failure patients positively correlated to NT-proBNP level, which relates to severity and prognosis of heart failure. The lambda level strongly correlated to NT-proBNP level, whereas the kappa/lambda ratio negatively correlated to NT-proBNP level.

[0060] (5-2) The kappa and lambda levels of heart failure patients positively correlated to the SCF level.

(5-3) The kappa and lambda levels of myocarditis patients increased, whereas the kappa/lambda ratio was unchanged.

(5-4) The kappa and lambda levels of myocarditis patients positively correlated to NT-proBNP level, whereas the kappa/lambda ratio negatively correlated to NT-proBNP level.

[0061] (5-5) The kappa and lambda levels of myocarditis patients concomitant with infection with hepatitis C virus increased. Of these, the lambda level considerably increased, and the kappa/lambda ratio considerably decreased.

(5-6) The kappa and lambda levels of cardiomyopathy patients positive to infection with hepatitis C virus increased, whereas the kappa/lambda ratio tended to decrease.

[0062] (5-7) Severe heart failure patients exhibited a lambda level higher than that of slight heart failure patients. The kappa/lambda ratio of severe heart failure patients decreased. Thus, the lambda level and the kappa/lambda ratio of heart failure patients reflect the severity of heart failure.

(5-8) The lambda level in heart failure concomitant with infection with hepatitis C virus further elevated but the kappa/lambda ratio lowered.

Example 3

**[0063]** Heart failure patients (48) were investigated as follows. (1) The free kappa light chain level, free lambda light chain level, and kappa/lambda ratio of heart failure patients (48) were compared with those of healthy subjects (17). The study has revealed that, in heart failure patients, the free lambda light chain level increases, and the kappa/lambda ratio decreases.

**[0064]**

[Table 6]

	n	kappa	lambda	kappa/lambda
Heart failure	48	32.7±0.5	80.2±57.5*	0.46±0.12**
Healthy subjects	17	27.9±5.3	43.6±8.8	0.66±0.16

Mean ± standard deviation, t-test, \* p<0.05, \*\*p<0.0001

**[0065]** (2) The relationship between survival time and FLC levels in heart failure patients (48) was investigated. The study has revealed that a negative correlation is found between the survival time and the lambda chain level.

**[0066]**

[Table 7]

X		Survival time (day) (y)		
		r	r <sup>2</sup>	p
Lambda	y=-6.2x+1010	-0.48	0.23	p<0.05

**[0067]** (3) The relationship between the free kappa light chain level, free lambda light chain level, and kappa/lambda ratio and the NT-proBNP (product of Roche Diagnostics) level, which serves as an index for heart failure and myocardial disorders was investigated.

The study has revealed that the kappa chain and lambda chain levels exhibit positive correlation to the NT-proBNP level but negative correlation to the kappa/lambda ratio. Thus, these FLC levels can serve as an index for heart failure and myocardial disorders.

**[0068]**

[Table 8]

X		NT-proBNP (pg/mL) (y)		
		r	r <sup>2</sup>	p
Kappa	y=246x-3290	0.40	0.16	p<0.005
Lambda	y=54.7x+341	0.49	0.24	p<0.0005
Kappa/lambda	y=-18800x+13300	-0.36	0.13	p<0.05

**[0069]** (4) The relationship between the free kappa light chain level, free lambda light chain level, and kappa/lambda ratio and the pulmonary arterial pressure determined by means of cardiac catheterization was investigated.

The kappa chain and lambda chain levels exhibited positive correlation to the mean pulmonary arterial pressure.

**[0070]**

[Table 9]

X		Mean pulmonary arterial pressure (mmHg) (y)		
		r	r <sup>2</sup>	p
Kappa	0.71x+8.7	0.54	0.30	p<0.05
Lambda	0.22x+14	0.48	0.23	p<0.05

**[0071]** (5) The relationship between the free kappa light chain level, free lambda light chain level, and kappa/lambda

EP 2 267 449 A1

ratio and the cardiac output determined by means of cardiac catheterization was investigated.

The kappa chain and lambda chain levels exhibited significant negative correlation to the cardiac output, thereby revealing that these FLC levels relate to heart functions.

[0072]

5

[Table 10]

X		Cardiac output (L/m <sup>2</sup> ) (y)		
		r	R <sup>2</sup>	p
Kappa	y=-0.4x+4.19	-0.78	0.61	p<0.0001
Lambda	y=-0.1x+3.74	-0.59	0.35	p<0.005

10

[0073] (6) The relationship between the free kappa light chain level, free lambda light chain level, and kappa/lambda ratio and the left ventricular internal dimension in systole (LVDs), left ventricular internal dimension in diastole (LVDd), and ejection fraction (EF) serving as an index for cardiac systolic function, which were determined by means of an ultrasonic cardiogram was investigated.

15

The lambda chain level was found to exhibit significant positive correlation to LVDd and LVDs and thus is conceived to be related to dimension of left ventricular cavity. The kappa/lambda ratio was found to exhibit negative correlation to LVDd and LVDs and positive correlation to EF and thus is conceived to be related to dimension of left ventricular cavity and cardiac systolic function.

20

[0074]

[Table 11]

X		LVDd (mm) (y)		
		R	r <sup>2</sup>	P
Kappa	y=0.41x+62.8	0.25	0.06	p=0.24
Lambda	y=0.33x+51.3	0.58	0.33	p<0.005
Kappa/lambda	y=-98x+119	-0.72	0.51	p<0.0001

25

30

X		LVDd (mm) (y)		
		R	r <sup>2</sup>	P
Kappa	y=0.42x+52.1	0.25	0.06	p=0.24
Lambda	y=0.33x+40.8	0.57	0.32	p<0.005
Kappa/lambda	y=-102x+110	-0.73	0.53	p<0.0001

35

40

X		EF (%) (y)		
		R	r <sup>2</sup>	P
Kappa/lambda	y=67.4x-0.14	0.59	0.35	p<0.005

45

[0075] (7) The correlation of the free kappa light chain level, free lambda light chain level, and kappa/lambda ratio to the CRP level serving as an index for inflammatory reaction was investigated.

The kappa chain and lambda chain levels were found to exhibit significant positive correlation to the CRP level and reflect inflammatory reaction.

50

[0076]

[Table 12]

X		CRP (mg/dL) (y)		
		R	r <sup>2</sup>	p P
Kappa	y=0.3x-6.89	0.5	0.25	p<0.05

55

EP 2 267 449 A1

(continued)

5

X		CRP (mg/dL) (y)		
		R	r <sup>2</sup>	p P
Lambda	y=0.95x-4.54	0.46	0.21	p<0.05
Kappa/lambda	y=-7.8+5.73	-0.16	0.02	p=0.46

10

**[0077]** (8) The correlation of the free kappa light chain level, free lambda light chain level, and kappa/lambda ratio to the type IV collagen level, serving as an index for fibrogenesis was investigated.

The kappa chain and lambda chain levels were found to exhibit positive correlation to the type IV collagen level, and the kappa/lambda ratio exhibit negative correlation to the type IV collagen level. Thus, FLC levels were found to be related to a marker for fibrogenesis.

15

**[0078]**

[Table 13]

20

X		Type IV collagen (ng/mL) (y)		
		R	r <sup>2</sup>	P
Kappa	y=0.48x+20.5	0.62	0.39	p<0.0001
Lambda	y=1.1x+89.3	0.78	0.61	p<0.0001
Kappa/lambda	y=-250x+291	-0.38	0.14	p<0.01

25

**[0079]** (9) Paraffin sections of cardiac muscular specimens of heart failure patients were stained with hematoxylin-eosin, and the diameters of cardiac muscular cells were measured, and the relationship between cell diameter and FLC level was investigated.

The cardiac muscular cell diameter was found to exhibit negative correlation to the kappa/lambda ratio.

**[0080]**

30

[Table 14]

35

x		Cardiac muscular cell diameter (μm) (y)		
		R	r <sup>2</sup>	P
Kappa/lambda	y=-31.8x+44.9	-0.63	0.39	p<0.005

40

**[0081]** (10) The relationship between the FLC levels and the activity of mast cells was investigated. RNA was extracted from cardiac muscular tissue specimens which had been sampled during surgery of a heart failure patient and preserved in a frozen state. Through real-time PCR, expression of mRNAs of stem cell factor (SCF), metalloprotease (MMP-9), chymase, and tryptase was investigated. The amount of each expressed mRNA was corrected by a housekeeping gene, GAPDH.

The kappa/lambda ratio was found to exhibit positive correlation to the expression amount of SCF, MMP-9, chymase, and tryptase and is conceived to be related to activation of mast cells.

45

**[0082]**

[Table 15]

50

x		SCF/GAPDH (y)		
		R	r <sup>2</sup>	P
Kappa/lambda	y=12900x-4590	0.46	0.21	p<0.05

55

x		MMP-9/GAPDH (y)		
		R	r <sup>2</sup>	P
Kappa/lambda	y=651x-231	0.44	0.19	p<0.05

(continued)

x		Chymase/GAPDH (y)		
		R	r <sup>2</sup>	P
Kappa/lambda	y=349x-124	0.43	0.19	p<0.05

x		Tryptase/GAPDH (y)		
		R	r <sup>2</sup>	P
Kappa/lambda	y=1050x-373	0.43	0.18	p<0.05

**[0083]** (11) Distribution of bound kappa chains and lambda chains in the cardiac muscular tissue of a heart failure patient was investigated.

Specifically, slices of 5 micrometers square were cut out of a paraffin block of a formalin-fixed specimen of the heart suffering heart failure. Each slice was deparaffinized with xylene, dehydrated with alcohol, and treated with hydrogen peroxide for inactivating endogenous peroxidase.

The heart specimens were incubated with Mouse monoclonal antibodies (products of Yamasa corporation, Chiba) to a kappa chain and to a lambda chain, serving as a primary antibody, overnight at 4°C, and immunostained by means of a Vectastain ABC kit (Burlingame, CA, U.S.A.) for mouse IgG, followed by color development with DAB.

As shown in Fig. 1, kappa chains and lambda chains were observed in leucocytes in the cardiac muscular tissue suffering heart failure. Thus, leucocytes infiltrating the cardiac muscular tissue were found to produce kappa chains and lambda chains.

**[0084]** (12) Urine FLC levels of heart failure patients (20) were determined, and the determined values were compared with those of healthy subjects.

In heart failure patients, the urine kappa chain and lambda chain levels were significantly higher than those of the healthy subjects.

**[0085]**

[Table 16]

	n	kappa (mg/L)	lambda (mg/L)	kappa/lambda
Healthy subjects	29	2.14±2.34	2.79±3.49	0.82±0.29
Heart failure	20	5.74±7.56*	7.10±8.75*	0.82±0.29

\*p<0.05 with respect to healthy subjects

**[0086]** (13) Correlation of urine FLC levels to blood FLC levels in heart failure patients (20) was investigated.

Significant positive correlation was observed between the urine kappa chain level, lambda chain level, and kappa/lambda ratio and the blood kappa chain level, lambda chain level, and kappa/lambda ratio.

**[0087]**

[Table 17]

	N	r	r <sup>2</sup>	P
Urine kappa (y) : blood kappa (x)	y=0.165x-6.28	0.45	0.20	p<0.05
Urine lambda (y) : blood lambda (x)	y=0.0936x-3.5	0.58	0.34	p<0.01
Urine kappa/lambda (y) : blood kappa/lambda (x)	y=0.922x+0.156	0.65	0.42	p<0.005

**[0088]**

(14) Results of Example 3

(14-1) In heart failure patients, the lambda chain level increased, and the kappa/lambda ratio decreased.

(14-2) The lambda chain level related to prognosis (survival time) of heart failure patients.

(14-3) The kappa chain and lambda chain levels correlated to NT-proBNP level, which is a marker for heart failure and for myocardial disorders, and to CRP level, which is a marker for inflammation.

(14-4) The kappa chain and lambda chain levels related to cardiac functions, and the lambda chain level and the kappa/lambda ratio correlated to dimension of intracardiac cavity.

(14-5) The kappa chain and lambda chain levels and the kappa/lambda ratio correlated to the level of fibrogenesis marker.

(14-6) The kappa/lambda ratio reversely correlated to hypertrophy of myocardial cells.

(14-7) The kappa/lambda ratio correlated to the level of mast-cell-related mediators.

(14-8) Leucocytes infiltrating the cardiac muscular tissue suffering heart failure produced kappa chains and lambda chains.

(14-9) Urine kappa chain and lambda chain levels of heart failure patients increased.

(14-10) The urine kappa chain level, lambda chain level, and kappa/lambda ratio of heart failure patients correlated to the blood kappa chain level, lambda chain level, and kappa/lambda ratio.

Example 4

**[0089]** FLC levels of acute myocardial infarction patients were measured in the onset day, 24 hours after, and four weeks after, and these values were compared with those of healthy subjects.

The kappa chain levels of the patients significantly decreased on the day of myocardial infarction onset and 24 hours after. However, no significant difference was found in the lambda chain level and the kappa/lambda ratio between the patients and the healthy subjects.

Thus, the measurement of the free kappa light chain level is useful for the diagnosis of acute myocardial infarction and also for the differential diagnosis of acute myocardial infarction from myocarditis and heart failure.

**[0090]**

[Table 18]

	n	kappa (mg/L)	lambda (mg/L)	kappa/lambda
Healthy subjects	17	27.9±5.3	43.6±8.8	0.66±0.16
Acute heart infarction				
Onset day	70	23.6±8.3*	50.2±40.5	0.54±0.20
24 hours	65	22.5±6.8*	48.7±42.5	0.56±0.23
4 weeks	36	27.4±7.8	53.6±19.8	0.53±0.14

One-way analysis of variance: F=5.0; p<0.005

Intergroup comparison: Fisher's LDS, \*p<0.05 (vs. healthy subjects, 4 weeks after)

**[0091]** As described hereinabove, FLC level measurement is useful for the diagnosis of heart failure, cardiomyopathy, myocarditis, and acute myocardial infarction. Since the kappa chain and lambda chain levels increase and the kappa/lambda ratio decreases in patients suffering cardiomyopathy or myocarditis concomitant with infection with hepatitis C virus, FLC levels are useful for the diagnosis of infection of such patients with hepatitis C virus and other viruses.

**[0092]** In heart failure patients concomitant with HCV infection, the lambda level further increases and the kappa/lambda ratio decreases, as compared with heart failure patients not infected with HCV. Therefore, coexistence of infection in a patient can be diagnosed.

According to the present invention, the effects of a heart failure therapeutic drug and an anti-viral drug can be evaluated. Therefore, the invention is of great value in development of therapeutic agents for heart failure, cardiomyopathy, and myocarditis as well as viral infections.

**[0093]** Severe heart failure patients exhibited a lambda level higher than that of slight heart failure patients. The kappa/lambda ratio of severe heart failure patients decreased. Thus, the lambda level and the kappa/lambda ratio correlated to the severity of heart failure. In addition, since the kappa chain and lambda chain levels correlate to the severity of

heart failure and to the NT-proBNP level, which is a useful index for prognosis of heart failure, FLC level measurement is useful for evaluation of severity and prognosis of heart failure.

The content of European patent application no. 06 731 578.8 (parent case) including entire description, all claims and figures is incorporated herein by reference in its entirety.

5

### Claims

1. A method of detecting heart failure, wherein the method comprises:

10

determining a free immunoglobulin kappa light chain level and/or a free immunoglobulin lambda light chain level of a specimen; and  
comparing at least one value of the lambda chain level and a kappa/lambda ratio with that of a healthy subject.

15

2. The method according to claim 1, wherein, in heart failure patients, the free lambda light chain level increases and the kappa/lambda ratio decreases.

3. The method according to claim 1 or 2, wherein the free immunoglobulin light chain levels are determined through immunoassay.

20

4. The method according to any one of claims 1 to 3, wherein the specimen is a blood, serum, plasma or urine specimen.

5. Use of a kit for carrying out a method as recited in claim 1, the kit comprising at least one reagent from the group of reagents consisting of a) to d):

25

- a) an immobilized free immunoglobulin light chain-specific antibody,
- b) a labelled free immunoglobulin light chain-specific antibody,
- c) a free immunoglobulin light chain-specific antibody, and
- d) a secondary antibody which can specifically bind to an antibody as recited in c) above.

30

35

40


45

50

55

Fig. 1

**× 600**



Hematoxylin-eosin  
staining

**FLC kappa**

**FLC lambda**



EUROPEAN SEARCH REPORT

Application Number  
EP 10 00 9115

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
Y	<p>NAKAMURA MOTOYUKI ET AL: "Reversible restrictive cardiomyopathy due to light-chain deposition disease." MAYO CLINIC PROCEEDINGS. MAYO CLINIC FEB 2002 LNKD- PUBMED:11838655, vol. 77, no. 2, February 2002 (2002-02), pages 193-196, XP002607102 ISSN: 0025-6196 * page 193, column 1, lines 1-9 * * page 193, column 1, lines 32-34 * * page 194, column 1, lines 32-34 * * page 194, column 2, lines 2-4 * * page 194, column 2, lines 8-11 * * page 195, column 1, lines 11-13 * * page 195, column 1, lines 2-7 * * page 196, column 1, lines 2-5 *</p> <p style="text-align: center;">-----</p>	1-5	INV. G01N33/53
Y	<p>DUBREY S W ET AL: "The clinical features of immunoglobulin light-chain (AL) amyloidosis with heart involvement." QJM : MONTHLY JOURNAL OF THE ASSOCIATION OF PHYSICIANS FEB 1998, vol. 91, no. 2, February 1998 (1998-02), pages 141-157, XP002485302 ISSN: 1460-2725 * page 141, column 1, line 17 - page 141, column 2, line 7 * * page 143, column 2, line 31 - page 143, column 2, line 41 * * page 144, column 2, line 9 - page 145, column 1, line 1 *</p> <p style="text-align: center;">-----</p> <p style="text-align: center;">-/--</p>	1-5	<p>TECHNICAL FIELDS SEARCHED (IPC)</p> <p>G01N</p>
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
Munich		27 October 2010	Landré, Julien
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>&amp; : member of the same patent family, corresponding document</p>			

1  
EPO FORM 1503 03.02 (P04C01)



EUROPEAN SEARCH REPORT

Application Number  
EP 10 00 9115

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
Y	<p>SANCHORAWALA VAISHALI ET AL: "Pulsed low dose intravenous melphalan in patients with AL amyloidosis, ineligible for aggressive treatment with high-dose melphalan and stem cell transplantation." BLOOD, vol. 104, no. 11, Part 1, November 2004 (2004-11), pages 657A-658A, XP002485170 &amp; 46TH ANNUAL MEETING OF THE AMERICAN-SOCIETY-OF-HEMATOLOGY; SAN DIEGO, CA, USA; DECEMBER 04 -07, 2004 ISSN: 0006-4971 * abstract *</p>	1-5	
Y	<p>KATZMANN JERRY A ET AL: "Serum reference intervals and diagnostic ranges for free kappa and free lambda immunoglobulin light chains: Relative sensitivity for detection of monoclonal light chains" CLINICAL CHEMISTRY, AMERICAN ASSOCIATION FOR CLINICAL CHEMISTRY, WASHINGTON, DC, vol. 48, no. 9, 1 September 2002 (2002-09-01), pages 1437-1444, XP002378160 ISSN: 0009-9147 * abstract * * page 1438, column 1, line 16 - page 1438, column 1, line 21 * * page 1439, column 2, line 17 - page 1439, column 2, line 21 * ----- -/--</p>	1-5	
The present search report has been drawn up for all claims			
Place of search <b>Munich</b>		Date of completion of the search <b>27 October 2010</b>	Examiner <b>Landré, Julien</b>
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p>		<p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- &amp; : member of the same patent family, corresponding document</p>	

1  
EPO FORM 1503 03-82 (P04C01)



EUROPEAN SEARCH REPORT

Application Number  
EP 10 00 9115

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
Y	HSU J Y ET AL: "The clinicopathological spectrum of renal amyloidosis." ZHONGHUA YI XUE ZA ZHI = CHINESE MEDICAL JOURNAL; FREE CHINA ED OCT 1994, vol. 54, no. 4, October 1994 (1994-10), pages 230-239, XP002485171 ISSN: 0578-1337 * abstract *	1-5	
Y	----- BURKHARDT O ET AL: "[Diagnosis of multiple myeloma by demonstrating plasma cells in bronchoalveolar lavage.]" DMW DEUTSCHE MEDIZINISCHE WOCHENSCHRIFT, vol. 128, no. 38, 19 September 2003 (2003-09-19), pages 1951-1954, XP002485172 ISSN: 0012-0472 * abstract *	1-5	
Y	----- NAKANO T ET AL: "ELISAs for free human immunoglobulin light chains in serum: improvement of assay specificity by using two specific antibodies in a sandwich detection method" JOURNAL OF IMMUNOLOGICAL METHODS, ELSEVIER SCIENCE PUBLISHERS B.V.,AMSTERDAM, NL, vol. 293, no. 1-2, 1 October 2004 (2004-10-01), pages 183-189, XP004634285 ISSN: 0022-1759 * page 185, paragraph 2.5; table 2 * * page 183, column 2, line 7 - page 184, column 1, line 4 * -----	1-5	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (IPC)
Place of search Munich		Date of completion of the search 27 October 2010	Examiner Landré, Julien
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document	

1  
EPO FORM 1503 03.82 (P04C01)

**REFERENCES CITED IN THE DESCRIPTION**

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

**Patent documents cited in the description**

- EP 06731578 A [0093]

**Non-patent literature cited in the description**

- *J. Immunol. Methods*, 2003, vol. 275, 9-17 [0022] [0048]
- Koteika Koso. Kodansha Ltd, 20 March 1975 [0031]
- Zoku Seikagaku Jikken Koza 5, Men-eki Seikagaku Kenkyu-ho. Tokyo Kagaku Dojin Co., Ltd, 1986, 102-112 [0033]
- Zoku Rajio Imunoassei. Kodansha Ltd, 01 May 1979 [0035]
- Koso Men-eki Sokutei-ho. Igaku-Shoin Ltd, 15 December 1982 [0035]
- Rinsho Kensa no tameno Imunoassei -Gijutsu to Oyo-. The Clinical Pathology Press, 1983 [0035]
- Biotechnology Encyclopedia. CMC Publishing Co., Ltd, 09 October 1986 [0035]
- Immunochemical techniques. Methods in ENZYMOLOGY. vol. 70 [0036]
- Immunochemical techniques. Methods in ENZYMOLOGY. vol. 73 [0036]
- Immunochemical techniques. Methods in ENZYMOLOGY. vol. 74 [0036]
- Immunochemical techniques (Part D: Selected Immunoassay). Methods in ENZYMOLOGY. vol. 84 [0036]
- Immunochemical techniques (Part E: Monoclonal Antibodies and General Immunoassay Methods). Methods in ENZYMOLOGY. vol. 92 [0036]
- *J. Immunol.*, 2003, vol. 275, 9-17 [0047]
- *J. Histochem. Cytochem.*, 1974, vol. 22, 1084-91 [0047]

专利名称(译)	用于诊断心力衰竭的生物标志物及其用途		
公开(公告)号	<a href="#">EP2267449A1</a>	公开(公告)日	2010-12-29
申请号	EP2010009115	申请日	2006-04-11
[标]申请(专利权)人(译)	晃松森		
申请(专利权)人(译)	松森, AKIRA		
当前申请(专利权)人(译)	松森, AKIRA		
[标]发明人	MATSUMORI AKIRA		
发明人	MATSUMORI, AKIRA		
IPC分类号	G01N33/53		
CPC分类号	G01N33/6893 G01N33/6857 G01N2800/32		
优先权	2005114350 2005-04-12 JP		
其他公开文献	EP2267449B1		
外部链接	<a href="#">Espacenet</a>		

摘要(译)

本发明涉及以下方法和用于实施该方法的试剂盒：用于检测心力衰竭的方法，其特征在于该方法包括测定标本的游离免疫球蛋白κ轻链水平和/或游离免疫球蛋白λ轻链水平。并且将κ链水平，λ链水平和κ/λ比的至少一个值与健康受试者的值进行比较。

Fig. 1

