

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 September 2002 (26.09.2002)

PCT

(10) International Publication Number
WO 02/075318 A2

(51) International Patent Classification⁷: **G01N 33/68**,
C07D 277/62, G01N 33/58, C07D 279/18

of Aberdeen, University Medical Buildings, Foresterhill,
Aberdeenshire AB25 2ZD (GB).

(21) International Application Number: PCT/GB02/01318

(74) Agents: **KREMER, Simon, M.** et al.; Mewburn Ellis,
York House, 23 Kingsway, London, Greater London
WC2B 6HP (GB).

(22) International Filing Date: 20 March 2002 (20.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0106953.3 20 March 2001 (20.03.2001) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **THE
UNIVERSITY COURT OF THE UNIVERSITY OF
ABERDEEN** [GB/GB]; Regent Walk, Aberdeen, Ab-
erdeenshire AB24 3FX (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **WISCHIK, Claude,
Michel** [FR/GB]; Department of Mental Health, University
of Aberdeen, University Medical Buildings, Foresterhill,
Aberdeenshire AB25 2ZD (GB). **HARRINGTON,
Charles, Robert** [GB/GB]; Department of Mental Health,
University of Aberdeen, University Medical Buildings,
Foresterhill, Aberdeenshire AB25 2ZD (GB). **RICKARD,
Janet, Elizabeth** [GB/GB]; Department of Mental Health,
University of Aberdeen, University Medical Buildings,
Foresterhill, Aberdeenshire AB25 2ZD (GB). **HORSLEY,
David** [GB/GB]; Department of Mental Health, University

Published:

— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: NEUROFIBRILLARY LABELS

(57) Abstract: Disclosed are methods for determining the stage of neurofibrillary degeneration associated with a tauopathy in a subject believed to suffer from the disease, which methods comprise the steps of: (i) introducing into the subject a ligand capable of labelling aggregated paired helical filament (PHF) tau protein, (ii) determining the presence and/or amount of ligand bound to extracellular aggregated PHF tau in the medial temporal lobe of the brain of the subject, (iii) correlating the result of the determination made in (ii) with the extent of neurofibrillary degeneration in the subject. The methods can be used for pre-mortem diagnosis and staging of tauopathies such as Alzheimer's Disease. Preferred ligands include sulphonated-benzothiazole-like compounds and diaminophenothiazines. Novel ligands (e.g. sulphonated-benzothiazole-like compounds) are also provided. The method may also include the use of "blocking ligands" to block competing binding sites. In other aspects the invention provides in vitro methods for identifying ligands capable of labeling aggregated PHF tau protein, the methods comprising the steps of: (i) providing a first agent suspected of being capable of labeling aggregated PHF tau protein, (ii) contacting (a) a tau protein or a derivative thereof containing the tau core fragment bound to a solid phase so as to expose a high affinity tau capture site, with (b) a liquid phase tau protein or derivative thereof capable of binding to the solid phase tau protein or derivative, and (c) said selected first agent and (d) a second agent known to be tau-tau binding inhibitor, (iii) selecting first agent which fully or partially relieves the inhibition of binding of the liquid phase tau protein or derivative of (b) to the solid phase tau protein or derivative of (a) by the inhibitor (d). Ligands may also be tested to confirm that they are not themselves inhibitors.



WO 02/075318 A2

Neurofibrillary labelsField of the invention

5

The present invention concerns materials, methods and models relating generally to the labelling and detection of neurofibrillary tangles. In addition, it concerns the identification and development of ligands suitable for

10 neuropathological staging and their use in the diagnosis, prognosis or treatment of diseases such as Alzheimer's Disease (AD).

Background to the invention15 *Neuropathological staging and AD*

The neuropathological staging proposed by Braak (Braak, H et al. (1991), Acta. Neuropathol. 82, 239-259) provides the best available definition of progression of relatively pure neurofibrillary

20 degeneration of the Alzheimer-type which is diagnostic of AD (Wischnik et al. (2000), "Neurobiology of Alzheimer's Disease", Eds. Dawbarn et al., The Molecular and Cellular Neurobiology Series, Bios Scientific Publishers, Oxford). This staging is shown

25 schematically in terms of brain region in Fig 2B, and is based on a regular regional hierarchy of neurofibrillary tangle (NFT) distribution. Regions of the brain which appear earlier in the hierarchy have both more tangles and are affected in less severe cases than those later in the list.

30 *Relationship between AD, clinical dementia and neuropathological staging*

The provision of an effective pre-mortem assessment of Braak Stage would be of use in the assessment and treatment of AD, for which

35 the differential includes Lewy Body dementia, Parkinson's disease, various forms of fronto-temporal and cortico-basal degeneration, progressive supranuclear palsy and a range of rare neurological syndromes.

The original model proposed by Braak was essentially qualitative in nature, and was not linked to any implications about the threshold for development of clinical dementia and symptoms.

5

In terms of the appearance of clinical dementia by DSM-IV criteria, this corresponds statistically to the transition between Braak stages 3 and 4 (Fig 2c). The DSM-IV criteria (Diagnostic and Statistical Manual of Mental Disorders, 4th Edition, American Psychiatric Association, American Psychiatric Press, Washington DC (1994)) for the definition of dementia equate to an MMSE (Mini-Mental State Examination) cut-off point of about 18, and corresponds to a dementia prevalence of about 5% of the population over-65 years old (over-65's represent about 17% of the total population).

10

15

Gertz et al. ((1996) Eur. Arch. Psychiatry Clin. Neurosci. 246, 132-6)) studied cases followed from general practice to post-mortem, which were rigorously characterised at the clinical level using CAMDEX (Roth et al, 1988, "The Cambridge Examination for Mental Disorders of the Elderly (CAMDEX)" Cambridge University Press). These were staged post-mortem by the criteria of Braak and, after excluding all cases with any degree of vascular pathology found post-mortem, there remained uncertainty in about one third of cases at the point of transition. That is, about one third of cases with a clinical diagnosis of AD are actually at early Braak stages (stages 1-3), have vascular pathology, or have concomitant Lewy body pathology. Thus there exists a high degree of uncertainty, even in the best practice research setting. The predominant neuropathological substrate that is actually present when a routine clinical diagnosis of AD is made is even more uncertain.

20

25

30

It has recently been reported that a molecule (FDDNP, 2-(1-{6-[(2-[¹⁸F]fluoroethyl)(methyl)amino]-2-naphthyl}ethylidene)malononitrile) demonstrates increased relative retention time (RRT) in medial temporal lobe brain regions (hippocampus, entorhinal cortex and amygdala) following injection and PET imaging in cases with a

35

clinical and neuroradiological diagnosis of Alzheimer's disease (Shoghi-Jadid et al., Am. J. Geriatr. Psychiatr. 2002, 10:24-35).

Although binding to NFTs and amyloid plaques is discussed, no
5 binding to NFTs is shown, although the compound does bind with high affinity to synthetic beta-amyloid fibrils in vitro.

When cases were matched for corresponding disease severity by MMSE score with a neuropathological case series in which vascular
10 pathology was excluded, the RRT values reported by Shoghi-Jadid et al. were found to correlate with beta-amyloid plaque counts but not with measures of neurofibrillary tangle pathology, as shown below.

Spearman rank correlation coefficients:

15

	MTL AP	Global AP	MTL NFT	Global NFT
RRT	0.665**	0.654**	0.244	0.189
p	<0.01	<0.01	>0.1	>0.1

20 Pearson correlation coefficients:

	MTL AP	Global AP	MTL NFT	Global NFT
RRT	0.602*	0.596*	0.266	0.275
p	<0.05	<0.05	>0.3	>0.3

25

Where the parameters are defined as follows:

MTL AP	medial temporal lobe amyloid plaques
Global AP	average amyloid plaque load in 12 brain regions
30 MTLNFT	medial temporal lobe neurofibrillary tangles
Global NFT	average NFT load in 12 brain regions

However, beta-amyloid deposition is known to discriminate poorly between normal aging and Alzheimer's disease (see Figure 2d
35 herein), and beta-amyloid pathology does not provide a sound basis for neuropathological staging (Braak and Braak, 1991). Therefore, FDDNP-RRT does not provide a method for in vivo neuropathological staging of Alzheimer's disease.

Although it is possible that further refinement in clinical methods with particular reference to more specific neuropsychological indicators (e.g. split attention tasks, delayed matching to sample, etc.) may improve the accuracy of clinical diagnosis, an essential problem is to develop methods for the direct measurement of underlying neuropathology during life, in particular the extent of neurofibrillary degeneration of the Alzheimer-type.

10 *Progression of neurofibrillary degeneration and tau*

As described above, the tau-based pathology of AD is a major feature of the phenotype. It is highly correlated with extent of neuronal destruction (reviewed in Wischik et al. (2000) *loc cit*).

15

On a cellular basis, the formation of NFTs from Tau is believed to proceed as follows. In the course of their formation and accumulation, paired helical filaments (PHFs) first assemble as filaments within the cytoplasm, probably from early tau oligomers which become truncated prior to, and in the course of, PHF assembly (Refs 26,27). They then go on to form classical intracellular NFTs. In this state, PHFs consist of a core of truncated tau and a fuzzy outer coat containing full-length tau (Wischik et al. (2000) *loc. cit.*). The assembly process is exponential, consuming the cellular pool of normal functional tau and inducing new tau synthesis to make up the deficit (Ref 29). Eventually functional impairment of the neurone progresses to the point of cell death, leaving behind an extracellular NFT. Cell death is highly correlated with the number of extracellular NFTs (Bondareff, W. et al. (1993) *Arc. Gen. Psychiatry* 50: 350-6). As the outer neuronal membrane is damaged and NFTs are extruded into the extracellular space, there is progressive loss of the fuzzy outer coat of the neurone with corresponding loss of N-terminal tau immunoreactivity, but preservation of tau immunoreactivity associated with the PHF core (Figure 3; Ref 30).

35

In the process of aggregation, tau protein undergoes a conformational change in the repeat domain associated with a half-

repeat phase-shift (Refs 32,33). This creates a proteolytically-stable fragment which is identical to that found in the core of the paired helical filaments (PHFs) which constitute the neurofibrillary tangles characteristic of AD. By analogy with other protein aggregation systems, the process most likely involves an alpha-helix to beta-strand change in conformation (reviewed in Wischik et al. (2000) loc. cit.).

Generally speaking therefore, the aggregation of tau can be considered in 3 stages: intracellular oligomers; intracellular filaments (stage 1 of Fig 3); extracellular filaments (stages 2 & 3 of Fig 3).

However, to date, no definitive correlation has been established between these stages, which occur at the cellular level, and possibly at different rates and probabilities in different regions in the brain, and the progression of pathology according to the defined hierarchical system of Braak and Braak which, as discussed above, is the best available definition of progression of relatively pure neurofibrillary degeneration.

Invasive methods for assessing AD

Lumbar-puncture CSF measurements enable discrimination between AD and controls, and between AD and other neurological disorders, but lumbar-puncture is more invasive than nuclear medicine-based approaches, and carries a higher risk (Refs. 17 to 21). EEG-neurological diagnosis has also been developed (Refs 22-25), but in this regard there remains a need for cheap instrumentation which can be used at the point of clinician contact.

Neurofibrillary degeneration via brain atrophy - SPECT and PET

Numerous studies have shown that global brain atrophy and specific medial temporal lobe atrophy, particularly of the hippocampus, are closely linked to underlying neurofibrillary degeneration of the Alzheimer-type, and are valuable in the early diagnosis of AD (Refs 1-8).

However, although the diagnosis of AD by monitoring global brain atrophy represents a methodology which can be made to work in a research setting, there are difficulties in defining and measuring atrophy in specific brain regions, and likewise in the measurement of global neocortical atrophy. In any case, a diagnosis based on detectable atrophy may come too late for effective treatment.

There have been advances in diagnostic methodology in recent years, following the identification of diagnostic features in SPECT scans (Refs 9-12; characteristic patterns of perfusion defect detected by HMPAO SPECT), PET scans (Refs 13-15; metabolic defect detected by glucose metabolism profile) and MRI scans (Ref 16; global brain atrophy, specific patterns of lobar atrophy). Of these, the most generally accessible are MRI and SPECT, since PET is for the present time limited to centres which have local specialised cyclotron and radiochemistry capability for the preparation of short half-life injectable radioligands (Aberdeen, London, Cambridge in UK). Notably, the characteristic early stage temporo-parietal perfusion defect detected by HMPAO SPECT in patients with AD corresponds very closely to the pattern of tau pathology which can be detected biochemically (Figure 1). The biochemical changes precede overt neurofibrillary degeneration as seen by the appearance of NFTs (Figure 2; Mukaetova-Ladinska *et al.*, 2000 *Am. J. Pathol.* Vol. 157, No. 2, 623-636).

However, although MRI and SPECT scans are useful for detecting specific patterns of perfusion defects characteristic of AD, discrimination between different neuropathological stages, or between AD and other types of dementia, is difficult.

For instance, SPECT is useful for detection of a specific pattern of bilateral temporo-parietal perfusion defect that is characteristic of AD (Refs 9 - 11), which can be useful even at very early stage disease. However, SPECT changes discriminate neuropathological stages poorly (Ref 12). Furthermore, discrimination between AD and Lewy Body dementia is difficult. Both have a bilateral temporo-parietal perfusion defect, but only in the

latter does an occipital perfusion defect tend to be present. The same patterns of defect can be demonstrated using PET measurement of glucose metabolism (Refs 13 - 15), but the problem of distinguishing Lewy Body dementia persists in this modality.

5

Thus, as can be inferred from data in Ref 12, the probability of successful SPECT detection of cases at Braak stages 1&2 is 50%, and at stages 3&4 is 60%. It is only at stages 5&6 that 95% of cases become SPECT-positive. Conversely, cases detected as SPECT-positive could be at stages 1&2 (20%), 3&4 (20%), or 5&6 (60%). SPECT will therefore fail to detect 40 - 50% of the pre-stage 4 target population for early diagnosis and therapeutic intervention. In a further study (data not shown) it was shown that overall agreement between SPECT diagnosis and clinical diagnosis was of the order of 50%.

15

Thus, in developing a treatment aimed specifically at preventing neurofibrillary degeneration of the Alzheimer-type, there is a critical need to develop, in parallel, non-invasive means of selecting patients for treatment, and monitoring their response to the treatment, according to a defined and reproducible definition of disease progression.

20

Disclosure of the invention

25

Brief summary of the invention

30

35

The present inventors have used immunochemical properties (Refs 26, 27, 30) to distinguish intracellular tangles from extracellular tangles. Both the frequency of cases with tangles in these categories (i.e. probability) and their quantity (i.e. counts per mm²) were determined in a prospective case series and grouped into the regions of the brain known to represent stages in the progression of pathology according to the system of Braak and Braak.

As described in more detail below, these antibody studies demonstrate for the first time that by employing extracellular vs.

intracellular specificity, in defined brain regions, deposits of PHF-tau provide a basis for empirical staging of the neurofibrillary degeneration of AD.

- 5 Thus in one aspect, the present invention provides a method for determining the stage of neurofibrillary degeneration associated with tauopathy (e.g. AD) in a subject believed to suffer from the disease, which method comprises the steps of:
- (i) introducing into the subject a ligand capable of labelling
 - 10 aggregated PHF tau,
 - (ii) determining the presence and/or amount of ligand bound to extracellular aggregated PHF tau in the medial temporal lobe of the brain of the subject,
 - (iii) correlating the result of the determination made in (ii) with
 - 15 the extent of neurofibrillary degeneration in the subject.

As described in the introduction, progression of neurofibrillary degeneration is definitive of the neuropathological staging proposed by Braak, which in turn is the best available

20 neuropathological definition of progression of AD. The methods of the present invention can thus be used to provide an actual Braak stage result. In preferred embodiments they can be used to diagnose patients at early Braak stages (e.g. Braak stage 2) even before clinical symptoms may be readily apparent - such diagnosis

25 can be used to direct timely treatment and advice.

Interestingly, results shown in Gertz *et al.* (1996) *loc cit*, based on immunological detection of NFTs, but which did not distinguish extracellular and intracellular tangles, showed little difference

30 between the numbers detected in demented (generally Braak stage 4-6) and non-demented (generally Braak stage 1-3) subjects in medial temporal lobe structures (see Figure 1 and Table 2, page 134 therein; the relevant structures are labelled Pre alpha ent., CA1, Pri Ento.). Thus the correlation demonstrated by the present

35 invention is particularly surprising.

The invention further provides novel ligands for use in labelling tau aggregates, plus also novel screens for discovering such

ligands.

Some of the aspects of the invention discussed above will now be dealt with in more detail.

5

Choice of subject

Suitable subjects for the method may be selected on the basis of conventional factors. Thus the initial selection of a patient may
10 involve any one or more of: rigorous evaluation by experienced clinician; exclusion of non-AD diagnosis as far as possible by supplementary laboratory and other investigations; objective evaluation of level of cognitive function using neuropathologically validated battery.

15

Ligands

The ligand is capable of labelling aggregated PHF tau, the formation of which is discussed above. It may specifically or
20 preferentially bind such tau (preferentially with respect to competing binding sites present in the relevant region of the brain). Suitable ligands (including novel ligands) and methods of identifying them are discussed below.

25 More specifically, the disclosure that Braak staging can be assessed on the basis described herein has significant implications for the choice and/or development of ligands for use in diagnostic labelling. Immunological methods suffer from the drawback that antibodies do not readily cross the blood-brain barrier in a
30 quantitative manner, and furthermore, the method may be clinically unsuitable since adverse reactions may be triggered by the injection of antibodies into the body for this purpose. It is consequently difficult to discriminate between different stages of tau aggregation on the basis of differential patterns of
35 immunoreactivity in living subjects.

The present inventors have therefore investigated the critical chemical characteristics of compounds which bind to neurofibrillary

tangles. They provide herein a minimal chemical structure required for binding which has implications, *inter alia*, in the development and use of compounds as ligands of neurofibrillary tangles and such processes, uses and compounds form further aspects of the invention.

Preferred ligands, including novel ligands, are disclosed in more detail hereinafter, but may include in particular sulphonated-benzothiazole-like compounds (see e.g. Figure 4a) and diaminophenothiazines (see e.g. Figure 8) as well as other mimetic compounds sharing an appropriate minimal chemical structure with either of these. Compositions comprising, or consisting of, combinations of the ligands disclosed herein (preferably distinguishable ligands e.g. in terms of labelling) and/or combinations of ligands with blocking agents (see below) form various aspects of the invention.

Binding to extracellular tau

The determination of (ii) above is made based on extracellular aggregated tau. In general terms, for the purposes of the present invention, this may be determined from extracellular tangles (see e.g. Refs 26, 27 and Examples, Methods and Materials, Table).

It has previously been shown from histological studies that, during the course of aggregation, tau protein acquires binding sites for compounds such as thiazin red and thioflavin-S (Refs 26, 27). The binding site can be shown to exist within the tangle itself, and not in extraneous proteins (Ref 34). Thus both intracellular and extracellular tangles are labelled to some extent by such ligands, as judged histologically.

In general terms, the probability or amount of extracellular binding sites (as opposed to total binding sites, or intracellular sites) may be determined either by using ligands which are too large to gain ready intracellular access, or ligands which can act intracellularly, but at a defined (relatively lower) concentration at which extracellular action is favoured.

Large chelated ligands, such as those susceptible to detection by SPECT, could be expected to at least reach and bind appropriate extracellular targets. Compounds labelled directly for PET could potentially detect both intracellular or extracellular targets, with the latter being favoured at low concentration. Thus the work of the present inventors shows that both of these detection methods have potential in Braak staging, when used with an appropriate tangle-binding ligand. Nevertheless, in the light of the present disclosure, it will be appreciated that in order to conveniently assess Braak stage via NFT numbers it may be important to employ ligands which are not only capable of crossing the blood brain barrier and labelling specified extracellular or intracellular deposits of aggregated tau, but preferably can also retain this property when conjugated to further compounds.

However, for the avoidance of doubt, ligands may be visualised or detected by any suitable means, and the skilled person will appreciate that any suitable detection means as is known in the art could be substituted for these examples.

Enhancement of preferential tau binding

In one embodiment of the invention, steps (i) and/or (ii) of the method are performed in conjunction with (preferably preceded by) the further step of introducing into the subject a second ligand which labels competing (i.e. non-aggregated tau) binding sites present in the relevant region of the brain preferentially to the first ligand.

Thus the methods and other embodiments herein may include a step:

(i) introducing into the subject a blocking ligand which labels non-aggregated tau binding sites in the brain of the subject preferentially to the ligand capable of labelling aggregated PHF tau.

A competing binding site may be one which is provided by e.g.

amyloid plaques, such as may be present in the subject. By introducing such second ligands into the subject, the relative or effective concentration of first ligand available to bind to aggregated tau may be enhanced. Suitable second ligands (or
5 blocking compounds as they may be described herein) are described below, but they may in particular include benzthiazoles such as are shown in Figure 5, compounds 1B and 2. Another suitable blocking ligand may be FDDNP of Shoghi-Jadid et al., Am. J. Geriatr. Psychiatr. 2002, 10:24-35, discussed above.

10

Brain regions

The significance of the medial temporal lobe i.e. E2/Trans (Entorhinal cortex layer 2/transitional entorhinal cortex) and
15 E4/Hc (Entorhinal cortex layer 4 and hippocampus) regions, and also neocortical structures (F/T/P regions - frontal, temporal, parietal) of the brain are demonstrated in Figures 25, 27, and 29.

In one embodiment, the method comprises only analysing the data
20 based on extracellular NFTs in the medial temporal lobe.

If a further embodiment, both this region and the neocortical structure data is assessed. In the latter case it may be preferable to assess intracellular PHF deposits.

25

Thus the methods and other embodiments herein may include a further step:

(iib) additionally determining the presence and/or amount of ligand
30 bound to intracellular aggregated PHF tau in a neocortical structure of the brain of the subject,

This may be followed by:

35 (iii) correlating the result of the determination made in (ii) and optionally (iib) with the extent of neurofibrillary degeneration in the subject, and hence the AD state of the subject.

The ligand used for intracellular labelling may in principle be the same as that used for extracellular labelling, but preferably will be different and/or labelled distinctively (such that it can be distinguished by whatever imaging process is being used).

5

The additional steps may be particularly preferred for assessing or confirming neurofibrillary degeneration in subjects at Braak stage 2-6.

10 *Determination of neurofibrillary degeneration*

The determination may be of the presence of binding in a given area. This determination can then be related to a normal range of values for cases without any pathology (i.e. cases putatively at Braak Stage 1), or ranges of reference values which have been determined for successive Braak stages to determine the neuropathological stage corresponding to the given determination. The correlation may be done by means of a look-up table or graph, e.g. based on data corresponding to the Figures and Table 1 in Example 1 herein for density. Alternatively, a given determination may be related with reference to a given threshold value to a probability of a case being at a Stage more advanced than Stage 1 (e.g. based on data corresponding to the Figures herein for probability), and thereby giving a probability of correctly attributing a diagnosis of Alzheimer's disease.

25

Uses of the method

The determination may be as part of a method of diagnosis or prognosis. It may be used to select a patient for treatment, or to assess the effectiveness of a treatment or a therapeutic e.g. an inhibitor of tau-tau association administered to the subject.

30

Thus embodiments of the invention include:

35

A ligand which is capable of labelling extracellular aggregated PHF tau for use in a method of diagnosis or prognosis of AD in a subject believed to suffer from the disease, which method comprises

the steps of:

- (i) introducing into the subject a ligand capable of labelling aggregated PHF tau,
- (ii) determining the presence and/or amount of ligand bound to extracellular aggregated PHF tau in the of the medial temporal lobe of the brain of the subject,
- (iii) correlating the result of the determination made in (ii) with the extent of neurofibrillary degeneration in the subject, and hence the AD state of the subject.

10

Use of a ligand which is capable of labelling extracellular aggregated PHF tau in a method for preparing a diagnostic or prognostic reagent suitable for use in a method of determining the stage of neurofibrillary degeneration associated with AD in a subject believed to suffer from the disease, which method comprises the steps of:

15

- (i) introducing into the subject said reagent which is capable of labelling aggregated PHF tau,
- (ii) determining the presence and/or amount of the reagent bound to extracellular aggregated PHF tau in the medial temporal lobe of the brain of the subject,
- (iii) correlating the result of the determination made in (ii) with the extent of neurofibrillary degeneration in the subject.

20


- 25 In a still further aspect, the invention provides a kit for performing the uses and methods described above, the kit comprising one or more ligands or derivatives as provided herein, which are capable of binding to the aggregated molecules. It may include means for increasing the detectability of such compounds e.g. a technetium chelating group, plus optionally means to conjugate this to the ligand, plus optionally technetium. Where the kit comprises a derivative of a compound as disclosed herein, this may be e.g. fluoroscopically detectable, as discussed elsewhere in this description. The kit may also include means for detecting or visualising the ligand, e.g. where the ligand has an incorporated biotin group, the kit preferably includes an antibiotin antibody. Similarly, the kit may include means for detecting the inherent fluorescence of a compound, means for detecting photoactivatable
- 30
- 35

groups, further labelled antibodies, etc.

Various preferred ligands for use in the methods and other
embodiments of the present invention will now be discussed in more
5 detail. In each case, those skilled in the art will appreciate that
instead of administering ligands directly, they could be
administered in a precursor form, for conversion to the active form
by an activating agent present in, or administered to, the same
subject.

Sulphonated-benzothiazole-like ligands

Suitable ligands for use in this aspect of the present invention
are compounds of the formula:



wherein:

W is S, O, or NH;

exactly one of X, Y, and Z is CH or N;

the others of X, Y, and Z are CH;

M¹ is an alkali metal cation;


RL is a rigid linker group;

Ar¹ is an C₅₋₂₀aryl group;

n is an integer from 0 to 3; and,


each R^{BT} is a core substituent.

In one embodiment, each of X, Y, and Z is CH, and the compounds
have the following formula:




16


In one embodiment, X is N; Y and Z are each CH; and the compounds have the following formula:




5 In one embodiment, Y is N; X and Z are each CH; and the compounds have the following formula:



10 In one embodiment, Z is N; X and Y are each CH; and the compounds have the following formula:




In one embodiment, W is S, and the compounds have the following formula:




15

In one embodiment, W is O, and the compounds have the following formula:



In one embodiment, W is NH, and the compounds have the following formula:



5

In one embodiment, each of X, Y, and Z is CH; and W is S.

In one embodiment, each of X, Y, and Z is CH; and W is O.

In one embodiment, each of X, Y, and Z is CH; and W is NH.

10 In one embodiment, X is N; Y and Z are each CH; and W is S.

In one embodiment, X is N; Y and Z are each CH; and W is O.

In one embodiment, X is N; Y and Z are each CH; and W is NH.

In one embodiment, Y is N; X and Z are each CH; and W is S.

15 In one embodiment, Y is N; X and Z are each CH; and W is O.

In one embodiment, Y is N; X and Z are each CH; and W is NH.

In one embodiment, Z is N; X and Y are each CH; and W is S.


In one embodiment, Z is N; X and Y are each CH; and W is O.

20 In one embodiment, Z is N; X and Y are each CH; and W is NH.

The bicyclic group, comprising W, X, Y, and Z, may be denoted the "core group." When each of X, Y, and Z is CH, and W is S, the compound may be referred to as a benzothiazole compound, and may be considered to have, as a core group, a benzothiazole group. The "core substituents" may be then be referred to as "benzothiazole substituents."

25

A preferred ligand for use in this aspect of the present invention is a ligand compound of the formula (I):



wherein:

- 5 M^1 is an alkali metal cation;
- RL is a rigid linker group;
- Ar^1 is an C_{5-20} aryl group;
- n is an integer from 0 to 3; and,
- each R^{BT} is a independently benzothiazole substituent.

10

Both the rigid linker group, RL , and the aryl group, Ar^1 , are substantially planar. In addition, the rigid linker group, RL , and the aryl group, Ar^1 , together with the core group (e.g., benzothiazole group), form a compound which is substantially planar. By "substantially planar," it is meant that the moiety/compound has a high degree of planarity e.g. less than 5, 4, 3, 2 or 1° twist between the components, as quantified using standard chemical models and assumptions. Preferably the twist will be no greater than that of the compound of Figure 16.


20

In one embodiment, the compound has a compound length which is from about 14.7 AU to about 15.3 AU.


25 The present inventors have determined that compounds having the characteristics described above may be particularly suitable for the 'Braak staging' methods of the invention. Such compounds may be known in the art, or may be novel as described in more detail below.

30 The "compound length" is the distance between the two most distant aromatic ring atoms (denoted "reference atoms"). For example, for benzothiazole compounds, at the benzothiazole "end" of the molecule, the reference atom will be one of two atoms:

19



At the aryl "end" of the molecule, when Ar¹ is an aryl group having a phenyl core (see below), the reference atom will be one of three atoms:



Distances used herein may be computed using 'Chemical Database Service', Daresbury, and the Cambridge Structure Database, using 'Chemical structure search and retrieval software'. This data and software are available in the public domain.

In one embodiment, M is Li, Na, K, or Cs.

In one embodiment, M is Na or K.

In one embodiment, n is 0. In one embodiment, n is 1.

In one embodiment, n is 2. In one embodiment, n is 3.

In one embodiment, each R^{BT} is independently selected from:

C₁₋₄alkyl, hydroxy, C₁₋₄alkoxy, nitro, cyano, halo, and amino.

In one embodiment, each R^{BT} is independently selected from: -Me, -Et, -nPr, -iPr, -OH, -OMe, -OEt, -O(nPr), -O(iPr), -NO₂, -CN, -F, -Cl, -Br, -I, -NH₂, -NHMe, -NH₂, -NHMe, -NH₂, -NH(iPr), -NH(nPr), -NMe₂, -NEt₂, N(iPr)₂, and -N(nPr)₂.


In one embodiment, each R^{BT} is independently selected from: C₁₋₄alkyl. In one embodiment, each R^{BT} is selected from: -Me, -Et, -nPr, and -iPr. In one embodiment, each R^{BT} is -Me.

20


In one embodiment, n is 1 and R^{BT} is -Me, -Et, -nPr, or -iPr.

In one embodiment, n is 1 and R^{BT} is -Me.

5 In one embodiment, the compound has the following formula:




In one embodiment, the compound has the following formula:




10

In one embodiment, RL is a group of the formula:



wherein m is an integer from 0 to 4, and each R^{RL} is independently a rigid linker aryl substituent, and the compounds have the formula:



15

In one embodiment, m is 0. In one embodiment, m is 1.

In one embodiment, m is 2. In one embodiment, m is 3.

In one embodiment, m is 4.

20


In one embodiment, each R^{RL} is independently selected from: C_{1-4} alkyl, hydroxy, C_{1-4} alkoxy, nitro, cyano, halo, and amino.

In one embodiment, each R^{RL} is independently selected from: -Me, -Et, -nPr, -iPr, -OH, -OMe, -OEt, -O(nPr), -O(iPr), -NO₂, -CN, -F, -Cl, -Br, -I, -NH₂, -NHMe, -NHEt, -NH(iPr), -NH(nPr), -NMe₂, -NEt₂, N(iPr)₂, and -N(nPr)₂.

5


In one embodiment, each R^{RL} is independently selected from: C_1 - $_4$ alkyl.

In one embodiment, RL is a group of the formula:




10

In one embodiment, RL is a group of the formula:



wherein p is an integer from 0 to 3, and each R^{RL} is independently a
15 rigid linker aryl substituent, and the compounds have the formula:



In one embodiment, p is 0. In one embodiment, p is 1.

In one embodiment, p is 2. In one embodiment, p is 3.

20

In one embodiment, each R^{RL} is independently selected from: C₁₋₄alkyl, hydroxy, C₁₋₄alkoxy, nitro, cyano, halo, and amino.


In one embodiment, each R^{RL} is independently selected from: -Me, -Et, -nPr, -iPr, -OH, -OMe, -OEt, -O(nPr), -O(iPr), -NO₂, -CN, -F, -Cl, -Br, -I, -NH₂, -NHMe, -NHEt, -NH(iPr), -NH(nPr), -NMe₂, -

NEt₂, N(iPr)₂, and -N(nPr)₂.

In one embodiment, each R^{RL} is independently selected from: C₁₋₄alkyl.

5

In one embodiment, RL is a group of the formula:



The aryl group, Ar¹, is a C₅₋₂₀aryl group. The term "C₅₋₂₀aryl," as
 10 used herein, pertains to a monovalent moiety obtained by removing a
 hydrogen atom from an aromatic ring atom of a C₅₋₂₀aromatic
 compound, said compound having one ring, or two or more rings
 (e.g., fused), and having from 5 to 20 ring atoms, and wherein at
 least one of said ring(s) is an aromatic ring. Preferably, each
 15 ring has from 5 to 7 ring atoms. "C₅₋₂₀" denotes ring atoms,
 whether carbon atoms or heteroatoms.


Examples of C₅₋₂₀aryl groups which do not have ring heteroatoms
 (i.e., C₅₋₂₀carboaryl groups) include, but are not limited to, those
 20 derived from benzene (i.e., phenyl) (C₆), naphthalene (C₁₀),
 anthracene (C₁₄), phenanthrene (C₁₄), naphthacene (C₁₈), and pyrene
 (C₁₆).

Examples of C₅₋₂₀heteroaryl groups include, but are not limited to,
 25 C₅heteroaryl groups derived from furan (oxole), thiophene (thiole),
 pyrrole (azole), imidazole (1,3-diazole), pyrazole (1,2-diazole),
 triazole, oxazole, isoxazole, thiazole, isothiazole, oxadiazole,
 and oxatriazole; and C₆heteroaryl groups derived from isoxazine,
 pyridine (azine), pyridazine (1,2-diazine), pyrimidine (1,3-
 30 diazine; e.g., cytosine, thymine, uracil), pyrazine (1,4-diazine),
 triazine, tetrazole, and oxadiazole (furazan).


Examples of C₅₋₂₀heterocyclic groups (including C₅₋₂₀heteroaryl
 groups) which comprise fused rings, include, but are not limited
 35 to, C₉heterocyclic groups derived from benzofuran, isobenzofuran,

indole, isoindole, purine (e.g., adenine, guanine), benzimidazole;
 C_{10} heterocyclic groups derived from quinoline, isoquinoline,
 benzodiazine, pyridopyridine, quinoxaline; C_{13} heterocyclic groups
 derived from carbazole; C_{14} heterocyclic groups derived from
 5 acridine, xanthene, phenoxathiin, phenazine, phenoxazine,
 phenothiazine.

In one embodiment, Ar^1 is an aryl group having a phenyl core, and
 has the formula:



10 wherein q is an integer from 0 to 5; and each R^A is independently
 an aryl substituent; R^C , if present, is a reactive conjugating
 substituent, or R^C is, or contains, a detectable label; and the
 compound has the formula:



15 In one embodiment, R^C , if present, is a reactive conjugating
 substituent, and is a group which is suitable for conjugation to
 another molecule or chemical species.

20 In one embodiment, R^C , if present, is a reactive conjugating
 substituent, and is, or contains, a reactive functional group
 suitable for conjugation to another molecule by chemical reaction
 therewith, to form a covalent linkage therebetween. Examples of
 suitable reactive functional groups include active esters (e.g.,
 25 succinimidyl esters).

In one embodiment, R^C , if present, is a reactive conjugating
 substituent, and is, or contains, a moiety suitable for conjugation
 to another molecule by a strong non-covalent interaction. Examples

of such groups include biotin (for binding with molecules bearing avidin or streptavidin).

In one embodiment, R^C , if present, is a reactive conjugating substituent, and is, or contains, a moiety suitable for conjugation to another molecule by complex or chelate formation, e.g., a chelating group. Examples of such groups include groups which complex with or chelate, e.g., metal ions, e.g., technetium ions. Examples of such groups include diethylenetriaminepentaacetic acid.

In one embodiment, R^C , if present, is, or contains, a detectable label. Examples of detectable labels include, e.g., dyes, fluorescent markers, antigenic groups, stable and unstable isotopes, and positron-emitting carbon atoms. In one embodiment, R^C , if present, is, or contains, a detectable label comprising a stable isotope. In one embodiment, R^C , if present, is, or contains, a detectable label comprising an unstable isotope. In one embodiment, R^C , if present, is, or contains, ^{18}F . In one embodiment, R^C , if present, is, or contains, a detectable label comprising a positron-emitting carbon atom.

Further R^C substituents are discussed below.

In one embodiment, R^C is present, and is as defined above.


In one embodiment, q is 0. In one embodiment, q is 1. In one embodiment, q is 2. In one embodiment, q is 3. In one embodiment, q is 4. In one embodiment, q is 5.

In one embodiment, each R^A is independently selected from: $-\text{OH}$, $-\text{NH}_2$, $-\text{NHR}^1$, $-\text{NR}^1\text{R}^2$, $-\text{SO}_3\text{M}^2$, $\text{C}_{1-4}\text{alkyl}$, wherein R^1 and R^2 are each $\text{C}_{1-4}\text{alkyl}$, and M^2 is an alkali metal cation, as defined above.

In one embodiment, at least one R^A is $-\text{OH}$ or $-\text{NH}_2$.

25

In one embodiment, Ar^1 is an aryl group having an amino-substituted phenyl core, and has the formula:



wherein r is an integer from 0 to 4, and each R^{A} is independently
 5 an aryl substituent, as defined above.


In one embodiment, r is 0. In one embodiment, r is 1.

In one embodiment, r is 2. In one embodiment, r is 3.


In one embodiment, r is 4.

10

In one embodiment, r is 1 and Ar^1 is a group of the formula:




In one embodiment, the compound has the formula:




15

In one embodiment, the compound has the formula:



In one embodiment, Ar^1 is an aryl group having a hydroxy-substituted phenyl core, and has the formula:




wherein s is an integer from 0 to 4, and each R^A is independently an aryl substituent, as defined above, and R^C , is present, is a reactive conjugating substituent, or R^C is, or contains, a detectable label, as defined above.

In one embodiment, s is 0. In one embodiment, s is 1.


10 In one embodiment, s is 2. In one embodiment, s is 3.

In one embodiment, s is 4.


In one embodiment, Ar^1 is a group of the formula:



15 In one embodiment, Ar^1 is a group of the formula:




In one embodiment, Ar^1 is a group of the formula:




27


In one embodiment, Ar^1 is an aryl group having a naphthyl core, and has the formula:



wherein t is an integer from 0 to 3, u is an integer from 0 to 4,
 5 and each R^{A} is independently an aryl substituent, as defined above,
 and the compound has the formula:




In one embodiment, Ar^1 is an aryl group having a hydroxy-
 10 substituted naphthyl core, and has the formula:




wherein v is an integer from 0 to 2, u is an integer from 0 to 4,
 and each R^{A} is independently an aryl substituent.


15 In one embodiment, Ar^1 has the formula:



In one embodiment, the compound has the formula:



In one embodiment, the compound has the formula:



5

In one embodiment, the ligand is a compound as described below under the heading "*Preferred sulphonated-benzothiazole-like ligands.*"

- 10 Compounds of the type described above, e.g. of formula (I), for use in the diagnostic methods of the present invention, may be prepared by conventional means - see e.g. Reference 31.

15 All such compounds described herein (or derivatives thereof) having the appropriate formula, size, planarity, and activity may be referred to generally, but not limitatively, hereinafter as 'sulphonated benzothiazole-like compounds' or 'SB ligands'). Such compounds will generally be ligands of aggregated tau molecules, e.g. those found in paired helical filaments or neurofibrillary
20 tangles.

The ligands described herein can suitably be detected by incorporating a positron-emitting carbon into one of the methyl groups of the compound as disclosed herein, and detecting the
25 compound by use of positron emission tomography (PET) as is known in the art. Alternatively, or in addition, a technetium-containing

chelate can be incorporated into the compound (e.g. as in the R^c group of the compounds described herein), so that selective detection of extracellular tangles could be achieved. A preferred chelating group is R^c = diethylenetriaminepentaacetic acid.

5

The ligands may be conjugated, chelated, or otherwise associated, with other chemical groups, dyes, fluorescent markers, antigenic groups, therapeutic moieties, or any other entity which may aid in a prognostic, diagnostic or therapeutic application. For instance, where the ligand is attached to a dye or fluorescent group, the conjugate can be used as a label of aggregated tau or tau-like molecules. It can thus be used to label intracellular or extracellular tangles characteristic of AD.

10

15 *Phenothiazines*

The present inventors have previously identified another class of compounds, members of which disrupt the structure of PHFs, and reverse the proteolytic stability of the PHF core (WO 96/30766).

20

Diaminophenothiazine compounds described in WO 96/30766 are shown by the structures of Figure 8a. Formula (IV) in Figure 8a represents different resonance form of (II) included for clarity. Compounds (II)-(IV) are all oxidised forms while (I) is a reduced form. Such compounds (which may be referred to hereinafter as 'diaminophenothiazines' or 'phenothiazines') include, e.g. tolonium chloride and methylene blue. Examples are shown in Figure 8b. All of these are shown in the oxidised form, with all except thionine being in the form stabilised salts (thionine is shown as a neutral oxidised form).

25

30

Compounds which may be used in the methods described herein may be any having a formula shown in Figure 8a, wherein:

each of R₁, R₃, R₄, R₆, R₇ and R₉ is independently hydrogen, halogen, hydroxy, carboxy, substituted or unsubstituted alkyl, haloalkyl or alkoxy;

35

R₅ is hydrogen, hydroxy, carboxy, substituted or unsubstituted alkyl, haloalkyl or alkoxy; and,

R₁₀ and R₁₁ are independently hydrogen, hydroxy, carboxy, substituted or unsubstituted alkyl, haloalkyl or alkoxy; and pharmaceutically acceptable salts thereof.

5 In one embodiment:

each of R₁, R₃, R₄, R₆, R₇ and R₉ is independently hydrogen, halogen, hydroxy, carboxy, substituted or unsubstituted C₁₋₆alkyl, C₁₋₄haloalkyl, or C₁₋₆alkoxy;

10 R₅ is independently hydrogen, hydroxy, carboxy, substituted or unsubstituted C₁₋₆alkyl, C₁₋₄haloalkyl, or C₁₋₆alkoxy;

R₁₀ and R₁₁ are independently selected from hydrogen, hydroxy, carboxy, substituted or unsubstituted C₁₋₆alkyl, C₁₋₄haloalkyl, or C₁₋₆alkoxy.

15 The term "alkyl" as used in this respect refers to straight or branched chain groups, preferably having one to eight, more preferably one to six, carbon atoms. For example, "alkyl" may refer to methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, tert-pentyl, hexyl, isohexyl,
20 and the like. Suitable substituents for the substituted alkyl groups used in the invention include the mercapto, thioether, nitro, amino, aryloxy, halogen, hydroxyl, and carbonyl groups as well as aryl, cycloalkyl and non-aryl heterocyclic groups.

25 The terms "alkoxy" refers to groups as defined herein above as alkyl groups, as the case may be, which also carry an oxygen atom interposed between them and the substrate residue to which they are attached.

30 The term "haloalkyl" represents a straight or branched alkyl chain having from one to four carbon atoms with 1, 2 or 3 halogen atoms attached to it. Typical haloalkyl groups include chloromethyl, 2-bromomethyl, 1-chloroisopropyl, 3-fluoropropyl, 2,3-dibromobutyl, 3-chloroisobutyl, iodo-t-butyl, trifluoromethyl and the like.

35

The "halogen" represents fluoro, chloro, bromo or iodo.

Some of these phenothiazines possess one or more asymmetrically

substituted carbon atoms and therefore exist in racemic and optically active forms. The invention is intended to encompass the racemic forms of the compounds as well as any of the optically active forms thereof.

5

Acid addition salts may be formed between basic compounds of Figure 8a or 8b and inorganic acids, e.g. hydrohalic acids such as hydrochloric acid and hydrobromic acid, sulphuric acid, nitric acid, phosphoric acid etc., or organic acid, e.g. acetic acid, citric acid, maleic acid, fumaric acid, tartaric acid, methanesulphonic acid, p-toluenesulphonic acid etc.

10

In a particular preferred embodiment the present invention employs a phenothiazine wherein:

15

R_1 , R_3 , R_4 , R_6 , R_7 and R_9 are independently -H, -CH₃, -C₂H₅, or -C₃H₇; R_{10} and R_{11} are independently -H, -CH₃, -C₂H₅ or -C₃H₇; R_5 is -H, -CH₃, -C₂H₅, or -C₃H₇; or a pharmaceutically acceptable salt thereof.

20

The present inventors now teach herein that such phenothiazine compounds of this sort can bind to the PHFs at a specific site which appears, on the basis of its binding characteristics, to be distinct from that at which the sulphonated benzothiazole-like compounds, described above, can bind. The binding of the phenothiazine compounds to this site is thought to effect the inhibition of tau aggregation.

25

Phenothiazine compounds may be used in the methods and other embodiments described above, incorporating labels as appropriate.

30


When suitably labeled with a positron-emitting functional group (detectable by PET - see Figures 11, 11b, 12, 13) such compounds would serve as ligands for all tau aggregates, and would be capable of crossing the blood-brain-barrier (Ref 36) and entering cells.

35

In a further embodiment, in the light of the disclosure herein, it will be appreciated that the effect of, and particularly the progress of, therapy based on tau-tau binding inhibitors may be monitorable by use of SB ligands.

Blocking ligands

Preferably these are benzthiazoles of the formula:



wherein:

n is an integer from 0 to 4;

each R^{BT} is independently a blocking ligand benzothiazole substituent;

m is an integer from 0 to 4;

each R^P is independently a phenylene substituent;

each R is independently $-H$ or an amino substituent; and,

either:

R^N and X^- are both absent and the associated (tertiary) nitrogen atom is neutral;

or:

R^N is a benzothiazolino substituent and the associated (quaternary) nitrogen atom bears a positive charge, and X^- is a counter ion.

Preferred benzthiazoles include thioflavin T. As shown in the Examples below, such compounds (e.g. 1b or 2 in Figure 5) are displaced from NFTs by SB-ligands (e.g. 1a in Figure 5). However such compounds do bind preferentially to amyloid.

In one embodiment, n is 0. In one embodiment, n is 1.

In one embodiment, n is 2. In one embodiment, n is 3.

In one embodiment, n is 4. In one embodiment, n is 0, 1, or 2.

Examples of blocking ligand benzothiazole substituents, R^{BT} , include, but are not limited to, C_{1-4} alkyl groups, $-SO_3H$, and $-SO_3M^3$, wherein M^3 is a cation. In one embodiment, M^3 is an alkali metal cation. In one embodiment, M^3 is Li, Na, K, or Cs. In one embodiment, M^3 is Na or K. Examples of C_{1-4} alkyl groups include, but are not limited to, $-Me$, $-Et$, $-nPr$, and $-iPr$.

In one embodiment, each R^{BT} is independently a C_{1-4} alkyl group.

In one embodiment, each R^{BT} is selected from: -Me, -Et, -nPr, and -iPr. In one embodiment, each R^{BT} is -Me. In one embodiment, n is 1 and R^{BT} is -Me, -Et, -nPr, or -iPr. In one embodiment, n is 1 and R^{BT} is -Me.

In one embodiment, one of the R^{BT} groups is $-SO_3H$ or $-SO_3M^3$. In one embodiment, one of the R^{BT} groups is $-SO_3H$ or $-SO_3M^3$, and another of the R^{BT} groups is a C_{1-4} alkyl group. In one embodiment, n is 2 and one R^{BT} is a C_{1-4} alkyl group and one R^{BT} is $-SO_3H$ or $-SO_3M^3$. In one embodiment, n is 2 and one R^{BT} is -Me and one R^{BT} is $-SO_3H$ or $-SO_3M^3$.

In one embodiment, R^N and X^- are both absent and the associated (tertiary) nitrogen atom is neutral.

In one embodiment, R^N is a benzothiazolino substituent and the associated (quaternary) nitrogen atom bears a positive charge, and X^- is a counter ion. Examples of benzothiazolino substituents, R^N , include, but are not limited to, C_{1-4} alkyl groups. In one embodiment, R^N is -Me, -Et, -nPr, or -iPr. In one embodiment, R^N is -Me. Examples of counter ions include, but are not limited to, Cl^- , Br^- , and I^- . In one embodiment, R^N is -Me and X^- is Cl^- .

In one embodiment, m is 0. In one embodiment, m is 1.

In one embodiment, m is 2. In one embodiment, m is 3.

In one embodiment, m is 4.

Examples of phenylene substituents, R^P , include, but are not limited to, C_{1-4} alkyl groups.


In one embodiment, each R is -H, and the amino group is $-NH_2$. In one embodiment, one R is -H and one R is an amino substituent. In one embodiment, each R is an amino substituent. Examples of amino substituents include, but are not limited to, C_{1-4} alkyl groups. In one embodiment, the amino group is $-NH_2$, $-NHMe$, $-NHet$, $-NH(iPr)$, $-NH(nPr)$, $-NMe_2$, $-NEt_2$, $N(iPr)_2$, or $-N(nPr)_2$.

Preferred embodiments of blocking ligands are shown in Figure 5 as

compounds 1b and 2.

Preferred sulphonated-benzothiazole-like ligands

- 5 In one aspect of the present invention, the ligands used to label the aggregated tau, preferable extracellular aggregated tau present in NFTs, are compounds having the formula (II):



wherein:


- 10 M^1 is an alkali metal cation;
 n is an integer from 0 to 3;
 each R^{BT} is a independently benzothiazole substituent;
 m is an integer from 0 to 4;
 each R^{RL} is independently a rigid linker aryl substituent;
 15 s is an integer from 0 to 4;
 each R^{A} is independently an aryl substituent; and,
 R^{C} , if present, is a reactive conjugating substituent, or
 R^{C} is, or contains, a detectable label.
- 20 In various embodiments, M^1 , n , each R^{BT} , each R^{RL} , s , each R^{A} , and R^{C} are as described herein (e.g., under the heading "Sulphonated-benzothiazole-like ligands," above).

25 The rigid linker group, RL , and the aryl group, Ar^1 , together with the benzothiazole group, form a compound which is substantially planar, that is, has a high degree of planarity.


As shown herein, such compounds may be particularly effective when it is desired to incorporate a bulky R^{C} group in order to facilitate
 30 detection.

35

In one embodiment, the compound has the formula:




In one embodiment, the compound has the formula:




5

In one embodiment, the compound has the formula:




In one embodiment, the compound has the formula:




10 In various embodiments, s may be as discussed above.

In one embodiment, each RA is independently selected from the substituents given above in relation to formula (I).


In one embodiment, the compound has the formula:




In one embodiment, the compound has the formula:




5 In one embodiment, the compound has the formula:



In one embodiment, the compound has the formula:




10 In one embodiment, the compound has the formula:




In one embodiment, the compound has the formula:

37




In one embodiment, the compound has the formula:




Various R^C substituents are discussed elsewhere herein.

5


In one embodiment, the compound has the formula:




In one embodiment, the compound has the formula:




10 In one embodiment, the compound has the formula:



In one embodiment, the compound has the formula:



In one embodiment, the compound has the formula:



5

Some of these preferred compounds are shown in Figures 4a-c, and derivatives thereof.

Thus, according to one aspect, the present invention provides a compound represented by the formula shown in Figure 4a, or a derivative thereof e.g. wherein R^c is a conjugating group as described above. As shown in the Examples hereinafter, such derivatives (e.g. compound 4b) retain the appropriate binding activity.

15

The novel compounds disclosed (e.g. of formula (II)) herein are useful *inter alia* as synthetic ligands of neurofibrillary tangles, such as those characteristic of AD. The discovery of the minimum critical structure required for binding to these tangles thus provides for the possibility of designing high-affinity ligands which can be used to target the tangles, and can thus be used in the diagnosis, prognosis or treatment of diseases such as AD.

25

Such compounds will be referred to below as preferred SB ligands.

Mimetics of preferred SB ligands

In general, there are several steps commonly taken in the design of a mimetic from a compound having a given target property (in this case a preferred SB tau-tau aggregation ligand) of which the most important is that the particular parts of the compound that are critical and/or important in determining the target property are determined. The provision by the present inventors of the minimum critical structure required for high affinity binding to aggregated tau molecules has obviated this step.

10 The minimum critical structure of compound 4a can be modeled according to its physical properties, e.g. stereochemistry, bonding, size and/or charge, using data from a range of sources, e.g. spectroscopic techniques, X-ray diffraction data and NMR. Computational analysis, similarity mapping (which models the charge and/or volume of the ligand, rather than the bonding between atoms) and other techniques can be used in this modeling process.

In a variant of this approach, the three-dimensional structure of the preferred SB ligand and its binding partner are modeled. This can be especially useful where the ligand and/or binding partner change conformation on binding, allowing the model to take account of this in the design of the mimetic. A template molecule is then selected onto which chemical groups which mimic the minimum critical structure can be grafted. The template molecule and the chemical groups grafted on to it can conveniently be selected so that the mimetic is easy to synthesise, is likely to be pharmacologically acceptable, and does not degrade *in vivo*, while retaining the required biological activity. The mimetic or mimetics found by this approach can then be screened to see whether they have the target property, or to what extent they exhibit it. Further optimisation or modification can then be carried out to arrive at one or more final mimetics for further testing or optimisation, e.g. *in vivo* or clinical testing. Optimisation may include selecting a mimetic compound as described above, and contacting it with a preparation of aggregated tau molecules (e.g. preaggregated tau in solution, or bound to a solid phase, or isolated from PHFs - see W096/30766 and assays described below) and determining the extent to which the test substance(s) binds to the

aggregated tau molecules and/or displaces compound 4a from the molecules.

Methods of labelling aggregated tau

5

In one aspect, the present invention thus provides a method of labelling aggregated tau or tau-like molecules, comprising contacting the aggregated tau molecules with a preferred SB-ligand compound or derivative thereof as provided herein (e.g. of formula (II)) and detecting the presence of said compound or derivative. Methods of use may be performed e.g. by analogy to the use of the ligands given in Refs 26-34.

Where used herein, the term "tau protein" refers generally to any protein of the tau protein family. Tau proteins are characterised as being one among a larger number of protein families which co-purify with microtubules during repeated cycles of assembly and disassembly (Shelanski et al. (1973) Proc. Natl. Acad. Sci. USA, 70., 765-768), and are known as microtubule-associated-proteins (MAPs). Members of the tau family share the common features of having a characteristic N-terminal segment, sequences of approximately 50 amino acids inserted in the N-terminal segment, which are developmentally regulated in the brain, a characteristic tandem repeat region consisting of 3 or 4 tandem repeats of 31-32 amino acids, and a C-terminal tail.

"Tau like" molecules include, for instance, MAP2, which is the predominant microtubule-associated protein in the somatodendritic compartment (Matus, A., in "Microtubules" [Hyams and Lloyd, eds.] pp 155-166, John Wiley and Sons, NY). MAP2 isoforms are almost identical to tau protein in the tandem repeat region, but differ substantially both in the sequence and extent of the N-terminal domain (Kindler and Garner (1994) Mol. Brain Res. 26, 218-224). Nevertheless, aggregation in the tandem-repeat region is not selective for the tau repeat domain. Thus it will be appreciated that any discussion herein in relation to tau protein or tau-tau aggregation should be taken as relating also to tau-MAP2 aggregation, MAP2-MAP2 aggregation and so on.

The preferred SB ligand may be conjugated, chelated, or otherwise associated with, a further group or entity which has a diagnostic, prognostic or therapeutic purpose or effect, e.g. to a fluorescent group which thus enables visualisation of neurofibrillary tangles to which the ligand binds.

Diagnostic compositions and uses

Generally, a preferred SB ligand according to the present invention (e.g. of formula (II)) may be provided in an isolated and/or purified form, i.e. substantially pure. This may include being in a composition where it represents at least about 90% active ingredient, more preferably at least about 95%, more preferably at least about 98%. Such a composition may, however, include inert carrier materials or other pharmaceutically- and physiologically-acceptable excipients. A composition according to the present invention may include in addition to a preferred SB ligand as disclosed herein, one or more other molecules of diagnostic, prognostic or therapeutic use.

A preferred SB ligand substance according to the present invention, or a composition comprising such a ligand, may be provided for use in a method of diagnosis, prognosis or treatment of the human or animal body by therapy, especially in relation to a condition such as AD as described below.

In a further aspect, the present invention provides a method of diagnosis or prognosis, the method comprising administering to the mammal a diagnostically- or prognostically- effective amount of one or more preferred SB ligands as described herein. This aspect embraces such compounds for use in a method of diagnosis or prognosis. Both *in vitro* and *in vivo* uses are encompassed by this aspect. *In vitro* methods may be performed by (i) obtaining a sample of appropriate tissue from a subject; (ii) contacting the sample with the preferred SB ligand; (iii) detecting the amount and/or localisation of the preferred SB ligand bound to the sample (iv) correlating the result of (v) with the stage or severity of the

disease in the subject.

In a further aspect, the present invention provides the use of a preferred SB ligand or derivative as provided herein, in the
5 manufacture of a composition for the diagnosis, prognosis or therapy of a disease as described above.

The disease or condition may be e.g. AD, or an AD-like condition, or any other condition in which aggregated protein molecules are
10 implicated.

Notably it is not only Alzheimer's Disease in which tau protein (and aberrant function or processing thereof) may play a role. The pathogenesis of neurodegenerative disorders such as Pick's disease
15 and Progressive Supranuclear Palsy (PSP) appears to correlate with an accumulation of pathological truncated tau aggregates in the dentate gyrus and stellate pyramidal cells of the neocortex, respectively. Other dementias include fronto-temporal dementia (FTD); parkinsonism linked to chromosome 17 (FTDP-17);
20 disinhibition-dementia-parkinsonism-amyotrophy complex (DDPAC); pallido-ponto-nigral degeneration (PPND); Guam-ALS syndrome; pallido-nigro-luysian degeneration (PNLD); cortico-basal degeneration (CBD) and others (see Wischik *et al.* 2000, loc. cit, for detailed discussion - especially Table 5.1). All of these
25 diseases, which are characterized primarily or partially by abnormal tau aggregation, are referred to herein as "tauopathies".

Diagnostic compositions may comprise, in addition to one of the above SB-ligand derivatives, a diagnostically acceptable excipient,
30 carrier, buffer, stabiliser, or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the binding activity of the substance to aggregated tau, or the efficacy of any bioactive group linked to or otherwise associated with the substance. The precise nature of the
35 carrier or other material may depend on the route of administration, e.g. oral, intravenous, cutaneous or subcutaneous, nasal, intramuscular, intraperitoneal routes.

Diagnostic compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may include a solid carrier such as gelatin or an adjuvant. Liquid diagnostic compositions generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included.

- For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, the ligand will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.
- A composition as described above may be administered alone or in combination with other treatments, either simultaneously or sequentially, depending on the condition to be treated.

Identification of ligands

A further method for identifying ligands for aggregated tau requires a screening assay which can be used in a format which permits high through-put screening of chemical libraries to identify compounds with the required activity. Until now, no such method has been readily available. Preferred methods would not require pre-labelled compounds, since the labelling process may severely limit chemical search capacity.

In a further aspect of the present invention a method is provided which can be used as a high through-put screening assay, which is not limited by the need to have pre-labelled test substances. In preferred embodiments the method employs the following steps:

1. the high capacity generation of tau proteins in a form which

have undergone partial aggregation in the course of preparation;

2. using tau proteins prepared in this manner for testing putative ligands in the tau-tau binding assay provided in WO96/30766 to identify substances which have minimal or entirely absent activity as tau-aggregation inhibitors or enhance tau-tau binding at high concentrations;

3. testing putative ligands in the presence of an exemplary potent tau-aggregation inhibitor, such as DMMB, at an inhibitory concentration;

4. putative ligands can be identified by the property that they lack capacity to block tau-tau binding through the repeat domain, but block the inhibitory activity of a potent tau-aggregation inhibitor.

Thus the invention provides an *in vitro* method for identifying a ligand capable of labeling aggregated PHF tau protein, the method comprising the steps of:

(i) providing a first agent suspected of being capable of labeling aggregated PHF tau protein,

(ii) contacting (a) a tau protein or a derivative thereof containing the tau core fragment bound to a solid phase so as to expose a high affinity tau capture site (e.g. a truncated tau protein corresponding to the core fragment and terminating at Ala390 - dGA), with (b) a liquid phase tau protein or derivative thereof capable of binding to the solid phase tau protein or derivative (e.g. dGAE which terminates at Glu-391), and (c) said selected first agent and (d) a second agent known to be tau-tau binding inhibitor,

(iii) selecting a first agent which fully or partially relieves the inhibition of binding of the liquid phase tau protein or derivative of (b) to the solid phase tau protein or derivative of (a) by the inhibitor (d).

Agents satisfying (iii) may be provided as ligands.

Preferably the method is carried out in conjunction with (before, during, after) the following steps:

(ibis) contacting (a) a tau protein or a derivative thereof containing the tau core fragment bound to a solid phase so as to expose a high affinity tau capture site, (b) a liquid phase tau protein or derivative thereof capable of binding to the solid phase tau protein or derivative, with (c) said first agent and, (ibis.1) detecting inhibition of tau-tau binding as exhibited by inhibition of binding of the liquid phase tau protein or derivative of (b) to the solid phase tau protein or derivative of (a), (ibis.2) selecting a first agent which has minimal or absent activity as tau-tau binding inhibitors and/or optionally enhance tau-tau binding.

Agents satisfying (iii) and (ibis.2) may be provided as ligands.

The inhibitor is preferably a diaminophenathiozine as described above (most preferably DMMB). The compounds selected for screening may be any compound, including SB-ligands.

In preferred forms the liquid phase tau protein or derivative is prepared in a form which has undergone partial aggregation prior to exposure to the solid phase. Apart from that, the assay may be carried out broadly as described in WO96/30766 and summarised in more detail in the Examples below. Preferably alkaline or physiological conditions (e.g. PBS) are used for the binding steps, and results are detected immunologically.

These and other aspects of the present invention will become more apparent on reading the ensuing non-limiting Examples, in which embodiments of the invention will be described by way of example only. Reference is made to the accompanying figures, in which:

Brief description of the Figures

Figure 1 shows the regional distribution of PHF-tau, measured using antibodies mAb423 (A) or mAb7.51 after formic acid treatment of the PHF fraction (B), for 18 cases of AD. Taken from Mukaetova-Ladinska et al., (1993), Am. J. Pathol. 143, 565-578.

Figure 2 (a) shows the aggregation of tau molecules and the appearance of neurofibrillary tangles during the pathological stages of AD. Taken from (Mukaetova-Ladinska, E.B. *et al.* (2000) *Am. J. Pathol.* Vol. 157, No. 2, 623-636); (b) shows the

5 neuropathological staging proposed by Braak; (c) shows that the appearance of clinical dementia by DSM-IV criteria appears to correspond statistically to the transition between stages III and IV;

10 (d) shows levels of SDS-insoluble beta-amyloid protein isolated from control cases and cases with Alzheimer's disease, as reported in Harrington *et al.*, (*Am J Pathol* 1994; 145: 1472-1484). Although the mean level is higher in AD than in controls, 70% of AD cases overlap with levels of beta-amyloid found in control subjects.

15 Figure 3 shows a schematic representation of a neurofibrillary tangle (top), and the immunoreactivity changes that are observed during disease progression (bottom). Taken from Bondareff *et al.* (1994) *J. Neuropathol. Exp. Neurol.* 53, 2, 158-164.

20 Figure 4 shows the chemical structures of: the minimum critical structure which enables high affinity binding to aggregated tau molecules (compound 4a); a biotinylated version of compound 4a (compound 4b); and an R-substituted derivative of compound 4a (compound 4c), wherein R is any suitable substituent.

25 Figure 5 shows the chemical structures of: primulin (compound 1a); thioflavin-T (compound 1b); 2-(4-amino phenyl)-6-methyl-1-sulphonate benzothiazole (compound 2); thiazin red (compound 3a); and thiazin yellow (compound 3b).

30 Figure 6 shows the fluorescence peaks of primulin in solution (left), and when bound within a pure preparation of PHFs (right).

35 Figure 7 shows the fluorescence peaks of primulin bound to PHFs, in the absence (left) and presence (right) of citraconic anhydride. As is shown, citraconic anhydride has the effect of disrupting the orderly structure of PHFs and releasing free tau and free unbound primulin. Citraconic anhydride also has the effect of reversing

the charge on lysine residues and this may also play a role in releasing the primulin.

Figures 8a and 8b illustrate exemplary tau-tau binding inhibitors,
5 as disclosed in WO 96/30766.

Figure 9 shows the fluorescence of primulin bound to PHFs in the presence of alcian blue. The figure demonstrates that, in the presence of alcian blue, a disrupter of PHF structure (Ref 33),
10 there is no disturbance of the characteristic bound primulin fluorescence peak at 460nm.

Figure 10 shows the effect of various compounds ("MR", molar ratio of compound:tau) on tau-tau binding using tau proteins prepared
15 according the method provided in WO96/30766 and referred to herein as "Preparation 1".

Figure 11a show schematically the synthesis of [^{11}C]-labelled methylene blue. The reaction proceeds via N-methylation of the
20 amines with [^{11}C]iodomethane. HPLC can be used to purify the desired product from any by-products. Figure 11b shows a further synthesis based on thionin, NaH and CH_3I .

Figure 12 shows schematically the synthesis of [^{11}C]-labelled Azure
25 B. The reaction proceeds via N-methylation of the amines with [^{11}C]iodomethane. HPLC can be used to purify the desired product from any by-products.

Figure 13 shows schematically the synthesis of a [^{18}F]-labelled
30 derivative of compound 4a of Figure 4. The reaction proceeds via nucleophilic aromatic substitution whereby a nitro-group on a precursor compound is exchanged for a [^{18}F] fluoro group. HPLC can be used to purify the desired product from any by-products.

Figure 14 shows the structures Primulin, Benthothiazole Analogue
35 and Thiazin Yellow. The sizes of these molecules have been determined on the basis of C-C distances known from chrystal structures, and designated A and B for each molecule. The C-C

lengths are as follows:

Primulin	Minimum	14.78 AU
	Maximum	15.11 AU
5	Mean	14.95 AU

Analogue	Minimum	15.05 AU
	Maximum	15.26 AU
10	Mean	15.17 AU

Thiaz. Yellow	Minimum	15.73 AU
	Maximum	16.14 AU
	Mean	15.93 AU

15 Figures 15 and 16 illustrate the crystal structure of the 'B' part of the primulin structure (Soon-Beng Teo *et al.*, 1995, *Acta Crystallogr.*, Sect. C, 591.

20 Figures 17 and 18 illustrate the crystal structure of a compound denoted N2A (Gilardi, R.D., 1972, *Acta Crystallogr.*, Sect. B, 107).

Figures 19 and 20 show the crystal structure of a diazoaminobenzene (Gladkova & Kondrashev, 1972, *Kristallografiya* (41) 17 33.

25 Figures 21 and 22 illustrate how the molecule of Figures 15 and 16 crystallises in space.

30 Figure 23 shows a comparison of the mean, maximum and minimum extents of molecules which are active ligands (primulin and "analog"), and thiazin yellow (which is inactive as a ligand). The dimensions are given in angstrom units (AU).

35 Figure 24 shows a similar comparison for the basic benzothiazole nucleus (i.e. molecules 1b and 2 of Figure 5) and the diaminophenothiazines. These distances are carbon-to-carbon distances.

Figure 25 shows probability of extracellular tangles as a function of Braak staging. Stages 2 - 4 can be clearly distinguished from stage 1 on the basis of probability of extracellular tangles in E2/Trans and E4/HC.

5

Figure 26 shows probability of intracellular tangles as a function of Braak staging. Intracellular tangles provide a poor basis for discrimination of early stages in these regions, but a good basis for discriminating stages 4 and 5 using neocortical regions.

10

Figure 27 corresponds to Figure 25, but wherein cases with MMSE scores greater than 21 in the 12 months prior to death were selected. Similar results are obtained.

15

Figure 28 corresponds to Figure 26, but using cases with MMSE scores greater than 21 in the 12 months prior to death were selected. Similar results are obtained.

20

Figure 29 shows extracellular tangle densities (counts per mm^2) as a function of Braak staging. Similar results are obtained to those shown in Figure 25.

25

Figure 30 shows intracellular tangle densities (counts per mm^2) as a function of Braak staging.

30

Figure 31a shows a barely visible tangle visualised with thioflavin-T at 0.001% (arrowhead). In ifI suspensions such as this, tangles can be seen by blue fluorescence which is not distinct from that associated with binding of contaminants in the preparation. The bottom panel shows that blue tangle fluorescence produced by thioflavin-T at 0.001% is displaced by yellow tangle fluorescence produced by primulin at 0.001%. Figure 31b shows Electron-microscopy of PHFs labelled after Pronase digestion. (a) Chemical labelling with the biotinylated benzothiazole analogue shown as 4b in Figure 4. PHFs were deposited on a carbon-coated grid after Pronase digestion, and incubated briefly with compound 4b, and then incubated with an anti-biotin antibody preparation that had been conjugated with colloidal gold. Decoration of

35

isolated PHFs establishes that the compound 4b binds to the proteolytically stable PHF structure. (b) mAb 423 immunolabelling of isolated PHFs after Pronase digestion., followed by incubation with a gold-conjugated goat anti-mouse second antibody as described in Novak et al. 1993 (Novak M, Kabat J, Wischik CM (1993) "Molecular characterisation of the minimal protease-resistant tau unit of the Alzheimers' disease paired helical filament", EMBO J 12: 365-370). As shown in this paper, mAb 423 labels PHFs derived from intracellular tangles (which preserve N-terminal tau immunoreactivity in the fuzzy outer coat) poorly, but Pronase digested PHFs strongly. Likewise Mena et al. (1996) (Mena R, Edwards PC, Harrington CR, Mukaetova-Ladinska EB, Wischik CM, "Staging the pathological assembly of truncated tau protein into paired helical filaments in Alzheimer's disease", Acta Neuropathol 91: 633-641) shows that in intracellular tangles mAb 423 immunoreactivity is largely occluded, but can be exposed by formic acid pretreatment of sections.

Figure 32 shows "Preparation 2" of purified tau protein as described in Example 7 below.

Figure 33 shows a graphical plot of the results of a preparative run for dGA. "Purification-fold" is expressed as the ratio of specific immunoreactivity for each fraction (ie immunoreactivity / protein concentration) to specific immunoreactivity in the DE flow-through.

Figure 34 shows gel filtration chromatography of purified dGAE.

Apparent elution size in non-denaturing conditions:

1 - ~320 kD; 2 - ~80 kD; 3 - ~30 kD; 4 - ~10 kD

About 64% of mAb 7.51 immunoreactivity elutes in fractions corresponding to species of size > 15kD.

Figure 35 shows gel filtration chromatography of purified T40.

Apparent elution size in non-denaturing conditions:

1 - ~450 kD; 2 - ~160 kD; 3 - ~55 kD.

About 50% of mAb 499 immunoreactivity elutes in fractions corresponding to species of size > 60kD.

Figure 36 shows thionine activity against tau-tau binding in Preparations 1 & 2.

5 Figure 37 shows toloum chloride activity against tau-tau binding in Preparations 1 & 2.

Figure 38 shows DMMB activity against tau-tau binding in Preparations 1 & 2.

10

Figures 39a-c show that full-length tau protein (hT40) prepared according to the Preparation 2 protocol demonstrated minimal tau-tau binding activity when used in the aqueous-phase with dGA in the solid-phase(b). However, when hT40 was used in the solid-phase (c),
15 binding of dGAE was similar to that obtained for binding of dGAE to dGA in the solid-phase (a).

Figure 40 shows primulin and thiazin red have no inhibitory activity against tau-tau binding in Preparation 2, and indeed
20 enhance such binding at high concentrations.

Figure 41 shows blocking of inhibitory effects of 5 μ M DMMB on tau-tau binding in the presence of increasing concentrations of Primulin (a) and Thiazin Red (b), expressed as "molar excess"
25 relative to DMMB. Similar results are shown for 15 μ M DMMB in the presence of increasing concentrations of Primulin (c) and Thiazin Red (d).

Figure 42 shows attenuation and reversal of inhibition of Tau-Tau binding by DMMB in the presence of increasing molar excess of Primulin. For each graph, Tau-Tau binding is shown in the presence of constant concentrations of DMMB co-incubated with Primulin at 0x, 1x, 5x, 10x, 100x the DMMB concentrations shown. Inhibition of
30 Tau-Tau binding produced by DMMB is progressively attenuated and
35 reversed in the presence of increasing molar excess of Primulin.

Figure 43 shows a tau-Tau binding curve in the presence of 25 μ M DMMB and increasing Primulin molar excess as shown. Tau-Tau binding

can be modelled as follows:

$$\text{Binding} = (\text{B}_{\text{Max}} \times [\text{Prim}]) / (\text{Kd} + [\text{Prim}])$$

5 Where B_{Max} = 1.67
 Kd = 13.37
 r = 0.977 (observed vs predicted)

10 Figure 44 shows a tau-Tau binding curve in the presence of 5 μM DMMB and increasing Primulin molar excess as shown. Tau-Tau binding can be modelled as follows:

$$\text{Binding} = (\text{B}_{\text{Max}} \times [\text{Prim}]) / (\text{Kd} + [\text{Prim}])$$

15 Where B_{Max} = 1.38
 Kd = 13.86
 r = 0.927 (observed vs predicted)

20 Figure 45 shows a tau-Tau binding curve in the presence of 5 μM DMMB and increasing Thiazin Red molar excess as shown. Tau-Tau binding can be modelled as follows:

$$\text{Binding} = (\text{B}_{\text{Max}} \times [\text{TR}]) / (\text{Kd} + [\text{TR}])$$

25 Where B_{Max} = 1.64
 Kd = 17.45
 r = 0.915 (observed vs predicted)

Examples

30

Methods and materials

PHF-binding compounds

35 Compounds used herein were supplied by ICI Pharmaceuticals unless stated otherwise. Thioflavin-T and thiazine yellow were purchased from Fluka AG.

Quantitation of Fluorescence

Serial 16 μm sections are cut from the hippocampus of a case dying with clinically and neuropathologically confirmed AD. These
5 sections were stained with thioflavin-S at concentrations 0.01%, 0.001% or 0.0001% in water for 5-10 min, then washed in water, and mounted in Apathe's aqueous medium. In a second series of experiments, sections were cut from the hippocampus and nucleus
10 basalis of Meynert. These sections were stained with primulin at concentrations 0.1%, 0.01%, 0.001% and 0.00001% in water for 5-10 min, then washed in water, and mounted in Apathe's aqueous medium.

A Leitz fluorescence microscope fitted with a photo-multiplier tube (Model MPV-2) was used to quantitate fluorescence emission. Three
15 Leitz filter blocks were used as follows:

1. Filter block H2, code 513 417
Excitation range Band pass 390-490nm
Mirror RKP 510 (i.e. transmit below 510 nm)
20 Suppression filter LP 515 (i.e. reflect above 515 nm)
3. Filter block G, code 513 416
Excitation range Band pass 350-460 nm
Mirror RKP 510 (i.e. transmit below 510 nm)
25 Suppression filter LP 515 (i.e. reflect above 515 nm)
4. Filter block A, code 513 410
Excitation range UV band pass 340-380 nm
Mirror RKP 400 (i.e. transmit below 400 nm)
30 Suppression filter LP 430 (i.e. reflect above 430 nm)

Preparation of If I and II

IfI material was prepared as described by Wischik et al (1985) J
35 Cell Biol 100: 1905-1912.

IfII material was prepared as described by Wischik et al (1995)
Neurobiol Aging 16: 409-431. For experiments involving non-pronase

digested ifII an identical protocol was followed, omitting the pronase digestion step.

Spectrofluorimetry of ifII

5

These measurements were carried out in a Perkin-Elmer spectrofluorimeter (model MPF-3). A concentration of ligand of 0.00001% was routinely used for all measurements. Primulin was found to have an excitation peak at 370 nm and an emission peak at 515 nm. All measurements were therefore carried out at a standard excitation wavelength of 370 nm, and a constant slit width of 3 mm.

10

Competitive Binding Assay

IfI material was homogenised in a 0.2 ml glass homogeniser in PBS. To the suspension, test compounds were added to final concentrations ranging from 0.1% to 0.00001%. These were allowed to incubate for 5 min and primulin was added at equivalent or lower concentration. The suspensions were transferred to a glass slide, and examined by fluorescence microscopy across a range of fluorescence filter blocks, covering excitation and emission wavelengths between 380 nm and 570 nm. The end point sought in these observations was displacement of typical primulin fluorescence from tangle fragments.

20

25

Ligand Electronmicroscopy

PHFs derived from an ifI fraction were deposited on a carbon coated grid after pronase digestion, and incubated briefly with a preparation of biotinylated Primulin, and then incubated with an anti-biotin antibody that has been conjugated with colloidal gold by the method of Slot and Gueze (1981).

30

Succinylation and Chromatography of IfII

35

Washed ifII fractions were taken up in 8 M urea/50 mM borate (1 ml, pH 9) and sonicated, 1 ml succinic anhydride in acetone was added to a final concentration of 250 mM succinate in 4 ml, and the pH

was maintained at 8.5 with sodium hydroxide. The solution was clarified by centrifugation and applied to a Sephacryl S200 column equilibrated bicarbonate. The column eluate was monitored at either 230 or 280 nm.

5

Because succinylated fractions could not be visualised by Coomassie staining or silver staining of gels, bands were detected by autoradiography after specific chemical labelling of ifII fractions with Bolton-Hunter reagent (Amersham).

10

For photoaffinity labelling of PHF derived peptides, ifI or ifII fractions were pre-incubated with an I¹²⁵-labelled photolabile derivative.

15

The photolabelled fraction running at a Kav of 0.21 was concentrated by ultrafiltration through an Amicon YM2 membrane (10ml), digested with chymotrypsin (0.01 mg/ml) in 50 mM ammonium bicarbonate. Chymotryptic fragments for sequence analysis were isolated by Dr H.C. Thogersen by reverse phase HPLC using a C18 column, with a 0-100% acetyl nitrile gradient, with 0.1% trifluoroacetic acid. Chymotryptic peptides were sequenced.

20

Morphological Studies of PHFs in the presence of phenothiazines

25

For these experiments, ifII fractions were prepared as described above for electron microscopy. This material was either incubated directly with preparations of phenothiazines at final concentrations ranging between 0.1% and 0.0001% and then applied to carbon coated grids, and examined directly after LiPTA staining (1%). Alternatively, ifII suspensions were deposited on carbon coated grids, partially dried, and washed with solutions of phenothiazine. Such preparations were either stained directly with LiPTA or were processed further for immunoelectron microscopy using 6.423 as the primary antibody. Electron micrographs were recorded at nominal magnifications between 25,000 and 45,000.

30

35

Calculation of aggregated tau protein in the extracellular space expressed in µg/g of brain tissue as a function of Braak staging

Previously reported PHF-tau levels in pmol/g (P) and tangle counts per mm² (T) in a clinically and neuropathologically staged cohort (R.Y.K. Lai, et al., *Neurobiol Aging* 16, 433 (1995)) were used to derive an estimate of PHF-tau level per affected pyramidal cell (PC) in pg/cell using the same ELISA in human brain. The tangle count per mm² provides an estimate of the number of affected pyramidal cells within a volume 1 mm x 1 mm x 0.1 mm (0.0001 cm³), allowing that any tangle profile counted in a nominal 7 µm section could extend ~ 45 µm orthogonal to the section in either direction (S.M. Blinkov, I.I. Glezer, *The human brain in figures and tables, a quantitative handbook*; Plenum Press, NY, 1968, Table 204). The core PHF-tau level in pg/cm³ is 10 x P since the PHF-core tau fragment is 10 kD (C.M. Wischik, et al., *Proc. Natl. Acad. Sci. USA* 85, 4506 (1988)). From this, $PC = (P \times 10) / (T / 0.0001)$. At Braak stages 4-6 (H. Braak, E. Braak, *Acta Neuropathol.* 82, 239 (1991)), regional values for PC in grey matter were: frontal cortex, 0.13 ± 0.05 pg/cell; hippocampus, 0.60 ± 0.39 pg/cell; temporal cortex, 1.074 ± 0.44 ; entorhinal cortex 1.56 ± 0.63 pg/cell. These differences reflect anatomical differences, different regional rates of disease progression (C. Baner, H. Braak, P. Fischer, K. Jellinger, *Neurosci. Lett.* 162, 179 (1993), also Gertz et al., *Acta Neuropathol.* 95, 154 (1988)), and the degree to which tangle counts underestimate PHFs accumulating in dystrophic neuritis at more advanced stages of pathology (Lai et al, 1995, *loc cit*). The overall means provide an approximation for the PHF-levels per cell which would be relevant to AD. These are 0.37 ± 0.08 pg/cell for cases at Braak stages 1-3, and 1.08 ± 0.28 pg/cell for cases at Braak stages 4-6.

For the purpose of estimating extracellular aggregated PHF tau, shown below in the Table, and in Figure 26, 27, 29 and 31, tangles were counted as extracellular if mAb 423 immunoreactivity could be demonstrated after treatment of glass-mounted sections with 98% formic acid for 5 minutes prior to incubation with mAb 423 for 1 hr. For the avoidance of doubt, this methodology differs from that reported in Mena et al. (1996) where free-floating vibrotome sections were incubated briefly with formic acid, and then overnight with mAb 423. As shown in that report, this latter

overnight free-floating section protocol achieved maximal mAb 423 immunoreactivity in intracellular tangles, and showed that all intracellular tangles contain mAb 423 immunoreactivity, albeit in a state substantially occluded by the fuzzy outer coat of the PHF (see Figure 3). The purpose of the present protocol was to ensure maximum labelling of the Stage 3 and Stage 2 tangles illustrated in Figure 3. Some minor degree of labelling of intracellular tangles could not be entirely excluded, and counting attempts have also been made where a subjective discrimination has been attempted. However, the latter estimates do not agree with density or probability of labelling of tangles by mAb AT8 (Figures 28, 30, 32), which is solely intracellular, and shows completely different profiles with respect neuropathological stage from those revealed using mAb 423. For the purposes of the present calculations, therefore, counts of mAb 423-immunoreactive tangles were taken as substantially or entirely representative of extracellular tangle pathology at Stages 2 and 3 as shown in Figure 3, but substantially not Stage 1 of Figure 3. For the avoidance of doubt, the Stages referred to in Figure 3 are not Braak stages, but stages of degeneration of a single neurone containing a tangle.

The specific data shown in Table 1 was based on the following:

Number	BST	ME1T4	PC	PT4	REG3B	SE1T4
1	1.0000	0.3982	1.5600	0.6212	1.0000	0.3982
2	2.0000	6.7259	1.5600	10.4924	1.0000	2.7047
3	3.0000	14.9646	1.5600	23.3448	1.0000	2.9836
4	4.0000	33.6297	1.5600	52.4624	1.0000	10.9883
5	5.0000	44.3102	1.5600	69.1240	1.0000	13.0298
6	1.0000	0.0	0.6000	0.0	2.0000	0.0

58

	7	2.0000	1.3865	0.6000	0.8319	2.0000	0.4531
	8	3.0000	3.7169	0.6000	2.2302	2.0000	0.8060
5	9	4.0000	8.9384	0.6000	5.3630	2.0000	3.0048
	10	5.0000	23.9479	0.6000	14.3687	2.0000	4.0567
10	11	1.0000	0.0	0.6000	0.0	3.0000	0.0
	12	2.0000	0.0	0.6000	0.0	3.0000	0.0
	13	3.0000	0.0	0.6000	0.0	3.0000	0.0
15	14	4.0000	0.1293	0.6000	0.0776	3.0000	0.1293
	15	5.0000	2.2007	0.6000	1.3204	3.0000	1.0634

wherein:

20

BST is Braak Stage

ME1T4 is the extracellular tangle count

PC is an estimate of the PHF-tau concentration per cell (calculated as above)

25 PT4 is the PHF content ascribed to extracellular tangles (PC x ME1T4)

REG3B is the grouping of brain regions into 3 groups as per Figs 26 and 27 of the SE1T4 is the standard error of the extracellular tangle count

30

Example 1 - aggregated tau in Braak staging

Based on immunochemical properties (Refs 26, 27, 30), it is possible to distinguish intracellular tangles from extracellular
 35 tangles. Both frequency of cases with tangles in these categories (ie probability) and their quantity (ie counts per mm²) were determined in a prospective case series and grouped into the

regions known to represent stages in the progression of pathology according to the system of Braak and Braak

As shown in Figure 25, stages 2 - 4 can be clearly distinguished from stage 1 on the basis of probability of extracellular tangles in E2/Trans

and E4/HC. Also shown are the figures for F/T/P regions (neocortical regions - frontal, temporal, parietal).

Conversely, intracellular tangles provide a poor basis for discrimination of early stages in these regions, but a good basis for discriminating stages 4 and 5 using neocortical regions. Similarly, when cases with MMSE scores greater than 21 in the 12 months prior to death were selected, similar results were obtained. Again, similar results were obtained when tangle densities were determined.

These results can be converted into approximations for the quantity of aggregated tau protein in the extracellular space expressed in $\mu\text{g/g}$ of brain tissue as described in Materials and Methods above. The results are shown in Table 1. These are underestimates, as the tangle counts underestimate the quantity of aggregated tau protein.

Table 1: ESTIMATE PHF-TAU CONTENT IN ECT'S BY REGION AND STAGE

REGION	BST	PHF/TAU ($\mu\text{g/g}$)
E2/TRANS	1	0.62
	2	10.49
	3	23.34
	4	52.46
	5	69.12
HC/E4	1	0
	2	0.83
	3	2.23
	4	5.36

	5	14.36
F/T/P	1	0
	2	0
5	3	0
	4	0.08
	5	1.32

Table 1 shows quantity of aggregated tau protein in the extracellular space expressed in $\mu\text{g/g}$ of brain tissue as a function of Braak staging. The data was calculated as described in the materials and methods.

In summary, these results demonstrate that extracellular deposits of PHF-tau in medial temporal lobe structures provide a basis for empirical staging of the neurofibrillary degeneration of AD. Such staging could only be accomplished by radio-imaging methods provided suitable ligands could be created.

Example 2 - assessment of compounds binding within the aggregated repeat domain of PHF-core tau protein

A prototype compound was obtained as one component of the crude, commercially-available preparation of thioflavin-S was separated into ~20 distinct constituents by analytical thin-layer chromatography, and preparative chromatography. Tests showed that not all of the constituents were able to act as effective tangle ligands. Specifically, pure primulin (Figure 5, compound 1a) was found to label tangles, but the benzothiazole thioflavin-T (Figure 5, compound 1b) was much less effective, although it labelled amyloid preferentially.

Furthermore, compound 1a was found to displace compound 1b at tangles when the latter was introduced at 10-fold excess into crude tangle extracts.

A possible difference was postulated to be the sulphonate group at position 1 (Figure 5, compound 2 [2 - (4 - amino phenyl) - 6 -

5 methyl - 1 - sulphonate benzothiazole]). However, primulin (compound 1a) was found to displace this from tangles (though not amyloid). Therefore, tangle labelling is not due solely to the sulphonated benzothiazole structure, indicating that a longer aromatic structure is required.

10 Purified thiazin red (Figure 5, compound 3a) was found to compete with primulin at equivalent concentrations, whereas the compound 3b (thiazin yellow, Figure 5) did not. Therefore, an extended aromatic benzothiazole structure does not, *per se*, determine high binding affinity within tangles.

15 In order to define a minimum critical requirement for competitive binding, the sulphonated benzothiazole was extended by addition of a single phenyl group across a diamino-linkage. This compound (Figure 4, compound 4a), although not fluorescent, was found to compete out thiazin red and primulin fluorescence at equivalent concentrations. Compound 4a therefore defines the minimum critical structure required for high affinity binding within the tangle.

20 In order to prove that the binding site within the tangle was in fact the PHF itself, compound 4a was further extended with addition of a biotin group (Figure 4, compound 4b). Since this was still found to compete primulin and thiazin red, compound 4b preserved high affinity binding within the tangle. Furthermore, immunogold-conjugated anti-biotin antibody was found to label isolated PHFs pre-incubated with compound 4b, whereas no labelling was demonstrated without pre-incubation or pre-incubation with biotin alone (Fig 31b). Finally, when a photo-activated conjugate of the compound was prepared, it was possible to identify and sequence the labelled protein. This was found to be the same core tau fragment as that isolated from the core of the PHF, which comprises the repeat region of the tau protein.

35 In summary, these results demonstrate unequivocally that the binding site for compounds 4a and 4b is within the aggregated repeat domain of the tau protein of the PHF-core. Furthermore, they demonstrate that compound 4a can be used as a chelate for

addition of functional groups without disturbing ligand activity within the PHF core. Therefore, compound 4a could be used as chelate for addition of technetium or other imaging moiety to generate a ligand suitable for detecting e.g. extracellular tangles in AD.

Example 3 - determination of optimum dimensions of ligand molecules

Figure 14 shows three of the structures described above, along with their dimensions as indicated. For example, the C11-C1 distance and C10-C1 distance are shown for primulin, a benzothiazole analogue (denoted 'analog'), and 'thiazin yellow'.

Figure 15 and 16 illustrate the crystal structures of the 'B' part of the primulin structure (Soon-Beng Teo et al., 1995, Acta Crystallogr., Sect. C, 591. As can be seen from Figure 16, which is a 'side-on' view, the molecule is essentially flat, although it has a slight twist. The 'A' part of primulin can be computed from the same molecule. From this, one can derive measures of A+B, which provide an indication of the actual length of one of the active species of the present invention.

To compute the size of the "analog" shown in Figure 14, measurement A was used from the data of Figure 15, and measurement B was determined from a molecule denoted N2A and shown in Figures 17 and 18 (Gilardi, R.D., 1972, Acta Crystallogr., Sect. B, 107). As can be seen from the side-on view in Figure 18, this part of the molecule is completely flat. The same measurements apply to thiazin red, which is identical in its dimensions to the "analog".

The size of thiazin yellow (shown in Figure 14) was determined as follows. The 'A' part comes from the molecule of Figure 15 which was used for primulin, while the 'B' part comes from the molecule shown in Figures 19 and 20 (Gladkova et al., 1972, Kristallografiya 41). Again, part B of the molecule is completely flat, and the only difference with respect to the molecule shown in Figure 17 is the distance between the aromatic groups.

Figures 21 and 22 illustrate how the molecule of Figure 15 crystallises in space. As can be seen, the molecule forms an alternating 'herring-bone' pattern, and does not stack. In comparison, the crystal structure of methylene blue indicates that the molecules form stacks with alternating sheets of water molecules between the pi-bonded stacks.

Table 2 tabulates the minimum, maximum and mean dimensions for primulin ("PRIM"), the analog ("ANAL"), thiazin yellow, and the benzothiazole unit alone (i.e. structures 1b and 2 as shown in Figure 5). The corresponding methylene blue dimensions are given as 'MBCC' (carbon to carbon) and 'MBNN' (nitrogen to nitrogen):

	1	2	3	4	5	6
	PRIM	ANAL	THIAZY	BENZTHIA	MBCC	MBNN
Minimum	14.7830	15.0500	15.7270	8.7700	7.0849	9.9600
Maximum	15.1120	15.2610	16.1380	8.9550	7.4443	9.9600
Mean	14.9475	15.1680	15.9273	8.8625	7.3031	9.9600

Figure 23 shows a comparison of the mean, maximum and minimum extents of molecules which are active ligands (primulin and "analog"), and thiazin yellow (which is inactive as a ligand). The dimensions are given in angstrom units (AU). In Figure 24, a similar comparison is made for the basic benzothiazole nucleus (i.e. molecules 1b and 2 of Figure 5) and the diaminophenothiazines. These distances are carbon-to-carbon distances.

The above results illustrate that the molecules provided herein are substantially flat. There is, however, a fundamental difference in activity between ligands according to the present invention and other molecules discussed above. As is shown in the Figures, suitable ligands according to the invention comprise long, flat molecules with dimensions between 14.783 and 15.261 AU. On the other hand, a longer molecule, such as thiazin yellow, which exceeds these dimensions (mean 15.927 AU) does not serve as an effective ligand, even though it is flat. However, certain shorter, flat, molecules bind preferentially to amyloid.

Example 4 - PET using ligand molecules and inhibitors

Figures 11 to 13 indicate typical synthesis methods which could be used to convert either the diaminophenothiazines or the "analog" into positron emitting species.

Figure 11b in particular shows a method whereby thionin is treated with NaH followed by labelled methyl iodide to give methylene blue.

A similar procedure can be adopted for the synthesis of methylene blue starting from Azure A and Azure B. Other strong bases may also be used.

Other methods that may be used include HCl and labelled MeOH; labelled trimethyl phosphoric acid; labelled dimethylsulphoxide and Labelled formaldehyde. The chemistry of the syntheses and general methodology are all familiar to persons skilled in the art. These examples are given without any implied restriction as to ultimate methodology.

Example 5 - blocking ligands

Compounds such as thioflavin-T and -S strongly stain amyloid deposits. However Figure 31 demonstrates that such compounds can be displaced from tangles by primulin. Therefore these compounds may be used as blocking reagents to saturate binding sites which are not of interest without inhibiting the binding of ligands to aggregated tau.

Example 6 - comparison of ligand molecules and inhibitors

There appears to be a fundamental difference in activity of the molecules which are effective ligands, compared with those which are effective inhibitors of tau-tau binding. The benzothiazole molecule does not disrupt PHFs, nor indeed do any of the ligands, whereas the diaminophenothiazine series constitute PHF-disrupters and tau aggregation inhibitors.

Further investigations into the relationship between aggregation-dependent tau ligands and tau aggregation inhibitors were carried out using primulin. Primulin in solution has a fluorescence peak at 520 nm. This shifts to 470 nm when primulin is bound within a pure preparation of PHFs (Figure 6). Treatment of PHFs with citraconic anhydride, which has been shown to disrupt the structure of the PHF and liberate free tau (as well as reversing the charge on lysines), was found abolish the 470 nm fluorescence peak (Figure 7).

Therefore, binding by such compounds is dependent on the polymerised state of tau found in the PHF, but is not present in free tau.

Compounds have been identified which disrupt the structure of the PHF and reverse the proteolytic stability of the PHF core (see WO 96/30766). Examples of such compounds are shown in accompanying Figure 8. The present inventors have now identified that these compounds bind to tau at a specific binding site within the high affinity tau-tau binding domain. However, it is found that such compounds may not disrupt the binding of primulin to tau in aggregated tau molecules, as shown by the retention of the fluorescence peak of primulin at 470 nm in the presence of alcian blue (Figure 9).

Thus it appears that although alcian blue can inhibit tau-tau interactions, either the site of the inhibition, or perhaps the order of binding interaction at which it acts, are such as to leave binding site for SB-ligands extant. Thus compounds which act as ligands of aggregated tau do not appear to bind at the same site(s) as compounds which are tau-aggregation inhibitors, although they may still affect the inhibitory properties of those inhibitors (see Example 7 below).

This point was further examined by studying the potency of typical aggregated tau ligands as tau-aggregation inhibitors. It has been shown previously that tau aggregation inhibitors (e.g. diaminophenothiazines) can be identified on the basis of inhibition of tau-tau binding in a solid-phase assay (WO 96/30766). When tested in the same assay, primulin and thiazin red were found to be

weak inhibitors of tau-tau binding (Figure 10). Thus, although these compounds are potent ligands for tau within the PHF-core, they are at most weak inhibitors at the site required for inhibition of tau-tau binding.

5

Demonstration that compounds of the diaminophenothiazine-like class bind tau in the aggregated state is provided by the direct demonstration of disruption of PHF structure in the presence of sufficiently high concentrations, particularly of compounds such as methylene blue. Thus, compounds of the diaminophenothiazine-like class which are inhibitors of tau-tau binding can serve as aggregated-tau ligands at lower concentrations.

10

In summary, the inventors have found that it is possible to define two classes of binding site within the core-PHF tau aggregate. Both are potentially useful for the development of radiological imaging ligands:

15

(i) Sulphonated benzothiazole-like sites: compounds of this type, associated with suitable chelates such as technetium, may serve as ligands for extracellular tangles, due to their size and charge.

20

(ii) Diaminophenothiazine-like sites: such compounds, when suitably labelled with a positron-emitting functional group, would serve as ligands for all tau aggregates, and would be capable of crossing the blood-brain-barrier (Ref 36) and entering cells. Thus these compounds, and derivatives thereof, have potential use in the labelling of intracellular tangles, e.g. those present in the brains of AD patients, or intracellular tangles when used at lower concentration.

25

30

Example 7 - example assays for identifying further diagnostic ligands based on relief of inhibition

(i) Preparation of Tau protein in which partial aggregation has occurred.

35

The preparation ("Preparation 2") is shown schematically in Figure

32, and differs from earlier described methods (e.g. in WO96/30766- "Preparation 1" - shown in (ii) below).

The recombinant cDNA plasmids are those described in WO96/30766 the disclosure of which is herein incorporated by reference.

Briefly, tau cDNA was generated using standard protocols (Sambrook, Fritsch & Maniatis "Molecular cloning. A Laboratory Manual" (1989) Cold Spring Harbor Laboratory, N.Y.) from mRNA isolated from brain tissue of an Alzheimer patient whose tissue was obtained 3 h after death. The cDNA library was screened with synthetic 17-mer oligonucleotide probes derived from the sequence from part of a PHF core protein (Goedert et al. (1988) Proc. Natl. Acad. Sci. USA, 85, 4051-4055). Full length cDNA clones were subcloned into the EcoRI site of M13mp19 and site-directed mutagenesis used to introduce a NdeI site in the context of the initiator codon. Following cleavage with NdeI and EcoRI, the resulting cDNA fragments were subcloned downstream of the T7 RNA polymerase promoter into NdeI/EcoRI -cut expression plasmid pRK172 (McLeod et al. (1987) EMBO J., 6, 729-736). pRK172 is a derivative of pBR322 that is propagated at very high copy number in E. coli due to removal of the pBR322 copy number control region. The plasmid carries an ampicillin resistance gene for selection of recombinant clones.

cDNA constructs coding for truncated forms of tau were prepared from mRNA as described in Novak et al. (1993) EMBO J., 12, 365-370. The mRNA was used as a template for polymerase chain reaction (PCR) using specific oligonucleotide primers. The sense primer contained an NdeI site and the anti-sense, an EcoRI site. PCR fragments were subcloned into pRK172 as described above. The primers used for construction of dGAE are given in Figure 22. The authenticity of all DNA fragments used for expression was confirmed by full length sequencing of both strands.

Details for the construction of htau40 ("T40") cDNA are described in (Goedert et al. (1989), Neuron 3: 519-526). This sequence is the largest form of tau found in the CNS and encodes tau protein that contains both the 2 N-terminal inserts of 29 amino acids each

and an extra 31 amino acid repeat in the tubulin-binding domain. The DNA sequence and its predicted amino acid sequence are shown in Figure 21 (SEQ ID NO: 4).

5 Recombinant plasmids were used to transform *E. coli* BL21 (DE3) a strain used for prokaryotic expression which carries a chromosomal copy of the bacteriophage T7 RNA polymerase gene under control of the *lac* UV5 promoter (Studier and Moffat (1986), *J. Mol. Biol.* 189, 113-130). Exponentially growing cultures were induced with IPTG
10 (iso-propyl thiogalactoside) for 3h.

Large-scale purification (1 litre bacterial culture) of tau fragments was carried out as described by Goedert and Jakes (1990, *EMBO J.*, 9, 4225-4230), with minor modifications. Cells were
15 disrupted by rapid freezing of the cell pellet in liquid nitrogen. The pellets were then suspended in buffer containing 50 mM PIPES, 1 mM dithiothreitol (DTT) (pH 6.8). The thermostable proteins in the supernatant were dialysed against PIPES/DTT, then applied to a column containing phosphocellulose equilibrated in the same buffer.
20 Tau protein was eluted with a gradient of NaCl (0-0.5M) in the above buffer. Fractions were analysed by SDS-PAGE and both Coomassie staining and immunoblotting. Those fractions containing tau were pooled, dialysed against 25mM MES, 1 mM DTT (pH 6.25) and stored at -20°C at approximately 5 mg/ml. Protein concentrations
25 were measured by the Lowry method (Harrington CR(1990), "Lowry protein assay containing sodium dodecyl sulphate in microtitre plates for protein determinations on fractions from brain tissue", *Analytical Biochemistry* 186: 285-287).

30 Preparation 2 differs from Preparation 1 above in the following respects: (1) The concentration of cells at the sonication stage is increased by 5-fold. (2) A batchwise adsorption of non-tau proteins to DE52 is included. (3) The proteins are not subjected to heat treatment. (4) The final step involves concentration using
35 polyethylene glycol.

Escherichia coli is grown in 2xTY medium (Oxoid) supplemented with ampicillin (50 micrograms per ml) to late-logarithmic phase. Cells

from 5 litres of culture are harvested by centrifugation and the cell pellets frozen rapidly over liquid nitrogen. The pellets are taken up with 50 mM PIPES (pH 6.8) containing 1mM EDTA, 1mM dithiothreitol and 1mM phenylmethylsulphonylfluoride (PMSF) and bacteria lysed by sonication (2 x 3 min) at 4C. The mixture is centrifuged at 10,000 rpm for 20 min. The supernatant is rotated with 1 gram of Whatman DE52 for 3 hr at 4C. The mixture is separated on a column and the flow-through material that fails to bind to DE52 is incubated for 3 hr at 4C, with rotation, with 0.4 g of Whatman P11 (freshly regenerated according to manufacturer's recommendations). The column is washed with column buffer (50 mM PIPES, pH 6.8 containing 1mM EGTA, 5mM EDTA, 0.2mM MgCl₂, 5mM β -mercaptoethanol and 1mM PMSF). Tau protein is eluted stepwise with a 0.1 to 1 M gradient of KCl in column buffer. Fractions containing tau (determined by immunoassay) are pooled and dialysed against 80mM PIPES (pH6.8) containing 1mM EGTA, 1mM MgCl₂ and 5mM β -mercaptoethanol, using dialysis tubing with a molecular weight cut-off of 1,000. The dialysate is concentrated by applying polyethylene glycol 8000 to the outside of the sac for 2-3 hr. The final concentration of tau ranged from 3 to 10 mg per ml.

In a typical large scale preparative run, the specific immunoreactivity of tau is purified approximately 30- to 40-fold from the material that fails to bind to DE52. Approximately 60% of the tau is recovered in the final product with 10% failing to bind to P11 and the remainder in fractions ignored from the column.

Table 3 shows the details of a preparative run for dGA.

"Purification-fold" is expressed as the ratio of specific immunoreactivity for each fraction (ie immunoreactivity / protein concentration) to specific immunoreactivity in the DE flow-through.

	Vol (ml)	mg/ml	AU/ml	AU	AU/mg	Purif'n (-fold)	Recovery (%AU)
Total cells	200	26	40000	8000000	1538		
DE flow thro'	240	20	40000	9600000	2000	1.00	100%
P11 flow thro'	220	19	5000	1100000	263		11% in P11 flow thro'
Fractions (6ml each)	#1	1.48	5000	30000	3388	2	
	#2	0.98	20000	120000	20492	10	
	#3	1.60	80000	480000	50000	25	
	#4	2.69	200000	1200000	74294	37	
	#5	3.09	200000	1200000	64767	32	
	#6	3.31	200000	1200000	60459	30	62% in pooled fractions
	#7	2.71	200000	1200000	73746	37	
	#8	2.70	125000	750000	46296	23	
	#9	2.20	100000	600000	45537	23	20% in discarded fractions
	#10	1.83	80000	480000	43764	22	
Pure dGA	14.5	8.4	600000	8700000	71429	36	

Figure 33 is a graphical plot of the data from Table 3.

Table 4 summarises yields from typical preparative runs for the tau protein species: dGA, dGAE and hT40.

Tau preparation	Protein concentration (mg/ml) based on:		Immunoreactivity (mAb 7/51; AU/ml)	Protein yield (per 10 litres)
	abs at 280 nm	Protein assay (BSA ref)		
dGAE (711)	3.3	3.8	300,000	100 mg
dGA (1511)	8.1	8.4	600,000	122 mg
T40 (1311)	3.5	3.1	125,000	120 mg
<u>based on extinction coefficients</u>		<u>abs at 280 nm</u>		
BSA (reference)	10 mg/ml	6.6		
Tau protein	10 mg/ml	14.7		

Proteins were separated by running down a 50 x 1cm sepharose CL-6B gel filtration column equilibrated with PBS buffer and run at room temperature. A molecular weight standard curve was prepared for the column by running molecular weight markers over the range 12,400 to 200,000 down the column. A standard curve was prepared by plotting the \log_{10} of the Mr in Kd against V_e/V_o for each protein standard, where V_e is the elution volume for the standard and V_o is the void volume for the column determined with blue dextran.

T40 or dGAE were loaded in 0.5ml buffer containing 5% glycerol and collected in 1ml fractions. Presence of protein in the fractions was determined spectrophotometrically by the absorbance at 280nm. Presence of dGAE or T40 was detected by ELISA with monoclonal
5 antibody 7/51 or 499 respectively. ELISA assay was carried out in 96 well PVC plates as follows:

50µl samples of each fraction incubate 1hr at 37C, wash plate in 0.05% tween-20 and then block binding sites with 200µl PBS + 2% non fat milk powder for 1hr at 37C. Wash plate in 0.05% tween-20,
10 incubate with 50µl primary antibody diluted 1:10 in PBS + 2% non fat milk powder for 1hr at 37C. Wash plate in 0.05% tween-20, incubate with 50µl secondary antibody (goat anti-mouse IgG:HRP conjugate) diluted in PBS + 2% non fat milk powder for 1hr at 37C. Wash plate in 0.05% tween-20 then rinse in deionised water, add
15 50µl freshly prepared substrate (TMB [tetramethylbenzidine] in sodium acetate buffer pH5.0 with H₂O₂, freshly prepared) and read rate of change in OD 650 over 2 minutes.

The elution profile of purified dGAE and purified T40 is shown in
20 Figures 34 (dGAE) and 35 (hT40). Although both these fragments typically run at about 12 kD and 55 kD respectively, about 64% of mAb 7.51 immunoreactivity (dGAE) elutes in fractions corresponding to species of size > 15kD and about 50% of mAb 499 immunoreactivity (hT40) eluted in fractions corresponding to species of size > 60kD.
25 Thus the tau proteins are present, at least in part, in pre-aggregated form.

(ii) effect of tau aggregation inhibitors measured using different preparations of Tau protein

30

When the proteins dGA and dGAE were prepared as indicated above ("Preparation 2"), the properties of the tau-tau binding assay were altered relative to the properties obtained using the preparative method described in WO96/30766 ("Preparation 1").

35

The assay is performed using 96 well PVC plates (Falcon Cat. No. 353912 are used), and the following steps:

1. 50 µl dGA (~10 µg/ml) in carbonate buffer, incubate 1 h at 37°C.

(Carbonate buffer: 50 mM carbonate bicarbonate, pH 9.6 (Na₂CO₃ 1.59 g/l, NaHCO₃ 2.93 g/l))

5

2. Wash plate in 0.05% Tween-20.

3 200 µl PBS+2% Marvel, incubate 1 h at 37°C.

10 4. Rinse plate 2X in deionised water, then wash in 0.05% Tween-20.

5 50 µl dGAE (~10 µg/ml) plus drug in PBS+1% fish skin gelatin+0.05% Tween-20, incubate 1 h at 37°C.

15 6 Wash plate in 0.05% Tween-20.

7 50 µl antibody 423 (1:10 dilution in PBS+2% Marvel), incubate 1 h at 37°C.

20 8 Rinse plate 2X in deionised water, then wash in 0.05% Tween-20.

9 50 µl HRP-anti-mouse (1:1000 dilution in PBS+0.05% Tween-20), incubate 1 h at 37°C.

25 10 Wash plate in 0.05% Tween-20, then rinse 1X with deionised water.

11 50 µl substrate solution, read immediately initial rate over 2 min in plate reader at OD650.

30

(Substrate solution: 50 mM sodium acetate, pH 5.0 + TMB (1 ml/100 ml of a 10 mg/ml solution in DMSO) + H₂O₂ (10 µl/100 ml)).

The compounds thionine and toloum chloride were found to require higher concentrations to exert inhibitory effects in Preparation 2 than in Preparation 1. This is shown in Figures 36 and 37.

Furthermore, the compound dimethyl methylene blue (DMMB) was found to have a higher inhibitory potency in Preparation 2 than in

Preparation 1. This is shown in Figure 38.

As similar differences could be seen in Tau protein prepared by the Preparation 1 method, but which had been allowed to aggregate in vitro over time, the interpretation of this effect is as follows. Higher concentrations of compounds such as thionine and tolonium chloride are required since two effects have to be achieved in order to achieve maximal inhibition of tau-tau binding:

1. disruption of pre-existing aggregates in the aqueous-phase;
2. inhibition of binding of aqueous-phase species to the solid phase.

The greater potency of DMMB in the Preparation 2 assay can be explained by greater binding affinity at the site of action required for both inhibitory effects.

Full-length tau protein (hT40) prepared according to the Preparation 2 protocol demonstrated minimal tau-tau binding activity when used in the aqueous-phase with dGA in the solid-phase. However, when hT40 was used in the solid-phase, binding of dGAE was similar to that obtained for binding of dGAE to dGA in the solid-phase (see Figure 39a-c). The interpretation is that in hT40 aggregates formed in the aqueous-phase tau-tau binding has already occurred in or through domains required for the binding interaction with dGA in the solid-phase. When hT40 is first plated in the solid-phase, the binding to PVC unfolds the protein/aggregates in such a manner as to make the critical tau-tau binding sites available.

(iii) effect of tau aggregation inhibitors measured using different preparations of Tau protein

In the Preparation 2 assay format, potent ligands of the kind typified by Primulin and Thiazin Red have no inhibitory activity on tau-tau binding. This is shown in Figure 40 (cf. Figure 10). Indeed in this assay, these compounds enhance tau-tau binding at concentrations greater than 100 μ M (i.e. 100-fold molar ratio with

respect to tau protein).

DMMB typically reduces tau-tau binding (with tau present typically at 1 μM) to 23% at DMMB 5 μM and 17% at DMMB 15 μM relative to that seen in the absence of DMMB.

Unexpectedly, this inhibitory effect can be completely reversed by co-incubation in the presence of increasing concentrations of high affinity ligands typified by Primulin and Thiazin Red. This is shown in Figure 41.

Therefore, aggregated-tau ligands may be characterised functionally as compounds which do not themselves inhibit tau-tau binding, but block the inhibitory effects of potent inhibitors of tau-tau binding.

As can be seen in Figure 42, inhibition of Tau-Tau binding produced by DMMB is progressively attenuated and reversed in the presence of increasing molar excess of Primulin. A similar effect can be shown for Thiazin Red. This indicates that the maximum inhibitory effect of DMMB is reduced by these compounds, and hence that they are acting as non-competitive inhibitors of DMMB. One possible explanation might be that the ligands stabilise the tau aggregates used in the assay, for example in regions outside the critical binding domain required for DMMB activity, and hence prevent the inhibitory effect of DMMB on Tau-Tau binding.

Figures 43 - 45 show that for any given concentration of DMMB, there is quantitative enhancement of Tau-Tau binding in the presence of Primulin (43, 44) or Thiazin Red (45) which can be modelled by a standard Michaelis-Menten equation. This implies that the Tau-aggregation enhancement effect of these ligands is proportional to the fraction of ligand-binding sites occupied, presumably within the Tau aggregates introduced into the aqueous-phase of the assay. The mean B_{max} for both ligands is ~ 1.6 . That is, the maximum ligand effect is to produce 1.6-fold the Tau-Tau binding signal seen in the absence of any drug. The mean K_d for this effect is $\sim 15x$. That is, for any given concentration of DMMB

> 4 μ M, 50% of maximal enhancement of Tau-Tau binding can be seen when the ligand molar excess is 15-fold relative to the concentration of DMMB.

5 Example 8 - example assays for identifying further diagnostic ligands based on ligands provided herein

Having defined two classes of ligands as described above suitable for labelling PHFs in AD, further ligands can be developed using
10 the compounds/derivatives in screening assays. Furthermore, modelling methods can be based on the ligands already presented.

(i) Identification of novel ligands at the sulphonated-benzothiazole site.

15

Using a suitably labelled preparation of a known sulphonated benzothiazole, incubated with a preparation of aggregated tau molecules (e.g. preaggregated tau in solution, or bound to a solid phase, or highly enriched PHFs isolated from AD brain - see
20 WO96/30766) compounds suspected of being suitable ligands can be introduced, and their capacity to compete with the known ligand in such a way as to prevent binding within the PHF can be tested.

(ii) Identification of novel ligands at the phenothiazine site.

25

The tau-tau binding assay described in WO 96/30766 can be used as a preliminary screen to identify potential inhibitors at the tau-tau binding site. Likewise, a suitably-labelled preparation of known diaminophenothiazines, incubated with a aggregated tau as described
30 above, could be used to screen for other compounds which are suspected of being competitors at this PHF-binding site and thus potentially suitable PHF ligands.

The physical implementation of competitive assays is well known in
35 the art. It may include measurement of fluorescence, radioactivity or any other suitable reporting system which derives from sulphonated benzothiazole-like compounds or diaminophenothiazine-like compounds not bound to PHFs, i.e. those which remain in

solution.

References

- 5 1 DeToleda-Morrell, L. et al. (1997), *Neurobiology of Aging* 18, 5, 463-8;
- 2 De Leon et al. (1997), *Neurobiol. Of Aging*, 18, 1, 1-11;
- 3 Mori, E et al. (1997), *Am. J. Psychiatry* 154:1, p18;
- 4 Juottonen, K. (1998); *J. Neurol. Neurosurg. Psychiatry* 65, 322-
10 327;
- 5 Bobinski, M. et al. (1999), *Lancet* 353, p.38;
- 6 Fox, N.C. (1999) *Neurol.* 52, 1687-9;
- 7 Jack, C.R. et al. (1997) *Neurol.* 49: 786-794;
- 8 Fox, N. et al. (1996), *Brain* 119, 2001-7;
- 15 9 Johnson, K.A. et al. (1998), *Neurol.* 50, 1563-1571;
- 10 Perez-Tur, J. et al. (1999), *Neurol.* 53, 411-3;
- 11 Lehtovirta, M. et al. (1998) *J. Neurol. Neurosurg. Psychiatry* 64, 742-6;
- 12 Nagy, Zs et al. (1999), *Dement. Geriatr. Cogn. Disord.* 10, 109-
20 114;
- 13 Ishii, K. et al. (1998), *Neurol.* 51, 125-130;
- 14 Imamura, T et al. (1997), *Neurosci. Lett.* 235, 49-52;
- 15 Minoshima, S. et al. (1997), *Ann. Arol.* 42, 85-94;
- 16 Ibanez, V. et al. (1998), *Neurol.* 50, 1585-1593;
- 25 17 Wischik, C.W. et al. (2000) "Neurobiology of Alzheimer's Disease", Eds. Dawbarn et al., *The Molecular and Cellular Neurobiology Series*, Bios Scientific Publishers, Oxford).
- 18 Carretero, M.T. et al. (1995), *Dementia* 6, 281-5;
- 19 Villareal, D.T. et al. (1998), *Alzheimer's Dis. Rev.* 3, 142-152;
- 30 20 Marin, D.B. et al. (1998), *Artherosclerosis* 140, 173-180;
- 21 Kuller, L.H. et al. (1998), *Stroke* 29, 388-398;
- 22 Vargha-Khadem, F. et al. (1997), *Science* 277, p376;
- 23 Willingham, D.B. (1997), *Neuron* 18, 5-8;
- 24 Lakmache, Y. et al. (1995), *PNAS USA* 95, 9042-6;
- 35 25 Hodges, J.R. et al. (1999), *PNAS USA* 96, 9444-8;
- 26 Mena, R. et al. (1995), *Acta Neuropathol.* 89, 50-6;
- 27 Mena, R. et al. (1996), *Acta Neuropathol.* 91, 633-641;

- 28 (deleted)
- 29 Lai, R. *et al.* (1995) *Neurobiol. Aging* 16, 3, 433-445;
- 30 Bondareff, W. *et al.* (1994) *J. Neuropathol. Exp. Neurol.* 53, 2,
158-164;
- 5 31 Resch, J.F. *et al.* (1991) *Bioorg. Med. Chem. Lett.* 1, 10, 519-
522;
- 32 Novak, M. *et al.* (1993), *EMBO J.* 12, 1, 365-370;
- 33 Wischik, C.W. *et al.* (1996), *PNAS USA* 93, 11213-8;
- 34 Wischik C.W. *et al.* (1989), *Curr. Opin. Cell Biol.* 1, 115-122;
- 10 35 WO 96/30766;
- 36 Muller, T. (1992), *Acta Anat.* 144, 39-44.

Claims

- 1 A method for determining the stage of neurofibrillary
degeneration associated with a tauopathy in a subject believed to
5 suffer from the disease, which method comprises the steps of:
(i) introducing into the subject a ligand capable of labelling
aggregated paired helical filament (PHF) tau protein,
(ii) determining the presence and/or amount of ligand bound to
extracellular aggregated PHF tau in the medial temporal lobe of the
10 brain of the subject,
(iii) correlating the result of the determination made in (ii) with
the extent of neurofibrillary degeneration in the subject.
- 2 A method as claimed in claim 1 wherein the determination in step
15 (ii) is used to establish the density ligand binding.
- 3 A method as claimed in claim 1 or claim 2 wherein the
correlation in step (iii) is made by reference to historical data.
- 20 4 A method as claimed in any one of the preceding claims wherein
the tauopathy is Alzheimer Disease (AD).
- 5 A method as claimed in claim 4 wherein the extent of
neurofibrillary degeneration is related to the neuropathological
25 staging of the progression of AD according to the defined
hierarchical system shown in Figure 2c.
- 6 A method as claimed in any one of the preceding claims wherein
the ligand is capable of crossing the blood brain barrier.
30
- 7 a method as claimed in any one of the preceding claims wherein
the ligand is conjugated, chelated, or otherwise associated, with a
detectable chemical group.
- 35 8 A method as claimed in claim 7 wherein the ligand is labelled
for SPECT and is not capable taken up intracellularly.
- 9 A method as claimed in claim 8 wherein a ligand comprises a


technetium-chelating group.

10 A method as claimed in claim 7 wherein the ligand is labelled for positron emission tomography (PET).

5

11 A method as claimed in claim 10 wherein the ligand comprises a positron-emitting carbon, optionally incorporated into a methyl group present in the ligand.

10 12 A method as claimed in any one of the preceding claims wherein the ligand is a compound of the formula:



wherein:

W is S, O, or NH;

15

exactly one of X, Y, and Z is CH or N;

the others of X, Y, and Z are CH;

M¹ is an alkali metal cation selected from: Li, Na, K, or Cs.

RL is a rigid linker group;

20


Ar¹ is an C₅₋₂₀aryl group;

n is an integer from 0 to 3; and,

each R^{BT} is independently a core substituent.

13 A method as claimed in claim 12 wherein each of X, Y, and Z is
25 CH.

14 A method as claimed in claim 13 wherein the ligand is a compound of the formula:



30

wherein:

M¹ is an alkali metal cation selected from: Li, Na, K, or Cs.

RL is a rigid linker group;

Ar¹ is an C₅₋₂₀aryl group;

5 n is an integer from 0 to 3; and,

each R^{BT} is independently a benzothiazole substituent.

15 A method as claimed in any one of claims 12 to 14 wherein each
of the rigid linker group, RL, and the aryl group, Ar¹, are
10 substantially planar.

16 A method as claimed in any one of claims 12 to 15 wherein the
rigid linker group, RL, and the aryl group, Ar¹, together with the
core group, form a compound which is substantially planar.

15 17 A method as claimed in any one of claims 12 to 16 wherein the
twist is no greater than that of the compound of Figure 16.

18 A method as claimed in any one of claims 12 to 17 wherein the
20 compound has a compound length which is from about 14.7 AU to about
15.3 AU.

19 A method as claimed in any one of claims 12 to 18 wherein each
R^{BT} is independently C₁₋₄alkyl, hydroxy, C₁₋₄alkoxy, nitro, cyano,
25 halo, or amino.


20 A method as claimed in claim 19 wherein each R^{BT} is
independently -Me, -Et, -nPr, -iPr, -OH, -OMe, -OEt, -O(nPr),
-O(iPr), -NO₂, -CN, -F, -Cl, -Br, -I, -NH₂, -NH₂, -NHMe, -NH₂Et,
30 -NH(iPr), -NH(nPr), -NMe₂, -NEt₂, N(iPr)₂, or -N(nPr)₂.

21 A method as claimed in claims 19 wherein each R^{BT} is
independently C₁₋₄alkyl.


35 22 A method as claimed in any one of claims 19 to 21 wherein n is
1, and R^{BT} is independently -Me, -Et, -nPr, or -iPr.

81


23 A method as claimed in any one of claims 12 to 22 wherein the ligand has the following formula:



5 24 A method as claimed in claim 23 wherein the ligand has the following formula:



25 A method as claimed in any one of claims 12 to 24 wherein RL is
10 a group of the formula:




wherein:

m is an integer from 0 to 4, and

each R^{RL} is independently a rigid linker aryl substituent,

15 and the ligand has the formula:



26 A method as claimed in claim 25 wherein each R^{RL} is
independently C₁₋₄alkyl, hydroxy, C₁₋₄alkoxy, nitro, cyano, halo, or
20 amino.


27 A method as claimed in claim 26 wherein each R^{RL} is
independently -Me, -Et, -nPr, -iPr, -OH, -OMe, -OEt, -O(nPr),

82


-O(iPr), -NO₂, -CN, -F, -Cl, -Br, -I, -NH₂, -NH₂, -NHMe, -NH₂Et, -NH(iPr), -NH(nPr), -NMe₂, -NEt₂, N(iPr)₂, or -N(nPr)₂.

28 A method as claimed in claim 26 wherein each R^{RL} is
5 independently C₁₋₄alkyl.

29 A method as claimed in any one of claims 1 to 24 wherein RL is a group of the formula:




10 30 A method as claimed in any one of claims 1 to 24 wherein RL is a group of the formula:



wherein

p is an integer from 0 to 3, and

15 each R^{RL} is independently a rigid linker aryl substituent, and the compounds have the formula:




31 A method as claimed in claim 30 wherein each R^{RL} is
independently C₁₋₄alkyl, hydroxy, C₁₋₄alkoxy, nitro, cyano, halo, or
20 amino.

32 A method as claimed in claim 31 wherein each R^{RL} is
independently -Me, -Et, -nPr, -iPr, -OH, -OMe, -OEt, -O(nPr), -
O(iPr), -NO₂, -CN, -F, -Cl, -Br, -I, -NH₂, -NH₂, -NHMe, -NH₂Et, -
25 NH(iPr), -NH(nPr), -NMe₂, -NEt₂, N(iPr)₂, or -N(nPr)₂.

33 A method as claimed in claim 31 wherein each R^{RL} is independently C_{1-4} alkyl.

34 A method as claimed in any one of claims 12 to 24 and 30 to 33
5 wherein RL is a group of the formula:



35 A method as claimed in any one of claims 12 to 34 wherein Ar^1 is selected from groups derived from benzene (C_6), naphthalene (C_{10}),
10 anthracene (C_{14}), phenanthrene (C_{14}), naphthacene (C_{18}), and pyrene (C_{16}).

36 A method as claimed in any one of claims 12 to 34 wherein Ar^1 is selected from:

15 C_5 heteroaryl groups derived from furan (oxole), thiophene (thiole), pyrrole (azole), imidazole (1,3-diazole), pyrazole (1,2-diazole), triazole, oxazole, isoxazole, thiazole, isothiazole, oxadiazole, and oxatriazole; and

C_6 heteroaryl groups derived from isoxazine, pyridine (azine),
20 pyridazine (1,2-diazine), pyrimidine (1,3-diazine), pyrazine (1,4-diazine), triazine, tetrazole, and oxadiazole (furazan).

37 A method as claimed in any one of claims 12 to 34 wherein Ar^1 is selected from:

25 C_9 heterocyclic groups derived from benzofuran, isobenzofuran, indole, isoindole, purine, benzimidazole;


C_{10} heterocyclic groups derived from quinoline, isoquinoline, benzodiazine, pyridopyridine, quinoxaline;

C_{13} heterocyclic groups derived from carbazole; and,

30 C_{14} heterocyclic groups derived from acridine, xanthene, phenoxathiin, phenazine, phenoxazine, phenothiazine.

84


38 A method as claimed in claim 35 wherein Ar^1 is an aryl group having a phenyl core, and has the formula:



wherein

- 5 q is an integer from 0 to 5; and
 each R^{A} is independently an aryl substituent;
 R^{C} , if present, is a reactive conjugating substituent, or
 R^{C} is, or contains, a detectable label;
 and the compound has the formula:

10



- 39 A method as claimed in claim 38 wherein R^{C} is present and is a reactive conjugating substituent, and is, or contains, a reactive functional group suitable for conjugation to another molecule by chemical reaction therewith, to form a covalent linkage therebetween.

- 40 A method as claimed in claim 39 wherein R^{C} is present and is, or contains, an active ester.

- 41 A method as claimed in claim 40 wherein R^{C} is present and is, or contains, a succinimidyl ester.

- 42 A method as claimed in claim 39 wherein R^{C} is present and is a reactive conjugating substituent, and is, or contains, a moiety suitable for conjugation to another molecule by a strong non-covalent interaction.

43 A method as claimed in claim 42 wherein R^C is present and is, or contains, biotin.

44 A method as claimed in claim 39 wherein R^C is present and is a reactive conjugating substituent, and is, or contains, a moiety suitable for conjugation to another molecule by complex or chelate formation.

45 A method as claimed in claim 44 wherein R^C is present and is, or contains, a technetium-chelating group.

46 A method as claimed in claim 45 wherein R^C is present and is, or contains diethylenetriaminepentaacetic acid.

47 A method as claimed in claim 38 wherein R^C is present and is, or contains, a detectable label.

48 A method as claimed in claim 47 wherein R^C is present and is, or contains, a dye, a fluorescent marker, an antigenic group, a stable or an unstable isotope, or a positron-emitting carbon atom.

49 A method as claimed in claim 48 wherein R^C is present and is, or contains, ¹⁸F.

50 A method as claimed in claim 48 wherein R^C is present and is, or contains, a positron-emitting carbon atom.

51 A method as claimed in any one of claims 38 to 50 wherein each R^A is independently selected from:

-OH, -NH₂, -NHR¹, -NR¹R², -SO₃M², and C₁₋₄alkyl;


wherein:

R¹ and R² are each C₁₋₄alkyl, and

M² is an alkali metal cation selected from Li, Na, K, or Cs.

52 A method as claimed in claim 51 wherein at least one R^A is -OH or -NH₂.

53 A method as claimed in claim 52 wherein Ar^1 is an aryl group having an amino-substituted phenyl core, and has the formula:



wherein

5 r is an integer from 0 to 4,
and each R^A is independently an aryl substituent.

54 A method as claimed claim 53 wherein each R^A is independently selected from:


10 $-\text{OH}$, $-\text{NH}_2$, $-\text{NHR}^1$, $-\text{NR}^1\text{R}^2$, $-\text{SO}_3\text{M}^2$, and $\text{C}_{1-4}\text{alkyl}$;

wherein:


R^1 and R^2 are each $\text{C}_{1-4}\text{alkyl}$, and

M^2 is an alkali metal cation selected from Li, Na, K, or Cs.

15 55 A method as claimed in claim 54 wherein the ligand has the formula:




56 A method as claimed in claim 55 wherein the ligand has the formula:



87

57 A method as claimed in claim 52 wherein Ar^1 is an aryl group having a hydroxy-substituted phenyl core, and has the formula:



5 wherein


s is an integer from 0 to 4, and
each R^A is independently an aryl substituent, and
 R^C , if present, is a reactive conjugating substituent, or
 R^C is, or contains, a detectable label.

10

58 A method as claimed in claim 57 wherein the ligand is as defined in any one of claims 108 to 150.

59 A method as claimed in claim 35 wherein Ar^1 is an aryl group having a naphthyl core, and has the formula:


15



wherein


t is an integer from 0 to 3,
 u is an integer from 0 to 4, and
each R^A is independently an aryl substituent,
and the compound has the formula:

20



88

60 A method as claimed in claim 59 wherein Ar^1 is an aryl group having a hydroxy-substituted naphthyl core, and has the formula:



5 wherein

v is an integer from 0 to 2,

u is an integer from 0 to 4, and

each R^A is independently an aryl substituent.

10 61 A method as claimed in claim 59 or claim 60 wherein each R^A is independently selected from:


$-\text{OH}$, $-\text{NH}_2$, $-\text{NHR}^1$, $-\text{NR}^1\text{R}^2$, $-\text{SO}_3\text{M}^2$, and $\text{C}_{1-4}\text{alkyl}$;

wherein:

R^1 and R^2 are each $\text{C}_{1-4}\text{alkyl}$, and


15 M^2 is an alkali metal cation selected from: Li, Na, K, or Cs.

62 A method as claimed in claim 61 wherein the ligand has the formula:







20

63 A method as claimed in claim 62 wherein the ligand has the formula:



64 A method as claimed in any one of claims 1 to 11 wherein the
5 ligand is a compound of one of the following formulae:



10

wherein:

each of R_1 , R_3 , R_4 , R_6 , R_7 and R_9 is independently hydrogen,

halogen, hydroxy, carboxy, substituted or unsubstituted alkyl, haloalkyl, or alkoxy;

R₅ is independently hydrogen, hydroxy, carboxy, substituted or unsubstituted alkyl, haloalkyl, or alkoxy;

5 R₁₀ and R₁₁ are independently selected from hydrogen, hydroxy, carboxy, substituted or unsubstituted alkyl, haloalkyl, or alkoxy;

or a pharmaceutically acceptable salt thereof.

10 65 A method as claimed in claim 64 wherein:

each of R₁, R₃, R₄, R₆, R₇ and R₉ is independently hydrogen, halogen, hydroxy, carboxy, substituted or unsubstituted C₁₋₆alkyl, C₁₋₄haloalkyl, or C₁₋₆alkoxy;

15 R₅ is independently hydrogen, hydroxy, carboxy, substituted or unsubstituted C₁₋₆alkyl, C₁₋₄haloalkyl, or C₁₋₆alkoxy;

R₁₀ and R₁₁ are independently selected from hydrogen, hydroxy, carboxy, substituted or unsubstituted C₁₋₆alkyl, C₁₋₄haloalkyl, or C₁₋₆alkoxy.

20 66 A method as claimed in claim 65 wherein said C₁₋₆alkyl is selected from: methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, tert-pentyl, hexyl, and isohexyl.

25 67 A method as claimed in claim 65 or claim 66 wherein the substituents of said substituted C₁₋₆alkyl are selected from: mercapto, thioether, nitro, amino, aryloxy, halogen, hydroxyl, carbonyl, C₅₋₂₀aryl, C₁₋₆cycloalkyl, and non-aryl C₃₋₂₀heterocyclyl.

30 68 A method as claimed in any one of claims 65 to 67 wherein said C₁₋₄haloalkyl is selected from: chloromethyl, 2-bromomethyl, 1-chloroisopropyl, 3-fluoropropyl, 2,3-dibrombutyl, 3-chloroisobutyl, iodo-t-butyl, and trifluoromethyl.

35 69 A method as claimed in any one of claims 64 to 68 wherein the ligand is an acid addition salt formed between a compound described in said claims and an acid.

70 A method as claimed in claim 69 wherein the acid is an inorganic acid or an organic acid.

5 71 A method as claimed in claim 70 wherein the ligand is a chloride salt.

72 A method as claimed in claim 70 wherein said organic acid is selected from: acetic acid, citric acid, maleic acid, fumaric acid, tartaric acid, methanesulphonic acid, and p-toluenesulphonic acid.

10

73 A method as claimed in any one of claims 64 to 72 wherein:

R_1 , R_3 , R_4 , R_6 , R_7 and R_9 are independently -H, -CH₃, -C₂H₅, or -C₃H₇;

R_{10} and R_{11} are independently -H, -CH₃, -C₂H₅ or -C₃H₇;

15 R_5 is independently -H, -CH₃, -C₂H₅, or -C₃H₇.

74 A method as claimed in claim 73 wherein the ligand is shown in Figure 8b.

20 75 A method as claimed in any one of claims 64 to 74 wherein the ligand comprises a positron-emitting carbon.

76 A method as claimed in any one of the preceding claims wherein (ii) further comprises the step of additionally determining the presence and/or amount of a ligand bound to intracellular aggregated tau in a neocortical structure of the brain of the subject.

25

77 A method as claimed in claim 76 wherein the ligand used to bind to extracellular aggregated PHF tau in the medial temporal lobe and the ligand used to bind to intracellular aggregated PHF tau in the neocortical structure of the brain are labelled distinctively.

30

78 A method as claimed in claim 76 or claim 77 wherein the ligand used to bind intracellular aggregated tau is the ligand described in any one of claims 64 to 75.

35


79 A method as claimed in any one of the preceding claims wherein

the ligand binds aggregated PHF tau preferentially with respect to competing binding sites present in the relevant region of the brain.

80 A method as claimed in any one of the preceding claims wherein steps (i) and/or (ii) of the method are performed in conjunction with the further step of introducing into the subject a further blocking ligand which labels the competing binding sites present in the relevant region of the brain preferentially to the ligand used to bind aggregated PHF tau.

81 A method as claimed in claim 80 wherein the blocking ligand is [^{18}F]FDDNP.

82 A method as claimed in claim 80 wherein the blocking ligand is a benzthiazole of the formula:



wherein:

n is an integer from 0 to 4;

each R^{BT} is independently a blocking ligand benzothiazole substituent;

m is an integer from 0 to 4;

each R^{P} is independently a phenylene substituent;

each R is independently -H or an amino substituent; and,

either:

R^{N} and X^- are both absent and the associated (tertiary) nitrogen atom is neutral;

or:

R^{N} is a benzothiazolino substituent and the associated (quaternary) nitrogen atom bears a positive charge, and X^- is a counter ion.

83 A method as claimed in claim 82 wherein the blocking ligand is thioflavin-T.

84 A method as claimed in claim 82 wherein each R^{BT} , is independently C_{1-4} alkyl, $-SO_3H$, or $-SO_3M^3$, wherein M^3 is a cation.

85 A method as claimed in claim 84 wherein M^3 is an alkali metal cation selected from: Li, Na, K, or Cs.

86 A method as claimed in any one of claims 82 to 85 wherein n is 1, and R^{BT} is -Me, -Et, -nPr, or -iPr.

87 A method as claimed in claim 86 wherein n is 1, and R^{BT} is -Me.

88 A method as claimed in claim 84 wherein one of the R^{BT} groups is $-SO_3H$ or $-SO_3M^3$, and another of the R^{BT} groups is C_{1-4} alkyl.

89 A method as claimed in any one of claims 82 to 85 wherein n is 2, and one R^{BT} is C_{1-4} alkyl, and one R^{BT} is $-SO_3H$ or $-SO_3M^3$.

90 A method as claimed in claim 89 wherein n is 2, and one R^{BT} is -Me, and one R^{BT} is $-SO_3H$ or $-SO_3M^3$.

91 A method as claimed in any one of claims 82 to 90 wherein R^N and X^- are both absent and the associated (tertiary) nitrogen atom is neutral.

92 A method as claimed in any one of claims 82 to 90 wherein R^N is a benzothiazolino substituent and the associated (quaternary) nitrogen atom bears a positive charge, and X^- is a counter ion.

93 A method as claimed in any one of claims 82 to 90 wherein R^N is C_{1-4} alkyl.

94 A method as claimed in claim 93 wherein R^N is -Me, -Et, -nPr, or -iPr.

95 A method as claimed in any one of claims 82 to 90 wherein X^- is Cl^- , Br^- , and I^- .

96 A method as claimed in any one of claims 82 to 95 wherein R^P is

C₁₋₄alkyl.

97 A method as claimed in any one of claims 82 to 96 wherein each R is -H, and the amino group is -NH₂.

5

98 A method as claimed in any one of claims 82 to 96 wherein one R is -H and one R is an amino substituent.

99 A method as claimed in any one of claims 82 to 96 wherein each R is an amino substituent.

10


100 A method as claimed in claim 99 wherein each of the amino substituents is C₁₋₄alkyl.

101 A method as claimed in any one of claims 82 to 96 wherein the amino group, -NR₂, is -NH₂, -NHMe, -NH₂Et, -NH(iPr), -NH(nPr), -NMe₂, -NEt₂, N(iPr)₂, or -N(nPr)₂.

15


102 A method as claimed in claim 101 wherein the blocking ligand is a benzthiazole of the formula:

20



103 A method as claimed in claim 101 wherein the blocking ligand is a benzthiazole of the formula:

25



104 A method as claimed in any one of the preceding claims for use in the diagnosis or prognosis of a tauopathy in a subject believed to suffer from said disease.


30

105 A method as claimed in claim 104 wherein tauopathy is AD.

106 A ligand as described in any one of claims 1 to 79 for use
in a method of diagnosis or prognosis of a tauopathy in a subject
believed to suffer from said disease, said method being a method as
5 described in claim 104 or claim 105.

107 A ligand as described in any one of claims 1 to 79 for use
in a method for determinining the effectiveness of a treatment of a
subject with a tau-tau aggregation inhibitor to inhibit
10 neurofibrillary degeneration in that subject, the method comprising
use of a method as described in any one of claims 1 to 103.

108 A ligand of the formula:



15 wherein:

M^1 is an alkali metal cation;

n is an integer from 0 to 3;

each R^{BT} is a independently benzothiazole substituent;

m is an integer from 0 to 4;

20 each R^{RL} is independently a rigid linker aryl substituent;

s is an integer from 0 to 4;

each R^A is independently an aryl substituent; and,

R^C , if present, is a reactive conjugating substituent, or
25 R^C is, or contains, a detectable label.

109 A ligand as claimed in claim 108 wherein M^1 is Li, Na, K, or
Cs.

110 A ligand as claimed in claim 108 or claim 109 wherein each
30 R^{BT} is independently C_{1-4} alkyl, hydroxy, C_{1-4} alkoxy, nitro, cyano,
halo, or amino.

111 A ligand as claimed in claim 110 wherein each R^{BT} is independently -Me, -Et, -nPr, -iPr, -OH, -OMe, -OEt, -O(nPr), -O(iPr), -NO₂, -CN, -F, -Cl, -Br, -I, -NH₂, -NHMe, -NH₂Et, -NH(iPr), -NH(nPr), -NMe₂, -NEt₂, N(iPr)₂, or -N(nPr)₂.

5

112 A ligand as claimed in claim 110 wherein each R^{BT} is independently C₁₋₄alkyl.

113 A ligand as claimed in claim 112 wherein each R^{BT} is independently -Me, -Et, -nPr, or -iPr.

10

114 A ligand as claimed in claim 113 wherein each R^{BT} is -Me.


115 A ligand as claimed in claim 113 wherein n is 1, and R^{BT} is independently -Me, -Et, -nPr, or -iPr.

15

116 A ligand as claimed in claim 115 wherein n is 1, and R^{BT} is -Me.


117 A ligand as claimed in any one of claims 108 to 116 wherein the benzothiazole group has the following formula:

20



118 A ligand as claimed in claim 117 wherein the ligand has the following formula:

25



119 A ligand as claimed in any one of claims 108 to 118 wherein each R^{RL} is independently C₁₋₄alkyl, hydroxy, C₁₋₄alkoxy, nitro, cyano, halo, or amino.

30

120 A ligand as claimed in claim 119 wherein each R^{RL} is independently -Me, -Et, -nPr, -iPr, -OH, -OMe, -OEt, -O(nPr), -O(iPr), -NO₂, -CN, -F, -Cl, -Br, -I, -NH₂, -NH₂, -NHMe, -NH₂Et, -NH(iPr), -NH(nPr), -NMe₂, -NEt₂, N(iPr)₂, or -N(nPr)₂.

121 A ligand as claimed in claim 120 wherein each R^{RL} is independently C₁₋₄alkyl.

122 A ligand as claimed in any one of claims 108 to 121 wherein each R^A is independently selected from:

-OH, -NH₂, -NHR¹, -NR¹R², -SO₃M², and C₁₋₄alkyl;

wherein:

R¹ and R² are each C₁₋₄alkyl, and

M² is an alkali metal cation.

123 A ligand as claimed in claim 122 wherein M² is Li, Na, K, or Cs.

124 A ligand as claimed in any one of claims 108 to 123 wherein R^C is present and is a reactive conjugating substituent, and is, or contains, a reactive functional group suitable for conjugation to another molecule by chemical reaction therewith, to form a covalent linkage therebetween.

125 A ligand as claimed in claim 124 wherein R^C is present and is, or contains, an active ester.

126 A ligand as claimed in claim 125 wherein R^C is present and is, or contains, a succinimidyl ester.

127 A ligand as claimed in claim 124 wherein R^C is present and is a reactive conjugating substituent, and is, or contains, a moiety suitable for conjugation to another molecule by a strong non-covalent interaction.

128 A ligand as claimed in claim 127 wherein R^C is present and is, or contains, biotin.

129 A ligand as claimed in claim 124 wherein R^C is present and is
a reactive conjugating substituent, and is, or contains, a moiety
suitable for conjugation to another molecule by complex or chelate
5 formation.

130 A ligand as claimed in claim 129 wherein R^C is present and
is, or contains, a technetium-chelating group.

10 131 A ligand as claimed in claim 130 wherein R^C is present and
is, or contains diethylenetriaminepentaacetic acid.

132 A ligand as claimed in any one of claims 108 to 123 wherein
 R^C is present and is, or contains, a detectable label.

15


133 A ligand as claimed in claim 132 wherein R^C is present and
is, or contains, a dye, a fluorescent marker, an antigenic group, a
stable or an unstable isotope, or a positron-emitting carbon atom.

20 134 A ligand as claimed in claim 133 wherein R^C is present and
is, or contains, ^{18}F .

135 A ligand as claimed in claim 134 wherein R^C is present and
is, or contains, a positron-emitting carbon atom.


25

136 A ligand as claimed in any one of claims 108 to 135 wherein
the ligand has the formula:




99


137 A ligand as claimed in claim 136 wherein the ligand has the formula:




5 138 A ligand as claimed in claim 137 wherein the ligand has the formula:



10 139 A ligand as claimed in claim 137 wherein the ligand has the formula:




140 A ligand as claimed in claim 139 wherein the ligand has the formula:




100


141 A ligand as claimed in claim 140 wherein the ligand has the formula:




5 142 A ligand as claimed in claim 137 wherein the ligand has the formula:



10 143 A ligand as claimed in claim 142 wherein the ligand has the formula:




144 A ligand as claimed in claim 143 wherein the ligand has the formula:




15

101


145 A ligand as claimed in claim 142 wherein the ligand has the formula:




5 146 A ligand as claimed in claim 145 wherein the ligand has the formula:



10 147 A ligand as claimed in claim 145 wherein the ligand has the formula:




148 A ligand as claimed in any one of claim 147 wherein the ligand has the formula:



102

149 A ligand as claimed in claim 148 wherein the ligand has the formula:



5 150 A ligand as claimed in any one of claims 136 to 149 which is conjugated, chelated, or otherwise associated, with a detectable chemical group.

151 A diagnostic composition comprising as its active ingredient
10 at least 90%, 95%, or 98% of the ligand of any one of claims 108 to 150.

152 A diagnostic composition as claimed in claim 151 further
comprising a carrier material or other pharmaceutically- and
15 physiologically-acceptable excipient.

153 A method of labelling aggregated tau or tau-like molecules,
comprising contacting the aggregated molecules with a ligand or
composition of any one of claims 108 to 152.

20

154 An *in vitro* method of diagnosis or prognosis of a tauopathy
in a subject believed to suffer from the disease, the method
comprising (i) obtaining a sample of appropriate tissue from a
subject; (ii) contacting the sample with a ligand or composition of
25 any one of claims 108 to 152; (iii) detecting the amount and/or
localisation of the ligand bound to the sample (iv) correlating the
result of (iii) with the stage or severity of the disease in the
subject.

30 155 A method as claimed in claim 153 where the tauopathy is AD.

156 A ligand according to any one of claims 108 to 150 for use
in a method of diagnosis, prognosis or therapeutic treatment of the

human or animal body for a tauopathy.

157 A ligand as claimed in claim 156 where the tauopathy is AD.

5 158 A method for identifying a compound capable of binding aggregated tau protein, the method comprising the steps:

(i) providing a sample of aggregated tau protein to which a ligand as claimed in any one of claims 108 to 150 has been bound,

(ii) contacting the sample with the putative tau-binding compound,

10 (iii) determining the extent of displacement of the ligand from the aggregated tau protein by the putative tau-binding compound

(iv) correlating the result of the determination made in (iii) with the ability of the compound to bind aggregated tau protein.

15 159 A method as claimed in claim 158 wherein the aggregated tau protein is in solution or bound to a solid phase.

160 An *in vitro* method for identifying a ligand capable of labeling aggregated PHF tau protein, the method comprising the steps of:

20 (i) providing a first agent suspected of being capable of labeling aggregated PHF tau protein,

(ii) contacting (a) a tau protein or a derivative thereof containing the tau core fragment bound to a solid phase so as to

25 expose a high affinity tau capture site, with (b) a liquid phase tau protein or derivative thereof capable of binding to the solid phase tau protein or derivative, and (c) said selected first agent and (d) a second agent known to be tau-tau binding inhibitor,

30 (iii) selecting first agent which fully or partially relieves the inhibition of binding of the liquid phase tau protein or derivative of (b) to the solid phase tau protein or derivative of (a) by the inhibitor (d).

161 A method as claimed in claim 160 wherein step (ii) is carried out in conjunction with:

35 (i)bis) contacting (a) a tau protein or a derivative thereof containing the tau core fragment bound to a solid phase so as to expose a high affinity tau capture site, with (b) a liquid phase

tau protein or derivative thereof capable of binding to the solid phase tau protein or derivative, and (c) said first agent, and (ibis.1) detecting inhibition of tau-tau binding as exhibited by inhibition of binding of the liquid phase tau protein or derivative of (b) to the solid phase tau protein or derivative of (a), (ibis.2) selecting first agent which has minimal or absent activity as tau-tau binding inhibitors and/or optionally enhance tau-tau binding.

10 162 A method as claimed in claim 160 or claim 161 wherein the inhibitor is a diaminophenathiozine as described in any one of claims 64 to 74.

15 163 A method as claimed in claim 162 wherein the inhibitor is DMMB.

164 A method as claimed in any one of claims 160 to 163 wherein the liquid phase tau protein or derivative is prepared in a form which has undergone partial aggregation prior to exposure to the solid phase.

165 A method as claimed in claim 174 wherein said preparation of liquid phase tau protein or derivative comprises the steps of:
(i) sonicating said tau protein or derivative and/or
25 (ii) exposing said tau protein or derivative to PEG through a semi-permeable membrane.

166 A method as claimed in any one of claims 160 to 165, wherein the step of contacting said agent and liquid phase tau protein or derivative with said solid phase tau protein or derivative is carried out in an alkaline buffer of pH 9-10.

167 A method as claimed in any one of claims 160 to 165, wherein the step of contacting said agent and liquid phase tau protein or derivative with said solid phase tau protein or derivative is carried out in physiological buffer conditions.

168 A method as claimed in any one of claims 160 to 165, wherein
the step of contacting the or each agent and liquid phase tau
protein or derivative with said solid phase tau protein or
derivative is carried out in 50-400 mM sodium chloride or a salt or
5 salt mixture of comparable ionic strength, and in a pH range of 4-
10.

169 A method as claimed in any one of claims 160 to 168, wherein
said liquid phase tau protein or derivative is labeled.

170 A method as claimed in any one of claims 160 to 168 wherein
said liquid phase tau protein or derivative is immunologically
distinguishable from said solid phase tau protein or derivative,
and the binding is detected by antibodies.

171 A method as claimed in any one of claims 160 to 170 wherein
a truncated tau protein corresponding to the core fragment and
terminating at Ala390 (dGA) is plated on a solid phase in buffer
conditions unfavourable to tau-tau association,

172 A method as claimed in any one of claims 160 to 171 wherein
a truncated tau protein corresponding to the core fragment and
terminating at Glue-391 (dGAE) or a full-length protein is added in
liquid phase together with the or each agent

173 A method as claimed in any one of claims 158 to 172 further
comprising the step of formulating the agent as a diagnostic or
prognostic reagent.

174 A method as claimed in any one of claims 158 to 172 further
comprising the step of using the agent as a ligand in the method of
any one of claims 1 to 11.




Figure 1

2/54




Figure 2a

3/54




Figure 2b

4/54




Figure 2c




Figure 2d




Figure 3

7/54




Figure 4

8/54




Figure 5

9/54




Figure 5 cont ...




Figure 6

11/54




Figure 7

12/54







Figure 8a

13/54

Diamino phenothiazines with varying methyl groups





Figure 8b

14/54



(ACB = Alcian Blue, c14537 = primulin)

Figure 9

Tau-tau binding vs Molar Ratio (compound:tau)
Thionine vs PHF-ligands




Figure 10

16/54


Synthesis of [^{11}C] labelled Methylene Blue

Figure 11a

17/54




Figure 11b

18/54


Synthesis of [^{11}C] labelled Azure B

Figure 12

19/54



Synthesis of [^{18}F] labelled


Figure 13

20/54

Primulin



Benzothiazole analogue



Thiazin yellow




Figure 14

21/54




Figure 15

22/54




Figure 16

23/54




Figure 17




Figure 18

25/54




Figure 19




Figure 20

27/54




Figure 21

28/54




Figure 22

29/54

Comparison of dimensions of active and
inactive molecules tested
(maximum, mean, minimum)




Figure 23

30/54

Comparison of dimensions of
benzothiazole and diaminophenothiazine molecules
(maximum, mean, minimum)




Figure 24

31/54

PROBABILITY OF 423-T'S BY REGION AND STAGE




Figure 25

32/54

PROBABILITY AT8-T'S BY REGION AND STAGE




Figure 26

33/54

PROBABILITY OF 423-T'S BY REGION AND STAGE
MMSE > 21




Figure 27

34/54

PROBABILITY OF AT8-T'S BY REGION AND STAGE MMSE > 21




Figure 28

35/54

DENSITY OF 423-T'S BY REGION AND STAGE




Figure 29

36/54

DENSITY OF AT8-T'S BY REGION AND STAGE




Figure 30

37/54




Figure 31a

38/54



Figure 31b

39/54




Figure 32

40/54




Figure 33

41/54




Figure 34

42/54





Figure 35

43/54

Preparation 1

dGA 5 mcg/ml, dGAE 5 mcg/ml



Preparation 2

dGA 4 mcg/ml, dGAE 5 mcg/ml





Figure 36

44/54

Preparation 1

dGA 5 mcg/ml, dGAE 5 mcg/ml



Preparation 2

dGA 10 mcg/ml, dGAE 10 mcg/ml





Figure 37

45/54

Preparation 1



Preparation 2




Figure 38




Figure 39a and 39b

C




Figure 39c

48/54






Figure 40

49/54

(a)

Tau-Tau Binding at 5 mM DMMB



(b)

Tau-Tau Binding at 5 mM DMMB





Figure 41a and 41b

50/54

(c)

Tau-Tau Binding at 15 mM DMMB



(d)

Tau-Tau Binding at 15 mM DMMB




Figure 41c and 41d




Figure 42

52/54




Figure 43

53/54




Figure 44

54/54




Figure 45

专利名称(译)	神经原纤维标签		
公开(公告)号	EP1421389A2	公开(公告)日	2004-05-26
申请号	EP2002720102	申请日	2002-03-20
申请(专利权)人(译)	大学评议阿伯丁大学		
当前申请(专利权)人(译)	WISTA LABORATORIES LTD.		
发明人	WISCHIK, CLAUDE, MICHEL,DEP. OF MENTAL HEALTH HARRINGTON, CHARLES, ROBERT,DEP. OF MENTAL HEALTH RICKARD, JANET, ELIZABETH,DEP. OF MENTAL HEALTH HORSLEY, DAVID,DEP. OF MENTAL HEALTH		
IPC分类号	A61K49/00 C07D277/66 C07D279/20 C07D417/12 G01N33/53 G01N33/58 G01N33/68 C07D277/62 C07D279/18		
CPC分类号	C07D417/12 C07D277/66 C07D279/20 G01N33/583 G01N33/6896		
优先权	2001006953 2001-03-20 GB		
其他公开文献	EP1421389B1		
外部链接	Espacenet		

摘要(译)

公开了用于确定与被认为患有该疾病的受试者中的tau蛋白病相关的神经原纤维变性阶段的方法，该方法包括以下步骤：(i) 向受试者中引入能够标记聚集的成对螺旋长丝(PHF)的配体tau蛋白，(ii) 确定与受试者大脑内侧颞叶中细胞外聚集的PHF tau结合的配体的存在和/或量，(iii) 将(ii)中所做的测定结果与所述程度相关联。在受试者中的神经原纤维变性。该方法可用于早产儿诊断和分期治疗阿尔茨海默病等tau蛋白病。优选的配体包括碘化-苯并噻唑类化合物和二氨基吩噻嗪。还提供了新配体(例如碘化-苯并噻唑类化合物)。该方法还可包括使用“阻断配体”来阻断竞争绑定网站。在其他方面，本发明提供了用于鉴定能够标记聚集的PHF tau蛋白的配体的体外方法，该方法包括以下步骤：(i) 提供怀疑能够标记聚集的PHF tau蛋白的第一试剂，(ii) 接触(a)含有与固相结合tau核心片段的tau蛋白质或其衍生物，以暴露高亲和力tau捕获位点，具有(b)能够结合固相tau的液相tau蛋白质或其衍生物蛋白质或衍生物，和(c)所述选定的第一试剂和(d)已知为tau-tau结合抑制剂的第二试剂，(iii) 选择完全或部分缓解液相tau蛋白结合抑制的第一试剂或(b)的衍生物通过抑制剂(d)得到固相tau蛋白或(a)的衍生物。还可以测试配体以确认它们本身不是抑制剂。