



- (51) **International Patent Classification:**
G01N33/53 (2006.01)
- (21) **International Application Number:**
PCT/US2014/056653
- (22) **International Filing Date:**
19 September 2014 (19.09.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/880,765 20 September 2013 (20.09.2013) US
- (71) **Applicant:** ASTUTE MEDICAL, INC. [US/US]; Big 2, R. 645, 3550 General Atomics Court, San Diego, CA 92121 (US).
- (72) **Inventors:** ANDERBERG, Joseph; 470 Delage Court, Encinitas, CA 92024 (US). GRAY, Jeff; 417 Bay Meadows Way, Solana Beach, CA 92075 (US). McPHERSON, Paul; 1449 Elva Court, Encinitas, CA 92024 (US). NAKAMURA, Kevin; 2409 Newcastle Ave, Cardiff by the Sea, CA 92007 (US). KAMPF, James, Patrick; 5882 Gablewood Way, San Diego, CA 92130 (US). KWAN, Thomas; 3950 Mahaila Ave #v23, San Diego, CA 92122 (US).
- (74) **Agents:** WHITTAKER, Michael, A. et al; Acuity Law Group, PC, 12707 High Bluff Drive, Suite 200, San Diego, CA 92130 (US).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report (Art. 21(3))



WO 2015/042465 A1

(54) **Title:** METHODS AND COMPOSITIONS FOR DIAGNOSIS AND PROGNOSIS OF APPENDICITIS AND DIFFERENTIATION OF CAUSES OF ABDOMINAL PAIN

(57) **Abstract:** The present invention relates to methods and compositions for monitoring, diagnosis, prognosis, and determination of treatment regimens in appendicitis patients and in patients at risk for appendicitis. In particular, the invention relates to using assays that detect one or more biomarkers as diagnostic and prognostic biomarker assays in such patients.

**METHODS AND COMPOSITIONS FOR DIAGNOSIS AND PROGNOSIS OF
APPENDICITIS AND DIFFERENTIATION OF CAUSES OF ABDOMINAL PAIN**

[0001] The present application claims priority from U.S. Provisional Patent Application 61/880,765 filed September 20, 2013, which is hereby incorporated in its entirety including all tables, figures and claims.

BACKGROUND OF THE INVENTION

[0002] The following discussion of the background of the invention is merely provided to aid the reader in understanding the invention and is not admitted to describe or constitute prior art to the present invention.

[0003] Acute appendicitis is an inflammatory condition which typically results from a primary obstruction of the appendix lumen. Once obstructed, the appendix subsequently swells, increasing pressures within the lumen and the walls of the appendix, resulting in thrombosis and occlusion of the small vessels, and stasis of lymphatic flow. The causative agents of appendicitis include foreign bodies, trauma, intestinal worms, lymphadenitis, and, most commonly, calcified fecal deposits known as appendicoliths or fecaliths. Diagnosis is based on patient history, symptoms and physical examination. Typical appendicitis usually includes abdominal pain beginning in the region of the umbilicus for several hours, associated with anorexia, nausea or vomiting. The pain typically settles into the right lower quadrant.

[0004] A commonly used acronym for diagnosis is PALF: pain, anorexia, leukocytosis, and fever. Atypical histories lack this typical progression and may include pain in the right lower quadrant as an initial symptom. Atypical histories often require imaging with ultrasound and/or CT scanning. Blood tests for appendicitis are normal, and so not diagnostic, in about 50% of cases. These tests tend to be relatively simple. An abnormal rise in the number of white blood cells in the blood is a crude indicator of infection or inflammation going on in the body. Such a rise is not specific to appendicitis alone. If it is abnormally elevated, with a good history and examination findings pointing towards appendicitis, the likelihood of having the disease is higher. Imaging tests such as CT, while useful, expose the recipient to diagnostic levels of radiation.

[0005] In terms of biomarkers, C-reactive protein (CRP), an acute-phase response protein produced by the liver in response to inflammatory processes, as been used by clinicians. Likewise, other general inflammatory markers such as procalcitonin, Interleukin-6 (IL-6), Interleukin-8 (IL-8), high mobility group box-1 protein (HMGB1) S100A8/A9, etc., have been studied by clinicians. Like the number of white blood cells, however, these are not specific biomarkers of appendicitis, and so exhibit poor specificity in use. Leucine-rich alpha-2-glycoprotein (LRG) was recently suggested to be a more specific indicator of appendicitis in a pediatric population. Kentsis et al., Ann. Emerg. Med. 55: 62-70.e4. Epub 2009 Jun 25; Kharbanda et al., Academic Emerg. Med. 19: 56-62, 2012.

[0006] There remains a need in the art for a rapid, objective, clinically accurate, available diagnostic tool for aiding in the diagnosis and care of appendicitis.

BRIEF SUMMARY OF THE INVENTION

[0007] The present invention relates to the identification and use of diagnostic markers for appendicitis. The methods and compositions described herein can meet the need in the art for rapid, sensitive and specific diagnostic assay to be used in the diagnosis and differentiation of abdominal pain and the identification of appendicitis. In various aspects, the invention relates to materials and procedures for identifying markers that are associated with the diagnosis, prognosis, or differentiation of appendicitis in a patient; to using such markers in diagnosing and treating a patient and/or to monitor the course of a treatment regimen; to using such markers to identify subjects at risk for one or more adverse outcomes related to appendicitis; and for screening compounds and pharmaceutical compositions that might provide a benefit in treating or preventing such conditions.

[0008] In a first aspect, the invention discloses methods for determining a diagnosis or prognosis related to appendicitis, or for differentiating between causes of abdominal pain. As described herein, measurement of one or more biomarkers selected from the group consisting of 72 kDa type IV collagenase, Adiponectin, Advanced glycosylation end product-specific receptor, Alpha-2 macroglobulin, Alpha-2-HS-glycoprotein, Alpha-fetoprotein, Angiopoietin-1, Antileukoprotease, Apolipoprotein(a), Cancer Antigen 15-3, Cancer Antigen 19-9, Carbonic anhydrase 9, Carcinoembryonic antigen-related cell adhesion molecule 5, C-C motif chemokine 1, C-C motif chemokine 13, C-C motif

chemokine 15, C-C motif chemokine 17, C-C motif chemokine 19, C-C motif chemokine 20, C-C motif chemokine 21, C-C motif chemokine 22, C-C motif chemokine 23, C-C motif chemokine 24, C-C motif chemokine 26, C-C motif chemokine 27, C-C motif chemokine 3, C-C motif chemokine 4, C-C motif chemokine 7, C-C motif chemokine 8, Ceruloplasmin, Choriogonadotropin subunit beta, Collagenase 3, C-Peptide, Creatine Kinase-MB, C-X-C motif chemokine 10, C-X-C motif chemokine 11, C-X-C motif chemokine 13, C-X-C motif chemokine 16, C-X-C motif chemokine 5, C-X-C motif chemokine 6, C-X-C motif chemokine 9, Cystatin-C, Endothelial protein C receptor, Eotaxin, Epidermal growth factor receptor, Ferritin, Fibrinogen, Gastric inhibitory polypeptide, Glucagon, Glucagon-like peptide 1, Granulocyte-macrophage colony-stimulating factor, Growth-regulated alpha, beta, and gamma proteins, Heparin-binding growth factor 2, Hepatocyte growth factor, Immunoglobulin A, Immunoglobulin M, Immunoglobulin G1, Immunoglobulin G2, Immunoglobulin G3, Immunoglobulin G4, Insulin, Insulin-like growth factor-binding protein 1, Insulin-like growth factor-binding protein 2, Insulin-like growth factor-binding protein 3, Insulin-like growth factor-binding protein 4, Insulin-like growth factor-binding protein 5, Insulin-like growth factor-binding protein 6, Insulin-like growth factor-binding protein 7, Interferon alpha-2, Interferon gamma, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 receptor antagonist protein, Interleukin-1 receptor type I, Interleukin-1 receptor type II, Interleukin-1 1, Interleukin-12, Interleukin-12 subunit beta, Interleukin-13, Interleukin-15, Interleukin-2, Interleukin-20, Interleukin-21, Interleukin-23, Interleukin-28A, Interleukin-29, Interleukin-3, Interleukin-33, Interleukin-4, Interleukin-5, Interleukin-6 receptor subunit alpha, Interleukin-6 receptor subunit beta, Interleukin-7, Interleukin-9, Interstitial collagenase, Islet amyloid polypeptide, Keratin, type I cytoskeletal 19 (aa3 11-367), Kit ligand, Leptin, Leukemia inhibitory factor, Lymphotoxin, Lymphotoxin-alpha, Macrophage colony-stimulating factor 1, Macrophage metalloelastase, Macrophage migration inhibitory factor, Matrilysin, Metalloproteinase inhibitor 1, Metalloproteinase inhibitor 2, Metalloproteinase inhibitor 3, Metalloproteinase inhibitor 4, Myoglobin, Pancreatic prohormone, Peptide YY, Pro-epidermal growth factor, Pro-interleukin-16, Prolactin, Prostate-specific antigen, Protein S100-A12, Protransforming growth factor alpha, Secretory immunoglobulin A, Serum amyloid P-component, SL cytokine, Stromal cell-derived factor 1, Stromelysin-1, Thrombopoietin, Thymic stromal lymphopoietin, Tumor necrosis factor, Tumor necrosis factor ligand superfamily member 10, Tumor necrosis factor ligand superfamily member 6, Tumor necrosis factor receptor superfamily member

1A, Tumor necrosis factor receptor superfamily member 1B, Vascular endothelial growth factor receptor 1, Vascular endothelial growth factor receptor 2, Vascular endothelial growth factor receptor 3, von Willebrand Factor, and WAP four-disulfide core domain protein 2 (each referred to herein for convenience as an "appendicitis biomarker") can be used for diagnosis, prognosis, risk stratification, monitoring, categorizing and determination of further diagnosis and treatment regimens in patients having or suspected of having appendicitis. The appendicitis biomarkers of the present invention may be used, individually or in panels comprising a plurality of appendicitis biomarkers. The presence or amount of such marker(s) in a sample obtained from the subject can be used to rule in or rule out appendicitis, and to monitor subjects for improving or worsening conditions related to appendicitis.

[0009] In a first aspect, the present invention relates to methods for evaluating an appendicitis patient or a patient being evaluated for a possible diagnosis. These methods comprise performing an assay method that is configured to detect one or more biomarkers selected from the group consisting of 72 kDa type IV collagenase, Adiponectin, Advanced glycosylation end product-specific receptor, Alpha-2 macroglobulin, Alpha-2-HS-glycoprotein, Alpha-fetoprotein, Angiopoietin-1, Antileukoproteinase, Apolipoprotein(a), Cancer Antigen 15-3, Cancer Antigen 19-9, Carbonic anhydrase 9, Carcinoembryonic antigen-related cell adhesion molecule 5, C-C motif chemokine 1, C-C motif chemokine 13, C-C motif chemokine 15, C-C motif chemokine 17, C-C motif chemokine 19, C-C motif chemokine 20, C-C motif chemokine 21, C-C motif chemokine 22, C-C motif chemokine 23, C-C motif chemokine 24, C-C motif chemokine 26, C-C motif chemokine 27, C-C motif chemokine 3, C-C motif chemokine 4, C-C motif chemokine 7, C-C motif chemokine 8, Ceruloplasmin, Choriogonadotropin subunit beta, Collagenase 3, C-Peptide, Creatine Kinase-MB, C-X-C motif chemokine 10, C-X-C motif chemokine 11, C-X-C motif chemokine 13, C-X-C motif chemokine 16, C-X-C motif chemokine 5, C-X-C motif chemokine 6, C-X-C motif chemokine 9, Cystatin-C, Endothelial protein C receptor, Eotaxin, Epidermal growth factor receptor, Ferritin, Fibrinogen, Gastric inhibitory polypeptide, Glucagon, Glucagon-like peptide 1, Granulocyte-macrophage colony-stimulating factor, Growth-regulated alpha, beta, and gamma proteins, Heparin-binding growth factor 2, Hepatocyte growth factor, Immunoglobulin A, Immunoglobulin M, Immunoglobulin G1, Immunoglobulin G2, Immunoglobulin G3, Immunoglobulin G4, Insulin, Insulin-like growth factor-binding

protein 1, Insulin-like growth factor-binding protein 2, Insulin-like growth factor-binding protein 3, Insulin-like growth factor-binding protein 4, Insulin-like growth factor-binding protein 5, Insulin-like growth factor-binding protein 6, Insulin-like growth factor-binding protein 7, Interferon alpha-2, Interferon gamma, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 receptor antagonist protein, Interleukin-1 receptor type I, Interleukin-1 receptor type II, Interleukin-11, Interleukin-12, Interleukin-12 subunit beta, Interleukin-13, Interleukin-15, Interleukin-2, Interleukin-20, Interleukin-21, Interleukin-23, Interleukin-28A, Interleukin-29, Interleukin-3, Interleukin-33, Interleukin-4, Interleukin-5, Interleukin-6 receptor subunit alpha, Interleukin-6 receptor subunit beta, Interleukin-7, Interleukin-9, Interstitial collagenase, Islet amyloid polypeptide, Keratin, type I cytoskeletal 19 (aa311-367), Kit ligand, Leptin, Leukemia inhibitory factor, Lymphotoxin, Lymphotoxin-alpha, Macrophage colony-stimulating factor 1, Macrophage metalloelastase, Macrophage migration inhibitory factor, Matrilysin, Metalloproteinase inhibitor 1, Metalloproteinase inhibitor 2, Metalloproteinase inhibitor 3, Metalloproteinase inhibitor 4, Myoglobin, Pancreatic prohormone, Peptide YY, Pro-epidermal growth factor, Pro-interleukin-16, Prolactin, Prostate-specific antigen, Protein S100-A12, Protransforming growth factor alpha, Secretory immunoglobulin A, Serum amyloid P-component, SL cytokine, Stromal cell-derived factor 1, Stromelysin-1, Thrombopoietin, Thymic stromal lymphopoietin, Tumor necrosis factor, Tumor necrosis factor ligand superfamily member 10, Tumor necrosis factor ligand superfamily member 6, Tumor necrosis factor receptor superfamily member 1A, Tumor necrosis factor receptor superfamily member 1B, Vascular endothelial growth factor receptor 1, Vascular endothelial growth factor receptor 2, Vascular endothelial growth factor receptor 3, von Willebrand Factor, and WAP four-disulfide core domain protein 2, the results of which assay(s) is/are then correlated to the status of the patient. This correlation to status may include one or more of the following: diagnosis of acute appendicitis; indication of a prognosis resulting from acute appendicitis. For convenience, patients being evaluated in this manner are referred to herein as "appendicitis patients," whether or not the appendicitis diagnosis has been confirmed at the time of the evaluation.

[0010] In certain embodiments, the methods for evaluating a patient described herein are methods for risk stratification of the patient; that is, assigning a likelihood of one or more future changes in health status to the patient. In these embodiments, the assay result(s) is/are correlated to one or more such future changes. A level or a change in level

of one or more appendicitis biomarkers, which in turn is(are) associated with an increased probability of morbidity or death, is referred to as being "associated with an increased predisposition to an adverse outcome" in a patient.

[0011] In such risk stratification embodiments, preferably the likelihood or risk assigned is that an event of interest is more or less likely to occur within 180 days of the time at which the body fluid sample is obtained from the appendicitis patient. In particularly preferred embodiments, the likelihood or risk assigned relates to an event of interest occurring within a shorter time period such as 18 months, 120 days, 90 days, 60 days, 45 days, 30 days, 21 days, 14 days, 7 days, 5 days, 96 hours, 72 hours, 48 hours, 36 hours, 24 hours, 12 hours, or less. A risk at 0 hours of the time at which the body fluid sample is obtained from the appendicitis patient is equivalent to diagnosis of a current condition.

[0012] For a positive going marker, an increased likelihood of the occurrence of a diagnosis is assigned to the patient when the measured concentration is above the threshold (relative to the likelihood assigned when the measured concentration is below the threshold); alternatively, when the measured concentration is below the threshold, an increased likelihood of the nonoccurrence of a diagnosis may be assigned to the patient (relative to the likelihood assigned when the measured concentration is above the threshold). For a negative going marker, an increased likelihood of the occurrence of a diagnosis is assigned to the patient when the measured concentration is below the threshold (relative to the likelihood assigned when the measured concentration is above the threshold); alternatively, when the measured concentration is above the threshold, an increased likelihood of the nonoccurrence of a diagnosis may be assigned to the patient (relative to the likelihood assigned when the measured concentration is below the threshold).

[0013] In certain embodiments, a biomarker or panel of biomarkers is correlated to a condition or disease by merely its presence or absence. In other embodiments, a threshold level of a diagnostic or prognostic indicator can be established, and the level of the indicator in a patient sample can simply be compared to the threshold level. A variety of methods may be used by the skilled artisan to arrive at a desired threshold value for use in these methods. For example, for a positive going marker the threshold value may be determined from a population of patients not having acute appendicitis by selecting a concentration representing the 75th, 85th, 90th, 95th, or 99th percentile of an appendicitis

biomarker or biomarkers measured in such "normal" patients. Alternatively, the threshold value may be determined from a "diseased" population of patients by selecting a concentration representing the 75th, 85th, 90th, 95th, or 99th percentile of a biomarker or biomarkers measured in patients suffering from acute appendicitis.

[0014] Alternatively, the threshold value may be determined from a "diseased" population of appendicitis patients having a predisposition for an outcome such as death, worsening disease, *etc.*), by selecting a concentration representing the 75th, 85th, 90th, 95th, or 99th percentile of a biomarker or biomarkers measured in patients suffering from acute appendicitis and who later suffered from the outcome of interest.

[0015] In another alternative, the threshold value may be determined from a prior measurement of a biomarker or biomarkers in the same patient; that is, a temporal change in the level of a biomarker or biomarkers in the same patient may be used to assign a diagnosis or a prognosis to the patient. For example, a diagnostic indicator may be determined at an initial time, and again at a second time. In such embodiments, an increase in the marker from the initial time to the second time may be diagnostic of appendicitis or a given prognosis.

[0016] The foregoing discussion is not meant to imply, however, that the appendicitis biomarkers of the present invention must be compared to corresponding individual thresholds. Methods for combining assay results can comprise the use of multivariate logistical regression, loglinear modeling, neural network analysis, n-of-m analysis, decision tree analysis, calculating ratios of markers, *etc.* This list is not meant to be limiting. In these methods, a composite result which is determined by combining individual markers may be treated as if it is itself a marker; that is, a threshold may be determined for the composite result as described herein for individual markers, and the composite result for an individual patient compared to this threshold.

[0017] The ability of a particular test to distinguish two populations can be established using ROC analysis. For example, ROC curves established from a "first" subpopulation which has a particular disease (or which is predisposed to some outcome), and a "second" subpopulation which does not have the disease (or is not so predisposed) can be used to calculate a ROC curve, and the area under the curve provides a measure of the quality of the test. Preferably, the tests described herein provide a ROC curve area

greater than 0.5, preferably at least 0.6, more preferably 0.7, still more preferably at least 0.8, even more preferably at least 0.9, and most preferably at least 0.95.

[0018] In certain aspects, the measured concentration of one or more appendicitis biomarkers, or a composite of such markers, may be treated as continuous variables. For example, any particular concentration can be converted into a corresponding probability of existing disease, of a future outcome for the appendicitis patient, or mortality, of a SIRS classification, etc. In yet another alternative, a threshold that can provide an acceptable level of specificity and sensitivity in separating a population of appendicitis patients into "bins" such as a "first" subpopulation and a "second" subpopulation. A threshold value is selected to separate this first and second population by one or more of the following measures of test accuracy:

an odds ratio greater than 1, preferably at least about 2 or more or about 0.5 or less, more preferably at least about 3 or more or about 0.33 or less, still more preferably at least about 4 or more or about 0.25 or less, even more preferably at least about 5 or more or about 0.2 or less, and most preferably at least about 10 or more or about 0.1 or less;

a specificity of greater than 0.5, preferably at least about 0.6, more preferably at least about 0.7, still more preferably at least about 0.8, even more preferably at least about 0.9 and most preferably at least about 0.95, with a corresponding sensitivity greater than 0.2, preferably greater than about 0.3, more preferably greater than about 0.4, still more preferably at least about 0.5, even more preferably about 0.6, yet more preferably greater than about 0.7, still more preferably greater than about 0.8, more preferably greater than about 0.9, and most preferably greater than about 0.95;

a sensitivity of greater than 0.5, preferably at least about 0.6, more preferably at least about 0.7, still more preferably at least about 0.8, even more preferably at least about 0.9 and most preferably at least about 0.95, with a corresponding specificity greater than 0.2, preferably greater than about 0.3, more preferably greater than about 0.4, still more preferably at least about 0.5, even more preferably about 0.6, yet more preferably greater than about 0.7, still more preferably greater than about 0.8, more preferably greater than about 0.9, and most preferably greater than about 0.95;

at least about 75% sensitivity, combined with at least about 75% specificity;

a positive likelihood ratio (calculated as sensitivity/(1-specificity)) of greater than 1, at least about 2, more preferably at least about 3, still more preferably at least about 5, and most preferably at least about 10; or

a negative likelihood ratio (calculated as (1-sensitivity)/specificity) of less than 1, less than or equal to about 0.5, more preferably less than or equal to about 0.3, and most preferably less than or equal to about 0.1.

[0019] The term "about" in the context of any of the above measurements refers to +/- 5% of a given measurement.

[0020] Multiple thresholds may also be used to assess a patient. For example, a "first" subpopulation identified by an existing disease, predisposition to a future outcome for the appendicitis patient, predisposition to mortality, etc., and a "second" subpopulation which is not so predisposed can be combined into a single group. This group is then subdivided into three or more equal parts (known as tertiles, quartiles, quintiles, etc., depending on the number of subdivisions). An odds ratio is assigned to appendicitis patients based on which subdivision they fall into. If one considers a tertile, the lowest or highest tertile can be used as a reference for comparison of the other subdivisions. This reference subdivision is assigned an odds ratio of 1. The second tertile is assigned an odds ratio that is relative to that first tertile. That is, someone in the second tertile might be 3 times more likely to suffer one or more future changes in disease status in comparison to someone in the first tertile. The third tertile is also assigned an odds ratio that is relative to that first tertile.

[0021] In certain embodiments, the assay method is an immunoassay. Antibodies for use in such assays will specifically bind a full length appendicitis biomarker of interest, and may also bind one or more polypeptides that are "related" thereto, as that term is defined hereinafter. Numerous immunoassay formats are known to those of skill in the art. Preferred body fluid samples are selected from the group consisting of urine, blood, serum, saliva, tears, and plasma.

[0022] The foregoing method steps should not be interpreted to mean that the appendicitis biomarker assay result(s) is/are used in isolation in the methods described herein. Rather, additional variables or other clinical indicia may be included in the methods described herein. For example, a risk stratification, diagnostic, classification, monitoring, etc. method may combine the assay result(s) with one or more variables

measured for the appendicitis patient selected from the group consisting of demographic information (e.g., weight, sex, age, race), clinical variables (e.g., blood pressure, temperature, respiration rate), risk scores (Alvarado score, Pediatric Appendicitis Score, etc.). This list is not meant to be limiting.

[0023] When more than one marker is measured, the individual markers may be measured in samples obtained at the same time, or may be determined from samples obtained at different (e.g., an earlier or later) times. The individual markers may also be measured on the same or different body fluid samples. For example, one appendicitis biomarker may be measured in a serum or plasma sample and another appendicitis biomarker may be measured in a urine sample. In addition, assignment of a likelihood may combine an individual biomarker assay result with temporal changes in one or more additional variables.

[0024] In various related aspects, the present invention also relates to devices and kits for performing the methods described herein. Suitable kits comprise reagents sufficient for performing an assay for at least one of the described appendicitis biomarkers, together with instructions for performing the described threshold comparisons.

[0025] In certain embodiments, reagents for performing such assays are provided in an assay device, and such assay devices may be included in such a kit. Preferred reagents can comprise one or more solid phase antibodies, the solid phase antibody comprising antibody that detects the intended biomarker target(s) bound to a solid support. In the case of sandwich immunoassays, such reagents can also include one or more detectably labeled antibodies, the detectably labeled antibody comprising antibody that detects the intended biomarker target(s) bound to a detectable label. Additional optional elements that may be provided as part of an assay device are described hereinafter.

[0026] Detectable labels may include molecules that are themselves detectable (e.g., fluorescent moieties, electrochemical labels, eel (electrochemical luminescence) labels, metal chelates, colloidal metal particles, *etc.*) as well as molecules that may be indirectly detected by production of a detectable reaction product (e.g., enzymes such as horseradish peroxidase, alkaline phosphatase, *etc.*) or through the use of a specific binding molecule which itself may be detectable (e.g., a labeled antibody that binds to the second antibody, biotin, digoxigenin, maltose, oligohistidine, 2,4-dinitrobenzene, phenylarsenate, ssDNA, dsDNA, *etc.*).

[0027] Generation of a signal from the signal development element can be performed using various optical, acoustical, and electrochemical methods well known in the art. Examples of detection modes include fluorescence, radiochemical detection, reflectance, absorbance, amperometry, conductance, impedance, interferometry, ellipsometry, *etc.* In certain of these methods, the solid phase antibody is coupled to a transducer (*e.g.*, a diffraction grating, electrochemical sensor, etc) for generation of a signal, while in others, a signal is generated by a transducer that is spatially separate from the solid phase antibody (*e.g.*, a fluorometer that employs an excitation light source and an optical detector). This list is not meant to be limiting. Antibody-based biosensors may also be employed to determine the presence or amount of analytes that optionally eliminate the need for a labeled molecule.

DETAILED DESCRIPTION OF THE INVENTION

[0028] The present invention relates to methods and compositions for diagnosis, differential diagnosis, risk stratification, monitoring, classifying and determination of treatment regimens in patients diagnosed with, or at risk of, appendicitis. In various embodiments, a measured concentration of one or more biomarkers selected from the group consisting of 72 kDa type IV collagenase, Adiponectin, Advanced glycosylation end product-specific receptor, Alpha-2 macroglobulin, Alpha-2-HS-glycoprotein, Alpha-fetoprotein, Angiotensin-1, Antileukoprotease, Apolipoprotein(a), Cancer Antigen 15-3, Cancer Antigen 19-9, Carbonic anhydrase 9, Carcinoembryonic antigen-related cell adhesion molecule 5, C-C motif chemokine 1, C-C motif chemokine 13, C-C motif chemokine 15, C-C motif chemokine 17, C-C motif chemokine 19, C-C motif chemokine 20, C-C motif chemokine 21, C-C motif chemokine 22, C-C motif chemokine 23, C-C motif chemokine 24, C-C motif chemokine 26, C-C motif chemokine 27, C-C motif chemokine 3, C-C motif chemokine 4, C-C motif chemokine 7, C-C motif chemokine 8, Ceruloplasmin, Choriogonadotropin subunit beta, Collagenase 3, C-Peptide, Creatine Kinase-MB, C-X-C motif chemokine 10, C-X-C motif chemokine 11, C-X-C motif chemokine 13, C-X-C motif chemokine 16, C-X-C motif chemokine 5, C-X-C motif chemokine 6, C-X-C motif chemokine 9, Cystatin-C, Endothelial protein C receptor, Eotaxin, Epidermal growth factor receptor, Ferritin, Fibrinogen, Gastric inhibitory polypeptide, Glucagon, Glucagon-like peptide 1, Granulocyte-macrophage colony-stimulating factor, Growth-regulated alpha, beta, and gamma proteins, Heparin-binding

growth factor 2, Hepatocyte growth factor, Immunoglobulin A, Immunoglobulin M, Immunoglobulin G1, Immunoglobulin G2, Immunoglobulin G3, Immunoglobulin G4, Insulin, Insulin-like growth factor-binding protein 1, Insulin-like growth factor-binding protein 2, Insulin-like growth factor-binding protein 3, Insulin-like growth factor-binding protein 4, Insulin-like growth factor-binding protein 5, Insulin-like growth factor-binding protein 6, Insulin-like growth factor-binding protein 7, Interferon alpha-2, Interferon gamma, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 receptor antagonist protein, Interleukin-1 receptor type I, Interleukin-1 receptor type II, Interleukin-1 1, Interleukin-12, Interleukin-12 subunit beta, Interleukin-13, Interleukin-15, Interleukin-2, Interleukin-20, Interleukin-21, Interleukin-23, Interleukin-28A, Interleukin-29, Interleukin-3, Interleukin-33, Interleukin-4, Interleukin-5, Interleukin-6 receptor subunit alpha, Interleukin-6 receptor subunit beta, Interleukin-7, Interleukin-9, Interstitial collagenase, Islet amyloid polypeptide, Keratin, type I cytoskeletal 19 (aa311-367), Kit ligand, Leptin, Leukemia inhibitory factor, Lymphotoxin, Lymphotoxin-alpha, Macrophage colony-stimulating factor 1, Macrophage metalloelastase, Macrophage migration inhibitory factor, Matrilysin, Metalloproteinase inhibitor 1, Metalloproteinase inhibitor 2, Metalloproteinase inhibitor 3, Metalloproteinase inhibitor 4, Myoglobin, Pancreatic prohormone, Peptide YY, Pro-epidermal growth factor, Pro-interleukin-16, Prolactin, Prostate-specific antigen, Protein S100-A12, Protransforming growth factor alpha, Secretory immunoglobulin A, Serum amyloid P-component, SL cytokine, Stromal cell-derived factor 1, Stromelysin-1, Thrombopoietin, Thymic stromal lymphopoietin, Tumor necrosis factor, Tumor necrosis factor ligand superfamily member 10, Tumor necrosis factor ligand superfamily member 6, Tumor necrosis factor receptor superfamily member 1A, Tumor necrosis factor receptor superfamily member 1B, Vascular endothelial growth factor receptor 1, Vascular endothelial growth factor receptor 2, Vascular endothelial growth factor receptor 3, von Willebrand Factor, and WAP four-disulfide core domain protein 2 or one or more markers related thereto, are correlated to the status of the patient. As described herein, measurement of one or more biomarkers of the present invention may be used, individually or in panels comprising a plurality of biomarkers, in methods and compositions for the diagnosis, prognosis, or differentiation of abdominal pain in order to rule in or out appendicitis and/or a particular outcome. Such markers can be used in diagnosing and treating a subject and/or to monitor the course of a treatment regimen; for screening subjects for the occurrence or risk of a particular disease; and for screening

compounds and pharmaceutical compositions that might provide a benefit in treating or preventing such conditions.

[0029] For purposes of this document, the following definitions apply:

[0030] The term "subject" as used herein refers to a human or non-human organism. Thus, the methods and compositions described herein are applicable to both human and veterinary disease. Further, while a subject is preferably a living organism, the invention described herein may be used in post-mortem analysis as well. Preferred subjects are humans, and most preferably "patients," which as used herein refers to living humans that are receiving medical care for a disease or condition. This includes persons with no defined illness who are being investigated for signs of pathology. An "appendicitis patient" is a patient exhibiting symptoms consistent with appendicitis and being evaluated for its presence, absence, or outcome

[0031] Conditions within the differential diagnosis include gallbladder attack, kidney infection, pneumonia, rheumatic fever, diabetic ketoacidosis, ectopic pregnancy, twisted ovarian cyst, hemorrhaging ovarian follicle, urinary tract infection, ulcerative colitis, pancreatitis, intestinal obstruction, pelvic inflammatory disease, diverticulitis, carcinoma of the colon, and aortic aneurysm. In preferred embodiments, the biomarkers of the present invention distinguish appendicitis from one or more of these mimicking conditions.

[0032] Preferably, an analyte such as an appendicitis biomarker is measured in a sample. Such a sample may be obtained from a patient, such as an appendicitis patient. Preferred samples are body fluid samples.

[0033] The term "body fluid sample" as used herein refers to a sample of bodily fluid obtained for the purpose of diagnosis, prognosis, classification or evaluation of an appendicitis patient of interest, such as a patient or transplant donor. In certain embodiments, such a sample may be obtained for the purpose of determining the outcome of an ongoing condition or the effect of a treatment regimen on a condition. Preferred body fluid samples include blood, serum, plasma, urine, saliva, sputum, and pleural effusions. In addition, one of skill in the art would realize that certain body fluid samples would be more readily analyzed following a fractionation or purification procedure, for example, separation of whole blood into serum or plasma components.

[0034] The term "diagnosis" as used herein refers to methods by which the skilled artisan can estimate and/or determine the probability ("a likelihood") of whether or not a patient is suffering from a given disease or condition. In the case of the present invention, "diagnosis" includes using the results of an assay, most preferably an immunoassay, for an appendicitis biomarker of the present invention, optionally together with other clinical characteristics, to arrive at a diagnosis (that is, the occurrence or nonoccurrence) of a disease or condition. That such a diagnosis is "determined" is not meant to imply that the diagnosis is 100% accurate. Many biomarkers are indicative of multiple conditions. The skilled clinician does not use biomarker results in an informational vacuum, but rather test results are used together with other clinical indicia to arrive at a diagnosis. Thus, a measured biomarker level on one side of a predetermined diagnostic threshold indicates a greater likelihood of the occurrence of disease in the appendicitis patient relative to a measured level on the other side of the predetermined diagnostic threshold.

[0035] Similarly, a prognostic risk signals a probability ("a likelihood") that a given course or outcome will occur. A level or a change in level of a prognostic indicator, which in turn is associated with an increased probability of morbidity or mortality is referred to as being "indicative of an increased likelihood" of an adverse outcome in a patient.

[0036] As used herein, the term "relating a signal to the presence or amount" of an analyte reflects the following understanding. Assay signals are typically related to the presence or amount of an analyte through the use of a standard curve calculated using known concentrations of the analyte of interest. As the term is used herein, an assay is "configured to detect" an analyte if an assay can generate a detectable signal indicative of the presence or amount of a physiologically relevant concentration of the analyte. Because an antibody epitope is on the order of 8 amino acids, an immunoassay configured to detect a marker of interest will also detect polypeptides related to the marker sequence, so long as those polypeptides contain the epitope(s) necessary to bind to the antibody or antibodies used in the assay. The term "related marker" as used herein with regard to a biomarker such as one of the appendicitis biomarkers described herein refers to one or more fragments, variants, etc., of a particular marker or its biosynthetic parent that may be detected as a surrogate for the marker itself or as independent biomarkers. The term also refers to one or more polypeptides present in a biological

sample that are derived from the biomarker precursor complexed to additional species, such as binding proteins, receptors, heparin, lipids, sugars, *etc.*

[0037] In this regard, the skilled artisan will understand that the signals obtained from an immunoassay are a direct result of complexes formed between one or more antibodies and the target biomolecule (*i.e.*, the analyte) and polypeptides containing the necessary epitope(s) to which the antibodies bind. While such assays may detect the full length biomarker and the assay result be expressed as a concentration of a biomarker of interest, the signal from the assay is actually a result of all such "immunoreactive" polypeptides present in the sample. Expression of biomarkers may also be determined by means other than immunoassays, including protein measurements (such as dot blots, western blots, chromatographic methods, mass spectrometry, *etc.*) and nucleic acid measurements (mRNA quantitation). This list is not meant to be limiting. With regard to biomarkers which exist in one form as type-I, type-II, or GPTanchored membrane proteins, such membrane proteins typically comprise a substantial extracellular domain, some or all of which can be detected as soluble forms present in aqueous samples such as blood, serum, plasma, urine, *etc.*, either as cleavage products or as splice variants which delete an effective membrane spanning domain. Preferred assays detect soluble forms of these biomarkers.

[0038] The term "positive going" marker as that term is used herein refer to a marker that is determined to be elevated in patients suffering from a disease or condition, relative to those not suffering from that disease or condition. The term "negative going" marker as that term is used herein refer to a marker that is determined to be reduced in patients suffering from a disease or condition, relative to patients not suffering from that disease or condition.

[0039] Appendicitis biomarkers

[0040] The following table provides a list of the biomarkers of the present invention, together with the Swiss-Prot entry number for the human precursor. As noted above, these biomarkers are referred to for convenience herein as "appendicitis biomarkers."

<u>SwissProtNum</u>	<u>Preferred Name</u>
Q15848	Adiponectin
Q15109	Advanced glycosylation end product-specific receptor
P01023	Alpha-2 macro globulin
P02765	Alpha-2-HS-glycoprotein

<u>SwissProtNum</u>	<u>Preferred Name</u>
P02771	Alpha-fetoprotein
Q15389	Angiopoietin-1
P03973	Antileukoproteinase
P08519	Apolipoprotein(a)
P15941	Cancer Antigen 15-3
N/A	Cancer Antigen 19-9
Q16790	Carbonic anhydrase 9
P06731	Carcinoembryonic antigen-related cell adhesion molecule 5
P22362	C-C motif chemokine 1
Q99616	C-C motif chemokine 13
Q16663	C-C motif chemokine 15
Q92583	C-C motif chemokine 17
Q99731	C-C motif chemokine 19
P78556	C-C motif chemokine 20
000585	C-C motif chemokine 21
000626	C-C motif chemokine 22
P55773	C-C motif chemokine 23
000175	C-C motif chemokine 24
Q9Y258	C-C motif chemokine 26
Q9Y4X3	C-C motif chemokine 27
P10147	C-C motif chemokine 3
P13236	C-C motif chemokine 4
P80098	C-C motif chemokine 7
P80075	C-C motif chemokine 8
P00450	Ceruloplasmin
P01233	Choriogonadotropin subunit beta
P45452	Collagenase 3
P01308(aa57-87)	C-Peptide
P12277;P06732	Creatine Kinase-MB
P02778	C-X-C motif chemokine 10
014625	C-X-C motif chemokine 11
043927	C-X-C motif chemokine 13
Q9H2A7	C-X-C motif chemokine 16
P42830	C-X-C motif chemokine 5
P80162	C-X-C motif chemokine 6
Q07325	C-X-C motif chemokine 9
P01034	Cystatin-C
Q9UNN8	Endothelial protein C receptor
P51671	Eotaxin
P00533	Epidermal growth factor receptor
P02792;P02794	Ferritin
P02671;P02675;P02679	Fibrinogen
P09681	Gastric inhibitory polypeptide
P01275	Glucagon

<u>SwissProtNum</u>	<u>Preferred Name</u>
P01275(aa98-127;aa98-128)	Glucagon-like peptide 1
P04141	Granulocyte-macrophage colony-stimulating factor
P09341;P19875;P19876	Growth-regulated alpha, beta, and gamma proteins
P09038	Heparin-binding growth factor 2
P14210	Hepatocyte growth factor
N/A	Immunoglobulin A
n/a	Immunoglobulin M
n/a	Immunoglobulin G1
N/A	Immunoglobulin G2
N/A	Immunoglobulin G3
n/a	Immunoglobulin G4
P01308	Insulin
P08833	Insulin-like growth factor-binding protein 1
P18065	Insulin-like growth factor-binding protein 2
P17936	Insulin-like growth factor-binding protein 3
P22692	Insulin-like growth factor-binding protein 4
P24593	Insulin-like growth factor-binding protein 5
P24592	Insulin-like growth factor-binding protein 6
Q16270	Insulin-like growth factor-binding protein 7
P01563	Interferon alpha-2
P01579	Interferon gamma
P01583	Interleukin-1 alpha
P01584	Interleukin-1 beta
P18510	Interleukin-1 receptor antagonist protein
P14778	Interleukin-1 receptor type I
P27930	Interleukin-1 receptor type II
P20809	Interleukin-11
P29459;P29460	Interleukin-12
P29460	Interleukin-12 subunit beta
P35225	Interleukin-13
P40933	Interleukin-15
P60568	Interleukin-2
Q9NYY1	Interleukin-20
Q9HBE4	Interleukin-21
Q9NPF7;P29460	Interleukin-23
Q8IZJ0	Interleukin-28A
Q8IU54	Interleukin-29
P08700	Interleukin-3
095760	Interleukin-33
P05112	Interleukin-4
P05113	Interleukin-5
P08887	Interleukin-6 receptor subunit alpha
P40189	Interleukin-6 receptor subunit beta
P13232	Interleukin-7
PI5248	Interleukin-9

<u>SwissProtNum</u>	<u>Preferred Name</u>
P03956	Interstitial collagenase
P10997	Islet amyloid polypeptide
P08727	Keratin, type I cytoskeletal 19 (aa31 1-367)
P21583	Kit ligand
P41159	Leptin
P15018	Leukemia inhibitory factor
P47992	Lymphotoxin
P01374	Lymphotoxin-alpha
P09603	Macrophage colony-stimulating factor 1
P39900	Macrophage metalloelastase
P14174	Macrophage migration inhibitory factor
P09237	Matrilysin
P01033	Metalloproteinase inhibitor 1
P16035	Metalloproteinase inhibitor 2
P35625	Metalloproteinase inhibitor 3
Q99727	Metalloproteinase inhibitor 4
P02144	Myoglobin
P01298	Pancreatic prohormone
P10082	Peptide YY
P01133	Pro-epidermal growth factor
Q14005	Pro-interleukin- 16
P01236	Prolactin
P07288	Prostate-specific antigen
P80511	Protein S100-A12
P01135	Protransforming growth factor alpha
N/A	Secretory immunoglobulin A
P02743	Serum amyloid P-component
P49771	SL cytokine
P48061	Stromal cell-derived factor 1
P08254	Stromelysin-1
P40225	Thrombopoietin
Q969D9	Thymic stromal lymphopoietin
P01375	Tumor necrosis factor
P50591	Tumor necrosis factor ligand superfamily member 10
P48023	Tumor necrosis factor ligand superfamily member 6
P19438	Tumor necrosis factor receptor superfamily member 1A
P20333	Tumor necrosis factor receptor superfamily member 1B
P17948	Vascular endothelial growth factor receptor 1
P35968	Vascular endothelial growth factor receptor 2
P35916	Vascular endothelial growth factor receptor 3
P04275	von Willebrand Factor
Q14508	WAP four-disulfide core domain protein 2
P05231	Interleukin-6
P08253	72 kDa type IV collagenase

[0041] Marker Assays

[0042] In general, immunoassays involve contacting a sample containing or suspected of containing a biomarker of interest with at least one antibody that specifically binds to the biomarker. A signal is then generated indicative of the presence or amount of complexes formed by the binding of polypeptides in the sample to the antibody. The signal is then related to the presence or amount of the biomarker in the sample. Numerous methods and devices are well known to the skilled artisan for the detection and analysis of biomarkers. *See, e.g.*, U.S. Patents 6,143,576; 6,113,855; 6,019,944; 5,985,579; 5,947,124; 5,939,272; 5,922,615; 5,885,527; 5,851,776; 5,824,799; 5,679,526; 5,525,524; and 5,480,792, and *The Immunoassay Handbook*, David Wild, ed. Stockton Press, New York, 1994, each of which is hereby incorporated by reference in its entirety, including all tables, figures and claims.

[0043] The assay devices and methods known in the art can utilize labeled molecules in various sandwich, competitive, or non-competitive assay formats, to generate a signal that is related to the presence or amount of the biomarker of interest. Suitable assay formats also include chromatographic, mass spectrographic, and protein "blotting" methods. Additionally, certain methods and devices, such as biosensors and optical immunoassays, may be employed to determine the presence or amount of analytes without the need for a labeled molecule. *See, e.g.*, U.S. Patents 5,631,171; and 5,955,377, each of which is hereby incorporated by reference in its entirety, including all tables, figures and claims. One skilled in the art also recognizes that robotic instrumentation including but not limited to Beckman ACCESS®, Abbott AXSYM®, Roche ELECSYS®, Dade Behring STRATUS® systems are among the immunoassay analyzers that are capable of performing immunoassays. But any suitable immunoassay may be utilized, for example, enzyme-linked immunoassays (ELISA), radioimmunoassays (RIAs), competitive binding assays, and the like.

[0044] Antibodies or other polypeptides may be immobilized onto a variety of solid supports for use in assays. Solid phases that may be used to immobilize specific binding members include those developed and/or used as solid phases in solid phase binding assays. Examples of suitable solid phases include membrane filters, cellulose-based papers, beads (including polymeric, latex and paramagnetic particles), glass, silicon wafers, microparticles, nanoparticles, TentaGels, AgroGels, PEGA gels, SPOCC gels, and multiple-well plates. An assay strip could be prepared by coating the antibody or a

plurality of antibodies in an array on solid support. This strip could then be dipped into the test sample and then processed quickly through washes and detection steps to generate a measurable signal, such as a colored spot. Antibodies or other polypeptides may be bound to specific zones of assay devices either by conjugating directly to an assay device surface, or by indirect binding. In an example of the later case, antibodies or other polypeptides may be immobilized on particles or other solid supports, and that solid support immobilized to the device surface.

[0045] Biological assays require methods for detection, and one of the most common methods for quantitation of results is to conjugate a detectable label to a protein or nucleic acid that has affinity for one of the components in the biological system being studied. Detectable labels may include molecules that are themselves detectable (*e.g.*, fluorescent moieties, electrochemical labels, metal chelates, *etc.*) as well as molecules that may be indirectly detected by production of a detectable reaction product (*e.g.*, enzymes such as horseradish peroxidase, alkaline phosphatase, *etc.*) or by a specific binding molecule which itself may be detectable (*e.g.*, biotin, digoxigenin, maltose, oligohistidine, 2,4-dinitrobenzene, phenylarsenate, ssDNA, dsDNA, *etc.*).

[0046] Preparation of solid phases and detectable label conjugates often comprise the use of chemical cross-linkers. Cross-linking reagents contain at least two reactive groups, and are divided generally into homofunctional cross-linkers (containing identical reactive groups) and heterofunctional cross-linkers (containing non-identical reactive groups). Homobifunctional cross-linkers that couple through amines, sulfhydryls or react non-specifically are available from many commercial sources. Maleimides, alkyl and aryl halides, alpha-haloacyls and pyridyl disulfides are thiol reactive groups. Maleimides, alkyl and aryl halides, and alpha-haloacyls react with sulfhydryls to form thiol ether bonds, while pyridyl disulfides react with sulfhydryls to produce mixed disulfides. The pyridyl disulfide product is cleavable. Imidoesters are also very useful for protein-protein cross-links. A variety of heterobifunctional cross-linkers, each combining different attributes for successful conjugation, are commercially available.

[0047] In certain aspects, the present invention provides kits for the analysis of the described appendicitis biomarkers. The kit comprises reagents for the analysis of at least one test sample which comprise at least one antibody that binds an appendicitis biomarker. The kit can also include devices and instructions for performing one or more of the diagnostic and/or prognostic correlations described herein. Preferred kits will

comprise an antibody pair for performing a sandwich assay, or a labeled species for performing a competitive assay, for the analyte. Preferably, an antibody pair comprises a first antibody conjugated to a solid phase and a second antibody conjugated to a detectable label, wherein each of the first and second antibodies bind an appendicitis biomarker. Most preferably each of the antibodies are monoclonal antibodies. The instructions for use of the kit and performing the correlations can be in the form of labeling, which refers to any written or recorded material that is attached to, or otherwise accompanies a kit at any time during its manufacture, transport, sale or use. For example, the term labeling encompasses advertising leaflets and brochures, packaging materials, instructions, audio or video cassettes, computer discs, as well as writing imprinted directly on kits.

[0048] Antibodies

[0049] The term "antibody" as used herein refers to a peptide or polypeptide derived from, modeled after or substantially encoded by an immunoglobulin gene or immunoglobulin genes, or fragments thereof, capable of specifically binding an antigen or epitope. See, e.g. Fundamental Immunology, 3rd Edition, W.E. Paul, ed., Raven Press, N.Y. (1993); Wilson (1994; J. Immunol. Methods 175:267-273; Yarmush (1992) J. Biochem. Biophys. Methods 25:85-97. The term antibody includes antigen-binding portions, i.e., "antigen binding sites," (e.g., fragments, subsequences, complementarity determining regions (CDRs)) that retain capacity to bind antigen, including (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CHI domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CHI domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward et al., (1989) Nature 341:544-546), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR). Single chain antibodies are also included by reference in the term "antibody."

[0050] Antibodies used in the immunoassays described herein preferably specifically bind to an appendicitis biomarker of the present invention. The term "specifically binds" is not intended to indicate that an antibody binds exclusively to its intended target since, as noted above, an antibody binds to any polypeptide displaying the epitope(s) to which the antibody binds. Rather, an antibody "specifically binds" if its affinity for its intended target is about 5-fold greater when compared to its affinity for a non-target molecule

which does not display the appropriate epitope(s). Preferably the affinity of the antibody will be at least about 5 fold, preferably 10 fold, more preferably 25-fold, even more preferably 50-fold, and most preferably 100-fold or more, greater for a target molecule than its affinity for a non-target molecule. In preferred embodiments, preferred antibodies bind with affinities of at least about 10^7 M^{-1} , and preferably between about 10^8 M^{-1} to about 10^9 M^{-1} , about 10^9 M^{-1} to about 10^{10} M^{-1} , or about 10^{10} M^{-1} to about 10^{12} M^{-1} .

[0051] Affinity is calculated as $K_d = k_{\text{off}}/k_{\text{on}}$ (k_{off} is the dissociation rate constant, K_{on} is the association rate constant and K_d is the equilibrium constant). Affinity can be determined at equilibrium by measuring the fraction bound (r) of labeled ligand at various concentrations (c). The data are graphed using the Scatchard equation: $r/c = K(n-r)$: where r = moles of bound ligand/mole of receptor at equilibrium; c = free ligand concentration at equilibrium; K = equilibrium association constant; and n = number of ligand binding sites per receptor molecule. By graphical analysis, r/c is plotted on the Y-axis versus r on the X-axis, thus producing a Scatchard plot. Antibody affinity measurement by Scatchard analysis is well known in the art. See, e.g., van Erp *et al.*, *J. Immunoassay* 12: 425-43, 1991; Nelson and Griswold, *Comput. Methods Programs Biomed.* 27: 65-8, 1988.

[0052] The term "epitope" refers to an antigenic determinant capable of specific binding to an antibody. Epitopes usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. Conformational and nonconformational epitopes are distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents.

[0053] Numerous publications discuss the use of phage display technology to produce and screen libraries of polypeptides for binding to a selected analyte. See, e.g. Cwirla *et al.*, *Proc. Natl. Acad. Sci. USA* 87, 6378-82, 1990; Devlin *et al.*, *Science* 249, 404-6, 1990, Scott and Smith, *Science* 249, 386-88, 1990; and Ladner *et al.*, U.S. Pat. No. 5,571,698. A basic concept of phage display methods is the establishment of a physical association between DNA encoding a polypeptide to be screened and the polypeptide. This physical association is provided by the phage particle, which displays a polypeptide as part of a capsid enclosing the phage genome which encodes the polypeptide. The establishment of a physical association between polypeptides and their genetic material allows simultaneous mass screening of very large numbers of phage bearing different polypeptides. Phage displaying a polypeptide with affinity to a target bind to the target

and these phage are enriched by affinity screening to the target. The identity of polypeptides displayed from these phage can be determined from their respective genomes. Using these methods a polypeptide identified as having a binding affinity for a desired target can then be synthesized in bulk by conventional means. *See, e.g.*, U.S. Patent No. 6,057,098, which is hereby incorporated in its entirety, including all tables, figures, and claims.

[0054] The antibodies that are generated by these methods may then be selected by first screening for affinity and specificity with the purified polypeptide of interest and, if required, comparing the results to the affinity and specificity of the antibodies with polypeptides that are desired to be excluded from binding. The screening procedure can involve immobilization of the purified polypeptides in separate wells of microtiter plates. The solution containing a potential antibody or groups of antibodies is then placed into the respective microtiter wells and incubated for about 30 min to 2 h. The microtiter wells are then washed and a labeled secondary antibody (for example, an anti-mouse antibody conjugated to alkaline phosphatase if the raised antibodies are mouse antibodies) is added to the wells and incubated for about 30 min and then washed. Substrate is added to the wells and a color reaction will appear where antibody to the immobilized polypeptide(s) are present.

[0055] The antibodies so identified may then be further analyzed for affinity and specificity in the assay design selected. In the development of immunoassays for a target protein, the purified target protein acts as a standard with which to judge the sensitivity and specificity of the immunoassay using the antibodies that have been selected. Because the binding affinity of various antibodies may differ; certain antibody pairs (e.g., in sandwich assays) may interfere with one another sterically, etc., assay performance of an antibody may be a more important measure than absolute affinity and specificity of an antibody.

[0056] While the present application describes antibody-based binding assays in detail, alternatives to antibodies as binding species in assays are well known in the art. These include receptors for a particular target, aptamers, etc. Aptamers are oligonucleic acid or peptide molecules that bind to a specific target molecule. Aptamers are usually created by selecting them from a large random sequence pool, but natural aptamers also exist. High-affinity aptamers containing modified nucleotides can confer improved characteristics on the ligand, such as improved in vivo stability or improved delivery

characteristics. Examples of such modifications include chemical substitutions at the ribose and/or phosphate and/or base positions, and may include amino acid side chain functionalities.

[0057] Assay Correlations

[0058] The term "correlating" as used herein in reference to the use of biomarkers refers to comparing the presence or amount of the biomarker(s) in a patient to its presence or amount in persons known to suffer from, or known to be at risk of, a given condition; or in persons known to be free of a given condition. Often, this takes the form of comparing an assay result in the form of a biomarker concentration to a predetermined threshold selected to be indicative of the occurrence or nonoccurrence of a disease or the likelihood of some future outcome.

[0059] Selecting a diagnostic threshold involves, among other things, consideration of the probability of disease, distribution of true and false diagnoses at different test thresholds, and estimates of the consequences of treatment (or a failure to treat) based on the diagnosis. For example, when considering administering a specific therapy which is highly efficacious and has a low level of risk, few tests are needed because clinicians can accept substantial diagnostic uncertainty. On the other hand, in situations where treatment options are less effective and more risky, clinicians often need a higher degree of diagnostic certainty. Thus, cost/benefit analysis is involved in selecting a diagnostic threshold.

[0060] Suitable thresholds may be determined in a variety of ways. For example, one recommended diagnostic threshold for the diagnosis of acute myocardial infarction using cardiac troponin is the 97.5th percentile of the concentration seen in a normal population. Another method may be to look at serial samples from the same patient, where a prior "baseline" result is used to monitor for temporal changes in a biomarker level.

[0061] Population studies may also be used to select a decision threshold. Receiver Operating Characteristic ("ROC") arose from the field of signal detection theory developed during World War II for the analysis of radar images, and ROC analysis is often used to select a threshold able to best distinguish a "diseased" subpopulation from a "nondiseased" subpopulation. A false positive in this case occurs when the person tests positive, but actually does not have the disease. A false negative, on the other hand, occurs when the person tests negative, suggesting they are healthy, when they actually do

have the disease. To draw a ROC curve, the true positive rate (TPR) and false positive rate (FPR) are determined as the decision threshold is varied continuously. Since TPR is equivalent with sensitivity and FPR is equal to 1 - specificity, the ROC graph is sometimes called the sensitivity vs (1 - specificity) plot. A perfect test will have an area under the ROC curve of 1.0; a random test will have an area of 0.5. A threshold is selected to provide an acceptable level of specificity and sensitivity.

[0062] In this context, "diseased" is meant to refer to a population having one characteristic (the presence of a disease or condition or the occurrence of some outcome) and "nondiseased" is meant to refer to a population lacking the characteristic. While a single decision threshold is the simplest application of such a method, multiple decision thresholds may be used. For example, below a first threshold, the absence of disease may be assigned with relatively high confidence, and above a second threshold the presence of disease may also be assigned with relatively high confidence. Between the two thresholds may be considered indeterminate. This is meant to be exemplary in nature only.

[0063] In addition to threshold comparisons, other methods for correlating assay results to a patient classification (occurrence or nonoccurrence of disease, likelihood of an outcome, etc.) include decision trees, rule sets, Bayesian methods, and neural network methods. These methods can produce probability values representing the degree to which a patient belongs to one classification out of a plurality of classifications.

[0064] Measures of test accuracy may be obtained as described in Fischer *et al*, *Intensive Care Med.* 29: 1043-51, 2003, and used to determine the effectiveness of a given biomarker. These measures include sensitivity and specificity, predictive values, likelihood ratios, diagnostic odds ratios, and ROC curve areas. The area under the curve ("AUC") of a ROC plot is equal to the probability that a classifier will rank a randomly chosen positive instance higher than a randomly chosen negative one. The area under the ROC curve may be thought of as equivalent to the Mann-Whitney U test, which tests for the median difference between scores obtained in the two groups considered if the groups are of continuous data, or to the Wilcoxon test of ranks.

[0065] As discussed above, suitable tests may exhibit one or more of the following results on these various measures: a specificity of greater than 0.5, preferably at least 0.6, more preferably at least 0.7, still more preferably at least 0.8, even more preferably at least 0.9 and most preferably at least 0.95, with a corresponding sensitivity greater than

0.2, preferably greater than 0.3, more preferably greater than 0.4, still more preferably at least 0.5, even more preferably 0.6, yet more preferably greater than 0.7, still more preferably greater than 0.8, more preferably greater than 0.9, and most preferably greater than 0.95; a sensitivity of greater than 0.5, preferably at least 0.6, more preferably at least 0.7, still more preferably at least 0.8, even more preferably at least 0.9 and most preferably at least 0.95, with a corresponding specificity greater than 0.2, preferably greater than 0.3, more preferably greater than 0.4, still more preferably at least 0.5, even more preferably 0.6, yet more preferably greater than 0.7, still more preferably greater than 0.8, more preferably greater than 0.9, and most preferably greater than 0.95; at least 75% sensitivity, combined with at least 75% specificity; a ROC curve area of greater than 0.5, preferably at least 0.6, more preferably 0.7, still more preferably at least 0.8, even more preferably at least 0.9, and most preferably at least 0.95; an odds ratio different from 1, preferably at least about 2 or more or about 0.5 or less, more preferably at least about 3 or more or about 0.33 or less, still more preferably at least about 4 or more or about 0.25 or less, even more preferably at least about 5 or more or about 0.2 or less, and most preferably at least about 10 or more or about 0.1 or less; a positive likelihood ratio (calculated as sensitivity/(1 -specificity)) of greater than 1, at least 2, more preferably at least 3, still more preferably at least 5, and most preferably at least 10; and or a negative likelihood ratio (calculated as (1-sensitivity)/specificity) of less than 1, less than or equal to 0.5, more preferably less than or equal to 0.3, and most preferably less than or equal to 0.1

[0066] Additional clinical indicia may be combined with the appendicitis biomarker assay result(s) of the present invention. Other clinical indicia which may be combined with the appendicitis biomarker assay result(s) of the present invention includes demographic information (e.g., weight, sex, age, race), medical history (e.g., family history, type of surgery, pre-existing disease such as aneurism, congestive heart failure, preeclampsia, eclampsia, diabetes mellitus, hypertension, coronary artery disease, proteinuria, or renal insufficiency), risk scores (APACHE score, PREDICT score, TIMI Risk Score for UA/NSTEMI, Framingham Risk Score), etc.

[0067] Combining assay results/clinical indicia in this manner can comprise the use of multivariate logistical regression, loglinear modeling, neural network analysis, n-of-m analysis, decision tree analysis, etc. This list is not meant to be limiting.

[0068] Selecting a Treatment Regimen

[0069] Once a diagnosis is obtained, the clinician can readily select a treatment regimen that is compatible with the diagnosis. The skilled artisan is aware of appropriate treatments for numerous diseases discussed in relation to the methods of diagnosis described herein. See, e.g., Merck Manual of Diagnosis and Therapy, 17th Ed. Merck Research Laboratories, Whitehouse Station, NJ, 1999. In addition, since the methods and compositions described herein provide prognostic information, the markers of the present invention may be used to monitor a course of treatment. For example, improved or worsened prognostic state may indicate that a particular treatment is or is not efficacious.

[0070] One skilled in the art readily appreciates that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The examples provided herein are representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention.

[0071] Example 1. Immunoassay format

[0072] Analytes are measured using standard sandwich enzyme immunoassay techniques. A first antibody which binds the analyte is immobilized in wells of a 96 well polystyrene microplate. Analyte standards and test samples are pipetted into the appropriate wells and any analyte present is bound by the immobilized antibody. After washing away any unbound substances, a horseradish peroxidase-conjugated second antibody which binds the analyte is added to the wells, thereby forming sandwich complexes with the analyte (if present) and the first antibody. Following a wash to remove any unbound antibody-enzyme reagent, a substrate solution comprising tetramethylbenzidine and hydrogen peroxide is added to the wells. Color develops in proportion to the amount of analyte present in the sample. The color development is stopped and the intensity of the color is measured at 540 nm or 570 nm. An analyte concentration is assigned to the test sample by comparison to a standard curve determined from the analyte standards.

[0073] Example 2. Use of analyte as a marker for assessing patients for appendicitis

[0074] Patients from the emergency department (ED) classified as positive for appendicitis according to clinical diagnosis at the study site were selected as a "diseased" cohort ("Cohort 1"). Plasma and urine samples from each patient in Cohort 1 were

collected at the time of appendectomy. Cohort 2 was a normal population sampled separately. The concentrations of the analyte in these samples were measured by standard immunoassay methods using commercially available assay reagents. A receiver operating characteristic (ROC) curve was generated using the concentrations, and the performance of the analyte is assessed by the area under the ROC curve (AUC). The two-tailed *p*-value of the AUC for the analyte was also calculated to determine statistical significance. "Inc/Dec" indicates if the marker is increasing or decreasing in Cohort 1 relative to Cohort 2.

Table 1: Urine samples

Preferred Name	Units	Increasing/ Decreasing	p
Adiponectin	ng/ml	inc	8.68E-07
Advanced glycosylation end product-specific receptor	pg/ml	dec	2.82E-01
Alpha-2 macroglobulin	ug/ml	inc	1.51E-04
Alpha-2-HS-glycoprotein	ng/ml	inc	1.97E-02
Angiotensinogen-converting enzyme 1	ng/ml	dec	7.54E-01
Antileukoprotease	pg/ml	dec	9.51E-01
Apolipoprotein(a)	ng/ml	dec	6.85E-05
Carbonic anhydrase 9	ng/ml	inc	3.80E-01
C-C motif chemokine 1	pg/ml	inc	7.88E-01
C-C motif chemokine 13	pg/ml	dec	3.19E-01
C-C motif chemokine 15	pg/ml	dec	1.45E-01
C-C motif chemokine 17	pg/ml	inc	6.55E-01
C-C motif chemokine 19	pg/ml	dec	5.62E-02
C-C motif chemokine 20	pg/ml	dec	2.49E-01
C-C motif chemokine 21	pg/ml	inc	7.47E-01
C-C motif chemokine 22	pg/ml	inc	7.81E-01
C-C motif chemokine 23	ng/ml	inc	5.63E-01
C-C motif chemokine 24	pg/ml	inc	3.00E-02
C-C motif chemokine 26	pg/ml	dec	6.44E-01
C-C motif chemokine 27	pg/ml	inc	3.18E-01
C-C motif chemokine 3	pg/ml	dec	3.36E-01
C-C motif chemokine 4	pg/ml	inc	6.33E-01
C-C motif chemokine 7	pg/ml	inc	6.44E-01
C-C motif chemokine 8	pg/ml	inc	8.29E-01
Ceruloplasmin	ng/ml	dec	8.02E-02
C-Peptide	pg/ml	dec	3.35E-04
Creatine Kinase-MB	ng/ml	inc	7.18E-01
C-X-C motif chemokine 10	pg/ml	inc	6.59E-02
C-X-C motif chemokine 11	pg/ml	inc	6.35E-01
C-X-C motif chemokine 13	pg/ml	dec	6.57E-01
C-X-C motif chemokine 16	ng/ml	inc	4.61E-01

Preferred Name	Units	Increasing/ Decreasing	P
C-X-C motif chemokine 5	pg/ml	inc	8.20E-01
C-X-C motif chemokine 6	pg/ml	inc	8.93E-06
C-X-C motif chemokine 9	pg/ml	inc	2.25E-01
Cystatin-C	ng/ml	dec	2.01E-01
Endothelial protein C receptor	ng/ml	inc	1.26E-02
Eotaxin	pg/ml	inc	7.99E-01
Epidermal growth factor receptor	pg/ml	inc	7.11E-01
Ferritin	pg/ml	inc	5.59E-06
Fibrinogen	ng/ml	inc	7.56E-01
Gastric inhibitory polypeptide	pg/ml	inc	6.44E-01
Glucagon	pg/ml	dec	8.18E-01
Glucagon-like peptide 1	pg/ml	inc	8.18E-01
Granulocyte-macrophage colony-stimulating factor	pg/ml	dec	9.87E-02
Heparin-binding growth factor 2	pg/ml	dec	7.31E-01
Hepatocyte growth factor	ng/ml	dec	6.68E-01
Immunoglobulin A	ng/ml	inc	6.83E-01
Immunoglobulin M	ng/ml	inc	2.64E-08
Immunoglobulin G1	ng/ml	dec	1.37E-02
Immunoglobulin G2	ng/ml	dec	2.19E-01
Immunoglobulin G3	ng/ml	dec	7.25E-01
Immunoglobulin G4	ng/ml	inc	4.56E-03
Insulin	pg/ml	dec	4.31E-02
Insulin-like growth factor-binding protein 1	ng/ml	dec	2.00E-03
Insulin-like growth factor-binding protein 2	ng/ml	dec	0.00E+00
Insulin-like growth factor-binding protein 3	ng/ml	dec	1.96E-01
Insulin-like growth factor-binding protein 4	ng/ml	inc	8.52E-01
Insulin-like growth factor-binding protein 5	ng/ml	dec	6.96E-01
Insulin-like growth factor-binding protein 6	ng/ml	inc	1.17E-01
Insulin-like growth factor-binding protein 7	ng/ml	inc	2.36E-01
Interferon alpha-2	pg/ml	dec	2.86E-01
Interferon gamma	pg/ml	inc	8.31E-01
Interleukin-1 alpha	pg/ml	dec	5.10E-01
Interleukin-1 beta	pg/ml	dec	9.67E-01
Interleukin-1 receptor antagonist protein	pg/ml	dec	4.62E-02
Interleukin-1 receptor type I	pg/mL	inc	3.16E-01
Interleukin-1 receptor type II	pg/mL	inc	9.27E-01
Interleukin-11	pg/ml	dec	7.53E-03
Interleukin-1 2	pg/ml	inc	8.48E-01
Interleukin-1 2 subunit beta	ng/ml	dec	5.34E-02
Interleukin-1 3	pg/ml	dec	8.78E-01
Interleukin-1 5	pg/ml	dec	2.31E-01
Interleukin-2	pg/ml	dec	5.87E-01
Interleukin-20	pg/ml	dec	3.22E-01
Interleukin-2 1	pg/ml	dec	4.17E-01

Preferred Name	Units	Increasing/ Decreasing	p
Interleukin-23	pg/ml	inc	6.19E--01
Interleukin-28A	pg/ml	inc	4.52E--01
Interleukin-29	pg/ml	inc	3.74E--02
Interleukin-3	pg/ml	inc	3.48E--01
Interleukin-33	pg/ml	inc	2.14E--01
Interleukin-4	pg/ml	inc	5.71E--01
Interleukin-5	ng/ml	inc	8.08E--01
Interleukin-6 receptor subunit alpha	pg/ml	inc	6.70E--01
Interleukin-6 receptor subunit beta	pg/ml	dec	6.63E--01
Interleukin-7	pg/ml	dec	8.59E--01
Interleukin-9	pg/ml	inc	4.32E--01
Islet amyloid polypeptide	pg/ml	dec	8.78E--01
Kit ligand	pg/ml	dec	3.64E--01
Leptin	pg/ml	dec	3.21E--01
Leukemia inhibitory factor	pg/ml	dec	8.27E--01
Lymphotactin	pg/ml	inc	4.48E--01
Lymphotoxin- alpha	pg/ml	dec	6.44E--01
Macrophage colony-stimulating factor 1	pg/ml	inc	1.01E--01
Metalloproteinase inhibitor 1	pg/ml	inc	7.52E--01
Metalloproteinase inhibitor 2	pg/ml	inc	1.78E--01
Metalloproteinase inhibitor 3	pg/ml	inc	8.45E--01
Metalloproteinase inhibitor 4	pg/ml	dec	1.62E--01
Mix of Growth-regulated alpha, beta, and gamma proteins	pg/ml	inc	4.65E--01
Myoglobin	ng/ml	dec	1.14E--02
Pancreatic prohormone	pg/ml	dec	5.89E--01
Peptide YY	pg/ml	dec	8.18E--01
Pro-epidermal growth factor	pg/ml	inc	5.02E--02
Pro-interleukin- 16	pg/ml	dec	9.39E--01
Protein S100-A12	ng/ml	dec	2.49E--02
Protransforming growth factor alpha	pg/ml	inc	5.62E--02
Secretory immunoglobulin A	ng/ml	inc	9.18E--01
Serum amyloid P-component	ng/ml	inc	1.02E--04
SL cytokine	pg/ml	dec	7.11E--01
Stromal cell-derived factor 1	pg/ml	dec	7.03E--02
Thrombopoietin	pg/ml	dec	1.15E--02
Thymic stromal lymphopoietin	pg/ml	inc	1.81E--03
Tumor necrosis factor	pg/ml	dec	8.29E--01
Tumor necrosis factor ligand superfamily member 10	pg/ml	inc	1.24E--01
Tumor necrosis factor receptor superfamily member 1A	pg/ml	dec	5.55E--03
Tumor necrosis factor receptor superfamily member 1B	pg/ml	dec	4.11E--02
Vascular endothelial growth factor receptor 1	pg/ml	dec	1.11E--01

Preferred Name	Units	Increasing/ Decreasing	P
Vascular endothelial growth factor receptor 2	pg/ml	dec	6.61E-01
Vascular endothelial growth factor receptor 3	pg/ml	dec	9.11E-01
von Willebrand Factor	ng/ml	inc	2.52E-04

Table 2: Plasma samples

Preferred Name	Units	Increasing/ Decreasing	P
72 kDa type IV collagenase	pg/ml	inc	1.73E-14
Adiponectin	ng/ml	inc	8.92E-01
Advanced glycosylation end product-specific receptor	pg/ml	inc	3.24E-01
Alpha-2 macroglobulin	ug/ml	dec	1.52E-01
Alpha-2-HS-glycoprotein	ng/ml	inc	4.26E-04
Alpha-fetoprotein	pg/ml	inc	8.85E-01
Angiopoietin- 1	ng/ml	inc	1.75E-02
Antileukoproteinase	pg/ml	dec	3.89E-01
Apolipoprotein(a)	ng/ml	inc	3.51E-01
Cancer Antigen 15-3	U/ml	inc	9.18E-06
Cancer Antigen 19-9	U/ml	inc	6.13E-01
Carbonic anhydrase 9	ng/ml	inc	3.78E-02
Carcinoembryonic antigen-related cell adhesion molecule 5	pg/ml	dec	9.44E-01
C-C motif chemokine 1	pg/ml	dec	1.79E-01
C-C motif chemokine 13	pg/ml	inc	5.89E-03
C-C motif chemokine 15	pg/ml	dec	1.67E-01
C-C motif chemokine 17	pg/ml	inc	1.69E-01
C-C motif chemokine 19	pg/ml	inc	2.24E-03
C-C motif chemokine 20	pg/ml	dec	4.43E-01
C-C motif chemokine 21	pg/ml	dec	1.43E-02
C-C motif chemokine 22	pg/ml	dec	3.28E-06
C-C motif chemokine 23	ng/ml	inc	1.25E-08
C-C motif chemokine 24	pg/ml	inc	3.32E-01
C-C motif chemokine 26	pg/ml	inc	1.54E-01
C-C motif chemokine 27	pg/ml	dec	1.44E-02
C-C motif chemokine 3	pg/ml	dec	1.81E-01
C-C motif chemokine 4	pg/ml	inc	3.44E-04
C-C motif chemokine 7	pg/ml	inc	8.34E-01
C-C motif chemokine 8	pg/ml	dec	9.23E-01
Ceruloplasmin	ng/ml	dec	9.62E-02
Choriogonadotropin subunit beta	mU/ml	inc	2.52E-02
Collagenase 3	pg/ml	dec	4.64E-01
C-Peptide	pg/ml	dec	2.52E-05
Creatine Kinase-MB	ng/ml	inc	3.37E-01
C-X-C motif chemokine 10	pg/ml	inc	1.08E-01

Preferred Name	Units	Increasing/ Decreasing	P
C-X-C motif chemokine 11	pg/ml	dec	6.95E-02
C-X-C motif chemokine 13	pg/ml	dec	5.59E-03
C-X-C motif chemokine 16	ng/ml	dec	1.47E-02
C-X-C motif chemokine 5	pg/ml	dec	9.76E-01
C-X-C motif chemokine 6	pg/ml	inc	1.04E-01
C-X-C motif chemokine 9	pg/ml	inc	4.93E-01
Cystatin-C	ng/ml	inc	8.02E-07
Endothelial protein C receptor	ng/ml	dec	3.19E-01
Eotaxin	pg/ml	inc	1.20E-07
Epidermal growth factor receptor	pg/ml	dec	6.80E-01
Ferritin	pg/ml	dec	1.92E-04
Fibrinogen	ug/ml	dec	5.85E-02
Gastric inhibitory polypeptide	pg/ml	dec	0.00E+00
Glucagon	pg/ml	dec	8.82E-01
Glucagon-like peptide 1	pg/ml	dec	6.75E-04
Granulocyte-macrophage colony-stimulating factor	pg/ml	inc	2.07E-02
Heparin-binding growth factor 2	pg/ml	inc	7.22E-01
Hepatocyte growth factor	ng/ml	inc	0.00E+00
Immunoglobulin A	ng/ml	dec	4.10E-03
Immunoglobulin M	ng/ml	dec	5.79E-01
Immunoglobulin G1	ng/ml	dec	4.93E-01
Immunoglobulin G2	ng/ml	dec	3.03E-01
Immunoglobulin G3	ng/ml	dec	2.08E-04
Immunoglobulin G4	ng/ml	dec	4.64E-01
Insulin	pg/ml	dec	8.32E-05
Insulin-like growth factor-binding protein 1	ng/ml	dec	2.19E-10
Insulin-like growth factor-binding protein 2	ng/ml	dec	8.69E-01
Insulin-like growth factor-binding protein 3	ng/ml	inc	1.75E-01
Insulin-like growth factor-binding protein 4	ng/ml	dec	5.07E-01
Insulin-like growth factor-binding protein 5	ng/ml	dec	9.25E-04
Insulin-like growth factor-binding protein 6	ng/ml	inc	1.68E-07
Insulin-like growth factor-binding protein 7	ng/ml	dec	7.07E-05
Interferon alpha-2	pg/ml	inc	1.55E-01
Interferon gamma	pg/ml	inc	1.04E-02
Interleukin-1 alpha	pg/ml	dec	8.53E-01
Interleukin-1 beta	pg/ml	dec	1.45E-01
Interleukin-1 receptor type I	pg/mL	dec	1.60E-03
Interleukin-1 receptor type II	pg/mL	inc	7.29E-01
Interleukin-11	pg/ml	inc	5.15E-01
Interleukin-1 2	pg/ml	inc	5.94E-01
Interleukin-1 2 subunit beta	ng/ml	dec	9.09E-01
Interleukin-1 3	pg/ml	inc	4.61E-01
Interleukin-1 5	pg/ml	dec	7.13E-02
Interleukin-2	pg/ml	dec	1.90E-03

Preferred Name	Units	Increasing/ Decreasing	P
Interleukin-20	pg/ml	inc	5.51E-01
Interleukin-21	pg/ml	dec	9.01E-01
Interleukin-23	pg/ml	dec	5.67E-01
Interleukin-28A	pg/ml	dec	4.44E-01
Interleukin-29	pg/ml	dec	3.04E-01
Interleukin-3	pg/ml	inc	6.55E-01
Interleukin-33	pg/ml	dec	5.24E-01
Interleukin-4	pg/ml	dec	4.79E-01
Interleukin-5	ng/ml	inc	2.65E-01
Interleukin-6 receptor subunit alpha	pg/ml	dec	2.88E-01
Interleukin-6 receptor subunit beta	pg/ml	dec	5.47E-01
Interleukin-7	pg/ml	inc	7.57E-01
Interleukin-9	pg/ml	dec	5.16E-01
Interstitial collagenase	pg/ml	dec	1.78E-05
Islet amyloid polypeptide	pg/ml	inc	2.96E-07
Keratin, type I cytoskeletal 19 (aa3 11-367)	pg/ml	dec	9.39E-01
Kit ligand	pg/ml	inc	8.87E-01
Leptin	pg/ml	dec	8.44E-01
Leukemia inhibitory factor	pg/ml	dec	3.49E-01
Lymphotactin	pg/ml	dec	5.21E-02
Lymphotoxin- alpha	pg/ml	inc	6.48E-01
Macrophage colony-stimulating factor 1	pg/ml	dec	7.41E-04
Macrophage metalloelastase	pg/ml	inc	8.74E-01
Macrophage migration inhibitory factor	pg/ml	inc	5.75E-06
Matrilysin	pg/ml	dec	1.59E-01
Metalloproteinase inhibitor 1	pg/ml	dec	9.56E-01
Metalloproteinase inhibitor 2	pg/ml	inc	4.10E-09
Metalloproteinase inhibitor 3	pg/ml	inc	2.28E-01
Metalloproteinase inhibitor 4	pg/ml	inc	4.21E-01
Mix of Growth-regulated alpha, beta, and gamma proteins	pg/ml	inc	2.33E-01
Myoglobin	ng/ml	dec	5.72E-01
Pancreatic prohormone	pg/ml	inc	2.44E-13
Peptide YY	pg/ml	inc	1.32E-03
Pro-epidermal growth factor	pg/ml	inc	1.23E-02
Pro-interleukin- 16	pg/ml	inc	3.21E-01
Prolactin	pg/ml	dec	2.55E-11
Prostate-specific antigen	pg/ml	dec	7.49E-01
Protein S100-A12	ng/ml	dec	2.02E-04
Protransforming growth factor alpha	pg/ml	inc	3.50E-02
Serum amyloid P-component	ng/ml	dec	4.77E-01
SL cytokine	pg/ml	dec	1.27E-01
Stromal cell-derived factor 1	pg/ml	inc	8.60E-01
Stromelysin- 1	pg/ml	dec	2.76E-06

Preferred Name	Units	Increasing/ Decreasing	p
Thrombopoietin	pg/ml	inc	1.76E-04
Thymic stromal lymphopoietin	pg/ml	dec	5.75E-01
Tumor necrosis factor	pg/ml	dec	1.15E-01
Tumor necrosis factor ligand superfamily member 10	pg/ml	inc	1.03E-06
Tumor necrosis factor ligand superfamily member 6	pg/ml	dec	3.96E-01
Tumor necrosis factor receptor superfamily member 1A	pg/ml	dec	1.20E-07
Tumor necrosis factor receptor superfamily member 1B	pg/ml	dec	1.03E-02
Vascular endothelial growth factor receptor 1	pg/ml	inc	6.04E-07
Vascular endothelial growth factor receptor 2	pg/ml	dec	2.63E-03
Vascular endothelial growth factor receptor 3	pg/ml	inc	8.51E-01
von Willebrand Factor	ng/ml	dec	8.01E-01
WAP four-disulfide core domain protein 2	pg/ml	dec	4.13E-04

[0075] While the invention has been described and exemplified in sufficient detail for those skilled in this art to make and use it, various alternatives, modifications, and improvements should be apparent without departing from the spirit and scope of the invention. The examples provided herein are representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Modifications therein and other uses will occur to those skilled in the art. These modifications are encompassed within the spirit of the invention and are defined by the scope of the claims.

[0076] It will be readily apparent to a person skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention.

[0077] All patents and publications mentioned in the specification are indicative of the levels of those of ordinary skill in the art to which the invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

[0078] The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations which is not specifically disclosed herein. Thus, for example, in each instance herein any of the terms "comprising", "consisting essentially of" and "consisting of" may be replaced with either

of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

[0079] Other embodiments are set forth within the following claims.

We claim:

1. A method of diagnosing appendicitis in a subject, or assigning a likelihood of a future outcome to a subject diagnosed with appendicitis, comprising:

performing one or more assays configured to detect one or more biomarkers selected from the group consisting of 72 kDa type IV collagenase, Adiponectin, Advanced glycosylation end product-specific receptor, Alpha-2 macroglobulin, Alpha-2-HS-glycoprotein, Alpha-fetoprotein, Angiopoietin-1, Antileukoprotease, Apolipoprotein(a), Cancer Antigen 15-3, Cancer Antigen 19-9, Carbonic anhydrase 9, Carcinoembryonic antigen-related cell adhesion molecule 5, C-C motif chemokine 1, C-C motif chemokine 13, C-C motif chemokine 15, C-C motif chemokine 17, C-C motif chemokine 19, C-C motif chemokine 20, C-C motif chemokine 21, C-C motif chemokine 22, C-C motif chemokine 23, C-C motif chemokine 24, C-C motif chemokine 26, C-C motif chemokine 27, C-C motif chemokine 3, C-C motif chemokine 4, C-C motif chemokine 7, C-C motif chemokine 8, Ceruloplasmin, Choriogonadotropin subunit beta, Collagenase 3, C-Peptide, Creatine Kinase-MB, C-X-C motif chemokine 10, C-X-C motif chemokine 11, C-X-C motif chemokine 13, C-X-C motif chemokine 16, C-X-C motif chemokine 5, C-X-C motif chemokine 6, C-X-C motif chemokine 9, Cystatin-C, Endothelial protein C receptor, Eotaxin, Epidermal growth factor receptor, Ferritin, Fibrinogen, Gastric inhibitory polypeptide, Glucagon, Glucagon-like peptide 1, Granulocyte-macrophage colony-stimulating factor, Growth-regulated alpha, beta, and gamma proteins, Heparin-binding growth factor 2, Hepatocyte growth factor, Immunoglobulin A, Immunoglobulin M, Immunoglobulin G1, Immunoglobulin G2, Immunoglobulin G3, Immunoglobulin G4, Insulin, Insulin-like growth factor-binding protein 1, Insulin-like growth factor-binding protein 2, Insulin-like growth factor-binding protein 3, Insulin-like growth factor-binding protein 4, Insulin-like growth factor-binding protein 5, Insulin-like growth factor-binding protein 6, Insulin-like growth factor-binding protein 7, Interferon alpha-2, Interferon gamma, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 receptor antagonist protein, Interleukin-1 receptor type I, Interleukin-1 receptor type II, Interleukin-1 1, Interleukin-12, Interleukin-12 subunit beta, Interleukin-13, Interleukin-15, Interleukin-2, Interleukin-20, Interleukin-21, Interleukin-23, Interleukin-28A, Interleukin-29, Interleukin-3, Interleukin-33, Interleukin-4, Interleukin-5, Interleukin-6 receptor subunit alpha, Interleukin-6 receptor subunit beta, Interleukin-7, Interleukin-9, Interstitial collagenase, Islet amyloid polypeptide, Keratin, type I cytoskeletal 19 (aa311-367), Kit ligand, Leptin,

Leukemia inhibitory factor, Lymphotoxin, Lymphotoxin-alpha, Macrophage colony-stimulating factor 1, Macrophage metalloelastase, Macrophage migration inhibitory factor, Matrilysin, Metalloproteinase inhibitor 1, Metalloproteinase inhibitor 2, Metalloproteinase inhibitor 3, Metalloproteinase inhibitor 4, Myoglobin, Pancreatic prohormone, Peptide YY, Pro-epidermal growth factor, Pro-interleukin-16, Prolactin, Prostate-specific antigen, Protein S100-A12, Protransforming growth factor alpha, Secretory immunoglobulin A, Serum amyloid P-component, SL cytokine, Stromal cell-derived factor 1, Stromelysin-1, Thrombopoietin, Thymic stromal lymphopoietin, Tumor necrosis factor, Tumor necrosis factor ligand superfamily member 10, Tumor necrosis factor ligand superfamily member 6, Tumor necrosis factor receptor superfamily member 1A, Tumor necrosis factor receptor superfamily member 1B, Vascular endothelial growth factor receptor 1, Vascular endothelial growth factor receptor 2, Vascular endothelial growth factor receptor 3, von Willebrand Factor, and WAP four-disulfide core domain protein 2 on a body fluid sample obtained from the subject to provide one or more assay result(s); and

correlating the assay result(s) to the occurrence or nonoccurrence of appendicitis in the subject or likelihood of the future outcome to the subject.

2. A method according to claim 1, wherein the performing step comprises introducing the body fluid sample obtained from the subject into an assay instrument which (i) contacts the body fluid sample with one or more binding reagents corresponding to the biomarker(s) being assayed, wherein each biomarker which is assayed binds to its respective specific binding reagent in an amount related to its concentration in the body fluid sample, (ii) generates one or more assay results indicative of binding of each biomarker which is assayed to its respective specific binding reagent; and (iii) displays the one or more assay results as a quantitative result in a human-readable form.
3. A method according to claim 1 or 2, wherein the future outcome is mortality.
4. A method according to claim 1 or 2, wherein the subject is being evaluated for abdominal pain.

5. A method according to claim 1 or 2, wherein the correlating step comprises determining the concentration of each biomarker which is assayed, and individually comparing each biomarker concentration to a corresponding threshold level for that biomarker.
6. A method according to claim 5, wherein the assay instrument performs the correlating step, which comprises determining the concentration of each biomarker which is assayed, individually comparing each biomarker concentration to a corresponding threshold level for that biomarker, and displaying an indication of whether each biomarker does or does not exceed its corresponding threshold in a human-readable form.
7. A method according to claim 2, wherein a plurality of the biomarkers are measured, wherein the assay instrument performs the correlating step, which comprises determining the concentration of each of the plurality of biomarkers, calculating a single value based on the concentration of each of the plurality of biomarkers, comparing the single value to a corresponding threshold level and displaying an indication of whether the single value does or does not exceed its corresponding threshold in a human-readable form.
8. A method according to one of claims 1-7, wherein method provides a sensitivity or specificity of at least 0.7 for the identification of appendicitis when compared to normal subjects.
9. A method according to one of claims 1-7, wherein method provides a sensitivity or specificity of at least 0.7 for the identification of appendicitis when compared to subjects exhibiting symptoms that mimic appendicitis symptoms.
10. A method according to one of claims 1-9, wherein the sample is selected from the group consisting of blood, serum, and plasma.

11. A method for evaluating biomarker levels in a body fluid sample, comprising: obtaining a body fluid sample from a subject selected for evaluation based on a determination that the subject is experiencing symptoms indicative of possible acute appendicitis; and performing one or more analyte binding assays configured to detect one or more biomarkers selected from the group consisting of 72 kDa type IV collagenase, Adiponectin, Advanced glycosylation end product-specific receptor, Alpha-2 macro globulin, Alpha-2-HS-glycoprotein, Alpha-fetoprotein, Angiopoietin-1, Antileukoproteinase, Apolipoprotein(a), Cancer Antigen 15-3, Cancer Antigen 19-9, Carbonic anhydrase 9, Carcinoembryonic antigen-related cell adhesion molecule 5, C-C motif chemokine 1, C-C motif chemokine 13, C-C motif chemokine 15, C-C motif chemokine 17, C-C motif chemokine 19, C-C motif chemokine 20, C-C motif chemokine 21, C-C motif chemokine 22, C-C motif chemokine 23, C-C motif chemokine 24, C-C motif chemokine 26, C-C motif chemokine 27, C-C motif chemokine 3, C-C motif chemokine 4, C-C motif chemokine 7, C-C motif chemokine 8, Ceruloplasmin, Choriogonadotropin subunit beta, Collagenase 3, C-Peptide, Creatine Kinase-MB, C-X-C motif chemokine 10, C-X-C motif chemokine 11, C-X-C motif chemokine 13, C-X-C motif chemokine 16, C-X-C motif chemokine 5, C-X-C motif chemokine 6, C-X-C motif chemokine 9, Cystatin-C, Endothelial protein C receptor, Eotaxin, Epidermal growth factor receptor, Ferritin, Fibrinogen, Gastric inhibitory polypeptide, Glucagon, Glucagon-like peptide 1, Granulocyte-macrophage colony-stimulating factor, Growth-regulated alpha, beta, and gamma proteins, Heparin-binding growth factor 2, Hepatocyte growth factor, Immunoglobulin A, Immunoglobulin M, Immunoglobulin G1, Immunoglobulin G2, Immunoglobulin G3, Immunoglobulin G4, Insulin, Insulin-like growth factor-binding protein 1, Insulin-like growth factor-binding protein 2, Insulin-like growth factor-binding protein 3, Insulin-like growth factor-binding protein 4, Insulin-like growth factor-binding protein 5, Insulin-like growth factor-binding protein 6, Insulin-like growth factor-binding protein 7, Interferon alpha-2, Interferon gamma, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 receptor antagonist protein, Interleukin-1 receptor type I, Interleukin-1 receptor type II, Interleukin-1 1, Interleukin-12, Interleukin-12 subunit beta, Interleukin-13, Interleukin-15, Interleukin-2, Interleukin-20, Interleukin-21, Interleukin-23, Interleukin-28A, Interleukin-29, Interleukin-3, Interleukin-33, Interleukin-4, Interleukin-5, Interleukin-6 receptor subunit alpha, Interleukin-6 receptor subunit beta, Interleukin-7,

Interleukin-9, Interstitial collagenase, Islet amyloid polypeptide, Keratin, type I cytoskeletal 19 (aa3 11-367), Kit ligand, Leptin, Leukemia inhibitory factor, Lymphotoxin, Lymphotoxin- α , Macrophage colony-stimulating factor 1, Macrophage metalloelastase, Macrophage migration inhibitory factor, Matrilysin, Metalloproteinase inhibitor 1, Metalloproteinase inhibitor 2, Metalloproteinase inhibitor 3, Metalloproteinase inhibitor 4, Myoglobin, Pancreatic prohormone, Peptide YY, Pro-epidermal growth factor, Pro-interleukin-16, Prolactin, Prostate-specific antigen, Protein S100-A12, Protransforming growth factor α , Secretory immunoglobulin A, Serum amyloid P-component, SL cytokine, Stromal cell-derived factor 1, Stromelysin-1, Thrombopoietin, Thymic stromal lymphopoietin, Tumor necrosis factor, Tumor necrosis factor ligand superfamily member 10, Tumor necrosis factor ligand superfamily member 6, Tumor necrosis factor receptor superfamily member 1A, Tumor necrosis factor receptor superfamily member 1B, Vascular endothelial growth factor receptor 1, Vascular endothelial growth factor receptor 2, Vascular endothelial growth factor receptor 3, von Willebrand Factor, and WAP four-disulfide core domain protein 2 by introducing the body fluid sample obtained from the subject into an assay instrument which (i) contacts the body fluid sample with one or more binding reagents corresponding to the biomarker(s) being assayed, wherein each biomarker which is assayed binds to its respective specific binding reagent in an amount related to its concentration in the body fluid sample, (ii) generates one or more assay results indicative of binding of each biomarker which is assayed to its respective specific binding reagent; and (iii) displays the one or more assay results as a quantitative result in a human-readable form.

12. A method according to claim 11, wherein the assay result(s) are displayed as a concentration of each biomarker which is assayed.

13. A method according to claim 12, wherein the assay instrument further individually compares each biomarker concentration to a corresponding threshold level for that biomarker, and displays an indication of whether each biomarker does or does not exceed its corresponding threshold in a human-readable form.

14. A method according to claim 11, wherein a plurality of the biomarkers are measured, and wherein the assay results(s) comprise a single value calculated using a function that converts the concentration of each of the plurality of biomarkers into a single value.

15. A method according to claim 14, wherein the assay instrument further compares the single value to a corresponding threshold level and displays an indication of whether the single value does or does not exceed its corresponding threshold in a human-readable form.

16. A method for evaluating biomarker levels in a body fluid sample, comprising: obtaining a body fluid sample from a subject selected for evaluation based on a determination that the subject has been diagnosed with acute appendicitis; and performing one or more analyte binding assays configured to detect one or more biomarkers selected from the group consisting of 72 kDa type IV collagenase, Adiponectin, Advanced glycosylation end product-specific receptor, Alpha-2 macroglobulin, Alpha-2-HS-glycoprotein, Alpha-fetoprotein, Angiotensin-1, Antileukoproteinase, Apolipoprotein(a), Cancer Antigen 15-3, Cancer Antigen 19-9, Carbonic anhydrase 9, Carcinoembryonic antigen-related cell adhesion molecule 5, C-C motif chemokine 1, C-C motif chemokine 13, C-C motif chemokine 15, C-C motif chemokine 17, C-C motif chemokine 19, C-C motif chemokine 20, C-C motif chemokine 21, C-C motif chemokine 22, C-C motif chemokine 23, C-C motif chemokine 24, C-C motif chemokine 26, C-C motif chemokine 27, C-C motif chemokine 3, C-C motif chemokine 4, C-C motif chemokine 7, C-C motif chemokine 8, Ceruloplasmin, Choriogonadotropin subunit beta, Collagenase 3, C-Peptide, Creatine Kinase-MB, C-X-C motif chemokine 10, C-X-C motif chemokine 11, C-X-C motif chemokine 13, C-X-C motif chemokine 16, C-X-C motif chemokine 5, C-X-C motif chemokine 6, C-X-C motif chemokine 9, Cystatin-C, Endothelial protein C receptor, Eotaxin, Epidermal growth factor receptor, Ferritin, Fibrinogen, Gastric inhibitory polypeptide, Glucagon, Glucagon-like peptide 1, Granulocyte-macrophage colony-stimulating factor, Growth-regulated alpha, beta, and gamma proteins, Heparin-binding growth factor 2, Hepatocyte growth factor, Immunoglobulin A, Immunoglobulin M, Immunoglobulin G1, Immunoglobulin G2, Immunoglobulin G3, Immunoglobulin G4, Insulin, Insulin-like growth factor-binding protein 1, Insulin-like growth factor-binding protein 2, Insulin-like growth factor-binding protein 3, Insulin-like growth factor-binding protein 4, Insulin-like growth factor-binding protein 5, Insulin-like growth factor-binding protein 6, Insulin-like growth factor-binding protein 7, Interferon alpha-2, Interferon gamma, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 receptor antagonist protein, Interleukin-1 receptor type I, Interleukin-1 receptor type II, Interleukin-1 1, Interleukin-12, Interleukin-12 subunit beta, Interleukin-

13, Interleukin-15, Interleukin-2, Interleukin-20, Interleukin-21, Interleukin-23, Interleukin-28A, Interleukin-29, Interleukin-3, Interleukin-33, Interleukin-4, Interleukin-5, Interleukin-6 receptor subunit alpha, Interleukin-6 receptor subunit beta, Interleukin-7, Interleukin-9, Interstitial collagenase, Islet amyloid polypeptide, Keratin, type I cytoskeletal 19 (aa3 11-367), Kit ligand, Leptin, Leukemia inhibitory factor, Lymphotoxin, Lymphotoxin-alpha, Macrophage colony-stimulating factor 1, Macrophage metalloelastase, Macrophage migration inhibitory factor, Matrilysin, Metalloproteinase inhibitor 1, Metalloproteinase inhibitor 2, Metalloproteinase inhibitor 3, Metalloproteinase inhibitor 4, Myoglobin, Pancreatic prohormone, Peptide YY, Pro-epidermal growth factor, Pro-interleukin-16, Prolactin, Prostate-specific antigen, Protein S100-A12, Protransforming growth factor alpha, Secretory immunoglobulin A, Serum amyloid P-component, SL cytokine, Stromal cell-derived factor 1, Stromelysin-1, Thrombopoietin, Thymic stromal lymphopoietin, Tumor necrosis factor, Tumor necrosis factor ligand superfamily member 10, Tumor necrosis factor ligand superfamily member 6, Tumor necrosis factor receptor superfamily member 1A, Tumor necrosis factor receptor superfamily member 1B, Vascular endothelial growth factor receptor 1, Vascular endothelial growth factor receptor 2, Vascular endothelial growth factor receptor 3, von Willebrand Factor, and WAP four-disulfide core domain protein 2 by introducing the body fluid sample obtained from the subject into an assay instrument which (i) contacts the body fluid sample with one or more binding reagents corresponding to the biomarker(s) being assayed, wherein each biomarker which is assayed binds to its respective specific binding reagent in an amount related to its concentration in the body fluid sample, (ii) generates one or more assay results indicative of binding of each biomarker which is assayed to its respective specific binding reagent; and (iii) displays the one or more assay results as a quantitative result in a human-readable form.

17. A method according to claim 16, wherein the assay result(s) are displayed as a concentration of each biomarker which is assayed.

18. A method according to claim 17, wherein the assay instrument further individually compares each biomarker concentration to a corresponding threshold level for that biomarker, and displays an indication of whether each biomarker does or does not exceed its corresponding threshold in a human-readable form.

19. A method according to claim 16, wherein a plurality of the biomarkers are measured, and wherein the assay results(s) comprise a single value calculated using a function that converts the concentration of each of the plurality of biomarkers into a single value.
20. A method according to claim 19, wherein the assay instrument further compares the single value to a corresponding threshold level and displays an indication of whether the single value does or does not exceed its corresponding threshold in a human-readable form.
21. A method according to one of claims 16-19, wherein the subject is selected for evaluation of a mortality risk within a period selected from the group consisting of 21 days, 14 days, 7 days, 5 days, 96 hours, 72 hours, 48 hours, 36 hours, 24 hours, and 12 hours.
22. A method according to one of claims 11-21, wherein the plurality of assays are immunoassays performed by (i) introducing the body fluid sample into an assay device comprising a plurality of antibodies, at least one of which binds to each biomarker which is assayed, and (ii) generating an assay result indicative of binding of each biomarker to its respective antibody.
23. A system for evaluating biomarker levels, comprising:
a plurality of reagents which specifically bind for detection a plurality of biomarkers selected from the group consisting of 72 kDa type IV collagenase, Adiponectin, Advanced glycosylation end product-specific receptor, Alpha-2 macroglobulin, Alpha-2-HS-glycoprotein, Alpha-fetoprotein, Angiopoietin-1, Antileukoproteinasase, Apolipoprotein(a), Cancer Antigen 15-3, Cancer Antigen 19-9, Carbonic anhydrase 9, Carcinoembryonic antigen-related cell adhesion molecule 5, C-C motif chemokine 1, C-C motif chemokine 13, C-C motif chemokine 15, C-C motif chemokine 17, C-C motif chemokine 19, C-C motif chemokine 20, C-C motif chemokine 21, C-C motif chemokine 22, C-C motif chemokine 23, C-C motif chemokine 24, C-C motif chemokine 26, C-C motif chemokine 27, C-C motif chemokine 3, C-C motif chemokine 4, C-C motif chemokine 7, C-C motif chemokine 8, Ceruloplasmin, Choriogonadotropin subunit beta, Collagenase 3, C-Peptide, Creatine Kinase-MB, C-X-C motif chemokine 10, C-X-C motif chemokine 11, C-X-C motif chemokine 13, C-X-C motif chemokine 16, C-X-C motif chemokine 5, C-X-C motif chemokine 6, C-X-C motif chemokine 9, Cystatin-C,

Endothelial protein C receptor, Eotaxin, Epidermal growth factor receptor, Ferritin, Fibrinogen, Gastric inhibitory polypeptide, Glucagon, Glucagon-like peptide 1, Granulocyte-macrophage colony-stimulating factor, Growth-regulated alpha, beta, and gamma proteins, Heparin-binding growth factor 2, Hepatocyte growth factor, Immunoglobulin A, Immunoglobulin M, Immunoglobulin G1, Immunoglobulin G2, Immunoglobulin G3, Immunoglobulin G4, Insulin, Insulin-like growth factor-binding protein 1, Insulin-like growth factor-binding protein 2, Insulin-like growth factor-binding protein 3, Insulin-like growth factor-binding protein 4, Insulin-like growth factor-binding protein 5, Insulin-like growth factor-binding protein 6, Insulin-like growth factor-binding protein 7, Interferon alpha-2, Interferon gamma, Interleukin-1 alpha, Interleukin-1 beta, Interleukin-1 receptor antagonist protein, Interleukin-1 receptor type I, Interleukin-1 receptor type II, Interleukin-11, Interleukin-12, Interleukin-12 subunit beta, Interleukin-13, Interleukin-15, Interleukin-2, Interleukin-20, Interleukin-21, Interleukin-23, Interleukin-28A, Interleukin-29, Interleukin-3, Interleukin-33, Interleukin-4, Interleukin-5, Interleukin-6 receptor subunit alpha, Interleukin-6 receptor subunit beta, Interleukin-7, Interleukin-9, Interstitial collagenase, Islet amyloid polypeptide, Keratin, type I cytoskeletal 19 (aa311-367), Kit ligand, Leptin, Leukemia inhibitory factor, Lymphotoxin, Lymphotoxin-alpha, Macrophage colony-stimulating factor 1, Macrophage metalloelastase, Macrophage migration inhibitory factor, Matrilysin, Metalloproteinase inhibitor 1, Metalloproteinase inhibitor 2, Metalloproteinase inhibitor 3, Metalloproteinase inhibitor 4, Myoglobin, Pancreatic prohormone, Peptide YY, Pro-epidermal growth factor, Pro-interleukin-16, Prolactin, Prostate-specific antigen, Protein S100-A12, Protransforming growth factor alpha, Secretory immunoglobulin A, Serum amyloid P-component, SL cytokine, Stromal cell-derived factor 1, Stromelysin-1, Thrombopoietin, Thymic stromal lymphopoietin, Tumor necrosis factor, Tumor necrosis factor ligand superfamily member 10, Tumor necrosis factor ligand superfamily member 6, Tumor necrosis factor receptor superfamily member 1A, Tumor necrosis factor receptor superfamily member 1B, Vascular endothelial growth factor receptor 1, Vascular endothelial growth factor receptor 2, Vascular endothelial growth factor receptor 3, von Willebrand Factor, and WAP four-disulfide core domain protein 2;

an assay instrument configured to (i) receive a body fluid sample, (ii) contact the plurality of reagents with the body fluid sample and (iii) generate and quantitatively display in

human readable form one or more assay results indicative of binding of each biomarker which is assayed to a respective specific binding reagent in the plurality of reagents.

24. A system according to claim 23 wherein the reagents comprise a plurality of antibodies, at least one of which binds to each of the biomarkers which are assayed.

25. A system according to claim 24 wherein assay instrument comprises an assay device and an assay device reader, wherein the plurality of antibodies are immobilized at a plurality of predetermined locations within the assay device, wherein the assay device is configured to receive the body fluid sample such that the body fluid sample contacts the plurality of predetermined locations, and wherein the assay device reader interrogates the plurality of predetermined locations to generate the assay results.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US20 14/056653

A. CLASSIFICATION OF SUBJECT MATTER
IPC(8) - G01 N 33/53 (2014.01)
CPC - G01 N 2800/06 (2014.1 1)
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC(8) - A61K 39/395; C07K 16/00; C12Q 1/68; G01N 33/53, 33/543, 33/68 (2014.01)
 USPC - 435/7.1 , 287.2, 810; 436/501 , 804

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
 CPC - A61K 9/0053, 2039/505; C07K 16/18; G01N 2800/06; Y10S 435/81, 436/804 (2014.01 1) (keyword delimited)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 PatBase, Google Patents, Google, PubMed
 Search terms used: Appendicitis assay and biomarker and device and outcome and threshold and biomarker and risk of mortality.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2012/0028268 A1 (KENTSIS et al) 02 February 2012 (02.02.2012) entire document	1, 2, 4-7, 11-20, 23-25
Y		3, 21
Y	WO 2004/059293 A2 (BUECHLER et al) 15 July 2004 (15.07.2004) entire document	3
Y	US 2002/0150957 A1 (SLOTMAN) 17 October 2002 (17.10.2002) entire document	21
A	US 2013/0122528 A1 (TYRELL et al) 16 May 2013 (16.05.2013) entire document	1-7, 11-21, 23-25
A	US 2013/0171670 A1 (BAR-OR et al) 04 July 2013 (04.07.2013) entire document	1-7, 11-21, 23-25

Further documents are listed in the continuation of Box C.

* Special categories of cited documents:
 "A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier application or patent but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed
 "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
 "&" document member of the same patent family

Date of the actual completion of the international search 19 November 2014	Date of mailing of the international search report 30 DEC 2014
---	--

Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, Virginia 22313-1450 Facsimile No. 571-273-3201	Authorized officer: Blaine R. Copenheaver PCT Helpdesk: 571-272-4300 PCT OSP: 571-272-7774
---	---

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2014/056653

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 8-10, 22
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

- Remark on Protest**
- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
 - The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
 - No protest accompanied the payment of additional search fees.

专利名称(译)	用于诊断和预测阑尾炎和腹痛原因的方法和组合物		
公开(公告)号	EP3047270A1	公开(公告)日	2016-07-27
申请号	EP2014845927	申请日	2014-09-19
[标]申请(专利权)人(译)	阿斯图特医药公司		
申请(专利权)人(译)	ASTUTE MEDICAL , INC.		
当前申请(专利权)人(译)	ASTUTE MEDICAL , INC.		
[标]发明人	ANDERBERG JOSEPH GRAY JEFF MCPHERSON PAUL NAKAMURA KEVIN KAMPF JAMES PATRICK KWAN THOMAS		
发明人	ANDERBERG, JOSEPH GRAY, JEFF MCPHERSON, PAUL NAKAMURA, KEVIN KAMPF, JAMES, PATRICK KWAN, THOMAS		
IPC分类号	G01N33/53		
CPC分类号	G01N33/6893 G01N2800/06 G01N2800/52 G01N2800/60		
优先权	61/880765 2013-09-20 US		
其他公开文献	EP3047270A4		
外部链接	Espacenet		

摘要(译)

本发明涉及用于监测，诊断，预后和确定阑尾炎患者和具有阑尾炎风险的患者的治疗方案的方法和组合物。特别地，本发明涉及使用检测一种或多种生物标志物的测定法作为此类患者中的诊断和预后生物标志物测定法。