

(19)



(11)

EP 2 235 529 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
14.12.2016 Bulletin 2016/50

(51) Int Cl.:
G01N 33/53 (2006.01) **G01N 33/00** (2006.01)
G01N 33/68 (2006.01) **G01N 33/74** (2006.01)
G01N 33/574 (2006.01)

(21) Application number: **09700404.8**

(86) International application number:
PCT/US2009/030340

(22) Date of filing: **07.01.2009**

(87) International publication number:
WO 2009/089286 (16.07.2009 Gazette 2009/29)

(54) **DETERMINATION OF SFLT-1: ANGIOGENIC FACTOR COMPLEX**

BESTIMMUNG DES SFLT-1: ANGIOGENESEFAKTORKOMPLEXES

DÉTERMINATION D'UN COMPLEXE SFLT-1 : FACTEUR ANGIOGÉNIQUE

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK TR

(74) Representative: **Goodfellow, Hugh Robin**
Carpmaels & Ransford LLP
One Southampton Row
London WC1B 5HA (GB)

(30) Priority: **07.01.2008 US 19351 P**

(56) References cited:
WO-A1-2006/128553 WO-A2-01/85796
WO-A2-2005/077007 WO-A2-2006/001888
US-A1- 2007 010 442 US-A1- 2007 111 326

(43) Date of publication of application:
06.10.2010 Bulletin 2010/40

(73) Proprietor: **ORTHO-CLINICAL DIAGNOSTICS, INC.**
Raritan, NJ 08869 (US)

- **ERRICO MICHELA ET AL: "Identification of placenta growth factor determinants for binding and activation of Flt-1 receptor", JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY FOR BIOCHEMISTRY AND MOLECULAR BIOLOGY, INC, US, vol. 279, no. 42, 15 October 2004 (2004-10-15), pages 43929-43939, XP002560050, ISSN: 0021-9258, DOI: DOI:10.1074/JBC.M401418200 [retrieved on 2004-07-21]**
- **ERIKSSON A ET AL: "Placenta growth-factor-1 antagonizes VEGF-induced angiogenesis and tumor growth by the formation of functionally inactive P1GF-1/VEGF heterodimers", CANCER CELL, CELL PRESS, US, vol. 1, 1 February 2002 (2002-02-01), pages 99-108, XP002967596, ISSN: 1535-6108, DOI: DOI:10.1016/S1535-6108(02)00028-4**

- (72) Inventors:
- **BASHIRIANS, George**
Penfield, NY 14526 (US)
 - **BYRNE, Deborah**
Rochester, NY 14626 (US)
 - **GROULX, Sarah**
Ontario, NY 14519 (US)
 - **JACKSON, Sharon**
Rochester, NY 14623 (US)
 - **LANE-BROWN, Heather**
Honeoye Falls, NY 14472 (US)
 - **SANKARAN, Banumathi**
Pittsford, NY 14534 (US)
 - **SCALICE, Edward**
Penfield, NY 14526 (US)
 - **YANG, Karen**
Edison, NJ 08820 (US)
 - **ZHENG, Jian**
Raritan, NJ 08869 (US)

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 235 529 B1

- **HORNIG C ET AL:** "Detection and quantification of complexed and free soluble human vascular endothelial growth factor receptor-1 (sVEGFR-1) by ELISA", **JOURNAL OF IMMUNOLOGICAL METHODS**, ELSEVIER SCIENCE PUBLISHERS B.V.,AMSTERDAM, NL, vol. 226, no. 1-2, 24 June 1999 (1999-06-24), pages 169-177, XP004171370, ISSN: 0022-1759, DOI: DOI:10.1016/S0022-1759(99)00065-4
- **HORNIG ET AL.:** 'Release and complex formation of soluble VEGFR-1 from endothelial cells and biological fluids.' **LABORATORY INVESTIGATION** vol. 80, no. 4, April 2000, pages 443 - 454
- **CHRISTINGER HANS W ET AL:** "The crystal structure of placental growth factor in complex with domain 2 of vascular endothelial growth factor receptor-1", **JOURNAL OF BIOLOGICAL CHEMISTRY**, AMERICAN SOCIETY FOR BIOCHEMISTRY AND MOLECULAR BIOLOGY, US, vol. 279, no. 11, 12 March 2004 (2004-03-12), pages 10382-10388, XP002587110, ISSN: 0021-9258, DOI: 10.1074/JBC.M313237200

Description

FIELD OF THE INVENTION

5 **[0001]** The invention relates to immunoassays and *in vitro* diagnostics and particularly to sFlt-1:angiogenic factor complex determination including in preeclampsia.

BACKGROUND OF THE INVENTION

10 **[0002]** Preeclampsia is a syndrome of hypertension, edema, and proteinuria that affects 5 to 10% of pregnancies and results in substantial maternal and fetal morbidity and mortality. Preeclampsia accounts for 200,000 maternal deaths worldwide per year. Clinical symptoms of preeclampsia typically appear after the 20th week of pregnancy and are usually detected during routine evaluation of a woman's blood pressure and testing for the presence of protein in a sample of her urine. However, such clinical evaluation is ineffective for early diagnosis of the syndrome. Being able to evaluate the likelihood of developing preeclampsia, and/or being able to diagnose preeclampsia in an early stage of gestation, and/or being able to diagnose preeclampsia during a phase of the disease when clinical evaluation is uninformative, would allow early intervention and reduce the risk of medical complications and mortality for a pregnant woman and developing fetus.

20 **[0003]** Currently there are no known cures for preeclampsia. Preeclampsia can vary in severity from mild to life threatening. Maternal complications include renal failure, HELLP syndrome (hemolysis, elevated liver enzymes, and thrombocytopenia), liver failure, cerebral edema with seizures and rarely death. Potential fetal complications include low birth weight, prematurity and death. A mild form of preeclampsia can be treated with bed rest and frequent monitoring. For moderate to severe cases, hospitalization is recommended and blood pressure medication or anticonvulsant medications to prevent seizures are prescribed. If the condition becomes life threatening to the mother or the baby, the pregnancy is terminated and the baby is delivered pre-term.

25 **[0004]** Molecular mechanisms associated with preeclampsia have recently been reviewed (Mutter and Karumanchi, *Microvascular Research* 75:1-8, 2008). As stated by Mutter and Karumanchi, it is believed that endothelial dysfunction contributes to the clinical syndrome of preeclampsia (Roberts and Cooper, *Lancet* 357:53-56, 2001). Many of the symptoms of the disease may result from aberrant endothelial function (including arterial hyperreactivity to exogenous and endogenous stimuli, proteinuria related to increased glomerular permeability, cerebral edema and increased CNS permeability, as well as vascular thrombosis resulting in the HELLP syndrome)(Roberts, *Semin Reprod Endocrinol* 16:5-15, 1998; Roberts and Cooper, *Lancet* 357:53-56, 2001). As such, there has been an active search for circulating factors that cause or contribute to endothelial dysfunction.

30 **[0005]** Proper development of the fetus and placenta is mediated by several growth factors. One of these growth factors is vascular endothelial growth factor (VEGF). VEGF is an endothelial cell-specific mitogen, an angiogenic inducer, and a mediator of vascular permeability. VEGF has also been shown to be important for glomerular capillary repair. VEGF binds as a homodimer to one of two homologous membrane-spanning tyrosine kinase receptors - the fms-like tyrosine kinase receptor (Flt-1)(also known as vascular endothelial growth factor receptor 1 or VEGF-R1), and the kinase domain region receptor (Flk/KDR)(also known as vascular endothelial growth factor receptor 2 or VEGF-R2) which are differentially expressed in endothelial cells obtained from many different tissues. Flt-1, but not Flk/KDR, is highly expressed by trophoblast cells which contribute to placenta formation. Placenta growth factor (PlGF) is a VEGF family member that is also involved in placenta development. PlGF is expressed by cyto- and syncytiotrophoblasts and is capable of inducing proliferation, migration, and activation of endothelial cells. PlGF binds to as a homodimer to the Flt-1 receptor, but not the Flk/KDR receptor. Both PlGF and VEGF contribute to the mitogenic activity and angiogenesis that are critical for the developing placenta.

45 **[0006]** A soluble form of the Flt-1 receptor (sFlt-1) has been identified in a cultured medium of human umbilical vein endothelial cells and *in vivo* expression was subsequently demonstrated in placenta tissue. sFlt-1 is a splice variant of the Flt-1 receptor which lacks the transmembrane and cytoplasmic domains (He et al., *Mol Endocrinol* 13:537-545, 1999; Kendall and Thomas, *Proc Natl Acad Sci USA* 90:10705-10709, 1993).

50 **[0007]** Recent work by researchers at Beth Israel Deaconess Medical Center and Harvard Medical School has demonstrated increased placental production and maternal serum levels of sFlt-1 in patients with preeclampsia (Ahmad and Ahmed, *Circ Res* 95:884-891, 2004; Chaiworapongsa et al., *Am J Obstet Gynecol* 190:1541-1547, 2004; Koga et al., *J Clin Endocrinol Metab* 88:2348-2351, 2003; Maynard et al., *J Clin Invest* 111:649-658, 2003; Shibata et al., *J Clin Endocrinol Metab* 90:4895-4903, 2005). sFlt-1 is able to bind both VEGF and PlGF. Free in serum, it may diminish binding of these growth factors to their cognate receptors Flt-1 and Flk/KDR respectively (Kendall et al., *Biochem Biophys Res Commun* 226:324-328, 1996). In addition to VEGF and PlGF, the placenta is known to produce a number of other angiogenic factors, including the angiopoietins (Ang-1 and Ang-2) as well as their receptor Tie-2 (Dunk et al., *Am J Pathol* 156:2185-2199, 2000; Geva et al., *J Clin Endocrinol Metab* 87:4213-4224, 2002; Goldman-Wohl et al., *Mol Hum*

Reprod 6:81-87, 2000). Increased levels of sFlt-1 and decreased levels of VEGF and PlGF are found in serum of women with preeclampsia.

[0008] Recent attention has also focused on another factor, endoglin (Eng), a co-receptor for transforming growth factor β 1 and β 3, and a protein expressed in large quantities by the placenta in preeclampsia. The extracellular domain of endoglin may be shed and is found in the serum where it is referred to as soluble endoglin (sEng). Like sFlt-1, sEng is increased in maternal serum 2 to 3 months prior to the onset of disease (Levine et al., N Engl J Med 355:992-1005, 2006).

[0009] The identification of an imbalance of circulating angiogenic factors that precedes the onset of preeclampsia or its clinical manifestation will be useful in designing screening and/or diagnostic tests to identify patients at risk for preeclampsia. Such a test would be invaluable to clinicians who may offer close follow-up and therapeutic interventions early in the course of disease. Several retrospective studies using serum obtained from patients having been afflicted with preeclampsia have shown that sFlt-1 concentrations in serum are high as much as 5 to 6 weeks before any clinical findings are noted (Chaiworapongsa et al., J Matern Fetal Neonatal Med 17:3-18, 2005; Hertig et al., Clin Chem 50:1702-1703, 2004; Levine et al., N Engl J Med 350:672-683, 2004; McKeeman et al., Am J Obstet Gynecol 191:1240-1246, 2004). In addition, free VEGF and PlGF are low (Hertig et al., Clin Chem 50:1702-1703, 2004; Levine et al., N Engl J Med 350:672-683, 2004). A recent systematic review of the literature to assess if elevated sFlt-1 or decreased PlGF in the serum could accurately predict the onset of preeclampsia concluded that third trimester increases in sFlt-1 and decrease in PlGF are associated with preeclampsia but there is currently insufficient data to recommend these as screening tests (Widmer et al., Obstet Gynecol 109:168-180, 2007).

[0010] A need continues to exist for more efficient and/or more effective methods of predicting a woman's risk for developing preeclampsia or determining if a woman has preeclampsia. Predicting and/or detecting preeclampsia in an early stage of gestation and/or in a stage of the disease where clinical evaluation may be uninformative would be particularly advantageous.

BRIEF SUMMARY OF THE INVENTION

[0011] The present invention is defined according to the appended claims.

[0012] In one aspect, the invention relates to a method of determining the presence or amount of sFlt-1:PlGF complex. The method comprises:

- a) capturing the complex with capture agent that binds to PlGF, and detecting the complex with detector agent that binds to one or both of sFlt-1 and sFlt-1:PlGF complex; or
- b) capturing the complex with capture agent that binds to sFlt-1, and detecting the complex with detector agent that binds to one or both of PlGF and sFlt-1:PlGF complex; or
- c) capturing the complex with capture agent that binds to sFlt-1:PlGF complex, and detecting the complex with detector agent that binds to sFlt-1:PlGF complex, wherein capture agent and detector agent are capable of being bound simultaneously to the complex; or
- d) capturing the complex with capture agent that binds to sFlt-1:PlGF complex, and detecting the complex with detector agent that binds to one or both of PlGF and sFlt-1.

[0013] In another aspect, the invention relates to a method of guiding therapy of, or predicting, detecting, or monitoring, vascular, vascular-related, cancer, cancer-related, cardiac, cardiac-related, preeclampsia or preeclampsia-related disease in an individual. The method comprises:

- a) determining the presence or amount of sFlt-1:PlGF complex in a sample from an individual; and
- b)
 - i) comparing the result of the sFlt-1:PlGF determination with one or more reference values; or
 - ii) converting the result of the sFlt-1:PlGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the sFlt-1:PlGF complex transformed results with one or more reference values;

thereby guiding therapy of, or predicting, detecting, or monitoring, the disease.

[0014] In one embodiment, the method may further comprise:

- a) determining the presence or amount of one or more molecular entities in the sample; and
- b)
 - i) comparing the result of the molecular entities determination and the result of the sFlt-1:PlGF complex deter-

mination with the one or more reference values; or
ii) converting the result of the molecular entities determination and the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the molecular entities transformed results and the sFlt-1:PIGF transformed results with the one or more reference values.

[0015] In yet another embodiment, the method may further comprise:

- a) determining the values of one or more physical attributes of the individual; and
- b)

- i) comparing the physical attributes determination and the result of the molecular entities determination and the result of the sFlt-1:PIGF complex determination with the one or more reference values; or
- ii) converting the physical attributes determination and the result of the molecular entities determination and the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the physical attributes transformed results and the molecular entities transformed results and the sFlt-1:PIGF transformed results with the one or more reference values.

[0016] In an additional embodiment, the method may comprise:

- a) determining the values of one or more physical attributes of the individual; and
- b)

- i) comparing the physical attributes determination and the result of the sFlt-1:PIGF complex determination with the one or more reference values; or
- ii) converting the physical attributes determination and the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the physical attributes transformed results and the sFlt-1:PIGF transformed results with the one or more reference values.

[0017] Another aspect of the invention relates to a composition for determining the presence or amount of sFlt-1:PIGF complex. The composition comprises:

- a) capture agent that binds to PIGF and detector agent that binds to one or both of sFlt-1 and sFlt-1:PIGF complex; or
- b) capture agent that binds to sFlt-1 and detector agent that binds to one or both of PIGF and sFlt-1:PIGF complex; or
- c) capture agent that binds to sFlt-1:PIGF complex and detector agent that binds to sFlt-1:PIGF complex, wherein capture agent and detector agent are capable of being bound simultaneously to the complex; or
- d) capture agent that binds to sFlt-1:PIGF complex and detector agent that binds to one or both of PIGF and sFlt-1.

[0018] Additionally, the invention relates to a composition comprising a purified form of sFlt-1:PIGF complex.

BRIEF DESCRIPTION OF THE DRAWINGS

[0019]

Figure 1 illustrates an assay structure for determining sFlt-1:PIGF complex by way of PIGF capture and sFlt-1 detection;
Figure 2 shows a dose response curve for an sFlt-1:PIGF-1 complex assay;
Figure 3 shows a dot-plot of normal (0) and preeclampsia (1) patient sample results based on an sFlt-1:PIGF-1 complex assay;
Figure 4 shows a dot-plot of the computed ratios of sFlt-1:PIGF complex to free PIGF in samples from normal (0) and preeclampsia (1) patients.

DETAILED DESCRIPTION OF THE INVENTION

[0020] sFlt-1, as used herein, refers to the soluble form of the fms-like tyrosine kinase receptor (Flt-1). Flt-1 is also known as vascular endothelial growth factor receptor 1 (VEGFR-1). sFlt-1 (also known as sVEGFR-1) is a splice variant

of Flt-1 which lacks the transmembrane and cytoplasmic domains of Flt-1. The amino acid sequence of Flt-1, containing amino acids for the transmembrane and cytoplasmic domains, is provided by Genbank Accession No. P17948 (1338 amino acids). The soluble form of Flt-1 is missing amino acids 688-1338, and amino acids 657-687 differ, resulting in an amino acid sequence of sFlt-1 of 687 amino acids. The sFlt-1 isoform is described in Genbank Accession No. P17948.

[0021] PIGF, as used herein, refers to placenta growth factor. PIGF has four currently recognized isotopes, including PIGF-1, PIGF-2, PIGF-3, and PIGF-4, which arise from alternative splicing. As an example of a PIGF amino acid sequence is provided by Genbank Accession No. P49763 (221 amino acids). The PIGF-1 isoform is missing amino acids 132-203, resulting in an amino acid sequence of PIGF-1 of 149 amino acids. The PIGF-1 isoform is described in Genbank Accession No. P49763.

[0022] sFlt-1 binds to PIGF, and the result is referred to herein as the sFlt-1:PIGF complex.

[0023] With these terms in mind, in one aspect the disclosure relates to a method of determining the presence or amount of sFlt-1:PIGF complex. The method comprises:

- a) capturing the complex with capture agent that binds to PIGF, and detecting the complex with detector agent that binds to one or both of sFlt-1 and sFlt-1:PIGF complex; or
- b) capturing the complex with capture agent that binds to sFlt-1, and detecting the complex with detector agent that binds to one or both of PIGF and sFlt-1:PIGF complex; or
- c) capturing the complex with capture agent that binds to sFlt-1:PIGF complex, and detecting the complex with detector agent that binds to sFlt-1:PIGF complex, wherein capture agent and detector agent are capable of being bound simultaneously to the complex; or
- d) capturing the complex with capture agent that binds to sFlt-1:PIGF complex, and detecting the complex with detector agent that binds to one or both of PIGF and sFlt-1.

[0024] In another aspect, the disclosure relates to a method of guiding therapy of, or predicting, detecting, or monitoring, vascular, vascular-related, cancer, cancer-related, cardiac, cardiac-related, preeclampsia or preeclampsia-related disease in an individual. The method comprises:

- a) determining the presence or amount of sFlt-1:PIGF complex in a sample from an individual; and
- b)
 - i) comparing the result of the sFlt-1:PIGF determination with one or more reference values; or
 - ii) converting the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the sFlt-1:PIGF complex transformed results with one or more reference values;

thereby guiding therapy of, or predicting, detecting, or monitoring, the disease.

[0025] In one embodiment, the method may further comprise:

- a) determining the presence or amount of one or more molecular entities in the sample; and
- b)
 - i) comparing the result of the molecular entities determination and the result of the sFlt-1:PIGF complex determination with the one or more reference values; or
 - ii) converting the result of the molecular entities determination and the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the molecular entities transformed results and the sFlt-1:PIGF transformed results with the one or more reference values.

[0026] In yet another embodiment, the method may further comprise:

- a) determining the values of one or more physical attributes of the individual; and
- b)
 - i) comparing the physical attributes determination and the result of the molecular entities determination and the result of the sFlt-1:PIGF complex determination with the one or more reference values; or
 - ii) converting the physical attributes determination and the result of the molecular entities determination and the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the physical attributes transformed results and

the molecular entities transformed results and the sFlt-1:PIGF transformed results with the one or more reference values.

[0027] In an additional embodiment, the method may comprise:

- a) determining the values of one or more physical attributes of the individual; and
- b)

- i) comparing the physical attributes determination and the result of the sFlt-1:PIGF complex determination with the one or more reference values; or
 - ii) converting the physical attributes determination and the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the physical attributes transformed results and the sFlt-1:PIGF transformed results with the one or more reference values.

Extending the concept of the subject disclosure beyond the illustrated disclosure the disclosure can relate to a first molecular entity and a second molecular entity capable of binding to each other to form a molecular complex, wherein in the preferred embodiment sFlt-1 is an example of a first molecular entity and PIGF is an example of a second molecular entity. sFlt-1:PIGF complex is an example of a molecular complex. The subject disclosure is described in detail below in respect to certain preferred embodiments involving angiogenic receptor sFlt-1 and angiogenic factors PIGF and VEGF. These molecular entities are of particular interest in respect to preeclampsia and preeclampsia-related disease. Immunoassays - General Discussion Applicable to the Subject

[0028] Disclosure

[0029] Analytical assays comprising agents that bind specifically to a substance whose presence or amount is to be determined are typically referred to as binding assays. Immunoassays are generally thought of as binding assays that utilize antibodies or fragments of antibodies as capture agents and detector agents. Although the invention is described in reference to capture and detector antibodies, binding agents other than antibodies also are contemplated. Such binding agents, as in the case of capture and detector antibodies, bind specifically to a first molecular entity or second molecular entity involved in complex formation or to the complex comprising the first and second molecular entities.

[0030] Immunoassays can be categorized as competitive or non-competitive (sandwich) immunoassays. In a competitive immunoassay, a labeled form of the substance to be determined is placed in competition with the (unlabeled) native substance derived from a sample. The labeled form competes with the native substance for binding to a fixed amount of a capture antibody; the capture antibody being capable of binding to the labeled substance and the (unlabeled) native substance - though not simultaneously. Unknown concentrations of the substance derived from the sample are determined from detectable signal arising from the labeled form of the substance that is either bound to the capture antibody or from the labeled form of the substance that is free, i.e., not bound to the capture antibody.

[0031] In a non-competitive (sandwich) immunoassay, the substance to be determined is contacted with capture antibody and detector antibody, which antibodies typically bind to the substance at different epitopes or, in any case, are capable of being bound to the substance simultaneously. There can be multiple types of capture antibody that bind to different epitopes of the substance and/or multiple types of detector antibody that bind to different epitopes of the substance. Detector antibody is labeled and capture antibody typically is immobilized on a solid support or is capable of being immobilized thereon. Conventional labels include radioactive tags, enzymes, chromophores, fluorophores, stable free radicals and enzyme cofactors, inhibitors and allosteric effectors.

[0032] Immunoassay methods can be performed using any suitable format. They can be carried out in solution, in test devices where soluble and insoluble components can be separated, or in dry analytical elements. Such immunoassay formats include, for example, enzyme-linked immunosorbent assays (ELISA), immunometric assays, dot blot (also known as slot blot) assays, etc. The immunoassays can be heterogeneous or homogeneous. In heterogeneous immunoassays, bound components are separated from free, non-bound components - usually by way of one or more wash-steps. In homogeneous immunoassays, separation of bound from free components is not required.

[0033] Numerous publications relating to immunoassays and immunoassay methods are available ("Practical Immunoassay", Butt ed, Marcel Dekker, 1984; "Immunochemistry of Solid Phase-Immunoassay"; Butler, CRC Press, 1991; "Immunoassay", Law ed, Taylor & Francis, 1996; "Immunoassay", Eleftherios et al., Academic Press, 1996; "Principles and Practice of Immunoassay", Second Edition, Price and Newman eds, Macmillan, 1997; "Immunoassays in the Clinical Laboratory", Nakamura et al. eds, Alan R Liss, Inc, 1979; "Quantitative Enzyme Immunoassay", Engvall et al. eds, Blackwell Scientific Publications, 1978; Sommer et al., Clin Chem 32:1770-1774, 1986; "A Primer for Multilayer Immunoassay", Berke, American Chemical Society Conference Proceeding, p 303-312, Plenum Press, 1988; US 4,200,690; US 4,207,307; US 4,407,943; US 4,550,075; US 4,551,426; US 4,560,648; US 5,312,744; US 5,314,830; US 5,424,220; US 5,415,998; US 5,501,949; US 5,518,887; US 5,663,054; US 5,789,261; US 5,935,780; US 5,958,339; US 6,087,188;

US 6,096,563; US 6,121,006; US 6,143,575; US 6,395,503; US 6,878,515; US 6,838,250; US 4,372,745; US 4,670,381; US 4,483,921; US 4,517,288; US 4,822,747; US 4,824,778; US 4,829,012; US 4,839,299; US 4,847,194; US 4,847,195; US 4,853,335; US 4,855,226; US 4,857,453; US 4,857,454; US 4,859,610; US 4,863,876; US 4,868,106; US 4,868,130; US 4,879,219; US 5,776,933.

5 **[0034]** Details relating to dry analytical immunoassay elements may be found in the following publications and references cited therein: US 3,867,258; US 3,992,158; US 4,042,435; US 4,050,898; US 4,066,403; US 4,153,668; US 4,258,001; US 4,292,272; US 4,430,436.

10 **[0035]** A support can be any material that is insoluble, or can be made insoluble by a subsequent reaction. The support can be chosen for its intrinsic ability to attract and immobilize a capture agent. Alternatively, the support can have affixed thereto a linking agent that has the ability to attract and immobilize the capture agent. The linking agent can, for example, include a charged substance that is oppositely charged with respect to the capture agent itself or to a charged substance conjugated to the capture agent. In general, the linking agent can be any binding partner (preferably specific) that is immobilized on (attached to) the support and has the ability to immobilize the capture agent through a binding reaction. The linking agent enables the indirect binding of the capture agent to a support before the performance of the assay or during the performance of the assay. The support can be, for example, plastic, derivatized plastic, magnetic or non-magnetic metal, glass or silicon, including, for example, a test tube, microwell, microtiter well, membrane, sheet, bead, microparticle, chip, and other configurations known to those of ordinary skill in the art. A support can be porous or nonporous material that is insoluble or essentially insoluble in aqueous compositions. A support can exist in a wide range of different forms and configurations: vessel, tube, microtiter plate, sphere, microparticle, rod, strip, filter paper, chromatography paper, membrane, etc.

20 **[0036]** The methods of the present invention can be adapted for use in systems that utilize microparticle technology including automated and semi-automated systems wherein the solid support comprises a microparticle as described in US 5,006,309 and US 5,089,424.

25 **[0037]** In a preferred format, capture agent is linked to biotin and immobilized on the internal surface of a microwell by way of biotin binding to avidin, streptavidin, or other biotin-binding agent.

[0038] Numerous publications relating to binding of molecules to supports are available. For instance: US 4,624,930; US 5,061,640; US 4,945,042; US 4,885,255; US 5,362,624; US 5,277,589; US 5,268,306; US 5,376,557; US 5,858,803; US 5,126,241; US 5,362,655; US 5,437,981; US 5,792,606; US 5,863,740; US 5,935,780; US 6,391,571.

30 **[0039]** In heterogeneous immunoassays separation of free from bound components can be effected by passing suitable liquid wash solution across the support, filtering soluble, free components away from the support, immuno-precipitating free components, precipitation with substances such as polyethylene glycol or ammonium sulfate, magnetic separation or binding to a different support, and so on.

35 **[0040]** A detector agent can be directly detectable or indirectly detectable. It may comprise a label that is capable of emitting a signal directly or indirectly. Suitable labels are known in the art, and include, for example, horseradish peroxidase, alkaline phosphatase, fluorescent tags, detectable tracers, and so forth. A label may be a chemical entity that is capable of directly eliciting a detectable signal - usually an optical signal: absorbance, reflectance, luminescence, and fluorescence. A signal may be derived from radioactivity. It may be voltammetric or conductometric, etc. A directly detectable label that is intrinsically capable of producing a detectable signal may include organic and inorganic substances capable of fluorescence, or phosphorescence, such as but not limited to, fluorescein and derivatives thereof, N-(3-fluoranthyl)-maleimide, radionuclides, such as carbon 14, tritium, phosphorus 32; azo-oxo, azo-tetrazo, azine, oxazine, thiazine, quinoline, indamine, pyrone and pyrazolone dyes. In general, a label can be a chemical element, a compound, or protein.

40 **[0041]** A label that is detectable indirectly usually will require the presence of one or more additional substances for production of detectable signal. Such labels include enzymes that require the presence of a substrate(s), a cofactor(s), or a metal(s). For example, horseradish peroxidase is a label that requires an electron donor and an oxidizing agent in order to generate a signal: for example, luminol and hydrogen peroxide to generate chemiluminescence, or triarylimidazole and hydrogen peroxide to generate dye.

45 **[0042]** A capture and/or detector agent in reference to the preferred embodiments is an agent that is capable of binding with specificity to sFlt-1, VEGF, PlGF or complexes comprising sFlt-1 and angiogenic factor. In general, considering a first molecular entity and a second molecular entity capable of binding to form a complex comprising both molecular entities, capture agent or detector agent that is stated to bind or specifically bind to the first molecular entity, unless otherwise indicated, is capable of binding to free first molecular entity and to the first molecular entity when it is bound to the second molecular entity in the complex. Similarly, capture agent or detector agent that is stated to bind or specifically bind to the second molecular entity, unless otherwise indicated, is capable of binding to free second molecular entity and to the second molecular entity when it is bound to the first molecular entity in the complex. Capture agent or detector agent stated to bind to complex or specifically bind to complex does not bind or does not bind substantially to free first molecular entity or free second molecular entity.

55 **[0043]** Capture and detector agents include antibodies or fragments thereof, and chimeric antibodies comprising

antibody fragments derived from different biological origins, such as human/mouse, mouse/goat, human/goat. They include non-antibody proteins and peptides, such as, angiogenic factors, angiogenic receptors, and non-protein binding agents. Monoclonal or polyclonal antibodies or combinations of monoclonal and polyclonal antibodies represent preferred capture and detector agents.

5 **[0044]** The term "sample" or "biological sample" includes any quantity of a substance from a living thing or formerly living thing. Such living things include, but are not limited to, humans, mice, monkeys, rats, rabbits, horses, and other animals. Samples include, but are not limited to, blood, serum, urine, tears, cells, organs, tissues, bone, bone marrow, lymph, lymph nodes, synovial tissue, chondrocytes, synovial macrophages, endothelial cells, and skin.

10 Molecular Complex - Capture and Detection Applicable to the Subject Invention

[0045] Various immunoassay embodiments will be described symbolically in reference to determination of molecular complex FP comprising molecular entity F and molecular entity P.

15 **[0046]** In one kind of immunoassay embodiment, FP can be captured using immobilized or immobilizable capture agent specific for F. Capture agent is capable of binding to free F and to F in complex FP. "Free" F means F that is not bound to P though it may be bound to other molecular entities. Detection is carried out using detector agent, preferably labeled with horseradish peroxidase, which detector agent specifically binds to P and is capable of binding to free P and to P in complex FP. Similarly, "free" P means P that is not bound to F though it may be bound to other molecular entities. In another immunoassay embodiment, detection is carried out using detector agent, preferably labeled with horseradish peroxidase, which detector agent binds specifically to complex FP and does not bind or does not bind substantially to either free F or free P.

20 **[0047]** In an analogous immunoassay embodiment, FP can be captured using immobilized or immobilizable capture agent specific for P. Capture agent is capable of binding to free P and to P in complex FP. Detection is carried out using detector agent specific for F, which detector agent is capable of binding to free F and to F in complex FP. Or, detection is carried out using detector agent specific for complex FP, which detector agent does not bind or does not bind substantially to either free F or free P.

25 **[0048]** In another immunoassay embodiment, complex FP can be captured using immobilized or immobilizable capture agent that binds to complex FP, which capture agent does not bind or does not bind substantially to either free F or free P. Detection is carried out using detector agent also specific for complex FP, which detector agent does not bind or does not bind substantially to either free F or free P. In this embodiment capture and detector agent bind to different sites of complex FP or, in any case, are capable of being bound simultaneously to complex FP. Alternatively, detection can be carried out using detector agent specific for F, which detector agent is capable of binding to free F and to F in complex FP. Or, detection can be carried out using detector agent specific for P, which detector agent is capable of binding to free P and to P in complex FP. Or, detection can be carried out using detector agent specific for F (and capable of binding to free F and to F in complex FP) and detector agent specific for P (and capable of binding to free P and to P in complex FP).

30 **[0049]** In the above-described immunoassay embodiments, separation of non-bound from bound components is carried out where separation is desirable or necessary. For example, in the case where FP is captured using immobilized capture agent specific for F and capable of binding to free F and complex FP (thus capturing and immobilizing free F and complex FP) and wherein FP is detected using detector agent specific for P, which detector agent is capable of binding to free P and to complex FP, subsequent to capture, separation of bound components from non-bound components would be carried out - usually via a wash procedure. Thus, bound free F and bound FP are separated from non-bound free F, non-bound FP, and free P (P is not captured and therefore it is not bound). Detector agent is then introduced for binding to captured and immobilized complex FP without interference from free P - which is removed during the wash procedure.

35 **[0050]** Examples of antibodies which can be used for capture and detection of sFlt-1:PIGF complex include, but are not limited to: mouse monoclonal anti-sFlt-1 antibody RD-1-49560 (R&D Systems), 3661-M16 (R&D Systems) and 4449-M24 (R&D Systems); mouse monoclonal anti-sFlt-1 antibody RD-7-49566 (R&D Systems), mouse monoclonal anti-sFlt-1 antibody M14 (Ortho-Clinical Diagnostics), mouse monoclonal anti-PIGF antibody 37203 (R&D Systems), rat monoclonal anti-PIGF antibody 358932 (R&D Systems), rat monoclonal anti-PIGF antibody 358905 (R&D Systems), monoclonal antibody 358932 (R&D Systems) and 261 (Ortho Clinical Diagnostics).

40 **[0051]** In accordance with the subject invention, a composition is also provided for use in the immunoassay embodiments, i.e. for determining the presence or amount of sFlt-1:PIGF complex. The composition comprises:

- 45
- a) capture agent that binds to PIGF and detector agent that binds to one or both of sFlt-1 and sFlt-1:PIGF complex; or
 - b) capture agent that binds to sFlt-1 and detector agent that binds to one or both of PIGF and sFlt-1:PIGF complex; or
 - c) capture agent that binds to sFlt-1:PIGF complex and detector agent that binds to sFlt-1:PIGF complex, wherein capture agent and detector agent are capable of being bound simultaneously to the complex; or
 - d) capture agent that binds to sFlt-1:PIGF complex and detector agent that binds to one or both of PIGF and sFlt-1.
- 50

[0052] As in the methods of the subject invention, one or both of the capture and detector agents can be polyclonal antibodies, or one or both of the capture and detector agents can be monoclonal antibodies, or one or both of the capture and detector agents can be a combination of monoclonal and polyclonal antibodies.

5 Assay Calibration and Performance Verification Applicable to the Subject Invention

[0053] Immunoassays require calibration and/or verification that the target analyte is detected with specified precision and accuracy. In respect to an immunoassay for determining complex FP, compositions comprising a purified form of the molecular complex FP can be used for such purposes. In the context of the present invention the term "purified" means that complex FP is or has been separated from free F and free P. Separation of complex FP from free F and free P can be achieved using one or more well known methods such as: methods that rely on the affinity of the individual components for specific binding partners, for example, F binding to (preferably immobilized) P, P binding to (preferably immobilized) F, or binding of the individual components to other binding partners such as F binding to (preferably immobilized) anti-F antibody, P binding to (preferably immobilized) anti-P antibody, or FP binding to (preferably immobilized) anti-FP antibody, or binding of the individual components to other types of specific binding partners (affinity chromatography); or, by way of ion exchange chromatography, metal-ion exchange chromatography, ligand exchange chromatography, or hydrophobic chromatography; or, using methods that rely on differences in mass and/or size and/or differences in net electric charge of the complex and its component molecular entities. These include size-exclusion chromatographic methods, centrifugation methods, molecular sieving methods, for example, membrane filtration, and electrophoretic methods, and so on.

[0054] If the complex is stable during separation and subsequent introduction into a storage composition (typically a liquid formulation) - wherein stable means that it does not readily dissociate to form substantial amounts of free F and free P during these procedures - the composition comprising the complex can then be placed under suitable conditions to maintain stability over longer times. It can, for example, be stored at a low temperature less than or equal to about 4°C if it is in a liquid state, or the liquid composition can be frozen and maintained as a solid at a temperature below 0°C, or it can be lyophilized and maintained in a state at some appropriate temperature which could even be greater than about 20°C. Upon removal from storage, after having been thawed or reconstituted if necessary, it should be stable for an acceptable period of time under the conditions for which it will be used for assay calibration or verification.

[0055] Modified forms of the complex can be produced in which F and P are covalently bonded to each other (for example, sFlt-1 covalently bonded to PIGF). Such a covalently modified form of FP will be stable with respect to dissociation to form free F and free P. Preferably, modified, covalent complex FP and noncovalent complex FP have substantially the same or similar binding properties with respect to capture agent and detector agent. If a modified, covalent form of FP and unmodified, noncovalent form, or natural/native form of complex FP do not have substantially the same or sufficiently similar binding properties with respect to capture and detector agent, appropriate mathematical correction can be applied to compensate for the differences in binding properties, as is known in the analytical arts. In the context of the present invention the term "form" of FP shall be understood to include intact, noncovalent, natural/native FP purified/isolated from a suitable source (such as human serum, placental tissue or fluids, and so on), noncovalent FP prepared from native F and native P or native F and recombinant P or recombinant F and recombinant P and, modified, covalently bonded FP (F and P covalently bonded to each other) prepared from native F and native P or native F and recombinant P or recombinant F and recombinant P.

[0056] In accordance with this aspect of the subject invention, a composition is also provided which comprises a purified form of sFlt-1:PIGF complex. One or both of sFlt-1 and PIGF of the sFlt-1:PIGF complex may be native, or one or both of sFlt-1 and PIGF of the sFlt-1:PIGF complex may be recombinant.

45 Predicting, Detecting/Diagnosing, Monitoring Disease Applicable to the Subject Invention

[0057] It is conventional practice to determine the presence or amount of a single substance (marker/biomarker) in a biological sample of an individual in order to determine if the individual is afflicted with a particular disease (detecting/diagnosing), is likely to become afflicted (predicting), to determine progression of the disease (monitoring), or if the individual is responding to treatment (monitoring).

[0058] For example, determining the presence of glucose (qualitative assay) or the amount of glucose (quantitative assay) in a urine or serum sample of an individual often is carried out in order to confirm a diagnosis of diabetes or to monitor the efficacy of treatment of an individual diagnosed as diabetic. An optical signal - oftentimes a colorimetric signal in both qualitative and quantitative assays - will form in proportion to the amount of glucose in a sample. In one type of colorimetric assay, the amount or color intensity of dye that forms, dye density, can be determined visually in a qualitative manner. Alternatively, the amount or color intensity of dye can be determined more accurately and precisely using suitable instrumentation to measure and assign a numeric value to the dye density. Visually determined dye density produced with a biological sample of an individual is usually compared with a reference dye density scale or dye density

range comprising a scale or range of dye densities correlated with non-diabetics and diabetics. Quantified dye density is usually compared with a numeric reference value or numeric reference range of values similarly correlated.

[0059] Reference values or reference ranges with respect to a particular disease are typically derived from populations diagnosed as free of that particular disease (normal) and populations diagnosed as afflicted with that particular disease. The reference values or reference ranges can include values or measures associated with different stages of the disease. Sample derived results can be compared with these reference values or reference ranges and the disease thereby detected/diagnosed or monitored. In the case of multiple biomarkers alone or in conjunction with physical attributes (discussed below), reference values and reference value ranges can also be correlated with different forms, origins, causes, and so forth, of a disease, which forms, origins, causes, etc. are responsive to different therapies. Sample derived results can be compared with these reference values or reference ranges and therapy to treat an individual can thereby be informed or guided.

[0060] The terms "reference value", "reference range of values", and so forth, if not otherwise indicated, are intended to encompass quantitative or numeric reference population measures and qualitative or visual reference population measures, such as dye density scales, color or hue scales, and so forth.

[0061] The conventional practice of using a single biomarker for detecting and monitoring a disease has been and continues to be a valuable aid to clinicians and the patients they serve. Frequently, more than one biomarker is known to be associated with a disease. For example, glycosylated hemoglobin is also associated with diabetes. Determining the amount of glucose and glycosylated hemoglobin in a sample from an individual diagnosed as diabetic can provide more information about the state of the disease in the individual or the effect of treatment than either determination alone. The determination of multiple biomarkers not only improves diagnosis and monitoring of a disease but, it may also enable a clinician to determine the likelihood or probability that an individual will be afflicted with a disease or is in a stage of the disease where clinical evaluation is uninformative. The determination of multiple biomarkers can also be useful in guiding therapy - allowing a clinician to distinguish different forms or stages of a disease so as to permit treatment of an individual using appropriate therapeutic measures. The determination of multiple biomarkers can be used in conjunction with one or more physical attribute of the individual to further improve predictive utility, diagnostic utility, and selection of appropriate therapy.

[0062] Such additional multiple biomarkers are referred to herein as one or more molecular entities. In the case of sFlt-1:PIGF complex detection, such molecular entities include, but are not limited to, endoglin, soluble endoglin, creatinine, c-reactive protein, urine protein, placental protein 13, human chorionic gonadotropin hormone, or alpha-fetoprotein and liver enzymes, such as, aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase and gamma-glutamyl transferase. As used herein, VEGF can include VEGF₁₂₁, VEGF₁₄₅, VEGF₁₆₅, VEGF_{165b}, VEGF₁₆₇, VEGF₁₈₃, VEGF₁₈₆, VEGF₁₈₉, or VEGF₂₀₆.

[0063] In a presently preferred embodiment, sFlt-1:PIGF complex is detected and the disease is preeclampsia or preeclampsia-related and guiding therapy, predicting, detecting, or monitoring is within the first 15 weeks of gestation or within the first 20 weeks of gestation or within the first 30 or more weeks of gestation.

[0064] Mathematical operations and algorithms can be used to convert the amounts of two or more biomarkers into one or more transformed-results. Such mathematical operations can include computing a ratio of the molecular entities determination to the sFlt-1:PIGF complex determination or the sFlt-1:PIGF complex determination to the molecular entities determination. The transformed-result or -results can be more informative than the direct concentration-based results. Mathematical operations or algorithms that convert the determinations of one or more biomarkers in combination with one or more physical attributes of an individual into one or more transformed-results also are being employed to provide improvements in guiding therapy, predicting, detecting/diagnosing, and monitoring disease. Physical attributes include, but are not limited to: gestational age, race, gender, age, blood pressure and body mass index. These can be used individually or in any combination along with one or more biomarker.

[0065] Numerous patent and non-patent publications are available describing mathematical operations and algorithms and their use in clinical diagnostics. For example, US Patent Application Publ No 2006/381104, US Patent Application Publ No 2003/700672, US Patent Application Publ No 2003/410572, US Patent Application Publ No 2003/634145, US Patent Application Publ No 2005/323460, PCT Intl Publ No WO 2007/044860, US Patent Application Publ No 2002/330696, US Patent Application Publ No 2002/331127, US Patent Application Publ No 1997/912133, US Patent No 6,306,087, US Patent No 6,248,063 and US Patent No 5,769,074.

Example I

Assay for sFlt-1:PIGF Complex

[0066] In this Example an immunometric assay is described for determining the amount of sFlt-1:PIGF complex in a sample, and the utility of the sFlt-1:PIGF complex as a diagnostic indicator of preeclampsia is shown.

[0067] The immunoassay format depicted in Figure 1 was used to capture and detect sFlt-1:PIGF complex. sFlt-1:PIGF

EP 2 235 529 B1

complex in a sample is captured using biotinylated rat anti-PlGF antibody. sFlt-1:PlGF complex captured by anti-PlGF antibody is immobilized by way of biotin binding to streptavidin coated on the surface of a reaction vessel. Free sFlt-1 and sFlt-1:VEGF complex are not captured by biotinylated anti-PlGF capture agent and are removed along with other non-bound substances during a wash procedure. The complex is detected by way of luminescence using horseradish peroxidase (HRP)-labeled anti-sFlt-1 antibody. The luminescence signal generated from bound HRP-labeled anti-sFlt-1 antibody is directly proportional to the concentration of sFlt-1:PlGF present in the sample. Any free PlGF captured by the biotinylated anti-PlGF capture agent does not form a sandwich with the HRP-labeled anti-sFlt-1 antibody and is not measured.

[0068] Materials used in the immunoassay include:

1. Biotin Reagent: biotinylated anti-PlGF antibody at 2.25 µg/mL in a pH buffered aqueous composition. Biotinylated anti-PlGF antibodies capable of binding to sFlt-1:PlGF complex include but are not limited to Ortho-Clinical Diagnostic's mouse monoclonal antibody OCD-261 and rat monoclonal antibodies 358905 and 358932 available from R&D Systems, Minneapolis, Minnesota.

2. Conjugate Reagent: HRP-labeled anti-sFlt-1 antibody at 1.125 µg/mL in a pH buffered aqueous composition. HRP-labeled anti-sFlt-1 antibodies which do not interfere with binding of VEGF or PlGF to sFlt-1 include but are not limited to Ortho-Clinical Diagnostic's mouse monoclonal antibody clones 3661-M16 and 4449-M24 and R&D System's monoclonal antibody 49560.

3. Calibrators were prepared by adding purified sFlt-1:PlGF complex to horse serum (where "purified sFlt-1:PlGF complex" means, as noted above, the complex is or has been separated from free sFlt-1 and free PlGF). Ten fold molar excess of recombinant human PlGF obtained from R&D Systems (Catalog Number: 264-PG) was incubated for 20 min at 20°C followed by an overnight incubation at 2-8°C with full length sFlt-1 (687 amino acids) obtained from Scios (Scios Inc., Mountain View, California) in phosphate buffered saline (PBS) buffer pH 7.4. The sFlt-1:PlGF complex was separated from free PlGF by size exclusion chromatography on a Superdex 200 column equilibrated with phosphate buffer containing 0.25M ethanolamine at pH 8.5. The fractions containing the complex were pooled and concentrated using a centrifugal concentration unit. The concentrated pool of complex was then dialyzed into PBS pH 7.4.

The purified sFlt-1:PlGF complex was diluted to obtain different concentrations in horse serum (GIBCO, Catalog number 26050) to provide calibrators CR1 to CR10 spanning an sFlt-1:PlGF complex concentration range between 0.0 and 25,000 pg/mL.

4. Reaction vessels were streptavidin-coated microwells (SAC wells).

Immunoassay Protocol

[0069] The immunoassay protocol was as follows:

Sample (80µL) was combined with 80µL Biotin Reagent in a SAC well and the mixture was incubated for 21 minutes at 37°C. A wash-step using VITROS ECi Universal Wash Reagent followed the incubation. After the wash-step, Conjugate Reagent (160µL) was added to the SAC well and incubated for 21 minutes at 37°C. A second wash-step using the VITROS ECi Universal Wash Reagent followed the incubation. VITROS ECi Signal reagent comprising luminescence reagents was then introduced and incubated for 5 minutes. The luminescence signal was acquired following the incubation.

[0070] The analytical and clinical performance of the sFlt-1:PlGF complex assay was evaluated. Results are shown in Tables 1-6 and Figures 2-4. In Table 1 the signal responses, Analyzer Light Units (ALU), obtained using calibrator fluids (comprising sFlt-1:PlGF complex) are shown. The data from Table 1 is shown as a dose-response plot in Figure 2. Evidence confirming that only sFlt-1:PlGF complex (and not free sFlt-1 or free PlGF) was detected in the assay was provided by way of cross reactivity studies in which samples comprising different concentrations of free sFlt-1 or free PlGF (and no sFlt-1:PlGF complex) were used in place of samples comprising sFlt-1:PlGF complex in the assay. These results are presented in Tables 2 and 3 where it can be seen that free sFlt-1 and free PlGF do not register significant responses.

[0071] Serum samples obtained from women in the 2nd or 3rd trimester of pregnancy diagnosed as either having preeclampsia or not having preeclampsia (normal) were assayed for sFlt-1:PlGF complex using the above-described assay. The results are summarized in Table 4. A dot-plot of the results is shown in Figure 3 where it can be seen that the sFlt-1:PlGF complex concentrations found in the serum samples from the normal population are generally lower and clearly discriminated from the concentrations found in the serum samples from the population diagnosed as having preeclampsia. Receiver operating characteristic (ROC) analysis of the data was carried out. The area under ROC curve (Table 5) is 0.783 demonstrating that the sFlt-1:PlGF complex is a useful biomarker for detecting/diagnosing preeclampsia.

sia.

[0072] For each sample, the ratio of the amount of sFlt-1:PIGF complex present in a sample relative to the amount of free PIGF present in the sample was computed and compared with the amount of free PIGF as determined using a free PIGF assay. These results are provided in Table 4. A dot-plot of the ratio (sFlt-1:PIGF complex/free PIGF) for each sample versus clinical diagnosis is shown in Figure 4. The computed ratios for the normal population samples are generally smaller in magnitude and clearly discriminated from the computed ratios for the preeclampsia population samples. The area under ROC curve (Table 6) is 0.889 demonstrating that the ratio, sFlt-1:PIGF complex/free PIGF, is also a useful marker for preeclampsia.

Example II

ELISA Detection of sFlt-1:PIGF Complex Using anti-sFlt-1 Capture

[0073] The use of ELISA methods for determining sFlt-1:PIGF complex is described in this Example. In this assay method, free sFlt-1 and sFlt-1:PIGF complex are captured using anti-sFlt-1 antibody, and labeled anti-PIGF antibody is used to detect sFlt-1:PIGF complex.

Plate Coating

[0074] Costar 96-well microtiter high-binding plates were coated with 100 μ L/well of a 10 mM phosphate, 2 mM EDTA, pH 7.0 solution containing 1 μ g/mL of anti-sFlt-1 antibody 49560 (R&D Systems) and incubated for 18 hrs at 25°C. Wells were washed once (PBS with 0.05% TWEEN-20), followed by addition of 290 μ L/well of blocking buffer (1% bovine serum albumin in phosphate buffered saline, pH 7.0), then incubated for 1 hr at 25°C. Wells were aspirated and dried for 4 hrs in a low humidity incubator then sealed in air-tight bags prior to use.

ELISA Method

[0075] A series of samples containing a constant amount of recombinant sFlt-1 (purified recombinant sFlt-1, Scios; expressed in insect cells; corresponding to the 687 amino acid full-length soluble fms-like tyrosine kinase 1) and varying amounts of recombinant PIGF (149 amino acids; PIGF-1; expressed in E. coli; R&D Systems, Catalog Number: 264-PG/CF) were incubated together in sample diluent (blocker casein in PBS with 0.05% TWEEN-20, Pierce) to allow formation of a sFlt-1:PIGF complex. 100 μ L of each preincubated sample was added to a microtiter well. 100 μ L sample diluent was added to at least one well to serve as control. Wells were incubated for 15 minutes at 37°C with shaking. Non-bound substances were removed by washing the plate 6 times with wash buffer (PBS with 0.05% TWEEN-20). Total sFlt-1, free sFlt-1 and sFlt-1:PIGF complex were measured individually in anti-sFlt-1 antibody coated plates by addition of 100 μ L/well of specific HRP-labeled antibodies (1 μ g/mL in blocker casein in PBS with 0.05% TWEEN 20) as outlined in Table 7. After addition of each detector antibody, the plate was incubated for 15 minutes with shaking at 37°C. The plate was washed six times with wash buffer. 100 μ L of O-Phenylenediamine (OPD) substrate reagent was added to each well and incubated in the dark for 30 minutes at 25°C. 25 μ L stop solution was added to each well and the absorbance at 492nm was obtained. (HRP substrate, OPD reagent and stop solution were from Ortho-Clinical Diagnostics, Inc., Raritan, New Jersey, USA).

ELISA Results - Total sFlt-1

[0076] The absorbance at 492nm obtained with each sample is provided in Table 8. HRP-labeled anti-sFlt-1 antibody M14 (Ortho-Clinical Diagnostics) binds to both free sFlt-1 and sFlt-1:PIGF complex, therefore, free sFlt-1 and sFlt-1 in the sFlt-1:PIGF complex are detected. As shown in Table 8, the signal response for total sFlt-1 (sum of observed signals for free sFlt-1 and sFlt-1:PIGF complex) is constant, independent of PIGF concentration.

ELISA Results - Free sFlt-1

[0077] HRP-labeled anti-sFlt-1 antibody 49566, (R&D Systems) binds to free sFlt-1 and does not bind sFlt-1:PIGF complex, therefore, only free sFlt-1 is detected. A decrease in signal response is observed as the PIGF concentration increases from 0 to 2000 pg/mL (Table 8). This is expected: as PIGF concentration increases more sFlt-1:PIGF complex forms and the concentration of free sFlt-1 decreases.

EP 2 235 529 B1

ELISA Results - sFlt-1:PIGF Complex

5 [0078] HRP-labeled anti-PIGF antibody 358905 (R&D Systems) binds to PIGF and therefore only sFlt-1 bound in complex sFlt-1:PIGF is detected. Table 8 shows an increase in signal response as the PIGF concentration increases from 0 to 2000 pg/mL.

[0079] These results as provided in Table 8 demonstrate that total sFlt-1, free sFlt-1 and sFlt-1:PIGF complex can be measured in an ELISA format using anti-sFlt-1 antibodies to capture the molecular species comprising sFlt-1.

10 [0080] The presence or amount of sFlt-1:PIGF complex in a sample can be determined by capture of sFlt-1 and detection of PIGF as demonstrated, and the measurement so obtained, alone or in conjunction with additional chemical and/or physical biomarkers, can be used as a diagnostic indicator for preeclampsia.

ELISA Detection of sFlt-1:PIGF Complex Using anti-PIGF Capture

15 [0081] In another ELISA method, free PIGF and sFlt-1:PIGF complex are captured using anti-PIGF antibody. Labeled anti-sFlt-1 antibody is used to detect sFlt-1:PIGF complex.

Plate coating

20 [0082] Costar 96-well microtiter high-binding plates were coated with 100 μ L/well of a 10 mM phosphate, 2 mM EDTA, pH7.0 solution containing 0.8 μ g/mL of anti-PIGF antibody 358932 (R&D Systems) and incubated 18hrs at 25°C. Wells were washed once (PBS with 0.05% Tween-20), followed by addition of 290 μ L/well of blocking buffer (1% bovine serum albumin in phosphate buffered saline, pH 7.0), then incubated for 1 hr at 25°C. Wells were aspirated and dried for 4 hrs in a low humidity incubator, then sealed in air-tight bags prior to use.

ELISA Method

25 [0083] A series of samples as shown in Table 9, contain a constant amount of recombinant PIGF (SEQ ID NO:2; expressed in E. coli; R&D Systems, Catalog Number: 264-PG/CF) and varying amounts of recombinant sFlt-1 (687 amino acid full-length soluble fms-like tyrosine kinase 1; expressed in insect cells; Scios, Inc.) in a sample diluent (Blocker Casein in PBS with 0.05% TWEEN-20, Pierce) which are incubated together to allow formation of a sFlt-1:PIGF complex; the mixture is added to the coated wells and the plate is incubated with shaking at 37°C for 15 minutes. Non-bound substances are removed by washing the plate with a mild detergent based buffer. Total PIGF, free PIGF and sFlt-1:PIGF complex are measured individually in anti-PIGF antibody coated plates by addition of 100 μ L/well of specific HRP-labeled antibodies (1 μ g/mL in Blocker Casein in PBS with 0.05% TWEEN-20, Pierce) as outlined in Table 7. After addition of each detector antibody the plate is incubated for 15 minutes with shaking at 37°C. The plate is washed six times with wash buffer. A colorimetric reaction follows addition of 100 μ L/well of OPD HRP substrate reagent and incubation in the dark for 30 minutes at 25°C 25 μ L/well stop solution is added and the absorbance at 492nm is obtained. (HRP substrate O-Phenylenediamine reagent and stop solution were from Ortho-Clinical Diagnostics, Inc., Raritan, New Jersey, USA).

ELISA Results - Total PIGF

40 [0084] Table 9 lists the absorbance at 492nm for each sample following each specific reaction. HRP-labeled anti-PIGF antibody 358905 (R&D Systems) binds to both free PIGF and sFlt-1:PIGF complex, therefore, free PIGF and PIGF in the sFlt-1:PIGF complex are detected. The total PIGF response column in Table 9 shows a constant response for samples with a PIGF concentration of 2000 pg/mL, regardless of the concentration of sFlt-1.

ELISA Results - Free PIGF

50 [0085] HRP-labeled anti-PIGF antibody 37203 (R&D Systems) binds to free PIGF and does not bind sFlt-1:PIGF complex, therefore, only free PIGF is detected. Table 9 shows a decrease in signal response as the sFlt-1 concentration increases from 0 to 2000 pg/mL. The concentration of free PIGF molecules should decrease and the concentration of sFlt-1:PIGF complex should increase as free PIGF binds to sFlt-1.

ELISA Results - sFlt-1:PIGF Complex

55 [0086] HRP-labeled anti-sFlt-1 antibody M14 (Ortho-Clinical Diagnostics) binds to sFlt-1 and to sFlt-1:PIGF complex and therefore only PIGF captured as a sFlt-1:PIGF complex is detected. Table 9 shows an increase in signal response as the sFlt-1 concentration increases from 0 to 2000 pg/mL.

[0087] The results outlined in Table 9 demonstrate total PIGF, free PIGF and the sFlt-1:PIGF complex can be measured in an ELISA format using anti-PIGF antibodies to capture all PIGF species and to independently detect each. As increasing concentrations of sFlt-1 is added to a constant concentration of PIGF, the free PIGF response is decreased and the sFlt-1:PIGF complex response is increased as demonstrated by the change in absorbance at 492 nm.

[0088] The presence or amount of sFlt-1:PIGF complex in a sample can be determined by capture of PIGF and detection of sFlt-1 as demonstrated, and the measurement so obtained, alone or in conjunction with additional chemical and/or physical biomarkers, can be used as a diagnostic indicator for preeclampsia.

Example III

sFlt-1:PIGF Complex, Total sFlt-1, Free PIGF Gestation Age: Weeks 27-37

Total sFlt-1 and Free PIGF Assays

[0089] Assays of total sFlt-1 and free PIGF were carried out using R&D Systems Quantikine® Kits Catalog Nos. DVR100B and DPG00, respectively. The assays were conducted according to protocols described in the R&D Systems package inserts for these products.

sFlt-1:PIGF Complex Assay

[0090] As in Example I, the immunoassay format depicted in Figure 1 was used to capture and detect sFlt-1:PIGF complex.

[0091] Materials used in the immunoassay include:

1. Biotin Reagent: biotinylated anti-PIGF antibody (Ortho Clinical Diagnostics mouse monoclonal, OCD-261) at 6.0 ug/mL in a pH buffered aqueous composition. Other biotinylated anti-PIGF antibodies capable of binding to sFlt-1:PIGF complex are available from R&D Systems, Minneapolis, Minnesota.

2. Conjugate Reagent: HRP-labeled anti-sFlt-1 antibody at 4.5 ug/mL in a pH buffered aqueous composition. HRP-labeled anti-sFlt-1 antibodies which do not interfere with binding of VEGF or PIGF to sFlt-1 include but are not limited to Ortho-Clinical Diagnostic's mouse monoclonal antibody clones 3661-M16 and 4449-M24 and R&D Systems monoclonal 49560.

3. Calibrators were prepared by adding purified sFlt-1: PIGF complex to horse serum (where "purified sFlt-1:PIGF complex" means, as noted above, the complex is or has been separated from free sFlt-1 and free PIGF). Ten fold molar excess of recombinant human PIGF obtained from R&D systems (Catalog Number: 264-PG) was incubated for 20 min at 20°C followed by an overnight incubation at 2-8°C with full length sFlt-1 (687 amino acids)(Scios Inc., Mountain View, California USA) in PBS buffer pH 7.4. The sFlt-1:PIGF complex was separated from free PIGF by size exclusion chromatography on a Superdex 200 column equilibrated with phosphate buffer containing 0.25M ethanolamine at pH 8.5. The fractions containing the complex were pooled and concentrated using a centrifugal concentration unit. The concentrated pool of complex was then dialyzed into PBS pH 7.4.

The purified sFlt-1:PIGF complex was diluted at different concentrations in horse serum (GIBCO, Catalog number 26050) to generate Calibrators CR1 to CR10 spanning sFlt-1 Complex concentration of 0.0 to 25,000 pg/mL.

4. Reaction vessels were streptavidin-coated microwells (SAC wells).

Immunoassay Protocol

[0092] The immunoassay protocol was as follows:

[0093] Sample (80 µL) was combined with 35 µL Biotin Reagent and 35 µL Conjugate Reagent in a SAC well and the mixture was incubated for 21 minutes at 37°C. A wash-step using VITROS ECi Universal Wash Reagent followed the incubation. After the wash-step VITROS ECi Signal Reagent was then introduced and incubated for 5 minutes. The luminescence signal was acquired following the incubation.

[0094] Each patient sample was assayed in duplicate for PIGF-1 and sFlt-1 according to R&D Systems ELISA assay protocols. A single assay of each patient sample for sFlt-1:PIGF-1 complex was carried out using the sFlt-1:PIGF-1 complex assay procedure as described above. The mean of the R&D System assay duplicates and the sFlt-1:PIGF-1 complex assay results for each patient sample are reported in the top portion of Table 10. Gestational age at the time of sample collection is reported in column 2 of Table 10. Diagnoses established near the time of sample collection according to the American College of Obstetrics and Gynecology (ACOG) guidelines are reported in column 3. Areas under ROC curves were determined as provided in the bottom portion of Table 10: PIGF-1 assay (0.8796), sFlt-1 assay (0.7994), sFlt-1:PIGF complex assay (0.7654), sFlt-1/PIGF ratio (0.8519) and sFlt-1:PIGF complex/PIGF ratio (0.8642).

EP 2 235 529 B1

These data show that for gestational age 27 to 37.4, the clinical performance of sFlt-1:PIGF Complex assay on its own or as a ratio to free PIGF assay is comparable to that of an assay which measures total sFlt-1.

Example IV

sFlt-1:PIGF Complex, sFlt-1, PIGF

Gestation Age: Weeks 13-21

[0095] As evidenced in this Example, evaluation of the presence or amount of sFlt-1:angiogenic factor complex provides superior clinical utility for diagnosis and prediction of preeclampsia compared with total sFlt-1 or the ratio of total sFlt-1 to free PIGF in early stages of gestation. Similarly, the ratio of sFlt-1:PIGF complex to free PIGF has superior clinical utility for diagnosis and prediction of preeclampsia compared with free PIGF, total sFlt-1 or the ratio of total sFlt-1 to free PIGF.

[0096] As in Example III, each patient sample was assayed in duplicate for free PIGF and total sFlt-1 according to R&D Systems ELISA assay protocols and a single assay was carried out on each patient sample for sFlt-1:PIGF complex using the complex assay procedure. The results are shown in Table 11. Gestational age at the time of sample collection is reported in column 2 of Table 11. Diagnoses were established according to the ACOG guidelines and assignments of mild and severe preeclampsia were rendered based on ICD-9 code designations. These are listed in columns 3 and 4 of Table 11. Upon collection, the blood samples were archived for later assay; diagnoses having been made between weeks 21 and post partum.

[0097] Areas under ROC curves were determined as provided in Table 12. The results presented in the uppermost data panel (Mild or Severe PE) are based on 94 patient samples collected within weeks 13-21 of gestation. The diagnoses of mild and severe preeclampsia were grouped together as preeclampsia-positive during ROC analysis. The computed areas under the ROC curves follow: free PIGF assay (0.5745), total sFlt-1 assay (0.5029), sFlt-1:PIGF complex assay (0.6331), total sFlt-1/free PIGF ratio (0.5396) and sFlt-1:PIGF complex/free PIGF ratio (0.6392). These results indicate that total sFlt-1, free PIGF and the total sFlt-1/free PIGF ratio are not useful for predicting or detecting preeclampsia during early stages of gestation. However, the sFlt-1:PIGF complex and sFlt-1:PIGF complex/free PIGF ratio results suggest that they are clinically capable of discriminating between subjects who subsequently developed mild or severe preeclampsia from those who did not.

[0098] The results presented in the middle data panel (Severe PE, Mild PE = Normal) are based on the same 94 patient samples; however, during ROC analysis the diagnoses of mild preeclampsia were reclassified as normal. Areas under the ROC curves are as follows: free PIGF assay (0.7142), total sFlt-1 assay (0.5622), sFlt-1:PIGF complex assay (0.6942), total sFlt-1/free PIGF ratio (0.6771) and sFlt-1:PIGF complex/free PIGF ratio (0.7445). These results indicate the superior clinical capability of sFlt-1:PIGF complex on its own or in combination with free PIGF as a ratio, as compared with total sFlt-1 in predicting the onset of preeclampsia.

[0099] The results of ROC analysis provided in the lowermost data panel are based on only those patient samples collected within weeks 16-21 of gestation, the patients having been diagnosed as normal, with mild preeclampsia being categorized as normal in the analysis, or as having been diagnosed with severe preeclampsia. Areas under the ROC curves follow: free PIGF assay (0.7394), total sFlt-1 assay (0.5455), sFlt-1:PIGF complex assay (0.7636), total sFlt-1/free PIGF ratio (0.6848) and sFlt-1:PIGF complex/free PIGF ratio (0.8707). The results demonstrate the superior performance of sFlt-1:PIGF complex on its own or as a ratio, as compared to total sFlt-1 in predicting the onset of preeclampsia.

Table 1

Dose-Response Results		
sFlt-1: PIGF Complex Calibrator ID	sFlt-1:PIGF Complex Concentration (pg/mL)	Signal (ALU)
CR1	0	14.7
CR2	75.0	20.8
CR3	188	37.3
CR4	375	76.3
CR5	750	165
CR6	1500	365
CR7	3125	869
CR8	6250	1723

EP 2 235 529 B1

(continued)

Dose-Response Results		
sFlt-1: PIGF Complex Calibrator ID	sFlt-1:PIGF Complex Concentration (pg/mL)	Signal (ALU)
CR9	12500	3628
CR10	25000	7292

Table 2

Cross reactivity of Free sFlt-1				
sFlt-1 Concentration (pg/mL)	Sample ID	Signal (ALU)	Predicted * Concentration (pg/mL)	%Cross Reactivity
0	S RIB	15.36	29.33	N/A
0	S R1H	11.02	6.181	N/A
100	S R2	12.24	13.52	13.5
500	S R3	6.96	0.00	0.0
1000	S R4	11.57	9.614	1.0
2000	S R5	16.30	33.56	1.7
10000	S R6	22.29	57.23	0.6
20000	S R7	13.19	18.68	0.1
40000	S R8	17.78	39.84	0.1
80000	S R9	33.82	95.54	0.1
160000	S R10	60.84	174.5	0.1

Table 3

Cross Reactivity of Free PIGF				
PIGF Concentration (pg/mL)	Sample ID	Signal (ALU)	Predicted concentration (pg/mL)	% Cross Reactivity
0	P R1S	14.97	27.5	N/A
0	P RIB	18.60	43.2	N/A
56	P R2	11.56	9.57	17.1
112	P R3	12.49	14.9	13.3
446	P R4	10.24	1.04	0.2
1115	P R5	10.57	3.28	0.3
2231	P R6	10.45	2.48	0.1
11154	P R7	11.46	8.91	0.1
22307	P R8	21.67	54.9	0.2

EP 2 235 529 B1

Table 4

Patient results: sFlt-1:PlGF complex, free PlGF and
sFlt-1:PlGF complex/free PlGF ratio

Sample ID	Gestational Age at Collection (weeks)	Diagnosis	Classification (1=preeclampsia, 0=normal)	sFlt-1:PlGF Complex Concentration (pg/mL)	Free PlGF Concentration (pg/mL)	sFlt-1:PlGF Complex/Free PlGF Ratio
N47	23 3/4	preeclampsia	1	316	23.3	13.6
N49	35	preeclampsia	1	3553	72.4	49.1
N50	30 6/7	preeclampsia	1	1421	6.16	230.7
N51	21 1/2	preeclampsia w/HELLP	1	717	5	143.3
N52	33 4/7	preeclampsia	1	79	4.02	19.6
N53	25 3/7	preeclampsia w/HELLP	1	432	5	86.4
N54	29	preeclampsia w/HELLP	1	1093	5	218.6
N55	26 5/7	preeclampsia	1	2437	27.2	89.6
N56	33 5/7	preeclampsia	1	2031	158	12.9
N58	38 4/7	normal	0	6	74.5	0.1
N59	22	normal	0	170	129	1.3
N60	38 6/7	normal	0	1122	508	2.2
N61	36 2/7	normal	0	42	40.5	1.0
N62	29 3/7	gestational hypertension + preclampsia	1	5629	45.9	122.6
N63	30	IUGR + gestational hypertension	0	1332	6.25	213.1
N64	36	normal	0	629	32.1	19.6
N65	32 5/7	normal	0	281	36.5	7.7
N67	39	normal	0	5	5	1.0
N68	37 5/7	normal	0	446	320	1.4

EP 2 235 529 B1

Table 5

ROC Analysis	
W = Area Under Curve, SE = Standard Error, p = probability	
sFlt-1:PIGF Complex Assay	
W:	0.8000
SE:	0.1017
p:	0.0016

Table 6

ROC Analysis	
W = Area Under Curve, SE = Standard Error, p = probability	
sFlt-1:PIGF/free PIGF ratio	
W:	0.889
SE:	0.0886
p:	<0.0001

Table 7

Analyte Measured	Total sFlt-1	Free sFlt-1	sFlt-1:PIGF complex	Total PIGF	Free PIGF	sFlt-1:PIGF complex
Coated on ELISA well	sFlt-1 specific monoclonal antibody (49560, R&D Systems) binds to Free sFlt-1 and sFlt-1:PIGF Complex			PIGF specific monoclonal antibody (358932, R&D Systems) binds to Free PIGF and sFlt-1:PIGF Complex		
sample	Preincubated sample containing 2000 pg/mL sFlt-1 and 0-2000 pg/mL PIGF			Preincubated sample containing 2000 pg/mL PIGF and 0-2000 pg/mL sFlt-1		
HRP detection antibody (1 μ g/mL)	M14 Ortho-clinical diagnostics	49566 R&D Systems	358905 R&D Systems	358905 R&D Systems	37203 R&D systems	M14 Ortho-clinical diagnostics
Detection antibody characteristic	Binds to free sFlt-1 and sFlt-1:PIGF complex	Binds free sFlt-1 but not to sFlt-1:PIGF complex	Binds to free PIGF and sFlt-1:PIGF complex	Binds to free PIGF and sFlt-1:PIGF complex	Binds to free PIGF but not to sFlt-1:PIGF complex	Binds to free sFlt-1 and sFlt-1:PIGF complex

Table 8

Plate coated with 1 μ g/mL anti-sFlt-1 monoclonal antibody 49560, R&D Systems		Absorbance 492 nm		
sFlt (pg/mL)	PIGF (pg/mL)	Total sFlt-1	Free sFlt-1	sFlt-1:PIGF complex
0	0	0.104	0.095	0.006
2000	2000	2.163	1.077	0.734
2000	200	2.144	1.215	0.176

EP 2 235 529 B1

(continued)

Plate coated with 1µg/mL anti-sFlt-1 monoclonal antibody 49560, R&D Systems		Absorbance 492 nm		
sFlt (pg/mL)	PIGF (pg/mL)	Total sFlt-1	Free sFlt-1	sFlt-1:PIGF complex
2000	20	2.198	1.601	0.096
2000	2	2.246	1.895	0.070
2000	0	2.290	1.976	0.014

Table 9

Plate Coated with 0.8 µg/mL anti-PIGF rat monoclonal 358932, R&D Systems		Absorbance 492 nm		
PIGF (pg/mL)	sFlt-1 (pg/mL)	Total PIGF	Free PIGF	sFlt:PIGF complex
0	0	0.001	0.004	0.022
2000	2000	3.000	1.908	1.185
2000	200	3.000	2.087	0.164
2000	20	3.000	2.146	0.017
2000	2	3.000	2.289	0.008
2000	0	3.000	3.000	0.002

Table 10

sFlt-1:PIGF complex, sFlt-1, free PIGF and Ratios Gestation Age: Weeks 27-37							
Subject ID	Gestational Age At Collection (Weeks)	Preeclampsia (1 =Yes, 0=No)	RDS ELISA Free PIGF	RDS ELISA Total sFlt-1	sFlt-1:PIGF Complex	RDS ELISA Total sFlt-1/RDS ELISA Free PIGF Ratio	sFlt-1:PIGF Complex/RDS ELISA Free PIGF Ratio
1	34.3	1	247	3127	652	12.7	2.6
7	34.9	0	938	1854	189	2.0	0.2
2	29.7	1	41.3	68920	1430	1667.7	34.6
9	30.1	0	469	900	92.3	1.9	0.2
3	37.0	1	66.8	16079	1300	240.5	19.4
8	36.1	0	144	36830	5960	255.5	41.3
6	30.1	1	77.8	46935	2430	603.5	31.2
14	31.1	0	870	1714	193	2.0	0.2
11	27.0	1	39.3	6906	559	175.8	14.2
15	27.1	0	256	2602	195	10.2	0.8
10	34.6	1	351	1716	219	4.9	0.6
12	34.6	0	60.5	8462	855	139.9	14.1
13	37.0	1	212	9699	1870	45.7	8.8
38	37.4	0	311	4376	794	14.1	2.6
17	37.4	1	33.0	8303	357	251.9	10.8
19	37.4	0	184	4165	499	22.6	2.7
21	37.0	1	58.7	13435	1250	228.8	21.3

EP 2 235 529 B1

(continued)

sFlt-1:PIGF complex, sFlt-1, free PIGF and Ratios Gestation Age: Weeks 27-37								
Subject ID	Gestational Age At Collection (Weeks)	Preeclampsia (1 =Yes, 0=No)	RDS ELISA Free PIGF	RDS ELISA Total sFlt-1	sFlt-1:PIGF Complex	RDS ELISA Total sFlt-1/RDS ELISA Free PIGF Ratio	sFlt-1:PIGF Complex/RDS ELISA Free PIGF Ratio	
39	37.4	0	263	1487	380	5.7	1.4	
22	29.0	1	85.3	15223	2000	178.4	23.4	
34	28.6	0	1443	3290	478	2.3	0.3	
23	36.4	1	131	6256	1270	47.7	9.7	
24	36.6	0	1622	7180	826	4.4	0.5	
27	29.7	1	37.2	14040	998	377.9	26.9	
35	29.9	0	883	1790	332	2.0	0.4	
28	30.3	1	100	13350	1880	133.2	18.8	
5	30.0	0	627	5214	391	8.3	0.6	
29	31.3	0	1789	2422	433	1.4	0.2	
30	31.3	1	51.5	34730	1630	674.3	31.6	
31	31.3	1	43.6	4745	259	108.9	5.9	
33	31.4	0	1084	3080	602	2.8	0.6	
16	33.4	1	1460	2106	492	1.4	0.3	
18	32.6	0	1324	1687	260	1.3	0.2	
26	31.1	1	379	1530	374	4.0	1.0	
41	32.1	0	548	1454	229	2.7	0.4	
36	30.6	1	29.9	27790	934	930.3	31.3	
4	30.0	0	900	2016	506	2.2	0.6	
			W:	0.8796	0.7994	0.7654	0.8519	0.8642
			SE:	0.0604	0.0769	0.0826	0.0669	0.0669
			p:	<0.0001	<0.0001	0.0007	<0.0001	<0.0001
			n	36	36	36	36	36

Table 11

sFlt-1:PlGF-1 complex, sFlt-1, Free PlGF and Ratios

Gestation Age: Weeks 13-21

(continues on next 3 pages)

Specimen No.	Gestational Age at Collection (weeks)	Mild or Severe PE (ICD9 Codes 642.01, 642.41, 2652.43, 642.71, 642.91, 642.51)	Severe PE with delivery (ICD9 Code 642.51)	Conc (pg/mL)			RDS ELISA Total sFlt-1/RDS ELISA Free PlGF Ratio	sFlt-1:PlGF Complex/RDS ELISA Total sFlt-1/RDS ELISA Free PlGF Ratio
				RDS ELISA Free PlGF	RDS ELISA Total sFlt-1	sFlt-1:PlGF Complex		
URMC618	12.57	1	1	45.3	1742	3.4	38.5	0.1
URMC619	12.86	0	0	210	2155	81.6	10.3	0.4
URMC594	14.14	0	0	40.5	1421	59.2	35.1	1.5
URMC592	14.29	1	1	44.9	2402	142	53.6	3.2
URMC606	14.29	0	0	97.5	1910	72.2	19.6	0.7
URMC624	14.29	1	1	42.2	3208	256	76.1	6.1
URMC664	14.29	0	0	161	2066	51.5	12.8	0.3
URMC625	14.43	0	0	56.2	3169	61.1	56.4	1.1
URMC663	14.57	1	0	44.1	1712	53.4	38.8	1.2
URMC604	14.86	1	0	127	1179	7.7	9.3	0.1
URMC527	15.00	1	0	75.4	2234	68.3	29.6	0.9
URMC529	15.00	0	0	35.9	1378	100	38.4	2.8
URMC545	15.00	1	0	59.5	2053	46.1	34.5	0.8
URMC621	15.00	1	0	145	4200	186	29.0	1.3
URMC547	15.14	0	0	75.3	3440	123	45.7	1.6
URMC587	15.14	0	0	53.1	1367	70.7	25.7	1.3
URMC623	15.14	0	0	39.4	1768	31.5	44.9	0.8
URMC645	15.29	1	0	38.4	1860	41.8	48.5	1.1
URMC647	15.29	0	0	53.4	2805	109	52.5	2.0
URMC651	15.29	1	0	112	2853	103	25.5	0.9
URMC586	15.43	1	0	20.7	5201	119	251.6	5.8

EP 2 235 529 B1

5
10
15
20
25
30
35
40
45
50
55

Specimen No.	Gestational Age at Collection (weeks)	Mild or Severe PE (ICD9 Codes 642.01, 642.41, 2652.43, 642.71, 642.91, 642.51)	Severe PE with delivery (ICD9 Code 642.51)	Conc (pg/mL)			RDS ELISA Total sFlt-1/RDS ELISA Free PlGF Ratio	sFlt-1:PlGF Complex/RDS ELISA Total sFlt-1/RDS ELISA Free PlGF Ratio
				RDS ELISA Free PlGF	RDS ELISA Total sFlt-1	sFlt-1:PlGF Complex		
URMC653	15.43	0	0	92.2	1431	41.1	15.5	0.4
URMC601	15.57	1	0	53.1	1192	43.2	22.5	0.8
URMC613	15.57	1	1	101	1499	25.3	14.9	0.3
URMC668	15.57	0	0	108	2091	188	19.4	1.7
URMC570	15.71	0	0	39.9	939	-1.23	23.5	0.0
URMC603	15.71	0	0	59.1	2281	35.3	38.6	0.6
URMC614	15.71	0	0	206	1302	59.3	6.3	0.3
URMC568	15.86	1	0	160	1526	119	9.5	0.7
URMC666	15.86	1	0	76.9	1915	92.4	24.9	1.2
URMC638	16.00	0	0	96.0	5102	95.2	53.1	1.0
URMC515	16.14	1	0	94.2	2754	166	29.2	1.8
URMC517	16.14	0	0	99.8	2649	55.2	26.5	0.6
URMC577	16.14	1	0	225	2297	173	10.2	0.8
URMC579	16.14	0	0	101	3007	67.9	29.8	0.7
URMC633	16.14	1	0	92.2	2860	104	31.0	1.1
URMC692	16.14	1	0	26.8	1303	44.3	48.5	1.7
URMC533	16.29	1	0	143	1458	110	10.2	0.8
URMC534	16.29	0	0	105	3248	95.8	30.9	0.9
URMC574	16.29	1	0	157	4192	78.7	26.7	0.5
URMC590	16.29	0	0	120	1381	106	11.5	0.9
URMC634	16.29	0	0	266	1941	91.9	7.3	0.3
URMC684	16.29	1	0	118	1475	63.1	12.5	0.5
URMC685	16.29	0	0	89.0	931	16.2	10.5	0.2
URMC636	16.43	1	0	184	1961	73.8	10.7	0.4
URMC648	16.43	1	0	121	1786	76.3	14.7	0.6
URMC650	16.43	0	0	69.7	1749	35.6	25.1	0.5
URMC536	16.57	1	0	129	2358	130	18.3	1.0
URMC571	16.57	1	0	58.9	2575	643	43.7	10.9
URMC573	16.57	0	0	68.3	3304	68.3	48.4	1.0

EP 2 235 529 B1

5
10
15
20
25
30
35
40
45
50
55

Specimen No.	Gestational Age at Collection (weeks)	Mild or Severe PE (ICD9 Codes 642.01, 642.41, 2652.43, 642.71, 642.91, 642.51)	Severe PE with delivery (ICD9 Code 642.51)	Conc (pg/mL)			RDS ELISA Total sFlt-1/RDS ELISA Free PlGF Ratio	sFlt-1:PlGF Complex/RDS ELISA Free PlGF Ratio
				RDS ELISA Free PlGF	RDS ELISA Total sFlt-1	sFlt-1:PlGF Complex		
URMC589	16.57	1	1	15.1	2293	322	151.9	21.3
URMC669	16.57	1	1	47.8	2373	105	49.6	2.2
URMC676	16.57	0	0	63.7	2353	88.1	36.9	1.4
URMC509	16.71	1	0	93.8	429	24.3	4.6	0.3
URMC538	16.71	0	0	113	2999	118	26.6	1.0
URMC671	16.71	0	0	77.9	754	-0.422	9.7	0.0
URMC511	16.86	0	0	192	3341	136	17.4	0.7
URMC512	16.86	1	0	53.5	754	-1.44	14.1	0.0
URMC513	16.86	0	0	170	1152	68.9	6.8	0.4
URMC542	16.86	1	0	155	13220	258	85.1	1.7
URMC543	16.86	0	0	86.5	2299	84.1	26.6	1.0
URMC576	16.86	0	0	169	4417	140	26.2	0.8
URMC597	17.00	0	0	138	1960	99.9	14.2	0.7
URMC610	17.00	1	1	121	2389	339	19.8	2.8
URMC612	17.00	0	0	70.9	1920	52.8	27.1	0.7
URMC595	17.29	1	0	62.9	2355	100	37.4	1.6
URMC657	17.29	1	0	114	1212	92.1	10.7	0.8
URMC691	17.29	0	0	113	2191	104	19.4	0.9
URMC643	17.43	0	0	222	5476	86	24.7	0.4
URMC672	17.43	1	0	247	3125	123	12.7	0.5
URMC521	17.57	1	1	46.4	346	12.7	7.4	0.3
URMC560	17.57	0	0	211	1754	70.9	8.3	0.3
URMC642	17.57	1	1	153	2390	183	15.6	1.2
URMC522	17.71	0	0	247	3507	116	14.2	0.5
URMC551	17.71	1	1	251	4549	448	18.2	1.8
URMC659	17.86	0	0	97.4	1457	74.3	15.0	0.8
URMC539	18.00	1	0	129	697	33.9	5.4	0.3
URMC541	18.00	0	0	115	1317	127	11.5	1.1
URMC530	18.29	1	1	89.2	1222	142	13.7	1.6

EP 2 235 529 B1

5
10
15
20
25
30
35
40

Specimen No.	Gestational Age at Collection (weeks)	Mild or Severe PE (ICD9 Codes 642.01, 642.41, 2652.43, 642.71, 642.91, 642.51)	Severe PE with delivery (ICD9 Code 642.51)	Conc (pg/mL)			RDS ELISA Total sFlt-1/RDS ELISA Free PIGF Ratio	sFlt-1:PIGF Complex/RDS ELISA Free PIGF Ratio
				RDS ELISA Free PIGF	RDS ELISA Total sFlt-1	sFlt-1:PIGF Complex		
URMC678	18.29	1	0	254	1893	224	7.4	0.9
URMC630	18.43	1	0	204	2461	116	12.0	0.6
URMC679	18.43	0	0	445	2478	206	5.6	0.5
URMC531	18.57	0	0	148	3006	19.1	20.3	0.1
URMC564	18.71	1	1	64.9	1969	80.3	30.4	1.2
URMC565	18.71	0	0	71.6	2653	23.4	37.0	0.3
URMC640	18.71	0	0	161	1411	42.8	8.8	0.3
URMC632	18.86	0	0	166	2573	79.2	15.5	0.5
URMC639	18.86	1	0	67.8	435	-4.36	6.4	-0.1
URMC616	19.00	0	0	329	1538	51.2	4.7	0.2
URMC518	19.29	1	0	280	4100	212	14.6	0.8
URMC615	19.57	1	1	33.8	11565	391	342.6	11.6
URMC519	19.71	0	0	217	878	102	4.1	0.5
URMC505	20.43	0	0	279	856	71.9	3.1	0.3
URMC503	21.14	1	0	483	1762	156	3.6	0.3

Table 12

45
50
55

ROC Analysis					
	RDS ELISA Free PIGF	RDS ELISA Total sFlt-1	sFlt-1:PIGF Complex	RDS ELISA Total sFlt-1/RDS ELISA Free PIGF Ratio	sFlt-1:PIGF Complex/RDS ELISA Free PIGF Ratio
Mild or Severe PE vs. No PE (Gestation Age at Collection 13 - 21)					
W:	0.5745	0.5029	0.6331	0.5396	0.6392
SE:	0.0594	0.0602	0.0591	0.0599	0.0578
p:	0.1050	0.4805	0.0122	0.2543	0.0080

(continued)

Mild or Severe PE vs. No PE (Gestation Age at Collection 13 - 21)					
n:	94	94	94	94	94
Severe PE (Mild PE = Normal, Gestation Age at Collection 13 - 21))					
W:	0.7142	0.5622	0.6942	0.6771	0.7445
SE:	0.0819	0.0854	0.1029	0.0838	0.1004
p:	0.0045	0.2333	0.0296	0.0173	0.0075
n:	94	94	94	94	94
Severe PE (Mild PE = Normal, Gestation Age at Collection 16 - 21))					
W:	0.7394	0.5455	0.7636	0.6848	0.8707
SE:	0.1041	0.1073	0.1044	0.0963	0.0847
p:	0.0108	0.3360	0.0058	0.0274	<0.0001
n:	64	64	64	64	64

Claims

1. A method of determining the presence or amount of sFlt- 1 : PIGF complex, a complex of the soluble form of the fms-like tyrosine kinase receptor (sFlt-1) and placenta growth factor (PIGF), the method comprising:
 - (a) capturing the complex with capture agent that binds to PIGF, and detecting the complex with detector agent that binds to one or both of sFlt-1 and sFlt-1:PIGF complex; or
 - (b) capturing the complex with capture agent that binds to sFlt-1, and detecting the complex with detector agent that binds to one or both of PIGF and sFlt-1:PIGF complex; or
 - (c) capturing the complex with capture agent that binds to sFlt-1:PIGF complex, and detecting the complex with detector agent that binds to sFlt-1:PIGF complex, wherein capture agent and detector agent are capable of being bound simultaneously to the complex; or
 - (d) capturing the complex with capture agent that binds to sFlt-1:PIGF complex, and detecting the complex with detector agent that binds to one or both of PIGF and sFlt-1.
2. The method of claim 1 wherein capture agent is immobilized on a support or capable of being immobilized on a support.
3. The method of claim 2 wherein the support is a surface of a test tube, a microwell, a microtiter well, a membrane, a sheet, a bead, a microparticle, or a chip.
4. The method of claim 2 wherein capture agent is linked to biotin.
5. The method of claim 4 wherein capture agent linked to biotin is immobilized through binding to avidin, streptavidin, or other biotin-binding agent.
6. A method of guiding therapy of, or predicting, detecting, or monitoring preeclampsia or preeclampsia-related disease in an individual, said method comprising:
 - (a) determining the presence or amount of sFlt-1:PIGF complex in a sample from an individual; and
 - (b)
 - i) comparing the result of the sFlt-1:PIGF determination with one or more reference values; or

ii) converting the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the sFlt-1:PIGF complex transformed results with one or more reference values; thereby guiding therapy of, or predicting, detecting, or monitoring, the disease.

5

7. The method of claim 6 further comprising:

- (a) determining the presence or amount of one or more molecular entities in the sample; and
- (b)

10

i) comparing the result of the molecular entities determination and the result of the sFlt-1:PIGF complex determination with the one or more reference values; or

ii) converting the result of the molecular entities determination and the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the molecular entities transformed results and the sFlt-1:PIGF transformed results with the one or more reference values.

15

8. The method of claim 7 further comprising:

20

- (a) determining the values of one or more physical attributes of the individual; and
- (b)

i) comparing the physical attributes determination and the result of the molecular entities determination and the result of the sFlt-1: PIGF complex determination with the one or more reference values; or

ii) converting the physical attributes determination and the result of the molecular entities determination and the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the physical attributes transformed results and the molecular entities transformed results and the sFlt-1:PIGF transformed results with the one or more reference values.

25

30

9. The method of claim 6 further comprising:

- (a) determining the values of one or more physical attributes of the individual; and
- (b)

35

i) comparing the physical attributes determination and the result of the sFlt-1:PIGF complex determination with the one or more reference values; or

ii) converting the physical attributes determination and the result of the sFlt-1:PIGF complex determination into one or more transformed results, using one or more mathematical operations or one or more algorithms, and comparing the physical attributes transformed results and the sFlt-1:PIGF transformed results with the one or more reference values.

40

10. The method of claim 6 wherein the disease is preeclampsia or preeclampsia-related and guiding therapy, predicting, detecting, or monitoring is within the first 15 weeks of gestation or within the first 20 weeks of gestation or within the first 30 or more weeks of gestation.

45

11. The method of claim 7 wherein the one or more molecular entities is selected from free PIGF, free sFlt-1, free VEGF, sFlt-1:VEGF complex, endoglin, soluble endoglin, creatinine, c-reactive protein, liver enzymes, urine protein, placental protein 13, human chorionic gonadotropin hormone, alpha-fetoprotein, and autoantibodies to angiotensin II Type I receptor.

50

12. The method of claim 11 wherein VEGF is VEGF121, VEGF145, VEGF165, VEGF165b, VEGF167, VEGF183, VEGF186, VEGF189, or VEGF206.

55

13. The method of claim 7 wherein the one or more mathematical operations comprises computing a ratio of the molecular entities determination to the sFlt-1: PIGF complex determination or the sFlt-1:PIGF complex determination to the molecular entities determination.

14. The method of claim 8 or claim 9 wherein the one or more physical attributes is selected from gestational age, race, age, gender, blood pressure, and body mass index.

15. A composition for determining the presence or amount of sFlt-1:PIGF complex comprising:

(a) capture agent that binds to PIGF and detector agent that binds to one or both of sFlt-1 and sFlt-1:PIGF complex; or

(b) capture agent that binds to sFlt-1 and detector agent that binds to one or both of PIGF and sFlt-1 : PIGF complex; or

(c) capture agent that binds to sFlt-1:PIGF complex and detector agent that binds to sFlt-1: PIGF complex, wherein capture agent and detector agent are capable of being bound simultaneously to the complex; or

(d) capture agent that binds to sFlt-1: PIGF complex and detector agent that binds to one or both of PIGF and sFlt-1.

16. The method of claim 1 or the composition of claim 15 wherein one or both of capture agent and detector agent are polyclonal antibodies.

17. The method of claim 1 or the composition of claim 15 wherein one or both of capture agent and detector agent are monoclonal antibodies.

18. The method of claim 1 or the composition of claim 15 wherein one or both of capture agent and detector agent are a combination of monoclonal and polyclonal antibodies.

19. The method of claim 1 or claim 6 or the composition of claim 15 wherein PIGF is PIGF-1, PIGF-2, or PIGF-3.

Patentansprüche

1. Verfahren zur Bestimmung des Vorliegens bzw. der Menge von sFlt-1:PIGF-Komplex, einem Komplex aus der löslichen Form des fms-ähnlichen Tyrosinkinase-Rezeptors (sFlt-1) und Plazenta-Wachstumsfaktor (Placenta Growth Factor, PIGF), wobei das Verfahren Folgendes umfasst:

(a) Einfangen des Komplexes mit einem Einfangmittel, das an PIGF bindet, und Nachweisen des Komplexes mit einem Nachweismittel, das an sFlt-1 oder/und sFlt-1:PIGF-Komplex bindet; oder

(b) Einfangen des Komplexes mit einem Einfangmittel, das an sFlt-1 bindet, und Nachweisen des Komplexes mit einem Nachweismittel, das an PIGF oder/und sFlt-1:PIGF-Komplex bindet; oder

(c) Einfangen des Komplexes mit einem Einfangmittel, das an sFlt-1:PIGF-Komplex bindet, und Nachweisen des Komplexes mit einem Nachweismittel, das an sFlt-1:PIGF-Komplex bindet, wobei Einfangmittel und Nachweismittel gleichzeitig an den Komplex gebunden sein können; oder

(d) Einfangen des Komplexes mit einem Einfangmittel, das an sFlt-1:PIGF-Komplex bindet, und Nachweisen des Komplexes mit einem Nachweismittel, das an PIGF oder/und sFlt-1 bindet.

2. Verfahren nach Anspruch 1, wobei Einfangmittel auf einem Träger immobilisiert ist oder auf einem Träger immobilisiert werden kann.

3. Verfahren nach Anspruch 2, wobei es sich bei dem Träger um eine Oberfläche eines Reagensröhrchens, eines Microwell, eines Mikrotiter-Well, einer Membran, einer Folie, eines Kügelchens, eines Mikropartikels oder eines Chip handelt.

4. Verfahren nach Anspruch 2, wobei Einfangmittel mit Biotin verknüpft ist.

5. Verfahren nach Anspruch 4, wobei mit Biotin verknüpftes Einfangmittel durch Bindung an Avidin, Streptavidin oder ein anderes Biotin bindendes Mittel immobilisiert wird.

6. Verfahren zur Therapieführung bei oder zum Vorhersagen, Nachweisen oder Überwachen von Präeklampsie oder einer mit Präeklampsie verwandten Krankheit bei einem Individuum, wobei das Verfahren Folgendes umfasst:

(a) Bestimmen des Vorliegens bzw. der Menge von sFlt-1:PIGF-Komplex in einer Probe von einem Individuum;

und
(b)

- 5 i) Vergleichen des Ergebnisses der sFlt-1:PIGF-Bestimmung mit einem oder mehreren Referenzwerten; oder
ii) Umwandeln des Ergebnisses der sFlt-1:PIGF-Komplex-Bestimmung in ein oder mehrere transformierte Ergebnisse unter Verwendung einer oder mehrerer Rechenoperationen oder eines oder mehrerer Algorithmen und Vergleichen der transformierten sFlt-1:PIGF-Komplex-Ergebnisse mit einem oder mehreren Referenzwerten; wodurch die Therapieführung bei bzw. das Vorhersagen, Nachweisen oder Überwachen der Krankheit erfolgt.

7. Verfahren nach Anspruch 6, ferner umfassend:

- 15 (a) Bestimmen des Vorliegens bzw. der Menge eines oder mehrerer molekularer Gebilde in der Probe; und
(b)

- 20 i) Vergleichen des Ergebnisses der Bestimmung der molekularen Gebilde und des Ergebnisses der sFlt-1:PIGF-Komplex-Bestimmung mit dem einen bzw. den mehreren Referenzwerten; oder
ii) Umwandeln des Ergebnisses der Bestimmung der molekularen Gebilde und des Ergebnisses der sFlt-1:PIGF-Komplex-Bestimmung in ein oder mehrere transformierte Ergebnisse unter Verwendung einer oder mehrerer Rechenoperationen oder eines oder mehrerer Algorithmen und Vergleichen der transformierten Ergebnisse für die molekularen Gebilde und der transformierten sFlt-1:PIGF-Ergebnisse mit dem einen bzw. den mehreren Referenzwerten.

25 8. Verfahren nach Anspruch 7, ferner umfassend:

- (a) Bestimmen der Werte für ein oder mehrere physische Attribute des Individuums; und
(b)

- 30 i) Vergleichen der Bestimmung der physischen Attribute und des Ergebnisses der Bestimmung der molekularen Gebilde und des Ergebnisses der sFlt-1:PIGF-Komplex-Bestimmung mit dem einen bzw. den mehreren Referenzwerten; oder
ii) Umwandeln der Bestimmung der physischen Attribute und des Ergebnisses der Bestimmung der molekularen Gebilde und des Ergebnisses der sFlt-1:PIGF-Komplex-Bestimmung in ein oder mehrere transformierte Ergebnisse unter Verwendung einer oder mehrerer Rechenoperationen oder eines oder mehrerer Algorithmen und Vergleichen der transformierten Ergebnisse für die physischen Attribute und der transformierten Ergebnisse für die molekularen Gebilde und der transformierten sFlt-1:PIGF-Ergebnisse mit dem einen bzw. den mehreren Referenzwerten.

40 9. Verfahren nach Anspruch 6, ferner umfassend:

- (a) Bestimmen der Werte für ein oder mehrere physische Attribute des Individuums; und
(b)

- 45 i) Vergleichen der Bestimmung der physischen Attribute und des Ergebnisses der sFlt-1:PIGF-Komplex-Bestimmung mit dem einen bzw. den mehreren Referenzwerten; oder
ii) Umwandeln der Bestimmung der physischen Attribute und des Ergebnisses der sFlt-1:PIGF-Komplex-Bestimmung in ein oder mehrere transformierte Ergebnisse unter Verwendung einer oder mehrerer Rechenoperationen oder eines oder mehrerer Algorithmen und Vergleichen der transformierten Ergebnisse für die physischen Attribute und der transformierten sFlt-1:PIGF-Ergebnisse mit dem einen bzw. den mehreren Referenzwerten.

55 10. Verfahren nach Anspruch 6, wobei es sich bei der Krankheit um Präeklampsie handelt oder die Krankheit mit Präeklampsie verwandt ist und Therapieführung, Vorhersagen, Nachweisen oder Überwachen innerhalb der ersten 15 Schwangerschaftswochen oder innerhalb der ersten 20 Schwangerschaftswochen oder innerhalb der ersten 30 oder mehr Schwangerschaftswochen erfolgt.

11. Verfahren nach Anspruch 7, wobei das eine bzw. die mehreren molekulare(n) Gebilde aus freiem PIGF, freiem sFlt-

EP 2 235 529 B1

1, freiem VEGF, sFlt-1:VEGF-Komplex, Endoglin, löslichem Endoglin, Creatinin, c-reaktivem Protein, Leberenzymen, Harnprotein, Plazentaprotein 13, menschlichem Choriongonadotropin-Hormon, Alpha-Fetoprotein und Autoantikörpern gegen Angiotensin-II-Typ-I-Rezeptor ausgewählt ist bzw. sind.

- 5 12. Verfahren nach Anspruch 11, wobei es sich bei VEGF um VEGF121, VEGF145, VEGF165, VEGF165b, VEGF167, VEGF183, VEGF186, VEGF189 oder VEGF206 handelt.
- 10 13. Verfahren nach Anspruch 7, wobei die eine oder mehreren Rechenoperationen Berechnen eines Verhältnisses der Bestimmung der molekularen Gebilde zur sFlt-1:PIGF-Komplex-Bestimmung oder der sFlt-1:PIGF-Komplex-Bestimmung zur Bestimmung der molekularen Gebilde umfassen.
14. Verfahren nach Anspruch 8 oder Anspruch 9, wobei das eine bzw. die mehreren physische(n) Attribut(e) aus Gestationsalter, Rasse, Alter, Geschlecht, Blutdruck und Body-Mass-Index ausgewählt ist bzw. sind.
- 15 15. Zusammensetzung zur Bestimmung des Vorliegens bzw. der Menge von sFlt-1:PIGF-Komplex, umfassend:
- (a) Einfangmittel, das an PIGF bindet, und Nachweismittel, das an sFlt-1 oder/und sFlt-1:PIGF-Komplex bindet; oder
 - (b) Einfangmittel, das an sFlt-1 bindet, und Nachweismittel, das an PIGF oder/und sFlt-1:PIGF-Komplex bindet; oder
 - (c) Einfangmittel, das an sFlt-1:PIGF-Komplex bindet, und Nachweismittel, das an sFlt-1:PIGF-Komplex bindet, wobei Einfangmittel und Nachweismittel gleichzeitig an den Komplex gebunden sein können; oder
 - (d) Einfangmittel, das an sFlt-1:PIGF-Komplex bindet, und Nachweismittel, das an PIGF oder/und sFlt-1 bindet.
- 25 16. Verfahren nach Anspruch 1 oder Zusammensetzung nach Anspruch 15, wobei es sich bei Einfangmittel oder/und Nachweismittel um polyklonale Antikörper handelt.
17. Verfahren nach Anspruch 1 oder Zusammensetzung nach Anspruch 15, wobei es sich bei Einfangmittel oder/und Nachweismittel um monoklonale Antikörper handelt.
- 30 18. Verfahren nach Anspruch 1 oder Zusammensetzung nach Anspruch 15, wobei es sich bei Einfangmittel oder/und Nachweismittel um eine Kombination monoklonaler und polyklonaler Antikörper handelt.
- 35 19. Verfahren nach Anspruch 1 oder Anspruch 6 oder Zusammensetzung nach Anspruch 15, wobei es sich bei PIGF um PIGF-1, PIGF-2 oder PIGF-3 handelt.

Revendications

- 40 1. Procédé de détermination de la présence ou la quantité de complexe sFlt-1:PIGF, un complexe de la forme soluble du récepteur de la tyrosine kinase de type fms (sFlt-1) et du facteur de croissance placentaire (PIGF), le procédé comprenant les étapes consistant à :
- (a) capturer le complexe avec un agent de capture, qui se lie au PIGF, et détecter le complexe avec un agent de détection qui se lie au sFlt-1 ou au complexe sFlt-1:PIGF ou bien aux deux ; ou
 - (b) capturer le complexe avec un agent de capture, qui se lie au sFlt-1, et détecter le complexe avec un agent de détection qui se lie au PIGF ou au complexe sFlt-1:PIGF ou bien aux deux ; ou
 - (c) capturer le complexe avec un agent de capture, qui se lie au complexe sFlt-1:PIGF, et détecter le complexe avec un agent de détection qui se lie au complexe sFlt-1:PIGF, dans lequel l'agent de capture et l'agent de détection sont capables d'être liés simultanément au complexe ; ou
 - (d) capturer le complexe avec un agent de capture, qui se lie au complexe sFlt-1:PIGF, et détecter le complexe avec un agent de détection qui se lie au PIGF ou au sFlt-1 ou bien aux deux.
- 45
- 50
- 55 2. Procédé selon la revendication 1, dans laquelle l'agent de capture est immobilisé sur un support ou est capable d'être immobilisé sur un support.
3. Procédé selon la revendication 2, dans laquelle le support est une surface d'un tube à essai, un micro-puits, un puits de micro-titration, une membrane, une feuille, une bille, une microparticule ou une puce.

EP 2 235 529 B1

4. Procédé selon la revendication 2, dans laquelle l'agent de capture est lié à la biotine.
5. Procédé selon la revendication 4, dans laquelle l'agent de capture lié à la biotine est immobilisé au moyen d'une liaison à l'avidine, la streptavidine ou un autre agent de liaison à la biotine.

5

6. Procédé de guidage d'une thérapie de la pré-éclampsie, ou de prédiction, détection ou suivi de celle-ci, ou d'une maladie apparentée à la pré-éclampsie chez un individu, ledit procédé comprenant les étapes consistant à :

10

- (a) déterminer la présence ou la quantité de complexe sFlt-1:PIGF dans un échantillon provenant d'un individu ; et
(b)

15

- i) comparer le résultat de la détermination du sFlt-1:PIGF avec une ou plusieurs valeur(s) de référence ; ou
ii) convertir le résultat de la détermination du complexe sFlt-1:PIGF en un ou plusieurs résultat(s) transformé(s), en utilisant une ou plusieurs opération (s) mathématique(s) ou bien un ou plusieurs algorithme(s), et comparer les résultats transformés du complexe sFlt-1:PIGF avec une ou plusieurs valeur(s) de référence ; en guidant, de ce fait, une thérapie de la maladie ou la prédiction, la détection ou le suivi de celle-ci.

20

7. Procédé, selon la revendication 6, comprenant en outre les étapes consistant à :

- (a) déterminer la présence ou la quantité d'une ou plusieurs entité(s) moléculaire(s) dans l'échantillon ;
et
(b)

25

- i) comparer le résultat de la détermination des entités moléculaires et le résultat de la détermination du complexe sFlt-1:PIGF avec une ou plusieurs valeur(s) de référence ; ou

30

- ii) convertir le résultat de la détermination des entités moléculaires et le résultat de la détermination du complexe sFlt-1:PIGF en un ou plusieurs résultat(s) transformé(s), en utilisant une ou plusieurs opération (s) mathématique(s) ou bien un ou plusieurs algorithme(s), et comparer les résultats transformés des entités moléculaires et les résultats transformés du sFlt-1:PIGF avec l'une ou les plusieurs valeur(s) de référence.

8. Procédé, selon la revendication 7, comprenant en outre les étapes consistant à :

35

- (a) déterminer les valeurs d'un ou plusieurs attribut(s) physique(s) de l'individu ; et
(b)

40

- i) comparer la détermination des attributs physiques et le résultat de la détermination des entités moléculaires et le résultat de la détermination du complexe sFlt-1:PIGF avec l'une ou les plusieurs valeur(s) de référence ; ou

45

- ii) convertir la détermination des attributs physiques et le résultat de la détermination des entités moléculaires et le résultat de la détermination du complexe sFlt-1:PIGF en un ou plusieurs résultat(s) transformé(s), en utilisant une ou plusieurs opération (s) mathématique(s) ou bien un ou plusieurs algorithme(s), et comparer les résultats transformés des attributs physiques et les résultats transformés des entités moléculaires et les résultats transformés du sFlt-1:PIGF avec l'une ou les plusieurs valeur(s) de référence.

9. Procédé, selon la revendication 6, comprenant en outre les étapes consistant à :

50

- (a) déterminer les valeurs d'un ou plusieurs attribut(s) physique(s) de l'individu ; et
(b)

55

- i) comparer la détermination des attributs physiques et le résultat de la détermination du complexe sFlt-1:PIGF avec l'une ou les plusieurs valeur(s) de référence ; ou

- ii) convertir la détermination des attributs physiques et le résultat de la détermination du complexe sFlt-1:PIGF en un ou plusieurs résultat (s) transformé(s), en utilisant une ou plusieurs opération(s) mathématique(s) ou bien un ou plusieurs algorithme(s), et comparer les résultats transformés des attributs physiques et les résultats transformés du sFlt-1:PIGF avec l'une ou les plusieurs valeur(s) de référence.

10. Procédé selon la revendication 6, dans laquelle la maladie est une pré-éclampsie ou est apparentée à une pré-

EP 2 235 529 B1

éclampsie, et la thérapie de guidage, la prédiction, la détection ou le suivi a lieu dans les 15 premières semaines de gestation ou dans les 20 premières semaines de gestation ou dans les 30 premières semaines de gestation ou plus.

- 5
11. Procédé selon la revendication 7, dans laquelle l'une ou les plusieurs entité(s) moléculaire(s) est (sont) choisie(s) parmi les : PIGF libre, sFlt-1 libre, VEGF libre, complexe sFlt-1:VEGF, endogline, endogline soluble, créatinine, protéine c réactive, enzymes hépatiques, protéine de l'urine, protéine 13 placentaire, hormone chorionique gonadotrope humaine, alpha-foetoprotéine et auto-anticorps contre le récepteur de type I de l'angiotensine II.
- 10
12. Procédé selon la revendication 11, dans laquelle le VEGF est un VEGF121, VEGF145, VEGF165, VEGF165b, VEGF167, VEGF183, VEGF186, VEGF189 ou VEGF206.
13. Procédé selon la revendication 7, dans laquelle l'une ou les plusieurs opération(s) mathématique(s) comprend (comprennent) le calcul d'un rapport de la détermination des entités moléculaires à la détermination du complexe sFlt-1:PIGF ou de la détermination du complexe sFlt-1:PIGF à la détermination des entités moléculaires.
- 15
14. Procédé selon la revendication 8 ou la revendication 9, dans laquelle l'un ou les plusieurs attribut(s) physique(s) est (sont) choisi (s) parmi l'âge de gestation, la race, l'âge, le genre, la pression sanguine et l'indice de masse corporelle.
- 20
15. Composition destinée à déterminer la présence ou la quantité de complexe sFlt-1:PIGF, comprenant :
- (a) un agent de capture, qui se lie au PIGF, et un agent de détection qui se lie au sFlt-1 ou au complexe sFlt-1:PIGF ou bien aux deux ; ou
- 25
- (b) un agent de capture, qui se lie au sFlt-1, et un agent de détection qui se lie au PIGF ou au complexe sFlt-1:PIGF ou bien aux deux ; ou
- (c) un agent de capture, qui se lie au complexe sFlt-1:PIGF, et un agent de détection qui se lie au complexe sFlt-1:PIGF, dans lequel l'agent de capture et l'agent de détection sont capables d'être liés simultanément au complexe ; ou
- 30
- (d) un agent de capture, qui se lie au complexe sFlt-1:PIGF, et un agent de détection qui se lie au PIGF ou au sFlt-1 ou bien aux deux.
16. Procédé, selon la revendication 1, ou composition, selon la revendication 15, dans laquelle l'agent de capture ou l'agent de détection, ou les deux, sont des anticorps polyclonaux.
- 35
17. Procédé, selon la revendication 1, ou composition, selon la revendication 15, dans laquelle l'agent de capture ou l'agent de détection, ou les deux, sont des anticorps monoclonaux.
18. Procédé, selon la revendication 1, ou composition, selon la revendication 15, dans laquelle l'agent de capture ou l'agent de détection, ou les deux, sont une combinaison d'anticorps monoclonaux et polyclonaux.
- 40
19. Procédé, selon la revendication 1 ou la revendication 6, ou composition, selon la revendication 15, dans laquelle le PIGF est un PIGF-1, un PIGF-2 ou un PIGF-3.

45

50

55

Figure 1




Figure 2

sFlt-1:PIGF Complex Assay
Dose Response Curve




Figure 3

sFlt-1: PIGF Complex
sFlt-1: PIGF Complex vs. Classification





Figure 4

sFlt-1: PIGF Complex
sFlt-1: PIGF Complex/Free PIGF vs. Classification



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 4200690 A [0033]
- US 4207307 A [0033]
- US 4407943 A [0033]
- US 4550075 A [0033]
- US 4551426 A [0033]
- US 4560648 A [0033]
- US 5312744 A [0033]
- US 5314830 A [0033]
- US 5424220 A [0033]
- US 5415998 A [0033]
- US 5501949 A [0033]
- US 5518887 A [0033]
- US 5663054 A [0033]
- US 5789261 A [0033]
- US 5935780 A [0033] [0038]
- US 5958339 A [0033]
- US 6087188 A [0033]
- US 6096563 A [0033]
- US 6121006 A [0033]
- US 6143575 A [0033]
- US 6395503 B [0033]
- US 6878515 B [0033]
- US 6838250 B [0033]
- US 4372745 A [0033]
- US 4670381 A [0033]
- US 4483921 A [0033]
- US 4517288 A [0033]
- US 4822747 A [0033]
- US 4824778 A [0033]
- US 4829012 A [0033]
- US 4839299 A [0033]
- US 4847194 A [0033]
- US 4847195 A [0033]
- US 4853335 A [0033]
- US 4855226 A [0033]
- US 4857453 A [0033]
- US 4857454 A [0033]
- US 4859610 A [0033]
- US 4863876 A [0033]
- US 4868106 A [0033]
- US 4868130 A [0033]
- US 4879219 A [0033]
- US 5776933 A [0033]
- US 3867258 A [0034]
- US 3992158 A [0034]
- US 4042435 A [0034]
- US 4050898 A [0034]
- US 4066403 A [0034]
- US 4153668 A [0034]
- US 4258001 A [0034]
- US 4292272 A [0034]
- US 4430436 A [0034]
- US 5006309 A [0036]
- US 5089424 A [0036]
- US 4624930 A [0038]
- US 5061640 A [0038]
- US 4945042 A [0038]
- US 4885255 A [0038]
- US 5362624 A [0038]
- US 5277589 A [0038]
- US 5268306 A [0038]
- US 5376557 A [0038]
- US 5858803 A [0038]
- US 5126241 A [0038]
- US 5362655 A [0038]
- US 5437981 A [0038]
- US 5792606 A [0038]
- US 5863740 A [0038]
- US 6391571 B [0038]
- US 2006381104 A [0065]
- US 2003700672 A [0065]
- US 2003410572 A [0065]
- US 2003634145 A [0065]
- US 2005323460 A [0065]
- WO 2007044860 A [0065]
- US 2002330696 A [0065]
- US 2002331127 A [0065]
- US 1997912133 B [0065]
- US 6306087 B [0065]
- US 6248063 B [0065]
- US 5769074 A [0065]

Non-patent literature cited in the description

- **MUTTER ; KARUMANCHI.** *Microvascular Research*, 2008, vol. 75, 1-8 [0004]
- **ROBERTS ; COOPER.** *Lancet*, 2001, vol. 357, 53-56 [0004]
- **ROBERTS.** *Semin Reprod Endocrinol*, 1998, vol. 16, 5-15 [0004]
- **HE et al.** *Mol Endocrinol*, 1999, vol. 13, 537-545 [0006]

EP 2 235 529 B1

- **KENDALL ; THOMAS.** *Proc Natl Acad Sci USA*, 1993, vol. 90, 10705-10709 [0006]
- **AHMAD ; AHMED.** *Circ Res*, 2004, vol. 95, 884-891 [0007]
- **CHAIWORAPONGSA et al.** *Am J Obstet Gynecol*, 2004, vol. 190, 1541-1547 [0007]
- **KOGA et al.** *J Clin Endocrinol Metab*, 2003, vol. 88, 2348-2351 [0007]
- **MAYNARD et al.** *J Clin Invest*, 2003, vol. 111, 649-658 [0007]
- **SHIBATA et al.** *J Clin Endocrinol Metab*, 2005, vol. 90, 4895-4903 [0007]
- **KENDALL et al.** *Biochem Biophys Res Commun*, 1996, vol. 226, 324-328 [0007]
- **DUNK et al.** *Am J Pathol*, 2000, vol. 156, 2185-2199 [0007]
- **GEVA et al.** *J Clin Endocrinol Metab*, 2002, vol. 87, 4213-4224 [0007]
- **GOLDMAN-WOHL et al.** *Mol Hum Reprod*, 2000, vol. 6, 81-87 [0007]
- **LEVINE et al.** *N Engl J Med*, 2006, vol. 355, 992-1005 [0008]
- **CHAIWORAPONGSA et al.** *J Matern Fetal Neonatal Med*, 2005, vol. 17, 3-18 [0009]
- **HERTIG et al.** *Clin Chem*, 2004, vol. 50, 1702-1703 [0009]
- **LEVINE et al.** *N Engl J Med*, 2004, vol. 350, 672-683 [0009]
- **MCKEEMAN et al.** *Am J Obstet Gynecol*, 2004, vol. 191, 1240-1246 [0009]
- **WIDMER et al.** *Obstet Gynecol*, 2007, vol. 109, 168-180 [0009]
- **Practical Immunoassay.** Marcel Dekker, 1984 [0033]
- **BUTLER.** *Immunochemistry of Solid Phase-Immunoassay.* CRC Press, 1991 [0033]
- **Immunoassay.** Taylor & Francis, 1996 [0033]
- **ELEFThERIOS et al.** *Immunoassay.* Academic Press, 1996 [0033]
- **Principles and Practice of Immunoassay.** Macmillan, 1997 [0033]
- **Immunoassays in the Clinical Laboratory.** Alan R Liss, Inc, 1979 [0033]
- **Quantitative Enzyme Immunoassay.** Blackwell Scientific Publications, 1978 [0033]
- **SOMMER et al.** *Clin Chem*, 1986, vol. 32, 1770-1774 [0033]
- **A Primer for Multilayer Immunoassay.** **BERKE.** American Chemical Society Conference Proceeding. Plenum Press, 1988, 303-312 [0033]

专利名称(译)	sflt-1的测定：血管生成因子复合物		
公开(公告)号	EP2235529B1	公开(公告)日	2016-12-14
申请号	EP2009700404	申请日	2009-01-07
[标]申请(专利权)人(译)	奥索临床诊断有限公司		
申请(专利权)人(译)	邻临床诊断中，INC.		
当前申请(专利权)人(译)	邻临床诊断中，INC.		
[标]发明人	BASHIRIANS GEORGE BYRNE DEBORAH GROULX SARAH JACKSON SHARON LANE BROWN HEATHER SANKARAN BANUMATHI SCALICE EDWARD YANG KAREN ZHENG JIAN		
发明人	BASHIRIANS, GEORGE BYRNE, DEBORAH GROULX, SARAH JACKSON, SHARON LANE-BROWN, HEATHER SANKARAN, BANUMATHI SCALICE, EDWARD YANG, KAREN ZHENG, JIAN		
IPC分类号	G01N33/53 G01N33/00 G01N33/68 G01N33/74 G01N33/574		
CPC分类号	G01N33/689 G01N33/543 G01N33/566 G01N33/6872 G01N2333/475 G01N2333/515 G01N2333/71 G01N2333/91205 G01N2800/368 G01N2800/50 G01N2800/56		
优先权	61/019351 2008-01-07 US		
其他公开文献	EP2235529A4 EP2235529A1		
外部链接	Espacenet		

摘要(译)

提供了用于确定包含第一和第二分子实体的复合物的存在或量的方法，优选sFlt-1：PIGF复合物。确定复合物的存在或量可用于预测，检测，监测疾病或指导治疗疾病的方法，例如血管，血管相关疾病，心脏，心脏相关疾病，癌症，癌症相关疾病，先兆子痫和先兆子痫相关疾病。确定sFlt-1：血管生成因子复合物特别适用于预测和检测妊娠早期和疾病阶段的先兆子痫，其中临床评估可能是无效的。

Table 4
 Patient results: sFlt-1:PlGF complex, free PlGF and
 sFlt-1:PlGF complex/free PlGF ratio

Patient ID	Gestational Age at Collection (Weeks)	Diagnosis	Classification (Pre-eclampsia, Normal)	sFlt-1:PlGF Complex Concentration (pg/mL)	Free PlGF Concentration (pg/mL)	sFlt-1:PlGF Complex/Free PlGF Ratio
N47	31 3/4	pre-eclampsia	1	316	23.3	13.6
N49	35	pre-eclampsia	1	3553	72.4	49.1
N50	30 6/7	pre-eclampsia	1	1421	6.16	230.7
N51	21 1/3	pre-eclampsia w/HELLP	1	717	5	143.3
N52	33 4/7	pre-eclampsia	1	79	4.02	19.6
N53	25 3/7	pre-eclampsia w/HELLP	1	432	5	86.4
N54	29	pre-eclampsia w/HELLP	1	1093	5	218.6
N55	26 5/7	pre-eclampsia	1	2437	27.2	89.6
N56	33 5/7	pre-eclampsia	1	2031	158	12.9
N58	38 4/7	normal	0	6	74.5	0.1
N59	22	normal	0	170	129	1.3
N60	38 6/7	normal	0	1122	508	2.2
N61	36 2/7	normal	0	42	40.5	1.0
N62	29 3/7	gestational hypertension + pre-eclampsia	1	5629	45.9	122.6
N63	30	IUGR + gestational hypertension	0	1332	6.25	213.1
N64	36	normal	0	629	32.1	19.6
N65	32 5/7	normal	0	281	36.5	7.7
N67	39	normal	0	5	5	1.0
N68	37 5/7	normal	0	446	320	1.4