



(11)

EP 2 815 238 B1

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
08.04.2020 Bulletin 2020/15

(51) Int Cl.:
G01N 33/53 (2006.01) G01N 33/68 (2006.01)

(21) Application number: **13749422.5**

(86) International application number:
PCT/IB2013/051193

(22) Date of filing: **14.02.2013**

(87) International publication number:
WO 2013/121368 (22.08.2013 Gazette 2013/34)

(54) **A PROCESS FOR DETECTION AND OPTIONAL QUANTIFICATION OF AN ANALYTE**
VERFAHREN ZUR DETEKTION UND OPTIONALEN QUANTIFIZIERUNG EINES ANALYTEN
PROCÉDÉ DE DÉTECTION ET DE QUANTIFICATION FACULTATIVE D'UN ANALYTE

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(30) Priority: **15.02.2012 IN 566CH2012**

(43) Date of publication of application:
24.12.2014 Bulletin 2014/52

(73) Proprietors:
• **Biocon Limited**
Bangalore 560 100
Karnataka (IN)
• **Mylan GmbH**
8050 Zurich (CH)

(72) Inventors:
• **SENGUPTA, Nilanjan**
Bangalore 560 076 (IN)
• **UDIAVAR, Anita, Rao**
Domlur
Bangalore 560 071 (IN)
• **THANGAMUTHU, Senthil**
Bhavani
Erode (Dist)
Tamil Nadu 638 301 (IN)
• **NAYAK, Vivek, Gopal**
Jaraganahalli J P Nagar Post
Bangalore 560 078 (IN)
• **PILLAI, Ranjit, Ravindran**
Bangalore 560 029 (IN)
• **VASHISHTA, Laxmikant**
Bangalore 560 043 (IN)

• **RAMASWAMY, Shilpa, Govinda**
Hosur Taluk
Krishnagiri 635 126 (IN)
• **HEGDE, Vishika**
Rajarajeshwari Nagar
Bangalore 560 098 (IN)
• **MELARKODE, Ramakrishnan**
Bangalore 560 035 (IN)

(74) Representative: **Michalski Hüttermann & Partner**
Patentanwälte mbB
Speditionstraße 21
40221 Düsseldorf (DE)

(56) References cited:
US-A- 4 604 348 US-A- 5 641 640
US-A1- 2002 090 662 US-A1- 2008 182 796
US-A1- 2011 177 095

• **LEE JEAN W ET AL: "Bioanalytical Approaches to Quantify Total and Free Therapeutic Antibodies and Their Targets: Technical Challenges and PK/PD Applications Over the Course of Drug Development", THE AAPS JOURNAL, SPRINGER US, BOSTON, vol. 13, no. 1, 15 January 2011 (2011-01-15), pages 99-110, XP035719169, DOI: 10.1208/S12248-011-9251-3 [retrieved on 2011-01-15]**
• **MAPLE, L. ET AL.: 'Development and validation of ELISA for Herceptin detection in human serum' JOURNAL OF IMMUNOLOGICAL METHODS vol. 295, no. 1-2, 2004, pages 169 - 182, XP004701203**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 815 238 B1

- ROCHE, J. A. ET AL.: 'Unmasking Potential Intracellular Roles for Dysferlin through Improved Immunolabeling Methods' JOURNAL OF HISTOCHEMISTRY AND CYTOCHEMISTRY vol. 59, no. 11, 2011, pages 964 - 975, XP055293970

Description**TECHNICAL FIELD OF THE DISCLOSURE**

5 **[0001]** The present disclosure relates to a process of utilizing a unique acidic dissociation buffer in a single step ligand binding assay to quantify total amount of an analyte present in a biological matrix containing a soluble target of the analyte. More specifically, the disclosure relates to a process for the development of immunoassays using an ELISA format to reliably quantify the total analyte using this unique buffer. The disclosure can be extended to immunogenicity assays for the detection of anti-analyte antibodies in the presence of excess free analyte.

BACKGROUND OF THE DISCLOSURE

10 **[0002]** Immunoassays are primarily used for the detection and quantitation of a wide variety of analytes such as hormones, tumor antigens, bacterial or viral antigens, protein therapeutics etc. in biological samples. Quantitative immunoassays are critical for understanding the Pharmacokinetic profile of the therapeutic proteins in support of efficacy and safety studies. In addition, immunoassays are also used to quantitate the soluble targets of the therapeutic proteins and anti-therapeutic protein antibodies. For soluble protein targets or shed receptor targets of the analyte, two different forms of the analyte exist in the biological matrix: the free analyte and the analyte complexed with the soluble target. In order to get an accurate pharmacokinetic profile of the analyte, it is ideal to develop ligand binding assay methods aimed at quantitating the total analyte concentrations instead of only the free analyte concentrations.

15 **[0003]** A frequent problem often encountered in quantitative immunoassays is accurate estimation of the total analyte concentrations in the presence of an interfering substance (soluble target) which competes with the immunoassay reagents for binding to the analyte. This leads to an underestimation of the analyte and compromises the true assessment of the pharmacokinetic profile of the analyte. The interfering substance can be removed by pre-treatment of the samples. However this step leads to loss of sample and introduces additional variability.

20 **[0004]** US 4,604,348 discloses a process for determining the presence of an antigen or antibody in a serum containing such antigen or antibody in the form an immune complex. US 5,641,640 discloses a method for assaying for an analyte in a fluid sample in which a change in refractive index at a solid optical surface in contact with the sample is measured. US 2008/182796 relates to modified Cry3A toxins and methods of making and using the same, as well as methods of detecting these toxins. US 2002/0090662 relates to a method for determining the amount of an analyte in a fluid sample in the presence of an interfering substance, such as the amount of circulating Herceptin® in serum or plasma of cancer patients undergoing Herceptin® therapy.

STATEMENT OF THE DISCLOSURE

25 **[0005]** The present invention is defined by the process of claim 1, the method of claim 6 and the use of claim 7.

30 **[0006]** The present disclosure also relates to a process for detection and optional quantification of analyte in biological sample, said process comprising providing modified citrate buffer to the sample prior to carrying out assay for said detection and optional quantification of the analyte within the biological sample; a method of disassociating complex formed by interaction of analyte and at least one interfering substance as target for the analyte in a biological sample, said method comprising act of providing modified citrate buffer to the biological sample for disassociating the complex; an enzyme linked immunosorbent assay (ELISA) kit for detecting and optionally quantifying analyte in biological sample, said kit comprising components selected from group comprising modified citrate buffer, capture antibody, detection antibody, reagent for reacting with the detection antibody, microtitre plates, coating buffer, washing solution, and stopping solution or any combination thereof; and a modified citrate buffer for detecting and optionally quantifying presence of an analyte in a biological sample.

BRIEF DESCRIPTION OF THE ACCOMPANYING FIGURES

35 **[0007]** In order that the disclosure may be readily understood and put into practical effect, reference is made to exemplary embodiments as illustrated with reference to the accompanying figures. The figures together with a detailed description below, are incorporated in and form part of the specification, and serve to further illustrate the embodiments and explain various principles and advantages, in accordance with the present disclosure where:

40 **Figure 1** shows graphs depicting interference of recombinant HER2-ECD on a Trastuzumab calibration curve.

45 **Figure 1A** shows a graph depicting that spiking of HER2-ECD results in a dose-dependent locational shift of Trastuzumab calibration curves when diluted in phosphate buffered saline Tween20 (PBST).

Figure 1B shows a graph depicting that modified citrate buffer diluent nullifies the dose-dependent interference of HER2-ECD on a Trastuzumab calibration curve.

Figure 2 shows the comparison of different acidic buffers at pH 3.5, to nullify the interference of recombinant HER2-ECD on a Trastuzumab calibration curve. Particularly, **Figure 2** is a graph showing that use of modified citrate buffer diluent results in a minimal locational shift induced by 1 µg/mL HER2-ECD spiking in serum.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0008] The present disclosure relates to a process for detection and optional quantification of an analyte in a biological sample, said process comprising providing modified citrate buffer to the sample prior to carrying out assay for said detection and optional quantification of the analyte within the biological sample.

[0009] In an embodiment of the present disclosure, the process comprises:

- (a) providing modified citrate buffer to the sample for obtaining diluted biological sample; and
- (b) carrying out a conventional assay for detecting and optionally quantifying the analyte in the sample.

[0010] In another embodiment of the present disclosure, the biological sample comprises at least one interfering substance as a target for the analyte, and wherein said interfering substance and said analyte combine to form a complex.

[0011] In yet another embodiment of the present disclosure, the formation of the complex reduces amount of free analyte in the biological sample, thereby hindering the accurate detection and quantification of said analyte.

[0012] In still another embodiment of the present disclosure, the modified citrate buffer dissociates the complex and restores the free analyte within the sample to enable the accurate detection and quantification of said analyte.

[0013] In still another embodiment of the present disclosure, the modified citrate buffer comprises Citrate-HCl at concentration of about 125mM and Tween-20 at concentration of about 0.1% w/v, and wherein pH of the buffer ranges from 2.5 to 3.8, preferably 3.5.

[0014] In still another embodiment of the present disclosure, the biological sample is selected from group comprising blood, plasma, and serum or any combinations thereof.

[0015] In still another embodiment of the present disclosure, the analyte is selected from group comprising a hormone, a tumor antigen, a bacterial antigen, a viral antigen, a protein therapeutic and an antibody or any combinations thereof.

[0016] In still another embodiment of the present disclosure, the analyte is an antibody.

[0017] In still another embodiment of the present disclosure, the antibody is against human epidermal growth factor receptor 2 (HER-2).

[0018] In still another embodiment of the present disclosure, the interfering substance is an extracellular domain of HER-2 (HER-2-ECD).

[0019] In still another embodiment of the present disclosure, the assay is enzyme linked immunosorbent assay (ELISA).

[0020] In still another embodiment of the present disclosure, the enzyme linked immunosorbent assay is sandwich enzyme linked immunosorbent assay and is carried out in presence of components selected from group comprising a capture antibody, a detection antibody, a reagent for reacting with the detection antibody, a microtitre plate, a coating buffer, a washing solution and a stopping solution or any combinations thereof.

[0021] In still another embodiment of the present disclosure, the capture antibody is an antibody against the analyte within the biological sample.

[0022] In still another embodiment of the present disclosure, the detection antibody is labelled with an enzyme selected from group comprising horseradish peroxidase and alkaline phosphatase and ruthenium.

[0023] In still another embodiment of the present disclosure, the reagent for reacting with the labelled antibody is chromogenic substrate selected from ABTS (2,2'-Azinobis[3-ethylbenzothiazoline-6-sulfonic acid]-diammonium salt, TMB (3,3',5,5'-tetramethylbenzidine) and electro chemiluminescent compatible substrate.

[0024] In still another embodiment of the present disclosure, the coating buffer is phosphate buffered saline.

[0025] In still another embodiment of the present disclosure, the washing solution comprises phosphate buffered saline and Tween-20.

[0026] In still another embodiment of the present disclosure, the stopping solution is 0.1N sulphuric acid.

[0027] In still another embodiment of the present disclosure, the enzyme linked immunosorbent assay (ELISA) comprises:

- a. coating a solid surface with the capture antibody against the analyte; providing a modified citrate buffer to the biological sample for obtaining a diluted biological sample;
- b. adding the diluted sample to the antibody coated solid surface or support of step (a);
- c. incubating and washing the support to remove any unbound analyte;

- d. adding a labelled detection antibody against the analyte and washing the support to remove any unbound detection antibody; and
- e. adding a reagent for reacting with the labelled antibody for detecting and optionally quantifying the analyte in the sample.

5

[0028] The present disclosure further relates to a method of disassociating complex formed by interaction of analyte and at least one interfering substance as target for the analyte in a biological sample, said method comprising act of providing modified citrate buffer to the biological sample for dissociating the complex.

10

[0029] The present disclosure further relates to an enzyme linked immunosorbent assay (ELISA) kit for detecting and optionally quantifying an analyte in a biological sample, said kit comprising components selected from group comprising modified citrate buffer, a capture antibody, a detection antibody, reagent for reacting with the detection antibody, microtitre plates, coating buffer, washing solution, and stopping solution or any combination thereof.

[0030] The present disclosure further relates to a modified citrate buffer for detecting and optionally quantifying presence of an analyte in a biological sample.

15

[0031] In an embodiment of the present disclosure, the modified citrate buffer comprises citrate-HCl at concentration of about 125 mM and polysorbate-20 at concentration of about 0.1% w/v, and wherein pH of the buffer ranges from 2.5 to 3.8, preferably 3.5.

[0032] In another embodiment of the present disclosure, the biological sample comprises at least one interfering substance as a target for the analyte, and wherein said interfering substance and said analyte interact to form complex.

20

[0033] In yet another embodiment of the present disclosure, the analyte is a human epidermal growth factor receptor 2 (HER-2) antibody.

[0034] In still another embodiment of the present disclosure, the interfering substance is an extracellular domain of HER-2 (HER-2-ECD).

25

DEFINITION OF TERMS

[0035] Modified Citrate Buffer- A Modified Citrate Buffer according to the present disclosure is a citrate buffer comprising sodium citrate and citric acid also comprising polysorbate 20 and HCl. The HCl is added in order to bring down the pH of the buffer to a desired range of 2.5-3.8.

30

[0036] The present disclosure relates to a process of utilizing a unique acidic dissociation buffer in a single step ligand binding assay/analytical assay to accurately quantitate the total amount of an analyte present in the biological matrix containing a soluble target of the analyte (also referred to herein as interfering substance). Using this buffer resolves the problem of underestimation of the analyte due to the presence of the interfering substance, without introducing any pre-treatment of samples, thus preserving the simplicity and ease of immunoassays. The process comprises the following steps:

35

- a) Coating a solid surface such as a microtiter plate with an antibody specific to the analyte.
- b) Diluting the biological samples containing the analyte and the interfering substance in modified Citrate buffer and adding the diluted samples to the coated solid surface from the previous step.
- c) Detecting the capture antibody-analyte complex by adding a labeled antibody against the analyte.
- d) Adding a reagent capable of reacting with the labeled antibody and quantitating the signal.

40

[0037] In an embodiment of the present disclosure Modified Citrate buffer is prepared by modifying a citrate buffer by addition of a strong acid, i.e., the modified citrate buffer of the present disclosure is a combination of the salt of a weak acid (Sodium citrate) and a weak acid (citric acid) with the addition of a strong acid to obtain the required pH of between 2.5 to 3.8.

45

[0038] In one aspect, the modified citrate buffer contains polysorbate. In another aspect, the modified citrate buffer contains polysorbate-20.

[0039] In yet another embodiment of the present disclosure, the capture and the detection antibodies may be of monoclonal or polyclonal origin. The epitope on the analyte recognized by the detection antibody is different from the epitope recognized by the capture antibody. The label conjugated to the detection antibody may be an enzyme such as horse radish peroxidase, alkaline phosphatase or other enzymes. A substrate for the enzyme conjugated to the detection antibody is used as an indicator. Common substrates used are selected from ABTS (2,2'-Azinobis[3-ethylbenzothiazoline-6-sulfonic acid]-diammonium salt and TMB (3,3',5,5'-tetramethylbenzidine).

50

[0040] In still another embodiment, the present disclosure relates to a process of utilizing a unique acidic dissociation buffer in a single step ligand binding assay to quantitate total amount of analyte present in the serum matrix containing the interfering substance.

55

[0041] In still another embodiment, the disclosure discloses a method of accurately determining the levels of an anti-

HER2 antibody in the presence of an interfering substance, an extracellular domain (ECD) of HER2 receptors. The biological sample in which the accurate determination of the level of anti-HER2 antibody is sought may be serum or plasma derived from a cancer patient undergoing anti-HER2 therapy.

In still another embodiment of the present disclosure, accuracy (analytical recovery) of trastuzumab QCs made in metastatic breast cancer sera containing endogenous HER2-ECD is maximum when modified citrate buffer is used as diluent.

[0042] The present disclosure is illustrated by its application to the accurate analytical recovery of trastuzumab in serum spiked with recombinant HER2-ECD, the soluble target of trastuzumab. Over-expression of the human epidermal growth factor receptor 2 (HER-2/neu) occurs in 25% of human breast cancers and is associated with poor prognosis. Trastuzumab is a humanized monoclonal antibody against the extracellular domain (ECD) of HER-2/neu receptor and binding results in inhibition of cancer cell proliferation. Trastuzumab is approved for the treatment of HER-2/neu over-expressing Metastatic Breast Cancer. The concentration of serum trastuzumab should be maintained above 10 μ g/mL during therapy, in order to effectively inhibit the growth of HER-2/neu over-expressing breast cancer cells. Accurate quantitation of serum trastuzumab concentrations is hence critical during therapy.

[0043] In an embodiment of the present disclosure, the full length HER-2/neu protein is composed of a cytoplasmic domain, a transmembrane domain and an extracellular domain (ECD) that is shed from the surface of cells. ECD is shed into the blood of normal individuals and is often elevated in metastatic breast cancer patients. Trastuzumab administered for treatment of these patients exists in two different forms in the matrix, the free drug and the drug complexed with the ECD. Elevated levels of circulating endogenous ECD result in increased levels of complexed trastuzumab. Standard immunoassays for quantitating trastuzumab measure only free trastuzumab leading to the underestimation of the circulating total trastuzumab levels. The assay disclosed herein eliminates the problem of ECD interference and thus underestimation by utilizing a unique acid-dissociation buffer which dissociates the trastuzumab-ECD complex in a single step.

[0044] The present disclosure is further elaborated with the help of following examples and associated figures. However, these examples should not be construed to limit the scope of the present disclosure.

EXAMPLES

EXAMPLE 1:

Preparation of Modified Citrate Buffer

[0045] In the present disclosure, 13.04 g of Citric Acid Monohydrate and 18.48g of Sodium Citrate Dihydrate are added to 950 mL of Milli Q water and dissolved. The pH of the solution is adjusted to 3.5 by adding Concentrated HCl (~ 4 mL). The volume of the solution is made up to 1000ml and the solution is filtered using 0.22 Micron filter unit to obtain the modified citrate buffer.

EXAMPLE 2:

Single step ELISA for accurate quantitation of trastuzumab in the presence of interfering HER2- ECD.

[0046] In the present disclosure, 100 μ L of a monoclonal antibody against trastuzumab Complementary Determining region (CDR) is coated on 96 well microtiter plates in PBS buffer at a final concentration of about 1 μ g/mL and incubated overnight at about 4°C. Calibrators and Quality Controls (QC) are prepared using trastuzumab in serum samples spiked with or without recombinant HER2-ECD. Samples are diluted about 50 fold (minimum required dilution, MRD) in modified citrate buffer containing 125 mM citrate-HCl and 0.1% Tween-20 (polysorbate 20) at pH between 2.8 to 3.5, and added to the blocked antibody coated microtiter plates. The plates are incubated at 22°C for one hour and washed five times with PBST. After washing, about 100 μ L of Peroxidase conjugated polyclonal detection antibody against trastuzumab CDR (1:2500 in PBST) is added to each well and incubated at 22°C for one hour. The plates are washed seven times with PBST and about 100 μ L of chromogenic substrate TMB is added to each well. After an incubation of 15 minutes, the reaction is stopped by the addition of 0.1N Sulphuric Acid and the plate is read at 450nm with a background subtraction at 630nm.

EXAMPLE 3:

Investigation of interference of HER2-ECD on serum calibration curve of Trastuzumab and studying the Effect of Modified Citrate Buffer on the Same

[0047] In the present disclosure, in order to investigate the interference of HER2-ECD on the serum calibration curve of trastuzumab, trastuzumab calibration curves are made in pooled healthy female serum containing either 0 ng/mL, 250ng/mL and 500ng/mL of spiked recombinant HER2-ECD. Samples are made to undergo 50 fold MRD in appropriate diluents - Figure 1. The Trastuzumab calibration curve was done using the ELISA assay as described above.

[0048] Figure (1A) shows that increasing the amount of spiked recombinant HER2-ECD resulted in a locational shift of the Trastuzumab Calibration curves in PBST diluent. Figure (1B) shows that the interference of spiked HER2-ECD is overcome when the serum samples underwent the MRD in modified citrate buffer (pH 3.5) diluent containing polysorbate 20. Dilution of samples containing spiked HER2-ECD and trastuzumab in modified citrate buffer containing polysorbate 20 resulted in the dissociation of immune complex formed between the analyte (trastuzumab) and target (HER2-ECD).

EXAMPLE 4:

Determining the Optimal Acidic pH of Buffer for Disassociating Trastuzumab-HER2-ECD complex

[0049] In the present disclosure, to determine the optimal acidic pH required to dissociate the trastuzumab-HER2-ECD complex, an acidic buffer, Citrate is tested at three different pH values as assay diluents for MRD. PBST diluent is also used as a control. Trastuzumab calibration curves are made in serum without spiked HER2-ECD and trastuzumab QC samples (500 ng/mL in neat serum) are prepared in 500 ng/mL recombinant HER2-ECD spiked serum. Calibration standards and QCs are made to undergo MRD in each of the four diluents. Analytical recovery of QC samples in each diluent is back calculated from the calibration curve prepared in the corresponding diluent. The lower the percentage relative error of recovery of trastuzumab QC, the better the buffer. As shown in table 1 above, a pH of 2.5 to 3.8 is required to obtain maximum recovery of the trastuzumab QC samples made in HER2-ECD spiked serum, with a pH of 3.5 showing the best result. The Trastuzumab calibration curve was done using the ELISA assay as described above.

TABLE 1- Optimum pH of Assay Diluent for Maximum Analytical Recovery of Trastuzumab LLOQ-QC (Lower Limit of Quantitation Quality Control)

S. No	Buffer	pH	% Analytical Recovery of trastuzumab QC (500ng/ml) in 500ng/ml HER2 ECD spiked serum	% Relative error of recovery of trastuzumab QC (500 ng/mL) in 500 ng/mL HER2 ECD spiked serum
1	PBST	7.4	32.4	67.6
2	Citrate buffer+ 0.1 % polysorbate-20	4.5	46.5	53.5
3	Modified Citrate + 0.1% polysorbate-20	4.0	74.78	25.22
4	Modified Citrate + 0.1% polysorbate-20	3.8	90.29	9.71
5	Modified Citrate + 0.1% polysorbate-20	3.5	104.67	4.67
6	Modified Citrate + 0.1% polysorbate-20	3.0	108.41	8.41
7	Modified Citrate + 0.1% polysorbate-20	2.5	88.77	11.23

EP 2 815 238 B1

(continued)

S. No	Buffer	pH	% Analytical Recovery of trastuzumab QC (500ng/ml) in 500ng/ml HER2 ECD spiked serum	% Relative error of recovery of trastuzumab QC (500 ng/mL) in 500 ng/mL HER2 ECD spiked serum
8	Modified Citrate + 0.1% polysorbate-20	2.0	ND	ND

EXAMPLE 5:

Comparison of Buffers for Obtaining Maximum Analytical Recovery of Trastuzumab QC

[0050] In the present disclosure, various acidic buffers at pH 3.5 are compared for MRD of samples to assess the most suitable combination of salt and weak acid in order to nullify the interference by HER2-ECD and maximize the analytical recovery of QC samples. Trastuzumab calibration curve, trastuzumab QCs (500 ng/mL and 1500 ng/mL in neat serum) are made in 1 µg/mL HER2-ECD spiked serum and made to undergo 50 fold MRD in four different diluents. The ELISA assay also includes a control trastuzumab calibration curve made in serum without spiked HER2-ECD and diluted 50 fold in PBST. Analytical recovery of QC samples in each diluent is back calculated from the control trastuzumab calibration curve made in serum without spiked HER2-ECD and diluted 50 fold in PBST.

[0051] Figure 2 shows that usage of Modified citrate buffer diluent results in minimal calibration curve shift induced by 1 µg/mL HER2 -ECD spike, as compared to the control un-spiked serum calibration curve diluted in PBST.

[0052] In the present disclosure, dilution of samples in other acidic buffers of same pH (conventional citrate buffer without the addition of HCl and glycine-HCl) failed to nullify the HER2-ECD induced calibration curve shift as shown in Table 2 below. Maximum recovery of the Trastuzumab QC samples (500ng/mL and 1500 ng/mL in neat serum) is observed when modified citrate buffer diluent is used.

TABLE 2- Comparison of Various Buffers for Obtaining Maximum Analytical Recovery of Trastuzumab QC

S. No.	Buffer	% Analytical Recovery of Trastuzumab QCs in 1 µg/ml HER2-ECD spiked serum	
		LLOQ QC (500ng/ml)	LOW QC (1500ng/ml)
1	PBST	10.28	40.97
2	Glycine HCl, pH 3.5	Out of Range	9.82
3	Citrate buffer+ 0.1% polysorbate 20 pH 3.5	72.56	75.16
4	Modified Citrate buffer+ 0.1% polysorbate 20 pH 3.5	86.06	86.82

[0053] This emphasizes the criticality of the modified citrate buffer composition on trastuzumab QC recovery in the presence of HER2-ECD.

EXAMPLE 6:

HER2-ECD Tolerance Determination of the Modified Immunoassay using Modified Citrate Buffer Diluent

[0054] In the present disclosure, to determine the maximum concentration of shed HER2-ECD that can be tolerated by the improved immunoassay using modified citrate buffer diluent containing polysorbate 20, trastuzumab QCs (375 ng/mL, 1125 ng/mL and 7500 ng/mL) are made in serum samples containing increasing amounts of recombinant HER2-ECD. The calibration standards and QC samples are made to undergo a 50 fold MRD in modified citrate buffer containing polysorbate 20. The analytical recoveries of QCs are back-calculated from a trastuzumab calibration curve made in un-spiked serum. It is observed that trastuzumab QCs can be reliably recovered from samples containing up to 2 µg/mL of HER2- ECD as shown in Table 3 below. The assay used in this experiment is ELISA.

TABLE 3- HER2-ECD Tolerance Determination of the Modified Immunoassay using Modified Citrate Buffer Diluent

S. No.	Spiked HER2-ECD level (ng/ml)	% Analytical Recovery of Trastuzumab QCs in recombinant HER2-ECD spiked samples		
		375ng/ml	1125ng/ml	7500ng/ml
1	50	106.11	100.37	93.11
2	500	97.11	98.27	95.04
3	1000	96.45	94.49	94.64
4	2000	98.01	92.09	95.30

EXAMPLE 7:**Analytical Recovery of Trastuzumab QC in Metastatic Breast Cancer Patient Samples with Varying Levels of shed HER2-ECD**

[0055] In the present disclosure, analytical Recovery of trastuzumab QCs made in Metastatic Breast Cancer (MBC) serum samples containing varying levels of endogenous shed HER2-ECD is shown in Table 4 below. A trastuzumab calibration curve made in un-spiked pooled healthy female sera and trastuzumab QCs made in MBC serum samples are diluted 50 fold using either PBST or modified citrate buffer containing polysorbate 20. The analytical recoveries of QCs are back-calculated from the corresponding diluent calibration curve. As seen in table 4 below, usage of modified Citrate buffer diluent results in optimal Trastuzumab QC recoveries compared to PBST diluent. The assay performed is ELISA. The concentrations of endogenous shed HER2-ECD are evaluated in MBC patient samples by a commercially available ELISA kit.

TABLE 4: Analytical recovery of Trastuzumab QC in metastatic breast cancer patient samples with varying levels of shed HER2-ECD. Usage of modified Citrate buffer diluent resulted in optimal Trastuzumab QC recoveries compared to PBST diluent.

Patient No	Shed HER2-ECD level (ng/mL)	% Analytical Recovery of Trastuzumab QCs in metastatic breast cancer patients at LLOQ QC (500ng/mL)	
		PBST diluent	Modified Citrate buffer diluent containing polysorbate 20
1	33.7	95.02	99.93
2	56.8	87.37	97.02
3	77.5	86.23	100
4	170	83.02	103.32
5	211.6	74.22	95.3
6	580	67.06	106.17
7	716.9	ND	104.48
8	833.6	ND	95.09
9	4761.8	BQL	87.29

***ND: Not done; BQL: Below Quantitation Limit.**

[0056] Therefore, from the above examples, it is clear that the process of the present disclosure utilizing the modified citrate buffer is capable of accurately determining and quantifying analytes, such as HER-2 antibody in a biological sample which contains a soluble target for the analyte such as HER-2 ECD. It is observed that Trastuzumab QCs can be reliably recovered from samples containing up to 4.7 $\mu\text{g/mL}$ of HER2- ECD as shown in Table 4 above.

OTHER EMBODIMENTS

[0057] While a number of embodiments and examples of this invention are described herein, it is apparent that these embodiments and examples may be altered to provide additional embodiments and examples which use the processes of this invention. Therefore it will be appreciated that the scope of this invention is to be defined by the appended claims, rather than by the specific embodiments that have been represented by way of example above.

Claims

1. A process for detection and optional quantification of the antibody trastuzumab in a biological sample, wherein the biological sample comprises an extracellular domain of HER-2 (HER-2-ECD) as an interfering substance and HER-2-ECD is capable of forming a complex with the antibody trastuzumab, the process comprising

- (a) providing a citrate buffer to the biological sample, the citrate buffer comprising sodium citrate, citric acid, polysorbate-20 and HCl, having a pH from 2.5 to 3.8, wherein the citrate buffer comprises citrate-HCl at a concentration of about 125 mM and polysorbate-20 at a concentration of about 0.1% w/v, and
 (b) carrying out an assay for said detection and optional quantification of the analyte within the biological sample.

2. The process of any of the preceding claims, wherein the biological sample is blood, plasma, serum or any combination thereof.

3. The process of any of the preceding claims, wherein the assay is an enzyme linked immunosorbent assay (ELISA).

4. The process of claim 3, wherein the enzyme linked immunosorbent assay is a sandwich enzyme linked immunosorbent assay and is carried out in presence of a capture antibody, a detection antibody, a reagent for reacting with the detection antibody, a microtitre plate, a coating buffer, a washing solution, a stopping solution, or any combination thereof, wherein optionally:

- (a) the capture antibody is an antibody against the analyte within the biological sample, or
 (b) the detection antibody is labelled with an enzyme selected from group comprising horseradish peroxidase and alkaline phosphatase and ruthenium, or
 (c) the reagent for reacting with the labelled antibody is chromogenic substrate selected from ABTS (2,2'-Azinobis[3-ethylbenzothiazoline-6-sulfonic acid]-diammonium salt, TMB (3,3',5,5'-tetramethylbenzidine) and electro chemiluminescent substrate, or
 (d) the coating buffer is phosphate buffered saline, or
 (e) the washing solution comprises phosphate buffered saline and Tween-20, or
 (f) the stopping solution is 0.1 N sulphuric acid.

5. The process of claims 3 or 4, wherein the ELISA comprises:

- a. coating a solid surface with the capture antibody against the analyte;
 b. providing the citrate buffer to the biological sample for obtaining diluted biological sample;
 c. adding the diluted sample to the antibody coated support of step (a);
 d. incubating and washing the support to remove any unbound analyte;
 e. adding a labelled detection antibody against the analyte and washing the support to remove any unbound detection antibody; and
 f. adding reagent for reacting with the labelled antibody for detecting and optionally quantifying the analyte in the sample.

6. A method of disassociating a complex formed by interaction of the antibody trastuzumab and at least one interfering substance as target for the antibody in a biological sample, said method comprising providing a citrate buffer to the biological sample for dissociating the complex, wherein the citrate buffer comprises sodium citrate, citric acid, polysorbate-20 and HCl, has a pH from 2.5 to 3.8, and comprises citrate-HCl at a concentration of about 125 mM and polysorbate-20 at a concentration of about 0.1% w/v, wherein the interfering substance is HER-2-ECD, wherein HER-2-ECD is capable of forming a complex with the antibody trastuzumab.

7. The use of a citrate buffer comprising sodium citrate, citric acid, polysorbate-20 and HCl, having a pH from 2.5 to 3.8, and comprising citrate-HCl at a concentration of about 125 mM and polysorbate-20 at a concentration of about 0.1% w/v, for detecting and optionally quantifying the presence of the antibody trastuzumab in a biological sample, wherein the biological sample comprises an extracellular domain of HER-2 (HER-2-ECD), and wherein HER-2-ECD is capable of forming a complex with the antibody trastuzumab.

Patentansprüche

1. Verfahren zum Nachweis und optional zur Quantifizierung des Antikörpers Trastuzumab in einer biologischen Probe, wobei die biologische Probe eine extrazelluläre Domäne von HER-2 (HER-2-ECD) als Störsubstanz aufweist und HER-2-ECD in der Lage ist, einen Komplex mit dem Antikörper Trastuzumab zu bilden, wobei das Verfahren aufweist
- a) einen Citratpuffer für die biologische Probe bereitstellen, wobei der Citratpuffer Natriumcitrat, Zitronensäure, Polysorbat-20 und HCl aufweist und einen pH-Wert von 2,5 bis 3,8 hat, wobei der Citratpuffer Citrat-HCl in einer Konzentration von etwa 125 mM und Polysorbat-20 in einer Konzentration von etwa 0,1% w/v aufweist und
- b) einen Test für den Nachweis und die optionale Quantifizierung des Analyten in der biologischen Probe durchführen.
2. Verfahren nach einem der vorangehenden Ansprüche, wobei die biologische Probe Blut, Plasma, Serum oder eine Kombination davon ist.
3. Verfahren nach einem der vorangehenden Ansprüche, wobei der Test ein enzymgebundener Immunosorbens-Assay (ELISA) ist.
4. Verfahren nach Anspruch 3, wobei der enzymgebundene Immunosorbens-Assay ein Sandwich-enzymgebundener Immunosorbens-Assay ist und in Gegenwart eines Capture-Antikörpers, eines Nachweis-Antikörpers, eines Reagens zur Reaktion mit dem Nachweis-Antikörper, einer Mikrotiterplatte, eines Beschichtungspuffers, einer Waschlösung, einer Stopplösung oder einer beliebigen Kombination davon durchgeführt wird, wobei optional:
- a) der Capture-Antikörper ein Antikörper gegen den Analyten in der biologischen Probe ist oder
- b) der Nachweisantikörper mit einem Enzym markiert ist, das aus der Gruppe ausgewählt ist, die Meerrettichperoxidase und alkalische Phosphatase und Ruthenium aufweist, oder
- c) das Reagenz zur Reaktion mit dem markierten Antikörper ein chromogenes Substrat ist, das aus ABTS (2,2'-Azinobis[3-ethylbenzothiazolin-6-sulfonsäure]-Diammoniumsalz, TMB (3,3',5,5'-Tetramethylbenzidin) und elektrochemilumineszierendem Substrat ausgewählt ist, oder
- d) der Beschichtungspuffer phosphatgepufferte Kochsalzlösung ist oder
- e) die Waschlösung Phosphat-gepufferte Kochsalzlösung und Tween-20 aufweist oder
- f) die Stopplösung 0,1 N Schwefelsäure ist.
5. Verfahren nach Anspruch 3 oder 4, wobei der ELISA aufweist:
- a. eine feste Oberfläche mit dem Capture-Antikörper gegen den Analyten beschichten,
- b. den Citratpuffer für die biologische Probe zum Erhalten einer verdünnten biologischen Probe bereitstellen,
- c. die verdünnte Probe zu dem mit Antikörper beschichteten Träger von Schritt a) zugeben,
- d. den Träger inkubieren und waschen, um nicht-gebundenen Analyten zu entfernen,
- e. einen markierten Nachweis-Antikörper gegen den Analyten zugeben und den Träger waschen, um nicht-gebundenen Nachweis-Antikörper zu entfernen, und
- f. ein Reagenz zur Reaktion mit dem markierten Antikörper zugeben, um den Analyten in der Probe nachzuweisen und gegebenenfalls zu quantifizieren.
6. Verfahren zum Dissoziieren eines Komplexes, der durch Wechselwirkung des Antikörpers Trastuzumab und mindestens einer störenden Substanz als Ziel für den Antikörper in einer biologischen Probe gebildet wird, wobei das Verfahren aufweist, einen Citratpuffer zur biologischen Probe zum Dissoziieren des Komplexes bereit zu stellen, wobei der Citratpuffer Natriumcitrat, Zitronensäure, Polysorbat-20 und HCl aufweist, einen pH-Wert von 2,5 bis 3,8 hat und Citrat-HCl in einer Konzentration von etwa 125 mM und Polysorbat-20 in einer Konzentration von etwa 0,1% w/v aufweist.

EP 2 815 238 B1

wobei die störende Substanz HER-2-ECD ist, wobei HER-2-ECD in der Lage ist, einen Komplex mit dem Antikörper Trastuzumab zu bilden.

7. Verwendung eines Citratpuffers, der Natriumcitrat, Zitronensäure, Polysorbat-20 und HCl aufweist, einen pH-Wert von 2,5 bis 3,8 hat und Citrat-HCl in einer Konzentration von etwa 125 mM und Polysorbat-20 in einer Konzentration von etwa 0,1% w/v aufweist, zum Nachweisen und optional Quantifizieren der Gegenwart des Antikörpers Trastuzumab in einer biologischen Probe, wobei die biologische Probe eine extrazelluläre Domäne von HER-2 (HER-2-ECD) aufweist und wobei HER-2-ECD in der Lage ist, einen Komplex mit dem Antikörper Trastuzumab zu bilden.

Revendications

1. Procédé pour la détection et la quantification facultative de l'anticorps trastuzumab dans un échantillon biologique, dans lequel l'échantillon biologique comprend un domaine extracellulaire de HER-2 (HER-2-ECD) en tant que substance interférente et HER-2-ECD est susceptible de former un complexe avec l'anticorps trastuzumab, le procédé comprenant

(a) la fourniture d'un tampon de citrate à l'échantillon biologique, le tampon de citrate comprenant du citrate de sodium, de l'acide citrique, du polysorbate 20 et de l'HCl, ayant un pH de 2,5 à 3,8, dans lequel le tampon de citrate comprend du citrate-HCl à une concentration d'environ 125 mM et du polysorbate 20 à une concentration d'environ 0,1 % M/V, et

(b) l'exécution d'un dosage pour lesdites détection et quantification facultative de la substance à analyser à l'intérieur de l'échantillon biologique.

2. Procédé selon l'une quelconque des revendications précédentes, dans lequel l'échantillon biologique est du sang, du plasma, du sérum ou n'importe quelle combinaison de ceux-ci.

3. Procédé selon l'une quelconque des revendications précédentes, dans lequel le dosage est un dosage immunoenzymatique utilisant un antigène absorbé (ELISA).

4. Procédé selon la revendication 3, dans lequel le dosage immunoenzymatique utilisant un antigène absorbé est un dosage immunoenzymatique utilisant un antigène absorbé en sandwich et est effectué en présence d'un anticorps de capture, un anticorps de détection, un réactif pour réagir avec l'anticorps de détection, une plaque de microtitre, un tampon de revêtement, une solution de lavage, une solution d'arrêt, ou n'importe quelle combinaison de ceux-ci : dans lequel de manière facultative :

(a) l'anticorps de capture est un anticorps contre la substance à analyser à l'intérieur de l'échantillon biologique, ou

(b) l'anticorps de détection est étiqueté avec une enzyme sélectionnée dans le groupe comprenant de la peroxydase de raifort et de la phosphatase alcaline et du ruthénium, ou

(c) le réactif pour réagir avec l'anticorps étiqueté est un substrat chromogène sélectionné parmi l'ABTS (acide 2,2'-Azinobis[3-éthylbenzothiazoline-6-sulfonique]-sel de diammonium, TMB (3,3',5,5'-tétraméthylbenzidine) et un substrat électro-chimioluminescent, ou

(d) le tampon de revêtement est une solution saline tamponnée au phosphate, ou

(e) la solution de lavage comprend une solution saline tamponnée au phosphate et du Tween-20, ou

(f) la solution d'arrêt est de l'acide sulfurique 0,1 N.

5. Procédé selon les revendications 3 ou 4, dans lequel l'ELISA comprend :

a. le revêtement d'une surface solide avec l'anticorps de capture contre la substance à analyser ;

b. la fourniture du tampon de citrate à l'échantillon biologique pour obtenir un échantillon biologique dilué ;

c. l'ajout de l'échantillon dilué au support revêtu d'anticorps de l'étape (a) ;

d. l'incubation et le lavage du support pour enlever toute substance à analyser non liée ;

e. l'ajout d'un anticorps de détection étiqueté contre la substance à analyser et le lavage du support pour enlever tout anticorps de détection non lié ; et

f. l'ajout d'un réactif pour réagir avec l'anticorps étiqueté pour la détection et facultativement la quantification de la substance à analyser dans l'échantillon.

EP 2 815 238 B1

6. Procédé de dissociation d'un complexe formé par interaction de l'anticorps trastuzumab et au moins une substance interférente en tant que cible pour l'anticorps dans un échantillon biologique, ledit procédé comprenant la fourniture d'un tampon de citrate à l'échantillon biologique pour dissocier le complexe, dans lequel le tampon de citrate comprend du citrate de sodium, de l'acide citrique, du polysorbate 20 et de l'HCl, a un pH de 2,5 à 3,8, et comprend du citrate-HCl à une concentration d'environ 125 mM et du polysorbate 20 à une concentration d'environ 0,1 % M/V, dans lequel la substance interférente est du HER-2-ECD, dans lequel l'HER-2-ECD est susceptible de former un complexe avec l'anticorps trastuzumab.
7. Utilisation d'un tampon de citrate comprenant du citrate de sodium, de l'acide citrique, du polysorbate 20 et de l'HCl, ayant un pH de 2,5 à 3,8 et comprenant du citrate-HCl à une concentration d'environ 125 mM et du polysorbate 20 à une concentration d'environ 0,1 % M/V, pour la détection et facultativement la quantification de la présence de l'anticorps trastuzumab dans un échantillon biologique, dans lequel l'échantillon biologique comprend un domaine extracellulaire de HER-2 (HER-2-ECD) et dans lequel l'HER-2-ECD est susceptible de former un complexe avec l'anticorps trastuzumab.

5

10

15

20

25

30

35

40

45

50

55




FIG. 1A




FIG. 1B




FIG. 2

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 4604348 A [0004]
- US 5641640 A [0004]
- US 2008182796 A [0004]
- US 20020090662 A [0004]

专利名称(译)	用于检测和任选量化分析物的方法		
公开(公告)号	EP2815238A2	公开(公告)日	2014-12-24
申请号	EP2013749422	申请日	2013-02-14
[标]申请(专利权)人(译)	拜康有限公司 迈兰公司		
申请(专利权)人(译)	Biocon公司 MYLAN GMBH		
当前申请(专利权)人(译)	Biocon公司 MYLAN GMBH		
[标]发明人	SENGUPTA NILANJAN UDIAVAR ANITA RAO THANGAMUTHU SENTHIL NAYAK VIVEK GOPAL PILLAI RANJIT RAVINDRAN VASHISHTA LAXMIKANT RAMASWAMY SHILPA GOVINDA HEGDE VISHIKA MELARKODE RAMAKRISHNAN		
发明人	SENGUPTA, NILANJAN UDIAVAR, ANITA, RAO THANGAMUTHU, SENTHIL NAYAK, VIVEK, GOPAL PILLAI, RANJIT, RAVINDRAN VASHISHTA, LAXMIKANT RAMASWAMY, SHILPA, GOVINDA HEGDE, VISHIKA MELARKODE, RAMAKRISHNAN		
IPC分类号	G01N33/53 G01N33/68		
CPC分类号	G01N33/5306 G01N33/6854 C07K16/28 G01N33/54393 G01N33/5748 G01N2333/71		
优先权	566CHE2012 2012-02-15 IN		
其他公开文献	EP2815238B1 EP2815238A4		
外部链接	Espacenet		

摘要(译)

本公开涉及在存在分析物的可溶性靶标的情况下检测和任选地定量样品中的分析物的方法。可溶性目标与分析物形成复合物，因此可能会干扰确定总分析物浓度。本发明的方法利用独特的修饰的柠檬酸盐缓冲液来稀释包含分析物和可溶性靶标的样品，这继而有助于解离分析物-可溶性靶标复合物，从而使本发明的方法能够检测并任选地量化测量。分析物准确。

