

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 April 2002 (25.04.2002)

PCT

(10) International Publication Number
WO 02/33119 A2

- (51) International Patent Classification⁷: C12Q 1/68, A61K 39/395, 38/17, C12N 15/11, A61K 31/70, C07K 16/28, G01N 33/53, C12N 15/85, 5/10, C07K 7/00, C12N 15/52, C07K 14/435
- (21) International Application Number: PCT/EP01/12018
- (22) International Filing Date: 17 October 2001 (17.10.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
MI2000A002240 17 October 2000 (17.10.2000) IT
- (71) Applicant and
(72) Inventor: PIETRANGELO, Antonello [IT/IT]; Via San Martino Mugnano, 3/1, I-41100 Modena (IT).
- (74) Agent: GERVASI, Gemma; Notarbartolo & Gervasi, Corso di Porta Vittoria, 9, I-20122 Milan (IT).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 02/33119 A2

(54) Title: MUTATIONS IN THE FERROPORTIN 1 GENE ASSOCIATED WITH HEREDITARY HAEMOCHROMATOSIS

(57) Abstract: The present invention relates to mutations in the gene coding for ferroportin 1 associated with hereditary haemochromatosis and methods for the diagnosis of hereditary haemochromatosis based on the identification of such mutations.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 April 2002 (25.04.2002)

PCT

(10) International Publication Number
WO 02/33119 A2

- (51) International Patent Classification⁷: C12Q 1/68, A61K 39/395, 38/17, C12N 15/11, A61K 31/70, C07K 16/28, G01N 33/53, C12N 15/85, 5/10, C07K 7/00, C12N 15/52, C07K 14/435
- (21) International Application Number: PCT/EP01/12018
- (22) International Filing Date: 17 October 2001 (17.10.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
MI2000A002240 17 October 2000 (17.10.2000) IT
- (71) Applicant and
(72) Inventor: PIETRANGELO, Antonello [IT/IT]; Via San Martino Mugnano, 3/1, I-41100 Modena (IT).
- (74) Agent: GERVASI, Gemma; Notarbartolo & Gervasi, Corso di Porta Vittoria, 9, I-20122 Milan (IT).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 02/33119 A2

(54) Title: MUTATIONS IN THE FERROPORTIN 1 GENE ASSOCIATED WITH HEREDITARY HAEMOCHROMATOSIS

(57) Abstract: The present invention relates to mutations in the gene coding for ferroportin 1 associated with hereditary haemochromatosis and methods for the diagnosis of hereditary haemochromatosis based on the identification of such mutations.

Mutations in the ferroportin 1 gene associated with hereditary haemochromatosis

FIELD OF THE INVENTION

The present invention concerns mutations in the gene coding for ferroportin 1
5 associated with hereditary haemochromatosis and the identification of such mutations
as a diagnostic method for hereditary haemochromatosis.

STATE OF THE ART

Haemochromatosis is a hereditary pathology characterised by an excessive
accumulation of iron in the organism, which over time leads to lesions of different
10 organs and tissues, particularly liver, myocardium, pancreas, kidney, spleen, gonads
and skin. Idiopathic haemochromatosis is the most widespread hereditary disease in
the Western population (incidence 1: 300) and is characterised by a recessive
transmission. Recently this type of haemochromatosis has been associated with
mutations of the HFE gene, located on the short arm of chromosome 6. In a study
15 carried out on patients suffering from this pathology in fact, it was observed that 83%
of the analyzed subjects had a single point mutation at this gene (C282Y) (Feder et
al, Nat Genet 1996, 13: 399-408).

However, more recent studies have shown that in the Mediterranean population only
64% of patients suffering from hereditary haemochromatosis are homozygotes for the
20 C282Y mutation. This suggests that, in the southern European population in
particular, other genes besides HFE may be responsible for idiopathic
haemochromatosis (Piperno et al, Gastroenterology 1998, 114: 996-1002 and Borot
et al, Immunogenetics 1997, 45: 320-324).

The identification of the genetic modifications responsible for hereditary haemochromatosis is of great diagnostic and therapeutic importance. Up to now, the diagnosis of haemochromatosis occurs too late and is based on the clinical symptomatology that develops as a result of often irreversible tissue injury. Besides,
5 diagnosis of this pathology is made difficult by the fact that its symptoms are often similar to those of other pathologies characterised by altered iron homeostasis.

The development of genetic screening methods for the early diagnosis, at presymptomatic stage, of hereditary haemochromatosis would allow a timely phlebotomy intervention before damage to organs and tissues occurs.

10 Furthermore, the identification of genetic alterations associated with hereditary haemochromatosis and the understanding of the role that they play in the development of the pathology are of extreme importance for the setting up of new and improved therapeutic strategies.

SUMMARY OF THE INVENTION

15 The inventor has previously identified and characterised a family suffering from a form of non HFE-dependent haemochromatosis with a autosomal dominant inheritance pattern, (Pietrangelo et al, New Eng J Med 1999, 341: 725-732).

The inventor has now surprisingly found that the locus of this pathology is on the long arm of chromosome 2 (2q32) and that subjects suffering from this type of
20 haemochromatosis have a mutation at a codon located in the hexone 3 of the gene coding for ferroportin 1, that is situated in the same chromosomal region, which is not observed in subjects who are not suffering from the pathology. This mutation leads to the substitution of an amino acid in the ferroportin 1 molecule.

Therefore the present invention refers to a nucleic acid coding for a mutated ferroportin 1 characterised by a mutation of the codon coding for the amino acid corresponding to position 77 of SEQ. ID. No. 2, a mutated ferroportin 1 protein coded by said nucleic acid and methods for the *in vitro* diagnosis of hereditary
5 haemochromatosis based on the identification of said nucleic acid or said protein.

DESCRIPTION OF THE FIGURES

Figure 1: Outline of the strategy for the identification of the mutation by means of enzymatic digestion with MbolI, described in Example 3.

In detail:

10 Figure 1a shows the genomic DNA sequence in the diagnostic method, in which the framed GCC sequence is the codon coding for the amino acid alanine in position 77 of the *wild type* ferroportin that is mutated to the GAC codon in individuals suffering from hereditary haemochromatosis. The double bar (//) denotes the separation between hexone 3 and a segment of the intron 3.

15 Figure 1b shows the primers used in the PCR reaction of the diagnostic method described in Example 3b and the target genomic DNA sequence in which N stands for the C or A nucleotide. The forward primer has a mismatched nucleotide, which is underlined (A instead of G).

Figure 1c shows the amplified sequence from control individuals, in the absence of
20 the mutation.

Figure 1d shows instead the amplified DNA sequence from individuals suffering from the pathology in which, in one of the alleles, the GCC codon is mutated to the GAC codon and leads to the appearance of a consensus site for the Mbo II enzyme.

Figure 2: Results of the diagnostic analysis on healthy family members or on family members suffering from haemochromatosis.

In detail:

Figure 2a shows the relationship between the analyzed individuals (pedigree). The subjects suffering from haemochromatosis are represented in black, while the healthy ones are in white. The circles represent the female subjects and the squares males.

Figure 2b shows the restriction profiles, following digestion with Mbo II, of the DNA amplified from each individual.

As shown in Figure 2c, in the case of healthy subjects, having only the *wild type* sequence, following digestion with Mbo II, the 131 base pair amplified DNA is not digested. Since all the subjects suffering from the pathology are heterozygotes for the mutation, the amplified DNA from these patients is digested in a band of 131 base pairs (normal allele) and two bands of 94 and 37 base pairs (the latter is not visible in Figure 2b).

15 DETAILED DESCRIPTION OF THE INVENTION

As will be shown in detail in the examples which follow, the authors of the present invention have identified that the mutation of a particular codon located in the hexone 3 of the ferroportin 1 gene is associated with a form of hereditary haemochromatosis that is not dependent on the HFE gene.

20 The identified mutation leads to the expression of a mutated ferroportin 1 in which the amino acid alanine, in a position corresponding to position 77 of SEQ ID No: 2 (sequence listing enclosed), is replaced.

The inventor has surprisingly found that the above mutation is a sufficient indication of the presence of hereditary haemochromatosis not linked to the HFE gene, and that therefore its identification is useful for the early diagnosis of this pathology. Besides, the authors of the present invention have found that hereditary haemochromatosis is
5 correlated with the functional impairment of ferroportin 1.

Therefore, according to a first aspect the present invention relates to a nucleic acid coding for a mutated ferroportin 1 characterised in that it comprises a mutation of the codon coding for the amino acid corresponding to position 77 of SEQ ID NO: 2.

The term "nucleic acid coding for a ferroportin 1" means a genomic DNA, cDNA,
10 DNA, for example obtained through PCR, or mRNA coding for the amino acid sequence of SEQ ID NO: 2 or for an amino acid sequence having at least 90% and preferably at least 95% homology with said amino acid sequence. When the nucleic acid is DNA this may be a single or double helix.

In addition, the invention comprises a nucleic acid with a sequence which is
15 complementary to that of the aforesaid nucleic acid. For instance, this sequence can be an antisense sequence used to stop, the expression of the gene or of the mRNA in the cells.

The mutation, according to the invention, leads to the substitution, in the molecule of *wild-type* ferroportin 1 (GenBank accession number: AF231121) of the amino acid
20 corresponding to position 77.

The term "*wild-type* ferroportin 1" refers to a ferroportin 1 which carries out its normal and physiological role, in particular that is without mutations which alter its functionality. Besides, the numerical position of the amino acid has the sole purpose

of identifying it and may vary due to the presence of variations in the amino acid sequence of the protein, for instance with changes in the species taken into consideration or because of the presence of mutations or deletions in the regions upstream of said amino acid.

5 The inventor has found that the substitution of the alanine in position 77 with a molecule of aspartic acid results in a structural modification in ferroportin 1 which may be shown, for example, by prediction programs of protein secondary structure, namely "PHDsec" (Rost et al, J Mol Biol 232: 584-599, 1993) and "JRED" (Cuff et al, Proteins: Structure, Function Genetics and 34: 508-519). In particular, said
10 substitution determines the passage of the protein region that goes from amino acid 58 to amino acid 81 (LLLTAVYGLVVAGS VLVLGXIIIGD, SEQ ID NO: 8) from the alpha-helix to beta-layer configuration.

The importance of the amino acid in position 77 in determining the secondary structure of ferroportin 1 is also confirmed by its high conservation among different
15 animal species.

The substitution of alanine, a small and uncharged amino acid, with a charged and larger molecule such as aspartic acid, results in steric and electrostatic interactions that destabilize the alpha helix hydrogen bonds. Similarly, substitutions of alanine with charged amino acids such as, for instance, arginine, lysine or glutamic acid or with
20 amino acids of greater sterical hinderance, for instance histidine, can lead to a similar distorsion of the molecule.

Furthermore, the ferroportin 1 region corresponding to SEQ ID NO 8 contains a post-transductional modification site of the protein, the myristilation site GAIIIGD, that is

altered in the mutated protein according to the invention. As is known, myristilation is important in helping the interactions of polypeptides with membrane phospholipids.

Therefore, according to a second aspect, the invention relates to a mutated ferroportin 1 protein characterised by the substitution of the amino acid in a position
5 corresponding to position 77 of SEQ ID NO: 2 and coded by the nucleic acid as above.

Said substitution consists in the substitution of the amino acid alanine in a position corresponding to position 77, with an amino acid having different steric and/or electrostatic properties from those of alanine. Preferably, said amino acid is selected
10 from the group including arginine, lysine, glutamic acid or aspartic acid, among which aspartic acid is preferred.

Therefore, according to a particularly preferred application, the mutation present in the nucleic acid of the invention consists in the substitution of the codon coding for the amino acid in a position corresponding to position 77 of SEQ ID NO: 2, preferably
15 GCC, with a codon chosen from the group including GAC and GAU, among which GAC is preferred.

In a further aspect of the invention relates to peptides with an amino acid sequence of at least 6 amino acids comprising the amino acid corresponding to position 77 of SEQ ID NO: 2 and the amino acids immediately downstream and/or upstream of this. The
20 length and sequence of such peptides are selected on the basis of criteria known to a person skilled in the art according to the desired application, for instance, to stimulate the production in host animals of antibodies specific for the mutation or to obtain peptides and/or antibodies that specifically interact with the mutated epitope.

A preferred embodiment of such peptides is the peptide corresponding to SEQ ID NO: 8, in which Xaa is preferably aspartic acid.

The present invention also relates to nucleotide fragments of the aforesaid nucleic acid comprising the mutated codon, and oligonucleotides with a sequence of at least
5 9 nucleotides and preferably at least 15 nucleotides of the aforesaid nucleic acid including said codon.

Said fragments and oligonucleotides can be of RNA or DNA and, in the latter case, single or double helix. Preferably, the oligonucleotides of the invention are single helix.

10 Furthermore the invention includes nucleotide fragments and oligonucleotides with sequences complementary to those of the aforesaid fragments or oligonucleotides.

By "nucleotide fragment", according to the present invention, is meant a nucleic acid with a sequence corresponding to a partial sequence of the nucleic acid of the invention having a length exceeding 100 base pairs.

15 By "oligonucleotide" according to the present invention, is meant a fragment of the nucleic acid of the invention of a maximum length of 100 base pairs.

The nucleotide fragments and oligonucleotides of the invention may be obtained, for instance, by digestion of the nucleic acid of the invention, through amplification by PCR or synthesized using techniques known in the art.

20 The oligonucleotides and the DNA fragments of the invention are used for different purposes such as, for instance, the production of chimeric proteins or antibodies, the determination of the mutation of the invention for diagnostic purposes or the inactivation of the mutated gene for therapeutic purposes. A person skilled in the art

is able to choose fragments and oligonucleotides of having sequence and length suitable for the desired uses. For instance, should said fragments or oligonucleotides be used for the determination of the mutation of invention with hybridization techniques, they must be of a length and sequence so as to be capable of hybridizing
5 in a specific way, under stringent conditions, at a nucleic acid sequence comprising the mutated codon.

According to a preferred embodiment, the fragments and oligonucleotides of the invention are labelled, for instance, with radioisotopes, enzymes, biotin-avidin or other molecules which allow them to be visualized through specific assays.

10 Moreover, the invention relates to the peptides coded by such fragments and oligonucleotides.

The nucleic acid, of the invention, a fragment thereof comprising the mutation or a nucleic acid comprising such fragment can be advantageously used for the production of a recombinant mutated ferroportin 1, a fragment of it or a chimeric
15 protein including this fragment, in order to, for instance, study the functional characteristics of the mutated protein, for example through competition studies, or produce antibodies. To this purpose, said nucleic acid or fragment is inserted into an expression vector which, in turn, is introduced into a procaryotic or eucaryotic cell using techniques well known in the art such as transfection, transformation, infection
20 or intranuclear injection.

Vectors suitable for this purpose include plasmids, vectors of viral origin and yeast or mammalian artificial chromosomes.

Accordingly, in a further aspect, the invention relates to a recombinant vector comprising a nucleic acid or a DNA fragment according to the invention as well as to eucaryotic or procaryotic cells comprising said vector.

The nucleic acid according to the invention can be used for the preparation of eucaryotic cells, tissues or non-human animals comprising a transgene coding for the mutated ferroportin 1 of the invention. The transgene can be permanently inserted into the genome of the cell, tissue or animal or be present in extrachromosomal form. Said cells, tissues or non-human animals are useful as models to study the function of the gene and of the protein comprising the mutation of the invention, as well as their role in the onset of hereditary haemochromatosis. This study is of particular importance for the development of new therapeutic approaches for the treatment of hereditary haemochromatosis.

In a further aspect, the invention refers to a method for the *in vitro* diagnosis of hereditary haemochromatosis in a mammal, preferably Homo Sapiens, comprising the following steps:

- a) isolation of genomic DNA or RNA from a biological sample obtained from said mammal;
- b) testing for the presence, in said genomic DNA or RNA, of the mutation according to the invention,

Wherein the presence of said mutation is an indication that said mammal is suffering from hereditary haemochromatosis.

Preferably said biological sample is a sample of plasma, saliva, urine, faeces, amniotic fluid or tissue.

Prior to testing, the RNA is preferably transformed into complementary DNA (cDNA) through a reverse transcription reaction.

The genomic DNA or the cDNA are analyzed directly or following *in vitro* amplification polymerase chain reaction (PCR) (Saiki et al, Science 239: 487-491, 1988) or other
5 techniques, for instance, ligase chain reaction (LCR) (Wu et al, genomics 4: 560-569, 1989) strand displacement amplification (SDA) (Walker et al, PNAS USA 89: 392-396) or self-sustained sequence replication (3SR) (Fahy et al, PCR methods Appl. 1: 25-33, 1992).

Preferably, the genomic DNA or the cDNA is amplified through PCR using a pair of
10 oligonucleotides (primers) suitable for the amplification of a segment of said DNA comprising the codon coding for the amino acid corresponding to position 77 of SEQ ID NO: 2.

For instance, pairs of primers that can be used to amplify the cDNA are those with the nucleotide sequence of SEQ ID No: 3 and SEQ ID No: 4 whereas primers suitable for
15 the amplification of the genomic DNA are those with the nucleotide sequence of SEQ ID No: 5 and SEQ ID No: 6.

Numerous techniques, well known in the art, can be used to determine the presence of the mutation according to the invention in the genomic DNA or the cDNA. Suitable techniques are, for instance, techniques based on the use of restriction enzymes
20 (Kan et al, Lancet: 910-912, 1978), hybridization techniques with allele-specific oligonucleotide probes (Wallace et al, Nucl Acids Res 6: 3543-3557, 1978) among which, for instance, hybridization with oligonucleotides immobilized on filters (Saiki et al, PNAS USA 86: 6230-6234, 1989) or micro-chips (Chee et al, Science 274: 610-

614, 1996) and oligonucleotide arrays (Maskos et al, Nucl Acids Res 21: 2269-2270, 1993), allele-specific PCR (Newton et al Nucl Acid Res 17: 2503-2516, 1989), mismatch repair detection (MRD) (Faham and Cox Genome Res: 474-482, 1995), Single-strand conformational polymorphism analysis (Ravnik-Glavac et al, Hum. Mol. Gen. 3: 801, 1994), gel electrophoresis in denaturant gradient (Guldborg et al., Nucl. Acids Res. 22: 880, 1994), Hot Cleavage (Cotton et al Proc.Natl. Acad Ski USA 85: 4397, 1988), DNase (Youil et al, PNAS USA 92: 87-91, 1995) and RNase protection assay (Winter et al. Proc. Natl. Acad. Ski. USA, 82: 7575, 1985; Meyers et al, Science 230: 1242, 1985), allele specific primer extension (Syvanen et al, genomics 8: 684-692, 1990 and Syvanen et al, Hum Mutat 13: 1-10, 1999), genetic bit analysis (GBA) (Nikiforov et al Nucl Acid Res 22: 4167-4175, 1994), primer-ligation assay (OLA) (Landergeren et al, Science 241: 1077, 1988), allele specific ligation chain reaction (LCR) (Barrany PNAS USA 88: 189-193, 1991), gap-LCR (Abravaya et al Nucl Acids Res 23: 675-682, 1995) and sequencing techniques. Particularly preferred techniques for the determination of the mutation of the invention are techniques based on the use of restriction enzymes, allele specific PCR, hybridization and sequencing techniques. Therefore, according to a first preferred application, the testing for the presence of the mutation according to the invention in the analyzed DNA occurs using techniques based on the use of restriction enzymes and comprises the following steps:

- a) amplification of the genomic DNA or cDNA with a pair of oligonucleotides suitable for the selective amplification of a segment of said DNA including the codon coding for the amino acid corresponding to position 77 of SEQ ID NO: 2 and the concurrent introduction into the amplified DNA of such a mutation which, in the presence of the

mutation of the invention, creates the consensus sequence for a restriction site otherwise not present;

b) incubation of the amplified DNA with an enzyme which is able to recognize said restriction site; and

5 c) analysis of the size of the digestion products;

wherein the occurring of digestion is a sign of the presence of the mutation of the invention in the genomic or complementary DNA.

The analysis of the size of the digestion products may be carried out, for instance, through gel electrophoresis, using a marker of molecular weights, followed by
10 visualization of the DNA bands using, for example, ethidium bromide.

In order to test, for example, for the presence of the substitution of the GCC codon, coding for alanine in position 77, with the GAC codon, coding for an aspartic acid molecule, oligonucleotides can be used with the nucleotide sequence of SEQ ID NO: 6 and SEQ ID NO: 7. As will be shown in the examples that follow, these
15 oligonucleotides give rise, in the presence of the aforesaid substitution, to a fragment of amplified DNA with the sequence shown in Figure 1d which contains the consensus site for the Mbo II enzyme, GAAGACATCATCGGT, not present in the fragment of DNA amplified from the *wild-type* DNA (Fig.1c). The subsequent incubation of the product of the amplification reaction with Mbo II results in the
20 fragment being digested only if the mutation was present in the original sequence.

According to a further preferred application, the determination of the mutation according to the invention is performed through hybridization techniques in which

fragments of the nucleic acid of the invention or oligonucleotides specific for the mutation according to the invention are used.

Said fragments or oligonucleotides are capable of hybridizing, in a specific way, at a sequence of the nucleic acid of the invention comprising the mutated codon also

5 when said sequence is present together with many other sequences.

A person skilled in the art is able to select each time the hybridization conditions and the length and sequence of the fragments or the oligonucleotides most suitable for the particular hybridization technique used and the kind of DNA that is being analyzed (genomic or complementary DNA, amplified or cloned in appropriate vectors).

10 According to a further preferred application, the diagnostic method envisages the use of allele-specific PCR, in which the genomic or complementary DNA undergoes a PCR reaction in which oligonucleotides are used which are able to selectively amplify a segment of said DNA including the mutated codon and not the corresponding segment including the *wild-type* codon.

15 In addition the present invention also relates to nucleic acid fragments and oligonucleotides according to the invention to be used in the aforesaid methods. In particular, it refers to oligonucleotides with the nucleotide sequence corresponding to SEQ ID NO: 3, 4, 5, 6 and 7.

Within the scope of the present invention are also included diagnostic kits for the
20 identification, in an individual, of the mutation according to the invention. According to a particularly preferred application, said diagnostic kits comprise oligonucleotides with a nucleotide sequence corresponding to SEQ ID NO: 6 and 7 and the Mbo II enzyme.

The present invention also relates to a method for the *in vitro* diagnosis of hereditary haemochromatosis in mammals including testing for the presence, in a biological sample from said mammal, of a mutated ferroportin 1 protein according to the invention, wherein the identification of said protein is an indication that the individual
5 is suffering from hereditary haemochromatosis.

Perferably said testing is performed through immunological assays in which monoclonal or polyclonal antibodies are used which are able to discriminate between a molecule of mutated ferroportin, according to the invention, and a molecule of *wild-type* ferroportin.

10 Therefore the present invention also refers to monoclonal or polyclonal antibodies which are able to specifically recognize a molecule of mutated ferroportin 1 according to the invention, or on epitope thereof comprising the mutation. Such antibodies are obtained using methods well known in the art such as those described by Harlow and Lane in *Antibodies, A Laboratory Manual*, Cold Spring Harbour Laboratory 1988.

15 The antibodies of the invention are particularly useful, as well as diagnostic reagents, in studying the characteristics of the protein or for therapeutic purposes. For instance, said antibodies can be used to determine the exact tissular or cellular localization of the mutated protein, to study its biochemical characteristics or to purify it by immunoaffinity.

20 Moreover, since the presence in an individual of a gene bearing the mutation of the invention and of the ferroportin 1 coded by it is correlated with the onset of hereditary haemochromatosis it is very important to have means to stop expression of the gene or to inactivate the protein.

Therefore, the invention also relates to oligonucleotides, for instance antisense oligonucleotides, suitable for inhibiting the expression of the gene coding for mutated ferroportin 1 of the invention, and antibodies and polypeptides able to specifically alter the functionality of the mutated ferroportin 1 of the invention.

- 5 In addition, the invention relates to pharmaceutical compositions comprising said oligonucleotides, antibodies or peptides mixed with pharmaceutically acceptable excipients.

EXAMPLES

EXAMPLE 1

- 10 Identification of the chromosome and the chromosomal locus associated with non HFE-dependent hereditary haemochromatosis

A DNA sample was extracted from the peripheral blood of the proband and family members suffering from the pathology using Quiagen Blood Extraction Kit (Quiagen).

- The DNA extracted was then used for a genome wide-search that was performed by
15 using the ABI PRISM Linkage mapping set (Perkin Elmer, United States). Fluorescent oligonucleotides were used for the PCR reactions, under the conditions suggested by the manufacturer. An aliquot of each PCR reaction was then sequenced in an ABI PRISM 377 DNA sequencer and the results obtained were analyzed using
GENESCAN software. The assignment of the allele was carried out using
20 Genotyper™ software. Statistical analysis was performed based on a dominant autosomal disease with complete penetrance. The gene-disease frequency was fixed at 0.012 and all the marker alleles were considered to be equally frequent. Table 1

contains the lod score, i.e. the maximum probability of association of a specific marker with the disease.

Table 1:

Marker	Z at θ_{max}								
	.00	.01	.05	.1	.2	.3	.4	Zmax (θ_{max})	
D2S2257	.20	.20	.19	.16	.11	.06	.02	.20	(.000)
D2S364	$-\infty$	3.46	3.82	3.67	3.00	2.08	.99	3.82	(.040)
D2S350	$-\infty$	2.82	3.16	2.99	2.31	1.44	.53	3.16	(.040)
D2S152	5.88	5.78	5.38	4.85	3.74	2.51	1.16	5.88	(.000)
D2S118	5.99	5.89	5.49	4.96	3.84	2.60	1.23	5.99	(.000)
D2S280	$-\infty$	2.35	2.77	2.70	2.22	1.53	.70	2.78	(.060)
D2S315	3.46	3.41	3.17	2.87	2.22	1.48	.66	3.46	(.000)
D2S117	$-\infty$	2.69	3.09	3.00	2.47	1.71	.76	3.09	(.050)

The higher the score, expressed logarithmically, the lower the probability that the association of the disease with the specific markers used is casual. For example, a lod score of 1.0 denotes 1 chance in 10 that the result is due to chance; a lod score of

2, one chance in 100 and so on. The fact that the markers D2S118 (5.99) and D2S152 (5.88) gave very high scores shows that the gene associated with the disease is situated in the region delimited by these markers. Ferroportin 1 is situated in this chromosomal region. No other chromosomal region gave similar results.

5 EXAMPLE 2

Identification of the mutation

Blood samples were taken from the proband, from 15 family members suffering from the pathology and from 25 family members not suffering from the pathology. Total RNA was isolated from macrophages obtained from each sample by extraction in
10 guanidine-isothiocyanate and the complementary DNA was then prepared according to a standard protocol (400 ng of total RNA, 1 µg of oligodT, 1mM of dNT, 20 U of reverse transcribed AMV in 20 µl of reaction buffer; Promega)

The complete ferroportin 1 sequence was then amplified from the cDNA through a PCR reaction using the following oligonucleotide pairs:

15 Forward primer 5'-GCTCAGGGCGTCCGCTAGGCT-3'

(SEQ.ID.NO 3)

reverse primer: 5'-GGCTTACACCCTCATGTTCT-3'

(SEQ ID NO: 4)

In detail, 10 ml of the product of the reverse transcription reaction were amplified in
20 50 µl (final volume) of 1X reaction buffer containing 200 µM dNTP, 1.5 mM MgCl₂, 0.25 µg of each of the above described oligonucleotide, 2.6 units of enzyme. A 30

cycle program was used for the amplification reaction, each of which was characterised by the following heat profile:

94°C for 1 minute,

58°C for 40 seconds,

5 75°C for 5 minutes.

From the amplified ferroportin 1 cDNA, four partially overlying DNA fragments were then obtained through a new PCR, using the following oligonucleotide pairs:

1° fragment:

Forward primer: 5'-GCTCAGGGCGTCCGCTAGGCT-3'

10 Reverse primer: 5'-CAGACACCGCAAAGTGCCACA-3'

2° fragment:

Forward primer: 5'-ACCTCGCTGGTGGTACAG-3'

Reverse primer: 5'-CCGCAAGCAAAGAGCTTG-3'

3° fragment:

15 Forward primer: 5'-GGTGCTATCTCCAGTTCCTT-3'

Reverse primer: 5'-AGACGTA CTCCACGCACA-3'

4° fragment:

Forward primer: 5'-TCAGTCTCCTTTGTGGCA-3'

Reverse primer: 5'-GGCTTACACCCTCATGTTCT-3'

20 The four fragments obtained from the amplification were then electrophoretically separated on agarose gel, purified using the Jet Sorb kit (Genenco) and sequenced directly using the Rhodamine Sequence kit (Perkin Elmer, United States). Sequencing revealed the presence, in subjects suffering from the pathology, of the

substitution of a C with an A in position 230 of SEQ.ID No.1 (nucleotide in position 534 of the sequence having GenBank accession number: AF231121), that was not found in any of the control subjects. This substitution is located in hexone 3 of the ferroportin 1 gene and results in the substitution of alanine with aspartic acid at
5 position 77 of ferroportin 1.

EXAMPLE 3

Diagnostic method

a) genomic DNA of the proband, of 15 family members suffering from the pathology and 125 control individuals, including 100 healthy volunteers and 25 family members
10 not suffering from the pathology, was extracted from leukocytes obtained from blood samples of the subjects to be analyzed using a blood DNA extraction kit (Quiagen).

The DNA obtained was then amplified by PCR using a pair of oligonucleotides complementary to the intronic regions flanking hexone 3 and having the following sequence:

15 Forward primer: 5'-CCTTTTGATAAGGAAGCAACTTCC-3'

(SEQ ID NO: 5)

Reverse primer: 5'-CAGAGGTAGCTCAGGCATTGGTCC-3'

(SEQ ID NO: 6)

In detail, 200 ng of genomic DNA were amplified in 50 µl of 1X reaction buffer
20 containing 200 µM dNTPs, 1.5 mM MgCl₂, 20 pmoles of each oligonucleotide and 2.6 U of enzyme.

A 30 cycle program was used for the amplification reaction, each of which was characterised by the following heat profile:

94°C for 1 minute,

60°C for 1 minute,

5 72°C for 45 seconds.

The DNA obtained was then purified using the PCR Wizard kit (Promega) and sequenced in an automatic ABI Prism 377 sequencer (Perkin Elmer, United States), with the same pair of oligonucleotides used for the PCR reaction.

The substitution of a C with an A in hexone 3 was found in the subjects suffering
10 from the pathology, but not in the control subjects.

b) blood samples were obtained from the proband, from 15 family members suffering from the pathology and from the 125 control subjects described in example 2 and the genomic DNA was extracted using the blood DNA extraction kit (Biorad).

Since the mutation observed does not give rise to the appearance or disappearance
15 of any restriction site, the portion of hexone 3 containing the mutation was amplified by PCR using a pair of oligonucleotides with the following sequences:

Forward primer: 5'-GTGGCAGGGTCTGTTCTGGTCCTGGAAG-3'

(SEQ.ID NO.7)

Reverse primer: 5'-CAGAGGTAGCTCAGGCATTGGTCC-3'

20 (SEQ.ID NO.6)

As shown in Figure 1b, the forward primer has a mismatched nucleotide (A instead of G, underlined). Amplification with the aforesaid oligonucleotides gives rise, in the presence of the mutation, to an amplified DNA fragment, having the sequence shown

in Figure 1d, that contains the consensus site for the Mbo II enzyme, GAAGACATCATCGGT. On the contrary, in the absence of the mutation, a DNA fragment with the sequence shown in Figure 1d is obtained and which does not contain such a restriction site (Fig.1c).

5 Therefore, the subsequent incubation of the product of the amplification reaction with the Mbo II enzyme results in the digestion of the fragment only if the original sequence contained the mutation.

In detail, 10 µl of the extracted genomic DNA were amplified in a final 50 µl of 1x reaction buffer containing 200 µM dNTPs, 1.5 mM MgCl₂, 20 pmoles of each
10 oligonucleotide and 2.6 units of enzyme. A 30 cycle program was used for the amplification reaction, each cycle being characterised by the following heat profile:

94°C for 1 minute,

58°C for 1 minute,

72°C for 45 seconds.

15 10 µl of the product obtained from the amplification were then digested with the Mbo II enzyme (Geneco) in 1X reaction buffer for 3 hours at 37°C.

The fragments obtained from the digestion were then separated on a 12% polyacrylamide gel. The samples obtained from patients suffering from the pathology resulted in 3 bands of 131, 94 and 37 base pairs. On the contrary, all the control
20 samples gave had just one band of 131 base pairs. Figure 2 shows the results obtained from the proband, from 4 family members suffering from the pathology and from 3 healthy family members. As can be seen from the figure, digestion of the

fragment of 131 base pairs into two fragments of 94 and 37 base pairs was only observed in subjects suffering from the pathology.

Also comprised in the present invention is any therapeutic operation involving the substitution of the mutated gene with the *wild type* gene. Therefore, the mutated
5 ferroportin 1 molecule is a therapeutic target for all genic therapy operations aimed at the substitution of the mutated gene.

CLAIMS

1. A nucleic acid coding for a ferroportin 1 characterised in that it comprises a mutation of the codon coding for the amino acid corresponding to position 77 of SEQ. ID. NO. 2.
- 5 2. A nucleic acid as claimed in claim 1 characterised in that it is cDNA.
3. A nucleic acid as claimed in claim 1 characterised in that it is mRNA.
4. A nucleic acid as claimed in claim 1 characterised in that it is genomic DNA.
5. A nucleic acid as claimed in claim 1 characterised in that said amino acid is alanine.
- 10 6. A nucleic acid as claimed in claim 1 characterised in that said mutation leads to the substitution of said amino acid with an amino acid having different steric and/or electrostatic characteristics.
7. A nucleic acid as claimed in claim 6 characterised in that said amino acid having different steric and/or electrostatic characteristics is selected from the group
- 15 comprising arginine, lysine, glutamic acid or aspartic acid.
8. A nucleic acid as claimed in claim 7 characterised in that said amino acid is aspartic acid.
9. A nucleic acid as claimed in claim 8 characterised in that said mutation consists in the substitution of the GCC codon with the GAC'or GAU codon.
- 20 10. A nucleic acid as claimed in claim 9 characterised in that said mutation consists in the substitution of the GCC codon with the GAC codon.
11. A nucleic acid with a sequence which is complementary to that of the nucleic acid as claimed in claim 1.

12. A mutated ferroportin 1 coded by a nucleic acid as claimed in claim 1.
13. A peptide with a sequence of at least 6 aminoacids of a ferroportin as claimed in claim 12 comprising the amino acid corresponding to position 77 of SEQ ID NO: 2.
14. A fragment of the nucleic acid as in claim 1 comprising the codon coding for the amino acid corresponding to position 77 of SEQ. ID. NO. 2.
- 5 15. A fragment as claimed in claim 14 characterised in that it is labelled.
16. An oligonucleotide with a sequence of at least 9 nucleotides of the nucleic acid as claimed in claim 1 and comprising the codon coding for the amino acid corresponding to position 77 of SEQ. ID. NO. 2
- 10 17. An oligonucleotide as claimed in claim 16 characterised in that said sequence is of at least 15 nucleotides.
18. An oligonucleotide as claimed in claim 16 characterised in that it is labelled.
19. A fragment with a sequence which is complementary to that of the fragment as claimed in claim 14.
- 15 20. An oligonucleotide with a sequence complementary to that of the oligonucleotide as claimed in claim 16.
21. A peptide coded by a fragment as claimed in claim 14 or by an oligonucleotide as claimed in claim 16.
22. A recombinant vector comprising the nucleotide sequence of the nucleic acid as claimed in claim 1 or of the fragment as claimed in claim 14.
- 20 23. A cell transfected or transformed with the recombinant vector as claimed in claim 22.

24. A eucaryotic cell, tissue or non-human animal comprising a transgene coding for a protein as claimed in claim 12.

25. A method for the in vitro diagnosis of hereditary haemochromatosis in a mammal comprising the following steps:

- 5 a) isolation of genomic DNA or RNA from a biological sample obtained from said mammal;
- b) testing for the presence of the mutation as claimed in claim 1 in said genomic DNA or RNA.

Wherein the presence of said mutation is an indication that said mammal is
10 suffering from hereditary haemochromatosis.

26. A method as claimed in claim 25 characterised in that said mammal is a Homo sapiens.

27. A method as claimed in claim 25 characterised in that said biological sample is a sample of blood, plasma, saliva, urine, faeces, amniotic fluid or tissue.

15 28. A method as claimed in claim 25 characterised in that the isolated genomic DNA is amplified before said testing.

29. A method as claimed in claim 28 characterised in that the genomic DNA is amplified by PCR using a pair of oligonucleotides suitable for the amplification of a segment of said DNA comprising the codon coding for the amino acid corresponding
20 to position 77 of SEQ ID NO: 2.

30. A method as claimed in claim 29 characterised in that said oligonucleotides have the nucleotide sequence of SEQ.ID NO: 5 and SEQ.ID NO: 6.

31. A method as claimed in claim 25 characterised in that the RNA is transformed into cDNA before said testing.

32. A method as claimed in claim 31 also characterised in that the cDNA is amplified before said testing.

5 33. A method as claimed in claim 32 characterised in that the cDNA is amplified by PCR using a pair of oligonucleotides suitable for the amplification of a segment of said DNA comprising the codon coding for the amino acid corresponding to position 77 of SEQ ID NO: 2.

34. A method as claimed in claim 33 characterised in that said oligonucleotides have
10 the nucleotide sequence of SEQ.ID NO: 3 and SEQ.ID NO: 4.

35. A method as claimed in claim 25 characterised in that said testing is carried out using a technique selected from the group including techniques based on the use of restriction enzymes, hybridization techniques with allele-specific oligonucleotide probes, allele-specific PCR, mismatch repair detection, single-strand conformational
15 polymorphism analysis, gel electrophoresis in denaturant gradient, Hot Cleavage, DNase and RNase protection assay, allele specific primer extension, genetic bit analysis oligonucleotide-ligation assay, allele specific ligation chain reaction and sequencing techniques.

36. A method as claimed in claim 35 characterised in that said testing is carried out
20 using techniques based on the use of restriction enzymes, allele specific PCR, hybridization or sequencing techniques.

37. A method as claimed in claim 36 characterised in that said testing comprises the following steps:

- a) amplification of the genomic DNA or of cDNA with oligonucleotides suitable for the selective amplification of a segment of said DNA including the codon coding for the amino acid corresponding to position 77 of SEQ ID NO: 2 and the concurrent introduction into the amplified DNA of such a mutation that, together with the mutation
- 5 of the invention, creates the consensus sequence for a restriction site otherwise not present;
- b) incubation of the amplified DNA with an enzyme which is able to recognize said restriction site; and
- c) analysis of the size of the digestion products;

10 wherein the occurring of digestion is a sign of the presence of the mutation as claimed in claim 1 in the genomic DNA or cDNA.

38. A method as claimed in claim 37 characterised in that said oligonucleotides have the nucleotide sequence of SEQ ID NO: 6 and SEQ ID NO: 7 and that said enzyme is Mbo II.

15 39. An oligonucleotide with the nucleotide sequence of SEQ ID NO: 3, 4, 5, 6 or 7.

40. A diagnostic kit for the determination of the mutation as claimed in claim 1 in an individual comprising oligonucleotides with the nucleotide sequence of SEQ ID NO: 6 and SEQ ID NO: 7 and the Mbo II enzyme.

41. A method for the *in vitro* diagnosis of hereditary haemochromatosis in a mammal

20 including the testing for the presence of a ferroportin 1 protein as claimed in claim 12 in a biological sample obtained from said mammal, in which the presence of said protein is an indication that said mammal is suffering from hereditary haemochromatosis.

42. A method as claimed in claim 41 in which said identification is carried out using antibodies which are able to specifically recognize said mutated ferroportin 1.

43. Monoclonal or polyclonal antibodies which are able to specifically recognize a mutated ferroportin 1 protein as claimed in claim 12.

5 44. Antibodies for the specific inactivation of a mutated ferroportin 1 protein as claimed in claim 12.

45. A pharmaceutical composition comprising antibodies as claimed in claim 44.

46. Polypeptides for the specific inactivation of a mutated ferroportin 1 as claimed in claim 12.

10 47. A pharmaceutical composition including polypeptides as claimed in claim 46.

48. Oligonucleotides able to stop the expression of the gene coding for a mutated ferroportin 1 as claimed in claim 12.

49. A pharmaceutical composition including oligonucleotides as claimed in claim 48.

FIGURE 1





FIGURE 2



SEQUENCE LISTING

<170> PatentIn Ver. 2.1

<210> 1

<211> 1716

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(1716)

<400> 1

```

atg acc agg gcg gga gat cac aac cgc cag aga gga tgc tgt gga tcc 48
Met Thr Arg Ala Gly Asp His Asn Arg Gln Arg Gly Cys Cys Gly Ser
 1                    5                10                15

ttg gcc gac tac ctg acc tct gca aaa ttc ctt ctc tac ctt ggt cat 96
Leu Ala Asp Tyr Leu Thr Ser Ala Lys Phe Leu Leu Tyr Leu Gly His
                20                25                30

tct ctc tct act tgg gga gat cgg atg tgg cac ttt gcg gtg tct gtg 144
Ser Leu Ser Thr Trp Gly Asp Arg Met Trp His Phe Ala Val Ser Val
                35                40                45

ttt ctg gta gag ctc tat gga aac agc ctc ctt ttg aca gca gtc tac 192
Phe Leu Val Glu Leu Tyr Gly Asn Ser Leu Leu Leu Thr Ala Val Tyr
                50                55                60

ggg ctg gtg gtg gca ggg tct gtt ctg gtc ctg gga nnn atc atc ggt 240
Gly Leu Val Val Ala Gly Ser Val Leu Val Leu Gly Xaa Ile Ile Gly
 65                70                75                80

gac tgg gtg gac aag aat gct aga ctt aaa gtg gcc cag acc tcg ctg 288
Asp Trp Val Asp Lys Asn Ala Arg Leu Lys Val Ala Gln Thr Ser Leu
                85                90                95

gtg gta cag aat gtt tca gtc atc ctg tgt gga atc atc ctg atg atg 336
Val Val Gln Asn Val Ser Val Ile Leu Cys Gly Ile Ile Leu Met Met
                100                105                110

gtt ttc tta cat aaa cat gag ctt ctg acc atg tac cat gga tgg gtt 384
Val Phe Leu His Lys His Glu Leu Leu Thr Met Tyr His Gly Trp Val
                115                120                125

ctc act tcc tgc tat atc ctg atc atc act att gca aat att gca aat 432
Leu Thr Ser Cys Tyr Ile Leu Ile Ile Thr Ile Ala Asn Ile Ala Asn
                130                135                140

ttg gcc agt act gct act gca atc aca atc caa agg gat tgg att gtt 480
Leu Ala Ser Thr Ala Thr Ala Ile Thr Ile Gln Arg Asp Trp Ile Val
145                150                155                160

gtt gtt gca gga gaa gac aga agc aaa cta gca aat atg aat gcc aca 528
Val Val Ala Gly Glu Asp Arg Ser Lys Leu Ala Asn Met Asn Ala Thr
                165                170                175

ata cga agg att gac cag tta acc aac atc tta gcc ccc atg gct gtt 576
Ile Arg Arg Ile Asp Gln Leu Thr Asn Ile Leu Ala Pro Met Ala Val
                180                185                190

ggc cag att atg aca ttt ggc tcc cca gtc atc ggc tgt ggc ttt att 624

```

Gly	Gln	Ile	Met	Thr	Phe	Gly	Ser	Pro	Val	Ile	Gly	Cys	Gly	Phe	Ile		
		195					200					205					
tcg	gga	tgg	aac	ttg	gta	tcc	atg	tgc	gtg	gag	tac	gtc	ctg	ctc	tgg	672	
Ser	Gly	Trp	Asn	Leu	Val	Ser	Met	Cys	Val	Glu	Tyr	Val	Leu	Leu	Trp		
	210					215					220						
aag	gtt	tac	cag	aaa	acc	cca	gct	cta	gct	gtg	aaa	gct	ggt	ctt	aaa	720	
Lys	Val	Tyr	Gln	Lys	Thr	Pro	Ala	Leu	Ala	Val	Lys	Ala	Gly	Leu	Lys		
	225				230					235					240		
gaa	gag	gaa	act	gaa	ttg	aaa	cag	ctg	aat	tta	cac	aaa	gat	act	gag	768	
Glu	Glu	Glu	Thr	Glu	Leu	Lys	Gln	Leu	Asn	Leu	His	Lys	Asp	Thr	Glu		
			245						250					255			
cca	aaa	ccc	ctg	gag	gga	act	cat	cta	atg	ggt	gtg	aaa	gac	tct	aac	816	
Pro	Lys	Pro	Leu	Glu	Gly	Thr	His	Leu	Met	Gly	Val	Lys	Asp	Ser	Asn		
			260					265					270				
atc	cat	gag	ctt	gaa	cat	gag	caa	gag	cct	act	tgt	gcc	tcc	cag	atg	864	
Ile	His	Glu	Leu	Glu	His	Glu	Gln	Glu	Pro	Thr	Cys	Ala	Ser	Gln	Met		
		275					280					285					
gct	gag	ccc	ttc	cg	acc	ttc	cga	gat	gga	tgg	gtc	tcc	tac	tac	aac	912	
Ala	Glu	Pro	Phe	Arg	Thr	Phe	Arg	Asp	Gly	Trp	Val	Ser	Tyr	Tyr	Asn		
	290					295					300						
cag	cct	gtg	ttt	ctg	gct	ggc	atg	ggt	ctt	gct	ttc	ctt	tat	atg	act	960	
Gln	Pro	Val	Phe	Leu	Ala	Gly	Met	Gly	Leu	Ala	Phe	Leu	Tyr	Met	Thr		
	305				310				315						320		
gtc	ctg	ggc	ttt	gac	tgc	atc	acc	aca	ggg	tac	gcc	tac	act	cag	gga	1008	
Val	Leu	Gly	Phe	Asp	Cys	Ile	Thr	Thr	Gly	Tyr	Ala	Tyr	Thr	Gln	Gly		
			325						330					335			
ctg	agt	ggt	tcc	atc	ctc	agt	att	ttg	atg	gga	gca	tca	gct	ata	act	1056	
Leu	Ser	Gly	Ser	Ile	Leu	Ser	Ile	Leu	Met	Gly	Ala	Ser	Ala	Ile	Thr		
			340					345					350				
gga	ata	atg	gga	act	gta	gct	ttt	act	tgg	cta	cgt	cga	aaa	tgt	ggt	1104	
Gly	Ile	Met	Gly	Thr	Val	Ala	Phe	Thr	Trp	Leu	Arg	Arg	Lys	Cys	Gly		
		355					360					365					
ttg	gtt	cgg	aca	ggt	ctg	atc	tca	gga	ttg	gca	cag	ctt	tcc	tgt	ttg	1152	
Leu	Val	Arg	Thr	Gly	Leu	Ile	Ser	Gly	Leu	Ala	Gln	Leu	Ser	Cys	Leu		
	370					375					380						
atc	ttg	tgt	gtg	atc	tct	gta	ttc	atg	cct	gga	agc	ccc	ctg	gac	ttg	1200	
Ile	Leu	Cys	Val	Ile	Ser	Val	Phe	Met	Pro	Gly	Ser	Pro	Leu	Asp	Leu		
	385				390					395					400		
tcc	gtt	tct	cct	ttt	gaa	gat	atc	cga	tca	agg	ttc	att	caa	gga	gag	1248	
Ser	Val	Ser	Pro	Phe	Glu	Asp	Ile	Arg	Ser	Arg	Phe	Ile	Gln	Gly	Glu		
				405					410					415			
tca	att	aca	cct	acc	aag	ata	cct	gaa	att	aca	act	gaa	ata	tac	atg	1296	
Ser	Ile	Thr	Pro	Thr	Lys	Ile	Pro	Glu	Ile	Thr	Thr	Glu	Ile	Tyr	Met		
			420					425					430				
tct	aat	ggg	tct	aat	tct	gct	aat	att	gtc	ccg	gag	aca	agt	cct	gaa	1344	
Ser	Asn	Gly	Ser	Asn	Ser	Ala	Asn	Ile	Val	Pro	Glu	Thr	Ser	Pro	Glu		
		435					440						445				

tct gtg ccc ata atc tct gtc agt ctg ctg ttt gca ggc gtc att gct 1392
 Ser Val Pro Ile Ile Ser Val Ser Leu Leu Phe Ala Gly Val Ile Ala
 450 455 460

gct aga atc ggt ctt tgg tcc ttt gat tta act gtg aca cag ttg ctg 1440
 Ala Arg Ile Gly Leu Trp Ser Phe Asp Leu Thr Val Thr Gln Leu Leu
 465 470 475 480

caa gaa aat gta att gaa tct gaa aga ggc att ata aat ggt gta cag 1488
 Gln Glu Asn Val Ile Glu Ser Glu Arg Gly Ile Ile Asn Gly Val Gln
 485 490 495

aac tcc atg aac tat ctt ctt gat ctt ctg cat ttc atc atg gtc atc 1536
 Asn Ser Met Asn Tyr Leu Leu Asp Leu Leu His Phe Ile Met Val Ile
 500 505 510

ctg gct cca aat cct gaa gct ttt ggc ttg ctc gta ttg att tca gtc 1584
 Leu Ala Pro Asn Pro Glu Ala Phe Gly Leu Leu Val Leu Ile Ser Val
 515 520 525

tcc ttt gtg gca atg ggc cac att atg tat ttc cga ttt gcc caa aat 1632
 Ser Phe Val Ala Met Gly His Ile Met Tyr Phe Arg Phe Ala Gln Asn
 530 535 540

act ctg gga aac aag ctc ttt gct tgc ggt cct gat gca aaa gaa gtt 1680
 Thr Leu Gly Asn Lys Leu Phe Ala Cys Gly Pro Asp Ala Lys Glu Val
 545 550 555 560

agg aag gaa aat caa gca aat aca tct gtt gtt tga 1716
 Arg Lys Glu Asn Gln Ala Asn Thr Ser Val Val
 565 570

<210> 2
 <211> 571
 <212> PRT
 <213> Homo sapiens

<400> 2
 Met Thr Arg Ala Gly Asp His Asn Arg Gln Arg Gly Cys Cys Gly Ser
 1 5 10 15
 Leu Ala Asp Tyr Leu Thr Ser Ala Lys Phe Leu Leu Tyr Leu Gly His
 20 25 30
 Ser Leu Ser Thr Trp Gly Asp Arg Met Trp His Phe Ala Val Ser Val
 35 40 45
 Phe Leu Val Glu Leu Tyr Gly Asn Ser Leu Leu Leu Thr Ala Val Tyr
 50 55 60
 Gly Leu Val Val Ala Gly Ser Val Leu Val Leu Gly Xaa Ile Ile Gly
 65 70 75 80
 Asp Trp Val Asp Lys Asn Ala Arg Leu Lys Val Ala Gln Thr Ser Leu
 85 90 95
 Val Val Gln Asn Val Ser Val Ile Leu Cys Gly Ile Ile Leu Met Met
 100 105 110
 Val Phe Leu His Lys His Glu Leu Leu Thr Met Tyr His Gly Trp Val
 115 120 125
 Leu Thr Ser Cys Tyr Ile Leu Ile Ile Thr Ile Ala Asn Ile Ala Asn
 130 135 140
 Leu Ala Ser Thr Ala Thr Ala Ile Thr Ile Gln Arg Asp Trp Ile Val
 145 150 155 160
 Val Val Ala Gly Glu Asp Arg Ser Lys Leu Ala Asn Met Asn Ala Thr
 165 170 175
 Ile Arg Arg Ile Asp Gln Leu Thr Asn Ile Leu Ala Pro Met Ala Val
 180 185 190

Gly Gln Ile Met Thr Phe Gly Ser Pro Val Ile Gly Cys Gly Phe Ile
 195 200 205
 Ser Gly Trp Asn Leu Val Ser Met Cys Val Glu Tyr Val Leu Leu Trp
 210 215 220
 Lys Val Tyr Gln Lys Thr Pro Ala Leu Ala Val Lys Ala Gly Leu Lys
 225 230 235 240
 Glu Glu Glu Thr Glu Leu Lys Gln Leu Asn Leu His Lys Asp Thr Glu
 245 250 255
 Pro Lys Pro Leu Glu Gly Thr His Leu Met Gly Val Lys Asp Ser Asn
 260 265 270
 Ile His Glu Leu Glu His Glu Gln Glu Pro Thr Cys Ala Ser Gln Met
 275 280 285
 Ala Glu Pro Phe Arg Thr Phe Arg Asp Gly Trp Val Ser Tyr Tyr Asn
 290 295 300
 Gln Pro Val Phe Leu Ala Gly Met Gly Leu Ala Phe Leu Tyr Met Thr
 305 310 315 320
 Val Leu Gly Phe Asp Cys Ile Thr Thr Gly Tyr Ala Tyr Thr Gln Gly
 325 330 335
 Leu Ser Gly Ser Ile Leu Ser Ile Leu Met Gly Ala Ser Ala Ile Thr
 340 345 350
 Gly Ile Met Gly Thr Val Ala Phe Thr Trp Leu Arg Arg Lys Cys Gly
 355 360 365
 Leu Val Arg Thr Gly Leu Ile Ser Gly Leu Ala Gln Leu Ser Cys Leu
 370 375 380
 Ile Leu Cys Val Ile Ser Val Phe Met Pro Gly Ser Pro Leu Asp Leu
 385 390 395 400
 Ser Val Ser Pro Phe Glu Asp Ile Arg Ser Arg Phe Ile Gln Gly Glu
 405 410 415
 Ser Ile Thr Pro Thr Lys Ile Pro Glu Ile Thr Thr Glu Ile Tyr Met
 420 425 430
 Ser Asn Gly Ser Asn Ser Ala Asn Ile Val Pro Glu Thr Ser Pro Glu
 435 440 445
 Ser Val Pro Ile Ile Ser Val Ser Leu Leu Phe Ala Gly Val Ile Ala
 450 455 460
 Ala Arg Ile Gly Leu Trp Ser Phe Asp Leu Thr Val Thr Gln Leu Leu
 465 470 475 480
 Gln Glu Asn Val Ile Glu Ser Glu Arg Gly Ile Ile Asn Gly Val Gln
 485 490 495
 Asn Ser Met Asn Tyr Leu Leu Asp Leu Leu His Phe Ile Met Val Ile
 500 505 510
 Leu Ala Pro Asn Pro Glu Ala Phe Gly Leu Leu Val Leu Ile Ser Val
 515 520 525
 Ser Phe Val Ala Met Gly His Ile Met Tyr Phe Arg Phe Ala Gln Asn
 530 535 540
 Thr Leu Gly Asn Lys Leu Phe Ala Cys Gly Pro Asp Ala Lys Glu Val
 545 550 555 560
 Arg Lys Glu Asn Gln Ala Asn Thr Ser Val Val
 565 570

<210> 3
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:PCR primer

<400> 3
 gctcagggcg tccgctaggc t

<210> 4
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR primer

<400> 4
ggcttacacc ctcatgttct 20

<210> 5
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR primer

<400> 5
ccttttgata aggaagcaac ttcc 24

<210> 6
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR primer

<400> 6
cagaggtagc tcaggcattg gtcc 24

<210> 7
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR primer

<400> 7
gtggcagggt ctgttctggt cctggaag 28

<210> 98
<211> 24
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:peptide

<400> 8
Leu Leu Leu Thr Ala Val Tyr Gly Leu Val Val Ala Gly Ser Val Leu
1 5 10 15
Val Leu Gly Xaa Ile Ile Gly Asp
20
565 570

专利名称(译)	铁转运蛋白1基因突变与遗传性血色病相关		
公开(公告)号	EP1352092A2	公开(公告)日	2003-10-15
申请号	EP2001987815	申请日	2001-10-17
[标]申请(专利权)人(译)	PIETRANGELO ANTONELLO		
申请(专利权)人(译)	pietrangelo, 先生		
当前申请(专利权)人(译)	pietrangelo, 先生		
[标]发明人	PIETRANGELO ANTONELLO		
发明人	PIETRANGELO, ANTONELLO		
IPC分类号	A01K67/027 A61K38/00 A61K39/395 A61K48/00 A61P43/00 C07K14/47 C07K16/18 C12N1/15 C12N1/19 C12N1/21 C12N5/10 C12N15/09 C12Q1/68 C12Q1/6883 G01N33/50 G01N33/68 A61K38/17 C12N15/11 A61K31/70 C07K16/28 G01N33/53 C12N15/85 C07K7/00 C12N15/52 C07K14/435		
CPC分类号	A61K38/00 A61K48/00 A61P3/02 C07K14/47 C12Q1/6883 C12Q2600/156 C12Q2600/158 G01N33/5008 G01N33/502 G01N33/5091 G01N33/68		
优先权	102000900882061 2000-10-17 IT		
其他公开文献	EP1352092B1		
外部链接	Espacenet		

摘要(译)

本发明涉及编码与遗传性血色病相关的铁转运蛋白1的基因中的突变和基于这种突变的鉴定诊断遗传性血色病的方法。