

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 September 2002 (26.09.2002)

PCT

(10) International Publication Number
WO 02/074782 A2

(51) International Patent Classification⁷: C07K

(21) International Application Number: PCT/BR02/00041

(22) International Filing Date: 18 March 2002 (18.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PI0101088-3 19 March 2001 (19.03.2001) BR

(71) Applicants (*for all designated States except US*): **BIOLAB SANUS FARMACÊUTICA LTDA.** [BR/BR]; 5386 Dos Bandeirantes Av., Planalto Paulista, São Paulo 04071-900 (BR). **FUNDAÇÃO DE AMPARO À PESQUISA DO ESTADO DE SÃO PAULO - FAPESP** [BR/BR]; 1500 Pío Xi St., Alto Da Lapa, São Paulo 05468-901 (BR).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MARTINS DE CAMARGO, Antônio, Carlos** [BR/BR]; 55 Mário Ferraz St. Rm. 11, Jardim Europa, São Paulo 01453-010 (BR). **CALHETA VIEIRA PORTARO, Fernanda** [BR/BR]; Centro de Toxicologia Aplicada "Cat/Cepi", 1500 Vital Brasil Av., Butantã, São Paulo 05503-900 (BR). **FERRAGINI MURBACK, Alessandra** [BR/BR]; Centro de Toxicologia Aplicada "Cat/Cepi", 1500 Vital Brasil Av., Butantã, São Paulo 05503-900 (BR). **ALVES IANZER, Danielle** [BR/BR]; Centro de Toxicologia Aplicada "Cat/Cepi", 1500 Vital Brasil Av., Butantã, São Paulo 05503-900

(BR). **POLISELLI FARSKY, Sandra, Helena** [BR/BR]; Centro de Toxicologia Aplicada "Cat/Cepi", 1500 Vital Brasil Av., Butantã, São Paulo 05503-900 (BR). **PALMA, Mario, Sergio** [BR/BR]; Centro de Toxicologia Aplicada "Cat/Cepi", 1500 Vital Brasil Av., Butantã, São Paulo 05503-900 (BR).

(74) Agent: **LLC INFO CONNECTION LTDA.**; 60 Hermenegarda St., RM 403, Méier, 20710-010 Rio de Janeiro (BR).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ISOLATION AND PURIFICATION PROCEDURE OF VASOPEPTIDASE PEPTIDE INHIBITORS

(57) Abstract: The present invention patent refers to the isolation and purification of peptides secreted by serpent venom glands, specifically *Bothrops jararaca*; to the peptide thus obtained, as well as to the production procedures by genetic engineering techniques in procaryotic and eukaryotic systems; to the engineered peptide thus obtained; to the production of said peptide by chemical synthesis, as well as to the peptide resulting from this chemical processing. It also refers to the utilization of said peptides, obtained by different procedures, in distinct pharmaceutical compositions, and introduced into the organism by a variety of means, in order for them to act as inhibitors of vasopeptidases, and consequently reduce systemic arterial blood pressure, and show local vasodilating action.



WO 02/074782 A2

"ISOLATION AND PURIFICATION PROCEDURE OF VASOPEPTIDASE PEPTIDE INHIBITORS, SPECIFIC FOR THE CARBOXYLIC SITE OF THE ANGIOTENSIN CONVERTING ENZYME, SECRETED BY SERPENT VENOM GLANDS (BPPS), PARTICULARLY *Bothrops jararaca*, OR PRODUCED
5 ENDOGENOUSLY (EVASINS), PRESENTING VASODILATOR AND ANTI-HYPERTENSIVE ACTION; PROCEDURE FOR THE AMINO ACID SEQUENCE DETERMINATION OF PEPTIDE INHIBITORS SECRETED BY SERPENT VENOM GLANDS (BPPs), OR ENDOGENOUSLY PRODUCED (EVASINS); AMINO ACID SEQUENCING PROCEDURE OF THE BPPs BY DEDUCTION OF
10 THE cDNA OF THE PRECURSORS OF THESE MOLECULES EXPRESSED IN SERPENT TISSUES, PARTICULARLY *Bothrops jararaca*; AMINO ACID SEQUENCE DETERMINATION PROCEDURE OF THE EVASINS BY DEDUCTION OF THE cDNA OF THE PRECURSORS OF THESE MOLECULES EXPRESSED IN SERPENT TISSUES, PARTICULARLY *Bothrops*
15 *jararaca*; cDNA AMPLIFICATION PROCEDURE FROM SERPENT BRAIN AND/OR PANCREAS cDNA LIBRARIES, PARTICULARLY FROM *Bothrops jararaca*; SOLID PHASE SYNTHESIS PROCEDURE OF VASOPEPTIDASE PEPTIDE INHIBITORS WITH VASODILATOR AND ANTI-HYPERTENSIVE ACTION, OF VASOPEPTIDASE PEPTIDE INHIBITORS WITH ANTI-
20 HYPERTENSIVE ACTION; USE OF VASOPEPTIDASE PEPTIDE INHIBITORS WITH VASODILATOR AND ANTI-HYPERTENSIVE ACTION TO OBTAIN PHARMACEUTICAL COMPOUNDS; DETERMINATION PROCEDURE OF THE INHIBITORY ACTIVITY ON VASOPEPTIDASES AND OF THE BIOLOGICAL ACTIVITY ON SMOOTH MUSCLE, AND ON THE
25 MICROCIRCULATORY AND CARDIOVASCULAR SYSTEMS."

The invention refers to the isolation and purification procedures of vasopectidase peptide inhibitors, specific for the carboxylic site of the angiotensin converting enzyme, secreted by serpent venom glands (BPPs), or
30 endogenously produced (EVASINS), presenting anti-hypertensive and vasodilator actions. It also refers to the isolation and determination procedures of the amino acid sequence of the peptide inhibitors secreted by serpent venom glands (BPPs), or endogenously produced (EVASINS). It

also refers to the amino acid sequence determination procedure of BPPs by deduction of the cDNA of the precursors of these molecules, expressed in serpent tissues, particularly *Bothrops jararaca*; to the
5 amplification procedure of the cDNA from brain or pancreas cDNA libraries, particularly from *Bothrops jararaca*; to the solid phase synthesis procedure of vasopeptidase peptide inhibitors; to the use of vasopeptidase peptide inhibitors with vasodilator and anti-hypertensive action. It further
10 refers to the determination procedure of the inhibitory activity on vasopeptidases, and of the biological activities on smooth muscle, and on the microcirculatory and the cardiovascular systems.

Specifically, the invention refers to the isolation of
15 the peptides secreted by serpent venom glands (BPPs), or found endogenously, particularly in the brain of *Bothrops jararaca*. We suggest the generic name EVASINS (**endogenous vasopeptidases inhibitors**) for the endogenous peptides; to the peptides thus obtained, as well as to the procedure of
20 producing them by genetic engineering methods in prokaryote and eukaryote systems; to the engineered peptides thus obtained; to the production of said peptides by chemical synthesis as well as to the peptides resulting from the proteolytic processing. It also refers to the use of the
25 peptides obtained by the referred procedures in pharmaceutical compounds with inhibitory action on vasopeptidases, and consequently lowering blood pressure, and the use of the referred peptides to lower blood pressure.

30 Thus the invention refers to peptides with cardiovascular action, known as bradykinin potentiating peptides, secreted by serpent venom glands (BPPs), or produced in other tissues of *Bothrops jararaca* (EVASINS).

The serpent venom is composed of a complex mixture of biologically active proteins and peptides. Among known, characterized biologically active peptides, we list bradykinin potentiating peptides (BPPs), natriuretic factors (C-type natriuretic peptides, CNP), saraphotoxins (SRTx), among others. These bioactive peptides (hormones, neuropeptides, toxins, etc.) are generated by the hydrolytic action of enzymes on the proteic precursors of these peptides, which contain one or more bioactive peptides per molecule. Toxins secreted by the venom glands of various serpents, such as *Bothrops jararaca*, contain a large number of peptides with hypotensive action (BPPs), whose mechanism of action is related to its inhibitory effect on the angiotensin-converting enzyme (ACE), blocking the conversion of angiotensin I into angiotensin II (hypertensive), and preserving bradykinin (hypotensive).

The primary structure of these peptides allows their classification into two families: those peptides smaller than 7 amino acid residues, and those larger than 7 amino acid residues, showing a high degree of homology among them.

Research in this specific field showed that a large variety of applications can be found for toxins, particularly the venom toxins of serpents, or the modifications of some of these molecules, such as the bradykinin potentiating peptides (BPPs), which can be specific agents of anti-hypertensive action in mammals, including man.

The amino acid sequence of other bradykinin potentiating peptides, having high degree of homology with the amino acid sequence of BPPs, were not obtained from the venom of snakes. In fact, these were deducted from the nucleotide sequences of their precursors found in other serpent tissues, such as the brain, pancreas and spleen

tissue, using molecular biology techniques. These endogenous peptides, in their synthetic form, like the BPPs secreted from the venom gland of the *Bothrops jararaca*, acting on the angiotensin converting enzyme (ACE) of the vascular endothelium, preventing the formation of angiotensin II (hypertensive) and preserving the hypotensive action of bradykinin.

The angiotensin converting enzyme (ACE) is a peptidase with two catalytic sites (carboxy- or C-site, and amino- or N-site), which are mainly located to the cytoplasmic membrane of endothelial cells. The C-terminal active site is more specific for angiotensin I and bradykinin. The BPPs act by inhibiting the ACE and, up to this invention, the specificity of the inhibition of the C- or N-sites was unknown, and the result of this inhibition is the reduction of arterial blood pressure.

Inhibition of the ACE, not selective for a specific active sites, led to the development of the site-directed inhibitor called captopril.

The N site metabolizes the natural circulating peptide hormone (Ac - Ser - Asp - Lys - Pro), which regulates hematopoiesis. The carboxylic domain of the enzyme (C site) is more specific for conversion of angiotensin I into angiotensin II and inactivates bradykinin. Up to this moment, it was not known whether all the described BPPs presented any preference for the C site, turning them into more specific antihypertensive agents.

There still are a number of unknown toxins from the venoms of snakes, which may have the potential to act as anti-hypertensive agents. These molecules may not only inhibit specifically the C-site of the ACE, but even, and simultaneously, inhibit other peptidases of the vascular

endothelium, as the neutral endopeptidase (NEP) and the endothelin converting enzyme (ECE). The ECE inactivates bradykinin, natriuretic peptides, hypotensive agents, and generates endothelins (hypertensive). The ACE, NEP and ECE are known as vasopeptidases, and are responsible for most of the arterial blood pressure control. These double and/or triple inhibitors are not only more specific, but also more efficient because they act on the metabolism of endogenous molecules regulating the arterial blood pressure. The distinction between the two catalytic sites of the ACE has been well-defined in: Dive, V.; Cotton J.; Yiotakis, M. A.; Vassiliou, S.; Jiracek, J.; Vazeaux, G.; Chauvet, G.; Cuniasse, P.; Corvol; *RXP 407, a phosphinic peptide, is a potent inhibitor of angiotensin I converting enzyme able to differentiate between its two active sites.* Proc. Natl. Acad. Sci. USA, 13, 4330-4335, (1999).

It is known from the scientific literature that some peptides from the venom of Brazilian and Asian serpents, presenting hypotensive action, have already been isolated and sequenced (Ondetti & Cushman, Ann.Rev.Biochem. 51, 293 - 308, 1982), and that the antihypertensive efficiency of one of them has been demonstrated in humans (Gravas et al. N. Engl. J. Med. 291, 817 - 821, 1974).

Several BPPs have been deduced from the amino acid sequences of their precursors using molecular biology techniques (Murayama et. al., Proc. Natl. Acad. Sci. USA, 94, 1189 - 1193, 1997).

The pharmaceutical industry developed a new antihypertensive drug of non-peptidic nature and without any chemical analogies with endogenous molecules. It was called OMOATRILAT, displaying inhibition properties for both the ACE and the NEP, presenting long-term anti-hypertensive efficiency in rats with high concentration of circulating renin. Besides, this substance improves cardiac

performance, prolonging the survival of spontaneously hypertensive rats, SUR, a model for human essential hypertension. The OMAPATRILAT is the first site-directed inhibitor for two vasopectidases, the ACE and NEP, in
5 advanced phase of clinical trials.

The major differences between the OMAPATRILAT and the BPPs and EVASINS are:

- 1- the BPPs and EVASINS are of polypeptidic nature;
- 2- they are synthetic molecules identical or homolog
10 to peptides secreted by exocrine glands, or endogenously produced in vertebrates;
- 3- as ACE inhibitors, the BPPs and EVASINS of 8 to 13 amino acid residues show preference for the C-site, and also inhibit the NEP X;
- 15 4- they are degraded mainly by humoral and tissue proteolytic enzymes;

The US 55389991 patent, published July 23rd, 1996, and the US 5559135 patent, published September 24th, 1996, describe the inhibitor secreted by serpent venom glands -
20 PCA-W-P-R-P-E-I-P-P- SQ 20881 - an ACE inhibitor, which consequently reduces the arterial blood pressure in animals, and which showed efficient reduction of arterial blood pressure in hypertensive individuals. However, these patents do not describe or reveal the specificity of action
25 of this inhibitor on one or the two catalytic sites of the ACE. The patents do not show either the importance of other enzymes capable of regulating the arterial blood pressure, therefore no research was done on BPPs with higher specificity of action.

30 The present invention differs from the aforementioned patents by demonstrating specificity of action of the EVASINS on vasopectidases; it also establishes a clear relationship between the chemical structures of several dozens of these molecules with their activity on the

cardiovascular system. This invention also establishes one structural motif and the endogenous origin of the leader molecules proposed here, features not predicted in the references quoted above.

5 The same arguments mentioned above can be used to differentiate the present invention from the USPTO patents US 3819831, published June 25th, 1974, US 3714140, published January 30th, 1973, US 3832337, published August 27th, 1974, US 3849252, published November 19th, 1974, US
10 3947575, published March 30th, 1976, US 3973006, published August 3rd, 1976, US 4731439, published March 15th, 1988, US 4774318, published September 27th, 1988, and US 5550127, published August 27th, 1996.

 Aiming at solving the existing technical problems,
15 in order to obtain a final product of superior quality in terms of efficiency as anti-hypertensive agents, the present invention proposes, for the first time and in a novel manner, BPPs and EVASINS, presenting: (1) a defined structural motif; (2) specificity for the C-site of the
20 ACE; and (3) high inhibitory specificity for the NEP X.

 These novel specific anti-hypertensive agents for the C site of the ACE, presenting inhibitory action on vasopeptidase were isolated and sequenced from the venom of the Brazilian serpent *Bothrops jararaca*, or identified by
25 cloning and cDNA sequencing, obtained from the tissues of this serpent. These peptides, which can also be obtained by chemical synthesis, and which contain the same amino acid sequence as the natural peptides, show long-term anti-hypertensive action in rats. The use of these synthetic
30 peptides, which have the same sequence as the natural peptides, used as such, or chemically modified in order to maintain the characteristics described above, can be useful as therapeutic agents in the treatment of cardiovascular disorders or in local circulatory actions, particularly in

hypertensive disorders in humans and other cardiovascular disorders.

Twenty-two (22) BPPs found in the venom and in tissues of *Bothrops jararaca*, were sequenced by mass spectrometry, or their sequences were deduced from cDNA sequences of the precursors of those molecules expressed in the tissues of the serpent. The corresponding synthetic peptides were tested as inhibitors of the C-site of the recombinant ACE and of NEP, and also both as potentiating agents of the bradykinin contractile activity on isolated guinea pig ileum, and as bradykinin hypotensive activity in rats. They were found both as BPPs (secreted) and as EVASINS (endogenous). They can be divided in two groups according to the specificity of action on the C site of the ACE:

a) non-specific. One peptide, the pentapeptide <E K W A P (Pyroglutamyl(<E)-Lysine(K)-tryptophane(W)-Alanine(A)-Proline (P), which, although not presenting a defined selectivity for the C-site of the ACE, presents high potentiating activity of bradykinin, as observed on the preparation of isolated smooth muscle as well as by its hypotensive effect. It was found as BPP (secreted from the serpent venom glands) as well as EVASINS (endogenous, from the serpent brain tissue). It presents K_i values in the μM range, for both the N and C sites of the ACE. It also inhibits, with low affinity, the NEP with K_i values ranging from 50-150 μM . It potentiates the contractile action of bradykinin on isolated guinea pig ileum with concentrations that vary from 15-300 nM. The hypotensive effect of bradykinin was also potentiated by 60% in concentrations of 300 nM, showing a 5-fold (five) increase in the duration of the hypotensive effect, when compared to the duration of the effect of bradykinin alone.

b) specific. These peptides present molecular masses ranging from 1000 to 1700 Daltons, containing 8 to 13 amino acid residues. In this group, the most selective and effective potentiators of the contractile action of the bradykinin on isolated guinea pig ileum and on rat arterial blood pressure, were found as BPPs or EVASINS. They were chemically modified generating other peptides with similar pharmacological properties.

The table 1 below presents the synthetic oligonucleotides derived from the inhibitors described in group b) for this present invention:

Table 1

Formula Sequences

I	$pp^1aa^1aa^2aa^3P^4aa^5aa^6P^7P^8$
II	$pp^1aa^1aa^2aa^3aa^4P^5aa^6aa^7P^8P^9$
III	$pp^1aa^1aa^2aa^3aa^4aa^5P^6aa^7aa^8P^9P^{10}$
IV	$pp^1aa^1aa^2aa^3aa^4aa^5aa^6P^7aa^8aa^9P^{10}P^{11}$
V	$pp^1aa^1aa^2aa^3aa^4aa^5aa^6aa^7P^8aa^9aa^{10}P^{11}P^{12}$
VI	$pp^1aa^1aa^2aa^3aa^4aa^5aa^6aa^7aa^8P^9aa^{10}aa^{11}P^{12}P^{13}$

Where:

P is always proline. All others are always L-amino acids and are presented in the one-letter-code (see below).

pp^1 is the N-terminal and can be also pyroglutamyl (<E) or any other simple amino acid, generally non-basic.

aa^1 is a non-basic amino acid, generally W, S, G or N;

aa^2 is a non-acid amino acid, generally P, G, W or R;

aa^3 is a non-acid amino acid, generally P, A, R or W;

aa^4 is P for formula I and it is generally the amino acid T, P, G, H, R, W or E for all other oligopeptides;

aa⁵ is generally Q, N, P, or G for formula I;
 aa⁶ for formula II, aa⁷ for formula III, aa⁸ for formula IV
 and aa⁹ for formula V are non-basic amino acids,
 generally Q, N, P, or G.

5 aa⁶ for formula I, aa⁷ for formula II, aa⁸ for formula III,
 aa⁹ for formula IV and aa¹⁰ for formula V are always I, A
 or T.

	Acid amino acids -	D, E
10	Basic amino acids -	K, R
	Aromatic amino acids -	F, W, Y

	G-glycine	N-asparagine
	A-alanine	Q-glutamine
15	P-proline	D-aspartic acid
	V-valine	E-glutamic acid
	I-isoleucine	K-lysine
	L-leucine	R-arginine
	S-serine	F-phenylalanine
20	T-threonine	H-histidine
	W-tryptophan	Y-tyrosine
	<E- pyroglutamine	

Of particular interest in group (b) are the peptides
 of 8 to 13 amino acids presenting their general formula the
 25 carboxy-terminal motif of the oligopeptide:

STRUCTURAL FORMULA [P X¹ X² P P]

where X¹ can be any amino acid and X² is generally I, and
 the N-terminal amino acid is blocked, generally by <E.

The peptides described in the above table show
 30 higher selectivity for the C-site of the ACE as
 characterized by the Ki values, ranging from 2 nM to 100
 μM, while for the N site of the ACE these values were above

50 μM . These peptides are also inhibitors of NEP with K_i values ranging from 5 to 150 μM . All of them potentiate the contractile activity of bradykinin on the isolated guinea pig ileum, doubling the contractile effect of bradykinin, in concentrations varying from 5 to 300 nM. The hypotensive effect of bradykinin was also potentiated by 30-80% in concentrations varying from 10-700 nM, increasing the duration of this hypotensive effect from 5 to 15 fold, as compared to bradykinin alone.

10 The present invention proposes for the first time: an isolation and purification procedure for vasopeptidase peptide inhibitors with anti-hypertensive action, secreted by serpent venom glands, particularly *Bothrops jararaca*, comprising the following steps:

15 A - ISOLATION AND PURIFICATION OF BPPs FROM THE VENOM OF *Bothrops jararaca*:

- 800 to 1500 mg of total venom obtained from a venom pool of *Bothrops jararaca* were dissolved in 7.0 to 15 ml of deionized water and centrifuged at 1500 to 2000 rpm for 15 to 30 minutes; the supernatant was removed and applied to a 1.2 x 101 cm Sephadex G-25 M gel filtration column (25-80 μ , Sigma); the column was equilibrated with ammonium acetate buffer (30 to 50 mM, pH 5.0 to 6.0) at room temperature. The sample was added to the top of the column and the components were eluted at a flow rate of 1.0 to 2.0 mL/min. The absorbance profile for each aliquot at 214 nm defined the constitution of the pools, based on the bradykinin potentiating activity on guinea pig ileum.

30 B - PARTIAL ISOLATION OF THE POOL COMPONENTS SHOWING BRADYKININ POTENTIATING ACTIVITY:

- High performance liquid chromatography - HPLC

The components of the pools obtained in step A, showing bradykinin potentiating activity, were partially isolated by high performance liquid chromatography - HPLC (Merck-Hitachi model L-6200A), with the UV-vis detector set at 214 nm, and the reverse-phase column C-18/Beckman (5 μ , 4.6 x 250 mm). Solvent A was 0.5 to 1 % TFA (trifluoroacetic acid) in H₂O, and solvent B was a 5% to 60 % gradient of acetonitrile (ACN), at a flow rate of 0.2 to 0.5 mL/min. The gradients used varied with the sample. The peptides were obtained by manual collection of the absorbance peaks at 214 nm.

C - DETERMINATION OF THE MOLECULAR MASS AND OF THE PRIMARY STRUCTURE OF THE BRADYKININ POTENTIATING PEPTIDES BY MASS SPECTROMETRY (ES-MS-MS).

Another aspect of the invention is related to the procedure for the amino acid sequence determination of the BPPs purified from serpent venom, more specifically *Bothrops jararaca*, by mass spectrometry performed in the Micromass Quattro II mass spectrometer (ESMS-MS / Micromass), in the positive ionization mode, with an electrospray ion source (Micromass), and the Mass Lynx software (Micromass) used for data acquisition. The samples were dissolved in 50% H₂O/ACN with 0.1% formic acid, and injected with a constant flow of 5 μ L/min, by means of an injection pump. The data were acquired in the first quadrupole (ESMS), by scanning the ratio mass/charge (m/z) between 400 and 1600 with a scanning time of 5 seconds during the whole analysis process. Sequencing was performed on selected peptide, displaying a protonated ion, as characterized in the first quadrupole, followed by fragmentation by induced dissociation collision (IDC) with a pressure of

argonium of 3×10^{-3} , and finally, the data were acquired by scanning at the second quadrupole (ESMS-MS). The software Mass Lynx (Micromass) was used for data acquisition. The machine's parameters were

5

optimized using synthetic bradykinin as a standard. Characterization and sequencing can also be performed with the HPLC system coupled to the mass spectrometry system (LC-ESMS-MS), using the Hewlett-Packard HPLC model 1100 with an automatic injector, the UV-vis

10

detector set at 214 nm, and the reverse-phase column C-18/The Separations Group ($4.6 \times 250 \text{ mm}/5\mu$), for the isolation of total venom components. Solvents used were: Solvent A: 0,1% TFA/H₂O - Solvent B: 10% A/ACN solvent, with a flow rate of 0.6 mL/min. The gradient

15

used was:
t = 0 - 5 min: 0% B
t = 5 - 65 min: 0 - 60% B
t = 65 - 70 min: 60 - 100% B
t = 70 - 75 min: 100% B

20

t = 75 - 80 min: 100 - 0% B
Out of the 600 $\mu\text{L}/\text{min}$ flow used in the HPLC, 20 μl were automatically injected into the mass spectrometer, and the remaining 580 μl went to the UV-vis detector.

25

Mass spectrometry was accomplished in a Micromass Quattro II (ESMS - MS/Micromass) mass spectrometer in the positive ionization mode with an ion electrospray source (Micromass). The acquisition of the characterization and sequencing data was performed by

30

means of a software created to this end, the Mass Lynx software (Micromass).

- 5 - Another aspect of this invention is the determination of the amino acid sequence of the BPPs by deduction of the cDNA sequence of the precursors of these molecules expressed in serpent tissues, specifically *Bothrops jararaca*, through the following steps:
- 10 - Total RNA from the brain of a single *B. jararaca* specimen was isolated by the guanidine isothiocyanate - phenol - chloroform extraction method, and the messenger RNA was purified by passing the total RNA solution twice through a pre-packed oligo-dT cellulose column. The integrity of the purified mRNA was checked by agarose gel electrophoresis followed by hybridization with radioactive probes, composed of
- 15 sequences encoding the precursor of the BPPs and the CNP, previously identified by Northern blot assays. Preparation of the cDNA library in λ ZAP phages (Stratagene) was carried out using 5 μ g of this sample, following the manufacturer's instructions by common
- 20 and usual methods. The initial packaging of the recombinant phages generated a titer of approximately 2×10^5 pfu/mL, with less than 1% of non-recombinant clones. This library was immediately amplified, with a final titer of approximately 9×10^9 pfu/mL.
- 25 F - PCR AMPLIFICATION (Polymerase Chain Reaction):
- 30 - The goal of the initial approach was to amplify the cDNA of interest from *B. jararaca* brain cDNA libraries by PCR, using oligonucleotides specific for the cDNA sequence encoding the precursor of the BPPs and the CNP. The reactions were carried out with materials and methods described here and it was possible to clone only one fragment of approximately 250 base pairs from the total phage lysate from the serpent brain cDNA

library (there was no amplification of any other fragment from the serpent brain library). Complete sequencing of this cDNA insert showed a very high degree of similarity with the cDNA sequence identified from the venom gland library, except for the insertion of a few nucleotides (12 bases) in the segment that precedes the natriuretic peptide and some point mutations (4 residues substituted), of which three are silent (encoding the same amino acid residue) and the fourth was "conservative" (a substitution for an amino acid with the same chemical characteristics). In spite of the numerous amplification attempts, changing reaction conditions, and/or the polymerase and oligonucleotide used, it was not possible to clone other fragments from any of the libraries studied.

- Approximately 5×10^6 clones were independently analyzed using the segment encoding the natriuretic peptide as a template for the synthesis of radioactive probes used for the hybridization experiments, which made it possible to identify only 13 positive clones. The plasmid vectors containing the cDNA inserts were recovered by mini-preparations of DNA and then sequenced using oligonucleotides, which anneal to sequences adjacent to the plasmid vector's (pBluescript SK+) multiple cloning site (commercial primers T3 and T7).

- Clone selection by hybridization of the cDNA library from the brain of *B. jararaca* allowed the isolation of a cDNA encoding the precursor for the BPPs and the CNP. (Figure 1). By analogy to the BPPs isolated from the venom of *Bothrops jararaca*, 7 peptides were identified, whose sequences are listed among those in table 1). Based on these sequences, peptides were synthesized that presented the same pharmacological

properties as the venom peptides, as well as inhibition of ECA and NEP.

Figure 1 shows the primary sequence of the EVASINS precursor from *Bothrops jararaca* brain, deduced from the corresponding cDNA sequence. Abbreviations for the nucleotides: **A**-adenosine; **C**-cytidine; **G**-guanosine; **T**-thymidine. **CNP**- **Type C** natriuretic peptide. Nucleotide sequence of the cDNA encoding the precursor protein of the EVASINS and **CNP**, isolated from the cDNA library from *Bothrops jararaca* brain, and the deduced amino acid sequence. The amino acid sequences of the EVASINS are underlined and the **CNP** sequence is shown in bold.

Another aspect of the invention is the procedure for the synthesis of the peptides in solid phase, using two specific strategies:

A - Use of the tert-butyloxycarbonyl group (Boc) as a temporary protector for the amino group and of benzyl derivatives (Bzl) for the protection of most of the amino acid reactive side chains.

- The first step in this synthesis strategy is to remove the Boc group from the first amino acid residue, bound to the resin.

Removal of the Boc group occurs in 30% TFA in dichloromethane (DCM), containing 2% anisole, for 30 min. The resin is then washed successively with isopropanol containing 0.5% anisole, DCM and MeOH. For the next amino acid coupling, the amino group is deprotonated by treatment with TEA 10% or DIPEA 5% in DCM for 10 min. The peptidyl-resin is then washed with DCM, MeOH, DCM and with the solvent used for the coupling.

Coupling of the amino acid starts with an activation phase, usually accomplished with coupling agents: diisopropylcarbodiimide (DIC) or 2- (1H- benzotriazolyl)-1,1,3,3-tetramethylurone tetrafluoroborate (TBTU). Usually

the Boc-amino acid (carboxylic component, CC) and the coupling agent are used in a 3-fold molar excess, as compared to the amount of amino groups (amine component, CA) in the resin.

5 - When DIC is used, the proportion of the reagents CA:CC:DIC is 1:3:3. When TBTU is used, the proportion of the reagents CA:CC:TBTU:DIEA is 1:3:3:4. Final concentration of these reagents is between 0.05 and 0.1 M.

10 - When coupling Asn (asparagine) or Gln (glutamine), hydroxibenzotriazol (HOBt) is used with the coupling agent to prevent formation of nitrile from these amino acids. For these couplings the final proportion of the reagents CA:CC:TBTU:HOBt:DIEA is 1:3:3:3:5. The coupling reaction, which normally takes 2 hours, is monitored using the
15 ninidrine qualitative method, in which a blue staining of the peptidyl-resin is indicative of incomplete coupling. In this case, recoupling is generally performed changing the solvent used in the previous coupling reaction and/or the coupling agent itself. When the coupling phase is finished,
20 the next cycle is initiated with the deprotection of the amino group until the next residue is coupled. The process continues in cycles until the end of the elongation of the desired sequence.

25 At the end of the synthesis procedure, the peptide is cleaved from the resin, and its side chains are deprotected in a single step by treatment with anhydrous HF at 0° C for 60-90 min in the presence of p-cresol and dimethylsulfide (DMS) (5%, v/v of each) which act as suppressors for collateral reactions induced by carbocations released
30 during the HF treatment. When the peptide sequences contain Trp residues, ethanol-1,2-dithiol (EDT) is added to remove the formyl group that protects the side chain of this amino acid simultaneously to the cleavage. EDT is used in the same proportion as the other suppressors (5% v/v, of each).

After cleavage, the resin is washed with ethyl acetate and the peptide is eluted by washing the resin with 5% (v/v) acetic acid in water, and lyophilized. A white powder is obtained, generally amorphous.

5 B - Use of the base-labile protector 9-fluorenylmethoxycarbonyl (Fmoc) and tert-butyl derivatives (tBu).

- Differently from the previous strategy, in this case the amino group protector Fmoc is used, and removed from
10 the amino acid by treatment with a 20% piperidine solution in DMF for about 20 min. This treatment deprotonates the amino groups for the forthcoming coupling step. The coupling methods are the same as the Boc chemistry's and the final cleavage is performed using a weaker acid
15 solution, because both the protector's bond to the side chains (generally of the tert-butylic type) and the peptide's bond to the resin are more labile than in the Boc chemistry's. Usually, the resin is treated with 85-95% TFA (v/v) and a mixture of different suppressors of collateral
20 reactions for about 2 h. After this treatment, the peptide is precipitated with the resin in ice-cold ethylic ether, and centrifuged at a speed of 2,500g for 5 min. The supernatant is discarded and the pellet is resuspended in ethylic ether, vortexed and centrifuged. The procedure is
25 repeated 5 times and the resin/peptide mixture is dried under vacuum. Finally, the peptide is removed from the resin with 5% AcOH in water (v/v) and filtered. The filtered solution is lyophilized until an amorphous white powder is obtained.

30 Both strategies A and B are based on the following references: 1-Barany, G. & Merrifield, R.B. (Gross, E. & Meinhofer, J., Eds.) (1980), The Peptides: Analysis, Synthesis and Biology, vol. II, 1, Academic Press, New York. 2- Stewart, J.M. & Young, J.D. (1984), Solid Phase

Peptide Synthesis, Pierce Chemical Company, Rockford. 3-
Fields, G.B. (1997), Methods in Enzymology, Academic Press,
California; 3- Atherton, E. & Sheppard, R.C. (1989), Solid
Phase Peptide Synthesis: A Practical Approach, I.L.R.
5 Press, Oxford. 4- Fields, G.B. & Noble, R.L. (1990) Int. J.
Pep. Prot. Res. 35, 161.

Following the methods described in this invention, a
pool of isolated peptides is obtained, presenting
bradykinin potentiating activity, as tested by its
10 contractile activity on smooth muscle, caused by
bradykinin, and measured using a preparation of isolated
guinea pig ileum, as shown in Example 5.

To determine the bradykinin potentiating activity, a
log-dose response curve of the effect of bradykinin on the
15 isolated guinea pig ileum was plotted.

The hypotensive effect of the bradykinin potentiating
activity was tested in normotensive and hypertensive (SHR)
male and female Wistar rats, anesthetized with ethylic
ether.

20 In the present invention, the peptides of particular
interest are natural anti-hypertensive peptides. These will
eventually be used as such, or in association with other
substances. These synthetic molecules can be conjugated to
a variety of ligands, such as acid groups, like sulfonyl,
25 carboxyl and phosphoryl, or other groups, such as thiols,
olefins, dithio, azo or diazo compounds, aldehydes and
similar compounds.

BPPs bound to specific ligands (peptide aptamers), can
be used to identify target proteins, such as enzymes and
30 receptors. For example, specific cells can present target
proteins that may be recognized or affected by these
modified BPPs. Thus, the use of specific ligands conjugated
to the BPPs can drive these molecules preferentially to the
cells that possess the specific target proteins. These

ligands can be, for example, methylene blue, crystal violet, aspirin, saccharin or others.

In addition to these, several other compounds can be used as ligands, such as steroids, low density lipoproteins, growth factors, viral proteins, etc. Compounds containing such ligands may be preferentially identified by immunoglobulins or their fragments, recognizing them as antigens, wherever they are located. The immunoglobulins of interest are some of the subtypes of IgA, IgD, IgM, IgE and IgG. Immunoglobulins can be derived from any source, particularly from mice and humans. The immunoglobulins can be derived from hybridomas, such as transformed lymphocytes, or obtained by recombinant DNA methods.

Particularly, the variable part of the mice immunoglobulin can be linked to the human immunoglobulin, to form a chimeric immunoglobulin with low immunogenicity. Only the use of the Fab fragments, F(ab')₂, Fv, or similar, would be necessary.

Ligands can be coupled to the toxins in question, in order to help them enter the cells and identify the target protein(s) inside. Disulfide bonds can be used to release the toxins from the ligands as they are reduced inside the cell.

The compounds in question can be used *in vivo* and *in vitro*. For the *in vivo* use the compounds can be administered trans-mucosa, parenterally, or by injection, particularly intravenously. The dosage will possibly vary from around 1 µg to 10 mg, usually 0,5 mg/kg of body weight.

Depending on the means of administration and the synthetic toxin used, presenting the same amino acid sequence as the naturally occurring toxin, or a modified form, the compound may eventually be used at a very low

dosage. The compound may eventually be diluted in a physiologically acceptable medium, such as a phosphate-saline buffer, saline solution or other more convenient solution.

5 These compounds may eventually be used in cases of chronic or acute cardiovascular disturbances, in which the micro-circulation would be the target of the treatment. In particular, these compounds may eventually be used in human hypertensive disturbances and their consequences, such as
10 those causing vascular lesions, hyperplasia, etc. A possible recommended use might be in cases of acute pulmonary edema. They may eventually be used in modified or unmodified form, for pathophysiological diagnosis, where the identification or interaction of these toxins with the
15 target proteins (enzymes, receptors, etc.) can be useful.

The BPPs and EVASINS in this invention are used in cardiovascular pathologies.

The innovations presented in this invention, in general terms, consist in adding to the less than 20 known
20 BPPs, dozens of other synthetic peptides with amino acid sequences identical to the natural BPPs and EVASINS, or their chemically modified homologues, which present structural motifs defined here, as well as in characterizing their biological actions through their
25 inhibitory activity on the carboxylic site of the angiotensin converting enzyme (ACE) and also on NEP and ECE.

The peptides with these characteristics will be generically designated as BPPs (bradykinin potentiating
30 peptides) and EVASINS (endogenous vasopeptidase inhibitors). The BPPs and EVASINS from this invention correspond to the peptides secreted by the venom gland and to peptides produced in endogenous tissues of *Bothrops jararaca*.

The sequences of the BPPs were determined by mass spectrometry; the sequences of the EVASINS were determined by deduction of the nucleotide sequence of the cDNA encoding the precursors of these molecules, expressed in
5 serpents, specifically *Bothrops jararaca*.

EXAMPLES

Example 1:

- PARTIAL ISOLATION OF THE POOL COMPONENTS SHOWING BRADYKININ POTENTIATING ACTIVITY:

10 - High performance liquid chromatography - HPLC

The components of the peptide pools displaying bradykinin potentiating activity were partially isolated by high performance liquid chromatography - HPLC Merck-Hitachi model L-6200A, UV-vis detector set at 214 nm, with a
15 reverse phase column: C-18 / Beckman (4.6 x 250 mm/5 μ resin) in 0.5% TFA (trifluoroacetic acid) in H₂O, and a 5% to 60% gradient of acetonitrile (ACN) at a flow rate of 0.2 to 0.5 mL/min. The gradients used varied according to the sample. The peptides were obtained by manually
20 fractionating the eluate according to the shape of the absorbance peaks at 214 nm.

Purification of the anti-hypertensive peptides from the venom of the Brazilian serpent *Bothrops jararaca*.

The total venom, obtained from a venom pool of *Bothrops jararaca* (1200 mg) supplied by the Venom Section of
25 Instituto Butantan, was dissolved in 10 mL of deionized water and centrifuged at 800g for 30 min. The supernatant was collected and loaded onto a Sephadex G-25 (25-80 μ , Sigma) gel filtration column (1.2 x 101 cm). The column was
30 equilibrated with 30 mM ammonium acetate buffer, pH 5.5 at room temperature. The sample was loaded onto the top of the column and the components were eluted with a flow rate of 1.2 mL/min. The absorbance profile of each aliquot, at 214 nm, defined the constitution of the pools.

High performance liquid chromatography - HPLC

The HPLC equipment used was a Merck-Hitachi, model L-6200A UV vis detector set at 214 nm, with a reverse phase column: C-18 / Beckman (4.6 x 250 mm/5 μ resin) for the partial isolation of the pools with bradykinin potentiating activity, in 0.5% TFA (trifluoroacetic acid), and a 5% to 60% gradient of acetonitrile (ACN) at a flow rate of 0.5 mL/min.

Example 2:**10 DETERMINATION OF THE MOLECULAR MASS AND OF THE PRIMARY STRUCTURE OF THE BRADYKININ POTENTIATING PEPTIDES BY MASS SPECTROMETRY**

The Mass spectrometry experiments were performed in a Micromass Quattro II mass spectrometer (ESMS-MS/Micromass) in the positive ionization mode with an electrospray ion source (Micromass). The Mass Lynx software was used for data acquisition.

The mass spectrometer's parameters were optimized using synthetic bradykinin as a standard.

20 The samples were dissolved in 50% ACN/H₂O and 0.1% formic acid, and injected at a constant flow rate of 0.5 μ L/min, by means of an injection pump. The data were acquired at the first quadrupole (ESMS), by scanning of the mass/charge ratio (m/z) between 400 and 1600 using a scanning time of 5 seconds during the whole analysis procedure.

30 The peptide, with protonated ion as characterized at the first quadrupole, was sequenced and fragmented by collision of induced dissociation (CID) with a pressure of argonium gas of 3×10^{-3} , and, finally, the data were acquired by scanning at the second quadrupole (ESMS-MS).

HPLC system coupled to the mass spectrometer (LC-ESMS-MS).

The HPLC system used was a Hewlett-Packard model 1100, with automatic injector, UV-vis detector set at 214 nm, with a reverse phase column: C-18 / Beckman (4.6 x 250 mm/5 μ resin) for the isolation of components from total venom.

- 5 - Solvent A: 0.1% TFA/H₂O
 - Solvent B: 10% solvent A/ACN, with a flow rate of 0.6 mL/min.

The gradient used was:

t = 0 - 5 min: 0% B

10 t = 5 - 65 min: 0% - 60% B

t = 65 - 70 min: 60% - 100% B

t = 70 - 75 min: 100% B

t = 75 - 80 min: 100% - 0% B

15 The HPLC flow rate was 600 μ L/min, out of which 20 μ l were automatically injected into the mass spectrometer, while the remaining 580 μ l were sent to the UV-vis detector.

20 Mass spectrometry was carried out in a Micromass Quattro mass spectrometer (Micromass). Sequencing and characterization data were acquired using a software specially developed for this end.

Example 3:

Synthesis of peptides in solid phase

Two strategies were used for peptide synthesis.

25 The first, more traditional strategy, uses the tert-butyloxycarbonyl group (Boc) as a temporary protector for the amino group and benzyl derivatives (Bzl), for the protection of most of the reactive side chains of the amino acids.

30 The second and most recent strategy alternates between the base labile protector 9-fluorenylmethoxycarbonyl (Fmoc) and tert-butyl derivatives (tBu).

a) **The Boc/Bzl strategy** - The first step in this strategy is the removal of the Boc group from the first amino acid residue bound to the resin.

Removal of the Boc group occurs in 30% TFA in dichloromethane (DCM), containing 2% anisole, for 30 min. The resin is washed with isopropanol containing 0.5% anisole, DCM and MeOH. For the coupling of the forthcoming amino acid, the amino group is deprotonated (neutralized) with 10% TEA or 5% DIPEA in DCM for 10 min. The peptidyl-resin is then washed with DCM, MeOH, DCM and the solvent to be used in the coupling phase.

Coupling of the amino acid starts with an activation phase, usually accomplished with coupling agents: diisopropylcarbodiimide (DIC) or 2- (1H- benzotriazolyl)-1,1,3,3-tetramethylurone tetrafluoroborate (TBTU). Usually the Boc-amino acid (carboxylic component, CC) and the coupling agent are used in a 3-fold molar excess, as compared to the amount of amino groups (amine component, CA) in the resin.

When DIC is used, the proportion of the reagents CA:CC:DIC is 1:3:3. When TBTU is used, the proportion of the reagents CA:CC:TBTU:DIEA is 1:3:3:4.

Final concentration of these reagents is between 0.05 and 0.1 M. When coupling Asn (asparagine) or Gln (glutamine), hydroxibenzotriazol (HOBt) is used with the coupling agent, to prevent the formation of nitrile from these amino acids.

For these couplings the final proportion of the reagents CA:CC:TBTU:HOBt:DIEA is 1:3:3:3:5. The coupling reaction, which normally takes 2 hours, is monitored using the ninidrine qualitative method and a blue staining of the peptidyl-resin is indicative of incomplete coupling. In this case, recoupling is generally performed changing the

solvent used for the previous coupling and/or the coupling agent.

When the coupling phase is finished, the next cycle is initiated with the deprotection of the amino group until
5 the next residue is coupled. The process goes on cyclically until the end of the elongation of the desired sequence.

- At the end of the synthesis procedure, the peptide is cleaved from the resin and its side chains are deprotected in a single step by treatment with anhydrous HF at 0°C for
10 60-90 min in the presence of p-cresol and dimethylsulfide (DMS) (5%, v/v each) which act as suppressors for collateral reactions induced by carbocations released during the HF treatment.

When the peptide sequences contain Trp residues,
15 ethanol-1,2-dithiol (EDT) is added to remove the formyl group that protects the side chain of this amino acid simultaneously with the cleavage. EDT is used at the same proportion as the other suppressors (5% v/v, each). After cleavage, the resin is washed with ethyl acetate and the
20 peptide is eluted by washing the resin with 5% (v/v) acetic acid in water, and lyophilized. A white powder is obtained, generally amorphous.

b) The Fmoc/tBu strategy

- Differently from the previous strategy, the amino
25 group protector Fmoc is used, which is removed from the amino acid by treatment with a 20% piperidine solution in DMF for about 20 min. This treatment deprotonates the amino groups for the next coupling step. The coupling methods are the same as the Boc chemistry's and the
30 final cleavage is performed using a weaker acid solution because both the protector's bond to the side chains (generally of the tert-butylic type) and the peptide's bond to the resin are more labile than the Boc chemistry's. Usually, the resin is treated with 85-95%

TFA (v/v) and a mixture of different suppressors of collateral reactions for about 2 h. After this treatment, the peptide is precipitated with the resin in ice-cold ethylic ether, and centrifuged at 2,500g for 5 min. The supernatant is discarded and the pellet is resuspended in ethylic ether, vortexed and centrifuged. The procedure is repeated 5 times and the resin and peptide mixture is dried under vacuum. In the end, the peptide is released from the resin with 5% AcOH in water (v/v) and filtered. The filtered solution is lyophilized until an amorphous white powder is obtained.

- 1- Atherton, E. & Sheppard, R.C. (1989), Solid Phase Peptide Synthesis: A Practical Approach, I.L.R. Press, Oxford.
- 2- Fields, G.B. & Noble, R.L. (1990) Int. J. Pep. Prot. Res. 35, 161.

Example 4:

Peptide sequences deduced from the cDNA encoding the precursors of the BPPs.

Northern blot assays

Total RNA from several tissues of *Bothrops jararaca* was isolated using the extraction method with guanidine isothiocyanate - phenol - chloroform. Ten micrograms of total RNA from each of the serpent's tissues were separated by denaturing agarose gel electrophoresis (1.7% formaldehyde), and transferred by capillarity to nylon membranes. The RNA was blotted to the membrane at 80°C in a vacuum-oven for 1 hour followed by UV treatment in a Crosslinker (model RPN 2500 - Amersham), at 70.000 $\mu\text{J}/\text{cm}^2$ for 5 minutes.

The membranes were pre-hybridized at 42°C, over night (approximately 16 hours), in a buffer containing 50% formamide, 2.5 mM K_2PO_4 , 5 X Denhardt's solution, 50 $\mu\text{g}/\text{mL}$ herring sperm DNA and 10% dextran sulfate.

The cDNA fragments used as templates for the synthesis of radioactive probes were obtained by digestion of the clone containing the cDNA insert encoding the BPPs and CNP (clone NM87, Murayama *et al.*, Proc. Natl. Acad. Sci. USA, 5 94, 1189-1193, 1997), with the appropriate restriction enzymes (Sma I or Sma I + Bam HI, for the sequences encoding the BPPs and the CNP, respectively).

The excised inserts were separated by low melting agarose gel electrophoresis and the DNA was recovered by 10 phenol/chloroform extraction followed by ethanol precipitation.

The radioactive probes were prepared using approximately 25 ng of template DNA, obtained as described above, for each reaction, following the instructions of the 15 manufacturer of the RediPrime kit (Amersham), based on the incorporation of α [³²P]-dCTP in the presence of random hexanucleotides and polymerase (Klenow or T4 DNA polymerase). The non-incorporated radioactive nucleotides were separated from the probe in inverse molecular sieve 20 columns ("spin columns S-200" Pharmacia).

Hybridizations with the radioactive probes were carried out at 42°C, for 16 hours, after adding the radioactive probe to the pre-hybridization solution at a concentration of approximately 1.5×10^6 cpm/mL.

25 Finally, the membranes were washed at high stringency conditions, i.e., washed four times at 65°C with a 2X SSC; 0.1% SDS solution for 15 minutes each, and three times at 65°C in a 0.1X SSC; 0.1% SDS solution. The membranes were exposed to X-rays films (LS - Kodak) in appropriate 30 cassettes, for the appropriate time, depending on the strength of the radioactive signal observed for each membrane. For a better detection of the weaker signals, the cassettes, contained a pair of screens, the film and the membrane, and were kept at - 80°C.

Purification of the messenger RNA and preparation of the cDNA library from *B. jararaca* brain

Total RNA from the brain of *B. jararaca* was isolated using the guanidine isothiocyanate - phenol - chloroform extraction method.

The messenger RNA was purified by passing the total RNA solution twice through a pre-packed oligo-dT cellulose column (GibcoBRL). One aliquot of the purified messenger RNA was submitted to denaturing agarose gel electrophoresis (containing 1.7% formaldehyde) followed by staining with ethidium bromide. The RNA was transferred by capillarity to nylon membranes to confirm the integrity of the samples through hybridizations with probes specific for the sequences encoding the BPPs. The preparation of the cDNA library in λ ZAP phages was carried out using a cDNA preparation kit from Stratagene (La Jolla, CA). After cloning the double stranded cDNA inserts, obtained from 5 μ g of brain messenger RNA into the λ ZAP phages, these were packaged *in vitro* and titered to verify the cloning efficiency. This phage library was amplified and the aliquots were stored at -20°C and -80°C , in the presence of chloroform and DMSO, respectively.

Amplification of the cDNA from the total phage lysate

Using 10 μ l of the total phage lysate obtained for the brain cDNA library, PCR (Polymerase Chain Reaction) amplification was carried out using specific oligonucleotides, derived from the sequence of the cDNA encoding the precursor of the BPPs and the CNP from the venom gland. The phage lysate was initially incubated at 100°C for 5 minutes, then cooled down to 4°C and, finally, added to the other components of the amplification reaction, composed of 10 pmol of each primer, 2.5 U of Taq polymerase (Amersham), 200 μM dNTPs and 1X PCR buffer,

supplied with the enzyme. The reactions thus prepared were then submitted to a denaturing cycle at 94°C for 4°C and, then, to 35 more cycles of 1 min at 94°C, 1 min at 60°C, 1 min at 72°C. Subsequently, the reactions were kept at 4°C
5 until the PCR products were analyzed by agarose gel electrophoresis, after staining with ethidium bromide. The PCR products of the expected size, based on the cDNA sequence from the venom gland, were subcloned in a plasmid vector pCRscript SK+ (Stratagene), following the
10 manufacturer's instructions (by usual methods), and sequenced using oligonucleotides that anneals to adjacent regions of the vector's multiple cloning site (primers T3 and T7).

15 Selection and identification of the clone encoding the precursor of the BPPs:

The fragments obtained by digestion of clone NM87 with Sma I or Sma I and Bam HI (approximately 450 bp and 432 bp, positions 164 - 610 and 610 - 1044, respectively) were used as templates for the synthesis of radioactive probes ($\alpha^{32}\text{P}$)
20 using the random primer method (Rediprime kit/Amersham). The nitrocellulose membranes (Schleicher & Schuell) prepared from plates containing approximately 50 thousand phage plaques each, were submitted to hybridization with the radioactive probes in 6X SSPE (1X SSPE: 0.15 M NaCl, 15
25 mM NaH_2PO_4 , pH 7, 1 mM EDTA), 50% formamide, 0.1% SDS and 5X Denhardt's at 42°C for 16 hours. The membranes were then washed twice in 2X SSC/0.1% SDS and three times in 0.1X SSC/0.1% SDS, at 65°C, for 15 minutes each (1X SSC: 0.15 M NaCl, 15 mM sodium citrate, pH 7).

30 Positive phage plaques were identified by autoradiography and were isolated for the analysis of the DNA insert. *In vivo* excision of the phagemid pBluescript from the vector λ ZAP was carried out using the "helper phage", following the

instructions of the manufacturer, and the extremities of the insert were sequenced.

Sequencing of the DNA inserts:

The sequencing reactions were carried out following the chain termination method with dideoxy-nucleotides, using the kit Big Dye (Perkin Elmer), followed by the analysis in the ABI 310 automatic sequencer, following the manufacturer's instructions.

Total RNA from the brain of a single *B. jararaca* specimen was isolated by the guanidine isothiocyanate - phenol - chloroform extraction method, and the messenger RNA was purified by passing the total RNA solution once through a pre-packed oligo-dT cellulose column. The integrity of the purified mRNA of the sample was checked by agarose gel electrophoresis followed by hybridization with radioactive samples synthesized from de cDNA encoding the precursor of the BPPs and the CNP (identified by Northern blot assays).

Preparation of the cDNA library in λ ZAP phages (Stratagene) was carried out by common and usual methods using 5 μ g of this sample, following the manufacturer's instructions.

The initial packaging of the recombinant phages obtained generated a titer of approximately 2×10^5 pfu/mL, with less than 1% of non-recombinant clones. This library was immediately amplified, with a final titer of approximately 9×10^9 pfu/mL.

PCR (Polymerase Chain reaction) amplification

The goal of the initial approach used was to amplify the cDNA of interest from *B. jararaca* brain cDNA library by PCR, using oligonucleotides specific for the cDNA sequence encoding the precursor of the BPPs and the CNP. The reactions were carried out with materials and methods described herein. Only one fragment of approximately 250

base pairs was cloned from the total phage lysate containing the brain cDNA library (there was no amplification of any other fragment from the serpent brain or pancreas library). Complete sequencing of this cDNA insert showed a very high degree of similarity with the cDNA sequence identified in the venom gland library, except for the insertion of a few nucleotides (12 bases) in the segment that precedes the natriuretic peptide and some point mutations (4 residues substituted), of which three are silent (encoding the same amino acid residue) and the fourth was "conservative" (a substitution for an amino acid with the same chemical characteristics). In spite of the numerous amplification attempts changing reaction conditions and/or the polymerase and oligonucleotide used, it was not possible to clone other fragments from any of the libraries studied.

Approximately 5×10^6 clones were independently analyzed using the segment encoding the natriuretic peptide as a template for the synthesis of radioactive probes used for the hybridization experiments, which made it possible to identify 13 positive clones. The plasmid vectors containing the cDNA inserts were recovered by mini-preparations of DNA and sequenced using oligonucleotides that anneal to adjacent regions of the plasmid vector's (pBluescript SK+) multiple cloning site (commercial primers T3 and T7).

Screening of clones by hybridization of the cDNA library from the brain of *B. jararaca* allowed the isolation of a cDNA encoding the BPPs as part of a precursor protein for these peptides (figure 1). Seven peptides (the sequences of which are listed among those in table 1) were identified by analogy with the BPPs isolated from the venom of *Bothrops jararaca*. Based on these sequences, peptides were synthesized that presented the same pharmacological

properties as the venom peptides, as well as inhibition of ECA and NEP.

Example 5:

**Testing the bradykinin potentiating activity of the
5 BPPs and EVASINS:**

Biological assay on guinea pig ileum:

The potentiating activity of the pools of isolated peptides was tested on the contractile activity of bradykinin on smooth muscle, measured using a preparation
10 of isolated guinea pig ileum. Female guinea pigs with a body weight between 160 and 180 g were used. Before starting the assays, the ileum was kept in TYRODE solution at 37°C for 30 min. One of the ends of the isolated ileum segment, measuring 1.5 to 2.0 cm, was tied to a semi-ring
15 located at the bottom of a glass container with a capacity of 5 mL, containing the TYRODE physiological solution at 37°C under constant oxygen bubbling through a capillary. The other extremity of the ileum segment was tied to a previously calibrated registering arm. The tension on the
20 arm was of 1 g, and the ileum contractions were recorded by an REC101 recorder (Pharmacia Biotech).

The samples were prepared in deionized water at the time of use, and the volume used for the biological preparation did not exceed 0.2 mL.

25 In order to determine the bradykinin potentiating effect of the samples, a log-dose curve of the effect of bradykinin on the ileum was plotted. Bradykinin activity was determined by measuring the contractions of the isolated guinea pig ileum, and the potentiating activity
30 was expressed as a function of the increase in tissue response to a standard dose of bradykinin.

Pools and fractions were assayed individually and were added 30 seconds before the addition of a simple dose of bradykinin. The measured response was interpolated in the

linear section of the log dose-effect curve, giving the potentiating activity as a function of the raise in tissue response to a standard dose of bradykinin. TYRODE solution: 20 mL of stock solution I, 40 mL of the solution II, 1 mL of difenidramine solution (1 mg/mL), 1 mL of atropine solution (1 mg/mL), 5.60 mM D-glucose and H₂O to 1 L.

All reagents used in this experiment were of analytic grade.

Example 6:

10 **Effect on the systemic blood circulation and microcirculation**

The hypotensive potentiating activity effect of bradykinin was tested in normotensive and hypertensive male and female Wistar rats, anesthetized with ethylic ether; a cannulae was introduced in the femoral vein for drug administration and in the femoral artery for blood pressure recording. A Gould polygraph was used coupled to a Statham Gould physiological transducer. The arterial pressure variation values were obtained by integration of the areas delimited by the pressure base line, and by comparing them with the values obtained in the control experiment. Toxin effect tests were carried out in normotensive and hypertensive rats by intra-vital microscopy using the mesenteric circulation. The animals were anesthetized, and their body temperature maintained by a heated plate, while the mesentery was trans-illuminated and observed through lenses coupled to a television camera. The vase diameters of these animals were measured with a micrometric screw. Tested substances were administered by a constant flow peristaltic pump or injected *in bolus*.

25 **Example 7**

Enzymatic assays for the determination of ACE inhibition by the BPPs.

The wild-type human angiotensin converting enzyme (ACE) and two mutants, containing only one functional active site, were obtained through stable transfection of chinese hamster ovary cells with the ACE encoding cDNA. The
5 two ACE mutants were expressed as full-length proteins, presenting one of the catalytic sites, N- or C- terminal, inactivated by substitution of the zinc binding histidine residues by lysine residues.

The enzymatic assays were carried out at 25°C with the
10 substrate Mca-Ala-Ser-Asp-Lys-DpaOH, in a 50 mM Hepes buffer, pH 6.8, containing 200 mM NaCl and 10 μ M ZnCl₂. The reactions were continuously monitored by determining the raise in fluorescence at 390 nm emission wave length, and excitation at 340 nm, caused by the cleavage of the
15 substrate ($S = K_m$, 40 μ M) by ACE, in a fluorimeter Fluorolite 10000 (Dynatech). The bradykinin potentiating peptides (BPPs) (inhibitors, were pre-incubated with the enzyme 90 min before the addition of the substrate.

Example 8:

20 **Characterization of the effect of the BPPs *in vivo* - arterial blood pressure and microcirculation**

Wistar male rats with a body weight between 180 and 220 g were anesthetized with sodic pentobarbital (Hypnol® Cristalia, 50 mg/kg of body weight, intraperitoneally),
25 while the body temperature was maintained between 36.5°C and 37°C by a heated plate. After tracheotomy, the ventilation of the animals was controlled by a mechanical ventilator (Harvard Rodent Ventilator, model 683, Harvard Apparatus, South Natick, Mass, USA), under standard
30 conditions, as follows: frequency: 57-65 inspirations/min; passing volume: 2-2.5 mL; inspired oxygen fraction: 0.25-0.40. Cannulae were introduced in the right carotid artery and the jugular vein (PE-50 catheters, 58 mm ID, Portex, Hythe, UK) for the continuous monitoring of the arterial

blood pressure (TSE System Technical & CHM Scientific Equipment GMBH), and for the administration of liquids, respectively. The mesenteric bed was exposed. After trichotomy of the abdominal region, the skin was ruptured
5 and the mesentery was reached through an incision at the alba line. The mesentery was exposed under controlled temperature on the transparent area of the plate, which was placed on the microscope charriot. The temperature of the plate was kept at 37°C. The exposed tissue was kept moist
10 and heated by perfusion with Ringer-Locke solution, pH 7.2-7.4 containing 1% gelatin (154 mM NaCl; 5.6 mM KCl; 2 mM CaCl₂·2H₂O; 6 mM NaHCO₃; 5 mM glucose) at 37°C.

The study of mesenteric microcirculation *in situ* was carried out using an optical microscope (Akioskop-Carl
15 Zeiss, Germany) coupled to a colored image projection camera (JVC-TKC 6000), which transmits, simultaneously or not, the images obtained in the microscope to a computer and/or to a TV monitor. Image analysis software, MS Windows compatible, (Kontron KS 300, Kontron Bild Analyse-GMBH,
20 Germany), is installed in the computer allowing quantitative measurements of the stored images at fixed time intervals. A video tape recorder capable of recording the entire course of the experiment is connected to the TV monitor.

25 Thus, the images visualized during the experiment can be quantitatively evaluated through the fixed image on the computer's monitor, and at the same time the course of the experiment observed on the TV monitor can be stored by the video tape recorder for later analysis.

30 The optical microscope has a system of amplifying lenses (Optovar), localized between the lenses and the projection camera. The image transmitted to the monitors results from magnifications determined by the lenses, amplifying lenses and projection camera. In the studies

described here, magnification on the computer monitor was of 860 times and on the TV monitor, of 1530 times.

Determination of the red blood cells speed in the microcirculatory vessels (diameters between 15 and 30 μm) was carried out by the photometric method described by BORDERS & GRANGER (1984) and complemented by Davis (1987). Through an optical doppler velocimeter (Microcirculation Research Institute, Texas A&M College of Medicine), installed between the eye piece of the optical microscope and the image projection camera of the intravital microscopy equipment, the blood flow was determined *in situ* on the computer monitor. The equipment is composed of two sensors (Planar photodiodes and low capacitance), which are placed at the center of the image of the vessel under observation on the computer monitor. The sensors, at an established and known distance between them, are capable of detecting alterations in the light intensity produced by the passage of red blood cells and transform them into voltage signals, which are proportional to the speed of those cells.

The sensors determine the diameter of the vessel and the speed of the red blood cells at the center of the blood flow.

Experimental protocols

25 Surgical procedures

- 10 minutes for stabilization of the microcirculatory net and blood pressure.
- Administration of BPPs or equivalent volume of sterile saline solution (5 mL in 10 minutes). Immediately after administration of BPPs or saline solution, injection of bradykinin *in bolus* (3 $\mu\text{g}/300 \mu\text{l}$).

CLAIMS

1- Procedure for the isolation and purification of vasopeptidase peptide inhibitors, showing specificity for the carboxyl site of the angiotensin-converting enzyme, secreted by serpent venom glands (BPPs), particularly *BOTHROPS JARARACA*, or produced endogenously (EVASINS), having vasodilating and anti-hypertensive action, consisting of the following steps:

- a) 800-1500 mg of total venom, obtained from a pool of *B. jararaca* venom are dissolved in 7.0-15 ml deionized water followed by centrifugation at 1500-2000 rpm for 15 to 20 minutes; the supernatant is passed through a gel-filtration column, Sephadex G-25 (25-80 μ , Sigma), 1.2 x 101 cm; the column is equilibrated with 30-50 mM ammonium-acetate buffer, pH 5.0 to pH 6.0, at room temperature;
- b) the sample is to the top of the column and the components are eluted at a flow rate of 1.0 to 2.0 mL/min;
- c) the absorbance profile is obtained for each aliquot at 214 nm in order to define the constitution of the pools
- d) the components of the pools showing potentiating activity are partially purified by high performance liquid chromatography, for which 0.1 to 1.0 % TFA (trifluoro acid)/ H₂O and acetonitrile/H₂O (9:1) are used as solvents, and a gradient of 5% to 60% of solvent B, with a flow rate of 0.2 to 0.5 mL/min.
- e) the molecular mass and the primary structure of the bradykinin potentiating peptides are determined by mass spectrometry (ESMS-MS)

2- Procedure for the isolation and purification of vasopeptidase peptide inhibitors, showing specificity for the carboxyl site of the angiotensin-converting enzyme, secreted by serpent venom glands (BPPs), particularly *BOTHROPS JARARACA*, or produced endogenously (EVASINS),

having vasodilating and anti-hypertensive action, according to claim 1, characterized by the fact that in step (d) the HPLC Merck-Hitachi, model L-6200A is used, with the UV-vis detector set at 214 nm, and the reverse-phase column C-18/Beckman (5 μ , 4.6 x 250 mm). The gradients varied according to the sample.

3 - Procedure for the amino acid sequence determination of the BPPs, characterized by the fact that it was performed by mass spectrometry analysis, by dissolving the samples in 50% H₂O/ACN with 0.1% of formic acid and injecting them under constant flow of 5 μ L/min by means of an injection pump. The data are acquired at the first quadrupole (EM-MS), scanning the mass/charge ratio (m/z) between 400 and 1600, using a scanning time of 5 seconds during the whole analysis process. The sequencing is performed by selecting the peptide displaying a protonated ion, as characterized at the first quadrupole, followed by fragmentation by collision induced dissociation (CID) under 3×10^{-3} argonium gas pressure, and data acquisition by scanning the second quadrupole (ESMS-MS).

4- Procedure for the amino acid sequence determination of the BPPs, according to claim 3, characterized by the fact that it is performed in a mass spectrometer Micromass Quattro II (ES-MS-MS/Micromass) in the positive ionization mode with the electrospray ion source (Micromass) and the software Mass Lynx (Micromass) is used for data acquisition.

5 - Procedure for the amino acid sequence determination of the BPPs, characterized by the fact that the characterization and the sequencing can be performed by the HPLC system, coupled to a mass spectrometry system (LC-

ESMS-MS), where the Hewlett-Packard HPLC, model 1100 is used, equipped with an automatic injector, a UV/vis detector set at 214 nm. The reverse-phase column is a C-18/The Separations Group (5 μ m, 4.6 x 250 mm) for the
5 purification of total venom components.

6 - Procedure for the amino acid sequence determination of the EVASINS by deduction using the cDNA of the precursors of these molecules expressed in serpent tissues, specifically of *Bothrops jararaca*, consisting of the
10 following steps:

a) Isolation of total RNA from various *B. jararaca* tissues employing the isothiocyanide guanidine-phenol-chloroform extraction method; 1.0 to 5.0 micrograms of total RNA of each serpent tissue are submitted to denaturing agarose
15 eletrophoresis (1.0 to 2.5% formaldehyde), and transferred by capillarity to nylon membranes; the RNA is blotted to the membrane in a vacuum-oven at 80 °C for 1-2 hours.

b) the membrane is exposed to UV light at 70.000 J/cm² in a Crosslinker (RPN 2500 model - Amersham) for 30 to 50
20 minutes.

c) pre-hybridization is performed at 42°C overnight, or for 16 hours, in a 50% formamide solution, containing 2.5 mM K₂PO₄, pH 7.4, 5X Denhardt's solution, 50 μ g/ml herring sperm DNA, and 10% dextran sulfate

25 d) cDNA fragments, used as template strands for the synthesis of the radioactive probes, re obtained by digestion of the clone containing the cDNA insert coding for the BPPs and CNP precursor, clone NM87, with the restriction enzymes SmaI, and SmaI + BamHI, for the coding
30 sequences of EVASINS and CNP, respectively

e) the inserts are obtained by gel electrophoresis, using low melting point agarose,

- f) followed by extraction with phenol and chloroform, and ethanol precipitation.
- g) radioactive probes are prepared using approximately 50-100 ng of template DNA in a reaction based on the incorporation of α [³²P]-dCTP in the presence of random hexanucleotides and polymerase (Klenow or T4 DNA polymerase).
- h) spin-columns S-200 from Pharmacia are used to separate the probes from non-incorporated radioactive-labeled nucleotides
- i) pre-hybridizations are performed at 65°C, for 30 to 45 minutes using the radioactive labeled cDNA probes,
- j) and the radioactive labeled probes are added to the pre-hybridization medium at a concentration of approximately 1.5×10^6 cpm/ml.
- k) membranes are washed at high stringency conditions (4 washes at 65°C in 2 X SSC/0.1% SDS for 15 min each, and 3 washes at 65°C in 0.1 X SSC/0.1% SDS for 10 min each;
- l) membranes are exposed to X-ray films in appropriate cassettes and for the time needed, depending on the observed intensity of the radioactive signal of the membrane;
- m) the films are kept at -80°C.

7 - Procedure for the amino acid sequence determination of the EVASINS by deduction, using the cDNA of the precursors of these molecules expressed in serpent brain tissues, specifically of *Bothrops jararaca*, consisting of the following steps:

- a) Total RNA from *B. jararaca* brain is isolated using the guanidin isothiocyanate-phenol-chloroform extraction method,
- b) messenger RNA is purified by passing total RNA solution twice through a pre-packed oligo-dT cellulose column (Gibco/BRL).

- c) One aliquot of the purified messenger RNA is submitted to denaturing agarose gel electrophoresis (1.0-2.5% formaldehyde) and stained with ethidium bromide
- d) the RNA is transferred by capillarity to nylon membranes to confirm its integrity by hybridizations with probes specific for the region coding for the BPPs;
- 5 e) the cDNA library is constructed in λ ZAP phages using common techniques;
- f) the double stranded cDNA inserts, obtained from 5 μ g of brain messenger RNA are cloned into λ ZAP phages;
- 10 g) which are packaged *in vitro*; the titer is determined to verify the cloning efficiency;
- h) the phage library is amplified and aliquots are kept at -20°C and -80°C , in the presence of chloroform and DMSO, respectively.
- 15

8 - Procedure for the amino acid sequence determination of the EVASINS by deduction of the cDNA of the precursors of these molecules expressed in serpent brain tissues, specifically of *Bothrops jararaca*, consisting of the following steps:

20

- a) total phage lysate (10 μ l) from the brain cDNA library was employed for PCR (Polymerase Chain Reaction) amplification, using specific oligonucleotides, designed according to the cDNA sequence coding for the precursor of the BPPs and CNP from the venom gland. Incubation is at 42°C for 15 minutes, followed by cooling at 4°C ;
- 25 b) the other components are added to the amplification reaction, consisting of 10 pmol of each primer, 2.5 U of Taq polymerase (Amersham), 200 μM dNTPs and 1X PCR buffer, supplied with the enzyme;
- 30 c) reactions mixes, thus prepared, are submitted to a denaturing cycle at 94°C for 4°C and, subsequently, to 35

additional cycles of 1 min at 94°C, 1 min at 60°C, and 1 min at 72°C;

d) the reaction mixes are kept at 4°C until the PCR products are analysed by agarose gel electrophoresis, and
5 stained with ethidium bromide;

e) the PCR products are subcloned into the pCRscript SK+ (Stratagene) plasmid vector by usual methods, and are sequenced using oligonucleotides which anneal to sites adjacent to the vector's multiple cloning site (primers T3
10 and T7).

9 - Procedure for the amino acid sequence determination of the BPPs by deduction of the cDNA of the precursors of these molecules expressed in serpent brain tissues, specifically of *Bothrops jararaca*, consisting of the
15 following steps:

a) Nitrocellulose membranes prepared from plates containing approximately 50 thousand phage plaques each, are submitted to hybridization with radioactive probes in 6X SSPE (1X SSPE: 0.15 M NaCl, 15 mM NaH₂PO₄, pH 7, 1mM EDTA), 50%
20 formamide, 0.1% SDS and 5X Denhardt's at 42°C for 16 hours;

b) The membranes are washed twice in 2X SSC/0.1% SDS and three times in 0.1X SSC/0.1% SDS, at 65°C, for 15 minutes each wash (1X SSC: 0.15 M NaCl, 15 mM sodium citrate, pH
7);

25 c) After autoradiography phage plaques identified by the probes are isolated for the analysis of the DNA insert;

d) In order to analyse the inserts, their ends are sequenced after *in vivo* excision of the phagemid pBluescript from the vector λ ZAP, performed using the
30 "helper phage".

10 - Procedure according to claim 9, characterized by the fact that fragments obtained by digestion of clone NM87 with SmaI or SmaI and BamHI, approximately 450 bp and 432 bp, positions 164-610 and 610-1044, respectively, are used
5 for the synthesis of radioactive probes (α ³² P) by the random-priming method (Rediprime kit/Amersham).

11 - Procedure to determine the amino acid sequence of the BPPs by deducing from the cDNA of these precursor molecules, expressed in serpent brain tissue, specifically
10 *Bothrops jararaca*, characterized by the fact that they comprise the following steps:

a) Total RNA is isolated from the brain of a sole *B. jararaca* serpent, employing the isothiocyanide guanidine-phenol-chloroform extraction method, and the messenger RNA
15 is purified by one passage through the ready-packed oligo dT-cellulose column;

b) the integrity of the messenger RNAs of the preparation is verified by agarose gel eletrophoresis, followed by hybridization with the radiolabeled probes, synthesized
20 according to the cDNA sequence coding for the BPPs and CNP precursor;

c) a cDNA library is constructed by common techniques in λ ZAP phages (Stratagene), using 5 μ g of this preparation,
d) the library is amplified, and the final titer
25 determined, which should be around 9×10^9 pfu/ml.

12 - Procedure according to claim 11, characterized by the fact of obtaining sequenced DNA inserts.

13 - Procedure of amplifying the cDNA from serpent brain cDNA libraries, specially *Bothrops jararaca*, characterized
30 by the following steps:

- a) oligonucleotides specific for the cDNA sequence coding for the BPPs and CNP precursor are used;
- b) a fragment of approximately 250 base pairs is cloned from the total phage lysate of the serpent brain cDNA library;
- 5 c) complete sequence of this cDNA insert is determined;
- d) a high level of similarity is found to the cDNA sequence identified in the venom gland library;
- e) approximately 4×10^6 independent clones are analysed, employing a coding segment of the natriuretic peptide as a template for the synthesis of radioactive probes used in hybridizations and identification of the plasmid vectors containing the cDNA inserts for the DNA mini-preps, and the sequencing of the inserts, using
- 10 oligonucleotides (primers) which anneal to the regions adjacent to the multiple cloning site of the plasmid (pBluescript SK+) (primers T3 and T7).
- 15

14 - Procedure according to claim 13 characterized by the **fact that** the selection of clones through hybridization of the cDNA library of the *B. jararaca* brain allows the

20 isolation of a cDNA coding for the BPPs precursor within the precursor protein of these bioactive peptides.

15 - Procedure according to claim 14 characterized by the **fact that** by analogy to the EVASINS with the BPPs isolated from the venom of *B. jararaca*, six (6) peptide formulas are

25 identified.

16 - Procedure according to claim 14 characterized by the fact that the six (6) peptide formulas are:

Formulas	Sequences
I	pp ¹ aa ¹ aa ² aa ³ P ⁴ aa ⁵ aa ⁶ P ⁷ P ⁸
5 II	pp ¹ aa ¹ aa ² aa ³ aa ⁴ P ⁵ aa ⁶ aa ⁷ P ⁸ P ⁹
III	pp ¹ aa ¹ aa ² aa ³ aa ⁴ aa ⁵ P ⁶ aa ⁷ aa ⁸ P ⁹ P ¹⁰
IV	pp ¹ aa ¹ aa ² aa ³ aa ⁴ aa ⁵ aa ⁶ P ⁷ aa ⁸ aa ⁹ P ¹⁰ P ¹¹
V	pp ¹ aa ¹ aa ² aa ³ aa ⁴ aa ⁵ aa ⁶ aa ⁷ P ⁸ aa ⁹ aa ¹⁰ P ¹¹ P ¹²
VI	pp ¹ aa ¹ aa ² aa ³ aa ⁴ aa ⁵ aa ⁶ aa ⁷ aa ⁸ P ⁹ aa ¹⁰ aa ¹¹ P ¹² P ¹³

10 where:

P is always proline. The remaining are always L-amino acids, and are presented by the one-letter code (see bellow)*.

15 pp¹ is the N-terminus, and may be pyroglutamyl (<E) or a simple amino acid, usually not basic;

aa¹ is a non-basic amino acid, usually W, S, G or N

aa² is a non-acid amino acid, usually P, G, W or R;

aa³ is a non-acid amino acid, usually P, A, R or W

20 aa⁴ is P for formula I, and is an amino acid, usually T, P, G, H, R, W or E for the remaining oligopeptides;

aa⁵ is usually Q, N, P, or G for formula I;

aa⁶ for formula II, aa⁷ for formula III, aa⁸ for formula IV, and aa⁹ for formula V is a non-basic amino acid, usually Q, N, P, or G.

25 aa⁶ for formula I, aa⁷ for formula II, aa⁸ for formula III, aa⁹ for formula IV, and aa¹⁰ for formula V are always I or A or T.

*

acid amino acids D, E

30 basic amino acids K, R

Aromatic amino acids F, W, Y

G-glycine	N-asparagine
A-alanine	Q-glutamine
5 P-proline	D- aspartic acid
V-valine	E- glutamic acid
I-isoleucine	K-lysine
L-leucine	R-arginine
S-serine	F-phenylalanine
10 T-threonine	H-histidine
W-tryptophane	Y-tyrosine
<E- pyroglutamyl	

17 - Solid-phase synthesis procedure of vasopeptidase peptide inhibitors exhibiting vasodilating and anti-
15 **hypertensive action, consisting of the following steps:**

- a) the Boc group from the first amino acid residue bound to resin is removed in the presence of 30% TFA in dichloromethane (DCM), containing 2% anisol, during 30 min.
- b) a wash with 1/2% anisol, DCM and MeOH follows. For the
20 coupling of the next amino acid,
- c) the amino group is neutralized by treatment with 10% TEA or 5% DIPEA in DCM for 10 min;
- d) the peptidyl-resin is washed with DCM, MeOH, DCM and the solvent that will be used in the coupling phase;
- 25 e) the amino acid is coupled by activation using acylating agents based on diisopropylcarbodiimide (DIC) or 2- (1H-benzotriazolyl)-1,1,3,3-tetramethylurone tetrafluorborate (TBTU), and a 3-fold molar excess of the Boc-amino acid

(carboxylic component, CC) and of the coupling agent, as compared to the amount of amino groups (amine component, CA) in the resin; the coupling reaction is monitored for 2 hours, using the ninidrine quantitative method;

5 f) the following cycle is initiated by deprotecting the amino acid until the forthcoming residue enters the reaction, and the process continues in cycles until the end of the elongation of the desired sequence;

10 g) the peptide is cleaved from the resin, and its side chains are deprotected by treatment with anhydrous HF at 0° C for 60-90 min in the presence of p-cresol and dimethylsulfide (DMS) (5%, v/v of each) which act as suppressors for collateral reactions induced by carbocations released during the HF treatment;

15 h) the resin is washed with ethyl acetate and the peptide is extracted with 5% acetic acid (AcOH) in water (v/v), followed by lyophilization in order to obtain a white powder, which is usually amorphous.

18- Solid-phase synthesis procedure for the vasopeptidase
20 peptide inhibitors exhibiting vasodilating and anti-hypertensive action, according to claim 17, characterized by the fact that in step (e) the proportion of the reagents CA:CC:DIC be 1:3:3, when DIC is used, and 1:3:3:4 of CA:CC:TBTU:DIEA, when TBTU is used. The final concentration
25 of these reagents lies between 0.05 and 0.1 M.

19 - Solid-phase synthesis procedure for the vasopeptidase
peptide inhibitors exhibiting vasodilating and anti-hypertensive action, according to claim 17, characterized
by the fact that in step (e) hydroxybenzotriazol (HOBT) is
30 used together with the acylating compound in the coupling of Asn and Gln, in order to prevent the formation of nitril

of these amino acids. The final proportion of CA:CC:TBTU:HOBt:DIEA is 1:3:3:3:5.

20 - Solid-phase synthesis procedure for the vasopeptidase peptide inhibitors exhibiting vasodilating and anti-hypertensive action, according to claim 17, characterized by the fact that in step (g) ethane-1,2-dithiol (EDT) is added, such that the removal of the formyl group, which protects the side chain of this amino acid, occurs simultaneously to the cleavage in sequences containing Trp residues.

21- Solid-phase synthesis procedure for the vasopeptidase peptide inhibitors exhibiting vasodilating and anti-hypertensive action, consisting of the following steps:

- a) the aminic protection group Fmoc is removed from the amino acid, by treatment with 20% piperidine solution in DMF during approximately 20 minutes, thereby deprotonating the amino groups;
- b) the amino acid is couple by activation with acylating agents containing diisopropylcarbodiimide (DIC) or 2-(1H-benzotriazolyl)-1,1,3,3-tetramethylurone tetrafluorborate (TBTU), with a 3 molar excess of the Boc amino acid and the acylating agent, as compared to the amount of amino groups at the resin; the coupling reaction followed during 2 hours using the ninidrine qualitative method;
- c) cleavage is performed in a medium, which is weaker acidic than HF, using 85-95% TFA (v/v), and a mixture of different suppressor types of collateral reactions is added for approximately 2 hours;
- d) the peptide is precipitated in ice-cold ethyl-ether together with the resin and centrifuged at 8000 rpm for 5 minutes. The supernatant is discarded;

- e) the residue is resuspended in ethyl-ether, vortexed and centrifuged, and the procedure are repeated 5 times; a wash and a precipitation follows;
- f) the mixture of resin and peptide is dried in vacuum, and
- 5 g) the peptide is extracted from the resin with 5% AcOH in water v/v), and
- h) the filtrate is lyophilized to obtain an amorphous white powder.

22 - Solid-phase synthesis procedure of the vasopeptidase peptide inhibitors exhibiting vasodilation and anti-hypertensive action, according to claim 21, characterized by the fact that in step (c) the cleavage is performed in a weaker acidic medium than HF, because the binding of the protectors of the side chains (usually of the tert-butylic type), as well as the binding of the peptide to the resin, is weaker than the one observed in the Boc chemistry.

23 - VASOPEPTIDASES PEPTIDE INHIBITORS with anti-hypertensive and vasodilating action, characterized by the following formula:

20 Group (a) showing no specificity for the C site of ACE:

<E K W A P

Group (b) showing specificity for the C site of ACE:

- I pp¹aa¹aa²aa³P⁴aa⁵aa⁶P⁷P⁸
- II pp¹aa¹aa²aa³aa⁴P⁵aa⁶aa⁷P⁸P⁹
- 25 III pp¹aa¹aa²aa³aa⁴aa⁵P⁶aa⁷aa⁸P⁹P¹⁰
- IV pp¹aa¹aa²aa³aa⁴aa⁵aa⁶P⁷aa⁸aa⁹P¹⁰P¹¹
- V pp¹aa¹aa²aa³aa⁴aa⁵aa⁶aa⁷P⁸aa⁹aa¹⁰P¹¹P¹²
- VI pp¹aa¹aa²aa³aa⁴aa⁵aa⁶aa⁷aa⁸P⁹aa¹⁰aa¹¹P¹²P¹³

where:

- 30 P is always proline. The remaining are always L-amino acids and are presented in the one-letter code (see below*).
- pp¹ is the N-terminus, and may be pyroglutamyl (<E) or a simple amino acid, usually not basic;

aa¹ is a non-basic amino acid, usually W, S, G or N
 aa² is a non-acid amino acid, usually P, G, W or R;
 aa³ is a non-acid amino acid, usually P, A, R or W
 aa⁴ is P for formula I, and is an amino acid, usually T,
 5 P, G, H, R, W or E for the remaining oligopeptides;
 aa⁵ is usually Q, N, P, or G for formula I;
 aa⁶ for formula II, aa⁷ for formula III, aa⁸ for formula IV,
 and aa⁹ for formula V is a non-basic amino acid, usually Q,
 N, P, or G.
 10 aa⁶ for formula I, aa⁷ for formula II, aa⁸ for formula III,
 aa⁹ for formula IV, and aa¹⁰ for formula V are always I or A
 or T.

G-glycine	N-asparagine
A-alanine	Q-glutamine
15 P-proline	D- aspartic acid
V-valine	E- glutamic acid
I-isoleucine	K-lysine
L-leucine	R-arginine
S-serine	F-phenylalanine
20 T-threonin	H-histidine
W-tryptophane	Y-tyrosine

<E- pyroglutamyl


In which:

- The peptides described in the above Table have higher
 25 selectivity for the C-site of ACE, which is characterized
 by the K_i values between 2 nM and 100 μM, while the values
 where above 50 μM for the N-site of the ACE; these peptides
 are also inhibitors of NEP, showing K_i values between 5-150
 μM; all of them potentiate the contractile action of
 30 bradykinin on isolated guinea-pig ileum, doubling the
 contractile effect of bradykinin, when used in
 concentrations varying from 5-300 nM; the hypotensive
 effect of bradykinin was also potentiated by 30 to 80%,
 when concentrations between 10 and 700 nM were used,
 35 amplifying the duration of the hypotension between 5 and 15

fold, when compared to the duration of the effect of bradykinin alone.

24 - VASOPEPTIDASE PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action, according to claim 23, characterized by the fact that D and E are acidic amino acids, K and R are basic amino acids, and F, W, and Y are aromatic amino acids.

25 - VASOPEPTIDASE PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action, according to claim 23 (group b), characterized by the fact that they are peptides of 8 - 13 amino acids presenting a general formula, which contains the sequence motif at the carboxyl-terminus of the oligopeptide:



where X^1 can be any amino acid and X^2 usually is I, and the N-terminal amino acid is blocked, usually by <E.

26 - VASOPEPTIDASE PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action, according to claims 23 and 25, characterized by the fact that they show higher selectivity for the C-site of ACE [group (b)], which are characterized by K_i values between 2 nM a 100 μ M, while the values for the N-site of ACE are above 50 μ M.

27 - VASOPEPTIDASE PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action, according to claims 23 and 25, characterized by the fact that they are also inhibitors of NEP, showing K_i values between 5-150 μ M.

28- VASOPEPTIDASE PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action, according to claims 23 and 25, characterized by the fact that they potentiate the contractile action of bradykinin on isolated guinea-pig ileum, doubling the contractile effect of bradykinin, in concentrations between 5-300 nM.

29- VASOPEPTIDASE PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action, according to claims 23 and 25, characterized by the fact that they potentiate the hypotensive effect of bradykinin by 30-80% in concentrations between 10-700nM, amplifying the duration of this hypotension by 5- to 15-fold when compared to the duration of the effect by bradykinin alone.

30 - VASOPEPTIDASE PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action, according to claims 23 and 25, characterized by the fact that they can be conjugated to a vast variety of ligands: acid groups like sulfonyl, carboxyl, phosphoryl, or other groups like thiols and olefines, dithio, azo or diazo compounds, aldehyds or alike.

31- VASOPEPTIDASE PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action, according to claim 30, characterized by the fact that they are used in systemic cardiovascular disorders, in general or localized, where microcirculation is the affected target.

32 - VASOPEPTIDASE PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action, according to claim 31, characterized by the fact that they are used in human hypertensive disorders and their consequences.

33 - VASOPEPTIDASES PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action, according to claim 32, characterized by the fact that they are used in vascular lesions and hyperplasias.

5 34 - VASOPEPTIDASE PEPTIDE INHIBITORS with vasodilation and anti-hypertensive action in obtaining pharmaceutical compounds characterized by the fact that diluted in physiologically acceptable medium, like phosphate-saline buffer, saline solution or other more convenient carriers.

10 35 - Utilization of VASOPEPTIDASE PEPTIDE INHIBITORS *in vitro* and *in vivo*.

15 36 - Utilization of the VASOPEPTIDASE PEPTIDE INHIBITORS *in vivo*, via trans-mucosa, parenterally, or by injection, particularly intravenously for systemic action or localized action in tissue microcirculation.

37 - Utilization of the VASOPEPTIDASE PEPTIDE INHIBITORS *in vivo*, in dosages varying between approximately 1 μ g to 10 mg, usually 0.5 mg/kg of body weight.

专利名称(译)	血管肽酶抑制剂的分离和纯化方法		
公开(公告)号	EP1587819A2	公开(公告)日	2005-10-26
申请号	EP2002704503	申请日	2002-03-18
[标]申请(专利权)人(译)	生物实验萨纽斯药物有限公司 圣保罗州研究奖励基金会		
申请(专利权)人(译)	BIOLAB SANUS FARMACEUTICA LTDA FUNDACAO DE AMPARO—一个PESQUISA DO斯卡德圣保罗 - FAPESP		
当前申请(专利权)人(译)	BIOLAB SANUS FARMACEUTICA LTDA FUNDACAO DE AMPARO—一个PESQUISA DO斯卡德圣保罗 - FAPESP		
[标]发明人	MARTINS DE CAMARGO ANTONIO CARLOS CALHETA VIERIA PORTARO FERNANDA FERRAGINI MURBACK A CENT TOXICOLOGIA APLICADA ALVES IANZER D CENT DE TOXICOLOGIA APLICADA POLISELLI FARSKY S H CENT TOXICOLOGIA APLICA PALMA MARIO SERGIO CENT TOXICOLOGIA APLICADA MIRIAN AKEMI FURUIE HAYASHI CENT DE TOXICOLOGIA		
发明人	MARTINS DE CAMARGO, ANTÔNIO CARLOS CALHETA VIERIA PORTARO, FERNANDA FERRAGINI MURBACK, A,CENTRO.TOXICOLOGIA APLICADA ALVES IANZER, D.,CENTRO DE TOXICOLOGIA APLICADA POLISELLI FARSKY, S, H,CENTRO TOXICOLOGIA APLICA. PALMA, MARIO SERGIO,CENTRO TOXICOLOGIA APLICADA MIRIAN,AKEMI,FURUIE,HAYASHI,CENTRO DE TOXICOLOGIA		
IPC分类号	G01N27/62 A61K38/00 A61P9/08 A61P9/12 A61P43/00 B01J20/281 C07K1/10 C07K7/06 C07K7/08 C12N9/64 C12N9/99 C12N15/09 C12Q1/68 G01N27/447 G01N30/34 G01N30/46 G01N30/72 G01N30 /88 G01N33/53 G01N33/566 G01N33/60 C07K1/00		
CPC分类号	C12N9/6418 A61K38/00		
优先权	PI0101088 2001-03-19 BR		
外部链接	Espacenet		

摘要(译)

本发明专利涉及蛇毒腺分泌的肽的分离和纯化，特别是两种药物；由此获得的肽，以及原核和真核系统中通过基因工程技术的生产程序；由此获得的工程化肽；通过化学合成制备所述肽，以及由该化学处理产生的肽。它还指在不同的药物组合物中利用通过不同方法获得的所述肽，并通过各种方式引入生物体中，以使它们充当血管肽酶的抑制剂，从而降低全身动脉血压，并显示局部血管扩张作用。