



US007658911B2

(12) **United States Patent**
Dambinova

(10) **Patent No.:** **US 7,658,911 B2**
(45) **Date of Patent:** ***Feb. 9, 2010**

(54) **ANTIBODY ASSAY METHODS TO ASSESS RISK FOR TIA/STROKE**

(75) Inventor: **Svetlana A. Dambinova**, Atlanta, GA (US)

(73) Assignee: **CIS Biotech, Inc.**, Atlanta, GA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 109 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **11/339,452**

(22) Filed: **Jan. 25, 2006**

(65) **Prior Publication Data**

US 2006/0172342 A1 Aug. 3, 2006

Related U.S. Application Data

(63) Continuation of application No. 11/076,074, filed on Mar. 9, 2005, which is a continuation of application No. 09/922,011, filed on Aug. 2, 2001, now Pat. No. 6,896,872.

(60) Provisional application No. 60/301,297, filed on Jun. 27, 2001.

(51) **Int. Cl.**
A61K 49/00 (2006.01)
A61K 39/00 (2006.01)
G01N 33/53 (2006.01)
G01N 33/00 (2006.01)

(52) **U.S. Cl.** **424/9.1; 424/139.1; 435/7.1; 435/7.95**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,849,895	A	12/1998	Daggett et al.
6,316,611	B1	11/2001	Daggett et al.
6,896,872	B2	5/2005	Dambinova
2005/0181466	A1	8/2005	Dambinova et al.
2006/0024749	A1	2/2006	Dambinova

FOREIGN PATENT DOCUMENTS

WO	WO 02/12892	A2	2/2002
WO	WO 2006/081188	A1	8/2006
WO	WO 2006/135893	A2	12/2006

OTHER PUBLICATIONS

Dambinova et al. Journal of Stroke and Cerebrovascular Disease, 10: 192 [abstract 194 from The 2001 North American Stroke Meeting].*
Izykenova et al. Journal of Stroke and Cerebrovascular Disease, 10: 192 [abstract 195 from The 2001 North American Stroke Meeting].*
Greenland et al. Archives of Internal Medicine, 165:2454-2456, Nov. 2005.*
Oei et al. Circulation, 111: 570-575, 2005.*
DeGiorgio et al., Nature Medicine, 7(11):1189-1193, 2001.*

Castillo, J., et al., "Neuroexcitatory Amino Acids and Their Relation to Infarct Size and Neurological Deficit in Ischemic Stroke," *Stroke*, 27(6):1060-1065 (Jun. 1996).

Castillo, J., et al., "Progression of ischaemic stroke and excitotoxic aminoacids," *Lancet*, 349(9045):79-83 (Jan. 11, 1997).

Dambinova, S.A., et al., "Autoantibodies to Subtypes of Glutamate Receptors as a Hallmarks of Brain Damage: Diagnostic Significance for Paroxysmal Activity and Ischemia," *J. Higher Nervous Activity*, 47(2):151-156 (1997) (transl. of: *Zh. Vyssh. Nerv. Deiat. Im. I. P. Pavlova*, 47(2):439-446 (Mar.-Apr. 1997).

Dambinova, S.A., et al., "Blood test detecting autoantibodies to N-methyl-D-aspartate neuroreceptors for evaluation of patients with transient ischemic attack and stroke," *Clinical Chemistry*, 49(10):1752-1762 (2003).

Dambinova, S.A., et al., "Monitoring of Brain Spiking Activity and Autoantibodies to N-Terminus Domain of GluR1 Subunit of AMPA Receptors in Blood Serum of Rats with Cobalt-Induced Epilepsy," *J. Neurochem.*, 72(3):2088-2093 (Nov. 1998).

Dambinova, S.A., et al., "Multiple panel of biomarkers for TIA/stroke evaluation," *Stroke*, 33(5):1181-1182 (May 2002).

Dambinova, S.A., et al., "The presence of autoantibodies to N-terminus domain of GluR1 subunit of AMPA receptor in the blood serum of patients with epilepsy," *J. Neurological Sciences*, 152(1): 93-97 (Nov. 6, 1997).

Davalos, A., et al., "Duration of Glutamate Release After Acute Ischemic Stroke," *Stroke*, 28(4): 708-710 (Apr. 1997).

Davalos, A., et al., "Neurological Deterioration in Acute Ischemic Stroke: Potential Predictors and Associated Factors in the European Cooperative Acute Stroke Study (ECASS) 1," *Stroke*, 30(12): 2631-2636 (Dec. 1999).

Gahring, L.C., et al., "Autoantibodies to Neuronal Glutamate Receptors in Patients with Paraneoplastic Neurodegenerative Syndrome Enhance Receptor Activation," *Molec. Medicine*, 1(3): 245-253 (Mar. 1995).

Gray, J.J., et al., "Evaluation of a commercial latex agglutination test for detecting antibodies to cytomegalovirus in organ donors and transplant recipients," *J. Virol. Meth.*, 16(1-2): 13-19 (May 1987).

Gusev, E.L., et al., "Investigation of the Level of Autoantibodies to Glutamate Receptors in Blood Serum of Patients in Acute Period of Ischemic Stroke," *Zhurnal Nevrologii I Psikiatrii Imeni S. S. Korsakova*; 96(5):68-72 (1996) (in Russian; Abstract in English).

(Continued)

Primary Examiner—John D. Ulm
Assistant Examiner—Stacey MacFarlane
(74) *Attorney, Agent, or Firm*—Clark G. Sullivan; Arnall Golden Gregory LLP

(57) **ABSTRACT**

A methods, kits and compositions for diagnosing a central nervous system disorder, particularly transient ischemic attack or stroke, comprising measuring the level of NR2A and/or NR2B NMDA receptor or fragment thereof, in a biological sample from a human subject, and optionally measuring other biomarkers such as homocysteine and glutamate. The method is particularly useful for identifying individuals that are at risk for stroke, and for diagnosing stroke in an emergency room setting.

5 Claims, No Drawings

OTHER PUBLICATIONS

- Gusev, E.L., et al., "Neuroprotective Effects of Glycine for Therapy of Acute Ischaemic Stroke," *Cerebrovasc. Dis.*, 10(1):49-60 (Jan.-Feb. 2000).
- Hess, S.D., et al.; "Cloning and Functional Characterization of Human Heteromeric N-Methyl-D-Aspartate Receptors," *J. Pharmacol. Exp. Ther.*, 278(2): 808-816 (Aug. 1996); XP000676640; Abstract.
- Hill, M.D., et al., "Biochemical markers in acute ischemic stroke," *Canadian Medical Association Journal*; 162:1139-1140 (Apr. 18, 2000).
- Lipton, S.A., et al., "Neurotoxicity associated with dual actions of homocysteine at the N-methyl-D-aspartate receptor," *Proc. Natl. Acad. Sci. USA*, 94(11):5923-5928 (May 27, 1997).
- Meldrum, B.S., "Glutamate as a Neurotransmitter in the Brain: Review of Physiology and Pathology," *J. Nutrition*, 130:1007S-1015S (2000).
- Monyer, H., et al., "Heteromeric NMDA Receptors: Molecular and Functional Distinction of Subtypes," *Science*, 256(5060):1217-1221; (May 22, 1992).
- Pang, S.C., et al., "Early diagnosis of acute myocardial infarction by myoglobin latex agglutination test," *Jpn. Heart J.*, 29(5):631-638 (Sep. 1988).
- Rogers, S.W., "Autoantibodies to Glutamate Receptor GluR3 in Rasmussen's Encephalitis," *Science*, 265(5172):648-651 (Jul. 29, 1994).
- Senju, O., et al., "A new immuno quantitative method by latex agglutination—application for the determination of serum C-reactive protein (CRP) and its clinical significance," *J. Clin. Lab. Immunol.*, 19(2): 99-103 (Feb. 1986).
- Skoromets, A.A., et al.; [Autoantibodies to NMDA-type glutamate receptors in the blood of patients with acute ischemic and hemorrhagic stroke] *Zh. Nevrol. Psikiatr. Im. S. S. Korsakova [J. Neurological Sciences]*, 97(6):53-58 (1997) (In Russian, Abstract in English).
- Vila, N., et al., "Plasma homocysteine levels in patients with ischemic cerebral infarction," *Med. Clin. (Barcelona)* 110(16): 605-608 (May 9, 1998). (In Spanish) Provided as abstract in English.
- Vincent, A., et al., "Pathogenic autoantibodies to neuronal proteins in neurological disorders," *J. Neuroimmunology*, 100(1-2):169-180 (Dec. 1999).
- Yamamoto, T., et al., "CSF and ECF Glutamate Concentrations in Head Injured Patients," *Acta Neurochir. [Suppl]* 75:17-19 (1999).
- U.S. Appl. No. 11/076,974, Dambinova.
- U.S. Appl. No. 11/339,440, Dambinova.
- U.S. Appl. No. 11/338,447, Dambinova.
- U.S. Appl. No. 11/451,679, Dambinova.
- Dambinova, et al., "Autoantibodies to Subtypes of Glutamate Receptors as a Hallmarks of Brain Damage: Diagnostic Significance for Paroxysmal Activity and Ischemia." *Journal of Higher Nervous Activity*. 1997. vol. 47 (2), pp. 151-156.
- Gusev, et al., "Neuroprotective Effects of Glycine for Therapy of Acute Ischaemic Stroke." *Cerebrovasc Dis* 2000; 10:49-60.
- Dambinova, et al., "Monitoring of Brain Spiking Activity and Autoantibodies to N-Terminus Domain of GluR1 Subunit of AMPA Receptors in Blood Serum of Rats with Cobalt-Induced Epilepsy." *Journal of Neurochemistry*, Lippincott-Raven Publishers, Philadelphia.
- Dambinova, et al., "The presence of autoantibodies to N-terminus domain of GluR1 subunit of AMPA receptor in the blood serum of patients with epilepsy." *Journal of Neurological Sciences*, 152 (1997) 93-97.
- Meldrum, B.S., "Glutamate as a Neurotransmitter in the Brain: Review of Physiology and Pathology¹." *Journal of Nutrition*. 2000; 130:1007S-1015S.
- Monyer et al., "Heteromeric NMDA Receptors . . ." *Science* May 22, 1992; 256:1217-1221.
- Dambinova et al., "Blood Test Detecting Autoantibodies . . ." *Clinical Chemistry* 2003 49(10); 1181-1182.

* cited by examiner

ANTIBODY ASSAY METHODS TO ASSESS RISK FOR TIA/STROKE

RELATION TO PRIOR APPLICATIONS

This application is a continuation of and claims priority to U.S. Utility application Ser. No. 11/076,074, filed Mar. 3, 2005 (currently pending), which is a continuation of U.S. Utility application Ser. No. 09/922,011, filed Aug. 2, 2001 (now issued as U.S. Pat. No. 6,896,872), which claims priority to U.S. Provisional Application No. 60/301,297, filed Jun. 7, 2001, and to U.S. Utility application Ser. No. 09/632,749, filed Aug. 4, 2000, of which U.S. Ser. No. 09/922,011 was a continuation-in-part.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates generally to the diagnosis, management and therapy of central nervous system disorders such as stroke, transient ischemic attack, and traumatic brain injury. In particular, the invention relates to methods and kits for evaluating these central nervous system disorders, in order to better respond to episodes of focal cerebral ischemia, and to best manage the risk associated with future acute incidences.

2. Background Information

Stroke or "brain attack" is clinically defined as a rapidly developing syndrome of vascular origin that manifests itself in focal loss of cerebral function. In more severe situations, the loss of cerebral function is global. A stroke occurs when the blood supply to the part of the brain is suddenly interrupted (ischemic) or when a blood vessel in the brain bursts, spilling blood into the spaces surrounding the brain cells (hemorrhagic). The symptoms of stroke are easy to spot: sudden numbness or weakness, especially on one side of the body; sudden confusion or trouble speaking or understanding speech; sudden trouble seeing in one or both eyes; sudden trouble walking; dizziness; or loss of balance or coordination. (National Institute of Neurological Disorders and Stroke, 2001). Stroke is the most common devastating neurologic disease in the world, and the third leading cause of death in world after heart disease and cancer. Despite recent progress understanding stroke mechanisms, stroke management is still not optimal for a number of reasons.

The importance of promptly diagnosing a stroke after symptoms appear cannot be overstated. Delays in diagnosis and medical intervention beyond 3 hours after stroke onset may contribute to clinical deterioration and disability. An early diagnosis enables doctors to more effectively choose the emergency intervention such as anti-platelet or/and neuro-protective therapy, and also to make better prognoses of disease outcome. Successful treatment of stroke requires rapid state diagnosis. The delay in achieving an accurate and certain diagnosis wastes the limited amount of time available in which the brain can respond to reperfusion, and significantly increases the risk of hemorrhage after most of the permanent injury has occurred (Marler J. R. *Annl. Emergency Med.* 1999, 33: 450-451).

Unfortunately, however, many people who have a stroke either do not seek immediate medical care or suffer from delays in medical care even in countries where stroke care is advanced, such as the United States and Europe (Alberts M J, Hademenos G, Latchaw R E, et al. *JAMA* 2000; 23:3102-3109). Several clinical criteria can be employed to diagnose whether a patient is having a stroke, but even all these criteria do not always allow one to differentiate the episode from other disorders, such as epilepsy, syncope, and migraine

(Toole J.F. *Cerebrovascular Disorders*. 1999. Lippincott, Williams & Wilkins, New York, 5th Ed., 542 p). Moreover, progressing stroke is only partially predictable based on clinical and neuroimaging data that is currently available to neurologists.

Transient ischemic attack (TIA) is a short-lived episode of focal neurologic deficit which often precedes the cerebral infarction of a stroke. It occurs when the blood supply to part of the brain is briefly interrupted, and is typically accompanied by permanent brain damage (albeit less severe damage than normally results from a stroke). TIA symptoms, which usually occur suddenly, are similar to those of stroke but do not last as long. Most symptoms of a TIA disappear within an hour, although they may persist for up to 24 hours. Symptoms can include: numbness or weakness in the face, arm, or leg, especially on one side of the body; confusion or difficulty in talking or understanding speech; trouble seeing in one or both eyes; and difficulty with walking, dizziness, or loss of balance and coordination. (National Institute of Neurological Disorders and Stroke, 2001). Patients who have suffered a TIA have 9.5 times greater risk of having a future stroke than those who have not had a TIA, and about one third of patients who suffer a TIA will have an acute stroke in the future. (American Stroke Association, 2001). However, because the symptoms of TIA are short term, many patients do not recognize the event as a TIA or perceive the event as a warning of a potentially impending stroke.

Standard treatments to reduce the risk of future stroke include the use of antiplatelet agents, particularly aspirin. People with atrial fibrillation (irregular beating of the heart) may be prescribed anticoagulants. The most important treatable factors linked to TIAs and stroke are high blood pressure, cigarette smoking, heart disease, carotid artery disease, diabetes, and heavy use of alcohol. Lifestyle changes can often be implemented to reduce these factors. However, it is necessary to diagnose the TIA as a warning sign of impending stroke before such treatments can be administered. Therefore, a laboratory blood test to detect TIA or stroke, or the risk of suffering a TIA or stroke in the future, would be of tremendous benefit.

During the past 5 years a number of molecular and immunochemical assays have been evaluated for clinical use in neurology. (Schenone A. et.al. *Current Opinion in Neurology*. 1999, 12: 603-604; Honnorat J. J. *Neurol. Neurosurg. Psychiatry*. 1996, 61:270-278). At present, the Thrombogene V and two Thrombx tests are available for diagnosing stroke/thrombosis from Athena Diagnostic. These tests evaluate the frequent deep vein thrombosis and hypercoagulation states of patients to evaluate the need for intravenous anticoagulant therapy. The Thrombogene V test detects the Factor V Leiden mutation by Polymerase Chain Reaction (PCR) in the blood of patients. The other two tests monitor changes of different blood coagulation markers: antithrombin III protein C, factor IX, and anticardiolipin antibodies (IgG, IgM, IgA) by use of ELISA technique. These tests thus reveal the hypercoagulation state as a result of a thrombotic events, such as stroke.

Stroke can be related to different types of venous thromboembolisms, which are common disorders with considerable morbidity and potential for mortality (Anderson, D.; Wells, P. *Cur. Opinion in Hemat.* 2000, 7: 296-301). The biochemical marker: D-dimer, a breakdown product of a cross-linked fibrin blood clot that indicates the occurrence of plasmin mediated lysis of cross-linked fibrin, has been extensively evaluated for use in diagnostic tests for indicating acute venous thromboembolism. Indeed, a fully automated, semi-quantitative latex agglutination assays that uses turbimetric or

agglutination endpoints has been developed that provides results within 20 minutes with sensitivity between 89% and 95% (Roussi J.; Bentolila L.; Contribution of D-dimer determination in the exclusion of deep venous thrombosis in spinal cord injury patients. *Spinal Cord*, 1999; v.37: p. 548-552). Unfortunately, however, the presence of D-dimer may also be increased in other settings that result in fibrin generation, including recent surgery, hemorrhage, trauma, cancer, and pregnancy (Anderson D R., Wells P S.; *Thromb. Haemost.*; 1999; 82:878-886).

However, the foregoing tests do not elucidate upon the TIA/stroke mechanisms that are actually responsible for the damage associated with neurotoxic molecular events. It is necessary to find out specific and sensitive biomarkers which could be helpful to recognize initial brain damage and which could help to choose not only the appropriate anticoagulant treatment, but also emergency or regular neuroprotective therapy.

It is well known that two of the three leading causes of death, namely cardiovascular diseases and stroke, are the end result of atherosclerosis. Thus, it is not surprising that several biochemical markers implicated in thromboembolic processes are also reported to be associated with stroke and stroke risk. Among these are homocysteine, cholesterol and LDL (*Cerebrovascular Disorders* ed. by J. E. Toole. Lippincott Williams & Wilkins. 1999, pp. 34-35), which are also classified as risk factors to cardiovascular and cerebrovascular diseases. (Hankey G J., and Eikelboom J W. *Lancet*. 354: 407-413 (1999). Approximately one fourth of patients with symptomatic atherosclerosis have elevated plasma homocysteine levels caused by various factors. High levels of homocysteine may run in families with increased susceptibility to heart attack and stroke (Graham I. J. *Ir. Call. Phys. Surg.* 1995; 24: 25-30). Elevated plasma homocysteine may be a causal and modifiable risk-factor for ischemic stroke, but the results of previous studies have been conflicting (Deulofeu V N R, Chamorro A, Piera C. *Med Clin (Barc)*. 1998; 110: 605-608; Yamamoto T, Rossi S, Stiefel M F, Doppenberg E, Zauner A, Bullock R, Marmarou A. *Acta Neurochir. Suppl.* (Wien) 1999; 75:17-19).

The neurotoxic effect of excitatory amino acids (glutamate, aspartate) in the brain has also been well documented. The results of this work show a correlation between glutamate content in the blood and the severity of acute ischaemia (Castillo J, Dávalos A, Naveiro J, Noya M. *Stroke* 1996, 27:1060-1065; Castillo J, Dávalos A, Noya M. *Lancet*. 1997; 349:79-83). Cerebral damage and its association with progressing stroke has been attributed to increased glutamate release, or low glutamate reuptake, both in animals and in humans (Dávalos A, Castillo J, Serena J, Noya M. *Stroke* 1997; 28:708-710).

However, only 56% of patients with progressing stroke are reported to have high glutamate content in their blood serum (Dávalos A., Toni D., Iweins F., et al., 1999, 30: 2631-2636). Moreover, even though glutamate is considered the strongest biochemical predictor of progressing stroke (Dávalos A, and Castillo J. In Book: *Cerebrovascular Disease*. Current Med. Inc.: Philadelphia. 2000 Chapter 16, pp. 169-181), this marker remains non-specific for TIA. Glutamate changes have also been observed in the blood of patients with epilepsy and other nervous system disorders (Meldrum B S. J. *Nutrition*. 2000, 130:1007S-1015S).

Over the last three decades substantial progress has been made in elucidating the mechanisms by which cerebral ischemia leads to brain damage. The cellular and molecular mechanisms of cerebral ischemia abnormalities have been better defined through the role of glutamate and glutamate

receptors, one of the most distributed excitatory neuroreceptors in brain, in regulating of initial stages of brain damage. Indeed, numerous molecular investigators consider glutamate receptors to be one of the key biological receptors involved in the molecular mechanisms of TIA and stroke (Meldrum B S. J. *Nutrition*. 2000, 130:1007S-1015S). According to a leading hypothesis, ischemia-induced glutamate release activates these glutamate receptors. It has been shown that glutamate and homocysteine (the sulfinate analog of aspartate) activate the glutamate binding site of NMDA receptors and participate in neurotoxic processes (Lipton S. A., Kim W. K., Choi Y. B., Kumar S., et al. *PNAS*. 1997, 94: 5923-5928).

Glutamate receptors are divided into two main groups: ionotropic and metabotropic. The ionotropic neuroreceptors are ligand-gated ion channels that are subdivided into NMDA, AMPA and kainate receptor subtypes. There are four NR2 subunits: NR2A, NR2B, NR2C and NR2D, which is responsible for Ca^{2+} -permeability regulation. NMDA receptors can be modified by ischemia, resulting in changes of ion permeability and/or ion selectivity.

Recent research findings indicate that the blood of patients with CNS disorders other than TIA or stroke exhibit properties of autoimmunization to products of nerve cell degradation (Vincent A., Oliver L., Pallace J. *J Neuroimmun.* 1999; 100: 169-180). For example, a correlation between AMPA GluR1 autoantibodies and common epilepsy has been shown (Dambinova et al. *J. Neurol. Sci.* 1997; 152: 93-97; Dambinova et al. *J. Neurochem.* 1998; 71: 2088-2093), as has a correlation between AMPA GluR3 autoantibodies and Rasmussen's encephalitis (Rogers S W, Andrews P I, Gahring L C, et al. *Science*. 1995; 265:648-651; Twyman R E, Gahring L C, Spiess J, Rogers S W. *Neuron*. 1995; 14:755-762; Gahring L C, Twyman R I, Greenlee J E, Rogers S W. *Mol. Med.* 1995; 1:245-253).

In a similar vein, several researchers have reported an increase in NMDA receptor synthesis, the appearance of high levels of receptor antigen, and the generation of autoantibodies to the receptors during the initial stages of cerebral ischemia (Gusev et al., *J. Neurol. & Psych.* 1996, 5:68-72; Dambinova et al. *J. Neurol. Sci.* 1997, 152:93-97; Dambinova et al. *J. Neurochem.* 1998, 71: 2088-2093). Acting on this research, one company developed a laboratory kit (cerebral ischaemia (CIS)-test) that detects autoantibodies to the N-terminus domain of the NR2A subunit in the blood of patients with TIA or stroke (Gusev E. I., Skvortsova V I, Alekseev A A, Izykenova G A, Dambinova S A. S. *S Korsakov's J. Neurol. & Psych.* 1996; 5:68-72). The N-terminus domain of the NR2A subunit of NMDA receptors was selected as the immunoreactive epitope on the basis of molecular biological and experimental studies showing that this epitope is the most immunoreactive region of the receptor (Dambinova S A, Izykenova G A. *J. High Nervous Activity.* 1997; 47:439-446).

More recently, researchers have reported a correlation between the effectiveness of a stroke treatment regimen and the levels of autoantibodies to the NR2A and NR2B subunits of NMDA. In particular, these researchers have reported increased titers of autoantibodies to the NR2A and NR2B subunits of NMDA in the blood of patients severely affected by stroke, and a reduction of the autoantibodies, accompanied by an improvement in neurological function, during therapy by glycine—a non-specific agonist of NMDA receptors (Gusev et al. *Cerebrovascular Diseases*. 2000, 10: 49-60). Patients that responded positively to glycine had lower autoantibody titers than patients who were not treated, and had levels of autoantibodies that were close to the levels of autoantibodies in control subjects.

Unfortunately, the use of NR2A and NR2B autoantibodies in the diagnosis of stroke or TIA does not provide a real-time assessment of the damage being done by a stroke or TIA. Rather, because of the time the immune system requires to mount an immune response, and to generate NR2A and NR2B autoantibodies, methods that test for these antibodies at best provide a delayed assessment of the extent and severity of stroke or TIA.

Investigators from Canada (Hill M. D., Jackowski G., Bayer N., Lawrence M., Jaeschke R. *Can. Med. Assoc. J.* 2000, 163: 1139-1140) have proposed a new diagnostic laboratory assay for differentiating stroke subtype. They designed a preliminary prospective cohort study to test a panel of biochemical markers (neuron-specific enolase [NSE], myelin basic protein [MBP], S-100 [beta] protein and thrombomodulin [Tm]) in blood samples from patients with acute ischemic stroke. These markers were chosen because they cover important cellular components of the brain that might be damaged in acute stroke. The 4 biochemical markers were assayed using a standard ELISA technique.

The results of this investigation demonstrated elevated levels of NSE in 89% of the patients admitted in hospitals, Tm in 43%, MBP in 39% and S-100 [beta] in 32%. At least one of the markers was elevated on admission in 93% of the acute stroke patients. By stroke type, 100% of the patients with lacunar stroke, 100% of those with posterior circulation stroke and 90% of those with partial anterior circulation stroke had elevated NSE levels on admission. Conversely, none of the patients with lacunar stroke had an elevated S-100 [beta] level initially or subsequently. Peak levels of NSE, S-100 [beta] and MBP, but not of Tm, were significantly correlated with admission NIHSS scores ($p < 0.05$).

For stroke, 3 hours is an outside limit for administering appropriate therapies. The focus must change from extensive evaluation before any action to a well-planned acute emergency therapy developed using an appropriate diagnostic strategy. Every future advance to improve the outcome after TIA/stroke will depend on a fast initial response—within minutes and not hours (Marler J. R. *Ann. Emergency Med.* 1999, 33: 450-451). Therefore, it is especially important to develop a fast and simple method (within one hour) of detecting brain and blood biomarkers capable of recognizing the initial processes of TIA/stroke before irreparable ischemic damage ensues.

OBJECTS OF INVENTION

Therefore, it is an object of the invention to provide biochemical methods and kits for diagnosing central nervous system disorders such as TIA and stroke.

It is another object of the present invention to improve upon the accuracy of currently available methods for diagnosing TIA and stroke, and to more accurately diagnose TIA and stroke to the exclusion of other nervous system disorders or traumatic brain injury.

It is still another object of the present invention to provide methods of diagnosing stroke using biochemical markers that distinguish between hemorrhagic and ischemic stroke.

Still another object of the invention is to provide biochemical analyses of the extent and progression of TIA or stroke, or the infarction resulting from the TIA or stroke.

It is another object of the present invention to provide rapid biochemical methods and kits for diagnosing TIA and stroke, to provide real-time assessments of TIA or stroke, within a window of time that permits effective therapeutic intervention.

It is another object of the present invention to provide rapid and inexpensive biochemical methods and kits for diagnosing TIA and stroke, which can be performed at frequent intervals to monitor the progression of a TIA or stroke, or the effectiveness of therapy administered against TIA or stroke.

Still another object of the present invention is to provide diagnostic methods and kits for assessing the risk of incurring a TIA or stroke, and for monitoring the remission of risk factors for TIA or stroke.

Still another object of the invention is to provide a panel of rapid multiple panel of biomarkers for assessing the nature, severity and progression of TIA or stroke, and thereby to enable a more effective selection of intervention therapy.

SUMMARY OF THE INVENTION

It has unexpectedly been discovered that levels of circulating NMDA receptor proteins or fragments thereof can be assessed using diagnostic kits and processes, and that levels of these proteins or fragments can be used to clinically evaluate patients suffering from ischemic central nervous system disorders such as stroke or TIA. When analyzed in combination with other biomarkers for stroke and TIA, such as the thromboembolic marker homocysteine, or the excitatory amino acid glutamate, these proteins can diagnose the existence of a stroke with remarkable accuracy (generally greater than 89%). In contrast, the efficacy of single parameters for early diagnosis of stroke is 58% for glutamate, 66% for homocysteine, and 79% for NMDA receptors. The rapid evaluation of these neural ischemic biomarkers in an emergency room setting will greatly enhance the confidence of physicians when diagnosing stroke or TIA, and significantly improve the speed at which therapy against the stroke or TIA can be administered.

The biomarkers also yield extensive evidence about the nature of the stroke or TIA and the type therapy which should be administered. For example, the respective levels of biomarkers can be evaluated to determine whether the patient is suffering an ischemic or hemorrhagic stroke, or whether the patient is suffering from a traumatic brain injury. The data from the biomarkers can also be used to monitor or evaluate the progression of the ischemic episode, as well as the damage that has resulted as a consequence of the ischemia. High levels of all parameters reflect the neurological deficit and may be also used for prognosis of disease outcome. Moreover, a relationship has been observed between the respective levels of the biomarkers and the degree of thromboembolic and neurotoxicity in brain processes under the stroke. Once again, these relationships can be put to extensive use when evaluating the choice of emergency therapy in short time frames, such as anti-platelet and neuroprotective therapy. The data can be used independently of other diagnostic strategies, but preferably forms an integral part of a comprehensive diagnostic strategy employing conventional diagnostic techniques.

The data obtained from the NMDA biomarkers, especially when combined with data from other biomarkers such as glutamate and homocysteine, can also be used to monitor the efficacy of a treatment regime. It has surprisingly been found that the NMDA biomarkers provide real time evidence of neurotoxicity, and that reductions in levels of circulating NR1A or NR2A NMDA receptors or fragments thereof correspond well with reductions in neurotoxic mechanisms. By obtaining data at appropriate intervals using rapid laboratory techniques such as latex agglutination, one is able to monitor the progression of the episode in response to the therapeutic regime.

A latex agglutination technique has also been developed which dramatically increases the speed of diagnosis obtained by the methods of this invention, and thereby improves the effectiveness of the methods in emergency-room settings. The technique can be adapted for use in the detection of NMDA receptors, homocysteine, glutamate, or any other suitable biomarker against central nervous system disorders. Using the latex agglutination technique, one is able to provide real-time biochemical diagnosis and monitoring of TIA/stroke patients (within about 30 minutes), and thereby dramatically improve, the effectiveness of response to TIA/stroke. This is surprising because these biomarkers are naturally occurring and, in contrast to viruses for which latex agglutination methods were originally developed, show much lower strengths of association with their corresponding antibodies.

This semi-quantitative method gives reliable data quickly in a format that is simple for interpretation. Surprisingly, the technique shows greater accuracy than even well established methods based upon HPLC and ELISA. The application of the latex agglutination technique to the analysis of brain biomarkers for stroke will decrease the cost of analysis, provide the opportunity to monitor real-time progress of a treatment procedure, and allow physicians to determine the efficacy of medication administered in the treatment of TIA or stroke.

The methods of the present invention also can be employed in a non-emergency setting, when evaluating the risk that an individual will suffer a stroke or TIA. In addition, based upon results showing an increased risk of suffering TIA or stroke, prevention therapy can be administered, and the effectiveness of the therapy monitored using the methods of the present invention.

The invention also relates to indirect methods for measuring levels of NR2A and NR2B NMDA receptor proteins or fragments thereof. Thus, analytical techniques can be used to evaluate indirect measures of NR2A and NR2B NMDA receptor proteins or fragments thereof, such as autoantibodies specific for the proteins, or cDNA that encodes for the proteins.

Additional advantages of the invention will be set forth in part in the description which follows, and in part will be obvious from the description, or may be learned by practice of the invention. The advantages of the invention will be realized and attained by means of the elements and combinations particularly pointed out in the appended claims. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

DETAILED DESCRIPTION OF THE INVENTION

The present invention may be understood more readily by reference to the following detailed description of preferred embodiments of the invention and the Examples included therein. Before the present methods and techniques are disclosed and described, it is to be understood that this invention is not limited to specific analytical or synthetic methods as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

DEFINITIONS AND USE OF TERMS

As used in this specification and in the claims which follow, the singular forms "a," "an" and "the" include plural referents

unless the context clearly dictates otherwise. Thus, for example, reference to "a fragment" includes mixtures of fragments, reference to "an cDNA oligonucleotide" includes more than one oligonucleotide, and the like.

An analogue of a protein, peptide, or polypeptide means a protein, peptide, or polypeptide that contains one or more amino acid substitutions, deletions, additions, or rearrangements. For example, it is well known in the art of protein biochemistry that an amino acid belonging to a grouping of amino acids having a particular size or characteristic (such as charge, hydrophobicity, and hydrophilicity) can often be substituted for another amino acid without altering the activity of the protein, particularly in regions of the protein that are not directly associated with biological activity. Thus, an analogue of an NMDA receptor or fragment thereof is useful in the present invention if it includes amino acid substitutions, deletions, additions or rearrangements at sites such that antibodies raised against the analogue are still specific against the NMDA receptor or fragment.

Preferably, an NMDA analogue has at least 80%, 85%, 90%, or 95% amino acid identity with naturally occurring NMDA. Amino acid identity is defined by a analogy comparison between the analogue and naturally occurring NMDA. The two amino acid sequences are aligned in such a way that maximizes the number of amino acids in common along the length of their sequences; gaps in either or both sequences are permitted in making the alignment in order to maximize the number of common amino acids. The percentage amino acid identity is the higher of the following two numbers: (1) the number of amino acids that the two polypeptides have in common with the alignment, divided by the number of amino acids in the NMDA analogue or fragment thereof, multiplied by 100, or (2) the number of amino acids that the two polypeptides have in common with the alignment, divided by the number of amino acids in naturally occurring NMDA or fragment thereof, multiplied by 100.

NMDA derivatives, and derivatives of NMDA fragments, include naturally occurring NMDA and NMDA analogues and fragments thereof that are chemically or enzymatically derivatized at one or more constituent amino acids, including side chain modifications, backbone modifications, and N- and C-terminal modifications, by for example acetylation, hydroxylation, methylation, amidation, phosphorylation or glycosylation. The term also includes NMDA salts such as zinc NMDA and ammonium NMDA.

A protein or peptide is measured "directly" in the sense that the protein or peptide is itself measured in the biological sample, as opposed to some other indirect measure of the protein or peptide such as autoantibodies to the protein or peptide, or cDNA associated with the expression of the protein or peptide.

The term "antibody" is intended to be synonymous with "immunoglobulin." As used herein, the term "antibody" is meant to include both the native antibody, and biologically active derivatives of antibodies, such as, for example, Fab', F(ab')₂ or Fv as well as single-domain and single-chain antibodies. A biologically active derivative of an antibody retains the ability to bind antigen.

60 General Discussion

The present disclosure describes diagnostic and therapeutic applications that result from the realization that genetic or accidental increase of NMDA receptors synthesis in the brain reflects a neurological ischemic deficit, and may be used for early diagnoses of stroke or TIA. NMDA receptors that are abnormally expressed in the brain are quickly metabolized and, following penetration of the blood brain barrier, these

metabolic destruction products enter the circulatory system. The immune system recognizes these peptides and protein fragments as foreign antigens and responds by generating autoantibodies to them.

In one aspect of the present invention, the correlation between increased NMDA receptor synthesis, and the appearance of high levels of the receptors in blood sera of individuals during the initial stages of cerebral ischemia, is used for diagnostic and therapeutic applications. Experiments in rats with focal ischemia have demonstrated that NR2A mRNA expression in the cortex and hippocampus can be measured within two hours of the onset of the ischemic episode, and thus provide an opportunity for real time measurement of ischemic processes and damage resulting therefrom. At the same time, meaningful expression of NR2C and NR2D mRNA is not observed in brain structures that showed no changes in NR1 mRNA expression in rat ischemic brain. These changes in NR2-receptor mRNA expression in the early stages of ischemia are observed prior to morphological evidence of neuronal damage or appearance of autoantibodies to them in blood serum specimens.

Thus, in one aspect the present invention provides a method for diagnosing a central nervous system disorder comprising measuring the level of NR2A and/or NR2B NMDA receptor or fragment thereof in a biological sample. Elevated levels of NR2A and NR2B NMDA receptors are specific to brain injury, and are expressed in ischemic brain tissue at higher rates than other NMDA receptors, and thus are uniquely suited for assessing ischemic brain episodes such as TIA or stroke. Baseline levels for determining whether the measured levels are elevated, and hence indicative of a central nervous system disorder, can be obtained from population norms or, preferably, from a patient's own test history.

The biological sample tested for the receptor or fragment can be derived from blood, urine, blood plasma, blood serum, cerebrospinal fluid, saliva, perspiration, or brain tissue. In a preferred embodiment, the biological sample is a blood sample. In an even more preferred embodiment the biological sample is a blood sample diluted to a ratio of from about 1:2 to about 1:32 (v:v).

Immunoassay techniques are generally preferred for measuring the proteins or peptides of the present invention, as discussed in greater detail herein, although other analytical techniques are also available as known to those skilled in the art, such as HPLC. The amino acid sequences of the NR2A and NR2B subunits, and antigenic fragments thereof, are recited in SEQ ID NOS. 1, 2, 3, 10, 11, and 12, and any fragment of these sequences can be employed in methods for directly detecting the receptors as long as sufficient antigenicity is maintained. However, when using immunoassays it has been found that the antigenic determinants are concentrated in the N-terminal domain of the NR2A and/or NR2B NMDA receptor, and that antibodies raised against the N-terminal domains and fragments thereof should be employed for optimal test results. The inventors have sequenced the amino acid chain of the N-terminal domains for these receptors, and set forth the sequences as SEQ ID NOS. 2 and 11, respectively, at the end of this document.

In a preferred embodiment, other biomarkers of central nervous system disorders are also measured to improve the accuracy of the diagnosis, and to provide further information about the nature, severity, or progression of the disorder. Particularly useful markers are directly implicated in the NMDA receptor pathway, and include naturally occurring agonists and antagonists of the NMDA receptors. An exemplary antagonist is glycine. Exemplary agonists include glutamate, polyglutamate, aspartate, polyaspartate,

homocysteine, and polyhomocysteine. A particularly preferred agonist for measuring the activity of the receptors is glutamate or polyglutamate.

In another embodiment, thromboembolic biomarkers are measured to obtain a simultaneous reading of the likelihood for clotting in the brain. Exemplary thromboembolic biomarkers include homocysteine or polyhomocysteine.

Titers of higher than 2.63 for combined levels of NR2A and NR2B, especially when combined with titers higher than 3.34 for glutamate and/or 2.23 for homocysteine, are remarkably predictive of the occurrence of stroke and typically justify immediate therapeutic intervention for the TIA or stroke or risk of stroke. These titers can be translated into absolute concentrations by reference to the examples hereof.

The methods of the present invention are preferably performed by directly measuring the levels of NR2A and/or NR2B biomarkers in a selected biological sample, using immunoassay techniques employing antibodies raised against the biomarkers, or through quantitative techniques such as HPLC. However, it is also possible to measure the presence of the NR2A and/or NR2B biomarkers indirectly. This can be done by directly measuring autoantibodies of the biomarkers, or by directly measuring the cDNA nucleic acid intermediates involved in expression of these biomarkers. If autoantibodies are measured, they are preferably measured using one or more antigenic fragments of the NR2A and/or NR2B receptors as the target of the antibody, as opposed to a whole NR2A and/or NR2B protein. Healthy persons generally have NR2A autoantibodies in an amount of about 1.0-2.0 ng/ml. Healthy persons generally have NR2A cDNA levels of about 1.0-1.5 pg/ml.

Latex Agglutination and Other Diagnostic Techniques

A number of immunoassays can be employed in accordance with the principles of the present invention. Examples include radioimmunoassays, enzyme immunoassays (e.g. ELISA), immunofluorescence, immunoprecipitation, latex agglutination, hemagglutination, and histochemical tests. A particularly preferred method, however, because of its speed and ease of use, is latex agglutination.

Latex agglutination assays have been described in Beltz, G. A. et al., in *Molecular Probes: Techniques and Medical Applications*, A. Albertini et al., eds., Raven Press, New York, 1989, incorporated herein by reference. In the latex agglutination assay, antibody raised against a particular biomarker is immobilized on latex particles. A drop of the latex particles is added to an appropriate dilution of the serum to be tested and mixed by gentle rocking of the card. With samples lacking sufficient levels of the biomarkers, the latex particles remain in suspension and retain a smooth, milky appearance. However, if biomarkers reactive with the antibody are present, the latex particles clump into visibly detectable aggregates.

An agglutination assay can also be used to detect biomarkers wherein the corresponding antibody is immobilized on a suitable particle other than latex beads, for example, on gelatin, red blood cells, nylon, liposomes, gold particles, etc. The presence of antibodies in the assay causes agglutination, similar to that of a precipitation reaction, which can then be detected by such techniques as nephelometry, turbidity, infrared spectrometry, visual inspection, colorimetry, and the like.

The term latex agglutination is employed generically herein to refer to any method based upon the formation of detectable agglutination, and is not limited to the use of latex as the immunosorbent substrate. While preferred substrates for the agglutination are latex based, such as polystyrene and polypropylene, particularly polystyrene, other well-known substrates include beads formed from glass, paper, dextran,

and nylon. The immobilized antibodies, may be covalently, ionically, or physically bound to the solid-phase immunoadsorbent, by techniques such as covalent bonding via an amide or ester linkage, ionic attraction, or by adsorption. Those skilled in the art will know many other suitable carriers for binding antibodies, or will be able to ascertain such, using routine experimentation.

Thus, in one embodiment, the method of measuring the NR2A and/or NR2B NMDA receptor, fragment thereof, or other biomarker is by latex agglutination comprising:

(i) contacting the biological sample with poly- or monoclonal antibodies bound on an agglutinating carrier to target biomarkers for a sufficient time period and under conditions to promote agglutination; and

(ii) reading a signal generated from the agglutination; wherein the amount of signal detected correlates to the titer of biomarkers present in the sample.

The reaction is preferably read macroscopically against a dark background for a sufficient time period. The method preferably yields a clinically useful reading within about 30 minutes or less.

It has been experimentally found that latex beads having a mean diameter of from about 0.25 to about 0.4 μm are particularly preferred in the practice of this invention. The poly- or monoclonal antibodies are preferably present in a ratio with the latex beads of about 1:1.

Latex beads having the foregoing characteristics can be prepared generally by adding antibodies to the target biomarker to a carrier solution that contains a 1% concentration (by weight) of latex beads, until the concentration of the antibodies in the carrier solution reaches about 2 mg/ml, and allowing the ingredients a sufficient time to covalently link, typically about 1 hour, in the presence of a linking agent such as glutaraldehyde.

Conventional methods can be used to prepare the antibodies. For example, by using a peptide of a NMDA protein, polyclonal antisera or monoclonal antibodies can be made using standard methods. A mammal, (e.g., a mouse, hamster, or rabbit) can be immunized with an immunogenic form of the peptide (preferably the NR2A and/or NR2B receptor, an antigenic determinant of the NR2A and/or NR2B receptor, or an analogue or derivative thereof) which elicits an antibody response in the mammal. Techniques for conferring immunogenicity on a peptide include conjugation to carriers or other techniques well known in the art. For example, the peptide can be administered in the presence of adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassay procedures can be used with the immunogen as antigen to assess the levels of antibodies. Following immunization, antisera can be administered and, if desired, polyclonal antibodies isolated from the sera.

To produce monoclonal antibodies, antibody producing cells (lymphocytes) can be harvested from an immunized animal and fused with myeloma cells by standard somatic cell fusion procedures thus immortalizing these cells and yielding hybridoma cells. Such techniques are well known in the art, (e.g., the hybridoma technique originally developed by Kohler and Milstein (Nature 256, 495-497 (1975)) as well as other techniques such as the human B-cell hybridoma technique (Kozbor et al., Immunol. Today 4, 72 (1983)), the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al. Monoclonal Antibodies in Cancer Therapy (1985) Allen R. Bliss, Inc., pages 77-96), and screening of combinatorial antibody libraries (Huse et al., Science 246, 1275 (1989)). Hybridoma cells can be screened immunochemically for production of antibodies specifically reac-

tive with the peptide and the monoclonal antibodies can be isolated. Therefore, the invention also contemplates hybridoma cells secreting monoclonal antibodies with specificity for NR2A or NR2B NMDA proteins or fragments thereof as described herein.

When the NR2A and/or NR2B receptors are detected indirectly, by measuring the cDNA expression of the NR2A and/or NR2B receptors, the measuring step in the present invention may be carried out by traditional PCR assays such as cDNA hybridization, Northern blots, or Southern blots. These methods can be carried out using oligonucleotides encoding the polypeptide antigens of the invention. Therefore, in one embodiment the methods are performed employing oligonucleotides that encode the amino acid sequence of SEQ ID NO: 2, which is preferably represented by nucleotides 371-1978 of SEQ ID NO: 6. More preferably, the nucleic acid construct comprises a oligonucleotide consisting of a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 3, which is preferably represented by oligonucleotides 1790-1852 of SEQ ID NO: 7.

Thus, in one embodiment the methods of this invention include measuring an increase of NR2A and/or NR2B cDNA expression by contacting the total DNA isolated from a biological sample with oligonucleotide primers attached to a solid phase, for a sufficient time period. In another preferred embodiment, NR2A and/or NR2B cDNA expression is measured by contacting an array of total DNA bound to a solid matrix with a ready-to-use reagent mixture containing oligonucleotide primers for a sufficient time period. Expressed NR2A cDNA is revealed by the complexation of the cDNA with an indicator reagent that comprises a counterpart oligonucleotide to the cDNA attached to a signal-generating compound. The signal-generating compound is preferably selected from the group consisting of horseradish peroxidase, alkaline phosphatase, urinase and non-enzyme reagents. The signal-generating compound is most preferably a non-enzyme reagent.

In a preferred embodiment, the solid phase is a polymer matrix. More preferably, the polymer matrix is polyacrylate, polystyrene, or polypropylene. In one preferred embodiment the solid phase is a microplate. In another preferred embodiment, the solid phase is a nitrocellulose membrane or a charged nylon membrane.

As mentioned above, the methods of performing the present invention also may be performed by measuring the levels of autoantibodies specific for the NR2A and/or NR2B subunits. These autoantibodies may be measured by any suitable immunoassay such as, for example, a radioimmunoassay, an immunofluorescence assay, an enzyme-linked immunosorbent assay (ELISA), an immunocytochemical assay, and immunoblotting. In a preferred embodiment, the antigen to which the anti-NR2A and/or NR2B autoantibodies bind is a polypeptide or protein fragment of the N-terminal domain of the NR2A and/or NR2B receptor. More preferably, the antigen comprises a polypeptide or protein fragment of amino acid SEQ ID NO:2, 3, 4, 11, 12, or 13, or a suitable analogue or derivative thereof.

Thus, in yet another embodiment the methods of the present invention are performed by measuring the levels of anti-NR2A and/or anti-NR2B autoantibodies, by contacting a biological sample with a polypeptide or protein fragment of the NR2A and/or NR2B receptor (preferably the N-terminal domain) (or an analogue or derivative thereof) attached to a solid phase, for a sufficient time period and under conditions to allow a complex to form between any NR2A and/or NR2B autoantibodies which may be present in the sample and the polypeptide or protein fragment, contacting the complex with

an indicator reagent comprising a secondary antibody specific for the species of the mammal attached to a signal-generating compound (or for the polypeptide or protein fragment); and measuring the signal generated. The peptide can be obtained directly from biological samples, by recombinant DNA techniques, or by direct chemical synthesis. The signal-generating compound is preferably selected from horseradish peroxidase, alkaline phosphatase, and urinase. More preferably, the signal-generating compound is horseradish peroxidase. Most preferably, the indicator reagent is rabbit anti-human IgG attached to horseradish peroxidase. The amount of signal detected is correlated to the amount of anti-NR2A and/or NR2B autoantibodies present in the biological sample.

In this method it is preferred that the solid phase be a polymer matrix. More preferably, the polymer matrix is selected from the group consisting of polyacrylate, polystyrene, and polypropylene. In one preferred embodiment the solid phase is a microplate. In another preferred embodiment, the solid phase is a nitrocellulose membrane or a charged nylon membrane.

The immunosorbent of the present invention for measuring levels of autoantibody can be produced as follows. A fragment of the receptor protein is fixed, preferably by covalent bond or an ionic bond, on a suitable carrier such as polystyrene or nitrocellulose. If the standard polystyrene plate for immunological examinations is employed, it is first subjected to the nitration procedure, whereby free nitrogroups are formed on the plate surface, which are reduced to amino groups and activated with glutaric dialdehyde serving as a linker. Next the thus-activated plate is incubated with about 2 to 50 nM of the target peptide for the purpose of chemically fixing the respective immunogenic fragment of the receptor protein for a time and at a temperature sufficient to assure fixation (i.e. for about 16 hours at 4° C.).

The amount of protein below 2 nM affects adversely the reliability of the findings, whereas its amount exceeding 50 nM is inexpedient due to an increase in the nonspecific binding of autoantibodies with the immunosorbents. The plate is then washed with an aqueous solution of sodium boron hydride and an aqueous solution of sodium chloride, vacuum-dried, enclosed in a hermetically sealed package, and put under storage at 4° C.

It is also practicable to produce the immunosorbent by fixing the respective fragment of the receptor protein on nitrocellulose strips by virtue of ionic interaction. The respective fragment of the receptor protein isolated from the mammals' brain is applied to nitrocellulose and incubated for 15 min at 37° C. Then nitrocellulose is washed with a 0.5% solution of Tween-20, and the resultant immunosorbent is dried at room temperature and stored in dry place for one year period.

Emergency Room Diagnosis and Prognosis

As mentioned above, the methods of the present invention are especially well suited for use in emergency room settings. There are two reasons for this. First, the method is extremely useful in an emergency room setting because NR2A and NR2B NMDA receptor levels are elevated at a very early stage of ischemic insult, and thus provide a real time indication of neurotoxic events. This is in contrast to autoantibodies which require that an immune response first be mounted by the insulted organism.

The second reason the method is useful in an emergency room setting is the speed and ease with which the latex agglutination procedure can be employed. Using the latex agglutination processes described herein, one is able to turn laboratory results around often in less than 30 or even 20 minutes. Thus, using the methods of the present invention real-time

data can be obtained that permits a therapeutic response within the window for an effective response to stroke.

Therefore, in one embodiment the invention provides a method for diagnosing the existence of a central nervous system disorder such as TIA or stroke, further comprising withdrawing the biological sample from a human subject, wherein the biological sample is withdrawn within three hours of the onset of symptoms of the central nervous system disorder. In still another embodiment of the invention, the amount of time elapsed between withdrawing the biological sample from the subject, and detecting or measuring the presence or quantity of the NR2A and/or NR2B NMDA receptor, is less than about one hour, 45 minutes, or 30 minutes.

One of the principal advantages of the present invention is the ability to distinguish ischemic episodes such as stroke from other brain injuries such as traumatic brain injury. Thus, in another embodiment, the invention provides a method for diagnosing the existence of TIA or stroke further comprising evaluating from the level of NR2A and/or NR2B NMDA receptor whether the brain injury is a traumatic brain injury or stroke/TIA, and administering traumatic brain injury or stroke/TIA therapy as appropriate.

Another advantage of the methods of the present invention which is extremely useful in an emergency room setting, is the ability to determine from the test data the type of stroke involved. In particular, if a stroke is suspected, the method will help diagnose whether the stroke is an ischemic or hemorrhagic insult. Thus, in another embodiment the invention provides a method for diagnosing the existence of TIA or stroke further comprising, when the diagnosis confirms a stroke, evaluating from the level of NR2A and/or NR2B NMDA receptor whether the stroke is ischemic or hemorrhagic and administering ischemic or hemorrhagic stroke therapy as appropriate.

Another advantage of the present invention is the ability to evaluate infarction volume and extent of neurotoxicity from NMDA expression data NMDA receptor expression research in an animal model of middle carotid artery occlusion has been employed to demonstrate such correlation. Thus, in still another embodiment the invention provides a method for diagnosing the existence of TIA or stroke further comprising, if TIA and/or stroke is confirmed, evaluating from the level of NR2A and/or NR2B NMDA receptor cranial infarct volume, and administering therapy appropriate to the infarct volume.

Moreover, one can periodically repeat the procedure, to provide continuous monitoring of a patient's state as interventional therapy is administered, to monitor the effectiveness of a particular therapeutic regime. In this embodiment, it is preferable for the mammal to be concurrently undergoing treatment for the disorder. More preferably, the samples are collected at intervals from about 20 min to about 1 month. Even more preferably, the interval is from about 20 min. to about 2 hours. Most preferably the samples are collected at an interval of about 30 minutes. Thus, in still another embodiment the invention provides a method for diagnosing the progression of TIA or stroke further comprising detecting or measuring the presence or quantity of a NR2A and/or NR2B NMDA receptor in a biological sample one or more additional times, at a frequency of less than about 6 hours.

Primary Care Physician Setting

In another application the method is used in a clinical setting to determine an individual's risk of stroke, or to monitor the effectiveness of risk reduction therapies. As mentioned above, a number of therapies can be employed to reduce the risk of stroke in an individual. The use of antiplatelet agents,

particularly aspirin, is a standard treatment for patients at risk for stroke. People with atrial fibrillation (irregular beating of the heart) may be prescribed anticoagulants. The most important treatable factors linked to TIAs and stroke are high blood pressure, cigarette smoking, heart disease, carotid artery disease, diabetes, and heavy use of alcohol. Medical help is available to reduce and eliminate these factors. Lifestyle changes such as eating a balanced diet, maintaining healthy weight, exercising, and enrolling in smoking and alcohol cessation programs can also reduce these factors. When these therapies are administered it is desirable to determine the effectiveness of the therapy.

Therefore, in one embodiment the invention provides a method for evaluating an individual's risk for TIA or stroke comprising measuring levels of NR2A and/or NR2B NMDA receptors or fragments thereof in a biological sample from the individual, and comparing the levels to a baseline level. In one embodiment the baseline levels are derived from population averages. In another embodiment the baseline levels are derived from the individual's own medical history.

In another embodiment the method is performed more than once to monitor the reduction or increase in risk for stroke or TIA, optionally in conjunction with the administration of risk reduction therapy. In one embodiment the method is performed at a frequency of from about one week to about six months. In another embodiment the method is performed at a frequency of from about one month to about three months.

In a particularly preferred embodiment other biomarkers are also measured to assess the risk for stroke or TIA. Particularly preferred biomarkers for risk of stroke or TIA are glutamate and homocysteine.

Novel Kits of the Present Invention

In another embodiment the invention provides kits for diagnosing central nervous system disorders such as TIA, stroke, and traumatic brain injury. NR2A and/or NR2B antibodies or antigens may be incorporated into immunoassay diagnostic kits depending upon whether autoantibodies or NMDA receptors are being measured. A first container may include a composition comprising an antigen or antibody preparation. Both antibody and antigen preparations should preferably be provided in a suitable titrated form, with antigen concentrations and/or antibody titers given for easy reference in quantitative applications.

The kits may also include an immunodetection reagent or label for the detection of specific immunoreaction between the provided antigen and/or antibody, as the case may be, and the diagnostic sample. Suitable detection reagents are well known in the art as exemplified by radioactive, enzymatic or otherwise chromogenic ligands, which are typically employed in association with the antigen and/or antibody, or in association with a second antibody having specificity for first antibody. Thus, the reaction is detected or quantified by means of detecting or quantifying the label. Immunodetection reagents and processes suitable for application in connection with the novel methods of the present invention are generally well known in the art.

The reagents may also include ancillary agents such as buffering agents and protein stabilizing agents, e.g., polysaccharides and the like. The diagnostic kit may further include where necessary agents for reducing background interference in a test, agents for increasing signal, apparatus for conducting a test, calibration curves and charts, standardization curves and charts, and the like.

In a more particular aspect the invention relates to a rapid multiple panel containing antibodies to the thromboembolic and neurotoxicity biomarkers glutamate, homocysteine and

NMDA receptors that employs latex agglutination. Thus, in one embodiment the invention provides a kit for diagnosing central nervous system disorders comprising: (1) an agglutinating immunosorbent for NR2A and/or NR2B NMDA receptors or fragments thereof, and (2) a control such as saline or a known concentration of NR2A and/or NR2B receptor or fragment thereof. In a more preferred embodiment the kit further comprises an agglutinating immunosorbent for another biomarker for TIA/stroke, such as an agonist or antagonist of NR2A and/or NR2B, a thromboembolic marker, or more particularly glutamate or polyglutamate, and/or an agglutinating homocysteine or polyhomocysteine. The agglutinating immunosorbent is preferably of the type discussed in greater detail above.

In another embodiment the invention relates to a kit for detecting NR2A and/or NR2B receptors or fragments thereof that does not employ latex agglutination. Thus, in another embodiment the invention provides a kit for diagnosing central nervous system disorders comprising: (1) an immunosorbent for NR2A and/or NR2B NMDA receptors or fragments thereof, and (2) an indicator reagent comprising secondary antibodies attached to a signal generating compound. The secondary antibodies can be specific for the receptor or fragment, or for the primary antibodies in the immunosorbent. In a preferred embodiment the kits further comprise an immunosorbent for glutamate or polyglutamate, and/or an immunosorbent for homocysteine or polyhomocysteine, and secondary antibodies against the glutamate and/or homocysteine, or to the primary antibodies on the immunosorbents against the glutamate or homocysteine. The immunosorbent preferably comprises anti-antibodies for the biomarkers bound to a solid support.

In another aspect the present invention relates to a test-kit that relies upon PCR amplification for measuring NR2A and/or NR2B levels. Thus, in another embodiment the invention provides a kit comprising: (a) one or more oligonucleotide primers (preferably of SEQ ID NO: 8) attached to a solid phase, (b) indicator reagent attached to a signal-generating, compound capable of generating a detectable signal from oligonucleotides, and (c) a control sample (i.e. template cDNA). The reagents may also include ancillary agents such as buffering agents, polymerase agents, and the like. The diagnostic kit may further include, where necessary, other members of the signal-producing system of which system the detectable group is a member (e.g., enzyme and non-enzyme substrates), agents for reducing background interference in a test, agents for increasing the signal, apparatus for conducting a test, and the like.

In another embodiment of test-kit comprises (a) a solid phase to which biological fluids for receiving total DNA including NR2A cDNA could be attached, (b) oligonucleotide primers, preferably in a ready-to-use PCR buffer, and (c) a control sample (i.e. template cDNA). Ancillary agents as described above may similarly be included.

In another embodiment the invention provides a diagnostic kit for detecting NR2A and/or NR2B autoantibodies comprising (a) a polypeptide of the N-terminal domain of the NR2A and/or NR2B receptor, fragment thereof, or analog or derivative thereof, (b) an indicator reagent comprising a secondary antibody specific for the autoantibody or the polypeptide attached to a signal-generating compound; and (c) a control sample, such as a known concentration of NR2A and/or NR2B polyclonal antibodies. The reagents may also include ancillary agents such as buffering agents and protein stabilizing agents, e.g., polysaccharides and the like. The diagnostic kit may further include, where necessary, other members of the signal-producing system of which system the

detectable group is a member (e.g., enzyme and non-enzyme substrates), agents for reducing background interference in a test, agents to increase the signal, apparatus for conducting a test, calibration and standardization information or instructions, and the like.

Novel Compositions of the Invention

The methods of the present invention rely upon a series of novel compositions which themselves form a part of the invention. Thus, in one series of embodiments the invention provides an isolated polypeptide fragment of the NR2A and/ or NR2B NMDA receptor, comprising:

1. An antigenic determinant of the NR2A NMDA receptor,
2. An antigenic determinant of the NR2B NMDA receptor,
3. The N-terminal domain of the NR2A NMDA receptor,
4. The N-terminal domain of the NR2B NMDA receptor,
5. SEQ ID NO. 2,
6. SEQ ID NO. 3,
7. SEQ ID NO. 4,
8. SEQ ID NO. 11,
9. SEQ ID NO. 12, and
10. SEQ ID NO. 13,

or an antigenic fragment, analog, or derivative thereof. In another series of embodiments the invention provides any, of the foregoing polypeptides linked covalently to a distinct antigenic determinant, such as human serum albumin. In still another series of embodiments the invention provides any of the foregoing polypeptides linked to any of the immunosorbent materials discussed above. The immunosorbent can be in the form of a bead for latex agglutination, in the size ranges discussed above, or in the form of a synthetic plate for conventional immunoassay analysis. The polypeptide can be linked to the immunosorbent using any conventional means of linkage, including covalent linkage, ionic linkage, and adsorption.

In another series of embodiments the present invention relates to the novel monoclonal and polyclonal antibodies specific for and/or raised against the foregoing polypeptides, including the foregoing polypeptides linked to distinct antigenic determinants. Thus, in one embodiment the invention provides non-human antibodies against any of the foregoing peptides or polypeptides or antigenic fragment, analog, or derivative thereof. In another embodiment the invention provides immunosorbents to which such antibodies are linked.

In another series of embodiments the present invention provides oligonucleotides that encode the foregoing peptides and polypeptides and fragments, analogs, and derivatives thereof, and to recombinant expression vectors that include such oligonucleotides. Such oligonucleotides include, without limitation, the oligonucleotides defined by SEQ ID NO:6, 7, 14, and 15, and fragments thereof which encode antigenic determinants.

In still another embodiment the present invention relates to isolated oligonucleotide sequences that are useful in the cDNA PCR analytical techniques of the present invention. Thus, the invention further provides oligonucleotides comprising the nucleotide sequences of SEQ ID NOS:7, 8, 9, 15, 16, and 17.

BRIEF DESCRIPTION OF THE SEQUENCE LISTINGS

The features, aspects, and advantages of the present invention will become better understood with regard to the following sequence listings where, in the sequence the recited amino acid position numbering reflects that used throughout this document.

SEQ ID NO:1. shows the full-length amino acid sequence of the mature NR2A receptor subunit, as follows:

SEQUENCE LISTING
 PEPTIDE
 Homo sapiens glutamate receptor. ionotropic, N-methyl D-aspartate 2A
 Science 256:1217-1221 (1992)
 NCBI/NM 000833.2

1	11	21	31	41	51	60
1	MGRLGYWTL	VLPALLVWRD	PAQNAAAEKG	PPALNIAVLL	GHSHDVTERE	LRNLWGPEQA
61	TGLPLDVNVV	ALLMNRDPK	SLITHVCDLM	SGARIHGLVF	GDDTDQEA	QMLDFISSQT
121	FIPILGIHGG	ASMIMADKDP	TSTFFQFGAS	IQQQATVMLK	IMQDYDWHVF	SLVTTIFPGY
181	RDFISFIKTT	VDNSFVGWDM	QNVITLDTSF	EDAKTQVQLK	KIHSSVILLY	CSKDEAVLIL
241	SEARSLGLTG	YDFFWIVPSL	VSGNTELPK	EFPSGLISVS	YDDWDYSLEA	RVRDGLGILT
301	TAASSMLEKF	SYIPEAKASC	YQAEKPEPT	LHTLHQFMVN	VTWDGKDLF	TEEGYQVHPR
361	LVVIVLNKDR	EWKVGKVEN	QTLSLRHAVW	PRYKSPSDCE	PDDNHSIVT	LEEAPFVIVE
421	DIDPLTETCV	RNTVPCRKFV	KINNSTNEGM	NVKKCKGFC	IDILKLSRT	VKFTYDLYLV
481	TNGKHGKKN	NVWNGMIGEV	VYQRAVAVG	SLTINEERSE	VVDFSVPFVE	TGISVMVRS
541	NGTVSPSAFL	EPFSASVWVM	MFVMLLIVSA	IAVFVFEYFS	PVGYNRNLAK	GKAPHGPSFT
601	IGKAIWLLWG	LVFNNSVPVQ	NPKGTTSKIM	VSVWAPFAVI	FLASYTANLA	AFMIQEEFVD
661	QVTGLSDKKF	QRPHDYSPPF	RFGTVPNGST	ERNTRNNYPY	MHQYMRFNQ	RGVEDALVSL
721	KTGKLDAFIY	DAAVLNKAG	RDEGCKLVTI	SGYIFASTG	YGIALQKQSP	WKRQIDLALL
781	QFVGDGEMEE	LETLWLTGIC	HNEKNEVMSS	QLDIDNMAGV	FYMLAAAMAL	SLITFIWEHL
841	FYWKLRFCFT	GVCSDRPGLL	FSISRGIYSC	IHGWHIEEK	KSPDFNLTGS	QSNMLKLLRS

- continued

901 AKNISMSNM NSSRMDSPKR ATDFIQRGS L IVDMVS DKGN LIYSDNRSFQ GKDSIFGDNM 960
 961 NELQTFVANR HKDNLSNYVF QGQHPLTLNE SNPNTVEVAV STESKGNRSP RQLWKKSMES 1020
 1021 LRQDSL NQNP VSQRDEKTAE NRTHSLKSPR YLPEEVAHSD ISETSSRATC HREPDNNKNH 1080
 1081 KTKDNFKRSM ASKYPKDCSD VDRTYMKTKA SSPRDKIYTI DGEKRPSFHL DPPQFVENIT 1140
 1141 LPENVGF PDT YQDHNENFRK GDSTLPMNRN PLHNEDGLPN NDQYKLYAKH FTLKDKGSPH 1200
 1201 SEGSDRYRQN STHCRSCLSN LPTYSGHFTM RSPFKCDA CL RMGNLYDIDE DQMLQETG NP 1260
 1261 ATREEVYQOD WSQNNALQFQ KNKLRINRQH SYDNILDKPR EIDL SRPSRS ISLKDRERLL 1320
 1321 EGNLYGSLFS VPSSKLLGNK SSLFPQGLE D SKRSKSLLPD HASDNPF LHT YGDDQRLVIG 1380
 1381 RCPSPDYKHS LPSQAVNDSY LRSLSRSTAS YCSRDSR GHS DVYISEHVMP YAANKNTMYS 1440
 1441 TPRVLNCSN RRVYKMP SI ESDV

SEQ ID NO:2. shows the amino acid sequence of the ²⁰ auto-antigenic region of the N-terminal domain of the NR2A subunit, as follows:

SEQ ID NO:2
 HOMO SAPIENS
 PAQNAAA EKG P PALNIAVLL GHSHDVTERE LRNLWGPEQA 60
 61 TGLPLDVNVV ALLMNRDPK SLITHVCDLM SGARIHGLVF GDDTDQEA VA QMLDFISSQT 120
 121 FIPILGIHGG ASMIMADKDP TSTFFQFGAS IQQQATVMLK IMQDYDWHVF SLVTTIPPGY 180
 181 RDFSIFIKTT VDNFSV GWM QNVITLDTSF EDAKTQVQLK KIHSSVILLY CSKDEAVLIL 240
 241 SEARSLGLTG YDFFWIVPSL VSGNTE LIPK EPPSGLISVS YDDWDYSLEA RVRDGLGILT 300
 301 TAASSMLEKF SYIPEAKASG YGQAEKPE TP LHTLHQFMVN VTWDGKDL SF TEEGYQVHPR 360
 361 LVVIVLNKDR EWEKVGK WEN QTLSLRHAVW PRYKSPSDCE PDDNHL SIVT LEEAPFVIVE 420
 421 DIDPLTETCV RNTVPCRKFV KINNSTNEGM NVKKCKGFC IDILKKSRT VKFTYDLYLV 480
 481 TNGKHGKKNV NVWNGMIGEV VYQRAVMAVG SLTINEERSE VVDFSVPFVE TGISVMVSR S 540
 541 NGTVSPSAFL EPPSAS

SEQ ID NO:3; shows a 21 amino acid antigenic peptide, corresponding to a fragment of the NR2A N-terminal domain. another such peptide (21 amino acids derived from the NR2A sequence and an N-terminal Cys for attachment to a carrier protein), as follows:

Artificial Sequence
 CNGMIGEVVYQRAVMAVGLTI

SEQ ID NO:3
 Homo sapiens
 NGMIGEVVYQRAVMAVGLTI

50 Full
 Base Count Origin

Homo sapiens glutamate receptor, ionotropic, N-methyl D-aspartate 2A (GRIN 2A) mRNA

SEQ ID NO:4. shows a 22 amino acid antigenic peptide, corresponding to a fragment of the NR2A N-terminal domain. another such peptide, modified by an N-terminal Cys for attachment to a carrier protein):

55 SEQ ID NO:5. shows the Oligonucleotide position numbering used throughout in reference to NR2A oligonucleotide sequences, as follows:

SEQ ID NO:5
 Science 256:1217-1221 (1992)
 May 22, 1992
 NIGB/NM_000833
 1 atcatgggac cgggtgagcg ctgagaatcg cggccgcagc catcagcctt ggagatgacc
 61 aggagcggcc actgctgaga actatgtgga gagaggctgc gagcctgct gcagagcctc

-continued

121 cggctgggat agccgcccc cgtggggcg atgcggacag cgcgggacag ccaggggagc
 181 gegctggggc cgcagcatgc gggaaaccgc taaaccgggt ggctgctgag gcggccgaga
 241 tgctcgtgcg cgcagcgcgc cccaactgcat cctcgacctt ctcgggctac agggaccgtc
 301 agtggcgact atgggcagag tgggctattg gaccctgctg gtgctgccgg cccttctggt
 361 ctggcgcggt ccggcgcgca gcgcggcggc ggagaagggt cccccgcgc taaatattgc
 421 ggtgatgctg ggtcacagcc acgacgtgac agagcgcgaa ctctgaacac tgtggggccc
 481 cgagcaggcg gcggggctgc ccttggaact gaactggta gctctgctga tgaaccgac
 541 egacccaag agcctcatca cgcacgtgtg cgacctcatg tcgggggac gcatccacgg
 601 cctcgtgttt ggggacgaca cggaccagga ggccgtagcc cagatgctgg atttatctc
 661 cccccacacc ttcgtcccca tcttgggcat tcatgggggc gcatctatga tcatggctga
 721 caaggatccg acgtctacct tcttcagtt tggagcgtcc atccagcagc aagccacgg
 781 catgctgaag atcatgcagg attatgactg gcatgtcttc tcctgggtga ccactatctt
 841 ccttggtac aggaattca tcagcttctg caagaccaca gtggacaaca gctttgtggg
 901 ctgggacatg cagaatgtga tcacactgga cacttccttt gaggatgcaa agacacaagt
 961 ccagctgaag aagatccact cttctgtcat cttgctctac tgttccaaag acgaggtgt
 1021 tctcattctg agtggggccc gctcccttg cctcaccggg tatgatttct tctggattgt
 1081 cccagcctg gtctctggga acacggagct catccaaaa gagtttccat cgggactcat
 1141 tctgtctccc tacgatgact gggactacag cctggaggcg agagtgggg acggcattgg
 1201 catcctaacc accgctgcat cttctatgct ggagaagttc tctacatcc ccgaggccaa
 1261 gcccagctgc tacgggcaga tggagaggcc agaggctccg atgcacacct tgcaccatt
 1321 tatggtcaat gttacatggg atggcaaaga cttatccttc actgaggaag gctaccaggt
 1381 gacccccagg ctggtggtga ttgtgctgaa caaagaccgg gaatgggaaa aggtgggcaa
 1441 gtgggagaac catacgtgca gctgaggca cgcctgtggt cccaggtaca agtccttctc
 1501 cgactgtgag ccgatgaca accatctcag catcgtcacc ctggaggagg cccattctg
 1561 catcgtggaa gacatagacc cctgaccga gacgtgtgtg aggaacaccg tgccatgtcg
 1621 gaagtctgca aaaatcaaca attcaaccaa tgaggggatg aatgtgaaga aatgctgcaa
 1681 ggggttctgc attgatattc tgaagaagct tccagaact gtgaagtta cttacgacct
 1741 ctatctgggt accaatggga agcatggcaa gaaagttaac aatgtgtgga atggaatgat
 1801 cgggtgaagt gtctatcaac gggcagtcac ggcagttggc tcgctacca tcaatgagga
 1861 acgttctgaa gtggtgact tctctgtgcc ctttgggaa acgggaatca gtgtcatggt
 1921 ttcaagaagt aatggcaccg tctcaccttc tgettttcta gaaccattca gcgcctctgt
 1981 ctgggtgatg atgtttgtga tgctgctcat tgtttctgcc atagctgttt ttgtcttga
 2041 atacttcagc cctgttggat acaacagaaa cttagccaaa gggaaagcac cccatgggccc
 2101 tctttttaca attgaaaaag ctatatggct tctttggggc ctggtgttca ataactccgt
 2161 gctgtccag aatcctaag ggaccaccag caagatcatg gtatctgtat gggccttctt
 2221 cgctgtcata ttctggcta gctacacagc caatctggct gccttcata tccaagagga
 2281 atttgtggac caagtgaccg gcctcagtga caaaaagttt cagagacctc atgactattc
 2341 cccacctttt cgatttggga cagtgccata tggaaagcag gagagaaca ttcggaataa
 2401 ctatccctac atgcatcagt acatgaccaa atttaacag aaaggagtag aggacgcctt
 2461 ggtcagcctg aaaacgggga agctggagc tttcatctac gatgccgag tcttgaatta

-continued

2521 caaggctggg agggatgaag gctgcaagct ggtgaccatc gggagtgggt acatctttgc
 2581 caccaccggt tatggaattg cccttcagaa aggctctcct tggaagaggc agatcgacct
 2641 ggcttctgct cagtttctgg gtgatgggta gatggaggag ctggagacct tgtggctcac
 2701 tgggatctgc cacaacgaga agaacgaggt gatgagcagc cagctggaca ttgacaacat
 2761 gggggcgta ttctacatgc tggtgcccgc catggccctt agcctcatca ccttcatctg
 2821 ggagacacct ttctactgga agctgctgct ctgtttcacc ggcgtgtgct ccgaccggcc
 2881 tgggttctgc ttctccatca gcaggggcat ctacagctgc attcatggag tgcacattga
 2941 agaaaaaag aagtctccag acttcaatct gacgggatcc cagagcaaca tggtaaaact
 3001 cctccggtca gccaaaaaca tttccagcat gtccaacatg aactcctcaa gaatggactc
 3061 acccaaaaaga gctgctgact tcatccaaag aggttccctc atcatggaca tggtttcaga
 3121 taaggggaat ttgatgtact cagacaacag gtcctttcag gggaaagaga gcatttttgg
 3181 agacaacatg aacgaactcc aaacatttgt ggccaaccgg cagaaggata acctcaataa
 3241 ctatgtattc cagggacaac atcctcttac tctcaatgag tccaacccta acacggtgga
 3301 ggtggcctg agcacaagaat ccaaagcgaa ctctagacct cggcagctgt ggaagaaatc
 3361 cgtggattcc ataccgagg attcactatc ccagaatcca gtctcccaga gggatgaggc
 3421 aacagcagag aataggacct actccctaaa gagccctagg tatcttccag aagagatggc
 3481 ccaactctgac atttcagaaa cgtcaaatcg ggccacgtgc cacagggaac ctgacaacag
 3541 taagaaccac aaaaccaagg acaactttaa aaggtcagtg gcctccaaat accccaagga
 3601 ctgtagttag gtcgagcgca cctacctgaa aaccaaatac agctccccta gagacaagat
 3661 ctacactata gatggtgaga aggagcctgg tttccactta gatccacccc agtttgttga
 3721 aaatgtgacc ctgcccgaga acgtggactt cccggacccc taccaggatc ccagtgaaaa
 3781 cttccgcaag ggggactcca cgctgccaat gaaccggaac cccttgcata atgaagaggg
 3841 gctttccaac aacgaccagt ataaactcta ctccaagcac ttcaacctga aagacaaggg
 3901 ttcccgcac agtgagacca gcgagcgata ccggcagaac tccaagcact gcagaagctg
 3961 cctttccaac atgcccacct attcaggcca cttcaccatg aggtcccctt tcaagtgcga
 4021 tgctgctctg cggatgggga acctctatga catcgatgaa gaccagatgc ttcaggagac
 4081 aggtaaccca gccaccgggg agcaggtcta ccagcaggac tgggcacaga acaatgccct
 4141 tcaattacaa aagaacaagc taaggattag ccgtcagcat tcctacgata acattgtcga
 4201 caaacctagg gagctagacc ttagcaggcc ctcccggagc ataagcctca aggacagggg
 4261 acggttctg gagggaaatt tttacggcag cctgtttagt gtcccctcaa gcaaaactct
 4321 ggggaaaaaa agctcccttt tcccccaagg tctggaggac agcaagagga gcaagtctct
 4381 cttgccagac cacacctccg ataacccttt cctccactcc cacagggatg accaacgctt
 4441 ggttattggg agatgccctt cggaccctta caaacctcg ttgccatccc aggcggtgaa
 4501 tgacagctat cttcggctgt ccttgaggtc aacggcatcg tactgttcca gggacagctc
 4561 gggccacaat gatgtgtata tttcggagca tgttatgcct tatgctgcaa ataagaataa
 4621 tatgtactct acccccagg ttttaaatc ctgcagcaat agacgcgtgt acaagaaaat
 4681 gcctagtatc gaatctgatg tttaaaaatc ttccattaat gttttatcta tagggaaata
 4741 cacgtaatgg ccaatgttct ggagggtaaa tgttgatgt ccaatagtgc cctgctaaga
 4801 ggaagaagat gtagggaggt attttgttgt tgttgttgtt ggctcttttg cacacggctt
 4861 catgccataa tcttccactc aaggaatctt gtgaggtgtg tgctgagcat ggcagacacc

-continued

4921 agataggatga gtccttaacc aaaaataact aactacataa gggcaagtct ccgggacatg
 4981 cctactgggt atgttgcaa taatgatgca ttggatgcca atggatgatg tatgatttcc
 5041 tatattccaa attccattaa ggtcagccca ccatgtaatt ttctcatcag aaatgcctaa
 5101 tggtttctct aatacagaat aagcaatag gtgtgcatgt aaacctgaca cagacaaaat
 5161 aaaaacagtt aagaatgcat ctgcactgta gtcggatttg aacatgtgca agagattagg
 5221 aagtttggct cgtaaacagtt tcagctttct tggatgcoct tccatcacag ccagggctca
 5281 cccaagaac tccaggctcc cctaaagaat agcaaatcag tgtgtctgtg atgactgtgc
 5341 taccttcatt atagtccatt tccaagacac atctggagcc aaaggccga gggaccctca
 5401 ggtggggaga gctacaggaa tctctttgga tggatgatgt tgtttctctc taccctcggc
 5461 ttcgatggtc ttgttcagag ctgcataaac taacacattt atgtctccga gatctaagtg
 5521 tggatctctt gtctgtgaca cagtggccat ttagtattat cccgaagacg cctatgtacg
 5581 taagtttga tttctctctt tctggatgt actcagggtt gtatagtatc tgttaccctt
 5641 tccctccag agtaaccata actcgttccg tttccaaaca gccatggtgg tgtccaatta
 5701 gctgtgtatc gctcttccca gagttgtaa tgtggtgaca tgcaccaaca gccgtatgtg
 5761 tactgtgatc tgtaagaagt acaatgccat ctgtctgccg aaggctagca tggttttagg
 5821 tttatcttcc ttcacatcca gaaattctgt tggacactca cttccacccc aaactcctca
 5881 aatcaaaaagc cttcaaaaac cgaggcactc ttggatctac cctgagtatc ctccaaactg
 5941 tggatagcag ttagtgagac aagcaatttc tccctctga gttattctct ctgttgggtg
 6001 caaaccactt catagcacca acagagatgt agggaaaatt cctcaaagta tttgtcattt
 6061 ctgagtcgcc tgcattatcc cattcttatt ctctcaaac ctgtgcata atgacatgaa
 6121 atgatatcca tttttttttt aagttagaaa cagagagggg aatacttatg catggggagc
 6181 ctgttagcac agtgcctgcc acaaaaacaa gtgccccga caagatagtt gctatgttat
 6241 gacactttct cagatcagga tttctagtt taaaaattaa atatcataaa acg

40

SEQ ID NO:6. shows the oligonucleotide sequence of the auto-antigenic region of the N-terminal domain of the NR2A subunit, as follows:

SEQ ID NO:6

N-terminal nucleotide sequence

371 ccggcgccga gcgcggcggc ggagaagggt cccccgcgc
 taaatattgc
 421 ggtgatgctg ggtcacagcc acgacgtgac agagcgcgaa
 cttcgaacac tgtggggccc
 481 cgagcaggcg gcggggctgc ccctggacgt gaacgtggta
 gctctgctga tgaaccgcac
 541 cgaccccaag agcctcatca cgcacgtgtg cgacctcatg
 tccggggcac gcatccacgg
 601 cctcgtgttt ggggacgaca cggaccagga ggccgtagcc
 cagatgctgg attttatctc
 661 ctccacaccc ttctgcccc tcttgggcat tcatgggggc
 gcatctatga tcatggctga
 721 caaggatccg acgtctacct tcttccagtt tggagcgtcc
 atccagcagc aagccacggt
 781 catgctgaag atcatgcagg attatgactg gcatgtcttc
 tccctggtga ccaactatctt

- continued

841 cectggctac agggaattca tcagettcgt caagaccaca
gtggacaaca gctttgtggg

901 ctgggacatg cagaatgtga tcacactgga cacttccttt
gaggatgcaa agacacaagt

961 ccagctgaag aagatccact cttctgtcat cttgctctac
tgttccaaag acgaggctgt

1021 tctcattctg agtgaggccc gctcccttgg cctcaccggg
tatgatttct tctggattgt

1081 ccccagcttg gtctctggga acacggagct catcccaaaa
gagtttccat cgggactcat

1141 ttctgtctcc tacgatgact gggactacag cctggaggcg
agagtgaggg acggcatgtg

1201 catcctaacc accgctgcat cttctatgct ggagaagttc
tctacatcc ccgaggccaa

1261 ggccagctgc tacgggcaga tggagaggcc agaggtccc
atgcacacct tgcaccatt

1321 tatggtcaat gttacatggg atggcaaaga cttatccttc
actgaggaag gctaccagg

1381 gcaccccagg ctggtgtgga ttgtgctgaa caaagaccgg
gaatgggaaa aggtgggcaa

1441 gtgggagaac catacgtga gcttgaggca cgccgtgtgg
cccaggtaaa agtccctctc

1501 cgactgtgag ccgatgaca accatctcag catcgtcacc
ctggaggagg cccattcgt

1561 catcgtagaa gacatagacc cctgaccga gacgtgtgtg
aggaacaccg tgccatgtcg

1621 gaagtctgtc aaaatcaaca attcaaccaa tgaggggatg
aatgtgaaga aatgctgcaa

1681 ggggttctgc attgatattc tgaagaagct ttccagaact
gtgaagttaa cttacgacct

1741 ctatctggtg accaatggga agcatggcaa gaaagttaac
aatgtgtgga atggaatgat

1801 cggatgaagtg gtctatcaac gggcagtcac ggcagttggc
tcgctcacca tc aatgagga

1861 acgttctgaa gtggtggact tctctgtgac ctttgtagaa
acgggaatca gtgcatggt

1921 ttcaagaagt aatggcaccg tctcaccttc tgcttttota
gaaccattca gcgcctct

SEQ ID NO: 7 shows a 62 oligonucleotide fragment target, as follows:

SEQ ID NO: 7
atggaatgatcggatgaagtgggtctatcaacgggcagtcgagtgaggc
tcgctcaccatc

SEQ ID NO: 8 shows one oligonucleotide primer, as follows:

SEQ ID NO: 8
agcatggcaagaaagttaaca

SEQ ID NO: 9 shows a second oligonucleotide primer, as follows:

SEQ ID NO: 9
acgttctgaagtgtggactt

SEQ ID NO: 10. shows the full-length amino acid sequence of the mature NR2B receptor subunit, as follows:

Peptide

Homo sapiens glutamate receptor, ionotropic, N-methyl D-aspartate 2B

Biochim. Biophys. Acta 1260:105-108(1995).

sequence NME2_HUMAN (Q13224)

1	11	21	31	41	51		
1	MKPRAECCSP	KFWLVLA	VLSRARSQK	SPPSIGIAVI	LVGTSDEVAI	KDAHEKDDFH	60

- continued

61 HLSVVPVEL VAMNETDPKS IITRICDLMS DRKIQGVVFA DDTDQEAIQ ILDFISAQTL 120
 121 TPILGIHGGG SMIMADKDES SMFFQFGPSI EQQASVMLNI MEEYDWIYFS IVTTYFPGYQ 180
 181 DFVNKIRSTI ENSFVGWELE EVLLLDMSLD DGDSKIQNQL KKLQSP⁵⁵ILL YCTKEEATYI 240
 241 FEVANSVGLT GYGYTWIVPS LVAGD⁵⁵DTVP AEPPTGLISV SYDEWDYGLP ARVRDGIAII 300
 301 TTAASDMLSE HSF⁵⁵PIPEPKSS CYN⁵⁵THEKRIY QSNMLNRYLI NVTFEGRNLS FSE⁵⁵DGYQMHP 360
 361 KLVIILLNKE RKWERV⁵⁵GKWK DKSLQMKY⁵⁵YV WPRMCPETEE QEDD⁵⁵HLSIVT LEEAPFVIVE 420
 421 SVDPLSGTCM RNTVPCQKRI VTENKTDEEP GYIKKCKGF CIDILK⁵⁵KISK SVKFTYDLYL 480
 481 V⁵⁵TNGKHGKKI NGT⁵⁵WNGMIGE VVMKRAYMAV GSLTINEERS EVVDFSV⁵⁵PFI ETGISVMVSR 540
 541 SNGTVSPSAF LEPFSADV⁵⁵WV MMFVMLLIVS AVAVFVFEYF SPVGYNRCLA DGREPGG⁵⁵PFS 600
 601 TIGKAIWLLW GLVFNNSV⁵⁵VP QNPKGTTSKI MVS⁵⁵VWAPFAV IFLASYTANL AAFMIQE⁵⁵EYV 660
 661 DQVSGLSDKK FQRPNDF⁵⁵SPP FRFGTVPNGS TERNIRN⁵⁵NYA EMHAYMGKFN QRGVDDALLS 720
 721 LKTGKLD⁵⁵AFI YDAAVLN⁵⁵YMA GRDEGCKLVT IGSGKVF⁵⁵FAST GYGIAIQKDS GWKRQVDLAI 780
 781 LQLFGDGEME ELEALWLTGI CHNEKNEVMS SQLDIDN⁵⁵MAG VFYMLGAAMA LSLITFICEH 840
 841 LFYWQFRHCF MGVCSGKPGM VFSISRGIYS CIHGVAIEER QSV⁵⁵MNSPTAT MN⁵⁵NTHSNILR 900
 901 LLRTAKNMAN LSGVNGSPQS ALDFIRRESS VYDISEH⁵⁵RRS FTHSDCKSYN NPPCEENLFS 960
 961 DYISEVERTF GNLQLKDSNV YQDHYHHHR PHSIGSASSI DGLYDCNPP FTTQ⁵⁵SRSISK 1020
 1021 KPLDIGLPSS KHSQ⁵⁵LSDLYG KFSFKSDRY⁵⁵S GHDDLIRSDV SDISTHTVTY GNIEGNA⁵⁵AKR 1080
 1081 RKQYKDSLK KRPASAKSRR EPDEIELAYR RRP⁵⁵PRSPDHK RYFRDKBGLR DFYLDQ⁵⁵FRTK 1140
 1141 ENSPHWEHVD LTDIYKERSD DFKRDSVSGG GPCTNRSHIK HGTGDKHGVV SGVPAPWEKN 1200
 1201 LTNVEWEDRS GGNFCRSCPS KLHNYSTTVT GQNSGRQACI RCEACKKAGN LYDISEDNSL 1260
 1261 QELDQPAAPV AVT⁵⁵SNASTTK YPQSP⁵⁵NTSKA QKKNR⁵⁵NKLR QHSYDTFVDL QKEEAALAPR 1320
 1321 SVSLKDKGRF MDGSPYAHMF EMSAGESTFA NNKSSVPTAG HHHHNP⁵⁵GGG YMLSKSLY⁵⁵PD 1380
 1381 RVTQNPFIFT FGDDQCLLHG SKSYFFRQPT VAGASKARPD FRALVTNKPV VSALHGAVPA 1440
 1441 RFQKDCICGN QSNPCVPNNK N⁵⁵PRAFNGSSN GHVYEKLS⁵⁵SI

SEQ ID NO: 11. shows the amino acid sequence of the auto-antigenic region of the N-terminal domain of the NR2B subunit, as follows:

45

SEQ ID NO: 11

Homo sapiens

RSQK SPPSIGIAVI LVGTSDEVAI KDAHEKDDPH 50 60

61 HLSVVPVEL VAMNETDPKS IITRICDLMS DRKIQGVVFA DDTDQEAIQ ILDFISAQTL 120121 TPILGLHGGG SMIMADKDES SMFFQFGPSI EQQASVMLNI MEEYDWIYFS IVTTYFPGYQ 180181 DFVNKIRSTI ENSFVGWELE EVLLLDMSLD DGDSKIQNQL KKLQSP⁵⁵ILL YCTKEEATYI 240241 FEVANSVGLT GYGYTWIVPS LVAGD⁵⁵DTVP AEPPTGLISV SYDEWDYGLP ARVRDGIAII 300301 TTAASDMLSE HSF⁵⁵PIPEPKSS CYN⁵⁵THEKRIY QSNMLNRYLI NVTFEGRNLS FSE⁵⁵DGYQMHP 360361 KLVIILLNKE RKWERV⁵⁵GKWK DKSLQMKY⁵⁵YV WPRMCPETEE QEDD⁵⁵HLSIVT LEEAPFVIVE 420421 SVDPLSGTCM RNTVPCQKRI VTENKTDEEP GYIKKCKGF CIDILK⁵⁵KISK SVKFTYDLYL 480481 V⁵⁵TNGKHGKKI NGT⁵⁵WNGMIGE VVMKRAYMAV GSLTINEERS EVVDFSV⁵⁵PFI ETGISVMVSR 540541 SNGTVSPSAF LEPFSAD

65

SEQ ID NO:12; shows a 20 amino acid antigenic peptide fragment of the NR2B subunit, as follows:

SEQ ID NO:12
Homo sapiens
GYIKKCKCKGF CIDILKKISK

SEQ ID NO:13 shows a 21 amino acid sequence of an antigenic fragment of the NR2B subunit modified by an N-terminal Cys for attachment to a carrier protein, as follows:

SEQ ID NO:13
Artificial Sequence (21 aminoacids)
CGYIKKCKCKGF CIDILKKISK.

5
Full

Base Count Origin

SEQ ID NO:14 shows the oligonucleotide position numbering used throughout in reference to NR2B oligonucleotide sequences, as follows:

SEQ. NO. 14
Homo sapiens glutamate receptor, ionotropic, N-methyl D-aspartate
2B mRNA
1 ttgaatttgc atctcttcaa gacacaagat taaaacaaaa tttacgctaa attggatttt
61 aaattatctt ccgttcattt atccttcgctc tttcttatgt ggatatgcaa gcgagaagaa
121 gggactggac attcccaaca tgctcactcc cttaatctgt ccgtctagag gtttggettcc
181 tacaaccaa gggagtcgac gaggtaaga tgaagcccag agcggagtgc tgttctccca
241 agttctgggt ggtgttgccc gtctcggccc tgtcaggcag cagagctcgt tctcagaaga
301 gcccccccag cattggcatt gctgtcatcc tcgtgggccc ttcgacgag gttggccatca
361 aggatgcca cgagaaagat gatttcacc atctctccgt ggtaccccgg gtggaactgg
421 tagccatgaa tgagaccgac ccaagagca tcatcaccg catctgtgat ctcatgtctg
481 accggaagat ccagggggtg gtgtttgctg atgacacaga ccaggaagcc atgccccaga
541 tcctcgattt catttcagca cagactctca ccccgatcct gggcatccac gggggctcct
601 ctatgataat ggcagataag gatgaatcct ccatgttctt ccagtttggc ccatcaattg
661 aacagcaagc ttccgtaatg ctcaacatca tggagaata tgactggtac atcttttcta
721 tcgtcaccac ctatttcctt ggctaccagg actttgtaaa caagatccgc agcaccattg
781 agaatagctt tgtgggctgg gagctagagg aggtcctcct actggacatg tccttggacg
841 atggagattc taagatccag aatcagctca agaaactca aagccccatc attcttcttt
901 actgtaccaa ggaagaagcc acctacatct ttgaagtggc caactcagta gggctgactg
961 gctatggcta cacgtggatc gtgcccagtc tgggtggcagg ggatacagac acagtgcctg
1021 cggagtccc cactgggctc atctctgat catatgatga atgggactat ggctccccg
1081 ccagagtgag agatggaatt gccataatca ccaactgctgc ttctgacatg ctgtctgagc
1141 acagcttcat cctgagccc aaaagcagtt gttacaacac ccacgagaag agaacttacc
1201 agtccaatat gctaaatagg tatctgatca atgtcacttt tgaggggagg aatttgcctt
1261 tcagtgaaga tggctaccag atgcaccga aactgggtgat aattcttctg aacaaggaga
1321 ggaagtggga aagggtgggg aagtggaaa acaagtccct gcagatgaag tactatgtgt
1381 ggccccgaat gtgtccagag actgaagagc aggaggatga ccactctgagc attgtgacct
1441 tggaggaggc accatttctc attgtggaaa gtgtggacc tctgagtgga acctgcatga
1501 ggaacacagt cccctgccc aaaagcatag tcaactgagaa taaaacagac gaggagccgg
1561 gttacatcaa aaaatgctgc aaggggttct gtattgacat ccttaagaaa atttctaaat
1621 ctgtgaagtt cacctatgac cttacctgg ttaccaatgg caagcatggg aagaaaatca
1681 atggaacctg gaatggtatg attggagagg tggatcatgaa gagggcctac atggcagtgg
1741 gctcactcac catcaatgag gaacgatcgg aggtggctga cttctctgtg cccttcatag
1801 agacagccat cagtgtcatg gtgtcacgca gcaatgggac tgtctcactc tctgcctctc

-continued

1861 tagagocatt cagcgetgac gtatgggtga tgatgtttgt gatgctgctc atcgtctcag
 1921 ccgtggctgt ctttgtcttt gagtacttca gccctgtggg ttataacagg tgccctcgtg
 1981 atggcagaga gcctgggtga ccctctttca ccctcggcaa agctatattg ttgctctggg
 2041 gtctgggtgt taacaactcc gtacctgtgc agaaccctaaa ggggaccacc tccaagatca
 2101 tgggtgctcagt gtgggccttc tttgctgtca tcttctcggc cagctacact gccaaacttag
 2161 ctgccttcat gatccaagag gaatatgtgg accaggtttc tggcctgagc gacaaaaagt
 2221 tccagagacc taatgacttc tcacccctt tccgctttgg gaccgtgcc aacggcagca
 2281 cagagagaaa tattcgaat aactatgcag aaatgcatgc ctacatggga aagtcaacc
 2341 agaggggtgt agatgatgca ttgctctccc tgaaaacagg gaaactggat gccttcatct
 2401 atgatgcagc agtgctgaac tatatggcag gcagagatga aggctgcaag ctggtgacca
 2461 ttggcagtgt gaaggtcttt gcttccactg gctatggcat tgccatccaa aaagattctg
 2521 ggtggaagcg ccaggtggac cttgctatcc tgcagctctt tggagatggg gagatggaag
 2581 aactggaagc tctctggctc actggcattt gtcacaatga gaagaatgag gtcatgagca
 2641 gccagctgga cattgacaac atggcagggg tcttctacat gttggggcg gccatggctc
 2701 tcagcctcat caccttcatc tgogaacacc ttttctattg gcagttccga cattgcttta
 2761 tgggtgtctg ttctggcaag cctggcatgg tcttctccat cagcagaggt atctacagct
 2821 gcatccatg ggtggcgatc gaggagcgcc agtctgtaat gaactcccc accgcaacca
 2881 tgaacaacac aactccaac atcctgcgcc tgetgcgcac ggccaagaac atggctaacc
 2941 tgtctggtgt gaatggctca ccgcagagcg ccctggactt catccgacgg gagtcatccg
 3001 tctatgacat ctcagagcac cgcgcagct tcacgcattc tgaactgcaa tctacaaca
 3061 acccgccctg tgaggagaac ctcttcagtg actacatcag tgaggtagag agaacgttcg
 3121 ggaacctgca gctgaaggac agcaacgtgt accaagatca ctaccacat caccaccggc
 3181 cccatagat tggcagtgcc agctccatcg atgggctcta cgactgtgac aaccaccctt
 3241 tcaccacca gtccaggtcc atcagcaaga agccctgga catcggcctc cctcctcca
 3301 agcacagcca gctcagtgac ctgtaacgca aattctcctt caagagcgac cgctacagtg
 3361 gccacgacga cttgatccgc tccgatgtct ctgacatctc aaccacacc gtcacctatg
 3421 ggaacatcga gggcaatgcc gccaaaggcc gtaagcagca atataaggac agcctgaaga
 3481 agcggcctgc ctccgccaag tcccgcaggg agtttgacga gatcgagctg gcctaccgtc
 3541 gccgaccgcc ccgctcccct gaccacaagc gctacttcag ggacaaggaa gggctacggg
 3601 acttctacct ggaccagttc cgaacaaagg agaactcacc cactgggag cacgtagacc
 3661 tgaccgacat ctacaaggag cggagtgatg actttaagcg cgactccatc agcggaggag
 3721 gccctgtac caacaggtct cacatcaagc acgggacggg cgacaaacac ggcgtggtca
 3781 gcgggtacc tgcaccttg gagagaacc tgaccaacgt ggagtgaggag gaccggtccg
 3841 ggggcaactt ctgcccagc tgtccctcca agctgcacaa ctactccacg accgtgacgg
 3901 gtcagaactc gggcagcgag gcgtgcatcc ggtgtgaggc ttgcaagaaa gcaggcaacc
 3961 tgtatgacat cagtggagac aactcctgc aggaactgga ccagccggct gcccagtg
 4021 cgtgacgctc aaacgcctcc accactaagt accctcagag cccgactaat tccaaggccc
 4081 agaagaagaa ccggaacaaa ctgcgcccgc agcactccta cgacaccttc gtggacctgc
 4141 agaaggaaga agccgcctg gcccgcgca gcgtaagcct gaaagacaag ggcgattca
 4201 tggatgggag cccctacgcc cacatgtttg agatgctcagc tggcgagagc acctttgcca

-continued

4261 acaacaagtc ctcagtgccc actgcccggac atcaccacca caacaacccc ggccggcgggt
 4321 acatgctcag caagtgcctc taccctgacc gggtcacgca aaacctttc atccccactt
 4381 ttggggacga ccagtgcttg ctccatggca gcaaatccta cttcttcagg cagccccagg
 4441 tggcgggggc gtcgaaagcc aggccggact tccgggcctt tgtcaccaac aagccgggtg
 4501 tctcggccct tcatggggcc gtgccagccc gttccagaa ggacatctgt atagggaaac
 4561 agtccaaccc ctgtgtgctt aacaacaaaa accccagggc tttcaatggc tccagcaatg
 4621 ggcattgtta tgagaaactt tctagtattg agtctgatgt ctgagtgagg gaacagagag
 4681 gttaaggttg gtacgggagg gtaaggctgt gggtcgctgt atgcgcatgt cacggagggt
 4741 gacgggggtg aacttggttc ccatttctc cttctctgtt ttaatttatt tatgggatcc
 4801 tggagtctgt gttcctactg ggggcaaccc tggtgaccag caccatctct cctcctttc
 4861 acagttctct ccttcttccc cccgctgtca gccattctct tccccatgag atgatgccat
 4921 gggccctctc agcagggggg ggtagagcgg agaaaggaag ggctgcatgc gggcttctc
 4981 ctgggtgtag agagctcctt gatatcctct ttgagtgaag ctgggagaac caaaaagagg
 5041 ctatgtgagc acaaaggtag cttttcccaa actgatcttt tcatttaggt gaggaagcaa
 5101 aagcatctat gtgagaccat ttagcacact gcttgtgaaa ggaagaggc tctggctaaa
 5161 ttcattgctc ttagatgaca tctgtctagg aatcatgtgc caagcagagg ttgggaggcc
 5221 atttgtgttt atatataagc ccaaaaatgc ttgettcaac cccatgagac tcgatagtgg
 5281 tgggtgaacag aaccaaggtt cattgggtggc agagtggatt cttgaacaaa ctggaaagta
 5341 cgttatgata gtgtcccccg gtgccttggg gacaagagca ggtggattgt gcgtgcatgt
 5401 gtgttcacgc acatttgac ccatgtgtag tcagggtgct caagagaagg caaccttgac
 5461 tctttctatt gtttcttcca atatcccaa gcagtgatgat tgtttggctt atatacagac
 5521 agagatggcc atgtattacc tgaattttgg ctgtgtctcc cttcatcctt ctggaataag
 5581 gagaatgaaa attcttgata aagaagattc tgtggctctaa acaaaaaaag gcggtgagca
 5641 atcctgcaag aacaaggtac ataaacaagt cctcagtggt tggcaattgt ttcaaccagt
 5701 ttgaaccaag aactttccag gaaggctaaa gggaaaccga attttcacag ccatgattct
 5761 tttgccaca cttgggagca aaagattcta caaagctctt ttgagcattt agactctcga
 5821 ctggccaagg tttggggaag aacgaagcca cctttgaaga agtaaggagt cgtgtatggt
 5881 agggtaagtg agagaggggg atgtttccaa tgettgtatc ccttctact taacctgaag
 5941 ctagacgagc aggtctcttc cccccaaaac tgattacaac tgctacagag cagacagtta
 6001 agagaaatga gcttgacctt taagagaaat gagctgact ccatgagtgc agctctggag
 6061 gtacgaaaag aggggaagag acttggaat gggagacggg ggcagagagg gaccctccac
 6121 cacctctttg ggctggctc cctgggaatg tgacttgagc ccagagtga cactcttggt
 6181 agaagccctt ctacctctct gcaacacctt gttccctct cagattgtac cattgag

55

SEQ ID NO:15 shows a 60 oligonucleotide fragment target, as follows:

SEQ ID NO:15
 g gttacatcaa aaaatgctg aaggggttct gtattgacat

ccttaagaaa atttctaaa

SEQ ID NO:16 shows one oligonucleotide primers (21 nucleotides), as follows:

SEQ ID NO:16
 tcactgagaa taaaacagac g

60

SEQ ID NO:17 shows one oligonucleotide primers (21 nucleotides), as follows:

SEQ ID NO:17
 t cacctatgac ctttacctgg

65

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds claimed herein are made and evaluated, and are intended to be purely exemplary of the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.) but some errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in ° C. or is at room temperature, and pressure is at or near atmospheric.

Example 1

Preparation of Polyclonal Antibodies (IgG) to Glutamate and Homocysteine

Glutamate (polyglutamate, 10 aminoacids) or homocysteine (polyhomocysteine, 10 aminoacids) alone will not generate antibodies when injected into an animal. Therefore, polyglutamate and polyhomocysteine were conjugated with human serum albumin for the immunization to obtain polyclonal antibodies. For glutaraldehyde conjugation, polyglutamate or homocysteine (10 mg) and 40 mg bovine serum albumin (BSA, Sigma, St. Louis, Mo.) were incubated for 2 hr at room temperature in 4 ml of PBS containing 5% glutaraldehyde. The reaction was stopped by adding glycine to a final concentration of 0.2 M, and the conjugate was dialyzed against PBS.

Rabbits were given initial injections of 1 mg of conjugated glutamate (polyglutamate) or homocysteine (polyhomocysteine) in complete Freund's and subsequent increased doses of injections (2 mg) in incomplete Freund's adjuvant at successive 2 week intervals. All injections were given subcutaneously. The immunization period lasted for 110 days. Antibodies (IgG) were affinity purified according to standard procedures (Warr, G. W., Purification of antibodies, In: Antibody as a Tool, Eds., Marchalonis, J. J., and G. W. Warr, J. Wiley, UK, pp. 59-96 (1982)) and were shown to be selective for glutamate or homocysteine by ELISA assay.

Example 2

Preparation of Polyclonal Antibodies (IgG) to NR2A Receptor Peptide

Using computer analysis of the hydrophobicity profile of human NR2A and NR2B NMDA receptors to predict the antigenic determinants in the protein structure, we selected fragments corresponding to the N-terminal sequence of human NR2A and NR2B receptor peptides for synthesis. The fragments corresponded to the N-terminal sequence of the NR2A and NR2B receptors, represented by SEQ ID NO: 1 and SEQ ID NO: 2 for the NR2A and NR2B receptors, respectively. The peptide fragments were reproduced using solid-phase synthesis, and had a purity ranging from 90% to 98%. The peptide sequences were verified by amino acid analysis after acid hydrolysis. A mixture of NR2A and NR2B peptides (1:1) was conjugated with human serum albumin for the immunization to obtain polyclonal antibodies. For glutaraldehyde conjugation, 10 mg of the mixture of peptides and 40 mg human serum albumin (Sigma, St. Louis, Mo.) were incubated for 1.45 hr at room temperature in 4 ml of PBS containing 5% glutaraldehyde. The reaction was stopped by

adding glycine to a final concentration of 0.2 M, and the conjugate was dialyzed against PBS.

Rabbit polyclonal antibodies were raised against the NR2A-B peptides. Rabbits were given initial injections of 1 mg of conjugated peptides in complete Freund's adjuvant and subsequent injections (0.5 mg) in incomplete Freund's adjuvant at successive 2 week intervals. Antibodies were affinity purified according to standard procedures (Warr, G. W., Purification of antibodies, In: Antibody as a Tool, Eds., Marchalonis, J. J., and G. W. Warr, J. Wiley, UK, pp. 59-96 (1982)) and were shown to be selective for NR2A and NR2B NMDA receptors using an ELISA assay.

Example 3

Preparation of Latex Beads Containing Biomarker Antibodies

Three different sensitized latex beads containing IgG against glutamate, homocysteine and NR2A-B receptor peptides were prepared using two types of blue polystyrene latex beads (diameter, 0.25 and 0.4 μm; Sigma, St. Louis, Mo.) as follows. A 1% suspension of latex beads in 50 mM PBS (1 ml, pH 7.0) was mixed with an equal volume of corresponding IgG (2 mg/ml) and incubated on a shaker at room temperature for 2 hours. The mixture was then washed twice with PBS by centrifugation at 9,500 g for 5 min. The pellet was suspended in PBS containing 1% BSA overnight at 4° C. After being washed twice with PBS, the sensitized latex beads were resuspended in latex diluent (50 mM PBS with 1% BSA) at a concentration of 0.4% and stored at 4° C. until used.

Preliminary experiments with latex agglutination (LA) alone were performed to identify problems and to select the most desirable latex particle size. Two types of commercial latex beads were coated with antibodies at various concentrations. Tests were initially performed with the corresponding amino acid or NR2A and NR2B receptor peptides as controls. Particle size and IgG concentration were found to be the primary factors affect the sensitivity of the test. The most desirable particle size was found to be 0.25 μm (blue latex) because particles of this size agglutinated each amino acid and peptide specifically. Higher IgG concentrations showed higher sensitivities. Using blue latex bead coated with 2 mg of IgG per ml, agglutination could be observed within 30 min.

Example 4

Latex Agglutination Analysis of Blood Serum Specimens

Blood samples (5 ml) were collected using standard venipuncture clinical protocol, from patients with TIA, stroke and brain injury (n=30) and examined at the laboratory of CIS Biotech, Inc. in Atlanta (Ga., USA). None of the patients had been treated with anticoagulants, and serum samples were obtained from the clotted blood. All specimens were free of visible lipids, white blood cells, platelets, fibrin, mucus or other contaminants that could cause "false positive" reactions. Platelets, white blood cells, mucus and fibrin were removed by centrifugation. Lipids were removed by filtration.

Specimens to be tested within 72 hours after collection were stored at 2-8° C. For longer storage periods, -20° C. or colder is recommended.

The semi-quantitative analysis of glutamate, homocysteine and NR2A-B receptor peptides in the serum samples is basi-

cally a three step process: serum sample dilution, reaction of latex beads with serum samples, and product analysis.

In previous experiments serial dilutions of the serum samples from 1:4 to 1:64 in saline containing 4% glycerol for better agglutination were performed. The highest dilution in which agglutination was observed corresponded to the sample titer.

Two 25 μ l aliquots of coated latex beads containing the corresponding IgG were layered on a double-concave slide (Fisher Sci., Norcross, Ga.), one with 25 μ l of the serum sample in serial dilution to be tested and one with 25 μ l of PBS as a negative control. After gentle mixing with vortex, agglutination was judged macroscopically against a dark background. A negative reaction corresponded to a homogeneous lactescent background with no agglutination; a positive reaction corresponded to a clearly visible agglutination against the black background and weakly visible agglutination on a slightly lactescent background.

The highest dilution at which agglutination occurs gives the titer of the sample. To obtain the approximate titer in μ g/ml we used the following calculation:

$$\text{Titer } \mu\text{g/ml} = A \times D$$

where A is the test sensitivity, and D is the highest dilution at which agglutination occurs.

Example 5

Description of Patients

Patients observed in trials (n=68) included 9 with pre-stroke, 9 with TIA (mean age 52.0 \pm 3.0), 31 with acute ischemic stroke (mean age 54.7 \pm 1.4) and 11 with mild brain injury (mean age 53.0 \pm 4.4). Clinical evaluation of patients by neuroimaging (CT, MRI, arteriography, Doppler ultrasonography, EEG), detailed physical and neurologic examination and laboratory tests was performed. Patients with TIA were characterized by contra lateral weakness, dysphasia, transient blurring of vision or blindness, abnormal pulsation of the common carotid arteries, microemboli confined to the ipsilateral retina. Untreated patients with pre-stroke demonstrated altered state of consciousness, severe headache, nausea and vomiting, visual disturbances, and focal neurological deficit, with some patients experiencing seizures.

The N-Score rating scale reported in "MCA Infarction" (Orgogozo, 1986) was used for evaluating the neurologic deficit in patients with acute cerebral stroke. The total score of acute cerebral stroke clinical manifestation differentiated severe patients (n=9, 11-35 scores) from patients with mild (n=12, 36-55 scores) and moderate patients (n=10, 60-90 scores). Most patients with acute cerebral ischaemia (61.3%) suffered ischemia in the carotid artery of left hemisphere. Arterial hypertension and cerebral atherosclerosis etiologically corresponded in all patients.

The patients with ischemia were divided into groups based on the differences between TIA, pre-stroke and acute ischemic pathogenic mechanisms. The clinical diagnosis was established on the basis of routine observations which included detailed neurological examination and neuroimaging.

ing. Groups of TIA (n=9) and pre-stroke patients with chronic cerebral blood insufficiency (n=9) were identified by neurophysiological investigations.

Example 6

Detection of Glutamate and Homocysteine in the Blood of Patients

Glutamate and homocysteine content were measured by standard high performance liquid chromatography (HPLC) according to methods described (Perry I. J., Refsum H., Morris R. W., Ebrahim S. B., Ueland P. M., Shaper A. G. *Lancet*. 1995, 346:1395-1398; Yamamoto T., Rossi S., Stiefel M., Doppenberg E., Zauner A., Bullock R., Marmarou A. *Acta Neurochir. Suppl.* 1999, 75: 17-19). The limits of the normal range were 165.0 μ mol/L for glutamate (Table 1) and 8.0 μ mol/L for homocysteine (Table 2). Elevated glutamate and homocysteine amounts were detected in the blood of patients with acute stroke. However, approximately 66% of these patients had additional risk factors indicative of atherosclerotic processes such as high cholesterol and LDL levels (Denisenko T. V., Skuliabin D., Gromov I., Cherkas Yi., Iluchina A., Dambinova S. A., 1998. *Vopr. Med. Khimii*. 44, 584-590, in Russian).

Abnormal glutamate and homocysteine plasma concentrations were observed more frequently in patients with TIA than in patients with acute stroke. The positive predictive efficiency of plasma glutamate for TIA patients was 56%. The positive predictive efficiency of plasma homocysteine for TIA patients was 66%. Baseline concentrations for glutamate and homocysteine are 160 μ mol/L and 10 μ mol/L, respectively. Routine treatment for TIA was found to consistently decrease the glutamate and homocysteine levels in the blood of patients (data not shown).

In patients with pre-stroke, slightly elevated levels homocysteine were observed; levels of glutamate were unchanged (Tables 1, 2). In patients with traumatic brain injury (TBI), glutamate levels were observed that were nearly twice the glutamate levels in healthy individuals; levels of homocysteine were up to 57% higher.

TABLE 1

Group	Glutamate concentration in the blood of patients detected by HPLC					
	Total N	Glutamate μ mol/L	HPLC predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	165.0 \pm 28.2	19	67.8	9	32.2
TIA	9	200.0 \pm 11.7	4	44.4	5	55.6
Pre-stroke	9	163.7 \pm 10.4	5	55.6	4	44.4
Acute stroke	31	172.1 \pm 20.6	13	41.9	18	58.1
TBI	11	305.0 \pm 28.8	4	36.4	7	63.6

We also compared homocysteine concentrations in the blood of patients with TIA and pre-stroke to homocysteine concentrations in the blood of patients who have had stroke onsets. We observed that homocysteine content in the blood of patients depended on stage of the stroke, but that homocysteine concentration did not correlate with the severity of the cerebral ischemia. A significant decrease in homocysteine levels in patients with acute stroke was observed after emergency therapy (data not shown).

Latex agglutination was also employed to detect TIA/stroke biomarkers in the blood serum of patients. The titer of

plasma glutamate determined by latex agglutination was 3.34±0.25 in the group of healthy volunteers. Homocysteine and glutamate trends observed using HPLC were similarly observed for different groups of patients observed by using the LA technique (Tables 3, 4). Thus, increased levels of glutamate and homocysteine were similarly observed in the blood of patients with TIA and acute stroke using LA.

With respect to predictive efficiency, however, LA showed a surprising improvement over HPLC. For example, the LA method improved the positive predictive efficiency of patients with TIA and acute stroke on the basis of glutamate content to more than 63% (Tables 1, 3). The negative predictive value for healthy patients was similarly improved when using the LA technique (Tables 3, 4). The predictive value of the LA technique in the group of patients with TBI was identical to the predictive value using HPLC.

TABLE 2

Homocysteine concentration in the blood of patients detected by HPLC						
Group	N	Homocysteine μmol/L	HPLC predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	8.0 ± 1.7	20	71.4	8	28.6
TIA	9	10.8 ± 1.3	3	33.3	6	66.4
Pre-stroke	9	9.0 ± 1.2	4	44.4	5	55.6
Acute stroke	31	11.5 ± 1.1	11	35.5	20	64.5
TBI	11	12.6 ± 2.1	4	36.4	7	63.6

TABLE 3

Detection of glutamate in the blood of patients by latex agglutination						
Group	N	Glutamate Titer	LA predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	3.34 ± 0.25	22	78.6	6	21.4
TIA	9	4.52 ± 0.38	3	33.3	6	66.4
Pre-stroke	9	3.57 ± 0.32	4	44.4	5	55.6
Acute stroke	31	4.34 ± 0.47	11	35.5	20	64.5
TBI	11	5.12 ± 0.62	4	36.4	7	63.6

TABLE 4

Detection of homocysteine the blood of patients by latex agglutination						
Group	N	Homocysteine Titer	LA predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	2.23 ± 0.21	21	75.0	7	25.0
TIA	9	3.95 ± 0.37	3	33.3	6	66.4
Pre-stroke	9	2.89 ± 0.12	4	44.4	5	55.6
Acute stroke	31	4.01 ± 0.41	10	32.3	21	67.7
TBI	11	4.74 ± 0.38	4	36.4	7	63.6

The Detection of NR2A-B in the Blood of Patients

The excessive activation and damage of NMDA receptors is the result of glutamate, aspartate and homocysteine neurotoxicity. Autoantibodies to have been detected in previous work in the blood of patients with TIA and pre-stroke, supporting our hypothesis that cerebral ischemia causes neuronal damage and the appearance of autoantibodies to NMDA receptor subunits (Gusev E. I., Skvortsova V. I., Alekseev A. A., Izykenova G. A., Dambinova S. A. S. S Korsakov's J. Neurol. & Psych. 1996, 5:68-72; Dambinova S. A., Izykenova G. A. J. High Nervous Activity. 1997, 47: 439-446).

The titer of NR2A-B receptor peptides in the blood of healthy volunteers determined by LA was 2.63±0.92. Using the LA technique, we observed an increase in the test efficiency in the group of healthy persons up to 89% (Table 5). We also observed an improvement in the sensitivity of the LA test over ELISA. For example, patients with pre-stroke had slightly increased levels of NR2A-B receptor peptides over healthy volunteers when tested by ELISA, but had nearly double the level of NR2A-B receptor peptides when measured by LA (Table 5, 6). We detected high levels of NR2A-B receptor peptides using both ELISA and LA in the blood of patients with TIA and acute stroke, and observed comparable levels of predictive efficiency for each test.

Patients with TIA received routine treatment to improve brain circulation. Upon receiving treatment, NR2A-B levels decreased to levels corresponding to those observed for the healthy individuals as the patient's state normalized. As mentioned earlier, glutamate and homocysteine contents also decreased during treatment, but it never reached the levels observed in healthy individuals.

TABLE 5

Detection of NR2A-B receptor peptides in the blood of patients by latex agglutination						
Group	N	NR2A-B Titer	LA predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	2.63 ± 0.92	25	89.3	3	10.7
TIA	9	7.34 ± 0.43	2	22.2	7	77.8
Pre-stroke	9	4.21 ± 0.26	2	22.2	7	77.8
Acute stroke	31	5.20 ± 1.71	4	9.7	27	87.1
TBI	11	3.99 ± 0.44	2	18.8	9	81.8

Completely different profiles of NR2A-B were revealed in the blood of patients with acute ischemic stroke. In the blood of patients (n=8) with severe cerebral ischaemia (30.4±3.2 Orgogozo scores) NR2A-B receptor peptides titer was 4 times higher than that for control group of healthy individuals. The peptides titer for patients with mild to moderate ischemic stroke (n=22, 49-62 Orgogozo scores) was slightly elevated in comparison with those with TIA. The tendency of slight decreases in NR2A-B receptor peptide levels was observed to the end of 30 days of patients' routine treatment, correlating with improvement in the neurological state.

TABLE 6

Detection of NR2A-B receptor peptides in the blood of patients by ELISA						
Group	Total N	NR2A-B Ng/ml	ELISA assay results			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	18.2 ± 2.1	20	71.4	8	28.6
TIA	9	66.6 ± 4.1	2	22.2	7	77.8
Pre-stroke	9	23.7 ± 1.9	3	33.3	6	66.7
Acute stroke	31	73.4 ± 6.5	5	16.1	26	83.9
TBI	11	54.3 ± 4.9	3	27.3	8	72.7

It is necessary notice that efficiency of both laboratory assays to detect the NR2A-B receptor peptides in the blood of patients with TIA/stroke and traumatic brain injury have been determined as 78 and 82% correspondingly.

The simultaneous detection of all brain damage biomarkers: glutamate, homocysteine and NR2A-B receptor peptides in the blood patients by latex agglutination allowed to diagnose rapidly TIA/stroke with efficiency up to 85-89%. The simultaneously increased levels of all biomarkers in the blood reflect the neurological deficit and may be used also for prognosis of diseases outcome. The relation between these biomarkers is showing the degree of thromboembolic and neurotoxicity, involvement in brain processes underlying the ischemia. That fact is very important for choosing the strategy of emergency therapy in short time.

Using the latex agglutination technique allowed us significantly cut off the time of blood analysis from 3-8 hours in ELISA or HPLC to 30 min in L.A. This RMP semi-quantitative test demonstrated the fast, simple for interpretation and reliable data.

Example 8

Identification of cDNA Sequence Encoding Antigenic Determinants of NMDA Receptors

It was necessary to first determine the cDNA sequence coding the immunological fragment of NMDA receptors responsible for the appearance of autoantibodies appearance. To find the most active peptide fragment of NMDA receptors a standard molecular biology procedure was used. Immunopositive phage GT11 containing cDNA coding NMDA receptors was isolated from a human cDNA library using autoantibodies to NMDA receptors isolated from blood samples of patients with severe cerebral ischemia or polyclonal antibodies to the NR2A receptor. An *E. Coli* bacterial system was employed to express the phage GT11 cDNA (600 bp). The expression product was transferred to a MBmp11 vector and a restriction map was constructed by use of a standard restrictases' kit. Three unique sites of the cDNA fragment (PstI, BamHI, and PsaI) were revealed, and the 5'-3' oligonucleotide sequence orientation using KpnI, BamHI and EcoR1 was deduced. The oligonucleotide (target cDNA) obtained was sequenced and compared to the sequence of the NR2A glutamate receptor (SEQ ID NO: 5) from the NCBI library. The target cDNA corresponded to the N-terminal domain of the NR2A receptor (620 bp) of SEQ ID NO: 6, namely SEQ ID NO: 7. Primers for this target nucleotide were designed. All the oligonucleotides were prepared by the phosphoramidite method on an Applied Biosystem 394 synthesizer and were purified by reverse-phase high-pressure liquid

chromatography (HPLC). The oligonucleotides used for detection and capture were synthesized with an amine arm at the 5' end.

Example 9

PCR Analysis of Blood Serum Specimens

Blood samples (5 ml collected by venipuncture) from patients with TIA and pre-stroke (n=30) were collected according to standard clinical protocol and examined at the Department of Neurology of Human Brain Institute, St. Petersburg Russia. The blood specimens were used for total DNA isolation or applied on FTA paper circles.

The quantitative analysis of NR2A cDNA expression in the serum samples is basically a three step process: Total DNA isolation and purification from sera of individuals; specific cDNA coding NR2A receptor amplification; and product analysis.

The total DNA isolated by DNAzol (Mol. Res. Center, Inc., Cincinnati, Ohio) or bound on FTA blood staining collection cards (Life Technologies, Inc., Gaithersburg, Md.) serves as a template for the polymerase chain reaction (PCR). In the first variant, the PCR assay uses a set of specially designed primers (50 pmol), immobilized on solid matrix of microplates and amplifies a specific cDNA sequence (620 bp) coding the NR2A glutamate receptor. In a second variant, the PCR assay uses a master ready-to-use buffer and amplifies cDNA bond on FTA paper. Following amplification, the quantity of a product is determined by enzyme or non-enzyme color reaction with a substrate.

Using the DNAzol reagent for DNA isolation, the whole blood of each individual (0.5 ml) was combined with 1 ml DNAzol (Mol. Res. Center, Inc., Cincinnati, Ohio) for 5 min at room temperature and lysed (Mackey K. et al. Mol. Biotechnol. 9: 1-5 (1997). The organic phase (0.4 ml) of each sample was transferred to a clean tube and 0.4 ml isopropanol was added. The mixture was incubated for 5 min at room temperature and centrifugated at 6,000 g for 6 minutes. The pellet was washed in 0.5 ml DNAzol and centrifugated at the same conditions. The total DNA pellet was mixed with 1 ml of 75% ethanol and centrifuged at 6,000 g for 5 minutes. Then the DNA pellet was diluted in 200 l of 8 mM NaOH and incubated at room temperature for 5 min followed by vortexing. Alkaline DNA solution was then neutralized with 160 l of 0.1 M HEPES, pH 7.4.

Immobilization of oligonucleotide probes (primers, SEQ ID NO:8) was performed as follows. A total of 100 l of 3xPBS buffer containing the primers (150 nM) was dropped into each well of a 96-well microtiter plate (Fisher Sci., Suwanee, Ga.). After incubation for 2 h at 37° C. or overnight at room temperature, the plate was washed three times with 1xPBS buffer containing 0.05% (w/vol) Tween 20. The oligonucleotide-coated plates were stable for 2 months at 4° C.

Direct PCR reactions were performed in a final volume of 50 l (Sisk R.B. in book: Molecular diagnostics: for the clinical laboratorian. Ed. by Coleman W.B., and Tsongalis G.J. Humana Press Inc., Totowa, N.J. 1997, pp.103-121). The total DNA (5 l), isolated from blood samples of individuals, to oligonucleotide-coated plate in duplicates and 45 l of master ready-to-use buffer containing 1 l TaKaRa Z-Taq DNA polymerase (TaKaRa Biomedicals, Otsu, Shiga, Japan) 101AMV/Tfl 5x reaction buffer, 1 l dNTP mix (Promega, Madison, Wis.) 2 l of 25 mM MgSO₄ were added and sealed. The 30-thermal cycles (98° C.—5 s, 66° C.—2 sec) amplification using programmable Gene Cyclor thermocycler (Bio-Rad Lab., Hemphstead, UK) for 20 minutes was performed. Then

501 of PicoGreen reagent (Mol. Probes, Inc., Eugene, Oreg.) were added to each PCR products and mixed on a shaker (BioTechniques 20:676 (1996)). Samples were incubated 5 min at room temperature, protected from light. After incubation the fluorescence of the samples was measured using a fluorescence microplate reader (Mol. Device, Sunnyvale, Calif.) and standard fluorescein wavelengths (excitation 480 nm, emission 520 nm). The fluorescence value of the reagent blank was subtracted from that of each of the samples, and the data was employed to generate five-point standardization curves of fluorescence versus DNA concentration, from 25 pg/ml to 25 ng/ml reaction of control target cDNA (50 ng/ml stock) with the same Pico Green reagent.

The other method of total DNA isolation is follows. Whole blood was spotted onto FTA paper and lysed, and samples of DNA immobilized within the matrix of the stain card were punched into a 3 mm ($\frac{1}{8}$ ") diameter paper (1 mm or 2 mm Harris Micro-Punch™) and amplified directly by the amplification mix (Mackey K. et al. Mol. Biotechnol. 9: 1-5 (1997)).

The FTA Bloodstain Card is divided into 4 circles for at least 4 different 120 1 samples of EDTA collected whole blood. Samples of blood were dried at room temperature for at least 1 hour. A circle was drawn with a #2 pencil around each blood to visualize where the blood had been spotted after the FTA paper processing. The FTA Bloodstain Card was then placed in a small plastic tray and 50 ml of FTA Purification Reagent was added and incubated on a shaker for 5 minutes. FTA Purification Reagent was replace 3 times with 25-50 ml of the fresh solution and shaken for an additional 5 minutes. Then 25-50 ml of TE-4 (10 mM Tris-HCl pH 8.0; 0.1 mM EDTA pH 8.0) was added and the mixture incubated twice on a shaker for 5 minutes. The FTA Bloodstain Card was allowed to air dry completely during 2 hours at room temperature. The samples were then punched from the cards using a 3 mm diameter punch or the Harris Micro-Punch (1.2 mm or 2.0 mm), and transferred into corresponding microplate wells. PCR was then performed using the above-described procedure using regular PCR microplates and a ready-to-use buffer containing primers.

Patients (n=30, the age of 44-77) were divided into two groups. The first group of patients (n=12) were diagnosed with TIA in the carotid circulatory system, according to the following neurological criteria. Neural dysfunction was localized to a specific vascular distribution; the duration of the attack was usually less that 15 minutes and never exceeded 24 hours; and the patients did not have abnormal neurologic signs between attacks. The second pre-stroke group (n=18) were diagnosed with TIA in the vertebral-basilar circulatory system. The second group of patients was subdivided on the basis of compensation or non-compensation of neurological deficit. The third group (n=12) included patients with migraine and epilepsy.

The control group of healthy individuals (n=20) showed a level of NR2A cDNA expression of 1.2 0.11 pg/ml. The first group demonstrated slightly elevated levels of NR2A cDNA expression of 1.7 0.13 pg/ml. The patients with compensation of neurological deficit from the second group showed a level of NR2A cDNA expression of 1.8 1.4 pg/ml. At the same time, the patients without compensation of neurological deficit that possessed more severe symptoms of TIA showed levels of NR2A cDNA expression of 3 times the levels seen in healthy individuals. Patients suffering migraine and epilepsy

did not show any increase of NR2A cDNA expression when compared with the control group.

Example 10

Immunological Analysis of Blood Serum Specimens

Blood samples (10 ml, collected by venipuncture) from patients with cerebral ischemia (n=70), and healthy individuals (n=200), collected according to standard clinical protocols, were examined at the Neurology Hospital of Russian Medical Academy (Moscow, Russia). The blood specimens were centrifugated (4000 g, 5 min, +4° C.) and the collected serum stored at -70° C. for further analysis.

Computer analysis was employed to predict the antigenic determinants in the NR2A receptor protein structure based on hydrophobicity profile (Hopp, T. P. and K. R. Woods, *Proc. Natl. Acad. Sci. USA* 6: 3824-3828 (1981)) and antigenicity (Welling, G. W., et al., *FEBS Lett.* 188: 215-218 (1985)). Based upon this analysis, the N-terminal sequence of the NR2A NMDA receptor was synthesized. This synthetic peptide, which corresponded to amino acid sequence (494-514) (Grandy, D K., et al., *Proc. Natl. Acad. Sci. U.S.A.* 86: 9762-9766(1989) (SEQ ID NO:3) of human NR2A NMDA, was produced by solid-phase synthesis in a NPS-400 semi-automated synthesizer (Neosystem Lab, France) on MBHA resin using the BOC/Bzl strategy for the first two amino acids. The peptides were purified by preparative HPLC on a DELTAPAC™C18 column (Waters Chromatography, Milford, Mass.) in a H₂O/acetonitrile/0.015 TFA system. The purity of the peptides was determined by analytical HPLC and ranged from 90% to 98%. The peptide sequence was verified by amino acid analysis after acid hydrolysis. This peptide was used in immunoassays of blood serum from patients and healthy individuals.

A quantitative analysis of the level of NR2A autoantibodies in serum samples was performed by enzyme-linked immunosorbent assay (ELISA) (Ngo, T. T. and H. M. Lenhoff, *FEBS Lett.* 116: 285-288 (1980)). The diluted blood sera (1:50) and polyclonal antibodies to the NR2A peptide as a standard (0.01 ng/ml-400 ng/ml) were applied to the immunosorbent. The plate was incubated for 1 h at 25° C. and then washed by 0.05 M phosphate buffer, pH 7.4, containing 0.05% of TWEEN-20™. Rabbit antibodies to the human immunoglobulin labeled with horseradish peroxidase were added (Sigma, St. Louis, Mo.; 1:1000), and the plate was incubated for 1 h at 25° C. After incubation the wells were washed twice in the same buffer. The reaction was revealed by o-phenylenediamine in 0.05 M citrate buffer, pH 4.3 monitored at 490 nm on a microplate reader (BioRad, UK). The titer of NR2A autoantibodies in blood serum was determined by ELISA using a standard curve of the absorbance units of NR2A autoantibodies versus their concentration in a micro-titer well plate.

The synthetic peptide corresponding to the NR2A NMDA glutamate receptors (3 µg) were immobilized on a nitrocellulose membrane (0.45 µm, Shleicher-Shuell, Germany) in phosphate-buffered saline (PBS), pH 7.4, then washed 2-3 times in the same buffer. Membranes with immobilized peptide were incubated with the diluted serum (1:50) of cerebral ischemia patients and other subjects for 1 h at 25° C., and then rinsed 4 times with the PBS buffer. Secondary rabbit anti-human immunoglobulins conjugated with horseradish peroxidase (Sigma, St. Louis, Mo.; 1:1000) were incubated with the membrane for 1 h at 25° C., then washed 4 times with PBS. The development of brown color was registered and then quantitated by densitometry.

To provide a positive control or standard, rabbit polyclonal antibodies were raised against NR2A synthetic peptide corresponding to amino acid sequence predicted from the cloned human NR2A protein (Science 256: 1217-1221 (1992); SEQ ID NO:1). For glutaraldehyde conjugation, 10 mg of peptide and 40 mg of human serum albumin (Sigma, St. Louis, Mo.) were incubated for 1.45 h at room temperature in 4 ml of PBS containing 5% glutaraldehyde. The reaction was stopped by adding glycine to a final concentration of 0.2 M, and the conjugate was dialyzed against PBS. Rabbits were given initial injections of 1 mg of conjugated peptide in complete Freund's and subsequent injections of 0.5 mg of peptide in incomplete Freund's adjuvant at successive 2 week intervals. Antibodies were affinity purified according standard procedure (Warr, G. W., Purification of antibodies, In: *Antibody as a Tool*, Eds., Marchalonis, J. J., and G. W. Warr, J. Wiley, UK, pp. 59-96 (1982)) and were shown to be selective for the NR2A NMDA glutamate receptor using Western blot analysis.

The patients (men, n=30; women, n=40; age of 40-75) were admitted in the hospital within no more than six hours after the onset of an ischemic episode. All patients were divided into three groups according to the severity of the stroke: The first group had moderate ischemic stroke (n=25), manifested by moderate focal deficit (>60—Orgogozo scale). The second group had severe stroke (n=30), manifested by mild disorders of consciousness, severe headache, meningeal signs, and pronounced focal deficit (30-60—Orgogozo scale). The third group had extremely severe stroke (n=15), accompanied by stupor-coma, signs of brain edema, autonomic dysfunction, and severe focal deficit (<30—Orgogozo scale).

The level of NR2A autoantibodies was measured in the blood serum of healthy persons (n=200, age 35-75) as a control, and ranged from 0.3-1.5 ng/ml. The NR2A autoantibody level in the 55 patients of the first and second groups was significantly greater than that in the control group (p<0.01). Levels of NR2A autoantibodies were monitored every three hours during the first day, and then up to 5th day after stroke. The level of NR2A autoantibodies in the blood serum of patients with severe stroke was significantly higher than that in the blood serum of patients with moderate stroke, especially in the 9-12 hours after the onset of a stroke (p<0.05). The tendency for NR2A autoantibodies level to decrease to the control level on the first day of stroke was registered in group of patients with good neurological recovery (90, 50, 5 units on Orgogozo scale). It can be concluded

that the dynamic changes in NR2A autoantibodies level may predict a recovery period of patients after ischemic stroke.

Example 11

SPRIA Assay of Autoantibodies

The solid-phase radioimmunoassay (SPRIA) of autoantibodies is performed as follows: a 10% acetic acid solution is added for one minute to the Cooker microtiter microplates (available from Dynatech Co., USA) for activation, whereupon 0.1 ml of the blood serum under analysis (diluted 1:40) is applied to the microplates and subjected to incubation for four hours at 25° C. Then the microplate are washed with a 0.14 M sodium chloride solution and 0.1 ml of a mixture of the respective fragment of the mammal's brain protein labeled by 125I in the presence of nonlabelled one. The plates are incubated for 20 hours at 4° C. On completion of incubation, the microplates are washed with a 0.14M sodium chloride solution, after which each of the wells of the microplates is cut off and placed in gamma-counting vials.

Example 12

ELISA Assay of Autoantibodies

The enzyme-linked immunosorbent assay (ELISA) of autoantibodies is carried out as follows: the samples of the blood serum diluted 1:40 or 1:50 are applied to the respective immunosorbent. Then the plate carrying the immunosorbent is incubated for 30 min at 37° C., whereupon the wells of the plate are washed with a 0.05 M phosphate buffer, containing 0.05% of Tween-20. Rabbit antibodies to human immunoglobulin labeled with horseradish peroxidase (conjugate) are added thereto, and the plate is reincubated for 35 min at 37° C., then washed by the aforementioned buffer and distilled water. The reaction with conjugate is determined by adding chromogen, i.e., orthophenylenediamine in the presence of 30% hydrogen peroxide. The intensity of color development is evaluated by using the rider (available Multiskan microplate rider) at the 492 nm wavelength.

It will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the scope or spirit of the invention. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 17

<210> SEQ ID NO 1

<211> LENGTH: 1464

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Met Gly Arg Leu Gly Tyr Trp Thr Leu Leu Val Leu Pro Ala Leu Leu
1 5 10 15

Val Trp Arg Asp Pro Ala Gln Asn Ala Ala Ala Glu Lys Gly Pro Pro
20 25 30

-continued

Ala Leu Asn Ile Ala Val Leu Leu Gly His Ser His Asp Val Thr Glu
35 40 45

Arg Glu Leu Arg Asn Leu Trp Gly Pro Glu Gln Ala Thr Gly Leu Pro
50 55 60

Leu Asp Val Asn Val Val Ala Leu Leu Met Asn Arg Thr Asp Pro Lys
65 70 75 80

Ser Leu Ile Thr His Val Cys Asp Leu Met Ser Gly Ala Arg Ile His
85 90 95

Gly Leu Val Phe Gly Asp Asp Thr Asp Gln Glu Ala Val Ala Gln Met
100 105 110

Leu Asp Phe Ile Ser Ser Gln Thr Phe Ile Pro Ile Leu Gly Ile His
115 120 125

Gly Gly Ala Ser Met Ile Met Ala Asp Lys Asp Pro Thr Ser Thr Phe
130 135 140

Phe Gln Phe Gly Ala Ser Ile Gln Gln Gln Ala Thr Val Met Leu Lys
145 150 155 160

Ile Met Gln Asp Tyr Asp Trp His Val Phe Ser Leu Val Thr Thr Ile
165 170 175

Phe Pro Gly Tyr Arg Asp Phe Ile Ser Phe Ile Lys Thr Thr Val Asp
180 185 190

Asn Ser Phe Val Gly Trp Asp Met Gln Asn Val Ile Thr Leu Asp Thr
195 200 205

Ser Phe Glu Asp Ala Lys Thr Gln Val Gln Leu Lys Lys Ile His Ser
210 215 220

Ser Val Ile Leu Leu Tyr Cys Ser Lys Asp Glu Ala Val Leu Ile Leu
225 230 235 240

Ser Glu Ala Arg Ser Leu Gly Leu Thr Gly Tyr Asp Phe Phe Trp Ile
245 250 255

Val Pro Ser Leu Val Ser Gly Asn Thr Glu Leu Ile Pro Lys Glu Phe
260 265 270

Pro Ser Gly Leu Ile Ser Val Ser Tyr Asp Asp Trp Asp Tyr Ser Leu
275 280 285

Glu Ala Arg Val Arg Asp Gly Leu Gly Ile Leu Thr Thr Ala Ala Ser
290 295 300

Ser Met Leu Glu Lys Phe Ser Tyr Ile Pro Glu Ala Lys Ala Ser Cys
305 310 315 320

Tyr Gly Gln Ala Glu Lys Pro Glu Thr Pro Leu His Thr Leu His Gln
325 330 335

Phe Met Val Asn Val Thr Trp Asp Gly Lys Asp Leu Ser Phe Thr Glu
340 345 350

Glu Gly Tyr Gln Val His Pro Arg Leu Val Val Ile Val Leu Asn Lys
355 360 365

Asp Arg Glu Trp Glu Lys Val Gly Lys Trp Glu Asn Gln Thr Leu Ser
370 375 380

Leu Arg His Ala Val Trp Pro Arg Tyr Lys Ser Phe Ser Asp Cys Glu
385 390 395 400

Pro Asp Asp Asn His Leu Ser Ile Val Thr Leu Glu Glu Ala Pro Phe
405 410 415

Val Ile Val Glu Asp Ile Asp Pro Leu Thr Glu Thr Cys Val Arg Asn
420 425 430

Thr Val Pro Cys Arg Lys Phe Val Lys Ile Asn Asn Ser Thr Asn Glu
435 440 445

-continued

Gly Met Asn Val Lys Lys Cys Cys Lys Gly Phe Cys Ile Asp Ile Leu
 450 455 460

Lys Lys Leu Ser Arg Thr Val Lys Phe Thr Tyr Asp Leu Tyr Leu Val
 465 470 475 480

Thr Asn Gly Lys His Gly Lys Lys Val Asn Asn Val Trp Asn Gly Met
 485 490 495

Ile Gly Glu Val Val Tyr Gln Arg Ala Val Met Ala Val Gly Ser Leu
 500 505 510

Thr Ile Asn Glu Glu Arg Ser Glu Val Val Asp Phe Ser Val Pro Phe
 515 520 525

Val Glu Thr Gly Ile Ser Val Met Val Ser Arg Ser Asn Gly Thr Val
 530 535 540

Ser Pro Ser Ala Phe Leu Glu Pro Phe Ser Ala Ser Val Trp Val Met
 545 550 555 560

Met Phe Val Met Leu Leu Ile Val Ser Ala Ile Ala Val Phe Val Phe
 565 570 575

Glu Tyr Phe Ser Pro Val Gly Tyr Asn Arg Asn Leu Ala Lys Gly Lys
 580 585 590

Ala Pro His Gly Pro Ser Phe Thr Ile Gly Lys Ala Ile Trp Leu Leu
 595 600 605

Trp Gly Leu Val Phe Asn Asn Ser Val Pro Val Gln Asn Pro Lys Gly
 610 615 620

Thr Thr Ser Lys Ile Met Val Ser Val Trp Ala Phe Phe Ala Val Ile
 625 630 635 640

Phe Leu Ala Ser Tyr Thr Ala Asn Leu Ala Ala Phe Met Ile Gln Glu
 645 650 655

Glu Phe Val Asp Gln Val Thr Gly Leu Ser Asp Lys Lys Phe Gln Arg
 660 665 670

Pro His Asp Tyr Ser Pro Pro Phe Arg Phe Gly Thr Val Pro Asn Gly
 675 680 685

Ser Thr Glu Arg Asn Ile Arg Asn Asn Tyr Pro Tyr Met His Gln Tyr
 690 695 700

Met Thr Arg Phe Asn Gln Arg Gly Val Glu Asp Ala Leu Val Ser Leu
 705 710 715 720

Lys Thr Gly Lys Leu Asp Ala Phe Ile Tyr Asp Ala Ala Val Leu Asn
 725 730 735

Tyr Lys Ala Gly Arg Asp Glu Gly Cys Lys Leu Val Thr Ile Gly Ser
 740 745 750

Gly Tyr Ile Phe Ala Ser Thr Gly Tyr Gly Ile Ala Leu Gln Lys Gly
 755 760 765

Ser Pro Trp Lys Arg Gln Ile Asp Leu Ala Leu Leu Gln Phe Val Gly
 770 775 780

Asp Gly Glu Met Glu Glu Leu Glu Thr Leu Trp Leu Thr Gly Ile Cys
 785 790 795 800

His Asn Glu Lys Asn Glu Val Met Ser Ser Gln Leu Asp Ile Asp Asn
 805 810 815

Met Ala Gly Val Phe Tyr Met Leu Ala Ala Ala Met Ala Leu Ser Leu
 820 825 830

Ile Thr Phe Ile Trp Glu His Leu Phe Tyr Trp Lys Leu Arg Phe Cys
 835 840 845

Phe Thr Gly Val Cys Ser Asp Arg Pro Gly Leu Leu Phe Ser Ile Ser
 850 855 860

Arg Gly Ile Tyr Ser Cys Ile His Gly Val His Ile Glu Glu Lys Lys

-continued

865	870	875	880
Lys Ser Pro Asp Phe Asn Leu Thr Gly Ser Gln Ser Asn Met Leu Lys	885	890	895
Leu Leu Arg Ser Ala Lys Asn Ile Ser Asn Met Ser Asn Met Asn Ser	900	905	910
Ser Arg Met Asp Ser Pro Lys Arg Ala Thr Asp Phe Ile Gln Arg Gly	915	920	925
Ser Leu Ile Val Asp Met Val Ser Asp Lys Gly Asn Leu Ile Tyr Ser	930	935	940
Asp Asn Arg Ser Phe Gln Gly Lys Asp Ser Ile Phe Gly Asp Asn Met	945	950	955
Asn Glu Leu Gln Thr Phe Val Ala Asn Arg His Lys Asp Asn Leu Ser	965	970	975
Asn Tyr Val Phe Gln Gly Gln His Pro Leu Thr Leu Asn Glu Ser Asn	980	985	990
Pro Asn Thr Val Glu Val Ala Val Ser Thr Glu Ser Lys Gly Asn Ser	995	1000	1005
Arg Pro Arg Gln Leu Trp Lys Lys Ser Met Glu Ser Leu Arg Gln	1010	1015	1020
Asp Ser Leu Asn Gln Asn Pro Val Ser Gln Arg Asp Glu Lys Thr	1025	1030	1035
Ala Glu Asn Arg Thr His Ser Leu Lys Ser Pro Arg Tyr Leu Pro	1040	1045	1050
Glu Glu Val Ala His Ser Asp Ile Ser Glu Thr Ser Ser Arg Ala	1055	1060	1065
Thr Cys His Arg Glu Pro Asp Asn Asn Lys Asn His Lys Thr Lys	1070	1075	1080
Asp Asn Phe Lys Arg Ser Met Ala Ser Lys Tyr Pro Lys Asp Cys	1085	1090	1095
Ser Asp Val Asp Arg Thr Tyr Met Lys Thr Lys Ala Ser Ser Pro	1100	1105	1110
Arg Asp Lys Ile Tyr Thr Ile Asp Gly Glu Lys Glu Pro Ser Phe	1115	1120	1125
His Leu Asp Pro Pro Gln Phe Val Glu Asn Ile Thr Leu Pro Glu	1130	1135	1140
Asn Val Gly Phe Pro Asp Thr Tyr Gln Asp His Asn Glu Asn Phe	1145	1150	1155
Arg Lys Gly Asp Ser Thr Leu Pro Met Asn Arg Asn Pro Leu His	1160	1165	1170
Asn Glu Asp Gly Leu Pro Asn Asn Asp Gln Tyr Lys Leu Tyr Ala	1175	1180	1185
Lys His Phe Thr Leu Lys Asp Lys Gly Ser Pro His Ser Glu Gly	1190	1195	1200
Ser Asp Arg Tyr Arg Gln Asn Ser Thr His Cys Arg Ser Cys Leu	1205	1210	1215
Ser Asn Leu Pro Thr Tyr Ser Gly His Phe Thr Met Arg Ser Pro	1220	1225	1230
Phe Lys Cys Asp Ala Cys Leu Arg Met Gly Asn Leu Tyr Asp Ile	1235	1240	1245
Asp Glu Asp Gln Met Leu Gln Glu Thr Gly Asn Pro Ala Thr Arg	1250	1255	1260
Glu Glu Val Tyr Gln Gln Asp Trp Ser Gln Asn Asn Ala Leu Gln	1265	1270	1275

-continued

Phe Gln Lys Asn Lys Leu Arg Ile Asn Arg Gln His Ser Tyr Asp
 1280 1285 1290

Asn Ile Leu Asp Lys Pro Arg Glu Ile Asp Leu Ser Arg Pro Ser
 1295 1300 1305

Arg Ser Ile Ser Leu Lys Asp Arg Glu Arg Leu Leu Glu Gly Asn
 1310 1315 1320

Leu Tyr Gly Ser Leu Phe Ser Val Pro Ser Ser Lys Leu Leu Gly
 1325 1330 1335

Asn Lys Ser Ser Leu Phe Pro Gln Gly Leu Glu Asp Ser Lys Arg
 1340 1345 1350

Ser Lys Ser Leu Leu Pro Asp His Ala Ser Asp Asn Pro Phe Leu
 1355 1360 1365

His Thr Tyr Gly Asp Asp Gln Arg Leu Val Ile Gly Arg Cys Pro
 1370 1375 1380

Ser Asp Pro Tyr Lys His Ser Leu Pro Ser Gln Ala Val Asn Asp
 1385 1390 1395

Ser Tyr Leu Arg Ser Ser Leu Arg Ser Thr Ala Ser Tyr Cys Ser
 1400 1405 1410

Arg Asp Ser Arg Gly His Ser Asp Val Tyr Ile Ser Glu His Val
 1415 1420 1425

Met Pro Tyr Ala Ala Asn Lys Asn Thr Met Tyr Ser Thr Pro Arg
 1430 1435 1440

Val Leu Asn Ser Cys Ser Asn Arg Arg Val Tyr Lys Lys Met Pro
 1445 1450 1455

Ser Ile Glu Ser Asp Val
 1460

<210> SEQ ID NO 2
 <211> LENGTH: 536
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Pro Ala Gln Asn Ala Ala Ala Glu Lys Gly Pro Pro Ala Leu Asn Ile
 1 5 10 15

Ala Val Leu Leu Gly His Ser His Asp Val Thr Glu Arg Glu Leu Arg
 20 25 30

Asn Leu Trp Gly Pro Glu Gln Ala Thr Gly Leu Pro Leu Asp Val Asn
 35 40 45

Val Val Ala Leu Leu Met Asn Arg Thr Asp Pro Lys Ser Leu Ile Thr
 50 55 60

His Val Cys Asp Leu Met Ser Gly Ala Arg Ile His Gly Leu Val Phe
 65 70 75 80

Gly Asp Asp Thr Asp Gln Glu Ala Val Ala Gln Met Leu Asp Phe Ile
 85 90 95

Ser Ser Gln Thr Phe Ile Pro Ile Leu Gly Ile His Gly Gly Ala Ser
 100 105 110

Met Ile Met Ala Asp Lys Asp Pro Thr Ser Thr Phe Phe Gln Phe Gly
 115 120 125

Ala Ser Ile Gln Gln Gln Ala Thr Val Met Leu Lys Ile Met Gln Asp
 130 135 140

Tyr Asp Trp His Val Phe Ser Leu Val Thr Thr Ile Phe Pro Gly Tyr
 145 150 155 160

Arg Asp Phe Ile Ser Phe Ile Lys Thr Thr Val Asp Asn Ser Phe Val

-continued

165				170				175							
Gly	Trp	Asp	Met	Gln	Asn	Val	Ile	Thr	Leu	Asp	Thr	Ser	Phe	Glu	Asp
			180								185				190
Ala	Lys	Thr	Gln	Val	Gln	Leu	Lys	Lys	Ile	His	Ser	Ser	Val	Ile	Leu
			195								200				205
Leu	Tyr	Cys	Ser	Lys	Asp	Glu	Ala	Val	Leu	Ile	Leu	Ser	Glu	Ala	Arg
			210				215								220
Ser	Leu	Gly	Leu	Thr	Gly	Tyr	Asp	Phe	Phe	Trp	Ile	Val	Pro	Ser	Leu
							230				235				240
Val	Ser	Gly	Asn	Thr	Glu	Leu	Ile	Pro	Lys	Glu	Phe	Pro	Ser	Gly	Leu
			245								250				255
Ile	Ser	Val	Ser	Tyr	Asp	Asp	Trp	Asp	Tyr	Ser	Leu	Glu	Ala	Arg	Val
			260								265				270
Arg	Asp	Gly	Leu	Gly	Ile	Leu	Thr	Thr	Ala	Ala	Ser	Ser	Met	Leu	Glu
			275				280								285
Lys	Phe	Ser	Tyr	Ile	Pro	Glu	Ala	Lys	Ala	Ser	Cys	Tyr	Gly	Gln	Ala
			290				295				300				
Glu	Lys	Pro	Glu	Thr	Pro	Leu	His	Thr	Leu	His	Gln	Phe	Met	Val	Asn
			305				310				315				320
Val	Thr	Trp	Asp	Gly	Lys	Asp	Leu	Ser	Phe	Thr	Glu	Glu	Gly	Tyr	Gln
			325								330				335
Val	His	Pro	Arg	Leu	Val	Val	Ile	Val	Leu	Asn	Lys	Asp	Arg	Glu	Trp
			340								345				350
Glu	Lys	Val	Gly	Lys	Trp	Glu	Asn	Gln	Thr	Leu	Ser	Leu	Arg	His	Ala
			355				360								365
Val	Trp	Pro	Arg	Tyr	Lys	Ser	Phe	Ser	Asp	Cys	Glu	Pro	Asp	Asp	Asn
			370				375				380				
His	Leu	Ser	Ile	Val	Thr	Leu	Glu	Glu	Ala	Pro	Phe	Val	Ile	Val	Glu
			385				390				395				400
Asp	Ile	Asp	Pro	Leu	Thr	Glu	Thr	Cys	Val	Arg	Asn	Thr	Val	Pro	Cys
			405								410				415
Arg	Lys	Phe	Val	Lys	Ile	Asn	Asn	Ser	Thr	Asn	Glu	Gly	Met	Asn	Val
			420								425				430
Lys	Lys	Cys	Cys	Lys	Gly	Phe	Cys	Ile	Asp	Ile	Leu	Lys	Lys	Leu	Ser
			435				440								445
Arg	Thr	Val	Lys	Phe	Thr	Tyr	Asp	Leu	Tyr	Leu	Val	Thr	Asn	Gly	Lys
			450				455				460				
His	Gly	Lys	Lys	Val	Asn	Asn	Val	Trp	Asn	Gly	Met	Ile	Gly	Glu	Val
			465				470				475				480
Val	Tyr	Gln	Arg	Ala	Val	Met	Ala	Val	Gly	Ser	Leu	Thr	Ile	Asn	Glu
			485								490				495
Glu	Arg	Ser	Glu	Val	Val	Asp	Phe	Ser	Val	Pro	Phe	Val	Glu	Thr	Gly
			500								505				510
Ile	Ser	Val	Met	Val	Ser	Arg	Ser	Asn	Gly	Thr	Val	Ser	Pro	Ser	Ala
			515								520				525
Phe	Leu	Glu	Pro	Phe	Ser	Ala	Ser								
			530				535								

<210> SEQ ID NO 3

<211> LENGTH: 21

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

-continued

Asn Gly Met Ile Gly Glu Val Val Tyr Gln Arg Ala Val Met Ala Val
1 5 10 15

Gly Ser Leu Thr Ile
 20

<210> SEQ ID NO 4
<211> LENGTH: 22
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: modified amino acid sequence

<400> SEQUENCE: 4

Cys Asn Gly Met Ile Gly Glu Val Val Tyr Gln Arg Ala Val Met Ala
1 5 10 15

Val Gly Ser Leu Thr Ile
 20

<210> SEQ ID NO 5
<211> LENGTH: 6293
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

atcatgggac cgggtgagcg ctgagaatcg cggccgcagc catcagccct ggagatgacc 60
aggagcggcc actgctgaga actatgtgga gagaggctgc gagccctgct gcagagcctc 120
cggctgggat agccgcccc cgtgggggcg atgcccagc cgcgggacag ccaggggagc 180
gcgctggggc cgcagcatgc gggaacccgc taaacccggt ggctgctgag gcggccgaga 240
tgctcgtgcg cgcagcgcgc cccactgcat cctcgacctt ctgggctac agggaccgtc 300
agtggcgact atgggcagag tgggctattg gaccctgctg gtgctgccgg cccttctggt 360
ctggcgcggt cgggcgccga gcgcggcggc ggagaagggt cccccgcgc taaatattgc 420
ggtgatgctg ggtcacagcc acgacgtgac agagcgcgaa cttcgaacac tgtggggccc 480
cgagcaggcg gcggggctgc ccctggacgt gaacgtggtg gctctgctga tgaaccgcac 540
cgacccaag agcctcatca cgcacgtgtg cgacctcatg tccggggcac gcatccacgg 600
cctcgtgttt ggggacgaca cggaccagga ggccgtagcc cagatgctgg attttatctc 660
ctcccacacc ttcgtcccca tcttgggcat tcatgggggc gcatctatga tcatggctga 720
caaggatccg acgtctacct tcttcagtt tggagcgtcc atccagcagc aagccacggt 780
catgctgaag atcatgcagg attatgactg gcatgtcttc tccctggtga ccactatctt 840
ccctggctac agggaattca tcagcttctg caagaccaca gtggacaaca gctttgtggg 900
ctgggacatg cagaatgtga tcacactgga cacttccttt gaggatgcaa agacacaagt 960
ccagctgaag aagatccact cttctgtcat cttgctctac tgttccaaag acgaggtgt 1020
tctcattctg agtgaggccc gctcccttgg cctcaccggg tatgatttct tctggattgt 1080
ccccagcttg gtctctggga acacggagct catcccacaaa gagtttccat cgggactcat 1140
ttctgtctcc tacgatgact gggactacag cctggaggcg agagtgaggg acggcattgg 1200
cactctaacc accgctgcat cttctatgct ggagaagttc tctacatcc ccgaggccaa 1260
ggccagctgc tacgggcaga tggagaggcc agaggtcccg atgcacacct tgcaccatt 1320
tatggtcaat gttacatggg atggcaaaga cttatccttc actgaggaag gctaccaggt 1380
gcaccccagg ctggtggtga ttgtgctgaa caaagaccgg gaatgggaaa aggtgggcaa 1440
gtgggagaac catacgtgta gcctgaggca cgcctgtgtg cccaggtaca agtccttctc 1500

-continued

cgactgtgag	ccggatgaca	accatctcag	catcgtcacc	ctggaggagg	ccccattcgt	1560
catcgtggaa	gacatagacc	ccctgaccga	gacgtgtgtg	aggaacaccg	tgccatgtcg	1620
gaagtctgtc	aaaatcaaca	attcaaccaa	tgaggggatg	aatgtgaaga	aatgctgcaa	1680
ggggttctgc	attgatattc	tgaagaagct	ttccagaact	gtgaagtta	cttacgacct	1740
ctatctggtg	accaatggga	agcatggcaa	gaaagttaac	aatgtgtgga	atggaatgat	1800
cggatgaagt	gtctatcaac	gggcagtcac	ggcagttggc	tcgctcacca	tcaatgagga	1860
acgttctgaa	gtggtggaat	tctctgtgcc	ctttgtggaa	acgggaatca	gtgtcatggt	1920
ttcaagaagt	aatggcaccg	tctcaccttc	tgcttttcta	gaaccattca	gcgccctctgt	1980
ctgggtgatg	atgtttgtga	tgctgtctcat	tgtttctgcc	atagctgttt	ttgtctttga	2040
atacttcagc	cctgttggat	acaacagaaa	cttagccaaa	gggaaagcac	cccatgggcc	2100
ttcttttaca	attgaaaaag	ctatatggct	tctttggggc	ctggtgttca	ataactccgt	2160
gcctgtccag	aatcctaaa	ggaccaccag	caagatcatg	gtatctgtat	gggccttctt	2220
cgctgtcata	ttcctggcta	gctacacagc	caatctggct	gccttcatga	tccaagagga	2280
atgtgtggac	caagtgaccg	gcctcagtga	caaaaagttt	cagagacctc	atgactattc	2340
cccacctttt	cgatttggga	cagtgccctaa	tggaagcacg	gagagaaaca	ttcggaataa	2400
ctatccctac	atgcatcagt	acatgaccaa	atttaacacg	aaaggagtag	aggacgcctt	2460
ggctcagcctg	aaaacgggga	agctggacgc	tttcatctac	gatgccgcag	tcttgaatta	2520
caaggctggg	agggatgaag	gctgcaagct	ggtgaccatc	gggagtgggt	acatctttgc	2580
caccaccggg	tatggaattg	cccttcagaa	aggctctcct	tggaagaggc	agatcgacct	2640
ggccttgctt	cagtttgtgg	gtgatggtga	gatggaggag	ctggagacc	tgtggctcac	2700
tgggatctgc	cacaacgaga	agaacgaggt	gatgagcagc	cagctggaca	ttgacaacat	2760
ggcggggcgt	ttctacatgc	tggtgcccgc	catggccctt	agcctcatca	ccttcatctg	2820
ggagcacctc	ttctactgga	agctgcgctt	ctgtttcacg	ggcgtgtgct	ccgaccggcc	2880
tgggttgcct	ttctccatca	gcaggggcat	ctacagctgc	attcatggag	tgcaattga	2940
agaaaagaag	aagtctccag	actcaaatct	gacgggatcc	cagagcaaca	tgtaaaaact	3000
cctccggtca	gccccaaaa	tttccagcat	gtccaacatg	aactcctcaa	gaatggactc	3060
acccaaaaga	gctgctgact	tcatocaaag	aggttccttc	atcatggaca	tggtttcaga	3120
taaggggaat	ttgatgtact	cagacaacag	gtcctttcag	gggaaagaga	gcatttttgg	3180
agacaacatg	aacgaactcc	aaacatttgt	ggccaaccgg	cagaagata	acctcaataa	3240
ctatgtattc	cagggacaac	atcctcttac	tctcaatgag	tccaacccta	acacggtgga	3300
ggtggccgtg	agcacagaat	ccaaagcgaa	ctctagacc	cggcagctgt	ggaagaaatc	3360
cgtggattcc	atacgcagag	attcactatc	ccagaatcca	gtctcccaga	gggatgaggc	3420
aacagcagag	aataggacc	actccctaaa	gagccctagg	tatcttcag	aagagatggc	3480
ccactctgac	atttcagaaa	cgtcaaatcg	ggccacgtgc	cacagggaac	ctgacaacag	3540
taagaaccac	aaaaccaagg	acaacttta	aaggtcagtg	gcctccaaat	acccaagga	3600
ctgtagtgag	gtcagcgcga	cctacctgaa	aaccaaatca	agctccccta	gagacaagat	3660
ctacactata	gatggtgaga	aggagcctgg	tttccactta	gatccacccc	agtttgttga	3720
aaatgtgacc	ctgcccagaga	acgtggactt	cccggacccc	taccaggatc	ccagtgaaaa	3780
cttccgcaag	ggggactcca	cgctgccaat	gaaccggaac	cccttgcata	atgaagaggg	3840

-continued

gctttccaac aacgaccagt ataaactcta ctccaagcac ttcaccttga aagacaaggg	3900
tccccgcac agtgagacca gcgagcgata cggcgagaac tccacgcact gcagaagctg	3960
cctttccaac atgcccacct attcaggcca cttcaccatg aggtcccct tcaagtgcga	4020
tgctgctg cggatgggga acctctatga catcgatgaa gaccagatgc ttcaggagac	4080
aggtaaccca gccaccgggg agcaggctta ccagcaggac tgggcacaga acaatgcct	4140
tcaattacaa aagaacaagc taaggattag ccgtcagcat tctacgata acattgtcga	4200
caaacctagg gagctagacc ttagcaggcc ctcccggagc ataagcctca aggacaggga	4260
acggcttctg gagggaaatt tttacggcag cctgtttagt gtcccctcaa gcaaaacttc	4320
ggggaaaaaa agctcccttt tcccccaagg tctggaggac agcaagagga gcaagtctct	4380
cttgccagac cacacctccg ataacccttt cctccactcc cacagggatg accaacgctt	4440
ggttattggg agatgcccc cggaccctta caaacactcg ttgccatccc aggcggtgaa	4500
tgacagctat ctccggtcgt ccttgaggtc aacggcatcg tactgttcca gggacagctg	4560
gggccacaat gatgtgata tttcggagca tgttatgect tatgtgcaa ataagaataa	4620
tatgtactct acccccaggg ttttaaattc ctgcagcaat agacgcgtgt acaagaaaat	4680
gcctagtatc gaatctgatg tttaaaaatc ttccattaat gttttatcta tagggaaata	4740
cacgtaatgg ccaatgttct ggagggtaaa tgttgatgt ccaatagtgc cctgctaaga	4800
ggaagaagat gtagggaggt attttgtgt tgttgttgtt ggctcttttg cacacggctt	4860
catgccataa tcttcactc aaggaatcct gtgaggtgtg tgcagcat gccagacacc	4920
agataggta gtccttaacc aaaaataact aactacataa gggcaagtct cgggacatg	4980
cctactgggt atgttgcaa taatgatgca ttggatgcca atggtgatgt tatgatttcc	5040
tatattccaa attccattaa ggtcagccca ccatgtaatt ttctcatcag aaatgcctaa	5100
tggtttctct aatacagaat aagcaatatg gtgtgcatgt aaacctgaca cagacaaaat	5160
aaaaacagtt aagaatgcat ctgcactgta gtcggatttg aacatgtgca agagattagg	5220
aagtttgct cgtaacagtt tcagctttct tgttatgect tccatcacag cccaggctca	5280
ccccagaac tccaggctcc cctaaagaat agcaaatcag tgtgttcgtg atgactgtgc	5340
taccttcatt atagttcatt tccaagacac atctggagcc aaaggcccga gggaccctca	5400
ggtggggaga gctacagaa tctctttgga tgttgatgtg tgtttctctc taccctcggc	5460
ttcgatggtc ttgttcagag ctgcataaac taacacattt atgtctccga gatctaagtg	5520
tggatcttct gctctgaca cagtggccat tgtagtttat cccgaagacg cctatgtacg	5580
taagtttga tttctcct tctggatg actcagggtt gtatagtatc tgttaccct	5640
tcctcccag agtaaccata actcgttccg tttcaaaca gccatggtgg tgtccaatta	5700
gctgtgtatc gctcttcca gagttgtaa tgtggtgaca tgcaccaaca gccgatgtg	5760
tactgtgatc tgtaagaagt acaatgcat ctgtctgccc aaggtagca tggttttagg	5820
tttatcttcc ttcacatca gaaattctgt tggacactca ctccacccc aaactcctca	5880
aatcaaaagc cttcaaaaca cgaggcactc ttggatctac cctgagtatc ctccaaactg	5940
tggatacagt ttagtgagac aagcaatttc tcccttctga gttattctct ctgttggtgg	6000
caaacactt catagcacca acagagatgt aggaaaaatt cctcaaagta tttgtcattt	6060
ctgagtcgcc tgcattatcc cattcttatt ctctcaaac ctgtgcatat atgacatgaa	6120
atgatatcca tttttttttt aagttagaaa cagagagggg aatacttatg catggggagc	6180
ctgttagcac agtgctgcc acaaaaacaa gtgccccga caagatagtt gctatgttat	6240

-continued

gacactttct cagatcagga ttttctagtt taaaaattaa atatcataaa acg 6293

<210> SEQ ID NO 6
 <211> LENGTH: 1608
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

ccggcgccga gcgcgccggc ggagaagggc ccccccgcgc taaatattgc ggtgatgctg 60
 ggtcacagcc acgacgtgac agagcgcgaa cttcgaacac tgtggggccc cgagcaggcg 120
 gcggggctgc ccttgacgt gaacgtggta gctctgctga tgaaccgcac cgacccaag 180
 agcctcatca cgcacgtgtg cgacctcatg tccggggcac gcataccagg cctcgtgttt 240
 ggggacgaca cggaccagga ggccgtagcc cagatgctgg atttatctc ctcccacacc 300
 ttcgtcccca tcttgggcat tcatgggggc gcatactatga tcatggctga caaggatccg 360
 acgtctacct tcttccagtt tggagcgtcc atccagcagc aagccacggt catgctgaag 420
 atcatgcagg attatgactg gcatagtctc tccttggtga ccaactatctt ccttgctac 480
 agggaattca tcagcttctg caagaccaca gtggacaaca gctttgtggg ctgggacatg 540
 cagaatgtga tcactctgga cacttccttt gaggatgcaa agacacaagt ccagctgaag 600
 aagatccact cttctgtcat cttgctctac tgttccaaag acgaggetgt tetcattctg 660
 agtgaggccc gtcctcttgg cctcaccggg tatgatttct tctggattgt cccagcttg 720
 gtctctggga acacggagct catcccaaaa gagtttccat cgggactcat ttctgtctcc 780
 tacgatgact gggactacag cctggaggcg agagtgagg acggcattgg catcctaacc 840
 accgctgcat cttctatgct ggagaagtcc tcctacatcc ccgaggccaa ggcagctgc 900
 tacgggcaga tggagaggcc agaggtccc atgcacacct tgcacccatt tatggtcaat 960
 gttacatggg atggcaaaaga cttatccttc actgaggaag gctaccaggt gcaccccagg 1020
 ctggtggtga ttgtgctgaa caaagaccgg gaatgggaaa aggtgggcaa gtgggagAAC 1080
 catacgtga gcttagggca cgcctgtgtg cccaggtaca agtccttctc cgactgtgag 1140
 ccggatgaca accatctcag catcgtcacc ctggaggagg cccattcgt catcgtggaa 1200
 gacatagacc ccttagccga gacgtgtgtg aggaacaccg tgccatgtcg gaagttcgtc 1260
 aaaatcaaca attcaaccaa tgaggggatg aatgtgaaga aatgctgcaa ggggttctgc 1320
 attgatattc tgaagaagct ttccagaact gtgaagtta cttacgacct ctatctggtg 1380
 accaatggga agcattggca gaaagttaac aatgtgtgga atggaatgat cggtgaagtg 1440
 gtctatcaac gggcagtcac ggcagttggc tcgctcacca tcaatgagga acgttctgaa 1500
 gtggtgact tctctgtgcc ctttgtggaa acgggaatca gtgtcatggt tcaagaagt 1560
 aatggcaccg tctcaacctc tgcctttota gaaccattca gcgcctct 1608

<210> SEQ ID NO 7
 <211> LENGTH: 62
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

atggaatgat cggatgaagtg gtctatcaac gggcagtcac ggcagttggc tcgctcacca 60
 tc 62

<210> SEQ ID NO 8

-continued

<211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: oligonucleotide primer

 <400> SEQUENCE: 8

 agcatggcaa gaaagttaac a 21

<210> SEQ ID NO 9
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: oligonucleotide primer

 <400> SEQUENCE: 9

 acgttctgaa gtggtggact t 21

<210> SEQ ID NO 10
 <211> LENGTH: 1480
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 10

 Met Lys Pro Arg Ala Glu Cys Cys Ser Pro Lys Phe Trp Leu Val Leu
 1 5 10 15

 Ala Val Leu Ala Val Ser Gly Ser Arg Ala Arg Ser Gln Lys Ser Pro
 20 25 30

 Pro Ser Ile Gly Ile Ala Val Ile Leu Val Gly Thr Ser Asp Glu Val
 35 40 45

 Ala Ile Lys Asp Ala His Glu Lys Asp Asp Phe His His Leu Ser Val
 50 55 60

 Val Pro Arg Val Glu Leu Val Ala Met Asn Glu Thr Asp Pro Lys Ser
 65 70 75 80

 Ile Ile Thr Arg Ile Cys Asp Leu Met Ser Asp Arg Lys Ile Gln Gly
 85 90 95

 Val Val Phe Ala Asp Asp Thr Asp Gln Glu Ala Ile Ala Gln Ile Leu
 100 105 110

 Asp Phe Ile Ser Ala Gln Thr Leu Thr Pro Ile Leu Gly Ile His Gly
 115 120 125

 Gly Ser Ser Met Ile Met Ala Asp Lys Asp Glu Ser Ser Met Phe Phe
 130 135 140

 Gln Phe Gly Pro Ser Ile Glu Gln Gln Ala Ser Val Met Leu Asn Ile
 145 150 155 160

 Met Glu Glu Tyr Asp Trp Tyr Ile Phe Ser Ile Val Thr Thr Tyr Phe
 165 170 175

 Pro Gly Tyr Gln Asp Phe Val Asn Lys Ile Arg Ser Thr Ile Glu Asn
 180 185 190

 Ser Phe Val Gly Trp Glu Leu Glu Glu Val Leu Leu Leu Asp Met Ser
 195 200 205

 Leu Asp Asp Gly Asp Ser Lys Ile Gln Asn Gln Leu Lys Lys Leu Gln
 210 215 220

 Ser Pro Ile Ile Leu Leu Tyr Cys Thr Lys Glu Glu Ala Thr Tyr Ile
 225 230 235 240

 Phe Glu Val Ala Asn Ser Val Gly Leu Thr Gly Tyr Gly Tyr Thr Trp
 245 250 255

 Ile Val Pro Ser Leu Val Ala Gly Asp Thr Asp Thr Val Pro Ala Glu

-continued

260					265					270					
Phe	Pro	Thr	Gly	Leu	Ile	Ser	Val	Ser	Tyr	Asp	Glu	Trp	Asp	Tyr	Gly
	275						280					285			
Leu	Pro	Ala	Arg	Val	Arg	Asp	Gly	Ile	Ala	Ile	Ile	Thr	Thr	Ala	Ala
	290					295					300				
Ser	Asp	Met	Leu	Ser	Glu	His	Ser	Phe	Ile	Pro	Glu	Pro	Lys	Ser	Ser
305					310					315					320
Cys	Tyr	Asn	Thr	His	Glu	Lys	Arg	Ile	Tyr	Gln	Ser	Asn	Met	Leu	Asn
				325					330					335	
Arg	Tyr	Leu	Ile	Asn	Val	Thr	Phe	Glu	Gly	Arg	Asn	Leu	Ser	Phe	Ser
		340						345					350		
Glu	Asp	Gly	Tyr	Gln	Met	His	Pro	Lys	Leu	Val	Ile	Ile	Leu	Leu	Asn
		355					360					365			
Lys	Glu	Arg	Lys	Trp	Glu	Arg	Val	Gly	Lys	Trp	Lys	Asp	Lys	Ser	Leu
	370					375					380				
Gln	Met	Lys	Tyr	Tyr	Val	Trp	Pro	Arg	Met	Cys	Pro	Glu	Thr	Glu	Glu
385						390					395				400
Gln	Glu	Asp	Asp	His	Leu	Ser	Ile	Val	Thr	Leu	Glu	Glu	Ala	Pro	Phe
				405					410					415	
Val	Ile	Val	Glu	Ser	Val	Asp	Pro	Leu	Ser	Gly	Thr	Cys	Met	Arg	Asn
			420						425				430		
Thr	Val	Pro	Cys	Gln	Lys	Arg	Ile	Val	Thr	Glu	Asn	Lys	Thr	Asp	Glu
		435					440					445			
Glu	Pro	Gly	Tyr	Ile	Lys	Lys	Cys	Cys	Lys	Gly	Phe	Cys	Ile	Asp	Ile
	450					455					460				
Leu	Lys	Lys	Ile	Ser	Lys	Ser	Val	Lys	Phe	Thr	Tyr	Asp	Leu	Tyr	Leu
465						470					475				480
Val	Thr	Asn	Gly	Lys	His	Gly	Lys	Lys	Ile	Asn	Gly	Thr	Trp	Asn	Gly
			485						490					495	
Met	Ile	Gly	Glu	Val	Val	Met	Lys	Arg	Ala	Tyr	Met	Ala	Val	Gly	Ser
			500					505					510		
Leu	Thr	Ile	Asn	Glu	Glu	Arg	Ser	Glu	Val	Val	Asp	Phe	Ser	Val	Pro
		515					520					525			
Phe	Ile	Glu	Thr	Gly	Ile	Ser	Val	Met	Val	Ser	Arg	Ser	Asn	Gly	Thr
	530					535					540				
Val	Ser	Pro	Ser	Ala	Phe	Leu	Glu	Pro	Phe	Ser	Ala	Asp	Val	Trp	Val
545						550					555				560
Met	Met	Phe	Val	Met	Leu	Leu	Ile	Val	Ser	Ala	Val	Ala	Val	Phe	Val
			565						570					575	
Phe	Glu	Tyr	Phe	Ser	Pro	Val	Gly	Tyr	Asn	Arg	Cys	Leu	Ala	Asp	Gly
			580						585				590		
Arg	Glu	Pro	Gly	Gly	Pro	Ser	Phe	Thr	Ile	Gly	Lys	Ala	Ile	Trp	Leu
		595					600					605			
Leu	Trp	Gly	Leu	Val	Phe	Asn	Asn	Ser	Val	Pro	Val	Gln	Asn	Pro	Lys
	610					615					620				
Gly	Thr	Thr	Ser	Lys	Ile	Met	Val	Ser	Val	Trp	Ala	Phe	Phe	Ala	Val
	625					630					635				640
Ile	Phe	Leu	Ala	Ser	Tyr	Thr	Ala	Asn	Leu	Ala	Ala	Phe	Met	Ile	Gln
			645						650					655	
Glu	Glu	Tyr	Val	Asp	Gln	Val	Ser	Gly	Leu	Ser	Asp	Lys	Lys	Phe	Gln
			660						665					670	
Arg	Pro	Asn	Asp	Phe	Ser	Pro	Pro	Phe	Arg	Phe	Gly	Thr	Val	Pro	Asn
		675					680							685	

-continued

Gly Ser Thr Glu Arg Asn Ile Arg Asn Asn Tyr Ala Glu Met His Ala
 690 695 700

Tyr Met Gly Lys Phe Asn Gln Arg Gly Val Asp Asp Ala Leu Leu Ser
 705 710 715

Leu Lys Thr Gly Lys Leu Asp Ala Phe Ile Tyr Asp Ala Ala Val Leu
 725 730 735

Asn Tyr Met Ala Gly Arg Asp Glu Gly Cys Lys Leu Val Thr Ile Gly
 740 745 750

Ser Gly Lys Val Phe Ala Ser Thr Gly Tyr Gly Ile Ala Ile Gln Lys
 755 760 765

Asp Ser Gly Trp Lys Arg Gln Val Asp Leu Ala Ile Leu Gln Leu Phe
 770 775 780

Gly Asp Gly Glu Met Glu Glu Leu Glu Ala Leu Trp Leu Thr Gly Ile
 785 790 795 800

Cys His Asn Glu Lys Asn Glu Val Met Ser Ser Gln Leu Asp Ile Asp
 805 810 815

Asn Met Ala Gly Val Phe Tyr Met Leu Gly Ala Ala Met Ala Leu Ser
 820 825 830

Leu Ile Thr Phe Ile Cys Glu His Leu Phe Tyr Trp Gln Phe Arg His
 835 840 845

Cys Phe Met Gly Val Cys Ser Gly Lys Pro Gly Met Val Phe Ser Ile
 850 855 860

Ser Arg Gly Ile Tyr Ser Cys Ile His Gly Val Ala Ile Glu Glu Arg
 865 870 875 880

Gln Ser Val Met Asn Ser Pro Thr Ala Thr Met Asn Asn Thr His Ser
 885 890 895

Asn Ile Leu Arg Leu Leu Arg Thr Ala Lys Asn Met Ala Asn Leu Ser
 900 905 910

Gly Val Asn Gly Ser Pro Gln Ser Ala Leu Asp Phe Ile Arg Arg Glu
 915 920 925

Ser Ser Val Tyr Asp Ile Ser Glu His Arg Arg Ser Phe Thr His Ser
 930 935 940

Asp Cys Lys Ser Tyr Asn Asn Pro Pro Cys Glu Glu Asn Leu Phe Ser
 945 950 955 960

Asp Tyr Ile Ser Glu Val Glu Arg Thr Phe Gly Asn Leu Gln Leu Lys
 965 970 975

Asp Ser Asn Val Tyr Gln Asp His Tyr His His His His Arg Pro His
 980 985 990

Ser Ile Gly Ser Ala Ser Ser Ile Asp Gly Leu Tyr Asp Cys Asp Asn
 995 1000 1005

Pro Pro Phe Thr Thr Gln Ser Arg Ser Ile Ser Lys Lys Pro Leu
 1010 1015 1020

Asp Ile Gly Leu Pro Ser Ser Lys His Ser Gln Leu Ser Asp Leu
 1025 1030 1035

Tyr Gly Lys Phe Ser Phe Lys Ser Asp Arg Tyr Ser Gly His Asp
 1040 1045 1050

Asp Leu Ile Arg Ser Asp Val Ser Asp Ile Ser Thr His Thr Val
 1055 1060 1065

Thr Tyr Gly Asn Ile Glu Gly Asn Ala Ala Lys Arg Arg Lys Gln
 1070 1075 1080

Gln Tyr Lys Asp Ser Leu Lys Lys Arg Pro Ala Ser Ala Lys Ser
 1085 1090 1095

-continued

```

<210> SEQ ID NO 11
<211> LENGTH: 531
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

Arg Ser Gln Lys Ser Pro Pro Ser Ile Gly Ile Ala Val Ile Leu Val
1          5          10          15
Gly Thr Ser Asp Glu Val Ala Ile Lys Asp Ala His Glu Lys Asp Asp
          20          25          30
Phe His His Leu Ser Val Val Pro Arg Val Glu Leu Val Ala Met Asn
          35          40          45
Glu Thr Asp Pro Lys Ser Ile Ile Thr Arg Ile Cys Asp Leu Met Ser
          50          55          60
Asp Arg Lys Ile Gln Gly Val Val Phe Ala Asp Asp Thr Asp Gln Glu
65          70          75          80
Ala Ile Ala Gln Ile Leu Asp Phe Ile Ser Ala Gln Thr Leu Thr Pro
          85          90          95
Ile Leu Gly Ile His Gly Gly Ser Ser Met Ile Met Ala Asp Lys Asp
          100          105          110
Glu Ser Ser Met Phe Phe Gln Phe Gly Pro Ser Ile Glu Gln Gln Ala
          115          120          125
Ser Val Met Leu Asn Ile Met Glu Glu Tyr Asp Trp Tyr Ile Phe Ser
          130          135          140
Ile Val Thr Thr Tyr Phe Pro Gly Tyr Gln Asp Phe Val Asn Lys Ile
145          150          155          160
Arg Ser Thr Ile Glu Asn Ser Phe Val Gly Trp Glu Leu Glu Glu Val
          165          170          175
Leu Leu Leu Asp Met Ser Leu Asp Asp Gly Asp Ser Lys Ile Gln Asn
          180          185          190
Gln Leu Lys Lys Leu Gln Ser Pro Ile Ile Leu Leu Tyr Cys Thr Lys
          195          200          205
Glu Glu Ala Thr Tyr Ile Phe Glu Val Ala Asn Ser Val Gly Leu Thr
          210          215          220
Gly Tyr Gly Tyr Thr Trp Ile Val Pro Ser Leu Val Ala Gly Asp Thr
225          230          235          240
Asp Thr Val Pro Ala Glu Phe Pro Thr Gly Leu Ile Ser Val Ser Tyr
          245          250          255
Asp Glu Trp Asp Tyr Gly Leu Pro Ala Arg Val Arg Asp Gly Ile Ala
          260          265          270
Ile Ile Thr Thr Ala Ala Ser Asp Met Leu Ser Glu His Ser Phe Ile
          275          280          285
Pro Glu Pro Lys Ser Ser Cys Tyr Asn Thr His Glu Lys Arg Ile Tyr
          290          295          300
Gln Ser Asn Met Leu Asn Arg Tyr Leu Ile Asn Val Thr Phe Glu Gly
305          310          315          320
Arg Asn Leu Ser Phe Ser Glu Asp Gly Tyr Gln Met His Pro Lys Leu
          325          330          335
Val Ile Ile Leu Leu Asn Lys Glu Arg Lys Trp Glu Arg Val Gly Lys
          340          345          350
Trp Lys Asp Lys Ser Leu Gln Met Lys Tyr Tyr Val Trp Pro Arg Met
          355          360          365
Cys Pro Glu Thr Glu Glu Gln Glu Asp Asp His Leu Ser Ile Val Thr
          370          375          380

```


-continued

Leu Glu Glu Ala Pro Phe Val Ile Val Glu Ser Val Asp Pro Leu Ser
 385 390 395 400

Gly Thr Cys Met Arg Asn Thr Val Pro Cys Gln Lys Arg Ile Val Thr
 405 410 415

Glu Asn Lys Thr Asp Glu Glu Pro Gly Tyr Ile Lys Lys Cys Cys Lys
 420 425 430

Gly Phe Cys Ile Asp Ile Leu Lys Lys Ile Ser Lys Ser Val Lys Phe
 435 440 445

Thr Tyr Asp Leu Tyr Leu Val Thr Asn Gly Lys His Gly Lys Lys Ile
 450 455 460

Asn Gly Thr Trp Asn Gly Met Ile Gly Glu Val Val Met Lys Arg Ala
 465 470 475 480

Tyr Met Ala Val Gly Ser Leu Thr Ile Asn Glu Glu Arg Ser Glu Val
 485 490 495

Val Asp Phe Ser Val Pro Phe Ile Glu Thr Gly Ile Ser Val Met Val
 500 505 510

Ser Arg Ser Asn Gly Thr Val Ser Pro Ser Ala Phe Leu Glu Pro Phe
 515 520 525

Ser Ala Asp
 530

<210> SEQ ID NO 12
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

Gly Tyr Ile Lys Lys Cys Cys Lys Gly Phe Cys Ile Asp Ile Leu Lys
 1 5 10 15

Lys Ile Ser Lys
 20

<210> SEQ ID NO 13
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: amino acid sequence of an antigenic fragment of
 the NR2B subunit modified by an N-terminal Cys for attachment to a
 carrier protein

<400> SEQUENCE: 13

Cys Gly Tyr Ile Lys Lys Cys Cys Lys Gly Phe Cys Ile Asp Ile Leu
 1 5 10 15

Lys Lys Ile Ser Lys
 20

<210> SEQ ID NO 14
 <211> LENGTH: 6237
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

ttgaatttgc atctcttcaa gacacaagat taaacaaaaa tttacgctaa attggatttt 60

aaattatctt ccgttcattt atccttcgctc tttcttatgt ggatattgcaa gcgagaagaa 120

gggactggac attccaaca tgctcaactcc cttaattctgt ccgtotagag gtttgcttc 180

tacaaaccaa gggagtcgac gagttgaaga tgaagcccag agcggagtgc tgttctocca 240

agttctggtt ggtggtggcc gtcttgccg tgtcaggcag cagagctcgt tctcagaaga 300

-continued

gccccccag cattggcatt gctgtcatcc tcgtgggcac ttccgacgag gtggccatca	360
aggatgcca cgagaaagat gatttcacc atctctccgt ggtaccccg gtggaactgg	420
tagccatgaa tgagaccgac ccaagagca tcatacccg catctgtgat ctcatgtctg	480
accggaagat ccagggggtg gtgtttgctg atgacacaga ccaggaagcc atcgcccaga	540
tcctcgattt catttcagca cagactctca ccccgatcct gggcatccac gggggctcct	600
ctatgataat ggcagataag gatgaatcct ccatgttctt ccagtttggc ccatcaattg	660
aacagcaagc ttocgtaatg ctcaacatca tggaagaata tgactgttac atcttttcta	720
tcgtcaccac ctatttcctt ggctaccagg actttgtaaa caagatccgc agcaccattg	780
agaatagctt tgtgggctgg gagctagagg aggtcctcct actggacatg tccttggacg	840
atggagattc taagatccag aatcagctca agaaacttca aagccccatc attcttcttt	900
actgtacca ggaagaagcc acctacatct ttgaagtggc caactcagta gggctgactg	960
gctatggcta cacgtggatc gtgcccagtc tgggtggcagg ggatacagac acagtgcctg	1020
cggagtccc cactgggctc atctctgtat catatgatga atgggactat ggctccccg	1080
ccagagtgag agatggaatt gccataatca ccaactgctgc ttctgacatg ctgtctgagc	1140
acagcttcat ccttgagccc aaaagcagtt gttacaacac ccacgagaag agaacttacc	1200
agtccaatat gctaaatagg tatctgatca atgtcacttt tgaggggagg aatttgcctt	1260
tcagtgaaga tggctaccag atgcaccga aactggtgat aattcttctg aacaaggaga	1320
ggaagtggga aagggtggg aagtggaaag acaagtcctt gcagatgaag tactatgtgt	1380
ggccccgaat gtgtccagag actgaagagc aggaggatga ccatctgagc attgtgacct	1440
tggaggaggc accatttctt attgtgaaa gtgtggacct tctgagtgga acctgcatga	1500
ggaacacagt cccctgccc aaacgcatag tcaactgagaa taaaacagac gaggagccgg	1560
gttacatcaa aaaatgctgc aaggggttct gtattgacat ccttaagaaa atttctaat	1620
ctgtgaagtt cacctatgac ctttacctgg ttaccaatgg caagcatggg aagaaaatca	1680
atggaacctg gaatggtatg attggagagg tggatcatgaa gagggcctac atggcagtgg	1740
gctcactcac catcaatgag gaacgatcgg aggtggtcga cttctctgtg cccttcatag	1800
agacaggcat cagtgtcatg gtgtcacgca gcaatgggac tgtctcactt tctgccttct	1860
tagagccatt cagcgtgac gtatgggtga tgatgtttgt gatgctgctc atcgtctcag	1920
ccgtggctgt ctttctctt gagtacttca gcctgtggg ttataacagg tgcctcgtg	1980
atggcagaga gcctgtgga ccctcttca ccatcgcaa agctatttgg ttgctctggg	2040
gtctggtggt taacaactcc gtacctgtgc agaaccctaaa ggggaccacc tccaagatca	2100
tggtgtcagt gtgggccttc tttgctgca tcttctggc cagctacact gccaaactag	2160
ctgccttcat gatccaagag gaatatgtgg accaggttct tggcctgagc gacaaaaagt	2220
tccagagacc taatgacttc tcacccctt tccgcttgg gaccgtgcc aacggcagca	2280
cagagagaaa tattcgcaat aactatgagc aaatgcacgc ctacatggga aagttaacc	2340
agaggggtgt agatgatgca ttgctctccc tgaaaacagg gaaactggat gccttcatct	2400
atgatgcagc agtgcgaac tatatggcag gcagagatga aggctgcaag ctggtgacca	2460
ttggcagtgg gaaggtcttt gcttccactg gctatggcat tgccatcaa aaagattctg	2520
ggtggaagcg ccaggtggac cttgctatcc tgcagctctt tggagatggg gagatggaag	2580
aactggaagc tctctggctc actggcattt gtcacaatga gaagaatgag gtcagagca	2640

-continued

gccagctgga cattgacaac atggcagggg tcttctacat gttgggggcg gccatggctc	2700
tcagcctcat caccttcato tgcgaacacc ttttctattg gcagttccga cattgcttta	2760
tgggtgtctg ttctggcaag cctggcatgg tcttctccat cagcagaggt atctacagct	2820
gcattccatgg ggtggcgatc gaggagcgc agtctgtaat gaactcccc accgcaacca	2880
tgaacaacac aactccaac atcctgcgcc tgctgcgcac ggccaagaac atggctaacc	2940
tgtctgggtg gaatggctca ccgagagcg ccctggactt catccgacgg gagtcatccg	3000
tctatgacat ctacagacac cgccgcagct tcacgcatc tgactgcaaa tcctacaaca	3060
accgcctctg tgaggagaac ctcttcagtg actacatcag tgaggtagag agaacgttcg	3120
ggaacctgca gctgaaggac agcaactgtt accaagatca ctaccaccat caccaccggc	3180
cccatagtat tggcagtgcc agctccatcg atgggctcta cgactgtgac aaccacccct	3240
tcaccacca gtccaggtcc atcagcaaga agcccctgga catcggcctc cctcctcca	3300
agcacagcca gctcagtgac ctgtacggca aattctcctt caagagcgac cgctacagtg	3360
gccacgacga cttgatccgc tccgatgtct ctgacatctc aaccacaccc gtcacctatg	3420
ggaacatcga gggcaatgcc gccaaaggcc gtaagcagca atataaggac agcctgaaga	3480
agcggcctgc ctccgccaag tcccgcaggg agtttgacga gatcgagctg gcctaccgtc	3540
gccgaccgcc ccgctcccc gaccacaagc gctacttcag ggacaaggaa gggctacggg	3600
acttctacct ggaccagttc cgaacaaagg agaactcacc ccaactgggag cacgtagacc	3660
tgaccgacat ctacaaggag cggagtgatg actttaagcg cgactccatc agcggaggag	3720
ggcctctgac caacaggtct cacatcaagc acgggacggg cgacaaaac gccgtggtca	3780
gcggggtacc tgcaccttgg gagaagaacc tgaccaactg ggagtgggag gaccggtccg	3840
ggggcaactt ctgccgcagc tgtcctcca agctgcacaa ctactccacg acggtgacgg	3900
gtcagaactc gggcaggcag gcgtgcatcc ggtgtgaggc ttgcaagaaa gcaggcaacc	3960
tgtatgacat cagtgaggac aactccctgc aggaactgga ccagccggct gccccagtgg	4020
cggtgacgtc aaacgcctcc accactaagt accctcagag cccgactaat tccaaggccc	4080
agaagaagaa ccggaacaaa ctgcccgggc agcaactccta cgacaccttc gtggacctgc	4140
agaaggaaga agcccctctg gccccgcgca gcgtaagcct gaaagacaag gcccgattca	4200
tggatgggag cccctacgcc cacatgtttg agatgtcagc tggcgagagc acctttgcca	4260
acaacaagtc ctacgtgccc actgcccggc atcaccacca caacaacccc ggcggcgggt	4320
acatgctcag caagtgcctc taccctgacc gggtcacgca aaacccttc atccccactt	4380
ttggggacga ccagtctctg ctccatggca gcaaatccta cttcttcagg cagcccacgg	4440
tggcgggggc gtcgaaagcc aggcgggact tccgggccct tgtcaccaac aagccggtgg	4500
tctcggccct tcattggggc gtgccagccc gtttccagaa ggacatctgt ataggaacc	4560
agtccaaccc ctgtgtgctc aacaacaaa accccagggc tttcaatggc tccagcaatg	4620
ggcatgttta tgagaaactt tctagtattg agtctgatgt ctgagtgagg gaacagagag	4680
gttaaggtgg gtacgggagg gtaaggctgt gggtcgctgt atgcgcatgt cacggagggt	4740
gacgggggtg aacttggctc ccatttgctc ctttcttgtt ttaatttatt tatgggatcc	4800
tggagttctg gttcctactg ggggcaaccc tggtgaccag caccatctct cctcctttc	4860
acagttctct ccttcttccc cccgctgtca gccattcctg ttcccagatg atgatgcat	4920
gggccctctc agcaggggag ggtagagcgg agaaaggaag ggctgcatgc gggcttctc	4980
ctggtgtgga agagctcctt gatatcctct ttgagtgaag ctgggagaac caaaaagagg	5040

-continued

```

ctatgtgagc acaaaggtag cttttcccaa actgatcttt tcatttaggt gaggaagcaa 5100
aagcatctat gtgagaccat ttagcacact gcttgtgaaa ggaaagaggc tctggctaaa 5160
ttcatgtgct ttagatgaca tctgtctagg aatcatgtgc caagcagagg ttgggaggcc 5220
atttgtgttt atataaagc ccaaaaatgc ttgcttcaac cccatgagac tcgatagtgg 5280
tggtgaacag aaccaaggt cattggtggc agagtggatt cttgaacaaa ctggaaagta 5340
cgttatgata gtgtccccg gtgccttggg gacaagagca ggtggattgt gcgtgcatgt 5400
gtgttcacgc acacttgac ccatgtgtag tcaggtgcct caagagaagg caacctgac 5460
tctttctatt gtttcttca atatcccaa gcagtgtgat tgtttggctt atatacagac 5520
agagatggcc atgtattacc tgaattttgg ctgtgtctcc cttcatcctt ctggaataag 5580
gagaatgaaa attcctgata aagaagattc tgtggtctaa acaaaaaaag gcggtgagca 5640
atcctgcaag aacaaggtac ataaacaagt cctcagtggt tggcaattgt ttcaaccagt 5700
ttgaaccaag aactttccag gaaggctaaa gggaaaccga attttcacag ccatgattct 5760
tttcccaca cttgggagca aaagattcta caaagctctt ttgagcattt agactctcga 5820
ctggccaagg tttggggaag aacgaagcca cctttgaaga agtaaggagt cgtgtatggt 5880
agggtaagtg agagaggggg atgtttccaa tgctttgatc ccttcttact taacctgaag 5940
ctagacgagc aggtctcttc cccccaaaac tgattacaac tgctacagag cagacagtta 6000
agagaaatga gcttgacctt taagagaaat gagctgcaact ccatgagtgc agctctggag 6060
gtacgaaaag aggggaagag acttggaat gggagacggg ggcagagagg gacctccac 6120
cacctctttg ggcttgctc cctgggaatg tgacttgagc ccagagtgaa cactcttgg 6180
agaagccctt ctacctctc gcaacacctt gtttccctct cagattgtac cattgag 6237

```

```

<210> SEQ ID NO 15
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 15

```

```

ggttacatca aaaaatgctg caaggggttc tgtattgaca tccttaagaa aatttctaaa 60

```

```

<210> SEQ ID NO 16
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer

```

```

<400> SEQUENCE: 16

```

```

tcaactgagaa taaaacagac g 21

```

```

<210> SEQ ID NO 17
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer

```

```

<400> SEQUENCE: 17

```

```

tcacctatga cctttacctg g 21

```

85

What is claimed is:

1. A method for aiding in the diagnosis that a human is having an ischemic episode comprising:

- a) obtaining a test sample of human blood, serum, or plasma from a human subject;
- b) analyzing the obtained test sample for the amount of an NR2 antibody that binds NR2A N-terminal domain fragment peptide having SEQ ID NO:3;
- c) comparing the result of step (b) with a corresponding reference amount of said NR2 antibody, wherein the corresponding reference amount is derived from a population of apparently healthy human subjects; and
- d) if the amount of NR2 antibody in the sample exceeds the reference amount, diagnosing the human as having an increased probability that the human is having an ischemic episode.

2. The method of claim 1 wherein the NR2 antibody is measured by immunoassay.

3. The method of claim 1 wherein the NR2 antibody is detected or measured by agglutination comprising:

- a) contacting the test sample with NR2 antigen bound on an agglutinating carrier for sufficient time and under con-

86

ditions to promote agglutination between the NR2 antigen and NR2 antibody in said test sample;

- b) measuring the signal generated from the agglutination; and
- c) correlating the strength of said signal to the titer of NR2 antibody present in said sample.

4. The method of claim 1 wherein levels of NR2 antibody is detected or are measured by a process comprising:

- a) contacting the test sample with a NR2 antigen capture agent for a time sufficient and under conditions to form a complex between said NR2 antigen capture agent and NR2 antibody in said test sample;
- b) contacting the complex with an indicator reagent comprising a secondary antibody attached to a signal generating compound; and
- c) measuring the strength of said signal.

5. The method of claim 1, wherein said step (b) comprises immunochemically analyzing said obtained test sample in an immunoassay, and said corresponding reference amount in step (c) is immunochemically derived using the same immunoassay in step (b).

* * * * *

专利名称(译)	用于评估TIA /中风风险的抗体测定方法		
公开(公告)号	US7658911	公开(公告)日	2010-02-09
申请号	US11/339452	申请日	2006-01-25
申请(专利权)人(译)	CIS BIOTECH , INC.		
当前申请(专利权)人(译)	CIS BIOTECH , INC.		
[标]发明人	DAMBINOVA SVETLANA A		
发明人	DAMBINOVA, SVETLANA A.		
IPC分类号	A61K49/00 A61K39/00 G01N33/00 G01N33/53 G01N33/68 G01N33/94		
CPC分类号	G01N33/6893 G01N33/9406 G01N2800/52 G01N2800/2871 G01N2333/70571		
优先权	60/301297 2001-06-27 US		
其他公开文献	US20060172342A1		
外部链接	Espacenet USPTO		

摘要(译)

用于诊断中枢神经系统疾病，特别是短暂性脑缺血发作或中风的方法，试剂盒和组合物，包括测量来自人受试者的生物样品中的NR2A和/或NR2B NMDA受体或其片段的水平，并任选地测量其他生物标志物如同型半胱氨酸和谷氨酸。该方法对于识别有中风风险的个体以及在急诊室环境中诊断中风特别有用。

TABLE 5

Detection of NR2A-B receptor peptides in the blood of patients by latex agglutination

Group	Total N	NR2A-B Titer	LA predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	2.63 ± 0.92	25	89.3	3	10.7
TIA	9	7.34 ± 0.43	2	22.2	7	77.8
Pre-stroke	9	4.21 ± 0.26	2	22.2	7	77.8
Acute stroke	31	5.20 ± 1.71	4	9.7	27	87.1
TBI	11	3.99 ± 0.44	2	18.8	9	81.8