



US 20030170720A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2003/0170720 A1**  
van der Kuyl et al. (43) **Pub. Date: Sep. 11, 2003**

(54) **MEANS AND METHODS FOR TREATMENT EVALUATION** Sep. 28, 2001 (EP)..... 0120373.2

**Publication Classification**

(76) Inventors: **Antoinette Cornelia van der Kuyl**,  
Loosdrecht (NL); **Marion Cornelissen**,  
Loosdrecht (NL)

(51) **Int. Cl.<sup>7</sup>** ..... **G01N 33/53**  
(52) **U.S. Cl.** ..... **435/7.1**

Correspondence Address:

**TRASK BRITT**  
**P.O. BOX 2550**  
**SALT LAKE CITY, UT 84110 (US)**

(57) **ABSTRACT**

The invention provides a method for determining whether a treatment is effective in changing the status of a certain set of target cells, such as a tumor, in a patient. This method implies obtaining a sample from a patient after initiation of a treatment, and determining whether said sample comprises an expression product of at least one marker gene. Preferably, said sample is a blood sample. In one aspect, said expression product is expressed by a peripheral blood mononuclear cell. Said marker gene may be a gene involved in the generation, maintenance and/or breakdown of blood vessels (angiogenesis). A method of the invention is very suitable to determine within a few days if a certain treatment against Kaposi's Sarcoma is successful. Moreover, this method is suitable for determining the presence of angiogenesis and/or tumor cells in a patient.

(21) Appl. No.: **10/055,728**

(22) Filed: **Jan. 23, 2002**

**Related U.S. Application Data**

(60) Provisional application No. 60/325,722, filed on Sep. 28, 2001.

(30) **Foreign Application Priority Data**

Jan. 23, 2001 (EP)..... 01200228.3

Figure 1




Figure 2

CATGGATGTGCACGATGGCAAGGTGGTGTCCACCCACGAGCAGGTCCTT  
CGCACCAAGAACTGAGGCTGCCAGCCCCGCTCAGGCCTAGGAGGCCCC  
CCGTGTGGACACAGATCCCCTGGAAGATCCCTCTCCTGCCAAGCACT  
TCACAGCTGGACCCCTGCTTCACCCTCACCCCTCCTGGCAATCAATACAG  
CTTCATTATCTGAGTTGCTAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

Figure 3

ATCTACCAGCTCATGATGCAGTGCTGGCAGCAGGAGCGTGCCACCGCCC  
CAAGTTCGCTGACATCGTCAGCATCCTGGACAAGCTCATTCGTGCCCTG  
ACTCCCTCAAGACCOCTGGCTGACTTTGACCCCGCGTGTCTATCCGGCTC  
CCCAGCACGAGCGGCCTCGGAGGGGGTGCCCTTCGACCGGTGTCCGAGT  
GGCTGGAGTCCATCAAGATGCAGCAGTATACGGAGCACTTC

Figure 4

CATGCTTGACATACCTACCAGTATTATTCCTGACGACACATATACATATG  
AGAATATACCTTATTTATTTTTGTGTAGGTGTCTGCCTTCACAAATGTCAT  
TGTCTACTCCTAGAAGAACCATACTCAATTTTTGTTTTGAGTACTGT  
ACTATCCTGTAATAATCTTAAGCAGGTTTGTTCAGCACTGATGGAAA  
ATACCAGTGTTGGGTTTTTTTTAGTTGCCAACAGTGTATGTTTGCTGAT  
TATTTATGACCTGAAATAATATATTTCTTCTTAAGAAGACATTTTGTTAC  
ATAAGGATGACTTTTTTATACAATGGAATAAATTATGGCATTTCATTTG

Figure 5

CATGCATCACGGATCAATAGACTGTA~~CTT~~ATTTTCCAATAAAAATTTCAA  
ACTTTGTA~~CT~~TT

Figure 6

AACTTGCCCTGTGCCTGTGTCCCCATGCTAGGGGCGGAGGGGTCTTTTC  
CTTCTTCTTTCTACCTACCCCTTTTCTCTTGGCCAGGGGCCTCGTATCCT  
ACCTTTCCTTGTCCCCTGGGCTGGCTGCACAGAGGATTGCCCTTCTCTTT  
TCAGAGCTGGCCCTCGATGCCAAATTAGCATTTAGTATTTTGCTCAAAGTC  
TAAGGGACC

Figure 7

CATGTTGCATATCAGGGTGCTCAAGGATTGGAGAGGAGACAAAACCAGG  
AGCAGCACAGTGGGGACATCTCCCGTCTCAACAGCCCAGGCCTATGGGG  
GCTCTGGAAGGATGGGCCAGCTTGCAGGGGTTGGGGAGGGAGACATCCA  
GCTTGGGCTTTCCCCTTTGGAATAAAC~~CA~~TTGGTCTGTCACAAAAAAAAAA  
AAAAAAAAAAAAAAAA

Figure 8

CATGCCCTGTTCAAGCTACTCCCACTCCCGGCTGTCATTCAGAAAAAAT  
AAATGTTCTAATAAGCTCCAAAAAAAAAAAAAAAAAAAAAAAAAAAA

Figure 9

CATGGATCAATCAGTGTGATTAGCTTTCTCAGCAGACATTGTGCCATATG  
TATCAAATGACAAATCTTTATTGAATGGTTTTGCTCAGCACCACCTTTTAA  
TATATTGGCAGTACTTATTATATAAAAGGTAAACCAGCATTCTCAAAAAAA  
AAAAAAAAAAAA

Figure 10

CATGGAGGGTGCCAACAGCATCTTTTCCGGGTTCTGCTCTTTCCAGAT  
ATGGAGGCCTGACCTGTGGGCTGCTTACATCCACCCCGGCTCCCCCTGC  
CAGCAACGCTCACTCTACCCCAACACCACCCCTTGCCCAGCCAATGCAC  
ACAGTAGGGCTTGGTGAATGCTGCTGAGTGAATGAGTAAATAAACTCTTC  
AAGGCC

**Figure 11**

CGGCTGGACACGTCGGAGGTGGTCTTCAACAGCAAGGAGCAAGGCTCCT  
GGGGCCGCGAGGAGCGCGGGCCGGGCGTTCCCTTTCCAGCGCGGGCAGCC  
CTTCGAGGTGCTCA'CATCGCGTCAGACGACGGOTTCAAGGCCGTGGTTG  
GGGACGCCCAGTACCACCACTTCCGCC

**Figure 12**

CACAACCTCGACTACTACAAGAAGACAACCAACGGCCGGCTGCCCGTGAA  
GTGGATGGCGCCTGAGGCATTATTTGACCGAGTCTACACTCACCAGAGTG  
ACGTCTGGTCCTTTGGGGTCCTGCTCTGGGAGATCTTCACGCTGGGGGGC  
TCCCGTACCCCGGCATCCCTGTGGAGGAGCTTTCAAGCTGCTGAAGGA  
GGGC

**Figure 13**

CATGGAGCAGCGCCCTGTTCCGGGGGCAGCCAGTGACCCAGCCCCACC  
AATGGGCCTCCAGAGACCCAGGAACAATAAAATGTCTTCTCCCACC

Figure 14

CATGCTGCACTGAGAACTGCATTTAGTAGCATTGTGCATCCAGCCG  
GAAGTTAAAGCACACTTACTTTATTCACCTATTTTATAATAAACGTTCTT  
GCTGCTGTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

Figure 15

CATGCCACAGGAGAATTCGGGGATTGAGTTTCTCTGAATAGCATATAT  
ATGATGCATCGGATAGGTCATTATGATTTTTACCATTTGACTTACATAA  
TGAAAACCAATTCATTTAAATATCAGATTATTATTTGTAAGTGTGGAA  
AAAGCTAATTGTAGTTTCATTATGAAGTTTCCCAATAAACAGGTATTC  
TAAACTTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

Figure 16

CATGCTGTGCGGAACTGCGTCAGGGCAAATGTACAGCAGGATTTCCCC  
AACCAGCTCCATCATCACAGACACAGAGGGCTGCAGGGGAGGCCTGCC  
ACTGTTTTGTCGACTCTGCCCTCCTCTGGCAGCATAGATCCTTAGGTGCTC  
AATAAAGGTGTGCTGTATTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

Figure 17

CATGCTCCATTGCCAGACTCTTGCTGGGAGCCCGTCCAGAATGTCCTCC  
CAATAAACTCCATCCTATGACGCAAAAAAAAAAAAAAAAAAAAAAAAAA ..

Figure 18

CATGACAGCGGCAATCTTTCTTTGGTCAAAGTTTTCTGTTATTTGCT  
TGTCATATTCGATGTACTTTAAGGTGTCTTTATGAAGTTGCTATTCTGGC  
AATAAACTTTTAGACTTTAAAAAAAAAAAAAAAAAAAAAAAAA

Figure 19





Figure 20



## MEANS AND METHODS FOR TREATMENT EVALUATION

### CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] Under the provisions of 35 U.S.C. 119(e), priority is claimed from U.S. Provisional Patent Application Serial No. 60/325,722, filed Sep. 28, 2001.

### TECHNICAL FIELD

[0002] The invention relates to the field of medicine. The invention particularly relates to the fields of molecular biology and detection methods.

### BACKGROUND

[0003] Recent advances in the knowledge of molecular processes in a cell and techniques to study these processes have resulted in improved methods of typing and treating diseases. Understanding of the underlying molecular diversity of tumors has, for instance, already led to a better understanding of the diversity of response to treatment of morphologically similar tumors. Improved typing influences the way tumor patients are being treated. A drawback of the current methods of treatment is, however, that it takes a relatively long time to determine whether a treatment given to a patient is actually effective. This impedes the optimization of dosages and/or schedules with which treatment is given. Moreover, it also slows down the possibility of adjusting the treatment regimen all together. For instance, adjustment of therapy is currently only possible when macroscopic analysis of tumor cells in the body indicates that the therapy given is not effective. Macroscopic changes typically need several weeks to manifest themselves and equipment to measure such changes is often not readily available.

### DISCLOSURE OF THE INVENTION

[0004] The present invention provides a method for determining whether a treatment is effective in changing a status of a certain set of target cells in an individual comprising obtaining a sample from the individual after initiation of the treatment and determining whether the sample comprises an expression product of at least one marker gene. In one embodiment of the invention the set of target cells comprises a tumor cell. By changing a status of a set of target cells is meant herein that at least one property of the set of target cells is altered. For instance, the amount of the target cells may be changed. The amount may either be increased or decreased. Alternatively, the activity of the target cells may be altered. The activity may be a replication activity. As another example, the activity may be an activity involved with angiogenesis. Alternatively, the activity may be an apoptotic activity.

[0005] It was found that tumor cells and/or surrounding tissue respond, on a molecular level, very quickly to an effective treatment. This response can be detected by measuring an expression product of a marker gene. Marker gene expression products are indicative for a response to treatment. Marker genes are typically genes that are expressed by said set of target cells, for instance tumor cells, and/or surrounding tissue. However, marker genes can also be expressed in non-tumor target cells in other compartments of the body, for instance blood cells and/or cardiovascular cells.

[0006] Alternatively, marker gene expression can be initiated upon treatment given to the individual. Marker gene expression products are responsive to treatment given to a patient. A response can be an alteration in the relative amounts of expression product. However, it can also be an alteration in absolute presence or absence of expressed product such as RNA and/or protein.

[0007] According to the invention, a sample which is obtained from a patient may comprise at least one of the target cells. This is particularly suitable for detecting circulating tumor cells which have released themselves from a tumor and are circulating in the blood of a patient.

[0008] Alternatively, the target cells may be non-tumor cells. In another embodiment of the invention, the sample does not comprise any target cells. However, the sample may comprise another, non-target cell. Expression, or change of expression, of at least one marker gene by the non-target cell is indicative for the status of a certain set of target cells. The non-target cell preferably comprises a peripheral blood mononuclear cell, as is described below. In yet another embodiment, the sample does not comprise any cell at all. For instance, an expression product of a marker gene, produced by a target cell or non-target cell elsewhere in an individual's body, may be present in the sample at detectable levels.

[0009] With a method of the invention it is possible to determine whether a treatment is effective in said individual. This can be done while a treatment is given or shortly after said treatment. Thus it is possible for instance to adjust treatment schedule, dosages and type on a patient per patient basis. It is preferred that said sample is obtained within a week of initiation of treatment. More preferably, said sample is obtained within two days of initiation of treatment. With a method of the invention it is possible to evaluate treatment effectiveness almost immediately after initiation of said treatment. A method of the invention thus offers a good opportunity for determining whether treatment adjustments are required.

[0010] A marker gene preferably comprises a gene involved in the generation, maintenance and/or breakdown of blood vessels (angiogenesis). Classes of genes involved in the process of angiogenesis encompass among others receptors, ligands and signaling molecules. Tumor cells are dependent on the growth of new blood vessels to maintain expansion of tumor mass. On the one hand, blood vessels are required to carry nutrients to the site of the tumor, whereas on the other hand waste material needs to be transported from the tumor. In the present invention it has been shown that expression products from genes involved in the generation, maintenance and breakdown of a blood vessel are among the first to respond to anti-tumor treatments. Such genes are therefore very suitable marker genes of the invention. In one embodiment said marker gene comprises a sequence as depicted in table 1 or 2. In another embodiment said marker gene comprises a sequence as depicted in FIGS. 1-18, or a part or analogue thereof. In a preferred embodiment said marker gene comprises a TIE 1 sequence, a Sialoadhesin or Siglec 1 sequence, a sequence as depicted in FIG. 8 or 17, or a part or analogue thereof.

[0011] A change in the level of expression product of a marker gene is indicative for whether a treatment is effective or not. For instance, the level of expression product of a

marker gene can be enhanced in a sample when a treatment is effective, alternatively expression product of a marker gene can be reduced. Thus, preferably, expression product of a marker gene is quantified. The level of expression product in a sample can vary due to changes in the expression of a marker gene. However, it is also possible that the level changes due to a change in type of cells comprising said expression product in said sample, for instance due to treatment related cell death at the site of the body where the sample is obtained. Considering that the level of expression product of marker genes can vary from patient to patient, it is preferred that a method of the invention further comprises comparing the level of expression product of said marker gene with a reference. Preferably said reference comprises the same type of tumor cells prior to, or in the absence of, said treatment. Preferably, said tumor cells are derived from the same patient. The difference in the level of expression product of a marker gene in an effective and a non-effective treatment can be very large. In the extreme cases the level of expression product can range from detectable to not detectable. Marker genes displaying such zero to one relation in expression product levels are preferred in the present invention. A zero to one relation can be used to design relatively simple test systems. A zero to one relation is of course dependent on the detection system used to detect expression product of a marker gene. Very sensitive expression detection systems will typically detect expression product where a less sensitive systems detects no expression product. An expression product can be RNA or a part thereof, transcribed from said marker gene or a translated protein or a part thereof. A person skilled in the art is well capable of designing the most appropriate expression detection system to practice this preferred embodiment of the invention.

[0012] A part of an RNA or DNA molecule is defined herein as an RNA or DNA sequence, comprising at least 50 nucleotides. A part and/or an analogue of an expression product is defined herein as a part and/or analogue that can be detected using essentially the same kind of detection method as said expression product, although the sensibility of detection may differ. An analogue of an RNA or DNA molecule is defined herein as an RNA or DNA sequence which is essentially the same as a particular RNA or DNA sequence. However, a nucleotide mutation, replacement, alteration, addition and/or deletion may have taken place naturally and/or performed artificially, without essentially altering the detection of said analogue as compared with the detection of said particular RNA or DNA sequence. A person skilled in the art is well able to determine whether a given RNA or DNA sequence is an analogue of a particular RNA or DNA sequence, using techniques known in the art.

[0013] In a preferred embodiment said tumor comprises Kaposi's Sarcoma. Kaposi's Sarcoma is a disease of proliferating blood vessels and therefore very much suited for identifying marker genes involved in angiogenesis. According to the invention, changes in angiogenesis factors are among the first marker events as a result of treatment. Kaposi's Sarcoma (KS) manifests itself clinically by reddish skin lesions. Kaposi's Sarcoma is a multicentric, malignant neoplastic vascular proliferation characterized by the development of bluish-red cutaneous nodules, usually on the lower extremities, most often on the toes or feet, and slowly increasing in size and number and spreading to more proximal areas. The tumors have endothelium-lined channels and vascular spaces admixed with variably sized aggregates of

spindle-shaped cells, and often remain confined to the skin and subcutaneous tissue, but widespread visceral involvement may occur. Kaposi's Sarcoma occurs spontaneously in Jewish and Italian males in Europe and the United States. An aggressive variant in young children is endemic in some areas of Africa. A third form occurs in about 0.04% of kidney transplant patients. There is also a high incidence in AIDS patients. (From Dorland, 27th ed & Holland et al., Cancer Medicine, 3d ed, pp2105-7)

[0014] Kaposi's Sarcoma is aggressive in HIV infected individuals. The angiogenic mechanism causing the lesions results from the interplay of viral and cellular gene expression and is poorly understood in terms as to which genes are involved and what controls their expression. The angiogenic proliferation in KS involves mechanisms likely to be universal in angiogenesis. The central role of angiogenesis in Kaposi's Sarcoma is clearly illustrated by the French name for this tumor: angiosarcomatose kaposi. Because of said central role of angiogenesis in Kaposi's Sarcoma, determination of marker genes involved in angiogenesis is very suitable to determine whether a treatment of Kaposi's Sarcoma is effective.

[0015] In the present invention, gene expression patterns of Kaposi's Sarcoma were examined with a method called serial analysis of gene expression (SAGE) Velculescu et al. (1995) Science 270; 484-487). This method allows the quantitative and simultaneous analysis of a large number of transcripts. SAGE is based on two principles. First, a short nucleotide sequence TAG (14 base pairs) contains sufficient information to uniquely identify a transcript, provided it is isolated from a defined position within the transcript. Second, concatenation of short sequence TAG's allows the efficient analysis of transcript in a serial manner by sequencing of multiple TAG's within a single clone.

[0016] Briefly, in this method a biotinylated oligo (dT) primer is used to synthesize cDNA from mRNA, and after digestion with a restriction enzyme, the most 3' terminus (near the poly-A tail) is isolated. These 3' fragments of cDNA are ligated to linkers and cleaved with a type II restriction enzyme to release short sequence (14 bp) of the original cDNA (TAG's). The TAG's are ligated to diTAG's and PCR amplified. These di-TAG's are then ligated to form long concatamers, which are cloned and sequenced. In this way, one sequence reaction yields information about the distribution of many different mRNA's. Finally, the calculation of the abundance of different TAG's and the matching of the TAG's in Genbank are done using the necessary computer software.

[0017] In another aspect the invention provides the use of a nucleic acid comprising a sequence as depicted in FIG. 1-18 and/or table 1 or 2, or a part or analogue thereof, in an expression product detection method. Preferably said nucleic acid comprises a TIE 1 sequence, a Sialoadhesin or Siglec 1 sequence, a sequence as depicted in FIG. 8 or 17, or a part or analogue thereof. Expression of a marker gene in an individual can be detected by determining whether said nucleic acid or part or analogue is able to hybridize with nucleic acid, preferably RNA, in a sample of said individual. If hybridisation takes place, it is indicative of expression of a marker gene in said individual. Of course, as is known by a person skilled in the art, a coding strand of DNA/RNA is capable of hybridizing with the complementary strand of a

corresponding doublestranded nucleic acid sequence. Hence, a complementary strand of a certain coding strand is particularly suitable for detection of expression of said coding strand. For instance, a complementary strand of a coding strand as depicted in **FIG. 1-18** and/or table 1 or 2 is suitable for detection of expression of a gene comprising said coding strand.

[0018] In yet another aspect, the invention provides the use of a proteinaceous molecule capable of specifically binding a protein encoded by a nucleic acid comprising a sequence as depicted in **FIG. 1-18** and/or table 1 or 2, Or a part or analogue thereof, in a detection method. Preferably, said proteinaceous molecule is capable of specifically binding a protein encoded by a nucleic acid comprising a TIE 1 sequence, a Salioadhesin or Siglec 1 sequence, a sequence as depicted in **FIG. 8** or **17**, or a part or analogue thereof. In one embodiment of the invention, said uses are directed toward determining the presence of a site of angiogenesis in an individual. In another embodiment of the invention, said uses are directed toward determining the presence of a tumor cell in an individual. The presence of a tumor cell in an individual can be determined because said tumor cell typically expresses marker genes that can be detected by an expression product detection method. For instance, an antibody, or analogue thereof, specifically directed against an expression product of said marker gene can be generated. Said antibody or analogue is suitable for determination of an expression product of said marker gene in a sample. To determine the presence of a tumor cell in an individual, a sample from said individual can be incubated with said antibody. If said sample contains an expression product of said marker gene, said antibody will bind. Binding can be demonstrated by techniques known in the art, like for instance ELISA. If binding of said antibody is demonstrated, one can conclude that said sample contains an expression product of said marker molecule. The presence of an expression product of said marker molecule can indicate the presence of a tumor cell in an individual, since said marker molecule is expressed by tumor cells. There are of course many more alternative techniques to detect an expression product with use of a proteinaceous binding molecule, which are well known in the art and need no further discussion here. Thus, proteins expressed by a tumor cell can be detected by a proteinaceous molecule capable of specifically binding a protein encoded by a nucleic acid comprising a sequence as depicted in **FIG. 1-18** and/or table 1 or 2, like for instance a TIE 1 sequence, a Salioadhesin and/or Siglec 1 sequence, or a part or analogue thereof. Likewise, the presence of a site of anglogenesis in an individual can be determined by detecting an expression product of a marker gene.

[0019] In another embodiment, a use of a nucleic acid comprising a sequence as depicted in **FIG. 1-18** and/or table 1 or 2, or a use of a proteinaceous molecule capable of specifically binding a protein encoded by a nucleic acid comprising a sequence as depicted in **FIG. 1-18** and/or table 1 or 2, or a part or analogue thereof, are directed toward determining whether a treatment is effective in changing the status of a certain set of target cells in an individual. In a preferred embodiment said nucleic acid comprises a TIE 1 sequence, a Salioadhesin or Siglec 1 sequence, a sequence as depicted in **FIG. 8** or **17**, or a part or analogue thereof In another preferred embodiment said proteinaceous molecule is capable of specifically binding a protein encoded by a

nucleic acid comprising a TIE 1 sequence, a Salioadhesin or Siglec 1 sequence, a sequence as depicted in **FIG. 8** or **17**, or a part or analogue thereof. More preferably, said uses are directed toward determining whether a treatment is effective in counteracting a tumor in an individual. In one embodiment of the invention, said tumor comprises Kaposi's Sarcoma.

[0020] In one aspect the invention provides the use of a nucleic acid comprising a sequence as depicted in table 1 or 2 and/or **FIG. 1-18** as an indicator for angiogenesis. In a preferred embodiment the invention provides the use of a nucleic acid comprising a TIE 1 sequence, a Salioadhesin or Siglec 1 sequence, a sequence as depicted in **FIG. 8** or **17**, or a part or analogue thereof as an indicator for angiogenesis. For instance, a nucleic acid comprising a sequence as depicted in **FIG. 1-18** and/or table 1 or 2 can be used as detection marker for the process of angiogenesis in the course of regenerative treatment. Changes in the expression level of the detection marker indicate active growth of blood vessels (i.e. angiogenesis) as was meant to induce with the regenerative treatment course. In a preferred embodiment such application is in the field of heart and coronary disease aimed at generation of new blood supply to affected organs by means of new blood vessels. Likewise, the treatment of tumors with anti-angiogenesis drugs can be monitored by changes in expression levels of detection marker genes as depicted in **FIG. 1-18** and/or table 1 or 2, such as a TIE 1 sequence, a Salioadhesin and/or Siglec 1 sequence.

[0021] In another aspect the invention provides the use of a nucleic acid comprising a sequence as depicted in **FIG. 1-18** and/or table 1 or 2 as detection marker for tumor cells. In yet another aspect the invention provides the use of a proteinaceous molecule encoded by a nucleic acid comprising a sequence as depicted in **FIG. 1-18** and/or table 1 or 2 or a proteinaceous molecule capable of binding a protein encoded by a nucleic acid comprising a sequence as depicted in **FIG. 1-18** and/or table 1 or 2 as detection marker for tumor cells.

[0022] With a method of the invention it is possible to monitor a specific status of an individual. The presence of a disease—or danger of developing one—can be determined by determining whether or not a sample of an individual comprises an expression product of a marker gene. This means that also the absence of a marker gene in a sample can be indicative for the presence of a disease, or for danger of developing a disease. This is possible for any disease, as long as the disease involves an altered expression pattern of at least one marker gene. Preferably the presence of a marker gene in a sample is determined.

[0023] Additionally, a healing process can be followed as well. For instance, recovery of damaged tissue can involve an increasing amount of expression product of a marker gene over time. It is however also possible that recovery of damaged tissue involves a decreasing amount of expression product of a marker gene over time. Samples taken at different time intervals provide information about the amount of expression product which is generated at different time points. An altered amount of a specific expression product found in samples during a period of time is indicative of the amount of tissue cells generated. Likewise, a decreasing amount of an expression product found in samples in a specific time-period can indicate a certain—

either beneficial or harmful—process. For instance, said process may involve the development or the treatment of disease. An important application is a treatment of heart and coronary disease. A method of the invention is very suitable for monitoring the generation of new cardiac tissue.

[0024] Thus, one aspect of the invention provides a method of diagnosis, in particular a method for determining whether an individual comprises a tumor cell and/or a site of angiogenesis, comprising obtaining a sample from an individual, and determining whether said sample comprises an expression product of at least one marker gene. Preferably, said marker gene comprises a sequence as depicted in table 1 or 2 and/or FIG. 1-18, or a part or analogue thereof. More preferably said marker gene comprises a TIE 1 sequence, a Saliadhesin or Siglec 1 sequence, a sequence as depicted in FIG. 8 or 17, or a part or analogue thereof. With a method of the invention it is possible to detect tumor cells that have released themselves from the tumor and are elsewhere in the body. In a preferred embodiment such detection is performed in the blood of a person detecting circulating tumor cells. These circulating tumor cells can be used for primary identification of presence of a tumor somewhere in the body and also for identification of the risk of metastasis of the tumor to other places in the body next to the primary location of the body. Likewise, a method of the invention is suitable for determining a site of angiogenesis in an individual. Angiogenesis is an indicator for different aspects. For instance, an increased level of angiogenesis indicates the presence of tumor cells, or the healing of damaged tissue, like for instance recovery of heart and coronary disease.

[0025] Since an angiogenic process is now easily monitored by a method of the invention, it is likewise easy to determine whether a certain treatment is effective in altering an angiogenic process. For instance, if a certain treatment is effective in counteracting an angiogenic process, the amount of an expression product of a marker gene involved in angiogenesis decreases as well over time. In the art, many drugs are known for anti-angiogenic treatment. Thus, one embodiment of the invention provides a method for determining whether a treatment is effective in altering an angiogenic process in an individual comprising obtaining a sample from said individual after initiation of said treatment, and determining whether said sample comprises an expression product of at least one marker gene. Preferably, said marker gene comprises a sequence as depicted in table 1, 2, and/or FIGS. 1-18, or a part or analogue thereof. More preferably, said marker gene comprises a TIE 1 sequence, a Saliadhesin or Siglec 1 sequence, a sequence as depicted in FIG. 8 or 17, or a part or analogue thereof. In one embodiment, said treatment comprises counteracting angiogenesis in said individual. In yet another embodiment said treatment involves the use of at least one of the following drugs: 2ME2, Angiostatin, Angiozyme, Anti-VEGF RhuMAB, Apra (CT-2584), Avicine, Benefin, BMS275291, Carboxyamidotriazole, CC4047, CC5013, CC7085, CDC801, CGP-41251 (PKC 412), CM101, Combretastatin A-4 Prodrug, EMD 121974, Endostatin, Flavopiridol, Genistein (GCP), Green Tea Extract, IM-862, ImmTher, Interferon alpha, Interleukin-12, Iressa (ZD1839), Marimastat, Metastat (Col-3), Neovastat, Octreotide, Paclitaxel, Penicillamine, Photofrin, Photopoint, PI-88, Prinomastat (AG-3340), PTK787

(ZK22584), R0317453, Solimastat, Squalamine, SU 101, SU 5416, SU-6668, Suradista (FCE 26644), Suramin (Metaret), Tetrathiomolybdate, Thalidomide, TNP-470, and/or Vitaxin. However, the artisan can think of more drugs that can be used during said treatment.

[0026] In a preferred embodiment, a sample of a method of the invention is a blood sample. Although the location of for instance an angiogenic process can be a tumor or a part of the skin, a blood sample is preferred, among other things because it is much easier to obtain. A blood sample is also often easier to investigate, requiring less expensive and/or specific equipment. Quite surprisingly, we have found that the expression of certain marker genes by hemopoietic cells, like peripheral blood mononuclear cells (PBMC), can be indicative for a process occurring somewhere else in an individual's body. For instance, the presence, or alteration in amount, of an expression product of a marker molecule in PBMC can indicate the presence of a tumor somewhere in the body. In example 10 it is for instance shown that a TIE 1 sequence (tag 15, table 3) or a Saliadhesin or Siglec 1 sequence (tag 82, table 4) are both upregulated in skin tumor and in PBMC cells in a Kaposi's Sarcoma patient. Additionally, example 8 shows that the absence of expression product of a Keratin 14 sequence (tag 7, table 3) in a blood sample of said patient, whereas Keratin 14 is overexpressed in tumor cells, indicates that said sample was not contaminated with tumor cells. Likewise, the expression of certain marker genes by PBMC can provide another diagnostic indication. The presence or absence of an expression product of a marker gene in PBMC provides adequate information about different aspects and/or processes of an individual's body. Preferably, the amount of expression product in a non-hemopoietic cell is compared with a reference value. This way an indication is obtained about an increment or decrement of expression in said hemopoietic cell.

[0027] In one aspect the invention therefore provides a method for determining whether an individual comprises a non-hemopoietic tumor cell and/or a site of angiogenesis, said method comprising determining whether a hemopoietic cell from said patient comprises an altered amount of an expression product of a marker gene as compared with a reference value. Preferably, said marker gene comprises a gene involved in angiogenesis. More preferably, said gene comprises a sequence as depicted in table 1 or 2, and/or FIGS. 1-18, or a part or analogue thereof. Most preferably, said gene comprises a TIE 1 sequence, a Saliadhesin or Siglec 1 sequence, a sequence as depicted in FIG. 8 or 17, or a part or analogue thereof. In one embodiment of the invention said hemopoietic cell comprises a peripheral blood mononuclear cell.

[0028] In one aspect the invention provides a method of the invention, wherein said expression product is expressed by a PBMC. Preferably, said expression product comprises a TIE 1 sequence, a Saliadhesin or Siglec 1 sequence, a sequence as depicted in FIG. 8 or 17, or a part or analogue thereof. As these sequences are involved in angiogenesis, the invention also provides a use of a PBMC expressed Keratin 14 sequence, TIE 1 sequence, Saliadhesin or Siglec 1 sequence, a sequence as depicted in FIG. 2, 8 or 17, or a part or analogue thereof, as an indicator for angiogenesis. Likewise, these sequences are involved in the presence of tumor cells. Therefore, a use of a PBMC expressed Keratin 14 sequence, TIE 1 sequence, Saliadhesin or Siglec 1

sequence, a sequence as depicted in FIG. 2, 8, or 17, or a part or analogue thereof, for determining the presence of a tumor cell in an individual, is also herewith provided. Additionally, the invention also provides an isolated Keratin 14 sequence, TIE 1 sequence, Sialoadhesin or Siglec 1 sequence, a sequence as depicted in FIG. 2, 8, or 17, or a part or analogue thereof, for use in a diagnostic method. A diagnostic method can be carried out using a diagnostic kit. Therefore, a diagnostic kit comprising a nucleic acid comprising a sequence as depicted in FIG. 1-18 and/or table 1 or 2, or a part or analogue thereof, and/or a proteinaceous molecule capable of specifically binding a protein encoded by said nucleic acid or said part or analogue, is also herewith provided. Preferably, said kit comprises a suitable means of detection. In one embodiment a diagnostic kit of the invention is provided comprising a Keratin 14 sequence, and/or a TIE 1 sequence, and/or a Sialoadhesin or Siglec 1 sequence, and/or a sequence as depicted in FIG. 2, 8 or 17, or a part or analogue thereof.

[0029] A diagnostic kit of the invention is particularly useful for carrying out a method of the invention. In yet another embodiment the invention therefore provides a use of a diagnostic kit of the invention for determining whether a treatment is effective in changing the status of a certain set of target cells in an individual and/or altering an angiogenic process in an individual. Additionally, the invention provides a use of a diagnostic kit of the invention for determining whether an individual comprises a tumor cell and/or a site of angiogenesis.

[0030] With a marker gene of the invention it is possible to screen for drugs directed against a disease for which said marker gene is indicative. There are many methods available in the art for screening for a specific drug activity. For instance, cells can be incubated with different potential drug compounds, and an expression pattern of a marker gene in said cells before and after exposure to each potential drug compound can be compared. A specific difference in an expression pattern after exposure to a particular potential drug compound shows that said compound is a suitable candidate for the development of a medicament. The invention therefore provides in one embodiment a use of an expression product of a gene comprising a sequence as depicted in FIG. 1-18, table 1 or table 2 as a drug target. Preferably, said sequence is a Sialoadhesin or Siglec 1, TIE 1, and/or Keratin 14 sequence. A compound capable of altering the activity of Sialoadhesin or Siglec 1, TIE 1, Keratin 14, and/or the expression of Sialoadhesin or Siglec 1, TIE 1, and/or Keratin 14 in a cell is also herewith provided. Said compound is particularly suitable for the preparation of a medicament.

[0031] The present invention is further explained in more detail by the following examples, which do not limit the invention in any way.

## EXAMPLES

### Example 1

[0032] In this example a selection of samples for analysis of expression profiles is made.

[0033] A 31-year old man was demonstrated to be HIV-1 seropositive in February 1997. The initial CD4 cell count was  $25 \times 10^9/l$ . The patient presented within two months a

mucocutaneous Herpes simplex infection and an extrapulmonary Cryptococcosis for which specific medication was given. The HIV-1 RNA load at presentation was 15,000 copies/ml and increased to 33,000 copies/ml in three months. Then antiretroviral therapy was started with zidovudine, lamivudine and indinavir. Immediately after start therapy the HIV-1 RNA load dropped below detection limit. In November 1997 the patient presented with gradual appearance of an increasing number of violaceous skin lesions that clinically resembled Kaposi's Sarcoma. The diagnosis was confirmed by histological examination of one of the lesions. At start of the chemotherapy (bleomycin, vincristine and adriamycin intravenously) KS had progressed to about 150 cutaneous lesions. The interval between the courses of chemotherapy was three weeks and stopped after the fifth course. Several lesions had disappeared by three weeks of therapy and complete remission was gradually reached after one year.

[0034] During chemotherapy several biopsies were taken. The first biopsy was obtained 24 hours after start chemotherapy (named KS1) and the second biopsy after 48 hours (named KS2). All biopsies were flash-frozen in liquid nitrogen immediately after surgical removal and stored at  $-80^\circ C$ . Diagnosis of Kaposi's Sarcoma was confirmed histopathologically.

[0035] Control SAGE libraries KS3 and KS4 were made from frozen material taken at autopsy from two AIDS patients with Kaposi's Sarcoma, both of which died in 1986 without having had any form of chemotherapy or retroviral treatment.

### Example 2

[0036] The expression profiles of the biopsy samples were determined using the SAGE technology. All biopsies were cut with a microtome in 15-20  $\mu m$  sections and transferred to a tube containing TRIzol. RNA isolation with TRIzol was performed according to the manufacturer's instructions. Poly (A) RNA was obtained using the Micro-FastTrack<sup>TM</sup> 2.0 mRNA Isolation Kit. cDNA preparation and the subsequent steps were performed as described by Velculescu. Primary analysis of the sequence results was performed using software especially designed for SAGE by the Bioinformatics Laboratory of the Academic Medical Centre, Amsterdam (van Kampen et al. USAGE: a web-based approach towards the analysis of SAGE data. Bioinformatics, in press). The libraries were also analysed using the Human Transcriptome Map (HTM), a program developed in the AMC, which maps TAG's onto human chromosomes (Caron et al. The Human Transcriptome Map reveals a clustering of highly expressed genes in chromosomal domains. Submitted for publication).

[0037] We sequenced ~47,000 TAG's from the four biopsies, 47,298 TAGS from KS 1 library, 46,671 from the KS 2 library, 49,335 TAG's from the KS3 library, and 48,814 TAG's from the KS4 library. TAG lists (i.e. individual TAG's plus the number of appearance) were compared with each other in USAGE, TAG sequences with the highest counts were identified with the amct2g database available in USAGE (which is an improved TAG identification compared with the SAGEmap database from CGAP (available

from GenBank). Secondly, TAG lists were mapped to chromosome locations with the HTM program, and at the same time compared with specific TAG lists (e.g. vascular endothelium, publicly available), and with a compilation of all TAG lists in the SAGEmap database (designated "All" in HTM) TAG's belonging to genes specifically up regulated in KS3 and KS 4 identified (Table 1). Nucleotide 15 was determined from the original diTAG list in USAGE. The TAG sequence of 15 nt. was checked with GenBank (BLAST) to confirm its identification. A few TAG's were eliminated because of ambiguity in the 15th nucleotide, or because of misidentification.

#### Example 8

[0038] Result of the analysis showing the identifiable TAG's derived from known genes with increased expression in Kaposi's Sarcoma SAGE libraries KS3 and KS4 compared to libraries KS1 and KS2. The TAG numbers are first normalized to a level of 100.000 total TAG's sequenced per library to allow comparison between the libraries and comparison with other libraries in the public domain.

[0039] The sequence catg precedes each TAG sequence given in column 2 of table 1.

TABLE 1

Overview of identifiable TAG's over-expressed in SAGE libraries KS3 and KS4.				
No.	TAG sequence (5' -> 3')	Unigene no.	ID	overexpression factor <sup>1</sup>
1.	ccccagtggc	Hs171596	EphA2	3
2.	cttgacatacc	Hs171695	Dual specificity phosphatase	3
3.	catcacggatc	Hs82112	IL1 receptor, type 1 <sup>2</sup>	10-30
4.	ggccaaaggcc	Hs78436	EphB1	>2
5.	ttgcatatcag	Hs82237	AT group D protein	10-15
6.	ccctgttcage	Hs78824	Tie 1 <sup>2</sup>	2-5
7.	gatcaatcagt	Hs16530	Small ind. cytokine A18	10-20
8.	gagggtgccaa	Hs898	Complement comp. 1Q $\beta$	5-10
9.	taaacctgctg	Hs99923	galectin 7	3-10
10.	gtggccagagg	Hs1420	FGFR3	2-5
11.	tctggcccage	Hs183	DARC (Duffy blood group)	8-10
12.	caggtcgctac	Hs75066	Translin	2-6
13.	gagcagcggcc	Hs112408	Psoriasis (S100 A7)	>20 (specific)
14.	acttatatgc	Hs76152	Decorin	2-10
15.	caggcctggcc	Hs74649	Cytochrome C oxydase subunit VIc	2-4
16.	gtgcggaggac	Hs181062	Serum amyloid A1	5-14
17.	acagcgcaat	Hs74316	Desmoplakin	5-10
18.	gatgtgcacga	Hs117729	Keratin 14	10-14
19.	caggtttcata	Hs24395	Small ind. cytokine, B14 (BRAK)	5-10
20.	aactctgacc	Hs93675	Decidual protein induced by progesterone <sup>2</sup>	3-10

<sup>1</sup>TAG numbers of appearance were normalized to library sizes of 100.000 TAG's.

<sup>2</sup>Identified as Pan Endothelial Markers by St. Croix et al., Genes expressed in human tumor endothelium. Science 289: 1197-1202, 2000.

#### Example 4

[0040] Result of the analysis showing the non-identifiable TAG's derived from EST's of genes with unknown function with increased expression in Kaposi's Sarcoma SAGE libraries KS3 and KS4 compared to libraries KS1 and KS2. The TAG numbers are first normalized to a level of 100.000 total TAG's sequenced per library to allow comparison between the libraries and comparison with other libraries in the public domain.

[0041] The sequence catg precedes each TAG sequence given in column 2 of table 2.

TABLE 2

Overview of identifiable TAG's over-expressed in SAGE libraries KS3 and KS4.				
No.	TAG sequence (5' -> 3')	Unigene no.	ID	overexpression factor <sup>1</sup>
1.	aaatcaataca	Hs94953	EST	4-10
2.	tggtactggc	Hs108741	EST	4-10
3.	tctgcactgag	Hs173789	EST	2-4
4.	caggctgctgg	Hs60440	EST	4-30
5.	atgacagatgg	Hs13775	EST	5-10
6.	gcacaacaaga	Hs236510	EST	3-10
7.	ccacaggagaa	Hs23579	EST	4-10
8.	ctgtcggaac	Hs46987	EST	2-10
9.	gatggctgct	Hs18104	EST	4-20
10.	ctccattgcca	Hs31869	EST	2-10
11.	acctcactgg	Hs112457	EST	Unique <sup>2</sup>

<sup>1</sup>TAG numbers of appearance were normalized to library sizes of 100.000 TAG's

<sup>2</sup>This TAG does not appear in any other SAGE library than our own libraries and seems to be a unique new indicator gene for angiogenesis.

#### Example 5

[0042] Kaposi's sarcoma skin tissue was obtained from the same two AIDS patients mentioned in example 1 from whom SAGE libraries KS3 and KS4 were made. Both patients were homosexual men and were infected at the beginning of the HIV-1 epidemic in Europe. Patient 1 born in Indonesia was demonstrated to be HIV-1 positive in 1982. In February 1985 a histological examination confirmed the diagnosis of Kaposi's sarcoma. He died 13 month later and postmortem examination revealed morphological variants of visceral KS. Patient 2 presented in February 1984 at the

Academic Medical Centre with progressive KS skin lesions. During follow-up the KS progressed to the intestines, oropharynx, lung, tongue, sinus piriformis and lymph nodes. In March 1986 the patient died and autopsy took place. The biopsies of said two patients were named KS3 and KS4.

[0043] Normal adult breast skin tissue was obtained as discarded tissue from reduction mammoplasties (obtained from the department of plastic Surgery of our hospital). RNA isolated of three breast reductions was used to construct the normal skin expression profile library.

[0044] The expression profiles of the biopsy samples were determined using the SAGE technology as described in example 2. We sequenced ~47,000 TAG's from the four biopsies, 49,335 TAG's from the KS3 library, and 48,814 TAG's from the KS4 library. TAG lists (i.e. individual TAG's plus the number of appearance) were compared with each other in USAGE, TAG sequences with the highest

determined from the original diTAG list in USAGE. The TAG sequence of 15 nt. was checked with GenBank (BLAST) to confirm its identification. A few TAG's were eliminated because of ambiguity in the 15th nucleotide, or because of misidentification.

#### Example 6

[0045] Result of the analysis showing the identifiable TAG's derived from known genes with increased expression in Kaposi's Sarcoma SAGE libraries KS3 and KS4 compared to the public libraries of National Center for Biotechnology Information. The TAG numbers are first normalized to a level of 100.000 total TAG's sequenced per library to allow comparison between the libraries and comparison with other libraries in the public domain. The sequence catg precedes each TAG sequence given in column 2 of table 3.

TABLE 3

Overview of identifiable tag's overexpressed in SAGE libraries KS3 and KS4				
Tag number	Tag sequence	Unigene no.	ID	overexpressed
TAG007	gatgtgcacga	Hs117729	Keratin 14	10-14
TAG010	ccccagtcggc	Hs171596	Eph A2 (angiogenesis)	3
TAG011	cttgacatacc	Hs171695	Dual specificity phosphatase	3
TAG012	catcacggatc	Hs82112	IL1 receptor, type 1*	10-30
TAG013	ggccaagggcc	Hs78436	Sorting nexin 17	>2
TAG014	ttgcataatcag	Hs82237	AT group D protein	10-15
TAG015	ccctgttcagc	Hs78824	Tie 1*(angiogenesis)	2-5
TAG016	gatcaatcagt	Hs16530	Small ind. cytokine A18	10-20
TAG017	gagggtgcca	Hs898	Complement comp. 1QB	5-10
TAG018	taaactctctg	Hs99923	galectin 7 (specific)	3-10
TAG019	gtggccaagg	Hs1420	FGFR3 (activated in carcinomas, angiogenesis)	2-5
TAG020	tctggcccagc	Hs183	DARC (Duffy blood group)	8-10
TAG021	cagctcgtctac	Hs75066	Translin(involved in translocations)	2-6
TAG022	gagcagcgccc	Hs112408	Psoriasin (S100 A7)	>20 (specific)
TAG033	acttattatgc	Hs76152	Decorin (connective tissue)	2-10
TAG034	caggcctggcc	Hs288761	Hypothetical protein FLJ21749	2-4
TAG035	gtgcccaggac	Hs181062	Serum amyloid A1	5-14
TAG036	acagcggcaat	Hs74316	Desmoplakin	5-10
TAG037	caggtttcata	Hs24395	Small ind. cytokine, B14 (BRAK)	5-10
TAG038	aactctgaccc	Hs93675	Decidual protein induced by progesterone*	3-10

\*Identified as Pan Endothelial Markers by St. Croix et al., Genes expressed in human tumor endothelium. Science 289: 1197-1202, 2000.

counts were identified with the amct2g database available in USAGE (which is an improved TAG identification compared with the SAGEmap database from CGAP (available from GenBank). Secondly, TAG lists were mapped to chromosome locations with the HTM program, and at the same time compared with specific TAG lists (e.g. vascular endothelium, publicly available), and with a compilation of all TAG lists in the SAGEmap database (designated "All" in HTM) TAG's belonging to genes specifically up regulated in KS3 and KS 4 identified (Table 1). Nucleotide 15 was

#### Example 7

[0046] Result of the analysis showing the non-identifiable TAG's derived from EST's of genes with unknown function with increased expression in Kaposi's Sarcoma SAGE libraries KS3 and KS4 compared to libraries KS1 and KS2. The TAG numbers are first normalized to a level of 100.000 total TAG's sequenced per library to allow comparison between the libraries and comparison with other libraries in the public domain. The sequence catg precedes each TAG sequence given in column 2 of table 4.

TABLE 4.

Overview of identifiable tag's over-expressed in SAGE libraries KS3 and KS4				
Tag number.	Tag sequence (5' > 3')	Unigene no.	ID	overexpressed
TAG023	aaatcaataca	Hs94953	EST	4-10
TAG024	tggttaactggc	Hs108741	EST	4-10
TAG025	tctgcactgag	Hs173789	EST	2-4
TAG026	caggctgctgg	Hs60440	EST	4-30
TAG027	atgacagatgg	Hs13775	EST	5-10
TAG028	gcacaacaaga	Hs236510	EST	3-10
TAG029	ccacaggagaa	Hs23579	EST	4-10
TAG030	ctgtgcggaac	Hs46987	EST	2-10
TAG031	gatggctgcct	Hs18104	EST	4-20
TAG032	ctccattgcca	Hs31869	EST	2-10
TAG004	acctccactgg	Hs112457	EST	Unique*

\*This tag does not appear in any other SAGE library than our own libraries and seems to be unique new indicator gene for angiogenesis.

[0047] The overexpressed TAG's listed in tables 3 and 4 are the same as in tables 1 and 2, respectively. This shows that the expression pattern after treatment is comparable with the expression pattern of healthy individuals with normal expression patterns

#### Example 8

[0048] Using an RT-PCR based method we were able to determine that TAG 11 (table 2) 1 TAG 004 (table 4) indeed represents a differently expressed gene. RNA was isolated from a KS lesion and the first strand cDNA synthesis was primed with an oligo(dT) primer with a 5' M13 tail (5'CTA GTT GTA AAA CGA CGG CCA G-(T)<sub>24</sub>3'). Ten microliter total RNA was used, plus primer and 5  $\mu$ l RT-mix (50 mM Tris, pH 8.3, 75mM KCl, 3 mM MgCl<sub>2</sub>, 10 mM DTT), 80 mM dNTPs and 20 units RNAsin were added, followed by an incubation for 3 minutes at 65° C. and chilled on ice. The RT reaction starts by adding 5 units AMV RT followed by an incubation of 45 minutes at 42° C. For the PCR we used a 19-base TAG-specific primer (which consisted of 11 nt identified in the sage with a 5' NLAIII restriction site and 5 mosine nucleotides to increase the annealing temperature of the primers) and the -21M13 primer. The RT-mix was added to 80  $\mu$ l PCR mixture containing the 100 ng of each primers (-21m13 PRIMER: 5' GTAAAA CGA CGG CCA GT 3' and 5'III IIC ATG ACC TCC ACT GG 3'), 5 mM Tris pH8.3), 20 mM KCl, 0.1 mg BSA per ml, dNTPs (0.1 mM each), 2,4 mM MgCl<sub>2</sub>, and 2 units Taq polymerase. After incubation of 5 minutes at 94° C., the reaction was subjected to 35 cycles of amplification in a thermocycler (9700 Perkin-Elmer). A cycle included denaturation for 1 minute at 95° C., annealing for 1 minute at 55° C. and extension for 2 minutes at 72° C. The last cycle was followed by 72° C. incubation for 10 minutes.

[0049] The amplified fragment was cloned in to an AT plasmid InvitroGen) and subsequently the insert was sequenced using the dye terminator sequencing kit from Applied Biosystems Inc. The fragment appeared to have a length of 102 base pairs and the sequence analysis of the fragment revealed the sequence as depicted in FIG. 1. This sequence was identical to an EST sequence identified from human foetal heart (GenBank acc. #AI217565 and others), which in turn matched a predicted exon on chromosome 19. A relation with angiogenesis has not been described previously and is new

#### Example 9

[0050] Confirmation of the Identity of Tag Sequences

[0051] A sequence consisting of 15 nucleotides should be enough to identify a particular specific mRNA or gene. To confirm the identity of the tags we developed a RT\*PCR using an oligo24dT primer for the RT-reaction. A 5'primer containing the tag sequence itself is used for second strand synthesis. The oligo24dT primer is extended at the 5'site extended with a -21M13 sequence. In the PCR following the RT-reaction -21M13 primer is used for the amplification together with the 5'primer containing the tag sequence. The 5'primer containing the tag sequence is extended with 5 Inosines at its 5'site to enlarge the binding capacity of this primer. The sequence of the amplified fragment can be determined to confirm that this is the gene as identified by the tag sequence. The RT-PCR reactions to confirm the tags were performed on the same tissue samples that were used to prepare the expression profiles of tag sequences.

[0052] Procedure

[0053] Common buffers used throughout the experiments:

[0054] 10x RT buffer:

[0055] 500 mM TRIS, pH 8.3

[0056] 750 mM KCL

[0057] 30 mM MgCl<sub>2</sub>

[0058] 100 mM DTT

[0059] 10x PCR buffer:

[0060] 200 mM TRIS pH 8.3

[0061] 500 mM KCL

[0062] 1 mg/ml BSA

[0063] 1 ml TRIzol reagent (Invitrogen Life Technologies, cat. no. 15596) is added to 10-100 mg tissue or approximately 10<sup>7</sup> cells immediately (tissue is sliced 14  $\mu$ m thick by microtome).

[0064] The Total RNA isolation of the samples is performed according to the manufacturers protocol as follows:

[0065] Add 0.2 ml of Chloroform (Merck) and shake the tube vigorously by hand for 15 seconds.

[0066] Incubate for 5 minutes at RT.

[0067] Centrifuge the sample at no more than 12,000 $\times$ g for 15 minutes at 4° C.

[0068] Transfer 600  $\mu$ l of the colourless upper aqueous layer to a new tube. The lower organic layer should be red.

[0069] Ad 0.5 ml isopropyl alcohol (Merck) and mix.

[0070] Incubate at room temperature for 10 minutes.

[0071] Centrifuge at no more than 12,000 $\times$ g for 15 minutes at 4° C.

[0072] Discard the supernatant and wash the RNA pellet with 1 ml 80% ethanol by vortexing and centrifuge at no more than 7,500 $\times$ g at 4° C. for 5 minutes and discard the supernatant.

[0073] Place the tube at 56° C. for 3 minutes to dry the pellet and proceed with the Poly A<sup>+</sup> RNA isolation as described in the next section. The Poly A<sup>+</sup> mRNA isolation was performed according to the manual of Micro Fastrack® 2.0 Poly A<sup>+</sup>mRNA 2.0 isolation kit as provided by the manufacturer (Invitrogen Corporation, Carlsbad USA; cat no K1520). Subsequently the isolated poly A<sup>+</sup>RNA was used as input for analysis in a RT-PCR reaction. The RT-PCR reactions started with the following mixture of ingredients:

21M13POLYT primer (100 ng/ $\mu$ l)	1.25 $\mu$ l
10 $\times$ RT buffer	2.0 $\mu$ l
100 mM dNTP (Pharmacia)	0.8 $\mu$ l
20 U RNAsin (Roche)	0.3 $\mu$ l
dH <sub>2</sub> O (Baker)	0.65 $\mu$ l

[0074] Add 10  $\mu$ l of Poly A<sup>+</sup> mRNA dilution to 5  $\mu$ l of RT-mix.

[0075] To anneal the primer to the template incubate the reaction mixture at 65° C. for 5 minutes followed by cooling down to room temperature.

[0076] Add 5  $\mu$ l 1U/ $\mu$ l AMV-RT to the reaction mixture and perform the Reverse Transcription by incubating at 42° C. for 45 minutes.

[0077] After the Reverse Transcription immediately incubate the mixture at 95° C. for 5 minutes to stop the reaction. Then let the reaction cool down to room temperature.

[0078] Add 80  $\mu$ l of PCR-mix to each reaction mixture (total volume is 100  $\mu$ l). Prepare the PCR-mix per reaction as follows:

5' primer (100 ng/ $\mu$ l), see table 5	1.0 $\mu$ l
21M13 (100 ng/ $\mu$ l)	1.0 $\mu$ l
10 $\times$ PCR buffer	8.0 $\mu$ l
100 mM MgCl <sub>2</sub>	2.1 $\mu$ l
Amplitaq 5 U/ $\mu$ l (Perkin Elmer)	0.4 $\mu$ l
dH <sub>2</sub> O (Baker)	67.5 $\mu$ l

[0079] PCR amplification was performed in a 9700 DNA thermal cycler (Perkin Elmer) according to the following program.

[0080] 5 minutes 95° C.

[0081] 1 minute 95° C.; 1 minute 55° C.; 2 minutes 72° C., for 35 cycles

[0082] 10 minutes 72° C.

[0083] The TA-cloning of the RT-PCR products was performed according to the manual of TOPO TA Cloning® kit as provided by the manufacturer (Invitrogen Corporation, Carlsbad USA; cat no K4600) using the pCR® II-TOPO® Dual promoter vector and the TOP10 One Shot® Cells.

[0084] Screening of the clones was performed using PCR with SP6 and T7 primers. As follows:

[0085] Resuspend the colony in 50  $\mu$ l dH<sub>2</sub>O (Baker). Add 1  $\mu$ l of this bacteria suspension to 10  $\mu$ l of PCR mixture.

[0086] Prepare the PCR-mixture per reaction as follows:

SP6 (100 ng/ $\mu$ l)	0.10 $\mu$ l
T7 (100 ng/ $\mu$ l)	0.10 $\mu$ l
10 $\times$ PCR buffer	1.00 $\mu$ l
100 mM MgCl <sub>2</sub>	0.20 $\mu$ l
100 mM dNTP's (Pharmacia)	0.08 $\mu$ l
Amplitaq 5 U/ $\mu$ l (Perkin Elmer)	0.04 $\mu$ l
dH <sub>2</sub> O (Baker)	7.48 $\mu$ l

[0087] PCR amplification was performed in a 9700 DNA thermal cycler (Perkin Elmer) according to the following program.

[0088] 5 minutes 95° C.

[0089] 30 seconds 95° C.; 30 seconds 55° C.; 1 minutes 72° C., for 25 cycles 10 minutes 72° C.

[0090] Run 5  $\mu$ l of the Colony-PCR product on a 1.5% agarose 1 $\times$ TBE gel stained with Ethidium Bromide.

[0091] Visualise the amplification products on a UV-illuminator to identify insert-containing clones.

[0092] Clones containing insert were sequenced from both directions using SP6 and T7 primers and the ABI Prism Big-Dye terminator cycle sequencing kit (Perkin Elmer Applied Biosystems, Foster City, Calif. USA).

TABLE 5

Primers used for Tag Confirmation		
Tag name	Sequence	ID
-21M13POLYT	CTA GTT GTA AAA CGA GGG CCA GTT RT-primer TTT TTT TTT TTT TTT TTT TTT T	
TAG007	III IIC ATG GAT GTG CAC G	keratin 14
TAG010	III IIC ATG CCC CAG TCG GC	ephrin A1 (angiogenesis)
TAG011	III IIC ATG CTT GAC ATA CC	dual specificity phosphatase
TAG012	III IIC ATG CAT CAC GGA TC	IL1 receptor, type 1

TABLE 5-continued

Tag name	Primers used for Tag Confirmation	
	Sequence	ID
TAG013	III IIC ATG GGC CAA AGG CC	Sorting Nexin 17 (SNX17)
TAG014	III IIC ATG TTG CAT ATC AG	AT group D protein
TAG015	III IIC ATG CCC TGT TCA GC	Tie 1 (angiogenesis)
TAG016	III IIC ATG GAT CAA TCA GT	small ind. Cytokine A18
TAG017	III IIC ATG GAG GCT GCC AA	complement comp. 1Q beta
TAG018	III IIC ATG TAA ACC TGC TG	galectin 7 (11 nt tag)
TAG019	III IIC ATG GTG GCC AGA GG	FGFRS (11 nt tag)
TAG020	III IIC ATG TCT GGC CCA GC	DARC
TAG021	III IIC ATG CAG GTC GCT AC	Translin
TAG022	III IIC ATG GAG CAG GGC CC	Psoriasin (S100 A7) (11 nt tag)
TAG033	III IIC ATG ACT TAT TAT GC	Decorin
TAG034	III IIC ATG CAG GCC TGG CC	Hypothetical protein FLJ21749
TAG035	III IIC ATG GTG CGG AGG AC	Serum amyloid
TAG036	III IIC ATG ACA GCG GCA AT	Desmoplakin
TAG037	III IIC ATG CAG GTT TCA TA	Small ind. Cytokine, B14 (BRAK)
TAG038	III IIC ATG AAC TCT GAC CC	Decidual protein induced by progesterone
TAG023	III IIC ATG AAA TCA ATA CA	EST Unigene no. Hs94953
TAG024	III IIC ATG TGG TAA CTG GC	EST Unigene no. Hs108741
TAG025	III IIC ATG TCT GCA CTG AG	EST Unigene no. Hs173789
TAG026	III IIC ATG CAG GCT GCT GG	EST Unigene no. Hs60440
TAG027	III IIC ATG ATG ACA GAT GG	EST Unigene no. Hs13775
TAG028	III IIC ATG GCA CAA CAA GA	EST Unigene no. Hs236510
TAG029	III IIC ATG CCA CAG GAG AA	EST Unigene no. Hs23579 (PIG)
TAG030	III IIC ATG CTG TGC GGA AC	EST Unigene no. Hs46987
TAG031	III IIC ATG GAT GGC TGC CT	EST Unigene no. Hs18104
TAG032	III IIC ATG CTC CAT TGC CA	Hs31869 siglec-1 or sialoadhesin
TAG004	III IIC ATG ACC TCC ACT GG	EST Unigene no. Hs1124557

**[0093]** Results of Confirmation of Tag Sequences

**[0094]** Of 18 tag sequences that were analysed with the protocol as described in this example 14 were confirmed with sequences analysis using the RT-PCR with the oligo24dT primer and the tag-based primer as described above (tags designated 004, 007, 011, 012, 014, 015, 016, 017, 022, 025, 029, 030, 032, 036 in table 5). Four tag sequences could not be confirmed based on this method (designated 010, 013, 018, 019). This was probably due to the fact that the tag-based primer was not specific enough or that the polyA tail of the mRNA was not long enough. For one tag an alternative more specific 5'primer was designed

to perform a RT-PCR together with -21M13POLYT (tag designated 004). For tag010 a complete specific primer set was designed to perform as well the Reverse Transcription as the amplification. Other tags were confirmed by using a specific RT-PCR primer set followed by a Nested PCR with a specific nested primer set (tags designated 013, 018, 019). Sequence results of all confirmations are listed below and in the figures. The tag sequence in the mRNA sequence if present is shown in bold fonts.

**[0095]** TAG004 (EST AI217565, Genbank Number BE466728)

[0096] Confirmed with protocol from this example.

CATGACCTCCACTGGAAGAGGGGGCTAGCGTGAGCGCTGATTCTCAACCT  
 ACCATAACTCTTTCTGCTCAGGAAGTCCAATAAAAACATTTTCCATCCA  
 (FIG. 1)

[0097] TAG007 (Keratin 14, Genbank Number NM\_008578)

[0098] Confirmed with protocol from this example.

**CATGGATGTGCACGATGGCAAGGTGGTGTCACCCACGAGCAGGTCTTT**  
 CGCACCAAGAACTGAGGCTGCCAGCCCCGTCAGGCCTAGGAGGCCCC  
 CCGTGTGGACACAGATCCCACTGGAAGATCCCTCTCTGCCCCAAGCACT  
 TCACAGCTGGACCCTGCTTACCCTCACCCCTCCTGGCAATCAATACAG  
 CTTTATTATCTGAGTTGCTAAAAAAAAAAAAAAAAAAAAAAAAAAAA  
 (FIG. 2)

[0099] TAG010 (Ephrin A2, Genbank Number XM\_002088)

[0100] The RT-PCR with the 5'primer designed on the catg-site (the original primer shown in table 5) gave no confirmation. Most probably the tag-based primer is not specific enough. Ephrin A2 specific primers were used for confirmation. The tag sequence is not included in the Ephrin specific RT-PCR.

ATCTACCAGCTCATGATGCAGTGCTGGCAGCAGGAGCGTCCACCGCCC  
 CAAGTTCGCTGACATCGTCAGCATCTGGACAAGCTCATTCTGCCCCCTG  
 ACTCCCTCAAGACCCTGGCTGACTTTGACCCCGCGTGTCTATCCGGCTC  
 CCCAGCACGAGCGGCTCGGAGGGGGTGCCCTTCCGCACGGTGTCCGAGT  
 GGCTGGAGTCCATCAAGATGCAGCAGTATACGGGACCTTC  
 (FIG. 3)

[0101] TAG011 (Dual Specificity Phosphatase, Genbank Number XM\_003720)

[0102] Confirmed with protocol from this example.

**CATGCTTGACATACTACCAGTATTATCCCGACGACATATACATATG**  
 AGAATATACCTTATTTATTTTGTGTAGGTGTCTGCCTTCACAAATGTCAT  
 TGTCTACTCCTAGAAGAACCATACTCAATTTTTGTTTTGAGTACTGT  
 ACTATCCTGTAATATATCTTAAAGCAGGTTGTTTTTCAGCACTGATGGAAA  
 ATACCAGTGTGGGTTTTTTTTTAGTTGCCAACAGTGTATGTTTTGCTGAT  
 TATTTATGACCTGAAATAATATATTTCT-  
 TCTTCTAAGAAGACATTTGTGTAC  
 ATAAGGATGACTTTTTTATACAATGGAATAAATTATGGCATTCTATTG  
 (FIG. 4)

[0103] TAG012 (IL1 Receptor, Type 1, Genbank Number XM\_002686)

[0104] Confirmed with protocol from this example.

**CATGCATCACGGATCAATAGACTGTACTTATTTTCCAATAAAAATTTTCAA**  
 ACTTTGTACTGTT  
 (FIG. 5)

[0105] TAG013 (ephrin B1, genbank numbers XM02585, BC002524)

[0106] The RT-PCR with the 5'primer designed on the catg-site (the original primer shown in table 5) gave no confirmation. Most probably the tag-based primer is not specific enough. Ephrin B1 specific primers were used for confirmation. The tag sequence is included in the 3'primer of the Ephrin B1 specific RT-PCR fragment. The Nested PCR fragment is shown here and is just located upstream of the tag sequence.

AACTTGCCCTGTGCTGTGTCCTCCCATGCTAGGGGCGGAGGGTCTTTTC  
 CTTCTTCTTTCTACCTACCCCTTTTCTCTTGCCAGGGGCTCGTATCCT  
 ACCTTTCTTGTCCCCTGGGCTGGCTGCACAGAGATTGCCCTTCTCTTT  
 TCAGAGCTGGCCCTCGATGCCAAATTAGCATTTAGTATTTTGTCTCAAAGTC  
 TAAGGGACC  
 (FIG. 6)

[0107] TAG014 (AT Group D Protein, Genbank Numbers XM\_006184, AF230388)

[0108] Confirmed with protocol from this example.

**CATGTTGCATATCAGGGTGCTCAAGGATTGGAGAGGAGACAAAACAGG**  
 AGCAGCACAGTGGGGACATCTCCCGTCTCAACAGCCCCAGGCCTATGGGG  
 GCTCTGGAAGGATGGGCCAGCTTGCAGGGGTTGGGGAGGAGACATCCA  
 GCTTGGGCTTTCCCTTTGGAATAAACCATTTGGTCTGTACAAAAAAAAAA  
 AAAAAAAAAAAAAAAAAA  
 (FIG. 7)

[0109] TAG015 (TIE 1, Genbank Number XM-002037)

[0110] Confirmed with protocol from this example.

CATGCCCTGTTTCTAGCTACTCCCACTCCCGGCTGTCTTTCAGAAAAAAT  
 AAATGTTCTAATAAGCTCCAAAAAAAAAAAAAAAAAAAAAAAAAAAA  
 (FIG. 8)

[0111] TAG016 (Small ind. Cytokine A18, Genbank Numbers XM\_008451, Y13710, AF111198)

[0112] Confirmed with protocol from this example.

**CATGGATCAATCAGTGTGATTAGCTTCTCAGCAGACATTGTCCATATG**  
 TATCAAATGACAAATCTTTATGAAATGGTTTTGCTCAGCACCACCTTTTAA  
 TATATTGGCAGTACTTATTATATAAAAGGTAAACCAGCATTCTCAAAAAA  
 AAAAAAAAAA  
 (FIG. 9)

[0113] TAG017 (Complement Comp. 1Q Beta, Genbank Number XM\_010666)

[0114] Confirmed with protocol from this example.

**CATGGAGGGTGCCAA**CAGCATCTTTTCCGGGTCTCTGCTCTTTCCAGAT  
 ATGGAGGCCTGACCTGTGGGTGCTTCACATCCACCCCGGCTCCCCTGC  
 CAGCAACGCTCACTCTACCCCAACACCACCCCTTGCCAGCAATGCAC  
 ACAGTAGGCTTGGTGAATGCTGCTGAGTGAATGAGTAAATAAACCTTC  
 AAGGCC  
 (FIG. 10)

[0115] TAG018 (Galectin 7, Genbank Numbers NM\_002307, U06648)

[0116] The RT-PCR with the 5'primer designed on the catg-site (the original primer shown in table 5) gave no confirmation. Most probably the tag-based primer is not specific enough. Galectin 7 specific primers were used for confirmation. The 5'primer used in the Galectin 7 specific RT-PCR contains the tag sequence. The tag sequence is included in the 5'primer of the Galectin 7 specific RT-PCR fragment. The Nested PCR fragment is shown here and is just located downstream of the tag sequence.

CGGCTGGACACGTCGGAGGTGGTCTTCAACAGCAAGGAGCAAGGCTCCT  
 GGGGCCGCGAGGAGCGCGGGCCGGGCGTTCTTCCAGCGCGGGCAGCC  
 CTTTCGAGGTGCTCATCATCGCGTCAGACGACGGCTTCAAGGCCGTGGTTG  
 GGGACGCCCAGTACCACCACTTCCGCC  
 (FIG. 11)

[0117] TAG019 (FGFR3, Genbank Numbers NM\_022965, NM\_000142)

[0118] The RT-PCR with the 5'primer designed on the catg-site (the original primer shown in table 5) gave no confirmation. Most probably the tag-based primer is not specific enough. FGFR3 specific primers were used for confirmation. The tag sequence is not included in the Ephrin specific RT-PCR. The Nested PCR fragment is shown here and is located upstream of the tag sequence.

CACAACCTCGACTACTACAAGAAGACAACCAACGGCCGGCTGCCCGTAA  
 GTGGATGGCGCCTGAGGCATTATTTGACCGAGTCTACACTCACCAGAGTG  
 ACGTCTGGTCTTTGGGGTCTGCTCTGGGAGATCTTACGCTGGGGGGC  
 TCCCCGTACCCCGGCATCCCTGTGGAGGAGCTCTCAAGCTGCTGAAGGA  
 GGGC  
 (FIG. 12)

[0119] TAG022 (Psoriasis (S100 A7), Genbank Number XM\_048120)

[0120] Confirmed with protocol from this example.

**CATGGAGCAGCGCCTGTT**CCGGGGCAGCCAGTGACCCAGCCCCACC  
 AATGGCCCTCCAGAGACCCAGGAACAATAAATGTCTTCTCCACC  
 (FIG. 13)

[0121] TAG025 EST Unigene no. Hs173789, genbank numbers XM\_018404, AL137262)

[0122] Confirmed with protocol from this example.

**CATGCTGCACTGAGAA**ACTGCATTTTCAGTAGCATTTGTCATCCAGCCG  
 GAAGTTAAAGCACACTTACTTTATTCACCTATTTTATAATAAACGTTCTT  
 GCTGCTGTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA  
 (FIG. 14)

[0123] TAG029 (PIG, Genbank Numbers XM\_011453, AJ251830)

[0124] Confirmed with protocol from this example.

**CATGCCACAGGAGA**AATTCGGG-  
 GATTTGAGTTTCTCTGAATAGCATATAT  
 ATGATGCATCGGATAGGTCATTAT-  
 GATTTTTTACCAITTCGACITACATAA  
 TGAAAACCAAITCAITTTAAATATCA-  
 GATTATATTTTAAAGTTGTGAA  
 AAAGCTAATGTAGTTTTTCAITAT-  
 GAAGTTTTCCCAATAAACCCAGGTATTC  
 TAAACTTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA  
 (FIG. 15)

[0125] TAG030 (EST Unigene no. Hs46987, genbank numbers DG151190, BG057289, BE858276, AV681759, BE503169)

[0126] Confirmed with protocol from this example.

**CATGCTGTGCGGA**ACTGCGTCAGGGCAAATGTACAGCAGGATTTCCCC  
 AACCCAGCTCCATCATCACAGACACAGAGGGTGCAGGGGAGGCCTGCC  
 ACTGTTTTGTCGACTCTGCCCTCCTCTGGCAGCATAGATCCTTAGGTGCTC  
 AATAAAGGTGTGCTGTATTGAAAAAAAAAAAAAAAAAAAAAAAAAAAA  
 (FIG. 16)

[0127] TAG032 (SialoAdhesin, Also Called Siglec 1, Genbank Number XM\_016245)

[0128] Confirmed with protocol from this example.

**CATGCTCCATTGCC**AGACTCTTGCTGGGAGCCCGTCCAGAATGCTCTCC  
 CAATAAACTCCATCCTATGACGCAAAAAAAAAAAAAAAAAAAAAAAAA  
 (FIG. 17)

[0129] TAG036 (Desmoplakin, Genbank Numbers XM\_004463, NM\_004415, AF139065)

[0130] Confirmed with protocol from this example.

**CATGACAGCGGCA**ATCTTTTCTTTGGTCAAAGTTTCTGTTTATTTTGTCT  
 TGTATATTCGATGTACTTTAAGGTGCTTTTATGAAGTTTGTCTATCTGGC  
 AATAAACTTTTAGACTTTAAAAAAAAAAAAAAAAAAAAAAAAAAAA  
 (FIG. 18)

## Example 10

[0131] Determination of the Gene Expression Levels of the Tag Sequences in Skin Samples

[0132] To get a feeling for the use of the tag sequences as markers for angiogenesis process skin samples with (5 different samples) and without (2 control samples) Kaposi's Sarcoma lesions were analysed for the expression level of the genes identified by the tag sequences.

[0133] Procedure

[0134] 1 ml TRIzol reagent (Invitrogen Life Technologies, cat. no. 15596) is added to 10-100 mg tissue or approximately  $10^7$  cells immediately (tissue is sliced  $14\ \mu\text{m}$  thick by microtome). The Total RNA isolation of the samples is performed according to the manufacturers protocol as follows:

[0135] Add 0.2 ml of Chloroform (Merck) and shake the tube vigorously by hand for 15 seconds.

[0136] Incubate for 5 minutes at RT.

[0137] Centrifuge the sample at no more than  $12,000\times g$  for 15 minutes at  $4^\circ\text{C}$ .

[0138] Transfer  $600\ \mu\text{l}$  of the colourless upper aqueous layer to a new tube. The lower organic layer should be red.

[0139] Add 0.5 ml isopropyl alcohol (Merck) and mix.

[0140] Incubate at room temperature for 10 minutes.

[0141] Centrifuge at no more than  $12,000\times g$  for 15 minutes at  $4^\circ\text{C}$ .

[0142] Discard the supernatant and wash the RNA pellet with 1 ml 80% ethanol by vortexing and centrifuge at no more than  $7,500\times g$  at  $4^\circ\text{C}$ . for 5 minutes and discard the supernatant.

[0143] Place the tube at  $56^\circ\text{C}$ . for 3 minutes to dry the pellet and proceed with the DNase treatment as described in the next section. To make sure no genomic DNA exists in the Total RNA isolate we perform a DNase treatment. Protection of RNA against RNase activity of DNase I is done by the addition of RNasin to the DNase reaction. The DNase treatment was performed as follows:

[0144] After TRIzol Total RNA isolation resuspend the pellet in  $88\ \mu\text{l}$   $\text{dH}_2\text{O}$ .

[0145] Add sequentially to the RNA solution:

[0146]  $10\ \mu\text{l}$  10x DNase Buffer (Ambion)

[0147]  $1\ \mu\text{l}$  40U/ $\mu\text{l}$  RNasin (Roche)

[0148]  $1\ \mu\text{l}$  10 U/ $\mu\text{l}$  DNase I (Roche)

[0149] Incubate at  $37^\circ\text{C}$ . for 1 hr.

[0150] Raise sample volume to  $200\ \mu\text{l}$  by adding  $100\ \mu\text{l}$   $\text{dH}_2\text{O}$ .

[0151] Add  $200\ \mu\text{l}$  cold Phenol/Chloroform pH8 (PC8) and mix thoroughly.

[0152] Centrifuge full speed for 5 minutes at room temperature in a microcentrifuge.

[0153] Transfer the aqueous top layer to a new microcentrifuge tube add sequentially:

[0154]  $3\ \mu\text{l}$  glycogen (Roche)

[0155]  $100\ \mu\text{l}$  10 M ammonium acetate

[0156]  $700\ \mu\text{l}$  100% ethanol

[0157] Mix thoroughly and centrifuge at  $25,000\times g$  for 15 minutes at  $4^\circ\text{C}$ . and decant the supernatant.

[0158] Wash twice by adding  $700\ \mu\text{l}$  80 % ethanol. Mix thoroughly and centrifuge at  $25,000\times g$  for 5 minutes at  $4^\circ\text{C}$ . and decant the supernatant.

[0159] Dry the pellet for 5 minutes at  $56^\circ\text{C}$ . and resuspend in  $11\ \mu\text{l}$   $\text{dH}_2\text{O}$ .

[0160] Dilute  $1\ \mu\text{l}$  of the RNA isolate in  $140\ \mu\text{l}$   $\text{dH}_2\text{O}$  and calculate the RNA concentration and yield of the isolate through  $\text{OD}_{260}$  measurement. The yield of DNase treated Total RNA should at least be  $13.5\ \mu\text{g}$  for the determination of the complete TAG expression profile.

[0161] Prepare a solution of the RNA isolate with a concentration of  $50\ \text{ng}/\mu\text{l}$ . This is the starting RNA solution for the series of dilution for the TAG specific RT-PCR.

[0162] Per sample of DNase treated Total RNA 18 TAG specific RT-PCR/Nested PCR reactions are performed in series of dilution of the RNA. For each sample five control RT-PCR's will be performed in series of dilution namely HIV-1 IGAG (SK39/145), GAPDH+RT (in duplo) and no-RT reaction (in duplo). The following dilution scheme was used to derived the samples in serial dilution that are analysed with the RT-PCR:

[0163] 1.  $270\ \mu\text{l}$   $50\ \text{ng}/\mu\text{l}$  DNase treated Total RNA  
DNase treated.  $10\ \mu\text{l}$   $50\ \text{ng}/\mu\text{l}$  solution as input in RT-PCR of each primerset= $500\ \text{ng}$ .

[0164] 2.  $27\ \mu\text{l}$   $50\ \text{ng}/14$  Total RNA 10x diluted= $270\ \mu\text{l}$   
 $5\ \text{ng}/\mu\text{l}$ .  $10\ \mu\text{l}$   $5\ \text{ng}/\mu\text{l}$  solution as input in RT-PCR of each primerset= $50\ \text{ng}$ .

[0165] 3.  $27\ \mu\text{l}$   $5\ \text{ng}/\mu\text{l}$  Total RNA 10x diluted  $270\ \mu\text{l}$   $0.5\ \text{ng}/\mu\text{l}$ .  
 $10\ \mu\text{l}$   $0.5\ \text{ng}/\mu\text{l}$  solution as input in RT-PCR of each primerset= $5\ \text{ng}$ .

[0166] 4.  $27\ \mu\text{l}$   $0.5\ \text{ng}/\mu\text{l}$  Total RNA 10x diluted= $270\ \mu\text{l}$   
 $0.05\ \text{ng}/\mu\text{l}$ .  $10\ \mu\text{l}$   $0.05\ \text{ng}/\mu\text{l}$  solution as input in RT-PCR of each primerset= $0.5\ \text{ng}$ .

[0167] 5.  $27\ \mu\text{l}$   $0.05\ \text{ng}/\mu\text{l}$  Total RNA 10x diluted= $270\ \mu\text{l}$   
 $0.005\ \text{ng}/\mu\text{l}$ .  $10\ \mu\text{l}$   $0.005\ \text{ng}/\mu\text{l}$  solution as input in RT-PCR of each primerset= $0.05\ \text{ng}$ .

[0168] 6.  $27\ \mu\text{l}$   $0.005\ \text{ng}/\mu\text{l}$  Total RNA 10x diluted= $270\ \mu\text{l}$   
 $0.0005\ \text{ng}/\mu\text{l}$ .  $10\ \mu\text{l}$   $0.0005\ \text{ng}/\mu\text{l}$  solution as input in RT-PCR of each primerset= $0.005\ \text{ng}$ .

[0169] 7.  $27\ \mu\text{l}$   $0.0005\ \text{ng}/\mu\text{l}$  Total RNA 10x diluted= $270\ \mu\text{l}$   
 $0.00005\ \text{ng}/\mu\text{l}$ .  $10\ \mu\text{l}$   $0.00005\ \text{ng}/\mu\text{l}$  solution as input in RT-PCR of each primerset= $0.0005\ \text{ng}$ .

[0170] 8.  $27\ \mu\text{l}$   $0.00005\ \text{ng}/\mu\text{l}$  Total RNA 10x diluted= $270\ \mu\text{l}$   
 $0.000005\ \text{ng}/\mu\text{l}$ .  $10\ \mu\text{l}$   $0.000005\ \text{ng}/\mu\text{l}$  solution as input in RT-PCR of each primerset= $0.00005\ \text{ng}$ .

[0171] TAG specific RT-PCR on Total RNA series of dilution using AMV-RT was performed as follows. The reverse Transcription Reactions of all the TAGs on DNase treated Total RNA in series of dilution are performed in 96 wells PCR plates. Ten  $\mu\text{l}$  of each Total RNA dilution is used as input for the RT PCR. The reaction volume of the Reverse Transcription is 20  $\mu\text{l}$  and contains dNTP's,  $\text{MgCl}_2$  and RNasin.

[0172] Prepare the RT-mix per reaction as follows:

3' primer (100 ng/ $\mu\text{l}$ ) see table 6	1.25 $\mu\text{l}$
10 $\times$ RT buffer	2.0 $\mu\text{l}$
100 mM dNTP (Pharmacia)	0.8 $\mu\text{l}$
20 U RNasin (Roche)	0.3 $\mu\text{l}$
$\text{dH}_2\text{O}$ (Baker)	0.65 $\mu\text{l}$

[0173] Add 10  $\mu\text{l}$  of Total RNA dilution to 5  $\mu\text{l}$  of RT-mix.

[0174] To anneal the primer to the template incubate the reaction mixture at 65° C. for 5 minutes followed by cooling down to room temperature.

[0175] Add 5  $\mu\text{l}$  1U/ $\mu\text{l}$  AMV-RT to the reaction mixture and perform the Reverse Transcription by incubating at 42° C. for 45 minutes.

[0176] After the Reverse Transcription immediately incubate mixture at 95° C. for 5 minutes to stop the reaction. Then let the reaction cool down to room temperature.

[0177] Add 80  $\mu\text{l}$  of PCR-mix to each reaction mixture (total volume is 100  $\mu\text{l}$ ). Prepare the PCR-mix per reaction as follows:

5' primer (100 ng/ $\mu\text{l}$ ) see table 6	1.0 $\mu\text{l}$
10 $\times$ PCR buffer	8.0 $\mu\text{l}$
100 mM $\text{MgCl}_2$	1.9 $\mu\text{l}$
Amplitaq 5 U/ $\mu\text{l}$	0.4 $\mu\text{l}$
$\text{dH}_2\text{O}$ (Baker)	68.7 $\mu\text{l}$

[0178] PCR amplification is performed in a 9700 DNA thermal cycler (Perkin Elmer) according to the following program.

[0179] 5 minutes 95 ° C.

[0180] 1 minute 95° C.; 1 minute 55° C.; 2 minutes 72° C., for 35 cycles 10 minutes 72° C.

[0181] Subsequent to this first round of amplification a second round, nested, amplification was performed. TAG specific second round nested PCR on RT-PCR product was performed as follows:

[0182] Add 5  $\mu\text{l}$  of the TAG RT-PCR product in 45  $\mu\text{l}$  of the Nested-PCR mix.

[0183] Prepare the Nested-PCR mix per reaction as follows:

5' nested primer (100 ng/ $\mu\text{l}$ ), see table 7	0.5 $\mu\text{l}$
3' nested primer (100 ng/ $\mu\text{l}$ ), see table 7	0.5 $\mu\text{l}$
10 $\times$ PCR buffer	5.0 $\mu\text{l}$
100 mM $\text{MgCl}_2$	1.25 $\mu\text{l}$
100 mM dNTP (Pharmacia)	0.4 $\mu\text{l}$
Amplitaq 5 U/ $\mu\text{l}$	0.2 $\mu\text{l}$
$\text{dH}_2\text{O}$ (Baker)	37.15 $\mu\text{l}$

[0184] The combination of primers used for amplification of the genes identified by the tags and the length of the amplified fragment is given in table 8.

[0185] PCR amplification is performed in a 9700 DNA thermal cycler (Perkin Elmer) according to the following program.

[0186] 5 minutes 95° C.

[0187] 1 minute 95° C.; 1 minute 55° C.; 2 minutes 72° C., for 25 cycles

[0188] 10 minutes 72° C.

[0189] Run 10  $\mu\text{l}$  of the Nested-PCR product on a 1.5% agarose 1xTBE gel stained with Ethidium Bromide.

[0190] Visualization of the TAG amplified fragments in the dilution series on a UV-illuminator reveals the level of expression by determining the highest dilution still giving a positive signal.

TABLE 6

RT-PCR primer design for first round amplification.

Primer	Sequence
5' TAG004GENE	GGC CTT TAA CAC CCC GTT CCT
3' TAG004GENE	TGG TAG GTT GAG AAT CAG CGC TCA
5' TAG007GENE-N	AGG AGA CCA AAG GTC GCT ACT GCA
3' TAG007GENE	CAG TTC TTG GTG CGA AGG ACC T
5' TAG010GENE	ATC TAC CAG CTC ATG ATG CAG TGC T
3' TAG010GENE	GAA GTG CTC CGT ATA CTG CTG CAT
5' TAG011GENE	AGT GGG TAC ATC AAG TCC ATC TGA
3' TAG011GENE	CAC TGG TAT TTT CCA TCA GTG CT
5' TAG012GENE	TAA AGT TGT CCT GCT TGA GCT GGA
3' TAG012GENE	GGC ACG TGA GCC TCT CTT TGC AGT
5' TAG013GENE	CTC TAC CCC AGA GGA ATT TAC AGA
3' TAG013GENE	GGG CCA GAC CAA ACA CAG ACC TCT
5' TAG014GENE	GGC AAC AAG CAG AAG GCG GTC A
3' TAG014GENE	TGA TCT TGA GCT GCA GCT GCT CCT
5' TAG015GENE	GAA TGT GCT GGT CGG AGA GAA
3' TAG015GENE	TGG GGC AGC TTT TCA TAG AGC T
5' TAG016GENE	TTC TCT GCC TGC CCA GCA TCA TGA
3' TAG016GENE	TCA GGC ATT CAG CTT CAG GTC GGT
5' TAG017GENE	GTC TCT ACT ACT TCA CCT ACC A
3' TAG017GENE	TGT TGG GGG TAG AGT GAG CGT TGC T
5' TAG018GENE	GCA GGT TCC ATG TAA ACC TGC TGT
3' TAG018GENE	CTG CTC AGA AGA TCC TCA CGG AGT
5' TAG019GENE	GTG ACC GAG GAC AAC GTG ATG AAG A
3' TAG019GENE	CAT GAT CAT GTA CAG GTC GTG TGT
5' TAG022GENE	TGA GCA ACA CTC AAG CTG AGA G
3' TAG022GENE	TCT CTG GAG GCC CAT TGG T
5' TAG025GENE	ATG GGG TCA GGA ACA TCT GGC AGA
3' TAG025GENE	TCC GGC TGG ATG ACA AAT GCT ACT
5' TAG029GENE	CTC AGG TTT ATC TGG GCT CTA TCA
3' TAG029GENE	TCA TAA TGA CCT ATC CGA TGC AT
5' TAG030GENE	CCT GCA AAG ATA GGA GAG GCT CCA

TABLE 6-continued

RT-PCR primer design for first round amplification.

Primer	Sequence
3'TAG030GENE	ATT GAG CAC CTA AGG ATC TAT GCT
5'TAG032GENE	TGC GAA TCA GGG ACC AAC AGG AGA
3'TAG032GENE	TTG GGA GGA CAT TCT GGA CGG GCT
5'TAG036GENE	ATT TAG CAG TAG TTC TAT TGG GCA
3'TAG036GENE	ACT GAT TAG CAC TTC AGA CGC ACT

[0191]

TABLE 7

Nested-PCR primer design.

Primer	Sequence
5'TAG004GENE-2	CAT CGA CAA ATT GCG ATC T
3'TAG004GENE-2	CGC TAG CCC CCT CTT CCA GT
5'TAG007GENE-2.1	AGG AGA TGA TTG GCA GCG T
3'TAG007GENE-2	GGA GGA GGT CAC ATC TCT GGA T
5'TAG010GENE-2	CCA AGT TCG CTG ACA TCG T
3'TAG010GENE-2	TGC TGG GGA GCC GGA TAG ACA
5'TAG011GENE-2	GAA GAG AAA GGA CTC AGT GT
3'TAG011GENE-2	AGA TAT ATT TAC ACG ATA GT
5'TAG012GENE-2	AAA TCC AAG ACT ATG AGA
3'TAG012GENE-2	CTT AGT GGC TGG TGA CAG T
5'TAG013GENE-2	AAC TTG CCC TGT GCC TGT GT
3'TAG013GENE-2	GGT CCC TTA GAC TTT GAG CA
5'TAG014GENE-2	CTT CTG CGA GCT GCA TCT CA
3'TAG014GENE-2	TGC AGT GAC AGC TCC GTC T
5'TAG015GENE-2	AGA GGA GGT TTA TGT GAA GA
3'TAG015GENE-2	ACT ATC TCC CAA AGA AGG ACT
5'TAG016GENE-2	TGT CCT CGT CTG CAC CAT
3'TAG016GENE-2	ATG TAT TTC TGG ACC CAC T
5'TAG017GENE-2	GTC ACC TTC TGT GAC TAT GCC T
3'TAG017GENE-2	ACA GGT CAG GCC TCC ATA TCT
5'TAG018GENE-2	CGG CTG GAC ACG TCG GA
3'TAG018GENE-2	GGC GGA AGT GGT GGT ACT
5'TAG019GENE-2	CAC AAC CTC GAC TAC TAC A
3'TAG019GENE-2	GCC CTC CTT CAG CAG CTT
5'TAG022GENE-2	TTC ACA AAT ACA CCA GAC GTG AT
3'TAG022GENE-2	GGG CGC TGC TCC ATG GCT CTG CT
5'TAG025GENE-2	TGC CTA GAA AGG GGT GGC T
3'TAG025GENE-2	TTC TCA GTG CAG ACA TGT GGC T
5'TAG029GENE-2	CAG GCT TCT GAT AGT TTG CAA CT
3'TAG029GENE-2	TAT GCT ATT CAG AGA AAC T
5'TAG030GENE-2	TCT AAT GCA TGT AGA AGC T
3'TAG030GENE-2	AGG GCA GAG TCG ACA AAA CAG T
5'TAG032GENE-2	TCT TGA GTG GGC TAG TGA CT
3'TAG032GENE-2	AGT CTG GCA ATG GAG CAT GA
5'TAG036GENE-2	TGC TAT ACC TTG ACT TCA T
3'TAG036GENE-2	TCC AAG TGT ACT GCT TAT
5'TAG036GENE-2.1	CTA GTA GTC AGT TGG GAG T
3'TAG036GENE-2.1	AGC CAG AAC AGC CTT TAC T

[0192]

TABLE 8

Primer combinations and amplified fragment length in the Tag specific Nested PCR reactions

Name	Sequence	Primer	PCR fragment
TAG004	5'TAG004gene-2	3'TAG004gene-2	182
TAG007	5'TAG007gene-2.1	3'TAG007gene-2	211
TAG010	5'TAG010gene-2	3'TAG010gene-2	108
TAG011	5'TAG011gene-2	3'TAG011gene-2	239
TAG012	5'TAG012gene-2	3'TAG012gene-2	197

TABLE 8-continued

Primer combinations and amplified fragment length in the Tag specific Nested PCR reactions

Name	Sequence	Primer	PCR fragment
TAG013	5'TAG013gene-2	3'TAG013gene-2	212
TAG014	5'TAG014gene-2	3'TAG014gene-2	243
TAG015	5'TAG015gene-2	3'TAG015gene-2	131
TAG016	5'TAG016gene-2	3'TAG016gene-2	219
TAG017	5'TAG017gene-2	3'TAG017gene-2	185
TAG018	5'TAG018gene-2	3'TAG018gene-2	175
TAG019	5'TAG019gene-2	3'TAG019gene-2	204
TAG022	5'TAG022gene-2	3'TAG022gene-2	238
TAG025	5'TAG025gene-2	3'TAG025gene-2	183
TAG029	5'TAG029gene-2	3'TAG029gene-2	141
TAG030	5'TAG030gene-2	3'TAG030gene-2	179
TAG032	5'TAG032gene-2	3'TAG032gene-2	223
TAG036	5'TAG036gene-2	3'TAG036gene-2	191

[0193] Results

[0194] The results of determination of the expression levels of the genes identified by the tag sequences are depicted in FIG. 19. The data clearly indicate that a number of the genes identified by the tag sequences have a higher expression in the skin samples with Kaposi's Sarcoma lesions compared to normal skin: tag007, tag010, tag012, tag013, tag014, tag015, tag016, tag017, tag022, tag029, tag030, tag032 and tag036.

Example 11

[0195] Determination of the Gene Expression Levels of the Tag Sequences in Peripheral Blood Mononuclear Cell (PBMC) Samples

[0196] To get a feeling for the use of the tag sequences as markers for angiogenesis process in samples not from the location of the angiogenesis process, i.e. the Kaposi's Sarcoma in the skin or another tumour in the body but at an accessible sample from the blood (PBMC's) the expression of 5 tag identified genes was determined in PBMC samples. The PBMC samples were from the blood of patients with (4 different samples) and without (2 control samples) Kaposi's Sarcoma lesions and were analysed for the expression level of 5 tag identified genes in PBMC's.

[0197] The procedure of the example was identical as described in example 10, with the exception that PBMC samples were used (approximately 10 million cells per sample, ranging from 2.5 to 50 million) instead of skin biopsies. The genes that were analysed are identified by tag007, tag017, tag010, tag013, tag015, tag029 and tag032. The results of the analysis are depicted in FIG. 20. It is clear from these data that the elevated expression in the tumour sites of the genes identified by tag015 (TIE 1) and tag032 (Sialoadhesin or Siglec 1) is paralleled in the blood cell fraction, i.e. PBMC.

[0198] Results

[0199] The data clearly show that the over expression of the gene identified by tag007 (Keratin 14) in skin samples (see example 5) is not paralleled in blood. In contrast, in the samples tested in this example the gene identified by tag007

is not expressed at all in the blood compartment. This shows that no tumour cells expressing tag007 are present in blood. As a consequence, measurement of tag007 in the blood could be a good indicator for the presence of these tumour cells in the blood, and thus a marker for circulating cancer cells that can cause metastasis.

[0200] The up regulation of genes identified by tag015 (TIE 1) and tag032 (Saliadhesin or Siglec 1) in skin samples is clearly paralleled in the blood. These two tags are higher expressed in blood from patients with tumours compared to healthy individuals. This up-regulation is due to up-regulation of expression in typical blood cells. The up-regulation cannot be due to the presence of tumour cells that express tag015 and tag032 in the blood, because the absence of expression of the gene identified by tag007 in the blood shows that no tumour cells are present in blood, as explained above. This means that measurement of expression of tag015 and/or tag032 in the blood indicates the presence of a tumour somewhere else in the body. Furthermore, measurement of expression of genes identified by tag015 and/or tag032 during anti-tumour therapy and/or anti-angiogenesis therapy can be used to monitor the efficacy of this treatment.

[0201] Conclusions

[0202] The paralleled up-regulation of the genes identified with tag015 (TIE 1) and tag032 (Saliadhesin or Siglec 1) in both the tumour and the blood, enables the monitoring of the efficacy of a therapy aimed at decreasing the growth of a tumour, in particular anti-angiogenic tumour treatment. This follows the reasoning that if these two genes are markers in PBMC for blood vessel formation in a tumour in another site in the body, these two markers in blood will also decrease with the decrease of this blood vessel growth at the tumour site.

[0203] The genes identified by tag007 (Keratin 14), tag015 (TIE 1) and tag032 (Saliadhesin or Siglec 1) have different expression in the blood of patients with a tumour compared to normal individuals. Therefore these genes, in particular the expression thereof, can be used to screen a population at risk for the presence of tumour in individual members of that population. All the genes identified by tags in this study that have changed expression levels comparing normal to tumour tissue, i.e. tag007, tag001, tag012, tag013, tag014, tag015, tag016, tag017, tag022, tag029, tag030, tag032 and tag036 are encoding potential target molecules for therapeutic compound with anti-angiogenic effects applicable in tumour treatment, and/or these genes encode potential target molecules that can be potential target molecules for therapeutic compounds that stimulate the growth of blood vessel, for instance in the treatment of heart and coronary disease.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0204] FIG. 1: Sequence involved in angiogenesis. A change of expression of this sequence after a certain treatment indicates that said treatment is effective. This sequence is identical to an EST sequence identified from human foetal heart (GenBank acc. #AI217565 and others), which in turn matches a predicted exon on chromosome 19. A relation with angiogenesis has not been described previously.

[0205] FIG. 2-18: Sequences which are identified by name and Genbank numbers (NCBI database). Other identification can be found in tables 1-4 (Unigene numbers) that can be found in the SAGE databases of NCBI.

[0206] FIG. 19. Mean expression levels and standard deviation depicted as log dilution factor that still is positive starting with 500 ng of total RNA isolated from 5 skin samples with Kaposi's Sarcoma (light bars) and 2 control, normal skin samples (dark bars).

[0207] FIG. 20. Mean expression levels and standard deviation depicted as log dilution factor that still is positive starting with 500 ng of total RNA isolated from 4 PBMC samples of patients with Kaposi's Sarcoma (light bars) and 2 control, normal PBMC samples (dark bars).

What is claimed is:

1. A method of determining whether a treatment is effective in changing a status of a certain set of target cells in an individual comprising: obtaining a sample from said individual after initiating said treatment; and determining whether said sample comprises an expression product of at least one marker gene.

2. The method according to claim 1, wherein said target cells comprise a tumor cell.

3. The method according to claim 1 or 2, wherein said sample comprises at least one of said target cells.

4. The method according to any one of claims 1-3, wherein said sample is obtained within one week of initiating said treatment.

5. The method according to any one of claims 1-4, wherein said sample is obtained within two days of initiating said treatment.

6. The method according to any one of claims 1-5, wherein said marker gene comprises a gene involved in the generation, maintenance and/or breakdown of blood vessels.

7. The method according to any one of claims 1-6, wherein said marker gene comprises a sequence as depicted in Table 1 or Table 2.

8. The method according to any one of claims 1-7, wherein said marker gene comprises a sequence selected from the group consisting of a sequence depicted in FIG. 1 through 18 or a part or analogue thereof.

9. The method according to any one of claims 1-8, wherein expression of said marker gene is quantified.

10. The method according to any one of claims 1-9, further comprising comparing expression of said marker gene with a reference value.

11. The method according to any one of claims 2-10, wherein said tumor comprises Kaposi's Sarcoma.

12. A method detecting an expression product of a marker gene comprising: obtaining a sample from an individual;

introducing a nucleic acid to said sample, said nucleic acid selected from the group consisting of a sequence as depicted in FIG. 1-18, a sequence as depicted in Table 1 and a sequence as depicted in Table 2, or a part or analogue thereof to said sample; and

determining whether said nucleic acid hybridizes in said sample.

**13.** A method of detecting an expression product of a marker gene comprising:

incubating a proteinaceous molecule to a sample from an individual, said proteinaceous molecule capable of specifically binding a protein encoded by a nucleic acid selected from the group consisting of a sequence as depicted in **FIG. 1-18**, a sequence as depicted in Table 1 and a sequence as depicted in Table 2, or a part or analogue thereof; and

detecting binding between said proteinaceous molecule and said protein.

**14.** The method according to claim 12 or claim 13, further comprising determining the presence of a tumor cell in an individual.

**15.** The method according to claim 12 or claim 13, further comprising determining the presence of a site of angiogenesis in an individual.

**16.** The method according to claim 12 or claim 13, further comprising determining whether a treatment is effective in changing the status of a certain set of target cells in an individual.

**17.** The method according to any one of claim **12-16**, further comprising determining whether a treatment is effective in counteracting a tumor in said individual.

**18.** The method according to claim 14 or **17**, wherein said tumor comprises Kaposi's Sarcoma.

**19.** A method for determining whether an individual possesses a tumor cell and/or a site of angiogenesis, comprising:

obtaining a sample from said individual; and

determining whether said sample comprises an expression product of at least one marker gene.

**20.** The method according to claim 19, wherein said marker gene comprises a sequence selected from the group consisting of a sequence as depicted in **FIG. 1-18**, a sequence as depicted in Table 1, a sequence as depicted in Table 2, or a part or analogue thereof.

**21.** A method of determining whether an individual possesses a non-hemopoietic tumor cell and/or a site of angiogenesis, said method comprising determining whether a hemopoietic cell from said individual comprises an altered amount of an expression product of a marker gene as compared with a reference value.

**22.** The method according to claim 21, wherein said marker gene comprises a gene involved in angiogenesis.

**23.** The method according to claim 21 or **22**, wherein said gene comprises a sequence selected from the group consisting of a sequence as depicted in **FIG. 1-18**, a sequence as depicted in Table 1, a sequence as depicted in Table 2, or a part or analogue thereof.

**24.** The method according to any one of claims **21-23**, wherein said hemopoietic cell comprises a peripheral blood mononuclear cell.

**25.** A method of determining whether a treatment is effective in altering an angiogenic process in an individual comprising:

obtaining a first sample from said individual before initiating said treatment;

obtaining a second sample from said individual after initiating said treatment; and

comparing expression of an expression product of at least one marker gene in said first sample and said second sample.

**26.** The method according to claim 25, wherein said treatment comprises counteracting angiogenesis in said individual.

**27.** The method according to claim 25 or **26**, wherein said marker gene comprises a sequence selected from the group consisting of a sequence as depicted in **FIG. 1-18**, a sequence as depicted in Table 1, a sequence as depicted in Table 2, or a part or analogue thereof.

**28.** The method according to any one of claims **25-27**, wherein said treatment involves the use of at least one drug selected from the group consisting of 2ME2, Angiostatin, Angiozyme, Anti-VEGF RhuMab, Apra (CT-2584), Avicine, Benefin, BMS275291, Carboxyamidotriazole, CC44047, CC5013, CC7085, CDC801, CGP-41251 (PKC 412), CM101, Combretastatin A-4 Prodrug, EMD 121974, Endostatin, Flavopiridol, Genistein (GCP), Green Tea Extract, IM-862, ImmTher, Interferon alpha, Interleukin-12, Iressa (ZD 1839), Marimastat, Metastat (Col-3), Neovastat, Octreotide, Paclitaxel, Penicillamine, Photofrin, Photopoint, PI-88, Prinomastat (AG-3340), PTK787 (ZK22584), R0317453, Solimastat, Squalamine, SU 101, SU 5416, SU-6668, Suradista (FCE 26644), Suramin (Metaret), Tetrathiomolybdate, Thalidomide, TNP-470, and Vitaxin.

**29.** The method according to any one of claims **1-11**, **19-20**, or **23-26**, wherein said sample is a blood sample.

**30.** The method according to any one of claims **1-11**, **19-20**, or **24-27**, wherein said sample comprises a peripheral blood mononuclear cell.

**31.** The method according to any one of claims **1-11**, or **19-30**, wherein said expression product comprises one of a TIE 1 sequence, a Salioadhesion or Siglec 1 sequence, a sequence as depicted in **FIG. 8** or **FIG. 17**, or a part of analogue thereof.

**32.** A method of detecting angiogenesis comprising detecting peripheral blood mononuclear cell expression of at least one of Keratin 14 sequence, TIE 1 sequence, a Salioadhesion or Siglec 1 sequence, a sequence as depicted in **FIG. 2**, **FIG. 8** or **FIG. 17**, or a part or analogue thereof.

**33.** A method of determining the presence of a tumor cell in an individual comprising:

obtaining a sample from said individual; and

detecting the level of peripheral blood mononuclear cell expression of at least one of a Keratin 14 sequence, TIE 1 sequence, a Salioadhesion or Siglec 1 sequence, a sequence as depicted in **FIG. 2**, **FIG. 8** or **FIG. 17**, or an analogue thereof.

**34.** A method of diagnosing presence of disease comprising comparing expression of an isolated Keratin 14 sequence, TIE 1 sequence, a Salioadhesion or Siglec 1 sequence, a sequence as depicted in **FIG. 2**, **FIG. 8** or **FIG. 17**, or an analogue thereof in an individual to a reference value.

**35.** A diagnostic kit comprising a nucleic acid comprising a sequence selected from the group consisting of a sequence as depicted in **FIGS. 1-18**, Table 1, Table 2, or a part or analogue thereof, and a proteinaceous molecule capable of specifically binding a protein encoded by said nucleic acid or said part or analogue thereof.

**36.** The diagnostic kit according to claim 35, further comprising at least one of a Keratin 14 sequence, a TIE 1 sequence, a Salioadhesion or Siglec 1 sequence, a sequence as depicted in **FIG. 2**, **FIG. 8** or **FIG. 17**, or an analogue thereof.

**37.** A method of determining whether a treatment is effective in changing the status of a certain set of target cells in an individual and/or altering an angiogenic process in an individual, said method comprising:

providing the diagnostic kit according to claim 35 or **36**;

obtaining a sample from said individual; and

detecting the presence of an expression product of at least one marker gene in said sample.

**38.** A method of determining whether an individual possesses a tumor cell and/or a site of angiogenesis, said method comprising:

providing the diagnostic kit according to claim 35 or **36**;

obtaining a sample from said individual; and

quantifying an expression product of at least one marker gene in said sample.

**39.** A method for identifying desired drug activity comprising: determining an expression pattern of a marker gene in cells;

incubating said cells with an expression product of a gene comprising a sequence as depicted in **FIG. 1-18**, Table 1 or Table 2; and

detecting an alteration in said expression pattern of said marker gene after said incubating.

**40.** A compound capable of altering the activity of at least one of Salioadhesion or Siglec 1, TIE 1, Keratin 14, and the expression of at least one of Salioadhesion or Siglec 1, TIE 1 and Keratin 14 in a cell.

**41.** A method of preparing a medicament comprising:

identifying a compound capable of altering the activity of at least one of Salioadhesion or Siglec 1, TIE 1, Keratin 14, and the expression of at least one of Salioadhesion or Siglec 1, TIE 1 and Keratin 14 in a cell; and

incorporating said identified compound into a medicament.

\* \* \* \* \*

专利名称(译)	用于治疗评估的手段和方法		
公开(公告)号	<a href="#">US20030170720A1</a>	公开(公告)日	2003-09-11
申请号	US10/055728	申请日	2002-01-23
[标]申请(专利权)人(译)	范德KUYL ANTOINETTE CORNELIA CORNELISSEN MARION		
申请(专利权)人(译)	范德KUYL ANTOINETTE CORNELIA CORNELISSEN MARION		
当前申请(专利权)人(译)	范德KUYL ANTOINETTE CORNELIA CORNELISSEN MARION		
[标]发明人	VAN DER KUYL ANTOINETTE CORNELIA CORNELISSEN MARION		
发明人	VAN DER KUYL, ANTOINETTE CORNELIA CORNELISSEN, MARION		
IPC分类号	C12Q1/68 G01N33/53		
CPC分类号	C12Q2600/158 C12Q1/6886		
优先权	2001020373 2001-09-28 EP 2001200228 2001-01-23 EP 60/325722 2001-09-28 US		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

本发明提供了一种用于确定治疗是否有效改变患者中某组靶细胞(例如肿瘤)的状态的方法。该方法意味着在开始治疗后从患者获得样品,并确定所述样品是否包含至少一种标记基因的表达产物。优选地,所述样品是血液样品。在一个方面,所述表达产物由外周血单核细胞表达。所述标记基因可以是参与血管生成,维持和/或分解(血管生成)的基因。本发明的方法非常适于在几天内确定针对卡波西氏肉瘤的某种治疗是否成功。此外,该方法适用于确定患者中血管生成和/或肿瘤细胞的存在。

Figure 1

